

GROTHENDIECK RINGS OF QUEER LIE SUPERALGEBRAS

SHIFRA REIF

ABSTRACT. We determine the Grothendieck rings of the category of finite-dimensional modules over Queer Lie superalgebras via their rings of characters. In particular, we show that the ring of characters of the Queer Lie supergroup $Q(n)$ is isomorphic to the ring of Laurent polynomials in x_1, \dots, x_n for which the evaluation $x_{n-1} = -x_n = t$ is independent of t . We thus complete the description of Grothendieck rings for all classical Lie superalgebras.

1. INTRODUCTION

The Grothendieck group is an invariant of an abelian category defined to be the span of all objects, modulo the relation $[A] + [C] = [B]$ for every exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$. When the category is a tensor category, the Grothendieck group inherits a ring structure via $[A] \cdot [B] = [A \otimes B]$. For finite-dimensional modules over semisimple Lie algebras, the Grothendieck ring is isomorphic to the ring of characters due to the highest weight description of simple modules, c.f. e.g. [SV, Prop. 4.2]. For Lie superalgebras, the ring of characters is the quotient of the Grothendieck ring by the relation identifying a module with its parity shift.

In this paper, we determine the ring of characters for queer Lie superalgebras. These algebras are one of the strange series in Kac's classification of simple Lie superalgebras [K], and is the remaining classical series for which the Grothendieck ring is described. For the queer Lie supergroup $Q(n)$, the description takes the following form.

Theorem 1. *The ring of characters of $Q(n)$ is isomorphic to the ring*

$$(1.1) \quad J_n := \left\{ f \in \mathbb{Q}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]^{S_n} : f|_{x_1 = -x_2 = t} \text{ is independent of } t \right\}.$$

This theorem is a natural extension of the description of Grothendieck rings for the other classical Lie algebras and superalgebras: The ring of S_n -invariant Laurent polynomials is the ring of characters of the Lie algebra $\mathfrak{gl}(n)$. Sergeev and Veselov showed in [SV] that such an evaluation condition characterizes the ring of characters for basic-classical Lie superalgebras. In [IRS], Im, Serganova and the author determine the Grothendieck ring for periplectic Lie superalgebras.

For basic-classical Lie superalgebras, the ring of characters is isomorphic to the ring of supercharacters. For the queer Lie superalgebra this is not the case. In fact, Cheng showed that the supercharacter of a finite-dimensional nontrivial simple $\mathfrak{q}(n)$ -module is always zero [Che].

To prove that all characters satisfy invariance conditions, we use restriction to low-rank subalgebras. To show the converse, namely that every invariant function is a linear combination of characters, we use a basis of the ring of characters which consists of Euler characteristic of certain cohomologies introduced by Penkov and Serganova [PS]. We first prove the result for the queer Lie supergroup $Q(n)$, and then extend it to the other queer Lie supergroups

and carry it over to Lie superalgebras. We also give a description of J_n in terms of invariance under the Weyl groupoid.

Acknowledgments. The author is thankful to Maria Gorelik and Vera Serganova for interesting conversations. This project is partially supported by ISF Grant No. 1221/17.

2. THE QUEER LIE SUPERALGEBRA AND ITS FINITE-DIMENSIONAL REPRESENTATIONS

We recall the structure and representation theory of the queer Lie superalgebras. One of their main difficulty is that they do not possess an invariant bilinear form, moreover their Cartan subalgebras are not purely even.

2.1. Preliminaries. The family of queer Lie superalgebras consist of: the subalgebra $\mathfrak{q}(n)$ of $\mathfrak{gl}(n|n)$ consisting of matrices of the form $\begin{bmatrix} A & B \\ B & A \end{bmatrix}$ where A, B are $n \times n$ matrices,

$$\mathfrak{sq}(n) := \left\{ \begin{bmatrix} A & B \\ B & A \end{bmatrix} \in \mathfrak{q}(n) : \text{tr} B = 0 \right\},$$

$\mathfrak{pq}(n) := \mathfrak{q}(n)/\mathbb{C}I$ and $\mathfrak{psq}(n) := \mathfrak{sq}(n)/\mathbb{C}I$. The even parts of $\mathfrak{q}(n)$ and $\mathfrak{sq}(n)$ are isomorphic to $\mathfrak{gl}(n)$, whereas the even parts of $\mathfrak{pq}(n)$ and $\mathfrak{psq}(n)$ are isomorphic to $\mathfrak{pgl}(n)$.

The Cartan subalgebra $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathfrak{h}_1$ of $\mathfrak{q}(n)$ and $\mathfrak{sq}(n)$ consists of matrices of the form $\begin{bmatrix} A & B \\ B & A \end{bmatrix}$ where A and B are diagonal $n \times n$ matrices. For $i = 1, \dots, n$, let $H_i = E_{ii} + E_{n+i, n+i}$. Let $\varepsilon_1, \dots, \varepsilon_n$ denote the basis of \mathfrak{h}_0^* dual to H_1, \dots, H_n . For $\mathfrak{pq}(n)$ and $\mathfrak{psq}(n)$, \mathfrak{h}_0^* and $\varepsilon_1, \dots, \varepsilon_n$ denote their images under the natural projection.

The set of roots is $\Phi = \{\varepsilon_i - \varepsilon_j : i \neq j\}$ where for each root there is an odd root vector and an even root vector. We choose the standard set of positive roots:

$$\Phi_0^+ = \Phi_1^+ = \{\varepsilon_i - \varepsilon_j : i < j\}.$$

The Weyl group W is isomorphic to S_n . We denote the denominators by

$$R_{\bar{0}} := \prod_{\alpha \in \Phi_0^+} (1 - e^{-\alpha}), \quad R_{\bar{1}} := \prod_{\alpha \in \Phi_1^+} (1 + e^{-\alpha}), \quad R = \frac{R_{\bar{0}}}{R_{\bar{1}}}.$$

Let $\rho_0 := (n-1)\varepsilon_1 + (n-2)\varepsilon_2 + \dots + \varepsilon_{n-1}$. Note that $e^{\rho_0} R_{\bar{0}}$ is W -anti-invariant and $e^{\rho_0} R_{\bar{1}}$ is W -invariant.

2.2. Finite-dimensional modules and their characters. We briefly recall the construction of highest weight $\mathfrak{q}(n)$ -modules, see [CW, Sec. 1.5.4] for more details. Given $\lambda \in \mathfrak{h}_0^*$, one defines a symmetric bilinear form on \mathfrak{h}_1 by $\langle v, w \rangle_\lambda := \lambda([v, w])$. Let $\mathfrak{h}'_1 \subseteq \mathfrak{h}_1$ be a maximal isotropic subspace with respect to $\langle \cdot, \cdot \rangle_\lambda$ and define $\mathfrak{h}' = \mathfrak{h}_0 \oplus \mathfrak{h}'_1$. The one dimensional \mathfrak{h}_0 -module $\mathbb{C}v_\lambda$, defined by $hv_\lambda = \lambda(h)v_\lambda$, extends to an \mathfrak{h}' -module by letting $\mathfrak{h}'_1.v_\lambda = 0$. Define the induced \mathfrak{h} -module

$$W_\lambda := \text{Ind}_{\mathfrak{h}'}^{\mathfrak{h}} \mathbb{C}v_\lambda.$$

A simple finite-dimensional $\mathfrak{q}(n)$ -module $L(\lambda)$ has W_λ as its highest weight space. Moreover, a highest-weight module $L(\lambda)$ is finite-dimensional if and only if $\lambda = \sum_{i=1}^n \lambda_i \varepsilon_i$ where $\lambda_i - \lambda_{i+1} \in \mathbb{Z}_+$ for every $i = 1, \dots, n-1$, and $\lambda_i = \lambda_{i+1}$ implies $\lambda_i = 0$ ([Pe, Thm. 4] and [CW, Thm 2.18]). We denote the set of these highest weights by $\tilde{\Lambda}_n$. Any finite-dimensional

simple $\mathfrak{q}(n)$ -module is a highest weight module. A $\mathfrak{q}(n)$ -module can be integrated to the Lie supergroup $Q(n)$ if and only if $\lambda \in \tilde{\Lambda}_n \cap \mathbb{Z}_n$. Let $\Lambda_n := \tilde{\Lambda}_n \cap \mathbb{Z}_n$.

Given a finite-dimensional $\mathfrak{q}(n)$ -module M with weight space decomposition $M = \bigoplus_{\lambda \in \mathfrak{h}_0} M_\lambda$ where

$$M_\lambda = \{v \in M \mid hv = \lambda(h)v \text{ for all } h \in \mathfrak{h}_0\},$$

we define the character of M to be

$$\text{ch}M := \sum_{\lambda \in \mathfrak{h}_0} (\dim M_\lambda) e^\lambda.$$

For a Lie superalgebra \mathfrak{g} , we denote by $J(\mathfrak{g})$ the ring of all characters of finite-dimensional modules. We shall use the notation $x_i := e^{\varepsilon_i}$, $i = 1, \dots, n$.

An element in the ring of characters is also called a *virtual character*. Note that the ring of characters is isomorphic to the quotient of Grothendieck ring by the ideal generated by $[M] - [\Pi M]$ where Π is the parity shift functor.

2.3. The Euler characteristic $E(\lambda)$. Let $\lambda \in \Lambda_n$. Note that W_λ is a module over the Cartan subgroup H that can be extended naturally to the Borel subgroup B corresponding to Φ^+ . We have the higher cohomology modules

$$H^i(\lambda) := H^i(G/P, \mathcal{L}(W_\lambda)).$$

Here P is the largest parabolic subgroup of G to which the B -module W_λ can be lifted and $\mathcal{L}(W_\lambda)$ denote the G -equivariant $\mathcal{O}_{G/P}$ -module induced by the P -module W_λ . Since $H^i(\lambda)$ is finite dimensional and is zero for large enough i , the Euler characteristic

$$E(\lambda) := \sum_{i \geq 0} (-1)^i [H^i(\lambda)]$$

belongs to $J(\mathfrak{q}(n))$.

One has

$$E(\lambda) = 2^{\lfloor \frac{l(\lambda)}{2} \rfloor} R^{-1} \cdot \sum_{w \in W} (-1)^{l(w)} w \left(\frac{e^\lambda}{\prod_{\beta \in \Phi^+(\lambda)} (1 + e^{-\beta})} \right).$$

Recall the definition of Schur's P -function p_λ for $\lambda \in \Lambda_n$:

$$(2.1) \quad p_{\lambda,n} = \sum_{w \in S_n/S_\lambda} w \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right)$$

where S_n/S_λ denotes the set of minimal length coset representatives for the stabilizer of λ in S_n . Then by [PS, Prop. 1], $E(\lambda)$ is a scalar multiple of p_λ . By [B, Cor. 4.26], $\{E(\lambda) \mid \lambda \in \Lambda_n\}$ forms a basis to the character ring.

3. THE RING OF CHARACTERS FOR QUEER LIE SUPERGROUPS.

The following lemma shows that the ring of characters of $Q(n)$ and $SQ(n)$ are contained in J_n (see (1.1)). The proof is similar to [SV, Prop. 4.3] except one replaces supercharacters by characters.

Lemma 2. *Let M be a module over $Q(n)$ or $SQ(n)$ then $\text{ch}M$ is in J_n .*

Proof. The S_n -symmetry follows from restricting the representation of f to the even part which is either $GL(n)$ or $SL(n)$. For the evaluation property, restrict M to $\mathfrak{g}_{12} := \mathfrak{h}_0 \oplus \mathfrak{g}_{\bar{1}, \varepsilon_1 - \varepsilon_2} \oplus \mathfrak{g}_{\bar{1}, \varepsilon_2 - \varepsilon_1}$. Then M has composition factors of the form $e^\lambda + e^{\lambda - (\varepsilon_1 - \varepsilon_2)}$ (typical case) or e^μ where $(\mu, \varepsilon_1 + \varepsilon_2) = 0$ (atypical case). The former is equal to $x_1^{\lambda_1} \cdots x_n^{\lambda_n} \left(1 + \frac{x_2}{x_1}\right)$ and the evaluation $x_1 = -x_2 = t$ equals zero. The latter is of the form $e^{\mu_1(\varepsilon_1 + \varepsilon_2) + \mu_3\varepsilon_3 + \cdots + \mu_n\varepsilon_n}$ and the evaluation gives equals to $-e^{\mu_3\varepsilon_3 + \cdots + \mu_n\varepsilon_n}$. \square

Note that since an element $f \in J_n$ is a W -invariant, the evaluation $f|_{x_{n-1} = -x_n = t}$ is also independent of t .

Definition 3. The evaluation map $\text{ev} : J_n \rightarrow J_{n-2}$ is defined by $\text{ev}(f) = f|_{x_{n-1} = -x_n}$.

3.1. The kernel of the evaluation map. The following proposition is a generalization of the fact that the kernel of the evaluation map of $GL(m|n)$ is spanned by characters of Kac modules [HR, Thm. 17].

Proposition 4. *The kernel of the evaluation map is in $J(Q(n))$.*

Proof. Suppose that $\text{ev}(f) = 0$ for $f \in J_n$. Then f is divisible by $x_{n-1} + x_n$ and by the W -invariance, also divisible by $\prod_{i \neq j} (x_i + x_j) = \prod_{i \neq j} (e^{\varepsilon_i} + e^{\varepsilon_j})$. So

$$f = \prod_{i \neq j} (e^{\varepsilon_i} + e^{\varepsilon_j}) \cdot g.$$

Since f and $\prod_{i \neq j} (e^{\varepsilon_i} + e^{\varepsilon_j})$ are W -invariant, so is g and by theory of symmetric functions, $g = \sum^{\text{finite}} a_\lambda s_\lambda$ where $a_\lambda \in \mathbb{Q}$ and s_λ is the Schur Laurent polynomial

$$s_\lambda = e^{-\rho_0} R_0^{-1} \cdot \sum_{w \in W} (-1)^{l(w)} w(e^{\lambda + \rho_0})$$

where $\lambda = (\lambda_1, \dots, \lambda_n)$ and $\lambda_1 \geq \dots \geq \lambda_n$. Now

$$\begin{aligned} f &= \sum_{\lambda}^{\text{finite}} a_\lambda \prod_{i \neq j} (e^{\varepsilon_i} + e^{\varepsilon_j}) e^{-\rho_0} R_0^{-1} \cdot \sum_{w \in W} (-1)^{l(w)} w(e^{\lambda + \rho_0}) \\ &= \sum_{\lambda}^{\text{finite}} a_\lambda \prod_{i \neq j} (1 + e^{-(\varepsilon_i - \varepsilon_j)}) R_0^{-1} \cdot \sum_{w \in W} (-1)^{l(w)} w(e^{\lambda + \rho_0}) \\ &= \sum_{\lambda}^{\text{finite}} a_\lambda E(\lambda + \rho_0). \end{aligned}$$

Note that the last equality uses the fact that $\lambda + \rho_0$ has a trivial stabilizer in W . Since $\lambda + \rho_0 \in \Lambda_n$, f is a linear combination of virtual characters and the assertion follows. \square

3.2. Surjectivity of the evaluation map.

Proposition 5. *The evaluation map $\text{ev} : J(Q(n)) \rightarrow J(Q(n-2))$ is surjective.*

To prove the proposition, we give for every $p_{\lambda, n-2} \in J(Q(n-2))$ a Schur function $p_{\lambda, n}$ such that $\text{ev}(p_{\lambda, n}) = p_{\lambda, n-2}$. For the convenience of the reader, we first present the argument on an example.

Example 6. Suppose $\overset{\circ}{\lambda} = (3, 1)$. Let $\lambda = (3, 1, 0, 0)$. Let us show that $\text{ev}(p_{\lambda,4}) = p_{\overset{\circ}{\lambda},2}$. One has

$$x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} = x_1^3 x_2 \frac{1 + x_1^{-1} x_2}{1 - x_1^{-1} x_2} \cdot \frac{1 + x_1^{-1} x_3}{1 - x_1^{-1} x_3} \cdot \frac{1 + x_1^{-1} x_4}{1 - x_1^{-1} x_4} \cdot \frac{1 + x_2^{-1} x_3}{1 - x_2^{-1} x_3} \cdot \frac{1 + x_2^{-1} x_4}{1 - x_2^{-1} x_4}.$$

Now, if $w \in S_4/S_\lambda$ is such that the terms $1 + x_3^{-1} x_4$ or $1 + x_4^{-1} x_3$ appear, then the evaluation of $w \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right)$ is zero. This means that for the evaluation to be nonzero $w^{-1}(\{3, 4\}) = \{3, 4\}$, which in this example means that $w \in S_2 := \{1, (1\ 2)\}$. Then

$$\begin{aligned} \text{ev}(p_{\lambda,4}) &= \text{ev} \sum_{w \in S_4/S_\lambda} w \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) \\ &= \text{ev} \sum_{w \in S_2} w \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) \\ &= \sum_{w \in S_2} w \left(\text{ev} \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) \right) \\ &= \sum_{w \in S_2} w \left(\text{ev} \left(x_1^3 x_2 \frac{1 + x_1^{-1} x_2}{1 - x_1^{-1} x_2} \cdot \frac{1 + x_1^{-1} x_3}{1 - x_1^{-1} x_3} \cdot \frac{1 + x_1^{-1} x_4}{1 - x_1^{-1} x_4} \cdot \frac{1 + x_2^{-1} x_3}{1 - x_2^{-1} x_3} \cdot \frac{1 + x_2^{-1} x_4}{1 - x_2^{-1} x_4} \right) \right) \\ &= \sum_{w \in S_2} w \left(\text{ev} \left(x_1^3 x_2 \frac{1 + x_1^{-1} x_2}{1 - x_1^{-1} x_2} \cdot \frac{1 + x_1^{-1} t}{1 - x_1^{-1} t} \cdot \frac{1 - x_1^{-1} t}{1 + x_1^{-1} t} \cdot \frac{1 + x_2^{-1} t}{1 - x_2^{-1} t} \cdot \frac{1 - x_2^{-1} t}{1 + x_2^{-1} t} \right) \right) \\ &= p_{\overset{\circ}{\lambda},2}. \end{aligned}$$

Proof of Proposition 5. Let $\overset{\circ}{\lambda} \in \mathbb{Z}_+^{n-2}$ and extend it to $\lambda \in \mathbb{Z}_+^n$ by adding zeros, namely $\{\overset{\circ}{\lambda}_1, \dots, \overset{\circ}{\lambda}_{n-2}\} \cup \{0\} = \{\lambda_1, \dots, \lambda_n\}$. We show that $\text{ev}(p_\lambda) = p_{\overset{\circ}{\lambda}}$. Since Schur P -functions span $J(Q(n-2))$, this will imply surjectivity.

Let $k \in \{1, \dots, n\}$ be such that $\lambda_k = \lambda_{k+1} = 0$ (such k may not be unique). Denote $\sigma_k = (k\ n-1)(k+1\ n) \in S_n$ and let ev_k be a map on J_n such that $\text{ev}_k(f) = f|_{x_k = -x_{k+1}}$. Then by symmetry $\text{ev} = \sigma_k \circ \text{ev}_k$. Now, if $w \in S_n/S_\lambda$ is such that $w^{-1}(\{k, k+1\}) \subseteq \{\{i, j\} \mid \lambda_i \neq \lambda_j\}$ then

$$\text{ev}_k \left(w \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) \right) = 0.$$

Denote by A the set of $w \in S_n/S_\lambda$ for which $w^{-1}(\{k, k+1\}) \subseteq \{i, j \mid \lambda_i = \lambda_j\}$. By the definition of Λ^n , $\lambda_i = \lambda_j$ only if $\lambda_i = 0$. Then, up to S_λ , $w^{-1}(\{k, k+1\}) = \{k, k+1\}$. Hence, w preserves $\{k, k+1\}$ and consequently w preserves $\{1, \dots, n\} / \{k, k+1\}$. In particular, w

commutes with ev_k and A is the permutation group of $\{1, \dots, n\} / \{k, k+1\}$. Thus

$$\begin{aligned}
\text{ev}(p_\lambda) &= \sigma_k \circ \text{ev}_k(p_{\lambda, n}) \\
&= \sigma_k \circ \text{ev}_k \sum_{w \in S_n / S_\lambda} w \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) \\
&= \sigma_k \circ \text{ev}_k \sum_{w \in A} w \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) \\
&= \sigma_k \sum_{w \in A} w \left(\text{ev}_k \left(x^\lambda \prod_{\substack{1 \leq i < j \leq n \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) \right) \\
&= \sigma_k \sum_{w \in A} w \left(\text{ev}_k \left(x^\lambda \prod_{\substack{i < j, i, j \notin \{k, k+1\} \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \cdot \prod_{\substack{i < k \\ \lambda_i \neq 0}} \frac{1 + x_i^{-1} x_k}{1 - x_i^{-1} x_k} \cdot \frac{1 + x_i^{-1} x_{k+1}}{1 - x_i^{-1} x_{k+1}} \cdot \prod_{\substack{i > k \\ \lambda_i \neq 0}} \frac{1 + x_k^{-1} x_i}{1 - x_k^{-1} x_i} \cdot \frac{1 + x_{k+1}^{-1} x_i}{1 - x_{k+1}^{-1} x_i} \right) \right) \\
&= \sigma_k \sum_{w \in A} w \left(x^{\hat{\lambda}} \prod_{\substack{i < j, i, j \notin \{k, k+1\} \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) \\
&= \sum_{w \in S_{n-2} / S_{\hat{\lambda}}} w \left(x^{\hat{\lambda}} \prod_{\substack{1 \leq i < j \leq n-2 \\ \lambda_i > \lambda_j}} \frac{1 + x_i^{-1} x_j}{1 - x_i^{-1} x_j} \right) = p_{\hat{\lambda}, n-2}
\end{aligned}$$

as required. \square

3.3. The ring of characters. We can now prove the main theorem.

Proof of Theorem 1. We prove the theorem by induction on n . For $n = 0$, $J_n = \mathbb{C}$ by definition. For $n = 1$, $Q(1)$ admits a 1-dimensional representation whose character is a scalar multiple of $e^{m\epsilon_1}$ for every $m \in \mathbb{Z}$. Then $J(Q(1)) = \mathbb{C}[x_1^{\pm 1}] = J_1$.

Suppose that $J(Q(n-2)) = J_{n-2}$. Recall the evaluation map $\text{ev} : J_n \rightarrow J_{n-2}$ and that $J(Q(n)) \subseteq J_n$. By Proposition 5, $\text{ev}(J(Q(n-2))) = J(Q(n)) = J_{n-2}$. So, any element in J_n is a sum of an element from $J(Q(n))$ and an element in the kernel of ev . By Proposition 4, the kernel is also contained in $J(Q(n))$ and the assertion is obtained. \square

Corollary 7. *The ring of characters of $SQ(n)$ is also isomorphic to J_n . Indeed, since $SQ(n)$ is a subgroup of $Q(n)$, every $Q(n)$ -module restricts to $SQ(n)$ and so $J(SQ(n)) \supseteq J(Q(n)) = J_n$. On the other hand, by Lemma 2, $J(SQ(n)) \subseteq J_n$ and we obtained the equality. In fact, the restriction of simple $Q(n)$ -modules to $SQ(n)$ is known by [GS, Cor. 4.6].*

Corollary 8. *The rings of characters of $PQ(n)$ and $PSQ(n)$ are equal to*

$$J_n \cap \text{span} \left\{ x_1^{\lambda_1} \cdots x_n^{\lambda_n} \mid \sum_{i=1}^n \lambda_i = 0 \right\}.$$

Indeed, a module over $Q(n)$ (resp. $SQ(n)$) lifts to a module over $PQ(n)$ (resp. $PSQ(n)$) if and only if the identity matrix acts by zero. This holds if and only if all the weights λ of

the module satisfy that $\sum_{i=1}^n \lambda_i = 0$. When the module is simple, this is true if and only if one weight λ of the module satisfies that $\sum_{i=1}^n \lambda_i = 0$.

Remark 9. The category of finite-dimensional $Q(n)$ -modules admits a subcategory of polynomial representations, namely those which appear as composition factors in the tensor algebra of the natural representations. The Grothendieck of this subcategory is the set of all polynomials in J_n . This ring is isomorphic to the center of $U(\mathfrak{q}(n))$ and was described in [Pa, Thm. 2.11] (see also [CW, A.3.4]).

4. THE RING OF CHARACTERS FOR QUEER LIE SUPERALGEBRAS

We first describe the ring of characters for the category of finite-dimensional modules whose weights half-integer. The simple modules in this category were studied in [CK].

Proposition 10. *Let J_n be as in (1.1) and $J_{\frac{1}{2},n} = (\text{span} \{x_1^{k_1} \cdots x_n^{k_n} \mid k_1, \dots, k_n \in \frac{1}{2} + \mathbb{Z}\})^{S_n}$. In particular, the ring of characters of the category of finite-dimensional half-integer weights is isomorphic to*

$$J_n \oplus \prod_{1 \leq i < j \leq n} (x_i + x_j) J_{\frac{1}{2},n}.$$

Proof. We need to show that the character of any module M of a half-integer weights is divisible by $(x_i + x_j)$ for every $i \neq j$. Note that $\alpha = \varepsilon_i - \varepsilon_j$ is also an even root. Restrict M to the corresponding $\mathfrak{sl}(2)$ -triple \mathfrak{s}_α . Since the weights are not integers, zero is not a weight and the restriction of \mathfrak{s}_α is a direct sum of $\mathfrak{sl}(2)$ -strings of even length. In particular, as an \mathfrak{h}_0 -module, $M = \bigoplus_{\lambda \in B} N_\lambda$ where B is a finite subset of \mathfrak{h}_0 and $\text{ch} N_\lambda = e^\lambda + e^{\lambda - \alpha} = x^\lambda + x^\lambda x_i^{-1} x_j = (x_i + x_j) x^\lambda x_i^{-1}$. \square

Similarly to $J_{\frac{1}{2},n}$, we can define the vector space

$$J_{a,n} := (\text{span} \{x_1^{k_1} \cdots x_n^{k_n} \mid k_1, \dots, k_n \in a + \mathbb{Z}\})^{S_n}$$

for any $a \in \mathbb{C}/\mathbb{Z}$.

Proposition 11. *The ring of characters of finite-dimensional $\mathfrak{q}(n)$ -modules is isomorphic to*

$$J_n \oplus \bigoplus_{a \in \mathbb{C}/\mathbb{Z}} \prod_{1 \leq i < j \leq n} (x_i + x_j) J_{a,n}.$$

Proof. Let $L(\lambda)$ be a simple highest weight module for which $\lambda \in a + \mathbb{Z}^n$. Suppose $a \neq \frac{1}{2}$, then, $\lambda_i \neq -\lambda_j$ for every $i \neq j$ and $L(\lambda)$ is a typical module. By, [Pe, Thm. 2] (see also [CW, Thm. 2.5.2.]),

$$\text{ch} L(\lambda) = 2^{\frac{l(\lambda) + \delta(\lambda)}{2}} R \cdot \sum_{w \in W} (-1)^{l(w)} e^{w(\lambda)}.$$

In particular $\text{ch} L(\lambda)$ is divisible by $e^{\rho_0} \prod_{i < j} (1 + e^{-\varepsilon_i + \varepsilon_j}) = \prod_{i < j} (x_i + x_j)$.

For $a = \frac{1}{2}$, the characters are divisible by $\prod_{i < j} (x_i + x_j)$ by the previous proposition and the assertion follows. \square

Remark 12. Note that $\mathfrak{q}(n)$ and $\mathfrak{sq}(n)$ admit the same rings of characters by a similar argument as in Corollary 7.

Corollary 13. *Similarly to Corollary 8,*

$$J(\mathfrak{pq}(n)) = J(\mathfrak{q}(n)) \cap \text{span} \left\{ x^\lambda \mid \lambda \in \mathbb{Z}^n, \sum_{i=1}^n \lambda_i = 0 \right\}$$

$$J(\mathfrak{spq}(n)) = J(\mathfrak{sq}(n)) \cap \text{span} \left\{ x^\lambda \mid \lambda \in \mathbb{Z}^n, \sum_{i=1}^n \lambda_i = 0 \right\}.$$

5. THE WEYL GROUPOID FOR $\mathfrak{q}(n)$.

We describe the polynomial invariants of certain affine action of the super Weyl groupoid \mathfrak{W} of $\mathfrak{q}(n)$. We follow [SV, Sec. 9] (for the periplectic case, see [IRS, Sec. 5.4]).

We identify \mathfrak{h}_0^* with its dual using the scalar product in which $(\varepsilon_i, \varepsilon_j) = \delta_{ij}$. Let \mathfrak{T} be a groupoid with the set of objects $\{[\varepsilon_i + \varepsilon_j], [-\varepsilon_i - \varepsilon_j] : i < j\}$ and the set of morphisms $\text{Mor}_{\mathfrak{T}}([\alpha], [\beta]) = \emptyset$ if $\alpha \neq \pm\beta$, and $\text{Mor}_{\mathfrak{T}}([\alpha], [\pm\alpha])$ is of cardinality 1. For every object $[\alpha]$, we denote by r_α the unique morphism in $\text{Mor}_{\mathfrak{T}}([\alpha], [-\alpha])$, and impose the relation $r_\alpha r_{-\alpha} = \text{id}_{[-\alpha]}$.

The Weyl group $W = S_n$ acts on $\Delta(\mathfrak{g}_0)$ and we have the naturally defined homomorphism Φ from W to the group of autoequivalences of \mathfrak{T} . This yields the semidirect product groupoid $\tilde{\mathfrak{T}} = W \ltimes \mathfrak{T}$. The objects of $\tilde{\mathfrak{T}}$ are the same as the objects of \mathfrak{T} . The morphisms in $\tilde{\mathfrak{T}}$ are generated by r_α and $w_\alpha^{w\alpha} \in \text{Mor}_{\tilde{\mathfrak{T}}}([\alpha], [w\alpha])$, where $w \in W$ and the following diagrams commute:

$$\begin{array}{ccc} [\alpha] & \xrightarrow{(uw)_\alpha^{uw\alpha}} & [uw\alpha] \\ & \searrow w_\alpha^{w\alpha} & \nearrow u_\alpha^{uw\alpha} \\ & & [w\alpha] \end{array} \quad \begin{array}{ccc} [\alpha] & \xrightarrow{(w)_\alpha^{w\alpha}} & [w\alpha] \\ r_\alpha \downarrow & & \downarrow r_{-w\alpha} \\ [-\alpha] & \xrightarrow{(w^{-1})_\alpha^{-w\alpha}} & [-w\alpha] \end{array}$$

The Weyl groupoid is defined as

$$\mathfrak{W} := W \amalg \tilde{\mathfrak{T}},$$

where W is considered as a groupoid with a single point base $[W]$.

Example 14. For $\mathfrak{q}(2)$, let s be the nontrivial element of S_2 . Then \mathfrak{W} takes the following form

$$s \circlearrowleft [S_2] \quad r_{\varepsilon_2 - \varepsilon_1, \varepsilon_1 - \varepsilon_2} \left(\begin{array}{c} [\varepsilon_1 - \varepsilon_2] \\ s \left(\begin{array}{c} \nearrow s \\ \downarrow \\ \searrow s \end{array} \right) \\ [\varepsilon_2 - \varepsilon_1] \end{array} \right) r_{\varepsilon_2 - \varepsilon_2, \varepsilon_1 - \varepsilon_2}$$

Let \mathfrak{H} denote the category of all affine subspaces of \mathfrak{h} with morphisms given by affine transformations. For any $\alpha = \pm(\varepsilon_i - \varepsilon_j)$, let

$$\Pi_\alpha = \{(x_1, \dots, x_n) \in \mathbb{C}^n : x_i = -x_j\}.$$

Note that $\Pi_\alpha = \Pi_{-\alpha}$. Define $\tau_\alpha \in \text{Mor}_{\mathfrak{H}}(\Pi_\alpha, \Pi_\alpha)$ by the formula

$$\tau_\alpha(x_1, \dots, x_n) = (x_1, \dots, x_i + 1, \dots, x_j - 1, \dots, x_n).$$

Define the functor $F : \mathfrak{W} \rightarrow \mathfrak{H}$ by setting

$$F([\alpha]) = \Pi_\alpha, \quad F([W]) = \mathbb{C}^n, \quad F(r_\alpha) = \tau_\alpha, \quad F(w) = w, \quad F(w_{\alpha, w\alpha}) = \text{Res}_{\Pi_\alpha} w.$$

A function f on \mathfrak{h} is called \mathfrak{W} -invariant if for any $\varphi \in \text{Mor}_{\mathfrak{W}}(A, B)$

$$F(\varphi)^* \text{Res}_{FB} f = \text{Res}_{FA} f.$$

The description of $J(Q(n))$ can be formulated as follows:

Theorem 15. *The reduced Grothendieck ring $J(Q(n))$ of finite-dimensional representations of the supergroup $Q(n)$ is isomorphic to the ring $(\mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}])^{\mathfrak{W}}$ of invariants under the Weyl groupoid \mathfrak{W} as defined above.*

REFERENCES

- [B] J. Brundan, *Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra $\mathfrak{q}(n)$* , Adv. Math. **182**, No. 1, 28-77 (2004).
- [Che] S.J. Cheng, *Supercharacters of queer Lie superalgebras* Journal of Mathematical Physics **58**, 061701 (2017).
- [Cho] S. Cho, *A new Littlewood–Richardson Rule for Schur P -functions*, Trans. Amer. Math. Soc. **365** (2013), no. 2, 939–972.
- [CK] S.J. Cheng, J.H. Kwon, *Finite-dimensional half-integer weight modules over queer Lie superalgebras*. Commun. Math. Phys. 346, 945–965 (2016)
- [CW] S.J. Cheng, W. Wang, *Dualities and Representations of Lie Superalgebras*, Graduate Studies in Mathematics 144, AMS, 2
- [DS] M. Duflo, V. Serganova, *On associated varieties for Lie superalgebras*, arXiv:0507198.
- [GS] N. Grantcharov, V. Serganova, *Extension Quiver for Lie Superalgebra $\mathfrak{q}(3)$* , SIGMA **16** (2020), 141, 32 pages.
- [HR] C. Hoyt, S. Reif, *Grothendieck rings for Lie superalgebras and the Duflo-Serganova functor*, Algebra Number Theory **12** no. 9, 2167—2184, (2018).
- [I] V. N. Ivanov, *Interpolation analogues of Schur Q -functions*, , Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **307** (2004), no. Teor. Predst. Din. Sist. Komb. i Algoritm. Metody. 10, 99–119, 281–282. MR2050689 (2004m:05268)
- [IRS] M.S. Im, S. Reif, V. Serganova, *Grothendieck rings of periplectic Lie superalgebras*, arXiv:1906.01948, to appear in Math. Res. Lett..
- [K] V.G. Kac, *Lie superalgebras*, Advances in Math. **26**, No. 1 (1977), 8–96.
- [KW] V.G. Kac, M. Wakimoto, *Representations of affine superalgebras and mock theta functions*, Adv.Math. **300** (2016) 17-70.
- [M] I.G. Macdonald, *Symmetric functions and Hall polynomials*, second ed., Oxford Mathematical Monographs, Oxford Science Publications, The Clarendon Press Oxford University Press, New York 1995, With contributions by A. Zelevinsky. MR883894 (96h:05207).
- [Pa] N. Pragacz, *Algebraic-geometric applications of Schur S - and Q -polynomials*, Topics in invariant theory (Paris, 1989/1990), 130–191, Lecture Notes in Math. **1478**, Springer, Berlin, 1991.
- [Pe] I. Penkov, *Characters of typical irreducible finite-dimensional irreducible $\mathfrak{q}(n)$ -modules*, Lett. Math. Phys. **40** (1997), 147–158.
- [PS] I. Penkov, V. Serganova, *Cohomology of G/P for classical complex Lie supergroups G and characters of some atypical G -modules*, Annales de l’Institut Fourier, Tome **39** (1989) no. 4, pp. 845-873.
- [SV] A. N. Sergeev, A. P. Veselov, *Grothendieck rings of basic classical Lie superalgebras*, Ann. of Math. (2) **173** 663—703, (2011).