

# DIAGRAMS AND IRREGULAR CONNECTIONS ON THE RIEMANN SPHERE

JEAN DOUÇOT

ABSTRACT. We define a diagram associated to any algebraic connection on a vector bundle on a Zariski open subset of the Riemann sphere, generalizing previous constructions to the case when there are several irregular singularities. The construction relies on applying the Fourier–Laplace transform to reduce to the case where there is only one irregular singularity at infinity, and then using the definition of Boalch–Yamakawa in that case. We prove the diagram is invariant under the symplectic automorphisms of the Weyl algebra, so that there are several readings of the same diagram corresponding to connections with different formal data, usually on different rank bundles.

## 1. INTRODUCTION

This work takes place in line with a series of works by several authors establishing links between symplectic moduli spaces of meromorphic connections on the Riemann sphere and graphs (or doubled quivers). The main horizon, as outlined in [9], would be to develop a theory of Dynkin-like diagrams for nonabelian Hodge spaces, in the perspective of their classification. Recall that any irreducible meromorphic connection lives in a finite dimensional symplectic moduli space  $\mathcal{M}_{DR}$  and the wild nonabelian Hodge theorem of Biquard–Boalch/Sabbah shows such moduli spaces admit hyperkahler metrics and are diffeomorphic to moduli spaces  $\mathcal{M}_{Dol}$  of meromorphic Higgs bundles. Further, the Riemann–Hilbert–Birkhoff correspondence implies these spaces also have a simple explicit description as a wild character variety  $\mathcal{M}_B$  (cf. [10] and references there).

Relations between meromorphic connections and graphs have a long history. The case of regular singular connections was related by Crawley-Boevey/Shaw [13, 14, 15] to star-shaped graphs, in their work on the Deligne–Simpson problem, asking when the associated moduli spaces are non-empty. In the case of irregular connections, the link with graphs was extensively studied by Boalch [5, 6, 8, 12], partly motivated by the desire to understand better the origin of the symmetries of the Painlevé equations studied by Okamoto in a series of works [30]. The symmetry groups found by Okamoto are indeed Weyl groups of some affine Dynkin diagrams [34], so a natural question was to find a geometric explanation for the appearance of those diagrams. Since the Painlevé equations are obtained as isomonodromic deformation equations for rank two connections on  $\mathbb{P}^1$ , with a given irregular type for each Painlevé equation, a natural idea was thus to try to associate a graph to a meromorphic connection. This was first done in [5, 6] focussing mainly on the “simply-laced case” of connections with only one unramified irregular pole of order at most 3, together with possible simple poles. This led to a generalization of the star-shaped graphs, the supernova graphs, involving an arbitrary complete multipartite graph. The definition of the graph was generalized to the case where the irregular pole is still unramified but has arbitrary order, in [5] Appendix C, where a conjecture was made relating the corresponding moduli spaces to Nakajima quiver varieties. This “quiver modularity conjecture” was proved in 2014 by Hiroe–Yamakawa [22]: a large class of quiver varieties are thus isomorphic to moduli spaces  $\mathcal{M}_{DR}^*$  of meromorphic connections. Beyond this case most of the moduli spaces are not directly related to Nakajima quiver varieties, so a different approach was needed.

Recently, Boalch and Yamakawa suggested a more general definition [12], for the case where there is at most one irregular pole, but with arbitrary structure; it is allowed to be twisted, i.e. to feature ramified exponential factors. The result is no longer a graph but a generalization they called a diagram: it is a graph where each edge has a multiplicity that may be negative. They showed the basic features of the story still hold and in particular that the dimension of

the moduli space is determined by the Cartan matrix of the diagram and the dimension vector, as for quiver varieties (cf. Thm. 6.1 below).

In this work, we extend this story to the general case of connections with an arbitrary number of irregular singularities of arbitrary formal type on the Riemann sphere. A specific motivation is to better understand the relation between Painlevé equations and quivers for the case of the third Painlevé equation, whose standard Lax pair features two second order poles and doesn't fit in the framework of previous works. We do this by implementing the strategy outlined in [12]. The idea is to apply the Fourier transform to reduce to the case where there is only one singularity at infinity. The diagram will then correspond both to the initial connection and to its Fourier transform. One may actually interpret in this way chapter 12 of Malgrange's book [28] which essentially corresponds to a bipartite graph whose vertices can be interpreted both in terms of the initial  $\mathcal{D}$ -module and of its Fourier transform. This class of examples was further extended from the bipartite case to the multipartite case in the simply-laced story [5, 6, 8]. They are symmetric under a larger class of transformations containing the Fourier transform: a natural action of  $SL_2(\mathbb{C})$  on the Weyl algebra. We will show that our general definition of diagram is invariant under this full  $SL_2(\mathbb{C})$  symmetry group, and thus independent of any choices involved in reducing to the set-up of [12].

In all those cases, the graphs associated to meromorphic connections feature several properties which make them useful objects in the study of their moduli spaces. For example, several formal data give rise to the same diagram: there are several ways of reading the same diagram as coming from different formal data, e.g. with different ranks or numbers of singularities. A remarkable fact is that in these cases the moduli spaces corresponding to those different readings are all isomorphic. The diagrams thus enable us to find isomorphisms between these wild character varieties, and we expect they will be a useful tool for the classification.

We make an essential use of the stationary phase formula which relates the formal data of a connection to those of its Fourier-Laplace transform: it says basically that their exponential factors are related by a Legendre transform. The formula was actually first established by Laumon [25] in a  $\ell$ -adic context, then proven in the complex setting by Fang and Sabbah independently [35, 17], and is already present in [28]. An essential feature is that the Fourier transform acts on the formal data in a non-local way, e.g. it sends all singularities at finite distance to infinity. This will allow us to reduce to the case with only one singularity. Basically, what happens is that  $SL_2(\mathbb{C})$  transformations act as the corresponding homographies on the *Fourier sphere* of [6], i.e. on the coefficients of the slope 2 terms at infinity, with singularities at finite distance corresponding to this coefficient being equal to infinity. Therefore, a generic rotation of the Fourier sphere ensures that all exponential factors are at infinity.

Some of the properties of the diagrams in the particular cases considered in previous works remain true in the more general setting. There are still several readings of the diagram depending on which point of the Fourier sphere we rotate to infinity. The dimension of the wild character variety is obtained from the diagram by a formula involving its Cartan matrix.

In [20], Hiroe defines quivers associated to unramified irregular connections on with an arbitrary number of irregular singularities. Our approach is many in respects different from his since he does not use the Fourier-Laplace transform to define the graph, and our diagrams are not the same as his. Furthermore, particular cases of our diagrams have been independently obtained by Xie [36] in the context of 3d mirror symmetry for some Argyres-Douglas theories. It would be interesting to better understand whether the different readings of our diagrams may be related to dualities of quantum field theories.

**1.1. Summary of results.** Our main result is to define a diagram in the general case of connections with any number of poles of any type on the Riemann sphere.

**Theorem 1.1.** *Given any algebraic connection  $(E, \nabla)$  on a Zariski open subset of the affine line  $\mathbb{C} = \mathbb{P}^1 \setminus \infty$ , there is a uniform way to define a diagram  $\Gamma$  and a dimension vector  $\mathbf{d} \in \mathbb{N}^N$  such that the wild character variety  $\mathcal{M}_B$  determined by  $(E, \nabla)$  has dimension*

$$(1.1) \quad \dim(\mathcal{M}_B) = 2 - (\mathbf{d}, \mathbf{d})$$

where  $(, )$  is the symmetric bilinear form determined by the Cartan matrix  $C = 2 - B$  of  $\Gamma$ .

To state the definition of the diagram, we review briefly the singularity data associated to an algebraic connection on a Zariski open subset of the affine line. Each point  $a \in \mathbb{P}^1$  determines a collection  $\mathcal{I}_a$  of circles  $\langle q \rangle$ . In brief each circle  $\langle q \rangle$  is the Riemann surface of a germ of a function of the form

$$(1.2) \quad q = \sum_{i=0}^s a_i z_a^{-\alpha_i/\beta},$$

on an open sector at  $a$ , where  $a_i \in \mathbb{C}$ ,  $a_i \neq 0$  and  $\alpha_i, \beta \geq 1$  are integers, in some local coordinate  $z_a$  at the singularity  $a \in \mathbb{P}^1$ . Some numbers attached to each circle  $\langle q \rangle$ ,  $q = \sum a_i z_a^{-\alpha_i/\beta}$  are as follows:

- slope( $q$ ) is the largest rational number  $\alpha_i/\beta$  present in  $q$ ,
- The ramification degree  $\text{ram}(q)$  is the lowest possible integer  $\beta \geq 1$  in  $q$ ,
- The irregularity  $\text{Irr}(q) \in \mathbb{N}$  is the product  $\text{ram}(q)\text{slope}(q)$ .

By definition an *irregular class*  $\Theta$  at  $a$  is the choice of a finite number of circles  $\langle q_i \rangle \subset \mathcal{I}_a$ , each with an integer multiplicity  $n_i \geq 1$ . An irregular class can be written as a formal sum:

$$\Theta = n_1 \langle q_1 \rangle + \cdots + n_m \langle q_m \rangle.$$

Any algebraic connection  $(E, \nabla)$  on a Zariski open subset of the affine line  $\mathbb{C} = \mathbb{P}^1 \setminus \infty$  determines an irregular class at each singularity; in brief the horizontal sections of the connection in the vicinity of  $a$  are linear combinations of terms featuring the exponentials  $e^{q(z)}$ . The irregularity of any irregular class  $\Theta = \sum n_i \langle q_i \rangle$  is  $\text{Irr}(\Theta) = \sum n_i \text{Irr}(q_i)$ , and the rank of an irregular class is  $\sum n_i \text{ram}(q_i)$ . Any circles  $\langle q_1 \rangle, \langle q_2 \rangle$  at  $a$  determine an irregular class  $\text{Hom}(\langle q_1 \rangle, \langle q_2 \rangle)$  of rank equal to the product  $\text{ram}(q_1) \text{ram}(q_2)$ .

The singularity data  $(\Theta, \mathcal{C})$  at  $a$  consists of an irregular class  $\Theta = \sum n_i \langle q_i \rangle$  together with conjugacy classes  $\mathcal{C}_i \subset GL_{n_i}(\mathbb{C})$ , for each circle in  $\Theta$ . They are the “formal monodromy classes” determined by the connection. Two connections are formally isomorphic at  $a$  if and only if they have the same singularity data at  $a$ .

Our definition builds on the definition of [12], which applies to the case where there is just one singularity at infinity. Suppose we have a connection  $(E, \nabla)$  on the Riemann sphere with just one pole, at  $\infty$ . Let  $(\Theta, \mathcal{C})$  be the singularity data of the connection at  $\infty$ . Let  $N_c$  be the set of circles present in  $\Theta$ . The diagram of [12] is built by gluing together a core diagram  $\Gamma_c$  and legs. The core diagram has nodes  $N_c$  and there is one leg for each node of the core diagram, determined by the conjugacy class  $\mathcal{C}_i$ . The edge multiplicities  $B_{ij}$  for the core diagram are as follows:

**Definition 1.2** ([12]). Write  $\Theta = \sum n_i \langle q_i \rangle$ . For any  $i, j \in N_c$ :

- If  $i \neq j$  then

$$(1.3) \quad B_{ij} = A_{ij} - \beta_i \beta_j,$$

where  $A_{ij} = \text{Irr}(\text{Hom}(\langle q_i \rangle, \langle q_j \rangle))$  and  $\beta_i = \text{ram}(q_i)$ ,

- If  $i = j$  then

$$(1.4) \quad B_{ii} = A_{ii} - \beta_i^2 + 1.$$

This definition was extended in [12] §2.1 to the case of connections with further tame poles on  $\mathbb{C}$ . The aim of this paper is to extend the definition to the case of connections with any number of poles of any type. For example the definition of [12] §2.1 covered the case of Painlevé III using an alternative Lax representation (with two tame poles and a pole at  $\infty$  with irregular class  $\langle z^{1/2} \rangle$ ) but it did not cover the standard representation, with two poles of order two. We will see how the same diagram appears directly from the standard representation of Painlevé III using our general definition.

In the general case, the diagram  $\Gamma(E, \nabla) = \Gamma(\Theta, \mathcal{C})$  that we associate to the connection  $(E, \nabla)$  only depends the singularity data  $(\Theta, \mathcal{C})$  at all of the singular points. It has again the

following structure: it consists of a core diagram  $\Gamma_c$  to which are then glued legs encoding certain conjugacy classes.

Let  $\mathcal{I} = \bigcup_{a \in \mathbb{P}^1} \mathcal{I}_a$  be the disjoint union of all the circles at all the points of the Riemann sphere. There is a map  $\pi : \mathcal{I} \rightarrow \mathbb{P}^1$  taking a circle to the point it lies over. As before the formal type of the connection determines an integer  $n_i$  and a conjugacy class  $\mathcal{C}_i \subset GL_{n_i}(\mathbb{C})$  for each circle  $i \subset \mathcal{I}$ , so that  $n_i = 0$  for all but a finite number of circles at each singular point.

We then modify this formal data as follows: if  $i$  is a tame circle at finite distance ( $\pi(i) \in \mathbb{C}$  and  $q_i = 0$ ) then replace  $n_i$  by the integer

$$m_i = \text{rank}(A - 1)$$

where  $A \in \mathcal{C}_i$ . Thus  $A = 1 + uv$  for a surjective map  $u : \mathbb{C}^{n_i} \rightarrow \mathbb{C}^{m_i}$  and an injective map  $v : \mathbb{C}^{m_i} \rightarrow \mathbb{C}^{n_i}$ . Then replace  $\mathcal{C}_i$  by the conjugacy class  $\check{\mathcal{C}}_i$  of  $(1 + uv)$  in  $GL_{m_i}(\mathbb{C})$ ; this class is called the *child* of  $\mathcal{C}_i$  in [8, appendix]. If  $\pi(i) = \infty$  or if  $i$  is not tame ( $q_i \neq 0$ ) then we do no modification:  $m_i = n_i$  and  $\check{\mathcal{C}}_i = \mathcal{C}_i$ . This defines modified formal data  $(\check{\Theta}, \check{\mathcal{C}})$ , where  $\check{\Theta}$  is the collection of circles with multiplicities  $m_i \geq 1$ . The core nodes  $N_c$  of our diagram are given by this collection of circles:

$$N_c = \{\text{circles } i \subset \mathcal{I} \mid m_i \geq 1\}.$$

We glue the leg determined by the conjugacy class  $\check{\mathcal{C}}_i$  onto the node  $i \in N_c$  as before. The remaining step to define the diagram is to define the integers  $B_{ij} = B_{ji}$  for  $i, j \in N_c$ :

**Definition 1.3.** Suppose  $i, j \in N_c$  are active circles and write  $a_i = \pi(i), a_j = \pi(j) \in \mathbb{C} \cup \infty$ , so that  $i = \langle q_i \rangle, j = \langle q_j \rangle$  are circles at  $a_i, a_j$  respectively. Let  $\alpha_i = \text{Irr}(q_i), \beta_i = \text{ram}(q_i)$  and similarly for  $j$ . If  $a_i = a_j$  let us denote by  $B_{i,j}^\infty$  the number determined by the expressions (1.3) and (1.4), applied to the circles  $i, j$ .

- (1) If  $a_i = a_j = \infty$  then  $B_{ij} = B_{ij}^\infty$ .
- (2) If  $a_i = \infty$  and  $a_j \neq \infty$  then  $B_{ij} = B_{ji} = \beta_i(\alpha_j + \beta_j)$ .
- (3) If  $a_i \neq \infty, a_j \neq \infty$  and  $a_i \neq a_j$  then  $B_{ij} = B_{ji} = 0$ .
- (4) If  $a_i = a_j \neq \infty$  then  $B_{ij} = B_{ji} = B_{i,j}^\infty - \alpha_i\beta_j - \alpha_j\beta_i$ .

We will prove directly that this does indeed define a diagram (i.e.  $B_{ii}$  is an even integer) in §4, independently of the proof suggested in [12] for the special case of one irregular singularity.

The basic idea leading to this definition goes as follows: we want to reduce to the situation of [12] by applying the Fourier-Laplace transform. The Fourier-Laplace transform already plays an important role in previous constructions, e.g. it is underlying the different readings of the quiver in [5, 6], and often allows to pass from one Lax representation to another for the Painlevé equations. The Fourier-Laplace transform induces a formal Fourier transform at the level of the formal data [4, 26]. More explicitly, the stationary phase formula [28, 17, 35] relates the global formal data of a connection to those of its Fourier-Laplace transform. A remarkable fact is that the Fourier-Laplace transform acts on the formal data in a non-local way. We use this property to send all active circles to infinity and recover the setting of [12]. The non-trivial point is to ensure that the diagram we obtain is independent of the choices made along the way.

Let us sketch the main ideas a little more precisely. Connections on Zariski open subsets of the affine line are closely related to modules over the Weyl algebra  $A_1 := \mathbb{C}[z]\langle \partial_z \rangle$ . There is a natural action of the group  $SL_2(\mathbb{C})$  on the Weyl algebra: a matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

in  $SL_2(\mathbb{C})$  induces an automorphism of the Weyl algebra by

$$(z, \partial_z) \mapsto (az + b\partial_z, cz + d\partial_z).$$

Of particular significance among those transformations is the Fourier-Laplace transform, corresponding to the matrix

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

The stationary phase formula [17, 35] implies that those symplectic transformations induce an action of  $SL_2(\mathbb{C})$  on modified singularity data  $(\check{\Theta}, \check{\mathcal{C}})$  of connections. The main result of this work, ensuring the diagram is independent of the choices, leading to our definition, is the following:

**Theorem 1.4.** *The diagram is invariant under symplectic transformations: if  $A \in SL_2(\mathbb{C})$ , then*

$$\Gamma(A \cdot (\check{\Theta}, \check{\mathcal{C}})) = \Gamma(\check{\Theta}, \check{\mathcal{C}}).$$

The diagram is thus an invariant under the action on  $SL_2(\mathbb{C})$ . As a consequence, a given diagram can be read in different ways, as coming from connections with different formal data related by a symplectic transformation.

**1.2. Plan of the paper.** Here is a brief outline of the structure of the paper. In section 2 we recall some facts about formal data of meromorphic connections, which we formulate in terms of local systems on a large collection of circles corresponding to the exponential factors of the connection. There are some subtleties involving the relation between connections and  $\mathcal{D}$ -modules, which lead to the introduction of the modified formal data. In section 3, we discuss the action of  $SL_2(\mathbb{C})$ . We review the stationary phase formula which relates the formal data of a connection to those of its Fourier transform. In section 4, we study the diagram associated to a connection with just one singularity at infinity defined in [12] and show that it is invariant under symplectic transformations. This is the crucial fact that will lead to the definition of the diagram in the general case. To this purpose, we find an explicit formula for the number of edges and loops of the diagram which is of independent interest. We also determine the monomials appearing in the Legendre transform of any exponential factor. The definition of the diagram in the general setting is given in section 5, where we also discuss how the different readings of the diagram are generalized to our setting. In section 6, we look at the dimension of the wild character variety, and show that it is given from the graph by the formula (1.1). Finally, in section 7 we discuss several examples of diagrams, many of them related to Painlevé equations. To facilitate the reading, a few somewhat lengthy proofs are relegated to appendices.

**Acknowledgements.** I thank my advisor Philip Boalch for his constant support and many useful discussions.

## 2. FORMAL DATA OF IRREGULAR CONNECTIONS

In this section, we describe the data encoding the formal type of an irregular connection on the Riemann sphere at each of its singular points. Among the different existing approaches to the formal classification of meromorphic connections, the one we will use is the geometric description of [11] close to the point of view of Deligne and Malgrange [16, 28], in terms of graded local systems on the circles of directions around the singular points. We refer the reader to [11] for more details.

**2.1. The exponential local system.** Let  $\Sigma$  be an algebraic curve, and  $a \in \Sigma$ . We consider the real blow-up  $\pi_a : \widehat{\Sigma}_a \rightarrow \Sigma$  of  $\Sigma$  at  $a$ , such that  $\partial_a := \pi^{-1}(a)$  is the circle of directions around  $a$ .

For  $a \in \Sigma$ , let  $z_a$  be a local coordinate at  $a$ . We define a local system of sets (i.e. a cover)  $\mathcal{I}$  on the circle  $\partial_a$  in the following way: if the open subset  $U \subset \partial_a$  is a germ of angular sector at  $a$ , the sections of  $\mathcal{I}_a$  on  $U$  are functions of the form

$$q_U(z_a) = \sum_{i=1}^s b_i z_a^{-i/r},$$

where  $r, s \in \mathbb{N}$ ,  $b_i \in \mathbb{C}$  for  $i = 1, \dots, s$ , for some determination on  $U$  on the  $r$ -th root  $z^{1/r}$ . It is also possible to give an intrinsic definition of  $\mathcal{I}_a$  independent of the choice of a local coordinate, see [11, Remark 3].

As a topological space, the cover  $\mathcal{I}_a$  is a disjoint union of circles. To any polynomial  $q \in z_a^{-1/r} \mathbb{C}[z_a^{-1/r}]$  in  $z_a^{1/r}$  corresponds a connected component of  $\mathcal{I}_a$ , homeomorphic to a circle, that we will denote  $\langle q \rangle_a$ , or simply  $\langle q \rangle$  when there will be no ambiguity about the point  $a$  which is considered. The function  $z_a \mapsto q(z_a)$  on the punctured disk  $D_a^\times$  around  $a$  is multivalued: if  $r$  is the smallest possible denominator for the exponents appearing in  $q$ , the cover  $\langle q \rangle \rightarrow \partial_a$  is of order  $r$ ; each leaf of the cover above  $U \subset \partial_a$  corresponding to a determination of the  $r$ -th root  $z_a^{1/r}$ . The number  $r =: \text{ram}(q)$  is the *ramification index* of  $q$ , its degree  $s$  as a polynomial in  $z_a^{1/r}$  is its *irregularity*  $\text{Irr}(q) := s$ . The quotient  $s/r$  is the *slope* of  $q$ . The circle  $\langle 0 \rangle$  is the *tame circle*.

If  $I \in \pi_0(\mathcal{I}_a)$  is a connected component of  $\mathcal{I}_a$ , there is no unique  $q$  such that  $I = \langle q \rangle_a$ : if  $\omega$  is a  $r$ -th root of unity with  $r = \text{ram}(q)$ , one has  $\langle q(z_a) \rangle = \langle q(\omega z_a) \rangle$ . The set of polynomials  $q'$  such that  $I = \langle q' \rangle_a$  is the Galois orbit of  $q$  under the action of the group  $\mu_r$  of  $r$ -th roots of unity. We thus have

$$\mathcal{I}_a = \bigsqcup \langle q \rangle_a.$$

Let us fix a direction  $p \in \partial_a$ , and denote by  $(\mathcal{I}_a)_p := \pi_a^{-1}(p)$  the fibre of  $\mathcal{I}_a$  at  $p$ . The monodromy of  $\mathcal{I}_a$  is an automorphism  $\rho_a : (\mathcal{I}_a)_p \rightarrow (\mathcal{I}_a)_p$ .

All this was local at a point  $a \in \Sigma$ . At the global level, we define the global exponential local system to be

$$\mathcal{I} = \bigsqcup_{a \in \Sigma} \mathcal{I}_a.$$

As a topological space,  $\mathcal{I}$  is thus a large collection of circles.

## 2.2. Graded local systems.

**Definition 2.1.** An  $\mathcal{I}_a$ -graded local system is a local system  $V^0 \rightarrow \partial_a$  of finite-dimensional vector spaces endowed with a point-wise grading

$$V_p^0 = \bigoplus_{i \in (\mathcal{I}_a)_p} V_{p,i}^0,$$

where  $V_p^0$  is the fibre of  $V^0$  at  $p$ , for each point  $p \in \partial_a$ , such that the grading is locally constant.

If we fix a base point  $p \in \partial_a$ , let  $\rho \in \text{Aut}((\mathcal{I}_a)_p)$  be the monodromy of  $\mathcal{I}_a$  and  $\hat{\rho} \in GL(V_p^0)$  the monodromy of  $V$ , this implies that

$$\hat{\rho}(V_{p,i}^0) = V_{p,\rho(i)}^0.$$

The data of its monodromy determines the local system  $V^0$  up to isomorphism. More precisely, the isomorphism class of an  $\mathcal{I}_a$ -graded local system  $V^0$  corresponds to the conjugacy class of  $\hat{\rho}$  under graded automorphisms of  $V_p^0$ .

Let  $V^0 \rightarrow \partial_a$  be an  $\mathcal{I}_a$ -graded local system. For  $i \in (\mathcal{I}_a)_p$ , let  $n_i := \dim V_{i,p}^0$ . If  $I = \langle q \rangle$  is a circle in  $\mathcal{I}_a$  with ramification order  $r$ , the fibre  $\langle q \rangle_p$  of the restriction  $\langle q \rangle \rightarrow \partial_a$  has  $r$  elements. Let  $V_{\langle q \rangle}^0 \rightarrow \partial$  be the restriction of  $V^0$  to its  $\langle q \rangle$ -graded part. Then all graded pieces of  $V_{\langle q \rangle}^0 \rightarrow \partial$  have the same dimension  $n_{\langle q \rangle}$ , that is  $n_{\langle q \rangle} = n_i$  for all  $i \in (V_{\langle q \rangle}^0)_p$ . The integer  $n_{\langle q \rangle}$  is the *multiplicity* of  $\langle q \rangle$ .

**Definition 2.2.** A (local) irregular class at  $a$  is a function  $\pi_0(\mathcal{I}_a) \rightarrow \mathbb{N}$ .

The *irregular class* of  $V^0$  is the function

$$\Theta_a(V^0) : \pi_0(\mathcal{I}_a) \rightarrow \mathbb{N}, \\ \langle q \rangle \mapsto n_{\langle q \rangle}.$$

The *active circles* the circles  $\langle q \rangle$  such that  $n_{\langle q \rangle} \neq 0$ .

The classical Fabry-Hukuhara-Turritin-Levelt theorem on the formal classification of meromorphic connections is formulated in this language in the following way:

**Theorem 2.3** ([16, 27]). *The category of connections on the formal punctured disk is equivalent to the category of  $\mathcal{I}_0$ -graded local systems.*

The basic idea behind this is that the active circles  $\langle q_i \rangle$  of the  $\mathcal{I}_0$ -graded local system associated to the formal connection correspond to the exponential terms  $e^{q_i}$  in the horizontal sections of the Hukuhara-Turritin normal form.

It is possible to view an  $\mathcal{I}_a$ -graded local system  $V^0 \rightarrow \partial$  as a local system on  $\mathcal{I}_a$ , such that the following diagram commutes:

$$\begin{array}{ccc} V^0 & \longrightarrow & \mathcal{I} \\ & \searrow & \downarrow \pi \\ & & \partial \end{array}$$

The isomorphism class of a local system  $V^0 \rightarrow \mathcal{I}_a$  with irregular class  $\Theta_a$  is given by the collection  $\mathcal{C} = (\mathcal{C}_{\langle q \rangle})$ , where  $\mathcal{C}_{\langle q \rangle} \subset GL_{n(\langle q \rangle)}(\mathbb{C})$  of the conjugacy classes of its monodromies around its active circles.

In all that follows, we consider an algebraic connection  $(E, \nabla)$  on a Zariski open subset  $\mathbb{A}^1 \setminus \{a_1, \dots, a_m\}$  of the affine line  $\mathbb{A}^1 = \mathbb{P}^1 \setminus \infty$ . From the formal classification,  $(E, \nabla)$  defines a global formal local system  $V^0 \rightarrow \mathcal{I}$ , with support on the singular points. Its isomorphism class is given by the pair  $(\Theta, \mathcal{C})$ , with  $\Theta : \pi_0(\mathcal{I}) \rightarrow \mathbb{N}$  the global irregular class, and  $\mathcal{C}$  the collection of the formal monodromies at each active circle. We will refer to the pair  $(\Theta, \mathcal{C})$  as the *formal data* of the connection  $(E, \nabla)$ .

**2.3. Modified formal data.** To arrive at our definition of the diagrams, we will need to use the Fourier-Laplace transform. However, the category which the Fourier-Laplace transform acts on is not the category of connections, but the category of modules on the Weyl algebra  $A_1 = \mathbb{C}[z]\langle \partial_z \rangle$ . This is the source of an important subtlety, pertaining to the fact that more formal data are needed to describe a  $\mathcal{D}$ -module than a connection. We take this into account by considering modified formal data, which we now describe.

We first consider the local situation at a singularity  $a \in \Sigma$ . Let  $V_{\langle 0 \rangle_a}^0$  be a local system on the tame circle  $\langle 0 \rangle_a$ , with monodromy  $T$ . We associate to it another local system  $\check{V}_{\langle 0 \rangle_a}^0 \rightarrow \langle 0 \rangle_a$  defined by  $\check{V}_{\langle 0 \rangle_a}^0 := \text{Im}(T - \text{Id})$ . Its rank is  $\text{rank}(T - \text{Id})$  and it has monodromy  $T|_{\text{Im}(T - \text{Id})}$ . Notice that the data of  $\check{V}_{\langle 0 \rangle_a}^0$  together with the rank of  $V_{\langle 0 \rangle_a}^0$  is enough to reconstruct the isomorphism class of  $V_{\langle 0 \rangle_a}^0$ .

Now, let  $V^0 \rightarrow \mathcal{I}_a$  be a formal local system at  $a$ . We define the *modified formal local system*  $\check{V}^0 \rightarrow \mathcal{I}$  associated to  $V^0$  to be the local system on  $\mathcal{I}_a$  such that  $\check{V}_{\langle 0 \rangle_a}^0$  is obtained from  $V_{\langle 0 \rangle_a}^0$  as described above, and  $\check{V}_{\langle q \rangle_a}^0 = V_{\langle q \rangle_a}^0$  for any other connected component  $\langle q \rangle_a$ .

Turning to the global situation for  $\Sigma = \mathbb{P}^1$ , if  $V^0 \rightarrow \mathcal{I}_a$  is a global formal local system, applying at each point at finite distance the previous construction defines a modified global formal local system  $\check{V}^0 \rightarrow \mathcal{I}$ .

**Definition 2.4.** Let  $(E, \nabla)$  an algebraic connection on a Zariski open subset  $\mathbb{A}^1 \setminus \mathbf{a}$  of the affine line,  $V^0 \rightarrow \mathcal{I}$  its formal local system and  $(\Theta, \mathcal{C})$  its (global) formal data. Its global modified formal data  $(\check{\Theta}, \check{\mathcal{C}})$  correspond to the isomorphism class of the global modified local system  $\check{V}^0 \rightarrow \mathcal{I}$  obtained by taking the (local) modified formal local system at each point  $a \neq \infty$  at finite distance, and keeping the non-modified formal local system  $V^0 \rightarrow \mathcal{I}_\infty$  at infinity.

Notice that  $\check{V}^0$  satisfies the condition  $\text{rank}(\check{V}_a^0) \leq \text{rank} \check{V}_\infty^0$  since for each  $a \neq \infty$  passing to the modified formal local system lowers the rank. If  $\check{V} \rightarrow \mathcal{I}$  is a formal local system, with isomorphism class  $\check{\mathcal{C}}$ , we say that  $V$ , or  $\check{\mathcal{C}}$  is *compatible* if it satisfies this condition. If there exists a connection  $(E, \nabla)$  on a Zariski open subset of the affine line such that  $\check{\mathcal{C}}$  is its modified formal data, we say that  $\check{\mathcal{C}}$  is *effective*.  $\check{\mathcal{C}}$  cannot be effective if it is not compatible.

If we know the modified formal local system  $\check{V}^0$  of a connection  $(E, \nabla)$ , since we keep the non-modified local system  $V_\infty^0 \rightarrow \mathcal{I}_\infty$  at infinity, we know the rank of  $(E, \nabla)$ . It is given by

$$\text{rank}(E, \nabla) = \sum_{\langle q \rangle \in \pi_0(\mathcal{I}_\infty)} n_{\langle q \rangle} \text{ram}(q).$$

In turn, this enables us to reconstruct from  $\check{V}^0 \rightarrow \mathcal{I}$  the isomorphism class of the non-modified local system  $V^0 \rightarrow \mathcal{I}$ . The data of  $(\check{\Theta}, \check{\mathcal{C}})$  is thus equivalent to the data of  $(\Theta, \mathcal{C})$ .

*Remark 2.1.* To a module  $M$  on the Weyl algebra one can associate a modified formal local system  $V \rightarrow \mathcal{I}$ , which encodes part of its formal data. This follows from the description of  $\mathcal{D}$ -modules given in [28, chapter IV], if we define the regular piece  $\check{V}^0 \rightarrow \langle 0 \rangle_a$  to correspond to the vanishing cycles of the regular part of the  $\mathcal{D}$ -module at  $a$ . If  $(E, \nabla)$  is an algebraic connection on a Zariski open subset  $U = \mathbb{A}^1 \setminus \mathbf{a}$  of the affine line, passing to the modified formal local system corresponds to taking the modified formal local system of its *minimal extension* (see [1, §5.5])  $M$  of  $(E, \nabla)$  along  $\mathbf{a}$  to  $\mathbb{A}^1$ , which is a module on the Weyl algebra.

**2.4. Summary of data.** In the rest of the article, we fix an algebraic connection  $(E, \nabla)$  on  $\mathbb{A}^1 \setminus \mathbf{a}$ , with modified irregular class  $\check{\Theta} : \pi_0(\mathcal{I}) \rightarrow \mathbb{N}$ , and modified conjugacy classes  $\check{\mathcal{C}}$ . We fix the following notations.

- We denote by  $a_1, \dots, a_m$  be the singularities at finite distance, so that  $\mathbf{a} = \{a_1, \dots, a_m\}$
- For  $k = 1, \dots, m$  let  $\langle q_1^{(k)} \rangle_{a_k}, \dots, \langle q_{r_k}^{(k)} \rangle_{a_k} \in \pi_0(\mathcal{I}_{a_k})$  denote the irregular active circles at  $a_k$ ,  $n_i^{(k)} \in \mathbb{N}$  their respective multiplicities, and  $\beta_j^{(k)}$  their respective ramification orders, and  $\alpha_j^{(k)}$  their irregularities, in an order such that the slopes satisfy

$$\frac{\alpha_1^{(k)}}{\beta_1^{(k)}} \geq \dots \geq \frac{\alpha_{r_k}^{(k)}}{\beta_{r_k}^{(k)}}.$$

- For  $k = 1, \dots, m$  let  $m_k$  be the multiplicity of the tame circle  $\langle 0 \rangle_{a_k}$  at  $a_k$  in the *modified* formal local system.
- At infinity, let  $\langle q_1^{(\infty)} \rangle_{\infty}, \dots, \langle q_{r_{\infty}}^{(\infty)} \rangle_{\infty} \in \pi_0(\mathcal{I}_{a_k})$  denote the active circles (the tame circle is included), their  $\beta_j^{(\infty)}$  respective ramification orders, and  $\alpha_j^{(\infty)}$  their irregularities, in an order such that the slopes satisfy

$$\frac{\alpha_1^{(\infty)}}{\beta_1^{(\infty)}} \geq \dots \geq \frac{\alpha_{r_{\infty}}^{(\infty)}}{\beta_{r_{\infty}}^{(\infty)}}.$$

The rank of any connection having these modified formal data is  $\sum_{j=1}^{r_{\infty}} n_i^{(\infty)} \beta_j^{(\infty)}$ .

### 3. ACTION OF $SL_2(\mathbb{C})$

We now discuss the action of the Fourier-Laplace transform on modules on the Weyl algebra, and the stationary phase formula relating the formal data of an  $A_1$ -module and its Fourier transform.

**3.1. The stationary phase formula.** The Fourier transform is the automorphism of the Weyl algebra  $A_1 = \mathbb{C}[z]\langle \partial_z \rangle$  defined by  $z \mapsto -\partial_z$ ,  $\partial_z \mapsto z$ . It induces a transformation on modules on the Weyl algebra: the Fourier-Laplace transform  $F \cdot M$  of a  $\mathbb{C}[z]\langle \partial_z \rangle$ -module  $M$  is the  $\mathbb{C}[\xi]\langle \partial_{\xi} \rangle$ -module obtained by setting:  $z \mapsto -\partial_{\xi}$ ,  $\partial_z \mapsto \xi$  (we use a dual variable  $\xi$  for the image in this paragraph). In particular, if  $M = \mathcal{D}_z / \mathcal{D}_z p$  with  $p \in \mathbb{C}[z]\langle \partial_z \rangle$ , one has  $F \cdot M = \mathcal{D}_{\xi} / \mathcal{D}_{\xi} p'$ , where  $p'$  is obtained from  $p$  by replacing  $z$  by  $-\partial_{\xi}$  and  $\partial_z$  by  $\xi$ .

If  $(E, \nabla)$  is an algebraic connection on a Zariski open subset  $\Sigma^{\circ} \subset \mathbb{C}$ , let  $M$  its minimal extension, and  $M' = F \cdot M$ . There exists [2] a meromorphic connection  $(E', \nabla')$  on a Zariski open subset  $\Sigma'^{\circ} \subset \mathbb{C}$  such that  $M'$  is the minimal extension of  $(E, \nabla)$ . It is the Fourier transform of  $(E, \nabla)$  and we denote it by  $F \cdot (E, \nabla)$ .

The stationary phase formula [17, 35] states that the formal data of the Fourier transform  $(E', \nabla')$  are determined by the formal data of  $(E, \nabla)$ .

**Theorem 3.1.** *There exists a bijection, that we also denote by  $F$ , from the set of effective formal data to itself, such that if  $(E, \nabla)$  is a connection of a Zariski open subset of  $\mathbb{C}$ ,  $(\check{\Theta}, \check{\mathcal{C}})$*



its modified formal data,  $(E', \nabla')$  is the Fourier transform of  $(E, \nabla)$  and  $(\check{\Theta}', \check{\mathcal{C}}')$  its modified formal data, the following diagram commutes:

$$\begin{array}{ccc} (E, \nabla) & \xrightarrow{F} & (E', \nabla') \\ \downarrow & & \downarrow \\ (\check{\Theta}, \check{\mathcal{C}}) & \xrightarrow{F} & (\check{\Theta}', \check{\mathcal{C}}') \end{array}$$

This map  $F$  is the formal Fourier transform. More precisely, the active exponents of  $\check{\Theta}'$  are related to the active exponents of  $\check{\Theta}$  by a Legendre transform.

Let us describe the formal Fourier transform and express it in our framework of local systems on  $\mathcal{I}$ . To this end, we formulate the Legendre transform as a homeomorphism sending collection of circles  $\mathcal{I}$  to the dual collection  $\mathcal{I}'$ , obtained by replacing the variable  $z$  by the variable  $\xi$ . The Legendre transform takes different forms depending on the circles in  $\pi_0(\mathcal{I})$ . It is therefore necessary to distinguish several types of circles as follows:

**Definition 3.2.** Any circle in  $\mathcal{I}$  belongs to one of the following five families:

- (1) The *pure circles* at infinity, of the form  $\langle \alpha z \rangle_\infty$ , with  $\alpha \in \mathbb{C}$ . There are of slope 1 if  $\alpha \neq 0$ , and 0 otherwise.
- (2) Other circles of slope  $\leq 1$  at infinity, of the form  $\langle \alpha z + q \rangle_\infty$ , with  $\alpha \in \mathbb{C}$ , and  $q \neq 0$  of slope  $< 1$ ,
- (3) Circles  $\langle q \rangle_\infty$  of slope  $> 1$  at infinity,
- (4) Irregular circles at finite distance  $\langle q \rangle_a$ , with  $q \neq 0$ ,  $a \in \mathbb{C} = \mathbb{P}^1 \setminus \{\infty\}$ .
- (5) The tame circles  $\langle 0 \rangle_a$ ,  $a \in \mathbb{C}$ .

We will denote by  $\mathcal{I}_1, \dots, \mathcal{I}_5 \subset \mathcal{I}$  the corresponding collections of circles. In a similar way, we denote by  $\mathcal{I}'_1, \dots, \mathcal{I}'_5 \subset \mathcal{I}'$  the dual collections. The Legendre transform  $\mathcal{L}$  yields homeomorphisms

$$\begin{aligned} \mathcal{L} : \mathcal{I}_1 &\rightarrow \mathcal{I}'_5, \\ \mathcal{L} : \mathcal{I}_2 &\rightarrow \mathcal{I}'_4, \\ \mathcal{L} : \mathcal{I}_3 &\rightarrow \mathcal{I}'_3, \\ \mathcal{L} : \mathcal{I}_4 &\rightarrow \mathcal{I}'_2, \\ \mathcal{L} : \mathcal{I}_5 &\rightarrow \mathcal{I}'_1, \end{aligned}$$

that we will all denote by  $\mathcal{L}$  with a slight abuse of notation.

Let us first describe the Legendre transform in some detail for circles of type 4 at  $z = 0$ , that is irregular circles at 0. The other cases are easily deduced from this one by changes of variable. In this paragraph, we set  $\partial := \partial_0$  the circle of directions at  $0 \in \mathbb{P}^1_z$ ,  $\mathcal{I} := \mathcal{I}_0$  the corresponding exponential local system,  $\partial'$  the circle of directions at  $\xi = \infty$  in  $\mathbb{P}^1_\xi$ , and  $\mathcal{I}' \rightarrow \partial'$  the corresponding exponential local system.

**Definition 3.3.** Let  $U \subset \partial$  be a germ of sector, and  $q_U \in \mathcal{I}(U)$  that is non-zero. For  $z \in U$ , we set

$$\phi(z) := \frac{dq}{dz}(z).$$

Choosing  $U$  small enough,  $\phi$  induces a biholomorphism between  $U$  and a sector  $U' \subset \partial'$ . For  $\xi \in U'$ , we set

$$\tilde{q}_{U'}(\xi) := q_U(\phi^{-1}(\xi)) - \xi \phi^{-1}(\xi).$$

This yields a section  $\tilde{q}_{U'} \in \mathcal{I}'(U')$ , called the Legendre transform of  $q_U$ .

The Legendre transform yields an homeomorphism between  $\mathcal{I}$  and  $\mathcal{I}'$ , the inverse homeomorphism being given by  $q_U(z) = \tilde{q}_{U'}(\phi(z)) + \phi(z)z$ . It follows from a straightforward computation that the image of an irregular circle  $I = \langle q \rangle$  by the Legendre transform is a circle  $I' = \langle \tilde{q} \rangle$  with ramification order  $r + s$  and slope  $s/(r + s)$ .

The general case of circles of type 2,3,4 is similar to the case of a circle of type 4 at  $z = 0$ . Let us summarize the properties of the Legendre transform for the different types of circles.

- Type 4: if  $a \in \mathbb{P}^1 \setminus \{\infty\}$  is a point at finite distance, the Legendre transform  $\mathcal{L}$  yields an homeomorphism between a circle of type 4 of the form  $\langle q \rangle_a$  with slope  $\alpha/\beta \neq 0$  at  $a$  and the circle of type 2  $\langle -a\xi + \tilde{q} \rangle_\infty$  with  $\tilde{q}$  of slope  $\alpha/(\alpha + \beta) < 1$ , once again determined by the same system of equations as before. The circle  $\langle -a\xi + \tilde{q} \rangle_\infty$  thus has slope 1 if  $a \neq 0$ , and slope  $< 1$  if  $a = 0$ .

$$\begin{array}{ccc} \langle q \rangle_0 & \xrightarrow[1:1]{\mathcal{L}} & \langle \tilde{q} \rangle_\infty \\ \beta:1 \downarrow \pi & & \alpha+\beta:1 \downarrow \pi \\ \partial_0 & & \partial'_\infty \end{array}$$

This is obtained in a straightforward way from the special case  $a = 0$  that we just detailed, by using the relation  $z_a = z - a$  between the local coordinate  $z_a$  at  $a$  and the coordinate  $z$  on  $\mathbb{P}^1$ . The factor  $\tilde{q}$  is given by the same system of equations as previously, with  $z$  replaced by  $z_a$ .

- Type 2: conversely, if  $\langle -ax + q \rangle_\infty$ , with  $a \in \mathbb{C}$  and  $q_1$  of slope  $\alpha/\beta < 1$  is a circle of type 2, the Legendre transform induces an homeomorphism between  $\langle q \rangle_\infty$  and a circle of type 4  $\langle \tilde{q} \rangle_a$  at the point  $a \in \mathbb{P}^1 \setminus \{\infty\} = \mathbb{C}$ , with slope  $\alpha/(\beta - \alpha)$ . We thus have

$$\begin{array}{ccc} \langle q \rangle_\infty & \xrightarrow[1:1]{\mathcal{L}} & \langle \tilde{q} \rangle_a \\ \beta:1 \downarrow \pi & & \beta-\alpha:1 \downarrow \pi \\ \partial_0 & & \partial'_\infty \end{array}$$

- Type 3: If  $\langle q \rangle$  is a circle at infinity of type type 3, of slope  $\alpha/\beta > 1$ ,  $\mathcal{L}$  induces a homeomorphism of  $\langle q \rangle$  on the circle  $\langle \tilde{q} \rangle$  of slope  $\alpha/\alpha - \beta > 1$ , of type 3. The situation is the following:

$$\begin{array}{ccc} \langle q \rangle_\infty & \xrightarrow[1:1]{\mathcal{L}} & \langle \tilde{q} \rangle_\infty \\ \beta:1 \downarrow \pi & & \alpha-\beta:1 \downarrow \pi \\ \partial_0 & & \partial'_\infty \end{array}$$

We now deal with circles of type 1 and 5. Let  $a \in \mathbb{P}^1 \setminus \{\infty\}$ , and consider the circles  $\langle 0 \rangle_a$  and  $\langle -a\xi \rangle_\infty$ . We define an homeomorphism  $\phi$  between the circles of directions  $\partial_a$  and  $\partial_\infty$  as follows, for any argument  $\theta \in \partial_a$ , we draw the half-line  $l_\theta$  starting from  $a$  with direction  $\theta$ , and let  $\phi(\theta) \in \partial_\infty$  be the direction at which  $l_\theta$  approaches  $z = \infty$ . Notice that this inverts the sense of rotation, in agreement with Malgrange [28, e.g. p. 98]. Replacing  $z$  by the dual coordinate  $\xi$  to identify  $\partial_\infty$  to  $\partial'_\infty$ , we get an homeomorphism  $\phi' : \partial_a \rightarrow \partial'_\infty$  which lifts to an homeomorphism  $\mathcal{L} : \langle 0 \rangle_a \rightarrow \langle -a\xi \rangle_\infty$ , as pictured on the diagram

$$\begin{array}{ccc} \langle 0 \rangle_a & \xrightarrow[\cong]{\mathcal{L}} & \langle -a\xi \rangle_\infty \\ \downarrow \pi & & \downarrow \pi \\ \partial_0 & \xrightarrow{\phi'} & \partial'_\infty \end{array}$$

The Legendre transform induces a transformation of the local systems on  $\mathcal{I}$ : if  $V \rightarrow \mathcal{I}$  is a local system, we define  $\tilde{V} := \mathcal{L}_*(V)$  such that the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\mathcal{L}_*} & \tilde{V} \\ \downarrow & & \downarrow \\ \mathcal{I} & \xrightarrow{\mathcal{L}} & \mathcal{I}' \end{array}$$

The previous construction accounts for what happens to the exponential factors in the Laplace method, but the square root term in the Gaussian integral remains to be taken into account. A way to do this is to define for any circle  $\langle q \rangle$  of type 2,3,4 a local system  $W_{\langle \tilde{q} \rangle} \rightarrow \langle \tilde{q} \rangle$  of one-dimensional vector spaces. In the case where  $\langle q \rangle$  is a circle of type 2 at zero, the definition can be formulated as follows.

**Definition 3.4.** Let  $\langle q \rangle_0$  be an irregular circle at  $z = 0$ ,  $\langle \tilde{q} \rangle_\infty$  its image by the Legendre transform. Let  $r = \text{ram}(q)$  and  $s = \text{Irr}(q)$ . Let  $\zeta$  a variable such that  $\zeta^{r+s} = 1/\xi$  so that the Legendre transform  $\tilde{q}$  is a polynomial in  $\zeta^{-1}$ , and the circle  $\langle \tilde{q} \rangle$  can be parametrized by the directions in the disk  $D_\zeta$ . We define  $W_{\langle \tilde{q} \rangle} \rightarrow \langle \tilde{q} \rangle$  to be the rank one local system with étalé space  $W_{\langle \tilde{q} \rangle}$  is  $\langle \tilde{q} \rangle \times \mathbb{C}$ , and a flat section of which given by any determination of the germ of the function  $\zeta \mapsto (\partial^2 q / \partial z^2)^{-1/2}(\zeta)$  on the disk  $D_\zeta$ .

**Lemma 3.5.** *If  $q$  has slope  $\alpha/\beta$ , with  $\beta = \text{ram}(q)$ , then the monodromy of  $W_{\langle \tilde{q} \rangle}$  is  $(-1)^\alpha$ .*

*Proof.* One has  $q(z) \sim z^{-\alpha/\beta}$ , hence  $\partial^2 q / \partial z^2 \sim z^{-(\alpha+2\beta)/\beta}$ . In terms of the coordinate  $\zeta$ , the Legendre transform implies  $\zeta \sim z^\beta$ , hence  $\partial^2 q / \partial z^2 \sim \zeta^{\alpha+2\beta}$ , hence  $(\partial^2 q / \partial z^2)^{-1/2} \sim \zeta^{-(\alpha+2\beta)/2}$ . Therefore, when going around  $\zeta = 0$  in  $D_\zeta$ ,  $(\partial^2 q / \partial z^2)^{-1/2}$  gets multiplied by  $\exp(-2i\pi \times \frac{\alpha+2\beta}{2}) = (-1)^\alpha$ . Notice this does not depend on the choice of determination of the square root  $(\partial^2 q / \partial z^2)$ , so that  $W_{\langle \tilde{q} \rangle}$  is well defined up to isomorphism.  $\square$

The definition of  $W_{\langle q \rangle}$  in the general case is similar, and the lemma remains true.

**Definition 3.6.** Let  $\check{V}$  be a compatible local system on  $\mathcal{I}$ , with isomorphism class  $(\check{\Theta}, \check{\mathcal{C}})$ . We set  $F \cdot \check{V} := \mathcal{L}_*(\check{V}) \otimes W$ , provided it is compatible. It is a local system on  $\mathcal{I}'$ . Let  $(\check{\Theta}', \check{\mathcal{C}}')$  be its isomorphism class. We define  $F \cdot (\check{\Theta}, \check{\mathcal{C}}) := (\check{\Theta}', \check{\mathcal{C}}')$ .

The operation  $F$  on modified formal local systems on  $\mathcal{I}$  is the counterpart at the level of formal data of the Fourier-Laplace transform of  $A_1$ -modules, that is it is such that the theorem 3.1 is true.

*Proof.* It seems that this result was first understood by Malgrange. The case of irregular circles has been independently proven by Fang [17] and Sabbah [35]. An alternative proof was also given later by Graham-Squire [19]. To see that our formulation is indeed equivalent to the one in [35], it suffices to take the monodromy of the local systems. The case of regular circles can be extracted from Malgrange [28, see p129]: the Fourier transform exchanges the local system of regular micro-solutions at finite distance and the local system of regular solutions at infinity.  $\square$

*Remark 3.1.* The definition of the formal Fourier transform  $F \cdot (\check{\Theta}', \check{\mathcal{C}}')$  might still make sense if it is not compatible. We will rather consider that it is undefined in this case. If  $F \cdot (\check{\Theta}', \check{\mathcal{C}}')$  is effective, the theorem guarantees that  $F \cdot (\check{\Theta}', \check{\mathcal{C}}')$  is also effective. If  $(\check{\Theta}', \check{\mathcal{C}}')$  is compatible but not effective,  $(\check{\Theta}', \check{\mathcal{C}}')$  may be undefined (see [2]).

If  $F \cdot (\check{\Theta}, \check{\mathcal{C}}) = (\check{\Theta}', \check{\mathcal{C}}')$  the irregular class  $\check{\Theta}'$  only depends on  $\check{\Theta}$ , so we set  $F \cdot \check{\Theta} = \check{\Theta}'$ . We will also set  $F \cdot \langle q \rangle := \mathcal{L}(\langle q \rangle) \in \pi_0(\mathcal{I})$  for any circle  $\langle q \rangle$ . If  $(\check{\Theta}, \check{\mathcal{C}})$  are effective formal data,  $\langle q \rangle$  is an active circle of  $\check{\Theta}$  if and only if  $F \cdot \langle q \rangle$  is an active circle of  $\check{\Theta}'$ . Explicitly, if  $\check{\Theta} = n_1 \langle q_1 \rangle + \dots + n_r \langle q_r \rangle$ , we have

$$\check{\Theta}' = F \cdot \check{\Theta} = n_1 F \cdot \langle q_1 \rangle + \dots + n_r F \cdot \langle q_r \rangle.$$

From the stationary phase formula we obtain a formula for the rank of the Fourier-Laplace transform of a connection with formal data  $(\Theta, \mathcal{C})$ .

**Lemma 3.7.** *Let  $(\check{\Theta}, \check{\mathcal{C}})$  be modified formal data, and  $(\check{\Theta}', \check{\mathcal{C}}') := F \cdot (\check{\Theta}, \check{\mathcal{C}})$ . The rank of a connection with modified formal data  $(\check{\Theta}', \check{\mathcal{C}}')$ , if it exists, is*

$$(3.1) \quad \sum_k \left( m_k + \sum_{i=1}^{r_k} n_i^{(k)} (\alpha_i^{(k)} + \beta_i^{(k)}) \right) + \sum_{i, \text{slope} > 1} n_i^{(\infty)} (\alpha_i^{(\infty)} - \beta_i^{(\infty)}),$$

where the second sum is on all active circles  $\langle q_i^{(\infty)} \rangle_\infty$  at infinity with slope  $\alpha_i^{(\infty)} / \beta_i^{(\infty)} > 1$ .

*Proof.* From the stationary phase formula the active circles of  $V'$  at infinity are the following:

- Images by Legendre transform of irregular circles  $\langle q^{(k)i} \rangle_{a_k}$ ,  $k = 1, \dots, m$  i.e circles  $\langle -a_k \xi + \tilde{q}^{(k)j} \rangle_\infty$ , with ramification  $\alpha_i^{(k)} + \beta_i^{(k)}$  and multiplicity  $n_i^{(k)}$ .

- Images of the tame circles  $\langle 0 \rangle_{a_k}$ ,  $k = 1, \dots, m$ : pure circles  $\langle -a_k \xi \rangle_\infty$ , with multiplicity  $m_k$ .
- Images of circles of slope  $> 1$  at infinity among the  $\langle q_i^{(\infty)} \rangle$ ,  $i = 1, \dots, r_\infty$ , of the form  $\langle \tilde{q}_i^{(\infty)} \rangle$ , with slope  $\alpha_i^{(\infty)} - \beta_i^{(\infty)}$  and multiplicity  $n_i^{(\infty)}$ .

Adding the contribution of those circles to the rank gives the result.  $\square$

*Remark 3.2.* Notice that this is consistent with the formula given by Malgrange [28, p. 79] for the rank of the Laplace transform of modules over the Weyl algebra.

**3.2. Symplectic transformations.** The Fourier-Laplace transform is part of a larger group of transformations acting on modules over the Weyl algebra. Indeed, to any matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

in  $SL_2(\mathbb{C})$ , we can associate an automorphism of the Weyl algebra  $A_1 = \mathbb{C}[z]\langle \partial_z \rangle$  given by  $z \mapsto az + b\partial_z$ ,  $\partial_z \mapsto cz + d\partial_z$ . This induces an action of the group  $SL_2(\mathbb{C})$  of symplectic transformations on modules over the Weyl algebra (see [28]).

The group of symplectic transformations is generated by three types of elementary transformations:

- The Fourier-Laplace transform  $F$ , corresponding to the matrix  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ .
- Twists at infinity  $T_\lambda$ , for  $\lambda \in \mathbb{C}$ , corresponding to the matrix  $\begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix}$ .
- Scalings  $S_\lambda$ , for  $\lambda \in \mathbb{C}^*$ , corresponding to the matrix  $\begin{pmatrix} \lambda^{-1} & 0 \\ 0 & \lambda \end{pmatrix}$ .

The geometric interpretation of twists and scalings is the following. The twist  $T_\lambda$  corresponds to taking the tensor product with the rank one module  $(\mathbb{C}[z], \partial_z + \lambda z)$ , and the scaling  $S_\lambda$  corresponds to do the change of variable  $z \mapsto z/\lambda$  on  $\mathbb{P}^1$ .

As for the Fourier transform, any element of  $SL_2(\mathbb{C})$  induces a transformation on connections on Zariski open subsets on the affine line, and a corresponding formal transformation (that we will also denote by  $A$ ) on compatible formal data, such that the diagram

$$\begin{array}{ccc} (E, \nabla) & \xrightarrow{A} & (E', \nabla') \\ \downarrow & & \downarrow \\ (\check{\Theta}, \check{\mathcal{C}}) & \xrightarrow{A} & (\check{\Theta}', \check{\mathcal{C}}') \end{array}$$

commutes. To show this, it is enough to check this for elementary transformations. We have already dealt with the case of the Fourier transform, and for the twists and scalings we have:

**Proposition 3.8.** *Let  $(E, \nabla)$  a connection on a Zariski open subset of the affine line,  $\check{V} \rightarrow \mathcal{I}$  its modified formal local system, and  $(\check{\Theta}, \check{\mathcal{C}})$  its isomorphism class.*

- Let  $t_\lambda : \mathcal{I}_\infty \rightarrow \mathcal{I}_\infty$  be the homeomorphism of  $\mathcal{I}_\infty$  defined by  $q_U(1/z) \mapsto q_U(1/z) + \frac{\lambda}{2}z^2$ , where  $q_U$  is a section of  $\mathcal{I}_\infty$  over an open sector  $U \subset \partial_\infty$ . We extend  $t_\lambda$  to an homeomorphism of  $\mathcal{I}$  by having it act trivially on  $\bigsqcup_{a \in \mathbb{C}} \mathcal{I}_a$ . Then the modified formal local system associated to  $T_\lambda \cdot (E, \nabla)$  is isomorphic to  $(t_\lambda)_* \check{V}$ .
- For  $\lambda \in \mathbb{C}^*$ , let  $s_\lambda : \mathcal{I} \rightarrow \mathcal{I}$  be the homeomorphism of  $\mathcal{I}$  defined in the following way: for  $a \in \mathbb{C}$ , if  $q_{U_a}(z - a)$  is a section of  $\mathcal{I}_a$  on the open sector  $U \subset \partial_a$ , its image by  $s_\lambda$  is the section  $q_{U_a/\lambda}(\lambda(z - a/\lambda))$  over the sector  $U_a/\lambda \subset \partial_{a/\lambda}$  which is the image of  $U$  by  $z \mapsto z/\lambda$ . If  $q_U(1/z)$  is a section of  $\mathcal{I}_\infty$ , its image is  $q_{U/\lambda}(\frac{1}{\lambda z})$ . The modified formal local system associated to  $S_\lambda \cdot (E, \nabla)$  is isomorphic to  $(s_\lambda)_* \check{V}$ .

We thus define  $T_\lambda \cdot (\check{\Theta}, \check{\mathcal{C}})$  as the isomorphism class of  $(t_\lambda)_* \check{V}$  for  $\lambda \in \mathbb{C}$ , and  $S_\lambda \cdot (\check{\Theta}, \check{\mathcal{C}})$  as the isomorphism class of  $(s_\lambda)_* \check{V}$  for  $\lambda \in \mathbb{C}^*$ .

*Proof.* The twist  $T_\lambda$  consists of tensoring  $(E, \nabla)$  with the rank one connection  $(\mathcal{O}, d + \lambda z dz)$ , having a third order pole at infinity and no other singularity. Therefore,  $T_\lambda$  only modifies the formal data of  $\nabla$  at infinity. Since  $\lambda z dz = -\lambda \frac{dz_\infty}{z_\infty^3}$ , if  $A$  is the matrix of the formalization at infinity  $\widehat{\nabla}_\infty$  in some basis, i.e.  $\widehat{\nabla}_\infty$  corresponds  $\partial_{z_\infty} + A$  in this basis, then the matrix of the formalization of  $T_\lambda \cdot \nabla$  at infinity in the same basis is  $A - \frac{\lambda}{z_\infty^3}$ . Integrating this, we obtain that the exponential factors at infinity of  $T_\lambda \cdot (E, \nabla)$  are obtained from the exponential factors of  $(E, \nabla)$  by adding  $\frac{\lambda}{2z_\infty^2}$ . The formal monodromies do not change.

The scaling  $S_\lambda$  corresponds to making the change of variable  $z' = \lambda z$ . The singularities of  $S_\lambda \cdot (E, \nabla)$  are thus the images of the ones of  $(E, \nabla)$  by  $z \mapsto z/\lambda$ , and the exponential factors of  $S_\lambda \cdot (E, \nabla)$  are obtained from the ones of  $(E, \nabla)$  by expressing them as a function of  $z'$ , i.e. by substituting  $z$  with  $\lambda z'$ .  $\square$

As for the Fourier transform, if  $A \cdot (\check{\Theta}, \check{\mathcal{C}}) = (\check{\Theta}', \check{\mathcal{C}}')$  the irregular class  $\check{\Theta}'$  only depends on  $\check{\Theta}$ , so we set  $A \cdot \check{\Theta} = \check{\Theta}'$ . For any circle  $\langle q \rangle$ , there is also a well-defined circle  $A \cdot \langle q \rangle \in \pi_0(\mathcal{I})$  such that, if  $(\check{\Theta}, \check{\mathcal{C}})$  are effective formal data,  $\langle q \rangle$  is an active circle of  $\check{\Theta}$  if and only if  $A \cdot \langle q \rangle$  is an active circle of  $\check{\Theta}'$ . If  $\check{\Theta} = n_1 \langle q_1 \rangle + \dots + n_r \langle q_r \rangle$ , we have

$$\check{\Theta}' = A \cdot \check{\Theta} = n_1 A \cdot \langle q_1 \rangle + \dots + n_r A \cdot \langle q_r \rangle.$$

The circle  $A \cdot \langle q \rangle$  can be determined in practice by factorizing  $A$  as a product of elementary operations.

#### 4. INVARIANCE OF THE DIAGRAM FOR ONE IRREGULAR SINGULARITY

In this section, we discuss the properties of the diagram associated by [12] to an irregular class at infinity. We show that the diagram is invariant under Fourier-Laplace transform, and under  $SL_2(\mathbb{C})$  transformations. To do this, we compute an explicit formula for the number of edges and loops of the diagram, as well as a formula for the terms appearing in the Legendre transform of an exponential factor.

**4.1. Diagram for one irregular singularity.** Let  $(E, \nabla)$  be an algebraic connection on the affine line. The only singularity is at infinity. Its formal local system  $V^0 \rightarrow \mathcal{I}$  (which is also the modified formal local system) has support on  $\mathcal{I}_\infty$ . Let  $(\Theta_\infty, \mathcal{C}_\infty)$  denote its formal data. Let us fix a direction  $d \in \partial_i nfty$ . We set  $G := GL(V_d)$  and  $H := \text{GrAut}(V_d^0)$  the set of graded automorphisms of  $V_d$ . Recall [11] that the conjugacy class  $\mathcal{C}_\infty$  of the monodromy of  $V^0 \rightarrow \partial_\infty$  is an element of a twist  $H(\partial)$  of  $H$ . The wild character variety is in this case a multiplicative symplectic quotient

$$(4.1) \quad \mathcal{M}_B(E, \nabla) = \mathcal{M}_B(\Theta_\infty, \mathcal{C}_\infty) = \text{Hom}_S(V^0) //_{\mathcal{C}} H,$$

where

$$(4.2) \quad \text{Hom}_S(\Theta_\infty) = \mathcal{A}(V^0) // G,$$

is a twisted quasi-Hamiltonian  $H$ -space, itself obtained by symplectic reduction with respect to  $G$  from the twisted quasi-Hamiltonian  $G \times H$ -space

$$(4.3) \quad \mathcal{A}(V^0) = H(\partial) \times G \times \prod_{d \in \mathbb{A}} \text{Sto}_d(V^0).$$

Let us denote by  $\langle q_1 \rangle, \dots, \langle q_r \rangle$  the active circles, and  $\beta_i := \text{ram}(q_i)$ . In this case, recall that we have from [12] the definition of a diagram, whose core is given by:

**Definition 4.1.** The *core diagram*  $\Gamma_c^{BY}(\Theta_\infty)$  associated to the irregular class  $\Theta_\infty$  has a set of nodes  $N$  given by the set of active circles  $\langle q_1 \rangle, \dots, \langle q_r \rangle$ , and for  $i, j = 1, \dots, r$ , the number of arrows  $B_{ij}$  between  $\langle q_i \rangle$  and  $\langle q_j \rangle$  is given by

- If  $i \neq j$  then

$$(4.4) \quad B_{ij} = A_{ij} - \beta_i \beta_j,$$

where  $A_{ij} = \text{Irr}(\text{Hom}(\langle q_i \rangle, \langle q_j \rangle))$ .

- If  $i = j$ , the number of oriented loops at  $\langle q_i \rangle$  is given by

$$(4.5) \quad B_{ii} = A_{ii} - \beta_i^2 + 1.$$

The numbers of edges/loops may be negative. One has  $B_{ij} = B_{ji}$  and we will show that  $B_{ii}$  is always an even number, so we can group the arrows two by two to get an unoriented diagram. The motivation for this definition comes from counting the number of appearances of blocks between graded parts of  $V^0$  associated to the different active circles in the explicit presentation of the wild character variety. The positive terms in  $B_{ij}$  correspond to the matrix blocks in  $\mathcal{A}(V^0)$ , and the negative terms in the  $B_{ij}$  correspond to the relations given by the quasi-Hamiltonian reduction.

The core diagram only depends on the irregular class  $\Theta_\infty$ , but it does not take into account the monodromies of the active circles. This can be done by adding legs to the core diagram encoding the conjugacy classes of the monodromies. For each circle  $I \in \pi_0(\mathcal{I}_\infty)$  in the support of  $\Theta_\infty$ , with multiplicity  $n(I)$ , let  $\mathcal{C}_I \subset GL_{n(I)}(\mathbb{C})$  be the corresponding conjugacy class. Recall e.g. from [8, def. 9.2] that a choice of marking of  $\mathcal{C}_I$  determines a leg  $\mathbb{L}_I$ , which comes together with its dimension vector  $\mathbf{d}_L \in \mathbb{Z}^{N_L}$ , and vector of labels  $\mathbf{q}_L \in (\mathbb{C}^*)^{N_L}$ , where  $N_L$  denotes the set of vertices of  $\mathbb{L}$ .

**Definition 4.2.** Let us choose for each active circle  $I$  a special marking of the conjugacy class  $\mathcal{C}_I$ . The *full diagram*  $\Gamma^{BY}(\Theta_\infty, \mathbf{C}_\infty)$  associated to  $(\Theta_\infty, \mathbf{C}_\infty)$  is the diagram obtained by gluing for each active circle  $I$  the leg  $\mathbb{L}(I)$  to the corresponding vertex  $v_I$  of  $\Gamma_c^{BY}(\Theta_\infty, \mathbf{C}_\infty)$ .

The full diagram inherits in this way a dimension vector  $\mathbf{d} \in \mathbb{Z}^N$ , as well as a vector of multiplicative labels  $\mathbf{q} \in (\mathbb{C}^*)^N$ .

This recipe only enables us to define a diagram for connections with only one singularity at infinity. Our goal is to define a diagram for the general case, that is for connections with an arbitrary number of singularities. The task is not straightforward: although it is possible to draw a diagram corresponding to the formal data at each singularity, it is not clear that there is a natural way to somehow glue those diagrams together to define a global diagram having good properties. In other words, the way in which the different singularities should “interact” is not clear.

The idea, already present in [6], is thus to use the Fourier transform to reduce to the case where all active circles are at infinity, where we know how to draw the diagram. Indeed, the Fourier transform sends to infinity all active circles at finite distance. However, the Fourier transform also sends to finite distance some of the active circles at infinity, and we are back to the problem we started with. For this reason, we need before taking the Fourier transform to apply some operation to prevent the active circles at infinity to go to finite distance. This can be achieved by using more general  $SL_2(\mathbb{C})$  transformations on modules over the Weyl algebra. As we shall see, any matrix in an open dense subset of  $SL_2(\mathbb{C})$  sends all active circles to infinity.

We thus want to define the diagram associated to a connection as the diagram associated to its image under a generic  $SL_2(\mathbb{C})$  transformation. For this to be well defined however, the diagram must not depend on the choice of  $SL_2(\mathbb{C})$  transformation to bring all active circles to infinity, in other words the diagram has to be invariant under  $SL_2(\mathbb{C})$  transformations. The goal of this section is to show that it is indeed the case. The main step will be to show that the diagram is invariant under Fourier transform.

**4.2. An explicit formula for the number of edges.** Let  $\langle q \rangle$  and  $\langle q' \rangle$  be two exponential factors at infinity, with ramification orders  $\beta$ , and  $\beta'$ , with slopes  $\alpha/\beta$  and  $\alpha'/\beta'$ . We set

$$q = \sum_{j=0}^p b_j z_\infty^{-\alpha_j/\beta}, \quad q' = \sum_{j=0}^{p'} b'_j z_\infty^{-\alpha'_j/\beta'}$$

the expression for  $q$  and  $q'$  where we write only the monomials with non-zero coefficients, i.e.  $b_j \neq 0$  and  $b'_j \neq 0$ ,  $\alpha_0 > \dots > \alpha'_p$  and  $\alpha'_0 > \dots > \alpha'_p$ . Because of ramification, a monomial appearing both in  $q$  and  $q'$  can give rise to Stokes arrows associated to the difference  $q - q'$ , since there are several leaves on the covers  $I := \langle q \rangle$  and  $I' := \langle q' \rangle$ . The leaves of  $\langle q \rangle$  correspond to the

images  $q$  under the action of the Galois group isomorphic to  $\mathbb{Z}/\beta\mathbb{Z}$  arising from ramification, i.e. to the polynomials

$$q_i = \sum_{j=0}^p b_j \omega^{-\alpha_j} z_\infty^{-\alpha_j/\beta}, \quad i = 0, \dots, \beta - 1,$$

with  $\omega = e^{2i\pi/\beta}$ . In a similar way, the leaves of  $\langle q' \rangle$  correspond to the polynomials

$$q'_i = \sum_{j=0}^{p'} b'_j \omega'^{-\alpha'_j} z_\infty^{-\alpha'_j/\beta'}, \quad i = 0, \dots, \beta' - 1,$$

with  $\omega' = e^{2i\pi/\beta'}$ .

Let  $r$  be the biggest integer such that  $\sum_{j=0}^{r-1} b_j z_\infty^{-\alpha_j/\beta}$  and  $\sum_{j=0}^{r-1} b'_j z_\infty^{-\alpha'_j/\beta'}$  have the same Galois orbit.

We set

$$q_c := \sum_{j=0}^{r-1} b_j z_\infty^{-\alpha_j/\beta}, \quad q'_c := \sum_{j=0}^{r-1} b'_j z_\infty^{-\alpha'_j/\beta'},$$

and

$$q_d := \sum_{j=r}^p b_j z_\infty^{-\alpha_j/\beta}, \quad q'_d := \sum_{j=r}^{p'} b'_j z_\infty^{-\alpha'_j/\beta'},$$

We get a decomposition

$$q = q_c + q_d, \quad q' = q'_c + q'_d,$$

of  $q$  and  $q'$  as the sum of a *common part*  $q_c$  and a *different part*  $q_d$ . Replacing  $q$  or  $q'$  by another element of their Galois orbit if necessary, we may assume that  $q_c = q'_c$ . In the case where  $q$  and  $q'$  do not have the same leading term, then  $r = 0$  and we have  $q_c = q'_c = 0$  and  $q = q_d$ ,  $q' = q'_d$ . In particular, this happens when  $q$  and  $q'$  do not have the same slope. Otherwise,  $q$  and  $q'$  have a non-zero common part, in particular they have the same slope  $\alpha_0/\beta = \alpha'_0/\beta'$ . The slope of  $q_d$  is  $\alpha_r/\beta$ , and the slope of  $q'_d$  is  $\alpha'_r/\beta'$ . We use the usual notation  $(\cdot, \cdot)$  to denote the greatest common divisor. With those notations now set, we are in position to state the formula for the number of edges between  $I$  and  $I'$ .

**Lemma 4.3.** • Assume that  $\alpha_r/\beta \geq \alpha'_r/\beta'$ . Then the number of edges between  $I = \langle q \rangle$  and  $I' = \langle q' \rangle$  is

$$B_{I,I'} = (\beta' - (\alpha'_0, \beta'))\alpha_0 + ((\alpha'_0, \beta') - (\alpha'_0, \alpha'_1, \beta'))\alpha_1 + \dots + ((\alpha'_0, \dots, \alpha'_{r-2}, \beta') - (\alpha'_0, \dots, \alpha'_{r-1}, \beta'))\alpha_{r-1} + (\alpha'_0, \dots, \alpha'_{r-1}, \beta')\alpha_r - \beta\beta'.$$

• In particular, if  $q$  and  $q'$  have no common parts and  $\alpha/\beta \geq \alpha'/\beta'$ , then

$$B_{I,I'} = \beta'(\alpha - \beta).$$

We have a similar result for the number of loops at a circle  $\langle q \rangle$ .

**Lemma 4.4.** Let  $q = \sum_{j=0}^p b_j z_\infty^{-\alpha_j/\beta}$  be an exponential factor of slope  $\alpha_0/\beta > 1$  as before.

• One has

$$(4.6) \quad B_{I,I} = (\beta - (\alpha_0, \beta))\alpha_0 + ((\alpha_0, \beta) - (\alpha_0, \alpha_1, \beta))\alpha_1 + \dots + ((\alpha_0, \dots, \alpha_{p-1}, \beta) - (\alpha_0, \dots, \alpha_p, \beta))\alpha_p - \beta^2 + 1.$$

• In particular, if  $(\alpha, \beta) = 1$ , then we have

$$(4.7) \quad B_{I,I} = (\beta - 1)(\alpha - \beta - 1).$$

The proofs are a bit lengthy and are given in appendix.

**Lemma 4.5.** The integer  $B_{I,I}$  is even.

*Proof.* Let us first assume that  $\beta$  is odd. Then all greatest common divisors appearing in the formula for  $B_{I,I}$  are odd, so differences between two consecutive g.c.d.s are even, and all terms involving those differences are even. The sum of the remaining terms is  $-\beta^2 + 1$  which is even, so the result follows.

Now, let us consider the case when  $\beta$  is even. Let us consider the sequence of greatest common divisors  $\beta, (\alpha_0, \beta), \dots, (\alpha_0, \dots, \alpha_p, \beta)$ . The first element  $\beta$  is even, the last one  $(\alpha_0, \dots, \alpha_p, \beta)$  is odd, and the sequence consists first of even integers until  $(\alpha_0, \dots, \alpha_{k_0}, \beta)$  where  $k_0$  is the smallest index  $k$  such that  $\alpha_k$  is odd. Starting from this element, all g.c.d.s in the sequence are odd. As a consequence, all the terms involving differences of g.c.d.s in the formula for  $B_{I,I}$  are even except  $((\alpha_0, \dots, \alpha_{k_0-1}, \beta) - (\alpha_0, \dots, \alpha_{k_0}, \beta))\alpha_{k_0}$  which is odd. The sum of the remaining terms  $-\beta^2 + 1$  is odd, and the conclusion follows.  $\square$

This proves what was a statement in [12, p.3], and guarantees that we have a diagram with unoriented edges.

**4.3. Form of the Legendre transform.** In this paragraph, we compute the Legendre transform of an exponential factor  $q$  at infinity as explicitly as possible. We determine the monomials appearing (with non-zero coefficients) in the Legendre transform.

**Proposition 4.6.** *Let  $q = \sum_{j=0}^p b_j z^{\alpha_j/\beta}$ , with  $b_j \neq 0$ , be an exponential factor at infinity with ramification  $\beta$ . We set  $\alpha := \alpha_0$ , so the slope of  $q$  is  $\alpha/\beta > 1$ . Then the exponents of  $\xi$  possibly appearing with non-zero coefficients in its Legendre transform  $\tilde{q}$  are of the form  $\frac{\alpha - k_1(\alpha - \alpha_1) - \dots - k_p(\alpha - \alpha_p)}{\alpha - \beta}$ , with  $k_1, \dots, k_p \geq 0$ . More precisely, if we set  $E := \{\gamma \in \mathbb{N} \mid \exists k_1, \dots, k_p \in \mathbb{N}, \gamma = k_1(\alpha - \alpha_1) + \dots + k_p(\alpha - \alpha_p)\}$  the Legendre transform has the form*

$$(4.8) \quad \tilde{q}(\xi) = \sum_{\substack{\gamma \in E \\ \alpha - \gamma > 0}} \tilde{b}_\gamma \xi^{\frac{\alpha - \gamma}{\alpha - \beta}},$$

where the sum is restricted to the terms such that the exponent  $\frac{\alpha - \gamma}{\alpha - \beta}$  is positive. Furthermore, the coefficients  $\tilde{b}_{(\alpha - \alpha_i)}$  are non-zero for  $i \geq 1$ .

We refer the reader to the appendix for the proof.

**4.4. Invariance of the diagram under Fourier-Laplace transform.** We now prove the invariance of the diagram under Fourier transform.

**Theorem 4.7.** *Let  $\Theta_\infty : \pi_0(\mathcal{I}_\infty) \rightarrow \mathbb{N}$  an irregular class infinity, such that the slopes of all active circles are  $> 1$ . Then the core diagram  $\Gamma_c^{BY}(\Theta_\infty)$  is invariant under Fourier-Laplace transform.*

*Proof.* The proof comes from combining lemma 4.3 giving the number of edges of the diagram, and lemma 4.6 giving the structure of the Legendre transform. This allows us to compute the number of edges in the diagrams  $\Gamma_c(\Theta_\infty)$  and  $\Gamma_c(\Theta'_\infty)$ , where  $\Theta'_\infty := F \cdot \Theta_\infty$ , and check they are the same. Keeping the same notations as before, let  $I := \langle q \rangle$  and  $I' := \langle q' \rangle$  be two distinct active circles of  $\Theta_\infty$ . We want to compute the number of edges between their images  $\tilde{I} := \langle \tilde{q} \rangle$  and  $\tilde{I}' := \langle \tilde{q}' \rangle$ . Lemma 4.6 implies that  $\tilde{q}$  is of the form

$$\tilde{q} = \sum_{\gamma \in E} \tilde{b}_\gamma \xi_\infty^{-\frac{\alpha - \gamma}{\alpha - \beta}},$$

with  $E = \{\gamma \in \mathbb{N} \mid \gamma < \alpha, \exists k_1, \dots, k_p \geq 0, \gamma = k_1(\alpha_0 - \alpha_1) + \dots + k_p(\alpha_0 - \alpha_p)\}$ , and similarly

$$\tilde{q}' = \sum_{\gamma' \in E'} \tilde{b}'_{\gamma'} \xi_\infty^{-\frac{\alpha' - \gamma'}{\alpha' - \beta'}},$$

with  $E' = \{\gamma' \in \mathbb{N} \mid \gamma' < \alpha', \exists k_1, \dots, k_p \geq 0, \gamma' = k_1(\alpha'_0 - \alpha'_1) + \dots + k_p(\alpha'_0 - \alpha'_p)\}$ . Let us denote by  $0 = \gamma_0 < \dots < \gamma_N$  the distinct elements of  $E$  and  $0 = \gamma'_0 < \dots < \gamma'_{N'}$  those of  $E'$ . We first determine the common parts and the different parts of  $\tilde{q}$  and  $\tilde{q}'$ . The integers  $\gamma$  and  $\gamma'$  containing only terms common to  $q$  and  $q'$ , i.e. of the form  $\gamma = k_1(\alpha_0 - \alpha_1) + \dots + k_{r-1}(\alpha_0 - \alpha_{r-1})$  and



$\gamma' = k_1(\alpha'_0 - \alpha'_1) + \dots + k_{r-1}(\alpha'_0 - \alpha'_{r-1})$ , belong to the part common to  $\tilde{q}$  and  $\tilde{q}'$ . Let us denote by  $E_c \subset E$  and  $E'_c \subset E'$  the corresponding subsets. Furthermore, the respective leading terms of the different parts of  $\tilde{q}_d$  and  $\tilde{q}'_d$  correspond to the first integers  $\gamma$  where the factors  $(\alpha - \alpha_r)$  and  $(\alpha' - \alpha'_r)$  appear, which are  $\alpha - \alpha_r =: \gamma_R$ , and  $\alpha' - \alpha'_{r+1} =: \gamma'_R$  with  $R \leq N$ . It follows that  $\tilde{q}_d$  is of degree  $\frac{\alpha_r}{\alpha - \beta}$  and  $\tilde{q}'_d$  is of degree  $\frac{\alpha'_r}{\alpha' - \beta'}$ . Since we have assumed  $\alpha_r/\beta > \alpha'_r/\beta'$ , we have  $\frac{\alpha_r}{\alpha - \beta} > \frac{\alpha'_r}{\alpha' - \beta'}$ . From lemma 4.3, the number of edges between  $\langle \tilde{q} \rangle$  and  $\langle \tilde{q}' \rangle$  is

$$(4.9) \quad B_{\tilde{I}, \tilde{I}'} = ((\alpha' - \beta') - p'_0)\alpha_0 + \sum_{k=1}^{R-1} (p'_{k-1} - p'_k)(\alpha - \gamma_k) + p_R(\alpha - \gamma_R) - (\alpha - \beta)(\alpha' - \beta'),$$

where  $p'_k = (\alpha'_0, \dots, \alpha'_0 - \gamma_k, \alpha' - \beta')$ .

In this expression, let us examine the g.c.d.s  $p'_k$  and the differences  $p'_{k-1} - p'_k$ . Notice that one has  $p'_{k-1} = p'_k$  as soon as any decomposition  $\gamma_k = k_1(\alpha'_0 - \alpha'_1) + \dots + k_p(\alpha'_0 - \alpha'_p)$  of  $\gamma_k$  only contains factors  $(\alpha - \alpha_j)$  already present in some  $p_l$  with  $l \leq k - 1$ . In other words, we can have  $p'_{k-1} > p'_k$  only when in  $\gamma_k$  a new factor  $(\alpha - \alpha_j)$  appears, that is when  $\gamma_k = \alpha - \alpha_j$  for some  $j = 1, \dots, p$ . The distinct values of  $p'_k$  are thus the g.c.d.s  $(\alpha'_0, \dots, \alpha'_l, \alpha' - \beta') = (\alpha'_0, \dots, \alpha'_l, \beta')$ . Therefore, the non-zero terms involving g.c.d.s in the formula for  $B_{\langle \tilde{q} \rangle, \langle \tilde{q}' \rangle}$  are exactly the same as those for  $B_{I, I'}$ . The sum of the remaining terms is  $\alpha'\beta - \beta\beta' = \beta(\alpha' - \beta')$  for  $B_{I, I'}$ , and  $(\alpha' - \beta')\alpha_0 - (\alpha - \beta)(\alpha' - \beta') = \beta(\alpha' - \beta')$  for  $B_{\tilde{I}, \tilde{I}'}$ : they are also the same. Finally, we have the equality  $B_{I, I'} = B_{\tilde{I}, \tilde{I}'}$  of the numbers of edges.

It remains to deal with the case of loops at  $\langle q \rangle$  and  $\langle \tilde{q} \rangle$ . In a similar way as the previous case, the non-zero differences of g.c.d.s appearing in the formula for  $B_{\tilde{I}, \tilde{I}'}$  are exactly the same as those appearing in the formula for  $B_{I, I'}$ . The sum of the remaining terms for  $B_{I, I'}$  is  $\beta(\alpha - \beta) + 1$ , and this expression is invariant under Fourier transform since  $\alpha \mapsto \alpha$  and  $\beta \mapsto \alpha - \beta$ . We thus have  $B_{I, I'} = B_{\tilde{I}, \tilde{I}'}$ , which completes the proof of the invariance of the diagram.  $\square$

**4.5. Invariance of the diagram under symplectic transformations.** The goal of this section is to show:

**Theorem 4.8.** *Let  $\Theta_\infty$  be an irregular class at infinity. There is an open dense subset  $S_V \subset SL_2(\mathbb{C})$  such that for all  $A \in S_V$ ,  $A \cdot \Theta_\infty$  only has a singularity at infinity. Furthermore, we have  $\Gamma_c^{BY}(\Theta_\infty) = \Gamma_c^{BY}(A \cdot \Theta_\infty)$ .*

To prove this, the idea is to decompose the elements of  $SL_2(\mathbb{C})$  as products of elementary operations.

**Lemma 4.9.** *Let  $\Theta_\infty$  be an irregular class at infinity. Then*

- For  $\lambda \in \mathbb{C}$ ,  $\Gamma_c(T_\lambda \cdot \Theta_\infty) = \Gamma_c^{BY}(\Theta_\infty)$ .
- For  $\lambda \in \mathbb{C}^*$ ,  $\Gamma_c(S_\lambda \cdot \Theta_\infty) = \Gamma_c^{BY}(\Theta_\infty)$ .

*Proof.* The twist  $T_\lambda$  induces via  $t_\lambda$  a bijection between the active circles of  $\Theta_\infty$  and  $T_\lambda \cdot \Theta_\infty$ . The differences of exponential factors  $q_U(z_\infty) - q'_U(z_\infty)$  are invariant under a twist  $T_\lambda$ , so the number of Stokes arrows between two active circles and their images by  $t_\lambda$  are the same. This proves the first part of the lemma. In a similar way,  $S_\lambda$  induces via  $s_\lambda$  a bijection between the active circles of  $\Theta_\infty$  and  $S_\lambda \cdot \Theta_\infty$ . The degree of a difference  $q_U(z_a) - q'_U(z_a)$  of exponential factors at  $a \in \mathbb{C}$  is the same as its image  $q_U(\lambda(z_a/\lambda)) - q'_U(\lambda(z_a/\lambda))$  by  $s_\lambda$ . This is also true for exponential factors at infinity. Therefore, the number of Stokes arrows between two active circles and their images by  $s_\lambda$  are the same, and the conclusion follows.  $\square$

Since we have already seen that the Fourier transform preserves the diagram when all circles are of slope  $> 1$ , i.e. when  $F \cdot \Theta_\infty$  only has a singularity at infinity, all elementary transformations preserve the core diagram. However, deducing the theorem from this by factoring matrices of  $SL_2(\mathbb{C})$  as products of elementary transformations is not entirely straightforward. Indeed, if  $A \in SL_2(\mathbb{C})$  admits a factorization  $A = A_k \dots A_1$  as a product of elementary transformations, it may occur that for some  $1 \leq i \leq k$ ,  $A_i \dots A_1 \cdot \Theta_\infty$  has singularities at finite distance. Therefore,

we have to show that  $A$  always admits a factorization such that this doesn't happen. We will do this in several steps.

**Lemma 4.10.** *Let  $\Theta_\infty$  be an irregular class at infinity. There exists  $A \in SL_2(\mathbb{C})$  such that all active circles of  $A \cdot \Theta_\infty$  have slope no greater than 2. More precisely, there exists such a matrix of the form  $FT_\lambda$ , for some  $\lambda \in \mathbb{C}$ .*

*Proof.* Let  $\langle q \rangle_\infty$  be an irregular circle at infinity. Its image by  $T_\lambda$  is  $\langle q + \frac{\lambda}{2}z^2 \rangle$ . The term  $\frac{\lambda}{2}z^2$  is of slope 2. There are thus two possibilities: either  $q$  has slope strictly greater than 2 and  $\text{slope}(q) = \text{slope}(q + \frac{\lambda}{2}z^2)$ , or  $q$  has slope less than 2 and  $\text{slope}(q + \frac{\lambda}{2}z^2) = 2$ , if we choose  $\lambda$  such that  $-\lambda/2$  is not equal to the coefficient of slope 2 of  $q$ . We then apply the Fourier transform. In the first case, if we denote  $\text{slope}(T_\lambda \cdot \langle q \rangle) = p/q > 2$ , with  $p, q \in \mathbb{N}$ , one has  $p - q > q$ , so from the stationary phase formula  $\text{slope}(FT_\lambda \cdot \langle q \rangle) = \frac{p}{p-q} < 2$ . In the second case we have  $\text{slope}(FT_\lambda \cdot \langle q \rangle) = \text{slope}(T_\lambda \cdot \langle q \rangle) = 2$ . As a consequence, if we choose  $\lambda$  such that  $-\lambda/2$  is not equal to any of the coefficients of slope 2 of the active circles of  $\Theta_\infty$ , then  $FT_\lambda \cdot \Theta_\infty$  only has active circles at infinity of slope no greater than 2.  $\square$

Let  $\langle q \rangle$  be a circle at infinity of slope  $\leq 2$ . It is clear that this circle remains at infinity when a twist or a scaling is applied. It remains at infinity under Fourier transform if and only if its slope is strictly greater than 1. In particular, it remains at infinity if its coefficient of slope 2 is non-zero.

**Proposition 4.11.** *Let  $q$  be an exponential factor at infinity of slope  $\leq 2$ . We write  $q = -\frac{\lambda}{2}z^2 + q_{<2}$  with  $q_{<2}$  of slope  $< 2$ . Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{C})$ . Let  $\langle q' \rangle$  be the image of  $\langle q \rangle_\infty$  by  $A$ . Then  $\langle q' \rangle$  is also at infinity unless  $\text{slope}(q_{<2}) \leq 1$  and  $c\lambda + d = 0$ . In this case, writing  $q' = -\frac{\lambda'}{2}z^2 + q'_{<2}$  with  $q'_{<2}$  of slope  $< 2$ , the coefficients  $\lambda$  and  $\lambda'$  are related by*

$$(4.10) \quad \lambda' = \frac{a\lambda + b}{c\lambda + d}.$$

To prove this, we begin with the case of elementary transformations.

**Lemma 4.12.** *The proposition is true if  $A$  is an elementary transformation*

*Proof.* The twist  $T_\mu = \begin{pmatrix} 1 & \mu \\ 0 & 1 \end{pmatrix}$  acts on  $q$  according to

$$T_\mu(q) = q - \frac{\mu}{2}z^2 = -\frac{\lambda + \mu}{2}z^2 + q_{<2}.$$

This amounts to  $\lambda \mapsto \lambda + \mu$ , so the property is verified. The scaling  $S_\mu = \begin{pmatrix} \mu^{-1} & 0 \\ 0 & \mu \end{pmatrix}$  sends  $q$  to

$$S_\mu(q) = q(z/\mu) = -\frac{\lambda\mu^2}{2}z^2 + q_{<2}(z/\mu).$$

This amounts to  $\lambda \mapsto \lambda\mu^2$ , so the property is again verified. The Fourier transform  $F = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  acts on  $q$  according to the Legendre transform: computing it explicitly we find

$$F(q) = \mathcal{L}q = \frac{1}{2\lambda}x^2 + \tilde{q}_0,$$

where  $\tilde{q}_0$  has slope  $< 2$ . This amounts to  $\lambda \mapsto -\frac{1}{\lambda}$  and the property is once again verified.  $\square$

The idea is then to factorize  $A$  as a product of elementary operations, using the following lemma.

**Lemma 4.13.** *Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{C})$ , and  $h_A : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  be the corresponding homography  $z \mapsto \frac{az+b}{cz+d}$ .*

- *If  $h_A(\infty) = \infty$ , then  $A$  admits a factorization as a product of elementary operations of the form  $A = S_\nu T_\rho$ , with  $\nu \in \mathbb{C}^*$ ,  $\rho \in \mathbb{C}$ .*

- Otherwise, if  $h_A(\infty) \neq \infty$ ,  $A$  admits a factorization of the form

$$(4.11) \quad A = T_\mu F S_\nu T_\rho,$$

with  $\mu \in \mathbb{C}$ ,  $\nu \in \mathbb{C}^*$ ,  $\rho \in \mathbb{C}$ .

*Proof.* We have  $h_A(\infty) = \frac{a}{c} \in \mathbb{P}^1$ , so  $h_A$  fixes infinity if and only if  $c = 0$ . This implies  $a \neq 0$  and  $d = a^{-1}$ , and we have  $A = S_\nu T_\rho$  with  $\nu = a$ ,  $\rho = b/a$ . When  $h_A(\infty) \neq \infty$ , we can reduce to the case where  $h_A$  fixes infinity. Let  $\mu := h_A(\infty) = \frac{a}{c}$ . We introduce

$$B_\mu := \begin{pmatrix} 0 & -1 \\ 1 & -\mu \end{pmatrix} = F^{-1} T_{-\mu},$$

such that the corresponding homography  $h_{B_\mu}$  sends  $\mu$  to  $\infty$ . Then the homography associated to  $B_\mu A$  sends  $\infty$  to  $\infty$ . This implies that  $B_\mu A$  has the form

$$B_\lambda A = \begin{pmatrix} \nu & \sigma \\ 0 & \nu^{-1} \end{pmatrix},$$

with  $\nu = -c$  and  $\sigma = -d$ , so that  $B_\mu A = S_\nu T_\rho$  with  $\rho = \sigma/\mu$ . Finally we get  $A = B_\mu^{-1} S_\nu T_\rho = T_\mu F S_\nu T_\rho$ .  $\square$

*Proof of the proposition.* If  $c = 0$  then  $d \neq 0$  and  $c\lambda + d = d \neq 0$ . In this case  $A$  factorizes as  $A = S_\nu T_\rho$ , so the circle  $\langle q' \rangle$  is at infinity. Now if  $c \neq 0$ , from the lemma  $A$  factorizes as  $A = T_\mu F S_\nu T_\rho$ . Let  $\langle q'' \rangle := S_\nu T_\rho \cdot \langle q \rangle_\infty$ , and write  $q'' = -\frac{\lambda''}{2} + q''_{<2}$  with the slope of  $q''_{<2}$  being  $< 2$ . The circle  $\langle q' \rangle$  is at infinity if and only if  $\langle q'' \rangle$  is not sent to finite distance by the Fourier transform, i.e. using the stationary phase formula if we don't have  $\lambda'' = 0$  and  $\text{slope}(q''_{<2}) \leq 1$ .

From the Legendre transform  $\text{slope}(q''_{<2}) = \text{slope}(q_{<2})$ . We have  $S_\nu T_\rho = \begin{pmatrix} -c & -d \\ 0 & -c^{-1} \end{pmatrix}$ , so  $\lambda'' = -\frac{c\lambda + d}{c}$ . Therefore we have  $\lambda'' \neq 0$  if and only if  $c\lambda + d \neq 0$ . It follows from this that, if we do not have  $c\lambda + d = 0$  and  $\text{slope}(q_{<2}) \leq 1$ , then at each step of the factorization the images of  $\langle q \rangle$  remain at infinity. Since at each elementary operation the coefficient  $\lambda$  is transformed according to the corresponding homography, the conclusion follows.  $\square$

We are now in position to prove the theorem.

*Proof of the theorem.* Let  $\Theta_\infty$  be an irregular class at infinity. Up to applying a well-chosen symplectic transformation, we may assume that all active circles of  $\Theta_\infty$  have slope less than 2. We denote by  $-\lambda_i/2, i = 1, \dots, r$  the coefficients of  $z^2$  of these active circles. It follows from the previous proposition that for  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{C})$ ,  $A \cdot \Theta_\infty$  only has a singularity at infinity if and only if  $c\lambda_i + d \neq 0$  for  $i = 1, \dots, m$ . This corresponds to an open dense subset of  $SL_2(\mathbb{C})$ . Furthermore  $A$  has a factorization  $A = T_\lambda F S_\mu T_\nu$ , such that when applying successively the elementary operations,  $\Theta_\infty$  always remains with only one singularity at infinity. At each step, the diagram doesn't change, and the conclusion follows.  $\square$

*Remark 4.1.* It follows from this analysis that one may view the active circles at finite distance as circles at infinity of slope  $\leq 2$  for which the coefficient of slope 2 is infinite. We may call the copy of  $\mathbb{P}^1$  where those coefficients live the *Fourier sphere*, as in [6]. Acting with symplectic transformations on circles at infinity of slope  $\leq 2$  amounts to acting on the Fourier sphere with the corresponding homographies. It is then clear that for a generic rotation none of the coefficients will be at infinity, i.e. there will only be one singularity at infinity.

## 5. DIAGRAMS FOR GENERAL CONNECTIONS

**5.1. Definition of the diagram.** To define the diagram in the the general case where there are several irregular singularities, the idea is to use the Fourier-Laplace transform to reduce to the situation where there is only one singularity at infinity studied in the previous section. The invariance of the diagram under  $SL_2(\mathbb{C})$  will ensure that this construction is well defined.

**Theorem 5.1.** *Let  $(E, \nabla)$  an algebraic connection on a Zariski open subset of  $\mathbb{A}^1$  with modified formal data  $(\check{\Theta}, \check{\mathcal{C}})$ . Then for any  $A$  in an open dense subset of  $SL_2(\mathbb{C})$ , the local system  $A \cdot (\check{\Theta}, \check{\mathcal{C}})$  only has support at infinity, and the diagram  $\Gamma^{BY}(A \cdot (\check{\Theta}, \check{\mathcal{C}}))$  is independent of  $A$ .*

*Proof.* It is easy to find a matrix  $A \in SL_2(\mathbb{C})$  such that  $A \cdot (\check{\Theta}, \check{\mathcal{C}})$  has support at infinity. For example we may take  $A$  of the form  $A = FT_\lambda$ , with  $\lambda \in \mathbb{C}$ : it suffices to choose the coefficient  $\lambda$  of the twist such that all the coefficients of the terms of slope 2 in the active circles of  $T_\lambda \cdot (\check{\Theta}, \check{\mathcal{C}})$  at infinity are non-zero. Applying then the Fourier transform, this guarantees that the active circles at infinity of  $T_\lambda \cdot \check{\Theta}$  remain at infinity while the active circles at finite distance are sent to infinity. The results of the previous section now imply that  $BA \cdot (\check{\Theta}, \check{\mathcal{C}})$  has support at infinity and that  $\Gamma^{BY}(BA \cdot (\check{\Theta}, \check{\mathcal{C}})) = \Gamma^{BY}(A \cdot (\check{\Theta}, \check{\mathcal{C}}))$  for  $B$  in an open dense subset of  $SL_2(\mathbb{C})$ .  $\square$

**Definition 5.2.** Let  $(E, \nabla)$  a connection on a Zariski open subset of the affine line,  $(\Theta, \mathcal{C})$  its formal data, and  $(\check{\Theta}, \check{\mathcal{C}})$  the corresponding modified formal data. We define the diagram associated to  $(E, \nabla)$  as

$$\Gamma(E, \nabla) = \Gamma(\Theta, \mathcal{C}) = \Gamma(\check{\Theta}, \check{\mathcal{C}}) := \Gamma^{BY}(A \cdot (\check{\Theta}, \check{\mathcal{C}}))$$

for any  $A \in SL_2(\mathbb{C})$  such that  $A \cdot (\check{\Theta}, \check{\mathcal{C}})$  has support at infinity.

**5.2. Direct formula for the core diagram.** The definition of the diagram involves a choice of  $SL_2(\mathbb{C})$  transformation. The explicit formulas for the number of edges and our understanding of the Legendre transform allow us to give a direct expression of the core diagram, which we may then take as the definition of the core diagram as in the introduction.

Let  $I = \langle q \rangle, I' = \langle q' \rangle$  be circles at points  $a, a' \in \mathbb{P}^1$ . Using similar notations as before, we write  $\beta := \text{ram}(q), \beta' := \text{ram}(q')$  and

$$q = \sum_{i=0}^k b_i z_a^{-\alpha_i/\beta}, \quad q' = \sum_{i=0}^{k'} b'_i z_{a'}^{-\alpha'_i/\beta'},$$

where the coefficients  $b_i, b'_i \in \mathbb{C}$  are non-zero, and set  $\alpha := \alpha_0 = \text{Irr}(q), \alpha' := \alpha'_0 = \text{Irr}(q')$ . If  $q$  and  $q'$  are both at infinity, that is if  $a = a' = \infty$ , then we know from a previous lemma the number of edges  $B_{\langle q \rangle, \langle q' \rangle}$ , which depends only on the degrees of the terms of  $q$  and  $q'$ . Let us denote by  $B_{\langle q \rangle, \langle q' \rangle}^\infty$  this formula.

**Theorem 5.3.** *Let  $I = \langle q \rangle_a$  and  $I' = \langle q' \rangle_{a'}$  be active circles. If  $a = a'$ , denoting as before by  $\alpha_r/\beta$  and  $\alpha'_r/\beta'$  the slopes of the different parts of  $q$  and  $q'$ , we assume that  $\frac{\alpha_r}{\beta} \geq \frac{\alpha'_r}{\beta'}$ . The number of edges  $B_{I, I'}$  between the vertices associated to  $I$  and  $I'$  following:*

- (1) *If  $a = a' = \infty$  then  $B_{I, I'} = B_{I, I'}^\infty$ .*
- (2) *If  $a = \infty$  and  $a' \neq \infty$  then  $B_{I, I'} = \beta(\alpha' + \beta')$ .*
- (3) *If  $a \neq \infty, a' \neq \infty$  and  $a \neq a'$  then  $B_{I, I'} = 0$ .*
- (4) *If  $a = a' \neq \infty$  then  $B_{I, I'} = B_{I, I'}^\infty - \alpha\beta' - \alpha'\beta$ .*

*Proof.* To prove this, we apply a generic  $SL_2(\mathbb{C})$  twist at infinity acting on exponential factors at infinity as  $q \mapsto q + \lambda z^2$ , followed by Fourier transform. From our study of the Legendre transform we obtain the form of the images of  $q$  and  $q'$  under this process, then we apply the formula for the number of edges at infinity. Let us first deal with the cases where  $q \neq q'$ .

- (1) This case is just a consequence of the fact that the twist and the Fourier transform leave  $B_{\langle q \rangle, \langle q' \rangle}^\infty$  invariant.
- (2) Let  $q_1$  and  $q'_1$  be the images of  $q, q'$ . If  $\frac{\alpha}{\beta} \leq 2$ , then  $q_1$  has ramification  $\beta$  and slope  $2 = 2\beta/\beta$ . If  $\frac{\alpha}{\beta} > 2$ , then  $q_1$  has ramification  $\alpha - \beta$  and slope  $\frac{\alpha}{\alpha - \beta}$ . On the other hand,  $q'_1$  is of the form

$$q'_1 = -a'z + b'z^{\alpha'/(\alpha' + \beta')} + \dots$$

for some  $b' \in \mathbb{C}, b' \neq 0$ , so  $q'_1$  has ramification  $\alpha' + \beta'$  and slope 1 (if  $a \neq 0$ ), or  $\frac{\alpha'}{\alpha' + \beta'}$  (if  $a = 0$ ). When  $\frac{\alpha}{\beta} \leq 2$ , it follows from the formula for  $B^\infty$  that

$$B_{\langle q \rangle, \langle q' \rangle} = B_{\langle q_1 \rangle, \langle q'_1 \rangle}^\infty = (\alpha' + \beta')(2\beta - \beta) = \beta(\alpha' + \beta').$$

When  $\frac{\alpha}{\beta} > 2$  the same formula gives

$$B_{\langle q \rangle, \langle q' \rangle} = B_{\langle q_1 \rangle, \langle q'_1 \rangle}^\infty = (\alpha' + \beta')(\alpha - (\alpha - \beta)) = \beta(\alpha' + \beta').$$

so both cases yield the expected result.

(3) In this case  $q_1$  and  $q'_1$  are of the form

$$q_1 = -az + bz^{\alpha/(\alpha+\beta)} + \dots, \quad q'_1 = -a'z + b'z^{\alpha'/(\alpha'+\beta')} + \dots,$$

that is  $q_1$  and  $q'_1$  have respective ramification orders  $\alpha + \beta$  and  $\alpha' + \beta'$ , have both slope 1 and non common part. The formula for  $B^\infty$  thus gives

$$B_{\langle q \rangle, \langle q' \rangle} = B_{\langle q_1 \rangle, \langle q'_1 \rangle}^\infty = (\alpha' + \beta')((\alpha + \beta) - (\alpha + \beta)) = 0.$$

(4) If  $a = a'$ ,  $q_1$  and  $q'_1$  are of the form

$$q_1 = -az + \tilde{q}, \quad q'_1 = -az + \tilde{q}',$$

with  $\tilde{q}$  having ramification  $\alpha + \beta$  and slope  $\frac{\alpha}{\alpha+\beta}$ , and  $\tilde{q}'$  having ramification  $\alpha' + \beta'$  and slope  $\frac{\alpha'}{\alpha'+\beta'}$ . Let us first assume that  $q$  and  $q'$  have a common part. Then so do  $q_1$  and  $q'_1$ . The slopes or their different parts are  $\frac{\alpha_r}{\alpha+\beta} \geq \frac{\alpha'_r}{\alpha'+\beta'}$ . The form of  $q_1$  and  $q'_1$  is given by the lemma giving the form of the Legendre transform. The slopes or their different parts are  $\frac{\alpha_r}{\alpha+\beta} \geq \frac{\alpha'_r}{\alpha'+\beta'}$  and the formula for  $B_{\langle q_1 \rangle, \langle q'_1 \rangle}^\infty$  gives

$$\begin{aligned} B_{\langle q \rangle, \langle q' \rangle} &= B_{\langle q_1 \rangle, \langle q'_1 \rangle}^\infty \\ &= ((\alpha' + \beta') - (\alpha'_0, \alpha' + \beta'))\alpha_0 + \dots + (\alpha'_0, \dots, \alpha'_{r-1}, \alpha' + \beta')\alpha_r - (\alpha + \beta)(\alpha' + \beta') \\ &= ((\alpha' + \beta') - (\alpha'_0, \beta'))\alpha_0 + \dots + (\alpha'_0, \dots, \alpha'_{r-1}, \beta')\alpha_r - (\alpha\alpha' + \beta\beta' + \alpha'\beta + \alpha\beta') \\ &= B_{\langle q \rangle, \langle q' \rangle}^\infty - \alpha\beta' - \alpha'\beta \\ &= B_{\langle q \rangle, \langle q' \rangle}^\infty - 2\alpha\beta', \end{aligned}$$

where in the last step we have used that  $\frac{\alpha}{\beta} = \frac{\alpha'}{\beta'}$  since  $q$  and  $q'$  have a common part. Now assume that  $q$  and  $q'$  have no common part. Then the slope of  $\tilde{q}$  and  $\tilde{q}'$  are respectively  $\frac{\alpha}{\alpha+\beta} \geq \frac{\alpha'}{\alpha'+\beta'}$  (the order is preserved by the Legendre transform), so from the formula for  $B_{\langle \tilde{q} \rangle, \langle \tilde{q}' \rangle}^\infty$  we have

$$B_{\langle q \rangle, \langle q' \rangle} = -\beta(\alpha' + \beta').$$

On the other hand, we have  $B_{\langle \tilde{q} \rangle, \langle \tilde{q}' \rangle}^\infty = \beta'(\alpha - \beta)$ , so we again have  $B_{\langle q \rangle, \langle q' \rangle}^\infty = B_{\langle q \rangle, \langle q' \rangle}^\infty - \alpha\beta' - \alpha'\beta$  as expected.

To complete the proof, it remains to deal with the case of loops. We check that one has  $B_{\langle q \rangle, \langle q \rangle} = B_{\langle q \rangle, \langle q \rangle}^\infty - 2\alpha\beta$  when  $a \neq \infty$ , so the formula for  $B_{\langle q \rangle, \langle q' \rangle}$  remains valid when  $q = q'$ .  $\square$

**5.3. Fundamental representations of the diagrams.** In this section, we discuss how a diagram can be “read” in different ways corresponding to connection on bundles of different ranks with different formal data, which we call the fundamental representations of the diagram. This generalizes the  $k$ -partite cases considered in [6].

The idea is the following. We have seen that the diagram associated to a meromorphic connection is invariant under the action of  $SL_2(\mathbb{C})$ . Each orbit under this action contains different formal data, with in general different numbers of singularities.

Let  $\langle q \rangle_\infty$  be the circle at infinity associated with an exponential factor  $q$ . We have seen that  $\langle q \rangle$  can be sent to finite distance by a symplectic transformation if and only if it is of the form

$$(5.1) \quad q = \lambda z^2 + q',$$

where  $q'$  is of slope  $\leq 1$ . Furthermore, when this is the case, the position of the singularity at finite distance that we obtain is determined by the coefficient of  $z$  in  $q$ , i.e. the number  $\mu \in \mathbb{C}$  such that

$$(5.2) \quad q = \lambda z^2 + \mu z + q'',$$

where  $q''$  has slope  $< 1$ .

Let us consider formal data at infinity  $(\Theta_\infty, \mathcal{C}_\infty)$  at infinity. Let  $N \subset \pi_0(\mathcal{I}_\infty)$  be the subset of active exponents, i.e. the support of  $\Theta_\infty$ .  $N$  is the set of vertices of the core diagram associated to  $\Theta_\infty$ . We can partition  $N$  according to the coefficients of  $z^2$  and  $z$  in the exponential factors. Let  $N_\infty \subset N$  be the set of exponential factors in  $N$  which are not of the form (5.1) for any  $\lambda \in \mathbb{C}$ . For  $\lambda \in \mathbb{C}$ , let  $N_\lambda \subset N$  be the set of exponential factors of the form (5.1). For  $\lambda, \mu \in \mathbb{C}$ , let  $N_{\lambda, \mu} \subset N_\lambda$  be the set of exponential factors of the form (5.2).

There is a finite number of coefficients  $\lambda \in \mathbb{C}$  such that  $N_\lambda$  is non empty, which we denote  $\lambda_1, \dots, \lambda_k$ . We set  $N_i := N_{\lambda_i}$  for  $i = 1, \dots, r$ . Then, for each  $i$ , there is again a finite number of coefficients  $\mu \in \mathbb{C}$  such that  $N_{\lambda_i, \mu}$  is non-empty, which we denote  $\mu_{i,1}, \dots, \mu_{i,s_i}$ , and we set  $N_{i,j} := N_{\lambda_i, \mu_{i,j}}$ . For each pair  $i, j$ , let  $t_{i,j} := \text{Card } N_{i,j}$ , and  $q_{i,j,k}$ ,  $k = 1, \dots, t_{i,j}$  be the exponential factors of slope  $< 1$  such that the elements of  $N_{i,j}$  are

$$(5.3) \quad \lambda_i z^2 + \mu_{i,j} z + q_{i,j,k}.$$

Let  $\beta_{i,j,k} = \text{ram } q_{i,j,k}$  be the ramification order of  $q_{i,j,k}$ , and  $\alpha_{i,j,k}/\beta_{i,j,k}$  its slope. We thus have the following partition of the set of active circles:

$$N = N_\infty \cup N_1 \cup \dots \cup N_r.$$

There are  $r + 1$  fundamental representations of diagram: the generic representation, corresponding to  $\Theta$ , and  $r$  other representations, depending on which of the sets  $N_1, \dots, N_r$  we choose to send to finite distance by a symplectic transformation. The generic representation corresponds to symplectic transformations  $A \in SL_2(\mathbb{C})$  such that all circles remain at infinity under  $A$ . From the previous discussion of the action of  $SL_2(\mathbb{C})$  on the coefficients of  $z^2$  in the exponential factor, if

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

the condition for this is  $2c\lambda - d \neq 0$ .

For each  $i = 1, \dots, r$  the  $i$ -th representation of the diagram corresponds to acting on  $\Theta_\infty$  with  $A \in SL_2(\mathbb{C})$  such that  $2c\lambda_i - d = 0$ . Such a symplectic transformation sends all circles in  $N_i$  to finite distance. The formal data that we obtain have  $s_i = \text{Card}(N_i)$  poles at finite distance corresponding to the coefficients  $\mu_{i,1}, \dots, \mu_{i,s_i}$  (and whose position depends on  $A$ ). For each  $j = 1, \dots, s_i$ , the vertices in  $N_{i,j}$  then correspond to the active circles of  $A \cdot \Theta_\infty$  at the pole associated to  $\mu_{i,j}$ . It follows from the Legendre transformation that the image  $A \cdot \langle q_{i,j,k} \rangle$  of the circle  $\langle q_{i,j,k} \rangle$  has ramification order  $\beta_{i,j,k} - \alpha_{i,j,k}$  and slope  $\alpha_{i,j,k}/(\beta_{i,j,k} - \alpha_{i,j,k})$ . This generalizes the different readings of the diagram considered in [5, 6, 8]. The situation considered in those works is the simply laced case where all circles at infinity are of the form

$$q = \lambda z^2 + \mu z,$$

with  $\lambda, \mu \in \mathbb{C}$ . In particular  $N_\infty$  is empty. In this situations, each circle can be sent by a symplectic transformation to a tame circle at finite distance.

## 6. DIMENSION OF THE WILD CHARACTER VARIETY

**6.1. The wild character variety.** Let us quickly review the quasi-Hamiltonian description of wild character variety  $\mathcal{M}_B(E, \nabla)$  in the general case with several singularities.

Let us set  $G := GL_n(\mathbb{C})$  where  $n$  is the rank of  $(E, \nabla)$ , and keeping the previous notations let  $a_1, \dots, a_m \in \mathbb{P}^1$  be the singular points of any connection with set of formal data  $(\Theta, \mathcal{C})$ . Let  $V^0 \rightarrow \mathcal{I}$  be the (mon-modified) local system on  $\mathcal{I}$  giving rise to  $V$ . For  $k = 1, \dots, m$ , we consider the restriction  $V_{a_k}^0 \rightarrow \partial_{a_k}$  of the local system system  $V^0$ , that we now see as a local system over the circles  $\partial_a$ ,  $a \in \mathbb{C}$ . Let us choose a direction  $d \in \partial_{a_k}$ , and denote by  $\rho_k \in GL(V_d^0)$  the monodromy of  $V_{a_k}^0$ . As before, we denote by  $\langle q_1^{(k)} \rangle, \dots, \langle q_{r_k}^{(k)} \rangle \in \pi_0(\mathcal{I}_{a_k})$  the active circles at  $a_k$ ,  $n_i^{(k)} \in \mathbb{N}$  their respective multiplicities, and  $\beta_j^{(k)}$  their respective ramification orders. Let  $H_k := \text{GrAut}(V_{a_k}^0) \simeq \prod_{i=1}^{r_k} GL_{n_i^{(k)}}(\mathbb{C})^{b_i^{(k)}}$ . The monodromy  $\rho_k$  is an element of the  $H_k$ -torsor

$H_k(\partial_{a_k})$  (see [11]). Explicitly, an element of  $H_k(\partial_{a_k})$  is of the form

$$h^{(k)} = \text{diag}(h_1^{(k)}, \dots, h_{r_k}^{(k)}),$$

where  $h_i^{(k)}$  is a square diagonal block matrix of size  $\beta_i^{(k)}$  of the form

$$h_i^{(k)} = \begin{pmatrix} 0 & \dots & 0 & * \\ * & & & 0 \\ & \ddots & & \vdots \\ & & * & 0 \end{pmatrix},$$

with each block in  $GL_{n_i^{(k)}}(\mathbb{C})$ .

The wild character variety  $\mathcal{M}_B(E, \nabla) = \mathcal{M}_B(\Theta, \mathcal{C})$  is obtained as a symplectic reduction of the twisted quasi-Hamiltonian manifold

$$\text{Hom}_{\mathbb{S}}(V) \simeq \mathcal{A}(V_{a_1}^0) \otimes \dots \otimes \mathcal{A}(V_{a_m}^0) // G,$$

where  $\otimes$  is the quasi-Hamiltonian fusion operation. Here, each piece  $\mathcal{A}(V_{a_k}^0)$  is a twisted quasi-Hamiltonian  $H_k \times G$ -space:

$$\mathcal{A}(V_{a_k}^0) = H(\partial_{a_k}) \times G \times \prod_{d \in \mathbb{A}_{a_k}} \text{Sto}_d,$$

where  $\mathbb{A}_{a_k}$  is the set of singular directions of the connection at  $a_k$ , and  $\text{Sto}_d$  is the Stokes group associated to the singular direction  $d \in \mathbb{A}_{a_k}$ . Hence,  $\text{Hom}_{\mathbb{S}}(E, \nabla)$  is a twisted quasi-Hamiltonian  $\mathbf{H} \times G$ -space, where  $\mathbf{H} = H_1 \times \dots \times H_m$ .

The formal monodromies  $\rho(a_k) \in H(\partial_{a_k})$  determine twisted conjugacy classes  $\mathcal{C}(\partial_{a_k}) \subset H(\partial_{a_k})$  which do not depend on the choice of direction  $d \in \partial_{a_k}$  and the choice of isomorphism  $V_d^0 \simeq \mathbb{C}^{\sum_j n_j^{(k)} \beta_j^{(k)}}$ . Finally, the wild character variety is the twisted quasi-Hamiltonian reduction of  $\text{Hom}_{\mathbb{S}}(E, \nabla)$  at the twisted conjugacy class  $\mathcal{C} := \prod_{k=1}^m \mathcal{C}(\partial_{a_k})$ , i.e.

$$(6.1) \quad \mathcal{M}_B(\Theta, \mathcal{C}) = \text{Hom}_{\mathbb{S}}(E, \nabla) //_{\mathcal{C}} \mathbf{H}.$$

**6.2. Formula for the dimension.** Let  $\Gamma$  be a diagram, possibly with loops, negative edges or negative loops. Let  $N$  be its set of vertices and  $k$  its cardinal. A dimension vector associated with  $\Gamma$  is an element  $\mathbf{d} = (d_i)_{i \in N} \in \mathbb{N}^N$ . Let  $A \in M_k(\mathbb{Z})$  be the adjacency matrix of the diagram. The Cartan matrix of the diagram is defined by

$$C = 2Id - A$$

We denote by  $(\cdot, \cdot)$  the bilinear form on  $\mathbb{C}^k$  with matrix  $C$ .

We have the following result:

**Theorem 6.1.** *The dimension of the wild character variety  $\mathcal{M}_B(\Theta, \mathcal{C})$ , when it is non-empty, is given by*

$$(6.2) \quad \dim \mathcal{M}_B(\Theta, \mathcal{C}) = 2 - (\mathbf{d}, \mathbf{d}),$$

where  $(\cdot, \cdot)$  is the bilinear form associated with the diagram  $\Gamma(\Theta, \mathcal{C})$ , and  $\mathbf{d}$  is the dimension vector of  $\Gamma(\Theta, \mathcal{C})$ .

The proof is given in appendix C.

**Corollary 6.2.** *If  $(\Theta, \mathcal{C})$  are the formal data of a connection  $(E, \nabla)$ , and  $A \in SL_2(\mathbb{C})$ , then the character varieties  $\mathcal{M}_B(\Theta, \mathcal{C})$  and  $\mathcal{M}_B(A \cdot (\Theta, \mathcal{C}))$  have the same dimension.*

*Proof.* By construction of the diagram  $V$  and  $A \cdot (\Theta, \mathcal{C})$  have the same diagram  $\Gamma$ . Therefore we have  $\dim \mathcal{M}_B(\Theta, \mathcal{C}) = 2 - (\mathbf{d}, \mathbf{d}) = \dim \mathcal{M}_B(A \cdot (\Theta, \mathcal{C}))$ .  $\square$


## 7. EXAMPLES OF DIAGRAMS

Let us now give a few examples of diagrams. We first review a few known cases to see how we recover the diagrams from our more general approach, then discuss some new cases.

**Complete bipartite case.** Let us consider the case of a connection with a second order poles at infinity, together with simple poles  $a_1, \dots, a_m$  at finite distance, with a global modified irregular class of the form

$$\check{\Theta} = k_1 \langle \alpha_1 z \rangle_\infty + \dots + k_n \langle \alpha_n z \rangle_\infty + l_1 \langle 0 \rangle_{a_1} + \dots + l_m \langle 0 \rangle_{a_m},$$

with  $k_i \in \mathbb{N}$ ,  $l_j \in \mathbb{N}$ , and  $\alpha_i \in \mathbb{C}$  two by two distinct,  $a_i \in \mathbb{C}$  two by two distinct. There are  $n$  active circles of slope 1 at infinity. The diagram is the following:



It is a complete bipartite graph, one part having  $n$  vertices and one part having  $m$  vertices. If  $n = 1$  or  $m = 1$  this is the star-shaped case.


**Simply laced case.** An important case is the one where there is only one unramified irregular pole at infinity of order 3, together with simple poles at finite distance. This is the case extensively studied in [5, 6, 8], giving rise in those works to simply-laced supernova graphs, with core any complete multipartite graph. We can check that our definition gives the same diagrams.

In this case, the circles at infinity are unramified and have slope  $\leq 2$ , i.e. they are of the form  $\langle q_{i,j} \rangle_\infty$  with  $q_{i,j} = \lambda_i z^2 + \mu_{i,j} z$ , for  $i = 1, \dots, k$ ,  $j = 1, \dots, s_i$ , where the coefficients  $\lambda_1, \dots, \lambda_k$  are all different, as well as  $\mu_{i,1}, \dots, \mu_{i,s_i}$  for all  $i$ . The simple poles correspond to tame circles  $\langle 0 \rangle_{a_1}, \dots, \langle 0 \rangle_{a_t}$ , where  $a_1, \dots, a_t \in \mathbb{C}$ . Let  $I_0 := \{\langle 0 \rangle_{a_1}, \dots, \langle 0 \rangle_{a_t}\}$  and for  $i = 1, \dots, k$ ,  $I_i := \{\langle \lambda_i z^2 + \mu_{i,1} z \rangle, \dots, \langle \lambda_i z^2 + \mu_{i,s_i} z \rangle\}$ . The sets  $I_0, \dots, I_k$  constitute a partition of the set of active circles. The core diagram has the following structure two active circles are either linked by no edge in the diagram when they belong to the same set  $I_k$ , otherwise they are linked by exactly one edge. The core diagram that we obtain is thus a  $k+1$ -partite graph. The diagram coincides with the one considered in [6, 8].

The standard rank two Lax representations of the Painlevé V and Painlevé IV equation fit into this setting. The representation for Painlevé V has one irregular singularity at infinity of order 2, together with two simple poles at finite distance. The modified irregular class is of the form

$$\check{\Theta} = \langle \alpha z \rangle_\infty + \langle \beta z \rangle_\infty + \langle 0 \rangle_a + \langle 0 \rangle_b,$$


with  $\alpha \neq \beta \in \mathbb{C}$ ,  $a \neq b \in \mathbb{C}$ . This gives the following diagram:



The standard representation for Painlevé IV has one irregular singularity at infinity of order 3, together with one simple pole at finite distance. The modified irregular class is of the form

$$\check{\Theta} = \langle \alpha z^2 \rangle_\infty + \langle \beta z^2 \rangle_\infty + \langle 0 \rangle_a,$$

with  $\alpha \neq \beta \in \mathbb{C}$ ,  $a \in \mathbb{C}$ . This gives the following diagram:






**Other supernova graphs.** More general supernova graphs correspond to the case where there is one irregular singularity at infinity with all active circles being unramified, together with simple poles at infinity. This is the case considered in [5, appendix C] and [22]. If the active circles at infinity are  $\langle q_1 \rangle, \dots, \langle q_r \rangle$  we have  $B_{ij} = \deg(q_i - q_j) - 1$ .

The standard Painlevé II Lax pair fits into this framework. The (modified or not) irregular class is of the form  $\Theta = \Theta_\infty = \langle \alpha z^3 \rangle_\infty + \langle \beta z^3 \rangle_\infty$  with  $\alpha \neq \beta$ . The diagram is



**Painlevé III.** The standard Painlevé III Lax pair corresponds to a rank 2 connection with 2 irregular singularities. The active circles are two irregular circles of slope 1 at infinity and 2 irregular circles of slope 1 at  $z = 0$ , say  $\langle \lambda_1 z \rangle_\infty, \langle \lambda_2 z \rangle_\infty, \langle \mu_1 z^{-1} \rangle_0, \langle \mu_2 z^{-1} \rangle_0$ , with  $\lambda_1 \neq \lambda_2, \mu_1 \neq \mu_2$ . With no loss of generality, up to applying a twist by a rank one connection on the trivial bundle on  $\mathbb{P}^1$ , we may assume that  $\mu_2 = 0$ , so that we have the active circle  $\langle 0 \rangle_0$ , and that the tame circle has trivial formal monodromy in the non-reduced formal local system, so that it has multiplicity zero in the associated modified formal local system. The diagram associated to the connection is given by




with dimension vector  $\mathbf{d} = (1, 1, 1)$ . This is exactly the same diagram as the one in [12].

**Degenerate Painlevé V.** The representation of Painlevé III known as degenerate Painlevé V corresponds as for Painlevé V to a rank 2 connection with one order two pole at infinity and two simple poles at finite distance [23, p. 34]. The difference is that the leading term of the connection is nilpotent (this is the meaning of the word degenerate in this context) which implies there is just one ramified active circle at infinity of slope  $1/2$ . The modified irregular class is of the form:

$$\check{\Theta} = \langle \alpha z^{1/2} \rangle_\infty + \langle 0 \rangle_a + \langle 0 \rangle_b,$$

and this gives the diagram




computed in [12]. This is the same diagram as for the standard Painlevé III Lax pair. The two irregular classes indeed correspond to different representations of the diagram.

**Degenerate Painlevé III.** The degenerate Painlevé III system admits, as for the standard Painlevé III representation, a representation corresponding to a rank 2 connection with two order two poles [29]. The difference is that the leading term of the connection matrix at one of the poles is nilpotent. Again, this corresponds to having one active circle with slope  $1/2$ . The modified irregular class has the following form:

$$\check{\Theta} = \langle z^{1/2} \rangle_\infty + \langle \alpha z^{-1} \rangle_0 + \langle \beta z^{-1} \rangle_0$$

with  $\alpha, \beta \in \mathbb{C}, \alpha \neq \beta$ .

Up to performing a twist at zero, we may assume that  $\beta = 0$ , so that we have the tame circle at 0, and that this circle has trivial formal monodromy, so that the corresponding piece of the modified formal local system has multiplicity 0. This leads to the following diagram, with one negative loop at each vertex, and 4 edges between the two vertices.




The Cartan matrix is

$$C = \begin{pmatrix} 4 & -4 \\ -4 & 4 \end{pmatrix}.$$

**Doubly degenerate Painlevé III.** The doubly degenerate Painlevé III system admits a Lax representation corresponding to two irregular singularities of order two, each with nilpotent leading term leading to an active circle of slope  $1/2$  [29]. The modified irregular class is of the form

$$\check{\Theta} = \langle \alpha z^{1/2} \rangle_{\infty} + \langle \beta z^{-1/2} \rangle_0$$

with  $\alpha, \beta$  nonzero complex numbers. This gives the following diagram, where the number on the edges correspond to their multiplicities.




The Cartan matrix is

$$C = \begin{pmatrix} 8 & -6 \\ -6 & 4 \end{pmatrix},$$

**Flaschka-Newell Lax pair for Painlevé II.** The Flaschka-Newell Lax pair for Painlevé II [18], also known as degenerate Painlevé IV, corresponds to a rank 2 connection with one irregular singularity at infinity with one active circle of slope  $3/2$ , together with one simple pole at finite distance. The modified irregular class is of the form

$$\check{\Theta} = \langle \alpha z^{3/2} \rangle_{\infty} + \langle 0 \rangle_0.$$

with  $\alpha \neq 0$ . This gives the diagram



that is the same diagram as for Painlevé II, as found in [12]. Again, the two irregular classes correspond to different fundamental representations of the diagram.

**Painlevé I.** The standard Lax pair for the Painlevé I equation corresponds to a rank 2 connection with just one irregular singularity at infinity, with one active circle of slope  $5/2$ . The modified irregular class is of the form

$$\check{\Theta} = \langle \alpha z^{5/2} \rangle_{\infty},$$

with  $\alpha \neq 0$ . The corresponding diagram has one vertex and one loop.



**H3 surfaces.** The Painlevé equations correspond to the simplest examples of non-trivial wild character varieties, since their moduli spaces are 2-dimensional. The corresponding nonabelian Hodge spaces  $\mathfrak{M}$  are thus complete hyperkahler manifolds of real dimension four, so are examples of H3 surfaces in the terminology of [9] to designate the two-dimensional wild character varieties. Apart from the cases corresponding to Painlevé equations, there are 3 other known H3 surfaces whose standard representation correspond to fuchsian connection with 3 regular singularities. Since they have no deformation parameters, they do not give rise to an isomonodromy system. Thanks to our more general theory, we are now able to associate a diagram to all H3 surfaces. They are represented on fig. 1. The reader may check that in each case we have  $2 - (\mathbf{d}, \mathbf{d}) = 2$  as expected. Notice that for Painlevé II, IV, V, VI, the diagram is exactly the affine Dynkin diagram corresponding to its Okamoto symmetries [30, 32, 31, 33, 29],




FIGURE 1. Diagrams associated to all known H3 surfaces. The names of the surfaces are as in [9]. All unspecified multiplicities are equal to 1.

**h-Painlevé systems.** It is possible to define a notion of h-Painlevé systems  $hP_k^{(n)}$ , where h stands for higher or hyperbolic or Hilbert. Higher Painlevé systems  $hP_k^{(n)}$  were introduced by Boalch [5, 6]: for each integer  $n$  there is a rank  $2n$  Lax representation giving rise to a  $2n$ -dimensional higher Painlevé moduli space. All higher Painlevé systems of a given number have the same diagram, but the dimension vector is a function of  $n$ . More precisely, the diagram associated to a given higher Painlevé system is obtained by taking the original diagram with all multiplicities scaled by  $n$ , then adding a leg of length one, with its end vertex having multiplicity 1, as on fig. 2 for higher Painlevé IV,V,VI. The higher Painlevé diagrams are hyperbolic Dynkin diagrams since they are obtained by adding a vertex to an affine Dynkin diagram. In this sense they can be seen as the next simplest examples after the affine case. The higher Painlevé moduli spaces are (conjecturally) related to Hilbert schemes on  $n$  points on the corresponding H3 surface (see [6]). The same recipe for Painlevé III yields the  $hP_3^{(n)}$  diagram.




FIGURE 2. Diagrams for higher Painlevé systems  $hP_{VI}^{(n)}$ ,  $hP_V^{(n)}$ ,  $hP_{IV}^{(n)}$ , and  $hP_3^{(n)}$ .

**Some 4-dimensional cases.** Another point of view on higher dimensional isomonodromy systems consists in looking at degenerations of an equation coming via isomonodromy from a fuchsian connection. This approach is used in [24] to list some representations of some 4-dimensional isomonodromy systems: they are defined there as the systems coming via degeneration from the isomonodromic deformation equations associated to connections with regular singularities giving rise to 4-dimensional moduli spaces. Whereas for the usual 2-dimensional Painlevé equations there is only one possible choice of formal data for fuchsian connections with non-trivial admissible deformations and 2-dimensional moduli spaces, in the 4-dimensional case there are 4 possible choices. The full degeneration scheme is represented at [24, p. 40]. Each of the 4 fuchsian formal data gives rise via degeneration to a family of 4-dimensional Painlevé type equations.

The two points of view on higher-dimensional isomonodromy systems are actually not independent. The elements of the fourth degeneration family of 4-dimensional Painlevé-type equations in the degeneration scheme of [24] (called there matrix Painlevé systems) correspond to the higher Painlevé equations in the sense of [6] in the 4-dimensional case, and the master diagram  $hP_6^{(2)}$  at the top of the coalescence cascade first appeared in the context of Painlevé systems in the list of 4-dimensional hyperbolic examples in [5, p. 12]. This notion of higher Painlevé systems is however more general than just this 4-dimensional case.

On figure 3 are drawn the diagrams associated by our approach to the 4-dimensional isomonodromy systems listed in [24] whose Lax representations feature several irregular singularities. These diagrams are to be contrasted with the shapes in [21, p.235].

*Remark 7.1.* Notice that the diagram associated to the Painlevé-type equation denoted  $H_{III}^{\text{Mat}}(D_6)$  in [24] fits in the higher Painlevé picture in the sense of Boalch: it is  $hP_3^{(2)}$ , obtained from the Painlevé III diagram by taking all multiplicities equal to 2 and adding a vertex with multiplicity 1. Beware that the usual terminology “matrix Painlevé equations” [3] differs from that of [24].

*Remark 7.2.* It is interesting to compare our diagrams with those of Hiroe [22] and Hiroe and Oshima [21]. In [20], Hiroe defines diagrams associated to meromorphic connections with several unramified irregular singularities. When there is only one irregular singularity, these diagrams coincide with the ones of [6] and in turn with ours, however when there are several irregular singularities they differ from ours. Hiroe also introduces the notion of *shape*, and in their approach the moduli spaces are classified by these shapes. When there are several irregular singularities, the shape differs from the diagram. This approach however does not allow to define diagrams for the degenerate and doubly degenerate Painlevé equations.

Name in [24]	Mod. irreg. class	Diagram
$H_{\text{Gar}}^{2+2+1}$	$\langle \alpha z \rangle_\infty + \langle \beta z \rangle_\infty + \langle \gamma z^{-1} \rangle_0 + \langle 0 \rangle_1$	
$H_{\text{Gar}}^{3+2}$	$\langle \alpha z^2 \rangle_\infty + \langle \beta z^2 \rangle_\infty + \langle \gamma z^{-1} \rangle_0$	
$H_{\text{FS}}^{A_3}$	$2\langle \alpha z \rangle_\infty + \langle \beta z \rangle_\infty + 2\langle \gamma z^{-1} \rangle_0$	
$H_{\text{Gar}}^{3/2+1+1+1}$	$\langle \alpha z^{1/2} \rangle_\infty + \langle 0 \rangle_a + \langle 0 \rangle_b + \langle 0 \rangle_c$	
$H_{\text{Ss}}^{D_4}$	$3\langle \alpha z \rangle_\infty + \langle \beta z \rangle_\infty + 2\langle \gamma z^{-1} \rangle_0$	
$H_{\text{III}}^{\text{Mat}}(D_6)$	$2\langle \alpha z \rangle_\infty + 2\langle \beta z \rangle_\infty + 2\langle \gamma z^{-1} \rangle_0$	

FIGURE 3. Diagrams for the Lax representations of 4-dimensional Painlevé-type equations of [24] featuring several irregular singularities.

#### APPENDIX A. FORMULA FOR THE NUMBER OF EDGES

We now prove the explicit formula for the number of edges (lemma 4.3). Lets us recall the formula:

**Lemma A.1.** • Assume that  $\alpha_r/\beta \geq \alpha'_r/\beta'$ . Then the number of edges between  $\langle q \rangle$  and  $\langle q' \rangle$  is

$$B_{\langle q \rangle, \langle q' \rangle} = (\beta' - (\alpha' \beta'))\alpha_0 + ((\alpha'_0, \beta') - (\alpha'_0, \alpha'_1, \beta'))\alpha_1 + \cdots + ((\alpha'_0, \dots, \alpha'_{r-2}, \beta') - (\alpha'_0, \dots, \alpha'_{r-1}, \beta'))\alpha_{r-1} + (\alpha'_0, \dots, \alpha'_{r-1}, \beta')\alpha_r - \beta\beta'$$

• In particular, if  $q$  and  $q'$  have no common parts and  $\alpha/\beta \geq \alpha'/\beta'$ , then

$$B_{\langle q \rangle, \langle q' \rangle} = \beta'(\alpha - \beta).$$

*Proof.* To compute the number of edges between  $\langle q \rangle$  and  $\langle q' \rangle$  we have to determine the local system  $\text{Hom}(\langle q \rangle, \langle q' \rangle)$  as well as its irregularity. For this we have to look at the differences  $q_i - q'_j$ , with  $i = 0, \dots, \beta - 1, j = 0, \dots, \beta' - 1$  between all possible leaves  $\langle q \rangle$  of  $\langle q' \rangle$ , find which connected components of  $\mathcal{I}$  they fall into and what the irregularity of those circles is. The subtlety is that the degree of  $q_i - q_j$  depends in general of  $i$  and  $j$ , so that the circles will not have the same irregularity.

Let  $\mu$  be the smallest common multiple of  $\beta$  and  $\tilde{\beta}$ . We set

$$\mu = k\beta, \quad \mu = k'\beta'.$$

Let us also denote by  $\delta$  the greatest common divisor of  $\beta$  and  $\beta'$ . We have

$$\beta = k'\delta, \quad \beta' = k\delta.$$

For  $0 \leq i \leq r$ , let  $\gamma_i$  be the integer such that

$$\frac{\alpha_i}{\beta} = \frac{\alpha'_i}{\beta'} = \frac{\gamma_i}{\delta}.$$

Any difference  $q_i - q'_j$  belongs to a circle  $I = \langle q_i - q'_j \rangle$  which is a connected component of  $\text{Hom}(\langle q \rangle, \langle q' \rangle)$ . However, several differences  $q_i - q'_j$  belong to the same connected component: each difference  $q_i - q_j$  corresponds to exactly one leaf of such a circle. This implies the following formula:

$$(A.1) \quad \text{Irr}(\text{Hom}(\langle q \rangle, \langle q' \rangle)) = \sum_{i=0}^{\beta-1} \sum_{j=0}^{\beta'-1} \text{slope}(q_i - q'_j).$$

Indeed, regrouping the terms of the sum according to the connected components, to each connected component  $I$  of ramification order  $r$  and irregularity  $s$  correspond  $r$  differences  $q_i - q'_j$  in the sum, with slope equal to  $\text{slope}(q_i - q'_j) = s/r$ . The total contribution of those terms is thus equal to  $s = \text{Irr}(I)$ .

The computation can be simplified by noticing the following fact: for any  $k \in \mathbb{Z}$ ,

$$\text{slope}(q_i - q_j) = \text{slope}(q_{i+k} - q'_{j+k}),$$

where the index  $i+k$  is seen as an element of  $\mathbb{Z}/\beta\mathbb{Z}$ , and  $j+k$  is seen as an element of  $\mathbb{Z}/\beta'\mathbb{Z}$ . Under this shifting action of  $\mathbb{Z}$ , the set  $\mathbb{Z}/\beta\mathbb{Z} \times \mathbb{Z}/\beta'\mathbb{Z}$  is partitioned into  $\delta$  orbits, each having cardinal  $\mu$ . Furthermore, the differences

$$q_0 - q'_j, \quad j = 0, \dots, \delta - 1,$$

belong to distinct orbits and thus yield one representative of each orbit. The formula (A.1) therefore becomes

$$\begin{aligned} \text{Irr}(\text{Hom}(\langle q \rangle, \langle q' \rangle)) &= \mu \sum_{j=0}^{\delta-1} \text{slope}(q_0 - q'_j) \\ &= \sum_{j=0}^{\delta-1} \deg_{z^{-1/\mu}}(q_0 - q'_j). \end{aligned}$$

The task is thus reduced to computing the degree of the differences  $q_0 - q'_j$ ,  $j = 0, \dots, \delta - 1$  as polynomials in  $z^{-1/\mu}$ . It is the exponent of the largest monomial having different coefficients in  $q_0$  and  $q'_j$ .

The common part  $q_c = q'_c$  is given by

$$q_c = q'_c = \sum_{i=0}^r b_i z_\infty^{-\alpha_i/\beta} = \sum_{i=0}^r b_i z_\infty^{-\alpha'_i/\beta'} = \sum_{i=0}^r b_i z_\infty^{-\gamma_i/\delta} = \sum_{i=0}^r b_i z_\infty^{-kk'\gamma_i/\mu}.$$

We will start by determining the number of indices  $j = 0 \in \{0, \dots, \delta - 1\}$  such that this degree is the maximal possible degree  $kk'\gamma_0$ , then the number of indices for which it is  $kk'\gamma_1$ , etc.

Let us thus begin by computing the number of differences with degree  $kk'\gamma_0$ . We consider the coefficient of  $z_\infty^{kk'\gamma_0/\mu}$  in the difference  $q_0 - q'_j$ ,  $j = 0, \dots, \delta - 1$ : the corresponding term is

$$a_0(z_\infty^{kk'\gamma_0/\mu} - e^{2i\pi j\gamma_0/\delta} z_\infty^{kk'\gamma_0/\mu}) = a_0(1 - e^{2i\pi j\gamma_0/\delta}) z_\infty^{kk'\gamma_0/\mu}.$$

The factor  $1 - e^{2i\pi j\gamma_0/\delta}$  is zero if and only if  $j$  is an integer multiple of  $\delta/(\gamma_0, \delta)$ . There are thus  $(\gamma_0, \delta)$  differences  $q_0 - q'_j$  having a degree strictly less than  $kk'\gamma_0$ . The  $\delta - (\gamma_0, \delta)$  other differences  $q_0 - q'_j$  have degree  $kk'\gamma_0$ . Each one of these contributes to  $kk'\gamma_0$  Stokes arrows from  $\langle q \rangle$  to  $\langle q' \rangle$ .

Next, we compute the number of differences  $q_0 - q'_j$ ,  $j = 0, \dots, \delta - 1$ , whose degree is  $kk'\gamma_1$ . In  $q_0 - q_j$  the monomial of degree  $kk'\gamma_1$  is

$$a_1(1 - e^{2i\pi j\gamma_1/\delta}) z_\infty^{kk'\gamma_1/\mu}.$$

As previously, this term is non-zero when  $j$  is an integer multiple of  $\delta/(\gamma_1, \delta)$ . It follows that  $q_0 - q'_j$  has degree strictly less than  $kk'\gamma_1$  when  $j$  is both a multiple of  $\delta/(\gamma_1, \delta)$  and of  $\delta/(\gamma_0, \delta)$ , i.e. is a multiple of their lowest common multiple  $\text{lcm}\left(\frac{\delta}{(\gamma_0, \delta)}, \frac{\delta}{(\gamma_1, \delta)}\right)$ . Since we have the equality

$$\text{lcm}\left(\frac{\delta}{(\gamma_0, \delta)}, \frac{\delta}{(\gamma_1, \delta)}\right) = \frac{\delta}{(\gamma_0, \gamma_1, \delta)},$$

we conclude that the number of differences  $q_0 - q'_j$ ,  $j = 0, \dots, \delta - 1$  having degree strictly less than  $kk'\gamma_1$  is equal to  $(\gamma_0, \gamma_1, \delta)$ . Therefore, the number of differences having degree equal to  $kk'\gamma_1$  is  $(\gamma_0, \delta) - (\gamma_0, \gamma_1, \delta)$ . Each of these differences gives rise to  $kk'\gamma_1$  Stokes arrows from  $\langle q \rangle$  to  $\langle q' \rangle$ .

The same reasoning can be carried out for the next terms in  $q_c = q'_c$ . By induction, we find that for  $1 \leq i \leq r$ , the number of differences  $q_0 - q'_j$ ,  $j = 0, \dots, \delta - 1$  with degree  $kk'\gamma_i$  is equal to the difference  $(\gamma_0, \dots, \gamma_{i-1}, \delta) - (\gamma_0, \dots, \gamma_i, \delta)$ , and each of those circles gives rise to  $kk'\gamma_i$  Stokes arrows between  $\langle q \rangle$  and  $\langle q' \rangle$ .

There only remain  $(\gamma_0, \dots, \gamma_{r-1}, \delta)$  differences having degree strictly less than  $kk'\gamma_{r-1}$ . For those differences, we have  $q_{c,0} - q'_{c,j} = 0$ , so only the different parts matter:  $q_0 - q'_j = q_{d,0} - q'_{d,j}$ . Now, the leading term of  $q_d$  is  $b_r z_\infty^{-\alpha_r/\beta} = b_r z_\infty^{-k\alpha_r/\mu}$ , so  $q_d$  has degree  $k\alpha_r$  as a polynomial in  $z^{-1/\mu}$ , and  $q'_d$  has degree  $k'\alpha'_r$  in  $z^{-1/\mu}$ . It follows that the degree of  $q_0 - q'_j$  is  $\max(k\alpha_r, k'\alpha'_r)$ , since we have assumed that  $\alpha_r/\beta \geq \alpha'_r/\beta'$ , the maximum is  $k\alpha_r$ . Each of these thus accounts for  $k\alpha_r$  Stokes arrows from  $\langle q \rangle$  to  $\langle q' \rangle$ .

Adding all contributions, and subtracting the  $\beta\beta'$  arrows appearing in the definition of  $B_{\langle q \rangle, \langle q' \rangle}$ , we get the following expression for the number of edges between  $\langle q \rangle$  and  $\langle q' \rangle$ :

$$B_{\langle q \rangle, \langle q' \rangle} = (\delta - (\gamma_0, \delta))kk'\gamma_0 + ((\gamma_0, \delta) - (\gamma_0, \gamma_1, \delta))kk'\gamma_1 + \dots + ((\gamma_0, \dots, \gamma_{r-2}, \delta) - (\gamma_0, \dots, \gamma_{r-1}, \delta))kk'\gamma_{r-1} + (\gamma_0, \dots, \gamma_{r-1}, \delta)k\alpha_r - \beta\beta'.$$

Since  $k\delta = \beta'$ ,  $k\gamma_i = \alpha'_i$ , and  $k'\gamma_i = \alpha_i$ , this yields the desired formula.

The case where  $q$  and  $q'$  have no common part corresponds to having  $r = 0$ . In the formula, all terms corresponding to the common part disappear, there only remains  $B_{\langle q \rangle, \langle q' \rangle} = \beta'(\alpha - \beta)$ .  $\square$

The similar formula for the number of loops is

**Lemma A.2.** *Let  $q = \sum_{j=0}^p b_j z_\infty^{-\alpha_j/\beta}$  be an exponential factor of slope  $\alpha_0/\beta > 1$  as before.*

- One has

(A.2)

$$B_{q,q} = (\beta - (\alpha_0, \beta))\alpha_0 + ((\alpha_0, \beta) - (\alpha_0, \alpha_1, \beta))\alpha_1 + \dots + ((\alpha_0, \dots, \alpha_{p-1}, \beta) - (\alpha_0, \dots, \alpha_p, \beta))\alpha_p - \beta^2 + 1.$$

- Otherwise, if  $(\alpha, \beta) = 1$ , then we have

(A.3)

$$B_{q,q} = (\beta - 1)(\alpha - \beta - 1).$$

*Proof.* The proof is similar to the case of two different circles. As previously we have

$$\text{Irr}(\text{End}(\langle q \rangle)) = \sum_{i=0}^{\beta-1} \sum_{j=0}^{\beta-1} \text{slope}(q_i - q_j) = \sum_{j=0}^{\beta-1} \text{slope}(q_0 - q_j).$$

Among those  $\beta$  differences, we determine how many have degree  $\alpha_0, \dots, \alpha_p$ , as polynomials in  $z^{-1/\beta}$ , and the remaining differences will have degree 0. We find that the number of differences with degree  $\alpha_0$  is  $\beta - (\alpha_0, \beta)$ , the number of differences with degree  $\alpha_1$  is  $(\alpha_0, \beta) - (\alpha_0, \alpha_1, \beta)$ , etc. One has  $(\alpha_0, \dots, \alpha_{p-1}, \beta) - (\alpha_0, \dots, \alpha_p, \beta)$  differences with degree  $\alpha_p$ , and the  $(\alpha_0, \dots, \alpha_p, \beta)$  remaining differences belong to connected components that are copies of  $\langle 0 \rangle$ . Each difference of degree  $\alpha_i$  accounts for  $\alpha_i$  Stokes arrows. The total number of (positive) Stokes arrows is thus

$$(\beta - (\alpha_0, \beta))\alpha_0 + ((\alpha_0, \beta) - (\alpha_0, \alpha_1, \beta))\alpha_1 + \dots + ((\alpha_0, \dots, \alpha_{p-1}, \beta) - (\alpha_0, \dots, \alpha_p, \beta))\alpha_p.$$

To this we must subtract a number of arrows equal to  $\beta(\beta - 1) + (\beta - 1) = \beta^2 - 1$ , which gives the desired formula.

When  $\alpha$  and  $\beta$  are relatively prime, only the first term remains: the number of positive Stokes arrows is  $(\beta - 1)\alpha$ , and the conclusion follows.  $\square$

## APPENDIX B. FORM OF THE LEGENDRE TRANSFORM

We now prove the formula for the levels of the Legendre transform (lemma 4.6).

**Lemma B.1.** *Let  $q = \sum_{j=0}^p b_j z^{\alpha_j/\beta}$ , with  $b_i \neq 0$ , be an exponential factor at infinity. We set  $\alpha := \alpha_0$ , so the slope of  $q$  is  $\alpha/\beta > 1$ . Then the exponents of  $\xi$  possibly appearing with non-zero coefficients in its Legendre transform  $\tilde{q}$  are of the form  $\frac{\alpha - k_1(\alpha - \alpha_1) - \dots - k_p(\alpha - \alpha_p)}{\alpha - \beta}$ , with  $k_1, \dots, k_p \geq 0$ . More precisely, if we set  $E := \{\gamma \in \mathbb{N} \mid \exists k_1, \dots, k_p \in \mathbb{N}, \gamma = k_1(\alpha - \alpha_1) + \dots + k_p(\alpha - \alpha_p)\}$  the Legendre transform has the form*

$$(B.1) \quad \tilde{q}(\xi) = \sum_{\substack{\gamma \in E \\ \alpha - \gamma > 0}} \tilde{b}_\gamma \xi^{\frac{\alpha - \gamma}{\alpha - \beta}},$$

where the sum is restricted to the terms such that the exponent  $\frac{\alpha - \gamma}{\alpha - \beta}$  is positive. Furthermore, the coefficients  $\tilde{b}_{(\alpha - \alpha_i)}$  are non-zero for  $i \geq 1$ .

*Proof.* The proof consists in computing the Legendre transform directly from the system of equations by which it is defined, in a way explicit enough to find the order of the terms which appear. The first equation of the system is

$$(B.2) \quad \frac{dq}{dz} = \xi,$$

it is interpreted as determining  $\xi$  as a function of  $z$ . We will show that this implies  $\xi$  is of the form

$$(B.3) \quad z = \sum_{\gamma \in E} c_\gamma \xi^{\frac{\beta - \gamma}{\alpha - \beta}}.$$

The second equation of the system then yields  $\frac{d\tilde{q}}{d\xi} = -z$ , and the lemma follows by integrating.

Equation (B.2) has a solution  $z(\xi)$  in the field of Puiseux series in the variable  $\xi$  which is unique once we fix a choice of  $\alpha - \beta$ -th root. To show (B.3), we thus can take this form of solution as an ansatz, and check that it gives a unique solution for the coefficients of this expression: this will automatically be the solution  $z(\xi)$ . So let us assume (B.3). We set

$$\frac{dq}{dz} = \sum_{j=0}^p b'_j z^{\frac{\alpha_j - \beta}{\beta}}.$$

This implies

$$\xi = \frac{dq}{dz} = \sum_{j=0}^p b'_j \left( \sum_{\gamma \in E} c_\gamma \xi^{\frac{\beta - \gamma}{\alpha - \beta}} \right)^{\frac{\alpha_j - \beta}{\beta}}.$$



We develop this expression to identify the coefficients. We have

$$\begin{aligned}
 \xi &= \sum_{j=0}^p b'_j \left( \sum_{\gamma} c_{\gamma} \xi^{\frac{\beta-\gamma}{\alpha-\beta}} \right)^{\frac{\alpha_j-\beta}{\beta}} \\
 &= \sum_{j=0}^p b'_j \xi^{-\frac{\alpha_j-\beta}{\alpha-\beta}} \left( \sum_{\gamma} c_{\gamma} \xi^{\frac{-\gamma}{\alpha-\beta}} \right)^{\frac{\alpha_j-\beta}{\beta}} \\
 &= \sum_{j=0}^p b''_j \xi^{1-\frac{\alpha-\alpha_j}{\alpha-\beta}} \left( 1 + \sum_{\gamma \neq 0} c'_{\gamma} \xi^{\frac{-\gamma}{\alpha-\beta}} \right)^{\frac{\alpha_j-\beta}{\beta}} \\
 &= \sum_{j=0}^p b''_j \xi^{1-\frac{\alpha-\alpha_j}{\alpha-\beta}} \left( \sum_{(l_{\gamma})_{\gamma \neq 0}} \prod_{\{\gamma, l_{\gamma} \neq 0\}} A_{l_{\gamma}}^{(j)} c'_{\gamma}{}^{l_{\gamma}} \xi^{\frac{-l_{\gamma}\gamma}{\alpha-\beta}} \right) \\
 &= \sum_{j, (l_{\gamma})} b''_j \left( \prod_{\{\gamma, l_{\gamma} \neq 0\}} A_{l_{\gamma}}^{(j)} c'_{\gamma}{}^{l_{\gamma}} \right) \xi^{\left( 1 - \frac{(\alpha-\alpha_j) + \sum_{\gamma} l_{\gamma}\gamma}{\alpha-\beta} \right)} \\
 &= \sum_{\delta \geq 0} d_{\delta} \xi^{1-\frac{\delta}{\alpha-\beta}},
 \end{aligned}$$

with

$$(B.4) \quad d_{\delta} = \sum_{\substack{j, (l_{\gamma}) \\ \sum_{\gamma} l_{\gamma}\gamma + (\alpha-\alpha_j) = \delta}} b''_j \left( \prod_{\{\gamma, l_{\gamma} \neq 0\}} A_{l_{\gamma}}^{(j)} c'_{\gamma}{}^{l_{\gamma}} \right).$$

In the calculation, we have set  $b''_j = b'_j c_0^{(\alpha_j-\beta)/\beta}$ ,  $c'_{\gamma} = c_{\gamma}/c_0$ , and the  $A_{l_{\gamma}}^{(j)}$  are the combinatorial coefficients appearing in the series expansion of the term  $(1 + \dots)^{(\alpha_j-\beta)/\beta}$ .

The crucial point in this computation is that the integers  $\delta = \sum_{\gamma} l_{\gamma}\gamma + (\alpha - \alpha_j)$  still belong to the set of exponents  $E$ . This is the reason why our ansatz is correct. This enables us to find the coefficients  $c_{\gamma}$  by induction:

- For  $\delta = 0$ , we have  $d_0 = b''_0$ . By comparing the terms in  $\xi$ , we find  $1 = d_0 = b''_0$ , then  $c_0$ . This is where the choice of  $\alpha - \beta$ -th root takes place.
- Let  $\delta \in E$ . Assume that we know all  $c_{\gamma}$  for  $\gamma \in E$  such that  $\gamma \leq \delta$ . Let  $\delta'$  be the smallest element of  $E$  strictly greater than  $\delta$ . Equation (B.2) gives  $d_{\delta'} = 0$ . But  $\delta'$  is a sum of terms featuring the  $c'_{\gamma}$  for  $\gamma \leq \delta'$ , and of a single term featuring  $d'_{\delta'}$ , equal to

$$b''_0 A_1^{(0)} c'_{\delta_1} = \frac{\alpha_0 - \beta}{\beta} c'_{\delta'}$$

since we have  $A_1^{(0)} = \frac{\alpha_0 - \beta}{\beta} \neq 0$ . All the other terms being known by induction hypothesis, this determines  $c'_{\delta'}$  and hence  $c_{\delta'}$ , in a unique way.

It remains to see that the first subleading term associated to each  $\alpha_i$ , that is  $\tilde{b}_{(\alpha-\alpha_i)}$ , is non-zero. For this we look at the coefficient  $\delta_{(\alpha-\alpha_j)}$ . Equation (B.3) implies that  $\delta_{(\alpha-\alpha_j)} = 0$ . On the other hand  $\delta_{(\alpha-\alpha_j)}$  is given by (B.4) with  $\delta = \alpha - \alpha_j$ . Let us write down this equation more explicitly. There are only two decompositions  $\alpha - \alpha_j$  of the form

$$\alpha - \alpha_j = \sum_{\gamma \in E} l_{\gamma}\gamma + (\alpha - \alpha_k),$$

with  $l_{\gamma} \geq 0, k \in \{0, \dots, p\}$ :

$$\alpha - \alpha_j = 0 + (\alpha - \alpha_j), \text{ and } \alpha - \alpha_j = 1 \times (\alpha - \alpha_j) + \underbrace{(\alpha - \alpha_0)}_{=0}.$$

Equation (B.4) thus yields

$$0 = d_{\alpha-\alpha_j} = b_j'' \times 1 + b_0'' A_{\alpha-\alpha_j}^{(j)} c'_{\alpha-\alpha_j}.$$

Since  $b_0'', b_j''$  and  $A_{\alpha-\alpha_j}^{(j)}$  are non-zero, this implies that  $c'_{\alpha-\alpha_j} \neq 0$ , and the conclusion follows.  $\square$

### APPENDIX C. COMPUTATION OF THE DIMENSION

We now turn to the proof of the formula for the dimension of the wild character variety (theorem 6.1):

**Theorem C.1.** *The dimension of the wild character variety  $\mathcal{M}_B(\Theta, \mathcal{C})$ , when it is non-empty, is given by*

$$(C.1) \quad \dim \mathcal{M}_B(\Theta, \mathcal{C}) = 2 - (\mathbf{d}, \mathbf{d}),$$

where  $(\cdot, \cdot)$  is the bilinear form associated with the diagram  $\Gamma(\Theta, \mathcal{C})$ , and  $\mathbf{d}$  is the dimension vector of  $\Gamma(\Theta, \mathcal{C})$ .

To prove this, we are going to use the twisted quasi-hamiltonian description of the wild character variety given in [11], to compute its dimension. Keeping previous notations, the wild character variety  $\mathcal{M}_B(V)$  is obtained as a symplectic reduction of the twisted quasi-Hamiltonian space

$$\mathrm{Hom}_{\mathbb{S}}(V) \simeq \mathcal{A}(V_{a_1}^0) \otimes \cdots \otimes \mathcal{A}(V_{a_m}^0) // G.$$

Here, each piece  $\mathcal{A}(V_{a_k}^0)$  is a twisted quasi-Hamiltonian  $H(\partial_{a_k}) \times G$ -space:

$$\mathcal{A}(V_{a_k}^0) = H(\partial_{a_k}) \times G \times \prod_{d \in \mathbb{A}_{a_k}} \mathrm{Sto}_d,$$

where  $\mathbb{A}_{a_k}$  is the set of singular directions of the connection at  $a_k$ , and  $\mathrm{Sto}_d$  is the Stokes group associated to the singular direction  $d \in \mathbb{A}_{a_k}$ . Hence,  $\mathrm{Hom}_{\mathbb{S}}(E, \nabla)$  is a twisted quasi-Hamiltonian  $\mathbf{H} \times G$ -space, where  $\mathbf{H} = H_1 \times \cdots \times H_m$ , with  $H_i = \mathrm{GrAut}(V_{a_i}^0)$ .

The formal monodromies  $\rho_k \in H(\partial_{a_k})$  determine twisted conjugacy classes  $\mathcal{C}(\partial_{a_k}) \subset H(\partial_{a_k})$  which do not depend on the choice of direction  $d \in \partial_{a_k}$  and the choice of isomorphism  $V_d^0 \simeq \mathbb{C}^{\sum_j n_i^{(k)} \beta_i^{(k)}}$ ; where the  $n_i^{(k)}$  denote the multiplicities of the active circles at  $a_i$ , and  $\beta_i^{(k)}$  their ramifications (the tame circle is included). The wild character variety is the twisted quasi-Hamiltonian reduction of  $\mathrm{Hom}_{\mathbb{S}}(E, \nabla)$  at the twisted conjugacy class  $\mathcal{C} := \prod_{k=1}^m \mathcal{C}(\partial_{a_k})$ , i.e.

$$(C.2) \quad \mathcal{M}_B(V) = \mathrm{Hom}_{\mathbb{S}}(E, \nabla) //_{\mathcal{C}} \mathbf{H}.$$

This description enables us to compute the dimension of  $\mathcal{M}_B(E, \nabla)$ . One has

$$(C.3) \quad \dim \mathcal{A}(V_{a_k}^0) = \dim G + \dim H(\partial_{a_k}) + \dim \prod_{d \in \mathbb{A}_{a_k}} \mathrm{Sto}_d.$$

The dimension of  $G$  is  $\dim G = n^2$ . One has  $\dim H(\partial_{a_k}) = \sum_i \beta_i^{(k)} n_i^{(k)2}$ . The dimension of the product of the Stokes groups can be expressed as a function of the number of Stokes arrows between the active circles (see [7]):

$$(C.4) \quad \dim \prod_{d \in \mathbb{A}_{a_k}} \mathrm{Sto}_d = \sum_{1 \leq i, j \leq r_k} n_i^{(k)} n_j^{(k)} B_{i,j}^+,$$

where  $B_{i,j}^+$  denotes the number of (positive) Stokes arrows between  $\langle q_i^{(k)} \rangle$  and  $\langle q_j^{(k)} \rangle$  in the Stokes diagram corresponding to the formal local system  $V_{a_i}^0 \rightarrow \partial_{a_i}$ .

The dimension of  $\mathrm{Hom}_{\mathbb{S}}(V)$  is

$$\dim \mathrm{Hom}_{\mathbb{S}}(V) = \sum_{i=1}^m \dim \mathcal{A}(V_{a_k}^0) - 2 \dim G + 2 \dim Z(G),$$

and the dimension of the wild character variety is then, taking into account the symplectic reduction at the conjugacy classes  $\mathcal{C}(\partial_{a_k})$  is given by

$$(C.5) \quad \dim \mathcal{M}_B(V) = \sum_{k=1}^m \left( \dim \mathcal{A}(V_{a_k}^0) + \dim \mathcal{C}(\partial_{a_k}) - 2 \dim H(\partial_{a_k}) \right) - 2 \dim G + 2 \dim Z(G).$$

We now are in position to compute the quantities  $D := 2 - (\mathbf{d}, \mathbf{d})$  and  $D' := \dim \mathcal{M}_B(V)$ . We recall our notations:

- Let  $a_1, \dots, a_m$  be the singularities at finite distance.
- For  $k = 1, \dots, m$ , let  $\langle q_1^{(k)} \rangle_{a_k}, \dots, \langle q_{r_k}^{(k)} \rangle_{a_k} \in \pi_0(\mathcal{I}_{a_k})$  denote the irregular active circles at  $a_k$ ,  $n_i^{(k)} \in \mathbb{N}$  their respective multiplicities, and  $\beta_j^{(k)}$  their respective ramification orders, and  $\alpha_j^{(k)}$  their irregularities, in an order such that the slopes satisfy

$$\frac{\alpha_1^{(k)}}{\beta_1^{(k)}} \geq \dots \geq \frac{\alpha_{r_k}^{(k)}}{\beta_{r_k}^{(k)}}.$$

- For  $k = 1, \dots, m$ , let  $m_k$  be the multiplicity of the tame circle  $\langle 0 \rangle_{a_k}$  in the modified formal local system at  $a_k$ .
- At infinity, let  $\langle q_1^{(\infty)} \rangle_{\infty}, \dots, \langle q_{r_{\infty}}^{(\infty)} \rangle_{\infty} \in \pi_0(\mathcal{I}_{a_k})$  denote the active circles (the tame circle is included),  $\beta_j^{(\infty)}$  their respective ramification orders, and  $\alpha_j^{(\infty)}$  their irregularities, in an order such that the slopes satisfy

$$\frac{\alpha_1^{(\infty)}}{\beta_1^{(\infty)}} \geq \dots \geq \frac{\alpha_{r_{\infty}}^{(\infty)}}{\beta_{r_{\infty}}^{(\infty)}}.$$

Here we will assume that all Stokes arrows come from the leading terms in the exponential factors. In the language of the previous section, this means that the different exponential factors have no common part. This entails no loss of generality, indeed in the computation of the number of Stokes arrows we have seen that the terms involving subleading terms do not change under Fourier transform, so that those term will give the same contribution to  $2 - (\mathbf{d}, \mathbf{d})$  and  $\dim \mathcal{M}_B(\Theta, \mathcal{C})$ .

Let us list the number of loops and edges between the different types of circles. From the formulas for the number of loops and edges, and the formulas the the transformation of slopes under Fourier transform, we get

- Loops at circles at infinity:  $B_{\langle q_i^{(\infty)} \rangle, \langle q_i^{(\infty)} \rangle} = (\beta_i^{(\infty)} - 1)(\alpha_i^{(\infty)} - \beta_i^{(\infty)} - 1)$ .
- Edges between different circles at infinity:  $B_{\langle q_i^{(\infty)} \rangle, \langle q_j^{(\infty)} \rangle} = \beta_j^{(\infty)}(\alpha_i^{(\infty)} - \beta_i^{(\infty)})$  if  $i < j$ .
- Loops at irregular circles at finite distance:  $B_{\langle q_i^{(k)} \rangle, \langle q_i^{(k)} \rangle} = (-\beta_i^{(k)} - 1)(\alpha_i^{(k)} + \beta_i^{(k)} - 1)$ .
- Edges between different irregular circles at a same pole at finite distance:  $B_{\langle q_i^{(k)} \rangle, \langle q_j^{(k)} \rangle} = -\beta_i^{(k)}(\alpha_j^{(k)} + \beta_j^{(k)})$  if  $i < j$ .
- Edges between the tame circle  $\langle 0 \rangle_{a_k}$  and an irregular circle at  $a_k$ :  $B_{\langle 0 \rangle_{a_k}, \langle q_i^{(k)} \rangle} = -\beta_i^{(k)}$ .
- Edges between circles at two different poles at finite distance: they are none, i.e.  $B_{\langle q_i^{(k)} \rangle, \langle q_j^{(l)} \rangle} = 0$
- Edges between an irregular circle at finite distance and a circle at infinity:  $B_{\langle q_i^{(k)} \rangle, \langle q_j^{(\infty)} \rangle} = \beta_j^{(\infty)}(\alpha_i^{(k)} + \beta_i^{(k)})$ .
- Edges between the tame circle  $\langle 0 \rangle_{a_k}$  and a circle at infinity:  $B_{\langle 0 \rangle_{a_k}, \langle q_i^{(\infty)} \rangle} = \beta_j^{(\infty)}$ .

For each circle  $I$ , let  $D_{\mathbb{L}_I}$  be the contribution to  $2 - (\mathbf{d}, \mathbf{d})$  of the leg  $\mathbb{L}_I$  associated to  $I$ . Putting everything together, this gives us the following lengthy expression for  $D$ :

(C.6)

$$\begin{aligned}
2 - (\mathbf{d}, \mathbf{d}) = & 2 + \sum_{i^{(\infty)}} \left( -2 + (\beta_i^{(\infty)} - 1)(\alpha_i^{(\infty)} - \beta_i^{(\infty)} - 1) \right) n_i^{(\infty)2} \\
& + \sum_{i^{(\infty)} < j^{(\infty)}} 2\beta_j^{(\infty)}(\alpha_i^{(\infty)} - \beta_i^{(\infty)})n_i^{(\infty)}n_j^{(\infty)} + \sum_k \sum_{i^{(k)}} \left( -2 + (-\beta_i^{(k)} - 1)(\alpha_i^{(k)} + \beta_i^{(k)} - 1) \right) n_i^{(k)2} \\
& + \sum_k \sum_{i^{(k)} < j^{(k)}} -2\beta_i^{(k)}(\alpha_j^{(k)} + \beta_j^{(k)})n_i^{(k)}n_j^{(k)} + \sum_k \sum_{i^{(k)}} -2\beta_i^{(k)}m_k n_i^{(k)} \\
& + \sum_k \sum_{i^{(k)} < j^{(\infty)}} 2\beta_j^{(\infty)}(\alpha_i^{(k)} + \beta_i^{(k)})n_i^{(k)}n_j^{(\infty)} + \sum_k \sum_{j^{(\infty)}} 2\beta_j^{(\infty)}m_k n_j^{(\infty)} - 2 \sum_k m_k^2 \\
& + \sum_{i^{(\infty)}} D(\mathbb{L}_{\langle q_{i^{(\infty)}} \rangle}) + \sum_k \sum_{i^{(k)}} D(\mathbb{L}_{\langle q_i^{(k)} \rangle}) + \sum_k D(\mathbb{L}_{\langle 0 \rangle_{a_k}}).
\end{aligned}$$

To compute the dimension of the wild character variety, we have to find the number of (positive) Stokes arrows at each singularity. We have

- Number of Stokes arrows between two irregular circles  $\langle q_i^{(k)} \rangle, \langle q_j^{(k)} \rangle$  with  $i < j$  at  $a_k$ :  $\beta_j^{(k)}\alpha_i^{(k)}$
- Number of Stokes loops at the irregular circle  $\langle q_i^{(k)} \rangle$ :  $\alpha_i^{(k)}(\beta_i^{(k)} - 1)$ .
- Number of Stokes arrows between the irregular circle  $\langle q_i^{(k)} \rangle$  and the tame circle  $\langle 0 \rangle_{a_k}$  at  $a_k$ :  $\alpha_i^{(k)}$ .

We also have the similar numbers of arrows at infinity.

Let  $n_k$  be the multiplicity of the tame circle  $\langle 0 \rangle_{a_k}$  in the non-modified local system  $V_{a_k}^0$ , i.e. the rank of the regular part of the connection at  $a_k$ . The dimension of the torsor  $H(\partial_{a_k})$  is

$$\dim H(\partial_{a_k}) = \sum_{i^{(k)}} n_i^{(k)2} \beta_i^{(k)} + n_k^2.$$

Therefore, we get from (C.3) the following expression for  $\dim \mathcal{A}(V_{a_k}^0)$ :

$$(C.7) \quad \dim \mathcal{A}(V_{a_k}^0) = n^2 + \sum_{i^{(k)}} (n_i^{(k)2} \beta_i^{(k)} + n_k^2) + \sum_{i^{(k)}} \alpha_i^{(k)}(\beta_i^{(k)} - 1)n_i^{(k)2} + \sum_{i^{(k)} < j^{(k)}} 2\beta_j^{(k)}\alpha_i^{(k)}n_i^{(k)}n_j^{(k)} + \sum_{i^{(k)}} 2\alpha_i^{(k)}n_i^{(k)}n_k.$$

Let  $\mathcal{C}_i^{(k)}$  be the twisted conjugacy class of the monodromy of the irregular circle  $\langle q_i^{(k)} \rangle$ , and  $\mathcal{C}_k^{\text{reg}}$  the conjugacy class of the monodromy of the tame circle in  $V_{a_k}^0$ , so that

$$\mathcal{C}(\partial_{a_k}) \simeq \prod_{i^{(k)}} \mathcal{C}_i^{(k)} \times \mathcal{C}_k^{\text{reg}} \subset H(\partial_{a_k}).$$

Using  $\dim Z(G) = 1$ , the formula (C.5) for the dimension of the wild character variety then becomes

$$\begin{aligned}
 \dim \mathcal{M}_B(V) = & 2 - 2n^2 \\
 & + \sum_k \left( n^2 - \sum_{i^{(k)}} (n_i^{(k)})^2 \beta_i^{(k)} + n_k^2 + \sum_{i^{(k)}} \alpha_i^{(k)} (\beta_i^{(k)} - 1) n_i^{(k)2} \right. \\
 & \left. + \sum_{i^{(k)} < j^{(k)}} 2\beta_j^{(k)} \alpha_i^{(k)} n_i^{(k)} n_j^{(k)} + \sum_{i^{(k)}} 2\alpha_i^{(k)} n_i^{(k)} n_k \right) \\
 \text{(C.8)} \quad & + n^2 - \sum_{i^{(\infty)}} n_i^{(\infty)2} \beta_i^{(\infty)} + \sum_{i^{(\infty)}} \alpha_i^{(\infty)} (\beta_i^{(\infty)} - 1) n_i^{(\infty)2} \\
 & + \sum_{i^{(\infty)} < j^{(\infty)}} 2\beta_j^{(\infty)} \alpha_i^{(\infty)} n_i^{(\infty)} n_j^{(\infty)} \\
 & + \sum_k \left( \sum_{i^{(k)}} \dim \mathcal{C}_i^{(k)} + \dim \mathcal{C}_k^{\text{reg}} \right) + \sum_{i^{(\infty)}} \dim \mathcal{C}_i^{(\infty)} + \dim \mathcal{C}_\infty^{\text{reg}}
 \end{aligned}$$

We have to compare these two expressions. First, we relate the contributions of the legs to the dimension of the conjugacy classes.

**Dimensions of conjugacy classes.** We have to deal with a subtlety: the leg  $\mathbb{L}_I$  which we glue to a circle  $I$  of ramification order  $\beta$  over the pole  $a \in \mathbb{P}^1$  of the core of the diagram corresponds to the conjugacy class of the monodromy of the local system  $V_I^0 \rightarrow I$ , whereas the twisted conjugacy class  $\mathcal{C}(I) \subset H(I)$  is the one of the monodromy of  $V_I^0 \rightarrow \partial_a$ , i.e. the local system downstairs on the circle of directions  $\partial_a$ .

$$\begin{array}{ccc}
 V_I^0 & \xrightarrow{\phi} & I \\
 & \searrow \psi & \downarrow \pi \\
 & & \partial_a
 \end{array}$$

Let us fix a direction  $d \in \partial_a$ , let  $d_0, \dots, d_{\beta-1}$  be its preimages by the cover  $\pi : I \rightarrow \partial_a$ . The fibre of  $V_I^0$  over  $d$  is the direct sum

$$V_{I,d}^0 = V_{I,d_0}^0 \oplus \dots \oplus V_{I,d_{\beta-1}}^0,$$

where  $V_{I,d_i}^0$  is the fibre of  $\phi : V_I^0 \rightarrow I$  over  $d_i$ , and the indices  $i$  are in  $\mathbb{Z}/\beta\mathbb{Z}$ . The monodromy of  $V_I^0 \rightarrow \partial_a$  is given by the collection of linear applications

$$\rho_i : V_{I,d_i}^0 \rightarrow V_{I,d_{i+1}}^0, \quad i = 0, \dots, \beta - 1.$$

The monodromy of  $V_I^0 \rightarrow I$  is then given by the product

$$\rho = \rho_{\beta-1} \dots \rho_0 : V_{I,d_0}^0 \rightarrow V_{I,d_0}^0$$

The group  $H_I := GL(V_{I,d_0}^0) \times \dots \times GL(V_{I,d_{\beta-1}}^0)$  acts on the monodromy by

$$(k_0, \dots, k_{\beta-1}) \cdot \rho_i = k_{i+1} \rho_i k_i^{-1},$$

where  $\mathbf{k} = (k_0, \dots, k_{\beta-1}) \in H_I$ . The twisted conjugacy class  $\mathcal{C}_I$  is the orbit of  $(\rho_0, \dots, \rho_{\beta-1})$  under this action. This induces on  $\rho$  the transformation

$$\mathbf{k} \cdot \rho = k_0 \rho k_0^{-1}.$$

which is just the action of  $GL(V_{I,d_0}^0)$  by conjugation, so that its orbit, which we will denote  $\tilde{\mathcal{C}}_I$ , is the conjugacy class of the monodromy of  $V_I^0 \rightarrow I$ . Furthermore, each fibre of the map

$$\begin{aligned}
 \mathcal{C}_I & \rightarrow \tilde{\mathcal{C}}_I \\
 (\rho_i) & \mapsto \rho
 \end{aligned}$$

is isomorphic to  $GL(V_{I,d_1}^0) \times \cdots \times GL(V_{I,d_{\beta-1}}^0)$ . It follows from this that the dimension of  $\mathcal{C}_I$  is

$$(C.9) \quad \dim \mathcal{C}_I = n_I^2(\beta - 1) + \dim \tilde{\mathcal{C}}_I,$$

where  $n_I$  is the dimension of each fibre  $V_{I,d_i}^0$ .

The legs of the diagram correspond to the reduced conjugacy classes  $\tilde{\mathcal{C}}_I$ . The dimension of a conjugacy class is obtained from the associated leg in the following way (see again [6]):

**Lemma C.2.** *Let  $\mathcal{C} \subset GL_n(\mathbb{C})$  a conjugacy class,  $\mathbb{L}$  the associated leg (after choosing a marking), with  $\mathbf{d}$  its dimension vector. The dimension of  $\mathcal{C}$  is given by*

$$(C.10) \quad \dim \mathcal{C} = 2n^2 - (\mathbf{d}, \mathbf{d}).$$

We can now compare the different terms appearing in the quantities  $D = 2 - (\mathbf{d}, \mathbf{d})$  and  $D' = \dim \mathcal{M}_B(V)$ . Let us consider the terms in  $D$  involving the multiplicities  $m_k$  of the tame circles: their contribution to  $D$  is

$$\begin{aligned} & \sum_k \sum_{i^{(k)}} -2\beta_i^{(k)} m_k n_i^{(k)} + \sum_k \sum_{j^{(\infty)}} 2\beta_j^{(\infty)} m_k n_j^{(\infty)} - 2 \sum_k m_k^2 \\ &= -2 \sum_k n_k(n - n_k) - 2m_k^2 + 2 \sum_k m_k n \\ &= \sum_k 2m_k n_k - 2m_k^2, \end{aligned}$$

where we use that  $n = n_k + \sum_{i^{(k)}} \beta_i^{(k)} n_i^{(k)}$  for each  $k$ , and the similar formula at infinity. We can recognize that  $2m_k n_k - 2m_k^2$  is exactly the contribution of the first edge of the leg associated to a special marking of the conjugacy class of the monodromy of the tame circle  $\langle 0 \rangle_k$ ; i.e. the edge corresponding to the eigenvalue 1. From this we deduce that

$$(C.11) \quad D(\mathbb{L}_{\langle 0 \rangle_{a_k}}) + 2m_k n_k - 2m_k^2 = \dim \mathcal{C}_k^{\text{reg}}.$$

We also have for each irregular circle at finite distance

$$D(\mathbb{L}_{\langle q_i^{(k)} \rangle}) = \dim \tilde{\mathcal{C}}_i^{(k)},$$

and similarly for the circles at infinity

$$D(\mathbb{L}_{\langle q_i^{(k)} \rangle}) = \dim \tilde{\mathcal{C}}_i^{(k)}.$$

This already gives us equalities between some terms appearing in  $D$  and  $D'$ . Let us deal with the remaining terms. First, for each pole  $a_k$  at finite distance, we have an equality between the following term in  $D$ , which is the contribution of the irregular circles at  $a_k$

$$\begin{aligned} & \sum_{i^{(k)}} \left( -2 + (-\beta_i^{(k)} - 1)(\alpha_i^{(k)} + \beta_i^{(k)} - 1) \right) n_i^{(k)2} + \sum_{i^{(k)} < j^{(k)}} -2\beta_j^{(k)} (\alpha_i^{(k)} + \beta_i^{(k)}) n_i^{(k)} n_j^{(k)} \\ &+ \sum_{i^{(k)}, j^{(\infty)}} 2\beta_j^{(\infty)} (\alpha_i^{(k)} + \beta_i^{(k)}) n_i^{(k)} n_j^{(\infty)}, \end{aligned}$$

and the following term in  $D'$

$$\begin{aligned} & n^2 - \sum_{i^{(k)}} (n_i^{(k)2} \beta_i^{(k)} + n_k^2) + \sum_{i^{(k)}} \alpha_i^{(k)} (\beta_i^{(k)} - 1) n_i^{(k)2} \\ &+ \sum_{i^{(k)} < j^{(k)}} 2\beta_j^{(k)} \alpha_i^{(k)} n_i^{(k)} n_j^{(k)} + \sum_{i^{(k)}} 2\alpha_i^{(k)} n_i^{(k)} n_k + n_i^{(k)2} (\beta_i^{(k)} - 1), \end{aligned}$$

where the term  $n_i^{(k)2} (\beta_i^{(k)} - 1)$  comes from equation (C.9). To obtain this equality, we notice that since  $n = \sum_{j^{(\infty)}} \beta_j^{(\infty)} n_j^{(\infty)}$ ,

$$\sum_{i^{(k)}, j^{(\infty)}} 2\beta_j^{(\infty)} (\alpha_i^{(k)} + \beta_i^{(k)}) n_i^{(k)} n_j^{(\infty)} = n \sum_{i^{(k)}, j^{(\infty)}} (\alpha_i^{(k)} + \beta_i^{(k)}) n_i^{(k)},$$

then we decompose on the both sides the total rank:  $n = \sum_{i^{(k)}} \beta_i^{(k)} n_i^{(k)} + n_k$ . It is then a somewhat tedious but straightforward check that we have the same terms on both sides.

It only remains to consider the contributions to  $D$  and  $D'$  of the circles at infinity: we have an equality between the following term in  $D$

$$\sum_{i^{(\infty)}} \left( -2 + (\beta_i^{(\infty)} - 1)(\alpha_i^{(\infty)} - \beta_i^{(\infty)} - 1) \right) n_i^{(\infty)^2} + \sum_{i^{(\infty)} < j^{(\infty)}} 2\beta_j^{(\infty)} (\alpha_i^{(\infty)} - \beta_i^{(\infty)}) n_i^{(\infty)} n_j^{(\infty)}$$

and the following term in  $D'$ :

$$\begin{aligned} & -2n^2 + n^2 - \sum_{i^{(\infty)}} n_i^{(\infty)^2} \beta_i^{(\infty)} + \sum_{i^{(\infty)}} \alpha_i^{(\infty)} (\beta_i^{(\infty)} - 1) n_i^{(\infty)^2} \\ & + \sum_{i^{(\infty)} < j^{(\infty)}} 2\beta_j^{(\infty)} \alpha_i^{(\infty)} n_i^{(\infty)} n_j^{(\infty)} + \sum_{i^{(\infty)}} n_i^{(\infty)^2} (\beta_i^{(\infty)} - 1), \end{aligned}$$

where once again the term  $n_i^{(\infty)^2} (\beta_i^{(\infty)} - 1)$  comes from (C.9). These equalities exhaust all terms appearing in  $D$  and  $D'$ , so this concludes the proof.

## REFERENCES

- [1] D. Arinkin. Fourier transform and middle convolution for irregular D-modules. *arXiv preprint arXiv:0808.0699*, 2008.
- [2] D. Arinkin. Rigid irregular connections on  $\mathbb{P}^1$ . *Compositio Mathematica*, 146(5):1323–1338, 2010.
- [3] S. P. Balandin and V. V. Sokolov. On the Painlevé test for non-abelian equations. *Phys. Lett. A*, 246(3-4):267–272, 1998.
- [4] S. Bloch and H. Esnault. Local Fourier transforms and rigidity for  $D$ -modules. *Asian Journal of Mathematics*, 8(4):587–606, 2004.
- [5] P. Boalch. Irregular connections and Kac-Moody root systems. *arXiv preprint arXiv:0806.1050*, 2008.
- [6] P. Boalch. Simply-laced isomonodromy systems. *Publications mathématiques de l’IHÉS*, 116(1):1–68, 2012.
- [7] P. Boalch. Geometry and braiding of Stokes data; fission and wild character varieties. *Annals of Mathematics*, pages 301–365, 2014.
- [8] P. Boalch. Global Weyl groups and a new theory of multiplicative quiver varieties. *Geometry & Topology*, 19(6):3467–3536, 2016.
- [9] P. Boalch. Wild character varieties, meromorphic Hitchin systems and Dynkin diagrams. *Geometry and Physics: Volume 2: A Festschrift in Honour of Nigel Hitchin*, 2:433, 2018.
- [10] P. Boalch. Topology of the Stokes phenomenon. *arXiv preprint arXiv:1903.12612*, 2019.
- [11] P. Boalch and D. Yamakawa. Twisted wild character varieties. *arXiv preprint arXiv:1512.08091*, 2015.
- [12] P. Boalch and D. Yamakawa. Diagrams for nonabelian Hodge spaces on the affine line. *Comptes Rendus. Mathématique*, 358(1):59–65, 2020.
- [13] W. Crawley-Boevey. Geometry of the moment map for representations of quivers. *Compositio Mathematica*, 126(3):257–293, 2001.
- [14] W. Crawley-Boevey. On matrices in prescribed conjugacy classes with no common invariant subspace and sum zero. *Duke Mathematical Journal*, 118(2):339–352, 2003.
- [15] W. Crawley-Boevey and P. Shaw. Multiplicative preprojective algebras, middle convolution and the Deligne–Simpson problem. *Advances in Mathematics*, 201(1):180–208, 2006.
- [16] P. Deligne. Letter to B. Malgrange. 1978.
- [17] J. Fang. Calculation of local Fourier transforms for formal connections. *Science in China Series A: Mathematics*, 52(10):2195–2206, 2009.
- [18] H. Flaschka and A. C. Newell. Monodromy- and spectrum-preserving deformations. I. *Comm. Math. Phys.*, 76(1):65–116, 1980.
- [19] A. Graham-Squire. Calculation of local formal Fourier transforms. *Arkiv för Matematik*, 51(1):71–84, 2013.
- [20] K. Hiroe. Linear differential equations on the Riemann sphere and representations of quivers. *Duke Mathematical Journal*, 166(5):855–935, 2017.
- [21] K. Hiroe and T. Oshima. A classification of roots of symmetric kac-moody root systems and its application. In *Symmetries, integrable systems and representations*, pages 195–241. Springer, 2013.
- [22] K. Hiroe and D. Yamakawa. Moduli spaces of meromorphic connections and quiver varieties. *Advances in Mathematics*, 266:120–151, 2014.
- [23] N. Joshi, A. Kitaev, and P. Treharne. On the linearization of the Painlevé III–VI equations and reductions of the three-wave resonant system. *Journal of mathematical physics*, 48(10):103512, 2007.
- [24] H. Kawakami, A. Nakamura, and H. Sakai. Part 2. degeneration scheme of 4-dimensional Painlevé-type equations. In *4-dimensional Painlevé-type equations*, pages 25–111. Mathematical Society of Japan, 2018.

- [25] G. Laumon. Transformation de Fourier, constantes d'équations fonctionnelles et conjecture de Weil. *Publications Mathématiques de l'IHÉS*, 65:131–210, 1987.
- [26] R. G. Lopez. Microlocalization and stationary phase. *Asian Journal of Mathematics*, 8(4):747–768, 2004.
- [27] B. Malgrange. La classification des connexions irrégulières à une variable. In *Mathematics and physics (Paris, 1979/1982)*, volume 37 of *Progr. Math.*, pages 381–399. Birkhäuser Boston, Boston, MA, 1983.
- [28] B. Malgrange. *Equations différentielles à coefficients polynomiaux*. Progress in mathematics. Birkhäuser, 1991.
- [29] Y. Ohyama, H. Kawamuko, H. Sakai, and K. Okamoto. Studies on the Painlevé equations. V. Third Painlevé equations of special type PIII (D7) and PIII (D8). *J. Math. Sci. Univ. Tokyo*, 13(2):145–204, 2006.
- [30] K. Okamoto. Studies on the Painlevé equations. *Annali di Matematica pura ed applicata*, 146(1):337–381, 1986.
- [31] K. Okamoto. Studies on the Painlevé equations. III: Second and fourth Painlevé equations, PII and PIV. *Mathematische Annalen*, 275(2):221–255, 1986.
- [32] K. Okamoto. Studies on the Painlevé equations II. Fifth Painlevé equation PV. *Japanese journal of mathematics. New series*, 13(1):47–76, 1987.
- [33] K. Okamoto. Studies on the Painlevé equations IV. Third painlevé equation PIII. *Funkcial. Ekvac.*, 30:305–32, 1987.
- [34] K. Okamoto. The Painlevé equations and the Dynkin diagrams. In *Painlevé transcendents*, pages 299–313. Springer, 1992.
- [35] C. Sabbah. An explicit stationary phase formula for the local formal Fourier-Laplace transform. In *Singularities I*, volume 474 of *Contemp. Math.*, pages 309–330. Amer. Math. Soc., Providence, RI, 2008.
- [36] D. Xie. 3d mirror for argyres-douglas theories. *arXiv preprint arXiv:2107.05258*, 2021.

LABORATOIRE DE MATHÉMATIQUES D'ORSAY, UNIVERSITÉ PARIS-SACLAY, FRANCE  
Email address: [jean.doucot@math.u-psud.fr](mailto:jean.doucot@math.u-psud.fr)