

On a tracial version of Haemers bound

Li Gao*

Sander Gribling†

Yinan Li‡

May 20, 2022

Abstract

We extend upper bounds on the quantum independence number and the quantum Shannon capacity of graphs to their counterparts in the commuting operator model. We introduce a von Neumann algebraic generalization of the fractional Haemers bound (over \mathbb{C}) and prove that the generalization upper bounds the commuting quantum independence number. We call our bound the tracial Haemers bound, and we prove that it is multiplicative with respect to the strong product. In particular, this makes it an upper bound on the Shannon capacity. The tracial Haemers bound is incomparable with the Lovász theta function, another well-known upper bound on the Shannon capacity. We show that separating the tracial and fractional Haemers bounds would refute Connes' embedding conjecture.

Along the way, we prove that the tracial rank and tracial Haemers bound are elements of the (commuting quantum) asymptotic spectrum of graphs (Zuiddam, *Combinatorica*, 2019). We also show that the inertia bound (an upper bound on the quantum independence number) upper bounds the commuting quantum independence number.

1 Introduction

1.1 Independence number from nonlocal games

Two players, Alice and Bob, attempt to convince a referee that a given graph G contains an independent set of size d . The referee sends two integers $i_A, i_B \in [d]$ to Alice and Bob, respectively, who then response vertices g_A and g_B satisfying that

- if $i_A = i_B$ then $g_A = g_B$;
- if $i_A \neq i_B$ then $\{g_A, g_B\} \in E(\overline{G})$.

Alice and Bob can decide a strategy beforehand, but they cannot communicate during the game. Such a nonlocal game is known as the independent set game [MR16, MRV16]. A classical strategy corresponds to the case that Alice and Bob only use shared (classical) randomness. A perfect classical strategy enables Alice and Bob to win the independent set game for graph G with certainty. It can be shown that a perfect classical strategy exists if and only if the graph G has an independent set of size d . Thus, the largest possible d for a graph G where a perfect classical strategy for the independent set game exists is exactly the independence number $\alpha(G)$ of G .

One may also consider quantum strategies, where Alice and Bob are allowed to share quantum entanglement and perform measurements on their own systems. A perfect quantum strategy enables

*Department of Mathematics, University of Houston, Houston, TX, 77204, USA. lgao12@uh.edu

†IRIF, Université de Paris, CNRS, Paris, France. gribling@irif.fr

‡Graduate school of mathematics, Nagoya University, Nagoya, Japan. Yinan.Li@maths.nagoya-u.ac.jp

Alice and Bob to win the independent set game for a graph G with a potentially larger d . The largest such d leads to the definition of quantum versions of the independence number of graphs. Note that different choices of the underlying model of quantum mechanics result in different sets of quantum strategies. In the tensor-product model, the shared entangled state is modeled by a bipartite state in a tensor product Hilbert space $\mathcal{H}_A \otimes \mathcal{H}_B$, and the measurements performed by Alice and Bob are modeled as observables on \mathcal{H}_A and \mathcal{H}_B , respectively. Define the quantum independence number of G , $\alpha_q(G)$, as the largest integer d for which there exists a perfect quantum strategy in the tensor-product model for the independent set game. One may also consider the commuting-operator model, where the shared entangled state is given by a state in a joint Hilbert space \mathcal{H} , and the measurements performed by Alice and Bob are modeled by commuting observables on \mathcal{H} . One similarly defines the commuting quantum independence number of G , $\alpha_{qc}(G)$, as the largest integer d for which there exists a perfect quantum strategy in the commuting-operator model for the independent set game. It is straightforward to see that for any graph G we have the chain of inequalities

$$\alpha(G) \leq \alpha_q(G) \leq \alpha_{qc}(G).$$

1.2 Quantum independence number: motivation and bounds

The quantum independence number $\alpha_q(G)$ has been extensively studied in the settings of nonlocal games [MSS13, CMSS14, MR16, MRV16], combinatorics [Rob13, WEA19], and zero-error quantum information theory [CLMW10, LMM⁺12, BBG13, BBL⁺15]. It has been shown that perfect quantum strategies for the independent set game can always be achieved by performing projective measurements on a shared maximally entangled state (of a finite dimension) [MR16, Theorem 2.1]. A natural question is whether we can determine $\alpha_q(G)$ for a given graph G . As a computational problem, determining whether $\alpha_q(G) \geq d$ for some integer d is NP-hard [Ji13]. In fact, any algorithm for determining whether $\alpha_q(G) \geq d$ leads to an algorithm for determining whether a given nonlocal game admits perfect quantum strategies achieved by performing projective measurements on a shared maximally entangled state (of a finite dimension) [MRV16]. Note that the latter family of problems includes the one which determines whether a synchronous game admits perfect quantum strategies (in the tensor-product model).

Another important reason to study the quantum independence number arises from the study of the zero-error capacity of classical channels, where the sender and the receiver are allowed to share an entangled state to assist their communication. It is known that the maximal number of zero-error messages one can send through a classical channel assisted by quantum entanglement is determined by its confusability graph [CLMW10]. This correspondence leads to the definition of the entanglement-assisted independence number $\alpha_*(G)$ of a (confusability) graph G . From this information-theoretic point of view, the quantum independence number of the confusability graph G can be equivalently interpreted as the maximal number of zero-error messages one can send through the corresponding classical channel assisted by performing projective measurements on a maximally entangled state [MR16, Section 5]. In information theory, it is also natural to consider the asymptotic zero-error rates, which leads to the definitions of the entanglement-assisted Shannon capacity $\Theta_*(G)$ and the quantum Shannon capacity $\Theta_q(G)$ of the confusability graph G :

$$\Theta_q(G) = \lim_{k \rightarrow \infty} \sqrt[k]{\alpha_q(G^{\boxtimes k})} \text{ and } \Theta_*(G) = \lim_{k \rightarrow \infty} \sqrt[k]{\alpha_*(G^{\boxtimes k})}.$$

From the definitions, we have

$$\alpha(G) \leq \alpha_q(G) \leq \alpha_*(G) \text{ and } \Theta(G) \leq \Theta_q(G) \leq \Theta_*(G),$$

where $\alpha(G)$ and $\Theta(G)$ denote the standard independence number and Shannon capacity of a graph G .

Note that in the vanishing-error setting¹, quantum entanglement is not helpful for enhancing the communication through classical channels [BSST02]. A natural question is whether a quantum advantage can be found in the zero-error setting. In [CLMW10, BBL⁺15], specific zero-error encoding schemes (using maximally entangled state and projective measurements) are proposed, which lead to lower bounds on $\alpha_q(G)$ (and thus also lower bounds on $\alpha_*(G)$). These lower bounds have been used to separate $\alpha(G)$ and $\alpha_q(G)$ for graphs constructed from proofs of the Kochen-Specker theorem [CLMW10, MSS13]. Moreover, separations of $\Theta(G)$ and $\Theta_q(G)$ can be shown by constructing geometric graphs whose Haemers bounds over certain finite fields are strictly smaller than these lower bounds [LMM⁺12, BBG13, BBL⁺15]. Here, the Haemers bound $\mathcal{H}(G; \mathbb{F})$ of a graph G over any field \mathbb{F} is an important classical upper bound on the Shannon capacity $\Theta(G)$ [Hae78, Hae79].

On the other hand, the celebrated Lovász theta function $\vartheta(G)$ [Lov79], which is another important upper bound on the Shannon capacity $\Theta(G)$, also upper bounds $\alpha_*(G)$ and $\Theta_*(G)$ [Bei10, DSW13]. Two central open problems in quantum zero-error information theory are to determine whether $\Theta_*(G) = \vartheta(G)$ and $\alpha_q(G) = \alpha_*(G)$ for all graphs G . Recently, it was observed that these two equalities cannot both hold [LZ21]: $\alpha_q(G)$ and $\Theta_q(G)$ are upper bounded by the fractional Haemers bounds $\mathcal{H}_f(G; \mathbb{R})$ and $\mathcal{H}_f(G; \mathbb{C})$ [Bla13, BC19] over \mathbb{R} and \mathbb{C} and there exist graphs G such that $\mathcal{H}_f(G; \mathbb{R}) \leq \mathcal{H}(G; \mathbb{R}) < \vartheta(G)$. Such an observation was made during the study of quantum asymptotic spectra of graphs, an important set of graph parameters which provide dual characterizations for the Shannon capacity and its quantum variants, initialized by Zuiddam [Zui19, Zui18].

1.3 Commuting quantum independence number: known and our results

The central problem in the theory of nonlocal games is to understand the differences between the two different models of quantum mechanics, the (finite-dimensional) tensor-product model and the commuting-operator model. A recent breakthrough (preprint) of Ji, Natarajan, Vidick, Wright and Yuen shows that these two models are not equivalent, which refutes the celebrated Connes' embedding conjecture [JNV⁺20].

Among graph-theoretic nonlocal games, the commuting quantum chromatic number and commuting quantum homomorphism have been extensively studied in the past decade. These games are defined in terms of the existence of perfect commuting quantum strategies for the graph coloring and homomorphism games, respectively [PT15, PSS⁺16, OP16, DP16]. In the seminal paper [PSS⁺16], a semidefinite programming (SDP) hierarchy is proposed which converges to the commuting quantum chromatic number. They also propose a new lower bound, the tracial rank, of the commuting quantum chromatic number. The tracial rank can be viewed as a C^* -algebraic generalization of the projective rank, which was introduced in [MSS13] for lower bounding the quantum chromatic number.² Tracial noncommutative polynomial optimization can be used to unify existing bounds and obtain SDP hierarchies of bounds on the commuting quantum independence number [GdLL18].

In this paper, we further extend recent results on quantum independence numbers to commuting quantum independence numbers. In particular, we prove that several known bounds on the quantum independence number and quantum Shannon capacity are also upper bounds on the commuting quantum independence number and commuting quantum Shannon capacity.

We first establish a characterization of the commuting quantum Shannon capacity, defined as

$$\Theta_{qc}(G) = \lim_{k \rightarrow \infty} \sqrt[k]{\alpha_{qc}(G^{\boxtimes k})},$$

¹The error probability goes to 0 when the number of uses of the channel goes to infinity [Sha48].

²In this paper, we evaluate the tracial rank and the projective rank on the complement of a graph G .

in terms of the asymptotic spectrum of graphs [Zui19, Zui18]. This can be done by exhibiting desired properties of the commuting quantum homomorphism. Such characterizations indicate that, to compute the Shannon capacity and its quantum variants, it is important to find graph parameters that are additive (with respect to disjoint union), multiplicative (with respect to strong product), normalized (the value of \overline{K}_d is d for any $d \in \mathbb{N}$) and monotonic (with respect to the corresponding homomorphism). Note that almost all well-known upper bounds on the Shannon capacities satisfy these four properties, including the Lovász theta function $\vartheta(G)$, fractional Haemers bound $\mathcal{H}_f(G; \mathbb{F})$ over any field \mathbb{F} , the projective rank $\overline{\xi}_f(G)$ and the fractional clique cover number $\overline{\chi}_f(G)$. We prove that the tracial rank $\overline{\xi}_{tr}(G)$ is another such graph parameter.

Our main contribution is a tracial version of the fractional Haemers bound (over \mathbb{C}) that upper bounds the commuting quantum independence number of graphs. We call this new bound the tracial Haemers bound. The bound is motivated by the study of the Shannon capacity (resp. quantum Shannon capacity) of graphs, where the fractional Haemers bound over any field (resp. over \mathbb{C} and \mathbb{R}) and Lovász theta function are incomparable elements in the asymptotic spectrum (resp. quantum asymptotic spectrum) of graphs. We show that the tracial Haemers bound is an element of the commuting quantum asymptotic spectrum of graphs, and it is incomparable with the Lovász theta function.

Our tracial Haemers bound is defined by generalizing a coordinate-free definition of the fractional Haemers bound due to Lex Schrijver [BC19, Remark after Prop. 7] to the von Neumann algebraic setting. When restricting to finite-dimensional von Neumann algebras, we show that it reduces to the fractional Haemers bound (over \mathbb{C}). We then prove that the tracial Haemers bound satisfies many properties of the fractional Haemers bound: it is additive, multiplicative, normalized and monotonic with respect to the commuting quantum homomorphism. This shows that the tracial Haemers bound is another element in the commuting quantum asymptotic spectrum of graphs. Thus, we provide a (potentially) better upper bound on the Shannon capacity and its quantum variants. We show the tracial Haemers bound of odd cycles equals their fractional clique cover number. This implies that the tracial Haemers bound and the Lovász theta function are incomparable. To compare the tracial Haemers bound with fractional Haemers bound, we show that the existence of a graph G for which $\mathcal{H}_{tr}(G) < \mathcal{H}_f(G; \mathbb{C})$ would refute Connes' embedding conjecture.

We now have the following inequality which illustrates the relations between independence numbers, Shannon capacities and elements in the asymptotic spectrum of graphs:

$$\begin{array}{ccccccc} \alpha(G) & \leq & \alpha_q(G) & \leq & \alpha_{qc}(G) & & \\ | \wedge & & | \wedge & & | \wedge & & \\ \Theta(G) & \leq & \Theta_q(G) & \leq & \Theta_{qc}(G) & \leq & \vartheta(G) \leq \overline{\xi}_{tr}(G) \\ & & & & & & \mathcal{H}_{tr}(G) \leq \mathcal{H}_f(G; \mathbb{C}) \\ & & & & & & \leq \overline{\xi}_f(G) \leq \overline{\chi}_f(G), \end{array}$$

where $\vartheta(G)$ is incomparable with $\mathcal{H}_{tr}(G)$ and with $\mathcal{H}_f(G; \mathbb{C})$, and $\mathcal{H}_f(G; \mathbb{C})$ is not known to be comparable with $\overline{\xi}_{tr}(G)$. Moreover, our tracial Haemers bound provides a unified way to describe almost all the above elements in the asymptotic spectrum of graphs (except the Lovász theta function, cf. Propositions 4.6 and 4.8 and Corollary 4.7).

To derive the desired properties of the tracial Haemers bound, we use an infinite-dimensional version of the rank-nullity theorem. This theorem can be established using an explicit representation of the C^* -algebra generated by two projections [RS89], which plays an important role in our proofs and has also been used in the study of tracial rank [DP16]. As a byproduct, we prove that the inertia bound for the independence number [Cve73] and the quantum independence number [WE18, WEA19], is also an upper bound for the commuting quantum independence number. For this, we first upper bound the commuting quantum independence number by the tracial packing number, an infinite-dimensional generalization of the projective packing number introduced in [Rob13]. Then

we upper bound the tracial packing number by the inertia bound using a characterization of the largest dimension of totally isotropic subspaces of an operator by allowing the ancillary system to be a von Neumann algebra with a tracial state. Although the inertia bound is only supermultiplicative (thus it fails to upper bound the commuting quantum Shannon capacity), it is incomparable with the Lovász theta function and tracial Haemers bound. In particular, the inertia bound allows us to identify the commuting quantum independence number of graphs for which the inertia bound equals the classical independence number: famous examples include odd cycles, perfect graphs, Kneser graphs, Andrasfai graphs, the Petersen graph, the Clebsch graph and many others.

The rest of this paper is organized as follows. In Section 2 we list some necessary definitions and useful lemmas in graph theory and operator algebras, which will be useful for proving our results. In Section 3 we establish asymptotic spectral characterizations for the C^* -algebraic and commuting quantum Shannon capacity of graphs, and prove that the tracial rank is an element in the commuting quantum asymptotic spectrum of graphs. In Section 4, we introduce the tracial Haemers bound, exhibit desired properties to show that it is an element in the commuting quantum asymptotic spectrum of graphs, and compare it with the Lovász theta function and the fractional Haemers bound. In Section 5, we prove the inertia bound for independence number and quantum independence number is also an upper bound for the commuting quantum independence number.

2 Preliminaries

2.1 Notation and preliminaries about graphs

We work with unweighted and undirected graphs with no loops, no parallel edges and finitely many vertices. For a graph G , $V(G)$ denotes its vertex set and $E(G) \subseteq \{\{g, g'\} : g, g' \in V(G) \text{ with } g \neq g'\}$ denotes its edge set. For $g \in V(G)$, let $N_G(g) = \{g' \in V(G) : \{g, g'\} \in E(G)\}$ be the set of vertices which is adjacent to g . The complement graph of G is the graph $\overline{G} = (V(G), E(\overline{G}))$, where $E(\overline{G}) = \{\{g, g'\} : g, g' \in V(G) \text{ with } g \neq g' \text{ and } \{g, g'\} \notin E(G)\}$. For $d \in \mathbb{N}$, let $[d] := \{1, \dots, d\}$ and K_d denotes the complete graph on d vertices with edge set $\{\{i, j\} : i, j \in [d] \text{ with } i \neq j\}$. By convention let $K_0 = (\emptyset, \emptyset)$ and $K_1 = ([1], \emptyset)$. Given two graphs G and H , their *disjoint union* is the graph $G \sqcup H$ with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. The *strong product* of two graphs G and H is the graph $G \boxtimes H$ with vertex set $V(G) \times V(H)$ and edge set

$$E(G \boxtimes H) = \left\{ \{(g, h), (g', h')\} : \begin{aligned} &\{g, g'\} \in E(G) \text{ and } \{h, h'\} \in E(H), \\ &\text{or } g = g' \in V(G) \text{ and } \{h, h'\} \in E(H), \\ &\text{or } \{g, g'\} \in E(G) \text{ and } h = h' \in V(H) \end{aligned} \right\}.$$

A *graph homomorphism* from G to H is an edge-preserving vertex map $f : V(G) \rightarrow V(H)$, i.e., a map such that $\{g, g'\} \in E(G)$ implies $\{f(g), f(g')\} \in E(H)$. The notation $G \rightarrow H$ is used to denote that there exists a graph homomorphism from G to H and the notation $G \leq_{loc} H$ means $\overline{G} \rightarrow \overline{H}$. The map $f : V(G) \rightarrow V(H)$ is an isomorphism if it is a bijection that preserves edges and non-edges. Graph isomorphism is an equivalence relation on the set of graphs and we let $\{\text{Graphs}\}$ denote the set of isomorphism classes of graphs. Note that the disjoint union \sqcup is an associative and commutative binary operation (addition) on $\{\text{Graphs}\}$ with identity element K_0 , and the strong product \boxtimes is an associative and commutative binary operation (multiplication) on $\{\text{Graphs}\}$ with identity element K_1 . Thus $\mathcal{G} = (\{\text{Graphs}\}, \sqcup, \boxtimes, K_0, K_1)$ is a *semiring*, called the semiring of graphs. The relation \leq_{loc} defines a preorder on \mathcal{G} , i.e., it satisfies $G \leq_{loc} G$, and $G \leq_{loc} H, H \leq_{loc} K$ implies $G \leq_{loc} K$ for any $G, H, K \in \mathcal{G}$. The following notions of semiring homomorphism and asymptotic spectrum of graphs play a central role in this paper.

Definition 2.1 (Asymptotic spectrum of graphs). *A spectral point ϕ is a \leq_{loc} -monotone semiring homomorphism from the semiring \mathcal{G} to the semiring $\mathbb{R}_{\geq 0}$. In particular, it satisfies the following conditions: For $G, H \in \{\text{Graphs}\}$:*

- (i) $\phi(\overline{K}_1) = 1$,
- (ii) $\phi(G \boxtimes H) = \phi(G)\phi(H)$,
- (iii) $\phi(G \sqcup H) = \phi(G) + \phi(H)$,
- (iv) $G \leq_{loc} H$ implies $\phi(G) \leq_{loc} \phi(H)$.

The asymptotic spectrum of graphs $\mathbf{X}(\mathcal{G}, \leq_{loc})$ is the set of all spectral points.

Remark 2.2. *A semiring homomorphism is also required to map K_0 to 0. This may not be well-defined for some explicit maps $\phi : \mathcal{G} \rightarrow \mathbb{R}_{\geq 0}$. By convention we set $\phi(K_0) = 0$ for all $\phi \in \mathbf{X}(\mathcal{G}, \leq_{loc})$.*

Known elements in $\mathbf{X}(\mathcal{G}, \leq_{loc})$ include the Lovász theta function ϑ [Lov79], the fractional Haemers bound $\mathcal{H}_f(G; \mathbb{F})$ over any field \mathbb{F} [Bla13, BC19], the projective rank $\overline{\xi}_f(G)$ [MR16] and the fractional clique cover number $\overline{\chi}_f(G)$ (cf. [Sch03, Eq. (67.112)]).

The following property of the theta function ϑ , introduced in [CMR⁺14], will be useful for us.

Proposition 2.3. *Given two graphs G and H , we write $G \leq_B H$ if there exist an integer $d \in \mathbb{N}$ and vectors $w \neq 0$, $w_{g,h} \in \mathbb{C}^d$ for every $g \in V(G)$ and $h \in V(H)$, satisfying the conditions:*

- (i) $\sum_{h \in V(H)} w_{g,h} = w$ for every $g \in V(G)$,
- (ii) $\langle w_{g,h}, w_{g',h'} \rangle = 0$ for all $\{g, g'\} \in E(\overline{G})$ and $(\{h, h'\} \in E(H) \text{ or } h = h')$,
- (iii) $\langle w_{g,h}, w_{g,h'} \rangle = 0$ for all $g \in V(G)$ and $h, h' \in V(H)$ with $h \neq h'$.

Then we have $\vartheta(G) \leq \vartheta(H) \iff G \leq_B H$.

The notion of graph homomorphism can be extended to the quantum setting via nonlocal games, as introduced in [MR16]. Let $M(d \times d', \mathbb{C})$ be the set of all $d \times d'$ matrices over \mathbb{C} . Let $M(d, \mathbb{C}) = M(d \times d, \mathbb{C})$ for short. A matrix $E \in M(d, \mathbb{C})$ is a projection if $E^2 = E = E^*$, where E^* denotes the conjugate transpose of E . Let I_d be the $d \times d$ identity matrix.

Definition 2.4 (Quantum homomorphism [MR16]). *We say there is a quantum homomorphism from a graph G to a graph H , denoted as $G \xrightarrow{q} H$, if, for some $d \in \mathbb{N}$, there exist projections $E_{g,h} \in M(d, \mathbb{C})$ for all $g \in V(G)$ and $h \in V(H)$ that satisfy*

- (i) $\sum_{h \in V(H)} E_{g,h} = I_d$ for any $g \in V(G)$,
- (ii) $E_{g,h}E_{g',h'} = 0$ if $\{g, g'\} \in E(G)$ and $(\{h, h'\} \in E(\overline{H}) \text{ or } h = h' \in V(H))$.

In addition we denote $G \leq_q H$ if $\overline{G} \xrightarrow{q} \overline{H}$.

Note that a graph homomorphism $f : V(G) \rightarrow V(H)$ defines a quantum homomorphism (with $d = 1$) by setting $E_{g,h} = \delta_{f(g),h}$. Hence, $\overline{G} \rightarrow \overline{H}$ implies $\overline{G} \xrightarrow{q} \overline{H}$ and thus $G \leq_{loc} H$ implies $G \leq_q H$. \leq_q is also a preorder on \mathcal{G} . We can analogously define

$$\mathbf{X}(\mathcal{G}, \leq_q) := \{\phi : \mathcal{G} \rightarrow \mathbb{R}_{\geq 0} : \phi \text{ is a } \leq_q \text{-monotone semiring homomorphism}\}$$

as the *quantum asymptotic spectrum of graphs*, this notion was first studied in [LZ21]. By the above observations it follows that $\mathbf{X}(\mathcal{G}, \leq_q) \subseteq \mathbf{X}(\mathcal{G}, \leq_{loc})$. Known elements in $\mathbf{X}(\mathcal{G}, \leq_q)$ include the Lovász theta function ϑ [Lov79], the fractional Haemers bound $\mathcal{H}_f(G; \mathbb{C})$ [Bla13, BC19]³ and the projective rank $\overline{\xi}_f(G)$ [MR16]. It remains unknown whether there are infinite many elements in $\mathbf{X}(\mathcal{G}, \leq_q)$.

³It has been shown in [LZ21] that $\mathcal{H}_f(G; \mathbb{C}) = \mathcal{H}_f(G; \mathbb{R})$ for any graph G .

2.2 A dual characterization for the (quantum) Shannon capacity of graphs

The main motivation to study the (quantum) asymptotic spectrum of graphs comes from its relevance to the study of (quantum versions of) independence numbers and the Shannon capacity of graphs. Recall that $\alpha(G)$ denotes the independence number of G , defined as the largest cardinality of an independent set of vertices of G , or, equivalently, $\alpha(G) = \max\{d : \overline{K}_d \leq_{loc} G\}$. Recall also that $\overline{\chi}(G)$ is the clique cover number of G (and equals the chromatic number $\chi(\overline{G})$ of \overline{G}), which can be equivalently defined as $\overline{\chi}(G) = \min\{d : G \leq_{loc} \overline{K}_d\}$. The inequalities $\alpha(G) \leq \vartheta(G) \leq \chi(\overline{G})$ are known as the sandwich theorem [Lov86, Knu94]. It follows directly from the definitions that for $\phi \in \mathbf{X}(\mathcal{G}, \leq_{loc})$ the sandwich theorem holds:

$$\alpha(G) \leq \phi(G) \leq \overline{\chi}(G) \quad \text{for any graph } G.$$

In addition, one can lift the sandwich theorem to asymptotic quantities. Let

$$\Theta(G) := \sup_k \sqrt[k]{\alpha(G^{\boxtimes k})} = \lim_{k \rightarrow \infty} \sqrt[k]{\alpha(G^{\boxtimes k})}$$

be the *Shannon capacity* of G [Sha56]. It is proved by Lovász [Lov75] (see also [Sch03, Theorem 67.17]) that

$$\overline{\chi}_f(G) = \inf_k \sqrt[k]{\overline{\chi}(G^{\boxtimes k})} = \lim_{k \rightarrow \infty} \sqrt[k]{\overline{\chi}(G^{\boxtimes k})}.$$

As an application of the definitions, for any $\phi \in \mathbf{X}(\mathcal{G}, \leq_{loc})$, we have

$$\Theta(G) \leq \phi(G) \leq \overline{\chi}_f(G).$$

Surprisingly, Zuiddam [Zui19] proved a stronger result, using Strassen's theory of asymptotic spectra (cf. [Str88, Zui18]):

Theorem 2.5 ([Zui19]). *For any graph G we have*

$$\Theta(G) = \min \{ \phi(G) : \phi \in \mathbf{X}(\mathcal{G}, \leq_{loc}) \}, \quad \overline{\chi}_f(G) = \max \{ \phi(G) : \phi \in \mathbf{X}(\mathcal{G}, \leq_{loc}) \}.$$

A follow-up work [LZ21] extends the above theorem to the quantum setting. The quantum independence number $\alpha_q(G)$ [MR16] and the quantum clique cover number $\overline{\chi}_q(G)$ [CMN+07] are quantum analogues of $\alpha(G)$ and $\overline{\chi}(G)$, defined as

$$\alpha_q(G) = \max\{d : \overline{K}_d \leq_q G\} \quad \text{and} \quad \overline{\chi}_q(G) = \min\{d : G \leq_q \overline{K}_d\}.$$

Then the *quantum Shannon capacity* is defined as

$$\Theta_q(G) := \sup_k \sqrt[k]{\alpha_q(G^{\boxtimes k})} = \lim_{k \rightarrow \infty} \sqrt[k]{\alpha_q(G^{\boxtimes k})}$$

and the asymptotic *quantum clique cover number* is defined as

$$\overline{\chi}_{q,\infty}(G) = \inf_k \sqrt[k]{\overline{\chi}_q(G^{\boxtimes k})} = \lim_{k \rightarrow \infty} \sqrt[k]{\overline{\chi}_q(G^{\boxtimes k})}.$$

The following quantum analogue of Theorem 2.5 holds.

Theorem 2.6 ([LZ21]). *For any graph G we have*

$$\Theta_q(G) = \min \{ \phi(G) : \phi \in \mathbf{X}(\mathcal{G}, \leq_q) \}, \quad \overline{\chi}_{q,\infty}(G) = \max \{ \phi(G) : \phi \in \mathbf{X}(\mathcal{G}, \leq_q) \}.$$

We point out that Theorems 2.5 and 2.6 follow from an abstract theorem for preordered semirings (cf. [Zui18, Theorem 2.13, Corollary 2.14-2.15]). For our purpose, it is sufficient to collect that \leq_{loc} and \leq_q satisfy the following properties:

Proposition 2.7. *Let $t \in \{loc, q\}$. The preorder \leq_t satisfies the following properties:*

- (i) $d \leq d'$ in $\mathbb{N} \iff \overline{K}_d \leq_t \overline{K}_{d'}$ in \mathcal{G} .
- (ii) If $G \leq_t H$ and $K \leq_t L$, then $G \sqcup K \leq_t H \sqcup L$ and $G \boxtimes K \leq_t H \boxtimes L$.
- (iii) For every $G, H \in \mathcal{G}$ with $H \neq K_0$, there exists a finite $d \in \mathbb{N}$, such that $G \leq_t \overline{K}_d \boxtimes H$.

Remark 2.8. These quantum analogues of $\alpha(G)$ and $\overline{\chi}(G)$ are defined by allowing (finite-dimensional) quantum strategies in corresponding nonlocal games. Another possible quantization is to consider entanglement-assisted zero-error information transmission, which is also studied in [LZ21]. A major open problem in zero-error quantum information theory is whether these two quantum notions are identical. In this paper we only focus on the nonlocal game setting.

2.3 Notation and preliminaries about operator algebras

We present necessary preliminary knowledge about operator algebra. See [Bla06, KR83] for a more detailed introduction. Let \mathcal{H} be a Hilbert space with inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$. We denote the norm on \mathcal{H} as $\|\cdot\|_{\mathcal{H}} := \sqrt{\langle \cdot, \cdot \rangle_{\mathcal{H}}}$ or simply $\|\cdot\|$ if there is no ambiguity. Let $B(\mathcal{H})$ be the space of bounded operators from \mathcal{H} to \mathcal{H} , equipped with the operator norm $\|X\|_{B(\mathcal{H})} = \sup\{\|X\eta\|_{\mathcal{H}} : \eta \in \mathcal{H}, \|\eta\| \leq 1\}$. We denote the identity operator in $B(\mathcal{H})$ by $I_{\mathcal{H}}$. We omit the subscript if it is clear in the context. For $X \in B(\mathcal{H})$, we write $\text{ran}(X) = \{X\eta : \eta \in \mathcal{H}\}$ (resp. $\text{cl}(\text{ran}(X))$) denote the range of X (resp. the closure of $\text{ran}(X)$) and $\text{ker}(X) = \{\eta \in \mathcal{H} : X\eta = 0\}$ denotes the kernel of X . The right support projection P_X of X is the projection onto $\text{ker}(X)^{\perp}$ and the left support projection Q_X of X is the projection onto $\text{cl}(\text{ran}(X))$. The right and left support projections satisfy the following relations:

$$\begin{aligned} XP_X &= X \text{ and } P_X Y = 0 \text{ whenever } XY = 0, \\ Q_X X &= X \text{ and } YQ_X = 0 \text{ whenever } YX = 0. \end{aligned} \tag{1}$$

A sequence of operators $\{T_n\}_{n \in \mathbb{N}}$ converges to T in the operator norm topology if $\|T_n - T\|_{B(\mathcal{H})} \rightarrow 0$. We denote this using $\lim_{n \rightarrow \infty} T_n = T$. A sequence of operators $\{T_n\}_{n \in \mathbb{N}}$ converges to T in strong operator topology (in short, SOT) if for any $\eta \in \mathcal{H}$, we have $\|T_n \eta - T \eta\|_{\mathcal{H}} \rightarrow 0$. We write $\text{sot-}\lim_{n \rightarrow \infty} T_n = T$. In general, the strong operator topology is weaker than the operator norm topology in $B(\mathcal{H})$, but they are the same if \mathcal{H} is finite dimensional.

A C^* -algebra \mathcal{A} is a norm-closed $*$ -subalgebra (closed under adjoint) of $B(\mathcal{H})$ for some Hilbert space \mathcal{H} . We denote the identity element of \mathcal{A} by $I_{\mathcal{A}}$ (or I if there is no confusion). A *state* τ on \mathcal{A} is a linear functional $\tau : \mathcal{A} \rightarrow \mathbb{C}$ that is *positive*, i.e., $\phi(X^*X) \geq 0$ for any $X \in \mathcal{A}$, and *unital*, i.e., $\phi(I_{\mathcal{A}}) = 1$. For example, a unit vector $\eta \in \mathcal{H}$ with $\|\eta\|_{\mathcal{H}} = 1$ gives a state $\tau_{\eta}(X) = \langle \eta, X\eta \rangle$. We call such states vector states. We say a state $\tau : \mathcal{A} \rightarrow \mathbb{C}$ is *tracial* if $\tau(XY) = \tau(YX)$ for all $X, Y \in \mathcal{A}$. The normalized trace $\text{tr} : M(d, \mathbb{C}) \rightarrow \mathbb{C}$ is a tracial state on the matrix algebra $M(d, \mathbb{C})$. We specify $\text{Tr}(X) = \sum_{i=1}^k X_{i,i}$ as the (unnormalized) trace of matrix X .

A von Neumann algebra \mathcal{M} is a SOT-closed $*$ -subalgebra of $B(\mathcal{H})$ for some \mathcal{H} . Since the strong operator topology is weaker than the norm topology in $B(\mathcal{H})$, every von Neumann algebra is a (unital) C^* -algebra. All C^* -algebras mentioned in this paper are unital (i.e., contain an identity element) and all mentioned von Neumann algebras are finite (i.e., admit a trace).

In contrast to C^* -algebras, a von Neumann algebra contains “enough” projections. That is, if $X \in \mathcal{M}$, then both its left support projection Q_X and right support projection P_X are also elements of \mathcal{M} . Moreover, a self-adjoint element $T = T^* \in \mathcal{M}$ admits a spectral decomposition in \mathcal{M} :

$$T = \int_{-\infty}^{\infty} \lambda dE_\lambda$$

where E_λ is the spectral projection of T onto the interval $(-\infty, \lambda]$. For a bounded Borel measurable function $f : \mathbb{R} \rightarrow \mathbb{R}$, the operator $f(T)$ by spectrum theorem is given by

$$f(T) = \int_{-\infty}^{\infty} f(\lambda) dE_\lambda .$$

If $T \in B(\mathcal{H})$ is normal (i.e., $TT^* = T^*T$), its left support projection Q_X and right support projection P_X coincide. If furthermore $T \geq 0$, then $T(T+\varepsilon I)^{-1} \rightarrow P_T$ as $\varepsilon \rightarrow 0$ in the strong operator topology.

Let $P, Q \in B(\mathcal{H})$ be two projections. We denote $P \vee Q$ as the (union) projection onto the closure of $\text{ran}(P) + \text{ran}(Q)$ and $P \wedge Q$ as the intersection projection onto $\text{ran}(P) \cap \text{ran}(Q)$. We have by functional calculus that (cf. [Bla06, I.5.2.1])

$$\text{ sot-} \lim_{n \rightarrow \infty} (P + Q)^{\frac{1}{n}} = P \vee Q , \quad \text{ sot-} \lim_{n \rightarrow \infty} (PQP)^n = P \wedge Q .$$

If $\text{ran}(P) \subseteq \text{ran}(Q)$, then $PQ = QP = P$ and $P \leq Q$ in the sense that $\langle \eta, P\eta \rangle \leq \langle \eta, Q\eta \rangle$ for every $\eta \in \mathcal{H}$. Moreover, by SOT-closedness, if P and Q belong to some von Neumann algebra \mathcal{M} , then $P \wedge Q$ and $P \vee Q$ also belong to \mathcal{M} .

A state $\tau : \mathcal{M} \rightarrow \mathbb{C}$ is normal if it is weak*-continuous, equivalently, for any monotone increasing net of positive operators $\{X_\alpha\} \subseteq \mathcal{M}$ with $X_\alpha \rightarrow X$, we have $\tau(X_\alpha) \rightarrow \tau(X)$. Vector states are always normal and normal states preserves the SOT-limit. As an example, for two projections $P, Q \in \mathcal{M}$ and a normal state $\tau : \mathcal{M} \rightarrow \mathbb{C}$ we have $\tau(P \wedge Q) = \lim_{n \rightarrow \infty} \tau((PQP)^n)$.

Let $\mathcal{A}_1 \subseteq B(\mathcal{H}_1)$ and $\mathcal{A}_2 \subseteq B(\mathcal{H}_2)$ be two C^* -algebras. Their direct sum $\mathcal{A}_1 \oplus \mathcal{A}_2$ is again a C^* -algebra and similarly for two von Neumann algebras $\mathcal{M}_1 \subseteq B(\mathcal{H}_1)$ and $\mathcal{M}_2 \subseteq B(\mathcal{H}_2)$. A *-homomorphism from \mathcal{A}_1 to \mathcal{A}_2 is a linear map $\pi : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ which preserves multiplication and the *-operation. We say \mathcal{A}_1 and \mathcal{A}_2 are *-isomorphic if there is a one-to-one and onto *-homomorphism from \mathcal{A}_1 to \mathcal{A}_2 . Let $\mathcal{A}_1 \odot \mathcal{A}_2$ denote the algebraic tensor product of \mathcal{A}_1 and \mathcal{A}_2 . The minimal C^* -algebraic tensor product $\mathcal{A}_1 \otimes_{\min} \mathcal{A}_2$ is the norm closure of $\mathcal{A}_1 \odot \mathcal{A}_2 \subseteq B(\mathcal{H}_1 \otimes \mathcal{H}_2)$. It does not depend on the choice of embeddings $\mathcal{A}_1 \subseteq B(\mathcal{H}_1)$ and $\mathcal{A}_2 \subseteq B(\mathcal{H}_2)$. Let ϕ_1 and ϕ_2 be states on \mathcal{A}_1 and \mathcal{A}_2 , respectively. The tensor product state $\phi_1 \otimes \phi_2 : \mathcal{A}_1 \odot \mathcal{A}_2 \rightarrow \mathbb{C}$ is defined by $(\phi_1 \otimes \phi_2)(X_1 \otimes X_2) = \phi_1(X_1)\phi_2(X_2)$ for $X_1 \in \mathcal{A}_1$, $X_2 \in \mathcal{A}_2$. It extends to a state on $\mathcal{A}_1 \otimes_{\min} \mathcal{A}_2$ (and also to the maximal tensor product $\mathcal{A}_1 \otimes_{\max} \mathcal{A}_2$, see [Bla06, II.9.3.5]). This state extension is the only property needed of the C^* -algebraic tensor product that we use in this paper and we therefore simply write $\mathcal{A}_1 \otimes \mathcal{A}_2$ for $\mathcal{A}_1 \otimes_{\min} \mathcal{A}_2$.

For two von Neumann algebras $\mathcal{M}_1 \subseteq B(\mathcal{H}_1)$ and $\mathcal{M}_2 \subseteq B(\mathcal{H}_2)$, the von Neumann algebra tensor product is $\mathcal{M}_1 \overline{\otimes} \mathcal{M}_2$ is the closure of $\mathcal{M}_1 \odot \mathcal{M}_2$ in the strong operator topology, i.e., $\mathcal{M}_1 \overline{\otimes} \mathcal{M}_2 = \overline{(\mathcal{M}_1 \odot \mathcal{M}_2)}^{\text{ sot }} \subseteq B(\mathcal{H}_1 \otimes \mathcal{H}_2)$. $\mathcal{M}_1 \overline{\otimes} \mathcal{M}_2$ is also independent of the choice of embeddings $\mathcal{M}_1 \subseteq B(\mathcal{H}_1)$ and $\mathcal{M}_2 \subseteq B(\mathcal{H}_2)$. Given two normal states ϕ_1 and ϕ_2 on \mathcal{M}_1 and \mathcal{M}_2 respectively, the tensor product state $\phi_1 \otimes \phi_2$ is also a normal state on $\mathcal{M}_1 \overline{\otimes} \mathcal{M}_2$. If there is no risk of confusion, we write $\mathcal{M}_1 \otimes \mathcal{M}_2$ for $\mathcal{M}_1 \overline{\otimes} \mathcal{M}_2$.

2.4 An infinite-dimensional rank-nullity theorem

Let \mathcal{H} be a Hilbert space and $S, T \subseteq \mathcal{H}$ be two subspaces of \mathcal{H} . When \mathcal{H} is finite dimensional, we have the following well-known identity

$$\dim(S) + \dim(T) = \dim(S + T) + \dim(S \cap T). \quad (2)$$

To see the above as a statement about ranks of operators, let P be the orthogonal projector on S , and Q the orthogonal projector on T . Then the above identity becomes

$$\text{rank}(P) + \text{rank}(Q) = \text{rank}(P \vee Q) + \text{rank}(P \wedge Q).$$

Moreover, if S and T are the row spaces of A and B in $B(\mathcal{H})$, respectively, we can compute the rank of AB^T by

$$\text{rank}(AB^T) = \dim(S) - \dim(S \cap T^\perp) = \text{rank}(P) - \text{rank}(P \wedge (I - Q)).$$

In the infinite-dimensional setting, the dimension of closed subspaces can be interpreted as the trace of their corresponding projections, a similar identity can be obtained as a consequence from Kaplansky's formula of projections (c.f. [KR86, Theorem 6.1.7]). This identity (and related results) will be a key tool to our proofs.

Lemma 2.9. *Let $\mathcal{M} \subseteq B(\mathcal{H})$ be a von Neumann algebra and let $P, Q \in \mathcal{M}$ be two projections. Suppose $\tau : \mathcal{M} \rightarrow \mathbb{C}$ is a normal tracial state. Then*

(i) $\tau(P) + \tau(Q) = \tau(P \vee Q) + \tau(P \wedge Q)$.

(ii) *Let $Y \in \mathcal{M}$ be the projection onto $\text{cl}(\text{ran}(QP))$. Then we have*

$$\tau(Q) \geq \tau(Y) = \tau(P) - \tau(P \wedge (I_{\mathcal{M}} - Q)).$$

(iii) *If additionally $P \wedge (I_{\mathcal{M}} - Q) = 0$, then $\tau(Y) = \tau(P)$. Moreover, for any $\varepsilon > 0$, there exists projection $P_\varepsilon \in \mathcal{M}$ such that $P_\varepsilon \leq P$, $\tau(P - P_\varepsilon) \leq \varepsilon$, and $\|P_\varepsilon(I_{\mathcal{M}} - Q)\| < 1$.*

Proof. Recall the two projections $P, Q \in \mathcal{M}$ are Murray-von Neumann equivalent, denoted by $P \sim Q$, if there exists a partial isometry $V \in \mathcal{M}$ such that $V^*V = P$ and $VV^* = Q$. It follows from the property of tracial states that $\tau(P) = \tau(Q)$ if $P \sim Q$. Then (i) follows from Kaplansky's formula of projections which states that for any two projections $P, Q \in \mathcal{M}$

$$P \vee Q - P \sim Q - P \wedge Q,$$

where the “ \sim ” operation is interpreted in the projection lattice of equivalence classes under \sim (see [KR83, Chapter 2]). For (ii), it is clear that $\text{ran}(Y) \subseteq \text{ran}(Q)$, hence $Y \leq Q$ and $\tau(Y) \leq \tau(Q)$. For the equality, let X denote the projector onto $\text{cl}(\text{ran}(PQ))$. It follows from [KR86, Proposition 6.1.6] that $X \sim Y$. Moreover, $\text{ran}(X)^\perp = \text{cl}(\text{ran}(PQ))^\perp = \ker(QP) = \ker P \oplus (\text{ran}(P) \cap \ker(Q))$. Hence

$$1 - X \sim (I_{\mathcal{M}} - P) + P \wedge (I_{\mathcal{M}} - Q), \quad Y \sim X \sim P - P \wedge (I - Q),$$

which proves (ii) by taking trace.

We give a detailed proof for (iii). Note that if $P \wedge (I_{\mathcal{M}} - Q) = 0$, then $\text{sot-}\lim_{n \rightarrow \infty} (P(I_{\mathcal{M}} - Q)P)^n \rightarrow 0$ (monotone decreasingly) and $\tau(Y) = \tau(P) - \tau(P \wedge (I_{\mathcal{M}} - Q)) = \tau(P)$. For any $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that $\tau((P(I_{\mathcal{M}} - Q)P)^{n_0}) \leq \varepsilon/2$. Choose $\delta > 0$ such that $(1 - \delta)^{n_0} \geq \frac{1}{2}$. Let

P_δ be the spectral projection of $P(I_{\mathcal{M}} - Q)P$ corresponding to the interval $(1 - \delta, 1]$. Thus $P_\delta \leq P$ and $(1 - \delta)P_\delta \leq P(I_{\mathcal{M}} - Q)P$. This implies

$$\frac{1}{2}\varepsilon \geq \tau((P(I_{\mathcal{M}} - Q)P)^{n_0}) \geq \tau((1 - \delta)^{n_0} P_\delta) = (1 - \delta)^{n_0} \tau(P_\delta).$$

Because $(1 - \delta)^{n_0} \geq 1/2$, we have

$$\tau(P_\delta) \leq \frac{\tau((P(I_{\mathcal{M}} - Q)P)^{n_0})}{(1 - \delta)^{n_0}} \leq \varepsilon.$$

Note that $\text{ran}(P_\delta) \subseteq \text{ran}(P(I_{\mathcal{M}} - Q)P) \subseteq \text{ran}(P)$. Then $P_\delta \leq P$ and we define the projection $P_\varepsilon = P - P_\delta$. We have $\tau(P - P_\varepsilon) = \tau(P_\delta) \leq \varepsilon$ and by the definition of P_δ ,

$$\|P_\varepsilon(I_{\mathcal{M}} - Q)\|^2 = \|P_\varepsilon(I_{\mathcal{M}} - Q)P_\varepsilon\| = \|(I_{\mathcal{M}} - P_\delta)P(I_{\mathcal{M}} - Q)P(I_{\mathcal{M}} - P_\delta)\| \leq 1 - \delta,$$

where the equality uses the fact that $P_\delta P = P_\delta$ and the inequality uses the fact that $I_{\mathcal{M}} - P_\delta$ is the spectral projection of $P(I_{\mathcal{M}} - Q)P$ corresponding to the interval $[0, 1 - \delta]$. \square

We remark that the above lemma can be alternatively proved via the representation theorem [RS89, Theorem 1.3] of the universal C^* -algebra generated by two projections. This representation theorem is also used in the study of the tracial rank [DP16].

3 C^* -algebraic and commuting quantum Shannon capacity and their asymptotic spectral characterizations

3.1 C^* -algebraic and commuting quantum homomorphisms

We first recall the definitions of graph homomorphism via C^* -algebras that were considered in [OP16]. They provide infinite-dimensional analogues of the previously mentioned notions of (quantum) graph homomorphism.

Definition 3.1. For graphs G and H , we say $G \xrightarrow{C^*} H$ if there exists a C^* -algebra $\mathcal{A}(G, H)$ containing projections $E_{g,h} \in \mathcal{A}(G, H)$ for every $g \in V(G)$ and $h \in V(H)$, satisfying the conditions:

- (i) $\sum_{h \in V(H)} E_{g,h} = I$ for any $g \in V(G)$,
- (ii) $E_{g,h} E_{g',h'} = 0$ if $\{g, g'\} \in E(G)$ and $(\{h, h'\} \in E(\overline{H})$ or $h = h' \in V(H))$.

Definition 3.2. For graphs G and H , we say $G \xrightarrow{q\mathcal{C}} H$ if there exists a C^* -algebra $\mathcal{A}(G, H)$ equipped with a tracial state τ and containing projections $E_{g,h} \in \mathcal{A}(G, H)$ for all $g \in V(G)$ and $h \in V(H)$, satisfying the conditions:

- (i) $\sum_{h \in V(H)} E_{g,h} = I$ for any $g \in V(G)$,
- (ii) $E_{g,h} E_{g',h'} = 0$ if $\{g, g'\} \in E(G)$ and $(\{h, h'\} \in E(\overline{H})$ or $h = h' \in V(H))$.

Note that Definition 3.1 reduces to Definition 3.2 if $\mathcal{A}(G, H)$ admits a tracial state. Definitions 3.1 and 3.2 reduce to the usual notion of graph homomorphism if $\mathcal{A}(G, H)$ has a 1-dimensional representation, and they reduce to the quantum homomorphism if $\mathcal{A}(G, H)$ has a finite-dimensional representation. Hence we have the following chain of implications:

$$G \rightarrow H \implies G \xrightarrow{q} H \implies G \xrightarrow{q\mathcal{C}} H \implies G \xrightarrow{C^*} H \implies G \xrightarrow{B} H \iff \vartheta(G) \leq \vartheta(H), \quad (3)$$

where the right most implication is shown in [OP16] and the last equivalence follows from Proposition 2.3.

These graph homomorphisms permit to define the corresponding C^* -algebraic and commuting quantum analogues of the independence number, clique cover number and asymptotic spectrum.

Definition 3.3. Let $t \in \{qc, C^*\}$ and G and H be graphs. Define

- $G \leq_t H$ if $\overline{G} \xrightarrow{t} \overline{H}$;
- $\alpha_t(G) := \max\{d : K_d \xrightarrow{t} \overline{G}\} = \max\{d : \overline{K_d} \leq_t G\}$, and $\Theta_t(G) := \lim_{k \rightarrow \infty} \sqrt[k]{\alpha_t(G^{\boxtimes k})}$;
- $\overline{\chi}_t(G) := \min\{d : \overline{G} \xrightarrow{t} K_d\} = \min\{d : G \leq_t \overline{K_d}\}$, and $\overline{\chi}_{t,\infty}(G) := \lim_{k \rightarrow \infty} \sqrt[k]{\overline{\chi}_t(G^{\boxtimes k})}$;
- $\mathbf{X}(\mathcal{G}, \leq_t) := \{\phi \in \text{Hom}(\mathcal{G}, \mathbb{R}_{\geq 0}) : \forall G, H \in \mathcal{G}, G \leq_t H \Rightarrow \phi(G) \leq \phi(H)\}$.

By Fekete's lemma [Fek23], both $\Theta_t(G)$ and $\overline{\chi}_{t,\infty}(G)$ are well-defined and equal $\sup_{k \in \mathbb{N}} \sqrt[k]{\alpha_t(G^{\boxtimes k})}$ and $\inf_{k \in \mathbb{N}} \sqrt[k]{\overline{\chi}_t(G^{\boxtimes k})}$, respectively. By Eq. (3), we obtain the chain of inclusions:

$$\mathbf{X}(\mathcal{G}, \leq_{C^*}) \subseteq \mathbf{X}(\mathcal{G}, \leq_{qc}) \subseteq \mathbf{X}(\mathcal{G}, \leq_q) \subseteq \mathbf{X}(\mathcal{G}, \leq_{loc}). \quad (4)$$

Our first result shows that \leq_{qc} and \leq_{C^*} define two new preorders on \mathcal{G} and satisfy similar properties as those in Proposition 2.7 for $t \in \{loc, q\}$.

Proposition 3.4. For graphs $G, H, K, L \in \mathcal{G}$ and $t \in \{qc, C^*\}$, we have

- (i) $G \leq_t G$,
- (ii) If $G \leq_t H$ and $H \leq_t K$, then $G \leq_t K$,
- (iii) $\overline{K_d} \leq_t \overline{K_{d'}}$ if and only if $d \leq d'$ in \mathbb{N} ,
- (iv) If $G \leq_t H$ and $K \leq_t L$, then $G \sqcup K \leq_t H \sqcup L$ and $G \boxtimes K \leq_t H \boxtimes L$,
- (v) If $H \neq K_0$, then there exists $d \in \mathbb{N}$ with $G \leq_t \overline{K_d} \boxtimes H$.

Proof. (i) is clear.

(ii) Let $\mathcal{A}(G, H)$ be a C^* -algebra containing a set of projections $\{E_{g,h} : g \in V(G), h \in V(H)\} \subseteq \mathcal{A}(G, H)$, and let $\mathcal{A}(H, K)$ be a C^* -algebra containing a set of projections $\{F_{h,k} : h \in V(H), k \in V(K)\} \subseteq \mathcal{A}(H, K)$, which are feasible solutions of $G \leq_{C^*} H$ and $H \leq_{C^*} K$, respectively. Define $A_{g,k} = \sum_{h \in V(H)} E_{g,h} \otimes F_{h,k}$ for $g \in V(G)$ and $k \in V(K)$ and let $\mathcal{A}(G, K) \subseteq \mathcal{A}(G, H) \otimes \mathcal{A}(H, K)$ be the C^* -algebra generated by $\{A_{g,k} : g \in V(G), k \in V(K)\}$. In [OP16, Prop. 4.6] it is shown that $\mathcal{A}(G, K)$ is a feasible solution of $G \leq_{C^*} K$. For $t = qc$, let in addition $\tau_{G,H}$ be a tracial state on $\mathcal{A}(G, H)$ and $\tau_{H,K}$ be a tracial state on $\mathcal{A}(H, K)$, respectively. Then $\tau_{G,K} = \tau_{G,H} \otimes \tau_{H,K}$ is a tracial state on $\mathcal{A}(G, K)$, since $\tau_{G,K}$ is a tracial state on $\mathcal{A}(G, H) \otimes \mathcal{A}(H, K)$ [Bla06, II.9.3.5].

(iii) By Eq. (3), $\overline{K_d} \leq_t \overline{K_{d'}}$ implies $\vartheta(\overline{K_d}) \leq \vartheta(\overline{K_{d'}})$ and thus $d \leq d'$. Conversely, $d \leq d'$ implies $\overline{K_d} \leq \overline{K_{d'}}$ by Proposition 2.7, which, using again Eq. (3), implies $\overline{K_d} \leq_t \overline{K_{d'}}$.

(iv) Let $\mathcal{A}(G, H)$ be a C^* -algebra containing projections $\{E_{g,h} : g \in V(G), h \in V(H)\}$ and $\mathcal{A}(K, L)$ be a C^* -algebra containing projections $\{F_{k,\ell} : k \in V(K), \ell \in V(L)\}$, which are feasible solutions of $G \leq_{C^*} H$ and $K \leq_{C^*} L$, respectively.

First we show $G \sqcup K \leq_{C^*} H \sqcup L$. For this, let $\mathcal{A}(G \sqcup K, H \sqcup L) \subseteq \mathcal{A}(G, H) \otimes \mathcal{A}(K, L)$ be the C^* -algebra generated by $\{A_{u,v} : u \in V(G \sqcup K), v \in V(H \sqcup L)\}$, where

$$A_{u,v} = \begin{cases} E_{u,v} \otimes I_{\mathcal{A}(K,L)} & \text{if } u \in V(G), v \in V(H), \\ I_{\mathcal{A}(G,H)} \otimes F_{u,v} & \text{if } u \in V(K), v \in V(L), \\ 0 & \text{otherwise.} \end{cases}$$

It is straightforward that $A_{u,v}$ is a projection for $u \in V(G \sqcup K)$ and $v \in V(H \sqcup L)$. For a fixed $u \in V(G \sqcup K)$, if $u \in V(G)$,

$$\sum_{v \in V(H \sqcup L)} A_{u,v} = \sum_{v \in V(H)} E_{u,v} \otimes I_{\mathcal{A}(K,L)} = I_{\mathcal{A}(G,H)} \otimes I_{\mathcal{A}(K,L)}.$$

Similarly, if $u \in V(K)$,

$$\sum_{v \in V(H \sqcup L)} A_{u,v} = \sum_{v \in V(L)} I_{\mathcal{A}(G,H)} \otimes F_{u,v} = I_{\mathcal{A}(G,H)} \otimes I_{\mathcal{A}(K,L)}.$$

Lastly, we show that if $\{u, u'\} \in E(\overline{G \sqcup K})$ and $(\{v, v'\} \in E(H \sqcup L)$ or $v = v')$, then $A_{u,v}A_{u',v'} = 0$ holds. Note that $\{v, v'\} \in E(H \sqcup L)$ or $v = v'$ implies that $v, v' \in V(H)$ or $v, v' \in V(L)$ (since there is no edge between $V(H)$ and $V(L)$). We distinguish the following cases:

- Either, $u, u' \in V(G)$ and $v, v' \in V(H)$, then $\{u, u'\} \in E(\overline{G})$ and thus $E_{u,v}E_{u',v'} = 0$, implying $A_{u,v}A_{u',v'} = E_{u,v}E_{u',v'} \otimes I_{\mathcal{A}(K,L)} = 0$.
- Or, $u, u' \in V(K)$ and $v, v' \in V(L)$, then $\{u, u'\} \in E(\overline{K})$ and thus $F_{u,v}F_{u',v'} = 0$, implying $A_{u,v}A_{u',v'} = I_{\mathcal{A}(G,H)} \otimes F_{u,v}F_{u',v'} = 0$.
- In all other cases, we have $A_{u,v} = 0$ or $A_{u',v'} = 0$, thus $A_{u,v}A_{u',v'} = 0$.

Thus we have shown $G \sqcup K \leq_{C^*} H \sqcup L$.

To show $G \sqcup K \leq_{qc} H \sqcup L$ one can use the above construction to build a C^* -algebra $\mathcal{A}(G \sqcup K, H \sqcup L)$ and the tracial state can be $\tau(G, H) \otimes \tau(K, L)$ as in (ii).

We now turn to show $G \boxtimes K \leq_{C^*} H \boxtimes L$. For this, let $\mathcal{A}(G \boxtimes K, H \boxtimes L) \subseteq \mathcal{A}(G, H) \otimes \mathcal{A}(K, L)$ be the C^* -algebra generated by $\{A_{(g,k),(h,l)} := E_{g,h} \otimes F_{k,l} : g \in V(G), h \in V(H), k \in V(K), l \in V(L)\}$. Then it is easy to verify that $\mathcal{A}(G \boxtimes K, H \boxtimes L)$ satisfies Definition 3.1. For $G \boxtimes K \leq_{qc} H \boxtimes L$, we use again the product tracial state $\tau(G, H) \otimes \tau(K, L)$ on $\mathcal{A}(G, H) \otimes \mathcal{A}(K, L)$.

(v) If $H \neq \overline{K_0}$, then there is a $d \in \mathbb{N}$ with $G \leq_{loc} \overline{K_d} \boxtimes H$ (see [Zui19]). Then the result follows using Eq. (3). \square

Using the above proposition, we can again invoke the abstract theorems for preordered semirings (cf. [Zui18, Theorem 2.13, Corollary 2.14-2.15]), that gave us Theorem 2.5 and Theorem 2.6. We obtain the following analogue for the cases $t \in \{qc, C^*\}$.

Theorem 3.5. *Let $t \in \{qc, C^*\}$ and $G \in \mathcal{G}$. We have*

$$\Theta_t(G) = \min\{\phi(G) : \phi \in \mathbf{X}(\mathcal{G}, \leq_t)\}, \quad \overline{\chi}_{t,\infty}(G) = \max\{\phi(G) : \phi \in \mathbf{X}(\mathcal{G}, \leq_t)\}. \quad (5)$$

3.2 Elements in the new quantum asymptotic spectra of graphs

It is proved in [OP16] that $G \leq_{C^*} H$ implies $G \leq_B H$, where the latter is equivalent to $\vartheta(G) \leq \vartheta(H)$. Thus, ϑ is a monotone for both preorders \leq_{C^*} and \leq_{qc} . Therefore,

$$\vartheta \in \mathbf{X}(\mathcal{G}, \leq_{C^*}) \subseteq \mathbf{X}(\mathcal{G}, \leq_{qc}).$$

We prove that the tracial rank introduced in [PSS+16] is an element of $\mathbf{X}(\mathcal{G}, \leq_{qc})$. In view of the inclusions in Eq. (4), we also obtain a new element in $\mathbf{X}(\mathcal{G}, \leq_q)$ and $\mathbf{X}(\mathcal{G}, \leq_{loc})$. In [PSS+16], the multiplicativity (w.r.t. strong product) and monotonicity (w.r.t. commuting quantum homomorphism) have been verified. We shall prove its additivity under disjoint union and show it is normalized. We first recall the definition of the tracial rank.

Definition 3.6. *We say that a graph G has a λ -tracial representation if there exists a C^* -algebra \mathcal{A} equipped with a tracial state τ and projections $E_g \in \mathcal{A}$ for $g \in V(G)$ which satisfy the conditions:*

- (i) $E_g E_{g'} = 0$ for all $\{g, g'\} \in E(\overline{G})$, and
- (ii) $\tau(E_g) = \frac{1}{\lambda}$ for all $g \in V(G)$.

The tracial rank of G is defined as

$$\overline{\xi}_{tr}(G) = \inf\{\lambda : G \text{ has a } \lambda\text{-tracial representation}\}. \quad (6)$$

Note that Definition 3.6 coincides with the one in [PSS+16] except we replace the graph by its complement. It is shown in [PSS+16, Theorem 6.11] that for every feasible solution $(\mathcal{A}, \{E_g\}, \tau)$, its GNS construction is also a feasible solution of $\overline{\xi}_{tr}$, and the second condition can be relaxed to

- (ii) $\tau(E_g) \geq \frac{1}{\lambda}$ for all $g \in V(G)$.

We use the above two formulations to show that the tracial rank is additive with respect to taking disjoint union.

Proposition 3.7. *For graphs $G, H \in \mathcal{G}$, we have $\overline{\xi}_{tr}(G \sqcup H) = \overline{\xi}_{tr}(G) + \overline{\xi}_{tr}(H)$.*

Proof. We first prove $\overline{\xi}_{tr}(G \sqcup H) \leq \overline{\xi}_{tr}(G) + \overline{\xi}_{tr}(H)$. Let $(\mathcal{A}_G, \{E_g\}_{g \in V(G)}, \tau_G)$ be a λ_G -tracial representation of $\overline{\xi}_{tr}(G)$ and let $(\mathcal{A}_H, \{F_h\}_{h \in V(H)}, \tau_H)$ be a λ_H -tracial representation of H , as given in Definition 3.6. Let $\mathcal{A} = \mathcal{A}_G \oplus \mathcal{A}_H$ be the direct sum of \mathcal{A}_G and \mathcal{A}_H , which is a C^* -algebra. \mathcal{A} contains projections $(E_g, 0)$ and $(0, F_h)$ for any $g \in V(G)$ and $h \in V(H)$. Define the linear map

$$\tau : \mathcal{A} \mapsto \mathbb{C}, \quad \tau((X, Y)) = \frac{\lambda_G}{\lambda_G + \lambda_H} \tau_G(X) + \frac{\lambda_H}{\lambda_G + \lambda_H} \tau_H(Y).$$

τ is clearly a tracial state on \mathcal{A} . Let $A_v \in \mathcal{A}$ for $v \in V(G) \cup V(H)$ by

$$A_v = \begin{cases} (E_v, 0) \in \mathcal{A}_G \oplus \mathcal{A}_H & \text{if } v \in V(G), \\ (0, F_v) \in \mathcal{A}_G \oplus \mathcal{A}_H & \text{if } v \in V(H). \end{cases}$$

Then each A_v is a projection and $A_v A_u = 0$ if $\{v, u\} \in E(\overline{G \sqcup H})$. Moreover, we have

$$\tau(A_v) = \begin{cases} \tau(E_v, 0) = \frac{\lambda_G}{\lambda_G + \lambda_H} \tau_G(E_v) = \frac{1}{\lambda_G + \lambda_H} & \text{if } v \in V(G), \\ \tau(0, F_v) = \frac{\lambda_H}{\lambda_G + \lambda_H} \tau_H(F_v) = \frac{1}{\lambda_G + \lambda_H} & \text{if } v \in V(H). \end{cases}$$

Hence $(\mathcal{A}', \{a_v\}_{v \in V(G) \cup V(H)}, \tau)$ is a $(\lambda_G + \lambda_H)$ -tracial representation of $G \sqcup H$, which implies $\bar{\xi}_{tr}(G \sqcup H) \leq \bar{\xi}_{tr}(G) + \bar{\xi}_{tr}(H)$.

We now prove $\bar{\xi}_{tr}(G \sqcup H) \geq \bar{\xi}_{tr}(G) + \bar{\xi}_{tr}(H)$. Let $(\mathcal{A}, \{A_v\}_{v \in V(G) \cup V(H)}, \tau)$ be a λ -tracial representation of $G \sqcup H$. Note that (i) implies that $A_g A_h = 0$ for every $g \in V(G)$ and $h \in V(H)$. Let \mathcal{M} be the von Neumann algebra generated by \mathcal{A} , τ extends to \mathcal{M} . Note that the projections $E = \bigvee_{v \in V(G)} A_v$ (the projection onto the closure of the range $\sum_{g \in V(G)} A_g$) and $F = \bigvee_{h \in V(H)} A_h$ (the projection onto the closure of the range $\sum_{u \in V(H)} A_u$) are elements of \mathcal{M} . We have $EF = 0$ and E and F commute with A_g for any $g \in V(G)$ and A_h for any $h \in V(H)$, respectively. Let \mathcal{A}_G be the C^* -algebra generated by $\{A_g : g \in V(G)\}$ with E being the identity element, and \mathcal{A}_H be the C^* -algebra generated by $\{A_h : h \in V(H)\}$ with F being the identity element. It is straightforward to see that $A_g A_{g'} = 0$ for all $\{g, g'\} \in E(\overline{G})$ and $A_h A_{h'} = 0$ for all $\{h, h'\} \in E(\overline{H})$.

We construct tracial states on \mathcal{A}_G and \mathcal{A}_H . Note that $\mathcal{A}_G, \mathcal{A}_H \subseteq \mathcal{M}$, thus the tracial state $\tau : \mathcal{M} \rightarrow \mathbb{C}$ is also a tracial linear map on \mathcal{A}_G and \mathcal{A}_H . Define $\tau_G : \mathcal{A}_G \rightarrow \mathbb{C}$ by $\tau_G(X) = \tau(X)/\tau(E)$ for any $X \in \mathcal{A}_G$ and $\tau_H : \mathcal{A}_H \rightarrow \mathbb{C}$ by $\tau_H(Y) = \tau(Y)/\tau(F)$ for any $Y \in \mathcal{A}_H$. Then τ_G and τ_H are tracial states on \mathcal{A}_G and \mathcal{A}_H , respectively. Note that

$$\tau_G(A_g) = \frac{\tau(A_g)}{\tau(E)} = \frac{1}{\lambda\tau(E)} \text{ for any } g \in V(G) \text{ and } \tau_H(A_h) = \frac{\tau(A_h)}{\tau(F)} = \frac{1}{\lambda\tau(F)} \text{ for any } h \in V(H).$$

This shows that $(\mathcal{A}_G, \{A_g\}_{g \in V(G)}, \tau_G)$ and $(\mathcal{A}_H, \{A_h\}_{h \in V(H)}, \tau_H)$ are $\lambda\tau(E)$ -tracial representation of G and $\lambda\tau(F)$ -tracial representation of H , respectively. Thus,

$$\bar{\xi}_{tr}(G) + \bar{\xi}_{tr}(H) \leq \lambda\tau(E) + \lambda\tau(F) \leq \lambda,$$

where the second inequality holds since $E + F \leq I_{\mathcal{A}}$. Taking the infimum over λ shows that $\bar{\xi}_{tr}(G) + \bar{\xi}_{tr}(H) \leq \bar{\xi}_{tr}(G \sqcup H)$. \square

Proposition 3.8. *The tracial rank is an element of $\mathbf{X}(\mathcal{G}, \leq_{qc})$.*

Proof. It is easy to verify that $\bar{\xi}_{tr}(K_1) = 1$. Then, by the additivity of $\bar{\xi}_{tr}$, we have $\bar{\xi}_{tr}(\overline{K}_d) = \bar{\xi}_{tr}(\bigsqcup_{i=1}^d K_1) = d$ for all $d \in \mathbb{N}$. Since tracial rank is also multiplicative and monotone, it is an element of $\mathbf{X}(\mathcal{G}, \leq_{qc})$. \square

4 Tracial Haemers bound

It is proved that the fractional Haemers bound and the projective rank are elements of the asymptotic spectrum $\mathbf{X}(\mathcal{G}, \leq_q)$ [LZ21], and we just proved that the tracial rank is an element of $\mathbf{X}(\mathcal{G}, \leq_{qc})$. Since the tracial rank can be viewed as an infinite-dimensional generalization of the projective rank [PSS⁺16], a natural question is whether there is an infinite-dimensional generalization of the fractional Haemers bound.

We define the tracial Haemers bound of G , denoted as $\mathcal{H}_{tr}(G)$. This new graph parameter is a von Neumann algebraic generalization of the fractional Haemers bound and we show that it is an element of $\mathbf{X}(\mathcal{G}, \leq_{qc})$. In other words, we show that it is normalized, \leq_{qc} -monotone, additive, and multiplicative. In particular, these also imply that $\mathcal{H}_{tr}(G)$ is an upper bound on $\alpha_{qc}(G)$ and $\Theta_{qc}(G)$.

4.1 Definition and examples

We start with recalling a coordinate-free definition of the fractional Haemers bound due to Lex Schrijver [BC19, Remark after Prop. 7]. The key part of this definition is the notion of a subspace representation of a graph.

Definition 4.1 (Subspace representation). *We say a graph G has a (d, r) -subspace representation (over \mathbb{C}) if there exist subspaces $S_g \subseteq \mathbb{C}^d$ for all $g \in V(G)$ satisfying*

- (i) $S_g \cap (\sum_{g' \in N_{\overline{G}}(g)} S_{g'}) = \{0\}$, where the summation is over all g' satisfying $\{g, g'\} \in E(\overline{G})$.
- (ii) $\dim(S_g) = r$ for all $g \in V(G)$;

The fractional Haemers bound (over \mathbb{C}) is then defined as

$$\mathcal{H}_f(G; \mathbb{C}) = \inf \left\{ \frac{d}{r} : G \text{ has a } (d, r)\text{-subspace representation over } \mathbb{C} \right\}. \quad (7)$$

In the infinite-dimensional setting, the trace of a projection is an analogue of the dimension of a subspace. This suggests to define the von Neumann-algebraic generalization of a subspace representation using projections. In order to do so, recall that the intersection of two subspaces and (the closure of) the sum of two subspaces corresponds to the wedge and the union of the projections, which can be expressed as strong-operator limits. Indeed, for two projections E, F in a von Neumann algebra $\mathcal{M} \subseteq B(\mathcal{H})$, the projection onto $\text{ran}(E) \cap \text{ran}(F)$ is $E \wedge F = \text{sot-}\lim_{n \rightarrow \infty} (EFE)^n$ and the projection onto $\text{cl}(\text{ran}(E) \cup \text{ran}(F))$ is $E \vee F = \text{sot-}\lim_{n \rightarrow \infty} (E + F)^{\frac{1}{n}}$. In light of Definition 4.1(i), it will be convenient to use the shorthand notation

$$E_{N_{\overline{G}}(g)} := \bigvee_{g' \in N_{\overline{G}}(g)} E_{g'} = \text{sot-}\lim_{n \rightarrow \infty} \left(\sum_{g' \in N_{\overline{G}}(g)} E_{g'} \right)^{1/n} \text{ for } g \in V(G). \quad (8)$$

We then have the following von Neumann-algebraic generalization of a subspace representation.

Definition 4.2 (Tracial subspace representation). *We say a graph G has a λ -tracial subspace representation if there exist a von Neumann algebra \mathcal{M} containing projections E_g for all $g \in V(G)$, and a normal tracial state $\tau : \mathcal{M} \rightarrow \mathbb{C}$, such that*

- (i) $E_g \wedge E_{N_{\overline{G}}(g)} = 0$.
- (ii) $\tau(E_g) = \frac{1}{\lambda}$ for all $g \in V(G)$.

We define the tracial Haemers bound of G as the infimum over all λ for which there exists a λ -tracial subspace representation of G .

Definition 4.3 (Tracial Haemers bound). *The tracial Haemers bound of a graph G is defined as*

$$\mathcal{H}_{tr}(G) := \inf \{ \lambda : G \text{ has a } \lambda\text{-tracial subspace representation} \}.$$

It is often convenient to work with a concrete representation $\mathcal{M} \subset B(\mathcal{H})$, and we will also refer to a tuple $(\mathcal{H}, \mathcal{M}, \{E_g\}, \tau)$ as a (concrete) tracial subspace representation of G . Note that for such a concrete von Neumann algebra, item (i) in Definition 4.2 can be equivalently formulated as

- (i) $S_g \cap \text{cl}(\sum_{g' \in N_{\overline{G}}(g)} S_{g'}) = \{0\}$, where $S_g = \text{ran}(E_g)$ for all $g \in V(G)$,⁴

which gives exactly correspondence to fractional Haemers bound. The next proposition shows a concrete tracial subspace representation is always achievable by GNS representation.

⁴The difference with item (i) of Definition 4.1 is the *closure* of $(\sum_{g' \in N_{\overline{G}}(g)} S_{g'})$. Note however that in finite-dimensional Hilbert spaces all linear subspaces are closed.

Proposition 4.4. *Every λ -tracial subspace representation $(\mathcal{M}, \{E_g\}, \tau)$ as in Definition 4.2 yields a λ -tracial subspace representation $(\mathcal{H}_\pi, \pi(\mathcal{M}), \pi(E_g), \phi_\tau)$ where $\pi : \mathcal{M} \rightarrow B(\mathcal{H}_\pi)$ is the GNS representation with respect to τ and ϕ_τ is the tracial vector state induced by τ .*

Proof. Note that the GNS representation π is normal because τ is a normal state. Thus, it preserves SOT limits, and hence $\pi(E_{N_{\overline{G}}(g)}) = \bigvee_{g' \in N_{\overline{G}}(g)} \pi(E_{g'})$. Then it follows from the condition (i) from Definition 4.2 that $\pi(E_g) \wedge \pi(E_{N_{\overline{G}}(g)}) = \pi(E_g \wedge E_{N_{\overline{G}}(g)}) = 0$ for every $g \in V(G)$. Moreover the GNS representation is such that $\phi_\tau(\pi(E_g)) = \tau(E_g)$ for all $g \in G$. \square

Remark 4.5.

a) In contrast to the definition of tracial rank [PSS⁺16] which uses C^* -algebras and tracial states, we use *von Neumann algebras* and *normal* tracial states in the definition. One reason is that the projection $E_{N_{\overline{G}}(g)}$ for each $g \in V(G)$ is an element in the von Neumann algebra generated by $\{E_g\}$, but not necessarily in the C^* -algebra generated by $\{E_g\}$. Moreover, using a normal tracial state τ ensures the following continuity: $\tau(P) = \lim_{k \rightarrow \infty} \tau(P_k)$ whenever $P = \text{sot-}\lim_{k \rightarrow \infty} P_k$. Note that the definition of the tracial rank uses only multiplication of operators and this is well-defined within in C^* -algebra (and therefore does not require normal states).

b) In the definition of a tracial subspace representation we may allow the equality in item (ii) to be an inequality: $\tau(E_g) \geq \frac{1}{\lambda}$ for all $g \in V(G)$. The proof is standard, we have included it in Appendix A for completeness.

We first show that the tracial Haemers bound is a proper infinite dimensional generalization of the fractional Haemers bound. It is clear that if a graph admits a (d, r) -subspace representation, then there is also a d/r -tracial subspace representation. The next proposition shows the converse also holds.

Proposition 4.6. *If a graph G has a λ -tracial subspace representation and the corresponding von Neumann algebra is finite-dimensional, then G has a (d, r) -subspace representation with $d/r \leq \lambda$.*

Proof. Let $(\mathcal{H}, \mathcal{M}, \{E_g\}, \tau)$ be a λ -tracial subspace representation of G , where \mathcal{M} is finite-dimensional. Thus \mathcal{M} is ($*$ -isomorphic to) a direct-sum of finite-dimensional matrix algebras: $\mathcal{A} = \bigoplus_{i=1}^{\ell} M(d_i, \mathbb{C})$. For each projection $E_g \in \mathcal{A}$, it can be uniquely represented as $E_g = \bigoplus_{i=1}^{\ell} E_g^i$, where $E_g^i \in M(d_i, \mathbb{C})$ is a projection for each $i \in [\ell]$. For the tracial state $\tau : \mathcal{A} \rightarrow \mathbb{C}$, there exists a probability distribution $p = (p_1, \dots, p_\ell) \in [0, 1]^\ell$ such that $\tau(\bigoplus_{i=1}^{\ell} X_i) = \sum_{i=1}^{\ell} \frac{p_i}{d_i} \text{Tr}(X_i)$. Note that since $\tau(E_g) = 1/\lambda$ for every $g \in V(G)$, we have that

$$\frac{1}{\lambda} = \sum_{i=1}^{\ell} \frac{p_i}{d_i} \text{Tr}(E_g^i) = \sum_{i=1}^{\ell} q_i \text{rank}(E_g^i) \text{ for all } g \in V(G),$$

where we set $q_i = p_i/d_i$ for each $i \in [\ell]$. Let

$$\begin{aligned} t^* = \max t \text{ s.t. } t \in [0, 1], \quad q \in \mathbb{R}_{\geq 0}^\ell, \quad \sum_{i=1}^{\ell} d_i q_i = 1, \\ \sum_{i=1}^{\ell} q_i \text{rank}(E_g^i) = t \quad \forall g \in V(G). \end{aligned} \tag{9}$$

It is clear that $t^* \geq 1/\lambda$. Since the coefficients of the constraints in the linear program described in Eq. (9) are all integers, the optimal value t^* is attained by a vector $(t^*, q_1, \dots, q_\ell)$ of rational

numbers. Write $q_i = m_i/d$ for some integers m_i ($i \in [\ell]$) and d . Let $E'_g = \bigoplus_{i=1}^{\ell} E_g^i \otimes I_{m_i}$ for each $g \in V(G)$ and observe that $E'_g \in M_d$ since $\sum_i d_i m_i = \sum_i d_i q_i d = d$. Note that

$$\text{rank}(E'_g) = \sum_{i=1}^{\ell} m_i \text{rank}(E_g^i) = dt^*.$$

Thus if we can show $\text{ran}(E'_g) \cap (\sum_{g' \in N_{\overline{G}}(g)} \text{ran}(E'_{g'})) = \{0\}$ for every $g \in V(G)$, then we have obtained a (d, dt^*) -subspace representation of G , where $d/(dt^*) = 1/t^* \leq \lambda$.

We are only left to prove that $\text{ran}(E'_g) \cap (\sum_{g' \in N_{\overline{G}}(g)} \text{ran}(E'_{g'})) = \{0\}$ (the closure can be omitted since these projections are finite dimensional). Note that for each $g \in V(G)$,

$$\text{ran}(E'_g) = \bigoplus_{i=1}^{\ell} \text{ran}(E_g^i) \otimes \mathbb{C}^{m_i} \subseteq (\bigoplus_{i=1}^{\ell} \text{ran}(E_g^i)) \otimes \mathbb{C}^m = \text{ran}(E_g) \otimes \mathbb{C}^m,$$

where $m \geq m_i$ for all $i \in [\ell]$. Since $\text{ran}(E_g) \cap (\sum_{g' \in N_{\overline{G}}(g)} \text{ran}(E_{g'})) = \{0\}$, we have

$$\text{ran}(E_g) \otimes \mathbb{C}^m \cap \left(\sum_{g' \in N_{\overline{G}}(g)} \text{ran}(E'_{g'}) \otimes \mathbb{C}^m \right) = \{0\}.$$

This concludes the proof. \square

As a corollary, we show that if we further restrict the von Neumann algebra in a tracial subspace representation to be commutative, we can get a fractional clique cover (or equivalently, a fractional coloring of the complement).

Corollary 4.7. *If a graph G has a λ -tracial subspace representation and the corresponding von Neumann algebra is commutative, then \overline{G} has a $d : r$ -coloring with $d/r \leq \lambda$.*

Proof. Assume that $(\mathcal{H}, \mathcal{M}, \{E_g\}, \tau)$ is a λ -tracial subspace representation and \mathcal{M} is commutative. Note that the $*$ -algebra \mathcal{M}_0 generated by the finite family of mutually commuting projection $\{E_g\}$ is always finite dimensional, since it is spanned by the monomials of the form $\prod_{g \in V(G)} E_g^{\varepsilon_g}$, where $\varepsilon_g \in \{0, 1\}$ for each $g \in V(G)$. Thus we may assume \mathcal{M} is finite-dimensional otherwise replace \mathcal{M} by the subalgebra \mathcal{M}_0 .

By the proof of Proposition 4.6, we can assume that $\{\text{ran}(E_g)\}$ forms a $d/r \leq \lambda$ subspace representation. For commuting operators $E, F \in \mathcal{M}$ we have $E \wedge F = EF$. Hence

$$E_g E_{N_{\overline{G}}(g)} = E_g \wedge E_{N_{\overline{G}}(g)} = 0.$$

Using that $E_{g'} \leq E_{N_{\overline{G}}(g)}$ for every $g' \in N_{\overline{G}}(g)$, this implies $E_g E_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$. This gives a commutative feasible solution of the projective rank, which can be transformed into a $d : r$ -coloring of \overline{G} using the proof of [PSS⁺16, Theorem 6.8]. \square

We show that the tracial Haemers bound is upper bounded by the tracial rank:

Proposition 4.8. *Let G be a graph. Then $\mathcal{H}_{tr}(G) \leq \overline{\xi}_{tr}(G)$.*

Proof. Let $(\mathcal{A}, \{E_g\}, \tau)$ be a feasible solution of $\overline{\xi}_{tr}(G)$ with value λ , as given in Definition 3.6, where τ is a faithful tracial vector state. Taking the GNS construction $\pi : \mathcal{A} \rightarrow B(\mathcal{H}_\pi)$ of τ , τ induces a tracial vector state ϕ_τ on the C^* -algebra $\pi(\mathcal{A})$, which naturally extends to a normal tracial state on the von Neumann algebra $\mathcal{M} = \overline{\pi(\mathcal{A})}^{\text{sof}}$. Moreover, since $E_g E_{g'} = 0$ for all $\{g, g'\} \in E(\overline{G})$, we have $\pi(E_g)\pi(E_{g'}) = 0$, which implies $\pi(E_g)\pi(E_{N_{\overline{G}}(g)}) = 0$ (and thus in particular $\pi(E_g) \wedge \pi(E_{N_{\overline{G}}(g)}) = 0$). This shows that $(\mathcal{H}_\pi, \mathcal{M}, \pi(E_g), \phi_\tau)$ is a tracial subspace representation of G with value λ . \square

As a first example, we compute the tracial Haemers bound of odd cycles. To do so, we follow the same strategy that has been used to show the analogous result for the fractional Haemers bound in [BC19].

Example 4.9. For all $d \in \mathbb{N}$, $\mathcal{H}_{tr}(C_{2d+1}) = d + \frac{1}{2}$.

Proof. Note that $\mathcal{H}_{tr}(C_{2d+1}) \leq \mathcal{H}_f(C_{2d+1}) = d + \frac{1}{2}$. Hence, we only need to show the lower bound. Let $(\mathcal{H}, \mathcal{M}, \{E_i\}_{i \in \mathbb{Z}_{2d+1}}, \tau)$ be a λ -tracial subspace representation of C_{2d+1} . First observe that we have $(E_0 \wedge E_1) \vee (E_0 \wedge E_{2d}) \leq E_0$ and using the graph structure of C_{2d+1} we have $E_1 \wedge E_{2d} = 0$ since $\{1, 2d\} \in E(C_{2d+1})$. Lemma 2.9 (i) then implies that

$$\frac{1}{\lambda} = \tau(E_0) \geq \tau(E_0 \wedge E_1) + \tau(E_0 \wedge E_{2d}). \quad (10)$$

We proceed to lower bound $\tau(E_0 \wedge E_1)$ and $\tau(E_0 \wedge E_{2d})$. Note that both are of the form $\tau(E_i \wedge E_{i+1 \bmod 2d+1})$. Using the symmetry of the odd cycle, it thus suffices to lower bound $\tau(E_0 \wedge E_1)$.

For any $k \in \mathbb{N}$ for which $2 \leq k \leq d-1$, let I_k be the independent set $I_k = \{3, 5, 7, \dots, 2k-1\}$. Let $E_{\{0,1\}} = E_0 \vee E_1$ be the projection onto $\text{cl}(\text{ran}(E_0) + \text{ran}(E_1))$ and $E_{I_k} = \bigvee_{i \in I_k} E_i$ be the projection onto $\text{cl}(\sum_{i \in I_k} \text{ran}(E_i))$. Since the edge $\{0, 1\}$ is not adjacent to any vertex in I_{d-1} , we have $\tau(E_{\{0,1\}} \vee E_{I_{d-1}}) \leq 1$. We can expand the left hand side using Lemma 2.9 (i) (iteratively) to show that

$$1 \geq \tau(E_{\{0,1\}} \vee E_{I_{d-1}}) = \tau(E_{\{0,1\}}) + \sum_{i \in I_{d-1}} \tau(E_i) = \tau(E_0) + \tau(E_1) - \tau(E_0 \wedge E_1) + \sum_{i \in I_{d-1}} \tau(E_i). \quad (11)$$

Here the first equality uses the fact that for every $2 \leq k \leq d-1$ we have $\tau(E_{\{0,1\}} \vee E_{I_k}) = \tau(E_{\{0,1\}} \vee E_{I_{k-1}}) + \tau(E_{2k-1})$, since vertex $2k-1$ is not adjacent with any vertices in $I_{k-1} \cup \{0, 1\}$. Combining Eq. (11) with the identity $\tau(E_i) = 1/\lambda$ for all $i \in \mathbb{Z}_{2d+1}$, shows

$$\tau(E_0 \wedge E_1) \geq \frac{d+1}{\lambda} - 1.$$

We can use this lower bound in Eq. (10) to obtain

$$\frac{1}{\lambda} = \tau(E_0) \geq \tau(E_0 \wedge E_1 + E_0 \wedge E_{2d}) = \tau(E_0 \wedge E_1) + \tau(E_0 \wedge E_{2d}) \geq \frac{2d+2}{\lambda} - 2.$$

Multiplying both sides by λ and rearranging gives $\lambda \geq d + \frac{1}{2}$. □

Corollary 4.10. The tracial Haemers bound and the Lovász theta function are incomparable.

Proof. The previous example shows that $\mathcal{H}_{tr}(C_5) = 2.5 > \sqrt{5} = \vartheta(C_5)$. On the other hand, let G be the complement of the Schläfli graph, $\mathcal{H}_{tr}(G) \leq \mathcal{H}_f(G; \mathbb{C}) \leq \mathcal{H}(G; \mathbb{C}) \leq 7 < 9 = \vartheta(G)$, where the second to last inequality was proved in [Hae79]. □

In the following subsections, we proceed to establish the monotonicity, additivity, normalization and multiplicativity of the tracial Haemers bound, along with some equivalent definitions. These enable us to conclude that $\mathcal{H}_{tr} \in \mathbf{X}(\mathcal{G}, \leq_{qc})$ and for any graph G ,

$$\alpha_{qc}(G) \leq \Theta_{qc}(G) \leq \mathcal{H}_{tr}(G).$$

4.2 Monotonicity of the tracial Haemers bound

We first introduce an equivalent formulation of a tracial subspace representation of a graph using two subspaces per vertex of G .

Proposition 4.11. *Let G be a graph. Then G has a λ -tracial subspace representation if and only if there exist a Hilbert space \mathcal{H} , a von Neumann algebra $\mathcal{M} \subseteq B(\mathcal{H})$ with a normal tracial state $\tau : \mathcal{M} \rightarrow \mathbb{C}$, containing projections E_g and F_g for all $g \in V(G)$, such that*

- (i) $\text{ran}(E_g) \cap \text{ran}(F_g)^\perp = \{0\}$ for any $g \in V(G)$,
- (ii) $F_g E_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$,
- (iii) $\tau(F_g) \geq \tau(E_g) \geq \frac{1}{\lambda}$ for all $g \in V(G)$.

Proof. Let $(\mathcal{H}, \mathcal{M}, \{E_g\}, \{F_g\}, \tau)$ be given as in Proposition 4.11. Let $S_g = \text{ran}(E_g)$ and $T_g = \text{ran}(F_g)$ for any $g \in V(G)$. We prove that $S_g \cap \text{cl}(\sum_{g' \in N_{\overline{G}}(g)} S_{g'}) = \{0\}$, which implies that $(\mathcal{H}, \hat{\mathcal{M}}, \{E_g\}, \tau)$ is a λ -tracial subspace representation, where $\hat{\mathcal{M}}$ is the von Neumann algebra generated by the set of projections $\{E_g\}$. Assume $x \in S_g \cap \text{cl}(\sum_{g' \in N_{\overline{G}}(g)} S_{g'})$, we show $x = 0$. To see this, note that $E_{g'} F_g = 0$ for any $\{g, g'\} \in E(\overline{G})$. Thus $T_g \perp S_{g'}$ for any $\{g, g'\} \in E(\overline{G})$ and therefore $T_g \perp \text{cl}(\sum_{g' \in N_{\overline{G}}(g)} S_{g'})$. This shows that $\text{cl}(\sum_{g' \in N_{\overline{G}}(g)} S_{g'}) \subseteq T_g^\perp$ and hence $x \in T_g^\perp$. Also, by assumption $x \in S_g$. However $S_g \cap T_g^\perp = \{0\}$, so $x = 0$.

Let $(\mathcal{H}, \mathcal{M}, \{E_g\}, \tau)$ be a λ -tracial subspace representation as given in Definition 4.2. For each $g \in V(G)$, let $F_g = I_{\mathcal{M}} - (\bigvee_{g' \in V(G)} E_{g'})$. Then F_g belongs to \mathcal{M} and we have $\text{ran}(E_g) \cap \text{ran}(F_g)^\perp = \{0\}$ by construction. Moreover we have $F_g E_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$ since the range of $E_{g'}$ is orthogonal to the range of F_g . It remains to show that $\tau(F_g) \geq \tau(E_g)$ for every $g \in V(G)$. Since $E_g \wedge (I - F_g) = 0$, Lemma 2.9 (ii) then shows that $\tau(F_g) \geq \tau(E_g)$ for any $g \in V(G)$. \square

We are ready to prove that \mathcal{H}_{tr} is \leq_{qc} -monotone.

Proposition 4.12. *For graphs $G, H \in \mathcal{G}$, we have that $G \leq_{qc} H$ implies $\mathcal{H}_{tr}(G) \leq \mathcal{H}_{tr}(H)$.*

Proof. Let $(\mathcal{H}_H, \mathcal{M}_H, \{E_h\}, \{F_h\}, \tau_H)$ be a λ -tracial subspace representation of H as given in Proposition 4.11. Let $\mathcal{H}_{G,H}$ be a Hilbert space, and let $\mathcal{A}(G, H) \subseteq B(\mathcal{H}_{G,H})$ be a C^* -algebra containing projections $\{P_{g,h} : g \in V(G), h \in V(H)\}$, with tracial state $\tau_{G,H} : \mathcal{A}(G, H) \rightarrow \mathbb{C}$, which is a feasible solution of $G \leq_{qc} H$. We first show that we can assume that $\mathcal{A}(G, H)$ is also a von Neumann algebra. Indeed, we can take the GNS representation π of $\mathcal{A}(G, H)$ with respect to $\tau_{G,H}$, and the von Neumann algebra $\mathcal{M}(G, H)$ generated by $\pi(\mathcal{A}(G, H))$, the image of the GNS construction. Then the tracial state τ' naturally extends to $\mathcal{M}(G, H)$. For ease of notation, we also use $\tau_{G,H}$ to denote the tracial state on $\mathcal{M}(G, H)$ and $\{P_{g,h} : g \in V(G), h \in V(H)\}$ to denote their images in $\mathcal{M}(G, H)$.

We construct a λ -tracial subspace representation of G . Set $A_g = \sum_{h \in V(H)} P_{g,h} \otimes E_h$ and $B_g = \sum_{h \in V(H)} P_{g,h} \otimes F_h$ for all $g \in V(G)$ and let \mathcal{M}_G be the von Neumann algebra generated by $\{A_g\}$ and $\{B_g\}$, which is a subalgebra of $\mathcal{M}(G, H) \overline{\otimes} \mathcal{M}_H$. Let $\tau_G = \tau_{G,H} \otimes \tau_H$ be the product tracial state on $\mathcal{M}(G, H) \overline{\otimes} \mathcal{M}_H$, thus also a tracial state on \mathcal{M}_G . We first verify that A_g and B_g are projections. Indeed,

$$A_g^2 = \sum_{h, h' \in V(H)} P_{g,h} P_{g,h'} \otimes E_h E_{h'} = \sum_{h \in V(H)} P_{g,h} \otimes E_h = A_g,$$

where the second equality uses the fact that $P_{g,h}P_{g,h'} = 0$ for all $h \neq h'$. The same argument works for B_g as well. We then lower bound the trace of A_g and B_g . Note that

$$\tau_G(A_g) = \sum_{h \in V(H)} \tau_{G,H}(P_{g,h}) \otimes \tau_H(E_h) \geq \frac{1}{\lambda} \sum_{h \in V(H)} \tau_{G,H}(P_{g,h}) = \frac{1}{\lambda},$$

where the last equality holds since $\sum_{h \in V(H)} P_{g,h} = I_{\mathcal{M}(G,H)}$. On the other hand,

$$\tau_G(B_g) = \sum_{h \in V(H)} \tau_{G,H}(P_{g,h}) \otimes \tau_H(F_h) \geq \sum_{h \in V(H)} \tau_{G,H}(P_{g,h}) \otimes \tau_H(E_h) = \tau_G(A_g).$$

We now verify that $B_g A_{g'} = 0$ if $\{g, g'\} \in E(\overline{G})$. Note that

$$B_g A_{g'} = \sum_{h, h' \in V(H)} P_{g,h} P_{g',h'} \otimes F_h E_{h'} = \sum_{\{h, h'\} \in E(\overline{H})} P_{g,h} P_{g',h'} \otimes F_h E_{h'} = 0,$$

where the second equality holds since $P_{g,h} P_{g',h'} = 0$ if $\{g, g'\} \in E(\overline{G})$ and $\{h, h'\} \in E(H)$ or $h = h'$ and the third equality holds since $F_h E_{h'} = 0$ if $\{h, h'\} \in E(\overline{H})$.

We are left to show that $\text{ran}(A_g) \cap \text{ran}(B_g)^\perp = \{0\}$ for every $g \in V(G)$. We first observe that

$$\text{ran}(A_g) = \oplus_{h \in V(H)} \text{ran}(P_{g,h} \otimes E_h) \text{ and } \text{ran}(B_g) = \oplus_{h \in V(H)} \text{ran}(P_{g,h} \otimes F_h),$$

since $(P_{g,h} \otimes E_h)(P_{g,h'} \otimes E_{h'}) = P_{g,h} P_{g,h'} \otimes E_h E_{h'} = 0$ and $(P_{g,h} \otimes F_h)(P_{g,h'} \otimes F_{h'}) = P_{g,h} P_{g,h'} \otimes F_h F_{h'} = 0$. Assume $x \in \text{ran}(A_g) \cap \text{ran}(B_g)^\perp$, that is, assume $A_g x = x$ and $B_g x = 0$. Note that

$$A_g B_g A_g = \sum_{h_1, h_2, h_3 \in V(H)} P_{g,h_1} P_{g,h_2} P_{g,h_3} \otimes E_{h_1} F_{h_2} E_{h_3} = \sum_{h \in V(H)} P_{g,h} \otimes E_h F_h E_h,$$

where the second equality use the fact that $P_{g,h}$ is a projection and $P_{g,h} P_{g,h'} = 0$ for any $h \neq h'$. Thus,

$$0 = \langle x, B_g x \rangle = \langle x, A_g B_g A_g x \rangle = \sum_{h \in V(H)} \langle x, (P_{g,h} \otimes E_h F_h E_h) x \rangle.$$

This implies that $\langle x, (P_{g,h} \otimes E_h F_h E_h) x \rangle = 0$ for any $h \in V(H)$ as $P_{g,h} \otimes E_h F_h E_h$ is a positive operator. Let $x_h = (P_{g,h} \otimes E_h) x \in \text{ran}(P_{g,h}) \otimes \text{ran}(E_h)$. We shall prove that $x_h = 0$ for all $h \in V(H)$, then $x = 0$ and we are done. For every fixed h , write the Schmidt decomposition of x_h as $\sum_i \mu_i y_i \otimes z_i$, where $\mu_i > 0$, $y_i \in \text{ran}(P_{g,h})$ and $z_i \in \text{ran}(E_h)$ for every i . Then the identity $\langle x, (P_{g,h} \otimes E_h F_h E_h) x \rangle = 0$ further implies

$$0 = \langle x_h, (P_{g,h} \otimes F_h) x_h \rangle = \sum_{i,j} \mu_i \langle y_i, P_{g,h} y_j \rangle \langle z_i, F_h z_j \rangle = \sum_i \mu_i \langle z_i, F_h z_i \rangle,$$

where the last equality holds since $P_{g,h} y_i = y_i$ for every i and $\langle y_i, y_j \rangle = 0$ for every $i \neq j$. Thus, $F_h z_i = 0$ for every i . Note that $\text{ran}(E_h) \cap \text{ran}(F_h)^\perp = \{0\}$, which implies $z_i = 0$ for all i . This concludes our proof. \square

4.3 Additivity of the tracial Haemers bound

In this subsection, we show additivity of $\mathcal{H}_{tr}(G)$. The subadditivity follows naturally from the original definition of tracial Haemers bound.

Proposition 4.13. *For graphs G and H we have $\mathcal{H}_{tr}(G \sqcup H) \leq \mathcal{H}_{tr}(G) + \mathcal{H}_{tr}(H)$.*

Proof. Let $(\mathcal{H}_G, \mathcal{M}_G, \{E_g\}, \{F_g\}, \tau_G)$ and $(\mathcal{H}_H, \mathcal{M}_H, \{\tilde{E}_h\}, \{\tilde{F}_h\}, \tau_H)$ be λ_G -tracial subspace representation of G and λ_H -tracial subspace representation of H , respectively, given in terms of Proposition 4.11. We construct a $(\lambda_G + \lambda_H)$ -tracial subspace representation of $G \sqcup H$. For $g \in V(G)$ define $A_g = (E_g, 0)$ and $B_g = (F_g, 0)$ and for $h \in V(H)$ let $A_h = (0, \tilde{E}_h)$ and $B_h = (0, \tilde{F}_h)$. Let $\mathcal{H} = \mathcal{H}_G \oplus \mathcal{H}_H$ and \mathcal{M} be the von Neumann algebra generated by $\{A_v : v \in V(G) \cup V(H)\}$ and $\{B_v : v \in V(G) \cup V(H)\}$. Note that $\mathcal{M} \subseteq \mathcal{M}_G \oplus \mathcal{M}_H$. Finally, for an element $(X, Y) \in \mathcal{M}$, define $\tau((X, Y)) = \frac{\lambda_G}{\lambda_G + \lambda_H} \tau_G(X) + \frac{\lambda_H}{\lambda_G + \lambda_H} \tau_H(Y)$. We verify that τ is a tracial state on \mathcal{M} . Note that $\tau((X^*X, Y^*Y)) = \frac{\lambda_G}{\lambda_G + \lambda_H} \tau_G(X^*X) + \frac{\lambda_H}{\lambda_G + \lambda_H} \tau_H(Y^*Y) \geq 0$, $\tau((I_{\mathcal{M}_G}, I_{\mathcal{M}_H})) = 1$ and

$$\begin{aligned} \tau(X_1 X_2, Y_1 Y_2) &= \frac{\lambda_G}{\lambda_G + \lambda_H} \tau_G(X_1 X_2) + \frac{\lambda_H}{\lambda_G + \lambda_H} \tau_H(Y_1 Y_2) \\ &= \frac{\lambda_G}{\lambda_G + \lambda_H} \tau_G(X_2 X_1) + \frac{\lambda_H}{\lambda_G + \lambda_H} \tau_H(Y_2 Y_1) = \tau(X_2 X_1, Y_2 Y_1). \end{aligned}$$

We show that $(\mathcal{H}, \mathcal{M}, \{A_v\}_{v \in V(G) \cup V(H)}, \{B_u\}_{u \in V(G) \cup V(H)}, \tau)$ is a $(\lambda_G + \lambda_H)$ -tracial subspace representation of $G \sqcup H$. First note that for $u \in V(G)$ we have $\tau(A_u) = \frac{\lambda_G}{\lambda_G + \lambda_H} \tau_G(E_u) + 0 \geq \frac{1}{\lambda_G + \lambda_H}$ and similarly for $v \in V(H)$, we have $\tau(A_v) \geq 0 + \frac{\lambda_H}{\lambda_G + \lambda_H} \tau_H(\tilde{E}_v) \geq \frac{1}{\lambda_G + \lambda_H}$. It is also straightforward that $\tau(B_v) \geq \tau(A_v)$ for every $v \in V(G) \cup V(H)$. We then verify that $B_v A_u = 0$ for $\{v, u\} \in E(\overline{G \sqcup H})$. If $v \in V(G)$ and $u \in V(H)$, $B_v A_u = (F_v, 0)(0, \tilde{E}_u) = 0$. A similar argument holds for $v \in V(H)$ and $u \in V(G)$. If $v, u \in V(G)$ and $\{v, u\} \in E(\overline{G})$, $B_v A_u = (F_v, 0)(E_u, 0) = (F_v E_u, 0) = 0$. A similar argument holds for $v, u \in V(H)$ and $\{v, u\} \in E(\overline{H})$. Finally we verify that $\text{ran}(A_u) \cap \text{ran}(B_u)^\perp = \{0\}$. We do so for $u \in V(G)$ and the argument for $u \in V(H)$ is similar. Note that $\text{ran}(A_u) = (\text{ran}(E_u), 0)$ and $\text{ran}(B_u) = (\text{ran}(F_u), 0)$. Then $\text{ran}(B_u)^\perp = (\text{ran}(F_u)^\perp, B(\mathcal{H}_H))$. Then $\text{ran}(A_u) \cap \text{ran}(B_u)^\perp = \{0\}$ and $\text{ran}(E_u) \cap \text{ran}(F_u)^\perp = \{0\}$. This concludes the proof. \square

To show superadditivity, we first strengthen the notion of a subspace representation to that of a projection representation by replacing the subspace intersection condition by a norm-inequality condition.

Definition 4.14 (Tracial projection representation). *We say a graph G has a λ -tracial projection representation if a von Neumann algebra \mathcal{M} containing projections E_g and F_g for all $g \in V(G)$, with a normal tracial state $\tau : \mathcal{A} \rightarrow \mathbb{C}$, such that*

- (i) $\|E_g(I_{\mathcal{A}} - F_g)\| < 1$ for any $g \in V(G)$,
- (ii) $F_g E_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$,
- (iii) $\tau(F_g) \geq \tau(E_g) \geq \frac{1}{\lambda}$ for all $g \in V(G)$.

As for Definition 4.2, the Hilbert \mathcal{H} in the above definition is unnecessary but we impose it for the convenience of discussion. Note that condition (i) above is a strengthening of the projection-intersection condition. Indeed, for two projections $P, Q \in B(\mathcal{H})$, if $\|PQ\| < 1$, then $P \wedge Q = 0$. If the underlying Hilbert space is finite dimensional, then the reverse direction also holds. In the infinite-dimensional setting, the reverse direction does not necessarily hold. Nevertheless, we shall prove that, for any graph G , the infimum over tracial subspace representations and the infimum over tracial projection representation are the same.

Proposition 4.15. *Let G be a graph. Then*

$$\mathcal{H}_{tr}(G) = \inf\{\lambda : G \text{ has a } \lambda\text{-tracial projection representation}\}.$$

Proof. We show equality between $\mathcal{H}_{tr}(G)$ and the parameter defined in this proposition. Note that the condition $\|E_g(I_{\mathcal{M}} - F_g)\| < 1$ implies $E_g \wedge (I_{\mathcal{M}} - F_g) = 0$. This shows that $\mathcal{H}_{tr}(G)$ is at most the parameter defined in this proposition. To prove the converse inequality, consider a tuple $(\mathcal{H}, \mathcal{M}, \{E_g\}, \{F_g\}, \tau)$ of G satisfying Proposition 4.11. Let $0 < \varepsilon \leq 1$. Lemma 2.9 (iii) implies that for every $g \in V(G)$, there is a projection $E_g^\varepsilon \in \mathcal{M}$ such that $E_g^\varepsilon \leq E_g$, $\|E_g^\varepsilon(I_{\mathcal{M}} - F_g)\| < 1$ and $\tau(E_g^\varepsilon) \geq \tau(E_g) - \varepsilon \geq \frac{1}{\lambda} - \varepsilon$. Let $\hat{\mathcal{M}}$ be the von Neumann algebra generated by $\{E_g^\varepsilon\}$ and $\{F_g\}$. It follows that $(\mathcal{H}, \hat{\mathcal{M}}, \{E_g^\varepsilon\}, \{F_g\}, \tau)$ is a $\frac{\lambda}{1-\varepsilon\lambda}$ -tracial projection representation of G . Letting ε tend to 0 and taking the infimum over λ concludes the proof. \square

With Proposition 4.15, we can further restrict the Hilbert space \mathcal{H} in Proposition 4.11, which will play a crucial role in the proof of superadditivity (and the proof of supermultiplicativity).

Proposition 4.16. *Let G be a graph. If there exists a λ -tracial projection representation of G , then there exists a λ -tracial subspace representation $(\mathcal{H}, \mathcal{M}, \{E_g\}, \{F_g\}, \tau)$ in the sense of Proposition 4.11, where*

$$\mathcal{H} = \text{ran}\left(\bigvee_{g \in V(G)} F_g\right).$$

Proof. Let $(\mathcal{H}, \mathcal{M}, \{E_g\}, \{F_g\}, \tau)$ be a λ -tracial projection representation of G and let F be the projection onto $\text{cl}(\sum_{g \in V(G)} \text{ran}(F_g))$. Let E'_g be the left support projection of FE_g (i.e. the projection onto the closure of $\text{ran}(FE_g)$) and set $F'_g = F_g$. Using the inclusions $\text{ran}(E'_g), \text{ran}(F'_g) \subseteq \text{ran}(F)$, we observe that F commutes with both F'_g and E'_g and that it acts as the identity on the algebra generated by the E'_g 's and F'_g 's. Let $\mathcal{H}' = \text{ran}(F)$ and let $\hat{\mathcal{M}}$ be the von Neumann algebra generated by $\{E'_g\}$ and $\{F'_g\}$, where F is the identity element. Define $\tau' : \hat{\mathcal{M}} \rightarrow \mathbb{C}$ as $\tau'(X) = \tau(X)/\tau(F)$ for $X \in \hat{\mathcal{M}}$. This is a tracial state on $\hat{\mathcal{M}}$. We show that $(\mathcal{H}', \hat{\mathcal{M}}, \{E'_g\}, \{F'_g\}, \tau')$ satisfies Proposition 4.11 with the same value.

We first verify that $F'_g E'_{g'} = 0$ for every $\{g, g'\} \in E(\overline{G})$. We have

$$F_g F E_{g'} = F_g E_{g'} = 0 \quad \text{for any } \{g, g'\} \in E(\overline{G}).$$

Using that E'_g is the left support projection on FE_g (in particular the property in Eq. (1)), this indeed implies that $F'_g E'_{g'} = F_g E_{g'} = 0$. Next we show $\text{ran}(E'_g) \cap \text{ran}(F_g)^\perp = \{0\}$. Let $v \in \text{ran}(E'_g) \cap \text{ran}(F_g)^\perp$, we show $v = 0$. By the definition of E'_g , there exists a sequence $\{u_i \in \mathcal{H}\}_{i \in \mathbb{N}}$ such that $FE_g u_i \rightarrow v$ as $i \rightarrow \infty$. On the other hand, $v \in \text{ran}(F_g)^\perp$ implies that $F_g v = 0$. Thus,

$$0 = F_g v = F_g \lim_{i \rightarrow \infty} FE_g u_i = \lim_{i \rightarrow \infty} F_g FE_g u_i = \lim_{i \rightarrow \infty} F_g E_g u_i.$$

In other words, $\|F_g E_g u_i\| \rightarrow 0$ as $i \rightarrow \infty$. We show this implies $\|E_g u_i\| \rightarrow 0$. Let $\|E_g(I_{\mathcal{A}} - F_g)\| = \delta_g < 1$ for any $g \in V(G)$. Then by the triangle inequality, for any $i \in \mathbb{N}$,

$$\|F_g E_g u_i\| \geq \|E_g u_i\| - \|(I_{\mathcal{A}} - F_g)E_g u_i\| \geq \|E_g u_i\| - \|E_g(I_{\mathcal{A}} - F_g)\| \|E_g u_i\| = (1 - \delta_g) \|E_g u_i\|.$$

Thus,

$$\lim_{i \rightarrow \infty} \|E_g u_i\| \leq (1 - \delta_g)^{-1} \lim_{i \rightarrow \infty} \|F_g E_g u_i\| = 0,$$

which implies $v = \lim_{i \rightarrow \infty} FE_g u_i = 0$. (The second inequality uses the submultiplicativity of the operator norm.)

Finally we show $\tau'(E'_g) = \tau(E'_g)/\tau(F) \geq \frac{1}{\lambda}$. Since $\tau(F) \leq 1$, it suffices to show $\tau(E'_g) = \tau(E_g)$. Since $\text{ran}(F)^\perp \subseteq \text{ran}(F_g)^\perp$, we have $\text{ran}(E_g) \cap \text{ran}(F)^\perp \subseteq \text{ran}(E_g) \cap \text{ran}(F_g)^\perp = \{0\}$. Applying Lemma 2.9 (ii) with $Q = F$ and $P = E_g$ therefore shows that for any tracial state $\tau : \mathcal{M} \rightarrow \mathbb{C}$ we have $\tau(E'_g) = \tau(E_g)$. \square

We are now ready to prove superadditivity of tracial Haemers bound.

Proposition 4.17. *For graphs G and H we have $\mathcal{H}_{tr}(G \sqcup H) \geq \mathcal{H}_{tr}(G) + \mathcal{H}_{tr}(H)$.*

Proof. Let $(\hat{\mathcal{H}}, \hat{\mathcal{M}}, \{\hat{E}_v\}, \{\hat{F}_v\}, \hat{\tau})$ be a λ -tracial projection representation of $G \sqcup H$ as formulated in Definition 4.14. Proposition 4.16 shows that we can obtain a λ -tracial subspace representation $(\mathcal{H}, \mathcal{M}, \{E_v\}, \{F_v\}, \tau)$ with $\mathcal{H} = \text{ran}(F)$, where F is the projection onto the closure of $\sum_{v \in V(G) \cup V(H)} \text{ran}(F_v)$. Using the graph structure of $G \sqcup H$ we see that the projections $\{E_g\}_{g \in V(G)}$ and $\{F_g\}_{g \in V(G)}$ satisfy

- (1) $\text{ran}(E_g) \cap \text{ran}(F_g)^\perp = \{0\}$ for any $g \in V(G)$,
- (2) $F_g E_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$.

Similar conditions hold for $\{E_h\}_{h \in V(H)}$ and $\{F_h\}_{h \in V(H)}$. Let E_G be the projection onto the closure of $\sum_{g \in V(G)} \text{ran}(E_g)$ and similarly let E_H be the projection onto the closure of $\sum_{h \in V(H)} \text{ran}(E_h)$. Let $\mathcal{H}_G = \text{ran}(E_G)$ and $\mathcal{H}_H = \text{ran}(E_H)$. Let \mathcal{M}_G (resp. \mathcal{M}_H) be the von Neumann algebra generated by the projections $\{E_g\}_{g \in V(G)}$ with identity element E_G (resp. $\{E_h\}_{h \in V(G)}$ with identity E_H). Thus $\mathcal{M}_G \subseteq B(\mathcal{H}_G)$ and $\mathcal{M}_H \subseteq B(\mathcal{H}_H)$. Let $\tau_G : \mathcal{M}_G \rightarrow \mathbb{C}$ be defined as $\tau_G(X) = \tau(X)/\tau(E_G)$ for $X \in \mathcal{M}_G$ and $\tau_H : \mathcal{M}_H \rightarrow \mathbb{C}$ be defined as $\tau_H(Y) = \tau(Y)/\tau(E_H)$ for $Y \in \mathcal{M}_H$. Since \mathcal{M}_G and \mathcal{M}_H are subsets of \mathcal{M} , τ_G and τ_H are tracial states on \mathcal{M}_G and \mathcal{M}_H respectively. Analogously to the first part of the proof of Proposition 4.11 one can show that $(\mathcal{H}_G, \mathcal{M}_G, \{E_g\}_{g \in V(G)}, \tau_G)$ and $(\mathcal{H}_H, \mathcal{M}_H, \{E_h\}_{h \in V(H)}, \tau_H)$ are $(\tau(E_G)\lambda)$ - and $(\tau(E_H)\lambda)$ -tracial subspace representations of G and H respectively. It follows that $\mathcal{H}_{tr}(G) + \mathcal{H}_{tr}(H) \leq \lambda(\tau(E_G) + \tau(E_H))$. If we can show that $\tau(E_G) + \tau(E_H) \leq 1$, then we indeed obtain $\mathcal{H}_{tr}(G \sqcup H) \geq \mathcal{H}_{tr}(G) + \mathcal{H}_{tr}(H)$.

To show that $\tau(E_G) + \tau(E_H) \leq 1$ we will apply Lemma 2.9 to E_G and E_H . To do so we are left to prove that $\text{ran}(E_G) \cap \text{ran}(E_H) = \{0\}$. Assume there is a nonzero vector $0 \neq u \in \text{ran}(E_G) \cap \text{ran}(E_H)$. Note that for any $g \in V(G)$, $h \in V(H)$, we have $F_g E_h = 0$ and $F_h E_g = 0$, since there are no edges between vertices in G and vertices in H . This shows that $F_h E_G = 0$ for any $h \in V(H)$ and $F_g E_H = 0$ for any $g \in V(G)$. Since $E_G u = u$ and $E_H u = u$, we have $F_v u = F_v E_H u = 0$ for every $v \in V(G)$ and $F_v u = F_v E_G u = 0$ for every $v \in V(H)$. Thus,

$$u \in \bigcap_{g \in V(G)} \text{ran}(F_g)^\perp \cap \bigcap_{h \in V(H)} \text{ran}(F_h)^\perp = \bigcap_{v \in V(G) \cup V(H)} \text{ran}(F_v)^\perp = \text{ran}(F)^\perp = \{0\},$$

where the last equality holds due to the assumption that $\mathcal{H} = \text{ran}(F)$. \square

By the additivity, we can easily show that the tracial Haemers bound is normalized:

Example 4.18. *Let \overline{K}_d be the empty graph with d vertices. Then $\mathcal{H}_{tr}(\overline{K}_d) = d$.*

Proof. It is easy to see that $\mathcal{H}_{tr}(\overline{K}_1) = 1$. Then $\mathcal{H}_{tr}(\overline{K}_d) = \mathcal{H}_{tr}(\sqcup_{i=1}^d \overline{K}_1) = \sum_{i=1}^d \mathcal{H}_{tr}(\overline{K}_1) = d$. \square

The normalization of the tracial Haemers bound, together with the monotonicity with respect to the commuting quantum homomorphism, implies that the tracial Haemers bound is an upper bound on the commuting quantum independence number.

Corollary 4.19. *For every graph G , we have $\alpha_{qc}(G) \leq \mathcal{H}_{tr}(G)$.*

Proof. By definition, if $\alpha_{qc}(G) = d$, then $\overline{K}_d \leq_{qc} G$. The monotonicity and normalization of \mathcal{H}_{tr} together show that $d = \mathcal{H}_{tr}(\overline{K}_d) \leq \mathcal{H}_{tr}(G)$. \square

4.4 Multiplicativity of the tracial Haemers bound

We now discuss the multiplicativity of $\mathcal{H}_{tr}(G)$. We first use Proposition 4.11 to prove that the tracial Haemers bound is submultiplicative.

Proposition 4.20. *For any graphs G and H , $\mathcal{H}_{tr}(G \boxtimes H) \leq \mathcal{H}_{tr}(G)\mathcal{H}_{tr}(H)$.*

Proof. Let $(\mathcal{H}_G, \mathcal{M}_G, \{E_g\}, \{F_g\}, \tau_G)$ and $(\mathcal{H}_H, \mathcal{M}_H, \{E'_h\}, \{F'_h\}, \tau_H)$ be tracial subspace representations of G and H , respectively, in the formulation of Proposition 4.11. Set $P_{g,h} = E_g \otimes E'_h$ and $Q_{g,h} = F_g \otimes F'_h$ for any $g \in V(G)$ and $h \in V(H)$. It is clear that $P_{g,h}$ and $Q_{g,h}$ are projections. Let $\mathcal{M}_{G,H}$ be the von Neumann algebra generated by the projections $\{P_{g,h}, Q_{g,h} : g \in V(G), h \in V(H)\}$. It follows that $\mathcal{M}_{G,H}$ is a subalgebra of $\mathcal{M}_G \otimes \mathcal{M}_H$, the latter of which is equipped with the product tracial state $\tau_G \otimes \tau_H$. Thus $\tau = \tau_G \otimes \tau_H$ is a tracial state on $\mathcal{M}_{G,H}$, and it is easy to see that $\tau(Q_{g,h}) \geq \tau(P_{g,h}) \geq \frac{1}{\lambda\mu}$ for any $g \in V(G)$ and $h \in V(H)$.

For $\{(g, h), (g', h')\} \in E(\overline{G \boxtimes H})$, $P_{g,h}Q_{g',h'} = E_g F_{g'} \otimes E'_h F'_{h'} = 0$ because $\{g, g'\} \in E(\overline{G})$ or $\{h, h'\} \in E(\overline{H})$. This verifies the orthogonality conditions between $P_{g,h}$ and $Q_{g',h'}$.

We now show that $\text{ran}(P_{g,h}) \cap \text{ran}(Q_{g,h})^\perp = \{0\}$ for every $g \in V(G)$ and $h \in V(H)$. Note that $\text{ran}(P_{g,h}) = \text{ran}(E_g) \otimes \text{ran}(E'_h)$ and

$$\text{ran}(Q_{g,h})^\perp = (\text{ran}(F_g) \otimes \text{ran}(F'_h))^\perp = \left(\mathcal{H}_G \otimes \text{ran}(F'_h)^\perp \right) \oplus \left(\text{ran}(F_g)^\perp \otimes \text{ran}(F'_h) \right).$$

Thus for $x \in \text{ran}(P_{g,h}) \cap \text{ran}(Q_{g,h})^\perp$, we have $x = y + z$, where $x \in \text{ran}(E_g) \otimes \text{ran}(E'_h)$, $y \in \mathcal{H}_G \otimes \text{ran}(F'_h)^\perp$ and $z \in \text{ran}(F_g)^\perp \otimes \text{ran}(F'_h)$. For any linear functional $\mathcal{L} : \text{ran}(F'_h) \rightarrow \mathbb{C}$, we have $(I_G \otimes \mathcal{L})(x) \in \text{ran}(E_g)$, $(I_G \otimes \mathcal{L})(y) = 0$ and $(I_G \otimes \mathcal{L})(z) \in \text{ran}(F_g)^\perp$. Thus we have $(I_G \otimes \mathcal{L})(x) = (I_G \otimes \mathcal{L})(z) \in \text{ran}(E_g) \cap \text{ran}(F_g)^\perp = \{0\}$. This implies $z = 0$ because \mathcal{L} is arbitrary. Similarly, we have $y = 0$ and hence $x = 0$. \square

To prove the supermultiplicativity of $\mathcal{H}_{tr}(G)$, we need to relax the conditions in the λ -tracial subspace (resp. projection) representation by modifying (i) and (iii) in Proposition 4.11 (resp. Proposition 4.15):

Proposition 4.21. *Let G be a graph. Then G has a λ -tracial subspace representation if and only if there exist a Hilbert space \mathcal{H} , a von Neumann algebra $\mathcal{M} \subseteq B(\mathcal{H})$ containing projections E_g and F_g for all $g \in V(G)$, with a normal tracial state $\tau : \mathcal{M} \rightarrow \mathbb{C}$, such that*

- (i) $F_g E_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$,
- (ii) $\tau(E_g) - \tau(X_g) \geq \frac{1}{\lambda}$ for all $g \in V(G)$, where X_g is the projection onto the subspace $\text{ran}(E_g) \cap \text{ran}(F_g)^\perp$.

Proof. We first prove that a tuple $(\mathcal{H}, \mathcal{M}, \{E_g\}, \{F_g\}, \tau)$ which satisfies Proposition 4.21 can be used to construct a λ -tracial subspace representation of G in terms of Proposition 4.11. Let $\tilde{E}_g = E_g - X_g$, for every $g \in V(G)$, where X_g is the projection onto the subspace $\text{ran}(E_g) \cap \text{ran}(F_g)^\perp$, and let $\tilde{F}_g = F_g$. Note that \tilde{E}_g is a projection since $\text{ran}(X_g) \subseteq \text{ran}(E_g)$. Let $\tilde{\mathcal{M}}$ be the von Neumann algebra generated by the projections \tilde{E}_g and \tilde{F}_g for all $g \in V(G)$. Then $\tilde{\tau} = \tau$ is a tracial state on $\tilde{\mathcal{M}} \subseteq \mathcal{M}$ and we observe that $\tilde{\tau}(\tilde{E}_g) = \tau(E_g) - \tau(X_g) \geq \frac{1}{\lambda}$.

It remains to show that (i) $\text{ran}(\tilde{E}_g) \cap \text{ran}(\tilde{F}_g)^\perp = \{0\}$ and (ii) $\tilde{F}_g \tilde{E}_{g'} = 0$ if $\{g, g'\} \in E(\overline{G})$. For (i), since $\text{ran}(\tilde{E}_g)$ is the orthogonal complement of $\text{ran}(X_g) = \text{ran}(E_g) \cap \text{ran}(F_g)^\perp$ in $\text{ran}(E_g)$, it is straightforward to see that $\text{ran}(\tilde{E}_g) \cap \text{ran}(\tilde{F}_g)^\perp = \{0\}$. For (ii), since $\text{ran}(\tilde{E}_{g'}) \subseteq \text{ran}(E_{g'}) \perp \text{ran}(F_g)$ for every $\{g, g'\} \in E(\overline{G})$, we have $\tilde{F}_g \tilde{E}_{g'} = 0$ as well. Thus, $(\mathcal{H}, \tilde{\mathcal{M}}, \{\tilde{E}_g\}, \{\tilde{F}_g\}, \tilde{\tau})$ is a λ -tracial subspace representation of G in terms of Proposition 4.11.

For the reverse direction, let $(\mathcal{H}, \mathcal{M}, \{E_g\}, \tau)$ be a feasible tuple for Definition 4.2. Let F_g be the projection onto $(\sum_{g' \in N_{\overline{G}}(g)} \text{ran}(E_{g'}))^\perp$. Then $F_g E_{g'} = 0$ for all $\{g, g'\} \in E(\overline{G})$ and $X_g = E_g \wedge (I_{\mathcal{M}} - F_g) = 0$ for all $g \in V(G)$. Since $\tau(E_g) - \tau(X_g) = \tau(E_g) = 1/\lambda$, we get a feasible solution of Proposition 4.21 with value λ . \square

Proposition 4.22. *Let G be a graph. Then $\mathcal{H}_{tr}(G)$ equals the infimum of λ for which there exist a Hilbert space \mathcal{H} , a von Neumann algebra $\mathcal{M} \subseteq B(\mathcal{H})$ containing projections E_g and F_g for all $g \in V(G)$, with a normal tracial state $\tau : \mathcal{M} \rightarrow \mathbb{C}$, such that*

- (i) $\|E_g(I_{\mathcal{M}} - F_g)\| < 1$ for any $g \in V(G)$,
- (ii) $F_g E_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$,
- (iii) $\frac{\tau(E_g)}{\tau(E) - \tau(X)} \geq \frac{1}{\lambda}$ for all $g \in V(G)$, where E and F are the projections onto the closures of the subspaces $\sum_{g \in V(G)} \text{ran}(E_g)$ and $\sum_{g \in V(G)} \text{ran}(F_g)$, respectively and X is the projection onto $\text{ran}(E) \cap \text{ran}(F)^\perp$.

Proof. Denote the infimum defined in the proposition by $\hat{\mathcal{H}}_{tr}(G)$. We first show $\hat{\mathcal{H}}_{tr}(G) \leq \mathcal{H}_{tr}(G)$. Let $(\mathcal{H}, \mathcal{M}, \{E_g\}, \{F_g\}, \tau)$ be a λ -tracial projection representation of G . Note that the projections E and X defined as in (iii) above commute. In particular, $E - X$ is the projection onto the orthogonal complement of $\text{ran}(E) \cap \text{ran}(F)^\perp$ in $\text{ran}(E)$ and therefore $\tau(E - X) \leq \tau(I_{\mathcal{M}}) = 1$. This shows that $\frac{\tau(E_g)}{\tau(E) - \tau(X)} \geq \tau(E_g) \geq \frac{1}{\lambda}$ for every $g \in V(G)$. Then it is clear that the same tuple $(\mathcal{H}, \mathcal{M}, \{E_g\}, \{F_g\}, \tau)$ satisfies the conditions of Proposition 4.22 with value λ . The inequality $\hat{\mathcal{H}}_{tr}(G) \leq \mathcal{H}_{tr}(G)$ follows by taking the infimum over all such λ .

Conversely, for every feasible solution $(\mathcal{H}, \mathcal{M}, \{E_g\}, \{F_g\}, \tau)$ of $\hat{\mathcal{H}}_{tr}(G)$ with value λ , we construct a λ -tracial subspace representation of G in terms of Proposition 4.21. Let \tilde{E}_g be the left support projection of $(E - X)E_g$ and \tilde{F}_g be the left support projection of $F_g(E - X)$ for every $g \in V(G)$. Note that $\text{cl}(\text{ran}((E - X)E_g)) \subseteq \text{ran}(E - X)$ and $\ker(F_g(E - X))^\perp \subseteq \text{ran}(E - X)$.⁵ Thus \tilde{E}_g and \tilde{F}_g commute with $E - X$ and we have $\tilde{E}_g(E - X) = \tilde{E}_g$ and $\tilde{F}_g(E - X) = \tilde{F}_g$. Let $\tilde{\mathcal{M}}$ be the von Neumann algebra generated by $\{\tilde{E}_g\}, \{\tilde{F}_g\}$ and $E - X$ (as the identity element). $\tilde{\mathcal{M}}$ is a subset of \mathcal{M} because $\tilde{E}_g, \tilde{F}_g \in \mathcal{M}$ are support projections. Define the tracial state $\tilde{\tau} : \tilde{\mathcal{M}} \rightarrow \mathbb{C}$ by $\tilde{\tau}(Y) = \frac{\tau(Y)}{\tau(E - X)}$. We verify that $(\mathcal{H}, \tilde{\mathcal{M}}, \{\tilde{E}_g\}, \{\tilde{F}_g\}, \tilde{\tau})$ is a λ -tracial subspace representation in terms of Proposition 4.21.

The key is to show that $F_g(E - X)E_{g'} = F_g E_{g'}$ for every $g, g' \in V(G)$. Indeed, $EE_{g'} = E_{g'}$ for any $g' \in V(G)$ since $E_{g'} \subseteq \bigvee_{g \in V(G)} E_g = E$. Also, $F_g X = 0$ because X is the projection onto $\text{ran}(E) \cap (\sum_{g \in V(G)} \text{ran}(F_g))^\perp$, which is a subspace of $\text{ran}(F_g)^\perp$ for any $g \in V(G)$. Thus, $F_g(E - X)E_{g'} = \tilde{F}_g(EE_{g'}) - (FX)E_{g'} = \tilde{F}_g E_{g'}$.

Next, we show that $\text{ran}(\tilde{E}_g) \cap \text{ran}(\tilde{F}_g)^\perp = \{0\}$ for every $g \in V(G)$. Let

$$v \in \text{ran}(\tilde{E}_g) \cap \text{ran}(\tilde{F}_g)^\perp = \text{cl}(\text{ran}((E - X)E_g)) \cap \ker(F_g(E - X)).$$

Then $v = \lim_{i \rightarrow \infty} (E - X)E_g u_i$ for some $\{u_i\}_{i \in \mathbb{N}} \subseteq \mathcal{H}$ and $F_g(E - X)v = 0$. Note that

$$\lim_{i \rightarrow \infty} F_g E_g u_i = \lim_{i \rightarrow \infty} (F_g(E - X))((E - X)E_g u_i) = F_g(E - X)v = 0.$$

Using condition (i) (i.e. $\|E_g(I_{\mathcal{M}} - F_g)\| = \delta_g < 1$) and the triangle inequality, we obtain

$$\|F_g E_g u_i\| \geq \|E_g u_i\| - \|(I_{\mathcal{M}} - F_g)E_g u_i\| \geq \|E_g u_i\| - \|E_g(I_{\mathcal{M}} - F_g)\| \|E_g u_i\| \geq (1 - \delta_g) \|E_g u_i\|.$$

⁵For the second, we know that $\ker(E - X) \subseteq \ker(F_g(E - X))$, thus $\ker(F_g(E - X))^\perp \subseteq \ker(E - X)^\perp = \text{ran}(E - X)$.

Therefore

$$\lim_{i \rightarrow \infty} \|E_g u_i\| \leq (1 - \delta_g)^{-1} \lim_{i \rightarrow \infty} \|F_g E_g u_i\| = 0$$

which implies $v = \lim_{i \rightarrow \infty} (E - X)E_g u_i = 0$.

We then verify that $\tilde{F}_g \tilde{E}_{g'} = 0$ for all $\{g, g'\} \in E(\overline{G})$. Note that \tilde{F}_g is the right support projection of $F_g(E - X)$, we have $F_g((E - X)E_{g'}) = 0$ since $F_g(E - X)E_{g'} = F_g E_{g'} = 0$ for $\{g, g'\} \in E(\overline{G})$. This implies $\tilde{F}_g \tilde{E}_{g'} = 0$ since $\tilde{E}_{g'}$ is the left support projection of $(E - X)E_{g'}$.

Finally we verify that $\tilde{\tau}(\tilde{E}_g) \geq \frac{1}{\lambda}$. We prove that $\tau(\tilde{E}_g) = \tau(E_g)$ for all $g \in V(G)$ and hence

$$\tilde{\tau}(\tilde{E}_g) = \frac{\tau(\tilde{E}_g)}{\tau(E - X)} = \frac{\tau(E_g)}{\tau(E - X)} \geq \frac{1}{\lambda}.$$

Using Lemma 2.9 (ii) (where we take $P = E_g$ and $Q = E - X$), it suffices to show that $\text{ran}(E_g) \cap \text{ran}(E - X)^\perp = \{0\}$. Let $v \in \text{ran}(E_g) \cap \text{ran}(E - X)^\perp$. Then we have $(E - X)v = 0$ and $v = E_g v = E v$ because $v \in \text{ran}(E_g) \subseteq \text{ran}(E)$, therefore $Xv = v$. This means that $v \in \text{ran}(X) = \text{ran}(E) \cap \text{ran}(F)^\perp$. On the other hand, note that $\text{ran}(F)^\perp \subseteq \text{ran}(F_g)^\perp$. We conclude that $v \in \text{ran}(E_g) \cap \text{ran}(F_g)^\perp$, which by assumption $\text{ran}(E_g) \cap \text{ran}(F_g)^\perp = \{0\}$. That completes the proof. \square

We use Propositions 4.15, 4.21 and 4.22 to prove super-multiplicativity of tracial Haemers bound.

Proposition 4.23. *For any graphs G and H , $\mathcal{H}_{tr}(G \boxtimes H) \geq \mathcal{H}_{tr}(G)\mathcal{H}_{tr}(H)$.*

Proof. Let $(\mathcal{H}, \mathcal{M}, \{E_{g,h}\}, \{F_{g,h}\}, \tau)$ be a λ -tracial projection representation of $G \boxtimes H$ as in Definition 4.14. Let E_g and F_g be the projections onto $\text{cl}(\sum_{h \in V(H)} \text{ran}(E_{g,h}))$ and $\text{cl}(\sum_{h \in V(H)} \text{ran}(F_{g,h}))$ for any fixed g , respectively. By construction we have that E_g and F_g belong to \mathcal{M} . Note that $F_{g,h} E_{g',h'} = 0$ for any $\{g, g'\} \in E(\overline{G})$ and any $h, h' \in V(H)$ because such $\{(g, h), (g', h')\} \in E(\overline{G \boxtimes H})$. This implies that $F_g E_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$. Let X_g be the projection onto the subspace $\text{ran}(E_g) \cap \text{ran}(F_g)^\perp$. Let $\mu^{-1} = \min\{\tau(E_g) - \tau(X_g) : g \in V(G)\}$. We obtained a feasible solution of $\mathcal{H}_{tr}(G)$ with value μ as given in Proposition 4.21. This shows that $\mathcal{H}_{tr}(G) \leq \mu$.

We now construct a λ/μ -tracial projection representation of H in terms of Proposition 4.22. Let $g_0 \in V(G)$ be such that $\mu^{-1} = \tau(E_{g_0}) - \tau(X_{g_0})$. Note that for $\{h, h'\} \in E(\overline{H})$, $E_{g_0,h} F_{g_0,h'} = 0$ and for $h \in V(H)$, $\|E_{g_0,h}(I_{\mathcal{M}} - F_{g_0,h})\| < 1$. Thus $(\mathcal{H}, \mathcal{M}, \{E_{g_0,h}\}, \{F_{g_0,h}\}, \tau)$ is a feasible solution of $\mathcal{H}_{tr}(H)$ where we use the formulation given in Proposition 4.22. To compute its value, we note that for every $h \in V(H)$ we have

$$\frac{\tau(E_{g_0,h})}{\tau(E_{g_0}) - \tau(X_{g_0})} = \mu \tau(E_{g_0,h}) \geq \frac{\mu}{\lambda}.$$

We conclude that for any $\lambda \geq \mathcal{H}_{tr}(G \boxtimes H)$, we have $\mathcal{H}_{tr}(G)\mathcal{H}_{tr}(H) \leq \mu \cdot \frac{\lambda}{\mu} = \lambda$. Taking the infimum over such λ gives that $\mathcal{H}_{tr}(G \boxtimes H) \geq \mathcal{H}_{tr}(G)\mathcal{H}_{tr}(H)$. \square

As we have shown that the tracial Haemers bound is multiplicative (Propositions 4.20 and 4.23), additive (Propositions 4.13 and 4.17), normalized (Example 4.18) and monotone with respect to \leq_{qc} (Proposition 4.12), we conclude the following:

Corollary 4.24. *The tracial Haemers bound is an element of $\mathbf{X}(\mathcal{G}, \leq_{qc})$.*

4.5 Comparing the fractional and tracial Haemers bounds

A natural question to ask is whether the fractional and tracial Haemers bounds can be strictly separated. Here we show a connection between this question and Connes' embedding conjecture. Connes' embedding conjecture (CEC), first proposed by Connes in his famous work [Con76], is a fundamental problem in the field of operator algebras (we give a precise formulation below). Over decades CEC has been shown to be equivalent to many important conjectures in various branches of operator algebras, mathematics and computer science. A disproof of CEC is announced in the preprint [JNV⁺20]. In [JNV⁺20] the authors take a computational complexity approach based on an equivalent formulation of CEC in terms of non-local games. Given the importance of CEC, finding an alternative proof of its failure is of great interest. We shall show that finding a graph G for which $\mathcal{H}_{tr}(G) \neq \mathcal{H}_f(G)$ would provide such an alternative proof. This is similar to [DP16, Corollary 3.10] which showed the analogous statement for a separation between projective rank and tracial rank.

CEC states that every von Neumann algebra (\mathcal{M}, τ) with a tracial state τ is embeddable into an ultra-product $(M_n)^\omega$ of matrix algebra M_n . In terms of Voiculescu's microstate formulation [Voi02], (\mathcal{M}, τ) is embeddable into $(M_n)^\omega$ if and only if for every finite family $\{x_1, \dots, x_n\} \subseteq \mathcal{M}$ of self-adjoint elements, any $\varepsilon > 0$ and integer $N \geq 1$, there is some $k \in \mathbb{N}$ and $k \times k$ self-adjoint matrices $X_1, \dots, X_n \in M_k$ such that for all $p \leq N$ and all $i_1, \dots, i_p \in [n]$,

$$|\mathrm{tr}(X_{i_1} X_{i_2} \cdots X_{i_p}) - \tau(x_{i_1} x_{i_2} \cdots x_{i_p})| < \varepsilon.$$

Here $\mathrm{tr} = \frac{1}{k} \mathrm{Tr}$ is the normalized trace. We say that (X_1, \dots, X_n) is an (N, ε) -microstate of (x_1, \dots, x_n) . We say a tuple (x_1, \dots, x_n) has a matricial microstate if it admits (N, ε) -microstate for all $N \geq 1$ and $\varepsilon > 0$. The following equivalent formulation is natural.

Proposition 4.25. *Let \mathcal{M} be a von Neumann algebra with a tracial state τ . The followings are equivalent.*

- (i) *Every finite family $\{x_1, \dots, x_n\}$ of self-adjoint elements in (\mathcal{M}, τ) admits a matricial microstate.*
- (ii) *Every finite family $\{e_1, \dots, e_n\}$ of **projections** in (\mathcal{M}, τ) admits a matricial microstate of **projections**, i.e., for all $N \geq 1$ and $\varepsilon > 0$, there is some $k \in \mathbb{N}$ and $k \times k$ **projections** $E_1, \dots, E_n \in M_k$ such that for all $p \leq N$ and all $i_1, \dots, i_p \in [n]$,*

$$|\mathrm{tr}(E_{i_1} E_{i_2} \cdots E_{i_p}) - \tau(e_{i_1} e_{i_2} \cdots e_{i_p})| < \varepsilon.$$

Proof. We first show (ii) \Rightarrow (i). Let $\{x_1, \dots, x_n\}$ be a family of self-adjoint elements. Let $M = \max_i \|x_i\|$. For any $\varepsilon > 0$ and positive integer N , there exist an integer L and self-adjoint elements $\tilde{x}_i = \sum_{j=1}^L \lambda_{i,j} e_{i,j}$ for every $i \in [n]$ of finite spectrum such that $\|x_i - \tilde{x}_i\| \leq \frac{\varepsilon}{NM^{N-1}}$, where the $e_{i,j}$'s are projections. Indeed, let $L = \lceil \frac{2M}{\varepsilon} \rceil = \lceil \frac{2NM^N}{\varepsilon} \rceil$, where $\lceil \cdot \rceil$ is the ceiling function. We can choose $\tilde{x}_i = f(x_i)$, where $f : [-M, M] \rightarrow \mathbb{R}$ is the step function:

$$f(t) = \frac{(2\ell - L)M}{L}, \quad \text{for } t \in \left[\frac{(2\ell - L)M}{L}, \frac{(2(\ell + 1) - L)M}{L} \right) \text{ and } \ell \in [L] \cup \{0\},$$

and $e_{i,j}$ is the spectral projection of x_i on the interval $\left[\frac{(2\ell - L)M}{L}, \frac{(2(\ell + 1) - L)M}{L} \right)$. Then for all $p \leq N$ and $i_1, \dots, i_p \in [n]$ we have

$$|\tau(x_{i_1} x_{i_2} \cdots x_{i_p}) - \tau(\tilde{x}_{i_1} \tilde{x}_{i_2} \cdots \tilde{x}_{i_p})| \leq \|x_{i_1} x_{i_2} \cdots x_{i_p} - \tilde{x}_{i_1} \tilde{x}_{i_2} \cdots \tilde{x}_{i_p}\| \leq \frac{\varepsilon}{NM^{N-1}} \cdot M^{N-1} \cdot N = \varepsilon.$$

By the condition (ii), there exist projection $\{E_{i,j}\}_{1 \leq i \leq n, 1 \leq j \leq m} \subseteq M_k$ such that for all $i_1, \dots, i_p \in [n]$ and $j_1, \dots, j_p \in [m]$,

$$|\mathrm{tr}(E_{i_1, j_1} E_{i_2, j_2} \cdots E_{i_p, j_p}) - \tau(e_{i_1, j_1} e_{i_2, j_2} \cdots e_{i_p, j_p})| < \frac{\varepsilon}{m^N M^N}.$$

Take $X_i = \sum_j \lambda_{i,j} E_{i,j}$ as self-adjoint elements in M_k . We have

$$\begin{aligned} & |\mathrm{tr}_k(X_{i_1} X_{i_2} \cdots X_{i_p}) - \tau(x_{i_1} x_{i_2} \cdots x_{i_p})| \\ & \leq |\mathrm{tr}_k(X_{i_1} X_{i_2} \cdots X_{i_p}) - \tau(\tilde{x}_{i_1} \tilde{x}_{i_2} \cdots \tilde{x}_{i_p})| + |\tau(x_{i_1} x_{i_2} \cdots x_{i_p}) - \tau(\tilde{x}_{i_1} \tilde{x}_{i_2} \cdots \tilde{x}_{i_p})| \\ & \leq \sum_{1 \leq j_1, j_2, \dots, j_p \leq m} |\lambda_{i_1, j_1} \lambda_{i_2, j_2} \cdots \lambda_{i_p, j_p}| \cdot |\mathrm{tr}(E_{i_1, j_1} E_{i_2, j_2} \cdots E_{i_p, j_p}) - \tau(e_{i_1, j_1} e_{i_2, j_2} \cdots e_{i_p, j_p})| + \varepsilon \\ & \leq \sum_{1 \leq j_1, j_2, \dots, j_p \leq m} M^p \frac{\varepsilon}{(mM)^N} + \varepsilon \\ & \leq \varepsilon + \varepsilon = 2\varepsilon \end{aligned}$$

Since ε is arbitrary, this proves (ii) \Rightarrow (i).

(i) \Rightarrow (ii) essentially follows from [DP16, Proposition 3.6]. Indeed, consider a family of projections $\{e_1, \dots, e_n\} \subseteq \mathcal{M}$. Let $C^*(e_1, \dots, e_n)$ be the C^* -algebra generated by $\{e_1, \dots, e_n\}$ and let $C^*(*_n \mathbb{Z}_2)$ be the universal C^* -algebra of n -projections (i.e. the full group C^* -algebra of n -fold free product of \mathbb{Z}_2 , see e.g. [Bla06]). Denote a_i as the projection generator of $C^*(*_n \mathbb{Z}_2)$. By the universality of $C^*(*_n \mathbb{Z}_2)$, there exists a $*$ -homomorphism $\pi : C^*(*_n \mathbb{Z}_2) \rightarrow C^*(e_1, \dots, e_n)$ such that $\pi(a_i) = e_i$ for all i . Moreover, $\tau \circ \pi$ induces a tracial state on $C^*(*_n \mathbb{Z}_2)$ such that for any $i_1, i_2, \dots, i_p \in \{1, \dots, n\}$

$$\tau(e_{i_1} e_{i_2} \cdots e_{i_p}) = \tau \circ \pi(a_{i_1} a_{i_2} \cdots a_{i_p}).$$

By [DP16, Proposition 3.6], for any (N, ε) , we have a $*$ -homomorphism $\pi_k : C^*(*_n \mathbb{Z}_2) \rightarrow M_k$ such that

$$|\tau \circ \pi(a_{i_1} a_{i_2} \cdots a_{i_p}) - \mathrm{tr} \circ \pi_k(a_{i_1} a_{i_2} \cdots a_{i_p})| \leq \varepsilon,$$

for all $p \leq N$ and all $i_1, \dots, i_p \in [n]$. Note that

$$\begin{aligned} \tau \circ \pi(a_{i_1} a_{i_2} \cdots a_{i_p}) &= \tau\left(\pi(a_{i_1}) \pi(a_{i_2}) \cdots \pi(a_{i_p})\right) = \tau(e_{i_1} e_{i_2} \cdots e_{i_p}) \\ \mathrm{tr} \circ \pi_k(a_{i_1} a_{i_2} \cdots a_{i_p}) &= \mathrm{tr}\left(\pi_k(a_{i_1}) \pi_k(a_{i_2}) \cdots \pi_k(a_{i_p})\right), \end{aligned}$$

and $\pi_k(a_1), \pi_k(a_2), \dots, \pi_k(a_n)$ are projections in M_k (since π_k is $*$ -homomorphism). Therefore, $(\pi_k(a_1), \pi_k(a_2), \dots, \pi_k(a_n))$ is an (N, ε) -micro state of (e_1, e_2, \dots, e_n) . \square

We prove that if $\mathcal{H}_{tr}(G) < \mathcal{H}_f(G)$ for some graph G , then every von Neumann algebra admitting a λ -tracial projection representation of G with $\lambda < \mathcal{H}_f(G)$ does not satisfy the ii) in above proposition. We need the following formulation of a (finite-dimensional) subspace representation, which can be seen as the finite-dimensional version of Proposition 4.21. We show that this formulation is equivalent to Definition 4.1 in Appendix B.

Proposition 4.26. *G has a (d, r) -subspace representation (over \mathbb{C}) if and only if there exist subspaces $S_g, T_g \subseteq \mathbb{C}^d$ for all $g \in V(G)$ satisfying*

- (i) $\dim(S_g) - \dim(S_g \cap T_g^\perp) \geq r$ for all $g \in V(G)$;
- (ii) $S_{g'} \subseteq T_g^\perp \forall \{g, g'\} \in E(\overline{G})$.

We also need the following lemma from Dykema and Paulsen (and its corollary). Here $\|x\|_{\tau,2} = \sqrt{\tau(x^*x)}$ denotes the L_2 -norm with respect to the tracial state τ .

Lemma 4.27 ([DP16, Lemma 3.12]). *Let G be a graph and \mathcal{M} be a von Neumann algebra equipped with a tracial state τ . For every $\varepsilon > 0$, there is a $\delta > 0$ such that if $\{E_g\}_{g \in V(G)}$ are projections in \mathcal{M} satisfying $\tau(E_g E_{g'}) < \delta$ for all $\{g, g'\} \in E(G)$, then there exist projections $\{\tilde{E}_g\}_{g \in V(G)}$ in \mathcal{M} satisfying $\tilde{E}_g \tilde{E}_{g'} = 0$ for all $\{g, g'\} \in E(G)$ and $\|E_g - \tilde{E}_g\|_{\tau,2} \leq \varepsilon$ for every $g \in V(G)$.*

Corollary 4.28. *Let G be a graph. For every $\varepsilon > 0$, there is a $\delta > 0$ such that if $\{E_g\}_{g \in V(G)}$ and $\{F_g\}_{g \in V(G)}$ are projections in a von Neumann algebra \mathcal{M} satisfying*

$$\tau(F_g E_{g'}) < \delta \quad \text{for all } \{g, g'\} \in E(G),$$

then there exist projections $\{\tilde{E}_g\}_{g \in V(G)}$ and $\{\tilde{F}_g\}_{g \in V(G)}$ in \mathcal{M} satisfying

$$\tilde{F}_g \tilde{E}_{g'} = 0 \quad \text{for all } \{g, g'\} \in E(G), \quad \text{and} \quad \|E_g - \tilde{E}_g\|_{\tau,2}, \|F_g - \tilde{F}_g\|_{\tau,2} \leq \varepsilon \quad \text{for every } g \in V(G).$$

Proof. Let G be a graph and let H be the bipartite graph constructed from G whose left and right vertex sets are two copies of the vertices of G . There is an edge between a vertex g in the left copy and a vertex g' in the right copy if and only if $\{g, g'\} \in E(G)$. For a vertex $g \in V(G)$ we assign E_g to the left copy of g and F_g to the right copy of g . The corollary then follows from applying Lemma 4.27 to H . \square

We then show the main result of this section.

Theorem 4.29. *If $\mathcal{H}_{tr}(G) < \mathcal{H}_f(G)$ for a graph G , then the von Neumann algebra (\mathcal{M}, τ) in any λ -tracial projection representation of G for $\lambda < \mathcal{H}_f(G)$ does not satisfy Connes' embedding conjecture. In particular, the projections $\{E_g, F_g\}_{g \in V(G)}$ in the λ -tracial projection representation (as given in Definition 4.14) do not admit a matricial microstate.*

Proof. Let \mathcal{M} be a von Neumann algebra with a normal tracial state τ . Assume that $\{e_g\}_{g \in V(G)}$ and $\{f_g\}_{g \in V(G)}$ are projections satisfying the conditions in Definition 4.14, i.e.

- (i) $\|e_g(I_{\mathcal{M}} - f_g)\| < 1$ for any $g \in V(G)$.
- (ii) $f_g e_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$,
- (iii) $\tau(e_g) \geq \frac{1}{\lambda}$ for all $g \in V(G)$.

We show by contradiction that $\{e_g\} \cup \{f_g, I_{\mathcal{M}} - f_g\}_{g \in V(G)}$ cannot have a matricial microstate. By condition (i), we have $\lim_{n \rightarrow \infty} \|(e_g(I_{\mathcal{M}} - f_g)e_g)^n\| = 0$. Then for every $\varepsilon > 0$, there exists a finite integer ℓ such that for any $g \in V(G)$ and $n \geq \ell$,

$$\tau((e_g(I_{\mathcal{M}} - f_g)e_g)^n) \leq \varepsilon.$$

Take the δ corresponding to $\varepsilon/(3\ell)$ in Corollary 4.28. Suppose $\{e_g\} \cup \{f_g, I_{\mathcal{M}} - f_g\}_{g \in V(G)}$ admits a $(3\ell, \varepsilon_2)$ -matricial microstate for $\varepsilon_2 := \min\{\delta, \varepsilon\}$. Then there exists $k \in \mathbb{N}$ and projections $E_g, F_g \subseteq M_k$ for $g \in V(G)$ satisfying

- (i) $\text{tr}((E_g(I_k - F_g)E_g)^\ell) \leq \varepsilon + \varepsilon_2$ for each $g \in V(G)$,
- (ii) $\text{tr}(F_g E_{g'}) \leq \varepsilon_2$ for any $\{g, g'\} \in E(\overline{G})$,

(iii) $\text{tr}(E_g) \geq \frac{1}{\lambda} - \varepsilon_2$ for all $g \in V(G)$.

We now use Corollary 4.28 (on the complement graph \overline{G}) to convert (ii) to an orthogonality condition. There exist projections (again of size $k \times k$) \tilde{E}_g, \tilde{F}_g for $g \in V(G)$ such that $\tilde{F}_g \tilde{E}_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$ and for all $g \in V(G)$ we have

$$\|E_g - \tilde{E}_g\|_{\text{tr},2} \leq \frac{\varepsilon}{3\ell}, \quad \|F_g - \tilde{F}_g\|_{\text{tr},2} \leq \frac{\varepsilon}{3\ell}.$$

Then we have

$$\begin{aligned} \text{tr}((\tilde{E}_g(I_k - \tilde{F}_g)\tilde{E}_g)^\ell) &\leq \text{tr}((E_g(I_k - F_g)E_g)^\ell) + \text{tr}(|(E_g(I_k - F_g)E_g)^\ell - (\tilde{E}_g(I_k - \tilde{F}_g)\tilde{E}_g)^\ell|) \\ &\leq \varepsilon + \varepsilon_2 + \left\| (E_g(I_k - F_g)E_g)^\ell - (\tilde{E}_g(I_k - \tilde{F}_g)\tilde{E}_g)^\ell \right\|_{\text{tr},2} \\ &\leq \varepsilon + \varepsilon + 3\ell \cdot \frac{\varepsilon}{3\ell} = 3\varepsilon, \end{aligned}$$

where the second inequality uses $\text{tr}(|A|) \leq \|A\|_{\text{tr},2}$ for the normalized trace “tr” and the third inequality uses the triangle inequality 3ℓ times and the fact that the norm of products of projections is at most 1.

This shows that for any $\varepsilon > 0$, there exist $\ell, k \in \mathbb{N}$ and $k \times k$ projections \tilde{E}_g, \tilde{F}_g for all $g \in V(G)$ satisfying:

- $\text{tr}((\tilde{E}_g(I_k - \tilde{F}_g)\tilde{E}_g)^\ell) \leq 3\varepsilon$ for any $g \in V(G)$,
- $\tilde{F}_g \tilde{E}_{g'} = 0$ for any $\{g, g'\} \in E(\overline{G})$.

Let $X_g = \tilde{E}_g \wedge (I_k - \tilde{F}_g)$ be the projection onto $\text{ran}(\tilde{E}_g) \cap \text{ran}(I_k - \tilde{F}_g)$. By monotonicity of the sequence $(\tilde{E}_g(I_k - \tilde{F}_g)\tilde{E}_g)^n \rightarrow X_g$, we have $\text{tr}(X_g) \leq \text{tr}((\tilde{E}_g(I_k - \tilde{F}_g)\tilde{E}_g)^\ell) \leq 3\varepsilon$. We show that this gives a feasible solution of $\mathcal{H}_f(G)$ in terms of Proposition 4.26. Let $S_g = \text{ran}(\tilde{E}_g)$ and $T_g = \text{ran}(\tilde{F}_g)$. Note that

$$\dim(S_g) - \dim(S_g \cap T_g^\perp) = k(\text{tr}(\tilde{E}_g - X_g)) \geq k(\text{tr}(E_g) - \text{tr}(|E_g - \tilde{E}_g|) - 3\varepsilon) \geq k\left(\frac{1}{\lambda} - \varepsilon_2 - \frac{\varepsilon}{3\ell} - 3\varepsilon\right).$$

Thus

$$\mathcal{H}_f(G) \leq \frac{1}{\frac{1}{\lambda} - 5\varepsilon} = \frac{\lambda}{1 - 5\lambda\varepsilon}.$$

Taking ε such that $\frac{\lambda}{1 - 5\lambda\varepsilon} < \mathcal{H}_f(G)$, we reach a contradiction. \square

5 Inertia upper bound on the commuting quantum independence number

We recall the infinite-dimensional version of the projective packing number.

Definition 5.1 (Tracial packing number). *Let G be a graph. A λ -tracial packing of G is a C^* -algebra \mathcal{A} containing projections $\{E_g\} \subseteq \mathcal{A}$ and equipped with a tracial state τ , such that $E_g E_{g'} = 0$ for all $\{g, g'\} \in E(G)$ and $\lambda = \sum_{g \in V(G)} \tau(E_g)$. The tracial packing number of G , which we denote by $\alpha_{\text{tr}}(G)$, is defined as the supremum of λ over all possible λ -tracial packing of G .*

Note that the value $\alpha_{\text{tr}}(G)$ in the above definition remains the same if we replace C^* -algebras by von Neumann algebras and replace tracial states by normal tracial states. When restricting to finite-dimensional C^* -algebras \mathcal{A} , the above definition gives the projective packing number $\alpha_p(G)$, which is an upper bound on $\alpha_q(G)$ [Rob13]. Similarly, the tracial packing number $\alpha_{\text{tr}}(G)$ upper bounds $\alpha_{qc}(G)$ (cf. [GdLL18, Proposition 9]).

Definition 5.2 (Inertia bound). *Let $A \in M_s$ be a Hermitian matrix. A subspace $S \subseteq \mathbb{C}^s$ is totally isotropic with respect to A if for any vector $v \in S$, $\langle v, Av \rangle = 0$. The inertia bound of A , denoted as $n(A)$, is the maximum dimension of totally isotropic subspaces of A .*

It was proved in [WE18, WEA19] that for any weighted adjacency matrix A of G , we have $\alpha_p(G) \leq n(W)$. We show this is also true for the tracial packing number. We first prove the following characterization of the inertia bound. Note that the original definition can be viewed as taking $\mathcal{M} = \mathbb{C}$.

Proposition 5.3. *Let $A \in M_s$ be a Hermitian matrix and let \mathcal{M} be a von Neumann algebra equipped with a normal tracial state τ . Then*

$$n(A) = \sup_E \{ \text{Tr} \otimes \tau(E) \mid E \in M_s \otimes \mathcal{M} \text{ projection s.t. for any } 0 \leq P \leq E, \text{Tr} \otimes \tau((A \otimes I_{\mathcal{M}})P) = 0 \}.$$

Proof. Let $S \subseteq \mathbb{C}^s$ be a totally isotropic subspace of A and let E_S be the projection onto S . Then the direction “ \leq ” follows from the fact that $E_S \otimes I_{\mathcal{M}}$ is feasible for the right hand side supremum. For the other direction, recall from [WE18, Lemma 1] that

$$n(A) = \text{Tr}(E_0) + \min\{\text{Tr}(E_+), \text{Tr}(E_-)\}.$$

where E_0 (resp. E_+ and E_-) is the projection onto the eigenspace of A of zero spectrum (resp. positive spectrum and negative spectrum). Without loss of generality, we assume $\text{rank}(E_+) = \text{Tr}(E_+) \geq \text{Tr}(E_-) = \text{rank}(E_-)$ and the smallest positive eigenvalue of A is $\lambda > 0$. Then $E_+ \otimes I_{\mathcal{M}}$ is the spectral projection of $A \otimes I_{\mathcal{M}}$ on $[\lambda, \infty)$. For any projection $E \in M_s \otimes \mathcal{M}$ such that

$$\text{Tr} \otimes \tau((A \otimes I_{\mathcal{M}})P) = 0, \forall 0 \leq P \leq E,$$

we have $(E_+ \otimes I_{\mathcal{M}}) \wedge E = 0$. Indeed, for any nonzero projection $P \leq E_+ \otimes I_{\mathcal{M}}$ and $P \leq E$,

$$\text{Tr} \otimes \tau((A \otimes I_{\mathcal{M}})P) \geq \lambda \text{Tr} \otimes \tau(P) > 0 = \text{Tr} \otimes \tau((A \otimes I_{\mathcal{M}})P).$$

which is a contradiction. Thus by Lemma 2.9 (i),

$$\begin{aligned} \text{Tr} \otimes \tau(E) &= \text{Tr} \otimes \tau(E) + \text{Tr} \otimes \tau(E_+ \otimes I_{\mathcal{M}}) - \text{Tr} \otimes \tau(E_+ \otimes I_{\mathcal{M}}) \\ &= \text{Tr} \otimes \tau(E \vee (E_+ \otimes I_{\mathcal{M}})) + \text{Tr} \otimes \tau(E \wedge (E_+ \otimes I_{\mathcal{M}})) - \text{Tr} \otimes \tau(E_+ \otimes I_{\mathcal{M}}) \\ &\leq \text{Tr} \otimes \tau(I_s \otimes I_{\mathcal{M}}) - \text{Tr} \otimes \tau(E_+ \otimes I_{\mathcal{M}}) = s - \text{Tr}(E_+) = \text{Tr}(E_0) + \text{Tr}(E_-) = n(A), \end{aligned}$$

where the inequality uses the fact that $E \wedge (E_+ \otimes I_{\mathcal{M}}) = 0$ and $E \vee (E_+ \otimes I_{\mathcal{M}})$ is a projection on $M_s \otimes \mathcal{M}$. Taking the supremum over all such E completes the proof. \square

Proposition 5.4. *Let G be a graph and let A be a weighted adjacency matrix of G . We have $\alpha_{qc}(G) \leq \alpha_{tr}(G) \leq n(A)$.*

Proof. We only need to prove the last inequality. Let $(\mathcal{M}, \{E_g\}, \tau)$ be a tracial packing of G . Define the projection $E = \sum_{g \in V(G)} e_g e_g^* \otimes E_g \in M_n \otimes \mathcal{M}$, where $n = |V(G)|$ and $\{e_g\}$ is a set of standard basis of \mathbb{C}^n . Note that

$$E(A \otimes I_{\mathcal{M}})E = \sum_{g, g' \in V(G)} \langle e_g, A e_{g'} \rangle e_g e_{g'}^* \otimes E_g E_{g'} = 0,$$

since $\langle e_g, A e_{g'} \rangle = 0$ for all $\{g, g'\} \notin E(G)$ and $E_g E_{g'} = 0$ for all $\{g, g'\} \in E(G)$. Then for any $0 \leq P \leq E$,

$$\text{Tr} \otimes \tau((A \otimes I_{\mathcal{M}})P) = \text{Tr} \otimes \tau((A \otimes I_{\mathcal{M}})EPE) = \text{Tr} \otimes \tau(E(A \otimes I_{\mathcal{M}})EP) = 0.$$

It follows from Proposition 5.3 that

$$\mathrm{Tr} \otimes \tau(E) = \sum_{g \in V(G)} \tau(E_g) \leq n(A),$$

which completes the proof. \square

Acknowledgments. We thank Monique Laurent for many interesting and helpful discussions and comments, and we thank the anonymous reviewers for many helpful comments. Part of this work was done when SG and YL were postdoctoral researchers at Centrum Wiskunde & Informatica (CWI) and QuSoft, the Netherlands, and when LG was a postdoctoral researcher at Zentrum Mathematik, Technische Universität München. SG’s research was partially supported by SIRTEQ-grant QuIPP. YL’s research was partially supported by MEXT Quantum Leap Flagship Program (MEXT Q-LEAP) Grant Number JPMXS0120319794.

References

- [BBG13] Jop Briët, Harry Buhrman, and Dion Gijswijt. Violating the Shannon capacity of metric graphs with entanglement. *Proceedings of the National Academy of Sciences*, 110(48):19227–19232, 2013. doi:10.1073/pnas.1203857110.
- [BBL⁺15] Jop Briët, Harry Buhrman, Monique Laurent, Teresa Piovosan, and Giannicola Scarpa. Entanglement-assisted zero-error source-channel coding. *IEEE Transactions on Information Theory*, 61(2):1124–1138, 2015. doi:10.1109/TIT.2014.2385080.
- [BC19] Boris Bukh and Christopher Cox. On a fractional version of Haemers’ bound. *IEEE Transactions on Information Theory*, 65(6):3340–3348, June 2019. doi:10.1109/TIT.2018.2889108.
- [Bei10] Salman Beigi. Entanglement-assisted zero-error capacity is upper-bounded by the Lovász ϑ function. *Physical Review A*, 82:010303, Jul 2010. doi:10.1103/PhysRevA.82.010303.
- [Bla06] Bruce Blackadar. *Operator algebras: theory of C^* -algebras and von Neumann algebras*, volume 122. Springer Science & Business Media, 2006.
- [Bla13] Anna Blasiak. *A graph-theoretic approach to network coding*. PhD thesis, Cornell University, 2013. URL: <https://ecommons.cornell.edu/bitstream/handle/1813/34147/ab675.pdf>.
- [BSST02] Charles H. Bennett, Peter W. Shor, John A. Smolin, and Ashish V. Thapliyal. Entanglement-assisted capacity of a quantum channel and the reverse Shannon theorem. *IEEE Transactions on Information Theory*, 48(10):2637–2655, 2002. doi:10.1109/TIT.2002.802612.
- [CLMW10] Toby S. Cubitt, Debbie Leung, William Matthews, and Andreas Winter. Improving zero-error classical communication with entanglement. *Physical Review Letters*, 104:230503, Jun 2010. doi:10.1103/PhysRevLett.104.230503.
- [CMN⁺07] Peter J. Cameron, Ashley Montanaro, Michael W. Newman, Simone Severini, and Andreas Winter. On the quantum chromatic number of a graph. *Electronic Journal of Combinatorics*, 14(1), 2007.

- [CMR⁺14] Toby S. Cubitt, Laura Mančinska, David E. Roberson, Simone Severini, Dan Stahlke, and Andreas Winter. Bounds on entanglement-assisted source-channel coding via the Lovász theta number and its variants. *IEEE Transactions on Information Theory*, 60(11):7330–7344, 2014. [arXiv:1310.7120](#), [doi:10.1109/TIT.2014.2349502](#).
- [CMSS14] André Chailloux, Laura Mancinska, Giannicola Scarpa, and Simone Severini. Graph-theoretical Bounds on the Entangled Value of Non-local Games. In Steven T. Flammia and Aram W. Harrow, editors, *9th Conference on the Theory of Quantum Computation, Communication and Cryptography (TQC 2014)*, volume 27 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 67–75, Dagstuhl, Germany, 2014. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik. URL: <http://drops.dagstuhl.de/opus/volltexte/2014/4807>, [doi:10.4230/LIPIcs.TQC.2014.67](#).
- [Con76] Alain Connes. Classification of injective factors Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$. *Annals of Mathematics*, pages 73–115, 1976. [doi:10.2307/1971057](#).
- [Cve73] D.M. Cvetković. Inequalities obtained on the basis of the spectrum of the graph. *Studia Sci. Math. Hungar.*, 8:433–436, 1973.
- [DP16] Kenneth J. Dykema and Vern I. Paulsen. Synchronous correlation matrices and Connes’ embedding conjecture. *Journal of Mathematical Physics*, 57(1):015214, 2016. [doi:10.1063/1.4936751](#).
- [DSW13] Runyao Duan, Simone Severini, and Andreas Winter. Zero-Error communication via quantum channels, noncommutative graphs, and a quantum Lovász number. *IEEE Transactions on Information Theory*, 59(2):1164–1174, Feb 2013. [doi:10.1109/TIT.2012.2221677](#).
- [Fek23] M. Fekete. Über die verteilung der wurzeln bei gewissen algebraischen gleichungen mit ganzzahligen koeffizienten. *Mathematische Zeitschrift*, 17(1):228–249, Dec 1923. [doi:10.1007/BF01504345](#).
- [GdLL18] Sander Gribling, David de Laat, and Monique Laurent. Bounds on entanglement dimensions and quantum graph parameters via noncommutative polynomial optimization. *Mathematical programming*, 170(1):5–42, 2018. [doi:10.1007/s10107-018-1287-z](#).
- [Hae78] Willem Haemers. An upper bound for the Shannon capacity of a graph. *Colloquia Mathematica Societatis János Bolyai*, 25:267–272, 1978.
- [Hae79] Willem Haemers. On some problems of Lovász concerning the Shannon capacity of a graph. *IEEE Transactions on Information Theory*, 25(2):231–232, 1979. [doi:10.1109/TIT.1979.1056027](#).
- [Ji13] Zhengfeng Ji. Binary constraint system games and locally commutative reductions. [arXiv:1310.3794](#), 2013. [arXiv:1310.3794](#).
- [JNV⁺20] Zhengfeng Ji, Anand Natarajan, Thomas Vidick, John Wright, and Henry Yuen. $\text{MIP}^* = \text{RE}$. [arXiv:2001.04383](#), January 2020. [arXiv:2001.04383](#).
- [Knu94] Donald E. Knuth. The sandwich theorem. *The Electronic Journal of Combinatorics*, 1(1):1, 1994.

- [KR83] Richard V. Kadison and John R. Ringrose. *Fundamentals of the theory of operator algebras. Volume I: Elementary theory*. Graduate Studies in Mathematics, 1983.
- [KR86] Richard V. Kadison and John R. Ringrose. *Fundamentals of the theory of operator algebras. Volume II: Advanced theory*. Graduate Studies in Mathematics, 1986.
- [LMM⁺12] Debbie Leung, Laura Mančinska, William Matthews, Māris Ozols, and Aidan Roy. Entanglement can increase asymptotic rates of zero-error classical communication over classical channels. *Communications in Mathematical Physics*, 311(1):97–111, Apr 2012. doi:[10.1007/s00220-012-1451-x](https://doi.org/10.1007/s00220-012-1451-x).
- [Lov75] László Lovász. On the ratio of optimal integral and fractional covers. *Discrete Mathematics*, 13(4):383–390, 1975. doi:[10.1016/0012-365X\(75\)90058-8](https://doi.org/10.1016/0012-365X(75)90058-8).
- [Lov79] László Lovász. On the Shannon capacity of a graph. *IEEE Transactions on Information theory*, 25(1):1–7, 1979. doi:[10.1109/TIT.1979.1055985](https://doi.org/10.1109/TIT.1979.1055985).
- [Lov86] László Lovász. *An Algorithmic Theory of Numbers, Graphs and Convexity*. Society for Industrial and Applied Mathematics, 1986. doi:[10.1137/1.9781611970203](https://doi.org/10.1137/1.9781611970203).
- [LZ21] Yinan Li and Jeroen Zuiddam. Quantum asymptotic spectra of graphs and non-commutative graphs, and quantum Shannon capacities. *IEEE Transactions on Information Theory*, 67(1):416–432, 2021. doi:[10.1109/TIT.2020.3032686](https://doi.org/10.1109/TIT.2020.3032686).
- [MR16] Laura Mančinska and David E. Roberson. Quantum homomorphisms. *Journal of Combinatorial Theory, Series B*, 118:228–267, 2016. doi:[10.1016/j.jctb.2015.12.009](https://doi.org/10.1016/j.jctb.2015.12.009).
- [MRV16] Laura Mančinska, David E. Roberson, and Antonios Varvisotis. On deciding the existence of perfect entangled strategies for nonlocal games. *Chicago Journal of Theoretical Computer Science*, 2016(5), April 2016.
- [MSS13] Laura Mančinska, Giannicola Scarpa, and Simone Severini. New separations in zero-error channel capacity through projective Kochen–Specker sets and quantum coloring. *IEEE Transactions on Information Theory*, 59(6):4025–4032, 2013. doi:[10.1109/TIT.2013.2248031](https://doi.org/10.1109/TIT.2013.2248031).
- [OP16] Carlos M. Ortiz and Vern I. Paulsen. Quantum graph homomorphisms via operator systems. *Linear Algebra and its Applications*, 497:23 – 43, 2016. doi:<https://doi.org/10.1016/j.laa.2016.02.019>.
- [PSS⁺16] Vern I. Paulsen, Simone Severini, Daniel Stahlke, Ivan G. Todorov, and Andreas Winter. Estimating quantum chromatic numbers. *Journal of Functional Analysis*, 270(6):2188 – 2222, 2016. doi:<https://doi.org/10.1016/j.jfa.2016.01.010>.
- [PT15] Vern I. Paulsen and Ivan G. Todorov. Quantum chromatic numbers via operator systems. *The Quarterly Journal of Mathematics*, 66(2):677–692, 2015. doi:[10.1093/qmath/hav004](https://doi.org/10.1093/qmath/hav004).
- [Rob13] David E. Roberson. *Variations on a Theme: Graph Homomorphisms*. PhD thesis, University of Waterloo, 2013. URL: <https://uwspace.uwaterloo.ca/handle/10012/7814>.

- [RS89] Iain Raeburn and Allan M. Sinclair. The C^* -algebra generated by two projections. *Mathematica Scandinavica*, 65(2):278–290, 1989. URL: <http://www.jstor.org/stable/24491975>.
- [Sch03] Alexander Schrijver. *Combinatorial optimization: Polyhedra and efficiency*, volume 24. Springer Science & Business Media, 2003.
- [Sha48] C. E. Shannon. A mathematical theory of communication. *Bell System Technical Journal*, 27(3):379–423, 1948. doi:<https://doi.org/10.1002/j.1538-7305.1948.tb01338.x>.
- [Sha56] Claude E. Shannon. The zero error capacity of a noisy channel. *Institute of Radio Engineers, Transactions on Information Theory*, IT-2(September):8–19, 1956. doi:[10.1109/TIT.1956.1056798](https://doi.org/10.1109/TIT.1956.1056798).
- [Str88] Volker Strassen. The asymptotic spectrum of tensors. *Journal für die Reine und Angewandte Mathematik. [Crelle's Journal]*, 384:102–152, 1988. doi:[10.1515/crll.1988.384.102](https://doi.org/10.1515/crll.1988.384.102).
- [Voi02] Dan Voiculescu. Free entropy. *Bulletin of the London Mathematical Society*, 34(3):257–278, 2002.
- [WE18] Pawel Wocjan and Clive Elphick. An inertial upper bound for the quantum independence number of a graph. arXiv:1808.10820, 2018. arXiv:[1808.10820](https://arxiv.org/abs/1808.10820).
- [WEA19] Pawel Wocjan, Clive Elphick, and Aida Abiad. Spectral upper bound on the quantum k -independence number of a graph. arXiv:1910.07339, 2019. arXiv:[1910.07339](https://arxiv.org/abs/1910.07339).
- [Zui18] Jeroen Zuiddam. *Asymptotic spectra, algebraic complexity and moment polytopes*. PhD thesis, University of Amsterdam, 2018. URL: <https://dare.uva.nl/search?identifier=9a8030e9-f708-4c95-9d50-f2a5919e75ed>.
- [Zui19] Jeroen Zuiddam. The asymptotic spectrum of graphs and the Shannon capacity. *Combinatorica*, 39:1173–1184, 2019. doi:[10.1007/s00493-019-3992-5](https://doi.org/10.1007/s00493-019-3992-5).

A Relaxing the equality in Definition 4.2

We first show a similar result as Definition 3.6: in the definition of a λ -tracial subspace representation we may relax the equality in (ii) to an inequality $\geq \frac{1}{\lambda}$. The proof strategy is similar to the one for tracial rank in [PSS⁺16, Theorem 6.11].

Proposition A.1. *Let G be a graph. If $\mathcal{M} \subseteq B(\mathcal{H})$ is a von Neumann algebra containing projections E_g for all $g \in V(G)$, with a normal tracial state $\tau : \mathcal{M} \rightarrow \mathbb{C}$, such that, for*

$$E_{N_{\overline{\mathcal{C}}}(g)} = \bigvee_{g' \in N_{\overline{\mathcal{C}}}(g)} E_{g'} = \text{sot-}\lim_{n \rightarrow \infty} \left(\sum_{g' \in N_{\overline{\mathcal{C}}}(g)} E_{g'} \right)^{1/n} \text{ for } g \in V(G),$$

we have

- (i) $\text{ran}(E_g) \cap \text{ran}(E_{N_{\overline{\mathcal{C}}}(g)}) = \{0\}$.
- (ii) $\tau(E_g) \geq \frac{1}{\lambda}$ for all $g \in V(G)$

then G has a λ -tracial subspace representation.

Proof. Define the scalars $c_g = \tau(E_g)$ for $g \in V(G)$. By (ii) we have $c_g \geq 1/\lambda$ for all $g \in V(G)$. Consider the von Neumann algebra $L_\infty(0, 1)$ represented on the Hilbert space $L_2(0, 1)$ with tracial state $\tau_2(f) = \int_{(0,1)} f(x) dx$, where dx is the Lebesgue measure. For a scalar $r \in (0, 1)$, let $P_r = I_{(0,r)}$ be the characteristic function on the open interval $(0, 1)$. Since $c_g \geq 1/\lambda$ for all $g \in V(G)$, we have $(\lambda c_g)^{-1} \leq 1$. For $g \in V(G)$, define $\widehat{E}_g = E_g \otimes P_{(\lambda c_g)^{-1}}$, which is a projection in $\mathcal{M} \overline{\otimes} L_\infty(0, 1) \subseteq B(\mathcal{H} \otimes L_2(0, 1))$. Similarly, let $\widehat{E}_{N_{\overline{G}}(g)}$ be the projection

$$\widehat{E}_{N_{\overline{G}}(g)} := \text{sot-} \lim_{n \rightarrow \infty} \left(\sum_{g' \in N_{\overline{G}}(g)} \widehat{E}_{g'} \right)^{1/n} \text{ for } g \in V(G).$$

Let $\tilde{\mathcal{M}} \subseteq \mathcal{M} \overline{\otimes} L_\infty(0, 1) \subseteq B(\mathcal{H} \otimes L^2(0, 1))$ be the von Neumann algebra generated by the set of projections $\{\widehat{E}_g : g \in V(G)\}$, equipped with the tracial state $\widehat{\tau} = \tau \otimes \tau_2$. Then, for $g \in V(G)$, we have

$$\widehat{\tau}(\widehat{E}_g) = \tau(E_g) \tau_2(P_{(\lambda c_g)^{-1}}) = c_g \cdot \frac{1}{\lambda c_g} = \frac{1}{\lambda}.$$

We are only left to prove $\text{ran}(\widehat{E}_g) \cap \text{ran}(\widehat{E}_{N_{\overline{G}}(g)}) = \{0\}$. Let $x \in \text{ran}(\widehat{E}_g) \cap \text{ran}(\widehat{E}_{N_{\overline{G}}(g)})$. Then there exist $y, y_{g'}^{(N)} \in \mathcal{H} \otimes L^2(0, 1)$ for all $g' \in N_{\overline{G}}(g)$ and $N \in \mathbb{N}$ such that

$$x = \widehat{E}_g y = (E_g \otimes P_{(\lambda c_g)^{-1}}) y$$

and

$$\lim_{N \rightarrow \infty} \left\| \left(\sum_{g' \in N_{\overline{G}}(g)} \widehat{E}_{g'} y_{g'}^{(N)} \right) - x \right\| = \lim_{N \rightarrow \infty} \left\| \left(\sum_{g' \in N_{\overline{G}}(g)} (E_{g'} \otimes P_{(\lambda c_{g'})^{-1}}) y_{g'}^{(N)} \right) - x \right\| = 0.$$

Note that for any linear functional (dual vector) $\mathcal{L} \in (L^2(0, 1))^*$, we have $(I_{\mathcal{A}} \otimes \mathcal{L})(x) \in \text{ran}(E_g)$. Moreover, let $z_{g'}^{(N)} = (I_{\mathcal{A}} \otimes \mathcal{L})((E_{g'} \otimes P_{(\lambda c_{g'})^{-1}}) y_{g'}^{(N)}) \in \mathcal{H}$ for every $g' \in N_{\overline{G}}(g)$, we have

$$\lim_{N \rightarrow \infty} \left\| \left(\sum_{g' \in N_{\overline{G}}(g)} z_{g'}^{(N)} \right) - (I_{\mathcal{A}} \otimes \mathcal{L})(x) \right\| \leq \|\mathcal{L}\| \cdot \lim_{N \rightarrow \infty} \left\| \left(\sum_{g' \in N_{\overline{G}}(g)} (E_{g'} \otimes P_{(\lambda c_{g'})^{-1}}) y_{g'}^{(N)} \right) - x \right\| = 0$$

Thus

$$(I_{\mathcal{A}} \otimes \mathcal{L})(x) \in \text{ran}(E_g) \cap \text{cl} \left(\sum_{g' \in N_{\overline{G}}(g)} \text{ran}(E_{g'}) \right) = \{0\},$$

for any $\mathcal{L} \in L^2(0, 1)^*$, which implies that $x = 0$. Thus we obtain a λ -tracial subspace representation of G . \square

B Equivalent formulation of the subspace representation of a graph

We prove the following equivalent formulation of the subspace representation of a graph (cf. Definition 4.1).

Proposition 4.26. *G has a (d, r) -subspace representation (over \mathbb{C}) if and only if there exist subspaces $S_g, T_g \subseteq \mathbb{C}^d$ for all $g \in V(G)$ satisfying*

- (i) $\dim(S_g) - \dim(S_g \cap T_g^\perp) \geq r$ for all $g \in V(G)$;

(ii) $S_{g'} \subseteq T_g^\perp \forall \{g, g'\} \in E(\overline{G})$.

Proof. Let $\{S_g\}_{g \in V(G)}$ be a (d, r) -subspace representation of G as given in Definition 4.1. Let $T_g = (\sum_{g' \in N_{\overline{G}}(g)} S_{g'})^\perp$. Then it is straightforward to see that $S_{g'} \subseteq T_g^\perp$ if $\{g, g'\} \in E(\overline{G})$ and $S_g \cap T_g^\perp = S_g \cap (\sum_{g' \in N_{\overline{G}}(g)} S_{g'}) = \{0\}$. Thus, $\{S_g\}_{g \in V(G)}$ and $\{T_g\}_{g \in V(G)}$ satisfy the conditions in Proposition 4.26.

On the other hand, let $\{S_g\}_{g \in V(G)}$ and $\{T_g\}_{g \in V(G)}$ satisfy the conditions in Proposition 4.26 with value (d, r) , we construct a (d, r) -subspace representation of G as given in Definition 4.1. For $g \in V(G)$, let \hat{S}_g be the orthogonal complement of $S_g \cap T_g^\perp$ in S_g . By (i), we know that $\dim(\hat{S}_g) \geq r$. Let \tilde{S}_g be a dimension- r subspace of \hat{S}_g . We are left to show that $\tilde{S}_g \cap (\sum_{g' \in N_{\overline{G}}(g)} \tilde{S}_{g'}) = \{0\}$. Note that

$$\left(\sum_{g' \in N_{\overline{G}}(g)} \tilde{S}_{g'} \right) \subseteq \left(\sum_{g' \in N_{\overline{G}}(g)} S_{g'} \right) \subseteq T_g^\perp,$$

where the first inclusion is due to the fact that $\tilde{S}_g \subseteq \hat{S}_g \subseteq S_g$ for any $g \in V(G)$ and the second inclusion is due to (ii). Since \tilde{S}_g is contained in the orthogonal complement of $S_g \cap T_g^\perp$ in S_g , we have $\tilde{S}_g \cap T_g^\perp = \{0\}$ and it follows that $\tilde{S}_g \cap (\sum_{g' \in N_{\overline{G}}(g)} \tilde{S}_{g'}) = \{0\}$. This shows that $\{\tilde{S}_g\}_{g \in V(G)}$ is a (d, r) -subspace representation of G . \square