

Certification of incompatible measurements using quantum steering

Shubhayan Sarkar,¹ Debashis Saha,¹ and Remigiusz Augusiak¹

¹*Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warsaw, Poland*

In recent years nonlocality has been extensively explored as a resource for self-testing—a device-independent way of certification of entangled quantum states and measurements performed on them. However, most of the efforts in designing self-testing schemes were concerned with entangled quantum states, leaving the problem of certification of quantum measurements largely unexplored. Here we address this problem, concentrating on a simpler, one-sided device-independent scenario. We propose a simple scheme for certification of any set of d -outcome projective measurements which do not share any common invariant subspace, termed here genuinely incompatible, and the maximally entangled state of two qudits. We also study robustness of our self-testing statements for a certain class of genuinely incompatible measurements including mutually unbiased bases.

Introduction.—The existence of non-local correlations, usually referred to as nonlocality, is one of the most intriguing features of quantum theory [1]. Formulated for the first time in a rigorous way by Bell in 1964 [2, 3], it is now considered to be one of the key resources accessible within the quantum world. In fact, Bell nonlocality lies at the heart of the field of device-independent (DI) quantum information processing which aims at designing protocols that do not require making assumptions on the devices used to perform them, with the most celebrated example being the DI quantum cryptography [4]. Another application of nonlocality which recently attracts considerable attention are the device-independent certification schemes [5–7], in which the statistics obtained from a device are used to certify relevant quantum properties such as the dimension of the underlying quantum system or whether it is entangled. The strongest among these certification schemes is referred to as self-testing [8], which aims to certify that the device operates on a specific quantum state and performs specific quantum measurements on it. Although recently, there has been a substantial progress in designing self-testing schemes for various quantum systems [9–14], the problem becomes increasingly difficult when one wants to go beyond quantum systems which are locally two-dimensional. In fact, most of the results are devoted to quantum systems which are locally qubits.

To reduce the complexity of the problem, some physically-motivated assumptions could be made on the scenario such as for instance that one of the measuring devices is trusted and performs known measurements. Such a scenario is referred to as one-sided-device-independent and the main resource it features which is used in particular for certification is quantum steering—another form of quantum nonlocality [15–17]. Recently, the possibilities this scenario offers for self-testing of quantum states and measurements were investigated in Refs. [18–22].

The steering scenario seems also to be perfectly suited for proposing certification schemes for quantum measurements, a problem that has been barely explored in the device-independent setting. In fact, most of the existing self-testing schemes based on non-locality are de-

voted to certification of entangled quantum states rather than quantum measurements. In particular, while there exists a self-testing scheme for any entangled bipartite state [23] of arbitrary local dimension, there is no general method for device-independent certification of any set of incompatible quantum measurements. Designing such a general scheme might in fact be an extremely difficult task as there exist sets of incompatible measurements which do not give rise to any non-locality [24]. At the same time, it is known that there is a one-to-one correspondence between quantum steering and measurement incompatibility [25–27].

In this work we address the problem of certification of quantum measurements within the one-sided device-independent scenario, concentrating on projective measurements. We provide a simple scheme for self-testing of families of d -outcome projective incompatible measurements together with the maximally entangled state of two qudits. Our schemes are designed for quantum observables which do not share a common invariant subspace (termed here genuinely incompatible) such as for instance those whose eigenvectors form mutually unbiased bases (MUBs) in d -dimensional Hilbert spaces. To this end, we use a family of steering inequalities inspired by those considered for instance in Refs. [28, 29]. We also discuss the case when the measurements to be certified are not genuinely incompatible and study robustness of our schemes for a particular choice of observables.

Preliminaries—Before presenting our results, we outline the scenario and introduce the relevant terminology.

Steering scenario. We consider a standard bipartite scenario exhibiting quantum steerability, however, for our purposes we formulate it in a way resembling the Bell scenario. Consider two spatially separated observers Alice and Bob who share a quantum state ρ_{AB} acting on $\mathcal{H}_A \otimes \mathcal{H}_B$ and perform quantum measurements on their shares of this state. Contrary to the device-independent setting here we additionally assume that Alice’s measuring device is trusted and performs well-known quantum measurements; Bob’s measurement device is untrusted. We also assume that the dimension of Alice’s Hilbert space is fixed, $\dim \mathcal{H}_A = d$.

We denote Alice's and Bob's measurements by $M_x = \{M_{a|x}\}$ and $N_y = \{N_{b|y}\}$, respectively, where $M_{a|x} \geq 0$ and $N_{b|y} \geq 0$ are the measurement operators that sum up to the identity; $x, y = 1, \dots, N$ and $a, b = 0, \dots, d-1$ denote measurement choices and outcomes, respectively. Correlations observed in this experiment are described by a collection of joint probability distributions $\{p(a, b|x, y)\}$, where $p(a, b|x, y) = \text{Tr}[(M_{a|x} \otimes N_{b|y})\rho_{AB}]$.

To detect that the state ρ_{AB} exhibits quantum steerability (or simply that it is steerable) one can use the following linear Bell-type inequalities

$$\mathcal{B}(p(a, b|x, y)) := \sum_{a, b, x, y} \alpha_{a, b|x, y} p(a, b|x, y) \leq \beta_L, \quad (1)$$

where β_L is the maximal value of $\mathcal{B}(p(a, b|x, y))$ over correlations admitting local hidden state models [15], or, equivalently, local hidden variable models in which Alice's probability distributions are quantum (see Appendix B). If violated, such an inequality implies that ρ_{AB} is steerable. The maximal value of $\mathcal{B}(p(a, b|x, y))$ achievable in quantum theory is denoted β_Q .

Incompatible measurements. Let us also recall the notion of incompatibility of quantum measurements. Consider two projective measurements M_x ($x = 1, 2$) and the associated hermitian observables. We call them incompatible or not jointly measurable if the associated observables do not commute. Below we introduce a more restrictive definition of incompatible measurements that not only requires them to be noncommuting but also not to possess a common invariant subspace.

We notice that for our purposes it will be convenient to express the correlations observed in the above experiment in terms of expectation values instead of probability distributions. Given that here we consider measurements of an arbitrary number of outcomes we appeal to the notion of generalized correlators (see, e.g., Refs. [30, 31]) which are generally complex numbers defined as the Fourier transform of $\{p(a, b|x, y)\}$,

$$\langle A_{k|x} B_{l|y} \rangle = \sum_{a, b=0}^{d-1} \omega^{ak+bl} p(a, b|x, y), \quad (2)$$

where $\omega = \exp(2\pi i/d)$ and $k, l \in \{0, 1, \dots, d-1\}$. In the quantum case these generalized correlators express as $\langle A_{k|x} B_{l|y} \rangle = \text{Tr}[\rho_{AB}(A_x^k \otimes B_{l|y})]$, where A_x are unitary operators whose eigenvalues are ω^i ($i = 0, \dots, d-1$), and the superscripts are operator powers, whereas $B_{l|y}$ are Fourier transforms of Bob's measurements operators, i.e., $B_{l|y} = \sum_b \omega^{bl} N_{b|y}$. The Fourier transform is invertible, and hence for a given y , $B_{k|y}$ fully determine Bob's measurement N_y . Thus, we can safely refer to the set $B_y = \{B_{k|y}\}$ as quantum measurement too. Notice that if N_y is projective, $B_{k|y}$ is k th power of a unitary observable, which, slightly abusing the notation, we also denote B_y , i.e., $B_{k|y} = B_y^k$ (see Appendix A for details).

Self-testing. Self-testing in the steering scenario was first defined in Ref. [18]. Inspired by [32], here we state a

more general version that does not assume the underlying state to be pure and measurements to be projective.

Consider again the above experiment with Alice and Bob performing measurements on a state ρ_{AB} and observing correlations $\{p(a, b|x, y)\}$; recall that A_i are projective and fixed, whereas Bob's measurements $B_i = \{B_{k|y}\}$ are arbitrary. Consider then a reference experiment involving a state $|\psi'\rangle_{AB}$ and measurements on Bob's side $B'_i = \{B'_{k|y}\}$ that we both want to certify. These experiments are equivalent, or, alternatively, $|\psi'\rangle_{AB}$ and B'_i are certified from $\{p(a, b|x, y)\}$ if there exists a unitary $U_B : \mathcal{H}_B \rightarrow \mathcal{H}_B$ such that

$$(\mathbb{1}_A \otimes U_B)\rho_{AB}(\mathbb{1}_A \otimes U_B^\dagger) = |\psi'\rangle\langle\psi'|_{AB'} \otimes \rho_{B''}, \quad (3)$$

$$U_B B_{k|y} U_B = B'_{k|y} \otimes \mathbb{1}_{B''}, \quad (4)$$

where \mathcal{H}_B decomposes as $\mathcal{H}_B = \mathcal{H}_{B'} \otimes \mathcal{H}_{B''}$, and $\rho_{B''}$ and $\mathbb{1}_{B''}$ is some state and the identity acting on $\mathcal{H}_{B''}$.

Results.—We begin by introducing a class of inequalities for testing nonclassicality (1) inspired by those considered for instance in Ref. [29]. Here we state them in the correlator picture as (see Appendix A)

$$\mathcal{B}_d = \sum_{i=1}^N \sum_{k=1}^{d-1} \langle A_i^k \otimes B_{k|i} \rangle \leq \beta_L, \quad (5)$$

where A_i are fixed d -outcome unitary observables whose eigenvalues are powers of ω , whereas $B_{k|i}$ ($i = 1, \dots, N$) are Fourier transforms of Bob's measurement operators that are arbitrary and act on a Hilbert space \mathcal{H}_B of unknown but finite dimension. Notice that steering inequalities in the correlator picture have already been considered, for instance in Ref. [33].

The maximal quantum value of \mathcal{B}_d is $\beta_Q = N(d-1)$ and it is achieved for instance by the maximally entangled state of local dimension d ,

$$|\phi_d^+\rangle = \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |ii\rangle, \quad (6)$$

and Bob's observables that satisfy $B_i = A_i^*$. Then, the maximal classical value of \mathcal{B}_d can be upper bounded as

$$\beta_L \leq \max_{\rho} \sum_{i=1}^N \sum_{k=1}^{d-1} \left| \langle A_i^k \rangle_{\rho} \right|. \quad (7)$$

It follows from this bound that the steering inequality (5) is non-trivial ($\beta_L < \beta_Q$) whenever the observables A_i do not share a common eigenvector. While this last fact is sufficient for the inequality (5) to exhibit quantum violation, it is not for deriving our self-testing statements. In fact, we need to restrict ourselves to a smaller class of quantum observables defined as follows.

Definition. Consider a set of N d -outcome unitary observables A_i acting on \mathbb{C}^d and obeying $A_i^d = \mathbb{1}$. We call them genuinely incompatible (GI) if there is no subspace $V \subset \mathbb{C}^d$ such that $\dim V < d$ and $A_i V \subseteq V$ for all i ; in other words, the only common invariant space of all A_i is the full space \mathbb{C}^d .

An excellent example of d genuinely incompatible (GI) observables are those whose eigenvectors form MUBs in \mathbb{C}^d (see Appendix C) such as for instance $X_d Z_d^k$ for prime d with $k = 0, \dots, d-1$; here, X_d and Z_d are d -dimensional generalizations of the qubit Pauli matrices, defined as $X_d|i\rangle = |i+1\rangle$ and $Z_d|i\rangle = \omega^i|i\rangle$, respectively, where $|i\rangle$ are the elements of the computational basis in \mathbb{C}^d and addition is mod d .

Let us now make a few observations regarding the above definition. First, genuinely incompatible observables do not share a common eigenvector, and thus give rise to a nontrivial inequality (5). The opposite implication is in general not true: observables A_i with no common eigenvector might not be genuinely incompatible. Second, any set of observables containing two GI measurements like X_d and Z_d is genuinely incompatible, however, there exist observables forming GI set while not being pairwise genuinely incompatible (see Appendix C for an example). Third, the following fact holds true for any set of GI observables (see Appendix C for a proof).

Lemma 1. *Consider a set of N d -outcome unitary observables A_i such that $A_i^d = \mathbb{1}_d$, and assume them to be GI. The following implication holds true for any normal matrix P on \mathbb{C}^d : if $[P, A_i] = 0$ for all i , then $P = \lambda \mathbb{1}_d$ for some $\lambda \in \mathbb{C}$.*

Now, we proceed to the main results of this work.

Theorem 1. *Assume that for a given set of GI unitary observables A_i obeying $A_i^d = \mathbb{1}_d$, the steering inequality (5) is maximally violated by a state ρ_{AB} acting on $\mathbb{C}^d \otimes \mathcal{H}_B$ and measurements B_i . Then, the following statements hold true for any d : (i) Bob's measurements are projective, (ii) Bob's Hilbert space decomposes as $\mathcal{H}_B = (\mathbb{C}^d)_{B'} \otimes \mathcal{H}_{B''}$ for some Hilbert space $\mathcal{H}_{B''}$, (iii) there exists a unitary transformation, $U_B : \mathcal{H}_B \rightarrow \mathcal{H}_B$, such that*

$$(\mathbb{1}_A \otimes U_B) \rho_{AB} (\mathbb{1}_A \otimes U_B^\dagger) = |\phi_d^+\rangle \langle \phi_d^+|_{AB'} \otimes \rho_{B''}, \quad (8)$$

where B'' denotes Bob's auxiliary system, and

$$\forall i, \quad U_B B_i U_B^\dagger = A_i^* \otimes \mathbb{1}_{B''}. \quad (9)$$

Proof. Here we present a sketch of the proof; the full version can be found in Appendix C. The maximal quantum value $\beta_Q = N(d-1)$ of the inequality (5) is the same as its algebraic value. Thus, if maximally violated it forces the state ρ_{AB} , and A_i and $B_{k|i}$ to satisfy

$$(A_i^k \otimes B_{k|i}) \rho_{AB} = \rho_{AB} \quad (10)$$

for all i and k . By applying $A_i^{d-k} \otimes B_{d-k|i}$ to both sides and using the fact that, by definition, $B_{d-k|i} = B_{k|i}^\dagger$, one finds that $B_{i|k}$ are all unitary, and thus each Bob's measurements is projective. We then denote $B_{k|i} = B_i^k$. Let us add that throughout our work we assumed for simplicity that Bob's reduced state $\rho_B = \text{Tr} \rho_{AB}$ is full rank [34].

Let us then observe that when ρ_{AB} achieves the maximal quantum violation of our inequality, so do all pure states $|\psi\rangle_{AB}$ belonging to its the support, meaning that

$$(A_i^k \otimes B_i^k) |\psi\rangle_{AB} = |\psi\rangle_{AB} \quad (11)$$

holds true to any i and k and $|\psi\rangle_{AB} \in \text{supp}(\rho_{AB})$. Next, we consider the Schmidt decomposition of this state,

$$|\psi\rangle_{AB} = \sum_{i=0}^{d-1} \lambda_i |e_i\rangle |f_i\rangle, \quad (12)$$

where the coefficients $\lambda_i \geq 0$ satisfy $\sum_i \lambda_i^2 = 1$, and the local vectors $|e_i\rangle \in \mathbb{C}^d$ and $|f_i\rangle \in \mathcal{H}_B$ are orthonormal. Notice that the number of terms in this decomposition is a consequence of the assumption that the dimension of Alice's Hilbert space is at most d .

Let us now make a few manipulations with (12). First, we observe that there exists a unitary U_B such that $|f_i\rangle = U_B |e_i^*\rangle$ for every i , where the asterisk denotes complex conjugation in the standard basis; thus

$$|\psi\rangle_{AB} = (P_A \otimes U_B) \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |e_i\rangle |e_i^*\rangle, \quad (13)$$

where P_A is a matrix diagonal in the $\{|e_i\rangle\}$ basis with eigenvalues $\sqrt{d} \lambda_i$; the fact that A_i are GI implies that P_A is full rank, $\text{rank}(P_A) = d$. We finally observe that the state appearing on the right hand side of (13) is the maximally entangled state (6); consequently

$$|\psi\rangle_{AB} = (P_A \otimes U_B) |\phi_d^+\rangle. \quad (14)$$

Let us also notice that Eq. (10) forces each observable B_i to have a block structure $B_i = \bar{B}_i \oplus E_i$, where \bar{B}_i is a $d \times d$ unitary whose eigenvalues are powers of ω and acts on the support of $\text{Tr}_A \psi_{AB}$, that is, the subspace spanned by the local vectors $|f_i\rangle$. Crucially, although the other block E_i is a proper observable too, it trivially acts on the state $|\psi\rangle_{AB}$, and hence does not give rise to the observed correlations. Thus, below we focus on \bar{B}_i .

Combining Eq. (10) for $k = 1$ with Eq. (14), we obtain

$$(A_i P_A \otimes \tilde{B}_i) |\phi_d^+\rangle = (P_A \otimes \mathbb{1}_B) |\phi_d^+\rangle,$$

where $\tilde{B}_i = U_B \bar{B}_i U_B^\dagger$. Using then the identity $R \otimes Q |\phi_d^+\rangle = R Q^T \otimes \mathbb{1} |\phi_d^+\rangle$ that holds for any matrices R and Q , and omitting the state $|\phi_d^+\rangle$, we arrive at a matrix equation $A_i P_A \tilde{B}_i^T = P_A$. After multiplying this relation by its hermitian conjugation and taking advantage of the facts that A_i and \tilde{B}_i are unitary and that $P_A \geq 0$, we can bring it to a simple condition for P_A :

$$[A_i, P_A] = 0 \quad (i = 1, \dots, N). \quad (15)$$

Given that A_i are genuinely incompatible, Lemma 1 tell us that P_A is proportional to $\mathbb{1}_d$. This means that for

any pure state from $\text{supp}(\rho_{AB})$ there exists a unitary U_B (different for a different state) such that

$$|\psi\rangle_{AB} = (\mathbb{1} \otimes U_B)|\phi_+\rangle. \quad (16)$$

Having Eq. (16), let us now go back to the mixed state ρ_{AB} and consider its eigendecomposition

$$\rho_{AB} = \sum_{s=1}^K p_s |\psi_s\rangle\langle\psi_s|_{AB}, \quad (17)$$

where $|\psi_s\rangle_{AB}$ and $p_s \geq 0$ are the eigenvectors and eigenvalues of ρ_{AB} , respectively. Clearly, every state in this decomposition admits the form (16), meaning that

$$|\psi_s\rangle_{AB} = \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |i\rangle |f_i^s\rangle \quad (18)$$

for some orthonormal bases $\{|f_i^s\rangle\}$ spanning the local supports of $|\psi_s\rangle_{AB}$, denoted by $V_s = \text{supp}(\text{Tr}_A(\psi_s))$. The key observation is that the conditions (10) along with the genuine incompatibility of A_i and the fact that $|\psi_s\rangle_{AB}$ are eigenvectors of ρ_{AB} force all these supports V_s to be pairwise orthogonal, which means that $\mathcal{H}_B = V_1 \oplus \dots \oplus V_K$. Since $\dim V_s = d$, this implies that $\mathcal{H}_B = (\mathbb{C}^d)_{B'} \otimes \mathcal{H}_{B''}$. Moreover, there exists a unitary U_B such that

$$U_B |f_i^s\rangle = |i\rangle_{B'} |s\rangle_{B''}. \quad (19)$$

Combining (17), (12) and (19) we arrive at (8) which together with Eq. (10) leads us to (9). \square

An immediate consequence of the above statement is a simple method of certification of projective measurements that are genuinely incompatible: to certify that certain observables A_i^* have been measured by the untrusted device at Bob's side, it is enough to check whether a corresponding inequality (5) constructed from these observables is maximally violated. In particular, any set of MUBs such as the one mentioned before can be certified in the one-sided device-independent way. To support this claim, the following lemma is proven in Appendix C.

Lemma 2. *Any two d -outcome unitary observables whose eigenbases are mutually unbiased, are GI.*

Weaker form of self-testing.—Let us briefly discuss the case when the set of observables on the trusted side are not GI, but still do not share a common eigenvector and can be used to violate the steering inequality (5) maximally. It can be concluded that when the observables on the trusted side share a common invariant subspace, the steering inequality (5) cannot be used to certify the full observable but only the part acting on this subspace. To give an example let us consider the case $d = 4$ and assume that Alice's observables are $A_1 = Z_4$ and

$$A_2 = \sum_{j=0}^1 (-1)^j (|+j\rangle\langle+j| + \mathbb{i}|-j\rangle\langle-j|), \quad (20)$$

where $|\pm_0\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}$, $|\pm_1\rangle = (|2\rangle \pm |3\rangle)/\sqrt{2}$. It can be checked that any matrix of the form $P = \lambda_1 \mathbb{1}_2 \oplus \lambda_2 \mathbb{1}_2$, where $\mathbb{1}_2$ is the 2×2 identity matrix and $\lambda_i \geq 0$ such that $\lambda_1^2 + \lambda_2^2 = 1/2$, commutes with both A_1 and A_2 from Eq. (20). Thus, if we observe the optimal value of the corresponding steering inequality (5), the proof of Theorem 1 tells us that the shared state is of the form

$$\lambda_1(|00\rangle + |11\rangle) + \lambda_2(|22\rangle + |33\rangle), \quad (21)$$

where in particular one of λ_i might vanish. In other words, maximal violation of our inequality certifies an entangled subspace. However, whether the whole measurement can be certified depends on whether P_A is full rank or not. For instance, if $\lambda_2 = 0$ in this example, then only the part of Bob's measurement which acts on the subspace $\text{span}\{|0\rangle, |1\rangle\}$ can be characterised and it is unitarily equivalent to $\Pi_2 A_i^* \Pi_2$ where $\Pi_2 = |0\rangle\langle 0| + |1\rangle\langle 1|$.

Robustness.—In practical scenarios it becomes relevant to study robustness of our self-testing result against experimental imperfections such as the presence of noise in the preparation device or imperfect measurements. This problem for various combinations of measurements on the trusted side can be addressed numerically using the approach of [35]. However, finding an analytical statement for any d is much more challenging. Here, assuming the underlying state to be pure and Bob's measurements to be projective, we address this issue and provide a way to find robustness bounds analytically for a class of genuinely d -outcome incompatible measurements representing MUBs of any dimension.

Theorem 2. *Consider a pair of genuinely incompatible observables $A_1 = X_d Z_d^l$ with $l \in \{0, 1, \dots, d-1\}$ and $A_2 = Z_d$, and assume that the following steering inequality*

$$\sum_{i=1}^2 \sum_{k=1}^{d-1} \langle A_i^k \otimes B_i^k \rangle \geq 2(d-1) - \varepsilon \quad (22)$$

is violated ε -close to the maximal violation by a state $|\psi\rangle_{AB} \in \mathbb{C}^d \otimes \mathcal{H}_B$ and observables B_i ($i = 1, 2$) acting on \mathcal{H}_B . Then, there exists a unitary operation $U_B : \mathcal{H}_B \rightarrow \mathcal{H}_B$ such that

$$\left\| (\mathbb{1} \otimes U_B)(\mathbb{1} \otimes B_i^k) |\psi\rangle - \mathbb{1} \otimes (B_i')^k |\phi_d^+\rangle \right\| \leq \sqrt{2\varepsilon} + 2\sqrt{d} \sqrt[4]{2\varepsilon} \quad (23)$$

and

$$\left\| B_i^k - (B_i')^k \right\|_2^2 \leq \sqrt{d} \sqrt{2\varepsilon} (1 + 4\sqrt{d} \sqrt{2\varepsilon}), \quad (24)$$

where $k = 0, \dots, d-1$ and B_i' are Bob's ideal observables, i.e., $B_i' = A_i^*$.

Proof. The proof is deferred to Appendix C. \square

Conclusions.—We propose a method for certification of a large class of d -outcome projective measurements, termed here genuinely incompatible, and the two-qudit maximally entangled state in the steering scenario. This is the first self-testing statement which applies to MUBs of any dimension; in particular it allows for robust self-testing of MUBs in Hilbert spaces of prime dimension. Moreover, unlike the previous approaches it does not require assuming that the underlying state is pure and measurements are projective but allows to deduce it from the observed violation. We further discuss the case when the measurements to be certified are not genuinely incompatible and find that depending on the state shared between the parties, only a part of the measurement could be characterised.

Some follow-up questions arise from our work. The

most important one would be whether the certification method presented here can be made fully device-independent, for instance, by using the approach of [36]. Another interesting avenue would be to find steering inequalities which could be used for self-testing of any non-maximally entangled state of arbitrary local dimension d with only two measurements on the untrusted side; a certification scheme for any pure state has recently been proposed [21], but it is a direct adaptation of the ‘projection method’ of [23] to the steering scenario.

I. ACKNOWLEDGEMENTS

This work is supported by the Polish National Science Center through the SONATA BIS project No. 2019/34/E/ST2/00369.

-
- [1] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, *Rev. Mod. Phys.* **86**, 419 (2014).
- [2] J. S. Bell, *Physics Physique Fizika* **1**, 195 (1964).
- [3] J. S. Bell, *Rev. Mod. Phys.* **38**, 447 (1966).
- [4] A. Acín, N. Brunner, N. Gisin, S. Massar, S. Pironio, and V. Scarani, *Phys. Rev. Lett.* **98**, 230501 (2007).
- [5] T. H. Yang and M. Navascués, *Phys. Rev. A* **87**, 050102 (2013).
- [6] R. Colbeck and R. Renner, *Nat. Comm.* **2**, 411 (2011).
- [7] S. Pironio, A. Acín, S. Massar, A. B. de la Giroday, D. N. Matsukevich, P. Maunz, S. Olmschenk, D. Hayes, L. Luo, T. A. Manning, and C. Monroe, *Nature* **464**, 1021 (2010).
- [8] D. Mayers and A. Yao, *Quantum Inf. Comput.* **4**, 273 (2004).
- [9] M. McKague, T. H. Yang, and V. Scarani, *J. Phys. A: Math. Theor.* **45**, 455304 (2012).
- [10] C. Bamps and S. Pironio, *Phys. Rev. A* **91**, 052111 (2015).
- [11] Y. Wang, X. Wu, and V. Scarani, *New J. Phys.* **18**, 025021 (2016).
- [12] I. Šupić, R. Augusiak, A. Salavrakos, and A. Acín, *New J. Phys.* **18**, 035013 (2016).
- [13] S. Sarkar, D. Saha, J. Kaniewski, and R. Augusiak, *npj Quantum Information* **7**, 151 (2021).
- [14] L. Mančinska, J. Prakash, and C. Schafhauser, “Constant-sized robust self-tests for states and measurements of unbounded dimension,” (2021), arXiv:2103.01729 [quant-ph].
- [15] H. M. Wiseman, S. J. Jones, and A. C. Doherty, *Phys. Rev. Lett.* **98**, 140402 (2007).
- [16] E. G. Cavalcanti, S. J. Jones, H. M. Wiseman, and M. D. Reid, *Phys. Rev. A* **80**, 032112 (2009).
- [17] M. T. Quintino, T. Vértesi, D. Cavalcanti, R. Augusiak, M. Demianowicz, A. Acín, and N. Brunner, *Phys. Rev. A* **92**, 032107 (2015).
- [18] I. Šupić and M. J. Hoban, *New J. Phys.* **18**, 075006 (2016).
- [19] A. Gheorghiu, P. Wallden, and E. Kashefi, *New J. Phys.* **19**, 023043 (2017).
- [20] S. Goswami, B. Bhattacharya, D. Das, S. Sasmal, C. Jebaratnam, and A. S. Majumdar, *Phys. Rev. A* **98**, 022311 (2018).
- [21] H. Shrotriya, K. Bharti, and L.-C. Kwek, *Phys. Rev. Research* **3**, 033093 (2021).
- [22] S.-L. Chen, C. Budroni, Y.-C. Liang, and Y.-N. Chen, *Phys. Rev. Lett.* **116**, 240401 (2016).
- [23] A. Coladangelo, K. T. Goh, and V. Scarani, *Nature Communications* **8** (2017).
- [24] F. Hirsch, M. T. Quintino, and N. Brunner, *Phys. Rev. A* **97**, 012129 (2018).
- [25] R. Uola, C. Budroni, O. Gühne, and J.-P. Pellonpää, *Phys. Rev. Lett.* **115**, 230402 (2015).
- [26] M. T. Quintino, T. Vértesi, and N. Brunner, *Phys. Rev. Lett.* **113**, 160402 (2014).
- [27] D. Cavalcanti and P. Skrzypczyk, *Phys. Rev. A* **93**, 052112 (2016).
- [28] M. Marciniak, A. Rutkowski, Z. Yin, M. Horodecki, and R. Horodecki, *Phys. Rev. Lett.* **115**, 170401 (2015).
- [29] P. Skrzypczyk and D. Cavalcanti, *Phys. Rev. A* **92**, 022354 (2015).
- [30] W. Son, J. Lee, and M. S. Kim, *Phys. Rev. Lett.* **96**, 060406 (2006).
- [31] Y.-C. Liang, C.-W. Lim, and D.-L. Deng, *Phys. Rev. A* **80**, 052116 (2009).
- [32] I. Šupić and J. Bowles, *Quantum* **4**, 337 (2020).
- [33] D. J. Saunders, S. J. Jones, H. M. Wiseman, and G. J. Pryde, *Nat. Phys.* **6**, 845 (2010).
- [34] In other words, we assume that the dimensions of Bobs measurement operators are the same as those of his reduced state ρ_B . Such an assumption is fully justified because we are able to make nontrivial statements on those parts of Bob’s measurement that act on $\text{supp}(\rho_B)$; what is beyond that support does not give rise to the observed nonclassical correlations.
- [35] S.-L. Chen, H.-Y. Ku, W. Zhou, J. Tura, and Y.-N. Chen, *Quantum* **5**, 552 (2021).
- [36] J. Bowles, I. Šupić, D. Cavalcanti, and A. Acín, *Phys. Rev. Lett.* **121**, 180503 (2018).

Appendix A: Correlation form of steering functionals

Let us consider the standard form of the steering functional, i.e.,

$$\sum_{b=0}^{d-1} \sum_{y=1}^N \text{Tr} \left(F_{b|y} \sigma_{b|y} \right), \quad (\text{A1})$$

where $\sigma_{b|y}$ form the assemblage and $F_{b|y}$ are some positive semi-definite operators acting on \mathbb{C}^d . Let us then consider a particular class of such functionals in which for every $y = 1, \dots, N$, the operators $F_{a|x}$ sum up to the identity, that is,

$$\sum_a F_{a|x} = \mathbb{1}, \quad (\text{A2})$$

and therefore form a quantum measurement. With the aid of Fourier transform we can represent each such a measurement by another set of d operators

$$A_{k|x} = \sum_{a=0}^{d-1} \omega^{ak} F_{a|x} \quad (\text{A3})$$

with $k = 0, \dots, d-1$ and $\omega = \exp(2\pi i/d)$, which for the purposes of this article we simply call generalized observables; in fact if $F_{a|x}$ forms a projective measurement, then $A_{k|x}$ are unitary observables with outcomes labelled by the powers of ω (see also below).

The inverse transform gives

$$F_{a|x} = \frac{1}{d} \sum_{k=0}^{d-1} \omega^{-ak} A_{k|x}. \quad (\text{A4})$$

Clearly, we can use the same approach to represent Bob's measurements $N_y = \{N_{b|y}\}_b$ in terms of generalized observables, that is,

$$B_{k|y} = \sum_{b=0}^{d-1} \omega^{bk} N_{b|y}, \quad (\text{A5})$$

and

$$N_{b|y} = \frac{1}{d} \sum_{k=0}^{d-1} \omega^{-kb} B_{k|y}. \quad (\text{A6})$$

It should also be noticed that the $B_{k|y}$ operators enjoy certain useful properties, that is, for any k and y :

$$B_{k|y}^\dagger = B_{d-k|y} = B_{-k|y} \quad (\text{A7})$$

and

$$B_{k|y}^\dagger B_{k|y} \leq \mathbb{1}_B, \quad (\text{A8})$$

where $\mathbb{1}_B$ is the identity acting on \mathcal{H}_B , and finally $B_{0|y} = \mathbb{1}$ for any y ; clearly, the same identities are satisfied by $A_{k|x}$. Importantly, the d operators $B_{k|y}$ fully determine Bob's measurements N_y . Slightly abusing the notation in what follows we denote $B_y = \{B_{k|y}\}_k$.

Let us now go back to the steering functionals (A1). By exploiting the fact that $\sigma_{b|y} = \text{Tr}_B[(\mathbb{1}_A \otimes N_{b|y})\rho_{AB}]$, it can be rewritten as a combination of expectation values of the above generalized observables,

$$\sum_{b=0}^{d-1} \sum_{y=1}^N \text{Tr} \left(F_{b|y} \sigma_{b|y} \right) = \frac{1}{d} \sum_{k=0}^{d-1} \sum_{y=1}^N \left\langle A_{k|y} \otimes B_{-k|y} \right\rangle. \quad (\text{A9})$$

We then further restrict the class of the steering functionals to the case when the operators $F_{a|x}$ form projective measurements for all x , meaning that they satisfy

$$F_{b|y}F_{b'|y} = \delta_{b,b'}F_{b|y}. \quad (\text{A10})$$

It is not difficult to see that under the above assumption, $A_{k|y}$ are all unitary operators with eigenvalues ω^{bk} ($b = 0, \dots, d-1$), and, moreover, $A_{k|y}$ for any y are simply the k th powers of $A_{1|y}$. Using this fact, denoting

$$A_{1|y} \equiv A_y, \quad (\text{A11})$$

dropping the $1/d$, and replacing $-k$ by k which can be done with no loss of generality as such a manipulation represents a simple relabelling the outcomes of Bob's measurements, we finally arrive at our steering functionals of the form

$$\langle \mathcal{B}_d \rangle = \sum_{k=1}^{d-1} \sum_{y=1}^N \langle A_y^k \otimes B_{k|y} \rangle, \quad (\text{A12})$$

where we also got rid of the term for $k = 0$ because the expectation value for this term is simply one. It is worth recalling here that while the dimension of Alice's observables is known to be d , Bob's measurements act on a Hilbert space of in principle unknown but finite dimension.

Let us make two comments here. The first one is that if Bob's measurements are projective too, the $B_{k|y}$ operators are also unitary for all k and y , and therefore $B_{k|y}$ in Eq. (A12) can be replaced by B_y^k .

Second, it should be stressed here that the form (A12) of the steering functional is particularly useful as far as self-testing is concerned because it is straightforward to find its maximal quantum value which simply equals its algebraic bound, i.e., $\beta_Q = N(d-1)$, as well as a quantum realization achieving this value. Precisely, such a realization consists of the maximally entangled state of two qudits

$$|\phi_+^d\rangle = \frac{1}{\sqrt{d}} \sum_i |ii\rangle \quad (\text{A13})$$

and Bob's measurement satisfying $B_i = A_i^*$ for any i .

On the other hand, any quantum realization achieving β_Q of our inequality for a given set of observables A_i , consisting of a mixed state ρ_{AB} and Bob's measurements B_i , must satisfy the following relations

$$A_y^k \otimes B_{k|y} \rho_{AB} = \rho_{AB} \quad (\text{A14})$$

for any y and k . This is because for maximal violation every term in Eq. (A12) satisfies

$$\text{Tr}[(A_y^k \otimes B_{k|y})\rho_{AB}] = 1, \quad (\text{A15})$$

which due to the facts that A_y are unitary and that $B_{k|y}$ satisfy (A8), directly implies (A14).

Appendix B: Bounding the maximal classical value of our functionals

In this section we derive an upper bound on the maximal classical value of the steering expression (7), and show that for any set of genuinely d -outcome incompatible measurements the corresponding steering inequality is non-trivial, that is, $\beta_L < \beta_Q$.

Let us begin by demonstrating how to derive the bound (7). To this end, we assume that the assemblage $\{\sigma_{b|y}\}$ admits a local hidden state model, that is,

$$\sigma_{b|y} = \sum_{\lambda} p(\lambda) p(b|y, \lambda) \rho_{\lambda}, \quad (\text{B1})$$

where λ is some random variable with some probability distribution $p(\lambda)$, $p(b|y, \lambda)$ a probability distributions according to which Bob returns the outcomes b given the measurement choice y and the hidden variable λ , and ρ_{λ} are some states acting on Alice's Hilbert space. In terms of the joint probability distributions such an LHS model expresses as

$$p(a, b|x, y) = \sum_{\lambda} p(\lambda) p(a|x, \rho_{\lambda}) p(b|y, \lambda), \quad (\text{B2})$$

where $p(a|x, \rho_\lambda)$ is the probability of obtaining the outcome a when measuring the observable A_x on the state ρ_λ . Then, in terms of the expectation values appearing in (A9), the above rewrites as

$$\langle A_{k|x} \otimes B_{l|y} \rangle_{\text{LHS}} = \sum_{\lambda} p(\lambda) \langle A_{k|x} \rangle_{\rho_\lambda} \langle B_{l|y} \rangle_{\lambda}, \quad (\text{B3})$$

where $\langle A_{k|x} \rangle_{\rho_\lambda} = \text{Tr}[A_{k|x} \rho_\lambda]$ with $A_{k|x}$ being operators representing the x th measurement performed by Alice [cf. Eq. (A3)]. Then, $\langle B_{l|y} \rangle_{\lambda}$ stands for the Fourier transform of $p(b|y, \lambda)$, i.e.,

$$\langle B_{l|y} \rangle_{\lambda} = \sum_b \omega^{bl} p(b|y, \lambda). \quad (\text{B4})$$

Now, our steering expression (A12) for LHS models reads

$$\langle \hat{\mathcal{B}}_d \rangle_{\text{LHS}} = \sum_{\lambda} p(\lambda) \sum_{y=1}^N \sum_{k=1}^{d-1} \langle A_{k|y} \rangle_{\rho_\lambda} \langle B_{-k|y} \rangle_{\lambda}, \quad (\text{B5})$$

and by noting that $|\langle B_{l|y} \rangle_{\lambda}| \leq 1$ for any choice of y, l and λ , its absolute value can be upper bounded as

$$|\langle \hat{\mathcal{B}}_d \rangle_{\text{LHS}}| \leq \sum_{\lambda} p(\lambda) \sum_{i=1}^N \sum_{k=1}^{d-1} |\langle A_i^k \rangle_{\rho_\lambda}|, \quad (\text{B6})$$

and further as

$$|\langle \hat{\mathcal{B}}_d \rangle_{\text{LHS}}| \leq \max_{\rho} \sum_{i=1}^N \sum_{k=1}^{d-1} |\langle A_i^k \rangle_{\rho}|, \quad (\text{B7})$$

where we took advantage of the fact that the right-hand side of the above inequality is a convex combination and that by optimizing it over states ρ acting on Alice's Hilbert space we cannot decrease its value. We thus obtain the bound (7).

A direct, but important for our considerations, corollary that stems from the above bound is that for LHS models with any set of genuinely incompatible measurements A_i , the right hand side of (A12) is smaller than $N(d-1)$, and thus $\beta_L < N(d-1)$, meaning that for any such a set our steering inequality is nontrivial. To see this explicitly let us assume that

$$\sum_{i=1}^N \sum_{k=0}^{d-1} |\langle A_i^k \rangle_{\rho}| = N(d-1), \quad (\text{B8})$$

which implies that $|\langle A_i^k \rangle_{\rho}| = 1$ for any i, k and some $|\psi\rangle \in \mathbb{C}^d$. Due to the fact that A_i^k are unitary operators, this means that $|\psi\rangle$ is their eigenvector, possibly with different eigenvalues. Consequently, these observables share a common subspace spanned by this vectors which contradicts that they are genuinely incompatible.

Appendix C: Self-testing

Here we present proofs of the lemmas and theorems stated in the main text.

Proof of Lemma 1

Lemma 1. Consider a set of N d -outcome unitary observables A_i obeying $A_i^d = \mathbb{1}_d$ that are genuinely incompatible. If for some nonzero normal matrix P acting on \mathbb{C}^d , $[P, A_i] = 0$ for every $i = 1, \dots, N$, then $P = \mu \mathbb{1}$ with $\mathbb{1}$ being the identity acting on \mathbb{C}^d and $\mu \in \mathbb{C}$.

Proof. The proof is by contradiction. Let us assume that the matrix P is not proportional to the identity. Since it is normal, we can decompose it as

$$P = \sum_i \lambda_i P_i, \quad (\text{C1})$$

where $\lambda_i \in \mathbb{C}$ are the eigenvalues of P which without any loss of generality we can assume to be all different, and P_i are the corresponding orthogonal eigenprojections. Here we do not exclude the possibility that one of the eigenvalues is zero; below we show that this is in fact impossible and that $\text{rank}(P) = d$.

Now, it is a known fact that if $[P, A_i] = 0$, then A_i is of block form

$$A_i = A_i^{(1)} \oplus \dots \oplus A_i^{(m)}, \quad (\text{C2})$$

where each block $A_i^{(j)}$ acts on $\text{supp}(P_j)$ and m is the number of different eigenvalues of P . For completeness, let us add here a short proof of this fact. To this aim, we rewrite $[P, A_i] = 0$ as

$$A_i P = P A_i. \quad (\text{C3})$$

We then sandwich this equation with P_m and P_n for $m \neq n$, which gives

$$P_m A_i P P_n = P_m P A_i P_n. \quad (\text{C4})$$

Noting that $P P_n = \lambda_n P_n$ and $P_m P = \lambda_m P_m$, we can rewrite the above as

$$(\lambda_n - \lambda_m) P_m A_i P_n = 0, \quad (\text{C5})$$

and since $\lambda_n \neq \lambda_m$ for $n \neq m$, we conclude that $P_m A_i P_n = 0$ for any pair $m \neq n$. Of course, this conclusion can be drawn for each A_i , and thus we obtain (C2).

The fact that all observables can be written as in (C2) implies that $\text{supp}(P_i)$ is an invariant subspace of every A_i of dimension lower than d , which contradicts the assumption. As a consequence, $\text{rank}(P) = d$. This completes the proof. \square

Exact self-testing

Let us now provide a full proof of our main result. We split it into two theorems. The first one is for pure states and projective measurements on Bob's side. The second theorem extends it to the case of mixed states and nonprojective measurements on Bob's side. We stress here that in our considerations we can assume that the local state ρ_B is full rank, or, in other words, that the Bob's measurements operators act on a Hilbert space of the same dimension as \mathcal{H}_B . Let us also add that in what follows all the considered observables are unitary and their eigenvalues are ω^i with $i = 0, \dots, d-1$.

Theorem 1. *Assume that the steering inequality (5) is maximally violated by a state $|\psi\rangle_{AB} \in \mathbb{C}^d \otimes \mathcal{H}_B$ and unitary d -outcome observables A_i, B_i ($i \in \{1, 2, \dots, N\}$) acting on, respectively, \mathbb{C}^d and \mathcal{H}_B such that the observables A_i on the trusted side are d -outcome genuinely incompatible. Then, the following statement holds true for any d : there exists a local unitary transformation on Bob side, $U_B : \mathcal{H}_B \rightarrow \mathcal{H}_B$, such that*

$$(\mathbb{1}_A \otimes U_B) |\psi\rangle_{AB} = |\phi_d^\dagger\rangle \quad (\text{C6})$$

and

$$\forall i, \quad U_B \bar{B}_i U_B^\dagger = A_i^*, \quad (\text{C7})$$

where \bar{B}_i is B_i projected onto the support of Bob's state ρ_B acting on \mathbb{C}^d .

Proof. The optimal quantum value $\beta_Q = N(d-1)$ of the steering inequality (5) is the same as its algebraic value. Thus, we have the following set of relations [cf. Eq. (A14)]

$$A_i^k \otimes B_i^k |\psi\rangle_{AB} = |\psi\rangle_{AB} \quad (\text{C8})$$

for all i and k , where due to the fact that Bob's measurements are projective, $B_{k|y} = B_y^k$. For clarity the rest of the proof is divided into three main steps.

Step 1. Let us first use these relations to prove that the fact that for A_i forming a set of genuinely incompatible measurements implies that $\text{rank}(\rho_A) = d$ with ρ_A being the reduced density matrix corresponding to Alice,

$\rho_A = \text{Tr}_B(|\psi\rangle\langle\psi|_{AB})$; in other words, any state maximally violating our inequality for genuinely incompatible measurements must be of maximal local rank. To this end, we assume that $\text{rank}(\rho_A) < d$ and consider (A14) for $k = 1$ when B_i is projective and ρ_{AB} is pure, which implies that

$$\Pi_A A_i \Pi_A \otimes B_i |\psi\rangle_{AB} = |\psi\rangle_{AB}, \quad (\text{C9})$$

where Π_A is the projector onto the support of ρ_A . Denoting then $\bar{A}_i \equiv \Pi_A A_i \Pi_A$ and owing to the fact that B_i are unitary and satisfy $B_i^d = \mathbb{1}$, we conclude that \bar{A}_i are unitary matrices on $\text{supp}(\rho_A)$ and satisfy $\bar{A}_i^d = \Pi_A$. It is then not difficult to see that since A_i are unitary, they must be of block form

$$A_i = \bar{A}_i \oplus A'_i \quad (\text{C10})$$

for some square unitary matrix A'_i of dimension $[d - \text{rank}(\rho_A)] \times [d - \text{rank}(\rho_A)]$. However, this implies that all the observables A_i have a common invariant subspace of dimension lower d , contradicting the assumption that they are genuinely incompatible on \mathbb{C}^d . As a consequence, $|\psi\rangle_{AB}$ is of full local rank, or equivalently $\text{rank}(\rho_A) = d$ which also means that $\text{rank}(\rho_B) = d$, where $\rho_B = \text{Tr}_A(|\psi\rangle\langle\psi|_{AB})$.

Step 2. Using a similar reasoning we now show that Bob's observables all split into a direct sum

$$B_i = \bar{B}_i \oplus E_i, \quad (\text{C11})$$

where $\bar{B}_i = \Pi_B B_i \Pi_B$ are $d \times d$ matrices acting on the support of ρ_B , whereas E_i are of unknown dimension and are defined outside $\text{supp}(\rho_B)$; in particular, $E_i |\psi\rangle_{AB} = 0$. Both \bar{B}_i and E_i are unitary and their spectra belong to $\{1, \dots, \omega^{d-1}\}$.

To prove the above statement, let us first notice that each B_i can be written in a block form,

$$B_i = \begin{pmatrix} \bar{B}_i & C_i \\ D_i & E_i \end{pmatrix}, \quad (\text{C12})$$

where, as before, $\bar{B}_i = \Pi_B B_i \Pi_B$ and $E_i = \Pi_B^\perp B_i \Pi_B^\perp$ where Π_B and Π_B^\perp are the projectors onto the support of ρ_B and its complement in \mathcal{H}_B , respectively. Let us then go back to Eq. (A14) for $k = 1$ and project it onto the support of ρ_B , that is,

$$A_i \otimes \Pi_B B_i \Pi_B |\psi\rangle_{AB} \equiv A_i \otimes \bar{B}_i |\psi\rangle_{AB} = |\psi\rangle_{AB}. \quad (\text{C13})$$

By applying $A_i^\dagger \otimes \bar{B}_i^\dagger$ to this identity and using the fact that A_i are unitary, we conclude that $\bar{B}_i^\dagger \bar{B}_i = \mathbb{1}_d$. In a fully analogous way we obtain from Eq. (A14) for $k = d - 1$ that $\bar{B}_i \bar{B}_i^\dagger = \mathbb{1}_d$ and thus all \bar{B}_i are unitary. On the other hand, by applying $A_i \otimes \bar{B}_i$ to Eq. (C13) $d - 1$ times, we obtain that $\bar{B}_i^d = \mathbb{1}_d$ and thus their spectra belong to $\{1, \dots, \omega^{d-1}\}$.

It is then direct to see that since both B_i and \bar{B}_i are unitary on the respective Hilbert spaces, the off-diagonal blocks in Eq. (C13) must vanish, $C_i = D_i = 0$, leading to the desired block form (C11). This finally implies that E_i must also be unitary and their eigenvalues are powers of ω .

Step 3. Let us then consider the Schmidt decomposition of $|\psi_{AB}\rangle$,

$$|\psi\rangle_{AB} = \sum_{i=0}^{d-1} \lambda_i |e_i\rangle |f_i\rangle, \quad (\text{C14})$$

where due to the fact that $\text{rank}(\rho_A) = d$, the Schmidt coefficient λ_i ($i = 0, \dots, d - 1$) are all positive and satisfy $\sum_i \lambda_i^2 = 1$. Moreover, the local vectors $|e_i\rangle$ and $|f_i\rangle$ form orthonormal bases in \mathbb{C}^d and \mathcal{H}_B , respectively.

We can now make a few manipulations with (C14). First, we observe that there exists a unitary U_B such that $|f_i\rangle = U_B |e_i^*\rangle$ for every i , where the asterisk denotes complex conjugation in the standard basis; thus

$$|\psi\rangle_{AB} = (P_A \otimes U_B) \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |e_i\rangle |e_i^*\rangle, \quad (\text{C15})$$

where P_A is a matrix diagonal in the $\{|e_i\rangle\}$ basis with eigenvalues $\sqrt{d} \lambda_i$; recall the fact that A_i are GI implies that P_A is full rank, $\text{rank}(P_A) = d$. We finally observe that the state appearing on the right hand side of (C15) is the maximally entangled state (A13); consequently

$$|\psi\rangle_{AB} = (P_A \otimes U_B) |\phi_d^+\rangle. \quad (\text{C16})$$

Now, combining Eqs. (C16) and (C8) for $k = 1$, we obtain

$$(A_i P_A \otimes \tilde{B}_i) |\phi_d^+\rangle = (P_A \otimes \mathbb{1}_B) |\phi_d^+\rangle, \quad (\text{C17})$$

where $\tilde{B}_i = U_B \bar{B}_i U_B^\dagger$. Using then the fact that $R \otimes Q |\phi_d^+\rangle = RQ^T \otimes \mathbb{1} |\phi_d^+\rangle$ holds for any two $d \times d$ matrices R and Q , and omitting the state $|\phi_d^+\rangle$, we obtain a set of matrix equation

$$A_i P_A \tilde{B}_i^T = P_A. \quad (\text{C18})$$

Let us then consider the Hermitian conjugation of the above equation

$$\tilde{B}_i^* P_A A_i^\dagger = P_A, \quad (\text{C19})$$

where we used the fact that P_A is self-adjoint. After multiplying Eq. (C18) by Eq. (C19) and using the fact that \tilde{B}_i are unitary, which implies that $\tilde{B}_i^T \tilde{B}_i^* = (\tilde{B}_i^\dagger \tilde{B}_i)^* = \mathbb{1}_d$, we arrive at

$$A_i P_A^2 A_i^\dagger = P_A^2. \quad (\text{C20})$$

Since A_i are unitary, the above implies a simple condition for the matrix P_A , $[A_i, P_A^2] = 0$, which due to the fact that $P_A \geq 0$ is equivalent to

$$[A_i, P_A] = 0 \quad (\text{C21})$$

for all i . Owing to the facts that $P_A \geq 0$ and that A_i are genuinely incompatible, the above lemma tell us that P_A is proportional to identity, proving Eq. (C6). Then, plugging this particular form of P_A into Eq. (C18) we obtain (C7), which completes the proof. \square

Let us now extend our self-testing statement to the case when the state shared by Alice and Bob is mixed which might for instance represent a cryptographic scenario with a malicious eavesdropper who keeps the purification $|\psi_{ABE}\rangle$ of ρ_{AB} . Moreover, we do not assume Bob's measurements to be projective.

Theorem 2. Assume that for a given set of GI unitary observables A_i obeying $A_i^d = \mathbb{1}_d$, the steering inequality (5) is maximally violated by a state ρ_{AB} acting on $\mathbb{C}^d \otimes \mathcal{H}_B$ and measurements B_i . Then, the following statements hold true for any d : (i) Bob's measurements are projective, that is, all operators $B_{k|y}$ are unitary such that $B_{k|y}^d = \mathbb{1}$, (ii) Bob's Hilbert space decomposes as $\mathcal{H}_B = (\mathbb{C}^d)_{B'} \otimes \mathcal{H}_{B''}$, and (iii) there exists a local unitary transformation on Bob side, $U_B : \mathcal{H}_B \rightarrow \mathcal{H}_B$, such that

$$(\mathbb{1}_A \otimes U_B) \rho_{AB} (\mathbb{1}_A \otimes U_B^\dagger) = |\phi_d^+\rangle \langle \phi_d^+|_{AB'} \otimes \rho_{B''}. \quad (\text{C22})$$

and

$$\forall i, \quad U_B B_i U_B^\dagger = A_i^* \otimes \mathbb{1}_{B''}, \quad (\text{C23})$$

where B'' denotes Bob's auxiliary system.

Proof. The departure point of the proof is the fact, already mentioned in Appendix A, that if for a given set of observables A_i , our steering inequality is maximally violated by ρ_{AB} and $B_{k|y}$, then

$$(A_i^k \otimes B_{k|i}) \rho_{AB} = \rho_{AB} \quad (\text{C24})$$

for every i and k .

Step 1. We begin by showing that Bob's measurements maximally violating our inequality are projective. Applying $A_i^{-k} \otimes B_{-k|i}$ to the above and exploiting the fact that $A_i^{-k} = (A_i^k)^\dagger$ and $B_{-k|i} = B_{k|i}^\dagger$, and using the fact that A_i are unitary, we arrive at

$$(\mathbb{1}_A \otimes B_{k|i}^\dagger B_{k|i}) \rho_{AB} = \rho_{AB}, \quad (\text{C25})$$

which directly implies that

$$B_{k|i}^\dagger B_{k|i} = \mathbb{1}_B, \quad (\text{C26})$$

where $\mathbb{1}_B$ is the identity on \mathcal{H}_B . This implies that $B_{k|i}$ are unitary for any choice of i and k , which through Eq. (A6) means that all $N_{b|y}$ are orthogonal projections, and thus Bob's measurements are projective. In what follows we substitute $B_{k|i} = B_i^k$.

Step 2. Let us now move on to proving the main result and consider a decomposition of ρ_{AB} maximally violating our inequality into pure states

$$\rho_{AB} = \sum_{s=1}^K p_s |\psi_s\rangle\langle\psi_s|_{AB}, \quad (\text{C27})$$

where $p_s \geq 0$ are such that $\sum_s p_s = 1$ and K is some positive integer. Clearly, we can assume that all $|\psi_s\rangle$ are linearly independent. In fact, with no loss of generality we can assume that the above is the eigendecomposition of ρ_{AB} , meaning that $|\psi_s\rangle$ are pairwise orthogonal, i.e., $\langle\psi_s|\psi_{s'}\rangle = \delta_{ss'}$.

Now, the fact that ρ_{AB} achieves the maximal quantum value of our inequality implies that each pure state in the decomposition (C27) achieves it too, and therefore any $|\psi_s\rangle_{AB}$ satisfies the following conditions

$$(A_i^k \otimes B_i^k) |\psi_s\rangle_{AB} = |\psi_s\rangle_{AB} \quad (\text{C28})$$

for any i and k . It follows from Theorem 1 that for any state satisfying such equations for genuinely incompatible observables A_i there exists a unitary U_B^s such that

$$(\mathbb{1} \otimes U_B^s) |\psi_s\rangle_{AB} = |\phi_+^d\rangle. \quad (\text{C29})$$

It is worth stressing here that the unitary operations U_B^s might be different for different $|\psi_s\rangle_{AB}$.

For further benefits we rewrite the above as

$$|\psi_s\rangle_{AB} = [\mathbb{1} \otimes (U_B^s)^\dagger] |\phi_+^d\rangle = \frac{1}{\sqrt{d}} \sum_i |i\rangle |f_i^s\rangle, \quad (\text{C30})$$

where the vectors $|f_i^s\rangle = (U_B^s)^\dagger |i\rangle$ form an orthonormal basis for any s . Moreover,

$$U_B^s \overline{B}_i^{(s)} (U_B^s)^\dagger = A_i^*, \quad (\text{C31})$$

where $\overline{B}_i^{(s)}$ denote Bob's observables B_i projected onto the local supports of $|\psi_s\rangle_{AB}$, that is,

$$\overline{B}_i^{(s)} = \Pi_B^s B_i \Pi_B^s, \quad (\text{C32})$$

where Π_B^s denotes the projector onto the support of $\rho_B^s = \text{Tr}_A[|\psi_s\rangle\langle\psi_s|_{AB}]$,

$$\text{supp}(\rho_B^s) \equiv V_s = \text{span}\{|f_0^s\rangle, \dots, |f_{d-1}^s\rangle\} \subset \mathcal{H}_B. \quad (\text{C33})$$

It is important to notice that as shown in the proof of Theorem 1, the $d \times d$ matrices $\overline{B}_i^{(s)}$ are unitary and their spectrum belongs to the set $\{1, \omega, \dots, \omega^{d-1}\}$. Moreover, from each such subspace Bob's observables B_i decompose as (see the proof of Theorem 1)

$$B_i = \overline{B}_i^{(s)} \oplus E_i^{(s)}, \quad (\text{C34})$$

where $E_i^{(s)}$ is a unitary operator too that acts on the complement of V_s in \mathcal{H}_B .

Our aim now is to prove that all the local subspaces V_s corresponding to the pure states $|\psi_s\rangle_{AB}$ are mutually orthogonal. To this end, let us focus on the first two vectors $|\psi_1\rangle_{AB}$ and $|\psi_2\rangle_{AB}$ from the decomposition (C27) and the corresponding subspaces V_1 and V_2 . It will be very convenient for us to represent the maximally entangled state $|\phi_+^d\rangle$ in Eq. (C29) in the eigenbasis of one of Alice's observables, say A_0 . Equivalently, due to the facts that $|\phi_+^d\rangle$ is invariant under the action of $U \otimes U^*$ for any unitary U and that U can be included on Bob's observables, without any loss of generality we can simply assume that the computational basis $\{|i\rangle\}$ is an eigenbasis of A_0 , that is,

$$A_0 |i\rangle = \omega^i |i\rangle \quad (i = 0, \dots, d-1). \quad (\text{C35})$$

By applying this fact to Eq. (C28) we see that both local bases $\{|f_i^1\rangle\}$ and $\{|f_i^2\rangle\}$ are in fact the eigenbases of B_0 , that is

$$B_0|f_i^s\rangle = \omega^{-i}|f_i^s\rangle \quad (s = 1, 2). \quad (\text{C36})$$

This imposes certain orthogonality relations between vectors from both bases. Namely, due to the fact that B_0 is unitary, vectors that correspond to its different eigenvalues must be orthogonal, that is,

$$\langle f_i^1 | f_j^2 \rangle = 0 \quad (i \neq j). \quad (\text{C37})$$

Thus, to prove that $V_1 \perp V_2$ it's enough to show that the above orthogonality holds true also when $s = t$. In order to do so, we can always decompose

$$|f_i^2\rangle = \alpha_i|f_i^1\rangle + \beta_i|g_i\rangle, \quad (\text{C38})$$

where $\alpha_i, \beta_i \in \mathbb{C}$ and $|\alpha_i|^2 + |\beta_i|^2 = 1$ and $|g_i\rangle$ is a normalized vector such that $\langle f_i^1 | g_i \rangle = 0$ for any i . Using then the orthogonality relations (C37) and the fact that $|f_i^1\rangle$ form an orthonormal set, it is not difficult to see from Eq. (C38) that

$$\beta_i \langle f_j^1 | g_i \rangle = 0 \quad (i, j = 0, \dots, d-1). \quad (\text{C39})$$

This means that either $\beta_i = 0$ or the vectors $|g_i\rangle$ for which $\beta_i \neq 0$ are orthogonal to the whole subspace V_1 .

Let us now go back to the conditions (C28) which for $k = 1$ can be rewritten as

$$(\mathbb{1}_A \otimes B_i)|\psi_s\rangle_{AB} = (A_i^\dagger \otimes \mathbb{1}_B)|\psi_s\rangle_{AB}. \quad (\text{C40})$$

Using then the Schmidt decompositions of $|\psi_s\rangle_{AB}$ given in Eq. (C30), the above gives

$$\sum_{k=0}^{d-1} |k\rangle \otimes (B_i|f_k^s\rangle) = \sum_{k=0}^{d-1} (A_i^\dagger|k\rangle) \otimes |f_k^s\rangle, \quad (\text{C41})$$

which directly implies a set of vector equations that for the two considered vectors $|\psi_1\rangle_{AB}$ and $|\psi_2\rangle_{AB}$ we state explicitly as

$$B_i|f_k^1\rangle = \sum_{m=0}^{d-1} \langle k | A_i^\dagger | m \rangle |f_m^1\rangle \quad (\text{C42})$$

and

$$B_i|f_k^2\rangle = \sum_{m=0}^{d-1} \langle k | A_i^\dagger | m \rangle |f_m^2\rangle, \quad (\text{C43})$$

where $k = 0, \dots, d-1$. We then use the decomposition (C38) in (C43), which leads us to

$$\alpha_k B_i|f_k^1\rangle + \beta_k B_i|g_k\rangle = \sum_{m=0}^{d-1} \alpha_m \langle k | A_i^\dagger | m \rangle |f_m^1\rangle + \sum_{m=0}^{d-1} \beta_m \langle k | A_i^\dagger | m \rangle |g_m\rangle. \quad (\text{C44})$$

Using then Eq. (C42) for $B_i|f_i^1\rangle$ and after some simple manipulations, the above gives

$$\sum_{m=0}^{d-1} (\alpha_k - \alpha_m) \langle k | A_i^\dagger | m \rangle |f_m^1\rangle = \sum_{m=0}^{d-1} \beta_m \langle k | A_i^\dagger | m \rangle |g_m\rangle - \beta_k B_i|g_k\rangle. \quad (\text{C45})$$

If we sandwich the above equation with $\langle f_n^1 |$ we obtain

$$(\alpha_k - \alpha_n) \langle k | A_i^\dagger | n \rangle = -\beta_k \langle f_n^1 | B_i | g_k \rangle, \quad (\text{C46})$$

where we used the fact that $|f_n^1\rangle \perp |g_i\rangle$ for any pair n, i (see above for explanation). Using finally Eq. (C39) together with the fact that B_i decompose into blocks given in Eq. (C34), meaning that they act invariantly on the subspace

spanned by $|f_n^1\rangle$, which can also be inferred from Eq. (C42), it is direct to see that the right-hand side of the above equations simply vanishes, and hence,

$$(\alpha_k - \alpha_n)\langle k|A_i^\dagger|n\rangle = 0 \quad (k, n = 0, \dots, d-1). \quad (\text{C47})$$

The key observation now is that the above system of equations is satisfied if, and only if the diagonal matrix $Q = \text{diag}[\alpha_0, \dots, \alpha_{d-1}]$ commutes with every A_i , that is,

$$[A_i^\dagger, Q] = 0 \quad (\text{C48})$$

for any i . However, as proven in Lemma 1 this is possible for genuinely incompatible observables A_i iff Q is proportional to the identity $\mathbb{1}_d$. This means that all α_i are equal; let us then denote $\alpha_i = \alpha$ for some $\alpha \in \mathbb{C}$.

The final step is to use the fact that the vectors $|\psi_1\rangle_{AB}$ and $|\psi_2\rangle_{AB}$ are orthogonal which together with Eq. (C37) gives

$$0 = \langle \psi_1 | \psi_2 \rangle = \frac{1}{d} \sum_k \langle f_k^1 | f_k^2 \rangle = \frac{1}{d} \sum_k \alpha_k = \alpha, \quad (\text{C49})$$

and so $\alpha_i = \alpha = 0$ for every i . If we plug this back to Eq. (C38) we see that $|f_i^2\rangle = \beta_i |g_i\rangle$ for some complex phase β_i . However, as already said $|g_i\rangle$ is orthogonal to $|f_i^1\rangle$ and hence $\langle f_i^2 | f_i^1 \rangle = 0$, which together with (C37) implies finally that the subspaces V_1 and V_2 are orthogonal. Clearly, we can apply the same argumentation to every pair of subspaces V_j and V_k , which allows us to conclude that they are all mutually orthogonal.

Step 3. We are now ready to prove Eqs. (9) and (C22). Indeed, the fact that the local supports V_s are mutually orthogonal implies that Bob's Hilbert space admits the following decomposition

$$\mathcal{H}_B = V_1 \oplus V_2 \oplus \dots \oplus V_K, \quad (\text{C50})$$

which given that $\dim V_s = d$, means in particular that $\mathcal{H}_B = (\mathbb{C}^d)_{B'} \otimes \mathcal{H}_{B''}$ for some Hilbert space $\mathcal{H}_{B''}$. Moreover, the fact that the vectors $|f_i^s\rangle$ span orthogonal subspaces for different s , implies that there exists a unitary operation $U_B : \mathcal{H}_B \rightarrow \mathcal{H}_B$ such that

$$U_B |f_i^s\rangle = |i\rangle_{B'} \otimes |s\rangle_{B''} \quad (\text{C51})$$

with $i = 0, \dots, d-1$ and $s = 1, \dots, K$. Thus,

$$(\mathbb{1}_A \otimes U_B) |\psi_s\rangle_{AB} = |\phi_+^d\rangle_{AB'} \otimes |s\rangle_{B''} \quad (\text{C52})$$

for every s , and therefore taking into account the decomposition (C27) we finally have

$$(\mathbb{1}_A \otimes U_B) \rho_{AB} (\mathbb{1}_A \otimes U_B^\dagger) = |\phi_+^d\rangle \langle \phi_+^d| \otimes \rho_{B''}, \quad (\text{C53})$$

where $\rho_{B''} = \sum_s p_s |s\rangle \langle s|_{B''}$. This is precisely Eq. (C22) we wanted to prove.

Having Eq. (C53) we can finally establish Eq. (C23). To this end, we notice that since $\mathcal{H}_B = (\mathbb{C}^d)_{B'} \otimes \mathcal{H}_{B''}$, we can $U_B B_i U_B^\dagger$ in the following block form

$$U_B B_i U_B^\dagger = \sum_{s,t=1}^K B_{s,t}^i \otimes |s\rangle \langle t|_{B''}, \quad (\text{C54})$$

where $B_{s,t}^i$ are $d \times d$ blocks acting on $(\mathbb{C}^d)_{B'}$, whereas $|s\rangle$ is the standard basis of $\mathcal{H}_{B''}$ and simultaneously the eigenbasis of $\rho_{B''}$. Plugging Eqs. (C53) and (C54) into Eq. (C24) for $k = 1$, we obtain the following matrix equation

$$\sum_{s,t} (A_i \otimes B_{s,t}^i) |\psi_+^d\rangle \langle \psi_+^d| \otimes p_t |s\rangle \langle t|_{B''} = |\psi_+^d\rangle \langle \psi_+^d| \otimes \sum_s p_s |s\rangle \langle s|_{B''}. \quad (\text{C55})$$

The off-diagonal terms on B'' subsystem imply $(A_i \otimes B_{s,t}^i) |\psi_+^d\rangle = 0$, which, given that A_i are unitary, directly give $B_{s,t}^i = 0$ for $s \neq t$. Then, the diagonal terms of (C55) lead us to

$$(A_i \otimes B_{s,s}^i) |\psi_+^d\rangle = |\psi_+^d\rangle, \quad (\text{C56})$$

which by virtue of $R \otimes Q|\phi_d^+\rangle = RQ^T \otimes \mathbb{1}|\phi_d^+\rangle$, which holds true for any two matrices R and Q , is equivalent to $B_{s,s}^i = A_i^*$. Substituting this back to Eq. (C54) one obtains

$$U_B B_i U_B^\dagger = \sum_{s=1}^K A_i^* \otimes |s\rangle\langle s|_{B''} = A_i^* \otimes \mathbb{1}_{B''}, \quad (\text{C57})$$

which is exactly Eq. (C23). This completes the proof. \square

Now, we move on to prove that any set of observables whose eigenvectors form mutually unbiased bases are genuinely incompatible.

Lemma 2. *Any two d -outcome observables whose eigenbases are mutually unbiased are genuinely incompatible.*

Proof. The proof is by contradiction. Let us begin by taking two d -dimensional mutually unbiased bases denoted by, $\{|s_i\rangle\}$ and $\{|t_j\rangle\}$ such that $|\langle s_i|t_j\rangle|^2 = 1/d$ for all $i, j \in \{0, 1, \dots, d-1\}$. Let us then construct observables from these bases,

$$A_1 = \sum_{i=0}^{d-1} \omega^{f(i)} |s_i\rangle\langle s_i|, \quad A_2 = \sum_{i=0}^{d-1} \omega^{g(i)} |t_i\rangle\langle t_i|, \quad (\text{C58})$$

where f and g are some permutations of the d -element set $\{0, \dots, d-1\}$, and assume that they share a common invariant subspace $V \subsetneq \mathbb{C}^d$ such that $\dim V < d$. This means that both can be written as $A_i = A'_i \oplus A''_i$, where A'_i acts on V , whereas A''_i on its complement in \mathbb{C}^d . Then, A_1 has eigenvectors that are orthogonal to eigenvectors of A_2 , which contradicts that the eigenvectors of A_1 and A_2 are mutually unbiased. \square

Exemplary set of genuinely incompatible observables which are not pairwise genuinely incompatible

For a note, we construct a set of genuinely incompatible measurements for $d = 4$ which are not pairwise genuinely incompatible. For this, we would require the underlying lemma.

Lemma 3. *Two invertible and diagonalizable matrices T and T' of dimension D , share a common non-trivial invariant subspace of dimension $d < D$ iff there exist d eigenvectors of T' which can be written as a linear combination of d eigenvectors of T .*

Proof. Let us first recall that any non-trivial invariant subspace of a matrix T is spanned by its eigenvectors which implies that if V_d is a d -dimensional common invariant subspace of T and T' , then V_d is spanned by d eigenvectors of T as well as d eigenvectors of T' . Since, eigenvectors of any matrix are linearly independent, this implies that there exist d eigenvectors of T' which can be written as linear combinations of d eigenvectors of T .

Also, if d eigenvectors of T' , denoted by $\{|e'_i\rangle\}_{i=1, \dots, d}$ can be written as a linear combination of d eigenvectors of T , denoted by $\{|e_i\rangle\}_{i=1, \dots, d}$, then there exist a common invariant subspace V_d of T and T' which is spanned by $\{|e'_i\rangle\}_{i=1, \dots, d}$. This completes the proof. \square

The example is stated below.

$$A_1 = \frac{1}{2} \begin{pmatrix} 1+i & 1-i & 0 & 0 \\ 1-i & 1+i & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2i \end{pmatrix}, \quad A_2 = \frac{1}{2} \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & -1+i & 1+i & 0 \\ 0 & 1+i & -1+i & 0 \\ 0 & 0 & 0 & -2i \end{pmatrix}, \quad A_3 = \frac{1}{2} \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 2i & 0 & 0 \\ 0 & 0 & -1-i & i-1 \\ 0 & 0 & i-1 & -1-i \end{pmatrix} \quad (\text{C59})$$

As discussed before, any non-trivial invariant subspace of a matrix T is spanned by its eigenvectors of T . Using this fact, any non-trivial invariant subspace of A_1 is spanned by the vectors $\{|+_{01}\rangle, |-_{01}\rangle, |2\rangle, |3\rangle\}$, any non-trivial invariant subspace of A_2 is spanned by the vectors $\{|0\rangle, |+_{12}\rangle, |-_{12}\rangle, |3\rangle\}$ and any non-trivial invariant subspace of A_3 is spanned by the vectors $\{|0\rangle, |1\rangle, |+_{23}\rangle, |-_{23}\rangle\}$ where $|\pm_{ij}\rangle = (|i\rangle \pm |j\rangle)/\sqrt{2}$.

Using Lemma 3, we can conclude that A_1 and A_3 have two common nontrivial invariant subspaces, spanned by $\{|0\rangle, |1\rangle\}$ and $\{|2\rangle, |3\rangle\}$. However, these two are not invariant subspaces of A_2 because it is not possible to represent an arbitrary vector from those subspaces as linear combinations of two of the eigenvectors of A_2 . Thus, we can conclude that there is no common invariant subspace shared between A_1, A_2 and A_3 .

Robust self-testing

Finally, we prove that for some particular choice of Alice's observables our self-testing statements are robust. However, for simplicity we assume here that the underlying state is pure and Bob's observables are projective.

Theorem 3. Consider a pair of genuinely d -outcome incompatible observables $A_1 = X_d Z_d^l$ with $l = 0, \dots, d-1$ and $A_2 = Z_d$, and assume that the following steering inequality

$$\sum_{i=1}^2 \sum_{k=1}^{d-1} \langle A_i^k \otimes B_i^k \rangle_{|\psi\rangle} \geq 2(d-1) - \varepsilon \quad (\text{C60})$$

is violated ε -close to the maximal violation by a state $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ and observables B_i ($i = 1, 2$) acting on \mathcal{H}_B . Then, for any $d \geq 2$ there exists a unitary operation $U_B : \mathcal{H}_B \rightarrow \mathcal{H}_B$ such that

$$\left\| (\mathbb{1} \otimes U_B)(\mathbb{1} \otimes B_i^k) |\psi\rangle - \mathbb{1} \otimes (B'_i)^k |\phi_d^+\rangle \right\| \leq \sqrt{2\varepsilon} + 2\sqrt{d} \sqrt[4]{2\varepsilon} \quad (\text{C61})$$

and

$$\|B_i^k - (B'_i)^k\|_2^2 \leq \sqrt{d} \sqrt{2\varepsilon} (1 + 4\sqrt{d} \sqrt{2\varepsilon}) \quad (\text{C62})$$

with $k = 0, \dots, d-1$, where B'_i are the ideal observables of Bob, i.e., $B'_i = A_i^*$, and $\|\cdot\|_2$ stands for the Hilbert-Schmidt norm.

Proof. The first step in proving the above statements is to observe that (C60) implies the following set of inequalities

$$\left| \langle \psi | A_i^k \otimes B_i^k | \psi \rangle \right| \geq \text{Re} \left(\langle \psi | A_i^k \otimes B_i^k | \psi \rangle \right) \geq 1 - \varepsilon \quad (\text{C63})$$

for any $i = 1, 2$ and $k = 1, \dots, d-1$.

The second observation we will exploit in our proof is that the shared state $|\psi\rangle$ can always be written in the basis of Z_d as

$$|\psi\rangle = \sum_{i=0}^{d-1} \alpha_i |i\rangle |b_i\rangle, \quad (\text{C64})$$

where $|b_i\rangle$ are some, not necessarily orthogonal vectors from \mathcal{H}_B , and α_i are non-negative numbers satisfying $\sum_{i=0}^{d-1} \alpha_i^2 = 1$.

Let us first exploit Ineq. (C63) for $i = 1$. Exploiting the fact that $(X_d Z_d^l)^k |i\rangle = \omega^{kl(i + \frac{k-1}{2})} |i+k\rangle$, where the addition is mod d , we obtain

$$\sum_i \alpha_i \alpha_{i+k} \text{Re} \left(\omega^{kl(i + \frac{k-1}{2})} \langle b_{i+k} | B_1^k | b_i \rangle \right) \geq 1 - \varepsilon \quad (\text{C65})$$

which by virtue of the facts that $\text{Re}(z) \leq |z|$ for any $z \in \mathbb{C}$ and that $\left| \omega^{kl(i + \frac{k-1}{2})} \langle b_{i+k} | B_1^k | b_i \rangle \right| \leq 1$ because B_1 is unitary and $|b_i\rangle$ are normalized, implies

$$\sum_i \alpha_i \alpha_{i+k} \geq 1 - \varepsilon \quad (\text{C66})$$

for any $k = 0, \dots, d-1$. By summing the above relation over k , one obtains

$$\sum_{i,k=0}^{d-1} \alpha_i \alpha_{i+k} = \left(\sum_{i=0}^{d-1} \alpha_i \right)^2 \geq d(1 - \varepsilon) \quad (\text{C67})$$

which gives

$$\sum_{i=0}^{d-1} \alpha_i \geq \sqrt{d} \sqrt{1 - \varepsilon}. \quad (\text{C68})$$

We can use this inequality to show that each α_i is " ε -close" to $1/\sqrt{d}$. To this end, let us write that

$$\begin{aligned} \sum_{i=0}^{d-1} \left(\alpha_i - \frac{1}{\sqrt{d}} \right)^2 &= \sum_{i=0}^{d-1} \alpha_i^2 - \frac{2}{\sqrt{d}} \sum_{i=0}^{d-1} \alpha_i + 1 \\ &= 2 \left(1 - \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} \alpha_i \right), \end{aligned} \quad (\text{C69})$$

where we used that $\sum_i \alpha_i^2 = 1$. Now, by using (C68) we can upper bound the above expression as

$$\begin{aligned} \sum_{i=0}^{d-1} \left(\alpha_i - \frac{1}{\sqrt{d}} \right)^2 &\leq 2(1 - \sqrt{1 - \varepsilon}) \\ &\leq 2\varepsilon, \end{aligned} \quad (\text{C70})$$

where we used $1 - \sqrt{1 - \varepsilon} \leq \varepsilon$, which in turn is a simple consequence of the fact that $\sqrt{1 - \varepsilon} \geq 1 - \varepsilon$ for any $0 \leq \varepsilon \leq 1$. This finally implies that for any i ,

$$\frac{1}{\sqrt{d}} - \sqrt{2\varepsilon} \leq \alpha_i \leq \frac{1}{\sqrt{d}} + \sqrt{2\varepsilon}. \quad (\text{C71})$$

In a similar manner we can prove that

$$\frac{1}{d} - \sqrt{2\varepsilon} \leq \alpha_i \alpha_{i+j} \leq \frac{1}{d} + \sqrt{2\varepsilon}. \quad (\text{C72})$$

for any $i, j = 0, \dots, d-1$. Precisely, let us consider the following expression

$$\begin{aligned} \sum_{i,j=0}^{d-1} \left(\alpha_i \alpha_{i+j} - \frac{1}{d} \right)^2 &= 2 - \frac{2}{d} \sum_{i,j=0}^{d-1} \alpha_i \alpha_{i+j} \\ &\leq 2 - \frac{2}{d} d(1 - \varepsilon) = 2\varepsilon, \end{aligned} \quad (\text{C73})$$

where we have used Eq. (C67). By noting that each term in the above sum must also be upper bounded by 2ε we arrive at (C72).

Let us now use conditions (C63) for $A_2 = Z_d$, which by using the fact that $Z_d^k|i\rangle = \omega^{ik}|i\rangle$ can be rewritten as

$$\sum_{i=0}^{d-1} \alpha_i^2 \operatorname{Re} \left(\omega^{ik} \langle b_i | B_2^k | b_i \rangle \right) \geq 1 - \varepsilon. \quad (\text{C74})$$

Using then the fact that $\operatorname{Re}(\omega^{ik} \langle b_i | B_2^k | b_i \rangle) \leq 1$ for any i, k , we have

$$\sum_{i \neq j} \alpha_i^2 + \alpha_j^2 \operatorname{Re} \left(\omega^{jk} \langle b_j | B_2^k | b_j \rangle \right) \geq 1 - \varepsilon, \quad (\text{C75})$$

which by virtue of $\sum_i \alpha_i^2 = 1$ implies

$$\alpha_j^2 \left[1 - \operatorname{Re} \left(\omega^{jk} \langle b_j | B_2^k | b_j \rangle \right) \right] \leq \varepsilon. \quad (\text{C76})$$

Using then the left-hand side inequality of Eq. (C71) and the fact the term in the square brackets is non-negative, we obtain

$$\frac{1}{d} \left[1 - \operatorname{Re} \left(\omega^{jk} \langle b_j | B_2^k | b_j \rangle \right) \right] \leq \varepsilon + \sqrt{2\varepsilon} \leq 2\sqrt{2\varepsilon}, \quad (\text{C77})$$

which, by rearranging terms gives

$$\operatorname{Re} \left(\omega^{jk} \langle b_j | B_2^k | b_j \rangle \right) \geq 1 - 2d\sqrt{2\varepsilon}. \quad (\text{C78})$$

Let us now consider the eigendecomposition of B_2 ,

$$B_2 = \sum_{i=0}^{d-1} \omega^i P_i, \quad (\text{C79})$$

where in general the orthogonal projections P_i may be of higher rank than one. By plugging it into (C78) one arrives at

$$\sum_{i=0}^{d-1} \text{Re} \left(\omega^{(i+j)k} \langle b_j | P_i | b_j \rangle \right) \geq 1 - 2d\sqrt{2\varepsilon}, \quad (\text{C80})$$

which after summing over $k = 0, \dots, d-1$ gives

$$\langle b_j | P_{-j} | b_j \rangle \geq 1 - 2d\sqrt{2\varepsilon}. \quad (\text{C81})$$

This inequality means that each vector $|b_j\rangle$ is ε -close to the subspace corresponding to the ω^{-j} outcome of B_2 . Let us then introduce vectors $|v_j\rangle = P_{-j}|b_j\rangle$ and their normalized versions $|\bar{v}_j\rangle = |v_j\rangle / \||v_j\rangle\|$. From the fact that the projections P_j are pairwise orthogonal we infer that $|\bar{v}_j\rangle$ are orthogonal as well. Moreover, Eq. (C81) implies that $\||v_j\rangle\| \geq 1 - 2d\sqrt{2\varepsilon}$, and so the norm of each $|v_j\rangle$ is close to unity.

Now, due to the fact that by the very construction each $|\bar{v}_j\rangle$ belongs to the subspace corresponding to P_{-j} , the observable B_2 can be written as

$$B_2 = \sum_{i=0}^{d-1} \omega^{-i} |\bar{v}_i\rangle\langle\bar{v}_i| \oplus B'_2, \quad (\text{C82})$$

where B'_2 is an operator whose support is orthogonal to all $|v_i\rangle$. Then, due to the fact that $|v_i\rangle$ are orthogonal one can find a unitary operation such that $U_B |\bar{v}_i\rangle = |d-i\rangle$. This operation brings B_2 to

$$U_B B_2 U_B^\dagger = \sum_{i=0}^{d-1} \omega^i |i\rangle\langle i| \oplus B''_2. \quad (\text{C83})$$

Denoting then $|b'_i\rangle = U_B |b_i\rangle$, where $|b_i\rangle$ are vectors appearing in the decomposition of $|\psi\rangle$ in Eq. (C64), we deduce from Eq. (C81) that

$$\langle b'_i | i \rangle \geq 1 - 2d\sqrt{2\varepsilon}. \quad (\text{C84})$$

Given conditions (C84) we can now move on to proving our main statements (C61) and (C62). To this end, we notice that for the unitary operation U_B introduced above, one has

$$\begin{aligned} \left\| (\mathbb{1} \otimes U_B B_i^k) |\psi\rangle - [\mathbb{1} \otimes (B_i^k)] |\phi_d^+\rangle \right\| &= \left\| (A_i^k \otimes U_B B_i^k) |\psi\rangle - [A_i^k \otimes (B_i^k)] |\phi_d^+\rangle \right\| \\ &= \left\| (A_i^k \otimes \tilde{B}_i^k) |\tilde{\psi}\rangle - |\tilde{\psi}\rangle + |\tilde{\psi}\rangle - |\phi_d^+\rangle \right\|, \end{aligned} \quad (\text{C85})$$

where the first equation is a consequence of the fact that A_i^k are unitary and that the vector norm is unitarily invariant, whereas the second equation has been obtained by adding and subtracting $|\tilde{\psi}\rangle$ with the latter defined as $|\tilde{\psi}\rangle = (\mathbb{1} \otimes U_B) |\psi\rangle$ and $\tilde{B}_i = U_B B_i U_B^\dagger$; in particular \tilde{B}_2 is given in Eq. (C83).

Using then the triangle inequality we arrive at

$$\left\| (A_i^k \otimes \tilde{B}_i^k) |\tilde{\psi}\rangle - |\tilde{\psi}\rangle + |\tilde{\psi}\rangle - |\phi_d^+\rangle \right\| \leq \left\| (A_i^k \otimes \tilde{B}_i^k) |\tilde{\psi}\rangle - |\tilde{\psi}\rangle \right\| + \left\| |\tilde{\psi}\rangle - |\phi_d^+\rangle \right\| \quad (\text{C86})$$

Let us now consider the first term of the right-hand side of the above inequality and write it as

$$\begin{aligned} \left\| (A_i^k \otimes \tilde{B}_i^k) |\tilde{\psi}\rangle - |\tilde{\psi}\rangle \right\| &= \left\| (A_i^k \otimes B_i^k) |\psi\rangle - |\psi\rangle \right\| \\ &= \left\{ 2 \left[1 - \text{Re} \left(\langle \psi | A_i^k \otimes B_i^k | \psi \rangle \right) \right] \right\}^{1/2} \\ &\leq \sqrt{2\varepsilon}, \end{aligned} \quad (\text{C87})$$

where to get the inequality we have used conditions (C63).

Let us then consider the second term appearing on the right-hand side of Eq. (C86) and rewrite it as

$$\begin{aligned}
\| |\tilde{\psi}\rangle - |\phi_d^+\rangle \| &= \{ 2 [1 - \text{Re}(\langle \tilde{\psi} | \phi_d^+ \rangle)] \}^{1/2} \\
&= \left\{ 2 \left[1 - \frac{1}{\sqrt{d}} \sum_i \alpha_i \text{Re}(\langle \bar{b}_i | i \rangle) \right] \right\}^{1/2} \\
&\leq \left\{ 2 \left[1 - \frac{1}{\sqrt{d}} (1 - 2d\sqrt{2\varepsilon}) \sum_i \alpha_i \right] \right\}^{1/2} \\
&\leq \left\{ 2 \left[1 - (1 - 2d\sqrt{2\varepsilon}) \sqrt{1 - \varepsilon} \right] \right\}^{1/2} \\
&\leq 2\sqrt{d} \sqrt[4]{2\varepsilon},
\end{aligned} \tag{C88}$$

where to obtain the first inequality we have used Eqs. (C84) and (C68), respectively, whereas the third inequality stems from two simple inequalities: $1 - \sqrt{1 - \varepsilon} \leq \varepsilon$ and $\varepsilon \leq \sqrt{2\varepsilon}$. By applying Eqs. (C86), (C87) and (C88) into (C85) we obtain (C61).

Let us eventually prove (C62). For this purpose, we first observe that

$$\left\| \tilde{B}_i^k - (B'_i)^k \right\|_2^2 = d \left\| [\tilde{B}_i^k - (B'_i)^k] |\phi_d^+\rangle \right\|^2. \tag{C89}$$

Then, the triangle inequality gives us

$$\begin{aligned}
\left\| \tilde{B}_i^k |\tilde{\psi}\rangle - (B'_i)^k |\phi_d^+\rangle \right\| &= \left\| \tilde{B}_i^k |\tilde{\psi}\rangle - (B'_i)^k |\phi_d^+\rangle + (B'_i)^k |\phi_d^+\rangle - \tilde{B}_i^k |\phi_d^+\rangle \right\| \\
&\geq \left\| [\tilde{B}_i^k - (B'_i)^k] |\phi_d^+\rangle \right\| - \left\| \tilde{B}_i^k (|\tilde{\psi}\rangle - |\phi_d^+\rangle) \right\|,
\end{aligned} \tag{C90}$$

which by taking into account the identity (C89) and the unitary invariance of the vector norm, leads us to

$$\left\| \tilde{B}_i^k - (B'_i)^k \right\|_2^2 \leq \sqrt{d} \left(\left\| \tilde{B}_i^k |\tilde{\psi}\rangle - (B'_i)^k |\phi_d^+\rangle \right\| + \| |\tilde{\psi}\rangle - |\phi_d^+\rangle \| \right). \tag{C91}$$

Using inequalities (C61) and (C88) we obtain (C62). \square