

Generalized Heisenberg algebra, realizations of the $\mathfrak{gl}(n)$ algebra and applications

Stjepan Meljanac^{*1}, Zoran Škoda^{†2}, and Rina Štrajn^{‡3}

¹Division of Theoretical Physics, Ruđer Bošković Institute, Bijenička cesta 54, 10002 Zagreb, Croatia

²Department of Teachers' Education, University of Zadar, Franje Tuđmana 24, 23000 Zadar, Croatia

³Department of Electrical Engineering and Computing, University of Dubrovnik, Ćira Carića 4, 20000 Dubrovnik, Croatia

Abstract

We introduce the generalized Heisenberg algebra appropriate for realizations of the $\mathfrak{gl}(n)$ algebra. Linear realizations of the $\mathfrak{gl}(n)$ algebra are presented and the corresponding star product, coproduct of momenta and twist are constructed. The dual realization and dual $\mathfrak{gl}(n)$ algebra are considered. Finally, we present a general realization of the $\mathfrak{gl}(n)$ algebra, the corresponding coproduct of momenta and two classes of twists. These results can be applied to physical theories on noncommutative spaces of the $\mathfrak{gl}(n)$ type.

^{*}meljanac@irb.hr

[†]zskoda@unizd.hr

[‡]rina.strajn@unidu.hr

1 Introduction

Noncommutative (NC) geometry appeared in theoretical physics as one possible approach towards the description of spacetime at the Planck scale [1, 2, 3, 4, 5]. The key assumption is noncommutativity between spacetime coordinates. The first model was proposed by Snyder in Ref. [3]. There are various models of NC spaces of Lie algebra type. Specially, an interesting model is the κ -Minkowski spacetime [4, 5, 6, 7, 8], where the parameter κ is usually interpreted as the quantum gravity scale and the spacetime coordinates close a Lie algebra. The κ -Poincaré quantum group [4, 5, 9, 10], as a possible quantum symmetry of the κ -Minkowski spacetime, allows for the study of deformed relativistic spacetime symmetries. Realizations of Lie algebras are important in the formulation of physical theories on NC spaces and the corresponding deformed symmetries. Particularly, realizations of NC coordinates are based on formal power series in the Heisenberg-Weyl algebra allowing one to simplify the methods of calculation on the deformed spacetime. Some physical applications of Lie algebra type NC spaces and the associated star products are presented in Refs. [11, 12, 13].

For certain Lie algebras, such as the orthogonal algebra $\mathfrak{so}(n)$, the Lorentz algebra $\mathfrak{so}(1, n - 1)$ and the $\mathfrak{gl}(n)$ algebra, the realizations are not well adapted due to the structure of their commutation relations. This was the motivation for introducing the generalized Heisenberg algebra and constructing an analogue of the Weyl realization of $\mathfrak{so}(n)$ and $\mathfrak{so}(1, n - 1)$ by formal power series in a semicompletion of the Heisenberg algebra [14]. This construction was applied to the extended Snyder model [15] and to the unification of the κ -Minkowski and extended Snyder spaces [16, 17].

In this paper we introduce another generalization of the Heisenberg algebra appropriate for realizations of the $\mathfrak{gl}(n)$ algebra. Specially, we consider linear realizations of the $\mathfrak{gl}(n)$ algebra and we use them to construct the corresponding star product, coproduct of momenta and twist as applications to a noncommutative space of $\mathfrak{gl}(n)$ type. Furthermore, we consider the dual realization and dual $\mathfrak{gl}(n)$ algebra, as well as a general realization of the $\mathfrak{gl}(n)$ algebra, coproduct of momenta and twists. These constructions were performed using methods proposed in Refs. [18, 19, 20, 21] and in a Lie deformed phase space from twists in the Hopf algebroid approach in Refs. [19, 22, 23, 24, 25].

The plan of paper is as follows. In section 2 we introduce the generalized Heisenberg algebra and present linear realizations of the $\mathfrak{gl}(n)$ algebra. In section 3 the star product, and in section 4 the coproduct of momenta and twist are constructed. In section 5 the dual realization and dual $\mathfrak{gl}(n)$ algebra are considered. In section 6 a general realization of the $\mathfrak{gl}(n)$ algebra, the corresponding coproduct of momenta and two classes of twists are presented. These results can be applied to physical theories on noncommutative spaces of the $\mathfrak{gl}(n)$ type.

2 Linear realizations of the $\mathfrak{gl}(n)$ algebra

We introduce the generalized Heisenberg algebra generated with n^2 coordinates $x_{\mu\nu}$ and momenta $p_{\mu\nu}$, $\mu, \nu = 1, \dots, n$, defined by

$$[x_{\mu\nu}, x_{\alpha\beta}] = 0, \quad [p_{\mu\nu}, p_{\alpha\beta}] = 0, \quad [p_{\mu\nu}, x_{\alpha\beta}] = -i\delta_{\mu\alpha}\delta_{\nu\beta} \quad (1)$$

Let us define the $\mathfrak{gl}(n)$ algebra generated with $\hat{x}_{\mu\nu}$, $\mu, \nu = 1, \dots, n$

$$[\hat{x}_{\mu\nu}, \hat{x}_{\lambda\rho}] = iu(\delta_{\mu\rho}\hat{x}_{\lambda\nu} - \delta_{\lambda\nu}\hat{x}_{\mu\rho}) = iuC_{\mu\nu,\lambda\rho,\alpha\beta}\hat{x}_{\alpha\beta}, \quad u \in \mathbb{R}, \quad (2)$$

where

$$C_{\mu\nu,\lambda\rho,\alpha\beta} = iu(\delta_{\mu\rho}\delta_{\lambda\alpha}\delta_{\nu\beta} - \delta_{\lambda\nu}\delta_{\mu\alpha}\delta_{\rho\beta}). \quad (3)$$

are structure constants. Note that for $u = 0$, $\hat{x}_{\mu\nu}$ commute among themselves.

One linear realization of $\hat{x}_{\mu\nu}$ is

$$\hat{x}_{\mu\nu} = x_{\mu\nu} + u x_{\mu\alpha} p_{\nu\alpha}, \quad (4)$$

$$\hat{x}_{\mu\nu} = x_{\alpha\beta} \varphi_{\alpha\beta,\mu\nu}, \quad (5)$$

where

$$\varphi_{\alpha\beta,\mu\nu} = \delta_{\alpha\mu}\delta_{\beta\nu} + u\delta_{\alpha\mu}p_{\nu\beta}. \quad (6)$$

Summation over repeated indices is understood and assumed throughout the whole paper.

Another linear realization of the $\mathfrak{gl}(n)$ algebra (2) is

$$\hat{x}_{\mu\nu} = x_{\mu\nu} - u x_{\alpha\nu} p_{\alpha\mu}, \quad (7)$$

$$\hat{x}_{\mu\nu} = x_{\alpha\beta} \varphi_{\alpha\beta,\mu\nu}, \quad (8)$$

where

$$\varphi_{\alpha\beta,\mu\nu} = \delta_{\alpha\mu}\delta_{\beta\nu} - u\delta_{\beta\nu}p_{\alpha\mu}. \quad (9)$$

This realization is related to the dual realization of the first realization (4) with $u \mapsto -u$, see section 5.

3 Star product

Action \triangleright is defined by

$$x_{\mu\nu} \triangleright f(x) = x_{\mu\nu} f(x), \quad (10)$$

$$p_{\mu\nu} \triangleright f(x) = -i \frac{\partial f(x)}{\partial x_{\mu\nu}}. \quad (11)$$

Using \triangleright , it follows

$$\hat{x}_{\mu\nu} \triangleright 1 = x_{\mu\nu}, \quad (12)$$

$$p_{\mu\nu} \triangleright e^{iqx} = q_{\mu\nu} e^{iqx}, \quad \text{where } qx = q_{\alpha\beta} x_{\alpha\beta}, \quad (13)$$

$$e^{ik_{\alpha\beta} \hat{x}_{\alpha\beta}} \triangleright e^{iq_{\gamma\delta} x_{\gamma\delta}} = e^{iJ_{\alpha\beta}(k,q,u)x_{\alpha\beta}}. \quad (14)$$

If $k_{\alpha\beta} = 0$, $J_{\mu\nu}(0, q, u) = q_{\mu\nu}$. If $q_{\mu\nu} = 0$, $J_{\mu\nu}(k, 0, u) = K_{\mu\nu}(k, u)$. If $u = 0$, $J_{\mu\nu}(k, q, 0) = k_{\mu\nu} + q_{\mu\nu}$.

From $J_{\mu\nu}(k, q, u)$, we can obtain the star product [18, 19]

$$e^{ikx} * e^{iqx} = e^{i\mathcal{D}(k,q,u)x}, \quad (15)$$

where

$$\mathcal{D}_{\mu\nu}(k, q, u) = J_{\mu\nu}(K^{-1}(k), q, u), \quad (16)$$

$$K_{\mu\nu}^{-1}(K(k)) = K_{\mu\nu}(K^{-1}(k)) = k_{\mu\nu}. \quad (17)$$

$J_{\mu\nu}(tk, q, u)$ is the unique solution of the partial differential equation [18, 20, 21]

$$\frac{\partial J_{\mu\nu}(tk, q, u)}{\partial t} = k_{\alpha\beta} \varphi_{\mu\nu, \alpha\beta}(J(tk, q, u)), \quad (18)$$

with boundary condition $J_{\mu\nu}(0, q, 0) = q_{\mu\nu}$. In our case

$$\frac{\partial J_{\mu\nu}(tk, q, u)}{\partial t} = k_{\alpha\beta}(\delta_{\alpha\mu}\delta_{\beta\nu} + u\delta_{\alpha\mu}J_{\beta\nu}(tk, q, u)) = k_{\mu\beta}(\delta_{\beta\nu} + uJ_{\beta\nu}(tk, q, u)). \quad (19)$$

The differential equation in matrix form is

$$\frac{\partial J(tk, q, u)}{\partial t} = k(I + uJ(tk, q, u)). \quad (20)$$

The solution in matrix form is

$$J(tk, q, u) = \frac{1}{u}(e^{uk} - I) + e^{uk}q, \quad (21)$$

or in terms of the components

$$J_{\mu\nu}(tk, q, u) = \left(e^{uk}\right)_{\mu\alpha} q_{\alpha\nu} + \frac{1}{u}(-I + e^{uk})_{\mu\nu}. \quad (22)$$

For $t = 0$

$$J_{\mu\nu}(0, q, u) = q_{\mu\nu}, \quad (23)$$

$$K_{\mu\nu}(k, u) = \frac{1}{u}(e^{uk} - I)_{\mu\nu}, \quad (24)$$

$$K_{\mu\nu}^{-1}(k, u) = \frac{1}{u}(\ln(I + uk))_{\mu\nu}. \quad (25)$$

Hence

$$\mathcal{D}_{\mu\nu}(tk, q, u) = k_{\mu\nu} + q_{\mu\nu} + uk_{\mu\alpha}q_{\alpha\nu}. \quad (26)$$

The star product is associative

$$(f * g) * h = f * (g * h). \quad (27)$$

4 Coproduct of momenta and twist

The coproduct of momenta is

$$\Delta p_{\mu\nu} = \mathcal{D}_{\mu\nu}(p \otimes 1, 1 \otimes p) = p_{\mu\nu} \otimes 1 + 1 \otimes p_{\mu\nu} + up_{\mu\alpha} \otimes p_{\alpha\nu}. \quad (28)$$

This coproduct is coassociative $(\Delta \otimes 1)\Delta = (1 \otimes \Delta)\Delta$.

In the Hopf algebroid approach [22, 23, 24, 25] we can write the twist [19]

$$\mathcal{F}^{-1} = : \exp(i(1 \otimes x_{\alpha\beta})(\Delta - \Delta_0)p_{\alpha\beta}) : \quad (29)$$

$$= \exp(-ip_{\alpha\beta} \otimes x_{\alpha\beta}) \exp(iK^{-1}(p)_{\gamma\delta} \otimes \hat{x}_{\gamma\delta}), \quad (30)$$

where $:$ denotes normal ordering, with the x s left from the p s. Using the coproduct Δp (28), we have

$$\mathcal{F}_1^{-1} =: \exp(iup_{\mu\nu} \otimes x_{\mu\alpha}p_{\nu\alpha}) : \quad (31)$$

and applying Theorem 1 [21] we get

$$\mathcal{F}_1^{-1} = \exp(i(\ln(I + up))_{\mu\nu} \otimes x_{\mu\alpha}p_{\nu\alpha}). \quad (32)$$

Note that

$$K_{\mu\nu}^{-1}(p, u) = \frac{1}{u}(\ln(I + up))_{\mu\nu}. \quad (33)$$

Here we introduce

$$Z_{\mu\nu} = (I + up)_{\mu\nu}, \quad (34)$$

and

$$L_{\mu\nu} = x_{\mu\alpha}p_{\nu\alpha}. \quad (35)$$

$L_{\mu\nu}$ generate the $\mathfrak{gl}(n)$ algebra, with properties

$$[Z_{\mu\nu}, \hat{x}_{\lambda\rho}] = -iu\delta_{\mu\lambda}Z_{\rho\nu}, \quad (36)$$

$$\Delta Z_{\mu\nu} = Z_{\mu\alpha} \otimes Z_{\alpha\nu}, \quad (37)$$

$$\Delta(\ln Z)_{\mu\nu} = (\ln Z)_{\mu\nu} \otimes 1 + 1 \otimes (\ln Z)_{\mu\nu}, \quad (38)$$

$$[L_{\mu\nu}, L_{\lambda\rho}] = i(\delta_{\mu\rho}L_{\lambda\nu} - \delta_{\lambda\nu}L_{\mu\rho}), \quad (39)$$

$$\Delta_0 L_{\mu\nu} = L_{\mu\nu} \otimes 1 + 1 \otimes L_{\mu\nu}, \quad (40)$$

$$\Delta_0 p_{\mu\nu} = p_{\mu\nu} \otimes 1 + 1 \otimes p_{\mu\nu}. \quad (41)$$

Hence, the twist can be written as

$$\mathcal{F}_1^{-1} = \exp(i(\ln Z)_{\mu\nu} \otimes L_{\mu\nu}). \quad (42)$$

It is easy to check that

$$\hat{x}_{\mu\nu} = m\mathcal{F}^{-1}(\mathfrak{s} \otimes 1)(x_{\mu\nu} \otimes 1) = x_{\mu\nu} + ux_{\mu\alpha}p_{\nu\alpha}, \quad (43)$$

where m is the multiplication map $m : A \otimes B \rightarrow AB$, and

$$\Delta p_{\mu\nu} = \mathcal{F} \Delta_0 p_{\mu\nu} \mathcal{F}^{-1} = \Delta_0 p_{\mu\nu} + up_{\mu\alpha} \otimes p_{\alpha\nu}, \quad (44)$$

where

$$\Delta_0 p_{\mu\nu} = p_{\mu\nu} \otimes 1 + 1 \otimes p_{\mu\nu}.$$

Twist \mathcal{F}_1 satisfies the Drinfeld cocycle condition

$$(1 \otimes \mathcal{F})(1 \otimes \Delta_0)\mathcal{F} = (\mathcal{F} \otimes 1)(\Delta_0 \otimes 1)\mathcal{F}. \quad (45)$$

The proof follows using the factorization property. Using $\Delta_0 L_{\mu\nu}$ (40), we get

$$(1 \otimes \Delta_0)\mathcal{F} = \mathcal{F}_{12}\mathcal{F}_{13} = \mathcal{F}_{13}\mathcal{F}_{12}, \quad (46)$$

$$(1 \otimes \mathcal{F})(1 \otimes \Delta_0)\mathcal{F} = \mathcal{F}_{23}\mathcal{F}_{12}\mathcal{F}_{13} = \mathcal{F}_{23}\mathcal{F}_{13}\mathcal{F}_{12}. \quad (47)$$

Using $\Delta(\ln Z)_{\mu\nu} = (\ln Z)_{\mu\nu} \otimes 1 + 1 \otimes (\ln Z)_{\mu\nu}$, we get

$$(\mathcal{F} \Delta_0 \otimes 1)\mathcal{F} = \mathcal{F}_{13}\mathcal{F}_{23}\mathcal{F}_{12} = \mathcal{F}_{23}\mathcal{F}_{13}\mathcal{F}_{12}. \quad (48)$$

Hence, the left and right side of (45) become $\mathcal{F}_{23}\mathcal{F}_{13}\mathcal{F}_{12}$.

5 Dual realization and dual $\mathfrak{gl}(n)$ algebra

Let us define the dual realization

$$\hat{y}_{\mu\nu} = m\tilde{\mathcal{F}}_1^{-1}(\triangleright \otimes 1)(x_{\mu\nu} \otimes 1) \quad (49)$$

$$= x_{\mu\nu} + ux_{\alpha\nu}p_{\alpha\mu} \quad (50)$$

$$= x_{\alpha\beta}\tilde{\varphi}_{\alpha\beta,\mu\nu}, \quad (51)$$

where

$$\tilde{\varphi}_{\alpha\beta,\mu\nu} = \delta_{\alpha\mu}\delta_{\beta\nu} - u\delta_{\beta\nu}p_{\alpha\mu}, \quad (52)$$

and $\tilde{\mathcal{F}}$ denotes interchange of the left and right sides in the \otimes product, $a \otimes b \rightarrow b \otimes a$. $\hat{y}_{\mu\nu}$ generate the dual $\mathfrak{gl}(n)$ algebra (2), with $u \mapsto -u$.

$$[\hat{y}_{\mu\nu}, \hat{y}_{\lambda\rho}] = -iu(\delta_{\mu\rho}\hat{y}_{\lambda\nu} - \delta_{\lambda\nu}\hat{y}_{\mu\rho}), \quad (53)$$

$$[Z_{\mu\nu}, \hat{y}_{\lambda\rho}] = -iu\delta_{\rho\nu}Z_{\mu\lambda}, \quad (54)$$

and

$$[\hat{x}_{\mu\nu}, \hat{y}_{\lambda\rho}] = 0. \quad (55)$$

The star product induced with the dual realization $\hat{y}_{\mu\nu}$ is constructed as in section 3. The partial differential equation for $\tilde{J}_{\mu\nu}(tk, q, u)$ is

$$\frac{\partial \tilde{J}_{\mu\nu}(tk, q, u)}{\partial t} = k_{\alpha\beta}\tilde{\varphi}_{\mu\nu,\alpha\beta}(\tilde{J}(tk, q, u)) \quad (56)$$

$$= k_{\mu\nu} + u\tilde{J}_{\mu\alpha}(tk, q, u)k_{\alpha\nu}. \quad (57)$$

The differential equation in matrix form is

$$\frac{\partial \tilde{J}(tk, q, u)}{\partial t} = (I + u\tilde{J}(tk, q, u))k. \quad (58)$$

The solution in matrix form is

$$\tilde{J}(tk, q, u) = \left(\frac{e^{utk} - I}{u} \right) + qe^{utk}, \quad (59)$$

and

$$\tilde{J}_{\mu\nu}(tk, q, u) = q_{\mu\alpha} \left(e^{utk} \right)_{\alpha\nu} + \left(\frac{e^{utk} - I}{u} \right)_{\mu\nu}. \quad (60)$$

For $t = 0$,

$$\tilde{J}_{\mu\nu}(0, q, u) = q_{\mu\nu}, \quad (61)$$

$$\tilde{K}_{\mu\nu}(k, u) = \frac{1}{u} \left(e^{uk} - I \right)_{\mu\nu} = K_{\mu\nu}(k, u), \quad (62)$$

$$\tilde{K}_{\mu\nu}^{-1}(k, u) = \frac{1}{u} \left(\ln(I + uk) \right)_{\mu\nu} = K_{\mu\nu}^{-1}(k, u). \quad (63)$$

Hence,

$$\tilde{D}_{\mu\nu}(k, q, u) = k_{\mu\nu} + q_{\mu\nu} + uk_{\alpha\nu}q_{\mu\alpha}. \quad (64)$$

The star product is associative.

The coproduct of $p_{\mu\nu}$ is

$$\tilde{\Delta}p_{\mu\nu} = p_{\mu\nu} \otimes 1 + 1 \otimes p_{\mu\nu} + up_{\alpha\nu} \otimes p_{\mu\alpha}, \quad (65)$$

where $\tilde{\Delta}$ denotes the interchange of the left and right side in the tensor product with respect to Δ , $a \otimes b \rightarrow b \otimes a$. The coproduct is coassociative.

The corresponding twist \mathcal{F}_2 in the Hopf algebroid approach [19] (sec.4) and the coproduct $\tilde{\Delta}p_{\mu\nu}$ (65) lead to

$$\mathcal{F}_2^{-1} = : \exp \left(iup_{\nu\alpha} \otimes x_{\mu\alpha}p_{\mu\nu} \right) : \quad (66)$$

$$= \exp \left(i \left(\ln(I + up) \right)_{\mu\nu} \otimes x_{\alpha\nu}p_{\alpha\mu} \right). \quad (67)$$

Note that

$$\tilde{K}_{\mu\nu}^{-1}(p, u) = \frac{1}{u} \left(\ln(I + up) \right)_{\mu\nu} = \frac{1}{u} \left(\ln Z \right)_{\mu\nu}. \quad (68)$$

We define

$$\tilde{L}_{\mu\nu} = x_{\alpha\nu}p_{\alpha\mu}. \quad (69)$$

$\tilde{L}_{\mu\nu}$ generate the $\mathfrak{gl}(n)$ algebra

$$[\tilde{L}_{\mu\nu}, \tilde{L}_{\lambda\rho}] = -i(\delta_{\mu\rho}\tilde{L}_{\lambda\nu} - \delta_{\lambda\nu}\tilde{L}_{\mu\rho}), \quad (70)$$

$$\Delta_0\tilde{L}_{\mu\nu} = \tilde{L}_{\mu\nu} \otimes 1 + 1 \otimes \tilde{L}_{\mu\nu}. \quad (71)$$

Hence, the twist can be written as

$$\mathcal{F}_2^{-1} = \exp \left(i \left(\ln Z \right)_{\mu\nu} \otimes \tilde{L}_{\mu\nu} \right). \quad (72)$$

It is easy to check that

$$\hat{y}_{\mu\nu} = m\mathcal{F}_2^{-1}(\triangleright \otimes 1)(x_{\mu\nu} \otimes 1) \quad (73)$$

$$= x_{\mu\nu} + ux_{\alpha\nu}p_{\alpha\mu}, \quad (74)$$

and

$$\tilde{\Delta}p_{\mu\nu} = \mathcal{F}_2\Delta_0p_{\mu\nu}\mathcal{F}_2^{-1} \quad (75)$$

$$= \Delta_0p_{\mu\nu} + up_{\alpha\nu} \otimes p_{\mu\alpha}. \quad (76)$$

Twist $\tilde{\mathcal{F}}_2$ satisfies the Drinfeld cocycle condition. From the duality property, it follows

$$\hat{x}_{\mu\nu} = m\tilde{\mathcal{F}}_2^{-1}(\triangleright \otimes 1)(x_{\mu\nu} \otimes 1) \quad (77)$$

$$= x_{\mu\nu} + ux_{\mu\alpha}p_{\nu\alpha}, \quad (78)$$

and

$$\Delta p_{\mu\nu} = \tilde{\mathcal{F}}_2\Delta_0p_{\mu\nu}\tilde{\mathcal{F}}_2^{-1}. \quad (79)$$

It is important to note that twists \mathcal{F}_1 and $\tilde{\mathcal{F}}_2$ are different, but lead to the same $\hat{x}_{\mu\nu}$ and the same coproduct Δ . Analogously, twists $\tilde{\mathcal{F}}_1$ and \mathcal{F}_2 are different, but they lead to the same $\hat{y}_{\mu\nu}$ and coproduct $\tilde{\Delta}$. Also, $\hat{y}_{\mu\nu}(-u)$ gives another linear realization of the $\mathfrak{gl}(n)$ algebra (2), different from $\hat{x}_{\mu\nu}$ (4). The dual realization with respect to $\hat{y}_{\mu\nu}(-u)$ is $\hat{x}_{\mu\nu}(-u)$, generating $\mathfrak{gl}(n)$, with $u \mapsto -u$.

6 General realization of the $\mathfrak{gl}(n)$ algebra, coproduct of momenta and two classes of twists

A general realization of the generators $\hat{x}_{\mu\nu}$ and $\hat{y}_{\mu\nu}$ generating the $\mathfrak{gl}(n)$ algebra (2) and the $\mathfrak{gl}(n)$ algebra (53) with $u \mapsto -u$, respectively, can be obtained using a similarity transformation on the coordinates $x_{\mu\nu}$ and momenta $p_{\mu\nu}$. Let us define new coordinates and momenta as

$$x'_{\mu\nu} = e^{x_{\alpha\beta}S_{\alpha\beta}(p,u)+T(p,u)}x_{\mu\nu}e^{-(x_{\alpha\beta}S_{\alpha\beta}(p,u)+T(p,u))}, \quad (80)$$

$$p'_{\mu\nu} = e^{x_{\alpha\beta}S_{\alpha\beta}(p,u)+T(p,u)}p_{\mu\nu}e^{-(x_{\alpha\beta}S_{\alpha\beta}(p,u)+T(p,u))} = \Lambda_{\mu\nu}(p, u). \quad (81)$$

Hence, using the inverse relations, we obtain

$$\hat{x}_{\mu\nu} = x'_{\alpha\beta}\varphi'_{\alpha\beta,\mu\nu}(p', u) + \chi'_{\mu\nu}(p', u), \quad (82)$$

$$\hat{y}_{\mu\nu} = x'_{\alpha\beta}\tilde{\varphi}'_{\alpha\beta,\mu\nu}(p', u) + \tilde{\chi}'_{\mu\nu}(p', u), \quad (83)$$

where $\varphi'_{\alpha\beta,\mu\nu}$, $\tilde{\varphi}'_{\alpha\beta,\mu\nu}$, $\chi'_{\mu\nu}$ and $\tilde{\chi}'_{\mu\nu}$ depend on $S(p, u)$ and $T(p, u)$. Note that, after inserting the realizations of $\hat{x}_{\mu\nu}$ (82) and $\hat{y}_{\mu\nu}$ (83), relations $[\hat{x}_{\mu\nu}, \hat{y}_{\lambda\rho}] = 0$ are satisfied. Furthermore $\Delta p'_{\mu\nu}$ is obtained from

$$\Delta p'_{\mu\nu} = \Delta(\Lambda_{\mu\nu}(p, u)) = \Lambda_{\mu\nu}(\Delta p, u)|_{p=\Lambda^{-1}(p', u)}, \quad (84)$$

i.e. expressing p as $\Lambda^{-1}(p', u)$. Similarly holds for $\tilde{\Delta}p'_{\mu\nu}$.

From $\Delta p'_{\mu\nu}$ and $\tilde{\Delta} p'_{\mu\nu}$ we easily obtain $\mathcal{D}'_{\mu\nu}(k, q, u)$ and $\tilde{\mathcal{D}}'_{\mu\nu}(k, q, u)$ that define the corresponding star products. Namely,

$$\Delta p'_{\mu\nu} = \mathcal{D}'_{\mu\nu}(p' \otimes 1, 1 \otimes p'), \quad (85)$$

$$\tilde{\Delta} p'_{\mu\nu} = \tilde{\mathcal{D}}'_{\mu\nu}(p' \otimes 1, 1 \otimes p'), \quad (86)$$

$$e^{ikx'} *_1 e^{iqx'} = e^{i\mathcal{D}'(k,q,u)x'} \quad (87)$$

$$e^{ikx'} *_2 e^{iqx'} = e^{i\tilde{\mathcal{D}}'(k,q,u)x'}, \quad (88)$$

$$\Delta p'_{\mu\nu}(\triangleright \otimes \triangleright)(e^{ikx'} \otimes e^{iqx'}) = \mathcal{D}'_{\mu\nu}(k, q, u)(e^{ikx'} \otimes e^{iqx'}), \quad (89)$$

$$\tilde{\Delta} p'_{\mu\nu}(\triangleright \otimes \triangleright)(e^{ikx'} \otimes e^{iqx'}) = \tilde{\mathcal{D}}'_{\mu\nu}(k, q, u)(e^{ikx'} \otimes e^{iqx'}), \quad (90)$$

$$\Delta_0 p'_{\mu\nu} = p'_{\mu\nu} \otimes 1 + 1 \otimes p'_{\mu\nu}. \quad (91)$$

Generally, $f *_1 g = g *_2 f$, for arbitrary $f(x')$ and $g(x')$. A consistency check is the relation between $\hat{x}_{\mu\nu}$ and $\Delta p'_{\alpha\beta}$

$$\hat{x}_{\mu\nu} = x'_{\mu\nu} + ix'_{\alpha\beta} m(\Delta - \Delta_0) p'_{\alpha\beta}(\triangleright \otimes 1)(x'_{\mu\nu} \otimes 1) + \chi'_{\mu\nu}(p', u), \quad (92)$$

and similarly for $\hat{y}_{\mu\nu}$.

The family of twists $\mathcal{F}_1(s)$ in the Hopf algebroid approach corresponding to the general realization of $\hat{x}_{\mu\nu}$ (82), for $\chi'_{\mu\nu}(p', u) = 0$, $T(p, u) = 0$ is

$$\mathcal{F}_1^{-1}(s) =: \exp\left(i((1-s)(1 \otimes x'_{\alpha\beta}) + sx'_{\alpha\beta} \otimes 1)(\Delta - \Delta_0)p'_{\alpha\beta}\right) :, \quad (93)$$

where $\Delta p'_{\mu\nu} = p'_{\mu\nu} \otimes 1 + 1 \otimes p'_{\mu\nu}$, and $s \in \mathbb{R}$. We point out that although these twists are different for different s , they define the same star product for all s . Namely,

$$\begin{aligned} e^{ikx'} *_1 e^{iqx'} &= m\mathcal{F}_1^{-1}(s)(\triangleright \otimes \triangleright)(e^{ikx'} \otimes e^{iqx'}) \\ &= m \exp\left(i((1-s)1 \otimes x'_{\alpha\beta} + s1 \otimes x'_{\alpha\beta})(\Delta - \Delta_0)p'_{\alpha\beta}\right)(\mathcal{D}'_{\alpha\beta}(k, q, u) - k_{\alpha\beta} - q_{\alpha\beta})(e^{ikx'} \otimes e^{iqx'}) \\ &= e^{ix_{\alpha\beta}\mathcal{D}'_{\alpha\beta}(k,q,u)} \end{aligned} \quad (94)$$

does not depend on s due to (89), the definition of normal ordering and the multiplication map m . Since $\Delta p'_{\mu\nu} = \mathcal{D}'(p' \otimes 1, 1 \otimes p')$, (85), the coproduct $\Delta p'_{\mu\nu} = \mathcal{F}_1(s)\Delta_0 p'_{\mu\nu}\mathcal{F}_1^{-1}(s)$ does not depend on the parameter s . Similarly, $\hat{x}_{\mu\nu} = m\mathcal{F}_1^{-1}(s)(\triangleright \otimes 1)(x'_{\mu\nu} \otimes 1)$ does not depend on s . Analogously, we can define the family of twists $\mathcal{F}_2(s)$ corresponding to the general realization of $\hat{y}_{\mu\nu}$ (83) with $\chi'_{\mu\nu}(p', u) = 0$ and $T(p, u) = 0$

$$\mathcal{F}_2^{-1}(s) =: \exp\left(i(1-s)(1 \otimes x'_{\alpha\beta} + sx'_{\alpha\beta} \otimes 1)(\Delta - \Delta_0)p'_{\alpha\beta}\right) :. \quad (95)$$

Note that $\mathcal{F}_1(s) = \tilde{\mathcal{F}}_2(1-s)$.

References

- [1] S. Doplicher, K. Fredenhagen, J. Roberts, ‘‘Spacetime quantization induced by classical gravity’’, Phys. Lett. **B331** 39-44 (1994).

- [2] S. Doplicher, K. Fredenhagen, J. Roberts, “The quantum structure of spacetime at the Planck scale and quantum fields”, *Comm. Math. Phys.* **172**, 187-220 (1995).
- [3] H. S. Snyder, “Quantized Space-Time”, *Phys. Rev.* **71** 38 (1947).
- [4] J. Lukierski, A. Nowicki, H. Ruegg, V. N. Tolstoy, “q-deformation of Poincaré algebra”, *Phys. Lett.* **B 264**, 331 (1991).
- [5] J. Lukierski, A. Nowicki, H. Ruegg, “New quantum Poincaré algebra and κ -deformed field theory”, *Phys. Lett.* **B 293** 344 (1992).
- [6] S. Majid, H. Ruegg, “Bicrossproduct structure of κ -Poincaré group and noncommutative geometry”, *Phys. Lett.* **B334** 348-354 (1994), arXiv:hep-th/9404107.
- [7] G. Amelino-Camelia, J. Lukierski, A. Nowicki, “Kappa-deformed covariant phase space and quantum gravity uncertainty relations”, *Phys. At. Nucl.* **61**, 1811-1815 (1998), arXiv:hep-th/9706031.
- [8] M. Daszkiewicz, J. Lukierski, M. Woronowicz, “Towards quantum noncommutative κ -deformed field theory”, *Phys. Rev.* **D77**, 105007 (2008), arXiv:0708.1561.
- [9] S. Majid, “Foundations of Quantum Group Theory”, Cambridge University Press, 1995.
- [10] V. Chari, A. Pressley, “A guide to quantum groups”, Cambridge University Press, Cambridge (1994).
- [11] D. Meljanac, S. Meljanac, S. Mignemi and R. Štrajn, “ κ -deformed phase spaces, Jordanian twists, Lorentz-Weyl algebra and dispersion relations”, *Phys. Rev.* **D 99** 126012 (2019), arXiv:1903.08679.
- [12] T. R. Govindarajan, K. S. Gupta, E. Harikumar, S. Meljanac, D. Meljanac, “Twisted Statistics in kappa-Minkowski Spacetime”, *Phys. Rev.* **D 77** 105010 (2008), arXiv:0802.1576.
- [13] T. Jurić, S. Meljanac, D. Pikutić, R. Štrajn, “Toward the classification of differential calculi on κ -Minkowski space and related field theories”, *JHEP* **1507** 055 (2015), arXiv:1502.02972.
- [14] S. Meljanac, T. Martinić-Bilać, S. Krešić-Jurić, “Generalized Heisenberg algebra applied to realizations of the orthogonal, Lorentz and Poincare algebras and their dual extensions”, *J. Math. Phys.* **61**, 051705 (2020), arXiv:2003.02726.
- [15] S. Meljanac, S. Mignemi, “Associative realizations of the extended Snyder model”, *Phys. Rev.* **D 102** 126011 (2020), arXiv:2007.13498.
- [16] S. Meljanac, S. Mignemi, “Unification of κ -Minkowski and extended Snyder spaces”, *Phys. Lett.* **B 814** 136117 (2021), arXiv:2101.05275.
- [17] S. Meljanac, S. Mignemi, “Associative realizations of κ -deformed extended Snyder model”, arXiv:2106.08131.

- [18] S. Meljanac, D. Meljanac, A. Samsarov, M. Stojić, “Kappa Snyder deformations of Minkowski spacetime, realizations and Hopf algebra”, *Phys. Rev. D* **83** 065009 (2011), arXiv:1102.1655.
- [19] S. Meljanac, D. Meljanac, A. Pachoł, D. Pikutić, “Remarks on simple interpolation between Jordanian twists”, *J.Phys.A* **50** 26, 265201 (2017), arXiv:1612.07984.
- [20] A. Borowiec, S. Meljanac, D. Meljanac and A. Pachoł, “Interpolations between Jordanian twists induced by coboundary twists”, *SIGMA* **15** 054 (2019), arXiv:1812.05535.
- [21] S. Meljanac, R. Štrajn, “Exponential formulas, normal ordering and the Weyl-Heisenberg algebra”, arXiv:2105.12593.
- [22] T. Jurić, S. Meljanac, R. Štrajn, “ κ -Poincaré-Hopf algebra and Hopf algebroid structure of phase space from twist”, *Phys. Lett. A* **377** 2472-2476 (2013), arXiv:1303.0994.
- [23] T. Jurić, D. Kovačević, S. Meljanac, “ κ -deformed phase space, Hopf algebroid and twisting”, *SIGMA* **10** 106, 18 pages (2014), arXiv:1402.0397.
- [24] J. Lukierski, D. Meljanac, S. Meljanac, D. Pikutić and M. Woronowicz, “Lie-deformed quantum Minkowski spaces from twists: Hopf algebraic versus Hopf-algebroid approach”, *Phys. Lett. B* **777** 1-7 (2018).
- [25] J. Lukierski, S. Meljanac and M. Woronowicz, “Quantum Twist-Deformed $D = 4$ Phase Spaces with Spin Sector and Hopf Algebroid Structures”, *Phys. Lett. B* **789** 82-87 (2019), arXiv:1811.07365.