

Relational quantum computing using only maximally mixed initial qubit states

Terry Rudolph¹ and Shashank Soyuz Virmani²

¹*Dept. of Physics, Imperial College London, London SW7 2AZ, tez@imperial.ac.uk*

²*Dept. of Mathematics, Brunel University London, Kingston Ln,
London, Uxbridge UB8 3PH, shashank.virmani@brunel.ac.uk*

(Dated: January 13, 2022)

In [1] we showed that the (non-destructive) 2-qubit singlet/triplet (s/t) measurement is universal for quantum computing, given only an initial Haar-twirled qubit ensemble of the form

$$\int d\Omega U(\Omega)^{\otimes 3N} (\rho_a^{\otimes N} \otimes \rho_b^{\otimes N} \otimes \rho_c^{\otimes N}) U^\dagger(\Omega)^{\otimes 3N}. \quad (1)$$

Here N grows polynomially in the computation size. The Bloch vectors of ρ_a, ρ_b, ρ_c needed only to satisfy the constraints that they are non-zero and linearly independent. Universality of s/t measurements under other assumptions was considered recently in [2]. Here we will show that in fact replacing (1) with even a resource of only maximally mixed single-qubits suffices to make the s/t measurement universal.¹

There are two steps to the procedure. The first is to note that we can efficiently prepare a state exponentially close to the maximally mixed state over the symmetric subspace of N qubits, ρ_N^{sym} . The second will be to convert several copies of ρ_N^{sym} to (1) using a procedure known as measurement-induced relative localisation [3].

Creating the maximally mixed state in the symmetric subspace: Suppose that we start with K qubits prepared in ρ_K^{sym} . We bring in a new qubit in the maximally mixed state and randomly pick pairs of qubits² to undergo a polynomial number of s/t measurements. If we only ever find triplet outcomes, we perform an exponentially good approximate projection [2] into the state ρ_{K+1}^{sym} . This occurs with probability $P(K) = (K+2)/(2K+2) > 1/2$. If we ever find a singlet outcome we discard those two qubits and the remaining $K-1$ qubits are left in ρ_{K-1}^{sym} .

We can interpret the protocol as a 1-d random walk process where we begin at $K=1$, and have a probability $P(K)$ of stepping to the right, and $1-P(K)$ of stepping to the left. The boundary at $K=0$ is absorbing (fail, restart) and let us consider our target is to create ρ_N^{sym} for some fixed N . The solution to this problem can be found in [5]. Note that the particle must eventually be absorbed

at one or the other boundaries. Eq. (2.7) of [5] for our case yields that the probability it is absorbed at the right hand boundary is $(N+1)/2N$, i.e. slightly higher than $1/2$. Thus we have finite probability of eventual success. To ensure the resources consumed (qubits/time steps) are polynomial we need to compute the conditional mean for the number of steps before stopping (absorption at a boundary). This can be found by solving the recurrence relations (3.1)-(3.3) in [5]. For starting at $K=1$ we find the expected number of steps before absorption is $(N^2+3N-4)/6$, which grows polynomially with N . (The method we have outlined may possibly be improved using a variant, with an appropriate input state, of the *splitting protocol* of [2].)

Relative Localization: The second step is to see how two sufficiently large maximally mixed symmetric states can be converted into an ensemble equivalent to a Haar-twirled product state over pure states with fixed overlap, i.e. a state of the form:

$$\int d\Omega U(\Omega)^{\otimes 2N} (|a\rangle \langle a|^{\otimes N} \otimes |b\rangle \langle b|^{\otimes N}) U^\dagger(\Omega)^{\otimes 2N}$$

for some arbitrary (but known) $|a\rangle, |b\rangle$ with $0 < |\langle a|b\rangle|^2 < 1$.

This can be done by using a “measurement induced localization” of the relative angle between initial (mixtures of) spin coherent states, in full analogy with the cases for optical phase, Bose-Einstein condensate phase, and particle position studied in detail in [3]. (If we can localize two such ensembles then we can use the same procedure to relationally localize further ensembles to the first two).

The basic intuition is simple: we start with two sources $\rho_{N+M}^{sym} \otimes \rho_{N+M}^{sym}$, as created in the first step, and interpret each as an ensemble of $N+M$ copies of a randomly selected pure state. We pair up M of the spins from each source and perform the singlet/triplet measurement on each pair, enabling us to get a good estimate of the overlap between the two (random) pure states. We then use the remaining $2N$ qubits for computation, under the assumption that the overlap is the estimated one [6]. As we now demonstrate, a fixed overall error across the $2N$ qubits requires M to grow only polynomially in N .

Because of the collective unitary freedom, we are free to decide that the first symmetric state is actually a source of $|0\rangle^{\otimes N+M}$, and the second state $|\theta\rangle^{\otimes N+M}$ is specified by the relative angle $\theta \in [0, \pi)$ its Bloch vector makes with the first source state, where θ has p.d.f

¹ Almost any resource of qubits can be turned into such by repeatedly performing the s/t measurement, keeping singlet outcomes, and discarding a member of each pair.

² More efficient choices than random pairings exist - for example interpreting the switches of the networks in [4] as s/t measurement locations.

$\sin(\theta)/2$. An s/t measurement on $|0\rangle \otimes |\theta\rangle$ gives a triplet outcome with probability $q = (1 + \cos^2(\theta/2))/2 = (3 + \cos(\theta))/4$. The total probability over the M measurements of obtaining n_1 triplet outcomes is

$$\begin{aligned} P(n_1) &= \int_0^\pi d\theta \binom{M}{n_1} q^{n_1} (1-q)^{M-n_1} \frac{\sin(\theta)}{2} \\ &= 2 \binom{M}{n_1} \int_{1/2}^1 dq q^{n_1} (1-q)^{M-n_1} \end{aligned}$$

This has the convenient interpretation that the probability of seeing a given number of triplets is described by a Bernoulli trial with a uniformly chosen q in the interval $[1/2, 1]$. Estimating θ corresponds to estimating q given the observed M, n_1 , so we will also write $|q\rangle := |\theta\rangle$.

Considering the function

$$\begin{aligned} T(a, b) &= \int_{1/2}^1 dq q^a (1-q)^b \\ &= \frac{a!b!}{(a+b+1)!} \frac{1}{2^{a+b+1}} \sum_{j=0}^a \binom{a+b+1}{j} \end{aligned}$$

we can use standard identities (e.g [here](#)) for partial sums of binomial coefficients to see that $T(a, b)$ is exponentially close (in $M = (a+b)$) to $\left(\binom{a+b}{a}(a+b+1)\right)^{-1}$ when $a > (a+b)/2$. Applying this to $P(n_1)$ we find that it is exponentially close to $2/(M+1)$, which means that with high probability on any given run of the procedure we will observe $n_1 > M/2$ triplet outcomes, and from now on, we consider only situations where this has occurred.

The probability density of q given n_1 triplet outcomes is (over the domain $q \in [1/2, 1]$):

$$\Pr(q|n_1, M) = \frac{q^{n_1} (1-q)^{M-n_1}}{T(n_1, M-n_1)},$$

from which we wish to bound the goodness of our estimated value of q (and hence θ). The mean and variance for this inference problem are given by

$$\begin{aligned} \mu &= \frac{T(n_1+1, M-n_1)}{T(n_1, M-n_1)} \approx \frac{n_1+1}{M+2} \\ \sigma^2 &= \frac{T(n_1+2, M-n_1)}{T(n_1, M-n_1)} \approx \frac{(n_1+1)(M+1-n_1)}{(M+2)^2(M+3)}, \end{aligned}$$

where \approx denotes exponential closeness. A simple upper bound on the variance is then $\sigma^2 < 1/M$.

Now, we are roughly in the following situation: we will operate as if $q = \mu$, i.e. the state of the second N qubits is $|\mu\rangle^{\otimes N}$ (by collective rotational freedom a state in the right semicircle of the XZ plane in the Bloch sphere), but with a low probability ($\leq 1/h^2$ by the [Chebyshev inequality](#)) the actual value of q could be further than $h\sigma$ from this. In later calculations we will pick $h = M^{1/6}$. The error we want to understand will ultimately arise from the trace distance between the esti-

mated state and the actual one, and so we wish to bound this.

To make things simpler we first ask, for any pair of q_1, q_2 with a fixed value of $|q_1 - q_2|$, what is the largest possible trace distance between the corresponding quantum states $|q_1\rangle, |q_2\rangle$? Elementary considerations [7] yield:

$$\| |q_1\rangle \langle q_1| - |q_2\rangle \langle q_2| \| \leq 2\sqrt{8|q_1 - q_2|} \quad (2)$$

We can use this to bound the overall error via:

$$\begin{aligned} &\| |\mu\rangle \langle \mu|^{\otimes N} - \int dq \Pr(q|n_1, M) |q\rangle \langle q|^{\otimes N} \| \\ &\leq \int dq \Pr(q|n_1, M) \| |\mu\rangle \langle \mu|^{\otimes N} - |q\rangle \langle q|^{\otimes N} \| \\ &\leq \sqrt{N-1} \int dq \Pr(q|n_1, M) \| |\mu\rangle \langle \mu| - |q\rangle \langle q| \| \\ &\leq 2\sqrt{8(N-1)} \int dq \Pr(q|n_1, M) \sqrt{|q - \mu|} \\ &\leq 2\sqrt{8(N-1)} \sqrt{\int dq \Pr(q|n_1, M) |q - \mu|} \\ &\leq 2\sqrt{8(N-1)} \sqrt{\frac{1}{h^2} + h\sigma} \leq 2\sqrt{8(N-1)} \sqrt{\frac{2}{M^{1/3}}} \end{aligned}$$

where the first inequality is the triangle inequality, the second is because for pure states it holds that $\|\psi^{\otimes N} - \phi^{\otimes N}\| \leq \sqrt{N-1}\|\psi - \phi\|$, the third is from the bound (2), the fourth is concavity of the square root, the fifth is from the largest probabilities (and $|q - \mu|$ values) consistent with the Chebyshev inequality, and the last is obtained by using $\sigma \leq 1/\sqrt{M}$ and picking $h = M^{1/6}$.

We deduce that given target overall error of ϵ , we can choose $M \sim (N/\epsilon^2)^3$, which is a polynomial cost.

Acknowledgements: SSV acknowledges thought provoking discussions with Nihaal Virmani. TR acknowledges interaction with one of two indistinguishable Virmanicles.

-
- [1] T. Rudolph and S. S. Virmani, *New Journal of Physics* **7**, 228 (2005).
 - [2] M. H. Freedman, M. B. Hastings, and M. S. Zini, (2021), arXiv:2105.04649 [quant-ph].
 - [3] H. Cable, P. L. Knight, and T. Rudolph, *Phys. Rev. A* **71**, 042107 (2005).
 - [4] A. Czumaj, in *Proceedings of STOC '15*, STOC '15 (2015) p. 703–712.
 - [5] M. A. El-Shehawey, *J. Phys A* **33**, 9005 (2000).
 - [6] We remark that the relative localisation will happen more efficiently using approximate total angular measurement protocols of [2] together with some form of global angular momentum inference scheme (see e.g. [8]).
 - [7] In brief: the state $|q\rangle$ has z component of its Bloch vector given by $z = 4q - 3$, so a given value of $|q_1 - q_2|$ constrains the Bloch vectors of $|q_1\rangle, |q_2\rangle$ to have a projection on the

z-axis to a fixed interval $4|q_1 - q_2|$. Positioning one end of this interval at either the north or south poles of the Bloch sphere yields the largest possible trace distance consistent with this projected value.

- [8] M. Fanizza, M. Rosati, M. Skotiniotis, J. Calsamiglia, and V. Giovannetti, Phys. Rev. Lett. **124**, 060503 (2020).