

Photon quantum mechanics in real Hilbert space

Margaret Hawton

Department of Physics, Lakehead University, Thunder Bay, ON, Canada, P7B 5E1

Classically, electromagnetic pulses are described by real fields that couple to charged matter and propagate causally. We will show here that real fields of the form used in standard classical electromagnetic theory have a quantum mechanical interpretation in which the probability density for a photon to be at \mathbf{x} is positive definite and operators representing all of the standard physical observables exist. A covariant alternative to the $\omega_k^{1/2}$ dependence that appears in most (but not all) presentations of quantum optics and quantum field theory is presented and real scalar one photon advanced and retarded potentials are derived.

I. INTRODUCTION

This work was originally motivated by mathematical proofs that a positive frequency field cannot be confined to a finite region of space. According to the Hegerfeldt theorem a positive frequency field localized in a finite region for an instant spreads immediately throughout space [1]. It has been shown explicitly in the case of one dimensional square wells [2] and for three dimensional position eigenvectors [3, 4] that this instantaneous localization is only apparent since it is due to destructive interference of intrinsically nonlocal counterpropagating waves. In algebraic quantum field theory (QFT), the Reeh-Schlieder theorem states that there are no local annihilation or creation operators [5]. However, confinement of real fields to a finite region is not a problem in classical electromagnetism (EM). It will be proved here that this use of real fields can be extended to photon quantum mechanics (QM).

The QM of electrons and other Fermions is well understood but a consistent first quantized theory of the photon has been elusive. Photons have two properties not shared with electrons that have made derivation of photon QM difficult - they are neutral and massless. While fields describing charged particles are intrinsically complex, neutral particles should be described by real fields. Reality of the photon wave function ensures that photons and antiphotons, being indistinguishable, are equally probable and that, after second quantization, their field operators become Hermitian. This property is problematic in a first quantized theory since the standard relativistic scalar product is zero for neutral particles. Also, while the Wigner little group describing massive particles is the set of spatial rotations, the Wigner little group for massless particles is cylindrically symmetrical. In their seminal paper titled "Localized states of elementary systems", Newton and Wigner assumed invariance under spherically symmetrical rotations and concluded that "for equations with zero mass .. with spin 1 (i.e. Maxwell's equations) we found that no localized states in the above sense exist. This is an unsatisfactory .. feature of our work" [6]. Both of these difficulties have been overcome [3, 4, 7-10] but here we will extend this work by formulating photon QM in terms of physically correct real fields and show that this leads to a significant

simplification of the mathematics.

In field theory, particles that transform into themselves are represented by real fields and Hermitian field operators [11]. These real fields can be written as linear combinations of the positive frequency particle terms, A^+ , and negative frequency antiparticle terms, A^- . The real fields $A_c = (A^+ + A^-)/\sqrt{2}$ and $A_s = (A^+ - A^-)/\sqrt{2}i$ are even and odd respectively under particle/antiparticle exchange (charge conjugation) where the subscripts c and s denote Fourier cosine and sine series respectively. The complex positive frequency function $A^+ = A_c + iA_s$ describes both of these potentials, so it will be used here to simplify the mathematics.

In Section II the photon equations of motion will be derived from the standard Lagrangian. The fields will be assumed to be real but for mathematical convenience these real fields will be identified with the real and imaginary parts of a complex positive frequency field. The zeroth component of the standard relativistic four-current will be interpreted as a positive definite photon number density and used to define a scalar product. Interaction with polarized matter will be included so that these equations can be applied to photon propagation in a nonabsorptive medium and emission and absorption of photons by localized sources and sinks. In Section III the real Hilbert space will be defined and operators for all the usual physical observables, including position, will be reviewed to provide a complete first quantized description of single-photon states. In Section IV the fields will be second quantized and in the final Section we will summarize and conclude.

II. REAL AND COMPLEX FIELDS, LAGRANGIAN AND SCALAR PRODUCT

In this Section four-vector notation and complex fields whose real and imaginary parts are even and odd under QFT charge conjugation will be defined. The photon equations of motion, four-current and positive definite number density will be derived from the real standard Lagrangian written in terms of complex fields. A scalar product will be defined in position space and Fourier transformed to momentum space to complete the Hilbert space.

Relativistic notation and SI units will be used. The contravariant space-time, wave vector and momentum four-vectors are $x = x^\mu = (ct, \mathbf{x})$, $k = (\omega_k/c, \mathbf{k})$ and $p = \hbar k$ where $kx = \omega_k t - \mathbf{k} \cdot \mathbf{x}$, the four-gradient is $\partial = (\partial_{ct}, -\nabla)$, the four-potential is $A(t, \mathbf{x}) = A^\mu = \left(\frac{\phi}{c}, \mathbf{A}\right)$ or $a(t, \mathbf{k}) = (a_0, \mathbf{a})$ and $a_\lambda(\mathbf{k})$ will denote a Lorentz invariant scalar describing a state with definite helicity, λ . The covariant four-vector corresponding to $U^\mu = (U_0, \mathbf{U})$ is $U_\mu = g_{\mu\nu}U^\nu = (U_0, -\mathbf{U})$ where $g_{\mu\nu} = g^{\mu\nu}$ is a 4×4 matrix with diagonal $(1, -1, -1, -1)$.

Positive and negative frequency four-potentials will be used here to describe photons and antiphotons respectively. In Fourier expanded form

$$A^+(x) = \sqrt{\frac{\hbar}{\epsilon_0}} \int_t \frac{d\mathbf{k}}{(2\pi)^3 \omega_k} a(\mathbf{k}) e^{-ikx} \quad (1)$$

$$A^-(x) = A^{+*}(x) \quad (2)$$

where $d\mathbf{k} \equiv d^3k$ is an infinitesimal volume in \mathbf{k} -space. This form of was selected because $\lim_{V \rightarrow \infty} \Delta\mathbf{n}/V = d\mathbf{k}/(2\pi)^3$ where $\Delta\mathbf{n}$ is the number of states and $\int d^4k \delta(\omega_k^2/c^2 - |\mathbf{k}|^2) = \int_t \frac{d\mathbf{k}}{(2\pi)^3 2\omega_k/c}$. Since kx and $\int_t \frac{d\mathbf{k}}{(2\pi)^3 \omega_k}$ are invariants, if $A(x)$ is a four-vector then $a(\mathbf{k})$ is a four-vector. For complex Fourier coefficients

$$a(\mathbf{k}) = \frac{a_R(\mathbf{k}) + ia_I(\mathbf{k})}{\sqrt{2}} \quad (3)$$

the positive frequency four-potential can be written as

$$A^+(x) = A_c(x) + iA_s(x) \quad (4)$$

where

$$A_c(x) = \frac{A^+(x) + A^-(x)}{\sqrt{2}} \quad (5)$$

$$= \sqrt{\frac{2\hbar}{\epsilon_0}} \int_t \frac{d\mathbf{k}}{(2\pi)^3 \omega_k} a_R(\mathbf{k}) \cos(kx),$$

$$A_s(x) = \frac{A^+(x) - A^-(x)}{\sqrt{2}i} \quad (6)$$

$$= \sqrt{\frac{2\hbar}{\epsilon_0}} \int_t \frac{d\mathbf{k}}{(2\pi)^3 \omega_k} a_I(\mathbf{k}) \sin(kx)$$

are even and odd with respect to interchange of photons and antiphotons (QFT charge conjugation). The real field is

$$2A(x) = A^+(x) + A^-(x) = A_c(x) + A_s(x). \quad (7)$$

The factor 2 is included for consistency with the definitions in QFT where division is by $2\omega_k$. The positive frequency electric and magnetic fields are $\mathbf{E}^+ = -\partial_t \mathbf{A}^+ - \nabla \phi^+$ and $\mathbf{B}^+ = \nabla \times \mathbf{A}^+$ and the antisymmetric Faraday tensor is

$$\mathcal{F}^{+\mu\nu} = \partial^\mu A^{+\nu} - \partial^\nu A^{+\mu} \quad (8)$$

where $\mathcal{F}^{+00} = \mathcal{F}^{+ii} = 0$, $\mathcal{F}^{+i0} = -\mathcal{F}^{+0i} = E_i^+/c$, $\mathcal{F}^{+ij} = -\mathcal{F}^{+ji} = \epsilon_{ijk} B_k^+$ and ϵ_{ijk} is the Levi-Civita symbol.

The Lagrangian describing two real fields can be written in complex form provided this field and its complex conjugate are treated as formally independent [12]. In the presence of a matter four-current density $J_m^{+\nu} + c.c.$, the real Lagrangian density will be written as

$$\mathcal{L} = \mathcal{L}_{std} + \mathcal{L}_{int} \quad (9)$$

$$\mathcal{L}_{std} = -\epsilon_0 (\mathbf{E}^+ \cdot \mathbf{E}^{+*} - c^2 \mathbf{B}^+ \cdot \mathbf{B}^{+*}) \quad (10)$$

$$= -\frac{1}{2} \epsilon_0 c^2 \mathcal{F}_{\mu\nu}^+ \mathcal{F}^{+\mu\nu*}$$

$$\mathcal{L}_{int} = -J_m^{+\nu*} A_\nu^+ - J_m^{+\nu} A_\nu^{+*} \quad (11)$$

where c is the speed of light and ϵ_0 is the dielectric permittivity. The Lagrange equations of motion are then

$$\partial_\mu \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu A_\nu^{+*})} \right] = \frac{\partial \mathcal{L}}{\partial A_\nu^{+*}} \quad (12)$$

where the momentum conjugate to A_ν^{+*} is

$$\Pi^{+\mu\nu} \equiv \frac{\partial \mathcal{L}}{\partial (\partial_\mu A_\nu^{+*})} = \epsilon_0 c^2 \mathcal{F}^{+\mu\nu}. \quad (13)$$

Eq. (12) then gives

$$\epsilon_0 c^2 \partial_\mu \mathcal{F}^{+\mu\nu} = \mu_0 J_m^{+\nu}, \quad (14)$$

that can be written as the Maxwell equations (ME)

$$\nabla \cdot \mathbf{D}^+ = \rho_m^+, \quad \nabla \times \mathbf{H}^+ - \partial_t \mathbf{D}^+ = \mathbf{J}_m^+ \quad (15)$$

where $\mathbf{D}^+ = \epsilon_0 \mathbf{E}^+$, $\mathbf{B}^+ = \mu_0 \mathbf{H}^+$ and $\mu_0 \epsilon_0 = 1/c^2$. Substitution of (8) in (14) gives the wave equation

$$\partial_\mu \partial^\mu A^{+\nu} - \nabla^2 A^{+\nu} = \partial^\nu \Lambda^+ + \mu_0 J_m^{+\nu} \quad (16)$$

where

$$\partial_\mu \partial^\mu = \partial_{ct}^2 - \nabla^2 \equiv \square. \quad (17)$$

The gauge is determined by

$$\Lambda^+ = c^{-2} \partial_t \phi^+ + \nabla \cdot \mathbf{A}^+. \quad (18)$$

In the Coulomb gauge $\nabla \cdot \mathbf{A}^+ = 0$, so there are no longitudinal modes and the scalar field satisfying $\nabla^2 \phi^+ = -\rho^+/\epsilon_0$ responds instantaneously to changes in charge density. Only the transverse modes propagate at the speed of light and are second quantized in quantum electrodynamics (QED) to allow creation and annihilation of physical photons. In the Lorenz gauge $\Lambda = 0$ inserted into (16) gives $\partial_\mu \partial^\mu A^{+\nu} = \mu_0 J_m^{+\nu}$. In this gauge all four components of A describing the scalar, longitudinal and transverse photon modes propagate at the speed of light and are second quantized in QED. Each complex equation in this paragraph is equivalent to two real equations; one for the even potentials and one for the odd potentials.

Using these complex fields a positive definite photon number density and a scalar product can be derived from the global phase change $A^+ \rightarrow e^{i\alpha} A^+$, $A^{+*} \rightarrow e^{-i\alpha} A^{+*}$ that is a symmetry of the free space Lagrangian \mathcal{L}_{std} . For an infinitesimal change in A , $\delta A^+ \simeq i\alpha A^+$ and $\delta A^{+*} \simeq -i\alpha A^{+*}$ so, using $\Pi^{+\mu\nu} = \epsilon_0 c^2 \mathcal{F}^{+\mu\nu}$, the Noether four-current density becomes

$$J^\mu(x) = \frac{i\epsilon_0 c^2}{2\hbar} [\mathcal{F}^{+\mu\nu*} A_\nu^+ - \mathcal{F}^{+\mu\nu} A_\nu^{+*}]. \quad (19)$$

Since $\partial^\mu \cos(kx) = -k^\mu \sin(kx)$ and $\partial_\mu \sin(kx) = k^\mu \cos(kx)$, differentiation of the complex potential A^+ with respect to any component of x^μ yields a factor $-i$ and changes $\cos(kx)$ to $-\sin(kx)$ and $\sin(kx)$ to $\cos(kx)$ so that $A_c + iA_s \rightarrow i\mathcal{F}_s + \mathcal{F}_c$ gives $\frac{1}{2}\mathcal{F}^{+*} A^+ + c.c. = \mathcal{F}_c A_c + \mathcal{F}_s A_s$ and

$$J^\mu(x) = \frac{\epsilon_0 c^2}{\hbar} (\mathcal{F}_c^{\mu\nu} A_{c\nu} + \mathcal{F}_s^{\mu\nu} A_{s\nu}). \quad (20)$$

If $J_m = 0$, J satisfies a continuity equation and the spatial integral of the number density $J^0(x)$ is conserved. The four-current density $J^\mu(x) \propto \mathcal{F}^{\mu\nu}(x) A_\nu(x)$ was first obtained in [13] where it was used to derive a Hermitian number density operator.

The four-current (19) or (20) is not gauge invariant due to its dependence on A . In the Coulomb gauge only transverse waves propagate, while in the Lorenz gauge longitudinal and transverse photons exist, but their contributions to the scalar product cancel in free space [4]. Writing $\mathcal{F}^{+0\nu}$ as an electric field and including only transverse photons with helicity $\lambda = \pm 1$, the scalar product at a fixed time t will be defined as

$$(A_1, A_2)_t \equiv \frac{i\epsilon_0}{2\hbar} \sum_{\lambda=\pm 1} \int_t dx \mathbf{A}_{1\lambda}^{+*}(x) \cdot \mathbf{E}_{2\lambda}^+(x) + c.c. \quad (21)$$

$$= \frac{\epsilon_0}{\hbar} \sum_{\lambda=\pm 1} \int_t dx [\mathbf{A}_{1c\lambda}(x) \cdot \mathbf{E}_{2c\lambda}(x) + \mathbf{A}_{1s\lambda}(x) \cdot \mathbf{E}_{2s\lambda}(x)]. \quad (22)$$

If $A_2 = A_1$ (21) reduces to the spatial integral of number density $J^0(t, \mathbf{x})$. Only free transverse photons are counted. In the presence of sources and sinks this photon number is not conserved.

Fourier transforms and the bra-ket notation will be defined nonrelativistically as in Schrödinger QM. At a fixed time

$$f(\mathbf{x}) \equiv \int \frac{d\mathbf{k}}{(2\pi)^3} \tilde{f}(\mathbf{k}) \exp(i\mathbf{k} \cdot \mathbf{x}), \quad (23)$$

$$\tilde{f}(\mathbf{k}) \equiv \int d\mathbf{x} f(\mathbf{x}) \exp(-i\mathbf{k} \cdot \mathbf{x}) \text{ and} \quad (24)$$

$$\langle f|g \rangle \equiv \int d\mathbf{x} f^*(\mathbf{x}) g(\mathbf{x}) = \int \frac{d\mathbf{k}}{(2\pi)^3} \tilde{f}^*(\mathbf{k}) \tilde{g}(\mathbf{k}) \quad (25)$$

$$\langle \mathbf{f} \cdot \mathbf{g} \rangle \equiv \sum_{j=1,3} \langle \mathbf{f}_j | \mathbf{g}_j \rangle. \quad (26)$$

Using this notation the scalar product (21) is

$$(A_1, A_2)_t = \frac{i\epsilon_0}{2\hbar} \sum_{\lambda=\pm 1} \langle \mathbf{A}_{1\lambda}^+ \cdot \mathbf{E}_{2\lambda}^+ \rangle + c.c. \quad (27)$$

$$= \frac{\epsilon_0}{\hbar} \sum_{\lambda=\pm 1} (\langle \mathbf{A}_{1c\lambda} \cdot \mathbf{E}_{2c\lambda} \rangle + \langle \mathbf{A}_{1s\lambda} \cdot \mathbf{E}_{2s\lambda} \rangle) \quad (28)$$

where from (1), using $a = (a_0, \mathbf{a})$ and $\mathbf{E}_\lambda = -\partial_t \mathbf{A}_\lambda$,

$$\mathbf{E}_\lambda^+(x) = i\sqrt{\frac{\hbar}{\epsilon_0}} \int_t \frac{d\mathbf{k}}{(2\pi)^3} \mathbf{a}_\lambda(\mathbf{k}) e^{-i\mathbf{k}x}. \quad (29)$$

By inspection of (1) and (29) the Fourier transforms of \mathbf{A}_λ and \mathbf{E}_λ are $\sqrt{\frac{\hbar}{\epsilon_0}} \mathbf{a}_\lambda(\mathbf{k}) / \omega_k$ and $i\sqrt{\frac{\hbar}{\epsilon_0}} \mathbf{a}_\lambda(\mathbf{k})$ respectively so, using the Parseval-Pancherel identity (25),

$$(A_1, A_2)_t = \sum_{\lambda=\pm 1} \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{\mathbf{a}_{1\lambda}(\mathbf{k})}{\omega_k} \cdot \mathbf{a}_{2\lambda}(\mathbf{k}) \quad (30)$$

$$= \sum_{\lambda=\pm 1} \left\langle \frac{\mathbf{a}_{1\lambda}}{\omega_k} \cdot \mathbf{a}_{2\lambda} \right\rangle. \quad (31)$$

Inspection of (21) to (31) shows that these expressions for the scalar product involve both the potential and the electric field rather than a single function. In the next Section we will show that $\langle \mathbf{E} \cdot \mathbf{A} \rangle$ implies a biorthogonal basis so this formulation is called biorthogonal QM [3, 14]. The relationship between the notation used here and that in [4] is that here $a(\mathbf{k})$ is a Fourier transform while in [4] $\mathbf{c}(\mathbf{k})$ is the probability amplitude for a covariantly normalized plane wave.

III. HILBERT SPACE AND OBSERVABLES

In this Section the Hilbert space and the momentum, position and angular momentum operators and their eigenvectors are defined. Covariant normalization that leads to expressions of the classical form is used. The probability amplitude to find a photon at \mathbf{x} on the t -hyperplane is calculated and it is verified that the Born rule is satisfied.

The real Hilbert space is the vector space of all A_c and A_s and their derivatives with the scalar product (28). For mathematical convenience the positive frequency potentials A^+ and scalar product (27) will be used. Since $A^{+\mu}(\mathbf{x}, t)$ is a four-vector, it follows from (1) that $a^\mu(\mathbf{k}, t)$ must be a four-vector. With the mutually orthogonal polarization unit vectors e^μ defined such that 0 is time-like, 1 and 2 are transverse and 3 is longitudinal, $e_0 = n^\mu = (1, 0, 0, 0)$, $\mathbf{e}_3(\mathbf{k}) = \mathbf{e}_\mathbf{k} = \mathbf{k}/|\mathbf{k}|$ and the definite helicity transverse unit vectors are

$$\mathbf{e}_\lambda = \frac{1}{\sqrt{2}} (\mathbf{e}_\theta + i\lambda \mathbf{e}_\phi) \quad (32)$$

for $\lambda = \pm 1$ where \mathbf{e}_θ , \mathbf{e}_ϕ and $\mathbf{e}_\mathbf{k}$ are \mathbf{k} -space spherical polar unit vectors on the t -hyperplane. Since e^μ is a

four-vector its coefficient in any expression for a^μ should be a Lorentz invariant scalar.

Momentum is an observable. It can be verified by substitution in (30) that the plane waves with definite momentum $\hbar\mathbf{k}'$ defined covariantly as

$$\mathbf{a}_{\mathbf{k}'\lambda'}(\mathbf{k}) = (2\pi)^3 \omega_k \delta(\mathbf{k} - \mathbf{k}') \mathbf{e}_{\lambda'}(\mathbf{k}') \quad (33)$$

are biorthogonal in the sense that

$$(A_{\mathbf{k}\lambda}, A_{\mathbf{k}'\lambda'}) = \delta_{\lambda\lambda'} (2\pi)^3 \omega_k \delta(\mathbf{k} - \mathbf{k}'). \quad (34)$$

This normalization of the plane wave basis is invariant as can be seen from $\int \frac{d\mathbf{k}}{\omega_k} \omega_k \delta(\mathbf{k} - \mathbf{k}') = 1$. In position space in the Heisenberg picture (HP) (1) gives

$$\mathbf{A}_{\mathbf{k}\lambda}^+(\mathbf{x}, t) = \sqrt{\frac{\hbar}{\epsilon_0}} e^{-ikx} \mathbf{e}_\lambda(\mathbf{k}). \quad (35)$$

In a general state A the probability amplitude for wave vector \mathbf{k} is

$$(A_{\mathbf{k}\lambda}, A) = a_\lambda(\mathbf{k}) \quad (36)$$

If ω_k is replaced with $\omega_k^{1/2}$ in (33) the noncovariant Newton-Wigner [6] normalization, $(A_{\mathbf{k}\lambda}, A_{\mathbf{k}'\lambda'})_{NW} = \delta_{\lambda\lambda'} (2\pi)^3 \delta(\mathbf{k} - \mathbf{k}')$, is obtained. The position space plane waves (35) are eigenvectors of the four-momentum operator

$$\hat{P} = (\hat{p}_0, \hat{\mathbf{p}}) = \left(\frac{1}{c} \hat{H}, -i\hbar\nabla \right) \quad (37)$$

with eigenvalues $\hbar(\omega_{k_j}/c, \mathbf{k})$. The Hamiltonian operator

$$\hat{H} = \hbar c \sqrt{-\nabla^2} \quad (38)$$

with eigenvalues $\hbar\omega_k$ generates unitary transformations according to the Schrödinger equation $i\hbar\partial_t A^+(t) = \hat{H}A^+(t)$. In \mathbf{k} -space the four momentum operator is $\hat{P} = (\hbar kc, \hbar\mathbf{k})$.

Position is also an observable. According to (24) the Fourier transform of the localized state $\delta(\mathbf{x} - \mathbf{x}')$ is the plane wave $\exp(-i\mathbf{k} \cdot \mathbf{x}')$ so covariant HP photon position eigenvectors should be of the form

$$\mathbf{a}_{x'\lambda'}(\mathbf{k}) = \mathbf{e}_{\lambda'}(\mathbf{k}) e^{ikx'}. \quad (39)$$

The projection of an arbitrary state A onto the $A_{x\lambda}$ basis,

$$\phi_\lambda^+(x) = (A_{x\lambda}, A)_t = \int_t \frac{d\mathbf{k}}{(2\pi)^3 \omega_k} a_\lambda(\mathbf{k}) e^{-ikx}, \quad (40)$$

has the mathematical form of a Lorentz invariant scalar potential that satisfies the zero mass Klein-Gordon (KG) equation. The photon position operator with commuting components can be derived by rotating $\mathbf{e}_1 + i\lambda\mathbf{e}_2$ about \mathbf{e}_2 by θ , then about \mathbf{e}_3 by ϕ to give $\mathbf{e}_\theta + i\lambda\mathbf{e}_\phi$ using the operator \hat{R} so that $\mathbf{e}_\theta + i\lambda\mathbf{e}_\phi = \hat{R}(\mathbf{e}_1 + i\lambda\mathbf{e}_2)$ and

$i\partial_{\mathbf{k}}$ transforms to $\hat{\mathbf{x}} = \hat{R}i\partial_{\mathbf{k}}\hat{R}^{-1}$ [15]. Alternatively it can be obtained by covariant differentiation [20]. It was originally obtained by brute force subtraction of the \mathbf{k} -space gradient of $\mathbf{e}_{\lambda_j}(\mathbf{k})$ to give [7]

$$\hat{\mathbf{x}} = i\partial_{\mathbf{k}} - i\alpha \frac{\mathbf{k}}{|\mathbf{k}|^2} + \frac{1}{|\mathbf{k}|^2} \mathbf{k} \times \hat{\mathbf{S}} - \hat{\lambda} \frac{\cos\theta}{k \sin\theta} \mathbf{e}_\phi \quad (41)$$

for position eigenvectors proportional to $(\omega_k)^\alpha$. The helicity λ photon position eigenvectors have a definite component of total angular momentum in the fixed but arbitrary direction \mathbf{e}_3 with indefinite spin and orbital contributions [15]. The total angular momentum operator is

$$\hat{\mathbf{J}} = \hat{\mathbf{x}} \times \hat{\mathbf{P}} + \hat{\mathbf{J}}_{int}, \quad (42)$$

$$\hat{\mathbf{J}}_{int} = \hbar\lambda \left(\frac{\cos\theta}{\sin\theta} \mathbf{e}_\theta + \mathbf{e}_\mathbf{k} \right). \quad (43)$$

where $\hat{\mathbf{J}}_{int}$ is the internal angular momentum operator and $\hbar\hat{\mathbf{x}} \times \hat{\mathbf{k}}$ describes external angular momentum. The position and angular momentum operators are reviewed in more detail in [4, 15] where rotation about \mathbf{k} through an Euler angle that gives a more general expression for \mathbf{J} is included.

Setting $\mathbf{a}_\lambda(\mathbf{k})$ equal to an $\alpha = 0$ position eigenvector $\mathbf{a}_{x'\lambda'}(\mathbf{k})$ in (40) with $\Delta t \equiv t - t'$ and $r \equiv |\mathbf{x} - \mathbf{x}'|$ an explicit expression for its time evolution can be obtained by integrating (40) to give

$$\phi_{x'\lambda'}^+(x) = \frac{1}{4\pi^2 r} \sum_{\gamma=\pm} \left[i\gamma\pi\delta(r - \gamma c\Delta t) + P \left(\frac{1}{r - \gamma c\Delta t} \right) \right] \quad (44)$$

for $\lambda' = \pm 1$. Both the real and imaginary parts of (44) satisfy the photon wave equation, but only its imaginary part is odd under QFT charge conjugation and couples to charged matter. Thus, if the real part of $\phi_\lambda(t, \mathbf{x})$ is discarded on physical grounds, (44) is reduced to its odd sine series imaginary part,

$$\phi_{sx'\lambda'}(x) = \frac{1}{4\pi r} [\delta(r + c\Delta t) - \delta(r - c\Delta t)] \quad (45)$$

that satisfies the homogeneous wave equation $\square\phi_\lambda(x) = 0$. Schweber [17] inverted \square and found that the unique Green's function solving $\square\phi_\lambda(x) = \delta(\mathbf{x} - \mathbf{x}')\delta(t - t')$ is $\frac{1}{4\pi r} [\delta(r + \Delta t) + \delta(r - c\Delta t)]$ where $t' = t - r < 0$ is the retarded time and $t' = t + r > 0$ is the advanced time. He concluded that the retarded potential is determined by boundary conditions and equals the sum of his unique contour integral independent Green's function and a solution to the homogeneous wave equations of the form (45). This retarded potential solution is important in classical EM and, according to (45), it can be applied to photon QM with the significant advantage that $\phi_\lambda(x)$ is a scalar.

In Schrödinger quantum mechanics, the non-square integrable δ -functions satisfy a completeness relation

and form a very convenient basis [18] so δ -localized photon states will be defined here. The usual approach in photon QM and QED is to introduce a factor $\omega_k^{1/2}$ into (39) to give $\mathbf{a}_{x'\lambda'}(\mathbf{k}) = \omega_k^{1/2} \mathbf{e}_{\lambda'}(\mathbf{k}') e^{ikx'}$ and the Newton-Wigner-like noncovariant normalization $(A_{\mathbf{x}\lambda}, A_{\mathbf{x}'\lambda'})_{NW} = \delta_{\lambda\lambda'} \delta(\mathbf{x} - \mathbf{x}')$. With this choice $\phi_\lambda^+(x) = \int_t \frac{d\mathbf{k}}{(2\pi)^3} \omega_k^{-1/2} a_\lambda(\mathbf{k}) e^{-ikx}$ is not covariant. Here, to retain covariance and avoid the inconvenient factor $\omega_k^{1/2}$, the time derivative of (40) that is the zeroth component of a four-vector will be used to define the localized basis. Since

$$\psi_\lambda^+(x) = i\partial_t \phi_\lambda^+(x) = \int_t \frac{d\mathbf{k}}{(2\pi)^3} a_\lambda(\mathbf{k}) e^{-ikx}, \quad (46)$$

with $kx = \omega_k t - \mathbf{k} \cdot \mathbf{x}$, for $a_\lambda(\mathbf{k}) = a_{x'\lambda'}(\mathbf{k})$, at $t = t'$,

$$\psi_{x'\lambda}^+(x) = \delta(\mathbf{x} - \mathbf{x}'). \quad (47)$$

Using (30) and expanding ψ_λ^+ in the δ -basis as

$$\psi_\lambda^+(t, \mathbf{x}) = \int_t d\mathbf{x}' \delta(\mathbf{x} - \mathbf{x}') \psi_\lambda^+(\mathbf{x}'), \quad (48)$$

it can be seen that $\psi_\lambda^+(x)$ is the probability amplitude for a photon to be in the state $\delta(\mathbf{x} - \mathbf{x}')$ on the t -hyperplane. (Only its odd sine-series part is truly localized.) Its integrated squared norm,

$$\begin{aligned} (A, A) &= \sum_{\lambda=\pm 1} \int d\mathbf{x} |\psi_\lambda^+(x)|^2 \\ &= \sum_{\lambda=\pm 1} \int d\mathbf{x} [|\psi_{c\lambda}(x)|^2 + |\psi_{s\lambda}(x)|^2], \end{aligned} \quad (49)$$

satisfies the Born rule. $\psi_{c\lambda}(x)$ and $\psi_{s\lambda}(x)$ are the real probability amplitudes for a λ -helicity photon to be at \mathbf{x} on the t -hyperplane. The λ -helicity \mathbf{x} -space probability density is

$$\rho_\lambda^\epsilon(x) = |\psi_{c\lambda}(x)|^2 + |\psi_{s\lambda}(x)|^2. \quad (50)$$

If (A, A) is finite, A is normalizable as $(A, A) = 1$. Since $a_\lambda(\mathbf{k})$ is the Fourier transform of $\psi_\lambda(x)$, using (25) the \mathbf{k} -space Born rule is

$$(A, A) = \sum_\lambda \int \frac{d\mathbf{k}}{(2\pi)^3} [|a_{c\lambda}(\mathbf{k})|^2 + |a_{s\lambda}(\mathbf{k})|^2]. \quad (51)$$

Covariance was maintained here by defining $\psi_\lambda^+(t, \mathbf{x})$ as the time-like component of a four-vector.

Only the odd $\psi_{s\lambda}$ density is localizable in a finite region and has the correct transformation properties under QFT charge conjugation [11]. The cosine solutions to the wave equation have been retained in (49) to (51) for generality because they satisfy ME and the requirement that A should be odd is not usually imposed.

Quantum mechanics requires state vectors to describe physical systems and operators representing observables

such that the only possible result of a measurement is one of their eigenvalues [18]. Eqs. (46) to (51) provide a scalar Schrödinger-like description of a photon with helicity λ in which $\psi_\lambda^+(t, \mathbf{x})$ is the probability amplitude for a photon to be at \mathbf{x} on the t -hyperplane and its FT $a_\lambda(\mathbf{k})$ is the probability amplitude for it to have momentum $\hbar\mathbf{k}$.

IV. SECOND QUANTIZATION

A first quantized photon cannot be created or destroyed - creation and annihilation of photons and the description of n -photon states requires second quantization. In QFT, QED and QO fields are second quantized by raising them to the status of operators. For an arbitrary first quantized state in which (1) and (7) are generalized to include a factor ω_k^α to accommodate the factor $\omega_k^{1/2}$ commonly used,

$$\widehat{\mathbf{A}}(x) = \sqrt{\frac{\hbar}{\epsilon_0}} \sum_{\lambda=\pm 1} \int_t \frac{d\mathbf{k}}{(2\pi)^3} (2\omega_k)^{\alpha-1} \left(\widehat{a}_{\mathbf{k}\lambda} e^{-ikx} + \widehat{a}_{\mathbf{k}\lambda}^\dagger e^{ikx} \right), \quad (52)$$

where the operator $\widehat{a}_{\mathbf{k}\lambda}$ annihilates a photon with wave vector \mathbf{k} and helicity λ and $\widehat{a}_{\mathbf{k}\lambda}^\dagger$ creates one. For plane wave states satisfying the commutation relations

$$\begin{aligned} [\widehat{a}_\lambda(\mathbf{k}), \widehat{a}_{\lambda'}(\mathbf{q})] &= 0, \quad [\widehat{a}_\lambda^\dagger(\mathbf{k}), \widehat{a}_{\lambda'}^\dagger(\mathbf{q})] = 0, \\ [\widehat{a}_\lambda(\mathbf{k}), \widehat{a}_{\lambda'}^\dagger(\mathbf{q})] &= \delta_{\lambda\lambda'} (2\pi)^3 (2\omega_k)^{1-2\alpha} \delta(\mathbf{k} - \mathbf{q}) \end{aligned} \quad (53)$$

it can be verified by substitution that the operators $\widehat{\mathbf{A}}_\lambda(x)$ and $\widehat{\mathbf{E}}_\lambda(x) = -\partial \widehat{\mathbf{A}}_\lambda(x)$ satisfy the commutation relations

$$\sum_{j=1,3} \left[\widehat{A}_{\lambda j}(t, \mathbf{x}), \widehat{E}_{\lambda' j}(t, \mathbf{x}') \right] = -\frac{i\hbar}{\epsilon_0} \delta_{\lambda\lambda'} \delta(\mathbf{x} - \mathbf{x}'). \quad (54)$$

on the t -hyperplane. The usual text book choice is $\alpha = \frac{1}{2}$ but $\alpha = 0$ for which (52) and (53) are covariant is used here and in [16].

In QED microcausality is a consequence of the commutation relations (54). (On a general spacelike hyperplane t is not necessarily equal to t' but time ordering is observer dependent.) Defining the one photon states

$$|\mathbf{A}_{x\lambda}\rangle = \widehat{\mathbf{A}}_{x\lambda} |0\rangle, \quad |\mathbf{E}_{x\lambda}\rangle = \widehat{\mathbf{E}}_{x\lambda} |0\rangle, \quad (55)$$

the vacuum expectation value of (54) multiplied by $i\epsilon_0/2\hbar$ is

$$\psi_{y\lambda}(x) = \frac{i\epsilon_0}{2\hbar} [\langle \mathbf{A}_{x\lambda} \cdot \mathbf{E}_{x'\lambda} \rangle + \langle \mathbf{A}_{x'\lambda} \cdot \mathbf{E}_{x\lambda} \rangle]. \quad (56)$$

It implies that creation at (t', \mathbf{x}') followed by annihilation at (t, \mathbf{x}) and creation at (t, \mathbf{x}) followed by annihilation at (t', \mathbf{x}') are equally likely. This provides a physical interpretation of the real probability amplitude and implies that we don't know whether the photon was created or destroyed at (t, \mathbf{x}) as discussed in [3, 4].

V. SUMMARY AND CONCLUSION

In its covariant $\alpha = 0$ version, photon quantum mechanics as described here preserves the classical form of the EM potential and fields when first quantized and second quantized. Only the interpretation need be changed - from real observable classical fields, to probability amplitudes, and then to operators that create and annihilate photons. The real potentials are even and odd under QFT charge conjugation, but only those that are odd can be localized in a finite region and coupled to charged matter. These even and odd fields are real and imagi-

nary parts of a positive frequency field whose use simplifies the mathematics and facilitates use of the standard Lagrangian and relativistic scalar product. Propagation of finite pulses is as in classical EM and mathematical techniques such as finite difference time domain (FDTD) [19] developed to handle problems in classical EM theory can be applied directly to single photons. Projection onto momentum and position bases gives a covariant Schrödinger-like description of photon QM. Eqs. (40) and (46) provide a new scalar description of single photon states with a well defined physical interpretation that may prove to be useful in applications.

-
- [1] G.C.Hegerfeldt, "Remark on causality and particle localization," *Phys. Rev. D* **10**, 3320-3321 (1974)
- [2] E. Karpov, G. Ordonez, T. Petrosky, I. Prigogine and G. Pronko, "Causality, delocalization, and positivity of energy," *Phys. Rev. A* **62**, 012103 (2000)
- [3] M. Hawton V. Debierre, "Maxwell meets Reeh-Schlieder: the quantum mechanics of neutral bosons," *Phys. Lett. A* **381**, 1926 (2017)
- [4] M. Hawton, "Maxwell quantum mechanics," *Phys. Rev. A* **100**, 012122 (2019)
- [5] H. Reeh and S. Schlieder, "Bemerkungen zur unitaräquivalenz von Lorentzinvarianten Feldern," *Nuovo Cimento* **22**, 1051 (1961)
- [6] T. D. Newton and E. P. Wigner, "Localized States for Elementary systems," *Rev. Mod. Phys.* **21**, 400 (1949)
- [7] M. Hawton, "Photon position operator with commuting components," *Phys. Rev. A* **59**, 954-959 (1999); "Photon wave functions in a localized coordinate space basis," *Phys. Rev. A* **59**, 3223-3227 (1999)
- [8] A. Mostafazadeh and F. Zamani, "Quantum Mechanics of Klein-Gordon Fields I: Hilbert Space, Localized States, and Chiral Symmetry," *Ann. Phys.* **321**, 2183 (2006)
- [9] H. Babaei and A. Mostafazadeh, "Quantum Mechanics of a Photon," *J. Math. Phys.* **58**, 082302 (2017)
- [10] M. Hawton and V. Debierre, "Photon position eigenvectors, Wigner's little group and Berry's phase", *J. Math. Phys.* **17**, 01602
- [11] M. Gell-Mann and A. Pais, "Behavior of Neutral Particles under Charge Conjugation," *Phys. Rev.* **97**, 1387-1389 (1958)
- [12] C. Cohen-Tannoudji, J. Dupont-Roc and G. Grynberg, *Photons and Atoms: Introduction to Quantum Electrodynamics*, Wiley-VCH (1997)
- [13] M. Hawton and T. Melde, "Photon number density operator $i\hat{E}\cdot\hat{A}$," *Phys. Rev. A* **51**, 4186 (1995)
- [14] D. C. Brody, "Biorthogonal Quantum Mechanics," *J. Math. Phys. A: Math. Theor.* **47**, 035305 (2014)
- [15] M. Hawton and W. E. Baylis, "Photon position operators and localized bases," *Phys. Rev. A* **64**, 012101 (2001); "Angular momentum and the geometrical gauge of localized photon states," **71**, 033816 (2005)
- [16] C. Itzykson and J. B. Zuber, *Quantum Field Theory 1st ed. McGraw-Hill, (1980)*; V. Debierre, La Fonction D'Onde du Photon en Principe et en Pratique, PhD thesis École Centrale de Marseille (2015) (in English)
- [17] S. L. Schweber, *Int. J. of Theor. Physics* **3**, 347-353 (1970)
- [18] C. Cohen-Tannoudji, B. Diu and F. Laloë, *Quantum Mechanics Volume one*, Wiley (1977)
- [19] Dong-Yeop Na and Weng Cho Chew, "Quantum Electromagnetic Finite-Difference Time-Domain Solver," *Quantum Rep.* **2**, 1-13 (2020)
- [20] M. Dobrski, M. Przanowski, J. Tosiek and F. J. Turubiates, "The geometrical interpretation of the photon position operator,arXiv:2104.04351 (2021)