

# Linear-time calculation of the expected sum of edge lengths in random projective linearizations of trees

Lluís Alemany-Puig\*  
 Universitat Politècnica de Catalunya,  
 Barcelona, Catalonia, Spain.

Ramon Ferrer-i-Cancho\*\*  
 Universitat Politècnica de Catalunya,  
 Barcelona, Catalonia, Spain.

*The syntactic structure of a sentence is often represented using syntactic dependency trees. The sum of the distances between syntactically related words has been in the limelight for the past decades. Research on dependency distances led to the formulation of the principle of dependency distance minimization whereby words in sentences are ordered so as to minimize that sum. Numerous random baselines have been defined to carry out related quantitative studies on languages. The simplest random baseline is the expected value of the sum in unconstrained random permutations of the words in the sentence, namely when all the shufflings of the words of a sentence are allowed and equally likely. Here we focus on a popular baseline: random projective permutations of the words of the sentence, that is, permutations where the syntactic dependency structure is projective, a formal constraint that sentences satisfy often in languages. Thus far, the expectation of the sum of dependency distances in random projective shufflings of a sentence has been estimated approximately with a Monte Carlo procedure whose cost is of the order of  $Zn$ , where  $n$  is the number of words of the sentence and  $Z$  is the number of samples; the larger  $Z$ , the lower the error of the estimation but the larger the time cost. Here we present formulae to compute that expectation without error in time of the order of  $n$ . Furthermore, we show that star trees maximize it, and devise a dynamic programming algorithm to retrieve the trees that minimize it.*

## 1. Introduction

A successful way to represent the syntactic structure of a sentence is a dependency graph (Nivre 2006) which relates the words of a sentence by pairing them with syntactic links as in Figure 1. Each link is directed and the arrow points from the *head word* to the *dependent word* (Figure 1). There are several conditions that are often imposed on the structure of dependency graphs (Nivre 2006). The first is *well-formedness*, namely, the graph is (weakly) connected. The second is *single-headedness*, that is, every word has at most one head. Another condition is *acyclicity*, that is, if two words, say  $w_i$  and  $w_j$ , are connected via following one or more directed links from  $w_i$  to  $w_j$  then there is no path of directed links from  $w_j$  to  $w_i$ . By definition, syntactic dependency trees always have a root vertex, that is, a vertex (word) with no head. The fourth condition is *projectivity*: often informally described as the situation where edges do not cross when drawn above the sentence and the root is not covered by any edge.

---

\* E-mail: lluis.alemany.puig@upc.edu.

\*\* E-mail: rferrericacho@cs.upc.edu.




Figure 1: Examples of sentences and their syntactic dependency structure. Here arc labels indicate dependency distances. The word within a rectangle is the root of the sentence, and the number on top of each edge denotes its length. a) Projective dependency tree (adapted from Groß and Osborne (2009)). b) Planar (but not projective) syntactic dependency structure (adapted from Groß and Osborne (2009)). c) Non-projective syntactic dependency structure (adapted from (Nivre 2009)).

When a dependency graph is well-formed, single-headed and acyclic, the graph is a directed tree, called syntactic dependency tree (Kuhlmann and Nivre 2006; Gómez-Rodríguez, Carroll, and Weir 2011). In addition, a syntactic dependency structure is projective if, for every vertex  $v$ , all vertices reachable from  $v$ , that is, the yield of  $v$ , form a continuous substring within the linear ordering of the sentence (Kuhlmann and Nivre 2006). Equivalently, a syntactic dependency structure is projective if the yield of each vertex of the tree forms a contiguous interval of positions in the linear ordering of the vertices. Kuhlmann and Nivre (2006) define an interval (with endpoints  $i$  and  $j$ ) as the set  $[i, j] = \{k \mid i \leq k \text{ and } k \leq j\}$ .

A linear arrangement of a graph is planar if it does not have edge crossings (Sleator and Temperley 1993; Kuhlmann and Nivre 2006). Then projectivity can be characterized as a combination of two properties: planarity and the fact that the root is not covered (Mel'čuk 1988). Planarity was, to the best of our knowledge, first thought of as *one-page embeddings* of trees by Bernhart and Kainen (1979). Figure 1 shows an example of a projective tree 1(a), a planar tree 1(b) and a non-projective tree 1(c) (see Bodirsky, Kuhlmann, and Möhl (2005) for further characterizations of syntactic dependency structures).

Here we represent the syntactic dependency structure of a sentence as a pair consisting of a rooted tree and a linear arrangement of its vertices. A rooted tree  $T^r$  is defined by a set of vertices  $V$ , a set of edges  $E$  and a root vertex  $r \in V$ , that is,  $T^r = (V, E; r)$ , (Figure 2(b)) to be a free tree  $T = (V, E)$  (Figure 2(a)) with one of its vertices, say  $r \in V$ , labeled as its root and with the edges oriented from  $r$  towards the leaves. A linear arrangement  $\pi$  (also called *embedding*) of an  $n$ -vertex graph  $G = (V, E)$  is a function that assigns every vertex  $u \in V$  to a position  $\pi(u)$ . Throughout this article, we assume that  $\pi(u) \in [n] = \{1, \dots, n\}$ . Linear arrangements are often seen as determined by the labeling of the vertices (Chung 1984; Kuhlmann and Nivre 2006), but here we consider that the labeling of a graph and  $\pi$  are independent. In order to clarify our notion of linear arrangement of a labeled graph  $G$ , we say that two linear arrangements  $\pi_1$  and  $\pi_2$  of  $G$  are equal if and only if, for every vertex  $u \in V$ , it holds that  $\pi_1(u) = \pi_2(u)$ .

In any linear arrangement  $\pi$  of a graph  $G = (V, E)$ , one can define properties on the graph's edges and on the arrangement as a whole. The length of an edge between vertices  $u$  and  $v$  is their distance in the linear arrangement, usually defined as

$$\delta_{uv}(\pi) = |\pi(u) - \pi(v)| \quad (1)$$




Figure 2: a) A free tree  $T = (V, E)$ . b) The tree  $T$  rooted at  $r = 4$ , yielding  $T^r = (V, E; 4)$ . c) Two different projective linear arrangements of  $T^r$ :  $\pi_1(5) = \pi_2(5) = 1$  (thus  $\pi_1^{-1}(1) = \pi_2^{-1}(1) = 5$ );  $\pi_1(8) = 6$ ,  $\pi_2(7) = 6$ .

Thus, the length of an edge in the arrangement is the number of vertices between its endpoints plus one as in previous studies (Iordanskii 1987; Shiloach 1979; Chung 1984; Hochberg and Stallmann 2003; Ferrer-i-Cancho 2004; Gildea and Temperley 2007, 2010; Ferrer-i-Cancho 2019). A less commonly used definition of edge length is (Hudson 1995; Hiranuma 1999; Eppler 2005; Liu, Xu, and Liang 2017)

$$\delta_{uv}^*(\pi) = |\pi(u) - \pi(v)| - 1 \quad (2)$$

Here we use  $D_\pi(G) = \sum_{uv \in E} \delta_{uv}(\pi)$  as the definition for the sum of edge lengths of  $G$  when it is linearly arranged by  $\pi$ , but we also derive some results for  $D_\pi^*(G) = \sum_{uv \in E} \delta_{uv}^*(\pi)$ .

There exists sizeable literature on the calculation of baselines for the sum of edge lengths on trees. Concerning the computation of the minimum baseline, Iordanskii (1987), and Hochberg and Stallmann (2003) independently devised an  $O(n)$ -time algorithm for planar (one-page) embedding of free trees. Gildea and Temperley (2007) sketched an algorithm for projective embeddings of rooted trees. Alemany-Puig, Esteban, and Ferrer-i-Cancho (2021) reviewed this problem and presented, to the best of our knowledge, the first  $O(n)$ -time algorithm. Polynomial-time algorithms for unconstrained embeddings were presented by Shiloach (1979), with complexity  $O(n^{2.2})$ , and later by Chung (1984), with complexities  $O(n^2)$  and  $O(n^\lambda)$ , where  $\lambda$  is any real number satisfying  $\lambda > \log 3 / \log 2$ . Concerning random baselines, the precursors are found in Zörnig's research on the distribution of the distance between repeats in a uniformly random arrangement of a sequence assuming that consecutive elements are at distance zero (Zörnig 1984) as in parallel research on syntactic dependency distances (Hudson 1995; Hiranuma 1999; Eppler 2005; Liu, Xu, and Liang 2017). Later, Ferrer-i-Cancho (2004, 2016) studied the expectation of the random variable  $D(T)$  defined as

$$D(T) = \sum_{uv \in E} \delta_{uv} \quad (3)$$

in uniformly random arrangements, where  $\delta_{uv}$  is a random variable defined over uniformly random unconstrained linear arrangements of the tree  $T$ , resulting in

$$\mathbb{E}[D(T)] = \frac{n^2 - 1}{3} \quad (4)$$

While there are constant-time formulae for the expectation of  $D(T)$  in unconstrained arrangements (Equation 4), a procedure to calculate the expected value of  $D(T)$

under projectivity is not forthcoming. Our primary goal is to improve the calculation of the expected sum of edge lengths in uniformly random projective arrangements with respect to the Monte Carlo method or random sampling method put forward by Gildea and Temperley (2007). Such a widely used procedure (Albert Park and Levy 2009; Futrell, Mahowald, and Gibson 2015; Kramer 2021) estimates the expectation of  $D$  of an  $n$ -vertex tree with an error that is negatively correlated with  $Z$ , the amount of arrangements sampled, while its cost is directly proportional to that amount, that is  $O(Zn)$ .

We improve upon these techniques by providing closed-form formulae for the expected value of  $D(T^r)$  in uniformly random projective arrangements of  $T^r$  that can be evaluated in  $O(n)$ -time. More formally, our goal in this article is to find closed-form formulae for  $\mathbb{E}_{\text{pr}}[D(T^r)]$ , the expectation of the random variable  $D(T^r)$  conditioned to the set of projective arrangements.

The structure of this article is the following. We first derive, in Section 2, an arithmetic expression for  $\mathbb{E}_{\text{pr}}[D(T^r)]$ , given by

### Theorem 1

Let  $T^r = (V, E; r)$  be a tree rooted at  $r \in V$ . The expected sum of edge lengths  $D(T^r)$  conditioned to uniformly random projective arrangements is

$$\mathbb{E}_{\text{pr}}[D(T^r)] = \frac{d_r(2n_r + 1) + n_r - 1}{6} + \sum_{u \in \Gamma_r} \mathbb{E}_{\text{pr}}[D(T_u^r)] \quad (5)$$

$$= \frac{1}{6} \left( -1 + \sum_{v \in V} n_v(2d_v + 1) \right) \quad (6)$$

where  $n_u$  denotes the number of vertices of the subtree  $T_u^r$  rooted at  $u \in V$ , that is,  $n_u = |V(T_u^r)|$ , and  $d_v$  is the out-degree of vertex  $v$  in the rooted tree. If  $d_r = 0$  then  $\mathbb{E}_{\text{pr}}[D(T^r)] = 0$ .

We then proceed to study which classes of trees of  $n$  vertices maximize and minimize  $\mathbb{E}_{\text{pr}}[D(T^r)]$  in Section 3 with the help of Equation 6. We characterize the maxima of  $\mathbb{E}_{\text{pr}}[D(T^r)]$ , and provide a dynamic programming algorithm to calculate its minima. Concerning maxima, the next theorem states that an  $n$ -vertex star tree rooted at its hub,  $\mathcal{S}_n^h$ , is the only maximum of  $\mathbb{E}_{\text{pr}}[D(T^r)]$ .

### Theorem 2

For any  $n$ -vertex rooted tree ( $n \geq 3$ ),  $\mathbb{E}_{\text{pr}}[D(T^r)] \leq \mathbb{E}_{\text{pr}}[D(\mathcal{S}_n^h)]$  with equality if, and only if,  $T^r = \mathcal{S}_n^h$ .

A tight upper bound of  $\mathbb{E}_{\text{pr}}[D(T^r)]$  is given by

### Corollary 1

Given any  $n$ -vertex rooted tree  $T^r = (V, E; r)$  rooted at  $r \in V$ , it holds that  $\mathbb{E}_{\text{pr}}[D(T^r)] \leq \mathbb{E}[D(T)] = (n^2 - 1)/3$  where  $\mathbb{E}[D(T)]$  is the expected sum of edge lengths in uniformly random (unconstrained) linear arrangements (Equation 4) and  $T$  is the free tree variant of  $T^r$ .

Section 4 compares our new method to calculate  $\mathbb{E}_{\text{pr}}[D(T^r)]$  exactly against the Monte Carlo estimation method using dependency treebanks and find that commonly used values of  $Z$  can yield a large relative error in the estimation on a single tree. We finally present some conclusions and propose future work in Section 5.




Figure 3: A tree  $T^r$  rooted at  $r$ . The children of the root are  $\Gamma_r = \{u_1, u_2, \dots, u_{d_r}\}$ , where  $d_r$  is the out-degree of  $r$ . Each  $T_{u_i}^r$ , for  $u_i \in \Gamma_r$ , denotes the subtree of  $T^r$  rooted at  $u_i$ .

## 2. Expected sum of edge lengths

We devote this section to characterize projective arrangements (Section 2.1) and to derive an arithmetic expression to calculate the sum of expected edge lengths in said arrangements (Section 2.2). We end this section with some instantiations of said expression for particular classes of trees (Section 2.3).

### 2.1 The number of random projective arrangements

The number of unconstrained arrangements of an  $n$ -vertex tree  $T$  is  $\mathbf{N}(T) = |\mathbf{P}(T)| = n!$ , where  $\mathbf{P}(T)$  denotes the set of all  $n!$  arrangements of  $T$ , hence  $\mathbf{N}(T)$  is independent from the tree structure. The number of projective arrangements of a tree, however, depends on its structure, in particular on the out-degree sequence of the tree, as is shown later in this section. Counting the amount of projective arrangements of a tree motivates a proper characterization that underpins the proof of Theorem 1. For this, we need to introduce some notation.

Henceforth we denote directed edges of a rooted tree  $T^r = (V, E; r)$  as  $uv = (u, v) \in E$ ; all edges are oriented towards the leaves. We denote the set of children of a vertex  $v \in V$  as  $\Gamma_v$ , and thus the out-degree of  $v$  is  $d_v = |\Gamma_v|$  in the rooted tree. In particular, we refer to the root's children as  $\Gamma_r = \{u_1, \dots, u_{d_r}\} \subset V$ . We denote the subtree of  $T^r$  rooted at  $u \in V$  as  $T_u^r$ ; we denote its size (in vertices) as  $n_u = |V(T_u^r)|$ ; notice that  $n_u \geq 1$ . We say that  $T_v^r$  is an immediate subtree of  $T_u^r$  if  $uv$  is an edge of the tree. Figure 3 depicts a rooted tree and the immediate subtrees of  $T^r$ .

The fact that subtrees span over intervals (Kuhlmann and Nivre 2006) is central to Proposition 1's proof. However, since intervals are associated to a fixed pair of starting and ending positions in a linear sequence, we use the term *segment* of a rooted tree  $T_u^r$  to refer to a real segment within the linear ordering containing all vertices of  $T_u^r$  (Figure 4); technically, that segment is an interval of length  $n_u$  whose starting and ending positions are unknown until the whole tree is fully linearized. Thus, a segment is a movable set of vertices within the linear ordering.

We provide a closed-form formula for the number of projective arrangements of a rooted tree,  $\mathbf{N}_{\text{pr}}(T^r)$ . This result helps us characterize said arrangements.

**Proposition 1**

Let  $T^r = (V, E; r)$  be a tree rooted at  $r \in V$ .

$$\mathbf{N}_{\text{pr}}(T^r) = (d_r + 1)! \prod_{u \in \Gamma_r} \mathbf{N}_{\text{pr}}(T_u^r) \quad (7)$$

$$= \prod_{v \in V} (d_v + 1)! \quad (8)$$

where  $d_v$  is the out-degree of vertex  $v$  in the rooted tree. If  $d_r = 0$  then  $\mathbf{N}_{\text{pr}}(T^r) = 1$ .

*Proof.* We can associate a set of segments to each vertex. The set of vertex  $u$  contains  $d_u + 1$  segments: one segment corresponds to  $u$  (the only segment of length 1), and the remaining  $d_u$  segments correspond to the immediate subtrees of  $T_u^r$ . We obtain a projective linear arrangement by permuting the elements of each set for all vertices. Therefore, a projective arrangement can be seen as being recursively composed of permutations of sets of segments. Such ‘recursion’ starts at the permutation of the set of segments associated to  $r$ . Note, then, that there are  $(d_r + 1)!$  possible permutations of the segments associated to vertex  $r$ . For a fixed permutation of the segments associated to  $r$ , there are  $\prod_{u \in \Gamma_r} \mathbf{N}_{\text{pr}}(T_u^r)$  different projective arrangements of its immediate subtrees  $T_u^r$ , hence the recurrence in Equation 7. Equation 8 follows upon unfolding the recurrence.  $\square$

The proof of Proposition 1 can be used to devise a simple procedure to generate projective arrangements uniformly at random, and another to enumerate all projective arrangements, of a rooted tree. As explained in previous articles (Gildea and Temperley 2007; Futrell, Mahowald, and Gibson 2015), the former method consists of first generating a uniformly random permutation of the  $d_v + 1$  segments associated to every vertex  $v \in V$  and, afterwards, constructing the arrangement using these permutations. When a tree is linearized using the permutations of the sets of segments, we say that each segment becomes an interval.

## 2.2 The expected sum of edge lengths in random projective arrangements

We first review the problem of computing the  $\mathbb{E}[D(T)]$  the expected value of  $D(T)$  in uniformly random unconstrained arrangements so as to introduce the methodology applied for  $\mathbb{E}_{\text{pr}}[D(T^r)]$ . The calculation requires two steps: first, the calculation of  $\mathbb{E}[\delta_{uv}]$ , the expected length of an edge, and second the calculation of  $\mathbb{E}[D(T)]$ ; henceforth we denote these value as  $\mathbb{E}[D]$  and  $\mathbb{E}[\delta]$  since they only depend on the size of  $T$ . For simplicity, we assume the definition of edge length in Equation 1.

The calculation of  $\mathbb{E}[\delta]$  requires the calculation of  $\mathbb{P}(\delta)$ , that is the probability that an edge linking two vertices has length  $\delta$ . This is actually the proportion of unconstrained linear arrangements such that the two vertices are at distance  $\delta$ ; said expected value and probability do not depend on the edge. There are  $\mathbf{N}(T) = n!$  unconstrained linear arrangements and  $2(n - d)(n - 2)!$  unconstrained arrangements where the pair of vertices are at distance  $\delta$ , hence

$$\mathbb{P}(\delta) = \frac{2(n - \delta)(n - 2)!}{n!} = \frac{2(n - \delta)}{n(n - 1)} \quad (9)$$

as expected from previous research (Zörnig 1984; Ferrer-i-Cancho 2004). Then, the expected length of an edge in an unconstrained random arrangement is (Zörnig 1984; Ferrer-i-Cancho 2004)

$$\mathbb{E}[\delta] = \sum_{\delta=1}^{n-1} \delta \mathbb{P}(\delta) = \frac{n+1}{3} \quad (10)$$

The second step is the calculation of  $\mathbb{E}[D]$ , the expected value of  $D$  in an unconstrained arrangement. Since a tree has  $n - 1$  edges,  $\mathbb{E}[D] = (n - 1)\mathbb{E}[\delta]$ , which gives Equation 4.

Given a random variable  $Y(T^r)$  whose probability space is that of all  $n!$  uniformly random (unconstrained) linear arrangements of  $T^r$ , we denote  $Y(T^r)$ 's expected value conditioned to projective arrangements with the subscript 'pr',  $\mathbb{E}_{\text{pr}}[Y(T^r)]$ ; more formally,  $\mathbb{E}_{\text{pr}}[Y(T^r)] = \mathbb{E}[Y(T^r) \mid \mathbf{P}_{\text{pr}}(T^r)]$ . To obtain an arithmetic expression for  $\mathbb{E}_{\text{pr}}[D(T^r)]$ , we follow again a two-step approach. First, we calculate the expected length of an edge in uniformly random projective arrangements. However, unlike the unconstrained case, the edge must be incident to the root. Second, we calculate the expected  $\mathbb{E}_{\text{pr}}[D(T^r)]$  applying the result of the first step. Before we proceed, we need to introduce some notation.

An edge connecting the root of tree ( $r$ ) with one of its children ( $u$ ) can be decomposed into an anchor (Shiloach 1979; Chung 1984) and its coanchor (Figure 4). Such decomposition is also found in (Gildea and Temperley 2007; Albert Park and Levy 2009) but using different terminology. In the context of projective linear arrangements, we define  $\alpha_{ru}$  as the length of the anchor, that is, the number of positions of the linear arrangement covered by the edge  $ru \in E$  in the segment of  $T_u^r$  including the end of the edge  $\pi(u)$ ; similarly, we define  $\beta_{ru}$  as the number of positions of the linear arrangement that are covered by that edge in segments other than that of  $T_u^r$  and  $r$ . Then the length of an edge connecting  $r$  with  $r$  is  $\delta_{ru}(\pi) = |\pi(r) - \pi(u)| = \alpha_{ru}(\pi) + \beta_{ru}(\pi)$ .

The next lemma shows that the expected value of  $\delta_{ru}$  depends only on the size of the whole tree ( $n_r$ ) and the size of the subtree rooted at the child ( $n_u$ ).

**Lemma 1**

Let  $T^r = (V, E; r)$  be a tree rooted at  $r \in V$ . Given an edge  $ru \in E$ , its anchor's expected length in uniformly random projective arrangements is

$$\mathbb{E}_{\text{pr}}[\alpha_{ru}] = \frac{n_u + 1}{2} \quad (11)$$

while the expected length of its coanchor is

$$\mathbb{E}_{\text{pr}}[\beta_{ru}] = \frac{n_r - n_u - 1}{3} \quad (12)$$

Therefore, the expected length of an edge  $ru \in E$  in such arrangements is

$$\mathbb{E}_{\text{pr}}[\delta_{ru}] = \mathbb{E}_{\text{pr}}[\alpha_{ru} + \beta_{ru}] = \frac{2n_r + n_u + 1}{6} \quad (13)$$

*Proof.* By the law of total expectation

$$\begin{aligned}\mathbb{E}_{\text{pr}}[\alpha_{ru}] &= \mathbb{E}_{\text{pr}}[\alpha_{ru} \mid \pi(u) < \pi(r)] \mathbb{P}_{\text{pr}}(\pi(u) < \pi(r)) \\ &\quad + \mathbb{E}_{\text{pr}}[\alpha_{ru} \mid \pi(u) > \pi(r)] \mathbb{P}_{\text{pr}}(\pi(u) > \pi(r))\end{aligned}\quad (14)$$

where  $\mathbb{P}_{\text{pr}}(\pi(u) < \pi(r))$  is the probability that  $u$  precedes  $r$  in a random projective linear arrangement and  $\mathbb{P}_{\text{pr}}(\pi(u) > \pi(r)) + \mathbb{P}_{\text{pr}}(\pi(u) < \pi(r)) = 1$ . As any projective linear arrangement such that  $u$  precedes  $r$  has a reverse projective arrangement where  $u$  follows  $r$ , then  $\mathbb{P}_{\text{pr}}(\pi(u) < \pi(r)) = 1/2$  and

$$\mathbb{E}_{\text{pr}}[\alpha_{ru}] = \frac{1}{2} (\mathbb{E}_{\text{pr}}[\alpha_{ru} \mid \pi(u) < \pi(r)] + \mathbb{E}_{\text{pr}}[\alpha_{ru} \mid \pi(u) > \pi(r)]) \quad (15)$$

Now, let  $q_u$  be relative position of vertex  $u$  in its segment in the (projective) linear arrangement (the  $i$ -th vertex of said segment is at relative position  $i$ ). If  $u$  precedes  $r$  in the linear arrangement ( $\pi(u) < \pi(r)$ ) as in Figure 4 then  $\alpha_{ru} = n_u - q_u + 1$ . If  $u$  follows  $r$  in the linear arrangement ( $\pi(r) < \pi(u)$ ),  $\alpha_{ru} = q_u$ . Applying these two results, we obtain

$$\mathbb{E}_{\text{pr}}[\alpha_{ru}] = \frac{1}{2} (\mathbb{E}_{\text{pr}}[n_u - q_u + 1 \mid \pi(u) < \pi(r)] + \mathbb{E}_{\text{pr}}[q_u \mid \pi(u) > \pi(r)]) \quad (16)$$

By symmetry,  $\mathbb{E}_{\text{pr}}[q_u \mid \pi(u) < \pi(r)] = \mathbb{E}_{\text{pr}}[q_u \mid \pi(u) > \pi(r)]$  and then  $\mathbb{E}_{\text{pr}}[\alpha_{ru}] = (n_u + 1)/2$ , hence Equation 11.

In order to calculate  $\beta_{ru}$ , we define  $s_{ru}$  as the number of intermediate segments between  $r$  and the segment of  $T_u^r$  in the linear arrangement. Therefore,  $\beta_{ru}$  can be decomposed in terms of the lengths of each of these segments as

$$\beta_{ru} = \sum_{i=1}^{s_{ru}} \beta_{ru}(i) \quad (17)$$

By the law of total expectation,

$$\mathbb{E}_{\text{pr}}[\beta_{ru}] = \sum_{s=1}^{d_r-1} \mathbb{E}_{\text{pr}}[\beta_{ru} \mid s_{ru} = s] \mathbb{P}_{\text{pr}}(s_{ru} = s) \quad (18)$$

where  $\mathbb{E}_{\text{pr}}[\beta_{ru} \mid s_{ru} = s]$  is the expectation of  $\beta_{ru}$  given that  $u$  and  $r$  are separated by  $s$  segments, and  $\mathbb{P}_{\text{pr}}(s_{ru} = s)$  is the probability that  $u$  and  $r$  are separated by  $s$  intermediate segments, both in uniformly random projective arrangements. On the one hand,

$$\mathbb{E}_{\text{pr}}[\beta_{ru} \mid s_{ru} = s] = \mathbb{E}_{\text{pr}}\left[\sum_{i=1}^s \beta_{ru}(i)\right] = s \mathbb{E}_{\text{pr}}[\beta_{ru}(i)] \quad (19)$$

where

$$\mathbb{E}_{\text{pr}}[\beta_{ru}(i)] = \frac{n_r - n_u - 1}{d_r - 1} \quad (20)$$




Figure 4: A permutation  $\tau$  of the segments associated to  $r$ . Each rectangle represents the segment of the root  $r$  and those of the subtrees rooted at  $u, v, w \in \Gamma_r$ , denoted as  $T_u^r, T_v^r, T_w^r$ . The representative vertices of the segments are, from left to right:  $u, r, v$  and  $w$ . The thick dashed line represents the anchor of edge  $ru$  (whose length is  $\alpha_{ru}$ ), and the dotted line represents the coanchor of edge  $ru$  (whose length is  $\beta_{ru}$ ).

is the average length of the segments excluding those of  $r$  and  $T_u^r$ . On the other hand,  $\mathbb{P}_{\text{pr}}(s_{ru} = s)$  is the proportion of projective linear arrangements where the segments of  $T_u^r$  and that of  $r$  are separated by  $s$  segments in the linear arrangement, that is

$$\mathbb{P}_{\text{pr}}(s_{ru} = s) = \frac{2(d_r - s)(d_r - 1)! \prod_{u \in \Gamma_r} \mathbf{N}_{\text{pr}}(T_u^r)}{(d_r + 1)! \prod_{u \in \Gamma_r} \mathbf{N}_{\text{pr}}(T_u^r)} = \frac{2(d_r - s)}{(d_r + 1)d_r} \quad (21)$$

Plugging Equations 19 and 21 into Equation 18, one finally obtains Equation 12,

$$\mathbb{E}_{\text{pr}}[\beta_{ru}] = 2 \frac{n_r - n_u - 1}{(d_r + 1)d_r(d_r - 1)} \sum_{s=1}^{d_r-1} s(d_r - s) = \frac{n_r - n_u - 1}{3} \quad (22)$$

□

Now we can derive an arithmetic expression for  $\mathbb{E}_{\text{pr}}[D(T^r)]$ .

*Proof of Theorem 1.* Consider the random variable  $D(T^r)$  over the probability space of uniformly random (unconstrained) linear arrangements of  $T^r$ , as defined above. This variable can be decomposed into two summations

$$D(T^r) = \sum_{u \in \Gamma_r} D(T_u^r) + \sum_{u \in \Gamma_r} \delta_{ru} \quad (23)$$

The first summation groups the edges by subtrees of  $T^r$ . The second summation groups the edges incident to the root  $r$ . Then, we can use linearity of expectation to obtain

$$\mathbb{E}_{\text{pr}}[D(T^r)] = \sum_{u \in \Gamma_r} \mathbb{E}_{\text{pr}}[D(T_u^r)] + \sum_{u \in \Gamma_r} \mathbb{E}_{\text{pr}}[\delta_{ru}] \quad (24)$$

The recurrence in Equation 5 follows easily from applying Lemma 1 to Equation 24, which gives,

$$\sum_{ru \in E} \mathbb{E}_{\text{pr}}[\delta_{ru}] = \frac{d_r(2n_r + 1)}{6} + \frac{1}{6} \sum_{u \in \Gamma_r} n_u = \frac{d_r(2n_r + 1) + n_r - 1}{6} \quad (25)$$

Equation 6 follows upon unfolding the recurrence.  $\square$

In the proof above we implicitly use our definition of edge length  $\delta_{uv}^*(\pi)$  (Equation 1). Nevertheless, the expression in Equation 6 can be easily adjusted to use different definitions of edge length, for example,  $\delta_{uv}^*(\pi)$  (Equation 2). It suffices to find an appropriate transformation of our definition of  $D(T^r)$  (Equation 3) into the one desired, namely  $D^*(T^r)$ . The next corollary gives the solution.

### Corollary 2

Let  $T^r = (V, E; r)$  be a tree rooted at  $r \in V$ . We have that

$$\mathbb{E}_{\text{pr}} [D^*(T^r)] = \frac{d_r(2n_r - 5) + n_r - 1}{6} + \sum_{u \in \Gamma_r} \mathbb{E}_{\text{pr}} [D^*(T_u^r)] \quad (26)$$

$$= \frac{1}{6} \left( 5 - 6n_r + \sum_{v \in V} n_v(2d_v + 1) \right) \quad (27)$$

*Proof of Corollary 2.* The fact that  $\mathbb{E}_{\text{pr}} [D^*(T^r)] = \mathbb{E}_{\text{pr}} [D(T^r)] - (n_r - 1)$ , transforms Equation 5 into Equation 26 immediately as well as Equation 6 into Equation 27 thanks to

$$\sum_{u \in \Gamma_r} \mathbb{E}_{\text{pr}} [D(T_u^r)] = \sum_{u \in \Gamma_r} (\mathbb{E}_{\text{pr}} [D^*(T_u^r)] + n_u - 1) = n_r - 1 - d_r + \sum_{u \in \Gamma_r} \mathbb{E}_{\text{pr}} [D^*(T_u^r)] \quad (28)$$

$\square$

It is easy to see that Equations 6 and 27 can both be evaluated in  $O(n)$ -time and  $O(n)$ -space, where  $n$  is the number of vertices of the tree: one only needs to compute the values  $n_v$  in  $O(n)$ -time, store them in  $O(n)$  space, and then evaluate the formula, also in  $O(n)$ -time using those values. In the analysis above, we are assuming that the values of  $d_v$  are already computed; depending on the data structure used to represent the tree, the cost of computing  $d_v$  might be relevant.

### 2.3 Formulae for classes of trees

For the following analyses, we define the hub of a rooted tree as the vertex of the underlying free tree that has the highest degree; we also use the term ‘leaf’ to refer to a leaf in the underlying free tree. We now instantiate Equations 6 and 8 for several classes of trees (Table 1). Star trees,  $\mathcal{S}_n$ , rooted at the hub,  $S_n^h$ , and at a leaf,  $S_n^l$ . Quasi-star trees,  $\mathcal{Q}_n$ , rooted at the hub,  $Q_n^h$ , at a leaf adjacent to the hub,  $Q_n^{hl}$ , at the leaf not adjacent to the hub  $Q_n^e$ , and at the only internal vertex that is not the hub,  $Q_n^b$ . Linear trees when rooted at a vertex at distance  $k \geq 0$  from one of the leaves,  $\mathcal{L}_n^k$ . Each class of tree is depicted in Figure 5. These classes trees are chosen for graph theoretic reasons. Linear trees minimize the variance of the degree (Ferrer-i-Cancho 2013); star graphs maximize it (Ferrer-i-Cancho 2013) and all their unconstrained linear arrangements are planar; concerning  $S_n^h$ , all its linear arrangements are projective; quasi star trees maximize the variance of the degree among trees whose set of unconstrained arrangements contains some non-planar arrangement (Ferrer-i-Cancho 2016).

We choose linear trees to illustrate how one can instantiate Equation 6. In order to ease this task, we rewrite Equation 6 using vectorial notation as

$$\mathbb{E}_{\text{pr}} [D(T^r)] = \frac{1}{6} \left( -1 + 2 \sum_{v \in V} n_v d_v + \sum_{v \in V} n_v \right) = \frac{1}{6} \left( -1 + 2 \vec{n}_v \cdot \vec{d}_v + \vec{n}_v \cdot \vec{1} \right) \quad (29)$$

For  $\mathcal{L}_n^0$ , we have

$$\begin{aligned} \vec{n}_v &= (n, n-1, n-2, \dots, 3, 2, 1) \\ \vec{d}_v &= (1, 1, 1, \dots, 1, 1, 0) \end{aligned}$$

Then

$$\mathbb{E}_{\text{pr}} [D(\mathcal{L}_n^0)] = \frac{1}{6} \left( -1 + 2 \sum_{i=2}^n i + \sum_{i=1}^n i \right) = \frac{(n-1)(n+2)}{4} \quad (30)$$

For  $\mathcal{L}_n^k$  with  $k > 0$ , we have

$$\begin{aligned} \vec{n}_v &= (1, 2, \dots, k-1, k, n, n-k-1, n-k-2, \dots, 2, 1) \\ \vec{d}_v &= (0, 1, \dots, 1, 1, 2, 1, \dots, 1, 0) \end{aligned}$$

and then

$$\mathbb{E}_{\text{pr}} [D(\mathcal{L}_n^k)] = \frac{1}{6} \left[ -1 + 2 \left( 2n + \sum_{j=2}^k j + \sum_{j=2}^{n-k-1} j \right) + \left( n + \sum_{j=1}^k j + \sum_{j=1}^{n-k-1} j \right) \right] \quad (31)$$

$$= \frac{(n-1)(3n+10) + 6k(k+1-n)}{12} \quad (32)$$

Regarding Equation 8 for  $\mathcal{L}_n^0$  and  $\mathcal{L}_n^k$  we have that

$$\mathbf{N}_{\text{pr}}(\mathcal{L}_n^0) = (0+1)! \prod_{i=1}^{n-1} (1+1)! = 2^{n-1} \quad (33)$$

$$\mathbf{N}_{\text{pr}}(\mathcal{L}_n^k) = (2+1)!(0+1)!(0+1)! \prod_{i=1}^{n-3} (1+1)! = 3 \cdot 2^{n-2} \quad (34)$$

### 3. Maxima and minima

In this section, we tackle the problem of computing the minima and characterizing the maxima of  $\mathbb{E}_{\text{pr}} [D(T^r)]$ . In particular, we give a closed-form formula for the maximum value of  $\mathbb{E}_{\text{pr}} [D(T^r)]$  and characterize the trees that maximize it, as well as outline a dynamic programming algorithm to compute the minima. Henceforth, we use  $\mathcal{T}_n$  to denote the set of  $n$ -vertex (unlabeled) rooted trees. Evidently, any tree that maximizes (resp. minimizes)  $\mathbb{E}_{\text{pr}} [D(T^r)]$  also maximizes (resp. minimizes)  $\mathbb{E}_{\text{pr}} [D^*(T^r)]$ , thus we

Table 1: Instantiations of  $\mathbf{N}_{\text{pr}}(T^r)$  and  $\mathbb{E}_{\text{pr}}[D(T^r)]$  for several classes of trees (Figure 5).

Class of tree	$T^r$	$\mathbf{N}_{\text{pr}}(T^r)$	$\mathbb{E}_{\text{pr}}[D(T^r)]$
Star	$\mathcal{S}_n^h$	$n!$	$(n^2 - 1)/3$
	$\mathcal{S}_n^l$	$2(n - 1)!$	$n(2n - 1)/6$
Quasi star ( $n \geq 4$ )	$\mathcal{Q}_n^h$	$2(n - 1)!$	$(2n^2 - 2n + 3)/6$
	$\mathcal{Q}_n^e$	$4(n - 2)!$	$(2n^2 - 2n + 3)/6$
	$\mathcal{Q}_n^b$	$6(n - 2)!$	$(2n^2 - 3n + 7)/6$
	$\mathcal{Q}_n^{hl}$	$4(n - 2)!$	$(2n^2 - 3n + 7)/6$
Linear	$\mathcal{L}_n^0$	$2^{n-1}$	$(n - 1)(n + 2)/4$
	$\mathcal{L}_n^k, (k > 0)$	$3 \cdot 2^{n-2}$	$[(n - 1)(3n + 10) + 6k(k + 1 - n)]/12$




Figure 5: Linear trees ( $\mathcal{L}_n$ ), star trees ( $\mathcal{S}_n$ ) and quasi-star trees ( $\mathcal{Q}_n$ ) of  $n$  vertices. Labels 0 and  $k$  in  $\mathcal{L}_n^k$  denote the distance of the labeled vertex from the same leaf. A circled dot marks a tree's root.

restrict our study to the former. Throughout this section, we use  $n_i$  to refer to the size of the subtree rooted at the  $i$ th child of the root for  $1 \leq i \leq d_r$ .

The following lemma shows that problem of constructing the  $n$ -vertex rooted trees  $T^r \in \mathcal{T}_n$  that maximize (resp. minimize)  $\mathbb{E}_{\text{pr}}[D(T^r)]$  has optimal substructure.

**Lemma 2**

Let  $\mathcal{T}_n$  be the set of all unlabeled rooted trees of  $n$  vertices.  $f(n)$ , the optimal value of  $\mathbb{E}_{\text{pr}}[D(T^r)]$  satisfies

$$f(n) = \text{opt}_{T^r \in \mathcal{T}_n} \{ \mathbb{E}_{\text{pr}}[D(T^r)] \} \tag{35}$$

$$= \text{opt}_{0 \leq d_r \leq n-1} \left\{ \frac{d_r(2n + 1) + n - 1}{6} + \text{opt}_{\substack{n_1 + \dots + n_{d_r} = n-1 \\ n_i \geq 1}} \left\{ \sum_{i=1}^{d_r} f(n_i) \right\} \right\} \tag{36}$$

*Proof.* We can construct an  $n$ -vertex optimal tree using optimal subtrees. We can obtain the right hand side of Equation 35 using the recurrence in Equation 5. An optimal  $n$ -

vertex tree is one whose cost value is optimal among all optimal  $n$ -vertex trees whose root has fixed degree  $d_r \in \{1, \dots, n-1\}$ . Given a fixed root degree  $d_r$  such that  $1 \leq d_r \leq n-1$ , an optimal  $n$ -vertex rooted tree can be built by constructing an optimal  $(n-1)$ -vertex forest of  $d$  rooted trees, each of size  $n_i$  and optimal among the  $n_i$ -vertex trees. Choosing the  $d_r$  trees to be  $n_i$ -optimal makes the sum of their costs,  $\sum_i f(n_i)$ , optimal.  $\square$

In subsequent paragraphs we use  $f_M(n)$  to denote the *maximization* variant and  $f_m(n)$  to denote the *minimization* variant of  $f(n)$ , respectively. Theorem 2 characterizes the maxima of  $\mathbb{E}_{\text{pr}}[D(T^r)]$ . Perhaps not so surprisingly, the only maximum of  $\mathbb{E}_{\text{pr}}[D(T^r)]$  is obtained by  $\mathcal{S}_n^h$ .

*Proof of Theorem 2.* We prove this by induction on  $n$  using the formalization of the optimum in Equation 35. The base cases can be easily obtained by an exhaustive enumeration of the (unlabeled) rooted trees of  $n$  vertices for some small  $n$ . Indeed, the only tree that maximizes  $\mathbb{E}_{\text{pr}}[D(T^r)]$  is  $\mathcal{S}_n^h$  for all  $n \leq 2$ .

For  $n \geq 3$ , it holds that  $\mathbb{E}_{\text{pr}}[D(\mathcal{S}_n^h)] = \mathbb{E}[D(\mathcal{S}_n^h)] = (n^2 - 1)/3$  (Equation 4). In order to prove that  $\mathcal{S}_n^h$  is maximum for larger  $n$ , it suffices to prove that

$$\mathbb{E}_{\text{pr}}[D(\mathcal{S}_n^h)] = \frac{n^2 - 1}{3} > \max_{0 \leq d_r \leq n-2} \left\{ \frac{d_r(2n+1) + n-1}{6} + L(n, d_r) \right\} \quad (37)$$

where

$$L(n, d_r) = \max_{\substack{n_1 + \dots + n_{d_r} = n-1 \\ n_i \geq 1}} \left\{ \sum_{i=1}^{d_r} f_M(n_i) \right\} \quad (38)$$

For this we need to know the maximum value of  $L(n, d_r)$ . Applying the induction hypothesis (each maximum subtree of a tree of  $n$  vertices is a star tree), we have that

$$L(n, d_r) = \max_{\substack{n_1 + \dots + n_{d_r} = n-1 \\ n_i \geq 1}} \left\{ \sum_{i=1}^{d_r} \frac{n_i^2 - 1}{3} \right\} = \frac{1}{3} \left( -d_r + \max_{\substack{n_1 + \dots + n_{d_r} = n-1 \\ n_i \geq 1}} \left\{ \sum_{i=1}^{d_r} n_i^2 \right\} \right) \quad (39)$$

Notice that any vector  $(n_1, \dots, n_i, \dots, n_{d_r})$ , such that  $1 \leq n_1 \leq \dots \leq n_{d_r}$ , can be transformed into another vector  $(n_1, \dots, n_i - 1, \dots, n_{d_r} + 1)$ , for any  $n_i \geq 2$ , such that the sum of squared components is strictly larger while the sum of the components remains constant. Therefore, the maximum sum of squares is obtained by choosing  $n_{d_r} = n - d_r$  and  $n_i = 1$  for  $1 \leq i < d_r$ , yielding

$$L(n, d_r) = \frac{1}{3} [-d_r + (n - d_r)^2 + (d_r - 1)] = \frac{(n - d_r)^2 - 1}{3} \quad (40)$$

and the theorem holds if, and only if

$$\frac{n^2 - 1}{3} > \max_{0 \leq d_r \leq n-2} \left\{ \frac{d_r(2n+1) + n-1}{6} + \frac{(n - d_r)^2 - 1}{3} \right\} \quad (41)$$

After rearranging the terms that do not depend on  $d$  to the left-hand side we obtain

$$-n + 1 > \max_{0 \leq d_r \leq n-2} \{d_r(2d_r - 2n + 1)\} \quad (42)$$

The right hand side of the inequality is maximized for  $d_r = n - 2$ . Since this last inequality holds true when  $n > 5/2$ , we are done.  $\square$

It is easy to see that  $\mathbb{E}_{\text{pr}} [D(T^r)]$  is bounded above by the expected value of  $D(T)$  over the space of uniformly random (unconstrained) linear arrangements.

*Proof of Corollary 1.* Due to Theorem 2 the maximum value of  $\mathbb{E}_{\text{pr}} [D(T^r)]$  is maximized by  $S_n^h$  which is exactly  $(n^2 - 1)/3$ , as shown in Table 1.  $\square$

We devised a dynamic programming algorithm based on Lemma 2 to calculate the trees minimizing  $\mathbb{E}_{\text{pr}} [D(T^r)]$ . The method to obtain said values and trees is outlined in Algorithm 3.1. That algorithm has two parameters:  $n$ , the number of vertices, and  $H$  a hash table whose keys are natural numbers  $k$  and the value associated to each key is a pair formed by  $f_m(k)$  and the  $k$ -vertex trees  $T^r$  that attain that  $f_m(k)$ . Notice that  $n$  is an input parameter, while  $H$  is an input/output parameter. In order to calculate the minimum  $n$ -vertex trees, the values of the parameters of the first call to Algorithm 3.1 are the value  $n$  and an empty  $H$ . Now, Algorithm 3.1 is a direct evaluation of Equation 35, that is, for every value of out-degree of the root  $d_r$  ( $1 \leq d_r \leq n - 1$ ), it computes the value  $f_m(n)$  by finding the partition of  $n - 1$  into  $d$  summands that minimizes

$$\frac{d_r(2n + 1) + n - 1}{6} + \underset{\substack{n_1 + \dots + n_{d_r} = n-1 \\ n_i \geq 1}}{\text{opt}} \left\{ \sum_{i=1}^{d_r} f_m(n_i) \right\} \quad (43)$$

where  $f_m(n_i)$  is calculated recursively and stored in the hash table  $H$ . Therefore, the algorithm's complexity heavily depends on the number of partitions of  $n - 1$ , denoted as  $p(n - 1)$ . The worst-case complexity of Algorithm 3.1, then, is  $\omega(n^c)$  (for any constant  $c \in \mathbb{R}$ ) due to the exponential nature of  $p(n)$  (Hardy and Ramanujan 1918). Finally, notice that the 'Modified Cartesian' product in line 19 of Algorithm 3.1 must ensure that no repeated trees are produced. Repeated trees arise in the standard Cartesian product due to the fact that some partitions may have repeated parts. As an example, consider the partition of 37 with two repeated parts (11, 13, 13); such parts have 1 and 2 non-isomorphic minimum trees respectively (Figure 6). Let  $t_{11} = \{T_A\}$  be the unique 11-vertex minimum tree, and  $t_{13} = \{T_B, T_C\}$  be the two 13-vertex minimum trees. The Cartesian product  $t_{11} \times t_{13} \times t_{13}$  produces four forests, two of them being isomorphic:  $(T_A, T_B, T_C)$  and  $(T_A, T_C, T_B)$ ; the other two forests are  $(T_A, T_B, T_B)$  and  $(T_A, T_C, T_C)$ . In order to obtain the unique  $n$ -vertex trees (up to isomorphism) that attain  $f_m(n)$ , we modify the standard Cartesian product. Let  $\{T^*\}^{(i)}$  be the list of  $n_i$ -vertex minimum trees. Thus, one element in the Cartesian product is obtained by choosing the trees in the  $j_1$ th,  $\dots$ ,  $j_{d_r}$ th positions of the lists, that is,

$$(\{T^*\}_{j_1}^{(1)}, \dots, \{T^*\}_{j_{d_r}}^{(d_r)}) \in \{T^*\}^{(1)} \times \dots \times \{T^*\}^{(d_r)} \quad (44)$$

where  $1 \leq j_i \leq |\{T^*\}^{(i)}|$  for all  $i \in [1, d_r]$ . Now, tree uniqueness is ensured by forcing indices of every pair of lists  $\{T^*\}^{(i)}, \{T^*\}^{(i+1)}$  such that  $n_i = n_{i+1}$  to be  $j_i \leq j_{i+1}$ .

In this article, we do not characterize the minima of  $\mathbb{E}_{\text{pr}} [D(T^r)]$  since, unlike the amount of its maxima, which shows a clear regularity, the amount of its minima varies with  $n$  in a non-monotonic fashion. Table 2 shows that the number of these minima oscillates between 1 and 2 for  $n \leq 20$ , and Figure 6 shows that the shape of the trees does not seem to fit into a simple class.

---

**Algorithm 3.1:** Calculate the minimum value and trees of  $\mathbb{E}_{\text{pr}} [D(T^r)]$ .

---

**Input:**  $n \in \mathbb{N}$ , the number of vertices of the trees.  $H$  is a hash table whose keys are natural numbers  $k \in \mathbb{N}$  and the value of each key is the pair formed by  $f_m(k)$  and the  $k$ -vertex trees  $T^r$  for which  $\mathbb{E}_{\text{pr}} [D(T^r)] = f_m(k)$ .

**Output:** The hash table  $H$  updated to contain the results for  $n' \leq n$ .

```

1 Function MINIMUM_E_PROJECTIVE( $n, H$ ) is
2   if hash table  $H$  contains key  $n$  then Stop.
3    $f_m \leftarrow (n^2 - 1)/3$  // The minimum value initialized at a maximum
4    $T^* \leftarrow \emptyset$  // The set of  $n$ -optimal trees
5   if  $n \leq 2$  then
6      $T^* \leftarrow \{\text{the only rooted tree } T^r \text{ of } n \text{ vertices}\}$ 
7      $f_m \leftarrow \mathbb{E}_{\text{pr}} [D(T^r)]$ 
8   else
9     for each  $d_r \in [1, n - 1]$  do
10       $\mathcal{P} \leftarrow$  The set of partitions of ' $n - 1$ ' in  $d_r$  summands.
11      for each partition  $P = \{n_1, \dots, n_{d_r}\} \in \mathcal{P}$  do
12        // Initialize the cost of the trees
13         $C \leftarrow (d_r(2n + 1) + n - 1)/6$ 
14        // Evaluate the partition recursively
15        for each  $n_i \in P$  do
16          // Calculate the  $n_i$ -minimum trees
17          MINIMUM_E_PROJECTIVE( $n_i, H$ )
18           $f_m^{(i)}, \{T^*\}^{(i)} \leftarrow H[n_i]$  // Retrieve the trees from  $H$ 
19           $C \leftarrow C + f_m^{(i)}$ 
20          if  $C > f_m$  then
21            Stop evaluating  $P$  and move on to the next partition
22          // Modified Cartesian product of all  $\{T^*\}^{(i)}$  (see text)
23           $F \leftarrow \{T^*\}^{(1)} \times \dots \times \{T^*\}^{(d)}$ 
24          //  $W$  is the set of  $(n, d, P)$ -optimal trees
25           $W \leftarrow$  add a root to every forest in  $F$ 
26          if  $C < f_m$  then  $f_m \leftarrow C, T^* \leftarrow W^*$ 
27          else if  $C = f_m$  then  $T^* \leftarrow T^* \cup W^*$ 
28   Store  $\langle f_m, T^* \rangle$  in  $H[n]$ 

```

---

#### 4. Exact versus approximate calculation of the expected sum of dependency distances.

In previous research, a random baseline for  $D$  under projectivity has been estimated with a Monte Carlo method that consists of averaging the values of  $D$  obtained via




Figure 6: The trees that minimize  $\mathbb{E}_{\text{pr}} [D(T^r)]$  for  $n \leq 20$ . Their values of  $\mathbb{E}_{\text{pr}} [D(T^r)]$  are given in Table 2. Roots are drawn atop each tree and each edge should be regarded as oriented away from the root.

Table 2: The columns indicate, from left to right, the number of (unlabeled) rooted trees (Sloane 1964-2021), the number of trees that minimize  $\mathbb{E}_{\text{pr}} [D(T^r)]$ , and the value of  $\mathbb{E}_{\text{pr}} [D(T^r)]$  for such trees. The trees yielding these values are displayed in Figure 6.

$n$	# trees	# opt trees	$\mathbb{E}_{\text{pr}} [D(T^r)]$	$n$	# trees	# opt trees	$\mathbb{E}_{\text{pr}} [D(T^r)]$
1	1	1	0	11	1,842	1	22
2	1	1	1	12	4,766	1	151/6
3	1	1	5/2	13	12,486	2	85/3
4	4	2	9/2	14	32,973	1	63/2
5	9	1	19/3	15	87,811	1	104/3
6	20	1	26/3	16	235,381	1	38
7	48	1	11	17	634,847	1	83/2
8	115	2	83/6	18	1,721,159	2	45
9	286	1	33/2	19	4,688,676	2	97/2
10	719	2	58/3	20	12,826,228	2	52

random sampling of  $Z$  random projective arrangements of a rooted tree (Gildea and Temperley 2007; Albert Park and Levy 2009; Gildea and Temperley 2010; Futrell, Mahowald, and Gibson 2015; Kramer 2021). That method has cost  $O(Zn)$ , where  $n$  is the cost of generating a random arrangement and computing  $D$  on it. Our method, on top of providing exact values, has a much lower complexity  $O(n)$  as there is no need for random sampling. We used the UD2.5 (de Marneffe et al. 2019) treebanks for Catalan, English and German for illustration purposes. English and German are selected because they have been used in previous research employing projective random baselines (Gildea and Temperley 2007; Albert Park and Levy 2009; Gildea and

Temperley 2010). Catalan is included as the native language of the present authors. For each sentence  $T^r$  in a treebanks, we calculated  $\mathbb{E}_{\text{pr}}[D(T^r)]$  in two ways. First, exactly with Equation 6 (denoted as  $\mathbb{E}$ ). Second, approximately by averaging the value of  $D$  obtained in  $Z = 10^i$ , for  $1 \leq i \leq 4$ , uniformly random projective arrangements (denoted as  $\tilde{\mathbb{E}}_{\text{pr}}[D(T^r)]$ ). Using these values, we calculated the relative error  $\varepsilon_{rel}$  of a tree as

$$\varepsilon_{rel} = \frac{\tilde{\mathbb{E}}_{\text{pr}}[D(T^r)] - \mathbb{E}_{\text{pr}}[D(T^r)]}{\mathbb{E}_{\text{pr}}[D(T^r)]} \quad (45)$$

where  $\tilde{\mathbb{E}}_{\text{pr}}[D(T^r)]$  is the Monte Carlo estimate of  $\mathbb{E}_{\text{pr}}[D(T^r)]$ .  $\varepsilon_{rel} > 0$  indicates that  $\tilde{\mathbb{E}}_{\text{pr}}[D(T^r)]$  underestimates  $\mathbb{E}_{\text{pr}}[D(T^r)]$ ;  $\varepsilon_{rel} < 0$  indicates overestimation error. Figure 7 shows the relative error as a function of sentence length (average, minimum, maximum and confidence interval).

In previous research, the value of  $Z$  is sometimes indicated, for example,  $Z = 10$  (Kramer 2021) and  $Z = 100$  (Futrell, Mahowald, and Gibson 2015) and sometimes not reported (Albert Park and Levy 2009; Gildea and Temperley 2010). When  $Z = 10$ , Figure 7 shows that the relative error peaks between  $n = 10$  and  $n = 20$  (close to the mean sentence length in English, Rudnicka (2018)) and tends to decay from then onwards. However, the confidence interval of the relative error clearly broadens for sufficiently large  $n$ , for example,  $n = 30$  or  $n = 40$  onwards. These behaviors smooth for  $Z = 100$  while the range of variation and the confidence interval are narrower. For  $Z = 10^4$ , the relative error reduces dramatically but it is important to bear in mind that our exact method is about  $10^4$  times faster than the corresponding Monte Carlo method.

## 5. Conclusions and future work

In this article, we have derived several simple closed-form formulae related to projective arrangements of rooted trees in Section 2. Firstly, we have given a simple closed-form formula to calculate the amount of *different* projective arrangements a rooted tree admits,  $N_{\text{pr}}(T^r)$  (Proposition 1). The proof reveals a straightforward way of enumerating such arrangements for an input rooted tree, and of generating this kind of arrangements uniformly at random without a rejection method. Secondly, and more importantly, we have provided a way of calculating, for any given rooted tree, the expected value of the sum of edge lengths over the space of uniformly random *different* projective arrangements,  $\mathbb{E}_{\text{pr}}[D(T^r)]$  (Theorem 1 and Corollary 2). This means that future studies in which such value is calculated approximately via random sampling of arrangements can now be calculated exactly and much faster. The  $O(Zn)$  Monte Carlo method to estimate true expectation with an error that tends to zero as  $Z$  tends to infinity can now be replaced by our fast  $O(n)$  with zero error. Moreover, these formulae can be instantiated in particular classes of trees (as shown in Section 2.3). In Section 3, we have characterized the trees that maximize  $\mathbb{E}_{\text{pr}}[D(T^r)]$  and proven that there exists a dynamic programming method to calculate the minima of  $\mathbb{E}_{\text{pr}}[D(T^r)]$ . A precise characterization of the minima should be the subject of future research. Finally, in Section 4, we have shown the advantage of an exact and fast calculation of  $\mathbb{E}_{\text{pr}}[D(T^r)]$  for future quantitative dependency syntax research.

The present article is a part of a research program on the calculation of random baselines for  $D$  via formulae or exact algorithms under formal constraints on linear arrangements that started about two decades ago with the unconstrained case (Zörnig 1984; Ferrer-i-Cancho 2004) (Equation 4). Here we have covered the projective case. In




Figure 7: The statistical properties of  $\varepsilon_{rel}$ , the relative error in the estimation of  $\mathbb{E}_{pr} [D(T^r)]$  with a Monte Carlo method (Equation 45), as a function of the size of the tree  $n$ . For all sentences of the same length, we show the average  $\varepsilon_{rel}$  (solid black line), its 99% confidence interval calculated with a bootstrap method (shaded gray region), and the maximum and minimum  $\varepsilon_{rel}$  (vertical blue bars). The end of each row indicates  $Z$ , that is the number of random projective arrangements used to estimate  $\mathbb{E}_{pr} [D(T^r)]$  (from  $Z = 10$  for the top row to  $Z = 10^4$  for the bottom row).

a forthcoming article, we will focus on planar linearizations of (free) trees – those in which there are no edge crossings – and obtain a closed-form formula, and a  $O(n)$ -time algorithm, to calculate  $\mathbb{E}_{\text{pl}} [D(T)]$ , the expected value of  $D(T)$  in uniformly random planar arrangements of a given (labeled free) tree  $T$  (Alemany-Puig and Ferrer-i-Cancho 2021).

Here we have focused on  $\mathbb{E}_{\text{pr}} [D(T^r)]$ , the expectation of  $D(T^r)$  on random arrangements of an individual tree. Future research should consider the problem of  $\mathbb{E} [\mathbb{E}_{\text{pr}} [D]]$ , the expectation of  $\mathbb{E}_{\text{pr}} [D]$  on ensembles of random trees of a fixed size  $n$ .  $\mathbb{E} [\mathbb{E}_{\text{pr}} [D]]$  is indeed the average value of  $\mathbb{E}_{\text{pr}} [D(T^r)]$  among all  $n$ -vertex rooted trees  $T^r$ . Although there are at least two ensembles possible, that is uniformly random *labeled* trees and uniformly random *unlabeled* trees, a closed-form formula (or algorithm) to calculate  $\mathbb{E} [\mathbb{E}_{\text{pr}} [D]]$  seems easier to obtain in the former.

Finally, our research paves the way to investigate the optimality of dependency distances of languages under projectivity. Recently, that optimality has been evaluated in 93 languages from 19 families with the help of a new score, defined as Ferrer-i-Cancho et al. (2020)

$$\Omega(T) = \frac{\mathbb{E} [D(T)] - D(T)}{\mathbb{E} [D(T)] - \min_{\pi} \{D(T)\}} \quad (46)$$

where the ‘min’ is taken over all unconstrained linear arrangements; finding such a minimum  $D$  is known as the Minimum Linear Arrangement problem (Garey and Johnson 1976; Shiloach 1979; Chung 1984). With the result in Theorem 1, one could replicate the aforementioned study under the projectivity constraint by re-defining the score as

$$\Omega_{\text{pr}}(T^r) = \frac{\mathbb{E}_{\text{pr}} [D(T^r)] - D(T^r)}{\mathbb{E}_{\text{pr}} [D(T^r)] - \min_{\pi \in \mathbf{P}_{\text{pr}}(T^r)} \{D(T^r)\}} \quad (47)$$

where the calculation of the minimum sum of edge lengths under the projectivity constraint is denoted as  $\min_{\pi \in \mathbf{P}_{\text{pr}}(T^r)} \{D(T^r)\}$  (Iordanskii 1987; Hochberg and Stallmann 2003; Gildea and Temperley 2007), and is linear-time computable (Alemany-Puig, Esteban, and Ferrer-i-Cancho 2021). Thanks to our article, such investigation of the optimality of dependency distances can be carried out reducing the computational cost and maximizing numerical precision with respect to an approach based on a Monte Carlo estimation of  $\mathbb{E}_{\text{pr}} [D(T^r)]$ .

### Acknowledgments

LAP is supported by Secretaria d’Universitats i Recerca de la Generalitat de Catalunya and the Social European Fund. RFC is also supported by the recognition 2017SGR-856 (MACDA) from AGAUR (Generalitat de Catalunya). RFC and LAP are supported by the grant TIN2017-89244-R from MINECO (Ministerio de Economía, Industria y Competitividad).

### References

- Albert Park, Y. and R. Levy. 2009. Minimal-length linearizations for mildly context-sensitive dependency trees. In *Proceedings of the 10th Annual Meeting of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies (NAACL-HLT) conference*, pages 335–343, Association for Computational Linguistics, Stroudsburg, PA, USA.
- Alemany-Puig, Lluís, Juan Luis Esteban, and Ramon Ferrer-i-Cancho. 2021. Minimum projective linearizations of trees in linear time. *Arxiv*.

- Alemany-Puig, Lluís and Ramon Ferrer-i-Cancho. 2021. Linear-time calculation of the expected sum of edge lengths in planar linearizations of trees. *Arxiv*.
- Bernhart, Frank and Paul C Kainen. 1979. The book thickness of a graph. *Journal of Combinatorial Theory, Series B*, 27(3):320–331.
- Bodirsky, Manuel, Marco Kuhlmann, and Mathias Möhl. 2005. Well-nested drawings as models of syntactic structure. In *Proceedings of the 10th Conference on Formal Grammar and 9th Meeting on Mathematics of Language*, pages 195–204.
- Chung, F.R.K. 1984. On optimal linear arrangements of trees. *Computers & Mathematics with Applications*, 10(1):43–60.
- Eppler, E. 2005. *The syntax of German-English code-switching*. Ph.D. thesis, University College, London. (unpublished).
- Ferrer-i-Cancho, R. 2013. Hubiness, length, crossings and their relationships in dependency trees. *Glottometrics*, 25:1–21.
- Ferrer-i-Cancho, R. 2016. Non-crossing dependencies: least effort, not grammar. In A. Mehler, A. Lücking, S. Banisch, P. Blanchard, and B. Job, editors, *Towards a theoretical framework for analyzing complex linguistic networks*. Springer, Berlin, pages 203–234.
- Ferrer-i-Cancho, Ramon. 2004. Euclidean distance between syntactically linked words. *Physical Review E*, 70(5):5.
- Ferrer-i-Cancho, Ramon. 2019. The sum of edge lengths in random linear arrangements. *Journal of Statistical Mechanics*, 2019(5):053401.
- Ferrer-i-Cancho, Ramon, Carlos Gómez-Rodríguez, Juan Luis Esteban, and Lluís Alemany-Puig. 2020. The optimality of syntactic dependency distances. *Arxiv*.
- Futrell, R., K. Mahowald, and E. Gibson. 2015. Large-scale evidence of dependency length minimization in 37 languages. *Proceedings of the National Academy of Sciences*, 112(33):10336–10341.
- Garey, Michael R. and D. S. Johnson. 1976. Some simplified np-complete graph problems. *Theoretical Computer Science*, pages 237–267.
- Gildea, Daniel and David Temperley. 2007. Optimizing grammars for minimum dependency length. In *Proceedings of the 45th Annual Meeting of the Association of Computational Linguistics*, pages 184–191, Association for Computational Linguistics, Prague, Czech Republic.
- Gildea, David and David Temperley. 2010. Do grammars minimize dependency length? *Cognitive Science*, 34(2):286–310.
- Gómez-Rodríguez, Carlos, John Carroll, and David Weir. 2011. Dependency Parsing Schemata and Mildly Non-Projective Dependency Parsing. *Computational Linguistics*, 37(3):541–586.
- Groß, T. and T. Osborne. 2009. Toward a practical dependency grammar theory of discontinuities. *SKY Journal of Linguistics*, 22:43–90.
- Hardy, G. H. and S. Ramanujan. 1918. Asymptotic formulae in combinatory analysis. *Proceedings of the London Mathematical Society*, s2-17(1):75–115.
- Hiranuma, S. 1999. Syntactic difficulty in English and Japanese: A textual study. *UCL Working Papers in Linguistics*, 11:309–322.
- Hochberg, Robert A. and Matthias F. Stallmann. 2003. Optimal one-page tree embeddings in linear time. *Information Processing Letters*, 87(2):59–66.
- Hudson, R. 1995. Measuring syntactic difficulty. *Unpublished paper*.
- Iordanskii, M. A. 1987. Minimal numberings of the vertices of trees — Approximate approach. In *Fundamentals of Computation Theory*, pages 214–217, Springer Berlin Heidelberg, Berlin, Heidelberg.
- Kramer, Alex. 2021. Dependency lengths in speech and writing: A cross-linguistic comparison via YouDePP, a pipeline for scraping and parsing YouTube captions. In *Proceedings of the Society for Computation in Linguistics*, volume 4.
- Kuhlmann, Marco and Joakim Nivre. 2006. Mildly non-projective dependency structures. In *Proceedings of the COLING/ACL 2006 Main Conference Poster Sessions*, July, pages 507–514.
- Liu, Haitao, Chunshan Xu, and Junying Liang. 2017. Dependency distance: A new perspective on syntactic patterns in natural languages. *Physics of Life Reviews*, 21:171–193.
- de Marneffe, Marie-Catherine, Filip Ginter, Yoav Goldberg, Jan Hajič, Christopher Manning, Ryan McDonald, Joakim Nivre, Slav Petrov, Sampo Pyysalo, Sebastian Schuster, Natalia Silveira, Reut Tsarfaty, Francis Tyers, and Dan Zeman. 2019. Universal dependencies. <https://universaldependencies.org/>. Accessed: 2019-09-30.
- Mel'čuk, Igor. 1988. *Dependency Syntax: Theory and Practice*. State University of New York Press, Albany, NY, USA.

- Nivre, Joakim. 2006. Constraints on non-projective dependency parsing. In *EACL 2006 - 11th Conference of the European Chapter of the Association for Computational Linguistics, Proceedings of the Conference*, pages 73–80.
- Nivre, Joakim. 2009. Non-projective dependency parsing in expected linear time. In *Proceedings of the Joint Conference of the 47th Annual Meeting of the ACL and the 4th International Joint Conference on Natural Language Processing of the AFNLP: Volume 1 - Volume 1, ACL '09*, pages 351–359, Association for Computational Linguistics, Stroudsburg, PA, USA.
- Rudnicka, Karolina. 2018. Variation of sentence length across time and genre: influence on the syntactic usage in English. In Richard J. Whitt, editor, *Diachronic corpora, genre, and language change*. pages 219–240.
- Shiloach, Yossi. 1979. A minimum linear arrangement algorithm for undirected trees. *SIAM Journal on Computing*, 8(1):15–32.
- Sleator, Daniel and Davy Temperley. 1993. Parsing English with a link grammar. In *Proceedings of the Third International Workshop on Parsing Technologies (IWPT'93)*, pages 277–292, ACL/SIGPARSE.
- Sloane, N. J. A. 1964-2021. The on-line encyclopedia of integer sequences – amount of unlabeled rooted trees. <https://oeis.org/A000081>. Accessed: 2020-11-23.
- Zörnig, P. 1984. The distribution of the distance between like elements in a sequence I. *Glottometrika*, 25(6):1–15.