

Projective representation of the Galilei group for classical and quantum-classical systems

A. D. Bermúdez Manjarres

*Universidad Distrital Francisco José de Caldas
Cra 7 No. 40B-53, Bogotá, Colombia
ad.bermudez168@uniandes.edu.co*

September 17, 2021

Abstract

A physically relevant unitary irreducible non-projective representation of the Galilei group is possible in the Koopman-von Neumann formulation of classical mechanics. This classical representation is characterized by the vanishing of the central charge of the Galilei algebra. This is in contrast to the quantum case where the mass plays the role of the central charge. Here we show, by direct construction, that classical mechanics also allows for a projective representation of the Galilei group where the mass is the central charge of the algebra. We extend the result to certain kind of quantum-classical hybrid systems.

1 Introduction

It was realized long time ago that classical mechanics can be stated in the same mathematical formalism of quantum mechanics, i.e., as a theory of operators acting on a Hilbert space [1, 2]. This operational formalism is known as the Koopman-von Neumann (KvN) theory.

Inspired by the KvN theory (though with a more general scope), a procedure to obtain classical dynamics as a unitary irreducible representation of the Galilei group was given in Ref [3]. An interesting aspect of this classical unitary representation is that the central charge of the Galilei algebra vanishes. This is, contrary to the quantum case, the classical unitary representation of the Galilei group considered in [3] was not projective. However, while the KvN rule makes unavoidable the vanishing of the central charge, classical dynamics does not demand it, a possibility that was overlooked.

Here we consider a variation of the KvN theory recently named as the Koopman-von Hove theory (KvH) [4, 5]. Although equally valid at the classical level, the KvH theory puts more emphasis on the phase of the classical Koopman wavefunctions.

We will show that the KvH leads to a projective representation of the Galilei group where the mass plays the role of the central charge of the theory, just as in quantum mechanics. This is, the KvH representation of the Galilei group is up to a phase.

Besides the purely classical results, the KvN and KvH formalism are of interest to contrast/compare quantum and classical mechanics and the study of quantization/dequantization rules [6, 7, 8, 9, 10]. The KvN and KvH theories are also of interest for the formulation of some quantum-classical hybrid theories [4, 5, 11, 12, 13, 14, 15, 16, 17, 18]. In this last regard, the knowledge of the non-relativistic space-time symmetries of the KvN theory was used to study the viability of quantum-classical hybrids of the Sudarshan type [19]. It was shown there that in Sudarshan hybrids, translational invariance does not equate to the conservation of the total linear momentum. It was concluded that Sudarshan hybrids were at odds with momentum conservation. Unfortunately, we will see that this conclusion was hastily made. We will show that the hybrid systems recently considered by Bondar, Gay-Balmaz, and Tronci [4] can be Galilei covariant and conserve the total momentum.

This work is organized as follows: in section 2 we review the KvN and KvH Hilbert space formulations of classical mechanics. We will focus on the construction of Hermitian operators from phase space functions.

In section 3 the elements of the Galilei algebra will be related to Hermitian operators constructed from the KvN and KvH. It is shown there that the KvH theory leads to a projective representation of the Galilei group.

In section 4 we analyze the Galilean covariance and momentum conservation of quantum-classical hybrid systems of the Sudarshan type obtained from the KvH formalism. We will see that the KvH theory leads naturally to a hybrid interaction term that conserves the total momentum and is not at odds with Galilean covariance.

2 The KvN and KvH mechanics

It is possible to associate a Hilbert \mathcal{H}_{cl} space to a given a phase space Γ . The procedure is as follows: let $d\omega$ be the usual symplectic measure of Γ , the vectors of \mathcal{H}_{cl} are then given by square integrable functions

$$\int_{\Gamma} |\psi|^2 d\omega < \infty. \quad (1)$$

The inner product on \mathcal{H}_{cl} can be defined as

$$\langle \psi_1 | \psi_2 \rangle = \int_{\Gamma} \psi_1^* \psi_2 d\omega. \quad (2)$$

From now on, we will specialize for the case of a single particle moving in R^3 , such that $d\omega = d\mathbf{q}d\mathbf{p}$ in Cartesian coordinates.

There are infinite ways to use phase space functions to define linear operators acting on \mathcal{H}_{cl} . The easiest method to define a Hermitian operator from a space space function is by multiplication as follows:

$$\hat{f}(\mathbf{q}, \mathbf{p})\psi(\mathbf{q}, \mathbf{p}) = f(\mathbf{q}, \mathbf{p})\psi(\mathbf{q}, \mathbf{p}). \quad (3)$$

The most important operators defined this way are the position and momentum operators

$$\begin{aligned} \hat{\mathbf{q}}f(\mathbf{q}, \mathbf{p}) &= \mathbf{q}f(\mathbf{q}, \mathbf{p}), \\ \hat{\mathbf{p}}f(\mathbf{q}, \mathbf{p}) &= \mathbf{p}f(\mathbf{q}, \mathbf{p}). \end{aligned} \quad (4)$$

Let us note that these so-defined position and momentum operators commute with each other $[q_i, p_j] = 0$; they do not have any uncertainty principle associated to them. Thus, they can be simultaneously measured, as they should in any classical theory. We also point out that $\hat{\mathbf{q}}$ and $\hat{\mathbf{p}}$ form a complete set of commuting observables since we are dealing with a single structureless particle.

There are other ways to define Hermitian operators from phase space functions. A rather natural way comes from using the Poisson bracket as follows:

$$\hat{F} = -i \{ \cdot, f(\mathbf{q}, \mathbf{p}) \}. \quad (5)$$

We will call the procedure given by Eq.(5) as the KvN rule. Using the the KvN rule on (\mathbf{q}, \mathbf{p}) we get

$$\begin{aligned} \hat{\lambda}_{\mathbf{q}} &= -i \{ \cdot, \mathbf{p} \} = -i \nabla_{\mathbf{q}}, \\ \hat{\lambda}_{\mathbf{p}} &= i \{ \cdot, \mathbf{q} \} = -i \nabla_{\mathbf{p}}. \end{aligned} \quad (6)$$

From their definition, we can write the following set of commutation relations

$$\begin{aligned} [\hat{q}_i, \hat{q}_j] &= [\hat{q}_i, \hat{p}_j] = [\hat{p}_i, \hat{p}_j] = 0, \\ [\hat{q}_i, \hat{\lambda}_{p_i}] &= [\hat{p}_i, \hat{\lambda}_{q_j}] = [\hat{\lambda}_{q_j}, \hat{\lambda}_{p_i}] = 0, \\ [\hat{q}_i, \hat{\lambda}_{q_j}] &= [\hat{p}_i, \hat{\lambda}_{p_j}] = i\delta_{ij}. \end{aligned} \quad (7)$$

The set $(\hat{\mathbf{q}}, \hat{\mathbf{p}}, \hat{\lambda}_{\mathbf{q}}, \hat{\lambda}_{\mathbf{p}})$ is irreducible on \mathcal{H}_{cl} in view of commutation relations (7) and Schur lemma.

Acting on the Hamiltonian function $H = \frac{\mathbf{p}^2}{2m} + V(\mathbf{q})$, the KvN rule gives the following operator

$$\hat{L} = -i \{ \cdot, H \} = \frac{1}{m} \hat{\mathbf{p}} \cdot \hat{\lambda}_{\mathbf{q}} - \nabla_{\mathbf{q}} V(\mathbf{q}) \cdot \hat{\lambda}_{\mathbf{p}}. \quad (8)$$

The operator \hat{L} is known as the Liouvillian, and it is of central importance since it governs time evolution. Consider the Liouville equation of classical statistical mechanics

$$\frac{\partial \rho}{\partial t} = - \{ \rho, H \} = -i \hat{L} \rho. \quad (9)$$

Due to the linearity in $\hat{\lambda}_{\mathbf{q}}$ and $\hat{\lambda}_{\mathbf{p}}$ of (9), the square integrable function defined by ¹ $\psi = \sqrt{\rho}e^{iS}$ also obeys the Liouville equation

$$\frac{\partial\psi}{\partial t} = -i\hat{L}\psi. \quad (10)$$

The equation (10) has the form of a Schrödinger equation, where \hat{L} takes the role of the ‘‘Hamiltonian’’ operator of the theory. Hence, classical statistical mechanics is equivalent to solving the Schrödinger-like equation (10). Particle mechanics and Hamilton equations are formally recovered with the use Klimontovich delta distributions, i.e., $\rho = \delta(\mathbf{p}(t), \mathbf{q}(t))$.

Another viable formula for assigning Hermitian operators to phase space functions is

$$\hat{F}^* = -i\{\cdot, f\} + f - \mathbf{p} \cdot \frac{\partial f}{\partial \mathbf{p}}. \quad (11)$$

After the convention used in [4, 5], we call the association (11) as the KvH rule². Applying the KvH rule on the Hamiltonian we get

$$\hat{L}^* = -i\{\cdot, H\} + H - \mathbf{p} \cdot \frac{\partial H}{\partial \mathbf{p}} = \hat{L} - \mathcal{L}, \quad (12)$$

where \hat{L} is the operator given by (8), and $\mathcal{L} = \mathbf{p} \cdot \frac{\partial H}{\partial \mathbf{p}} - H$ is the Lagrangian associated to H .

Acting on $\psi = \sqrt{\rho}e^{iS}$, the Schrodinger-like equation generated by \hat{L}^*

$$\frac{\partial\psi}{\partial t} = -i\hat{L}^*\psi, \quad (13)$$

splits into

$$\begin{aligned} \frac{\partial\rho}{\partial t} + \{\rho, H\} &= 0, \\ \frac{\partial S}{\partial t} + \{S, H\} &= \mathcal{L}. \end{aligned} \quad (14)$$

So, the KvH dynamics is equivalent to the classical Liouville equation, but, contrary to the KvN case, the KvH equation also contain information about the classical phase.

It is worth mentioning an important third possibility to associate Hermitian operators to phase space functions, although we will not use it in this work. Instead of the Poisson bracket, the star product of Wigner phase space representation of quantum mechanics can be used [20]. The use of the star product leads to quantum representation of the Galilei group.

¹A constant is needed here to make the exponent dimensionless. This constant is usually chosen to be \hbar^{-1} . In this work we set $\hbar=1$.

²There is a gauge freedom in the assignation of the Hermitian operator given by the KvH rule, see Ref [4], but we will stick with the equation (11) for the rest of this work.

3 The KvN and KvH Representations of the Galilei group

In Hamiltonian mechanics, the space-time transformations associated with the Galilei group are realized via canonical transformations. The generator functions associated with these canonical transformations are the linear momentum \mathbf{p} , the angular momentum $\mathbf{j} = \mathbf{r} \times \mathbf{p}$, the dynamic mass moment $\mathbf{g} = m\mathbf{q} - t\mathbf{p}$ and the Hamiltonian of the free particle $H_{free} = \frac{\mathbf{p}^2}{2m}$. They realize the Lie algebra of the Galilei group via the Poisson bracket. This is known as a canonical representation of a Lie group [21, 22].

The canonical representation of the Galilei group can be reinterpreted as a unitary representation in the context of the KvN theory as follows: first, the operator $\hat{\lambda}_{\mathbf{q}}$ is the generator of translation. Its action on an element of $\mathcal{H}(\Gamma)$ is given by

$$\exp \left[-i\mathbf{a} \cdot \hat{\lambda}_{\mathbf{q}} \right] \psi(\mathbf{q}, \mathbf{p}) = \psi(\mathbf{q} - \mathbf{a}, \mathbf{p}). \quad (15)$$

On the other hand, while $\hat{\lambda}_{\mathbf{p}}$ translate the momentum coordinates, $\exp \left[-i\mathbf{b} \cdot \hat{\lambda}_{\mathbf{p}} \right] \psi(\mathbf{q}, \mathbf{p}) = \psi(\mathbf{q}, \mathbf{p} - \mathbf{b})$, $\hat{\lambda}_{\mathbf{p}}$ is not an element of the Galilei algebra.

The other generators of the Galilei algebra can be obtained with the KvN rule as

$$\begin{aligned} \hat{\mathcal{J}}_i &= -i \{, j_i \} = \varepsilon_{ijk} \left(\hat{q}_j \hat{\lambda}_{q_k} + \hat{p}_j \hat{\lambda}_{p_k} \right), \\ \hat{\mathcal{G}} &= -i \{, \mathbf{g} \} = -t \hat{\lambda}_{\mathbf{q}} - m \hat{\lambda}_{\mathbf{p}}, \\ \hat{L} &= -i \{, H_{free} \} = \frac{1}{m} \hat{\mathbf{p}} \cdot \hat{\lambda}_{\mathbf{q}}. \end{aligned} \quad (16)$$

In the KvN theory $\hat{\mathcal{J}}$ is the generator of rotations, $\hat{\mathcal{G}}$ is the generator of Galilean boosts, and \hat{L} is the time translation operator for the case of a free particle. The above operators obey the commutation relations of the Galilei algebra.

$$\begin{aligned} \left[\hat{\lambda}_{q_i}, \hat{\lambda}_{q_j} \right] &= \left[\hat{\mathcal{G}}_i, \hat{\mathcal{G}}_j \right] = \left[\hat{\mathcal{J}}_i, \hat{L} \right] = \left[\hat{\lambda}_{q_i}, \hat{L} \right] = 0; \\ \left[\hat{\mathcal{J}}_i, \hat{\mathcal{J}}_j \right] &= i\varepsilon_{ijk} \hat{\mathcal{J}}_k; \quad \left[\hat{\mathcal{J}}_i, \hat{\lambda}_{q_k} \right] = i\varepsilon_{ijk} \hat{\lambda}_{q_k}; \\ \left[\hat{\mathcal{J}}_i, \hat{\mathcal{G}}_j \right] &= i\varepsilon_{ijk} \hat{\mathcal{G}}_k; \quad \left[\hat{\mathcal{G}}_i, \hat{L} \right] = i\hat{\lambda}_{q_i}, \end{aligned} \quad (17)$$

and

$$\left[\hat{\mathcal{G}}_j, \hat{\lambda}_{q_i} \right] = 0. \quad (18)$$

Equations (17) and (18) are a realization of the Galilei algebra in the very particular case of a vanishing central charge, as evidenced by the vanishing right-hand-side of (18). Since the set $\left\{ \hat{\mathbf{q}}, \hat{\mathbf{p}}, \hat{\lambda}_{\mathbf{q}}, \hat{\lambda}_{\mathbf{p}} \right\}$ is irreducible in $\mathcal{H}(\Gamma)$, and

taking into account the vanishing of the central charge, the above constitutes a unitary irreducible faithful representation of the Galilei group.

The remaining space-time transformations of the Galilei group are realized as unitary transformation given by

$$\begin{aligned}\exp\left[-i\theta\hat{\mathbf{n}}\cdot\hat{\mathcal{J}}\right]\psi(\mathbf{q},\mathbf{p}) &= \psi(\mathbf{q}-\theta\hat{\mathbf{n}}\times\mathbf{q},\mathbf{p}-\theta\hat{\mathbf{n}}\times\mathbf{p}), \\ \exp\left[i\mathbf{v}\cdot\hat{\mathcal{G}}\right]\psi(\mathbf{q},\mathbf{p}) &= \psi(\mathbf{q}-\mathbf{v}t,\mathbf{p}-m\mathbf{v}), \\ \exp\left[-it\hat{L}\right]\psi(\mathbf{q},\mathbf{p}) &= \psi\left(\mathbf{q}-\frac{t}{m}\mathbf{p},\mathbf{p}\right).\end{aligned}\quad (19)$$

We now proceed to investigate the effect of the KvH rules on the representation theory of the Galilei group. The KvH Liouvillian in the free particle case is given by

$$\hat{L}^* = -i\{H_{free}\} - H_{free} = \frac{\hat{\mathbf{p}}}{m}\cdot\hat{\lambda}_{\mathbf{q}} - \frac{\hat{\mathbf{p}}^2}{2m}.\quad (20)$$

The boost operator is

$$\hat{\mathcal{G}}_j^* = -i\{\mathbf{g}\} + \mathbf{g} - \mathbf{p}\cdot\frac{\partial\mathbf{g}}{\partial\mathbf{p}} = -t\hat{\lambda}_{\mathbf{q}} - m\hat{\lambda}_{\mathbf{p}} + m\mathbf{q}.\quad (21)$$

All the other generators are unaltered compared to the KvN case. It can be checked that the replacement $\hat{L}\rightarrow\hat{L}^*$ and $\hat{\mathcal{G}}\rightarrow\hat{\mathcal{G}}^*$ gives a realization of the Galilei algebra. However, contrary to the KvN case, now the central charge of the algebra does not vanish. Instead, we have the commutation relation

$$\left[\hat{\mathcal{G}}_j^*,\hat{\lambda}_{q_i}\right] = im\delta_{ij}.$$

We can see that the mass is the central charge in the KvH theory, just as in quantum mechanics. Hence, the unitary representation of the Galilei group given by the KvH theory is projective.

In the KvH theory, the unitary transformation associated to space translation and rotations are the same as in Eqs.(15) and (19). Galilean boosts and time translation are modified to include a phase factor as follows

$$\begin{aligned}\exp\left[i\mathbf{v}\cdot\hat{\mathcal{G}}^*\right]\psi(\mathbf{q},\mathbf{p}) &= e^{i(mq-\frac{m^2t}{2})}\psi(\mathbf{q}-\mathbf{v}t,\mathbf{p}-m\mathbf{v}), \\ \exp\left[-it\hat{L}^*\right]\psi(\mathbf{q},\mathbf{p}) &= e^{-i\frac{\mathbf{p}^2}{2m}t}\psi\left(\mathbf{q}-\frac{t}{m}\mathbf{p},\mathbf{p}\right).\end{aligned}\quad (22)$$

Finally, we mention that while $\hat{\lambda}_{\mathbf{q}}$ is the same in both theories, $\hat{\lambda}_{\mathbf{p}}$ is not. In the KvH theory we have $\hat{\lambda}_{\mathbf{p}}^* = \hat{\lambda}_{\mathbf{p}} + \mathbf{q}$ and $[\hat{\lambda}_{\mathbf{q}}^*,\hat{\lambda}_{\mathbf{p}}^*] = -i$.

3.1 Covariance of composite systems.

Let us now consider the case of two interacting particles. We can to construct a representation of the Galilei group on the space $\mathcal{H}_T = \mathcal{H}_{cl}^{(1)} \otimes \mathcal{H}_{cl}^{(2)}$ starting from the representation of the individual sectors. Except for the Liouvillian, the generators of the algebra acting on \mathcal{H}_T are given by the sum of the individual generators; for example, the total translation operator is given by $\hat{\lambda}_{\mathbf{q}total} = \hat{\lambda}_{\mathbf{q}_1} + \hat{\lambda}_{\mathbf{q}_2}$. All the commutation relations of the Galilei algebra not involving the Liouvillian are satisfied identically. On the other hand, the total Liouvillian is allowed to have an extra term representing the interaction between the particles. The allowed interaction term can then be deduced from considerations of Galilean covariance. Using the KvH rule on the Hamiltonian

$$H = \frac{\mathbf{p}_1^2}{2m_1} + \frac{\mathbf{p}_2^2}{2m_2} + V(\mathbf{q}_1, \mathbf{q}_2)$$

gives the Liouvillian

$$\begin{aligned} \hat{L}^* &= \frac{1}{m_1} \hat{\mathbf{p}}_1 \cdot \hat{\lambda}_{\mathbf{q}_1} + \frac{1}{m_2} \hat{\mathbf{p}}_2 \cdot \hat{\lambda}_{\mathbf{q}_2} \\ &\quad - \nabla_{\mathbf{q}_1} V \cdot \hat{\lambda}_{\mathbf{p}_1} - \nabla_{\mathbf{q}_2} V \cdot \hat{\lambda}_{\mathbf{p}_2} \\ &\quad - \frac{\hat{\mathbf{p}}_1^2}{2m_1} - \frac{\hat{\mathbf{p}}_2^2}{2m_2} + V. \end{aligned} \quad (23)$$

Translational invariance dictates that

$$\left[\hat{\lambda}_{\mathbf{q}total}, \hat{L}^* \right] = 0. \quad (24)$$

The necessary and sufficient condition for Eq (24) to be true is that the potential energy depend only on the relative position $V = V(\hat{\mathbf{q}}_1 - \hat{\mathbf{q}}_2)$. The above also takes care of the commutation relation with the Galilean boost

$$\left[\hat{\mathcal{G}}_{total}^*, \hat{L}^* \right] = \hat{\lambda}_{\mathbf{q}_1} + \hat{\lambda}_{\mathbf{q}_2}. \quad (25)$$

Finally, if V is a scalar then

$$\left[\hat{\mathcal{J}}_{total}, \hat{L} \right] = 0. \quad (26)$$

Notice that the condition of translational invariance is not the same as conservation of the total momentum $\hat{\mathbf{p}}_{total} = \hat{\mathbf{p}}_1 + \hat{\mathbf{p}}_2$. However, it is true that $V = V(\hat{\mathbf{q}}_1 - \hat{\mathbf{q}}_2)$ leads to

$$\left[\hat{\mathbf{p}}_{total}, \hat{L}^* \right] = 0. \quad (27)$$

4 Quantum-classical hybrids

It is possible to obtain the two-body Schrödinger equation from the Liouvillian (23). Following Klein [9], the procedure consist of two steps: (1) impose the condition $\partial_p \psi = 0$ for every momentum coordinate, (2) make the replacement the replacement $p \rightarrow -i\partial_q$, this correspond to the usual canonical quantization rule. For the Liouvillian (23), the result is

$$i\frac{\partial\psi}{\partial t} = -\frac{1}{2m_1}\nabla_{\mathbf{q}_1}^2\psi - \frac{1}{2m_2}\nabla_{\mathbf{q}_2}^2\psi + V\psi. \quad (28)$$

Instead of doing a full quantization of (23), we can obtain a hybrid theory by applying Klein's method to only one particle. This will result in a special case of an hybrid theory of the Sudarshan type [4, 5, 11]. To distinguish the classical degrees of freedom from the quantum ones, we keep the (\mathbf{q}, \mathbf{p}) notation for the classical particle and use \mathbf{x} and \mathbf{k} for the quantum coordinate and momentum, respectively. The quantum position and momentum operators obeys the usual Heisenberg commutation relations $[\hat{x}_i, \hat{k}_j] = i\delta_{ij}$. The hybrid Hilbert space is of the form $\mathcal{H}_T = \mathcal{H}_Q \otimes \mathcal{H}_{cl}$, where the vectors of the quantum sector \mathcal{H}_Q are given by normalizable linear combinations of the kets $|\mathbf{x}\rangle$.

In the quantum sector, and for a single spinless particle, the elements of the Galilei algebra in terms \mathbf{x} of \mathbf{k} are: \mathbf{k} itself, the angular momentum $\hat{\mathbf{j}} = \hat{\mathbf{x}} \times \hat{\mathbf{k}}$, the dynamic mass moment $\hat{\mathbf{g}} = m\hat{\mathbf{x}} - t\hat{\mathbf{k}}$, and the Hamiltonian $\hat{H}_Q = \frac{\hat{\mathbf{k}}^2}{2m}$, where m is the central charge of the algebra [23, 24, 25, 26]. In the quantum case, \mathbf{k} has the dual role of being the momentum operator and the generator of spatial translations.

Performing the partial quantization of (23) results in the hybrid total time-translation operator

$$\begin{aligned} \hat{L}_h &= \frac{1}{m_1}\hat{\mathbf{p}} \cdot \hat{\lambda}_{\mathbf{q}} - \frac{\hat{\mathbf{p}}^2}{2m_1} + \frac{1}{2m_2}\hat{\mathbf{k}}^2 \\ &\quad - \nabla_{\mathbf{q}}V \cdot \hat{\lambda}_{\mathbf{p}} + V. \end{aligned} \quad (29)$$

As in section 3.1, the commutation relations of the Galilei algebra of the composite systems not involving the time translation operator are all satisfied identically, and the ones that do involve \hat{L}_h give conditions for the allowed interaction terms. We emphasize here that in general Sudarshan hybrids, the conditions for Galilean covariance does not imply the conservation of quantities like the linear or the angular momentum. For example, an interaction term can be translational invariant without conserving the total linear momentum [19]. However, we will focus our attention on the particular interaction term given in (29).

A translational invariant potential $V = V(\hat{\mathbf{q}} - \hat{\mathbf{x}})$ guarantees the translational invariance of the hybrid Liouvillian (29)

$$[\hat{\lambda}_{\mathbf{q}_1} + \hat{\mathbf{k}}, \hat{L}_h] = 0, \quad (30)$$

and the conservation of the total momentum

$$\left[\hat{\mathbf{p}} + \hat{\mathbf{k}}, \hat{L}_h \right] = 0. \quad (31)$$

Rotational invariance is obtained if V is a scalar.

With $\hat{\mathcal{G}}_{hybrid} = m_2 \hat{\mathbf{x}} - t \hat{\mathbf{k}} - t \hat{\lambda}_{\mathbf{q}} - m_1 \hat{\lambda}_{\mathbf{p}} + m_1 \mathbf{q}$, it is immediate to check that the central charge of the algebra is the total mass

$$\left[\hat{\mathcal{G}}_{hybrid}, \hat{\lambda}_{\mathbf{q}_1} + \hat{\mathbf{k}} \right] = m_1 + m_2. \quad (32)$$

Hence, the hybrid representation of the Galilei group is projective.

It is worth emphasizing that it is the combination $-\nabla_{\mathbf{q}} V \cdot \hat{\lambda}_{\mathbf{p}} + V$ in (29) that allows translational invariance and the conservation of the linear momentum at the same time. Neither V nor $-\nabla_{\mathbf{q}} V \cdot \hat{\lambda}_{\mathbf{p}}$ do the trick on their own.

5 Conclusions

In quantum mechanics, the central charge of the Galilei algebra is the mass, and it plays the role of a superselection operator. In the quantum case, representations with a vanishing central charge are unphysical. We have shown that in the KvH representation of the Galilei group the mass is the central charge. It is interesting that classical mechanics allows, but does not require, projective unitary representations of the Galilei group. At the classical level, whether the central charge vanishes or not has no effect since the KvN and the KvH formalism are equally valid.

For hybrid systems, the KvH rule and the Klein quantization method lead to the specific interaction term $-\nabla_{\mathbf{q}} V \cdot \hat{\lambda}_{\mathbf{p}} + V$ that guarantees Galilean covariance and conservation of the total momentum.

References

- [1] B. O. Koopman, Proc. Natl. Acad. Sci. U.S.A. **17**, 315 (1931).
- [2] J. von Neumann, Ann. Math. **33**, 587 (1932).
- [3] A. D. Bermudez Manjarres, M. Nowakowski and D. Batic, Ann. Phys (NY), **416**, 168157 (2020).
- [4] D. I. Bondar, F. Gay-Balmaz and C. Tronci, Proc. R. Soc. A, **475**, (2019).
- [5] F. Gay-Balmaz and C. Tronci, Nonlinearity, **33** 5383 (2020).
- [6] D. I. Bondar, R. Cabrera, R. Lompay, M.Y Ivanov and H.A. Rabitz, Phys. Rev. Lett. **109**, 190403 (2012).
- [7] A. A. Abrikosov, Jr., E. Gozzi, and D. Mauro, Ann. Phys. (N.Y.) **317**, 24 (2005).

- [8] D. Mauro, Phys. Lett. A **315**, (2003), 28-35.
- [9] U. Klein, Quantum Stud.: Math. Found. 5, 219-227 (2018)
- [10] B. Kostant, Line bundles and the prequantized Schrödinger equation. In Colloquium on group theoretical methods in physics. Centre de Physique Théorique, Marseille, IV.1–IV.22, (1972).
- [11] E. C. G. Sudarshan, Pramana **6**, 117 (1976).
- [12] T. N. Sherry and E. C. G. Sudarshan, Phys. Rev. D **18**, 4580 (1978).
- [13] T. N. Sherry and E. C. G. Sudarshan, Phys. Rev. D **20**, 857 (1979).
- [14] S. R. Gautam, T. N. Sherry, and E. C. G. Sudarshan, Phys. Rev. D **20**, 3081 (1979).
- [15] C. Barceló, R. Carballo-Rubio, L.J Garay and R. Gomez-Escalante, Phys. Rev. A **86**, 042120 (2012).
- [16] A. Peres and D. R. Terno, Phys. Rev. A **63**, 022101 (2001).
- [17] M. Radonjic *et al*, Phys. Rev. A **89**, 024104 (2014).
- [18] H.R Jauslin, D. Sugny, Dynamics of mixed classical-quantum systems, geometric quantization and coherent states. In Mathematical horizons for quantum physics. Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore, no. 20, pp. 65–96. Singapore: World Scientific, (2010).
- [19] A. D. Bermúdez Manjarres and N. Marín-Medina, Phys. Rev. A. **102**, 042221 (2020).
- [20] M.D. Oliveira, M.C.B. Fernandes, F.C. Khanna, A.E. Santana, and J.D.M. Vianna, Ann. Phys (NY), **312** (2004) 492–510.
- [21] M. Pauri and G. Prosperi, J. Math. Phys. **9**, 1146 (1968) .
- [22] E.C.G. Sudarshan and N. Mukunda, “Classical Dynamics : A Modern Perspective”, (John Wiley & Sons, New York, 1974).
- [23] T. F. Jordan, Linear Operators for Quantum Mechanics , (John Wiley & Sons, New York; 1969).
- [24] L. E. Ballentine, Quantum Mechanics: A Modern Development, (World Scientific Publishing, Singapore; 1998).
- [25] J. -M. Lévy-Leblond, Galilei group and Galilean invariance, Group Theory and its Applications , Vol. 2, (Academic press: New York ; 1971).
- [26] J. M. Jauch, Foundation of Quantum Mechanics, (Addison-Wesley Publishing Co, Reading, Massachussetts; 1968).