

Bökstedt periodicity generator via K -theory

A. Fonarev and D. Kaledin*

Contents

1	Generalities.	4
1.1	Stabilization.	4
1.2	Additivization.	6
1.3	THH and K-theory.	8
2	The construction.	10
2.1	The object.	10
2.2	The trace computation.	11
2.3	The cohomology class.	12

Introduction.

In its most basic form, “Bökstedt periodicity” refers to the computation of the Topological Hochschild Homology $THH_*(k)$ of a prime field $k = \mathbb{F}_p$ with p elements; the answer is that $THH_*(k) \cong k[v]$ is the polynomial algebra in one generator v of homological degree 2. The computation was done by Bökstedt [Bö] in the same paper where Topological Hochschild Homology was first defined. Slightly later, in a wonderful series of papers [FPSVW], [JP], [PW] stemming from [P1], Pirashvili and others showed that $THH_*(k)$ can be identified with several other homology theories including Mac Lane Homology $HM_*(k)$, where the same periodicity has been earlier established by Breen [Br]. A good overview of the whole story as it was seen in 1998 can be found in [P2], and a more recent survey is available in [HN].

As of now, Bökstedt periodicity has numerous applications, including some very prominent ones such as [BMS], and several proofs none of which are elementary. The latter is somewhat surprising, given that the question is

*Both authors supported by the Basis Foundation, grant 18-1-6-95-1, Leader (Math).

so fundamental, and the answer is so miraculously simple. What happens is, one can easily construct a spectral sequence converging to $THH(k)$ whose E_2 -term can be read off the dual Steenrod algebra, and the sequence degenerates at E_p . However, if p is not 2, then the E_2 -term is rather large, and while it already contains v , there are many extra elements that are killed off by the differential d_{p-1} . So, the spectral sequence is highly non-degenerate. One has to prove this somehow, and then show that $THH(k) \cong k[v]$ as a k -algebra (this is also non-trivial since in the E_p term, $v^p = 0$). Roughly speaking, the existing proofs belong to one of the following two types.

- (i) “Topological” proofs that use Dyer-Lashof operations. This was the original approach of Bökstedt [Bö]. Recently, a pretty definitive proof along the same lines was given in [KN] that reduces the claim to the following amazing one-line observation: the dual Steenrod algebra for k is freely generated by one element of degree 1 as an E_2 -algebra. This yields Bökstedt periodicity almost immediately, and the observation itself follows from classical computations of Dyer-Lashof operations.
- (ii) “Representation-theoretic” proofs. These use the identification between THH and Mac Lane Homology, and then work with Mac Lane Cohomology instead of Homology, and use polynomial functors and modular representation theory of $GL(n)$. The original proof along these lines is [FLS], and it can be considerably simplified using more recent technology (see e.g. [Ka, Section 11]).

In our opinion, none of these proofs is so satisfactory as to make the other ones obsolete. If one just wants shortness, it is hard to beat [KN]. However, while the argument proves the theorem, it does not explain the miracle but moves it one step up. Why would the dual Steenrod algebra, an E_∞ -algebra by nature, become so simple when considered as an E_n -algebra for $n = 2$? One has a feeling that there should be a categorical reason for this, but at the moment, it is not clear, at least to us, what it might be. Also, the duality between Mac Lane Homology and Mac Lane Cohomology is not visible in the topological approach, and as far as we know, a possible cohomological counterpart to THH has not been properly studied.

In the representation-theoretic approach, it is the cohomology that plays the main role, and the non-degeneration of the spectral sequence acquires some sort of an explanation. At the end of the day, the main fact is that the category of strictly polynomial functors of some fixed degree has finite homological dimension. This line of reasoning actually leads quite far — for example, to the celebrated finiteness results of Friedlander and Suslin [FS]

and more recent strong results of Touzé [T]. However, there is a price to pay: while non-degeneration for cohomology is equivalent to non-degeneration for homology, this approach says absolutely nothing about multiplication in homology. We obtain a graded vector space isomorphism $THH_*(k) \cong k[v]$, but to check that it is multiplicative, one needs an additional argument.

One possible argument goes as follows. Firstly, for dimension reasons, it suffices to check that the generator v in $THH_2(k)$ is not nilpotent. To do this, it suffices to construct a multiplicative map $\varphi_* : THH_*(k) \rightarrow R_*(k)$ to some graded algebra $R_*(k)$ such that $\varphi_*(v)^n \neq 0$ for all $n \geq 0$. A tempting option is to take as $R_*(k)$ the periodic cyclic homology $HP_*(k)$, with the multiplicative map $\varphi_* : THH_*(k) \rightarrow HP_*(k)$ extracted from the cyclotomic structure map on $THH_*(k)$ of [NS] by the method suggested in [M]. We have $HP_*(k) \cong k((u))$, for a generator u of cohomological degree 2, and *a posteriori*, $\varphi(v) = u^{-1}$; were we to know this *a priori*, this would be it.

The argument in [Ka, Section 11] goes along the same lines but in a slightly different way. Instead of $HP_*(k)$, one takes $R_*(k) = \check{H}^*(\mathbb{Z}/p\mathbb{Z}, k)$, the Tate cohomology algebra of the cyclic group $\mathbb{Z}/p\mathbb{Z}$ with coefficients in the trivial module k . It also has a generator u of cohomological degree 2. To construct a multiplicative map φ_* , one uses yet another reinterpretation of $THH_*(k)$ — namely, the identification, essentially due to [JP], between $THH_*(A, M)$, for an algebra A and an A -bimodule M , and the *stabilization*, a.k.a. *additivization*, a.k.a. the first Goodwillie derivative of the cyclic nerve functor $|N^{cy}(F(A), F(M))|$, where $F(A)$ is the category of finitely generated free A -modules, and $F(M)$ is an $F(A)$ -bimodule corresponding to M . If $A = k$, then it is easy to write a functorial multiplicative map $\varphi : |N^{cy}(F(k), F(M))| \rightarrow C(M)$ to the p -th cyclic power $C(M) = (M^{\otimes p})^{\mathbb{Z}/p\mathbb{Z}}$ of a vector space M , and then one shows that the stabilization of $C(M)$ is the truncation $\tau^{\geq 0} \check{H}^*(\mathbb{Z}/p\mathbb{Z}, M^{\otimes p})$ of the Tate cohomology algebra. Stabilizing φ gives our map φ_* , and it remains to show that up to a non-zero constant, $\varphi_*(v) = u^{-1}$. This is a computation in homological degree 2, so it should not be too difficult.

However, there is a catch: it turns out that it *is* difficult. In the end, [Ka] resorts to a workaround: by plugging in some additional structure — namely, the Connes-Tsygan differential B on $THH_*(k)$ — one can further reduce the computation to something in homological degree 0. That one, fortunately, is indeed quite easy. However, the Connes-Tsygan differential in question only exists as a map of spectra, and it in fact vanishes on individual homotopy groups. This makes the resulting argument somewhat delicate, and requires rather strong stabilization technology to prove that everything is compatible with everything else.

The goal of this note is to suggest an alternative approach to proving that the map φ , above indeed sends the Bökstedt periodicity generator v to a non-trivial multiple of u^{-1} . We use a slightly different stabilization construction of THH : instead of cyclic nerves, we stabilize the so-called *K-theory with coefficients*, as suggested and done in [DM]. This is much simpler technically since cyclic nerves typically have higher homotopy groups, while *K-theory with coefficients* in our setting essentially reduces to K_0 (“essentially” means “after localization at p ”, and this is harmless since $THH(k)$ is p -local, see Lemma 1.3). We then show that φ extends to *K-theory with coefficients*, and exhibit an explicit degree-2 class v in the stabilization of $K_0(k, M)$ whose image $\varphi_2(v)$ is u^{-1} . This proves, yet again, Bökstedt periodicity (and shows that our explicit class is a multiplicative generator). The construction is made possible by a couple of additional simplifications, discussed below in Section 2. The actual element in *K-theory* that we construct is quite simple, it is a module of dimension 2 (and we need to exclude the case $p = 2$). Curiously, the construction depends on an additional parameter $\lambda \in k \setminus \{0, 1\}$, and we were *not* able to pinpoint a value for the parameter that makes things work; all we prove is that such a value exists.

The note is split into three parts. The first one is this introduction. Then Section 1 contains all the necessary generalities and preliminaries, and Section 2 is devoted to the actual construction. All the material in Section 1 is standard, but for lack of convenient references, we follow the exposition in [Ka, Section 4] for everything related to stabilization.

1 Generalities.

1.1 Stabilization. For any small category \mathfrak{l} , denote by $\mathrm{Ho}(I)$ the category of functors from I to the category Top_+ of pointed compactly generated topological spaces localized with respect to pointwise weak equivalences (where the localization can be constructed by any of the standard methods, e.g. by model embeddings of [DHKS], and if one wishes, one can replace Top_+ with the category of pointed simplicial sets). If we let $\mathrm{Ho} = \mathrm{Ho}(\mathrm{pt})$ be the localization of Top_+ itself, then any $X \in \mathrm{Ho}(I)$ defines a functor $\mathrm{Ho}(X) : I \rightarrow \mathrm{Ho}$, $i \mapsto X(i)$. A functor $\gamma : I' \rightarrow I$ between small categories induces a pullback functor $\gamma^* : \mathrm{Ho}(I) \rightarrow \mathrm{Ho}(I')$, $X \mapsto X \circ \gamma$.

Say that a category I is *half-additive* if it is pointed and has finite co-products, and for any half-additive small I , say that $X \in \mathrm{Ho}(I)$ is *additive* if for any $i, i' \in I$, the natural map

$$(1.1) \quad X(i \sqcup i') \rightarrow X(i) \times X(i')$$

is a weak equivalence. The simplest half-additive category is the category Γ_+ of finite pointed sets. In this case, $X \in \text{Ho}(\Gamma_+)$ is known as a Γ -space, and its additivity means being *special* in the sense of the Segal machine [S]. If for any integer $n \geq 0$ we denote by $n_+ \in \Gamma_+$ the set with n non-distinguished elements, then $X(1_+)$ for any special Γ -space X is a commutative monoid object in Ho , and $\pi_0(X(1_+))$ is an honest commutative monoid. The Γ -space X is called *group-like* if $\pi_0(X(1_+))$ is not just a monoid but a group. For any small category I , we say that $X \in \text{Ho}(\Gamma_+ \times I)$ is *additive* resp. *stable along* Γ_+ if it becomes additive resp. group-like after restriction to $\Gamma_+ \times i$ for any object $i \in I$. If I is half-additive, we have the smash-product functor $m : \Gamma_+ \times I \rightarrow I$ sending $n_+ \times i$ to the coproduct of n copies of i , $X \in \text{Ho}(I)$ is additive iff m^*X is additive along Γ_+ , and we say that X is *stable* if $m^*X \in \text{Ho}(\Gamma_+ \times I)$ is stable along Γ_+ .

For any half-additive small I , let $\text{Ho}^{st}(I) \subset \text{Ho}(I)$ be the full subcategory spanned by stable objects. Then the embedding $\text{Ho}^{st}(I) \rightarrow \text{Ho}(I)$ admits a left-adjoint *stabilization functor* $\text{Stab} : \text{Ho}(I) \rightarrow \text{Ho}^{st}(I)$. Again, the stabilization can be constructed by any of the standard methods; we follow the overview given in [Ka, Section 4]. Explicitly, one can extend $X : I \rightarrow \text{Top}_+$ to a functor $X^\Delta : \Delta^o I \rightarrow \text{Top}_+$ from simplicial objects in I by applying it pointwise and taking geometric realization. We then have

$$(1.2) \quad \text{Stab}(X)(i) \cong \text{colim}_n \Omega^n X^\Delta(m(\Sigma_n \times i)), \quad i \in I, X \in \text{Ho}(I),$$

where Σ_n is the standard simplicial n -sphere (that is, the finite pointed simplicial set obtained by taking the elementary n -simplex Δ_n and collapsing its $(n-1)$ -skeleton to the distinguished point). In particular, it is clear from (1.2) that Stab preserves finite products and finite homotopy limits. If I has a unital monoidal structure $- \otimes -$ that is distributive — that is, $(i_0 \sqcup i_1) \otimes i' \cong (i_0 \otimes i') \sqcup (i_1 \otimes i')$, and similarly in the second variable — then $\text{Ho}(I)$ acquires a convolution monoidal structure, $\text{Ho}(X) : I \rightarrow \text{Ho}$ is lax monoidal whenever $X \in \text{Ho}(I)$ is an algebra object, and in this case, $\text{Stab}(X)$ is also an algebra object in a natural way (see [Ka, Subsection 4.5]).

By the Segal machine, $\text{Ho}^{st}(\Gamma_+)$ is the category of connective spectra. In particular, it is an additive category, so that for any integer n and object $X \in \text{Ho}^{st}(\Gamma_+)$, we have the endomorphism $n \text{id} : X \rightarrow X$. If $n \geq 0$, then the induced endomorphism $n(X) : X(1_+) \rightarrow X(1_+)$ of the underlying infinite loop space $X(1_+)$ is immediately obvious from the Γ -space structure: it is given by the composition

$$(1.3) \quad X(1_+) \xrightarrow{\delta} X(1_+)^n \xrightarrow{e} X(n_+) \xrightarrow{\mu} X(1_+),$$

where δ is the diagonal map, μ is induced by the projection $n_+ \rightarrow 1_+$ sending all the non-distinguished elements in n_+ to the unique non-distinguished element in 1_+ , and e is inverse to the n -fold iteration of (1.2).

For any small I , the full subcategory $\mathrm{Ho}^{st}(\Gamma_+, I) \subset \mathrm{Ho}(\Gamma_+ \times I)$ spanned by objects stable along Γ_+ is a convenient model for the homotopy category of functors from I to connective spectra. The underlying functor to spaces is $\varepsilon_1^* X \in \mathrm{Ho}(I)$, where for any $n \geq 0$, we denote by $\varepsilon_n : I \rightarrow \Gamma_+ \times I$ the embedding onto $n_+ \times I$. For any $i \in I$, $\varepsilon_1(X)(i) = X(1_+ \times i)$ carries a structure of a group-like special Γ -space, and the functor $\mathrm{Ho}(\varepsilon_1^* X) : I \rightarrow \mathrm{Ho}$ lifts to a functor $\mathrm{Ho}^{st}(X) : I \rightarrow \mathrm{Ho}^{st}(\Gamma_+)$. The category $\mathrm{Ho}^{st}(\Gamma_+, I)$ is additive for any small I but this requires a proof; what is immediately obvious is a functorial endomorphism $n(X) : \varepsilon_1^* X \rightarrow \varepsilon_1^* X$, $n \geq 0$ obtained by replacing $X(1_+)$, $X(n_+)$ in (1.3) by $\varepsilon_1^* X$, $\varepsilon_n^* X$.

If I is half-additive, we have the functor $m^* : \mathrm{Ho}^{st}(I) \rightarrow \mathrm{Ho}^{st}(\Gamma_+, I)$. It is a fully faithful embedding, with a left-adjoint $\mathrm{Stab}_I : \mathrm{Ho}^{st}(\Gamma_+, I) \rightarrow \mathrm{Ho}^{st}(I)$ isomorphic to $\mathrm{Stab} \circ \varepsilon_1^*$ (see e.g. [Ka, Subsection 5.3], and specifically [Ka, Corollary 4.11]). Informally, one can also stabilize functors to connective spectra, and the underlying space of a spectral stabilization coincides with the space-level stabilization of the underlying space. For any stable X in $\mathrm{Ho}(I)$ and $i \in I$, $X(i) \cong \varepsilon_1^* m^* X(i)$ is canonically a connective spectrum, and if I is unital monoidal and $i = 1$ is the unit object, then $X(1)$ is a ring spectrum. Stabilization Stab_I commutes with finite products and finite homotopy limits, and by (1.3), for any $X \in \mathrm{Ho}^{st}(\Gamma_+, I)$ and integer $n \geq 0$, we have $\mathrm{Stab}(n(X)) = n(\mathrm{Stab}(X))$.

Example 1.1. An additive category I is certainly half-additive, so that all the stabilization machinery applies to $\mathrm{Ho}(I)$. If we further assume that I is Karoubi-closed, then we have the Dold-Kan equivalence $\Delta^o I \cong C_{\geq 0}(I)$ between simplicial objects in I and chain complexes in I concentrated in non-negative homological degrees. Then $m(\Sigma_n \times i)$ in (1.2) coincides with $B^n i$, the simplicial object that corresponds to i considered as a complex placed in homological degree n . Another simplification is that for any stable object $X \in \mathrm{Ho}^{st}(I)$, the map $n(m^* X)$ of (1.3) is simply induced by the endomorphism $n \mathrm{id} : \mathrm{Id} \rightarrow \mathrm{Id}$ of the identity functor $\mathrm{Id} : I \rightarrow I$. Indeed, since I is additive, the n -fold coproduct $m(n_+ \times i)$ of an object $i \in I$ with itself is canonically identified with the n -fold product i^n , and then the diagonal map δ in (1.3) is induced by the corresponding functorial map $\mathrm{id} \rightarrow m \circ \varepsilon_n$.

1.2 Additivization. For any commutative ring k , the stabilization construction has a k -linear homological counterpart. Namely, for any small

category I , let $\text{Fun}(I, k)$ be the abelian category of functors from I to the category $k\text{-mod}$ of k -modules, let $C_{\leq 0}(I, k)$ be the category of complexes in $\text{Fun}(\mathcal{E}, k)$ concentrated in non-negative homological degrees, and let $\text{Ho}(I, k)$ be the category obtained by inverting quasiisomorphisms in $C_{\leq 0}(I, k)$, so that $\text{Ho}(I, k)$ is the connective part of the standard t -structure on the derived category of $\text{Fun}(I, k)$. Alternatively, one can apply the Dold-Kan equivalence and interpret $C_{\leq 0}(I, k)$ as the category of functors from \mathcal{E} to simplicial k -modules, and invert pointwise weak equivalences. We then have the forgetful functor $\text{Ho}(I, k) \rightarrow \text{Ho}(I)$ sending a simplicial k -module to the underlying simplicial set.

If I is additive, then again, $X \in \text{Ho}(I, k)$ is additive iff (1.1) is a quasiisomorphism. For an additive $X \in \text{Ho}(\Gamma_+, k)$, the underlying Γ -space is automatically group-like, so that in the k -linear context, stability and additivity are one and the same. The embedding $\text{Ho}^{st}(I, k) \rightarrow \text{Ho}(I, k)$ of the full subcategory spanned by stable objects admits a left-adjoint stabilization, or additivization functor Stab . This is again compatible with monoidal structures, and can be explicitly computed by (1.2), where Ω^n now stands for the homological shift, and X^Δ is defined by taking X pointwise, then applying the Dold-Kan equivalence and totalizing the resulting bicomplex. The forgetful functor $\text{Ho}(I, k) \rightarrow \text{Ho}(I)$ commutes with stabilization. There are also alternative purely homological constructions of additivization, see [Ka, Subsection 4.6].

Example 1.2. Let k be a perfect field of some positive characteristic p , let I be the category of finite-dimensional k -vector spaces, and let $C \in \text{Ho}(I, k)$ be the *cyclic power functor* given by $C(M) = H^0(\mathbb{Z}/p\mathbb{Z}, M^{\otimes p})$, $M \in I$, where $\mathbb{Z}/p\mathbb{Z}$ acts by the longest cycle permutation. Then we have

$$(1.4) \quad \text{Stab}(C)(M) \cong \tau^{\leq 0} \check{H}^*(\mathbb{Z}/p\mathbb{Z}, M^{\otimes p}), \quad M \in \mathcal{E},$$

where \check{H}^* stands for Tate cohomology, and $\tau^{\leq 0}$ is the canonical truncation at 0 (also known as the “connective cover”). Roughly speaking — see [Ka, Lemma 5.1] for details — to compute the right-hand side of (1.4), one takes a projective resolution P_\bullet of the trivial $k[\mathbb{Z}/p\mathbb{Z}]$ -module k , lets P'_\bullet be the cone of the augmentation map $P_\bullet \rightarrow k$, and takes $H^0(\mathbb{Z}/p\mathbb{Z}, P'_\bullet \otimes M^{\otimes p})$ (up to a quasiisomorphism, this does not depend on the choice of P_\bullet). Then one checks that stabilizing $H^0(\mathbb{Z}/p\mathbb{Z}, P_\bullet \otimes M^{\otimes p})$ gives 0, and $\tau^{\leq 0} \check{H}^*(\mathbb{Z}/p\mathbb{Z}, M^{\otimes p})$ is stable. The cyclic power functor C is lax monoidal, so that its stabilization $\text{Stab}(C)$ is also lax monoidal, and $\text{Stab}(C)(k)$ carries a natural multiplication; this is the same multiplication as the usual multiplication in Tate

cohomology. If $p \geq 3$, then $\text{Stab}(C) \cong k[\varepsilon, u]$ is the free graded-commutative algebra with two generators ε, u of homological degrees $\deg \varepsilon = 1, \deg u = 2$.

Note that if in Example 1.2, one computes $\text{Stab}(C)$ by (1.2), then at n -th step, what we have is the complex $H^0(\mathbb{Z}/p\mathbb{Z}, P(n)_\bullet \otimes M^{\otimes p})$, where $P(n)_\bullet$ is obtained by applying the Dold-Kand equivalence to the p -fold pointwise tensor product $(B^n k)^{\otimes p}$, and shifting it by n . We have $P(n)_0 \cong k, P(n)_l = 0$ for $l > (p-1)n$, and all the $k[\mathbb{Z}/p\mathbb{Z}]$ -modules $P(n)_l, 1 \leq l \leq (p-1)n$ are projective. The only non-trivial homology group of the complex $P(n)_\bullet$ is k in the bottom degree $(p-1)n$. When one takes the colimit with respect to n , one gets the acyclic complex $P'_\bullet = \text{colim}_n P(n)_\bullet$ with $P'_0 \cong k$, and projective $k[\mathbb{Z}/p\mathbb{Z}]$ -modules $P_l, l \geq 1$.

1.3 THH and K-theory. Now fix a commutative ring k , a flat unital associative k -algebra A , and an A -bimodule M . Denote by $P(A)$ the category of finitely generated projective left A -modules, and let $\langle P(A), M \rangle$ be the category of pairs $\langle V, a \rangle, V \in P(A), a : V \rightarrow M \otimes_A V$ an arbitrary map. Then the (split) exact structure on $P(A)$ induces an exact category structure on $\langle P(A), M \rangle$, and one can consider the corresponding K -theory space. Following [DM], we denote it $K(A, M)$ and call it *K-theory of A with coefficients in M* . The fundamental result of [DM] then claims that we have a natural homotopy equivalence

$$(1.5) \quad THH(A, M) \cong \text{Stab}(K)(A, M),$$

where stabilization is taken with respect to the bimodule variable M (strictly speaking, this is not well-defined since the category of A -bimodules is large, but it is obvious from (1.2) that for any fixed M , it suffices to stabilize over the small category formed by finite sums $M^{\oplus n}, n \geq 0$). To obtain individual Topological Hochschild Homology groups $THH_*(A, M)$ one takes the homotopy groups $\pi_*(THH(A, M))$ of the space $THH(A, M)$. In the particular case $A = k, k$ a field, M is simply a k -vector space, and we have an obvious collection of functors

$$(1.6) \quad \begin{aligned} \langle P(k), M \rangle \times \langle P(k), M' \rangle &\rightarrow \langle P(k), M \otimes M' \rangle, \\ \langle V, a \rangle \times \langle V', a' \rangle &\mapsto \langle V \otimes V', a \otimes a' \rangle \end{aligned}$$

that turns $K(k, M)$ into a lax monoidal functor; the induced multiplication on $\text{Stab}(K)(k, k) \cong THH(k)$ is the standard multiplication on $THH(k)$. Note that if M is finite-dimensional, then $\langle P(k), M \rangle$ is the category of finite-dimensional modules over the tensor algebra $T^* M^*$ of the dual vector space

M . If we further assume that k is a finite field, there is the following additional simplification.

Lemma 1.3. *Let k be a finite field. Then the map $K(k, M) \rightarrow K_0(k, M)$ becomes an equivalence after stabilization with respect to M .*

Proof. Let $K_{\geq 1}(k, M)$ be the homotopy fiber of the augmentation map. Since stabilization commutes with finite homotopy limits, it suffices to prove that $K_{\geq 1}(k, -)$ has trivial stabilization $X(-) = \text{Stab}(K_{\geq 1})(k, -)$. Let $p = \text{char } k$, and consider the endomorphism $p(X) : X \rightarrow X$ of (1.3). Then since the category of k -vector spaces is additive and annihilated by p , we have $p(X) = 0$ by Example 1.1, so that X is a retract of the homotopy fiber X_p of the map $p(X) : X \rightarrow X$. Moreover, by their construction, K -theory spaces come from connective spectra, so we can obtain X by stabilizing the corresponding object $Y(-) \in \text{Ho}^{st}(\Gamma_+, k\text{-mod})$ along $k\text{-mod}$, and then X_p is the stabilization of the homotopy fiber Y_p of the map $p(Y) : Y \rightarrow Y$. Up to a shift, $Y_p(M) = K_{\geq 1}(k, M)/p$ is the truncated K -theory of the category $\langle P(k), M \rangle$ with coefficients in $\mathbb{Z}/p\mathbb{Z}$. This is a Noetherian Artinian abelian category, so by devissage, we have

$$(1.7) \quad K_{\geq 1}(k, M)/p \cong \bigoplus_V K_{\geq 1}(k(V))/p,$$

where the sum is over all the isomorphism classes of irreducible objects $V \in \langle p(k), M \rangle$, and $k(V)$ is the endomorphism field of the object V . But $k(V)$ is a finite field of characteristic p , so $K_{\geq 1}(k(V))/p = 0$ by [Q]. \square

Lemma 1.3 shows that for a finite field, computing $THH(k, M)$ by (1.5) reduces to stabilizing $K_0(k, M)$, a functor from k -vector spaces to discrete abelian groups. The values of this functor are rather large — as in (1.7), $K_0(k, M)$ for a finite-dimensional vector space M is the free abelian group generated by all isomorphism classes of irreducible finite-dimensional modules over T^*M^* . Already for $\dim M = 2$ this is a countable but huge set with no discernible structure. However, we can at least construct maps from $K_0(k, M)$ to other functors. What we need is a map

$$(1.8) \quad \varphi : K_0(k, M) \rightarrow C(M), \quad \langle V, a \rangle \mapsto \text{Tr}(a^p),$$

where $C(M)$ is the cyclic power functor of Example 1.2, $a^p : V \rightarrow V \otimes M^{\otimes p}$ is the p -fold iteration of the map a , and its trace $\text{Tr}(a^p) \in M^{\otimes p}$ lies in $C(M)$ by virtue of the cyclic symmetry property of traces ($\text{Tr } ab = \text{Tr } ba$). The map (1.8) is obviously functorial; moreover, if we equip its source with

the lax monoidal structure induced by (1.6), then it is a lax monoidal map with respect to the standard lax monoidal structure on $C(M)$ (checking this amounts to observing that $(a \otimes b)^p = a^p \otimes b^p$). Therefore we can apply the stabilization functor Stab and obtain a multiplicative map

$$\varphi_{\bullet} = \pi_{\bullet}(\text{Stab}(\varphi)) : THH_{\bullet}(k) \rightarrow \text{Stab}(C)(k) \cong k[\varepsilon, u],$$

where the right-hand side is as in Example 1.2. What we will do is construct an element $v \in THH_2(k)$ such that $\varphi_2(v) = u$.

2 The construction.

2.1 The object. Fix a finite field k of characteristic $p = \text{char } k \geq 3$. As in Subsection 1.3, let $P(k)$ be the category of finite-dimensional k -vector spaces, and for any $M \in P(k)$, let $\langle P(k), M \rangle$ be the exact category of pairs $\langle V, a \rangle$, $V \in P(k)$, $a : V \rightarrow V \otimes_k M$. For any pair $\langle V, a \rangle \in \langle P(k), M \rangle$, let $[V, a] \in K_0(k, M)$ be its class. Note that if $M = k$ is one-dimensional, the same devissage as in (1.7) provides an identification

$$(2.1) \quad K_0(k, k) \cong \mathbb{Z}[k] \cong \bigoplus_{\lambda \in k} \mathbb{Z} \cdot [k, \lambda].$$

Consider the endofunctor $\langle P(k), M \rangle \rightarrow \langle P(k), M \rangle$ sending $\langle V, a \rangle$ to $\langle V, -a \rangle$, and let $\iota : K_0(k, M) \rightarrow K_0(k, M)$ be the corresponding involution. Denote by $L(k, M) = K_0(k, M)/K_0(k, M)^{\iota}$ the quotient of the K_0 -group $K_0(k, M)$ by its subgroup of ι -invariants. Note that since p is odd, $\text{tr}(-a)^{\otimes p} = -\text{tr } a^{\otimes p}$, so that the map (1.8) factors as

$$(2.2) \quad K_0(k, M) \xrightarrow{q} L(k, M) \xrightarrow{\psi} C(M),$$

where q is the quotient map. Since as we saw in the proof of Lemma 1.3, the stabilization $\text{Stab}(K_0)_{\bullet}(k, M) \cong THH_{\bullet}(k, M)$ is p -local, and p is odd, we have a decomposition

$$(2.3) \quad THH_{\bullet}(k, k) \cong THH_{\bullet}(k, k)^{\iota} \oplus \text{Stab}(L)_{\bullet}(k, k),$$

and to find an element $v \in THH_2(k, k)$ with $\varphi(v) \neq 0$, it suffices to find an element $v \in \text{Stab}(L)_2(k, k)$ with $\psi(v) \neq 0$ (in fact, it is easy to see from (1.3) that ι acts on $THH_{\bullet}(k, k)$ by $-\text{id}$, so that $THH_{\bullet}(k, k)^{\iota}$ in (2.3) vanishes, but we will not need it).

In order to find such an element v , we can compute $\text{Stab}(L)_{\bullet}(k, k)$ by the colimit (1.2) in the form described in Example 1.1, and in fact, the desired element v appears already at the first step of the colimit.

To construct it, note that $B(k)([n]) \cong k^{\oplus n}$, $[n] \in \Delta$, and we have the face maps $\partial_n^i : B(k)([n]) \rightarrow B(k)([n-1])$, $0 \leq i \leq n$. To simplify notation, let $L([n]) = L(k, B(k)([n]))$, with the face maps $\partial_n^i : L([n]) \rightarrow L([n-1])$. Consider the projective line $\mathbb{P}^1(k)$ over k , with the line bundle $\mathcal{O}(2)$, and identify $H^0(\mathbb{P}^1(k), \mathcal{O}(2)) \cong \mathfrak{sl}_2(k)$, the Lie algebra of vector fields on $\mathbb{P}^1(k)$. For any point $p \in \mathbb{P}^1(k)$, evaluation at p gives a map $\text{ev}_p : \mathfrak{sl}_2(k) \rightarrow k$ whose kernel $\mathfrak{b}_p \subset \mathfrak{sl}_2(k)$ is a Borel subalgebra. Identify $\mathfrak{sl}_2(k) \cong \mathfrak{sl}_2(k)^*$ by the Killing form, and note that dually, we have maps $\text{ev}_p^* : \mathfrak{sl}_2(k) \rightarrow \mathfrak{b}_p^*$. Let $V = k^{\oplus 2}$ be the 2-dimensional k -vector space, and let

$$(2.4) \quad \alpha : \mathfrak{sl}_2(k) \otimes_k V \rightarrow V$$

be the standard $\mathfrak{sl}_2(k)$ -action on V .

Lemma 2.1. *For any element $\lambda \in k$, $\lambda \neq 0, 1$, there exists a unique isomorphism $B(k)([2]) \cong \mathfrak{sl}_2(k)$ such that $\partial_2^i : B(k)([2]) \rightarrow B(k)([1]) \cong k^{\oplus 2}$ for $i = 0, 1, 2, 3$ factors through the evaluation $\text{ev}_p^* : \mathfrak{sl}_2(k) \rightarrow \mathfrak{b}_p^*$ at $p = 0, 1, \infty, \lambda$. Moreover, if we let $a_\lambda : V \rightarrow V \otimes_k B(k)([2]) \cong V \otimes_k \mathfrak{sl}_2(k)$ be dual to (2.4) under this unique isomorphism, and let $v_\lambda = q([V, a_\lambda]) \in L([2])$ be the image of the class $[V, a] \in K_0(k, B(k)([2]))$ under the quotient map (2.2), then we have $\partial_2^i(v_\lambda) = 0$ for any $i = 0, 1, 2, 3$.*

Proof. The first claim is obvious: quadruples of one-dimensional subspaces $k \subset k^{\oplus 3}$ in generic position form a single $GL_3(k)$ -orbit, and quadruples $\{\text{Ker } \partial_n^i\}$ and $\{\text{Ker } \text{ev}_p^*\}$ are both in generic position (for any $\lambda \neq 0, 1$). For the second claim, note that $\partial_2^i([V, a])$, $i = 0, 1, 2, 3$ is then the class $[V, b_p]$, $p = 0, 1, \infty, \lambda$, where $b_p : V \rightarrow V \otimes_k B(k)([1]) \cong V \otimes_k \mathfrak{b}_p^*$ is dual to the action $\mathfrak{b}_p \otimes V \rightarrow V$ induced by (2.4). Since $\mathfrak{b}_p \subset \mathfrak{sl}_2(k)$ is a Borel subalgebra, V as a representation of \mathfrak{b}_p is an extension of 1-dimensional representations that both come by restriction via the quotient map $u : \mathfrak{b}_p \rightarrow k \cong \mathfrak{b}_p/\mathfrak{u}_p$, where $\mathfrak{u}_p \subset \mathfrak{b}_p$ is the nilpotent radical. By devissage, we then have

$$[V, b_p] = u^*[k, \zeta_0] + u^*[k, \zeta_1]$$

for some elements $\zeta_0, \zeta_1 \in k$, where $[k, -]$ is as in (2.1). Since elements in $\mathfrak{b}_p \subset \mathfrak{sl}_2(k) \subset \text{End}(V)$ are traceless, we moreover have $\zeta_1 = -\zeta_0$, and then $[V, b_p]$ is invariant under ι , so that $\partial_n^i(v) = q([V, b_p]) = 0$. \square

2.2 The trace computation. Let $\mathfrak{g} = \mathfrak{sl}_2(k)$ with the standard basis given by the matrices

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix},$$

and let x, y, z denote the basis in \mathfrak{g}^* dual to e, h, f . We would like to explicitly write down the element $t = \varphi([V, a])$ (recall that φ was defined in (1.8)). The space $C(\mathfrak{g}^*)$ has a basis indexed by necklaces of length p over the alphabet x, y, z . We will use noncommutative monomials in order to represent the corresponding basis vector. In order to write t down in the latter basis, one needs to compute the traces of all p -element products of e, h , and f . Consider such a product. We may assume that e or f occurs in it at least once (p is odd, so $\text{tr } h^p = \text{tr } h = 0$). Since $eh = e$ and $fh = -f$, dropping all the occurrences of h will only change the sign. Thus, we may assume that the factors are equal to either e or f . Since $e^2 = f^2 = 0$, both e and f are traceless, and

$$ef = e_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix},$$

we conclude that the trace of the original product is zero unless it was of the form

$$h \dots heh \dots hfh \dots heh \dots hfh \dots h$$

(in other words, in the corresponding necklace e appears at least once, every e is followed by a number of h 's and an f , and every f is followed by a number of h 's and an e), and the trace of the latter equals 1 up to a sign.

We conclude that

$$(2.5) \quad t = \sum \pm xy \dots yzy \dots yxy \dots yzy \dots y,$$

where the explicit signs will be computed later.

2.3 The cohomology class. In order to do the necessary computations, we will use the explicit form of stabilization via the Dold-Kan equivalence; namely, its first step. Let X_\bullet denote the simplicial group obtained from the complex

$$k[-1] = \dots \rightarrow 0 \rightarrow k \rightarrow 0.$$

Denote the standard bases in X_2, X_3 , and X_4 by $X_2 = \langle w_1, w_2 \rangle$, $X_3 = \langle v_1, v_2, v_3 \rangle$, and $X_4 = \langle u_1, u_2, u_3, u_4 \rangle$ respectively. The face maps $d_i : X_3 \rightarrow X_2$ are given by the matrices

$$d_0 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad d_1 = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad d_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \quad d_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix},$$

while the face maps $d_j : X_4 \rightarrow X_3$ are given by the matrices

$$d_0 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad d_1 = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad d_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$d_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \quad d_4 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

We would like to find an isomorphism $\tau : \mathfrak{g}^* \xrightarrow{\sim} X_3$ such that the face maps d_0, d_1, d_2, d_3 become dual to inclusions of Borel subalgebras (see Lemma 2.1). In other words, the inverse images of $\text{Ker } d_i$, $i = 0, \dots, 3$, should lie on the quadric $4ef + h^2$. A simple computation shows that

$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} \lambda(\lambda - 1) & 0 & 0 \\ \lambda^2 & 2\lambda & -\lambda \\ 0 & 0 & -(\lambda - 1) \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

is a family of such morphisms indexed by $\lambda \in \mathbb{A}^1(k) \setminus \{0, 1\}$. Its inverse (up to a scalar) is given by

$$(2.6) \quad \tau_\lambda(x) = -2v_1, \quad \tau_\lambda(y) = v_2 + \lambda(v_1 - v_2 + v_3), \quad \tau_\lambda(z) = 2\lambda v_3.$$

Extend τ_λ to $\lambda \in \{0, 1\}$ by the same formulas, and consider the family of traces $t_\lambda = C(\tau_\lambda)(t)$.

Proposition 2.2. *For some $\lambda \in \mathbb{A}^1(k) \setminus \{0, 1\}$ the element t_λ represents a non-trivial homology class in the normalized complex associated to the simplicial group $C(X_\bullet)$.*

The proof of this proposition will take the rest of the section.

Recall that by Lemma 2.1 t_λ lies in the normalized complex. Thus, it is enough to show that t_λ is not in the image of the map

$$d_0 : \bigcap_{i=1, \dots, 4} \text{Ker } d_i \rightarrow C(X_3).$$

Lemma 2.3. *One has $t_0 = t_1 = 0$.*

Proof. Recall that t is a linear combination of cyclic monomials of the form $xy \dots yzy \dots yxy \dots yzy \dots y$, and each of these monomials contains at least one z (see (2.5) and the discussion preceding it). Since $\tau_0(z) = 0$, we immediately conclude that $t_0 = 0$.

Let us now show that $t_1 = C(\tau_1)(t) = 0$. Recall that $\tau_1(x) = -2v_1$, $\tau_1(y) = v_1 + v_3$, $\tau_1(z) = 2v_3$. It is easy to see that it is enough to prove the following. Let $q > 1$ be an odd integer. Consider a length q binary word of the form $w = 0^{a_1}1^{b_1}0^{a_2}1^{b_2} \dots 0^{a_k}1^{b_k}$ such that $a_i, b_i > 0$ for $i = 1, \dots, k$. Let P denote the set of sequences $S_w = (s_1, s_2, \dots, s_{2l})$, $1 \leq s_1 < s_2 < \dots < s_{2l} \leq q$, $l > 0$, such that $w_{s_i} \neq w_{s_{i+1}}$ for all $i = 1, \dots, 2l - 1$. Given a pair (w, s) , $s \in S_w$, let $p(w, s) = g_1 g_2 \dots g_q$, where

$$g_i = \begin{cases} h, & \text{if } i \text{ does not appear in } s, \\ e, & \text{if } i \text{ appears in } s \text{ and } w_i = 0, \\ f, & \text{if } i \text{ appears in } s \text{ and } w_i = 1, \end{cases}$$

and let $t(w, s) = \text{tr } p(w, s)$. We claim that $\sum_{s \in S_w} t(w, s) = 0$.

We prove the latter by induction on l . First, let $l = 1$. Without loss of generality assume that b_1 is even. Then

$$S_w = \{(s_1, s_2) \mid 1 \leq s_1 \leq a_1, a_1 + 1 \leq s_2 \leq q\}.$$

The involution ι , which sends (s_1, s_2) to $(s_1, s_2 + 1)$ when $s_2 - a_1$ is odd, and to $s_2 - 1$ when $s_2 - a_1$ is even, acts freely on S_w . Moreover, $t(w, s) = -t(w, \iota s)$ since $p(w, s)$ and $p(w, \iota s)$ differ by exchanging adjacent h and f in the product.

Now assume that $l > 1$. Without loss of generality assume that b_1 is even. Then $S_w = S'_w \sqcup S''_w$, where S''_w denotes the set of those sequences containing an element from $\{a_1 + 1, a_1 + 2, \dots, a_1 + b_1\}$. Remark that such an element must be unique, and an argument similar to the one used for the base case shows that $\sum_{s \in S''_w} t(w, s) = 0$. It remains to observe that S'_w is naturally in bijection with $S_{w'}$, where $w' = 0^{a_1+a_2}1^{b_2}0^{a_3}1^{b_3} \dots 0^{a_k}1^{b_k}$, and traces are preserved under this bijection. The claim now follows from the inductive hypothesis. \square

Let us now look at the formal derivative $t'_\lambda(0)$. Since, $\tau_\lambda(z) = 2\lambda v_3$ and every monomial appearing in t contains at least one z , we conclude that the monomials contributing to $t'_\lambda(0)$ are of the form $xy^a zy^b$, where $a, b \geq 0$, $a + b = p - 2$. Remark that the coefficient in t in front of $xy^a zy^a$ equals $\text{tr}(eh^a fh^b) = (-1)^b$. Since $\tau_\lambda(z)$ is linear homogeneous in λ , $\tau_\lambda(y) = v_2 + \lambda(v_1 - v_2 + v_3)$, and $\tau_\lambda(x) = -2v_1$, we conclude that

$$t'_\lambda(0) = -4 \sum_{a=0}^{p-2} (-1)^a v_1 v_2^{p-2-a} v_3 v_2^a.$$

Before we prove the following lemma, let us introduce some convenient notation. Let N_k denote the set of length p necklaces in the alphabet $\{1, 2, \dots, a\}$. Given an element $w = w_1 w_2 \dots w_p \in N_4$, denote by u_w the basis vector in $C(X_4)$ corresponding to $u_{w_1} u_{w_2} \dots u_{w_p}$. Given a necklace $w \in N_3$, the basis vector v_w in $C(X_3)$ is defined similarly. We abuse the notation and define $d_2 : \{1, 2, 3, 4\} \rightarrow \{1, 2, 3\}$ by the rules $d_2(1) = 1$, $d_2(2) = d_2(3) = 2$, $d_2(4) = 3$. Then for any $w \in N_4$ one has $d_2(u_w) = v_{d_2(w)}$. The maps d_1 and d_3 are defined similarly.

We will need the following subsets of N_k . Let $N_k^{1,1}$ denote the set of those necklaces which contain exactly one bead equal to 1 and exactly one bead equal to k . Let $N_k^{+,1}$ denote the set of those necklaces which contain more than one bead equal to 1 and exactly one bead equal to k . The set $N_k^{1,+}$ is defined similarly. Finally, the set $N_k^{+,+}$ consists of those necklaces which contain more than one bead equal to 1 and more than one bead equal to k . One has the following identities:

$$(2.7) \quad \begin{aligned} d_2(N_4^{?,?}) \subseteq N_3^{?,?}, \quad d_1(N_4^{1,1}) \subseteq N_3^{1,1} \sqcup N_3^{+,1}, \quad d_1(N_4^{+,1}) \subseteq N_3^{+,1}, \\ d_3(N_4^{1,1}) \subseteq N_3^{1,1} \sqcup N_3^{1,+}, \quad d_3(N_4^{1,+}) \subseteq N_3^{1,+}. \end{aligned}$$

Finally, denote by $\bar{N}_k = N_k^{1,1} \sqcup N_k^{1,+} \sqcup N_k^{+,1} \sqcup N_k^{+,+}$ (these are all the necklaces containing at least one bead equal to 1 and at least one bead equal to k).

Lemma 2.4. *The element $t'(0)$ is not in the image of the map*

$$d_0 : \bigcap_{i=1, \dots, 4} \text{Ker } d_i \rightarrow C(X_3).$$

Proof. For convenience, we will work with $v = -t'_\lambda(0)/4$ instead. Precisely,

$$v = v_1 v_2^{p-2} v_3 - v_1 v_2^{p-3} v_3 v_2 + \dots + v_1 v_2 v_3 v_2^{p-3} - v_1 v_3 v_2^{p-2}.$$

Consider the element $u' \in C(X_4)$ given by

$$\begin{aligned} u' = & u_2 u_3^{p-2} u_4 - u_2 u_3^{p-3} u_4 u_3 + \dots + u_2 u_3 u_4 u_3^{p-3} - u_2 u_4 u_3^{p-2} - \\ & - u_1 u_3^{p-2} u_4 + u_1 u_3^{p-3} u_4 u_3 - \dots - u_1 u_3 u_4 u_3^{p-3} + u_1 u_4 u_3^{p-2}. \end{aligned}$$

It satisfies the equations $d_0(u') = v$, $d_1(u') = 0$, $d_2(u') = -v$, $d_3(u') = 0$, $d_4(u') = 0$. Thus, it is enough to show that there is no solution to the system of equations

$$(2.8) \quad d_0(u) = d_1(u) = d_3(u) = d_4(u) = 0, \quad d_2(u) = v.$$

Consider the subspace $\text{Ker } d_0$. Since $d_0(u_0) = 0$ and $d_0(u_i) = v_{i-1}$ for $i = 1, 2, 3$, we see that $\text{Ker } d_0$ is spanned by the basis vectors u_w for those $w \in \bar{N}_4$ which contain at least one bead equal to 1. Similarly, $\text{Ker } d_4$ is spanned by the basis vectors corresponding to necklaces containing at least one bead equal to 4. We conclude that $\bar{U} = \text{Ker } d_0 \cap \text{Ker } d_4$ is spanned by the basis vectors u_w , where w runs through the set \bar{N}_4 .

Recall that $U = \langle u_w \mid u_w \in \bar{N}_4 \rangle$, and put $V = \langle v_w \mid v_w \in \bar{N}_3 \rangle$. It follows from the identities (2.7) that $d_i(U) \subseteq V$ for $i = 1, 2, 3$. Since $v \in V$, it is enough to show that there exists no $u \in U$ such that $d_2(u) = v, d_1(u) = d_3(u) = 0$. In order to do that, we are going to construct an element $\nu \in V^*$ such that $\nu(v) \neq 0$, while $d_2^*(\nu)$ equals zero when restricted to $\text{Ker } d_2 \cap \text{Ker } d_3$.

Let $\langle u_w^\vee \mid w \in \bar{N}_4 \rangle$ and $\langle v_w^\vee \mid w \in \bar{N}_3 \rangle$ denote the bases in U^* and V^* dual to $\langle u_w \mid w \in \bar{N}_4 \rangle$ and $\langle v_w \mid w \in \bar{N}_3 \rangle$ respectively. Consider the map $p : X_4 \rightarrow [1]$ given by $p(1) = p(2) = 0, p(3) = p(4) = 1$. Let $w = w_1 w_2 \dots w_p$ be a word in the alphabet $\{1, 2, 3, 4\}$ with at least one occurrence of 1 and at least one occurrence of 4. Let $0 \leq i < p$ be such that $\sigma^i p(w)$ is lexicographically minimal, where σ denotes the cyclic permutation. If $1 \leq i_1 < i_2 < \dots < i_a \leq p$ are the indices of the letters w_i in w equal to 1, put

$$\gamma_1(w) = \frac{i_1 + i_2 + \dots + i_a}{a} - i \in k.$$

Similarly, if $1 \leq j_1 < j_2 < \dots < j_b \leq p$ are the indices of the letters w_j in w equal to 4, put

$$\gamma_4(w) = \frac{j_1 + j_2 + \dots + j_b}{b} - i \in k.$$

Finally, put $\Delta(w) = \gamma_1(w) - \gamma_4(w)$. Remark that both s_1 and s_4 are invariant under the cyclic action. Thus, s_1, s_4 , and Δ descend to maps from \bar{N}_4 to k .

Consider the element $\mu \in U^*$ given by

$$\mu = \sum_{w \in \bar{N}_4} \Delta(w) u_w^\vee.$$

We claim that $\mu(\text{Ker } d_1 \cap \text{Ker } d_3) = 0$. Indeed, pick sections $s_i : \bar{N}_3 \rightarrow \bar{N}_4$ of the maps $d_i, i = 1, 2, 3$. Then

$$\mu = \sum_{w \in \bar{N}_3} (\gamma_1(s_1(w)) d_1^* - \gamma_4(s_3(w)) d_3^*) v_w^\vee,$$

and the left hand side trivially vanishes on $\text{Ker } d_1 \cap \text{Ker } d_3$. Meanwhile, since $\delta(w)$ depends only on the positions the beads equal to 1 and 4, we see that

$\mu = d_2^* \nu$, where

$$\nu = \sum_{w \in \bar{N}_3} \Delta(s_2(w)) v_w^\vee.$$

Finally, let us compute $\nu(v)$. For $w \in N_4^{1,1}$ the number δ measures the clockwise distance from the bead equal to 1 to the bead equal to 4. Thus

$$\nu(v) = 1 - 2 + 3 - 4 + \dots + (p - 2) - (p - 1) = \frac{1 - p}{2} \neq 0.$$

□

The proof of Proposition 2.2 now trivially follows from Lemmas 2.3 and 2.4 and the fact that the parametric curve t_λ is of degree at most $p - 1$.

References

- [BFS] C.P. Bendel, E.M. Friedlander, and A. Suslin, *Infinitesimal 1-parameter subgroups and cohomology*, JAMS **10** (1997), 693–728.
- [BMS] B. Bhatt, M. Morrow, and P. Scholze, *Topological Hochschild homology and integral p-adic Hodge theory*, Publ. Math. IHES, **129** (2019), 199–310.
- [Bö] M. Bökstedt, *Topological Hochschild homology*, preprint, Bielefeld, 1985.
- [Br] L. Breen, *Extensions du groupe additif*, Publ. Math. IHES, **48** (1978), 39–125.
- [DM] B. Dundas and R. McCarthy, *Stable K-Theory and Topological Hochschild Homology*, Annals of Math. **140** (1994), 685–701.
- [DHKS] W. Dwyer, Hirschhorn, D. Kan, and J. Smith, *Homotopy Limit Functors on Model Categories and Homotopical Categories*, Mathematical Surveys and Monographs, **113**, AMS, Providence, RI, 2004.
- [FPSVW] Z. Fiedorowicz, T. Pirashvili, R. Schwänzl, R. Vogt, and F. Waldhausen, *Mac Lane homology and topological Hochschild homology*, Math. Ann. **303** (1995), 149–164.
- [FLS] V. Franjou, J. Lannes, and L. Schwartz, *Autour de la cohomologie de Mac Lane des corps finis*, Invent. Math. **115** (1994), 513–538.
- [FS] E. Friedlander and A. Suslin, *Cohomology of finite group schemes over a field*, Invent. Math. **127** (1997), 209–270.
- [HN] L. Hesselholt and Th. Nikolaus, *Topological cyclic homology*, [arXiv:1905.08984](https://arxiv.org/abs/1905.08984).
- [JP] M. Jibladze and T. Pirashvili, *Cohomology of algebraic theories*, J. of Algebra, **137** (1991), 253–296

- [Ka] D. Kaledin, *Trace theories, Bökstedt periodicity and Bott periodicity*, [arXiv:2004.04279](#).
- [KN] A. Krause and T. Nikolaus, *Bökstedt periodicity and quotients of DVRs*, [arXiv:1907.03477](#).
- [L] J.-L. Loday, *Cyclic Homology*, second ed., Springer, 1998.
- [M] A. Mathew, *Kaledin’s degeneration theorem and topological Hochschild homology*, [arXiv:1710.09045](#).
- [NS] T. Nikolaus and P. Scholze, *On topological cyclic homology*, *Acta Math.* **221** (2018), 203–409.
- [P1] T. Pirashvili, *New homology and cohomology of rings*, *Bull. Acad. Sci. Georgian SSR* **133** (1989), 477–480.
- [P2] T. Pirashvili, Chapter 13 in [L].
- [PW] T. Pirashvili and F. Waldhausen, *Mac Lane homology and topological Hochschild homology*, *J. Pure Appl. Alg.* **82** (1992), 81–99.
- [Q] D. Quillen, *On the Cohomology and K-Theory of the General Linear Groups Over a Finite Field*, *Ann. of Math.*, **96** (1972), 552–586.
- [S] G. Segal, *Categories and cohomology theories*, *Topology* **13** (1974), 293–312.
- [T] A. Touzé, *Universal classes for algebraic groups*, *Duke Math. J.* **151**, (2010), 219–249.

Affiliations (for both authors):

1. Steklov Mathematics Institute (main affiliation).
2. National Research University Higher School of Economics.

E-mail addresses: kaledin@mi-ras.ru, avfonarev@mi-ras.ru.