

VIRTUAL AND UNIVERSAL BRAID GROUPS, THEIR QUOTIENTS AND REPRESENTATIONS

V. BARDAKOV, I. EMEL'YANENKOV, M. IVANOV, T. KOZLOVSKAYA,
T. NASYBULLOV, A. VESNIN

ABSTRACT. In the present paper we study structural aspects of certain quotients of braid groups and virtual braid groups. In particular, we construct and study linear representations $B_n \rightarrow \mathrm{GL}_{n(n-1)/2}(\mathbb{Z}[t^{\pm 1}])$, $VB_n \rightarrow \mathrm{GL}_{n(n-1)/2}(\mathbb{Z}[t^{\pm 1}, t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}])$ which are connected with the famous Lawrence-Bigelow-Krammer representation. It turns out that these representations are faithful representations of crystallographic groups B_n/P'_n , VB_n/VP'_n , respectively. Using these representations we study certain properties of the groups B_n/P'_n , VB_n/VP'_n . Moreover, we construct new representations and decompositions of universal braid groups UB_n .

Keywords: virtual braid, universal braid, crystallographic group, representations by automorphisms, linear group.

Mathematics Subject Classification 2010: 20F36, 20F29, 57M27, 57M25.

CONTENTS

Introduction	2
1. Preliminaries	4
1.1. Braid groups and conjugating automorphism groups	4
1.2. The singular braid monoid	7
1.3. The virtual braid group and the welded braid group	7
1.4. Braid groups and crystallographic groups	9
2. Linear representation of B_n/P'_n	10
3. Linear representation of VB_n/VP'_n	13
4. Decomposition and linearity of VB_3	15
5. Universal braid group	18
5.1. Representations by automorphisms of a free group	19
5.2. Linear representations	24
5.3. Universal pure braid group	25
Acknowledgements	27
References	27

INTRODUCTION

Braid groups were introduced by E. Artin as a tool for working with classical knots and links. The famous Alexander theorem says that every classical link is equivalent to the closure of some braid, and the well-known Markov theorem describes braids which have equivalent closures.

In 1999 L. Kauffman introduced the virtual knot theory which generalizes the classical knot theory [33]. Virtual braid groups were introduced in the same paper as a tool for working with virtual knots and links. The analogues of the Alexander and the Markov theorems for virtual braids and links were formulated and proved by S. Kamada in [31]. Another form of the Markov theorem for virtual braids and links was introduced by Kauffman and Lambropoulou in [35].

Welded braid groups, unrestricted virtual braid groups, flat virtual braid groups, flat welded braid groups, Gauss virtual braid groups are quotients of virtual braid groups. These groups have various applications in algebra and topology, in particular, each of these groups corresponds to some knot theory, which has a geometric interpretation that is often very useful when studying applications. Welded and unrestricted virtual braid groups are used for working with welded links [25] and fused links [2, 41], respectively. Flat virtual braid groups were introduced in [34] for working with flat virtual knots and links, which were introduced in [33], and which model curves on surfaces. The notion of flat virtual links is the same to the notion of virtual strings introduced by V. Turaev in [48]. In the papers [30, 36] invariants for flat virtual knots and links were used for constructing invariants for virtual knots and links. Gauss virtual braid groups are used for studying Gauss virtual links, which are called “homotopy classes of Gauss words” by Turaev in [49], and called “free knots” by Manturov in [38]. All together classical braid groups, virtual braid groups, welded braid groups, unrestricted virtual braid groups, flat virtual braid groups, flat welded braid groups and Gauss virtual braid groups are called braid-like groups. Universal braid groups were introduced in [4]. The name “universal” in this paper is motivated by the fact that universal braid groups contain classical braid groups, and moreover these groups have singular braid groups and all braid-like groups as quotients. Hence the structure of these groups play an important role in the study of all braid-like groups.

One way of studying braid-like groups is to study representations of these groups by automorphisms of different algebraic systems. Using the representations one can prove that various algorithmic problems (the word problem, the conjugacy problem etc) can be solved in these groups. Also using representations of braid-like groups one can construct invariants for the corresponding knot theories [12, 13, 14].

One of the first representations of braid-like groups is the Artin’s representation $\varphi : B_n \rightarrow \text{Aut}(F_n)$ of the braid group B_n by automorphisms of the free group F_n with n generators. The Artin’s representation is

faithful. During the years a lot of representations of braid-like groups were investigated. For example, there are the Silver-Williams representation $\varphi_{SW} : VB_n \rightarrow \text{Aut}(F_n * \mathbb{Z}^{n+1})$ (see [46]), the Boden-Dies-Gaudreau-Gerlings-Harper-Nicas representation $\varphi_{BD} : VB_n \rightarrow \text{Aut}(F_n * \mathbb{Z}^2)$ (see [21]), the Kamada representation $\varphi_K : VB_n \rightarrow \text{Aut}(F_n * \mathbb{Z}^{2n})$ (see [15]), the representations $\varphi_M : VB_n \rightarrow \text{Aut}(F_n * \mathbb{Z}^{2n+1})$, $\tilde{\varphi}_M : VB_n \rightarrow \text{Aut}(F_n * \mathbb{Z}^n)$ of Bardakov-Mikhailchishina-Neshchadim (see [10, 11]). In the paper [8] there are representations of the flat virtual braid groups by automorphisms of the free group, which do not preserve forbidden relations.

The study of linear representations of braid groups was started when Burau constructed his famous representation $\varphi_B : B_n \rightarrow \text{GL}_n(\mathbb{Z}[t, t^{-1}])$ (see, for example, [20, Section 3]). For $n = 2, 3$ the Burau representation is known to be faithful, for $n \geq 5$ the Burau representation is known to have non-trivial kernel [18], for $n = 4$ the question on faithfulness of the Burau representation is still open (see some positive results, for example, in [17]). The Burau representation was generalized in 1961 by Gassner [27] to a related representation of the pure braid group $\varphi_G : P_n \rightarrow \text{GL}_n(\mathbb{Z}[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_n^{\pm 1}])$. It is not known whether the Gassner representation is faithful for $n > 3$. Some new results about representations of virtual braid groups can be found, for example, in [12, 14]. Certain linear representations of flat virtual braid groups and Gauss virtual braid groups are studied in [8].

Another way to study braid-like groups is to explore normal forms of elements in braid-like groups which usually (but not always) arise when a braid-like group can be written as a product (for example, direct product, semidirect product, free product etc). There are several normal forms of elements in classical braid groups which can be found, for example, in [32]. Attempts to construct a normal form for elements of the virtual braid groups were made in the paper [4]. Paper [7] contains results about the normal form of elements in unrestricted virtual braid groups and flat welded braid groups.

In the present paper we study structural aspects of certain quotients of braid groups and virtual braid groups using both representations and decompositions. In particular, we construct and study linear representations

$$B_n \rightarrow \text{GL}_{n(n-1)/2}(\mathbb{Z}[t^{\pm 1}]), \quad VB_n \rightarrow \text{GL}_{n(n-1)/2}(\mathbb{Z}[t^{\pm 1}, t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}])$$

which are connected with the famous Lawrence-Bigelow-Krammer representation. It turns out that these representations are faithful representations of crystallographic groups B_n/P'_n , VB_n/VP'_n , respectively. Using these representations we study certain properties of the groups B_n/P'_n , VB_n/VP'_n . Moreover, we construct new representations and decompositions of universal braid groups.

The paper is organized as follows. In Section 1 we recall definitions and basic properties of braid groups, virtual braid groups, welded braid groups and pure subgroups of these groups. In Section 2 we construct a faithful linear representation of the crystallographic group B_n/P'_n that is the quotient of the braid group B_n by the commutator subgroup $P'_n = [P_n, P_n]$ of

the pure braid group P_n (Theorem 2.1). The torsion in this group is studied in [28]. Using the constructed representation we describe all elements of order three in B_3/P'_3 (Proposition 2.1). Note that this group has only three-torsion. In Section 3 we construct a faithful linear representation of another crystallographic group VB_n/VP'_n which contains B_n/P'_n (Theorem 3.1). In Proposition 3.2 we demonstrate that VB_3/VP'_3 contains infinitely many elements of order two. Note that B_n/P'_n has no two-torsion. In Section 4 we show that VP_3 can be decomposed into the free product of infinite cyclic group and group that is a semi-direct product of free group of rank three and free group of rank two (Proposition 4.1). In Theorem 4.1 we prove that VB_3 is linear. In Section 5 we discuss the universal braid group UB_n introduced in [4], and its pure subgroup UP_n . In particular we construct several representations of UP_n by automorphisms of free groups which act faithfully on a subgroup T_n of UP_n . Also in Theorem 5.2 we demonstrate that the groups UP_2 and UP_3 admit decompositions into free products.

1. PRELIMINARIES

In this section we give necessary preliminaries. We use classical notations. If G is a group, and $a, b \in G$, then we denote by $a^b = b^{-1}ab$ the conjugate of a by b , and by $[a, b] = a^{-1}b^{-1}ab$ the commutator of the elements a, b . If X is a set, then we denote by $\text{Sym}(X)$ the set of all bijections from X to X . All actions are supposed to be right, i. e. if $f, g \in \text{Sym}(X)$, then for $x \in X$ we denote $fg(x) = g(f(x))$.

1.1. Braid groups and conjugating automorphism groups. For $n \geq 2$ the braid group B_n on n strands is defined as the group generated by $(n-1)$ elements $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ with defining relations

$$\begin{aligned} (1) \quad & \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, & i = 1, 2, \dots, n-2, \\ (2) \quad & \sigma_i \sigma_j = \sigma_j \sigma_i, & |i - j| \geq 2. \end{aligned}$$

Relations (1) and (2) are called *braid relations*. There is a homomorphism φ from B_n to the symmetric group S_n on n symbols $\{1, 2, \dots, n\}$, which maps the generator σ_i to the transposition $(i, i+1)$ for $i = 1, 2, \dots, n-1$. The kernel $\text{Ker}(\varphi)$ is called the *pure braid group* on n strands and is denoted by P_n . From the definition it is clear that we have the following short exact sequence:

$$1 \rightarrow P_n \rightarrow B_n \xrightarrow{\varphi} S_n \rightarrow 1.$$

It is known that the pure braid group P_n can be generated by the elements $a_{i,j}$ for $1 \leq i < j \leq n$, where

$$\begin{aligned} a_{i,i+1} &= \sigma_i^2, & i = 1, 2, \dots, n-1, \\ a_{i,j} &= \sigma_{j-1} \sigma_{j-2} \dots \sigma_{i+1} \sigma_i^2 \sigma_{i+1}^{-1} \dots \sigma_{j-2}^{-1} \sigma_{j-1}^{-1}, & i+1 < j \leq n. \end{aligned}$$

Also it is known that the pure braid group P_n can be decomposed into a semi-direct product $P_n = U_n \rtimes P_{n-1}$, where U_n is the free group with free generators $a_{1,n}, a_{2,n}, \dots, a_{n-1,n}$, and P_{n-1} is the pure braid group on $(n-1)$ strands. Here U_n is a normal subgroup of P_n . Similar, P_{n-1} is a semi-direct product of the free group U_{n-1} with the free generators $a_{1,n-1}, a_{2,n-1}, \dots, a_{n-2,n-1}$ and P_{n-2} , and so on. Therefore P_n has the following decomposition into the iterated semi-direct product (see [39]):

$$P_n = U_n \rtimes (U_{n-1} \rtimes (\dots \rtimes (U_3 \rtimes U_2) \dots)),$$

where $U_i \simeq F_{i-1}$ is the free group on $(i-1)$ generators for $i = 2, 3, \dots, n$.

The group B_n has a faithful representation by automorphisms of the free group $F_n = \langle x_1, x_2, \dots, x_n \rangle$, where the generator σ_i , $i = 1, 2, \dots, n-1$, defines the automorphism (which we denote by the same symbol $\sigma_i \in \text{Aut}(F_n)$) which acts on the generators x_1, x_2, \dots, x_n of F_n by the following rule

$$\sigma_i : \begin{cases} x_i \mapsto x_i x_{i+1} x_i^{-1}, \\ x_{i+1} \mapsto x_i, \\ x_l \mapsto x_l, \end{cases} \quad l \neq i, i+1.$$

By the Artin's theorem [20, Theorem 1.9] an automorphism β from $\text{Aut}(F_n)$ belongs to $B_n < \text{Aut}(F_n)$ if and only if β satisfies the following two conditions:

- 1) $\beta(x_i) = a_i^{-1} x_{\pi(i)} a_i$ for $1 \leq i \leq n$,
- 2) $\beta(x_1 x_2 \cdots x_n) = x_1 x_2 \cdots x_n$,

where π is a permutation from S_n and $a_i \in F_n$ for $i = 1, 2, \dots, n$.

The automorphism β of the free group F_n is called a *conjugating automorphism* (or permutation-conjugating by terminology from [25]) if it satisfies condition 1) from the list above. It is clear that every automorphism from $B_n < \text{Aut}(F_n)$ is a conjugating automorphism. The group of all conjugating automorphisms of F_n is denoted by C_n . It is known that this group is generated by the automorphisms σ_i for $i = 1, 2, \dots, n-1$ and automorphisms α_i for $i = 1, 2, \dots, n-1$, where α_i acts on the generators x_1, x_2, \dots, x_n of F_n by the following rule

$$\alpha_i : \begin{cases} x_i \mapsto x_{i+1}, \\ x_{i+1} \mapsto x_i, \\ x_l \mapsto x_l, \end{cases} \quad l \neq i, i+1.$$

Generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ of C_n satisfy the braid relations (1) and (2) of B_n . Also it is easy to see that the automorphisms $\alpha_1, \alpha_2, \dots, \alpha_{n-1}$ of C_n generate the symmetric group S_n in C_n which has the following defining relations

- (3) $\alpha_i \alpha_{i+1} \alpha_i = \alpha_{i+1} \alpha_i \alpha_{i+1}, \quad i = 1, 2, \dots, n-2,$
- (4) $\alpha_i \alpha_j = \alpha_j \alpha_i, \quad |i - j| \geq 2,$
- (5) $\alpha_i^2 = 1, \quad i = 1, 2, \dots, n-1.$

Moreover, the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \alpha_1, \alpha_2, \dots, \alpha_{n-1}$ of C_n satisfy the following mixed relations (see [25, 44]):

$$\begin{aligned} (6) \quad & \alpha_i \sigma_j = \sigma_j \alpha_i, & |i - j| \geq 2, \\ (7) \quad & \sigma_i \alpha_{i+1} \alpha_i = \alpha_{i+1} \alpha_i \sigma_{i+1}, & i = 1, 2, \dots, n-2, \\ (8) \quad & \sigma_{i+1} \sigma_i \alpha_{i+1} = \alpha_i \sigma_{i+1} \sigma_i, & i = 1, 2, \dots, n-2. \end{aligned}$$

Summarizing, the group C_n has generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \alpha_1, \alpha_2, \dots, \alpha_{n-1}$ with the defining relations (1)-(8).

For $1 \leq i \neq j \leq n$ denote by $\varepsilon_{i,j}$ the automorphism of F_n which acts on the generators of F_n by the rule

$$\varepsilon_{i,j} : \begin{cases} x_i \mapsto x_j^{-1} x_i x_j, & i \neq j, \\ x_l \mapsto x_l, & l \neq i. \end{cases}$$

The subgroup of $\text{Aut}(F_n)$ generated by $\varepsilon_{i,j}$ for $1 \leq i \neq j \leq n$ is called the *group of basis-conjugating automorphisms* and is denoted by Cb_n . It is clear that every generator $\varepsilon_{i,j}$ belongs to C_n , therefore the group Cb_n of basis-conjugating automorphisms is a subgroup of the group C_n of conjugating automorphisms. In [40] it is proved that Cb_n has the following defining relations:

$$\begin{aligned} (9) \quad & \varepsilon_{i,j} \varepsilon_{k,l} = \varepsilon_{k,l} \varepsilon_{i,j}, \\ (10) \quad & \varepsilon_{i,j} \varepsilon_{k,j} = \varepsilon_{k,j} \varepsilon_{i,j}, \\ (11) \quad & (\varepsilon_{i,j} \varepsilon_{k,j}) \varepsilon_{i,k} = \varepsilon_{i,k} (\varepsilon_{i,j} \varepsilon_{k,j}), \end{aligned}$$

where different letters denote different indices.

It is easy to see that the group C_n of conjugating automorphisms can be decomposed into a semi-direct product $C_n = Cb_n \rtimes S_n$, where the permutation group S_n is generated by the automorphisms $\alpha_1, \alpha_2, \dots, \alpha_{n-1}$. In [44] it is proved that the following relations hold:

$$\begin{aligned} \varepsilon_{i,i+1} &= \alpha_i \sigma_i^{-1}, \\ \varepsilon_{i+1,i} &= \sigma_i^{-1} \alpha_i, \\ \varepsilon_{i,j} &= \alpha_{j-1} \alpha_{j-2} \dots \alpha_{i+1} \varepsilon_{i,i+1} \alpha_{i+1} \dots \alpha_{j-2} \alpha_{j-1}, & \text{if } i < j, \\ \varepsilon_{j,i} &= \alpha_{j-1} \alpha_{j-2} \dots \alpha_{i+1} \alpha_i \varepsilon_{i,i+1} \alpha_i \alpha_{i+1} \dots \alpha_{j-2} \alpha_{j-1}, & \text{if } i < j. \end{aligned}$$

The structure of Cb_n was studied in [4, 5], where it was shown that Cb_n , $n \geq 2$, has a decomposition into an iterated semi-direct product

$$Cb_n = D_{n-1} \rtimes (D_{n-2} \rtimes (\dots \rtimes (D_2 \rtimes D_1) \dots)),$$

where D_i denotes a subgroup of Cb_n generated by the elements

$$\{\varepsilon_{i+1,1}, \varepsilon_{i+1,2}, \dots, \varepsilon_{i+1,i}, \varepsilon_{1,i+1}, \varepsilon_{2,i+1}, \dots, \varepsilon_{i,i+1}\},$$

where $i = 1, 2, \dots, n-1$. The elements $\{\varepsilon_{i+1,1}, \varepsilon_{i+1,2}, \dots, \varepsilon_{i+1,i}\}$ generate the free group of rank i in D_i , while the elements $\{\varepsilon_{1,i+1}, \varepsilon_{2,i+1}, \dots, \varepsilon_{i,i+1}\}$ generate the free abelian group of rank i in D_i .

The pure braid group P_n is a subgroup of Cb_n . The generators $a_{i,j}$ (for $1 \leq i < j \leq n$) of P_n can be written in Cb_n in the following form

$$a_{i,i+1} = \varepsilon_{i,i+1}^{-1} \varepsilon_{i+1,i}^{-1}$$

for $i = 1, 2, \dots, n-1$, and

$$\begin{aligned} a_{i,j} &= \varepsilon_{j-1,i} \varepsilon_{j-2,i} \cdots \varepsilon_{i+1,i} (\varepsilon_{i,j}^{-1} \varepsilon_{j,i}^{-1}) \varepsilon_{i+1,i}^{-1} \cdots \varepsilon_{j-2,i}^{-1} \varepsilon_{j-1,i}^{-1} \\ &= \varepsilon_{j-1,j}^{-1} \varepsilon_{j-2,j}^{-1} \cdots \varepsilon_{i+1,j}^{-1} (\varepsilon_{i,j}^{-1} \varepsilon_{j,i}^{-1}) \varepsilon_{i+1,j} \cdots \varepsilon_{j-2,j} \varepsilon_{j-1,j}, \end{aligned}$$

for $2 \leq i+1 < j \leq n$.

1.2. The singular braid monoid. *The Baez-Birman monoid* [3, 19] or *the singular braid monoid* SB_n is defined as the monoid with the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \sigma_1^{-1}, \sigma_2^{-1}, \dots, \sigma_{n-1}^{-1}, \tau_1, \tau_2, \dots, \tau_{n-1}$, and defining relations (1), (2) and additional relations

$$\begin{aligned} (12) \quad & \sigma_i \sigma_i^{-1} = \sigma_i^{-1} \sigma_i = 1, & i = 1, 2, \dots, n-1, \\ (13) \quad & \tau_i \tau_j = \tau_j \tau_i, & |i-j| \geq 1, \\ (14) \quad & \tau_i \sigma_j = \sigma_j \tau_i, & |i-j| \geq 1, \\ (15) \quad & \tau_i \sigma_i = \sigma_i \tau_i, & i = 1, 2, \dots, n-1, \\ (16) \quad & \sigma_i \sigma_{i+1} \tau_i = \tau_{i+1} \sigma_i \sigma_{i+1}, & i = 1, 2, \dots, n-2, \\ (17) \quad & \sigma_{i+1} \sigma_i \tau_{i+1} = \tau_i \sigma_{i+1} \sigma_i, & i = 1, 2, \dots, n-2. \end{aligned}$$

In the work [26] it was proved that the singular braid monoid SB_n is embedded into the group SG_n , which is called the singular braid group and has the same defining relations as SB_n .

1.3. The virtual braid group and the welded braid group. The *virtual braid group* VB_n on n strands, introduced in [33], is the group with the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \rho_1, \rho_2, \dots, \rho_{n-1}$, which is defined by the relations (1)-(2) and additional relations

$$\begin{aligned} (18) \quad & \rho_i \rho_{i+1} \rho_i = \rho_{i+1} \rho_i \rho_{i+1}, & i = 1, 2, \dots, n-2, \\ (19) \quad & \rho_i \rho_j = \rho_j \rho_i, & |i-j| \geq 2, \\ (20) \quad & \rho_i^2 = 1, & i = 1, 2, \dots, n-1, \\ (21) \quad & \sigma_i \rho_j = \rho_j \sigma_i, & |i-j| \geq 1, \\ (22) \quad & \rho_i \rho_{i+1} \sigma_i = \sigma_{i+1} \rho_i \rho_{i+1}, & i = 1, 2, \dots, n-2. \end{aligned}$$

The elements $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ generate the braid group B_n with the braid relations (1)-(2) in VB_n . Relations (18)-(20) are defining relations of the permutation group S_n , where the elements $\rho_1, \rho_2, \dots, \rho_{n-1}$ generate the permutation group S_n in VB_n , hence the virtual braid group VB_n is generated by the braid group B_n and the permutation group S_n . Relations (21)-(22) are called the mixed relations of the virtual braid group. Note that relations (22) are equivalent to the relations

$$\rho_{i+1} \rho_i \sigma_{i+1} = \sigma_i \rho_{i+1} \rho_i$$

for $i = 1, 2, \dots, n-1$. It was shown in [29] that the relations

$$(23) \quad \rho_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \rho_{i+1}, \quad \rho_{i+1} \sigma_i \sigma_{i+1} = \rho_i \sigma_{i+1} \sigma_i.$$

donot hold in VB_n . Relations (23) are usually called *forbidden relations*.

The *virtual pure braid group* introduced in [4] is the kernel $VP_n = \text{Ker}(\nu)$ of the homomorphism $\nu : VB_n \rightarrow S_n$ from the virtual braid group VB_n onto the symmetric group S_n given on generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \rho_1, \rho_2, \dots, \rho_{n-1}$ of VB_n by the following rule

$$\nu(\sigma_i) = \nu(\rho_i) = \rho_i, \quad \text{for } i = 1, 2, \dots, n-1.$$

From the definition it is clear that the following short exact sequence holds:

$$1 \rightarrow VP_n \rightarrow VB_n \xrightarrow{\nu} S_n \rightarrow 1.$$

Following [4] for $1 \leq i \neq j \leq n$ let us definte the elements $\lambda_{i,j}$ by the following formulas

$$\begin{aligned} \lambda_{i,i+1} &= \rho_i \sigma_i^{-1}, \\ \lambda_{i+1,i} &= \rho_i \lambda_{i,i+1} \rho_i = \sigma_i^{-1} \rho_i \end{aligned}$$

for $i = 1, 2, \dots, n-1$, and

$$\begin{aligned} \lambda_{i,j} &= \rho_{j-1} \rho_{j-2} \dots \rho_{i+1} \lambda_{i,i+1} \rho_{i+1} \dots \rho_{j-2} \rho_{j-1}, \\ \lambda_{j,i} &= \rho_{j-1} \rho_{j-2} \dots \rho_{i+1} \lambda_{i+1,i} \rho_{i+1} \dots \rho_{j-2} \rho_{j-1} \end{aligned}$$

for $1 \leq i < j-1 \leq n-1$. The following theorem is proved in [4].

Theorem 1.1. *The group VP_n admits a presentation with the generators $\lambda_{k,l}$ for $1 \leq k \neq l \leq n$ and the defining relations*

$$(24) \quad \lambda_{i,j} \lambda_{k,l} = \lambda_{k,l} \lambda_{i,j},$$

$$(25) \quad \lambda_{k,i} \lambda_{k,j} \lambda_{i,j} = \lambda_{i,j} \lambda_{k,j} \lambda_{k,i},$$

where different letters denote different indices.

The *welded braid group* WB_n on n strands introduced in [25] is the group with the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \alpha_1, \alpha_2, \dots, \alpha_{n-1}$, where the elements $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ generate the braid group B_n on n strands (i. e. satisfy relations (1)-(2)), the elements $\alpha_1, \alpha_2, \dots, \alpha_{n-1}$ generate the symmetric group S_n (i. e. satisfy relations (18)-(20) where we change the letter ρ by the letter α), and moreover, the following mixed relations hold:

$$(26) \quad \alpha_i \sigma_j = \sigma_j \alpha_i, \quad |i-j| \geq 2,$$

$$(27) \quad \sigma_i \alpha_{i+1} \alpha_i = \alpha_{i+1} \alpha_i \sigma_{i+1}, \quad i = 1, 2, \dots, n-2,$$

$$(28) \quad \sigma_{i+1} \sigma_i \alpha_{i+1} = \alpha_i \sigma_{i+1} \sigma_i, \quad i = 1, 2, \dots, n-2.$$

It was observed in [25] that WB_n is isomorphic to the group of conjugating automorphisms C_n . It was proved in [25] that the following relations symmetric to (27)

$$\sigma_{i+1} \alpha_i \alpha_{i+1} = \alpha_i \alpha_{i+1} \sigma_i,$$

hold in WB_n for $i = 1, 2, \dots, n - 2$, while the relations

$$\alpha_{i+1} \sigma_i \sigma_{i+1} = \sigma_i \sigma_{i+1} \alpha_i$$

do not hold.

Comparing defining relations (21)–(22) of VB_n with defining relations (26)–(28) of WB_n , one can see that the difference between WB_n and VB_n is given by relations (28). Therefore, there exists a homomorphism

$$\varphi_{VW} : VB_n \longrightarrow WB_n,$$

which maps σ_i to σ_i , and ρ_i to α_i for $i = 1, 2, \dots, n - 1$. Hence, WB_n is the homomorphic image of VB_n .

Recall that a group G is said to be *linear* if there exists a faithful representation of G into the general linear group $GL_m(k)$ for some integer $m \geq 2$ and a field k . It was shown in [18, 37] that the braid group B_n is linear for every $n \geq 2$. In [50] there were constructed linear representations of the virtual braid groups VB_n and the welded braid groups WB_n by matrices from $GL_n(\mathbb{Z}[t, t^{-1}])$. These representations extend the well-known Burau representation. In [5] there was constructed a linear representation of the group $C_n \simeq WB_n$ which is an extension (with additional conditions on parameters) of the well-known Lawrence-Bigelow-Krammer representation.

1.4. Braid groups and crystallographic groups. An interesting connection between braid groups and crystallographic groups was observed in [28]. Recall that a discrete and uniform subgroup G of $\mathbb{R}^n \rtimes O_n(\mathbb{R}) \subseteq \text{Aff}(\mathbb{R}^n)$ is said to be a *crystallographic group* of dimension n . If in addition G is torsion free, then G is called a *Bieberbach group* of dimension n . The correspondence between Bieberbach groups and fundamental groups of closed flat Riemannian manifold is presented in [51].

The following characterization of crystallographic groups was used in [28, Lemma 8] in order to demonstrate the relation between braid groups and crystallographic groups.

Lemma 1.1. *Let G be a group. Then G is a crystallographic group if and only if there exist an integer $n \in \mathbb{N}$ and a short exact sequence*

$$0 \rightarrow \mathbb{Z}^n \rightarrow G \rightarrow \Phi \rightarrow 1$$

such that the following conditions hold:

- (a) Φ is a finite group,
- (b) the integral representation $\theta : \Phi \rightarrow \text{Aut}(\mathbb{Z}^n)$, induced by conjugation on \mathbb{Z}^n and defined by $\theta(\varphi)(x) = \pi x \pi^{-1}$, where $x \in \mathbb{Z}^n$, $\varphi \in \Phi$ and $\pi \in G$ is such that $\zeta(\pi) = \varphi$, is faithful.

If G is a crystallographic group, then the integer n from Lemma 1.1 is called the dimension of G , the finite group Φ is called the holonomy group of G , and the integral representation $\theta : \Phi \rightarrow \text{Aut}(\mathbb{Z}^n)$ is called the holonomy representation of G .

Let $P'_n = [P_n, P_n]$ be the commutator subgroup of the pure braid group P_n . It was shown in [28] that the quotient B_n/P'_n is a crystallographic group for all $n \geq 3$. Moreover, it was proved that B_n/P'_n possesses torsion, and there is a one-to-one correspondence between the conjugacy classes of the finite-order elements of B_n/P'_n and the conjugacy classes of the elements of odd order of the symmetric group S_n .

The groups B_n/P'_n also arise in the study of pseudo-symmetric braided categories [42]. In this paper the authors consider the quotient, denoted by PS_n , of B_n by the normal subgroup generated by the relations

$$s_i s_{i+1}^{-1} s_i = s_{i+1} s_i^{-1} s_{i+1}, \quad i = 1, \dots, n-2,$$

and show that PS_n is isomorphic to B_n/P'_n .

2. LINEAR REPRESENTATION OF B_n/P'_n

Let $R = \mathbb{Z}[q^{\pm 1}, t^{\pm 1}]$ be the ring of Laurent polynomials with two variables q, t , and V be the free R -module with the basis $e_{i,j}$ for $1 \leq i < j \leq n$. The Lawrence-Bigelow-Krammer representation (shortly LBKR) is the linear representation of the braid group B_n by the automorphisms of V

$$\psi = \psi_{q,t} : B_n \rightarrow \text{Aut}(V)$$

which is given on the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ of B_n by the following rule

$$(29) \quad \psi(\sigma_k)(e_{i,j}) = \begin{cases} e_{i,j}, & k < i-1 \text{ or } k > j, \\ e_{i-1,j} + (1-q)e_{i,j}, & k = i-1, \\ tq(q-1)e_{i,i+1} + qe_{i+1,j}, & k = i < j-1, \\ tq^2e_{i,j}, & k = i = j-1, \\ e_{i,j} + tq^{k-i}(q-1)^2e_{k,k+1}, & i < k < j-1, \\ e_{i,j-1} + tq^{j-i}(q-1)e_{j-1,i}, & i < k = j-1, \\ (1-q)e_{i,j} + qe_{i,j+1}, & k = j. \end{cases}$$

Example 2.1. If $n = 3$, then the braid group B_3 is generated by σ_1, σ_2 , and the module V has the basis $\{e_{1,2}, e_{1,3}, e_{2,3}\}$. From formula (29) it follows that the linear maps $\psi(\sigma_1), \psi(\sigma_2), \psi(\sigma_1)^{-1}, \psi(\sigma_2)^{-1}$ can be presented by

the following matrices, which are written in the basis $e_{1,2}, e_{1,3}, e_{2,3}$ of V

$$\begin{aligned} [\psi(\sigma_1)] &= \begin{pmatrix} tq^2 & 0 & 0 \\ tq(q-1) & 0 & q \\ 0 & 1 & 1-q \end{pmatrix}, \\ [\psi(\sigma_2)] &= \begin{pmatrix} 1-q & q & 0 \\ 1 & 0 & tq^2(q-1) \\ 0 & 0 & tq^2 \end{pmatrix}, \\ [\psi(\sigma_1)]^{-1} &= \begin{pmatrix} t^{-1}q^{-2} & 0 & 0 \\ -1+2q^{-1}-q^{-2} & 1-q^{-1} & 1 \\ -q^{-1}+q^{-2} & q^{-1} & 0 \end{pmatrix}, \\ [\psi(\sigma_2)]^{-1} &= \begin{pmatrix} 0 & 1 & -q+1 \\ q^{-1} & 1-q^{-1} & -q+2-q^{-1} \\ 0 & 0 & t^{-1}q^{-2} \end{pmatrix}. \end{aligned}$$

For two complex numbers $q_0, t_0 \in \mathbb{C}$ denote by V_{q_0, t_0} the free module over the ring $\mathbb{Z}[q_0, t_0]$ with the same basis as V , and denote by ψ_{q_0, t_0} the specialization of $\psi_{q, t}$ obtained by putting $q = q_0, t = t_0$. The above specialization gives the linear representation $\psi_{q_0, t_0} : B_n \rightarrow \text{Aut}(V_{q_0, t_0})$. In particular, the map $\psi_{1, t} : B_n \rightarrow \text{Aut}(V_{1, t})$ is defined on the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ of B_n by the following rules

$$\psi_{1, t}(\sigma_k) : \begin{cases} e_{k, k+1} \mapsto te_{k, k+1}, \\ e_{i, j} \mapsto e_{s_k(i), s_k(j)}, \quad \text{if } (i, j) \neq (k, k+1) \text{ and } 1 \leq i < j \leq n, \end{cases}$$

where $s_k = (k, k+1)$ is the transposition from S_n . In this formula we assume that $e_{j, i} = e_{i, j}$ if $i < j$.

Lemma 2.1. *If $q = 1$ and $t = \pm 1$, then the image of B_n under the specialization $\psi_{1, \pm 1}$ of LBKR is isomorphic to the symmetric group S_n .*

Proof. It is enough to observe that $(\psi_{1, \pm 1}(\sigma_i))^2$ is the identity automorphism. This property immediately follows from the definition of $\psi_{1, \pm 1}$. \square

From the expressions of the generators $a_{i, j}$ ($1 \leq i < j \leq n$) of P_n introduced in Section 1.1 and direct calculations we have the following result.

Lemma 2.2. *The representation $\psi_{1, t} : B_n \rightarrow \text{Aut}(V_{1, t})$ acts on the generators $a_{i, j}$ ($1 \leq i < j \leq n$) of P_n by the following rules*

$$\psi_{1, t}(a_{ij}) : \begin{cases} e_{i, j} \mapsto t^2 e_{i, j}, \\ e_{k, l} \mapsto e_{k, l}, \quad \text{for } (i, j) \neq (k, l). \end{cases}$$

The following theorem is the main results of the present section.

Theorem 2.1. *The image of B_n under $\psi_{1, t}$ is isomorphic to B_n/P'_n .*

Proof. In Section 1.1 we have noticed that the following short exact sequence holds

$$1 \rightarrow P_n \rightarrow B_n \rightarrow S_n \xrightarrow{\pi} 1.$$

From this short exact sequence and the isomorphism

$$(B_n/P'_n)/(P_n/P'_n) = B_n/P_n$$

we have the following short exact sequence

$$1 \rightarrow P_n/P'_n \rightarrow B_n/P'_n \xrightarrow{\pi} S_n \rightarrow 1.$$

From this short exact sequence it follows that every element $b \in B_n/P'_n$ can be expressed in the form $b = p\lambda$, where $p \in P_n/P'_n$ and λ is a coset representative of P_n/P'_n in B_n/P'_n . In order to prove the statement of the theorem we have to prove that if b is non-trivial, then $\psi_{1,t}(b)$ is non-identity automorphism.

Let $b = p\lambda \neq 1$. If $\lambda \neq 1$, then $\pi(b) \neq 1$, and hence the composition of $\psi_{1,t}$ and specialization $t = 1$ send b to non-trivial automorphism. Therefore, $\psi_{1,t}(b)$ is non-trivial. If $\lambda = 1$ and $b \neq 1$, then b can be presented in the form

$$b = a_{1,2}^{\alpha_{1,2}} \left(a_{1,3}^{\alpha_{1,3}} a_{2,3}^{\alpha_{2,3}} \right) \dots \left(a_{1,n}^{\alpha_{1,n}} \dots a_{n-1,n}^{\alpha_{n-1,n}} \right) P'_n$$

for some integers $\alpha_{i,j} \in \mathbb{Z}$. From Lemma 2.2 it follows that the matrix of the automorphism $\psi_{1,t}(b)$ is the following diagonal matrix

$$[\psi_{1,t}(b)] = \text{diag}(t^{2\alpha_{1,2}}, t^{2\alpha_{1,3}}, t^{2\alpha_{2,3}} \dots t^{2\alpha_{1,n}} \dots t^{2\alpha_{n-1,n}}),$$

which is not the identity matrix. Hence, we have proved that if b is non-trivial, then its image $\psi_{1,t}(b)$ is non-trivial. \square

It was shown in [28] that the group B_n/P'_n possesses torsion, and there is a one-to-one correspondence between the conjugacy classes of the finite-order elements of B_n/P'_n and the conjugacy classes of the elements of odd order of the symmetric group S_n . The group B_3/P'_3 was considered separately in the paper [28]. Using the linear representation $\psi_{1,t}$ we obtain the following statement.

Proposition 2.1. *The group B_3/P'_3 has only 3-torsion. Every element which has one of the following forms*

$$a_{1,2}^\alpha a_{1,3}^\beta a_{2,3}^\gamma \sigma_1 \sigma_2, \quad a_{1,2}^\alpha a_{1,3}^\beta a_{2,3}^\gamma \sigma_2 \sigma_1,$$

where $\alpha, \beta, \gamma \in \mathbb{Z}$ are such that $\alpha + \beta + \gamma = -1$, has order 3. Any other non-trivial element has infinite order.

Proof. Since $\psi_{1,t}(B_n) \cong B_3/P'_3$, we can present elements of B_3/P'_3 by matrices in $\text{GL}_3(\mathbb{Z}[t^{\pm 1}])$. For $n = 3$ any element $b \in B_3/P'_3$ can be presented in the form $b = p\lambda$, where $p \in P_3/P'_3$ has the following form

$$p = \text{diag}(t^{2\alpha}, t^{2\beta}, t^{2\gamma}), \text{ for } \alpha, \beta, \gamma \in \mathbb{Z},$$

and λ is one of the following matrices

$$\begin{aligned} \lambda_0 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & \lambda_1 &= \begin{pmatrix} t & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, & \lambda_2 &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & t \end{pmatrix}, \\ \lambda_3 &= \begin{pmatrix} 0 & t & 0 \\ 0 & 0 & t \\ 1 & 0 & 0 \end{pmatrix}, & \lambda_4 &= \begin{pmatrix} 0 & 0 & 1 \\ t & 0 & 0 \\ 0 & t & 0 \end{pmatrix}, & \lambda_5 &= \begin{pmatrix} 0 & 0 & t \\ 0 & t & 0 \\ t & 0 & 0 \end{pmatrix}. \end{aligned}$$

Let us consider each λ_i separately. First, it is obvious that the element $p\lambda_0$ has infinite order. If the element $p\lambda_1$ has a finite order, then from the matrix form of λ_1 it is clear that this order has to be even. Since for an arbitrary integer k we have

$$(p\lambda_1)^{2k} = \text{diag}(t^{2k(2\alpha+1)}, t^{2k(\beta+\gamma)}, t^{2k(\beta+\gamma)}),$$

the element $p\lambda_1$ has infinite order. Further if the element $p\lambda_2$ has a finite order, then from the matrix form of λ_2 we see that this order is even. Since for an arbitrary integer k we have

$$(p\lambda_2)^{2k} = \text{diag}(t^{2k(\alpha+\beta)}, t^{2k(\alpha+\beta)}, t^{2k(2\gamma+1)}),$$

the element $p\lambda_2$ has infinite order. From the matrix representations it is clear that the order of the element $p\lambda_3$ is divisible by 3. For an arbitrary k we have the equality

$$(p\lambda_3)^{3k} = \text{diag}(t^{2k(\alpha+\beta+\gamma+1)}, t^{2k(\alpha+\beta+\gamma+1)}, t^{2k(\alpha+\beta+\gamma+1)}),$$

i. e. if $\alpha + \beta + \gamma + 1 = 0$, then $p\lambda_3$ has order 3, otherwise $p\lambda_3$ has infinite order. Using the same argument, the order of the element $p\lambda_4$ is divisible by 3, and for an arbitrary k we have

$$(p\lambda_4)^{3k} = \text{diag}(t^{2k(\alpha+\beta+\gamma+1)}, t^{2k(\alpha+\beta+\gamma+1)}, t^{2k(\alpha+\beta+\gamma+1)}),$$

i. e. if $\alpha + \beta + \gamma + 1 = 0$, then $p\lambda_4$ has order 3, otherwise $p\lambda_4$ has infinite order. It is easy to check that elements of the form $p\lambda_5$ have infinite order \square

In general case it is not difficult to prove the following statement.

Proposition 2.2. *If b is a non-trivial element in B_n/P'_n such that the permutation $\pi(b)$ is non-identity but fixes some points, then b has infinite order.*

3. LINEAR REPRESENTATION OF VB_n/VP'_n

In Section 1.3 we have noticed that the following short exact sequence holds

$$1 \rightarrow VP_n \rightarrow VB_n \rightarrow S_n \rightarrow 1.$$

From this short exact sequence and the isomorphism

$$(VB_n/VP'_n)/(VP_n/VP'_n) = VB_n/VP_n$$

we have the following short exact sequence

$$1 \rightarrow VP_n/VP'_n \rightarrow VB_n/VP'_n \rightarrow S_n \rightarrow 1,$$

where $VP_n/VP'_n = \mathbb{Z}^{n(n-1)}$ (since VP_n is generated by the elements $\lambda_{i,j}$ for $1 \leq i \neq j \leq n$), $S_n = \langle \rho_1, \rho_2, \dots, \rho_{n-1} \rangle$, and the group S_n acts on the generators of VP_n/VP'_n by the permutation of indices. From the equality $VB_n = VP_n \rtimes S_n$ it follows that $VB_n/VP'_n = (VP_n/VP'_n) \rtimes S_n$, and hence, VB_n/VP'_n is a crystallographic group.

In this section we construct an extension of the representation $\psi_{1,t}$ from the braid group B_n to the virtual braid group VB_n . Let U be the free module with the basis $e_{i,j}$, $1 \leq i < j \leq n$ over the ring $\mathbb{Z}[t^{\pm 1}, t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$ of Laurent polynomials with n variables t, t_1, \dots, t_{n-1} . Let us define the map $\Psi : VB_n \rightarrow \text{Aut}(U)$ on the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$, $\rho_1, \rho_2, \dots, \rho_{n-1}$ of VB_n by the following rules:

$$(30) \quad \Psi(\sigma_i) = \psi_{1,t}(\sigma_i),$$

$$(31) \quad \Psi(\rho_i) : \begin{cases} e_{i,i+1} \mapsto e_{i,i+1}, \\ e_{k,l} \mapsto e_{k,l}, & k \neq i, i+1, l \neq i, i+1, \\ e_{i,j} \mapsto t_i e_{i+1,j}, & i < j-1, \\ e_{i+1,j} \mapsto t_i^{-1} e_{i,j}, \\ e_{k,i} \mapsto t_i e_{k,i+1}, \\ e_{k,i+1} \mapsto t_i^{-1} e_{k,i}. \end{cases}$$

The following property can be easily verified using direct calculations.

Proposition 3.1. *The map $\Psi : VB_n \rightarrow \text{Aut}(U)$, defined by (30) and (31) is a representation. The images $\Psi(\sigma_i)$ and $\Psi(\rho_i)$ satisfy both forbidden relations (23) for VB_n .*

The following statement describes the image of Ψ .

Theorem 3.1. *The image $\Psi(VB_n)$ is isomorphic to VB_n/VP'_n .*

Proof. Using direct calculations it is easy to see that the image $\Psi(b)$ of every element $b \in VP'_n$ is the trivial automorphism. Hence, in order to prove the theorem we have to prove that the image of any elements $b \in VB_n \setminus VP'_n$ is non-trivial. If b does not belong to VP_n , then the image of b under Ψ is clearly non-trivial automorphism of U . If b is a non-trivial element of VP_n , then looking to the representative of b in bVP'_n it is easy to see that the matrix of $\Psi(b)$ is a non-identity diagonal matrix. \square

The following questions arise naturally from the paper [28] where the same questions are studied for classical linear groups.

Question 3.1. *What are the possible finite orders of elements from VB_n/VP'_n which do not belong to S_n ?*

Question 3.2. *Which Bieberbach subgroups belong to VB_n/VP'_n ?*

Let us consider the particular case $n = 3$. The virtual braid group VB_3 is generated by four elements $\sigma_1, \sigma_2, \rho_1, \rho_2$ and it has the following defining

relations

$$\sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2, \quad \rho_1\rho_2\rho_1 = \rho_2\rho_1\rho_2, \quad \rho_1^2 = \rho_2^2 = 1, \quad \rho_1\rho_2\sigma_1 = \sigma_2\rho_1\rho_2.$$

The images of the generators $\sigma_1, \sigma_2, \rho_1, \rho_2$ of VB_n under the representation $\Psi : VB_3 \rightarrow \text{GL}_3(\mathbb{Z}[t^{\pm 1}, t_1^{\pm 1}, t_2^{\pm 1}])$ are the following matrices

$$\begin{aligned} \Psi(\sigma_1) &= \begin{pmatrix} t & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, & \Psi(\sigma_2) &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & t \end{pmatrix}, \\ \Psi(\rho_1) &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & t_1 \\ 0 & t_1^{-1} & 0 \end{pmatrix}, & \Psi(\rho_2) &= \begin{pmatrix} 0 & t_2 & 0 \\ t_2^{-1} & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

The group VP_3 is generated by the elements

$$\begin{aligned} \lambda_{12} &= \rho_1\sigma_1^{-1}, & \lambda_{21} &= \sigma_1^{-1}\rho_1, & \lambda_{13} &= \rho_2\lambda_{12}\rho_2, \\ \lambda_{31} &= \rho_2\lambda_{21}\rho_2, & \lambda_{23} &= \rho_2\sigma_2^{-1}, & \lambda_{32} &= \sigma_2^{-1}\rho_2. \end{aligned}$$

Using direct calculations it is easy to see that the images of the elements $\lambda_{12}, \lambda_{21}, \lambda_{13}, \lambda_{31}, \lambda_{23}, \lambda_{32}$ under the representations Ψ are the following diagonal matrices

$$\begin{aligned} \Psi(\lambda_{12}) &= \text{diag}(t^{-1}, t_1, t_1^{-1}), & \Psi(\lambda_{21}) &= \text{diag}(t^{-1}, t_1^{-1}, t_1), \\ \Psi(\lambda_{13}) &= \text{diag}(t_1, t, t_1^{-1}), & \Psi(\lambda_{31}) &= \text{diag}(t_1^{-1}, t^{-1}, t_1), \\ \Psi(\lambda_{23}) &= \text{diag}(t_2, t_2^{-1}, t^{-1}), & \Psi(\lambda_{32}) &= \text{diag}(t_2^{-1}, t_2, t^{-1}), \end{aligned}$$

in particular, $\Psi(\lambda_{12}) \neq \Psi(\lambda_{21})$. Let us demonstrate that there are torsion elements in VB_3/VP_3' which do not belong to S_3 . Consider the element

$$p = \lambda_{12}^a \lambda_{21}^\alpha \lambda_{13}^b \lambda_{23}^c \lambda_{31}^\beta \lambda_{32}^\gamma d \rho_1,$$

where $\alpha, \beta, \gamma, a, b, c \in \mathbb{Z}$ and $d \in VP_3'$. Then

$$\Psi(p)^2 = \text{diag} \left(t^{-2(a+\alpha)} t_1^{2(b-\beta)} t_2^{-2(c-\gamma)}, t^{b-\beta-\gamma-c} t_1^{\beta-b} t_2^{\gamma-c}, t^{b-\beta-\gamma-c} t_1^{\beta-b} t_2^{2(\gamma-c)} \right).$$

From this formula it is clear that if $\alpha = -a$, $\beta = b$, and $\gamma = c = 0$, then $\Psi(p)^2 = E$. The same arguments lead to the following statement.

Proposition 3.2. *The group VB_3/VP_3' has infinitely many elements of order 2.*

4. DECOMPOSITION AND LINEARITY OF VB_3

Properties of the group VP_3 were studied in [9]. In particular, the certain decomposition of this group was found in [9]. In the present section we construct a simpler decomposition and prove that VP_3 , and hence VB_3 , is linear. The following statement is the first main result of this section.

Proposition 4.1. $VP_3 = (F_3 \rtimes F_2) * \mathbb{Z}$.

Proof. By Theorem 1.1, the group VP_3 has a presentation with six generators $\lambda_{12}, \lambda_{21}, \lambda_{13}, \lambda_{23}, \lambda_{32}, \lambda_{31}$, and six relations which can be write as the following conjugations:

$$(32) \quad \lambda_{12}(\lambda_{13}\lambda_{23}) = (\lambda_{23}\lambda_{13})\lambda_{12} \Leftrightarrow (\lambda_{13}\lambda_{23})^{\lambda_{12}^{-1}} = \lambda_{23}\lambda_{13},$$

$$(33) \quad \lambda_{13}(\lambda_{12}\lambda_{32}) = (\lambda_{32}\lambda_{12})\lambda_{13} \Leftrightarrow (\lambda_{32}\lambda_{13}^{-1})^{\lambda_{12}^{-1}} = \lambda_{13}^{-1}\lambda_{32},$$

$$(34) \quad \lambda_{31}(\lambda_{32}\lambda_{12}) = (\lambda_{12}\lambda_{32})\lambda_{31} \Leftrightarrow (\lambda_{32}\lambda_{31}^{-1})^{\lambda_{12}^{-1}} = \lambda_{31}^{-1}\lambda_{32},$$

$$(35) \quad \lambda_{21}(\lambda_{23}\lambda_{13}) = (\lambda_{13}\lambda_{23})\lambda_{21} \Leftrightarrow (\lambda_{23}\lambda_{13})^{\lambda_{21}^{-1}} = \lambda_{13}\lambda_{23},$$

$$(36) \quad \lambda_{23}(\lambda_{21}\lambda_{31}) = (\lambda_{31}\lambda_{21})\lambda_{23} \Leftrightarrow (\lambda_{31}\lambda_{23}^{-1})^{\lambda_{21}^{-1}} = \lambda_{23}^{-1}\lambda_{31},$$

$$(37) \quad \lambda_{32}(\lambda_{31}\lambda_{21}) = (\lambda_{21}\lambda_{31})\lambda_{32} \Leftrightarrow (\lambda_{31}\lambda_{32})^{\lambda_{21}^{-1}} = \lambda_{32}\lambda_{31}.$$

Note that equations (32) and (33) are also equivalent to the following commutation relations

$$(\lambda_{13}\lambda_{23})^{\lambda_{12}^{-1}\lambda_{13}^{-1}} = \lambda_{13}\lambda_{23}, \quad (\lambda_{32}\lambda_{13}^{-1})^{\lambda_{12}^{-1}\lambda_{13}^{-1}} = \lambda_{32}\lambda_{13}^{-1},$$

respectively. Let us define the following elements of VP_3 :

$$b_{11} = \lambda_{21}\lambda_{12}, \quad b_{12} = \lambda_{13}\lambda_{12}, \quad b_{21} = \lambda_{13}\lambda_{23}, \quad b_{22} = \lambda_{32}\lambda_{31}, \quad b_{23} = \lambda_{32}\lambda_{23},$$

and rewrite all relations (32)-(37) in order to express how the group $\langle b_{11}, b_{12} \rangle$ acts on the group $\langle b_{21}, b_{22}, b_{23} \rangle$ by conjugations:

$$\begin{aligned} b_{21}^{b_{12}^{-1}} &= (\lambda_{13}\lambda_{23})^{\lambda_{12}^{-1}\lambda_{13}^{-1}} \stackrel{(32)}{=} \lambda_{13}\lambda_{23} = b_{21}, \\ b_{22}^{b_{12}^{-1}} &= \lambda_{13}\lambda_{12} \cdot \lambda_{32}\lambda_{31} \cdot \lambda_{12}^{-1}\lambda_{13}^{-1} \stackrel{(34)}{=} \lambda_{13} \cdot \lambda_{31}\lambda_{32} \cdot \lambda_{13}^{-1} \\ &= \lambda_{13}\lambda_{23} \cdot \lambda_{23}^{-1}\lambda_{32}^{-1} \cdot \lambda_{32}\lambda_{31} \cdot \lambda_{32}\lambda_{23} \cdot \lambda_{23}^{-1}\lambda_{13}^{-1} = b_{22}^{b_{23}b_{21}^{-1}}, \\ b_{23}^{b_{12}^{-1}} &= (\lambda_{32}\lambda_{23})^{\lambda_{12}^{-1}\lambda_{13}^{-1}} = (\lambda_{32}\lambda_{13}^{-1} \cdot \lambda_{13}\lambda_{23})^{\lambda_{12}^{-1}\lambda_{13}^{-1}} \\ &\stackrel{(32),(33)}{=} \lambda_{32}\lambda_{13}^{-1} \cdot \lambda_{13}\lambda_{23} = b_{23}, \\ b_{21}^{b_{11}^{-1}} &= (\lambda_{13}\lambda_{23})^{\lambda_{12}^{-1}\lambda_{21}^{-1}} \stackrel{(32),(35)}{=} \lambda_{13}\lambda_{23} = b_{21}, \\ b_{22}^{b_{11}^{-1}} &= (\lambda_{32}\lambda_{31})^{\lambda_{12}^{-1}\lambda_{21}^{-1}} \stackrel{(34),(37)}{=} \lambda_{32}\lambda_{31} = b_{22}, \\ b_{23}^{b_{11}^{-1}} &= (\lambda_{32}\lambda_{13}^{-1} \cdot \lambda_{13}\lambda_{23})^{\lambda_{12}^{-1}\lambda_{21}^{-1}} \stackrel{(33),(32)}{=} (\lambda_{13}^{-1}\lambda_{32} \cdot \lambda_{23}\lambda_{13})^{\lambda_{21}^{-1}} \\ &= (\lambda_{13}^{-1}\lambda_{23}^{-1} \cdot \lambda_{23}\lambda_{31}^{-1} \cdot \lambda_{31}\lambda_{32} \cdot \lambda_{23}\lambda_{13})^{\lambda_{21}^{-1}} \\ &\stackrel{(35),(36),(37),(35)}{=} \lambda_{23}^{-1}\lambda_{13}^{-1} \cdot \lambda_{31}^{-1}\lambda_{23} \cdot \lambda_{31}\lambda_{23} \cdot \lambda_{32}\lambda_{31} \cdot \lambda_{13}\lambda_{23} \\ &= \lambda_{23}^{-1}\lambda_{13}^{-1} \cdot \lambda_{31}^{-1}\lambda_{32}^{-1} \cdot \lambda_{32}\lambda_{23} \cdot \lambda_{32}\lambda_{31} \cdot \lambda_{13}\lambda_{23} = b_{23}^{b_{22}b_{21}^{-1}}. \end{aligned}$$

To summarize, we have six calculations which show the following six relations:

$$(38) \quad b_{21}^{b_{12}^{-1}} = b_{21}, \quad b_{22}^{b_{12}^{-1}} = b_{22}^{b_{23}b_{21}^{-1}}, \quad b_{23}^{b_{12}^{-1}} = b_{23},$$

$$(39) \quad b_{21}^{b_{11}^{-1}} = b_{21}, \quad b_{22}^{b_{11}^{-1}} = b_{22}, \quad b_{23}^{b_{11}^{-1}} = b_{23}^{b_{22}b_{21}}.$$

Moreover, in the sequence of six calculations we made, we used exactly one of the relations (32)-(37) which was not yet used in the preceding ones. It means that the set of relations (38)-(39) is equivalent to the set of relations (32)-(37). In particular, the groups $F_2 = \langle b_{11}, b_{12} \rangle$, $F_3 = \langle b_{21}, b_{22}, b_{23} \rangle$ are free, and the five elements $b_{11}, b_{12}, b_{21}, b_{22}, b_{23}$ generate the semidirect product $F_3 \rtimes F_2$, where F_2 acts on F_3 by the way described in (32)-(37). Finally, the five generators $b_{11}, b_{12}, b_{21}, b_{22}, b_{23}$ together with any one of the generators λ_{ij} , for instance with λ_{13} , generate the full group VP_3 . Since the six relations describing the semidirect product structure form a complete set of relations of VP_3 , the group $\langle \lambda_{13} \rangle$ forms a free factor in VP_3 and we have the decomposition

$$VP_3 = (\langle b_{21}, b_{22}, b_{23} \rangle \rtimes \langle b_{11}, b_{12} \rangle) * \langle \lambda_{13} \rangle.$$

The proposition is proved. \square

Let us now prove that the group VP_3 is linear. Denote by $\mu : VB_n \rightarrow S_n$ the homomorphism defined on the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \rho_1, \rho_2, \dots, \rho_{n-1}$ of VB_n as follows:

$$\mu(\sigma_i) = 1, \quad \mu(\rho_i) = \rho_i, \quad i = 1, 2, \dots, n-1,$$

where $S_n = \langle \rho_1, \rho_2, \dots, \rho_{n-1} \rangle$. Denote by H_n the normal closure of B_n in VB_n . It is obvious that $\text{Ker}(\mu)$ coincides with H_n . Now, define the following elements:

$$x_{i,i+1} = \sigma_i, \quad x_{i+1,i} = \rho_i \sigma_i \rho_i,$$

and

$$\begin{aligned} x_{i,j} &= \rho_{j-1} \cdots \rho_{i+1} \sigma_i \rho_{i+1} \cdots \rho_{j-1} & \text{for } 1 \leq i < j-1 \leq n-1, \\ x_{j,i} &= \rho_{j-1} \cdots \rho_{i+1} \rho_i \sigma_i \rho_i \rho_{i+1} \cdots \rho_{j-1} & \text{for } 1 \leq i < j-1 \leq n-1. \end{aligned}$$

The symmetric group $S_n = \langle \rho_1, \rho_2, \dots, \rho_{n-1} \rangle$ acts on the elements $x_{i,j}$ for $1 \leq i \neq j \leq n$ by permutation of indices, i. e.,

$$\rho x_{i,j} \rho^{-1} = x_{\rho(i), \rho(j)}, \quad \rho \in S_n.$$

The following proposition is proved in [43] (see also [6]).

Proposition 4.2. 1) *The group H_n admits a presentation with the generators $x_{k,l}$ for $1 \leq k \neq l \leq n$, and the defining relations:*

$$(40) \quad x_{i,j} x_{k,l} = x_{k,l} x_{i,j},$$

$$(41) \quad x_{i,k} x_{k,j} x_{i,k} = x_{k,j} x_{i,k} x_{k,j},$$

where distinct letters stand for distinct indices.

2) The group VB_n is isomorphic to $H_n \rtimes S_n$ where S_n acts on the generators $x_{i,j}$ of H_n by permutation of indices.

Corollary 4.1. *The group H_2 is isomorphic to $\mathbb{Z} * \mathbb{Z}$ which is residually nilpotent.*

Now we are ready to prove the second main result of this section.

Theorem 4.1. *The group VB_3 is linear.*

Proof. Since $VB_3 = H_3 \rtimes S_3$, the group VB_3 is linear if and only if H_3 is linear. The group H_3 decomposes into a free product $G_1 * G_2$, where $G_1 = \langle x_{1,2}, x_{2,3}, x_{3,1} \rangle$ and $G_2 = \langle x_{1,3}, x_{3,2}, x_{2,1} \rangle$. The group G_i , $i = 1, 2$ is isomorphic to the 2-nd affine Artin-Tits group of type \mathcal{A} , also called circular braid group on 3 strands (see [1]). Since G_i can be embedded in B_4 (see [47]), it is linear. Since the free product of linear groups is linear, H_3 is linear. \square

Question 4.1. *What is the minimal dimension of the faithful linear representation of VP_3 ?*

Question 4.2. *What is the image of H_n under the representation Ψ ?*

5. UNIVERSAL BRAID GROUP

The universal braid group UB_n was defined in [4] as a group with generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$, $\tau_1, \tau_2, \dots, \tau_{n-1}$ which satisfy braid relations (1)-(2), commutativity relations

$$\tau_i \tau_j = \tau_j \tau_i, \quad |i - j| \geq 2,$$

and mixed relations

$$\tau_i \sigma_j = \sigma_j \tau_i \quad |i - j| \geq 2.$$

Recall [22] that the Artin's group of type I , denoted by A_I , is the group with generators a_i , $i \in I$ and defining relations

$$a_i a_j a_i \dots = a_j a_i a_j \dots, \quad i, j \in I,$$

where the words from the left and from the right hand side of the equality above consist of m_{ij} alternating letters a_i and a_j . The following statement is proved in [4].

Proposition 5.1. 1) *The elements $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ generate the braid group B_n in the universal braid group UB_n .*

2) *There are homomorphisms $\varphi_{US} : UB_n \rightarrow SG_n$, $\varphi_{UV} : UB_n \rightarrow VB_n$ and $\varphi_{UB} : UB_n \rightarrow B_n$.*

3) *The group UB_n is the Artin's group.*

Statement 2) of Proposition 5.1 is the reason why the group UB_n is called the universal braid group. The commutator subgroup of UB_n is studied in [23].

Denote by T_n the group generated by $\tau_1, \tau_2, \dots, \tau_{n-1}$ with the following defining commutativity relations

$$\tau_i \tau_j = \tau_j \tau_i, \quad |i - j| > 1.$$

It is clear that T_n forms a subgroup in the singular braid group SG_n and a subgroup in the universal braid group UB_n . By adding to T_n the relations $\tau_i^2 = 1$ for $i = 1, 2, \dots, n - 1$ we get the twin group or the flat braid group (see [16]).

Since the elements $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ generate a subgroup of UB_n which is isomorphic to the braid group B_n , the elements $\tau_1, \tau_2, \dots, \tau_{n-1}$ generate a subgroup of UB_n which is isomorphic to T_n , and the group UB_n is generated by the elements $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \tau_1, \tau_2, \dots, \tau_{n-1}$, in order to define some representation of UB_n we have to define a representation of $B_n < UB_n$, a representation of $T_n < UB_n$, and then match these two representations.

5.1. Representations by automorphisms of a free group. Further we consider several representations of T_n by automorphisms of free groups. The first one-parametric family of representations consists of the representations $\phi_k : T_n \rightarrow \text{Aut}(F_n)$ for $k \in \mathbb{Z}$, where ϕ_k acts on the generators $\tau_1, \tau_2, \dots, \tau_{n-1}$ of T_n by the following rule

$$\phi_k(\tau_i) : \begin{cases} x_i \rightarrow x_{i+1}^{-k} x_i x_{i+1}^k, \\ x_j \rightarrow x_j, \end{cases} \quad \text{if } j \neq i.$$

The second family of representations is the two-parametric family of representations $\psi_{k,l} : T_n \rightarrow \text{Aut}(F_n)$ for $k, l \in \mathbb{Z} \setminus \{0\}$, where $\psi_{k,l}$ acts on the generators $\tau_1, \tau_2, \dots, \tau_{n-1}$ of T_n by the following rule

$$\psi_{k,l}(\tau_i) : \begin{cases} x_i \rightarrow x_i^{-l} x_{i+1}^{-k} x_i x_{i+1}^k x_i^l, \\ x_{i+1} \rightarrow x_i^{-l} x_{i+1} x_i^l, \\ x_j \rightarrow x_j \end{cases} \quad \text{if } j \neq i, i + 1.$$

It is evidently that $\psi_{k,0} = \phi_k$.

Finally, the representation $\xi : T_n \rightarrow \text{Aut}(F_{3n-2})$, where the free group F_{3n-2} has the set of free generators

$$\{x_1, \dots, x_n, y_{1,1}, \dots, y_{1,n-1}, y_{2,1}, \dots, y_{2,n-1}\},$$

acts on the generators $\tau_1, \tau_2, \dots, \tau_{n-1}$ of T_n by the following rule

$$\xi(\tau_i) : \begin{cases} x_i \rightarrow x_i y_{i,1}, \\ x_{i+1} \rightarrow x_{i+1} y_{i,2}, \\ x_j \rightarrow x_j, & \text{if } j \neq i \\ y_{i,m} \rightarrow y_{i,m}, & \text{if } i = 1, 2 \text{ and } m = 1, \dots, n - 1. \end{cases}$$

From the definition it is clear that $\phi_k(\tau_i) = \varepsilon_{i,i+1}^k$ and $\psi_{k,l}(\tau_i) = \varepsilon_{i,i+1}^k \varepsilon_{i+1,i}^l$, therefore, representations ϕ_k and $\psi_{k,l}$ act from T_n to the group of basis conjugating automorphisms Cb_n .

Lemma 5.1. *Let w be an element in T_n , and $s \in \{1, 2, \dots, n-1\}$ be a minimal number such that w cannot be written in terms of generators $\tau_1, \tau_2, \dots, \tau_{n-1}$ without τ_s . If $k \neq 0$, then $\phi_k(w)(x_s) \neq x_s$.*

Proof. Every word w in the group T_n can be written in the normal form

$$w = \tau_{s_1}^{\alpha_1} \tau_{s_2}^{\alpha_2} \dots \tau_{s_r}^{\alpha_r},$$

where $\alpha_1, \alpha_2, \dots, \alpha_r$ are non-zero integers, and for all $i = 1, 2, \dots, r-1$ we have either $s_i < s_{i+1}$ or $s_i = s_{i+1} + 1$. If the word w is written in the normal form using generators $\tau_{s_1}, \tau_{s_2}, \dots, \tau_{s_r}$, then w cannot be written (even not in normal form) without using these generators. From this fact it follows that the word w from the formulation of the lemma can be written in terms of $\tau_s, \tau_{s+1}, \dots, \tau_{n-1}$, and the generator τ_s is used in the expression of w .

We will prove the lemma using specific induction on n . First, we will prove that the lemma is correct in the case when $s = n-1$, then we will assume that the lemma is correct for all numbers $s+1, s+2, \dots, n-1$, and using this conjecture we will prove that the lemma is correct for s .

The basis of the induction is obvious: if $s = n-1$, then $w = \tau_{n-1}^\alpha$ for $\alpha \neq 0$, and from the definition of ϕ_k it is clear that

$$\phi_k(w)(x_{n-1}) = \phi_k(\tau_{n-1}^\alpha)(x_{n-1}) \neq x_{n-1}.$$

Let us assume that the lemma is correct for all numbers $s+1, s+2, \dots, n-1$, and let us prove the lemma for s .

By contrary, suppose that w is a counter example with the minimal value $|w|$, where $|w|$ denotes the length of w , when it is written in terms of generators $\tau_1, \tau_2, \dots, \tau_{n-1}$. This assumption means that w cannot be written in terms of generators $\tau_1, \tau_2, \dots, \tau_{n-1}$ without τ_s and $\phi_k(w)(x_s) = x_s$. Since $s \in \{1, 2, \dots, n-1\}$ is a minimal number such that w cannot be written in terms of generators $\tau_1, \tau_2, \dots, \tau_{n-1}$ without τ_s , the element w can be written in the following form

$$(42) \quad w = w_1 \tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} \dots w_m \tau_s^{\alpha_m} w_{m+1},$$

where $\alpha_1, \alpha_2, \dots, \alpha_m$ are non-zero integers, and w_1, w_2, \dots, w_{m+1} are elements from T_n which can be written in terms of $\tau_{s+1}, \tau_{s+2}, \dots, \tau_{n-1}$.

From the definition of the representation ϕ_k it is clear that since w_{m+1} can be written in terms of generators $\tau_{s+1}, \tau_{s+2}, \dots, \tau_{n-1}$, and $\phi_k(w)(x_s) = x_s$ then for the element

$$\tilde{w} = w_{m+1} w w_{m+1}^{-1} = w_{m+1} w_1 \tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} \dots w_m \tau_s^{\alpha_m}$$

we have the equality $\phi_k(\tilde{w})(x_s) = x_s$. Since $|\tilde{w}| \leq |w|$, considering the element \tilde{w} instead of w we can assume that the equality $w_{m+1} = 1$ holds in (42), i. e.

$$(43) \quad w = w_1 \tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} \dots w_m \tau_s^{\alpha_m}.$$

Since the element w_1 can be written in terms of generators $\tau_{s+1}, \tau_{s+2}, \dots, \tau_{n-1}$ from the definition of the representation ϕ_k it is clear that

$$\phi_k(w_1 \tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} \dots w_m \tau_s^{\alpha_m})(x_s) = \phi_k(\tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} \dots w_m \tau_s^{\alpha_m})(x_s).$$

Since $|\tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} \dots w_m \tau_s^{\alpha_m}| \leq |w_1 \tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} \dots w_m \tau_s^{\alpha_m}| = |w|$, considering the element $\tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} \dots w_m \tau_s^{\alpha_m}$ instead of w we can assume that the equality $w_1 = 1$ holds in (42), i. e.

$$(44) \quad w = \tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} w_3 \dots w_m \tau_s^{\alpha_m}.$$

Consider the following elements in T_n :

$$\begin{aligned} u_1 &= w_2 w_3 w_4 \dots w_m, \\ u_2 &= w_3 w_4 \dots w_m, \\ u_3 &= w_4 \dots w_m, \\ &\vdots \\ u_{m-1} &= w_m, \\ u_m &= 1. \end{aligned}$$

From the definition of the representation ϕ_k and direct calculations it follows that

$$(45) \quad \phi_k(w)(x_s) = x_s^{(\phi_k(u_m)(x_{s+1}^{\beta_m}))(\phi_k(u_{m-1})(x_{s+1}^{\beta_{m-1}})) \dots (\phi_k(u_2)(x_{s+1}^{\beta_2}))(\phi_k(u_1)(x_{s+1}^{\beta_1}))},$$

where $\beta_i = k\alpha_i$ for $i = 1, 2, \dots, m$. Since $\phi_k(w)(x_s) = x_s$ and (from the definition of ϕ_k) the element

$$\left(\phi_k(u_m)(x_{s+1}^{\beta_m})\right) \left(\phi_k(u_{m-1})(x_{s+1}^{\beta_{m-1}})\right) \dots \left(\phi_k(u_2)(x_{s+1}^{\beta_2})\right) \left(\phi_k(u_1)(x_{s+1}^{\beta_1})\right)$$

can be written in terms of $x_{s+1}, x_{s+2}, \dots, x_n$, from equality (45) we conclude the equality

$$(46) \quad \left(\phi_k(u_m)(x_{s+1}^{\beta_m})\right) \left(\phi_k(u_{m-1})(x_{s+1}^{\beta_{m-1}})\right) \dots \left(\phi_k(u_2)(x_{s+1}^{\beta_2})\right) \left(\phi_k(u_1)(x_{s+1}^{\beta_1})\right) = 1.$$

From the definition of ϕ_k it is easy to see that for $i = 1, 2, \dots, m$ we have the equality

$$(47) \quad \left(\phi_k(u_i)(x_{s+1}^{\beta_i})\right) = \left(x_{s+1}^{\beta_i}\right)^{v_i},$$

where v_1, v_2, \dots, v_m are elements in F_n which can be written in terms of $x_{s+2}, x_{s+3}, \dots, x_n$. From the equality $u_m = 1$ we have

$$\left(\phi_k(u_m)(x_{s+1}^{\beta_m})\right) = x_{s+1}^{\beta_m}$$

and taking into account equality (47) we can rewrite equality (46) in the following form

$$(48) \quad x_{s+1}^{\beta_m} \left(x_{s+1}^{\beta_{m-1}}\right)^{v_{m-1}} \dots \left(x_{s+1}^{\beta_2}\right)^{v_2} \left(x_{s+1}^{\beta_1}\right)^{v_1} = 1.$$

Due to the fact that for $i = 1, 2, \dots, m$ the elements $v_2, v_3, \dots, v_m \in F_n$ can be written in terms of $x_{s+2}, x_{s+3}, \dots, x_n$, in order for the element $x_{s+1}^{\beta_m}$ in

equality (48) to be reduced with something, there must be a number t such that $\left(x_{s+1}^{\beta_t}\right)^{v_t} = x_{s+1}^{\beta_t}$ and

$$(49) \quad x_{s+1}^{\beta_m} \left(x_{s+1}^{\beta_{m-1}}\right)^{v_{m-1}} \cdots \left(x_{s+1}^{\beta_{t+1}}\right)^{v_{t+1}} \left(x_{s+1}^{\beta_t}\right)^{v_t} = 1.$$

From equality (49) it follows that

$$(50) \quad \begin{aligned} x_s &= x_s^{x_{s+1}^{\beta_m} \left(x_{s+1}^{\beta_{m-1}}\right)^{v_{m-1}} \cdots \left(x_{s+1}^{\beta_{t+1}}\right)^{v_{t+1}} \left(x_{s+1}^{\beta_t}\right)^{v_t}} \\ &= x_s^{\left(\phi_k(u_m)(x_{s+1}^{\beta_m})\right) \left(\phi_k(u_{m-1})(x_{s+1}^{\beta_{m-1}})\right) \cdots \left(\phi_k(u_t)(x_{s+1}^{\beta_t})\right)} \\ &= \phi_k(\tau_s^{\alpha_t} w_{t+1} \tau_s^{\alpha_{t+1}} w_{t+2} \cdots \tau_s^{\alpha_{m-1}} w_m \tau_s^{\alpha_m})(x_s) \end{aligned}$$

If $t \neq 1$, then

$$\left| \tau_s^{\alpha_t} w_{t+1} \tau_s^{\alpha_{t+1}} w_{t+2} \cdots \tau_s^{\alpha_{m-1}} w_m \tau_s^{\alpha_m} \right| < \left| \tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} w_3 \cdots w_m \tau_s^{\alpha_m} \right| = |w|$$

and equality (50) contradicts the fact that w is a counter example to the lemma with the minimal value $|w|$, hence we have $t = 1$, what means that

$$\phi_k(u_1)(x_{s+1}^{\beta_1}) = \left(x_{s+1}^{\beta_1}\right)^{v_1} = x_{s+1}^{\beta_1}.$$

This equality together with the definition of ϕ_k implies that the equality $\phi_k(u_1)(x_{s+1}) = x_{s+1}$ holds.

Let us calculate $\phi_k(w)(x_{s+1})$. From the definition of ϕ_k we have the following equality

$$\begin{aligned} \phi_k(w)(x_{s+1}) &= \phi_k(\tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} w_3 \cdots w_m \tau_s^{\alpha_m})(x_{s+1}) = \\ &= \phi_k(w_2 w_3 \cdots w_m)(x_{s+1}) = \phi_k(u_1)(x_{s+1}) = x_{s+1}. \end{aligned}$$

So, now we have the following equalities

$$(51) \quad w = \tau_s^{\alpha_1} w_2 \tau_s^{\alpha_2} w_3 \cdots w_m \tau_s^{\alpha_m},$$

$$(52) \quad \phi_k(w)(x_s) = x_s,$$

$$(53) \quad \phi_k(w)(x_{s+1}) = x_{s+1}.$$

Note that in equality (51) the number m can not be equal to 1, since if $m = 1$, then $w = \tau_s^{\alpha_1}$, and it is clear that $\phi_k(\tau_s^{\alpha_1})(x_s) \neq x_s$ for $\alpha_1 \neq 0$. Therefore $m > 1$.

Consider the element

$$\tilde{w} = \tau_s^{-\alpha_1} w \tau_s^{\alpha_1} = w_2 \tau_s^{\alpha_2} w_3 \cdots w_m \tau_s^{\alpha_m + \alpha_1}.$$

From equalities (52), (53) we have the equality

$$\phi_k(\tilde{w})(x_s) = x_s.$$

Since the element w_2 can be written in terms of generators $\tau_{s+1}, \tau_{s+2}, \dots, \tau_{n-1}$ from the definition of the representation ϕ_k it is clear that

$$x_s = \phi_k(w_2 \tau_s^{\alpha_2} w_3 \cdots w_m \tau_s^{\alpha_m + \alpha_1})(x_s) = \phi_k(\tau_s^{\alpha_2} w_3 \cdots w_m \tau_s^{\alpha_m + \alpha_1})(x_s)$$

and since $|\tau_s^{\alpha_2} w_3 \dots w_m \tau_s^{\alpha_m + \alpha_1}| < |w_2 \tau_s^{\alpha_2} w_3 \dots w_m \tau_s^{\alpha_m + \alpha_1}|$ we must have that the element $\tau_s^{\alpha_2} w_3 \dots w_m \tau_s^{\alpha_m + \alpha_1}$ doesn't depend on τ_s . It means that $m = 2$ and $\alpha_1 + \alpha_m = 0$, and hence

$$w = \tau_s^{\alpha_1} w_2 \tau_s^{-\alpha_1},$$

where the word w_2 can be written in terms of generators $\tau_{s+1}, \tau_{s+2}, \dots, \tau_n$ and $\phi_k(w_2)(x_{s+1}) = x_{s+1}$. By the induction conjecture the word w can be written in terms of $\tau_{s+2}, \tau_{s+3}, \dots, \tau_{n-1}$, therefore from the defining relations of T_n we have $w = \tau_s^{\alpha_1} w_2 \tau_s^{-\alpha_1} = w_2$, i. e. w can be written without using τ_s , what contradicts conditions of the lemma. \square

The following theorem is the main result of this section.

Theorem 5.1. *The following statements hold.*

- (1) *If $k \neq 0$, then the representation $\phi_k : T_n \rightarrow Cb_n$ is faithful.*
- (2) *If $k \neq 0$, then the representation $\psi_{k,k} : T_n \rightarrow Cb_n$ is faithful.*
- (3) *The representation $\xi : T_n \rightarrow \text{Aut}(F_{3n-2})$ is faithful for $n = 3$ and has non-trivial kernel for $n \geq 4$.*

Proof. (1) Let w be an element from $\text{Ker}(\phi_k)$. Since for all $s \in \{1, 2, \dots, n\}$ we have $\phi_k(w)(x_s) = x_s$, from Lemma 5.1 it follows that in the representation of w in terms of $\tau_1, \tau_2, \dots, \tau_{n-1}$ there are no generators $\tau_1, \tau_2, \dots, \tau_{n-1}$, hence, $w = 1$.

(2) From result of Collins [24] follows that the subgroup $\langle \sigma_1^2, \sigma_2^2, \dots, \sigma_{n-1}^2 \rangle$ of B_n has the presentation

$$\langle \sigma_1^2, \sigma_2^2, \dots, \sigma_{n-1}^2 \mid \sigma_i^2 \sigma_j^2 = \sigma_j^2 \sigma_i^2 \text{ for } |i - j| \geq 2 \rangle,$$

hence the map $\tau_i \rightarrow \sigma_i^2$ for $i = 1, 2, \dots, n - 1$ gives an isomorphism

$$T_n \rightarrow \langle \sigma_1^2, \sigma_2^2, \dots, \sigma_{n-1}^2 \rangle < P_n.$$

From the main result of the paper [45] it follows that if $k \neq 0$, then the map $\sigma_i^2 \mapsto \varepsilon_{i,i+1}^k \varepsilon_{i+1,i}^k = \psi_{k,k}(\tau_i)$ for $i = 1, 2, \dots, n - 1$ gives a faithful representation of P_n to $\text{Aut}(F_n)$. Since $\psi_{k,k}$ is the superposition of these two maps

$$T_n \rightarrow \langle \sigma_1^2, \sigma_2^2, \dots, \sigma_{n-1}^2 \rangle \rightarrow \text{Aut}(F_n),$$

the representation $\psi_{k,k}$ is faithful.

(3) For $n = 3$ the group $\xi(T_3)$ acts on the generator x_2 as the free group F_2 which is isomorphic to T_3 , therefore the representation $\xi : T_n \rightarrow \text{Aut}(F_{3n-2})$ is faithful for $n = 3$. If $n \geq 4$, then from direct calculations it is easy to see that the element

$$\tau_1 \tau_2 \tau_3 \tau_2^{-1} \tau_1^{-1} \tau_2 \tau_3^{-1} \tau_2^{-1}$$

belongs to the kernel of ξ . \square

Let us now consider extensions of the representations $\phi_k, \psi_{k,l}$ from the group T_n to the group UB_n ,

$$\begin{aligned}\tilde{\phi}_k &: UB_n \rightarrow \text{Aut}(F_n), & k \in \mathbb{Z}, \\ \tilde{\psi}_{k,l} &: UB_n \rightarrow \text{Aut}(F_n), & k, l \in \mathbb{Z}\end{aligned}$$

which act on the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ of UB_n as Artin representation presented in Section 1.1. The following statement says that despite the fact that according to Theorem 5.1 the representation $\phi_k : T_3 \rightarrow \text{Aut}(F_3)$ is faithful for $k \neq 0$, the extension $\tilde{\phi}_k : UB_3 \rightarrow \text{Aut}(F_3)$ is not faithful.

Proposition 5.2. *The representation $\tilde{\phi}_k : UB_3 \rightarrow \text{Aut}(F_3)$ has non-trivial kernel for any $k \in \mathbb{Z}$.*

Proof. Using direct calculations it is easy to see that the element

$$\sigma_1 \tau_1^k \sigma_2 \tau_2^k \sigma_1 \tau_1^k \tau_2^{-k} \sigma_2^{-1} \tau_1^{-k} \sigma_1^{-1} \tau_2^{-k} \sigma_2^{-1}$$

belongs to the kernel of $\tilde{\phi}_k$. □

Problem 5.1. *Is it true that $\tilde{\psi}_{k,k}$ is faithful for $k \neq 0$?*

At the end of this subsection we introduce a representation

$$\Psi_{k,l} : UB_n \rightarrow \text{Aut}(F_{n+1}),$$

where $F_{n+1} = \langle x_1, x_2, \dots, x_n, y \rangle$ which generalizes the representation $\tilde{\psi}_{k,l}$, and which acts on the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \tau_1, \tau_2, \dots, \tau_{n-1}$ of UB_n by the rules

$$\begin{aligned}\Psi_{k,l}(\sigma_i) &: \begin{cases} x_i \mapsto x_i x_{i+1} x_i^{-1}, \\ x_{i+1} \mapsto x_i, \\ x_j \mapsto x_j, & j \neq i, i+1, \\ y \mapsto y, \end{cases} \\ \Psi_{k,l}(\tau_i) &: \begin{cases} x_i \rightarrow y^{-1} x_i^{-l} x_{i+1}^{-k} x_i x_{i+1}^k x_i^l y, \\ x_{i+1} \rightarrow y^{-1} x_i^{-l} x_{i+1}^l y, \\ x_j \rightarrow x_j & \text{if } j \neq i, i+1, \\ y \mapsto y, \end{cases}\end{aligned}$$

for $i = 1, 2, \dots, n-1$. It is easy to see that $\Psi_{k,l}$ generalizes the representation $\tilde{\psi}_{k,l}$: if we put $y = 1$, then $\Psi_{k,l}$ goes to $\tilde{\psi}_{k,l}$.

Problem 5.2. *Do there exist some integers k, l such that the representation $\Psi_{k,l} : UB_n \rightarrow \text{Aut}(F_{n-1})$ is faithful for all $n > 2$?*

5.2. Linear representations. As a corollary from the previous section, in the present section we construct a linear representations of the groups T_n . In order to do it we use the Magnus approach for construction of linear representations from representations by automorphisms of free groups. This

approach is described, for example, in [8, Section 4.1]. In the present section we will just introduce the final representation we got.

Let M be the free module with the basis $\{e_1, e_2, \dots, e_{n-1}\}$ over the ring $R = \mathbb{Z}[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_n^{\pm 1}]$ of Laurent polynomials with variables t_1, t_2, \dots, t_n . Define the map $\Theta : T_n \rightarrow \text{Aut}(M)$ which acts on the generators $\tau_1, \tau_2, \dots, \tau_{n-1}$ by the following rule

$$\Theta(\tau_i) : \begin{cases} e_i \mapsto t_{i+1}^{-1} e_i + t_{i+1}^{-1} (t_i - 1) e_{i+1}, \\ e_j \mapsto e_j, \end{cases} \quad j \neq i.$$

The following statement obviously follows from the fact that the matrices of the elements $\Theta(\tau_i)$ in the basis e_1, e_2, \dots, e_n are triangular.

Proposition 5.3. *The image $\text{Im } \Theta$ is a solvable group. Hence, for $n > 2$ the representation Θ has a non-trivial kernel.*

5.3. Universal pure braid group. In this section we study universal pure braid groups on small amount of strands. Consider the homomorphism $\pi : UB_n \rightarrow S_n$ given on generators of UB_n by the rule $\pi(\sigma_i) = \pi(\tau_i) = (i, i+1)$ for $i = 1, 2, \dots, n-1$. The kernel $\text{Ker}(\pi)$ of this homomorphism is known as the *universal pure braid group*, denoted by UP_n . The following short exact sequence holds:

$$1 \rightarrow UP_n \rightarrow UB_n \xrightarrow{\pi} S_n \rightarrow 1.$$

It is obvious that P_n is a subgroup of UP_n . Hence, the last group contains all generators and relations of P_n .

Theorem 5.2. *The following properties hold:*

- 1) $UB_2 = B_2 * T_2 \cong F_2$, $UB_3 = B_3 * T_3 = B_3 * F_2$,
- 2) $UP_2 = F_3$, $UP_3 = P_3 * F_{12}$.

Proof. 1) Follows from the presentations of UB_2 and UB_3 .

2) We will find the presentation of $UP_3 = \text{Ker}(\pi)$ by the Reidemeister-Shreier method using the Shreier set $\{1, \sigma_1, \sigma_2, \sigma_1\sigma_2, \sigma_2\sigma_1, \sigma_1\sigma_2\sigma_1, \sigma_2\sigma_1\sigma_2\}$. Since the restriction of homomorphism π on $B_3 \subset UB_3$ has P_3 as a kernel,

it is enough to consider only generators τ_1, τ_2 of UB_3 . Hence we get

$$\begin{aligned}
S_{1,\tau_1} &= \tau_1 \cdot (\overline{\tau_1})^{-1} = \tau_1 \cdot \sigma_1^{-1}, \\
S_{1,\tau_2} &= \tau_2 \cdot (\overline{\tau_2})^{-1} = \tau_2 \cdot \sigma_2^{-1}, \\
S_{\sigma_1,\tau_1} &= \sigma_1 \tau_1 \cdot (\overline{\sigma_1 \tau_1})^{-1} = \sigma_1 \tau_1, \\
S_{\sigma_1,\tau_2} &= \sigma_1 \tau_2 \cdot (\overline{\sigma_1 \tau_2})^{-1} = \sigma_1 \tau_2 \sigma_2^{-1} \sigma_1^{-1}, \\
S_{\sigma_2,\tau_1} &= \sigma_2 \tau_1 \cdot (\overline{\sigma_2 \tau_1})^{-1} = \sigma_2 \tau_1 \sigma_1^{-1} \sigma_2^{-1}, \\
S_{\sigma_2,\tau_2} &= \sigma_2 \tau_2, \\
S_{\sigma_1 \sigma_2,\tau_1} &= \sigma_1 \sigma_2 \tau_1 \sigma_1^{-1} \sigma_2^{-1} \sigma_1^{-1}, \\
S_{\sigma_1 \sigma_2,\tau_2} &= \sigma_1 \sigma_2 \tau_2 \sigma_1^{-1}, \\
S_{\sigma_2 \sigma_1,\tau_1} &= \sigma_2 \sigma_1 \tau_1 \sigma_2^{-1}, \\
S_{\sigma_2 \sigma_1,\tau_2} &= \sigma_2 \sigma_1 \tau_2 \sigma_2^{-1} \sigma_1^{-1} \sigma_2^{-1}, \\
S_{\sigma_1 \sigma_2 \sigma_1,\tau_1} &= \sigma_1 \sigma_2 \sigma_1 \tau_1 \sigma_2^{-1} \sigma_1^{-1}, \\
S_{\sigma_1 \sigma_2 \sigma_1,\tau_2} &= \sigma_1 \sigma_2 \sigma_1 \tau_2 \sigma_2^{-1} \sigma_1^{-1} \sigma_2^{-1} \sigma_1^{-1}.
\end{aligned}$$

Since, T_3 is free, all of its subgroups are also free. Hence, we have the decomposition for UP_3 . The group UP_2 is its subgroup and it is generated by

$$a_{12} = \sigma_1^2, S_{1,\tau_1} = \tau_1 \sigma_1^{-1}, S_{\sigma_1,\tau_1} = \sigma_1 \tau_1.$$

We do not have relations between them. Hence, UP_2 is free of rank 3. \square

Corollary 5.1. *The following properties hold:*

- 1) UB_3 is linear;
- 2) UP_3 is residually nilpotent.

Proof. 1) Follows from the fact that the free product of two linear groups is linear.

2) Follows from the fact that the free product of two residually nilpotent groups that do not have torsion is residually nilpotent. \square

Remark 1. *If $n > 3$, then we can not say that UB_n is the free product of B_n and T_n . In this case we have mixed relations of the form*

$$[\sigma_i, \tau_j] = 1, \quad |i - j| > 1.$$

Also, if $n > 3$ we can not say that UP_n is the free product of P_n and a free group. In this case we have relations of the form

$$[\tau_i^2, \tau_j^2] = 1, \quad |i - j| > 1.$$

Question 5.1. *For $n > 3$ find a set of generators and defining relations of the group UP_n .*

ACKNOWLEDGEMENTS

The results were obtained during the program on knots and braid groups advised by V. Bardakov, T. Nasybullov and A. Vesnin in the frame of The First Workshop at the Mathematical Center in Akademgorodok supported by the Ministry of Science and Higher Education of the Russian Federation (agreement no. 075–2019–1675).

REFERENCES

- [1] M. Albar, D. Johnson, The centre of the circular braid group, *Math. Japonica*, V. 30, 1985, 641–645.
- [2] B. Audoux, P. Bellingeri, J.-B. Meilhan, E. Wagner, Extensions of some classical local moves on knot diagrams, *Michigan Math. J.*, V. 67, N. 3, 2018, 647–672.
- [3] J. C. Baez, Link invariants of finite type and perturbation theory, *Lett. Math. Phys.*, V. 26, N. 1, 1992, 43–51.
- [4] V. Bardakov, The virtual and universal braids, *Fund. Math.*, V. 184, 2004, 1–18.
- [5] V. Bardakov, Structure of a conjugating automorphism groups, *Algebra and Logic*, V. 42, N. 3, 2003, 287–303.
- [6] V. Bardakov, P. Bellingeri, Combinatorial properties of virtual braids, *Topology Appl.*, V. 156, N. 6, 2009, 1071–1082.
- [7] V. Bardakov, P. Bellingeri, C. Damiani, Unrestricted virtual braids, fused links and other quotients of virtual braid groups, *J. Knot Theory Ramifications*, V. 24, N. 12, 2015, 1550063.
- [8] V. Bardakov, B. Chuzhinov, I. Emel'yanenkov, M. Ivanov, E. Markhinina, T. Nasybullov, S. Panov, N. Singh, S. Vasyutkin, V. Yakhin, A. Vesnin, Representations of flat virtual braids which do not preserve the forbidden relations, *ArXiv:Math/2010.03162*.
- [9] V. Bardakov, R. Mikhailov, V. Vershinin, J. Wu, On the pure virtual braid group PV_3 , *Commun. in Algebra*, V. 44, N. 3, 2016, 1350–1378.
- [10] V. Bardakov, Yu. Mikhailchishina, M. Neshchadim, Representations of virtual braids by automorphisms and virtual knot groups, *J. Knot Theory Ramifications*, V. 26, N. 1, 2017, 1750003.
- [11] V. Bardakov, Yu. Mikhailchishina, M. Neshchadim, Virtual link groups, *Sib. Math. J.*, V. 58, N. 5, 2017, 765–777.
- [12] V. Bardakov, T. Nasybullov, Multi-switches and representations of braid groups, *ArXiv:Math/1907.09230*.
- [13] V. Bardakov, T. Nasybullov, Multi-switches and virtual knot invariants, *Topology and App.*, V. 293, 2021, 107552.
- [14] V. Bardakov, T. Nasybullov, Multi-switches, representations of virtual braids, and invariants of virtual links, *Algebra and Logic*, V. 59, N. 4, 2020, 341–345.
- [15] V. Bardakov, M. Neshchadim, On a representation of virtual braids by automorphisms, *Algebra and Logic*, V. 56, N. 5, 2017, 355–361.
- [16] V. Bardakov, M. Singh, A. Vesnin, Structural aspects of twin and pure twin groups, *Geom. Dedicata*, V. 203, 2019, 135–154.
- [17] A. Beridze, P. Traczyk, Burau representation for $n = 4$, *J. Knot Theory Ramifications*, V. 27, N. 3, 2018, 1840002.
- [18] S. Bigelow, Braid groups are linear, *J. Amer. Math. Soc.*, V. 14, N. 2, 2001, 471–486.
- [19] J. Birman, New points of view in knot theory, *Bull. Amer. Math. Soc.*, V. 28, N. 2, 1993, 253–287.
- [20] J. Birman, *Braids, links and mapping class group*, Princeton–Tokyo: Univ. press, 1974.
- [21] H. Boden, E. Dies, A. Gaudreau, A. Gerlings, E. Harper, A. Nicas, Alexander invariants for virtual knots, *J. Knot Theory Ramifications*, V. 24, N. 3, 2015, 1550009.

- [22] E. Brieskorn, K. Saito, Artin-Gruppen und Coxeter-Gruppen, *Invent. Math.*, V. 17, 1972, 245–271.
- [23] S. Dey, K. Gongopadhyay, Commutator subgroups of singular braid groups, *ArXiv:Math/1806.05902*.
- [24] D. J. Collins, Relations among the squares of the generators of the braid group, *Invent. Math.*, V. 117, 1994, 525–529.
- [25] R. Fenn, R. Rimányi, C. Rourke, The braid-permutation group, *Topology*, V. 36, N. 1, 1997, 123–135.
- [26] R. Fenn, E. Keyman, C. Rourke, The singular braid monoid embeds in a group, *J. Knot Theory Ramifications*, V. 7, N. 7, 1998, 881–892.
- [27] B. Gassner, On braid groups, *Abh. Math. Sem. Univ. Hamburg*, V. 25, 1961, 10–22.
- [28] D. Goncalves, J. Guaschi, O. Ocampo, A quotient of the Artin braid groups related to crystallographic groups, *J. Algebra*, V. 474, 2017, 393–423.
- [29] M. Goussarov, M. Polyak, O. Viro, Finite-type invariants of classical and virtual knots, *Topology*, V. 39, N. 5, 2000, 1045–1068.
- [30] M. Ivanov, A. Vesnin, F -polynomials of tabulated virtual knots, *J. Knot Theory Ramifications*, V. 29, N. 8, 2020, 2050054.
- [31] S. Kamada, Braid presentation of virtual knots and welded knots, *Osaka J. Math.*, V. 44, 2007, 441–458.
- [32] C. Kassel, V. Turaev, *Braid groups*, Springer, Berlin, 2008.
- [33] L. Kauffman, Virtual knot theory, *Eur. J. Comb.*, V. 20, N. 7, 1999, 663–690.
- [34] L. Kauffman, A survey of virtual knot theory, *Knots in Hellas '98 (Delphi)*, 143–202, *Ser. Knots Everything*, 24, World Sci. Publ., River Edge, NJ, 2000.
- [35] L. Kauffman, S. Lambropoulou, The L-Move and Virtual Braids, *J. Knot Theory Ramifications*, V. 15, N. 6, 2006, 773–811.
- [36] K. Kaur, M. Prabhakar, A. Vesnin, Two-variable polynomial invariants of virtual knots arising from flat virtual knot invariants, *J. Knot Theory Ramifications*, V. 27, N. 13, 2018, 1842015.
- [37] D. Krammer, Braid groups are linear, *Annals of Math.*, V. 155, N. 1, 2002, 131–156.
- [38] V. Manturov, H. Wang, Markov theorem for free links, *J. Knot Theory Ramifications*, V. 21, N. 13, 2012, 1240010.
- [39] A. Markoff, Foundations of the algebraic theory of braids, *Trudy Mat. Inst. Steklova*, V. 16, 1945, 1–54.
- [40] J. McCool, On basis-conjugating automorphisms of free groups, *Can. J. Math.*, V. 38, N. 6, 1986, 1525–1529.
- [41] T. Nasybullov, The classification of fused links, *J. Knot Theory Ramifications*, V. 25, N. 21, 2016, 1650076.
- [42] F. Panaite, M. Staic, A quotient of the braid group related to pseudosymmetric braided categories, *Pacific J. Math.*, V. 144, 2010, 155–167.
- [43] L. Rabenda, Traces de Markov sur les algèbres de Hecke singulières, mémoire de DEA, Université de Bourgogne (2007).
- [44] A. Savushkina, On the group of conjugating automorphisms of a free group, *Math. Notes*, V. 60, N. 1, 1996, 68–80.
- [45] V. Shpilrain, Representing braids by automorphisms, *Int. J. Algebra and Comput.*, V. 11, N. 6, 2001, 773–777.
- [46] D. Silver, S. Williams, Alexander groups and virtual links, *J. Knot Theory Ramifications*, V. 10, N. 1, 2001, 151–160.
- [47] C. Squier, On certain 3-generator Artin groups, *Trans. Amer. Math. Soc.*, V. 302, N. 1, 1987, 117–124.
- [48] V. Turaev, Virtual strings, *Ann. Inst. Fourier (Grenoble)*, V. 54, N. 7, 2004, 2455–2525.
- [49] V. Turaev, Topology of words, *Proc. Lond. Math. Soc.*, V. 95, N. 2, 2007, 360–412.

- [50] V. Vershinin, On homology of virtual braids and Burau representation, *J. Knot Theory Ramifications*, V. 10, N. 5, 2001, 795–812.
- [51] J. Wolf, *Spaces of constant curvature*, sixth edition, AMS Chelsea Publishing, V. 37, 2011.

Valeriy Bardakov^{*,†,‡} (bardakov@math.nsc.ru),
 Ivan Emel'yanenkov^{*} (i.emelianenkov@g.nsu.ru),
 Maxim Ivanov^{*} (m.ivanov2@g.nsu.ru),
 Tatyana Kozlovskaya[†] (t.kozlovskaya@math.tsu.ru),
 Timur Nasybullov^{*,†,‡} (timur.nasybullov@mail.ru),
 Andrei Vesnin^{*,†,‡} (vesnin@math.nsc.ru)

^{*} Novosibirsk State University, Pirogova 1, 630090 Novosibirsk, Russia,

[†] Tomsk State University, Lenin Ave. 36, 634050 Tomsk, Russia,

[‡] Sobolev Institute of Mathematics, Acad. Koptyug Ave. 4, 630090 Novosibirsk, Russia,