

Monads on Categories of Relational Structures

Chase Ford¹, Stefan Milius², and Lutz Schröder

Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany

Abstract

We introduce a framework for universal algebra in categories of relational structures given by finitary relational signatures and finitary or infinitary Horn theories, with the arity λ of a Horn theory understood as a strict upper bound on the number of premisses in its axioms; key examples include partial orders ($\lambda = \omega$) or metric spaces ($\lambda = \omega_1$). We establish a bijective correspondence between λ -accessible enriched monads on the given category of relational structures and a notion of λ -ary *algebraic theories* (i.e. with operations of arity $< \lambda$), with the syntax of algebraic theories induced by the relational signature (e.g. inequations or equations-up-to- ϵ). We provide a generic sound and complete derivation system for such *relational algebraic theories*, thus in particular recovering (extensions of) recent systems of this type for monads on partial orders and metric spaces by instantiation. In particular, we present an ω_1 -ary algebraic theory of metric completion. The theory-to-monad direction of our correspondence remains true for the case of κ -ary algebraic theories and κ -accessible monads for $\kappa < \lambda$, e.g. for finitary theories over metric spaces.

1 Introduction

Monads play an established role in the semantics of sequential and concurrent programming [20] – they encapsulate side-effects, such as statefulness, nontermination, nondeterminism, or probabilistic branching. The well-known correspondence between monads on the category of sets and algebraic theories [15] impacts accordingly on programming syntax, as witnessed, for example, in work on algebraic effects [22]: Operations of the theory serve as syntax for computational effects such as non-deterministic or probabilistic choice. The comparative analysis of programs or systems beyond two-valued equivalence checking, e.g. under behavioural preorders, such as similarity, or behavioural distances, involves monads based on categories beyond sets, such as the categories **Pos** of partial orders or **Met** of (1-bounded) metric spaces. This has sparked recent interest in presentations of such monads using suitable variants of the notion of algebraic theory. While it is, in principle, possible to work with equational presentations that encapsulate the additional structure within the signature [12], it seems at least equally natural to represent the additional structure (e.g. distance or ordering) within the judgements of the theory. Indeed, Mardare, Power, and Plotkin replace equations with equations-up-to- ϵ in their *quantitative algebraic theories* [19], which present monads on **Met**, and in our own previous work on behavioural preorders [8] as well as in our own recent work with Adámek [6], we have used *inequational theories* to present monads on **Pos**.

In the present paper, we introduce a generalized approach to such notions of algebraic theory: We work in categories of finitary relational structures (more precisely, the objects are sets interpreting a given signature of finitary relation symbols), axiomatized by Horn theories whose axioms are implications with possibly infinite sets of antecedents. We say that such a theory is λ -ary for a regular cardinal λ if all its axioms have less than λ antecedents. For instance, **Pos** can be presented by a finitary (i.e. ω -ary) Horn theory over a binary relation \leq , and **Met** by an ω_1 -ary Horn theory over binary relations $=_\epsilon$ ‘equality up to ϵ ’

¹ Chase Ford acknowledges support by the Deutsche Forschungsgemeinschaft (DFG) as part of the Research and Training Group 2475 “Cybercrime and Forensic Computing” (393541319/GRK2475/1-2019)

² Stefan Milius acknowledges by the Deutsche Forschungsgemeinschaft (DFG) under projects MI 717/5-2 and MI 717/7-1

indexed over rational numbers ϵ . We exploit that the models of a λ -ary Horn theory form a locally λ -presentable category \mathcal{C} [5] to give a syntactic characterization of λ -accessible monads on \mathcal{C} in terms of a notion of relational algebraic theory, in the sense that we prove a monad-theory correspondence. Following Kelly and Power [12], we use λ -presentable objects of \mathcal{C} as arities and as contexts of axioms; however, as indicated above, we provide a syntax by expressing axioms using the relational signature instead of necessarily using only equality. We give a sound and complete deduction system for the arising *relational logic* (which generalizes standard equational logic), thus obtaining an explicit description of the monad generated by a relational algebraic theory in the indicated sense. One consequence of our main result is that quantitative algebraic theories [19] induce ω_1 -accessible monads. More generally, presentations of ω_1 -presentable monads in our formalism may involve operations with countable non-discrete arities: indeed, we present an ω_1 -ary relational algebraic theory that defines the metric completion monad. We also take a glimpse at the more involved setting of κ -accessible monads on \mathcal{C} where $\kappa < \lambda$ (e.g. finitary monads on **Met**). We give a partial characterization of κ -presentable objects in this setting, and show that while the monad-to-theory direction of our correspondence fails for $\kappa < \lambda$, the theory-to-monad direction does hold. This implies that some salient quantitative algebraic theories induce finitary monads; e.g. the theory of quantitative join-semilattices [19].

Related Work We have already mentioned work by Kelly and Power on finitary monads [12] and by Mardare et al. on quantitative algebraic theories [19], as well as our own previous work [8] and our joint work with Adámek [6].

Power and Nishizawa [21] have extended the approach of Kelly and Power to deal with different enrichments of a category and the monads thereon, and obtain a correspondence between enriched Lawvere theories [23] and finitary enriched monads. More recently, Power and Garner [24] have provided a more thorough understanding of the equivalence between enriched finitary monads and enriched Lawvere theories as an instance of a free completion of an enriched category under a class of absolute colimits. Rosický [25] establishes a monad-theory correspondence for λ -accessible enriched monads and a notion of λ -ary enriched theory *à la* Linton [14], where arities of operations are given by pairs of objects. Like in the setting of Kelly and Power, relations (inequations, distances) are encoded in the arities. So the syntactic notion of theory is different from (and more abstract than) ours. Lucyshyn-Wright [16] establishes a rather general correspondence between monads and abstract theories in symmetric monoidal closed categories, parametric in a choice of arities, which covers several notions of theory and correspondences in the categorical literature under one roof.

Kurz and Velebil [13] characterize classical ordered varieties [7] (which are phrased in terms of inequalities) as precisely the exact categories in an enriched sense with a ‘suitable’ generator. In recent subsequent work, Adámek et al. [2] establish a correspondence of these varieties with enriched monads on **Pos** that are strongly finitary [11], i.e. their underlying functor is a left Kan-extension of the embedding of finite discrete posets into **Pos**.

The main distinguishing feature of our work relative to the above is the explicit syntactic description of the monad obtained from a theory via a sound and complete derivation system.

2 Preliminaries

We review the basic theory of locally presentable categories (see [5] for more detail) and of monads. We assume a modest familiarity with the elementary concepts of category theory [3] and with ordinal and cardinal numbers [9]. We write $\text{card } X$ for the cardinality of a set X

and, where κ is a cardinal, we write $X' \subseteq_{\kappa} X$ to indicate that $X' \subseteq X$ and $\text{card } X < \kappa$.

Locally Presentable Categories Fix a regular cardinal λ (i.e. an infinite cardinal which is not cofinal to any smaller cardinal). A poset (I, \leq) is λ -directed if each subset $I_0 \subseteq_{\lambda} I$ has an upper bound: there exists $u \in I$ such that $i \leq u$ for all $i \in I_0$. A λ -directed diagram is a functor whose domain is a λ -directed poset (viewed as a category); colimits of such diagrams are also called λ -directed. An object X in a category \mathcal{C} is λ -presentable if the covariant hom-functor $\mathcal{C}(X, -)$ preserves λ -directed colimits. That is, X is λ -presentable if for each directed colimit $(D_i \xrightarrow{c_i} C)_{i \in I}$ in \mathcal{C} , every morphism $m: X \rightarrow C$ factors through one of the c_i essentially uniquely: there exists $i \in I$ and $g: X \rightarrow D_i$ such that $m = c_i \cdot g$, and for all $g': X \rightarrow D_i$ such that $m = c_i \cdot g'$, there exists $j \geq i$ such that $D(i \rightarrow j) \cdot g = D(i \rightarrow j) \cdot g'$.

► **Definition 2.1.** A category \mathcal{C} is *locally λ -presentable* if it is cocomplete, its full subcategory $\text{Pres}_{\lambda}(\mathcal{C})$ given by the λ -presentable objects of \mathcal{C} is essentially small, and every $C \in \mathcal{C}$ is a λ -directed colimit of objects in $\text{Pres}_{\lambda}(\mathcal{C})$. When $\lambda = \omega$ (resp. ω_1), we speak of *locally finitely* (resp. *countably*) *presentable categories*. We call \mathcal{C} *locally presentable* if it is locally λ -presentable for some cardinal λ . A functor F on a locally presentable category is λ -accessible if it preserves λ -directed colimits. When $\lambda = \omega$ or ω_1 , we speak of *finitary* and *countably accessible functors*, respectively.

Reflective subcategories A full subcategory \mathcal{C}' of a category \mathcal{C} is *reflective* if the embedding $\iota: \mathcal{C}' \hookrightarrow \mathcal{C}$ is a right adjoint. In this case, we write $r_X: X \rightarrow RX$ (or just r if X is clear from the context) for the universal arrows; we call RX *the reflection of $X \in \mathcal{C}$* , r_X *the reflective arrow*, and the left adjoint R *the reflector*. The universal property of $r_X: X \rightarrow RX$ is as follows: For each morphism $f: X \rightarrow Y$ in \mathcal{C} where Y lies in \mathcal{C}' , there exists a unique morphism $f^{\sharp}: RX \rightarrow Y$ such that $f = f^{\sharp} \cdot r_X$. We call \mathcal{C}' *epi-reflective* if r_X is epi for all $X \in \mathcal{C}$. We will employ the following reflection theorem:

► **Theorem 2.2** [5, Cor. 2.48]. *If \mathcal{C}' is a full subcategory of a locally λ -presentable category \mathcal{C} and \mathcal{C}' is closed under limits and λ -directed colimits in \mathcal{C} , then \mathcal{C}' is reflective and locally λ -presentable.*

Monads A *monad* on a category \mathcal{C} is a functor $T: \mathcal{C} \rightarrow \mathcal{C}$ equipped with natural transformations $\eta: \text{Id} \rightarrow T$ (the *unit*) and $\mu: TT \rightarrow T$ (the *multiplication*) such that the diagrams below commute.

$$\begin{array}{ccc}
 T & \xrightarrow{T\eta} & TT & \xleftarrow{\eta T} & T \\
 & \searrow \text{id} & \downarrow \mu & \swarrow \text{id} & \\
 & & T & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 TTT & \xrightarrow{T\mu} & TT \\
 \mu T \downarrow & & \downarrow \mu \\
 TT & \xrightarrow{\mu} & T
 \end{array}$$

We call the monad (T, η, μ) λ -accessible if its underlying functor is λ -accessible.

► **Definition 2.3.** An *Eilenberg-Moore algebra* for the monad (T, η, μ) is a \mathcal{C} -morphism of the shape $a: TX \rightarrow X$ satisfying the following coherence laws:

$$\begin{array}{ccc}
 X & \xrightarrow{\eta} & TX \\
 & \searrow \text{id} & \downarrow a \\
 & & X
 \end{array}
 \qquad
 \begin{array}{ccc}
 TTX & \xrightarrow{\mu} & TX \\
 Ta \downarrow & & \downarrow a \\
 TX & \xrightarrow{a} & X
 \end{array}$$

A *homomorphism* from $a: TX \rightarrow X$ to an Eilenberg-Moore algebra $b: TY \rightarrow Y$ is a morphism $h: X \rightarrow Y$ in \mathcal{C} such that $h \cdot a = Th \cdot b$.

► **Notation 2.4.** For a functor $F: \mathcal{C} \rightarrow \mathcal{C}$, we write $\mathbf{Alg} F$ for the category of F -algebras and homomorphisms, i.e. $\mathbf{Alg} F$ has \mathcal{C} -morphisms of the shape $a: FA \rightarrow A$ as objects, and a homomorphism $(A, a) \rightarrow (B, b)$ is a \mathcal{C} -morphism $h: A \rightarrow B$ such that $h \cdot a = b \cdot Fh$.

3 Categories of Relational Structures

As indicated previously, we will study monads over base categories consisting of (single-sorted) relational structures. Specifically, we will restrict the relational signature to be finitary but allow infinitary Horn axioms. We proceed to recall basic definitions, examples, and results, in particular on closed structure and (local) presentability. In Section 3.1, we present new results on the partial characterization of (internally) λ -presentable objects in cases where the overall local presentability index of the category is greater than λ .

► **Definition 3.1.** (1) A *relational signature* is a set Π of *relation symbols* α, β, \dots together with a finite *arity* $0 < \text{ar}(\alpha) \in \omega$ for all $\alpha \in \Pi$. A Π -*edge* in a set S is a pair $e = \alpha(f)$ where $\alpha \in \Pi$ and $f: \text{ar}(\alpha) \rightarrow S$ is a function. For a map $g: S \rightarrow Y$, we write $g \cdot e = \alpha(g \cdot f)$. We extend this notation pointwise to sets E of edges: $g \cdot E = \{g \cdot e \mid e \in E\}$.

(2) A Π -*structure* X consists of an underlying set $|X|$ (or just X when no confusion is likely) and a set $\mathbf{E}(X)$ of Π -edges in $|X|$. If $\alpha(f) \in \mathbf{E}(X)$, we write $\alpha_X(f)$ or even $X \models \alpha(f)$.

(3) A *relation-preserving map* or briefly a *morphism* from X to a Π -structure Y is a function $g: |X| \rightarrow |Y|$ such that $g \cdot \mathbf{E}(X) \subseteq \mathbf{E}(Y)$. If g additionally is injective and *relation-reflecting*, i.e. whenever $g \cdot e \in \mathbf{E}(Y)$ for an edge e , then $e \in \mathbf{E}(X)$, then g is an *embedding*. We denote by $\mathbf{Str}(\Pi)$ the category of Π -structures and relation-preserving maps.

► **Notation 3.2.** Given an edge $\alpha(f)$ such that $f(i) := x_i$ for all $i \in \text{ar}(\alpha)$, we sometimes write $\alpha(x_1, \dots, x_{\text{ar}(\alpha)})$ (or just $\alpha(x_i)$) instead of $\alpha(f)$. We will pass between these presentations without further mention.

We are going to carve out full subcategories of $\mathbf{Str}(\Pi)$ by means of infinitary Horn axioms, whose **syntax** we recall next.

► **Definition 3.3.** Let Π be a relational signature, and λ a regular cardinal. We fix a set \mathbf{Var} of variables such that $\text{card}(\mathbf{Var}) = \lambda$. A λ -*ary Horn formula* over Π has the form

$$\Phi \implies \psi$$

where Φ is a set of Π -edges in \mathbf{Var} such that $\text{card} \Phi < \lambda$ and ψ is a $\Pi \cup \{=\}$ -edge in \mathbf{Var} , for a fresh binary relation symbol $=$. In case $\Phi = \{\varphi_1, \dots, \varphi_n\}$ is finite, we write $\varphi_1, \dots, \varphi_n \implies \psi$, and if $\Phi = \emptyset$, then we just write $\implies \psi$. A λ -*ary Horn theory* $\mathcal{H} = (\Pi, \mathcal{A})$ consists of a relational signature Π and a set \mathcal{A} of λ -ary Horn formulae over Π , the *axioms* of \mathcal{H} .

We fix a λ -ary Horn theory $\mathcal{H} = (\Pi, \mathcal{A})$ for the rest of the paper. We define the **semantics** of Horn formulae in a Π -structure X as follows. We denote by \bar{X} the $\Pi \sqcup \{=\}$ -structure obtained from X by putting $=_X := \{(x, x) \mid x \in X\}$. A *valuation* is a map $\kappa: \mathbf{Var} \rightarrow |X|$. We say that X *satisfies* a Horn formula $\Phi \implies \psi$ if whenever κ is a valuation such that $X \models \kappa \cdot \phi$ for all $\phi \in \Phi$, then $\bar{X} \models \kappa \cdot \psi$. Finally, X is a *model* of \mathcal{H} , or of \mathcal{A} , if X satisfies all axioms of \mathcal{H} . The full subcategory of $\mathbf{Str}(\Pi)$ spanned by the models of \mathcal{A} is $\mathbf{Str}(\Pi, \mathcal{A})$ (or $\mathbf{Str} \mathcal{H}$).

We have an obvious notion of *derivation* under \mathcal{H} over a given set Z (e.g. of variables or points in a structure): We extend \mathcal{H} to $(\Pi \cup \{=\}, \bar{\mathcal{A}})$ where $\bar{\mathcal{A}}$ consists of the axioms in \mathcal{A} and additional axioms stating that $=$ is an equivalence and that all relations in Π are closed

under $=$ in the obvious sense. Then we have a single (λ -ary) derivation rule for application of Horn axioms $(\Phi \implies \psi) \in \bar{\mathcal{A}}$ over Z :

$$\frac{\kappa \cdot \Phi}{\kappa \cdot \psi} (\kappa: \text{Var} \rightarrow Z).$$

We say that a set E of edges over Z *entails* an edge e over Z (*under \mathcal{H}*) if e is derivable from edges in E in this system. In case $Z = \text{Var}$ and $\text{card } E < \lambda$, the expression $E \implies e$ is in fact a Horn formula, and we then also say that \mathcal{H} *entails* $E \implies e$ if E entails e .

► **Assumption 3.4.** For technical convenience, we assume that the fixed Horn theory $\mathcal{H} = (\Pi, \mathcal{A})$ *expresses equality*. That is, there exists a set $Eq(x, y)$ of Π -edges in variables x, y such that \mathcal{H} entails $Eq(x, y) \implies x = y$ as well as $\implies \psi$ for all edges $\psi \in Eq(x, x)$ (where we use obvious notation for substitution; formally, $Eq(x, x) = g \cdot Eq(x, y)$ where $g(x) = g(y) = x$). Moreover, we assume that \mathcal{A} explicitly includes the (derivable) formulae $Eq(x_1, y_1) \cup \dots \cup Eq(x_{\text{ar}(\alpha)}, y_{\text{ar}(\alpha)}) \cup \{\alpha(x_1, \dots, x_{\text{ar}(\alpha)})\} \implies \alpha(y_1, \dots, y_{\text{ar}(\alpha)})$ saying that all relations $\alpha \in \Pi$ are closed under Eq (implying also that Eq is symmetric and transitive). This is w.l.o.g. as we can always extend a given Horn theory with an equality predicate axiomatized by the above conditions without changing its category of models; indeed we leave this predicate implicit in examples whose natural presentation does not include it.

► **Example 3.5.** We mention some key examples of Horn theories:

- (1) The category **Set** of sets and functions is specified by the trivial Horn theory (\emptyset, \emptyset) .
- (2) The category **Pos** of partially ordered sets (posets) and monotone maps is specified by the ω -ary Horn theory consisting of a single binary relation symbol \leq and the axioms

$$x \leq x; \quad x \leq y, y \leq z \implies x \leq z; \quad \text{and} \quad x \leq y, y \leq x \implies x = y.$$

This theory expresses equality (Assumption 3.4) via $Eq(x, y) = \{x \leq y, y \leq x\}$.

- (3) The theory \mathcal{H}_{Met} of *metric spaces* is the ω_1 -ary theory consisting of binary relation symbols $=_\epsilon$ for all $\epsilon \in \mathbb{Q} \cap [0, 1]$, and the axioms

$$\implies x =_0 x \quad \text{(Ref1)}$$

$$x =_0 y \implies x = y \quad \text{(Equal)}$$

$$x =_\epsilon y \implies y =_\epsilon x \quad \text{(Sym)}$$

$$\{x =_\epsilon y, y =_{\epsilon'} z\} \implies x =_{\epsilon+\epsilon'} z \quad \text{(Triang)}$$

$$x =_\epsilon y \implies x =_{\epsilon+\epsilon'} y \quad \text{(Up)}$$

$$\{x =_{\epsilon'} y \mid \mathbb{Q}_{\geq 0} \ni \epsilon' > \epsilon\} \implies x =_\epsilon y \quad \text{(Arch)}$$

where ϵ, ϵ' range over $\mathbb{Q} \cap [0, 1]$ (that is, the axioms mentioning ϵ, ϵ' are in fact axiom schemes representing one axiom for each ϵ, ϵ'). This theory expresses equality via $Eq(x, y) = \{x =_0 y\}$; in fact, even if we remove $=_0$, the remaining theory still expresses equality via $Eq(x, y) = \{x =_{1/n} y \mid n > 0\}$. The theory \mathcal{H}_{Met} specifies the category **Met** of 1-bounded metric spaces and non-expansive maps, in the sense that $\mathbf{Str}(\mathcal{H}_{\text{Met}})$ and **Met** are concretely isomorphic: $X \in \mathbf{Str}(\mathcal{H}_{\text{Met}})$ induces the 1-bounded metric space (X, d) given by $d(x, y) = \bigwedge \{\epsilon \mid x =_\epsilon y \in \mathbf{E}(X)\}$, and conversely a metric space (X, d) induces the \mathcal{H}_{Met} -model on X with edges $\{x =_\epsilon y \mid x, y \in X, d(x, y) \leq \epsilon\}$.

- (4) Let L be a complete lattice (for simplicity), and let $L_0 \subseteq L$ be meet-dense in L in the sense that $l = \bigwedge \{p \in L_0 \mid p \geq l\}$ for each $l \in L$; whenever $q \geq \bigwedge P$ for $q \in L_0$ and $P \subseteq L_0$

6 Monads on Categories of Relational Structures

such that $\bigwedge P \notin L_0$, then $q \geq p$ for some $p \in P$ (e.g. these conditions hold trivially for $L_0 = L$). Further, fix λ such that $|L_0| < \lambda$. Let \mathcal{H}_L be the λ -ary Horn theory with binary relation symbols α_p for all $p \in L_0$ and axioms

$$\begin{aligned} \{\alpha_p(x, y) \mid p \in P\} &\implies \alpha_q(x, y) && (P \subseteq L_0, q = \bigwedge P \in L_0) && \text{(Arch)} \\ \alpha_p(x, y) &\implies \alpha_q(x, y) && (p, q \in L_0, p \leq q) && \text{(Up)} \end{aligned}$$

where p, q range over L_0 . Then $\mathbf{Str}(\mathcal{H}_L)$ is concretely isomorphic to the category of L -valued relations, whose objects X are sets X equipped with map $P: X \times X \rightarrow L$, and whose morphisms $(X, P) \rightarrow (Y, Q)$ are maps $X \rightarrow Y$ such that $Q(f(x), f(y)) \leq P(x, y)$. (Of course, the previous example is essentially the special case $L = [0, 1]$, $L_0 = \mathbb{Q} \cap [0, 1]$ with some additional axioms.)

(5) A signature of partial operations is a set P of operation symbols f with assigned finite arities $\text{ar}(f)$. A (partial) P -algebra is then a set A and, for each $f \in P$, a partial function $f_A: A^{\text{ar}(f)} \rightarrow A$. A homomorphism of partial algebras is a map $h: A \rightarrow B$ such that whenever $f_A(x_1, \dots, x_{\text{ar}(f)})$ is defined, then $f_B(h(x_1), \dots, h(x_{\text{ar}(f)}))$ is defined and equals $h(f_A(x_1, \dots, x_{\text{ar}(f)}))$. The category of partial P -algebras and their homomorphisms is concretely isomorphic to the category of models of the ω -ary Horn theory consisting of relational symbols α_f of arity $\text{ar}(f) + 1$ for all $f \in P$ (with $\alpha_f(x_1, \dots, x_{\text{ar}(f)}, y)$ being understood as $f(x_1, \dots, x_{\text{ar}(f)}) = y$), and axioms

$$\{\alpha(x_1, \dots, x_{\text{ar}(f)}, y), \alpha(x_1, \dots, x_{\text{ar}(f)}, z)\} \implies y = z.$$

We proceed to recall some key aspects of the categorical structure of $\mathbf{Str}(\mathcal{H})$.

Reflection We first note

► **Proposition 3.6.** $\mathbf{Str}(\Pi, \mathcal{A})$ is a (full) epi-reflective subcategory of $\mathbf{Str}(\Pi)$.

Since $\mathbf{Str}(\Pi)$ is easily seen to be complete and cocomplete, it follows that $\mathbf{Str}(\Pi, \mathcal{A})$ is cocomplete and moreover closed under limits in $\mathbf{Str}(\Pi)$, and hence complete. We write

$$R: \mathbf{Str}(\Pi) \rightarrow \mathbf{Str}(\Pi, \mathcal{A}) \quad \text{and} \quad r_X: X \rightarrow RX$$

for the left adjoint of the inclusion $\mathbf{Str}(\Pi, \mathcal{A}) \hookrightarrow \mathbf{Str}(\Pi)$ (the *reflector*) and the corresponding (surjective) reflection maps, respectively. Explicitly, RX is constructed as follows. We define an equivalence \sim on X by $x \sim y$ if $\mathbf{E}(X)$ entails $x = y$ under \mathcal{H} (in the sense defined above), and let $q: X \rightarrow X/\sim$ denote the quotient map; then RX has underlying set X/\sim , and contains precisely the edges $q \cdot e$ such that $\mathbf{E}(X)$ entails e ; moreover, $r_X = q$ as a map.

Local presentability One easily checks

► **Lemma 3.7.** An object $(X, E) \in \mathbf{Str}(\Pi)$ is λ -presentable iff $\text{card } X < \lambda$ and $\text{card } E < \lambda$; the category $\mathbf{Str}(\Pi)$ is locally finitely presentable.

By Proposition 3.6 and since $\mathbf{Str}(\Pi, \mathcal{A})$ is easily seen to be closed under λ -directed colimits in $\mathbf{Str}(\Pi)$, we thus have

► **Proposition 3.8** [5, Example 5.27(3)]. $\mathbf{Str}(\mathcal{H})$ is locally λ -presentable.

The forgetful functor $\mathbf{Str}(\mathcal{H}) \rightarrow \mathbf{Set}$ preserves λ -directed colimits. Moreover, we have an easy characterization of λ -presentable objects:

► **Proposition 3.9.** For an \mathcal{H} -model X , the following are equivalent.

- (1) X is λ -presentable in $\mathbf{Str}(\Pi, \mathcal{A})$;
- (2) $X \cong R(Y, E)$ for some λ -presentable $(Y, E) \in \mathbf{Str}(\Pi)$;
- (3) $\text{card } |X| < \lambda$, and X is λ -generated, i.e. there exists $E \subseteq \mathbf{E}(X)$ such that $\text{card } E < \lambda$ and E entails every edge in $\mathbf{E}(X)$ under \mathcal{H} (equivalently, Ri is an isomorphism where $i: (|X|, E) \rightarrow X$ is the $\mathbf{Str}(\Pi)$ -morphism carried by id_X).

► **Remark 3.10.** For instance, every finite partial order is ω -presentable, and every countable metric space is ω_1 -presentable. We emphasize that the situation is more complicated for κ -presentable objects where $\kappa < \lambda$; we treat this case in more detail in Section 3.1. For instance, every finite metric space with rational distances (cf. Example 3.5) is finitely generated in the sense of Proposition 3.9 but not finitely presentable.

Closed monoidal structure The pointwise structure defines an *internal hom* functor:

► **Definition 3.11.** The *internal hom* of $X, Y \in \mathbf{Str}(\Pi)$ is the Π -structure $[X, Y]$ carried by $\mathbf{Str}(\Pi)(X, Y)$ with set of edges

$$\mathbf{E}([X, Y]) := \{e \mid \forall x \in X. \pi_x \cdot e \in \mathbf{E}(Y)\}$$

where $\pi_x: \mathbf{Str}(\Pi)(X, Y) \rightarrow Y$ is defined by $\pi_x(g) = g(x)$. For each $X \in \mathbf{Str}(\Pi)$, the assignment $Y \mapsto [X, Y]$ defines a (covariant) *internal hom functor*

$$[X, -]: \mathbf{Str}(\Pi) \rightarrow \mathbf{Str}(\Pi)$$

with the action on a morphism $m: Y \rightarrow Z$ given by post-composition: $[X, m](g) := m \cdot g$.

One easily checks that $[X, -]$ restricts to

$$[X, -]: \mathbf{Str}(\mathcal{H}) \rightarrow \mathbf{Str}(\mathcal{H}).$$

It turns out that the internal hom functor is always part of a closed symmetric monoidal structure on $\mathbf{Str}(\mathcal{H})$. Indeed, for $X, Y \in \mathbf{Str}(\Pi)$, we define $X \otimes Y$ as the structure with underlying set $X \times Y$ and edges

$$\{e \mid (\pi_1 \cdot e \text{ constant} \wedge \pi_2 \cdot e \in \mathbf{E}(Y)) \vee (\pi_2 \cdot e \text{ constant} \wedge \pi_1 \cdot e \in \mathbf{E}(X))\}.$$

where an edge (α, f) is *constant* if f is a constant map, and $\pi_1: X \times Y \rightarrow X$ and $\pi_2: X \times Y \rightarrow Y$ are the projection maps. We then have $\mathbf{Str}(\Pi)$ -morphisms $u_Y: Y \rightarrow [X, Y \otimes X]$, $u_Y(y)(x) = (y, x)$. It is straightforward to check that u_Y is a universal arrow. That is:

► **Proposition 3.12.** For every $X \in \mathbf{Str}(\Pi)$, $(-) \otimes X$ is a left adjoint of $[X, -]$.

► **Corollary 3.13.** For every $X \in \mathbf{Str}(\mathcal{H})$, $R((-) \otimes X)$ is a left adjoint of $[X, -]: \mathbf{Str}(\mathcal{H}) \rightarrow \mathbf{Str}(\mathcal{H})$.

That is, $\mathbf{Str}(\mathcal{H})$ is a closed symmetric monoidal category, with monoidal structure

$$X \otimes_{\mathcal{H}} Y = R(X \otimes Y).$$

We briefly refer to $\otimes_{\mathcal{H}}$ as the *Manhattan product*.

► **Example 3.14.** (1) In \mathbf{Pos} (Example 3.5(2)), the Manhattan product coincides with binary Cartesian product (so \mathbf{Pos} is Cartesian closed).

(2) In **Met** (Example 3.5(3)), the Manhattan product $(X, d_X) \otimes_{\mathcal{H}_{\text{Met}}} (Y, d_Y)$ is $X \times Y$ equipped with the well-known Manhattan metric d given by $d((x_1, y_1), (x_2, y_2)) = \min(d_X(x_1, x_2) + d_Y(y_1, y_2), 1)$ (while Cartesian products carry the supremum metric).

► **Definition 3.15.** A functor $F: \mathbf{Str}(\mathcal{H}) \rightarrow \mathbf{Str}(\mathcal{H})$ that preserves the pointwise structure on morphisms is called *enriched*. That is, we call F enriched if for all $X, Y \in \mathbf{Str}(\mathcal{H})$ and all edges $f: \text{ar}(\alpha) \rightarrow \mathbf{Str}(\mathcal{H})(X, Y)$ ($\alpha \in \Pi$), if $[X, Y] \models \alpha(f_i)$, then $[FX, FY] \models \alpha(F(f_i))$.

Internal local presentability For use of objects X as arities of operations, we will in fact need that the *internal* hom $[X, -]$ is λ -accessible, in which case we say that X is *internally λ -presentable*. For distinction, we sometimes say that X is *externally λ -presentable* if X is λ -presentable in the standard sense. Indeed, one can show using Proposition 3.9 that the class of λ -presentable objects in $\mathbf{Str}(\mathcal{H})$ is closed under the Manhattan tensor; it follows fairly straightforwardly that

► **Proposition 3.16.** *Every λ -presentable object in $\mathbf{Str}(\mathcal{H})$ is internally λ -presentable.*

Indeed this implies that $\mathbf{Str}(\mathcal{H})$ is *internally locally λ -presentable* [10].

3.1 Compact Horn Models

We have seen above that the category $\mathbf{Str}(\mathcal{H})$ (where \mathcal{H} is a λ -ary Horn theory) is (internally) locally λ -presentable, with a straightforward characterization of the (internally) λ -presentable objects (Propositions 3.9 and 3.16). We proceed to look at the rather less straightforward notion of internally κ -presentable objects in $\mathbf{Str}(\Pi, \mathcal{A})$ for $\kappa < \lambda$. The main scenario that motivates our interest in this case is that of finitary monads on categories that are internally locally λ -presentable only for some $\lambda > \omega$, such as metric spaces.

Further unfolding definitions. we have that an object X is internally κ -presentable if for every κ -directed colimit $(D_i \xrightarrow{c_i} C)_{i \in I}$, the canonical morphism

$$\text{colim}[X, D_i] \rightarrow [X, \text{colim } D_i]$$

is an isomorphism. We split this property into two parts: We say that X is *weakly κ -presentable* if the canonical morphism is always surjective, and *co-weakly κ -presentable* if the canonical morphism is always an embedding. Below, we give necessary and sufficient conditions for weak κ -presentability. Co-weak κ -presentability is a more elusive property; more concretely, it means roughly that X -indexed tuples of derivations in the given Horn theory can be synchronized into single derivations over X -indexed tuples of points. We give some examples below (Example 3.17).

► **Example 3.17.** We give some examples and non-examples of internally finitely presentable objects in locally ω_1 -presentable categories $\mathbf{Str}(\Pi, \mathcal{A})$.

(1) A metric space is internally finitely presentable iff it is finite and discrete. The ‘if’ direction has surprisingly complicated reasons: It holds only because over the reals, finite joins distribute over directed infima. On the other hand, no non-empty metric space is *externally* finitely presentable, as its hom-functor will fail to preserve the colimit of the directed chain $(D_i)_{i < \omega}$ of spaces D_i with underlying set $\{0, 1\}$ and metric $d(0, 1) = 1/(i + 1)$.

(2) In the category of L -valued relations for a complete lattice L in which binary joins fail to distribute over directed infima (such lattices exist), the two-element discrete space fails to be internally finitely presentable.

(3) Let L be as in the previous item, and assume additionally that there is $l \in L$ such that in the downset of l , finite joins do distribute over directed infima (again, such L exist). Take the Horn theory of L -valued relations, extended with an additional (two-valued) relation α and axioms

$$\alpha(x, y) \wedge \alpha(x', y') \implies x =_l x' \quad \alpha(x, y) \wedge \alpha(x', y') \implies y =_l y'.$$

Then the set $\{0, 1\}$ equipped with the discrete L -valued relation and $\alpha(0, 1)$ is internally finitely presentable.

We proceed to give the announced characterization of weakly finitely presentable objects.

► **Definition 3.18.** A *cover* (Y, E) , or just E , of $X \in \mathbf{Str}(\Pi, \mathcal{A})$ is a set E of edges in some set $Y \supseteq |X|$ such that all edges of X are implied by those in E under the Horn theory \mathcal{A} . That is, the underlying map $r_{(Y, E)}: Y \rightarrow |R(Y, E)|$ of the reflection composes with the inclusion $i: |X| \hookrightarrow Y$ to yield a morphism $r_{(Y, E)} \cdot i: X \rightarrow R(Y, E)$ (in $\mathbf{Str}(\Pi, \mathcal{A})$). Then X is κ -compact if for each cover (Y, E) of X there exist $E' \subseteq_\kappa E$ and a morphism $f: X \rightarrow R(Y, E')$ such that $r_{(Y, E)} \cdot i = Rj \cdot f$ where $j: (Y, E') \rightarrow (Y, E)$ is the $\mathbf{Str}(\Pi)$ -morphism carried by id_Y :

$$\begin{array}{ccc} & R(Y, E') & \\ & \nearrow f & \\ X & \xrightarrow{r_{(Y, E)} \cdot i} & R(Y, E) \\ & & \downarrow Rj \end{array} \quad (3.1)$$

► **Lemma 3.19.** *Every κ -compact object is κ -generated.*

► **Remark 3.20.** We will show that the weakly finitely presentable objects in $\mathbf{Str}(\Pi, \mathcal{A})$ are precisely the κ -compact objects with less than κ elements (Proposition 3.21). This characterization breaks under seemingly innocuous variations of the definition of κ -compactness:

(1) It is essential that the edges of a cover live over a superset Y of X . If we were to restrict covers to consist of edges over X (call such a cover an X -cover), then finite ω -compact objects in the arising relaxed sense would in general fail to be finitely presentable. E.g. take (Π, \mathcal{A}) to be the theory of metric spaces additionally equipped with a transitive relation α . Then the set $X = \{0, 2\}$ equipped with the discrete metric and the edge $\alpha(0, 2)$ satisfies the relaxed definition of compactness (every X -cover must contain the edge $\alpha(0, 2)$) but fails to be weakly finitely presentable: The colimit of the ω -chain of objects D_i with underlying set $\{0, 1, 1', 2\}$, distances $d(0, 1) = d(1', 2) = 1$, $d(1, 1') = 1/i$, and edges $\alpha(0, 1)$ and $\alpha(1', 2)$ is not weakly preserved by the hom-functor $\mathbf{Str}(\Pi, \mathcal{A})(X, -)$ (the obvious inclusion of X into the colimit fails to factorize through any of the D_i).

(2) Note that we do not require that the factorization f of $r_{(Y, E)} \cdot i$ in (3.1) equals $r_{(Y, E')} \cdot i$; i.e. f may rename elements of X into elements of Y that lie outside X . Let us refer to the natural-sounding strengthening of κ -compactness where we do require $f = r_{(Y, E')} \cdot i$ as *strong κ -compactness*; e.g. X is strongly ω -compact if every cover of X has a finite subcover. However, this notion is too strong, i.e. not every (weakly) finitely presentable object in $\mathbf{Str}(\Pi, \mathcal{A})$ is strongly ω -compact. As a counterexample, consider the same Horn theory as in the previous item but without the transitivity axiom for α . Then the same object X as in the previous item is weakly finitely presentable (even internally finitely presentable) but not strongly ω -compact, as witnessed by the cover $E = \{\alpha(0', 2)\} \cup \{0 =_{1/n} 0' \mid n > 0\}$.

► **Proposition 3.21.** *The following are equivalent for $X \in \mathbf{Str}(\Pi, \mathcal{A})$:*

- (1) X is weakly κ -presentable;
- (2) X is κ -compact, and $\text{card } |X| < \kappa$.

4 Relational Algebraic Theories

We next describe a framework of universal algebra for enriched κ -accessible monads on the internally locally λ -presentable category $\mathbf{C} = \mathbf{Str}(\mathcal{H})$ of \mathcal{H} -models, for $\kappa \leq \lambda$. We establish one direction of our theory-monad correspondence: We show that every theory in our framework induces a κ -accessible monad (Remark 4.12) whose algebras are precisely the models of the theory (Theorem 4.13). We address the converse direction in Section 5. We write \mathbf{C}_0 for the ordinary category underlying the closed monoidal category \mathbf{C} .

Following Kelly and Power [12], we use the internally λ -presentable objects in \mathbf{C} as the arities of operation symbols. The full subcategory $\mathbf{Pres}_\lambda(\mathbf{C})$ of internally λ -presentable objects is essentially small (Proposition 3.16); we fix a small subcategory \mathcal{P}_λ of internally λ -presentable \mathbf{C} -objects representing all such objects up to isomorphism. For all infinite $\kappa < \lambda$, the full subcategory $\mathcal{P}_\kappa \hookrightarrow \mathcal{P}_\lambda$ is given by the internally κ -presentable objects in \mathcal{P}_λ .

► **Definition 4.1.** Let $\kappa \leq \lambda$ be a regular cardinal. A κ -ary signature is a set Σ of operation symbols σ , each of which is equipped with an arity $\mathbf{ar}(\sigma) \in \mathcal{P}_\kappa$.

A Σ -algebra A consists of a \mathbf{C} -object, also denoted A , and a family of \mathbf{C} -morphisms

$$\sigma_A : [\mathbf{ar}(\sigma), A] \rightarrow A \quad (\sigma \in \Sigma)$$

A homomorphism from A to a Σ -algebra B is a morphism $h : A \rightarrow B$ in \mathbf{C} such that the diagram below commutes for all $\sigma \in \Sigma$.

$$\begin{array}{ccc} [\mathbf{ar}(\sigma), A] & \xrightarrow{\sigma_A} & A \\ h \cdot (-) \downarrow & & \downarrow h \\ [\mathbf{ar}(\sigma), B] & \xrightarrow{\sigma_B} & B \end{array}$$

We write $\mathbf{Alg} \Sigma$ for the category of Σ -algebras and homomorphisms. By a subalgebra of the Σ -algebra A , we understand a Σ -algebra B equipped with a homomorphism $h : B \hookrightarrow A$ whose underlying \mathbf{C} -morphism is an embedding.

Signatures and their algebras Fix a κ -ary signature Σ for the remainder of this section. The category $\mathbf{Alg} \Sigma$ can be presented as a category of functor algebras:

► **Definition 4.2.** The signature functor associated to Σ , $H_\Sigma : \mathbf{C} \rightarrow \mathbf{C}$, is given by

$$H_\Sigma = \coprod_{\sigma \in \Sigma} [\mathbf{ar}(\sigma), -].$$

The categories $\mathbf{Alg} \Sigma$ and $\mathbf{Alg} H_\Sigma$ are clearly isomorphic as concrete categories over \mathbf{C} , so the forgetful functor $\mathbf{Alg} \Sigma \rightarrow \mathbf{C}_0$ inherits all properties of the forgetful functor $\mathbf{Alg} H_\Sigma \rightarrow \mathbf{C}_0$. We collect a few basic consequences of this observation:

► **Remark 4.3.** (1) In general, the forgetful functor $\mathcal{U} : \mathbf{Alg} F \rightarrow \mathcal{C}$ from the category $\mathbf{Alg} F$ of F -coalgebras for a functor F on a category \mathcal{C} creates all limits in \mathcal{C} . It follows that $\mathbf{Alg} \Sigma$ has all limits, and the forgetful functor $\mathbf{Alg} \Sigma \rightarrow \mathbf{C}_0$ creates them.

(2) Since H_Σ is a colimit of κ -accessible functors $[\mathbf{ar}(\sigma), -]$, it is itself κ -accessible, so that the forgetful functor $\mathbf{Alg} H_\Sigma \rightarrow \mathbf{C}_0$ creates κ -directed colimits, and the same holds for the forgetful functor $\mathbf{Alg} \Sigma \rightarrow \mathbf{C}_0$.

(3) From the previous observation (which implies that H_Σ is also λ -accessible) and Proposition 3.8, we obtain by [5, Remark 2.75] (for λ -accessible functors F on locally λ -presentable categories, $\text{Alg } F$ is locally λ -presentable) that $\text{Alg } \Sigma$ is locally λ -presentable.

(4) Adámek [1] shows that for a λ -accessible functor F on a cocomplete category \mathcal{C} , the forgetful functor $\text{Alg } F \rightarrow \mathcal{C}$ is right adjoint. From (2) and cocompleteness of \mathbf{C}_0 (Section 3), we thus obtain that the forgetful functor $\text{Alg } \Sigma \rightarrow \mathbf{C}_0$ is right adjoint; that is, every object $X \in \mathbf{C}$ generates a free Σ -algebra $F_\Sigma X$.

Varieties of Σ -Algebras We now describe a syntax for specifying full subcategories of $\text{Alg } \Sigma$. As a first step, we introduce a notion of Σ -term, defined as usual in universal algebra:

► **Definition 4.4** (Σ -Terms; substitution). For $X \in \mathbf{C}$, we call its underlying set $|X|$ the set of *variables* in X . The set $T_\Sigma(X)$ of Σ -terms in X is defined inductively as follows:

- (1) Each variable in $|X|$ is a Σ -term in X ;
- (2) For each $\sigma \in \Sigma$ and each map $f: |\text{ar}(\sigma)| \rightarrow T_\Sigma(X)$, $\sigma(f)$ is a Σ -term in X .

We usually omit the signature Σ from the notation and speak simply of *terms* (in X). We employ standard syntactic notions: A *substitution* is a map $\tau: |Y| \rightarrow T_\Sigma(X)$, for $X, Y \in \mathbf{C}$. We extend τ to a map $\bar{\tau}$ on terms $t \in T_\Sigma(X)$ as usual. Formally, we define $\bar{\tau}(t)$ inductively by $\bar{\tau}(x) = \tau(x)$ for $x \in X$, and $\bar{\tau}\sigma(f) = \sigma(\bar{\tau} \cdot f)$ for $f: \text{ar}(\sigma) \rightarrow T_\Sigma(X)$. We will not further distinguish between τ and $\bar{\tau}$ in the notation, writing $\tau(t) = \bar{\tau}(t)$ and $\tau \cdot f = \bar{\tau} \cdot f$ for t, f as above. Moreover, the set $\text{sub}(t)$ of *subterms* of a term $t \in T_\Sigma(X)$ is defined as usual; formally, we simultaneously define $\text{sub}(t)$ and $\text{sub}(f)$ for $f: I \rightarrow T_\Sigma(X)$ (with I some index set or object) inductively by $\text{sub}(x) = \{x\}$ for $x \in X$; $\text{sub}(\sigma(f)) = \{\sigma(f)\} \cup \text{sub}(f)$ for $f: \text{ar}(\sigma) \rightarrow T_\Sigma(X)$; and $\text{sub}(f) = \bigcup_{i \in I} \text{sub}(f(i))$.

Note that term formation operates without regard for the relational structure. Consequently, the evaluation of terms in a given Σ -algebra may fail to be defined:

► **Definition 4.5.** Let A be a Σ -algebra. For an object $X \in \mathbf{C}$ and a relation-preserving assignment $e: X \rightarrow A$, the partial *evaluation map* $e^\#: T_\Sigma(X) \rightarrow A$ is inductively defined by

- (1) $e^\#(x) = e(x)$ for $x \in X$, and
- (2) $e^\#(\sigma(f))$ is defined for $\sigma \in \Sigma$ and $f: |\text{ar}(\sigma)| \rightarrow T_\Sigma(X)$ iff the following hold:
 - (a) $e^\# \cdot f(i)$ is defined for all $i \in \text{ar}(\sigma)$, and
 - (b) if $\alpha(g)$ is a Π -edge in $\text{ar}(\sigma)$, then $A \models \alpha(e^\# \cdot (f \cdot g))$.

In case $e^\#(\sigma(f))$ is defined, we put $e^\#(\sigma(f)) = \sigma_A(e^\# \cdot f)$.

As indicated previously, we phrase theories using the relations in Π :

► **Definition 4.6.** A Σ -relation $X \vdash \alpha(f)$ consists of a *context* $X \in \mathbf{C}$ and a Π -edge $\alpha(f)$ in $T_\Sigma(X)$. We say that $X \vdash \alpha(f)$ is κ -ary if $X \in \mathcal{P}_\kappa$. A Σ -algebra A *satisfies* $X \vdash \alpha(f)$ if, for each relation preserving assignment $e: X \rightarrow A$, $e^\# \cdot f(i)$ is defined for all $i \in X$, and $\alpha_A(e^\# \cdot f)$. A (κ -ary) *relational algebraic theory* (Σ, \mathcal{E}) consists of the (κ -ary) signature Σ and a set \mathcal{E} of κ -ary Σ -relations. It determines the subcategory $\text{Alg}(\Sigma, \mathcal{E})$ of $\text{Alg } \Sigma$ consisting of those Σ -algebras which satisfy each Σ -relation in \mathcal{E} . We refer to categories of the shape $\text{Alg}(\Sigma, \mathcal{E})$ as *varieties* of Σ -algebras.

► **Remark 4.7.** For $\mathbf{C}_0 = \mathbf{Pos}$, the above notion of variety of Σ -algebras corresponds precisely to what we have termed ‘varieties of coherent algebras’ in earlier work with Adámek [6].

► **Example 4.8.** Recall that a (1-bounded) metric space X is *complete* if every Cauchy sequence $(x_i)_{i \in \omega}$ of points in X has a limit in X . That is, if (x_i) satisfies the Cauchy property

$$\forall \epsilon > 0. \exists N_\epsilon \in \omega. \forall n, m \geq N_\epsilon (d(y_n, y_m) < \epsilon), \quad (4.1)$$

then there is a point $\lim(x_i) \in X$ with the property of a limit: for all $\epsilon > 0$ there is $N \in \omega$ such that $x_n =_\epsilon \lim(x_i)$ for all $n \geq N$. The full subcategory $\mathbf{CMS} \hookrightarrow \mathbf{Met}$ of complete metric spaces is specified by the relational algebraic theory described below. Thus, by Theorem 4.13 below, we recover the fact that \mathbf{CMS} is monadic over \mathbf{Met} . Furthermore, we obtain a completely syntactic ω_1 -ary description of the metric completion monad via the deduction system introduced later in this section.

The theory $\mathbb{T}_{\mathbf{CMS}}$ of complete metric spaces has Γ -ary *limit operations* \lim_Γ for all spaces $\Gamma \in \mathcal{P}_{\omega_1}$ of the form $\Gamma = \{x_i \mid i \in \omega\}$ where $(x_i)_{i \in \omega}$ is a Cauchy sequence in Γ . The axioms of $\mathbb{T}_{\mathbf{CMS}}$ then say precisely that $\lim_\Gamma(x_i)$ is a limit of (x_i) . Explicitly, for all Γ as above, we impose all axioms of the shape

$$\Gamma \vdash \lim_\Gamma(x_n) =_\epsilon x_k \quad (k \geq N_\epsilon) \quad \text{where } N_\epsilon \text{ is as in (4.1).}$$

We fix a variety $\mathcal{V} = \mathbf{Alg}(\Sigma, \mathcal{E})$ for the remainder of this section. We are going to see that \mathcal{V} is a reflective subcategory of $\mathbf{Alg} \Sigma$ by application of Theorem 2.2, i.e. we show that \mathcal{V} is closed under limits and κ -directed colimits in $\mathbf{Alg} \Sigma$. We state the second property separately:

► **Proposition 4.9.** \mathcal{V} is closed under κ -directed colimits in $\mathbf{Alg} \Sigma$.

Combining this with Remark 4.3(2), we obtain

► **Corollary 4.10.** The forgetful functor $V: \mathcal{V} \rightarrow \mathbf{C}$ is κ -accessible.

It is fairly straightforward to show that \mathcal{V} is also closed under products and subobjects, and hence under limits (in $\mathbf{Alg} \Sigma$). Thus, as announced, we have:

► **Proposition 4.11.** \mathcal{V} is a reflective subcategory of $\mathbf{Alg} \Sigma$.

► **Remark 4.12.** It follows that the forgetful functor $\mathcal{V} \rightarrow \mathbf{C}_0$ has a left adjoint, namely the composite $\mathbf{C} \xrightarrow{F_\Sigma} \mathbf{Alg} \Sigma \xrightarrow{R_\mathcal{V}} \mathcal{V}$, where F_Σ is the left adjoint of the forgetful functor $\mathbf{Alg} \Sigma \rightarrow \mathbf{C}$ (Remark 4.3(4)) and $R_\mathcal{V}$ is the reflector according to Proposition 4.11. We call the ensuing monad $\mathbb{T}_\mathcal{V}$ the *free-algebra monad* of \mathcal{V} ; by Corollary 4.10, $\mathbb{T}_\mathcal{V}$ is κ -accessible.

Indeed, \mathcal{V} is essentially the category of Eilenberg-Moore algebras of $\mathbb{T}_\mathcal{V}$: Using Beck's monadicity theorem, one can show that

► **Theorem 4.13.** The forgetful functor $\mathcal{V} \rightarrow \mathbf{C}_0$ is monadic.

► **Corollary 4.14.** Every κ -ary relational algebraic theory may be translated into an enriched κ -accessible monad, preserving categories of models.

Relational Logic We proceed to set up a system of rules for deriving relations among terms. The calculus will involve two forms of judgements, both mentioning a context $X \in \mathbf{Str}(\Pi)$ (not necessarily κ -presentable). By a *relational judgement*

$$X \vdash \alpha(t_1, \dots, t_{\text{ar}(\alpha)}),$$

where $t_1, \dots, t_{\text{ar}(\alpha)} \in T_\Sigma(X)$, we indicate that for every valuation of X that is *admissible*, i.e. satisfies the relational constraints specified in X , the terms t_i are defined, and the resulting

tuple of values is in relation α . We treat expressions $\alpha(t_1, \dots, t_{\text{ar}(\alpha)})$ notationally as edges over $T_\Sigma(X)$, in particular sometimes write them in the form $\alpha(f)$ for $f: \text{ar}(\alpha) \rightarrow T_\Sigma(X)$. Moreover, a *definedness judgement* of the form

$$X \vdash \downarrow t$$

states that t is defined for all admissible valuations of X . (We could encode $\downarrow t$ as $\phi(t, t)$ for any $\phi \in \text{Eq}(x, y)$ but for technical reasons we prefer to keep definedness judgement distinct from relational judgements.)

The rules of the arising system of *relational logic* are shown below:

$$\begin{aligned} (\text{Var}) \frac{}{X \vdash \downarrow x} \quad (x \in X) \quad (\text{Ctx}) \frac{}{X \vdash \alpha(x_1, \dots, x_{\text{ar}(\alpha)})} (X \models \alpha(x_1, \dots, x_{\text{ar}(\alpha)})) \\ (\text{Mor}) \frac{\{X \vdash \alpha(f_i(j)) \mid j \in \text{ar}(\sigma)\} \cup \{X \vdash \downarrow \sigma(f_i) \mid i \in \text{ar}(\alpha)\}}{X \vdash \alpha(\sigma(f_i))} ((f_i: \text{ar}(\sigma) \rightarrow T_\Sigma(X))_{i \in \text{ar}(\alpha)}) \\ (\text{E-Ar}) \frac{\{X \vdash \alpha(f \cdot g) \mid \alpha(g) \in \text{ar}(\sigma)\} \cup \{X \vdash \downarrow f(i) \mid i \in \text{ar}(\sigma)\}}{X \vdash \downarrow \sigma(f)} (f: \text{ar}(\sigma) \rightarrow T_\Sigma(X)) \\ (\text{I-Ar}) \frac{\{X \vdash \alpha(\tau \cdot f) \mid \alpha(f) \in \Delta\} \cup \{X \vdash \downarrow \tau(y) \mid y \in \Delta\}}{X \vdash \beta(c)} (+) \\ (\text{RelAx}) \frac{\{X \vdash \tau \cdot \varphi \mid \varphi \in \Phi\} \cup \{X \vdash \downarrow \tau(f(i)) \mid i \in \text{ar}(\alpha)\}}{X \vdash \alpha(\tau \cdot f)} (\Phi \implies \alpha(f) \in \mathcal{A}, \tau: \text{Var} \rightarrow T_\Sigma(X)) \\ (\text{Ax}) \frac{\{X \vdash \alpha(\tau \cdot f) \mid \alpha(f) \in \Delta\} \cup \{X \vdash \downarrow \tau(y) \mid y \in \Delta\}}{X \vdash \beta(\tau \cdot g)} (\Delta \vdash \beta(g) \in \mathcal{E}) \end{aligned}$$

Recall that both the arities of operations in Σ and the contexts of the κ -ary Σ -relations in \mathcal{E} are in \mathcal{P}_κ . We assume such a $\Delta \in \mathcal{P}_\kappa$ to be specified as $\Delta = R(Y, E)$ by a κ -presentable object $(Y, E) \in \mathbf{Str}(\Pi)$ (cf. Lemma 3.7, Proposition 3.9, Lemma 3.19, Proposition 3.21); by writing $\phi \in \Delta$ for an edge ϕ , we indicate that $\phi \in E$ (rather than just $\phi \in \mathbf{E}(\Delta)$). The rules (E-Ar) and (I-Ar) apply to every $\sigma \in \Sigma$, and rule (Mor) applies to every $\sigma \in \Sigma$ and every $\alpha \in \Pi$. The side condition (+) of (I-Ar) is the following: for some axiom $\Delta \vdash \gamma(g)$ of \mathcal{V} there is $\sigma(h) \in \text{sub}(g)$, where $h: \text{ar}(\sigma) \rightarrow T_\Sigma(\Delta)$, such that $\text{ar}(\sigma) \models \beta(k)$ and

$$c = \text{ar}(\beta) \xrightarrow{k} \text{ar}(\sigma) \xrightarrow{h} T_\Sigma(\Delta) \xrightarrow{\tau} T_\Sigma(X).$$

Rule (Mor) captures the fact that operations σ are interpreted as morphisms of type $[\text{ar}(\sigma), A] \rightarrow A$, a condition that relates to enrichment of the induced monad. Rule (E-Ar) states that operations are defined when all the constraints given by their arity are satisfied. Rules (RelAx) and (Ax) allow application of the axioms of the Horn theory and the variety, respectively, in both cases instantiated with a substitution. A general substitution rule is not included but admissible. Rule (I-Ar) captures that every axiom of the variety is understood as implying that (under the constraints of the context) all subterms occurring in it are defined, in the sense that the constraints in the arities of the relevant operations hold.

► **Remark 4.15.** Instantiating the above system of rules to the theory of partial orders yields essentially the ungraded version of our previous deduction system for graded monads on **Pos** [8], up to the above-mentioned coding of definedness judgements. At first glance, the instantiation to the theory of metric spaces appears to yield a system that differs in several respects from the existing system of quantitative algebra [19]; besides the mentioned

absence of a general substitution rule, this concerns most prominently the absence of a cut rule (included in [19]) in our system. These distinctions are only superficial: as mentioned above, the more general substitution rule is admissible in our system, and it follows from completeness (Theorem 4.19) that the cut rule is admissible as well.

► **Lemma 4.16.** *The following rules are admissible:*

$$\text{(Arity)} \quad \frac{X \vdash \downarrow \sigma(m)}{X \vdash \alpha(m \cdot f)} \quad (\text{ar}(\sigma) \models \alpha(f), \quad m: |\text{ar}(\sigma)| \rightarrow T_\Sigma(X)) \quad \text{(Subterm)} \quad \frac{X \vdash \alpha(f)}{X \vdash \downarrow u} \quad (u \in \text{sub}(f))$$

Constructing free algebras We now show that relational logic gives rise to a syntactic construction of free algebras in the variety \mathcal{V} .

The set $\mathcal{T}_\mathcal{V}(X)$ of *derivably \mathcal{V} -defined terms in X* consists of those terms $t \in T_\Sigma(X)$ such that $X \vdash \downarrow t$ is derivable. We equip $\mathcal{T}_\mathcal{V}(X)$ with the relations

$$\mathcal{T}_\mathcal{V}(X) \models \alpha(f) \iff X \vdash \alpha(f) \text{ is derivable} \quad (\alpha \in \Pi, f: \text{ar}(\alpha) \rightarrow \mathcal{T}_\mathcal{V}(X))$$

making it into a Π -structure. We write \sim for the relation on $\mathcal{T}_\mathcal{V}(X)$ given by *derivable equality*: that is, for all $s, t \in \mathcal{T}_\mathcal{V}(X)$ we put $s \sim t$ iff $X \vdash \varphi$ is derivable for all $\varphi \in \text{Eq}(s, t)$, which is clearly an equivalence relation. The \sim -equivalence class of $t \in \mathcal{T}_\mathcal{V}(X)$ is denoted by $[t]$. We pick a *splitting* $u: \mathcal{T}_\mathcal{V}(X)/\sim \rightarrow \mathcal{T}_\mathcal{V}(X)$ of the canonical quotient map $q: \mathcal{T}_\mathcal{V}(X) \rightarrow \mathcal{T}_\mathcal{V}(X)/\sim$, i.e. $q \cdot u = \text{id}$, so u picks representatives of \sim -equivalence classes. Then $\mathcal{T}_\mathcal{V}(X)/\sim$ carries the structure of a \mathbf{C} -object, with edges defined by $\mathcal{T}_\mathcal{V}(X)/\sim \models \alpha(f)$ iff $\mathcal{T}_\mathcal{V}(X) \models \alpha(u \cdot f)$. ('Only if' means that u is relation preserving.)

► **Definition 4.17.** The *algebra $\mathcal{F}X$ of defined terms in X* is the Σ -algebra obtained by equipping $\mathcal{T}_\mathcal{V}(X)/\sim$ with the operations $\sigma_{\mathcal{F}X}: [\text{ar}(\sigma), \mathcal{T}_\mathcal{V}(X)/\sim] \rightarrow \mathcal{T}_\mathcal{V}(X)/\sim$ well-defined by $f \mapsto [\sigma(u \cdot f)]$, where $u: \mathcal{F}\Gamma \rightarrow \mathcal{T}_\mathcal{V}(X)$ is the chosen splitting of $q: \mathcal{T}_\mathcal{V}(X) \rightarrow \mathcal{F}X$.

► **Theorem 4.18.** *For every $X \in \mathbf{C}$, $\mathcal{F}X$ is a free algebra in \mathcal{V} .*

► **Theorem 4.19** (Soundness and Completeness). *$X \vdash \alpha(f)$ is derivable iff every $A \in \mathcal{V}$ satisfies $X \vdash \alpha(f)$.*

► **Remark 4.20.** Consequently, our system instantiated to the theory of metric spaces and the system of quantitative algebra [19], which is also sound and complete, are deductively equivalent. Hence, our results thus far imply that every quantitative algebraic theory induces an ω_1 -accessible monad. Indeed this remains true if one admits operations of countable arity, as in our theory of complete metric spaces (Example 4.8). Due to non-discrete contexts in axioms, monads induced by quantitative algebraic theories (such as $x =_{1/2} y \vdash x =_0 y$) in general fail to be finitary. However, our results do imply that the induced monad is finitary if only discrete contexts are used; e.g. this holds for the theories of left-invariant barycentric algebras and of quantitative semi-lattices, respectively [19] (note for the latter that axiom (S4) can be omitted in [19, Def. 9.1]). We conjecture that monads induced by *continuous equation schemes* [19] are also finitary.

5 Enriched Accessible Monads

We proceed to establish the monad-to-theory direction of our correspondence; as already indicated, given our fixed λ -ary Horn theory \mathcal{H} , this works only for λ -accessible monads and λ -ary theories, but not for accessibility degrees $\kappa < \lambda$ as in the theory-to-monad direction. So let $\mathbb{T} = (T, \eta, \mu)$ be an enriched λ -accessible monad on \mathbf{C} . We proceed to extract a λ -ary

relational algebraic theory from \mathbb{T} . We first review the equivalence between monads and *Kleisli triples* (see, e.g., Moggi [20], and originally Manes [18, Exercise 12]).

► **Definition 5.1.** A *Kleisli triple* in \mathbf{C}_0 is a triple $(T, \eta, (-)^*)$ consisting of a mapping $T: \mathbf{C}_0 \rightarrow \mathbf{C}_0$ (of objects), a morphism $\eta_X: X \rightarrow TX$ for all $X \in \mathbf{C}_0$, and an assignment of a morphism $f^*: TX \rightarrow TY$ to every morphism $f: X \rightarrow TY$. This data is subject to the laws below for all $X \in \mathbf{C}_0$ and all morphisms $f: X \rightarrow TY$ and $g: Y \rightarrow TZ$:

$$\eta_X^* = \text{id}_X, \quad f^* \cdot \eta_X = f, \quad \text{and} \quad g^* \cdot f^* = (g^* \cdot f)^*. \quad (5.1)$$

► **Remark 5.2.** The mapping which assigns to each monad (T, η, μ) the Kleisli triple $(T, \eta, (-)^*)$ with $(-)^*$ defined by $f^* = TX \xrightarrow{Tf} TTY \xrightarrow{\mu_Y} TY$ for $f \in \mathbf{C}_0(X, TY)$ yields a bijective correspondence between monads and Kleisli triples on \mathbf{C}_0 .

► **Notation 5.3.** For each operation σ in a signature Σ , we have a term $\sigma(u_{\text{ar}(\sigma)})$, where $u_{\text{ar}(\sigma)}$ is the inclusion $\text{ar}(\sigma) \hookrightarrow T_\Sigma(\text{ar}(\sigma))$. By abuse of notation, we also write σ for $\sigma(u_{\text{ar}(\sigma)})$.

► **Definition 5.4.** The λ -ary signature $\Sigma_{\mathbb{T}}$ induced by \mathbb{T} is the disjoint union of the sets $|T\Gamma|$ ($\Gamma \in \mathcal{P}_\lambda$), where elements of $|T\Gamma|$ have arity Γ . The *variety* $\mathcal{V}_{\mathbb{T}}$ induced by \mathbb{T} is $\mathcal{V}_{\mathbb{T}} = \text{Alg}(\Sigma_{\mathbb{T}}, \mathcal{E}_{\mathbb{T}})$ where $\mathcal{E}_{\mathbb{T}}$ contains all axioms of the following shape, with $\Gamma \in \mathcal{P}_\lambda$:

- (1) $\Gamma \vdash \alpha(\sigma_1, \dots, \sigma_{\text{ar}(\alpha)})$ for all $\sigma_i \in T\Gamma$ such that $T\Gamma \models \alpha(\sigma_1, \dots, \sigma_{\text{ar}(\alpha)})$
- (2) $\Gamma \vdash f^*(\sigma) = \sigma(f)$ for all $\Delta \in \mathcal{P}_\lambda$, morphisms $f: \Delta \rightarrow T\Gamma$, and $\sigma \in |T\Delta|$.
- (3) $\Gamma \vdash \eta_\Gamma(x) = x$ for every $x \in \Gamma$.

Note that in the second item above, for every $x \in \Delta$ the operation symbol $f(x) \in |T\Gamma|$ is considered as a term according to Notation 5.3. Hence $\sigma(f)$ is a term, too.

We now show that \mathbb{T} is the free-algebra monad of its induced variety $\mathcal{V}_{\mathbb{T}}$. For each $X \in \mathbf{C}$, the \mathbf{C} -object TX carries a canonical Σ -algebra structure with each operation σ_{TX} being defined by $\sigma_{TX}(f) := f^*(\sigma)$. We call TX the *canonical algebra over X* .

► **Lemma 5.5.** *Every canonical algebra lies in $\mathcal{V}_{\mathbb{T}}$.*

► **Theorem 5.6.** *Each enriched λ -accessible monad \mathbb{T} is the free-algebra monad of its induced variety $\mathcal{V}_{\mathbb{T}}$, with the free algebra on X given by the canonical algebra TX .*

► **Remark 5.7.** We have thus shown that given a λ -ary Horn theory \mathcal{H} , we can translate λ -accessible monads on $\mathbf{Str}(\mathcal{H})$ back into λ -ary theories, preserving the notion of model. For example, every ω_1 -accessible monad on \mathbf{Met} is induced by an ω_1 -ary theory, as illustrated in Example 4.8. The situation is more complicated for κ -ary monads where $\kappa < \lambda$. E.g. we can generate a finitary monad on \mathbf{Met} from a single binary operation of type $\{(x, y) \in A^2 \mid d(x, y) < 1/2\} \rightarrow A$. This monad is not induced by any theory with operations of internally finitely presentable (i.e. discrete) arity, in particular neither by an ω -ary theory in our framework nor by a quantitative algebraic theory [19].

6 Conclusions

We have introduced the framework of *relational logic* for reasoning about algebraic structure on categories of (finitary) relational structures axiomatized by possibly infinitary Horn theories, such as partial orders or metric spaces. We have proved soundness and completeness of a generic algebraic deduction system, and we have shown that λ -ary relational algebraic theories are in correspondence with λ -accessible enriched monads when the underlying Horn

theory is also λ -ary (where ‘ λ -ary’ refers to the arity of operations for relational algebraic theories, and to the number of premisses in axioms for Horn theories). Our results allow for a straightforward specification also of infinitary constructions such as metric completion.

The theory-to-monad direction of the above-mentioned correspondence remains true for κ -ary relational algebraic theories and κ -accessible monads on categories of models of λ -ary Horn theories for $\kappa < \lambda$, e.g. when looking at monads and theories on metric spaces. One open end that we leave for future research is to obtain a more complete coverage of this case, which will require a substantial generalization of both the way arities of operations are defined (these can no longer be taken to be objects of the base category) and in the way the axioms of the theory are organized, likely using more topologically-minded approaches.

References

- 1 Jirí Adámek. Free algebras and automata realizations in the language of categories. *Comment. Math. Univ. Carolin.*, pages 589–602, 1974.
- 2 Jirí Adámek, Matěj Dostál, and Jirí Velebil. A categorical view of varieties of ordered algebras, 2021. [arXiv:2011.13839](https://arxiv.org/abs/2011.13839).
- 3 Jiří Adámek, Horst Herrlich, and George Strecker. *Abstract and Concrete Categories*. Wiley-Interscience, New York, 1990.
- 4 Jirí Adámek, Stefan Milius, Lurdes Sousa, and Thorsten Wißmann. Finitely presentable algebras for finitary monads. *Theory Appl. Categ.*, 34:1179–1195, 2019.
- 5 Jiří Adámek and Jiří Rosický. *Locally Presentable and Accessible Categories*. London Mathematical Society Lecture Note Series. Cambridge University Press, 1994.
- 6 Jiří Adámek, Chase Ford, Stefan Milius, and Lutz Schröder. Finitary monads on the category of posets, 2020. [arXiv:2011.14796](https://arxiv.org/abs/2011.14796).
- 7 Stephen Bloom. Varieties of ordered algebras. *J. Comput. System Sci.*, pages 200–212, 1976.
- 8 Chase Ford, Stefan Milius, and Lutz Schröder. Behavioural preorders via graded monads. In *Logic in Computer Science, LICS 2021*. IEEE, 2021. To appear. Preprint available as [arXiv e-print 2011.14339](https://arxiv.org/abs/2011.14339).
- 9 Thomas Jech. *Set Theory*. Springer, 2003.
- 10 Max Kelly. Structures defined by finite limits in the enriched context, I. *Cah. Topol. Géom. Différ. Catég.*, 23(1):3–42, 1982.
- 11 Max Kelly and Steve Lack. Finite-product-preserving functors, kan extensions, and strongly-finitary 2-monads. *Appl. Categorical Struct.*, 1(1):85–94, 1993.
- 12 Max Kelly and John Power. Adjunctions whose counits are coequalizers, and presentations of finitary enriched monads. *J. Pure Appl. Algebra*, pages 163–179, 1993.
- 13 Alexander Kurz and Jiří Velebil. Quasivarieties and varieties of ordered algebras: regularity and exactness. *Math. Struct. Comput. Sci.*, 27(7):1153–1194, 2017.
- 14 Fred Linton. An outline of functorial semantics. In B. Eckmann, editor, *Seminar on Triples and Categorical Homology Theory*, volume 80 of *Lecture Notes Math.*, pages 7–52. Springer, 1969.
- 15 Fred E.J. Linton. Some aspects of equational theories. In *Proc. Conf. on Categorical Algebra at La Jolla*, pages 84–94. Springer, 1966.
- 16 R. Lucyshyn-Wright. Enriched algebraic theories and monads for a system of arities. *Theory Appl. Categ.*, 31(5):101–137, 2016.
- 17 Saunders MacLane. *Categories for the Working Mathematician*. Springer, 2nd edition, 1998.
- 18 Ernest Manes. *Algebraic Theories*. Springer, 1976.
- 19 Radu Mardare, Prakash Panangaden, and Gordon Plotkin. Quantitative algebraic reasoning. In Martin Grohe, Eric Koskinen, and Natarajan Shankar, editors, *Logic in Computer Science, LICS 2016*, pages 700–709. ACM, 2016. URL: <https://doi.org/10.1145/2933575.2934518>, doi:10.1145/2933575.2934518.
- 20 Eugenio Moggi. Notions of computations and monads. *Inf. Comput.*, 93(1):55–92, 1991.
- 21 Koki Nishizawa and John Power. Lawvere theories enriched over a general base. *J. Pure Appl. Algebra*, 213(3):377–386, 2009.
- 22 Gordon Plotkin and John Power. Adequacy for algebraic effects. In *Foundations of Software Science and Computation Structures, FOSSACS 2001*, volume 2030 of *LNCS*, pages 1–24. Springer, 2001. doi:10.1007/3-540-45315-6_1.
- 23 John Power. Enriched lawvere theories. *Theory Appl. Categories*, pages 83–93, 1999.
- 24 John Power and Richard Garner. An enriched view on the extended finitary monad-Lawvere theory correspondence. *Logical Methods Comput. Sci.*, 14(1:16):23 pp., 2018.
- 25 Jiří Rosický. Metric monads, 2021. [arXiv:2012.14641](https://arxiv.org/abs/2012.14641).

A

 Details for Section 3

Details for Example 3.5 *L-valued relations:* Every \mathcal{H}_L -model (X, E) induces an L -valued relation P_E on X by

$$P_E(x, y) = \bigwedge \{p \in L_0 \mid \alpha_p(x, y) \in E\}.$$

Conversely, every L -valued relation P on a set X induces a \mathcal{H}_L -model (X, E_P) given by

$$E_P = \{\alpha_p(x, y) \mid p \in L_0, p \geq P(x, y)\}.$$

The given conditions on L_0 ensure precisely that these constructions are mutually inverse: By meet density of L_0 , it is straightforward to see that $P_{E_P} = P$. Moreover, given a \mathcal{H}_L -model (X, E) , we have $E_{P_E} = E$. Here, ‘ \supseteq ’ is trivial; we prove ‘ \subseteq ’. So suppose that

$$q \geq \bigwedge \{p \in L_0 \mid \alpha_p(x, y) \in E\};$$

we have to show $\alpha_q(x, y) \in E$. We distinguish the following cases:

- (1) If $r := \bigwedge \{p \in L_0 \mid \alpha_p(x, y) \in E\} \in L_0$, then by **(Arch)**, $\alpha_r(x, y) \in E$, and hence $\alpha_q(x, y) \in E$ by **(Up)**.
- (2) Otherwise, the assumptions on L_0 imply that there is $p \in L_0$ such that $\alpha_p(x, y) \in E$ and $q \geq p$; then $\alpha_q(x, y) \in E$ by **(Up)**.

Metric spaces: As indicated in the main text, this is essentially a special case of the one for L -valued relations: $L_0 := [0, 1] \cap \mathbb{Q} \subseteq [0, 1] =: L$ satisfy the relevant assumptions, so we have mutually inverse constructions as indicated above and in the main text. Under this correspondence, the axioms of metric spaces clearly correspond precisely to the axioms **(Refl)**, **(Equal)**, **(Sym)**, and **(Triang)** of \mathcal{H}_{Met} .

Proof of Proposition 3.9. (2) \implies (1): Generally, left adjoint functors (in this case, R) preserve λ -presentable objects if their right adjoint (in this case, the inclusion $\mathbf{Str}(\Pi, \mathcal{A}) \hookrightarrow \mathbf{Str}(\Pi)$) preserves λ -directed colimits (see e.g. [4, Lem. 2.4(1)]).

(1) \implies (3): The bound on the cardinality is as in the more general case (see the proof of Proposition 3.21). We can assume that $r_X = \text{id}_X$. Write X as the λ -directed colimit in $\mathbf{Str}(\Pi)$ of the objects $(|X|, E)$ such that $E \subseteq \mathbf{E}(X)$ and $\text{card } E < \lambda$. This colimit is preserved by the left adjoint R . Now apply λ -presentability of X to see that $r_X = \text{id}_X: X \rightarrow RX$ factorizes through Ri for one of the inclusions $i: (|X|, E) \rightarrow X$; that is, Ri is a split epimorphism. The claim follows once we show that Ri is also a monomorphism. To this end, consider the naturality square

$$\begin{array}{ccc} (|X|, E) & \xrightarrow{r_{(|X|, E)}} & R(|X|, E) \\ \downarrow i & & \downarrow Ri \\ X & \xrightarrow{r_X} & RX \end{array}$$

In this diagram, the underlying maps of i and r_X are bijective, so the underlying map of $r_{(|X|, E)}$ is injective, and hence also bijective since r is componentwise surjective. It follows that the underlying map of Ri is bijective, hence Ri is monic.

(3) \implies (2): Take $(Y, E) = (|X|, E)$. ◀

Proof of Proposition 3.12. We show that u_Y is indeed a universal arrow; that is, we show that u_Y is actually a morphism of the claimed type, and that u_Y has the following universal

property: Given a further $\mathbf{Str}(\Pi)$ -morphism $g: Y \rightarrow [X, Z]$, there exists a unique morphism $g^\sharp: Y \otimes X \rightarrow Z$ such that $[X, g^\sharp] = g$:

$$\begin{array}{ccc} Y & \xrightarrow{u_Y} & [X, Y \otimes X] & & Y \otimes X \\ & \searrow \forall g & \downarrow [X, g^\sharp] & & \downarrow \exists! g^\sharp \\ & & [X, Z] & & Z \end{array}$$

We prove the latter property first. We put

$$g^\sharp(y, x) = g(y)(x).$$

This is clearly the only map making the diagram commute, so we are only left to show that it is indeed a morphism. So let $e \in \mathbf{E}(Y \otimes X)$; we have to show $g^\sharp \cdot e \in \mathbf{E}(Z)$. We distinguish cases on e according to the definition of $Y \otimes X$:

- $\pi_1 \cdot e$ is constant, say with constant value y , and $\pi_2 \cdot e \in \mathbf{E}(X)$. Then $g^\sharp \cdot e = g(y) \cdot \pi_2 \cdot e \in \mathbf{E}(Z)$ because $g(y)$ is a morphism, and
- $\pi_2 \cdot e$ is constant, say with constant value x , and $\pi_1 \cdot e \in \mathbf{E}(Y)$. Then $g^\sharp \cdot e = \pi_x \cdot g \cdot e \in \mathbf{E}(Z)$ because g is a morphism.

We next show that u_Y has the claimed type; that is, we show that for $y \in Y$, $u_Y(y): X \rightarrow Y \otimes X$ is a morphism. So let $e \in \mathbf{E}(X)$. Then $u_Y(y) \cdot e \in \mathbf{E}(Y \otimes X)$ as required, since $\pi_1 \cdot u_Y(y) \cdot e$ is constant and $\pi_2 \cdot u_Y(y) \cdot e = e \in \mathbf{E}(X)$. Finally, we show that u_Y is a morphism. So let $e \in \mathbf{E}(Y)$, and let $x \in X$; we have to show that $\pi_x \cdot u_Y \cdot e \in \mathbf{E}(Y \otimes X)$. This holds because $\pi_2 \cdot \pi_x \cdot u_Y \cdot e$ is constant and $\pi_1 \cdot \pi_x \cdot u_Y \cdot e = e \in \mathbf{E}(Y)$. ◀

We have claimed that the internal hom restricts to $\mathbf{Str}(\mathcal{H})$, that is:

▶ **Lemma A.1.** $[X, Y] \in \mathbf{Str}(\Pi, \mathcal{A})$ for all $Y \in \mathbf{Str}(\Pi, \mathcal{A})$.

Proof of Lemma A.1. Let $\Phi \implies \beta(g)$ be an axiom and let $e: \mathbf{Var} \rightarrow \mathbf{Str}(\Pi)(X, Y)$ be a map such that $\alpha(e \cdot f) \in \mathbf{E}([X, Y])$ for all $\alpha(f) \in \Phi$. We proceed to show that $\beta(e \cdot g) \in \mathbf{E}([X, Y])$. That is, we verify that $\beta(e \cdot g)_x \in \mathbf{E}(Y)$ for all $x \in X$.

To this end, it suffices to show that $\beta(\xi_x \cdot g) \in \mathbf{E}(Y)$ for all $x \in X$, where $\xi_x: \mathbf{Var} \rightarrow Y$ is the assignment $v \mapsto e(v)(x)$. Indeed, we have

$$\xi_x \cdot g(i) = \xi_x(g(i)) = e(g(i))(x) = (e \cdot g)_x(i)$$

for all $i \in \mathbf{ar}(\beta)$. Thus, $\xi_x \cdot g = (e \cdot g)_x$. To conclude the proof, note that for each $\alpha(f) \in \Phi$ we have $\alpha((e \cdot f)_x) \in \mathbf{E}(Y)$ for all $x \in X$. Hence also $\alpha(\xi_x \cdot f) \in \mathbf{E}(Y)$ for all $x \in X$. Since Y is an \mathcal{A} -model, it follows that $\beta(\xi_x \cdot g) \in \mathbf{E}(Y)$ for all $x \in X$, as desired. ◀

Proof of Corollary 3.13. From Proposition A.2 below, using Lemma A.1. ◀

▶ **Proposition A.2.** Let $\iota: \mathcal{C}' \hookrightarrow \mathcal{C}$ be a full reflective subcategory, with reflector $R: \mathcal{C} \rightarrow \mathcal{C}'$, of a closed monoidal category $(\mathcal{C}, \otimes, I)$ such that $[C, D]$ lies in \mathcal{C}' for every $C, D \in \mathcal{C}'$. Then \mathcal{C}' is also closed monoidal, with monoidal product $R(X \otimes Y)$.

Proof. We have the following chain of bijections, natural in Y and Z (and omitting ι in front of X and Y):

$$\mathcal{C}'(R(Y \otimes X), Z) \cong \mathcal{C}(Y \otimes X, \iota Z) \cong \mathcal{C}(Y, [X, \iota Z]) \cong \mathcal{C}'(Y, [X, Z]),$$

where the last step uses that $[X, \iota Z]$ lies in \mathcal{C}' and \mathcal{C}' is full. ◀

Proof of Proposition 3.16. One extracts from [5, Proof of Proposition 2.23] that if F is a right adjoint functor between locally λ -presentable categories whose left adjoint preserves externally λ -presentable objects, then F is λ -accessible. We apply this to $F = [X, -]$ for X externally λ -presentable. Preservation of externally λ -presentable objects by the left adjoint $(-) \otimes X$ of F means precisely that the class of externally λ -presentable objects is closed under $\otimes_{\mathcal{H}}$ as claimed in the main body of the paper. By Proposition 3.9, this follows immediately from the corresponding statement for $\mathcal{H} = (\Pi, \emptyset)$. But the latter is clear by the description of externally λ -presentable objects in $\mathbf{Str}(\Pi)$ in terms of simple cardinality constraints. \blacktriangleleft

B Details for Section 3.1

Details for Example 3.17 (2): *Example of a complete lattice L where binary joins fail to distribute over directed infima:* Take L to consist of two copies of the reals with their usual orderings, with elements denoted $l(x)$ and $r(x)$, respectively, for $x \in \mathbb{R}$, and additionally a top element \top , a bottom element \perp , and an element b that satisfies $l(0) \leq b$, $r(0) \leq b$, and no other inequalities other than the ones entailed by \top and \perp being a top and a bottom element, respectively. Then $\bigwedge_n (l(1/n) \vee r(1/n)) = \top$ but $(\bigwedge_n l(1/n)) \vee (\bigwedge_n r(1/n)) = l(0) \vee r(0) = b$.
 (3): In the same L , there is also an element as prescribed in (3), namely $l = l(1)$.

Proof of Lemma 3.19. Given (X, E) κ -compact, apply κ -compactness to the cover E of (X, E) . \blacktriangleleft

We next verify the characterization of weakly κ -presentable objects in the locally λ -presentable category \mathbf{C} , where $\kappa < \lambda$ is an infinite cardinal, as precisely the κ -compact objects of cardinality $< \lambda$ (Proposition 3.21)

The main essence of the characterization is the captured inn the following lemmas:

► **Lemma B.1.** *Let $X \in \mathbf{Str}(\Pi, \mathcal{A})$.*

(1) *X is κ -compact iff for every morphism of the form $f: X \rightarrow R(Y, E)$ for $(Y, E) \in \mathbf{Str}(\Pi)$, there exist $E' \subseteq_{\kappa} E$ and $f': X \rightarrow R(Y, E')$ such that $f = Rj \cdot f'$, where $j: (Y, E') \rightarrow (Y, E)$ is the $\mathbf{Str}(\Pi)$ -morphism carried by id_Y .*

(2) *If $\text{card } |X| < \kappa$, then X is κ -compact iff for every morphism of the form $f: X \rightarrow R(Y, E)$ for $(Y, E) \in \mathbf{Str}(\Pi)$, there exist $Y' \subseteq_{\kappa} Y$, $E' \subseteq_{\kappa} E$ and $f': X \rightarrow R(Y', E')$ such that $f = Rj \cdot f'$, where $j: (Y', E') \rightarrow (Y, E)$ is the $\mathbf{Str}(\Pi)$ -morphism carried by the inclusion $Y' \hookrightarrow Y$.*

Proof. (1) ‘If’ is trivial; we prove ‘only if’. For ease of notation, we assume w.l.o.g. that $|X|$ and Y are disjoint; then their union $|X| \cup Y$ serves as a coproduct $|X| + Y$. Fix a map $s: |R(Y, E)| \rightarrow Y$ splitting the reflection $r_{(Y, E)}: (Y, E) \rightarrow R(Y, E)$; that is, $r_{(Y, E)} \cdot s = \text{id}_Y$ (s is not in general a morphism). We define a set E_f of edges over $|X| + Y$ by

$$E_f = \bigcup_{x \in |X|} \text{Eq}(x, s(f(x))).$$

We then have a morphism

$$h = ((Y, E) \xrightarrow{\text{inr}} (|X| + Y, E_f \cup E) \xrightarrow{r_{(|X|+Y, E_f \cup E)}} R(|X| + Y, E_f \cup E)).$$

Note that $r_{(|X|+Y, E_f \cup E)}$ merges every $x \in |X|$ with $e(f(x)) \in Y$, and indeed it is not difficult to see that $h^\sharp: R(Y, E) \rightarrow R(|X| + Y, E_f \cup E)$ is an isomorphism. In addition, the map $r_{(|X|+Y, E_f \cup E)} \cdot \text{inl}: X \rightarrow R(|X| + Y, E_f \cup E)$ carries a morphism, where inl is the

left-hand coproduct injection $|X| \rightarrow |X| + Y$; indeed, $r_{(|X|+Y, E_f \cup E)} \cdot \text{inl} = h^\# \cdot f$. By assumption, we thus have $E' \subseteq_\kappa E_f \cup E$ and a morphism $l: X \rightarrow R(|X| + Y, E')$ such that $Rj \cdot l = r_{(|X|+Y, E_f \cup E)} \cdot \text{inl}$ where $j: (|X| + Y, E') \rightarrow (|X| + Y, E_f \cup E)$ is the morphism carried by $\text{id}_{|X|+Y}$.

Next, we define the map

$$k = [s \cdot f, \text{id}_Y]: |X| + Y \rightarrow Y.$$

Then we have a morphism

$$p = r_{(Y, E' \cap E)} \cdot k: (|X| + Y, E') \rightarrow R(Y, E' \cap E).$$

To see this, let $(\alpha, g) \in E' \subseteq E_f \cup E$. The case where $(\alpha, g) \in E$ is trivial; the other case is that $(\alpha, g) \in Eq(x, s(f(x)))$ for some $x \in X$. Such an edge is preserved by p because $p(x) = r_{(Y, E' \cap E)} \cdot s \cdot f(x) = r_{(Y, E' \cap E)}(s(f(x))) = p(s(f(x)))$.

We thus have a diagram

$$\begin{array}{ccccc}
 & & R(|X| + Y, E') & & \\
 & \nearrow l & \downarrow Rj & \searrow p^\# & \\
 X & \xrightarrow{r \cdot \text{inl}} & R(|X| + Y, E_f \cup E) & & R(Y, E' \cap E) \\
 & \searrow f & \downarrow (h^\#)^{-1} & \swarrow Rm & \\
 & & R(Y, E) & &
 \end{array}$$

where $m: (Y, E' \cap E) \rightarrow (Y, E)$ is the morphism carried by id_Y . We are done once we show that the diagram commutes. The upper left triangle commutes by construction of l , and commutation of the lower left triangle has been noted above (in the equivalent form $h^\# \cdot r \cdot \text{inl} = f$). Commutation of the right-hand square is equivalent to $h^\# \cdot Rm \cdot p^\# = Rj$. This holds precisely when it holds when precomposed with the reflection $r_{(|X|+Y, E')}$; we compute as follows (in **Set**):

$$\begin{aligned}
 h^\# \cdot Rm \cdot p^\# \cdot r_{(|X|+Y, E')} &= h^\# \cdot Rm \cdot p && \text{univ. prop. of } r_{(|X|+Y, E')} \\
 &= h^\# \cdot Rm \cdot r_{(Y, E' \cap E)} \cdot k && \text{def. of } p \\
 &= h^\# \cdot r_{(Y, E)} \cdot m \cdot k && \text{def. of } R \text{ on morphisms} \\
 &= h^\# \cdot r_{(Y, E)} \cdot k && m \text{ carried by identity} \\
 &= h \cdot k && \text{univ. prop. of } r_{(Y, E)} \\
 &= h \cdot [s \cdot f, \text{id}_Y] && \text{def. of } k \\
 &= r_{(|X|+Y, E_f + E)} \cdot \text{inr} \cdot [s \cdot f, \text{id}_Y] && \text{def. of } h \\
 &\stackrel{(*)}{=} r_{(|X|+Y, E_f + E)} \cdot [\text{inl}, \text{inr}] && \\
 &= r_{(|X|+Y, E_f + E)} \cdot j && j \text{ carried by identity} \\
 &= Rj \cdot r_{(|X|+Y, E')} && \text{def. of } R \text{ on morphisms.}
 \end{aligned}$$

For the step marked with $(*)$ we consider coproduct components separately: the right-hand component is clear, and for the left-hand one use that $r_{(|X|+Y, E_f \cup E)}$ merges $\text{inr} \cdot s \cdot f$ and inl since $E_f \supseteq Eq(x, s(f(x)))$.

(2) Immediate from the first claim and Lemma B.2. \blacktriangleleft

► **Lemma B.2.** *Let $f: X \rightarrow R(Y, E)$, and suppose that $\text{card } |X|, \text{card } E < \kappa$. Then there exist $Y' \subseteq_\kappa Y$ such that all edges in E are edges over Y' , and $f': X \rightarrow R(Y', E)$ such that $f = Ri \cdot f'$ where $i: (Y', E) \hookrightarrow (Y, E)$ is the morphism carried by the inclusion $Y' \hookrightarrow Y$.*

Proof. Fix a splitting $s: |R(Y, E)| \rightarrow Y$ of the reflection $r_{(Y, E)}$ (s is not in general a morphism). Write Y_E for the set of elements of Y that appear in edges in E , and put

$$Y' = e \cdot f[X] \cup Y_E;$$

if Y' is empty, then add an arbitrary element of Y to Y' (if Y is also empty, then there is nothing to show). Then $\text{card } Y' < \kappa$, and since $Y' \neq \emptyset$, the inclusion map $i: Y' \hookrightarrow Y$ has a left inverse $q: Y \rightarrow Y'$, i.e. $q \cdot i = \text{id}_{Y'}$. Since $s(f(x)) \in Y'$ for all $x \in |X|$, we have $i \cdot q \cdot s \cdot f = s \cdot f$. Moreover, q is clearly a morphism $q: (Y, E) \rightarrow (Y', E)$. The arising morphism $f' = Rq \cdot f$ provides the claimed factorization of f :

$$\begin{aligned} Ri \cdot Rq \cdot f &= R(i \cdot q) \cdot f && \text{functoriality of } R \\ &= R(i \cdot q) \cdot r_{(Y, E)} \cdot s \cdot f && \text{since } r_{(Y, E)} \cdot s = \text{id} \\ &= r_{(Y, E)} \cdot i \cdot q \cdot s \cdot f && \text{def. of } R \text{ on morphisms} \\ &= r_{(Y, E)} \cdot s \cdot f && \text{since } e \cdot f = i \cdot q \cdot s \cdot f \\ &= f && \text{since } r_{(Y, E)} \cdot s = \text{id}. \quad \blacktriangleleft \end{aligned}$$

Proof of Proposition 3.21. (1) \Rightarrow (2): We first prove that $\text{card } |X| < \kappa$. Write X as the κ -directed colimit of all its subobjects of cardinality smaller than κ . This is possible by our assumptions that arities of relation symbols in Π are finite. Since X is weakly κ -presentable, we know that id_X factorizes through one of the colimit injections, say $i: X' \rightarrow X$ for X' with $\text{card } |X'| < \kappa$. That is, for some morphism $h: X \rightarrow X'$ we have $i \cdot h = \text{id}_X$, whence $X \cong X'$ has cardinality smaller than κ .

Second, we prove that X is κ -compact. Given a cover (Y, E) of X , with inclusion $i: |X| \hookrightarrow Y$, write (Y, E) as the κ -directed colimit in $\mathbf{Str}(\Pi)$ of all objects (Y, E') such that $E' \subseteq_{\kappa} E$. Since the reflector R is a left adjoint, it preserves all colimits, so $R(Y, E)$ is the κ -directed colimit of the $R(Y, E')$. By weak κ -presentability of X , the morphism $r_{(Y, E)} \cdot i: X \rightarrow R(Y, E)$ factors through one of the colimit injections, as required.

(2) \Rightarrow (1): Let $(D_i \xrightarrow{c_i} D)_{i \in I}$ be a directed colimit of objects $D_i = (Y_i, E_i)$ in $\mathbf{Str}(\Pi, \mathcal{A})$, and let $f: X \rightarrow D$ be a morphism. Let $(D_i \xrightarrow{\bar{c}_i} (Y, E))_{i \in I}$ be the colimit of the D_i in $\mathbf{Str}(\Pi)$; then $D = R(Y, E)$ and $c_i = r_{(Y, E)} \cdot \bar{c}_i = R(\bar{c}_i)$ (assuming w.l.o.g. that $R(Y_i, E_i) = (Y_i, E_i)$). By Lemma B.1, there exist $Y' \subseteq_{\kappa} Y$, $E' \subseteq_{\kappa} E$, and $f': X \rightarrow R(Y, E')$ such that $f = Rj \cdot f'$ where $j: (Y', E') \rightarrow (Y, E)$ is the morphism carried by the inclusion $Y' \hookrightarrow Y$. Now (Y', E') is κ -presentable in $\mathbf{Str}(\Pi)$; so there are i and $j': (Y', E') \rightarrow (Y_i, E_i)$ such that $j = \bar{c}_i \cdot j'$. This yields the desired factorization of f : For $f'' = Rj' \cdot f'$, we have

$$c_i \cdot f'' = R\bar{c}_i \cdot Rj' \cdot f' = Rj \cdot f' = f. \quad \blacktriangleleft$$

C Details for Section 4

We will now verify that the object assignment $I: \mathbf{Alg } \Sigma \rightarrow \mathbf{Alg } H_{\Sigma}$ forms a concrete isomorphism over \mathbf{C} . An element of $H_{\Sigma} X$ is a morphism $f: \text{ar}(\sigma) \rightarrow X$ in \mathbf{C} for some specified $\sigma \in \Sigma$; we denote this by the pair (σ, f) .

Indeed, it is easy to verify that the assignment $I: (A, \sigma_A) \mapsto (A, \alpha)$ which maps each Σ -algebra A to the H_{Σ} -algebra $\alpha: H_{\Sigma} A \rightarrow A$ defined by

$$\alpha(f: \text{ar}(\sigma) \rightarrow A) := \sigma_A(f)$$

is an isomorphism, and I preserves the \mathbf{C} -object underlying each Σ -algebra and the underlying morphism of each homomorphism in $\mathbf{Alg } \Sigma$. That is, for the forgetful functors

$$U: \mathbf{Alg } \Sigma \rightarrow \mathbf{C}_0 \quad \text{and} \quad \bar{U}: \mathbf{Alg } H_{\Sigma} \rightarrow \mathbf{C}_0,$$

we have $U = I \cdot \bar{U}$.

► **Proposition C.1.** *Alg Σ and Alg H_Σ are isomorphic as concrete categories over \mathbf{C} .*

Proof. Every H_Σ -algebra $\alpha: H_\Sigma A \rightarrow A$ induces a Σ -algebra A , on the same carrier A , with the operations σ_A ($\sigma \in \Sigma$) defined by

$$\sigma_A(f) := \alpha(\sigma, f) \quad \text{for every } f: \text{ar}(\sigma) \rightarrow A.$$

Conversely, each Σ -algebra A induces a H_Σ -algebra $\alpha_A: H_\Sigma A \rightarrow A$ defined by

$$\alpha(\sigma, f) = \sigma_A(f), \quad f \in [\text{ar}(\sigma), A].$$

These constructions are mutually inverse, and a \mathbf{C} -morphism $h: A \rightarrow B$ is a homomorphism $A \rightarrow B$ of Σ -algebras iff it is a homomorphism $(A, \alpha) \rightarrow (B, \beta)$ of the corresponding H_Σ -algebras. We restrict ourselves to showing the latter. To this end, suppose first that $h: A \rightarrow B$ is a morphism of Σ -algebras. Then $h \cdot \sigma_A(f) = \sigma_B(h \cdot f)$ for all $\sigma \in \Sigma$ and all morphisms $f: \text{ar}(\sigma) \rightarrow A$. Thus, for all morphisms $f: \text{ar}(\sigma) \rightarrow A$, we have

$$h \cdot \alpha(f) = h(\sigma_A(f)) = \sigma_B(h \cdot f) = \beta(h \cdot f) = \beta \cdot H_\Sigma(h)(f).$$

Conversely, if $h: A \rightarrow B$ is a morphism of H_Σ -algebras, then $h \cdot \alpha = \beta \cdot H_\Sigma(h)$. To conclude the proof, we compute

$$h \cdot \sigma_A(f) = h \cdot \alpha(f) = \beta \cdot H_\Sigma(h)(f) = \beta(h \cdot f) = \sigma_B(h \cdot f).$$

In short, we conclude that the assignment described above yields an isomorphism

$$I: \text{Alg } \Sigma \cong \text{Alg } H_\Sigma$$

which is moreover concrete over \mathbf{C} : indeed, I clearly preserves underlying sets of Σ -algebras, and it preserves the underlying maps of homomorphisms, as demonstrated above. ◀

Varieties of Σ -algebras

► **Lemma C.2.** *Let $h: A \rightarrow B$ be a homomorphism of Σ -algebras A, B , and let $e: X \rightarrow A$ be a relation-preserving assignment. Then, for all terms $t \in T_\Sigma(X)$, we have:*

- (1) $(h \cdot e)^\#(t)$ is defined whenever $e^\#(t)$ is defined, and $(h \cdot e)^\#(t) = h(e^\#(t))$.
- (2) if $(h \cdot e)^\#(t)$ is defined and h is an embedding, then $e^\#(t)$ is defined.

Proof of Lemma C.2. (1) We proceed by induction on $t \in T_\Sigma(X)$. In case t is a variable in context X , the desired statement is immediate: indeed, $(h \cdot e)^\#(t)$ is defined, and

$$(h \cdot e)^\#(t) = h \cdot e(t) = h(e(t)) = h(e^\#(t))$$

where we have only used the definition of $(-)^\#$. Now, suppose that t has the shape $\sigma(f)$ for some operation $\sigma \in \Sigma$ and some $f: |\text{ar}(\sigma)| \rightarrow T_\Sigma(X)$. If $e^\#(\sigma(f))$ is defined, then

- (a) $e^\# \cdot f(i)$ is defined for all $i \in \text{ar}(\sigma)$;
- (b) $\alpha_A(e^\# \cdot (f \cdot g))$ for all $\alpha(g) \in \mathbf{E}(\text{ar}(\sigma))$.

Then, applying the inductive hypothesis to the subterms $f(i)$ of t , we obtain that $(h \cdot e)^\#(f(i))$ is defined for all $i \in \text{ar}(\sigma)$. Moreover, by (2), and since h is a homomorphism, we see that

$\alpha_B((h \cdot e)^\# \cdot f \cdot g)$ for all edges $\alpha(g) \in \mathbf{E}(\mathbf{ar}(\sigma))$. Thus, $(h \cdot e)^\#(t)$ is defined. To conclude the proof, we compute

$$\begin{aligned}
(h \cdot e)^\#(\sigma(f)) &= \sigma_B((h \cdot e)^\# \cdot f) && \text{definition } (-)^\# \\
&= \sigma_B(h \cdot e^\# \cdot f) && \text{induction} \\
&= h(\sigma_A(e^\# \cdot f)) && h \text{ is a homomorphism} \\
&= h(e^\#(\sigma(f))) && \text{definition } (-)^\# \quad \blacktriangleleft
\end{aligned}$$

(2) Now suppose that h is an embedding; we proceed by induction as before. For the inductive step, suppose that $t \in T_\Sigma(X)$ is of the form $t = \sigma(f)$ for some $\sigma \in \Sigma$ and some map $f: \mathbf{ar}(\sigma) \rightarrow T_\Sigma(X)$, and assume that $(h \cdot e)^\#(\sigma(f))$ is defined. Then, by definition of $(-)^\#$, we have that

- (a) $(h \cdot e)^\#(f(i))$ is defined for all $i \in \mathbf{ar}(\sigma)$ and
- (b) $B \models \alpha((h \cdot e)^\# \cdot (f \cdot g))$ for all $\alpha(g) \in \mathbf{E}(\mathbf{ar}(\sigma))$.

Then, by induction, we have that $e^\#(f(i))$ is defined for all $i \in \mathbf{ar}(\sigma)$. Hence, by the first part of this proposition, we see that $(h \cdot e)^\#(f(i)) = h \cdot e^\#(f(i))$ for all $i \in \mathbf{ar}(\sigma)$. In particular, given an edge $\alpha(g) \in \mathbf{E}(\mathbf{ar}(\sigma))$, we have that $B \models \alpha(h \cdot e^\# \cdot (f \cdot g))$. Since h is relation reflecting, it follows that $A \models \alpha(e^\# \cdot (f \cdot g))$ as well. Thus, $e^\#(t)$ is defined, as desired.

Proof of Proposition 4.9. Immediate from Lemma C.3 below. ◀

► **Lemma C.3.** *Let $(D_i \xrightarrow{d_i} A)_{i \in I}$ be a κ -directed colimit in $\mathbf{Alg} \Sigma$, and let $X \vdash \alpha(f)$ be a Σ -relation. If each D_i satisfies $X \vdash \alpha(f)$, then D satisfies $X \vdash \alpha(f)$.*

Proof. Suppose that every D_i satisfies $X \vdash \alpha(f)$ and let $g: X \rightarrow A$ be a relation preserving assignment. Using that $(D_i \xrightarrow{d_i} A)_{i \in I}$ is a κ -directed colimit, it follows, since X is κ -presentable, that there exists $i \in I$ and a relation preserving map $\bar{g}: X \rightarrow D_i$ such that $g = d_i \cdot \bar{g}$. Since D_i satisfies $\Gamma \vdash \alpha(f)$, we have:

- (1) $(\bar{g})^\# \cdot f(i)$ is defined for all $i \in \mathbf{ar}(\alpha)$;
- (2) $D_i \models \alpha((\bar{g})^\# \cdot f)$.

Applying Lemma C.2 to the homomorphism $d_i: D_i \rightarrow A$ and the assignment $(\bar{g}): X \rightarrow D_i$, we see that

$$g^\#(f(i)) = (d_i \cdot \bar{g})^\#(f(i)) = d_i \cdot (\bar{g})^\#(f(i))$$

is defined for all $i \in \mathbf{ar}(\alpha)$ by (1). Furthermore, using that d_i is relation preserving, it follows from (2) that $D \models \alpha(d_i \cdot (\bar{g})^\# \cdot f)$. To conclude the proof, we use that $d_i \cdot (\bar{g})^\# = (d_i \cdot \bar{g})^\# = g^\#$ to see that $D \models \alpha(g^\# \cdot f)$. Hence D satisfies $X \vdash \alpha(f)$, as desired. ◀

► **Lemma C.4.** *Let $A = \prod_{i \in I} A_i$ be a product of algebras such that each A_i is an algebra in the variety \mathcal{V} . Then, for all relation preserving assignments $f: \Gamma \rightarrow A$ and all terms $t \in T_\Sigma(\Gamma)$: $f^\#(t)$ is defined iff $(\pi_i \cdot f)^\#(t)$ is defined for all $i \in I$.*

Proof. To this end, we proceed by induction on $t \in T_\Sigma(\Gamma)$. If t is a variable in context Γ , then there is nothing to show. Suppose that $t = \sigma(g)$ for some $\sigma \in \Sigma$ and some map $g: \mathbf{ar}(\sigma) \rightarrow T_\Sigma(\Gamma)$. Then, for all $i \in I$, we have that $(\pi_i \cdot f)^\#(\sigma(g))$ is defined and $A_i \models \alpha((\pi_i \cdot f)^\# \cdot g)$ since each A_i is an algebra in \mathcal{V} . ◀

Proof of Proposition 4.11. It suffices to show that \mathcal{V} is closed under subalgebras and products in $\mathbf{Alg} \Sigma$; this implies that \mathcal{V} is closed under limits in $\mathbf{Alg} \Sigma$. Since \mathcal{V} is closed under κ -directed colimits in $\mathbf{Alg} \Sigma$ by Proposition 4.9, we may then apply Theorem 2.2 to see that \mathcal{V} is a reflective subcategory of $\mathbf{Alg} \Sigma$, as required.

We first show that \mathcal{V} is closed under subalgebras. Given a subalgebra $h: B \hookrightarrow A$ of an algebra A in \mathcal{V} and an axiom $\Gamma \vdash \alpha(f)$, we verify that B satisfies this axiom. To this end, let $e: \Gamma \rightarrow B$ be a relation preserving assignment. Using that A is an algebra in \mathcal{V} , it follows that $(h \cdot e)^\#(f(i))$ is defined for all $i \in \Gamma$, and $A \models \alpha((h \cdot e)^\# \cdot f)$. Since h is an embedding, we may apply Lemma C.2 to see that $e^\#(f(i))$ is defined for all $i \in \text{ar}(\sigma)$ hence also $(h \cdot e)^\# = h \cdot e^\#$. Using the latter, we see that $A \models \alpha(h \cdot e^\# \cdot f)$ whence $B \models \alpha(e^\# \cdot f)$ since h is relation reflecting. It now follows that B satisfies $\Gamma \vdash \alpha(f)$, as desired.

To conclude, we now show that \mathcal{V} is closed under products. Let $A = \prod_{i \in I} A_i$ be a product of algebras in \mathcal{V} with projections $\pi_i: A \rightarrow A_i$; we are going to verify that A lies in \mathcal{V} . To this end, let $\Gamma \vdash \alpha(e)$ be an axiom of \mathcal{V} and let $f: \Gamma \rightarrow A$ be a relation preserving map. Since each A_i is an algebra in \mathcal{V} , we have, for all $i \in I$, that $(\pi_i \cdot f)^\#(e(j))$ is defined for all $j \in \text{ar}(\alpha)$, and $A_i \models \alpha((\pi_i \cdot f)^\# \cdot e)$. By Lemma C.2, we have that $(\pi_i \cdot f)^\# = \pi_i \cdot f^\#$ since each π_i is a homomorphism. By Lemma C.4, $f^\#(e(j))$ is defined for all $j \in \text{ar}(\alpha)$. Then, since the π_i are jointly relation reflecting, we have that $A \models \alpha(f^\# \cdot e)$. Hence A satisfies $\Gamma \vdash \alpha(f)$, as desired. \blacktriangleleft

Proof of Theorem 4.13. We are going use Beck's Monadicity Theorem [17, Thm. IV.7.1]. We have seen in Remark 4.12 that $V: \mathcal{V} \rightarrow \mathbf{C}$ has a left adjoint. Thus, it suffices to show that $V: \mathcal{V} \rightarrow \mathbf{C}$ creates coequalizers of V -split pairs. Let $f, g: A \rightarrow B$ be a V -split pair of homomorphisms in \mathcal{V} . That is, there exist relation preserving maps c, i, j as depicted in the commutative diagram below

$$\begin{array}{ccc}
 VA & \begin{array}{c} \xrightarrow{Vf} \\ \xrightarrow{Vg} \end{array} & VB & \xrightarrow{c} & C \\
 \curvearrowleft & & \curvearrowright & & \\
 & \xrightarrow{j} & & \xrightarrow{i} & \\
 & & & &
 \end{array}
 \quad \text{such that} \quad
 \begin{array}{l}
 c \cdot Vf = c \cdot Vg, \quad Vf \cdot j = \text{id}_{VB}, \\
 c \cdot i = \text{id}_C, \quad Vg \cdot j = i \cdot c.
 \end{array}$$

(This holds in \mathbf{C} , and the equations easily imply that c is an *absolute* coequalizer of Vf and Vg [17, Sec. VI.6], i.e. a coequalizer which is preserved by every functor; subsequently we will omit writing V .) We first show that C carries the structure of a Σ -algebra: indeed, for each $\sigma \in \Sigma$, we define $\sigma_C: [\text{ar}(\sigma), C] \rightarrow C$ by

$$\sigma_C(m) := c \cdot \sigma_B(i \cdot m).$$

Then $\sigma_C(f)$ is defined for all $f \in [\text{ar}(\sigma), A]$ since $i \cdot f \in [\text{ar}(\sigma), B]$, and σ_C is relation preserving because both c and σ_B are relation preserving. Hence $(C, (\sigma_C)_{\sigma \in \Sigma})$ defines a Σ -algebra, as claimed. We next show that $c: B \rightarrow C$ is a homomorphism: given a Γ -ary operation $\sigma \in \Sigma$ and a relation preserving map $m: \Gamma \rightarrow B$, we have

$$\begin{array}{ll}
 c \cdot \sigma_B(m) = c \cdot \sigma_B((f \cdot j) \cdot m) & f \cdot j = \text{id}_B \\
 = c \cdot f \cdot \sigma_A(j \cdot m) & f \text{ a homomorphism} \\
 = c \cdot g \cdot \sigma_A(j \cdot m) & c \cdot f = c \cdot g \\
 = c \cdot \sigma_B(g \cdot j \cdot m) & g \text{ a homomorphism} \\
 = c \cdot \sigma_B(i \cdot c \cdot m) & g \cdot j = i \cdot c \\
 = \sigma_C(c \cdot m) & \text{definition of } \sigma_C
 \end{array}$$

Thus, c is a homomorphism, as claimed. Moreover, the operations σ_C are the unique morphisms $[\mathbf{ar}(\sigma), C] \rightarrow C$ making c a homomorphism. Indeed, if c is a homomorphism and $m: \mathbf{ar}(\sigma) \rightarrow C$ is a relation preserving map, then

$$\sigma_C(m) = \sigma_C(c \cdot i \cdot m) = c \cdot \sigma_B(i \cdot m),$$

where we use that $c \cdot i = \text{id}_B$ in the first equality.

We next show that C is an algebra in \mathcal{V} . Indeed, we will prove that C satisfies every Σ -relation satisfied by B ; the claim will then follow because B is an algebra in \mathcal{V} . To this end, suppose that B satisfies $\Gamma \vdash \alpha(e)$ and let $m: \Gamma \rightarrow C$ be a relation preserving map. Then, the map $i \cdot m: \Gamma \rightarrow B$ is relation preserving whence, for all $l \in \mathbf{ar}(\alpha)$, $(i \cdot m)^\#(e(l))$ is defined and $B \models \alpha((i \cdot m)^\# \cdot e)$. Then, since c is a homomorphism, we know by Lemma C.2 that, for all $l \in \mathbf{ar}(\alpha)$, $(c \cdot i \cdot m)^\#(e(l))$ is defined hence also $m^\#(e(l))$ is defined since $c \cdot i = \text{id}$. Finally, $A \models \alpha(m \cdot e)$. Indeed, we have

$$m^\# = (\text{id} \cdot m)^\# = (c \cdot i \cdot m)^\# = c \cdot (i \cdot m)^\#,$$

and $A \models \alpha(c \cdot (i \cdot m)^\# \cdot e)$ because $B \models \alpha((i \cdot m)^\#)$ and c is relation preserving.

We have now seen that there is a unique Σ -algebra structure on C making c a homomorphism, and C is an algebra in \mathcal{V} . Thus, in order to conclude the proof, it suffices to show that c is a coequalizer of f and g in \mathcal{V} . We already know that c is a coequalizer in \mathbf{C} . Given a homomorphism $d: B \rightarrow D$ such that $d \cdot f = d \cdot g$ we therefore obtain a unique morphism $h: C \rightarrow D$ in \mathbf{C} such that $h \cdot c = d$. To complete the proof we need to show that h is a homomorphism. To see this consider the following diagrams, for every operation symbol σ :

$$d \cdot (-) \left[\begin{array}{ccc} [\mathbf{ar}(\sigma), B] & \xrightarrow{\sigma_B} & B \\ c \cdot (-) \downarrow & & \downarrow c \\ [\mathbf{ar}(\sigma), C] & \xrightarrow{\sigma_C} & C \\ h \cdot (-) \downarrow & & \downarrow h \\ [\mathbf{ar}(\sigma), D] & \xrightarrow{\sigma_D} & D \end{array} \right] d$$

The left-hand and right-hand parts clearly commutes, and the upper square and outside do since c and d are homomorphisms. Thus, the desired lower square commutes when precomposed by $c \cdot (-)$. The latter morphism is epimorphic since it is a coequalizer being the image of the absolute coequalizer c under the internal hom-functor $[\mathbf{ar}(\sigma), -]$. Hence, the desired square commutes. \blacktriangleleft

Proof of Corollary 4.14. We know that the assignment $\mathcal{V} \mapsto \mathbb{T}_{\mathcal{V}}$ of a κ -ary relational algebraic theory to its free-algebra monad \mathbb{T} preserves categories of models by Theorem 4.13, and $\mathbb{T}_{\mathcal{V}}$ is κ -accessible, as discussed in Remark 4.12. To conclude, we must show that $\mathbb{T}_{\mathcal{V}}$ is enriched. That is, given $\alpha \in \Pi$ and $(f_i: X \rightarrow Y)_{i \in \mathbf{ar}(\alpha)}$ such that $[X, Y] \models \alpha(f_i)$, we must show that $[T_{\mathcal{V}}X, T_{\mathcal{V}}Y] \models \alpha(T_{\mathcal{V}}f_i)$.

To this end, let $\alpha \in \Pi$ and $f_i: X \rightarrow Y$ be as above, and let $e: E \hookrightarrow T_{\mathcal{V}}X$ be the substructure of $T_{\mathcal{V}}X$ given by all $t \in T_{\mathcal{V}}X$ such that $T_{\mathcal{V}}Y \models \alpha(T_{\mathcal{V}}f_i(t))$. We first show that $\eta_X(x) \in E$ for all $x \in X$. We know that $[X, Y] \models \alpha(f_i)$, and this means that $Y \models \alpha(f_i(x))$ for all $x \in X$. Since $\eta_Y: Y \rightarrow T_{\mathcal{V}}Y$ is relation preserving, it follows that $T_{\mathcal{V}}Y \models \alpha(\eta_Y \cdot f_i(x))$. By naturality of η , the square below is commutative

$$\begin{array}{ccc} X & \xrightarrow{\eta_X} & T_{\mathcal{V}}X \\ f_i \downarrow & & \downarrow T_{\mathcal{V}}f_i \\ Y & \xrightarrow{\eta_Y} & T_{\mathcal{V}}Y \end{array}$$

whence $T_{\mathcal{V}}Y \models \alpha(T_{\mathcal{V}}f_i(\eta_X(x)))$. That is, $\eta_X(x) \in E$, as claimed.

Furthermore, E is closed under the operations of the algebra $T_{\mathcal{V}}X$ because each operation $\sigma_{T_{\mathcal{V}}X}: [\mathbf{ar}(\sigma), T_{\mathcal{V}}X] \rightarrow T_{\mathcal{V}}X$ is relation preserving. Indeed, let $\sigma \in \Sigma$ and let $h: \mathbf{ar}(\sigma) \rightarrow T_{\mathcal{V}}X$ be a relation preserving map such that $h[\mathbf{ar}(\sigma)] \subseteq E$; we are going to verify that $\sigma_{T_{\mathcal{V}}X}(h) \in E$. Note that we have the commutative square

$$\begin{array}{ccc} [\mathbf{ar}(\sigma), T_{\mathcal{V}}X] & \xrightarrow{\sigma_{T_{\mathcal{V}}X}} & T_{\mathcal{V}}X \\ T_{\mathcal{V}}f_i \cdot (-) \downarrow & & \downarrow T_{\mathcal{V}}f_i \\ [\mathbf{ar}(\sigma), T_{\mathcal{V}}Y] & \xrightarrow{\sigma_{T_{\mathcal{V}}Y}} & T_{\mathcal{V}}Y \end{array}$$

Thus, for the map $h: \mathbf{ar}(\sigma) \rightarrow T_{\mathcal{V}}X$, we have that

$$T_{\mathcal{V}}f_i(\sigma_{T_{\mathcal{V}}X}(h)) = \sigma_Y(T_{\mathcal{V}}f_i \cdot h) \quad \text{for every } i \in \mathbf{ar}(\alpha).$$

Now, using that $h[\mathbf{ar}(\sigma)] \subseteq E$, it follows that $T_{\mathcal{V}}Y \models \alpha(T_{\mathcal{V}}f_i(h(j)))$ for all $j \in \mathbf{ar}(\sigma)$. Thus, by definition of $\mathbf{E}([\mathbf{ar}(\sigma), T_{\mathcal{V}}Y])$, we have that $[\mathbf{ar}(\sigma), T_{\mathcal{V}}Y] \models \alpha(T_{\mathcal{V}}f_i \cdot h)$. This implies that $T_{\mathcal{V}}Y \models \alpha(\sigma_{T_{\mathcal{V}}Y}(T_{\mathcal{V}}f_i \cdot h))$ since $\sigma_{T_{\mathcal{V}}Y}$ is relation preserving. Applying the commutative square above, it follows that $T_{\mathcal{V}}Y \models \alpha(T_{\mathcal{V}}f_i \cdot \sigma_{T_{\mathcal{V}}X}(h))$ hence also $\sigma_{T_{\mathcal{V}}X}(h) \in E$, as claimed.

We conclude that E is a Σ -subalgebra of $T_{\mathcal{V}}X$ containing $\eta_X[X]$. Since $T_{\mathcal{V}}X$ is a free algebra of \mathcal{V} with the universal morphism η_X , it follows that $E = T_{\mathcal{V}}X$. In particular, $T_{\mathcal{V}}Y \models \alpha(T_{\mathcal{V}}f_i(t))$ holds for all $t \in T_{\mathcal{V}}X$, whence $\alpha(T_{\mathcal{V}}f_i)$ holds in $[T_{\mathcal{V}}X, T_{\mathcal{V}}Y]$, as desired. \blacktriangleleft

Relational Logic

Proof of Lemma 4.16. Admissibility of (Arity): Admissibility of the rule (Arity) is immediate: in any derivation of $X \vdash \downarrow\sigma(m)$, the last rule applied was (E-Ar). We may apply this rule just in case we have derivations of all desired relational judgements $X \vdash \alpha(f)$.

Admissibility of (Subterm): We proceed to show that whenever $X \vdash \alpha(f)$ is derivable, where $f: \mathbf{ar}(\alpha) \rightarrow T_{\Sigma}(X)$, then $X \vdash \downarrow u$ is derivable for all $u \in \mathbf{sub}(f)$; we do so by making a case distinction on the basis of whether $u \in \mathbf{sub}(f)$ is a variable (in X) or $u = \sigma(m)$ for some $\sigma \in \Sigma$ and some map $m: |\mathbf{ar}(\sigma)| \rightarrow T_{\Sigma}(X)$. If u is a variable, then $X \vdash \downarrow u$ is derivable via (Var). So let us suppose that $u = \sigma(m)$. Then, by application of (Arity) (which was shown to be admissible above), we have a derivation of $X \vdash \beta(m \cdot g)$ for all edges $\beta(g)$ such that $\mathbf{ar}(\sigma) \models \beta(g)$. Thus, by application of (E-Ar), we have a derivation of $X \vdash \downarrow\sigma(m)$, as desired. \blacktriangleleft

► Corollary C.5. *Let $\sigma \in \Sigma$ and let $m: |\mathbf{ar}(\sigma)| \rightarrow T_{\Sigma}(\Gamma)$. Then $\Gamma \vdash \downarrow\sigma(m)$ iff $\Gamma \vdash \alpha(m \cdot f)$ is derivable for all edges $\alpha(f)$ such that $\mathbf{ar}(\sigma) \models \alpha(f)$.*

Proof. Suppose that $\Gamma \vdash \downarrow\sigma(m)$ is derivable. Then $\Gamma \vdash \alpha(m \cdot f)$ is derivable for all edges $\alpha(f)$ in $|\mathbf{ar}(\sigma)|$ such that $\mathbf{ar}(\sigma) \models \alpha(f)$ by application of (Arity); this rule is admissible by Lemma 4.16. Conversely, if $\Gamma \vdash \alpha(m \cdot f)$ is derivable for all edges $\alpha(f)$ such that $\mathbf{ar}(\sigma) \models \alpha(f)$, then $\Gamma \vdash \downarrow\sigma(m)$ is derivable via (E-Ar), as desired. \blacktriangleleft

In **Remark 4.15** we have suggested that the relational logic enjoys an admissible substitution rule. We substantiate this claim now:

► Proposition C.6. *Let $X, Y \in \mathbf{C}$ and let $\tau: Y \rightarrow T_{\Sigma}(X)$ be a substitution. The following rules are admissible:*

$$\text{(SubsR)} \quad \frac{\{X \vdash \alpha(\tau \cdot f) \mid Y \models \alpha(f)\} \cup \{X \vdash \downarrow\tau(y) \mid y \in Y\}}{X \vdash \beta(\tau \cdot g)} \quad \frac{Y \vdash \beta(g)}{g: \mathbf{ar}(\beta) \rightarrow T_{\Sigma}(Y)} \quad (\beta \in \Pi)$$

$$\text{(SubsD)} \quad \frac{\{X \vdash \alpha(\tau \cdot f) \mid Y \models \alpha(f)\} \cup \{X \vdash \downarrow\tau(y) \mid y \in Y\} \quad Y \vdash \downarrow t}{X \vdash \downarrow\tau(t)} \quad (t \in T_\Sigma(Y))$$

Proof. Suppose that all premisses in $\{X \vdash \alpha(\tau \cdot f) \mid Y \models \alpha(f)\} \cup \{X \vdash \downarrow\tau(y) \mid y \in Y\}$ are derivable. We prove by simultaneous induction on derivations that whenever $Y \vdash \beta(g)$ is derivable, then $X \vdash \beta(\tau \cdot g)$ is derivable, and whenever $Y \vdash \downarrow t$ is derivable, then $X \vdash \downarrow\tau(t)$ is derivable.

As a base case, observe that if $Y \vdash \beta(g)$ was derived using (Ctx), then $Y \models \beta(g)$ whence $X \vdash \beta(\tau \cdot g)$ is derivable by assumption. Similarly, if $Y \models \downarrow y$ was derived using (Var), then $X \vdash \downarrow\tau(y)$ is derivable by assumption. As for the inductive step, we first note that corresponding to (l-Ar) is analogous to the case for (Ax) shown further below; the remaining cases are as follows.

(RelAx) Suppose that $Y \vdash \beta(g)$, where $g = \tau' \cdot f$, $\Phi \implies \beta(f)$ is some relational axiom and $\tau' : \text{Var} \rightarrow T_\Sigma(Y)$, is was derived in the last step from premisses:

- (a) $X \vdash \tau' \cdot \varphi$ for all $\varphi \in \Phi$;
- (b) $\downarrow\tau'(f(i))$ for all $i \in \text{ar}(\alpha)$.

By the induction hypothesis, $Y \vdash \tau \cdot \tau' \cdot \varphi$ for $\varphi \in \Phi$ and $Y \vdash \downarrow\tau(\tau'(f(i)))$ for $i \in \text{ar}(\alpha)$ are derivable. Thus $Y \vdash \beta(\tau \cdot \tau' \cdot f)$ is derivable by an application of the (RelAx) rule.

(E-Ar): Suppose that $Y \vdash \downarrow\sigma(f)$, for $\sigma \in \Sigma$ and $f : \text{ar}(\sigma) \rightarrow T_\Sigma(Y)$, was derived in the last step from premisses $Y \vdash \alpha(f \cdot g)$ for all $\alpha(g) \in \text{ar}(\sigma)$ and $X \vdash \downarrow f(i)$ for all $i \in \text{ar}(\alpha)$. By induction, $X \vdash \alpha(\tau \cdot f \cdot g)$ is derivable for all $\alpha(g) \in \text{ar}(\sigma)$, and $X \vdash \downarrow(\tau \cdot f)(i)$ is derivable for all $i \in \text{ar}(\sigma)$. Hence $X \vdash \downarrow\sigma(\tau \cdot f)$ is derivable via (E-Ar), and since $\tau(\sigma(f)) = \sigma(\tau \cdot f)$ we are done.

(Mor): Suppose that $Y \vdash \beta(g)$, where $g : \text{ar}(\beta) \rightarrow T_\Sigma(Y)$ has the form of an assignment $g(i) = \sigma(f_i)$ for some $\sigma \in \Sigma$ and $f_i : \text{ar}(\sigma) \rightarrow T_\Sigma(Y)$ for $i \in \text{ar}(\beta)$, was derived from premisses

- (a) $Y \vdash \beta(f_i(j))$ for all $j \in \text{ar}(\sigma)$;
- (b) $Y \vdash \downarrow\sigma(f_i)$ for all $i \in \text{ar}(\beta)$;

Applying the inductive hypothesis to these items, it follows immediately that $X \vdash \beta(\tau \cdot g)$ is derivable via (Mor).

(Ax): Finally, suppose that $Y \vdash \beta(\kappa \cdot k)$ was derived via (Ax) using an axiom $\Gamma \vdash \beta(k) \in \mathcal{E}$ and a substitution $\kappa : \Gamma \rightarrow T_\Sigma(Y)$ from premisses $Y \vdash \alpha(\kappa \cdot f)$ for all $\alpha(f) \in \mathbf{E}(\Gamma)$ and $Y \vdash \downarrow\kappa(z)$ for all $z \in \Gamma$. By induction, $X \vdash \alpha(\tau \cdot \kappa \cdot f)$ is derivable for all $\alpha(f) \in \mathbf{E}(\Gamma)$, and $X \vdash \downarrow(\tau \cdot \kappa)(z)$ is derivable for all $z \in \Gamma$. Thus, by application of (Ax), we have a derivation of $X \vdash \beta(\tau \cdot \kappa \cdot k)$. \blacktriangleleft

Constructing free algebras

- **Lemma C.7.** (1) *The relational structure on $\mathcal{F}X$ is independent of the choice of u .*
- (2) *$\sigma_{\mathcal{F}X}$ is defined on $[\text{ar}(\sigma), \mathcal{T}_v(X)/\sim]$ and independent of the choice of the splitting u .*

Proof. (1) Given another splitting v of q , we need to prove that $\mathcal{T}_v(X) \models \alpha(u \cdot f)$ iff $\mathcal{T}_v(X) \models \alpha(v \cdot f)$ for every $f : \text{ar}(\alpha) \rightarrow \mathcal{F}X$. Equivalently, $X \vdash \alpha(u \cdot f)$ is derivable iff so is $X \vdash \alpha(v \cdot f)$. Since u, v are splittings of q we have that for each $i \in \text{ar}(\alpha)$, $X \vdash \varphi(u \cdot g(i), v \cdot g(i))$ is derivable for every $\varphi(x, y) \in Eq(x, y)$. Since by Assumption 3.4, \mathcal{A} explicitly includes axioms stating that all relations are closed under $Eq(-, -)$, the claim follows by an application of the (RelAx) rule in each direction.

(2) For the first claim, we show that $\sigma(u \cdot g)$ is defined for all $g \in [\text{ar}(\sigma), \mathcal{F}X]$. Since g is relation preserving, it follows that for every edge $\alpha(f)$ such that $\text{ar}(\sigma) \models \alpha(f)$ we have

$\mathcal{F}X \models \alpha(u \cdot g \cdot f)$. That is, $X \vdash \alpha(u \cdot g \cdot f)$ is derivable. Hence $X \vdash \downarrow\sigma(u \cdot g)$ is derivable via (E-Ar), which implies $\sigma(u \cdot g) \in \mathcal{T}_{\mathcal{V}}(X)$.

We now show that $[\sigma(u \cdot g)]$ is independent of the choice of u . To this end, suppose that we are given another splitting v , and let $\varphi(x, y) \in Eq(x, y)$. By definition of \sim , it suffices to show that $X \vdash \varphi(\sigma(u \cdot g), \sigma(v \cdot g))$ is derivable. Since u, v are splittings of g , we have that for each $i \in \text{ar}(\sigma)$, $X \vdash \varphi(u \cdot g(i), v \cdot g(i))$ is derivable. The stated goal $X \vdash \varphi(\sigma(u \cdot g), \sigma(v \cdot g))$ follows by rule (Mor). \blacktriangleleft

► **Remark C.8.** The set $\mathcal{T}_{\mathcal{V}}(X)$ is a Σ -algebra under the operations given by the usual term formation: for every operation symbol σ in Σ we have

$$\sigma_{\mathcal{T}_{\mathcal{V}}(X)}: [\text{ar}(\sigma), \mathcal{T}_{\mathcal{V}}(X)] \rightarrow \mathcal{T}_{\mathcal{V}}(X) \quad \text{given by} \quad f \mapsto \sigma(f).$$

Indeed, for a relation preserving map $f: \text{ar}(\sigma) \rightarrow \mathcal{T}_{\mathcal{V}}(X)$ we have $X \vdash \alpha(f \cdot g)$ for every $\alpha(g) \in \text{ar}(\sigma)$. Thus $X \vdash \downarrow\sigma(f)$ by an application of the (E-Ar) rule.

► **Corollary C.9.** *The quotient map $q: \mathcal{T}_{\mathcal{V}}(X) \rightarrow \mathcal{F}X$ is a relation preserving Σ -algebra homomorphism.*

Proof. To see that q relation preserving, suppose $\mathcal{T}_{\mathcal{V}}(X) \models \alpha(f)$ for $f: \text{ar}(\alpha) \rightarrow \mathcal{T}_{\mathcal{V}}(X)$. By Lemma C.7(1), we may assume that the splitting u is chosen such that $u \cdot q \cdot f = f$. Therefore $\mathcal{T}_{\mathcal{V}}(X) \models \alpha(u \cdot q \cdot f)$ which gives $\mathcal{F}X \models \alpha(q \cdot f)$ as desired.

Given σ in Σ and $g: \text{ar}(\sigma) \rightarrow \mathcal{T}_{\mathcal{V}}(X)$ relation preserving, we may assume by Lemma C.7(2) that the splitting $u: \mathcal{F}X \rightarrow \mathcal{T}_{\mathcal{V}}(X)$ is chosen such that $u \cdot q \cdot g = g$. Then we have

$$\begin{aligned} q \cdot \sigma_{\mathcal{T}_{\mathcal{V}}(X)}(g) &= q(\sigma(g)) && \text{def. of } \sigma_{\mathcal{T}_{\mathcal{V}}(X)} \\ &= q(\sigma(u \cdot q \cdot g)) && \text{since } g = u \cdot q \cdot g \\ &= \sigma_{\mathcal{F}X}(q \cdot g). && \text{def. of } \sigma_{\mathcal{F}X} \end{aligned} \quad \blacktriangleleft$$

Proof of Theorem 4.18. Indeed, let $\Delta \vdash \beta(g)$ be an axiom of \mathcal{V} , where $g: \text{ar}(\beta) \rightarrow T_{\Sigma}(\Delta)$, and let $e: \Delta \rightarrow \mathcal{F}X$ be a relation preserving map; we are going to show that $e^{\#}(g(i))$ is defined for all $i \in \text{ar}(\beta)$ and that $\mathcal{F}X \models \beta(e^{\#} \cdot g)$. (This implies that $\mathcal{F}X$ satisfies $\Delta \vdash \beta(g)$, as required.) To this end, define a substitution $\tau: \Delta \rightarrow \mathcal{T}_{\mathcal{V}}(X)$ (again note $\mathcal{T}_{\mathcal{V}}(X) \subseteq T_{\Sigma}(X)$) as the map

$$\tau = |\Delta| \xrightarrow{e} \mathcal{F}X \xrightarrow{u} \mathcal{T}_{\mathcal{V}}(X).$$

For all edges $\alpha(f) \in \Delta$, we have that $\mathcal{F}X \models \alpha(e \cdot f)$ because e is relation-preserving. Equivalently, $\mathcal{T}_{\mathcal{V}}(X) \models \alpha(u \cdot e \cdot f)$, which means that $X \vdash \alpha(\tau \cdot f)$ is derivable. Hence also $X \vdash \beta(\tau \cdot g)$ is derivable via (Ax). Since q is relation preserving by Corollary C.9 we have $\mathcal{F}X \models \beta(q \cdot \tau \cdot g)$. We shall show below that for all r in $\text{sub}(g)$ we have that both $q \cdot \tau(r)$ and $e^{\#}(r)$ are defined and

$$q \cdot \tau(r) = e^{\#}(r). \tag{C.1}$$

(Recall from Definition 4.4 that we write τ also for the extension $\bar{\tau}: T_{\Sigma}(\Delta) \rightarrow T_{\Sigma}(X)$.) Thus, in particular $e^{\#}(g(i))$ is defined for all $i \in \text{ar}(\beta)$, and we obtain $\mathcal{F}X \models \beta(e^{\#} \cdot g)$ as desired.

To prove (C.1) we proceed by induction on r . For a variable x in $\text{sub}(g)$ we clearly have that both $q \cdot \tau$ and $e^{\#}$ are defined in x and

$$q \cdot \tau(x) = q \cdot u \cdot e(x) = e(x) = e^{\#}(x),$$

using the definitions of τ and $e^\#$ and that $q \cdot u = \text{id}$. For a term $\sigma(k)$ in $\text{sub}(g)$ we know by induction that both $q \cdot \tau(k(i))$ and $e^\#(k(i))$ are defined and $q \cdot \tau(k(i)) = e^\#(k(i))$ for all $i \in \text{ar}(\sigma)$. Then $\tau(\sigma(k))$ is a provably defined term, i.e. it lies in $\mathcal{T}_V(X)$, so that q is defined on it; indeed, since $X \vdash \beta(\tau \cdot g)$ is derivable, this follows from an application of the (Subterm) rule in Lemma 4.16 since $\sigma(k)$ in $\text{sub}(g)$ yields that $\tau(\sigma(k))$ lies in $\text{sub}(\tau \cdot g)$. Now we obtain

$$\begin{aligned} q \cdot \tau(\sigma(k)) &= q(\sigma(\tau \cdot k)) && \text{def. of } \tau = \bar{\tau} \\ &= \sigma_{\mathcal{F}X}(q \cdot \tau \cdot k) && \text{by Corollary C.9} \\ &= \sigma_{\mathcal{F}X}(e^\# \cdot k) && \text{by induction} \\ &= e^\#(\sigma(k)) && \text{def. of } e^\#. \end{aligned}$$

$\mathcal{F}X$ is free on X : We define $\eta_X: X \rightarrow \mathcal{F}X$ by $\eta_X(x) = [x]$ and prove that this is a universal map. First observe that η is clearly relation preserving since for every $X \models \alpha(f)$ we have $\mathcal{F}X \models \alpha(\eta_X \cdot f)$ by the rule (Ctx). Now given $f: X \rightarrow A$ relation preserving we define $\bar{f}: \mathcal{F}X \rightarrow A$ by

$$\bar{f}([t]) = f^\#(t) \quad \text{for all } t \in \mathcal{T}_V(X).$$

Then \bar{f} is well-defined by soundness (proved below), and we clearly have $\bar{f} \cdot \eta_X = f$.

(1) We prove that \bar{f} is relation preserving. Suppose that $\mathcal{F}X \models \alpha(g)$. Equivalently we have $\mathcal{T}_V(X) \models \alpha(u \cdot g)$, which in turn means that $X \vdash \alpha(u \cdot g)$ is derivable. By soundness, A satisfies $X \vdash \alpha(u \cdot g)$. In particular, we have $A \models \alpha(f^\# \cdot u \cdot g)$. Then we obtain $A \models \alpha(\bar{f} \cdot g)$ as desired since

$$f^\# \cdot u = \bar{f} \cdot q \cdot u = \bar{f}$$

because $f^\# \cdot q = \bar{f}$ by the definition of \bar{f} .

(2) Next we show that \bar{f} is Σ -algebra homomorphism. Indeed, for every operation symbol σ we show that the following square commutes:

$$\begin{array}{ccc} [\text{ar}(\sigma), \mathcal{F}X] & \xrightarrow{\sigma_{\mathcal{F}X}} & \mathcal{F}X \\ \bar{f} \cdot (-) \downarrow & & \downarrow \bar{f} \\ [\text{ar}(\sigma), A] & \xrightarrow{\sigma_A} & A \end{array}$$

Indeed, for every g in $[\text{ar}(\sigma), \mathcal{F}X]$ we have

$$\begin{aligned} \bar{f}(\sigma_{\mathcal{F}X}(g)) &= \bar{f}([\sigma(u \cdot g)]) && \text{def. of } \sigma_{\mathcal{F}X} \\ &= f^\#(\sigma(u \cdot g)) && \text{def. of } \bar{f} \\ &= \sigma_A(f^\# \cdot u \cdot g) && \text{def. of } f^\# \\ &= \sigma_A(\bar{f} \cdot q \cdot u \cdot g), && \text{since } \bar{f} \cdot q = f^\# \text{ by def.} \\ &= \sigma_A(\bar{f} \cdot g) && \text{since } q \cdot u = \text{id}. \end{aligned}$$

(3) Finally, we show that \bar{f} is unique. Indeed this follows from the required homomorphism property and the required extension property $\bar{f} \cdot \eta_X = f$, since by construction every element of $\mathcal{F}X$ is denoted by a well-founded term formed from the elements of X using the operations of Σ (formally, use well-founded induction on terms). \blacktriangleleft

► Lemma C.10. *Let $t \in T_\Sigma(X)$. Then, for the universal map $\eta_X: X \rightarrow \mathcal{F}X$, if $\eta_X^\#(t)$ is defined, then $t \in \mathcal{T}_V(X)$ and $\eta_X^\#(t) = [t]$.*

The latter means that $\eta^\#$ is the canonical quotient map $q: \mathcal{T}_\mathcal{V}(X) \rightarrow \mathcal{F}X$.

Proof. By induction on terms. If t is a variable in X , then $X \vdash \downarrow t$ is derivable via (Var) and $\eta_X^\#(t) = \eta_X(t) = [t]$. Inductively assume that the lemma holds for all $s \in \text{sub}(t)$, and let $t = \sigma(m)$ for some $\sigma \in \Sigma$ and some $m: \text{ar}(\sigma) \rightarrow T_\Sigma(X)$ such that $\eta_X^\#(\sigma(m))$ is defined. Then:

- (1) $\eta_X^\#(m(i))$ is defined for all $i \in \text{ar}(\sigma)$;
- (2) $\mathcal{F}X \models \beta(\eta_X^\# \cdot m \cdot g)$ for all $g: \text{ar}(\beta) \rightarrow \text{ar}(\sigma)$ such that $\text{ar}(\sigma) \models \beta(g)$.

Combining the inductive hypothesis with (1), we have that $X \vdash \downarrow m(i)$ for all $i \in \text{ar}(\sigma)$, and $\eta^\#(m(i)) = [m(i)]$; that is, $\eta_X^\# \cdot m = q \cdot m$.

In particular, by (2), we have that $\mathcal{F}X \models \beta(u \cdot \eta_X^\# \cdot m \cdot g)$ for all $g: \text{ar}(\beta) \rightarrow \text{ar}(\sigma)$ such that $\text{ar}(\sigma) \models \beta(g)$ and where $u: \mathcal{F}X \rightarrow \mathcal{T}_\mathcal{V}(X)$ is a splitting of the canonical quotient map q . By Lemma C.7(2), we may assume that u is chosen such that $u \cdot q \cdot m = m$. Now, we have

$$u \cdot \eta_X^\# \cdot m \cdot g = u \cdot q \cdot m \cdot g = m \cdot g,$$

whence $\mathcal{F}X \models \beta(m \cdot g)$. This means that $X \vdash \beta(m \cdot g)$ is derivable for all $\beta(g) \in \text{ar}(\sigma)$. Hence $X \vdash \downarrow \sigma(m)$ is derivable via (E-Ar), which means that $\sigma(m)$ lies in $\mathcal{T}_\mathcal{V}(X)$. Moreover, we have

$$\begin{aligned} \eta_X^\#(\sigma(m)) &= \sigma_{\mathcal{F}X}(\eta_X^\# \cdot m) && \text{def. of } \eta_X^\# \\ &= \sigma_{\mathcal{F}X}(q \cdot m) && \text{induction hypothesis} \\ &= q(\sigma(m)) && \text{by Corollary C.9} \\ &= [\sigma(m)]. \end{aligned} \quad \blacktriangleleft$$

► **Notation C.11.** Let A be a Σ -algebra, let $\tau: X \rightarrow T_\Sigma(Y)$ be a substitution, and let $e: X \rightarrow A$ be an assignment such that $e^\# \cdot \tau(x)$ is defined for all $x \in X$. We define

$$e_\tau = (X \xrightarrow{\tau} T_\Sigma(Y) \xrightarrow{e^\#} A).$$

► **Lemma C.12 (Substitution lemma).** Let A be a Σ -algebra in \mathcal{V} , let $\tau: X \rightarrow T_\Sigma(Y)$ be a substitution, and let $e: Y \rightarrow A$ be an assignment such that $e^\#(\tau(x))$ is defined for all $x \in X$. Then, for all $t \in T_\Sigma(X)$:

$$e^\#(\bar{\tau}(t)) \text{ is defined} \quad \text{iff} \quad e_\tau^\#(t) \text{ is defined};$$

in this case, $e^\#(\bar{\tau}(t)) = e_\tau^\#(t)$.

Proof. By induction on terms. If $t = x \in X$, then

$$\begin{aligned} e^\# \cdot \bar{\tau}(t) &= e^\# \cdot \tau(x) && \text{def. of } \bar{\tau} \\ &= e_\tau(x) && \text{def. of } e_\tau \\ &= e_\tau^\#(t) && \text{def. of } e_\tau^\#, \end{aligned}$$

with the reading of the equational steps including that the left side is defined iff the right side is.

Next, suppose that $t = \sigma(f)$ for some $\sigma \in \Sigma$ and some map $f: |\text{ar}(\sigma)| \rightarrow T_\Sigma(X)$. We want to show that $e^\#(\bar{\tau}(\sigma(f))) = e^\#(\sigma(\bar{\tau} \cdot f))$ is defined iff $e_\tau^\#(\sigma(f))$ is defined.

Observe that if $e^\#(\sigma(\bar{\tau} \cdot f))$ is defined, then by definition of $e^\#$ we have

- (1) $e^\#(\bar{\tau}(f(i)))$ is defined for all $i \in \text{ar}(\sigma)$;
- (2) $A \models \alpha(e^\# \cdot \bar{\tau} \cdot (f \cdot g))$ for all $\alpha(g) \in \text{ar}(\sigma)$.

Thus, by the inductive hypothesis, we have that $e_\tau^\#(f(i))$ is defined for all $i \in \text{ar}(\sigma)$, and $A \models \alpha(e_\tau^\# \cdot (f \cdot g))$ for all $\alpha(g) \in \text{ar}(\sigma)$. Thus, $e_\tau^\#(\sigma(f))$ is defined. The converse direction follows completely analogously (indeed, simply note that each implication is reversible). We conclude the proof with a computation in which all expressions are equi-defined:

$$\begin{aligned}
e_\tau^\#(\bar{\tau}(\sigma(f))) &= e_\tau^\#(\sigma(\bar{\tau} \cdot f)) && \text{def. of } \bar{\tau} \\
&= \sigma_A(e_\tau^\# \cdot \bar{\tau} \cdot f) && \text{def. of } e_\tau^\# \\
&= \sigma_A(e_\tau^\# \cdot f) && \text{induction hypothesis} \\
&= e_\tau^\#(\sigma(f)) && \text{def. of } e_\tau^\# \quad \blacktriangleleft
\end{aligned}$$

► **Corollary C.13.** *Let A, τ , and e be as in Lemma C.12 above. Then, for all $\alpha \in \Pi$ and all maps $f: |\text{ar}(\alpha)| \rightarrow T_\Sigma(X)$,*

$$A \models \alpha(e^\# \cdot \bar{\tau} \cdot f) \quad \text{iff} \quad A \models \alpha(e_\tau^\# \cdot f).$$

Proof of Theorem 4.19. Soundness (\Rightarrow): Let A be an algebra in \mathcal{V} and let $e: X \rightarrow A$ be a relation preserving assignment. We are going to show by simultaneous induction on derivations that

- (1) if $X \vdash \alpha(f)$ is derivable, then $e^\# \cdot f(i)$ is defined for all $i \in \text{ar}(\alpha)$, and $A \models \alpha(e^\# \cdot f)$;
- (2) if $X \vdash \downarrow t$ is derivable, then $e^\#(t)$ is defined.

For the base case of our inductive proof, first observe that if $X \vdash \alpha(f)$ is derivable via (Ctx), then each $f(i)$ is a variable, so $e^\# \cdot f(i) = e \cdot f(i)$ is defined, and moreover $X \models \alpha(f)$, so $A \models \alpha(e \cdot f)$ since $e: X \rightarrow A$ is relation preserving. Second, if $X \vdash \downarrow t$ is derivable via (Var), then t is a variable in X so that $e^\#(t) = e(t)$ is defined. We proceed by case distinction on the last rule applied in the derivation.

(RelAx): this follows from the fact that the given algebra A is carried by an \mathcal{H} -model, in combination with Lemma C.12 and Corollary C.13.

(E-Ar): Suppose that $X \vdash \downarrow t$ is derived via the rule (E-Ar) so that $t = \sigma(h)$ for some $\sigma \in \Sigma$ and some map $h: \text{ar}(\sigma) \rightarrow T_\Sigma(X)$. Then, by induction, we are given that $e^\#(h(i))$ is defined for all $i \in \text{ar}(\sigma)$, and $A \models \alpha(e^\# \cdot (h \cdot g))$ for all $\alpha(g) \in \text{ar}(\sigma)$. It follows immediately from the definition of $e^\#$ that $e^\#(\sigma(h))$ is defined, as required.

(Mor): Suppose that $X \vdash \alpha(f)$ is derived using the rule (Mor). Then, for some $\sigma \in \Sigma$, we have that $f(i) = \sigma(g_i)$ for all i , where $g_i: \text{ar}(\sigma) \rightarrow T_\Sigma(X)$, and the premisses

$$X \vdash \alpha(g_i(j)) \quad (j \in \text{ar}(\sigma)) \quad \text{and} \quad X \vdash \downarrow \sigma(g_i) \quad (i \in \text{ar}(\alpha))$$

of (Mor) are derivable. Then, by induction, $e^\#(\sigma(g_i))$ is defined for all $i \in \text{ar}(\alpha)$. By definition of $e^\#$, this means that $A \models \beta(e^\# \cdot (g_i \cdot k))$ for all $\beta(k) \in \text{ar}(\sigma)$; that is, $e^\# \cdot g_i \in [\text{ar}(\sigma), A]$ for all $i \in \text{ar}(\alpha)$. Moreover, by the induction hypothesis applied to the first item, we have that $A \models \alpha(e^\# \cdot g_i(j))$ for all $j \in \text{ar}(\sigma)$. In other words, $A \models \alpha(\pi_j \cdot (e^\# \cdot g_i))$ for all $j \in \text{ar}(\sigma)$ (recall that $\pi_j(e^\# \cdot g_i) = e^\# \cdot g_i(j)$); that is, $[\text{ar}(\sigma), A] \models \alpha(e^\# \cdot g_i)$. Now, using that σ_A is a relation-preserving map $[\text{ar}(\sigma), A] \rightarrow A$, it follows that $A \models \alpha(\sigma_A(e^\# \cdot g_i))$. But $\sigma_A(e^\# \cdot g_i) = e^\#(\sigma(g_i)) = e^\# \cdot f(i)$ for all $i \in \text{ar}(\alpha)$, so we have that $A \models \alpha(e^\# \cdot f)$, as desired.

(Ax): Now suppose that $X \vdash \alpha(f)$ is derivable via (Ax) so that for some axiom $\Delta \vdash \alpha(h)$, where $h: \text{ar}(\alpha) \rightarrow T_\Sigma(\Delta)$, and some substitution $\tau: \Delta \rightarrow T_\Sigma(X)$ we have $f = \tau \cdot h$. The premisses of this rule have the following shape:

- (1) $X \vdash \downarrow \tau(y)$ for all $y \in \Delta$;
- (2) $X \vdash \beta(\tau \cdot g)$ for all $\beta(g) \in \Delta$.

By the inductive hypothesis, derivability of these premisses implies that $e^\# \cdot \tau(y)$ is defined for all $y \in \Delta$, and the assignment $e_\tau = e^\# \cdot \tau: \Delta \rightarrow A$ is relation-preserving. Since A , being in \mathcal{V} , satisfies $\Delta \vdash \alpha(h)$, it follows that $A \models \alpha(e_\tau^\# \cdot h)$. By Corollary C.13, we obtain $A \models \alpha(e^\# \cdot \tau \cdot h)$, i.e. $A \models \alpha(e^\# \cdot f)$ as required.

(l-Ar): Finally, assume that $X \vdash \alpha(f)$ is derived using the rule (l-Ar). Then, for some substitution $\tau: \Delta \rightarrow T_\Sigma(X)$ and some \mathcal{V} -axiom $\Delta \vdash \beta(g)$, there exists $\sigma(h) \in \text{sub}(g)$ and $\alpha(k) \in \text{ar}(\sigma)$ such that

$$f = \text{ar}(\alpha) \xrightarrow{k} \text{ar}(\sigma) \xrightarrow{h} T_\Sigma(\Delta) \xrightarrow{\tau} T_\Sigma(X),$$

and the premisses

- (1) $X \vdash \gamma(\tau \cdot c)$ for all $\gamma(c) \in \Delta$;
- (2) $X \vdash \downarrow \tau(y)$ for all $y \in \Delta$

of (l-Ar) are derivable. Just as for the rule (Ax), the induction hypothesis implies that $e^\# \cdot \tau(x)$ is defined for all $x \in X$, and $e_\tau = e^\# \cdot \tau: X \rightarrow A$ is a relation-preserving assignment. Since A lies in \mathcal{V} , we know that A satisfies $\Delta \vdash \beta(g)$ so that $e_\tau^\#(g(i))$ is defined for all $i \in \text{ar}(\beta)$ and $A \models \beta(e_\tau^\# \cdot g)$. Since $\sigma(h) \in \text{sub}(g(i))$ for some $i \in \text{ar}(\beta)$, it follows from a routine induction on terms that $e_\tau^\#(\sigma(h))$ is defined (i.e. we use that definedness is inherited by subterms). Unwinding the definition of $e_\tau^\#$, we obtain that $e_\tau^\#(h(i))$ is defined for all $i \in \text{ar}(\sigma)$, and $A \models \gamma(e_\tau^\# \cdot h \cdot c)$ for all $\gamma(c) \in \text{ar}(\sigma)$. In particular, $A \models \alpha(e_\tau^\# \cdot h \cdot k)$. By Corollary C.13, this implies that $A \models \alpha(e^\# \cdot \tau \cdot h \cdot k)$; since $f = \tau \cdot h \cdot k$, this means that $A \models \alpha(e^\# \cdot f)$, as desired.

Completeness (\Leftarrow): Suppose that every algebra in \mathcal{V} satisfies $X \vdash \alpha(f)$. Then, in particular, $\mathcal{F}X$ satisfies $X \vdash \alpha(f)$. Hence, for the relation preserving assignment $\eta_X: X \rightarrow T_\Sigma(X)$ we have that $\eta_X^\#(f(i))$ is defined for all $i \in \text{ar}(\alpha)$, and $\mathcal{F}X \models \alpha(\eta^\# \cdot f)$. By Lemma C.10, we have that $\eta_X^\#(f(i)) = [f(i)]$ for all $i \in \text{ar}(\alpha)$. In particular, f factorizes as a map $f: \text{ar}(\alpha) \rightarrow \mathcal{F}_\mathcal{V}(X) \hookrightarrow T_\Sigma(X)$, and $\mathcal{F}_\mathcal{V}(X) \models \alpha(f)$. That is, $X \vdash \alpha(f)$ is derivable. \blacktriangleleft

D Details for Section 5

Proof of Lemma 5.5. Let $X \in \mathbf{C}$; we first show, for a given operation symbol σ of arity Γ , that $\sigma_{TX}: [\Gamma, TX] \rightarrow TX$ is relation preserving. Indeed, we have

$$\sigma_{TX} = ([\Gamma, TX] \xrightarrow{(-)^*} [T\Gamma, TX] \xrightarrow{\pi_x} TX),$$

where $(-)^*$ is the Kleisli extension of the monad T and $\pi_x(g) = g(x)$ (cf. Definition 3.11) is the evaluation map for $x = \sigma = \sigma(u_\Gamma)$ (Notation 5.3). The Kleisli extension $(-)^*$ is relation preserving since T is an enriched monad, and π_x clearly is relation preserving by the definition of the relational structure on $[T\Gamma, TX]$. Thus σ_{TX} is relation preserving.

Now we are going to verify that TX lies in $\mathcal{V}_\mathbb{T}$. First, observe that for all $\sigma \in \Sigma_\mathbb{T}$ and all relation preserving assignments $f: \text{ar}(\sigma) \rightarrow TX$, we have

$$f^\#(\sigma) = f^*(\sigma). \tag{D.1}$$

(On the left-hand side of the equation above, we view σ as a term according to Notation 5.3.) Indeed, where $|\text{ar}(\sigma)| = \{x_i \mid i \in \text{ar}(\sigma)\}$, we see that $f^\#(\sigma)$ is defined using that f is relation preserving: for every Π -edge $\alpha(g)$ in $\text{ar}(\sigma)$ we have $\alpha_{TX}(f^\# \cdot (u_{\text{ar}(\sigma)}) \cdot g) = \alpha_{TX}(f \cdot g)$. Furthermore, using only the definitions of $f^\#$ and of σ_{TX} , we have

$$f^\#(\sigma) = f^\#(\sigma(x_i)) = \sigma_{TX}(f^\#(x_i)) = \sigma_{TX}(f(x_i)) = \sigma_{TX}(f) = f^*(\sigma).$$

We will now see that TX satisfies all axioms of \mathcal{V}_T , as required. To this end, we make a case distinction on the axiom types described in Definition 5.4:

- (1) Given a Σ -relation $\Gamma \vdash \alpha(\sigma_i)$ such that $T\Gamma \models \alpha(\sigma_i)$ and a relation preserving map $f: \Gamma \rightarrow TX$, we first note that $TX \models \alpha(f^\#(\sigma_i))$ since $f^*: T\Gamma \rightarrow TX$ is relation preserving and $f^*(\sigma_i) = f^\#(\sigma_i)$ for all $i \in \text{ar}(\alpha)$ by (D.1). Thus, to see that TX satisfies $\Gamma \vdash \alpha(\sigma_i)$, it suffices to show that $f^\#(\sigma_i)$ is defined for all $i \in \text{ar}(\sigma)$. This follows immediately from (D.1).
- (2) Now, given $\Delta \in \mathcal{P}_\lambda$, a morphism $f: \Delta \rightarrow T\Gamma$, and an operation $\sigma \in T\Delta$, we verify that TX satisfies $\Gamma \vdash f^*(\sigma) = \sigma(f)$. To this end, let $m: \Gamma \rightarrow TX$ be a relation preserving assignment. Then $m^\#(f^*(\sigma))$ is defined; this is shown completely analogously as before. Furthermore, we have

$$\begin{aligned}
m^\#(f^*(\sigma)) &= m^* \cdot f^*(\sigma) && \text{by (D.1)} \\
&= (m^* \cdot f)^*(\sigma) && \text{by (5.1)} \\
&= \sigma_{TX}(m^* \cdot f) && \text{def. of } \sigma_{TX} \\
&= \sigma_{TX}(m^\# \cdot f) && \text{by (D.1)} \\
&= m^\#(\sigma(f)) && \text{def. of } m^\#.
\end{aligned}$$

- (3) Given $f: \Gamma \rightarrow TX$ relation preserving, we use (D.1) to obtain

$$f^\#(\eta_\Gamma(x)) = f^*(\eta_\Gamma(x)) = f(x) = f^\#(x). \quad \blacktriangleleft$$

Proof of Theorem 5.6. We first show that every canonical algebra TX has the universal property of a free algebra in \mathcal{V}_T with respect to the monad unit $\eta_X: X \rightarrow TX$. We split this into two parts:

- (1) Assume that $X = \Gamma \in \mathcal{P}_\lambda$ and let $f: \Gamma \rightarrow A$ be a relation preserving map; we proceed to show that there exists a unique homomorphism $\bar{f}: T\Gamma \rightarrow A$ such that $f = \bar{f} \cdot \eta_\Gamma$. To this end, let \bar{f} be the map defined by

$$\bar{f}(\sigma) := \sigma_A(f)$$

for all $\sigma \in T\Gamma$. We first prove that $\bar{f} \cdot \eta_\Gamma = f$; for every $x \in \Gamma$ we have

$$\begin{aligned}
\bar{f} \cdot \eta_\Gamma(x) &= \bar{f}(\eta_\Gamma(x)) \\
&= (\eta_\Gamma(x))_A(f) && \text{def. of } \bar{f} \\
&= (\eta_\Gamma(x))_A(f^\# \cdot u_\Gamma) && \text{def. of } f^\# \\
&= f^\#((\eta_\Gamma(x))(u_\Gamma)) && \text{def. of } f^\# \\
&= f^\#(x) && \text{since } A \text{ satisfies } \Gamma \vdash \eta_\Gamma(x) = x \\
&= f(x) && \text{def. of } f^\#.
\end{aligned}$$

Next we prove that \bar{f} is relation preserving. Indeed, given an edge $\alpha(\sigma_i)$ in $T\Gamma$, we have that A satisfies $\Gamma \vdash \alpha(\sigma_i)$ since it is an algebra in \mathcal{V}_T . In particular, $A \models \alpha_A(f^\#(\sigma_i))$. Moreover, we have

$$f^\#(\sigma_i) = f^\#(\sigma_i(x_i)) = (\sigma_i)_A(f(x_i)) = \bar{f}(\sigma_i)$$

for all $i \in \text{ar}(\alpha)$. Hence $A \models \alpha(\bar{f}(\sigma_i))$.

Now we show that \bar{f} is a homomorphism. To this end, let $\tau \in \Sigma_{\mathbb{T}}$. We are going to show that the following square commutes

$$\begin{array}{ccc} [\mathbf{ar}(\tau), T\Gamma] & \xrightarrow{\tau_{T\Gamma}} & T\Gamma \\ \bar{f} \cdot (-) \downarrow & & \downarrow \bar{f} \\ [\mathbf{ar}(\tau), A] & \xrightarrow{\tau_A} & A \end{array}$$

Indeed, for every relation preserving $m: \mathbf{ar}(\tau) \rightarrow T\Gamma$ we have

$$\begin{aligned} \bar{f}(\tau_{T\Gamma}(m)) &= \bar{f}(m^*(\tau)) && \text{def. of } \tau_{T\Gamma} \\ &= (m^*(\tau))_A(f) && \text{def. of } \bar{f} \\ &= f^\#(m^*(\tau)) && \text{def. of } f^\# \\ &= f^\#(\tau(m)) && A \text{ satisfies } \Gamma \vdash m^*(\tau) = \tau(m) \\ &= \tau_A(f^\# \cdot m) && \text{def. of } f^\# \\ &= \tau_A(\bar{f} \cdot m). \end{aligned}$$

For the last step we shall prove that $f^\#(m(x)) = \bar{f}(m(x))$ for every $x \in \mathbf{ar}(\tau)$. Indeed, using the definition of $f^\#$ we see that the operation symbol $\sigma = m(x)$, considered as the term $\sigma(u_\Gamma)$ as in Notation 5.3 satisfies

$$f^\#(\sigma(u_\Gamma)) = \sigma_A(f^\# \cdot u_\Gamma) = \sigma_A(f) = \bar{f}(\sigma).$$

As for the uniqueness, suppose that $\bar{f}: T\Gamma \rightarrow A$ is a homomorphism such that $\bar{f} \cdot \eta_\Gamma = f$. Then the above square commutes for $\mathbf{ar}(\tau) = \Gamma$ which applied to $\eta_\Gamma \in [\Gamma, T\Gamma]$ yields for every $\sigma \in |T\Gamma|$:

$$\begin{aligned} \bar{f}(\sigma) &= \bar{f}(\eta_\Gamma^*(\sigma)) && \text{by (5.1)} \\ &= \bar{f}(\eta_\Gamma^\#(\sigma)) && \text{by (D.1)} \\ &= \bar{f}(\sigma_{T\Gamma}(\eta_\Gamma)) && \text{def. of } \eta_\Gamma \\ &= \sigma_A(\bar{f} \cdot \eta_\Gamma) && \bar{f} \text{ homomorphism} \\ &= \sigma_A(f) && \text{since } \bar{f} \cdot \eta_\Gamma = f. \end{aligned}$$

(2) Now suppose that X is an arbitrary object in \mathbf{C} . Using that \mathbf{C} is locally λ -presentable, we may express X as a λ -directed colimit of λ -presentable objects, say $X = \text{colim } \Gamma_i$. Then, since T is λ -accessible, we see that $TX = \text{colim } T\Gamma_i$, and this lifts to a colimit in $\mathcal{V}_{\mathbb{T}}$ because the forgetful functor $\mathcal{V}_{\mathbb{T}} \rightarrow \mathbf{C}_0$ creates λ -directed colimits by Proposition 4.9 and Remark 4.3(2).

(3) To complete the proof we apply Remark 5.2. The given monad \mathbb{T} and the free algebra monad of $\mathcal{V}_{\mathbb{T}}$ share the same object assignment $X \mapsto TX$, and the family of morphisms η_X , as shown in the previous two items. It remains to prove that for every morphism $h: X \rightarrow TY$ the morphism $h^*: TX \rightarrow TY$ is a Σ -homomorphism. Then \mathbb{T} and the free algebra monad of $\mathcal{V}_{\mathbb{T}}$ also share the same operator $h \mapsto h^*$. Given $\sigma \in |T\Gamma|$ we shall prove that the following square commutes:

$$\begin{array}{ccc} [\Gamma, TX] & \xrightarrow{\sigma_{TX}} & TX \\ h^* \cdot (-) \downarrow & & \downarrow h^* \\ [\Gamma, TY] & \xrightarrow{\sigma_{TY}} & TY \end{array}$$

36 Monads on Categories of Relational Structures

Indeed, given $f: \Gamma \rightarrow TY$ we have

$$\begin{aligned} h^* \cdot \sigma_{TX}(f) &= h^* \cdot f^*(\sigma) && \text{def. of } \sigma_{TX} \\ &= (h^* \cdot f)^*(\sigma) && \text{by (5.1)} \\ &= \sigma_{TY}(h^* \cdot f) && \text{def. of } \sigma_{TY}. \end{aligned}$$

