

# THE MODULI STACK OF PRINCIPAL $\rho$ -SHEAVES AND GIESEKER-HARDER-NARASIMHAN FILTRATIONS

ANDRES FERNANDEZ HERRERO, TOMÁS L. GÓMEZ, AND ALFONSO ZAMORA

ABSTRACT. Given a smooth projective variety  $X$  and a connected reductive group  $G$  defined over a field of characteristic 0, we define a moduli stack of principal  $\rho$ -sheaves that compactifies the stack of  $G$ -bundles on  $X$ . We apply the theory developed in [HL14, AHLH19] to construct a moduli space of Gieseker semistable principal  $\rho$ -sheaves. This provides an intrinsic stack-theoretic construction of the moduli space of semistable singular principal bundles [Sch02, GLSS08].

Our second main result is the definition of a schematic Gieseker-Harder-Narasimhan filtration for  $\rho$ -sheaves, which induces a stratification of the stack by locally closed substacks. This filtration for a general reductive group  $G$  is a refinement of the canonical slope parabolic reduction previously considered at the level of points in [AAB02] and as a stratification of the stack in [GN18, GN16]. In an appendix, we apply the same techniques to define Gieseker-Harder-Narasimhan filtrations in arbitrary characteristic and show that they induce a stratification of the stack by radicial morphisms.

## CONTENTS

1. Introduction	2
1.1. Stack-theoretic approach vs. GIT	5
2. Preliminaries	7
2.1. Notation	7
2.2. Big open sets and torsion-free sheaves	8
2.3. The scheme of $G$ -reductions	10
3. Principal $\rho$ -sheaves	13
3.1. Principal $\rho$ -sheaves and the moduli stack $\text{Bun}_\rho(X)$	13
3.2. Filtrations of $\rho$ -sheaves	14
3.3. Filtrations of $\rho$ -sheaves in the split case	16
4. $\Theta$ -stratification on $\text{Bun}_\rho(X)$	19
4.1. The numerical invariant	19
4.2. Affine Grassmannians for $\rho$ -sheaves	20
4.3. Monotonicity of the numerical invariant	23
4.4. HN boundedness	26
4.5. $\Theta$ -stratification, semistability and leading term filtrations	33
5. Completeness of the stack $\text{Bun}_\rho(X)$	36
6. Moduli space for the semistable locus	41
6.1. The degree of a $\rho$ -sheaf	41

---

2020 *Mathematics Subject Classification.* 14D20, 14D23, 14F06, 14J60, 14L24.

*Key words and phrases.* Moduli spaces, principal bundles, stacks,  $\Theta$ -stratifications, Harder-Narasimhan filtration, Gieseker stability, principal  $\rho$ -sheaves.

6.2.	Boundedness of the semistable locus	43
6.3.	Moduli space for the semistable locus	44
7.	Generalization: $\rho$ -sheaves for products of general linear groups	45
8.	Gieseker-Harder-Narasimhan filtrations for $\rho$ -sheaves	47
8.1.	Lexicographic filtrations	48
8.2.	Multi-weighted parabolic reductions	49
8.3.	The GHN filtration for $\rho$ -sheaves	51
8.4.	The GHN filtration in families	53
9.	Examples	55
9.1.	Torsion-free sheaves	55
9.2.	Example of Gieseker-Harder-Narasimhan filtration	56
10.	Comparison with previous work	58
10.1.	Comparison with previous notions of semistability	58
10.2.	Comparison of the leading term filtration with previous notions in the literature	63
Appendix A.	Gieseker-Harder-Narasimhan filtrations in positive characteristic	65
References		67

## 1. INTRODUCTION

Fix a smooth projective variety  $X$  and a connected reductive group  $G$  defined over a field of characteristic 0 (see Appendix A and Theorem 3 for positive characteristic). When  $\dim X = 1$ , Ramanathan [Ram75] constructed a projective moduli space of semistable principal  $G$ -bundles on  $X$  using Geometric Invariant Theory (GIT). The stability condition in this case is analogous to the notion of slope stability for a vector bundle (introduced by Mumford), in the sense that it involves the degree and the rank of certain vector bundles.

If  $\dim X > 1$ , it is necessary to consider also degenerations of principal  $G$ -bundles in order to construct a proper moduli space. These degenerations are called principal  $\rho$ -sheaves, where  $\rho$  is a choice of representation of  $G$ . A principal  $\rho$ -sheaf is a triple  $(P, \mathcal{E}, \psi)$  consisting of a torsion-free sheaf  $\mathcal{E}$  on  $X$ , a principal  $G$ -bundle  $P$  defined on the open set  $U$  where  $\mathcal{E}$  is locally free, and an isomorphism  $\psi$  between the restriction  $\mathcal{E}|_U$  and the vector bundle associated to  $P$  using the representation  $\rho$  (see Definition 3.1 and Proposition 2.18). More generally, we allow  $\rho$  to be a faithful homomorphism from  $G$  into a product of general linear groups (see Definition 7.1, Definition 10.1, and Section 7)<sup>1</sup>. These  $\rho$ -sheaves have also been called in the literature “singular principal bundles” by Schmitt [Sch02, GLSS08] (in the case when  $\rho$  is valued in a special linear group), and “principal  $G$ -sheaves” by Gómez-Sols [GS05] (when  $\rho$  is the adjoint representation, not injective in general). We show that there is an algebraic stack  $\text{Bun}_\rho(X)$  that parametrizes principal  $\rho$ -sheaves on  $X$ .

When going from  $\dim X = 1$  to  $\dim X > 1$ , an additional consideration needs to be introduced in order to construct the moduli space. Instead of using slope semistability, we have to use a notion where the degree is substituted by the Hilbert polynomial. This is Gieseker semistability for torsion-free sheaves. An analogous notion of semistability was

---

<sup>1</sup>If we want our definition of stability to coincide with Ramanathan’s in the case of curves, then we have to require that the representation  $\rho$  is central (Definition 10.5).

defined in [GS05] and [GLSS08] for principal  $\rho$ -sheaves. Using the methods of GIT, Gómez-Sols [GS05], Schmitt [Sch02] and Gómez-Langer-Schmitt-Sols [GLSS08] constructed moduli spaces of semistable  $\rho$ -sheaves. We remark that the moduli space obtained will in general depend on the choice of representation  $\rho$ .

Alper, Halpern-Leistner and Heinloth [AHLH19] have recently introduced an alternative approach to GIT for the construction of moduli spaces. Their approach is intrinsic to the algebraic stack associated to the moduli problem of interest. In this article we employ their techniques instead of the GIT methods that were previously used in the literature.

In this direction, our first main result in this article is an intrinsic-stack theoretic construction of the moduli space of semistable  $\rho$ -sheaves, where  $\rho$  is any faithful representation of  $G$  into a product of general linear groups. In order to obtain a quasicompact moduli space, it is helpful to fix a tuple  $P^\bullet = (P^1, P^2, \dots, P^b)$  of rational polynomials  $P^i \in \mathbb{Q}[t]$  and restricts to the locus of  $\rho$ -sheaves where the  $i^{\text{th}}$  underlying sheaf has Hilbert polynomial  $P^i$ . We state the result here in the case when the representation  $\rho$  is central (Definition 10.5)<sup>2</sup>:

**Theorem 1.** *Fix a central faithful representation  $\rho : G \hookrightarrow \prod_i \mathrm{GL}_{n_i}$ . Then*

- (a) *The moduli stack  $\mathrm{Bun}_\rho(X)$  of  $\rho$ -sheaves admits a  $\Theta$ -stratification induced by the polynomial numerical invariant  $\nu$  defined in Subsection 4.1.*
- (b) *(Semistable reduction) The semistable locus  $\mathrm{Bun}_\rho(X)^{\mathrm{ss}}$  satisfies the existence part of the valuative criterion for properness.*
- (c) *For any choice of tuple of rational polynomials  $P^\bullet$ , the open substack  $\mathrm{Bun}_\rho(X)^{\mathrm{ss}, P^\bullet}$  of  $\rho$ -sheaves with tuple  $P^\bullet$  admits a proper good moduli space (in the sense of [Alp13]).*

If the representation  $\rho$  is valued in a single special linear group (resp.  $\rho$  is the adjoint representation), then we recover the moduli space constructed in [Sch02, GLSS08] (resp. [GS05]).

Another important aspect of the theory of semistable vector bundles on a curve is the existence of a canonical filtration for any unstable vector bundle. This is called the Harder-Narasimhan filtration, and it is such that the associated graded object is a direct sum of semistable sheaves. This is helpful because unstable vector bundles are then constructed as extensions of semistable bundles, thus allowing us to do recursive arguments to study properties of the moduli space, as done by Harder-Narasimhan [HN75] and Atiyah-Bott [AB83]. More generally, for principal  $G$ -bundles on a curve Behrend [Beh95] and Biswas-Holla [BH04] constructed the Harder-Narasimhan reduction to a parabolic subgroup  $P \subset G$ . An important property is that the associated principal  $L$ -bundle is semistable, where  $L = G/P$  is the Levi factor of  $P$ .

For torsion free sheaves over  $X$  with  $\dim X > 1$ , we can construct two different Harder-Narasimhan filtrations, depending on the stability condition we use, slope or Gieseker stability. They are respectively called slope-Harder-Narasimhan and Gieseker-Harder-Narasimhan.

For principal  $\rho$ -sheaves over  $X$  with  $\dim X > 1$  we only find in the literature the slope version of the Harder-Narasimhan filtration, constructed at the level of points by Anchouche-Azad-Biswas [AAB02] and in families by Gurjar-Nitsure [GN18, GN16]. Biswas and Zamora [BZ16] showed that the notion of Gieseker filtration is not well-behaved with respect to change of group. The strategy of considering the canonical filtration of the adjoint vector bundle does not yield in general a well-defined notion of Gieseker-Harder-Narasimhan reduction in higher dimensions. Therefore a new approach needs to be introduced.

Our second main result is the definition of a Gieseker-Harder-Narasimhan parabolic reduction in the context of  $\rho$ -sheaves.

<sup>2</sup>A more general modified statement is proven in the body of the document, see Theorem 7.2.

**Theorem 2.** *Every principal  $\rho$ -sheaf admits a uniquely defined multi-weighted filtration called the Gieseker-Harder-Narasimhan filtration (abbreviated GHN filtration), see Definition 8.10.*

*This filtration is schematic, in the sense that the universal stacks for relative GHN filtrations induce a locally closed stratification of  $\text{Bun}_\rho(X)$ .*

*If the group  $G$  is split, then the GHN filtration comes from a multi-weighted parabolic reduction defined over a big open subset (cf. Definition 2.1). The associated Levi  $\rho_{\bar{\chi}}$ -sheaf is semistable.*

If the representation  $\rho$  is central, then our GHN filtration refines the canonical slope reduction constructed in [AAB02]. In this case, the resulting stratification of the stack  $\text{Bun}_\rho(X)$  refines the canonical slope stratification on the stack of  $G$ -bundles that was recently constructed in [GN18].

Our main tools originate in some recent work on the theory of stacks. Alper, Halpern-Leistner and Heinloth [AHLH19] have recently developed a theory which produces moduli spaces for Artin stacks, generalizing results of Keel-Mori on Deligne-Mumford stacks. This can be combined with the theory of  $\Theta$ -stability and  $\Theta$ -stratifications by Halpern-Leistner [HL14] to produce a powerful stack-theoretic approach to the construction of moduli spaces and canonical filtrations (see [HL14, §5] and [HLHJ21, §2.5] for more details). Halpern-Leistner defines in [HL14] a notion of filtration of an object in an Artin stack. Given certain cohomology classes on the stack, one can define a numerical invariant on filtrations. An object in the stack is called semistable if the numerical function is non-positive for all filtrations. If the numerical invariant satisfies certain properties, then the open substack of semistable objects admits a *good moduli space* (as defined in [Alp13]). This is an intrinsic way of constructing the moduli space, in the sense that we do not need to choose a parameter space nor the action of a group.

Here are the main steps involved in this general approach:

- (Step 1) Define a (polynomial) numerical invariant  $\nu$  on the stack (Subsection 4.1).
- (Step 2) Prove that the numerical invariant  $\nu$  satisfies some monotonicity properties (Subsection 4.3).
- (Step 3) Show that  $\nu$  satisfies the HN boundedness condition (Subsection 4.4).
- (Step 4) Prove boundedness of the semistable locus (Subsection 6.2).

In this article we carry out this program for the moduli stack of principal  $\rho$ -sheaves, thus obtaining the intrinsic stack-theoretic construction of the moduli space described in Theorem 1. Furthermore, this theory provides a  $\Theta$ -stratification of  $\text{Bun}_\rho(X)$ , which yields a canonical filtration (called the leading term HN filtration) for each unstable object. We apply this construction iteratively to the centers of the  $\Theta$ -stratification to define the GHN reduction from Theorem 2.

In Appendix A we explain how to apply the same techniques to define the GHN filtration for  $\rho$ -sheaves in arbitrary characteristic.

**Theorem 3.** *Suppose that  $k$  is a field of arbitrary characteristic. Every principal  $\rho$ -sheaf admits a uniquely defined multi-weighted filtration called the GHN filtration, after perhaps passing to a finite purely inseparable extension of the field of definition.*

*Relative GHN filtrations are represented by a schematic morphism  $H \rightarrow \text{Bun}_\rho(X)$  that is surjective, radicial and of finite type. Each connected component of  $H$  is finite over its image.*

*If the group  $G$  is split, then the GHN filtration comes from a multi-weighted parabolic reduction defined over a big open subset. The associated Levi  $\rho_{\bar{\chi}}$ -sheaf is semistable.*

Here is an outline of the paper:

In Section 2 we discuss some of the necessary background on big open subschemes and torsion-free sheaves. We define the scheme of  $G$ -reductions (Definition 2.14), which is a version of the fiber bundle of  $G$ -reductions in the context of torsion-free sheaves.

In Section 3 we define the notion of principal  $\rho$ -sheaf (Definition 3.1) and define the stack  $\text{Bun}_\rho(X)$  (Definition 3.5) in the case of a representation  $\rho : G \rightarrow \text{GL}(V)$  into a single general linear group. Furthermore, we use the work of Heinloth [Hei17] to produce an explicit description of  $\Theta$ -filtrations on  $\text{Bun}_\rho(X)$  (Proposition 3.16).

Section 4 is the technical heart of the paper. We define a polynomial numerical invariant  $\nu$  on the stack  $\text{Bun}_\rho(X)$ . Following [HLHJ21], we define a higher dimensional analogue of the affine Grassmannian for  $\rho$ -sheaves and use it to prove the relevant monotonicity properties of  $\nu$ . We also prove the HN boundedness condition via an optimization argument on cones inside the vector space of cocharacters of  $G$  (see Proposition 4.26). This way we obtain a  $\Theta$ -stratification. In Subsection 4.5 we give descriptions of the semistable locus and the  $\Theta$ -stratification. We call the resulting canonical filtration of  $\rho$ -sheaves the “leading term HN filtration”.

In Section 5 we prove that the stack  $\text{Bun}_\rho(X)$  satisfies the existence part of the valuative criterion for properness. The key lemma is a result about extension of  $G$ -bundles over a DVR after a purely ramified extension (Proposition 5.2). The proof boils down to a description of Galois cohomology classes in [BT87] and the behavior of the Bruhat-Tits building under purely ramified extensions.

Section 6 includes a proof of boundedness of the semistable locus, and finalizes the intrinsic construction of the moduli space (Theorem 6.8) in the case of a representation  $\rho : G \rightarrow \text{GL}(V)$  into a single general linear group.

In Section 7 we generalize the results of previous subsections to the case when  $\rho$  is a faithful homomorphism into a product of general linear groups.

Section 8 introduces the notion of lexicographic filtration which provides a convenient framework for the construction of the Gieseker-Harder-Narasimhan filtration in Subsection 8.3. We end the section by defining a relative version of the Gieseker-Harder-Narasimhan filtration and proving that it induces a locally closed stratification of the stack  $\text{Bun}_\rho(X)$ .

In Section 9 we present several examples to illustrate the theory developed. In Subsection 9.1 we consider the case when  $G$  is a general linear group and  $\rho$  is the identity. In this case a  $\rho$ -sheaf is the same thing as a torsion-free sheaf. We describe explicitly the construction of our Gieseker-Harder-Narasimhan filtration in this case, and show that it coincides with the classical notion for torsion-free sheaves (Proposition 9.3). In Subsection 9.2 we work out explicitly the case of a  $\rho$ -sheaf for  $\text{SO}(7)$  and compare its leading term HN filtration with its Gieseker-Harder-Narasimhan filtration.

In Section 10 we compare our notions with previous ones in the literature. In Subsection 10.1 we introduce central representations (Definition 10.5). If  $\rho$  is central, then we prove that our notion of Gieseker semistability is compatible with previous notions in the literature, and our Gieseker-Harder-Narasimhan stratification of the stack refines the slope stratification constructed in [GN18, GN16].

In Appendix A we show how to apply our techniques to define the Gieseker-Harder-Narasimhan stratification in arbitrary characteristic. In this case the GHN filtrations induce a “stratification” of the stack by radicial morphisms of finite type.

**1.1. Stack-theoretic approach vs. GIT.** We end our introduction with a discussion comparing the stack theoretic approach in [HL14, AHLH19] with the usual construction of moduli spaces using Geometric Invariant Theory. We hope that this can provide some intuition for those readers that are more familiar with the GIT approach.

The construction of the moduli space of torsion free sheaves (and principal  $\rho$ -sheaves) follows the usual application of GIT. We first construct a scheme  $R$  parametrizing all objects together with an extra data (in this case an isomorphism between a fixed vector space  $W$  and the space of sections  $H^0(\mathcal{E}(n))$  for a chosen  $n \gg 0$  and the sheaf  $\mathcal{E}$  associated to each object). The group  $\mathrm{SL}(W)$  acts on the scheme  $R$ . We choose an ample line bundle on  $R$  together with a linearization of the action of  $\mathrm{SL}(W)$ . Mumford's GIT shows that there is an open subset  $R^{ss} \subset R$  which admits a *good quotient* [Ses72], and this is the moduli space of semistable torsion-free sheaves we are looking for.

To identify the subset  $R^{ss}$  of semistable points we use the Hilbert-Mumford criterion. For each point  $p \in R$  and for each one-parameter subgroup  $\lambda : \mathbb{G}_m \rightarrow \mathrm{SL}(W)$  we consider the weight  $\mu(\lambda, p)$  of the action of the one-parameter subgroup  $\lambda$  on the fiber of the limiting point  $\lim_{t \rightarrow 0} \lambda(t) \cdot p$ . The point  $p$  is semistable if and only if  $\mu(\lambda, p) \leq 0$  for all one-parameter subgroups  $\lambda$ . We remark that, in the usual applications of GIT to the construction of moduli spaces involving sheaves, the one-parameter subgroup  $\lambda$  produces a filtration of the sheaf, and the Hilbert-Mumford criterion becomes equivalent to checking certain numerical condition over all filtrations.

In [HL14], the one-parameter subgroups appearing in the Hilbert-Mumford criterion of GIT are replaced by *filtered objects*. For any algebraic stack  $\mathcal{M}$ , a filtration of a point  $p : \mathrm{Spec} k \rightarrow \mathcal{M}$  is a morphism

$$(1) \quad f : \Theta := [\mathbb{A}^1/\mathbb{G}_m] \rightarrow \mathcal{M}$$

and an identification  $f(1) \simeq p$ . Given  $f$ , there is a group homomorphism from  $\mathbb{G}_m$  to the automorphism group of the object  $f(0)$  in  $\mathcal{M}$ , which can be thought of as the graded object associated to the filtration  $f$ .

For understanding these notions it is useful to let  $\mathcal{M}$  be the moduli stack of torsion free sheaves on a projective scheme  $X$ . By the Rees construction (cf. Proposition 3.8), a decreasing  $\mathbb{Z}$ -indexed filtration

$$0 \subset \cdots \subset \mathcal{E}_{i+1} \subset \mathcal{E}_i \subset \cdots \subset \mathcal{E}$$

of a torsion free sheaf  $\mathcal{E}$  on  $X$  determines an  $\mathbb{A}^1$ -family  $\tilde{\mathcal{E}}$  of torsion free sheaves on  $X$ . We have  $\tilde{\mathcal{E}}|_{X_t} \cong \mathcal{E}$  for all  $t \neq 0$ , and  $\tilde{\mathcal{E}}|_{X_0} \cong \bigoplus \mathcal{E}_i/\mathcal{E}_{i+1}$ . The group  $\mathbb{G}_m$  has a natural action on the graded object  $\tilde{\mathcal{E}}|_{X_0}$ , with weight  $t^i$  on each piece  $\mathcal{E}_i/\mathcal{E}_{i+1}$ . The family  $\tilde{\mathcal{E}}$  gives a morphism  $\mathbb{A}^1 \rightarrow \mathcal{M}$  and the standard action of  $\mathbb{G}_m$  on  $\mathbb{A}^1$  lifts to  $\tilde{\mathcal{E}}$ . Hence this morphism descends to a map from the quotient stack  $[\mathbb{A}^1/\mathbb{G}_m]$ , providing a filtration  $f$  as in (1). One can define a function  $\nu$  which assigns a real number for each filtration  $f$  as in (1). This is done in a similar way to GIT: we choose a line bundle  $L$  on  $\mathcal{M}$ , and define  $\nu(f)$  to be the weight of the action of  $\mathbb{G}_m$  on the fiber of  $L|_{f(0)}$ .

In practice this approach has an important advantage: in the GIT case we are looking at the action of  $\mathbb{G}_m$  on  $\mathrm{SL}(W)$ . This produces a filtration on the vector space  $W \cong H^0(E(n))$ , which in turn produces a filtration on  $\mathcal{E}$ . To show that the numerical condition of the Hilbert-Mumford criterion given by  $\mu(\lambda, p)$  matches the stability condition on Hilbert polynomials is an involved calculation. On the other hand, if we follow the intrinsic approach of [HL14], we are directly looking at a filtration on  $\mathcal{E}$  and the action of  $\mathbb{G}_m$  on the associated graded sheaf. This makes it easier to obtain the stability condition.

In the case of principal  $G$ -bundles the advantage of the new approach is even more clear. In the GIT setup we are looking at all filtrations of  $W \cong H^0(E(n))$ , and it is not easy to show that it is enough to look at filtrations coming from 1-parameter subgroups of  $G$ . In contrast, in the theory of [HL14, Hei17] we only look at these subgroups from the very beginning.

Both GIT and [HL14] can be used to produce filtrations of unstable objects. We recall how this works in GIT. Let  $p \in R$  be an unstable point. Among all the 1-parameter

subgroups  $\lambda$  of  $\mathrm{SL}(W)$  with  $\mu(\lambda, p) > 0$ , we can find one for which this weight is maximal (after some normalization). This defines the Kempf filtration (cf. [Kem78]) of  $p$ . This filtration depends on the choices that were made in the GIT construction (the number  $n$ , the parameter space  $R$  and the linearization). For torsion free sheaves it can be shown that it is independent of the choice of  $n$  for  $n \gg 0$ , and it coincides with the Gieseker-Harder-Narasimhan filtration [GSZ15, Zam13]. Unfortunately, the computations become too involved in the case of principal  $G$ -bundles, so we do not obtain a Gieseker-Harder-Narasimhan filtration following this approach [Zam16].

In the theory in [HL14], given an unstable object  $p \in \mathcal{M}(k)$  we can also find a filtration  $f$  with  $f(1) \simeq p$  such that  $\nu(f)$  is maximal. This program was carried out for  $\Lambda$ -modules and pairs in [HLHJ21]. It should be pointed out that the filtration  $f$  which maximizes  $\nu$  does not have the property that the graded pieces of the filtration are semistable. But if we apply this procedure inductively we obtain the Gieseker-Harder-Narasimhan filtration.

In this article we show that a similar story applies in the case of principal  $\rho$ -sheaves. We use this to define what we call the Gieseker-Harder-Narasimhan reduction of an unstable  $\rho$ -sheaf. We refer the reader to Section 9 for examples.

**Acknowledgments:** This project was started during the workshop “Moduli problems beyond geometric invariant theory” at the American Institute of Mathematics. We would like to thank the American Institute of Mathematics and the organizers that made the workshop possible. We would like to thank V. Balaji, Federico Fuentes, Daniel Halpern-Leistner and Jochen Heinloth for helpful remarks.

This work is supported by grants CEX2019-000904-S and PID2019-108936GB-C21 (funded by MCIN/AEI/ 10.13039/501100011033), and NSF grants DMS-1454893 and DMS-2001071.

## 2. PRELIMINARIES

**2.1. Notation.** We will work over a field  $k$  of characteristic 0. We denote by  $\mathrm{Aff}_k$  the category of affine schemes over  $k$ . More generally, for any  $k$ -scheme  $T$  we let  $\mathrm{Aff}_T$  be the category of (absolutely) affine schemes equipped with a morphism to  $T$ . We write  $\mathrm{Sch}_k$  and  $\mathrm{Sch}_T$  to denote the category of all schemes over  $k$  and  $T$  respectively. Unless otherwise stated, all schemes will be understood to belong to  $\mathrm{Sch}_k$ . An undecorated product of schemes (e.g.  $X \times S$ ) is understood to be a fiber product over  $k$ . We might sometimes write  $X_S$  instead of  $X \times S$ .

All of the sheaves that we consider in this paper are quasicohherent. Whenever we write “ $\mathcal{O}_Y$ -module” we mean a quasicohherent  $\mathcal{O}_Y$ -module. In order to ease notation for pullbacks, we use the following convention. Whenever we have a map of schemes  $f : Y \rightarrow T$  and a sheaf  $\mathcal{G}$  on  $T$ , we will write  $\mathcal{G}|_Y$  to denote the pullback  $f^*\mathcal{G}$ , if the morphism  $f$  is clear from context.

We fix a smooth projective geometrically connected variety  $X$  over  $k$ . We denote by  $d$  the dimension of  $X$ . We also choose once and for all an ample line bundle  $\mathcal{O}(1)$  on  $X$ . For any coherent sheaf  $\mathcal{F}$  on  $X$ , we denote by  $P_{\mathcal{F}}$  the Hilbert polynomial of  $\mathcal{F}$  with respect to the line bundle  $\mathcal{O}(1)$ . If  $\mathcal{F}$  is a torsion-free sheaf on  $X$ , then this is a polynomial of degree  $d$  in the variable  $n$  which can be written in the form

$$P_{\mathcal{F}}(n) = \sum_{i=0}^d \frac{a_i}{i!} n^i$$

for some sequence of rational numbers  $a_i$  (see [HL97, Lemma 1.2.1]). We define the reduced Hilbert polynomial to be  $\bar{p}_{\mathcal{F}} := \frac{1}{a_d} P_{\mathcal{F}}$ . For every  $0 \leq i \leq d-1$ , we set  $\hat{\mu}_i(\mathcal{F}) := \frac{a_i}{a_d}$ . We call this the  $i^{\mathrm{th}}$  slope of  $\mathcal{F}$ . The number  $\hat{\mu}_{d-1}(\mathcal{F}) = \frac{a_{d-1}}{a_d}$  is simply called the slope of  $\mathcal{F}$ .

We define an ordering for polynomials where, for any two  $p_1, p_2 \in \mathbb{R}[n]$ , we write  $p_1 \geq p_2$  if  $p_1(n) \geq p_2(n)$  for  $n \gg 0$ .

Fix a reductive linear algebraic group  $G$  over  $k$ . We write  $k[G]$  to denote the coordinate ring of  $G$ , which has the structure of a commutative Hopf algebra. We will denote by  $\text{Bun}_G(X)$  the moduli stack of principal  $G$ -bundles on  $X$ . This is an algebraic stack that has affine diagonal and is locally of finite type over  $k$  (by [HR19, Thm. 1.2] taking  $Z = X$  and  $X = BG$ ).

Fix a  $k$ -vector space  $V$  of dimension  $r$ . Let us denote by  $\text{GL}(V)$  the group of linear automorphisms of  $V$ . We choose once and for all a faithful representation  $\rho : G \rightarrow \text{GL}(V)$ .

We denote by  $\text{SL}(V)$  the closed algebraic subgroup of  $\text{GL}(V)$  consisting of linear automorphisms  $g$  of  $V$  such that the  $r^{\text{th}}$  exterior power  $\wedge^r g$  induces the identity on  $\wedge^r V$ .

Set  $\Theta$  to be the quotient stack  $[\mathbb{A}_k^1 / \mathbb{G}_m]$ . Here by convention we always equip  $\mathbb{A}_k^1 = \text{Spec}(k[t])$  with the  $\mathbb{G}_m$ -action that gives the coordinate function  $t$  weight  $-1$ .

## 2.2. Big open sets and torsion-free sheaves.

**Definition 2.1.** *Let  $S$  be a  $k$ -scheme. An open subscheme  $U \subset X_S$  is said to be big (relative to  $S$ ) if for all points  $s \in S$  the complement  $X_s \setminus U_s$  of the fiber  $U_s$  is a closed subset of codimension at least 2 in  $X_s$ .*

**Lemma 2.2.** *Let  $S$  be a  $k$ -scheme. Let  $j : U \hookrightarrow X_S$  be a big open subset in  $X_S$ . Then the following are satisfied*

- (a) *For all points  $x \in X_S \setminus U$ , the local ring  $\mathcal{O}_{X_S, x}$  has depth at least 2.*
- (b)  *$U$  is scheme theoretically dense in  $X_S$ .*
- (c) *Let  $p : Y \rightarrow X_S$  be an affine morphism of finite type. Any section of  $p$  defined over  $U$  extends uniquely to a section defined over the whole  $X_S$ .*

*Proof.* (a) Let  $x \in X_S \setminus U$  be a point with image  $s$  in  $S$ . By assumption the point  $x$  has codimension at least 2 in  $X_s$ . Since  $X_s$  is smooth, this means that  $\mathcal{O}_{X_s, x}$  has depth at least 2. So there exists a regular sequence  $\bar{f}_1, \bar{f}_2$  of length 2 in  $\mathcal{O}_{X_s, x}$ . Lift this sequence to a pair of elements  $f_1, f_2 \in \mathcal{O}_{X_S, x}$ . Since  $\mathcal{O}_{X_S, x}$  is flat over  $\mathcal{O}_{S, s}$ , we can apply [Sta21, Tag 00ME] twice to conclude that  $f_1, f_2$  is a regular sequence. First we apply [Sta21, Tag 00ME] with  $N = M = \mathcal{O}_{X_S, x}$  and  $u = f_1$ . Then we apply this result a second time with  $N = M = \mathcal{O}_{X_S, x}/(f_1)$  and  $u = f_2$ .

(b) By definition, the scheme theoretic closure of  $U$  is cut out by the ideal sheaf of  $\mathcal{O}_{X_S}$  that locally consists of sections whose restriction to  $U$  vanishes. In other words, it is cut out by the kernel of the unit  $\mathcal{O}_{X_S} \rightarrow j_* j^* \mathcal{O}_{X_S}$ . By part (a) every point outside  $U$  has depth at least 2. Therefore the unit map  $\mathcal{O}_{X_S} \rightarrow j_* j^* \mathcal{O}_{X_S}$  is an isomorphism by [Sta21, Tag 0E9I]. We conclude that the scheme theoretic closure is cut out by the 0 ideal, i.e. it is the whole  $X_S$ .

(c) Uniqueness of the extension follows because  $U$  is scheme theoretically dense in  $X_S$  and the morphism  $Y \rightarrow X_S$  is separated. It remains to show existence. Since extensions are unique, any set of local extensions will glue. Therefore we can work Zariski locally. Let  $V$  be an affine open subset of  $X_S$ . Since  $Y$  is of finite type over  $X_S$ , the scheme  $Y_V$  can be viewed as a closed subscheme of  $\mathbb{A}_V^n$  for some  $n$ . Since  $U \cap V$  is scheme theoretically dense inside  $V$ , any extension of the section to  $\mathbb{A}_V^n$  will automatically factor through the closed subscheme  $Y_V$ . It therefore suffices to extend the composition  $p : U \cap V \rightarrow Y_{U \cap V} \hookrightarrow \mathbb{A}_{U \cap V}^n$  to a map  $\tilde{p} : V \rightarrow \mathbb{A}_V^n$ . This amounts to taking a section of  $\mathcal{O}_{U \cap V}^{\oplus n}$  and extending to a section of  $\mathcal{O}_V^{\oplus n}$ . This in turn follows from [Sta21, Tag 0E9I] applied to the sheaf  $\mathcal{O}_V^{\oplus n}$ . □



For the following definition, recall that the rank of a torsion-free sheaf on an integral scheme is the dimension of the generic fiber.

**Definition 2.3.** *Let  $S$  be a  $k$ -scheme. Let  $\mathcal{F}$  be a  $\mathcal{O}_{X_S}$ -module. We say that  $\mathcal{F}$  is a relative torsion-free sheaf on  $X_S$  if it is  $S$ -flat, finitely presented, and for all points  $s \in S$  the fiber  $\mathcal{F}|_{X_s}$  is torsion-free. We say that  $\mathcal{F}$  has rank  $h$  if all of the fibers  $\mathcal{F}|_{X_s}$  have rank  $h$ .*

**Remark 2.4.** *From now on we will drop the adjective “relative” whenever it is clear from context.*

**Lemma 2.5.** *Let  $\mathcal{F}$  be a torsion-free sheaf on  $X_S$ . Then, there exists a big open subset  $U \subset X_S$  such that  $\mathcal{F}|_U$  is locally-free.*

*Proof.* Let  $U$  denote the (maximal) open subset of  $X_S$  where  $\mathcal{F}$  is locally-free. Choose  $s \in S$ . Since  $\mathcal{F}$  is  $S$ -flat and finitely presented, we can apply [Sta21, Tag 0CZR] to conclude that  $U \cap X_s$  coincides with the set of points of  $X_s$  where the fiber  $\mathcal{F}|_{X_s}$  is locally-free. Since  $X_s$  is smooth, all points of codimension 1 are discrete valuation rings. Therefore the torsion-free sheaf  $\mathcal{F}|_{X_s}$  is automatically free at all codimension  $\leq 1$  points. We conclude that the closed complement of  $U \cap X_s$  has codimension at least 2, as desired.  $\square$

Next we recall the notion of determinant for torsion-free sheaves, as in [Kob87, 5.6]. Let  $Y$  be a  $k$ -scheme and  $\mathcal{G}$  be a finitely presented  $\mathcal{O}_Y$ -module. Let  $h$  be a nonnegative integer. There is a natural action of the symmetric group  $S_h$  of  $h$ -letters on the  $h$ -fold tensor  $\mathcal{G}^{\otimes h}$ , given by permuting the indexes. Let  $\text{sign} : S_h \rightarrow \{1, -1\}$  denote the sign character.

**Definition 2.6.** *Let  $Y$  and  $\mathcal{G}$  be as in the paragraph above. We denote by  $\mathcal{G}_{\text{Alt}, h}$  the  $\mathcal{O}_Y$ -submodule of  $\mathcal{G}^{\otimes h}$  that is locally generated over each affine open by sections of the form  $\sigma \cdot (x_1 \otimes x_2 \otimes \cdots \otimes x_h) - \text{sign}(\sigma) x_1 \otimes x_2 \otimes \cdots \otimes x_h$  with  $\sigma \in S_h$ . We denote by  $\bigwedge^h \mathcal{G}$  the quotient sheaf  $\mathcal{G}^{\otimes h} / \mathcal{G}_{\text{Alt}, h}$ .*

Note that  $\bigwedge^h \mathcal{G}$  is a finitely presented  $\mathcal{O}_Y$ -sheaf by definition. The formation of  $\bigwedge^h \mathcal{G}$  is compatible with arbitrary base-change, because taking pullbacks is right exact and commutes with tensor products.

**Definition 2.7.** *Let  $S$  be a  $k$ -scheme and let  $\mathcal{F}$  be a torsion-free sheaf of rank  $h$  on  $X_S$ . We define  $\det(\mathcal{F}) := \left( \bigwedge^h \mathcal{F} \right)^{\vee\vee}$ .*

For all morphisms of  $k$ -schemes  $T \rightarrow S$  there is a canonical morphism  $\det(\mathcal{F})|_{X_T} \rightarrow \det(\mathcal{F}|_{X_T})$ . It turns out that this is an isomorphism, as shown in the following lemma.

**Lemma 2.8.** *Let  $S$  be a  $k$ -scheme and let  $\mathcal{F}$  be a torsion-free sheaf on  $X_S$  such that the rank of the fibers is constant. Then:*

- (a) *Let  $j : U \hookrightarrow X_S$  be a big open subscheme such that  $\mathcal{F}$  is locally-free. Then  $\det(\mathcal{F}) = j_*(\det(\mathcal{F}|_U))$ , where  $\det(\mathcal{F}|_U)$  on the right hand side denotes the usual determinant of a locally-free sheaf.*
- (b)  *$\det(\mathcal{F})$  is a line bundle on  $X_S$ .*
- (c) *For all morphisms of  $k$ -schemes  $T \rightarrow S$ , the canonical morphism  $\det(\mathcal{F})|_{X_T} \rightarrow \det(\mathcal{F}|_{X_T})$  is an isomorphism.*

*Proof.* (a) By construction,  $\det(\mathcal{F})|_U$  coincides with the usual determinant of  $\mathcal{F}|_U$  as a locally-free sheaf. Therefore it suffices to show that the unit  $\det(\mathcal{F}) \rightarrow j_*(\det(\mathcal{F})|_U)$  is an isomorphism. By [Sta21, Tag 0E9I] it suffices to check that the sheaf  $\det(\mathcal{F}) = \text{Hom} \left( \left( \bigwedge^h \mathcal{F} \right)^\vee, \mathcal{O}_{X_S} \right)$  has depth at least 2 at all points in the complement of  $U$ . Notice that this follows from (2) in [Sta21, Tag 0AV5], since  $\mathcal{O}_{X_S}$  has depth at least 2 at all points in the complement of  $U$  by Lemma 2.2.

- (b) One can define a notion of determinant  $d(\mathcal{F})$  using a finite resolution by locally-free sheaves, as in [HL97, pg. 36,37]. By definition,  $d(\mathcal{F})$  is a line bundle that agrees with  $\det(\mathcal{F})$  on the open subset  $U$ . By the same proof as in part (a) applied to the line bundle  $d(\mathcal{F})$ , it follows that  $d(\mathcal{F}) \cong j_*(\det(\mathcal{F}|_U))$ . We conclude that  $\det(\mathcal{F}) = j_*(\det(\mathcal{F}|_U)) = d(\mathcal{F})$  is a line bundle.
- (c) This follows because the notion of determinant  $d(\mathcal{F})$  defined in [HL97, pg. 36,37] using a fixed choice of a resolution can be seen to be compatible with base-change, by pulling back the resolution. □

Recall that  $r$  is the dimension of the vector space  $V$  that we use to define the faithful representation of the group  $G$ . The following is an open substack of the algebraic stack of coherent sheaves on  $X$  as in [LMB00, 4.6.2].

**Definition 2.9.** We define  $\mathrm{Coh}_r^{tf}(X)$  to be the pseudofunctor from  $(\mathrm{Aff}_k)^{op}$  into groupoids defined as follows. For any affine scheme  $T \in \mathrm{Aff}_k$ , we set

$$\mathrm{Coh}_r^{tf}(X)(T) = [ \text{groupoid of } T\text{-flat relative torsion-free sheaves } \mathcal{F} \text{ of rank } r \text{ on } X_T ]$$

**Proposition 2.10.**  $\mathrm{Coh}_r^{tf}(X)$  is an algebraic stack with affine diagonal and locally of finite type over  $k$ .

*Proof.* Since the rank of a family of torsion-free sheaves is a locally constant function on the base,  $\mathrm{Coh}_r^{tf}(X)$  is an open substack of the stack  $\mathrm{Coh}^{tf}(X)$  that parametrizes all torsion-free sheaves. The proposition follows, because  $\mathrm{Coh}^{tf}(X)$  is an algebraic stack with affine diagonal and locally of finite type over  $k$  [HLHJ21, Prop. 2.5]. □

**Remark 2.11.** The stack  $\mathrm{Coh}_r^{tf}(X)$  contains the stack  $\mathrm{Bun}_r(X)$  of rank  $r$  vector bundles on  $X$  as an open substack.

**2.3. The scheme of  $G$ -reductions.** Let  $S$  be a  $k$ -scheme and let  $\mathcal{F} \in \mathrm{Coh}_r^{tf}(X)(S)$  be a rank  $r$  torsion-free sheaf on  $X_S$ . In this subsection we modify the construction in [Sch02] in order to define a scheme of  $G$ -reductions of  $\mathcal{F}$ . In our convention the representation  $\rho$  is the dual of the representation in [Sch02].

Let  $Y$  be a  $k$ -scheme. Let  $\mathcal{G}$  be a finitely presented  $\mathcal{O}_Y$ -module. We can form the universal graded associative algebra  $T(\mathcal{G}) := \bigoplus_{j=0}^{\infty} \mathcal{G}^{\otimes j}$ . There is a graded ideal that on each affine open is generated by sections of the form  $x \otimes y - y \otimes x$ . The quotient of  $T(\mathcal{G})$  by this ideal is the symmetric algebra  $\mathrm{Sym}^{\bullet}(\mathcal{G})$ . By definition  $\mathrm{Sym}^{\bullet}(\mathcal{G})$  is a graded commutative  $\mathcal{O}_Y$ -algebra that is locally finitely presented as an algebra.

Suppose that the sheaf  $\mathcal{G}$  admits a  $k[G]$ -comodule structure. This induces a comodule structure on  $T(\mathcal{G})$ , where we equip each graded component  $\mathcal{G}^{\otimes j}$  with the tensor comodule structure. The ideal used to form  $\mathrm{Sym}^{\bullet}(\mathcal{G})$  is a  $k[G]$ -subcomodule. It follows that the quotient  $\mathrm{Sym}^{\bullet}(\mathcal{G})$  acquires a  $k[G]$ -comodule structure that is compatible with the multiplication and the grading.

**Definition 2.12.** Let  $S$  be a  $k$ -scheme and let  $\mathcal{F}$  be a torsion-free sheaf of rank  $r$  on  $X_S$ . We set

$$E_V(\mathcal{F}) := \mathrm{Sym}^{\bullet}(\mathcal{F} \otimes_k V^{\vee}) \otimes_{\mathcal{O}_{X_S}} \mathrm{Sym}^{\bullet}(\det(\mathcal{F})^{\vee} \otimes_k \det(V))$$

We also define  $H(\mathcal{F}, V) := \underline{\mathrm{Spec}}_{X_S}(E_V(\mathcal{F}))$ . Note that the structure morphism  $H(\mathcal{F}, V) \rightarrow X_S$  is affine and of finite presentation.

The vector space  $V$  acquires a left  $k[G]$ -comodule structure  $V \rightarrow V \otimes k[G]$  via the representation  $\rho : G \rightarrow \mathrm{GL}(V)$ . We equip  $V^{\vee}$  with the dual right comodule structure  $c : V^{\vee} \rightarrow V^{\vee} \otimes k[G]$ . This induces a comodule structure  $\mathcal{F} \otimes_k V^{\vee} \rightarrow (\mathcal{F} \otimes_k V^{\vee}) \otimes k[G]$

given by  $\text{id}_{\mathcal{F}} \otimes c$ . By the discussion above, the graded algebra  $\text{Sym}^\bullet(\mathcal{F} \otimes_k V^\vee)$  inherits a right  $k[G]$ -comodule structure.

On the other hand,  $\det(V)$  acquires a left  $k[G]$ -comodule structure via the determinant representation of  $V$ . By post-composing with the antipode of the Hopf algebra  $k[G]$  (i.e. pre-composing the representation with inversion on  $G$ ), we get a right  $k[G]$ -comodule structure on  $\det(V)$ . In a similar way as above, this induces a right  $k[G]$ -comodule structure on the algebra  $\text{Sym}^\bullet(\det(\mathcal{F})^\vee \otimes_k \det(V))$ .

By taking the tensor comodule structure for the  $k[G]$ -comodules defined above, we can equip  $E_V(\mathcal{F})$  with a right  $k[G]$ -comodule structure

$$c_{E_V(\mathcal{F})} : E_V(\mathcal{F}) \rightarrow E_V(\mathcal{F}) \otimes_k k[G]$$

Since this comodule structure is compatible with the algebra multiplication, it translates into a left action of the algebraic group  $G$  on the scheme  $H(\mathcal{F}, V)$ .

Consider the subalgebra  $E_V(\mathcal{F})^G$  of  $G$ -invariants of  $E_V(\mathcal{F})$ . Recall that this is defined over each affine open as the  $\mathcal{O}_{X_S}$ -subalgebra of  $E_V(\mathcal{F})$  consisting of sections  $s$  such that  $c_{E_V(\mathcal{F})}(s) = s \otimes 1_{k[G]}$ . Since the characteristic of  $k$  is 0 and  $G$  is reductive, the algebra of invariants  $E_V(\mathcal{F})^G$  is a locally finitely presented  $\mathcal{O}_{X_S}$ -algebra and its formation is compatible with arbitrary base-change. This can be checked affine locally, and over each affine it follows from the existence of the Reynolds operator [MFK94, Chpt. 1, §1,2]. By definition, the scheme  $\underline{\text{Spec}}_{X_S}((E_V(\mathcal{F}))^G)$  is the affine GIT quotient  $H(\mathcal{F}, V)//G$  relative to  $X_S$ .

Let  $U$  be a big open subscheme of  $X_S$  where  $\mathcal{F}$  is locally-free. The restriction  $H(\mathcal{F}, V)|_U$  is, by definition, the scheme

$$H := \text{Hom}(\mathcal{F}|_U, V \otimes_k \mathcal{O}_U) \times \text{Hom}(\det(\mathcal{F}|_U)^\vee, \det(V^\vee) \otimes_k \mathcal{O}_U)$$

parametrizing pairs  $(f_1, f_2)$  of homomorphisms of vector bundles

$$f_1 : \mathcal{F}|_U \rightarrow V \otimes_k \mathcal{O}_U \quad \text{and} \quad f_2 : \det(\mathcal{F}|_U)^\vee \rightarrow \det(V^\vee) \otimes_k \mathcal{O}_U$$

This means that there is a universal pair  $(u_1, u_2)$  of morphisms

$$u_1 : \mathcal{F}|_H \rightarrow V \otimes_k \mathcal{O}_H \quad \text{and} \quad u_2 : \det(\mathcal{F}|_H)^\vee \rightarrow \det(V^\vee) \otimes_k \mathcal{O}_H$$

We take the  $r^{\text{th}}$  wedge of  $u_1$  to get a universal determinant morphism.

$$\wedge^r u_1 : \det(\mathcal{F}|_H) \rightarrow \det(V) \otimes_k \mathcal{O}_H$$

We set  $\delta_U := (\wedge^r u_1) \otimes u_2$ . Using the canonical trivializations  $\det(V) \otimes \det(V^\vee) \xrightarrow{\sim} k$  and  $\det(\mathcal{F}|_H) \otimes \det(\mathcal{F}|_H)^\vee \xrightarrow{\sim} \mathcal{O}_H$ , we can view this as a map  $\delta_U : \mathcal{O}_H \rightarrow \mathcal{O}_H$ . We can then form the map  $\delta_U - 1 : \mathcal{O}_H \rightarrow \mathcal{O}_H$  obtained by subtracting the identity of  $\mathcal{O}_H$ . Notice that  $\delta_U - 1$  cuts out the divisor on  $H$  given by the locus where  $u_1$  is an isomorphism and  $u_2$  is the determinant of the inverse dual of  $u_1$ . In particular the vanishing locus of  $\delta_U - 1$  is canonically isomorphic to the scheme parametrizing linear isomorphisms of vector bundles  $\text{Iso}(\mathcal{F}|_U, V \otimes_k \mathcal{O}_U)$ . This identification is given by forgetting  $u_2$ .

By construction, the composition  $\mathcal{O}_{H//G} \hookrightarrow \mathcal{O}_H \xrightarrow{\delta_U - 1} \mathcal{O}_H$  lands in the  $\mathcal{O}_U$ -subalgebra of invariants  $(E_V(\mathcal{F})|_U)^G$ . Therefore this composition induces a well-defined map  $\alpha_U : \mathcal{O}_{H//G} \rightarrow \mathcal{O}_{H//G}$  that cuts out an effective divisor on  $H//G$ . We denote this closed subscheme by  $(H//G)_{\alpha_U=0} \hookrightarrow H//G$ .

**Remark 2.13.** *Alternatively, we can think of the construction above as follows. The divisor  $D$  cut out by the equation  $\delta_U - 1$  on  $H$  is  $G$ -stable. Then  $(H//G)_{\alpha_U=0}$  is the unique closed subscheme in the GIT quotient  $H//G$  such that the preimage recovers  $D$ . It follows that  $(H//G)_{\alpha_U=0} = (H_{\delta_U=1})//G$ .*

**Definition 2.14.**  $\text{Red}_G(\mathcal{F})$  is defined to be the scheme theoretic closure of  $(H//G)_{\alpha_U=0}$  inside  $H(\mathcal{F}, V)//G$ . In other words,  $\text{Red}_G(\mathcal{F})$  is the scheme theoretic image of the composition

$$(H//G)_{\alpha_U=0} \hookrightarrow H//G \hookrightarrow H(\mathcal{F}, V)//G$$

where the second map  $H//G \hookrightarrow H(\mathcal{F}, V)//G$  is the open immersion that comes from identifying  $H//G \xrightarrow{\sim} (H(\mathcal{F}, V)//G)|_U$  via the compatibility of taking  $G$ -invariants with flat base-change.

**Remark 2.15.** It should be noted that the construction of  $\text{Red}_G(\mathcal{F})$  does not depend on the choice of  $U$ . This can be seen as follows. For any two big open subsets  $U_1$  and  $U_2$  of  $X_S$ , the intersection  $U_1 \cap U_2$  is also big. Since the element  $\alpha_U$  is canonically defined above, the restrictions of  $\alpha_{U_1}$  and  $\alpha_{U_2}$  to the subset  $U_1 \cap U_2$  agree with  $\alpha_{U_1 \cap U_2}$ . Notice that  $H$  is always flat over  $U$ , because it is locally given by a polynomial algebra. It follows that  $H//G$  is also flat over  $U$ , because taking  $G$  invariants is exact by the existence of the Reynolds operator. Since  $U$  is scheme theoretically dense in  $X_S$  by Lemma 2.2, it follows that the open  $(H//G)|_{U_1 \cap U_2}$  is scheme theoretically dense in  $(H//G)|_{U_1}$  and  $(H//G)|_{U_2}$ . In particular the closures of  $(H//G)_{\alpha_{U_1}=0}$  and  $(H//G)_{\alpha_{U_2}=0}$  both agree with the closure of  $(H//G)_{\alpha_{U_1 \cap U_2}=0}$ .

The following lemma is an immediate consequence of the constructions above.

**Lemma 2.16.** The scheme  $\text{Red}_G(\mathcal{F})$  is affine and of finite presentation over  $X_S$ . The formation of  $\text{Red}_G(\mathcal{F})$  is compatible with base-change. In other words, for any morphism of  $k$ -schemes  $T \rightarrow S$  there is a canonical isomorphism  $\text{Red}_G(\mathcal{F}) \times_{X_S} X_T \xrightarrow{\sim} \text{Red}_G(\mathcal{F}|_{X_T})$ .  $\square$

Let  $U$  be a big open subset of  $X_S$  where  $\mathcal{F}$  is locally-free. Since  $\mathcal{F}|_U$  is a vector bundle of rank  $r$  on  $U$ , we can naturally associate to it a  $\text{GL}(V)$ -bundle  $\mathcal{P}$  on  $U$ .

**Proposition 2.17.** Let  $\mathcal{F}$  and  $U$  be as above. Let  $\mathcal{P}$  denote the corresponding  $\text{GL}(V)$ -bundle on  $U$ . Then there is a canonical identification of  $U$ -schemes  $\text{Red}_G(\mathcal{F})|_U \xrightarrow{\sim} \mathcal{P}/G$ . Here the quotient on the right is formed via the inclusion  $\rho : G \hookrightarrow \text{GL}(V)$ .

*Proof.* Let  $H_{\delta_U=1}$  denote the closed subscheme of  $H$  cut out by the equation  $\delta_U - 1$ , as described in the construction of  $\text{Red}_G(\mathcal{F})$  above. By definition,  $\text{Red}_G(\mathcal{F})|_U$  is the affine GIT quotient  $(H_{\delta_U=1})//G$ . We have seen that  $H_{\delta_U=1}$  is canonically isomorphic to the  $U$ -scheme  $\text{Iso}(\mathcal{F}|_U, V \otimes_k \mathcal{O}_U)$  parametrizing isomorphisms of vector bundles. This identification respects the natural left  $G$ -actions defined on both schemes via the inclusions  $\rho : G \rightarrow \text{GL}(V)$ .

The map that takes an isomorphism  $\varphi : \mathcal{F}|_U \oplus \det(\mathcal{F})^\vee|_U \xrightarrow{\sim} (V \oplus \det(V^\vee)) \otimes_k \mathcal{O}_U$  to its inverse in  $\text{Hom}((V \oplus \det(V^\vee)) \otimes_k \mathcal{O}_U, \mathcal{F}|_U \oplus \det(\mathcal{F})^\vee|_U)$  induces an isomorphism of schemes

$$(-)^{-1} : \text{Iso}(\mathcal{F}|_U, V \otimes_k \mathcal{O}_U) \xrightarrow{\sim} \text{Iso}(V \otimes_k \mathcal{O}_U, \mathcal{F}|_U)$$

The right-hand side admits a right action of  $\text{GL}(V)$  given by precomposition. By definition  $\text{Iso}(V \otimes_k \mathcal{O}_U, \mathcal{F}|_U)$  is the total space of the  $\text{GL}(V)$ -bundle  $\mathcal{P}$  equipped with its right  $\text{GL}(V)$ -action. The inclusion  $\rho : G \hookrightarrow \text{GL}(V)$  induces a right  $G$ -action on  $\mathcal{P}$ . The inversion map  $(-)^{-1}$  above intertwines the left  $G$ -action on the left hand side and the right  $G$ -action on the right hand side.

Since the representation  $\rho$  is faithful, it follows that  $G$  acts properly and freely on  $\mathcal{P}$ . Therefore the GIT quotient  $\mathcal{P}//G$  and the quotient as étale sheaves  $\mathcal{P}/G$  coincide. This way we get the sought after canonical identification of  $\text{Red}_G(\mathcal{F})|_U \xrightarrow{\sim} \mathcal{P}/G$ .  $\square$

The following proposition explains the connection between the scheme  $\text{Red}_G(\mathcal{F})$  and  $G$ -reductions of structure group.

**Proposition 2.18.** *For any big open subset  $U \subset X_S$  where  $\mathcal{F}$  is locally-free, there is a bijection between the following two sets (A), (B):*

- (A) *The set of sections of the structure morphism  $\text{Red}_G(\mathcal{F}) \rightarrow X_S$ .*
- (B) *The set of  $G$ -reductions of structure group for the  $\text{GL}(V)$ -bundle  $\mathcal{P}$  corresponding to  $\mathcal{F}|_U$ .*

*Proof.*  $\text{Red}_G(\mathcal{F}) \rightarrow X_S$  is an affine morphism of finite type. Lemma 2.2 shows that the restriction of maps induces a bijection between sections  $\sigma : X_S \rightarrow \text{Red}_G(\mathcal{F})$  defined over the whole  $X_S$  and sections  $\sigma_U : U \rightarrow \text{Red}_G(\mathcal{F})$  defined over  $U$ . By Proposition 2.17, sections of  $\text{Red}_G(\mathcal{F})|_U \rightarrow U$  are in canonical correspondence with sections of  $\mathcal{P}/G \rightarrow U$  via the inversion map. A section of  $\mathcal{P}/G \rightarrow U$  is the same as a  $G$ -reduction of structure group for  $\mathcal{P}$ . This establishes the canonical bijection between (A) and (B).  $\square$

### 3. PRINCIPAL $\rho$ -SHEAVES

#### 3.1. Principal $\rho$ -sheaves and the moduli stack $\text{Bun}_\rho(X)$ .

**Definition 3.1.** *Let  $S$  be a  $k$ -scheme. A principal  $\rho$ -sheaf on  $X_S$  consists of a pair  $(\mathcal{F}, \sigma)$ , where*

- (1)  *$\mathcal{F}$  is a family of torsion-free sheaves of rank  $r$  on  $X_S$ .*
- (2)  *$\sigma$  is a section of the structure morphism  $\text{Red}_G(\mathcal{F}) \rightarrow X_S$ .*

**Remark 3.2.** *We will often omit the word “principal” and refer to these objects simply as  $\rho$ -sheaves.*

By Proposition 2.18, one can think of a  $\rho$ -sheaf over  $X_S$  as the data of

- (1) A torsion-free sheaf  $\mathcal{F}$  of rank  $r$  on  $X_S$ .
- (2) A reduction of structure group to  $G$  for the  $\text{GL}(V)$  bundle  $\mathcal{F}|_U$ , where  $U$  is (any) big open subscheme of  $X_S$  such that  $\mathcal{F}|_U$  is locally-free.

**Remark 3.3.** *Assume that the image of the representation  $\rho$  is contained in the subgroup  $\text{SL}(V) \subset \text{GL}(V)$ . Then our  $\rho$ -sheaves can be viewed as singular principal  $G$ -bundles [GLSS08] for the dual representation  $\rho^\vee$ . This readily follows from the description given above.*

**Definition 3.4.** *Let  $(\mathcal{F}_1, \sigma_1)$  and  $(\mathcal{F}_2, \sigma_2)$  be two  $\rho$ -sheaves on  $X_S$ . By construction, an isomorphism  $\varphi : \mathcal{F}_1 \xrightarrow{\sim} \mathcal{F}_2$  induces an isomorphism of  $G$ -reduction schemes  $\text{Red}_G(\mathcal{F}_1) \xrightarrow{\sim} \text{Red}_G(\mathcal{F}_2)$ . We say that  $\varphi$  is an isomorphism of the  $\rho$ -sheaves if the sections  $\sigma_1$  and  $\sigma_2$  are identified under this induced isomorphism.*

Using this notion of isomorphism, we endow the class of  $\rho$ -sheaves on  $X_S$  with the structure of a groupoid. By Lemma 2.16, the formation of  $\text{Red}_G(\mathcal{F})$  is compatible with base-change. In particular, it makes sense to pull back principal  $\rho$ -sheaves under a morphism of  $k$ -schemes  $T \rightarrow S$ .

**Definition 3.5.** *We define  $\text{Bun}_\rho(X)$  to be the pseudofunctor from  $(\text{Aff}_k)^{op}$  into groupoids defined as follows. For any affine scheme  $T \in \text{Aff}_k$ , we set*

$$\text{Bun}_\rho(X)(T) = [ \text{groupoid of } \rho\text{-sheaves } (\mathcal{F}, \sigma) \text{ on } X_T ]$$

There is a natural forgetful morphism  $\text{Bun}_\rho(X) \rightarrow \text{Coh}_r^{tf}(X)$  given by  $(\mathcal{F}, \sigma) \mapsto \mathcal{F}$ .

**Proposition 3.6.** *The forgetful morphism  $\text{Bun}_\rho(X) \rightarrow \text{Coh}_r^{tf}(X)$  is schematic, affine and of finite type. In particular  $\text{Bun}_\rho(X)$  is an algebraic stack with affine diagonal and locally of finite type over  $k$ .*

*Proof.* Let  $T \in \text{Aff}_k$ . Choose a morphism  $T \rightarrow \text{Coh}_r^{tf}(X)$  represented by a relative torsion-free sheaf  $\mathcal{F}$  on  $X_T$ . The fiber product  $\text{Bun}_\rho(X) \times_{\text{Coh}_r^{tf}(X)} T$  is the functor that parametrizes sections of the structure morphism  $\text{Red}_G(\mathcal{F}) \rightarrow X_T$ . Since we know that  $\text{Red}_G(\mathcal{F}) \rightarrow X_T$  is affine and of finite presentation, we can apply [Beh91, Prop. 4.4.1] (or [HR15, Lemma 2.3 (i)] + [HR19, Thm. 1.3 (ii)]) to conclude that  $\text{Bun}_\rho(X) \times_{\text{Coh}_r^{tf}(X)} T$  is represented by a scheme that is relatively affine and of finite type over  $T$ .  $\square$

Recall that we have an open immersion  $\text{Bun}_r(X) \hookrightarrow \text{Coh}_r^{tf}(X)$  from the stack  $\text{Bun}_r(X)$  of rank  $r$  vector bundles on  $X$ . The following shows that the same holds for  $\text{Bun}_\rho(X)$  and the stack of principal  $G$ -bundles.

**Proposition 3.7.** *There is a Cartesian square*

$$\begin{array}{ccc} \text{Bun}_G(X) & \hookrightarrow & \text{Bun}_\rho(X) \\ \downarrow \rho_* & & \downarrow \\ \text{Bun}_r(X) & \hookrightarrow & \text{Coh}_r^{tf}(X) \end{array}$$

where  $\text{Bun}_G(X)$  is the stack of principal  $G$ -bundles on  $X$  and  $\rho_*$  is the map on stacks induced by extension of the structure group via the representation  $\rho$ .

*Proof.* Let  $S \in \text{Aff}_k$ . The groupoid  $\text{Bun}_r(X)(S)$  consists of points  $\mathcal{F} \in \text{Coh}_r^{tf}(X)(S)$  where  $\mathcal{F}$  is locally-free on the entirety of  $X_S$ . Applying Proposition 2.18 with  $U = X_S$ , we see that the fiber product  $\text{Bun}_r(X) \times_{\text{Coh}_r^{tf}(X)} \text{Bun}_\rho(X)(S)$  parametrizes pairs  $(\mathcal{P}, \theta)$ , where  $\mathcal{P}$  is a  $\text{GL}(V)$ -bundle on  $X_S$  and  $\theta$  is a reduction of structure group to  $G$ . This groupoid is naturally isomorphic to the groupoid of  $G$ -bundles on  $X_S$ .  $\square$

**3.2. Filtrations of  $\rho$ -sheaves.** We will “destackify” the notion of  $\Theta$ -filtration [HL14, Hei17, AHLH19] and explain what it means in the context of  $\rho$ -sheaves. For this subsection we will work over a field extension  $K \supset k$ .

Let us recall the definition of a  $\Theta$ -filtration. We denote by  $\Theta_K$  the base-change  $[\mathbb{A}_K^1/\mathbb{G}_m]$ . The stack  $\Theta_K$  has a unique closed point 0 corresponding to the origin of  $\mathbb{A}_K^1$ . It also has an open point 1, which is the image of the  $K$ -point 1 in  $\mathbb{A}_K^1$ . Let  $\mathcal{M}$  be a stack over  $k$  and let  $p \in \mathcal{M}(K)$  be a  $K$ -point. A  $\Theta$ -filtration of the point  $p$  is a morphism of stacks  $f : \Theta_K \rightarrow \mathcal{M}$  along with an isomorphism  $f(1) \xrightarrow{\sim} p$ . The graded point  $f|_{[0/\mathbb{G}_m]}$  is called the associated graded of the filtration. We say that the filtration is nondegenerate if the action of  $\mathbb{G}_m$  on the object  $f(0)$  is nontrivial.

We start by giving a concrete description of  $\Theta$ -filtrations in the stack  $\text{Coh}_r^{tf}(X)$ .

**Proposition 3.8.** *Let  $\mathcal{F}$  be a  $K$ -point of  $\text{Coh}_r^{tf}(X)$ . A  $\Theta$ -filtration  $f : \Theta_K \rightarrow \text{Coh}_r^{tf}(X)$  of  $\mathcal{F}$  corresponds to a  $\mathbb{Z}$ -filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  by  $\mathcal{O}_{X_K}$ -subsheaves of  $\mathcal{F}$  satisfying the following properties*

- (1)  $\mathcal{F}_{m+1} \subset \mathcal{F}_m$ .
- (2)  $\mathcal{F}_m = 0$  for  $m \gg 0$  and  $\mathcal{F}_m = \mathcal{F}$  for  $m \ll 0$ .
- (3)  $\mathcal{F}_m/\mathcal{F}_{m+1}$  is torsion-free.

An isomorphism between two such filtrations is an automorphism of  $\mathcal{F}$  that identifies the filtrations. The graded object at the origin  $f|_{[0/\mathbb{G}_m]} : [0/\mathbb{G}_m] \rightarrow \text{Coh}_r^{tf}(X)$  is the associated graded sheaf  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m/\mathcal{F}_{m+1}$ .

*Proof.* By definition, a  $\Theta$ -filtration of  $\mathcal{F}$  consists of the following data

- (a) A  $\mathbb{G}_m$ -equivariant family  $\tilde{\mathcal{F}}$  of torsion-free sheaves on  $X \times \mathbb{A}_K^1$ .

(b) An identification  $\tilde{\mathcal{F}}|_{X \times 1} \xrightarrow{\sim} \mathcal{F}$ .

For any  $\mathbb{Z}$ -filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  satisfying (1) and (2) above, we can associate a  $\mathbb{G}_m$ -equivariant coherent sheaf on  $X \times \mathbb{A}_K^1$  via the Rees construction [HL14, Prop. 1.0.1]. Namely, we can set  $\tilde{\mathcal{F}} := \bigoplus_{m \in \mathbb{Z}} t^{-m} \mathcal{F}_m$ . Here  $\mathbb{G}_m$  acts on  $t^{-m} \mathcal{F}_m$  with weight  $m$ . By construction we have that  $\tilde{\mathcal{F}}|_{X \times (\mathbb{A}_K^1 \setminus \{0\})}$  is the constant family  $\mathcal{F}|_{X \times (\mathbb{A}_K^1 \setminus \{0\})}$ . All  $\mathbb{G}_m$ -equivariant coherent sheaves  $\tilde{\mathcal{F}}$  on  $X \times \mathbb{A}_K^1$  along with an isomorphism  $\tilde{\mathcal{F}}|_{X \times 1} \xrightarrow{\sim} \mathcal{F}$  arise by applying the Rees construction to a  $\mathbb{Z}$ -filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  satisfying conditions (1) and (2) above [HL14, Prop. 1.0.1].

The restriction  $\tilde{\mathcal{F}}|_{X \times 0}$  is the associated graded  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m / \mathcal{F}_{m+1}$ . So we see that  $\tilde{\mathcal{F}}$  is a family of torsion-free sheaves if and only if  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m / \mathcal{F}_{m+1}$  is torsion-free on  $X_K$ . This, in turn, is equivalent to condition (3) above.  $\square$

For any filtration  $\Theta_K \rightarrow \text{Bun}_\rho(X)$  we can post-compose with the forgetful morphism  $\text{Bun}_\rho(X) \rightarrow \text{Coh}_r^{tf}(X)$  to get a filtration  $\Theta_K \rightarrow \text{Coh}_r^{tf}(X)$ . This defines a functor  $F_K$  between the groupoid of  $\Theta$ -filtrations of  $K$ -points of  $\text{Bun}_\rho(X)$  and the groupoid of  $\Theta$ -filtrations of  $K$ -points of  $\text{Coh}_r^{tf}(X)$ .

**Proposition 3.9.** *The functor  $F_K$  defined above is faithful.*

*Proof.* This follows from [AHLH19, Lem. 3.7 (3)], which applies here because the forgetful morphism is affine by Proposition 3.6.  $\square$

This tells us that we can view filtrations of  $\text{Bun}_\rho(X)$  as a subcategory of filtrations in  $\text{Coh}_r^{tf}(X)$ . In other words, if  $(\mathcal{F}, \sigma)$  is a  $\rho$ -sheaf, then a filtration of  $(\mathcal{F}, \sigma)$  is the same thing as a  $\mathbb{Z}$ -filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  of  $\mathcal{F}$  (as in Proposition 3.8) satisfying some ‘‘compatibility conditions’’. We will often refer to  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  as the underlying associated filtration.

Our goal for the remainder of this section is to explain these ‘‘compatibility’’ conditions. First, we need to explain what is a filtration of  $\text{GL}(V)$ -bundles on any scheme  $U$ .

**Proposition 3.10.** *Let  $U$  be a scheme over  $K$ . Let  $\mathcal{F}$  be a  $\text{GL}(V)$ -bundle on  $U$ . Then the following two groupoids are equivalent.*

- (1) *The groupoid of  $\text{GL}(V)$ -bundles  $\mathcal{P}$  on  $U \times_K \Theta_K$  along with an isomorphism of the restriction  $\mathcal{P}|_{U \times_K 1}$  with  $\mathcal{F}$ .*
- (2) *The groupoid of decreasing  $\mathbb{Z}$ -weighted filtrations  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  of  $\mathcal{F}$  by vector subbundles  $\mathcal{F}_m$ . An isomorphism between two filtrations is just an automorphism of  $\mathcal{F}$  that identifies the filtrations.*

*Proof.* The equivalence is given by the Rees construction. The argument is the same as in Proposition 3.8, we omit the details.  $\square$

Now we are ready to give a first description of  $\Theta$ -filtrations for a  $\rho$ -sheaf. Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . A filtration of  $(\mathcal{F}, \sigma)$  will consist of a pair  $((\mathcal{F}_m)_{m \in \mathbb{Z}}, \mathcal{P})$ . Here  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  is a filtration of  $\mathcal{F}$ , as described in Proposition 3.8. Let  $U \hookrightarrow X_K$  be the maximal big open subset of  $X_K$  where the associated graded  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m / \mathcal{F}_{m+1}$  is locally-free. The second element of the pair  $\mathcal{P}$  is a  $G$ -bundle on  $U \times_K \Theta_K$  such that the associated  $\text{GL}(V)$ -bundle  $\rho_*(\mathcal{P})$  on  $U \times_K \Theta_K$  corresponds to the restricted filtration  $(\mathcal{F}_m|_U)_{m \in \mathbb{Z}}$  under the equivalence described in Proposition 3.10.

The filtration  $(\mathcal{F}_m|_U)_{m \in \mathbb{Z}}$  on  $U$  completely determines the underlying associated filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  on  $X_K$ . This is because each  $\mathcal{F}_m$  is saturated on  $\mathcal{F}$ , so it is determined in codimension 1 [Lan75, pg. 90-92]. We can describe  $\mathcal{F}_m$  more explicitly in terms of reflexive hulls. Let  $\tilde{\mathcal{F}}_m$  denote the pushforward of the vector bundle  $\mathcal{F}_m|_U$  under the open immersion  $U \hookrightarrow X_K$ . We have that  $\tilde{\mathcal{F}}_m$  is a subsheaf of the reflexive hull  $\mathcal{F}^{\vee\vee}$  of  $\mathcal{F}$ . Then

we can recover  $\mathcal{F}_m$  as the intersection  $\tilde{\mathcal{F}}_m \cap \mathcal{F}$  taken inside of  $\mathcal{F}^{\vee\vee}$ . In particular the filtration is completely determined by the  $G$ -bundle  $\mathcal{P}$  on  $U \times_K \Theta_K$ . We will, however, keep the underlying associated filtration as part of the notation.

A morphism between two such filtrations  $((\mathcal{F}_m)_{m \in \mathbb{Z}}, \mathcal{P})$  and  $((\mathcal{F}'_m)_{m \in \mathbb{Z}}, \mathcal{P}')$  consists of an isomorphism of  $G$ -bundles  $\varphi : \mathcal{P} \xrightarrow{\sim} \mathcal{P}'$  defined over some big open subset  $U \subset X_K$  where both  $\mathcal{P}$  and  $\mathcal{P}'$  are defined. We require that the associated isomorphism of  $\mathrm{GL}(V)$ -bundles on  $U \times_K \Theta_K$  given by  $\rho_*(\varphi) : (\mathcal{F}_m|_U)_{m \in \mathbb{Z}} \xrightarrow{\sim} (\mathcal{F}'_m|_U)_{m \in \mathbb{Z}}$  admits a (unique) extension to an isomorphism of filtrations  $\tilde{\varphi} : (\mathcal{F}_m)_{m \in \mathbb{Z}} \xrightarrow{\sim} (\mathcal{F}'_m)_{m \in \mathbb{Z}}$  defined over all of  $X_K$ . By the description of the filtrations as intersections of reflexive hulls, it follows that we only need to check that the isomorphism  $\rho_*(\varphi)|_{U \times_K 1}$  extends to an automorphism  $\mathcal{F} \xrightarrow{\sim} \mathcal{F}$ . This is automatically satisfied if  $\mathcal{F}$  is reflexive, but not otherwise.

**Definition 3.11.** *Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . We will denote by  $\mathrm{Filt}(\mathcal{F}, \sigma)$  the groupoid of pairs  $((\mathcal{F}_m)_{m \in \mathbb{Z}}, \mathcal{P})$ , with isomorphisms as described above.*

**Proposition 3.12.** *Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . There is a natural equivalence between the following two groupoids.*

- (1) *The groupoid of  $\Theta$ -filtrations  $f : \Theta_K \rightarrow \mathrm{Bun}_\rho(X)$  of the  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$ .*
- (2) *The groupoid  $\mathrm{Filt}(\mathcal{F}, \sigma)$ .*

*Proof.* The groupoid of  $\Theta$ -filtrations classifies  $\mathbb{G}_m$ -equivariant  $\rho$ -sheaves  $(\tilde{\mathcal{F}}, \tilde{\sigma})$  on  $X \times \mathbb{A}_K^1$  along with an isomorphism of the restriction  $(\tilde{\mathcal{F}}, \tilde{\sigma})|_{X \times 1}$  with the original  $(\mathcal{F}, \sigma)$ . So  $\tilde{\mathcal{F}}$  is a  $\Theta$ -filtration of  $\mathcal{F}$  and  $\tilde{\sigma}$  is a  $\mathbb{G}_m$ -equivariant section  $\tilde{\sigma} : X \times \mathbb{A}_K^1 \rightarrow \mathrm{Red}_G(\tilde{\mathcal{F}})$  extending  $\sigma : X \times 1 \rightarrow \mathrm{Red}_G(\mathcal{F})$ .

By Proposition 3.8, the bundle  $\tilde{\mathcal{F}}$  is obtained by applying the Rees construction to some  $\mathbb{Z}$ -filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$ . Let  $U$  be the maximal open in  $X_K$  where  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m / \mathcal{F}_{m+1}$  is locally-free. Then  $U \times_K \mathbb{A}_K^1$  is a big open in  $X \times \mathbb{A}_K^1$  and  $\tilde{\mathcal{F}}|_{U \times_K \mathbb{A}_K^1}$  is locally-free. By Proposition 2.18, the  $\mathbb{G}_m$ -equivariant section  $\tilde{\sigma}$  is equivalent to the data of a  $\mathbb{G}_m$ -equivariant  $G$ -reduction of the  $\mathrm{GL}(V)$ -bundle  $\tilde{\mathcal{F}}|_{U \times_K \mathbb{A}_K^1}$ . This is by definition a  $\mathbb{G}_m$ -equivariant  $G$ -bundle  $\mathcal{P}$  on  $U \times_K \mathbb{A}_K^1$  such that the associated  $\mathrm{GL}(V)$ -bundle  $\rho_*(\mathcal{P})$  on  $U \times_K \Theta_K$  recovers the restriction of the filtration  $(\mathcal{F}_m|_U)_{m \in \mathbb{Z}}$  under the equivalence in Proposition 3.10. This shows how to define the functor on objects, and it also shows that it is essentially surjective. The definition of the functor on morphisms and the fact that it is fully faithful follows from the definition of the isomorphisms in the groupoid  $\mathrm{Filt}(\mathcal{F}, \sigma)$ .  $\square$

**3.3. Filtrations of  $\rho$ -sheaves in the split case.** We will describe the elements of  $\mathrm{Filt}(\mathcal{F}, \sigma)$  more explicitly in the case when  $G_K$  is split. For that purpose, we need a better description of  $G$ -bundles on  $U \times_K \Theta_K$ . This is done in [Hei17, 1.F], which gives a group theoretic description in terms of parabolic reductions and a generalized Rees construction<sup>3</sup>.

We start by recalling some facts about parabolic subgroups of reductive groups. Let  $H$  be a reductive group over  $K$ . Let  $\lambda$  be a cocharacter  $\lambda : \mathbb{G}_m \rightarrow H$ . We can associate a corresponding parabolic subgroup  $P_\lambda$ . For any  $K$ -scheme  $T$ , its  $T$ -points are

$$P_\lambda(T) = \left\{ p \in H(T) \mid \lim_{x \rightarrow 0} \lambda(x) p \lambda(x)^{-1} \text{ exists} \right\}$$

Furthermore, the cocharacter  $\lambda$  induces a Levi decomposition  $P_\lambda = L_\lambda \ltimes U_\lambda$ , where

$$U_\lambda(T) = \left\{ u \in H(T) \mid \lim_{x \rightarrow 0} \lambda(x) u \lambda(x)^{-1} = 1_H \right\}$$

<sup>3</sup>We note that some of the discussion in [Hei17, 1.F] applies without the hypothesis that  $G_K$  is split. However, the results are more concrete under this assumption.



$$L_\lambda(T) = \left\{ l \in H(T) \mid \lim_{x \rightarrow 0} \lambda(x) l \lambda(x)^{-1} = l \right\}$$

**Definition 3.13.** Let  $H$  be a reductive linear algebraic group over  $K$ . Let  $Y$  be a  $K$ -scheme and let  $\mathcal{H}$  be a  $H$ -bundle on  $Y$ . A weighted parabolic reduction is a pair  $(\lambda, \mathcal{H}_\lambda)$ , where  $\lambda : \mathbb{G}_m \rightarrow H$  is a cocharacter and  $\mathcal{H}_\lambda$  is a reduction of structure group of  $\mathcal{H}$  to the parabolic subgroup  $P_\lambda$ .

Let  $\theta : H \rightarrow F$  be a homomorphism of reductive groups over  $K$ . For every cocharacter  $\lambda : \mathbb{G}_m \rightarrow H$ , we can compose with  $\theta$  to obtain a cocharacter  $\theta \circ \lambda$  of  $F$ . Notice that  $\theta$  restricts to a well defined homomorphism of parabolic subgroups  $\theta_P : P_\lambda \rightarrow P_{\theta \circ \lambda}$ . Suppose that we have an  $H$ -bundle  $\mathcal{H}$  defined on a  $K$ -scheme  $Y$ . Then we can define an  $F$ -bundle  $\theta_* \mathcal{H}$  via extension of structure group. If we have a parabolic reduction  $\mathcal{H}_\lambda$  of  $\mathcal{H}$  to the subgroup  $P_\lambda$ , then by extension of structure group using  $\theta_P$  we get a parabolic reduction of  $\theta_* \mathcal{H}$  to the subgroup  $P_{\theta \circ \lambda}$ . This way we can define a map  $\theta_*$  between weighted parabolic reductions of  $\mathcal{H}$  and weighted parabolic reductions of  $\theta_* \mathcal{H}$ .

Having set up the notation for weighted parabolic reductions, we proceed to explain how this notion relates to  $G$ -bundles on  $U \times_K \Theta_K$  following [Hei17]. Let  $U$  be a  $K$ -scheme. Suppose that  $\mathcal{G}$  is a  $G$ -bundle on  $U$ . We can equivalently view this as a  $G_K$ -bundle on  $U$ . Let  $(\lambda, \mathcal{G}_\lambda)$  be a weighted parabolic reduction of  $\mathcal{G}$ . The morphism  $\lambda : \mathbb{G}_m \rightarrow G_K$  can be used to define a homomorphism  $\text{conj}_\lambda : P_\lambda \times \mathbb{G}_m \rightarrow P_\lambda \times \mathbb{G}_m$  of group schemes over  $\mathbb{G}_m$  defined by  $(p, x) \mapsto (\lambda(x)p\lambda(x)^{-1}, x)$ . This extends to a homomorphism  $\text{gr}_\lambda : P_\lambda \times_K \mathbb{A}_K^1 \rightarrow P_\lambda \times_K \mathbb{A}_K^1$ . Let  $t \in \mathbb{A}_K^1$  be a closed point. The restriction of  $\text{gr}_\lambda$  to a slice  $P_\lambda \times \{t\}$  is the identity when  $t = 1$ , it is an isomorphism when  $t \neq 0$ , and it is the projection of the parabolic subgroup  $P_\lambda$  to its Levi  $L_\lambda$  when  $t = 0$ . Now we can define a  $P_\lambda$ -bundle on  $U \times_K \Theta_K$  as follows. Note that  $\mathcal{G}_\lambda \times_K \mathbb{A}_K^1$  is a  $P_\lambda \times_K \mathbb{A}_K^1$ -torsor on  $X \times \mathbb{A}_K^1$ . Using the homomorphism  $\text{gr}_\lambda$  we obtain, by extension of structure group, a new  $P_\lambda \times_K \mathbb{A}_K^1$ -torsor which is denoted  $(\mathcal{G}_\lambda \times_K \mathbb{A}_K^1) \times_{\mathbb{A}_K^1}^{\text{gr}_\lambda} (P_\lambda \times_K \mathbb{A}_K^1)$ . This bundle is  $\mathbb{G}_m$ -equivariant because the homomorphism  $\text{gr}_\lambda$  is  $\mathbb{G}_m$ -equivariant (with the adjoint action on  $P_\lambda$  and multiplication on  $\mathbb{A}_K^1$ ), and therefore we can take the quotient by  $\mathbb{G}_m$

$$\text{Rees}(\mathcal{G}_\lambda, \lambda) := \left[ \left( (\mathcal{G}_\lambda \times_K \mathbb{A}_K^1) \times_{\mathbb{A}_K^1}^{\text{gr}_\lambda} (P_\lambda \times_K \mathbb{A}_K^1) \right) / \mathbb{G}_m \right]$$

By construction, the restriction  $\text{Rees}(\mathcal{G}_\lambda, \lambda)|_{U \times 1}$  is canonically isomorphic to the  $P_\lambda$ -bundle  $\mathcal{G}_\lambda$ , and the restriction to  $U \times 0$  is a  $P_\lambda$ -bundle which admits a reduction of structure group to  $L_\lambda$ .

We can extend the structure group from  $P_\lambda$  to  $G_K$  in order to obtain a  $G_K$ -bundle  $\text{Rees}(\mathcal{G}_\lambda, \lambda) \times^{P_\lambda} G_K$ . By construction, the restriction  $\text{Rees}(\mathcal{G}_\lambda, \lambda) \times^{P_\lambda} G_K|_{U \times_K 1}$  is canonically isomorphic to the original  $G$ -bundle  $\mathcal{G}$ .

If  $G_K$  is split, then it turns out that all  $\Theta$ -filtrations of  $G$ -bundles on  $U$  arise this way.

**Proposition 3.14.** Suppose that  $G_K$  is split. Let  $U$  be a connected scheme of finite type over  $K$ . Let  $\mathcal{G}$  be a  $G$ -bundle on  $U$ . Then every  $G$ -bundle  $\mathcal{E}$  on  $U \times_K \Theta_K$  such that  $\mathcal{E}|_{U \times_K 1} \cong \mathcal{P}$  is of the form  $\text{Rees}(\mathcal{G}_\lambda, \lambda) \times^{P_\lambda} G_K$ , where

- (a)  $\lambda : \mathbb{G}_m \rightarrow G_K$  is a cocharacter that is uniquely determined up to conjugation.
- (b)  $\mathcal{G}_\lambda$  is a  $P_\lambda$ -reduction of  $\mathcal{G}$ .

In other words, all  $\Theta$ -filtrations of  $\mathcal{G}$  arise from weighted parabolic reductions of  $\mathcal{G}$ .

*Proof.* This is [Hei17, Lem. 1.13]. The lemma found there is stated for a connected smooth projective curve over  $K$ , but the same argument applies to any connected scheme of finite type over  $K$ .  $\square$

Suppose that  $G_K$  is split. Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf over  $X_K$ . Let  $U$  be a big open subset of  $X_K$  where  $\mathcal{F}$  is locally-free. Let  $\mathcal{G}$  denote the  $G_K$ -bundle on  $U$  corresponding to the

$\rho$ -sheaf  $(\mathcal{F}, \sigma)$ . By Proposition 3.14, all  $\Theta$ -filtrations of  $\mathcal{G}$  arise from the Rees construction applied to a weighted parabolic reduction  $(\lambda, \mathcal{G}_\lambda)$  of  $\mathcal{G}$ . We have an associated weighted parabolic reduction  $(\rho_*(\lambda), \rho_*(\mathcal{G}_\lambda))$  of the  $\mathrm{GL}(V)$ -bundle  $\mathcal{F}|_U$ . The Rees construction applied to this weighted parabolic reduction yields a filtration  $(\mathcal{F}'_m)_{m \in \mathbb{Z}}$  of  $\mathcal{F}|_U$ . This induces a filtration  $(\mathcal{F})_{m \in \mathbb{Z}}$  of  $\mathcal{F}$  as follows. Let  $\widetilde{\mathcal{F}}'_m$  denote the pushforward of  $\mathcal{F}'_m$  under the open inclusion  $U \hookrightarrow X_K$ . Note that  $\widetilde{\mathcal{F}}'_m$  is a subsheaf of the reflexive hull  $\mathcal{F}^{\vee\vee}$ . Then, we set  $\mathcal{F}_m := \widetilde{\mathcal{F}}'_m \cap \mathcal{F}^{\vee\vee}$ . It follows by construction that  $(\mathcal{F})_{m \in \mathbb{Z}}$  satisfies the properties in Proposition 3.8, so it is indeed a filtration of  $\mathcal{F}$ .

The pair  $((\mathcal{F}_m)_{m \in \mathbb{Z}}, \mathrm{Rees}(\mathcal{G}_\lambda, \lambda) \times^{P_\lambda} G)$  is an element in  $\mathrm{Filt}(\mathcal{F}, \sigma)$ . Conversely, it follows from our discussion above that every element of  $\mathrm{Filt}(\mathcal{F}, \sigma)$  arises this way. This is our sought after describing the filtrations of the  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$ . We state this formally with a definition and a proposition.

**Definition 3.15.** *Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . A weighted parabolic reduction of  $(\mathcal{F}, \sigma)$  consists of a pair  $(\lambda, \mathcal{G}_\lambda)$ , where*

- (a) *Implicit in the notation we have the choice of (any) big open subset  $U \subset X_K$  such that  $\mathcal{F}|_U$  is locally-free. We denote by  $\mathcal{G}$  the  $G_K$ -bundle on  $U$  corresponding to  $(\mathcal{F}, \sigma)$ .*
- (b)  *$(\lambda, \mathcal{G}_\lambda)$  is a weighted parabolic reduction of  $\mathcal{G}$ .*

*The filtration  $\mathcal{F}_m := \widetilde{\mathcal{F}}'_m \cap \mathcal{F}^{\vee\vee}$  of  $\mathcal{F}$  described in the discussion above is called the underlying filtration associated to the parabolic reduction  $(\lambda, \mathcal{G}_\lambda)$ . We say that  $(\lambda, \mathcal{G}_\lambda)$  is nondegenerate if  $\lambda$  is not the trivial cocharacter.*

Now we record the description of filtrations of  $\rho$ -sheaves obtained from the discussion above.

**Proposition 3.16.** *Suppose that  $G_K$  is split. Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . Then every filtration in  $\mathrm{Filt}(\mathcal{F}, \sigma)$  is of the form  $((\mathcal{F}_m)_{m \in \mathbb{Z}}, \mathrm{Rees}(\mathcal{G}_\lambda, \lambda) \times^{P_\lambda} G)$ , where  $(\lambda, \mathcal{G}_\lambda)$  is a weighted parabolic reduction of  $(\mathcal{F}, \sigma)$  with underlying associated filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$ .*

*Moreover the filtration  $((\mathcal{F}_m)_{m \in \mathbb{Z}}, \mathrm{Rees}(\mathcal{G}_\lambda, \lambda) \times^{P_\lambda} G)$  is nondegenerate if and only if  $(\lambda, \mathcal{G}_\lambda)$  is nondegenerate.  $\square$*

We can also give a concrete description of graded points of the stack  $\mathrm{Bun}_\rho(X)$ . A graded  $\rho$ -sheaf over  $X_K$  is defined to be a morphism  $[\mathrm{Spec}(K)/\mathbb{G}_m] \rightarrow \mathrm{Bun}_\rho(X)$ . By unraveling the definitions, this notion admits the following concrete description. A graded  $\rho$ -sheaf on  $X_K$  consists of a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  on  $X_K$ , plus a  $\mathbb{Z}$ -grading  $\mathcal{F} = \bigoplus_{m \in \mathbb{Z}} \overline{\mathcal{F}}_m$  of the torsion-free sheaf  $\mathcal{F}$  such that each summand  $\overline{\mathcal{F}}_m$  is also torsion-free. We get a well-defined  $\mathbb{G}_m$ -action on the scheme of  $G$ -reductions  $\mathrm{Red}_G(\mathcal{F})$  and we impose the condition that the section  $\sigma$  is  $\mathbb{G}_m$ -invariant. Hence we can think of morphisms  $g : [\mathrm{Spec}(K)/\mathbb{G}_m] \rightarrow \mathrm{Bun}_\rho(X)$  concretely as a graded pair  $(\bigoplus_{m \in \mathbb{Z}} \overline{\mathcal{F}}_m, \sigma)$ .

We end this section by explaining a concrete description of the associated graded of a filtration. Suppose that  $G_K$  is split. Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . Let  $f : \Theta_K \rightarrow \mathrm{Bun}_\rho(X)$  be a filtration of  $(\mathcal{F}, \sigma)$ . By Proposition 3.16  $f$  comes from a weighted parabolic reduction  $(\lambda, \mathcal{G}_\lambda)$  with underlying associated filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$ . We can restrict  $f$  to the point  $0 \in \Theta_K$  in order to obtain a graded  $\rho$ -sheaf  $f|_{[0/\mathbb{G}_m]} : [\mathrm{Spec}(K)/\mathbb{G}_m] \rightarrow \mathrm{Bun}_\rho(X)$ . The underlying graded torsion-free sheaf of  $f|_{[0/\mathbb{G}_m]}$  is the associated graded  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m / \mathcal{F}_{m+1}$ .

We can describe the  $G$ -reduction over  $U$  of the  $\rho$ -sheaf  $f|_{[0/\mathbb{G}_m]}$  as follows. Let  $P_\lambda = L_\lambda \times U_\lambda$  be the Levi decomposition defined by  $\lambda$ . Then over the big open subset  $U$ , the corresponding  $G$ -bundle is the one obtained by applying extension of structure group under the composition  $P_\lambda = L_\lambda \times U_\lambda \twoheadrightarrow L_\lambda \hookrightarrow G_K$  to the  $P_\lambda$ -bundle  $\mathcal{G}_\lambda$ .

#### 4. $\Theta$ -STRATIFICATION ON $\text{Bun}_\rho(X)$

In this section we define a polynomial numerical invariant  $\nu$  on the stack  $\text{Bun}_\rho(X)$ . This invariant  $\nu$  can be described in terms of reduced Hilbert polynomials. We prove that  $\nu$  induces a  $\Theta$ -stratification, as in [HL14, Defn. 2.1.2] [HLHJ21, Defn. 2.23].

**4.1. The numerical invariant.** We define a family of line bundles  $L_n$  on  $\text{Bun}_\rho(X)$  following [HLHJ21, §2].

**Definition 4.1.** *Let  $n \in \mathbb{Z}$ . We define  $M_n$  to be the line bundle on  $\text{Bun}_\rho(X)$  determined as follows. Let  $T \in \text{Aff}_k$  and let  $f : T \rightarrow \text{Bun}_\rho(X)$  be a morphism, represented by a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  on  $X_T$ . Let  $\pi_T : X_T \rightarrow T$  denote the structure morphism. Then, we set*

$$f^* M_n := \det (R\pi_{T*}(\mathcal{F}(n)))$$

Moreover, we define  $b_d := \bigotimes_{i=0}^d M_i^{(-1)^i \binom{d}{i}}$ . We set  $L_n$  to be the rational line bundle given by

$$f^* L_n := f^* M_n \otimes (f^* b_d)^{-\otimes \bar{p}_{\mathcal{F}}(n)}$$

**Remark 4.2.**  *$L_n$  is an element of the rational Picard group  $\text{Pic}(\text{Bun}_\rho(X)) \otimes \mathbb{Q}$  of the stack. This is what we mean by rational line bundle. We will henceforth omit the adjective “rational” and refer to  $L_n$  as a line bundle. This should not cause confusion, as all the arguments will apply to the rational Picard group in the same way as for the usual Picard group.*

$L_n$  is the restriction under the forgetful map  $\text{Bun}_\rho(X) \rightarrow \text{Coh}^{tf}(X)$  of the line bundle  $L_n$  on  $\text{Coh}^{tf}(X)$  defined in [HLHJ21, §2]. See the discussion therein for more details on this family of line bundles.

Next, we define a rational quadratic norm on graded  $\rho$ -sheaves (see [HLHJ21, §2.5, §4.1] for more details on rational quadratic norms).

**Definition 4.3.** *We define the rational quadratic norm  $b$  on graded points of  $\text{Bun}_\rho(X)$  as follows. Let  $g : [\text{Spec}(K)/\mathbb{G}_m] \rightarrow \text{Bun}_\rho(X)$  be represented by a graded  $\rho$ -sheaf  $(\bigoplus_{m \in \mathbb{Z}} \bar{\mathcal{F}}_m, \sigma)$ . Then we set*

$$b(g) := \sum_{m \in \mathbb{Z}} m^2 \cdot \text{rk}(\bar{\mathcal{F}}_m)$$

We define the polynomial numerical invariant  $\nu$  to be the one determined by the family of line bundles  $L_n$  and the rational quadratic norm  $b$ , as it was done for the case of pure sheaves in [HLHJ21, §4.1]. For our purposes, it is sufficient to know the value of the numerical invariant associated to a nondegenerate  $\Theta$ -filtration. We proceed to describe this next.

Let  $K \supset k$  be a field extension, and let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . Let  $f : \Theta_K \rightarrow \text{Bun}_\rho(X)$  be a nondegenerate filtration of  $(\mathcal{F}, \sigma)$ . By Proposition 3.9 this amounts to a filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  of the torsion-free sheaf  $\mathcal{F}$  satisfying some extra compatibility conditions. The restriction  $f|_{[0/\mathbb{G}_m]} \rightarrow \text{Bun}_\rho(X)$  yields a graded  $\rho$ -sheaf on  $X_K$ , whose underlying torsion-free sheaf is given by the associated graded  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m/\mathcal{F}_{m+1}$ . Then we define  $\nu(f)$  to be the polynomial in  $\mathbb{R}[n]$  defined by

$$(2) \quad \nu(f) := \frac{\text{wt}(L_n|_0)}{\sqrt{b(f|_{[0/\mathbb{G}_m]})}}$$

The same computation as in [HLHJ21, §4.1] shows that this polynomial numerical invariant  $\nu$  admits a description in terms of reduced Hilbert polynomials. Denote by  $P_{\mathcal{O}_X}$  the Hilbert polynomial of the structure sheaf of  $X$  with respect to  $\mathcal{O}(1)$ , which is a polynomial of degree  $d$ . Let  $\frac{A_d}{d!}$  be the leading coefficient of  $P_{\mathcal{O}_X}$ .

**Proposition 4.4.** *Let  $K \supset k$  be a field extension. Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . Choose a nondegenerate filtration  $f : \Theta_K \rightarrow \text{Bun}_\rho(X)$  of  $(\mathcal{F}, \sigma)$ . Suppose that  $f$  has underlying associated filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  of  $\mathcal{F}$ . Then, we have*

$$(3) \quad \nu(f) = \sqrt{A_d} \cdot \frac{\sum_{m \in \mathbb{Z}} m \cdot (\bar{p}_{\mathcal{F}_m/\mathcal{F}_{m+1}} - \bar{p}_{\mathcal{F}}) \cdot \text{rk}(\mathcal{F}_m/\mathcal{F}_{m+1})}{\sqrt{\sum_{m \in \mathbb{Z}} m^2 \cdot \text{rk}(\mathcal{F}_m/\mathcal{F}_{m+1})}}$$

*Proof.* The value of the denominator  $\sqrt{b(f|_{[0/\mathbb{G}_m]})}$  follows by definition. The numerator has been computed in [HLHJ21, §4.1]. The extra factor  $\sqrt{A_d}$  appears because we define the rank to be the dimension of the fiber of  $\mathcal{F}$  at the generic point, as opposed to the rank defined in [HLHJ21, §2.3] using the leading term of the Hilbert polynomial.  $\square$

Given a weighted parabolic reduction  $f = (\lambda, \mathcal{G}_\lambda)$  of a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$ , we denote by  $\nu(f)$  the value of the numerical invariant for the  $\Theta$ -filtration obtained by applying the Rees construction to  $f$ . If  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  is the underlying associated filtration of  $(\lambda, \mathcal{G}_\lambda)$ , then the same formula (3) applies.

**4.2. Affine Grassmannians for  $\rho$ -sheaves.** The setup for this subsection is as follows. Let  $S$  be a quasicompact  $k$ -scheme. Let  $\mathcal{F}$  be a family of torsion-free sheaves of rank  $r$  on  $X_S$ . Suppose that we are given an effective Cartier divisor  $D \hookrightarrow X_S$  that is flat over  $S$ . Set  $Q = X_S \setminus D$  (note that this is not a big open subset). The paper [HLHJ21] defines a notion of affine Grassmannian  $\text{Gr}_{X_S, D, \mathcal{F}}$ . We recall the definition and some properties proven in [HLHJ21, §3].

**Definition 4.5.**  $\text{Gr}_{X_S, D, \mathcal{F}}$  is the functor from  $\text{Aff}_S^{\text{op}}$  to sets given as follows. For every affine scheme  $T$  in  $\text{Aff}_S$ , the set  $\text{Gr}_{X_S, D, \mathcal{F}}(T)$  consists of equivalence classes of pairs  $(\mathcal{E}, \theta)$ , where

- (1)  $\mathcal{E}$  is a family of torsion-free sheaves on  $X_T$ .
- (2)  $\theta$  is morphism  $\theta : \mathcal{F}|_{X_T} \rightarrow \mathcal{E}$  such that the restriction  $\theta|_{Q_T}$  is an isomorphism.

Two such pairs  $(\mathcal{E}_1, \theta_1)$  and  $(\mathcal{E}_2, \theta_2)$  are considered equivalent if there is an isomorphism  $\mathcal{E}_1 \xrightarrow{\sim} \mathcal{E}_2$  that identifies the morphisms  $\theta_1$  and  $\theta_2$ .

Let  $j$  be the open immersion  $j : Q \hookrightarrow X_S$ . Choose  $T \in \text{Aff}_S$  and  $(\mathcal{E}, \psi) \in \text{Gr}_{X_S, D, \mathcal{F}}(T)$ . The unit  $\mathcal{E} \rightarrow j_{T*} j_T^* \mathcal{E}$  is a monomorphism. Hence we can use the isomorphism  $j_{T*} (j_T^*)^* \theta$  in order to view  $\mathcal{E}$  as a subsheaf of  $j_{T*} j_T^* \mathcal{F}|_{X_T}$ . In particular we see that  $\theta$  is automatically injective, and we have  $\mathcal{F}|_{X_T} \subset \mathcal{E} \subset j_{T*} j_T^* \mathcal{F}|_{X_T}$ . Note that there is an increasing sequence of sheaves

$$\mathcal{F}|_{X_T} \subset \mathcal{F}|_{X_T}(D_T) \subset \mathcal{F}|_{X_T}(2D_T) \subset \cdots \subset \mathcal{F}|_{X_T}(nD_T) \subset \cdots$$

We can use this to define a truncated version of  $\text{Gr}_{X, D, \mathcal{E}}$ .

**Definition 4.6.** Let  $N$  be a positive integer.  $\text{Gr}_{X, D, \mathcal{F}}^{\leq N}$  is the subfunctor of  $\text{Gr}_{X_S, D, \mathcal{F}}^{\leq N}$  defined as follows. For all  $T$  in  $\text{Aff}_T$ , we set

$$\text{Gr}_{X, D, \mathcal{F}}^{\leq N}(T) := \left\{ \begin{array}{l} \text{pairs } (\mathcal{E}, \theta) \text{ in } \text{Gr}_{X_S, D, \mathcal{F}}(T) \text{ such that} \\ \mathcal{E} \subset \mathcal{F}|_{X_T}(ND_T) \end{array} \right\}$$

For any rational polynomial  $P \in \mathbb{Q}[n]$ , we denote by  $\text{Gr}_{X_S, D, \mathcal{F}}^{\leq N, P}$  the open and closed subfunctor of  $\text{Gr}_{X_S, D, \mathcal{F}}^{\leq N}$  consisting of points  $(\mathcal{E}, \theta)$  where  $\mathcal{E}$  has Hilbert polynomial  $P$  on every fiber.

**Proposition 4.7** ([HLHJ21, Prop. 3.13]). *We have  $\text{Gr}_{X_S, D, \mathcal{F}} = \varinjlim_{N > 0} \text{Gr}_{X_S, D, \mathcal{F}}^{\leq N}$  as presheaves on  $\text{Aff}_S$ . For each  $N \geq 0$ ,*

$$\text{Gr}_{X_S, D, \mathcal{F}}^{\leq N} = \bigsqcup_{P \in \mathbb{Q}[n]} \text{Gr}_{X_S, D, \mathcal{F}}^{\leq N, P}$$

Each  $\mathrm{Gr}_{X_S, D, \mathcal{F}}^{\leq N, P}$  is represented by a projective scheme of finite presentation over  $S$ . This induces a presentation of  $\mathrm{Gr}_{X_S, D, \mathcal{F}}$  as an ind-projective ind-scheme over  $S$ .

Suppose now that we have a section  $\sigma : Q \rightarrow \mathrm{Red}_G(\mathcal{F})$  defined over  $Q$ . We use this datum to define a version of the affine Grassmannian for  $\rho$ -sheaves.

**Definition 4.8.** *The affine Grassmannian  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}$  is the functor from  $(\mathrm{Aff}_S)^{op}$  into sets defined as follows. For each affine scheme  $T$  in  $\mathrm{Aff}_S$ ,  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}(T)$  is the set of equivalence classes of the following data*

- (1) A  $\rho$ -sheaf  $(\mathcal{E}, \zeta)$  on  $X_T$ .
- (2) A morphism  $\theta : \mathcal{F}|_{X_T} \rightarrow \mathcal{E}$  such that the restriction  $\theta|_{Q_T}$  is an isomorphism. Moreover, we require that the following diagram induced by the isomorphism  $\theta|_{Q_T}$  commutes.

$$\begin{array}{ccc} Q_T & \xrightarrow{\zeta|_{Q_T}} & \mathrm{Red}_G(\mathcal{E}|_{Q_T}) \\ & \searrow \sigma|_{Q_T} & \uparrow \wr \\ & & \mathrm{Red}_G(\mathcal{F}|_{Q_T}) \end{array}$$

Two elements above are considered equivalent if there is an isomorphism of  $\rho$ -sheaves that is compatible with the morphism  $\theta$ .

There is a forgetful morphism  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma} \rightarrow \mathrm{Gr}_{X_S, D, \mathcal{F}}$  defined by  $(\mathcal{E}, \zeta, \theta) \mapsto (\mathcal{E}, \theta)$ .

**Proposition 4.9.** *The forgetful morphism witnesses  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}$  as a closed sub ind-scheme of  $\mathrm{Gr}_{X_S, D, \mathcal{F}}$ .*

Before proving Proposition 4.9, we first state a couple of lemmas.

**Lemma 4.10.** *Let  $Y \rightarrow S$  be a proper morphism of finite presentation. Let  $\mathcal{G}$  and  $\mathcal{F}$  be  $\mathcal{O}_Y$ -modules. Suppose that  $\mathcal{F}$  is finitely presented and  $S$ -flat. Fix a morphism  $\sigma : \mathcal{G} \rightarrow \mathcal{F}$ . Let  $Z_\sigma^{\mathcal{G}, \mathcal{F}}$  be the functor from  $k$ -schemes to sets that sends a  $k$ -scheme  $T$  to*

$$Z_\sigma(T) := \left\{ \begin{array}{l} \text{morphisms } g : T \rightarrow S \text{ such that} \\ (id_Y \times_S g)^* \sigma = 0 \end{array} \right\}$$

Then  $Z$  is represented by a closed subscheme of  $S$ .

*Proof.* By [Sta21, Tag 08K6], there is a separated  $S$ -scheme  $\mathrm{Hom}(\mathcal{G}, \mathcal{F})$  parametrizing morphisms  $\mathcal{G} \rightarrow \mathcal{F}$ . The two morphisms  $\sigma$  and  $0$  yield two sections  $f_\sigma, f_0 : S \rightarrow \mathrm{Hom}(\mathcal{O}_Y, \mathcal{F})$ . The functor  $Z_\sigma$  is represented by the locus where these two morphisms  $f_\sigma, f_0$  agree. This is a closed subscheme of  $S$ , because  $\mathrm{Hom}(\mathcal{G}, \mathcal{F}) \rightarrow S$  is separated.  $\square$

Let  $S$  be a  $k$ -scheme. Suppose that  $p : Y \rightarrow X_S$  is an affine morphism of finite presentation. Let  $s : Q \rightarrow Y$  be a section of  $p$  defined over the open complement  $Q = X_S \setminus D$ .

**Definition 4.11.** *We define  $\mathrm{Sect}_s^Y$  to be the functor from  $(\mathrm{Aff}_S)^{op}$  to sets given as follows. For  $T \in \mathrm{Aff}_S$ , let  $s_T : Q_T \rightarrow Y_T$  and  $p_T : Y_T \rightarrow X_T$  denote the base-changes. We define*

$$\mathrm{Sect}_s^Y(T) := \left\{ \begin{array}{l} \text{sections } f : X_T \rightarrow Y_T \text{ of } p_T \\ \text{such that } f|_{Q_T} = s_T \end{array} \right\}$$

**Lemma 4.12.** *The functor  $\mathrm{Sect}_s^Y$  is represented by a closed subscheme of  $S$ .*

*Proof.* After passing to an affine open cover, we can assume that  $S = \mathrm{Spec}(R)$ . Since everything is of finite presentation, we can use standard approximation results to reduce to the case when  $R$  is Noetherian. So  $X_S$  is Noetherian, and  $Y = \underline{\mathrm{Spec}}_{X_S}(\mathcal{A})$  for some finitely generated  $\mathcal{O}_{X_S}$ -algebra  $\mathcal{A}$ .

We can choose a finite open cover  $\{V_i\}$  of  $X_S$  and a finite set of sections on each  $V_i$  that generate  $\mathcal{A}|_{V_i}$  as a  $\mathcal{O}_{V_i}$ -algebra. By [Sta21, Tag 01PF] there exists a coherent subsheaf  $\mathcal{F} \subset \mathcal{A}$  that contains all of these sections. Serre vanishing implies that exists some positive integers  $N, m$  and a surjective map of sheaves  $\mathcal{O}_{X_S}^{\oplus m} \rightarrow \mathcal{F}(N)$ . Twisting by  $\mathcal{O}_X(-N)$ , we get a surjection  $\mathcal{O}_{X_S}(-N)^{\oplus m} \rightarrow \mathcal{F}$ . This induces a surjective map of  $\mathcal{O}_{X_S}$ -algebras  $\text{Sym}(\mathcal{O}_{X_S}(-N)^{\oplus m}) \rightarrow \mathcal{A}$ .

If we set  $\mathcal{E} := \mathcal{O}_{X_S}(-N)^{\oplus m}$ , then we have a closed immersion  $\iota : Y \hookrightarrow \underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})$ . Let us denote by  $s' : Q \rightarrow \underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})$  the composition of  $s : Q \rightarrow Y$  with the closed immersion  $\iota$ . There is a natural map of functors  $\text{Sect}_s^Y \rightarrow \text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})}$  given by composition with  $\iota$ .

The lemma is implied by the following two claims:

- (i) The functor  $\text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})}$  is represented by a closed subscheme of  $S$ .
- (ii) The map  $\text{Sect}_s^Y \rightarrow \text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})}$  is a closed immersion.

For claim (i), note that  $s'$  is just an element of  $\Gamma(Q, \mathcal{E})$ . Since  $j_* j^* \mathcal{E} = \varinjlim_{m \in \mathbb{Z}} \mathcal{E}(mD)$ , there exists a positive integer  $M$  such that  $s'$  extends to a section  $\sigma \in \Gamma(X_S, \mathcal{E}(MD))$ . Consider the composition  $\bar{\sigma} : \mathcal{O}_{X_S} \xrightarrow{\sigma} \mathcal{E}(MD) \rightarrow \mathcal{E}(MD) / \mathcal{E}$ . The functor  $\text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})}$  can be seen to be isomorphic to the functor  $Z_{\bar{\sigma}}$ , as in Lemma 4.10. The coherent sheaf  $\mathcal{E}(MD) / \mathcal{E}$  is  $S$ -flat by the slicing criterion for flatness [Sta21, Tag 046Y]. Hence Lemma 4.10 implies that  $Z_{\bar{\sigma}}$  is represented by a closed subscheme of  $S$ .

We now proceed to prove claim (ii). Let  $T$  be a  $S$ -scheme. For our purposes we can assume that  $T$  is affine and Noetherian. Choose  $T \rightarrow \text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})}$  represented by a section  $f$  of the structure morphism  $\underline{\text{Spec}}_{X_T}(\text{Sym } \mathcal{E}_T) \rightarrow X_T$ . We want to show that the fiber product  $\text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})} \times_{\text{Sect}_s^Y} T$  is represented by a closed subscheme of  $T$ . For any scheme  $W$ , we have

$$\text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})} \times_{\text{Sect}_s^Y} T(W) = \left\{ \begin{array}{l} \text{morphisms } h : W \rightarrow T \text{ such that} \\ (\text{id}_{X_T} \times h)^* f : X_W \rightarrow \underline{\text{Spec}}_{X_W}(\text{Sym } \mathcal{E}_W) \\ \text{factors through } Y_W \end{array} \right\}$$

Define  $B$  to be the fiber product

$$\begin{array}{ccc} B & \hookrightarrow & X_T \\ \downarrow & & \downarrow f \\ Y_T & \xrightarrow{\iota \times \text{id}_T} & \underline{\text{Spec}}_{X_T}(\text{Sym } \mathcal{E}_T) \end{array}$$

The morphism  $m : B \hookrightarrow X_T$  is a closed immersion. For any scheme  $W$ , the  $W$ -points of the fiber product can be alternatively described by

$$\text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})} \times_{\text{Sect}_s^Y} T(W) = \left\{ \begin{array}{l} \text{morphisms } h : W \rightarrow T \text{ such that} \\ (\text{id}_{X_T} \times h)^* m : B_W \hookrightarrow X_W \text{ is an isomorphism} \end{array} \right\}$$

Let us denote by  $\varphi : \mathcal{J} \hookrightarrow \mathcal{O}_{X_T}$  the inclusion of the  $\mathcal{O}_{X_T}$ -sheaf of ideals  $\mathcal{J}$  corresponding to the closed subscheme  $B \hookrightarrow X_T$ . The fiber product  $\text{Sect}_{s'}^{\underline{\text{Spec}}_{X_S}(\text{Sym } \mathcal{E})} \times_{\text{Sect}_s^Y} T$  is isomorphic to  $Z_{\varphi}$ . Hence it is represented by a closed subscheme of  $T$  by Lemma 4.10.  $\square$

*Proof of Proposition 4.9.* Let  $T \in \text{Aff}_S$  and let  $T \rightarrow \text{Gr}_{X_S, D, \mathcal{F}}$  be a morphism represented by a pair  $(\mathcal{E}, \theta)$ . By Definition 4.8 the restriction  $\theta|_{Q_T}$  over the open subset  $Q_T$  is an

isomorphism. This yields an identification  $\mathrm{Red}_G(\mathcal{F})|_{Q_T} \xrightarrow{\sim} \mathrm{Red}_G(\mathcal{E})|_{Q_T}$ . We have a section  $s$  induced by  $\sigma$  as follows

$$s : Q_T \xrightarrow{\sigma|_{Q_T}} \mathrm{Red}_G(\mathcal{F})|_{Q_T} \xrightarrow{\sim} \mathrm{Red}_G(\mathcal{E})|_{Q_T}$$

After unraveling the definitions, it can be seen that the fiber product  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma} \times_{\mathrm{Gr}_{X_S, D, \mathcal{F}}} T$  is naturally isomorphic to  $\mathrm{Sect}_s^{\mathrm{Red}_G(\mathcal{E})}$ . By Lemma 4.12, this is represented by a closed subscheme of  $T$ .  $\square$

Let  $P \in \mathbb{Q}[n]$  be a rational polynomial. For each  $N \geq 0$  we define the subfunctors

$$\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}^{\leq N} := \mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma} \cap \mathrm{Gr}_{X_S, D, \mathcal{F}}^{\leq N}$$

$$\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}^{\leq N, P} := \mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma} \cap \mathrm{Gr}_{X_S, D, \mathcal{F}}^{\leq N, P}$$

Both of these are closed sub ind-schemes of  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}$ . Propositions 4.7 and 4.9 imply that  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma} = \varinjlim_{N > 0} \mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}^{\leq N}$ . Furthermore each  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}^{\leq N, P}$  is represented by a scheme that is projective and of finite presentation over  $S$ .

There is a natural forgetful morphism  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma} \rightarrow \mathrm{Bun}_\rho(X)$  that takes a point  $(\mathcal{E}, \zeta, \theta)$  in  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}$  and forgets the isomorphism  $\theta$ . We can use this map to pullback the line bundle  $L_n$  to  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}$ . We will abuse notation and denote this restriction by  $L_n$  as well. The following corollary shows that the family  $L_n^\vee$  of dual line bundles on  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}$  is “asymptotically ample”

**Corollary 4.13.** *Let  $N \in \mathbb{Z}_{\geq 0}$  and  $P \in \mathbb{Q}[n]$ . There exists some integer  $m \gg 0$  such that for all  $n \geq m$ , the restriction  $L_n^\vee|_{\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}^{\leq N, P}}$  is  $S$ -ample.*

*Proof.* This follows from Proposition 4.9 and [HLHJ21, Prop. 3.19], which states that  $L_n^\vee$  is eventually ample on  $\mathrm{Gr}_{X_S, D, \mathcal{F}, \sigma}^{\leq N, P}$ .  $\square$

**4.3. Monotonicity of the numerical invariant.** In this subsection we prove that the invariant  $\nu$  on  $\mathrm{Bun}_\rho(X)$  is strictly  $\Theta$ -monotone and strictly  $S$ -monotone. We recall what these notions mean, see also [HL14, §5] or [HLHJ21, §2.5] for more details.

We need some setup first. Let  $R$  be a complete discrete valuation ring over  $k$ . Choose a uniformizer  $\pi$  of  $R$ . We define  $Y_{\Theta_R} := \mathbb{A}_R^1$  equipped with the  $\mathbb{G}_m$ -action that gives  $t$  weight  $-1$ . We have  $\Theta_R = [\mathbb{A}_R^1 / \mathbb{G}_m]$ . Note that  $\mathbb{A}_R^1$  contains a unique  $\mathbb{G}_m$ -invariant closed point cut out by the ideal  $(t, \pi)$ . We will denote this point by  $(0, 0)$ .

Define  $Y_{\overline{ST}_R} := \mathrm{Spec}(R[t, s]/(st - \pi))$ . We equip  $Y_{\overline{ST}_R}$  with the  $\mathbb{G}_m$ -action that gives  $s$  weight 1 and  $t$  weight  $-1$ . The isomorphism class of the  $\mathbb{G}_m$ -scheme  $Y_{\overline{ST}_R}$  is independent of the choice of uniformizer  $\pi$ . Define  $\overline{ST}_R := [Y_{\overline{ST}_R} / \mathbb{G}_m]$ . We will denote by  $(0, 0)$  the unique  $\mathbb{G}_m$ -fixed closed point in  $Y_{\overline{ST}_R}$ . We have that  $(0, 0)$  is cut out by the ideal  $(s, t)$ .

For any field  $\kappa$  and integer  $a \geq 1$ , we denote by  $\mathbb{P}_\kappa^1[a]$  the  $\mathbb{G}_m$ -scheme  $\mathbb{P}_\kappa^1$  equipped with the  $\mathbb{G}_m$ -action determined by the equation  $t \cdot [x : y] = [t^{-a}x : y]$ . We set  $0 = [0 : 1]$  and  $\infty = [1 : 0]$ .

With all this notation in place, we are ready to define the monotonicity properties that we will prove.

**Definition 4.14.** *A polynomial numerical invariant  $\nu$  on a stack  $\mathcal{M}$  is strictly  $\Theta$ -monotone (resp. strictly  $S$ -monotone) if the following condition holds.*

*Let  $R$  be any complete discrete valuation ring and set  $\mathfrak{X}$  to be  $\Theta_R$  (resp.  $\overline{ST}_R$ ). Choose a map  $\varphi : \mathfrak{X} \setminus (0, 0) \rightarrow \mathcal{M}$ . Then, after maybe replacing  $R$  with a finite DVR extension, there exists a reduced and irreducible  $\mathbb{G}_m$ -equivariant scheme  $\Sigma$  with maps  $f : \Sigma \rightarrow Y_{\mathfrak{X}}$  and  $\tilde{\varphi} : [\Sigma / \mathbb{G}_m] \rightarrow \mathcal{M}$  such that*

- (M1) The map  $f$  is proper,  $\mathbb{G}_m$ -equivariant and its restriction over  $Y_{\mathfrak{X}} \setminus (0, 0)$  induces an isomorphism  $f : \Sigma_{Y_{\mathfrak{X}} \setminus (0, 0)} \xrightarrow{\sim} Y_{\mathfrak{X}} \setminus (0, 0)$ .
- (M2) The following diagram commutes

$$\begin{array}{ccc} [(\Sigma_{Y_{\mathfrak{X}} \setminus (0, 0)}) / \mathbb{G}_m] & & \\ f \downarrow & \searrow \tilde{\varphi} & \\ \mathfrak{X} \setminus (0, 0) & \xrightarrow{\varphi} & \mathcal{M} \end{array}$$

- (M3) Let  $\kappa$  be a finite extension of the residue field of  $R$ . For any  $a \geq 1$  and any finite  $\mathbb{G}_m$ -equivariant morphism  $\mathbb{P}_{\kappa}^1[a] \rightarrow \Sigma_{(0, 0)}$ , we have  $\nu(\tilde{\varphi}|_{[\infty/\mathbb{G}_m]}) \geq \nu(\tilde{\varphi}|_{[0/\mathbb{G}_m]})$ .

Before we proceed with the proof of monotonicity, we need to record a rational filling condition for the stack  $\text{Bun}_{\rho}(X)$  in the spirit of [HLHJ21, §4.2]. Let  $Y$  denote either  $Y_{\Theta_R}$  or  $Y_{\overline{ST}_R}$  as described above.

**Proposition 4.15** ([HLHJ21, Lem. 4.7(i)]). *Let  $(\mathcal{E}, \zeta)$  be a  $\mathbb{G}_m$ -equivariant  $\rho$ -sheaf on  $X_{Y \setminus (0, 0)}$ . Then there exists*

- (1) A  $\mathbb{G}_m$ -equivariant family  $\mathcal{F}$  of torsion-free sheaves on  $X_Y$ .
- (2) A relative effective Cartier divisor  $D \hookrightarrow X_Y$  that is stable under the action of  $\mathbb{G}_m$ .
- (3) A  $\mathbb{G}_m$ -equivariant morphism  $\theta : \mathcal{F}|_{X_{Y \setminus (0, 0)}} \rightarrow \mathcal{E}$  such that the restriction  $\theta|_{(X_{Y \setminus (0, 0)}) \setminus (D_{Y \setminus (0, 0)})}$  is an isomorphism.

□

We are now ready to prove the main proposition of this subsection.

**Proposition 4.16.** *The invariant  $\nu$  on the stack  $\text{Bun}_{\rho}(X)$  is both strictly  $\Theta$ -monotone and strictly  $S$ -monotone.*

*Proof.* The argument is a variation of the one in [HLHJ21, §4.3] using our version of the affine Grassmannian. We let  $Y = Y_{\Theta_R}$  or  $Y_{\overline{ST}_R}$ . Suppose that we are given a morphism  $f : [(Y \setminus (0, 0)) / \mathbb{G}_m] \rightarrow \text{Bun}_{\rho}(X)$ . This consists of the data of a  $\rho$ -sheaf  $(\mathcal{E}, \zeta)$  on  $X_{Y \setminus (0, 0)}$  that is  $\mathbb{G}_m$ -equivariant with respect to the action of  $\mathbb{G}_m$  on  $Y \setminus (0, 0)$ . By Proposition 4.15, there is a  $\mathbb{G}_m$ -equivariant torsion-free sheaf  $\mathcal{F}$  on  $X_Y$ , a  $\mathbb{G}_m$ -stable  $Y$ -flat effective Cartier divisor  $D \hookrightarrow X_Y$  and a  $\mathbb{G}_m$ -equivariant monomorphism  $\theta : \mathcal{F}|_{X_{(Y \setminus (0, 0))}} \rightarrow \mathcal{E}$  such that  $\theta$  is an isomorphism away from  $D$ .

Set  $Q := X_Y \setminus D$ . The isomorphism  $\theta|_{Q_{Y \setminus (0, 0)}}$  induces a  $\mathbb{G}_m$ -equivariant identification  $\text{Red}_G(\mathcal{E})|_{Q_{Y \setminus (0, 0)}} \xrightarrow{\sim} \text{Red}_G(\mathcal{F})|_{Q_{Y \setminus (0, 0)}}$ . We have a  $\mathbb{G}_m$ -equivariant section induced by  $\zeta$ :

$$\sigma_{Q_{Y \setminus (0, 0)}} : Q_{Y \setminus (0, 0)} \xrightarrow{\zeta|_{Q_{Y \setminus (0, 0)}}} \text{Red}_G(\mathcal{E})|_{Q_{Y \setminus (0, 0)}} \xrightarrow{\sim} \text{Red}_G(\mathcal{F})|_{Q_{Y \setminus (0, 0)}}$$

The local ring  $\mathcal{O}_{Y, (0, 0)}$  at  $(0, 0)$  has depth 2. For any point  $u$  in the fiber  $Q_{(0, 0)}$ , the local ring  $\mathcal{O}_{Q, u}$  is flat over  $\mathcal{O}_{Y, (0, 0)}$ . It follows that  $\mathcal{O}_{Q, u}$  has depth at least 2 for all  $u \in Q_{(0, 0)}$ . The same type of argument as in Lemma 2.2 (c) shows that the  $\mathbb{G}_m$ -equivariant section  $\sigma_{Q_{Y \setminus (0, 0)}}$  extends uniquely to a  $\mathbb{G}_m$ -equivariant section  $\sigma : Q \rightarrow \text{Red}_G(\mathcal{F})$ . We now have all of the necessary data to form the affine grassmannian  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}$ . Since everything is  $\mathbb{G}_m$ -equivariant, the ind-scheme  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}$  acquires a natural  $\mathbb{G}_m$ -action.

The data of the  $\rho$ -sheaf  $(\mathcal{E}, \zeta)$  and the monomorphism  $\theta$  is by definition a  $\mathbb{G}_m$ -equivariant section  $s : Y \setminus (0, 0) \rightarrow \text{Gr}_{X_Y, D, \mathcal{F}, \sigma}$  of the structure morphism  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma} \rightarrow Y$ . We can summarize everything in the following diagram, where all of the maps are  $\mathbb{G}_m$ -equivariant:



$$\begin{array}{ccc}
& & \text{Gr}_{X_Y, D, \mathcal{F}, \sigma} \xrightarrow{\text{Forget}} \text{Bun}_\rho(X) \\
& \nearrow s & \downarrow \\
Y \setminus (0, 0) & \hookrightarrow & Y
\end{array}$$

By construction, the composition  $\text{Forget} \circ s$  is the original  $\mathbb{G}_m$ -equivariant morphism  $f : Y \setminus (0, 0) \rightarrow \text{Bun}_\rho(X)$ .

It follows from the definition of  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P}$  that the  $\mathbb{G}_m$ -action on  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}$  preserves each stratum  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P}$ . Since  $Y$  is affine and connected, the map  $s : Y \setminus (0, 0) \rightarrow \text{Gr}_{X_Y, D, \mathcal{F}, \sigma}$  factors through  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P}$  for some  $N \gg 0$  and  $P \in \mathbb{Q}[x]$ . Therefore, we have the following commutative diagram with  $\mathbb{G}_m$ -equivariant morphisms.

$$\begin{array}{ccc}
& & \text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P} \xrightarrow{\text{Forget}} \text{Bun}_\rho(X) \\
& \nearrow s & \downarrow \\
Y \setminus (0, 0) & \hookrightarrow & Y
\end{array}$$

Let  $\Sigma \subset \text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P}$  denote the scheme theoretic image of  $Y \setminus (0, 0)$  in  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P}$ . Note that  $\Sigma$  is a reduced  $\mathbb{G}_m$ -scheme with a natural structure morphism to  $Y$ . The map  $\Sigma \rightarrow Y$  is projective, because  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P}$  is projective over  $Y$  and  $\Sigma$  is a closed subscheme of  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P}$ . By construction the morphism  $\Sigma \rightarrow Y$  is  $\mathbb{G}_m$ -equivariant and restricts to an isomorphism over the open  $Y \setminus (0, 0)$ .

The composition  $\Sigma \rightarrow \text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P} \xrightarrow{\text{Forget}} \text{Bun}_\rho(X)$  restricts to the original  $\mathbb{G}_m$ -equivariant morphism  $f : Y \setminus (0, 0) \rightarrow \text{Bun}_\rho(X)$  over the open subset  $Y \setminus (0, 0) \subset \Sigma$ . We therefore obtain a morphism  $\tilde{f} : [\Sigma/\mathbb{G}_m] \rightarrow \text{Bun}_\rho(X)$  satisfying condition (M2) in Definition 4.14.

We are left to check condition (M3) in Definition 4.14. Choose an integer  $a \geq 1$ . Consider  $\mathbb{P}_\kappa^1$  equipped with the  $\mathbb{G}_m$ -action defined by  $t \cdot [x : y] = [t^{-a}x : y]$ . Suppose that we are given a finite  $\mathbb{G}_m$ -equivariant morphism  $\mathbb{P}_\kappa^1 \rightarrow \Sigma_{(0,0)}$ . By Corollary 4.13, there exists some  $M \gg 0$  such that for all  $n \geq M$  the restriction  $L_n^\vee|_{\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}^{\leq N, P}}$  is  $Y$ -ample. Therefore the restriction  $L_n|_{\mathbb{P}_\kappa^1}$  is anti-ample for  $n \gg 0$ . This means that  $L_n|_{\mathbb{P}_\kappa^1} \cong \mathcal{O}_{\mathbb{P}_\kappa^1}(-N_n)$  for some  $N_n > 0$ . Note that  $\mathcal{O}_{\mathbb{P}_\kappa^1}(-N_n)$  admits a unique  $\mathbb{G}_m$ -equivariant structure up to twisting by a character, and we have  $\text{wgt } \mathcal{O}_{\mathbb{P}_\kappa^1}(-N_n)|_\infty \geq \text{wgt } \mathcal{O}_{\mathbb{P}_\kappa^1}(-N_n)|_0$ . By definition, we have

$$\begin{aligned}
\nu \left( \tilde{f}|_{[\infty/\mathbb{G}_m]} \right) &= \frac{\text{wgt } L_n|_{\infty_i}}{\sqrt{b \left( \tilde{f}|_{[\infty_i/\mathbb{G}_m]} \right)}} \\
\nu \left( \tilde{f}|_{[0/\mathbb{G}_m]} \right) &= \frac{\text{wgt } L_n|_{0_i}}{\sqrt{b \left( \tilde{f}|_{[0_i/\mathbb{G}_m]} \right)}}
\end{aligned}$$

We have just shown that  $\text{wgt } L_n|_\infty \geq \text{wgt } L_n|_0$  for  $n \gg 0$ . Hence, we will be done if we can prove  $b \left( \tilde{f}|_{[\infty/\mathbb{G}_m]} \right) = b \left( \tilde{f}|_{[0/\mathbb{G}_m]} \right)$ . Let  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_{m,0}$  (resp.  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_{m,\infty}$ ) denote the underlying graded torsion-free sheaves of  $\tilde{f}|_{[0/\mathbb{G}_m]}$  (resp.  $\tilde{f}|_{[\infty/\mathbb{G}_m]}$ ). The norm  $b$  only depends on the ranks  $\text{rk}(\mathcal{F}_{m,0})$  (resp.  $\text{rk}(\mathcal{F}_{m,\infty})$ ) of each graded piece. Hence it suffices to show that  $\text{rk}(\mathcal{F}_{m,0}) = \text{rk}(\mathcal{F}_{m,\infty})$  for all  $m \in \mathbb{Z}$ . By construction, the graded  $\rho$ -sheaves  $\tilde{f}|_{[0/\mathbb{G}_m]}$  and  $\tilde{f}|_{[\infty/\mathbb{G}_m]}$  come from the affine Grassmannian  $\text{Gr}_{X_Y, D, \mathcal{F}, \sigma}$ . More

precisely, away from the divisor  $D$ , the underlying graded sheaves  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_{m,0}|_{X_\kappa \setminus D_\kappa}$  and  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_{m,0}|_{X_\kappa \setminus D_\kappa}$  are both isomorphic to the graded sheaf  $\mathcal{F}|_{X_{(0,0)} \setminus D_{(0,0)}}$  after scaling each weight by  $a$ . Since the rank is determined at the generic point of  $X_\kappa$ , this shows that  $\text{rk}(\mathcal{F}_{m,0}) = \text{rk}(\mathcal{F}_{m,\infty})$  for all  $m$ , as desired.  $\square$

**4.4. HN boundedness.** We start by recalling the definition of HN boundedness (see also [HL14, §5] or [HLHJ21, §2.5]).

**Definition 4.17** (HN Boundedness). *Let  $\mathcal{M}$  be an algebraic stack locally of finite type over  $k$ . Let  $\nu$  be a polynomial numerical invariant on  $\mathcal{M}$ . We say that  $\nu$  satisfies the HN boundedness condition if the following is true.*

*Choose an algebraically closed field extension  $K \supset k$ . Let  $S$  be a scheme of finite type over  $K$  and let  $g : S \rightarrow \mathcal{M}$  be a morphism. Then, there exists a quasicompact open substack  $\mathcal{U}_S \subset \mathcal{M}$  such that the following holds: For all closed points  $s \in S$  and all nondegenerate filtrations  $f : \Theta \rightarrow \mathcal{M}$  of the point  $g(s)$  with  $\nu(f) > 0$ , there exists another filtration  $f'$  of  $g(s)$  satisfying  $\nu(f') \geq \nu(f)$  and  $f'|_0 \in \mathcal{U}_S$ .*

In plain words, this says that for all bounded families and all destabilizing filtrations  $f$  of an element in the family, we can always find a more destabilizing filtration  $f'$  such that the associated graded  $f'|_0$  lies on some fixed bounded family.

In the case of  $\text{Bun}_\rho(X)$  we can reduce the proof of HN boundedness to a more concrete statement. This is the content of the following lemma.

**Lemma 4.18.** *In order to show that  $\nu$  satisfies HN boundedness on the stack  $\text{Bun}_\rho(X)$ , it is sufficient to prove the following statement.*

*(\*) Let  $K \supset k$  be an algebraically closed field extension of  $k$ . Let  $S$  be a scheme of finite type over  $K$ . Let  $g : S \rightarrow \text{Bun}_\rho(X)$  be a morphism represented by a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  on  $X_S$ . Then, there exists a real number  $C_S$  such that the following holds: For all closed points  $s \in S$  and all nondegenerate weighted parabolic reductions  $f = (\lambda, \mathcal{G}_\lambda)$  of the point  $(\mathcal{F}|_{X_s}, \sigma|_{X_s})$  with  $\nu(f) > 0$ , there exists another weighted parabolic reduction  $f' = (\lambda', \mathcal{G}_{\lambda'})$  of  $(\mathcal{F}|_{X_s}, \sigma|_{X_s})$  with underlying associated filtration  $(\mathcal{F}'_m)_{m \in \mathbb{Z}}$  such that the following are satisfied.*

- (1)  $\nu(f') \geq \nu(f)$ .
- (2) We have  $\widehat{\mu}_{d-1}(\mathcal{F}'_m/\mathcal{F}'_{m+1}) \geq C_S$  for all  $m \in \mathbb{Z}$ .

*Proof.* We can replace  $k$  by  $K$  in order to assume that the ground field is algebraically closed. Then, all reductive groups are automatically split. By Proposition 3.16, all filtrations of  $(\mathcal{F}|_{X_s}, \sigma|_{X_s})$  arise from weighted parabolic reductions of  $(\mathcal{F}|_{X_s}, \sigma|_{X_s})$ . Let  $\mathfrak{S}$  denote the set of all pairs  $(s, f')$ , where  $s \in S$  is a closed point and  $f'$  is a weighted parabolic reduction of  $(\mathcal{F}|_{X_s}, \sigma|_{X_s})$  satisfying (2) above. It suffices to show that the set of associated graded points  $\mathfrak{S}_2 := \{ f'|_0 \text{ for } (s, f') \in \mathfrak{S} \}$  is contained in a quasicompact open substack of  $\text{Bun}_\rho(X)$ .

Recall that we have a forgetful morphism  $\text{Forget} : \text{Bun}_\rho(X) \rightarrow \text{Coh}_r^{tf}(X)$  into the stack of rank  $r$  torsion-free sheaves on  $X$ . By Proposition 3.6, the morphism  $\text{Forget}$  is of finite type. Hence it is sufficient to prove that the set  $\mathfrak{S}_3 := \{ \text{Forget}(f'|_0) \text{ for } (s, f') \in \mathfrak{S} \}$  is contained in a quasicompact open substack of  $\text{Coh}_r^{tf}(X)$ . Let  $\mathfrak{S}'$  denote the set of pairs  $(s, (\mathcal{F}_m)_{m \in \mathbb{Z}})$ , where  $s \in S$  is a closed point and  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  is a decreasing  $\mathbb{Z}$ -filtration of the sheaf  $\mathcal{F}|_{X_s}$  such that  $\widehat{\mu}_{d-1}(\mathcal{F}_m/\mathcal{F}_{m+1}) \geq C_S$  for all  $m \in \mathbb{Z}$ . It is sufficient to prove that the set of sheaves

$$\mathfrak{S}'_1 = \left\{ \bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m/\mathcal{F}_{m+1} \quad \text{for } (s, (\mathcal{F}_m)_{m \in \mathbb{Z}}) \in \mathfrak{S}' \right\}$$

is contained in a quasicompact open substack of  $\text{Coh}_r^{tf}(X)$ . This follows from [HLHJ21, Lemma 5.4].  $\square$

To prove HN boundedness for  $\text{Bun}_\rho(X)$  in Proposition 4.26, we need to understand how the numerical invariant changes for parabolic subgroups associated to different filtrations. We will use Lemma 4.18. After replacing  $k$  with the extension  $K$ , we can assume that the ground field is algebraically closed (in particular all tori are automatically split). Fix a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  on  $X$ . Let  $j : U \hookrightarrow X$  be a big open subset such that  $\mathcal{F}|_U$  is locally-free. We will develop some setup and prove some preliminary lemmas before the proof of HN boundedness.

Let  $Z$  denote the maximal central torus of  $G$ . Let  $X_*(Z)$  denote the group of cocharacters of  $Z$ . We denote by  $X^*(Z)$  the dual group of characters of  $Z$ . Set  $X_*(Z)_\mathbb{R} := X_*(Z) \otimes \mathbb{R}$  and similarly  $X^*(Z)_\mathbb{R} := X^*(Z) \otimes \mathbb{R}$ . We fix once and for all a basis  $z_1, \dots, z_h$  of  $X_*(Z)$ .

Let  $f = (\lambda, \mathcal{G}_\lambda)$  be a weighted parabolic filtration of  $(\mathcal{F}, \sigma)$ , and let  $P_\lambda \subset G$  be the parabolic subgroup associated to  $\lambda$ . Choose a maximal torus  $T$  of  $P_\lambda$  containing the image of the cocharacter  $\lambda$ . Set  $T' := T \cap [G, G]$ , the intersection of  $T$  with the derived subgroup  $[G, G]$  of  $G$ . Denote the groups of cocharacters of  $T$  and  $T'$  by  $X_*(T)$  and  $X_*(T')$  respectively. Observe that there are natural inclusions  $X_*(Z)_\mathbb{R} \hookrightarrow X_*(T)_\mathbb{R}$  and  $X_*(T')_\mathbb{R} \hookrightarrow X_*(T)_\mathbb{R}$ . These induce a direct sum decomposition  $X_*(T)_\mathbb{R} = X_*(Z)_\mathbb{R} \oplus X_*(T')_\mathbb{R}$ .

Choose a Borel subgroup  $B \supset T$  contained in  $P_\lambda$ . This yields a set of simple roots  $\Delta = \{\alpha_1, \alpha_2, \dots, \alpha_l\} \subset X_*(T')_\mathbb{R}^\vee$ , where  $l$  is the rank of  $G$ , which form a basis for  $X^*(T')_\mathbb{R}$ . Denote by  $\omega_i^\vee \in X_*(T')_\mathbb{R}$  the fundamental coweight dual to the simple root  $\alpha_i$  and define the ordered basis  $\mathcal{B}$  of the vector space  $X_*(T)_\mathbb{R}$  as

$$\mathcal{B} = (z_1, z_2, \dots, z_h, \omega_1^\vee, \omega_2^\vee, \dots, \omega_l^\vee)$$

This induces a dual basis  $\mathcal{B}^\vee = (z_1^\vee, z_2^\vee, \dots, z_h^\vee, \alpha_1, \alpha_2, \dots, \alpha_l)$  on the dual vector space  $X^*(T)_\mathbb{R}$ .

The torus  $T \subset G$  acts on the representation  $V$ . Choose a decomposition of  $V$  as a direct sum of one-dimensional weight spaces

$$V = \bigoplus_{i=1}^r V_{\chi_i}$$

where  $(\chi_i)_{i=1}^r$  is an ordered  $r$ -tuple of characters in  $X^*(T)_\mathbb{R}$  with some possible repetitions.

**Definition 4.19.** *Using coordinates in the dual basis  $\mathcal{B}^\vee$ , we denote by  $M$  the  $(h+l) \times r$ -matrix with coefficients in  $\mathbb{Q}$  corresponding to the tuple  $(\chi_i)_{i=1}^r$ .*

The matrix  $M$  does not depend on the choice of torus  $T$  and Borel subgroup  $B$  (once we fix the basis of  $X_*(Z)$ ), because all torus-Borel pairs  $T \subset B$  are conjugate in  $G$ .

The parabolic  $P_\lambda$  determines a subset  $I_{P_\lambda} \subset \Delta$  consisting of those simple roots  $\alpha_i \in \Delta$  such that the root group  $U_{-\alpha_i}$  corresponding to  $-\alpha_i$  is not contained in  $P_\lambda$  (e.g. we have  $I_B = \Delta$  and  $I_G = \emptyset$ ). Let  $J_{P_\lambda} \subset \{1, 2, \dots, l\}$  be the set of indexes such that  $I_{P_\lambda} = \{\alpha_j \mid j \in J_{P_\lambda}\}$ .

**Definition 4.20.** *We define  $\mathcal{C}_\lambda$  to be the closed  $\mathbb{R}$ -span of the cone of  $P_\lambda$ -dominant coweights in  $X_*(T)_\mathbb{R}$ , with the origin removed:*

$$\mathcal{C}_\lambda = \left( \sum_{i=1}^h \mathbb{R}z_i + \sum_{j \in J_{P_\lambda}} \mathbb{R}^{\geq 0} \omega_j^\vee \right) \setminus \{0\}$$

For any cocharacter  $\delta \in \mathcal{C}_\lambda \cap X_*(T)$ , let  $P_\delta$  be its associated parabolic subgroup. Note that  $P_\delta \supset P_\lambda$ . To every such  $\delta$  we can associate a weighted parabolic reduction  $(\delta, \mathcal{G}_\delta)$ , where  $\mathcal{G}_\delta$  is the  $P_\delta$ -bundle obtained from the  $P_\lambda$ -bundle  $\mathcal{G}_\lambda$  via extension of structure group. The evaluation of the numerical invariant  $\nu$  on these weighted parabolic reductions  $(\delta, \mathcal{G}_\delta)$  defines a function

$$\nu : \mathcal{C}_\lambda \cap X_*(T) \rightarrow \mathbb{R}[n], \quad \delta \mapsto \nu(\delta, \mathcal{G}_\delta).$$

By the formula in Proposition 4.4, the polynomial  $\nu(\delta, \mathcal{G}_\delta)$  has degree less than or equal to  $d - 1$ .

**Definition 4.21.** We define  $\nu_{d-1} : \mathcal{C}_\lambda \cap X_*(T) \rightarrow \mathbb{R}$  to be the function that associates to each  $\delta$  the (leading) degree  $(d - 1)$ -coefficient of the polynomial  $\nu(\delta, \mathcal{G}_\delta)$ .

We denote by  $L_\lambda$  the unique Levi subgroup of the parabolic  $P_\lambda$  containing the maximal torus  $T$ , as described in Subsection 3.3. Let  $Z_\lambda$  be the maximal central torus of  $L_\lambda$ . For each  $i$ , let  $\chi_i^\lambda$  be the restriction of the character  $\chi_i$  to  $X_*(Z_\lambda)$ . Set  $V_{\chi_i^\lambda} = \bigoplus_{\{j | \chi_j^\lambda = \chi_i^\lambda\}} V_{\chi_j^\lambda}$ . We have a  $X^*(Z_\lambda)$ -grading on the vector space  $V$  with weight decomposition:

$$V = \bigoplus_{\{\chi_i^\lambda\}} V_{\chi_i^\lambda}$$

Here the index set  $\{\chi_i^\lambda\}$  runs over the set of all restrictions  $\chi_i^\lambda \in X^*(Z_\lambda)$  without repetitions, and  $Z_\lambda$  acts on  $V_{\chi_i^\lambda}$  with weight  $\chi_i^\lambda$ . Note that each  $V_{\chi_i^\lambda}$  is a representation of  $L_\lambda$ , because  $Z_\lambda$  is central in  $L_\lambda$ . Let  $\mathcal{G}_{L_\lambda}$  denote the  $L_\lambda$ -bundle on  $U$  obtained from  $\mathcal{G}_\lambda$  via the natural quotient morphism  $P_\lambda \rightarrow L_\lambda$ . Let  $\mathcal{G}_{L_\lambda} \times^{L_\lambda} V_{\chi_i^\lambda}$  denote the associated vector bundle on  $U$ .

**Definition 4.22.** We define  $\mathcal{E}_{\chi_i^\lambda}$  to be the reflexive extension  $j_* \left( \mathcal{G}_{L_\lambda} \times^{L_\lambda} V_{\chi_i^\lambda} \right)$  on  $X$ .

We will denote by  $H$  the hyperplane class in  $X$  corresponding to the ample line bundle  $\mathcal{O}(1)$ , and denote by  $c_1(\mathcal{E}_{\chi_i^\lambda})$  the first Chern class of  $\mathcal{E}_{\chi_i^\lambda}$ .

**Lemma 4.23.** Set  $c_i := \frac{c_1(\mathcal{E}_{\chi_i^\lambda}) \cdot H^{d-1}}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})}$  and  $c := \frac{c_1(\mathcal{F}) \cdot H^{d-1}}{r}$ . Then, we have

$$(4) \quad \nu_{d-1}(\delta) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\sum_{i=1}^r \langle \delta, \chi_i^\lambda \rangle \cdot (c_i - c)}{\sqrt{\sum_{i=1}^r (\langle \delta, \chi_i^\lambda \rangle)^2}}$$

In particular, the function  $\nu_{d-1}$  can be naturally interpreted as a scale-invariant continuous function on the whole cone  $\mathcal{C}_\lambda$  of dominant coweights with real coefficients.

*Proof.* For any  $\delta \in \mathcal{C}_\lambda \cap X_*(T)$ , denote by  $L_\delta$  the unique Levi subgroup of the parabolic  $P_\delta$  containing the maximal torus  $T$ . Note that  $L_\lambda \subset L_\delta$ . Denote by  $Z_\delta$  the maximal central torus of  $L_\delta$ .

Let  $\mathcal{G}_{L_\delta}$  be the  $L_\delta$ -bundle obtained from  $\mathcal{G}_\delta$  via the quotient  $P_\delta \rightarrow L_\delta$ . Observe that  $\mathcal{G}_{L_\delta}$  is obtained from  $\mathcal{G}_{L_\lambda}$  by extending the structure group via  $L_\lambda \hookrightarrow L_\delta$ . Let  $\mathcal{E}_U^\delta$  be the vector bundle on  $U$  obtained by extending the structure group via the inclusion  $L_\delta \hookrightarrow G \xrightarrow{\rho} \text{GL}(V)$ . For all  $\delta$  we have a canonical identification  $\mathcal{E}_U^\delta \cong \mathcal{E}_U^\lambda$ . Hence we will denote all such  $\mathcal{E}_U^\delta$  by the same notation  $\mathcal{E}_U$ . Let  $\mathcal{E}$  be the reflexive sheaf on  $X$  obtained by pushing forward  $\mathcal{E}_U$  under the open inclusion  $U \hookrightarrow X$ . Let  $(\mathcal{F}_m^\delta)_{m \in \mathbb{Z}}$  be the underlying filtration of  $\mathcal{F}$  associated to the weighted parabolic reduction  $(\delta, \mathcal{G}_\delta)$ . The description of the associated graded at the end of Subsection 3.3 shows that there is an identification  $(\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta) |_U \xrightarrow{\sim} \mathcal{E}_U$  of associated graded sheaves on  $U$ .

This induces a natural identification  $\bigoplus_{m \in \mathbb{Z}} (\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta)^{\vee \vee} \xrightarrow{\sim} \mathcal{E}$  of reflexive sheaves on  $X$ . We will describe a  $\mathbb{Z}$ -grading on  $\mathcal{E}$  determined by the action of the cocharacter  $\delta$ . The inclusion  $Z_\lambda \hookrightarrow T$  induces an inclusion  $X_*(Z_\lambda) \hookrightarrow X_*(T)$ . Recall that  $\chi_i^\lambda$  denotes the restriction of each character  $\chi_i$  to  $X_*(Z_\lambda)$ . We have a  $X^*(Z_\lambda)$ -grading on the vector space  $V$  with weight decomposition:

$$V = \bigoplus_{\{\chi_i^\lambda\}} V_{\chi_i^\lambda}$$

For any representative  $\chi_i^\lambda$  of  $[\chi_i^\lambda]$  we have  $V_{\chi_i^\lambda} = \bigoplus_{\{j \mid \chi_j^\lambda = \chi_i^\lambda\}} V_{\chi_j^\lambda}$ . Hence  $\dim(V_{\chi_i^\lambda}) = \#\{j \mid \chi_j^\lambda = \chi_i^\lambda\}$ . Since  $Z_\lambda$  is central in  $L_\lambda$ , the action of  $L_\lambda$  on  $V$  preserves each  $Z_\lambda$ -weight space  $V_{\chi_i^\lambda}$ , hence each  $V_{\chi_i^\lambda}$  is a representation of  $L_\lambda$ . This way we get the grading

$$\mathcal{E}_U = \bigoplus_{\{\chi_i^\lambda\}} \mathcal{E}_{U, \chi_i^\lambda}$$

where  $\mathcal{E}_{U, \chi_i^\lambda}$  is the vector bundle  $\mathcal{G}_{L_\lambda} \times^{L_\lambda} V_{\chi_i^\lambda}$  associated to the representation  $V_{\chi_i^\lambda}$  and  $Z_\lambda$  acts with weight  $\chi_i^\lambda$  on  $\mathcal{E}_{U, \chi_i^\lambda}$ . Note that  $\text{rk}(\mathcal{E}_{U, \chi_i^\lambda}) = \dim(V_{\chi_i^\lambda})$ . This grading extends uniquely to a grading of the reflexive extension

$$\bigoplus_{m \in \mathbb{Z}} (\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta)^{\vee\vee} \cong \mathcal{E} = \bigoplus_{\{\chi_i^\lambda\}} \mathcal{E}_{\chi_i^\lambda}$$

The cocharacter  $\delta$  acts on  $\mathcal{E}_U$  via its natural inclusion in  $Z_\delta$ . Using  $Z_\delta \hookrightarrow Z_\lambda$ , we can view  $\delta$  as a cocharacter of  $Z_\lambda$ . It follows from our discussion that  $\delta$  acts with weight  $\langle \delta, \chi_i^\lambda \rangle$  on  $\mathcal{E}_{U, \chi_i^\lambda}$ . This shows that

$$(\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta)|_U = \bigoplus_{\{\chi_i^\lambda \mid \langle \delta, \chi_i^\lambda \rangle = m\}} \mathcal{E}_{U, \chi_i^\lambda}$$

and therefore

$$(5) \quad (\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta)^{\vee\vee} = \bigoplus_{\{\chi_i^\lambda \mid \langle \delta, \chi_i^\lambda \rangle = m\}} \mathcal{E}_{\chi_i^\lambda}$$

Observe that we have equality of slopes  $\widehat{\mu}_{d-1}(\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta) = \widehat{\mu}_{d-1}((\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta)^{\vee\vee})$ . Taking the leading coefficient in the formula (3) in Proposition 4.4, we get

$$(6) \quad \nu_{d-1}(\delta) = \frac{\sqrt{A_d}}{(d-1)!} \cdot \frac{\sum_{m \in \mathbb{Z}} m \cdot (\widehat{\mu}_{d-1}(\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta) - \widehat{\mu}_{d-1}(\mathcal{F})) \cdot \text{rk}(\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta)}{\sqrt{\sum_{m \in \mathbb{Z}} m^2 \cdot \text{rk}(\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta)}}$$

Using the description of the quotients  $\mathcal{F}_m^\delta / \mathcal{F}_{m+1}^\delta$  in terms of the  $\mathcal{E}_{\chi_i^\lambda}$ , we can rewrite (6) as:

$$(7) \quad \begin{aligned} \nu_{d-1}(\delta) &= \frac{\sqrt{A_d}}{(d-1)!} \cdot \frac{\sum_{m \in \mathbb{Z}} \sum_{\{\chi_i^\lambda \mid \langle \delta, \chi_i^\lambda \rangle = m\}} m \cdot (\widehat{\mu}_{d-1}(\mathcal{E}_{\chi_i^\lambda}) - \widehat{\mu}_{d-1}(\mathcal{F})) \cdot \#\{j \mid \chi_j^\lambda = \chi_i^\lambda\}}{\sqrt{\sum_{m \in \mathbb{Z}} m^2 \cdot \#\{j \mid \langle \delta, \chi_i^\lambda \rangle = m\}}} \\ &= \frac{\sqrt{A_d}}{(d-1)!} \cdot \frac{\sum_{i=1}^r \langle \delta, \chi_i^\lambda \rangle \cdot (\widehat{\mu}_{d-1}(\mathcal{E}_{\chi_i^\lambda}) - \widehat{\mu}_{d-1}(\mathcal{F}))}{\sqrt{\sum_{i=1}^r (\langle \delta, \chi_i^\lambda \rangle)^2}} \end{aligned}$$

Observe that in the sum above we are indexing over all  $\chi_i^\lambda$  with repetitions. This is done in order to account for the ranks.

We can further rewrite this by using the Hirzebruch-Riemann-Roch theorem [Ful98, Ch. 15]. Let  $H$  denote the hyperplane class on  $X$  corresponding to the line bundle  $\mathcal{O}(1)$ . Let  $t_1$  be the degree 1 term of the Todd class of  $X$ . For any torsion-free sheaf  $\mathcal{A}$  on  $X$  with first Chern class  $c_1(\mathcal{A})$ , Hirzebruch-Riemann-Roch implies that

$$\widehat{\mu}_{d-1}(\mathcal{A}) = \frac{1}{A_d \text{rk}(\mathcal{A})} c_1(\mathcal{A}) \cdot H^{d-1} + \frac{1}{A_d} t_1 \cdot H^{d-1}$$

Using this formula for  $\mathcal{E}_{\chi_i^\lambda}$  and  $\mathcal{F}$  yields

$$\widehat{\mu}_{d-1}(\mathcal{E}_{\chi_i^\lambda}) - \widehat{\mu}_{d-1}(\mathcal{F}|_{X_s}) = \frac{1}{A_d} \cdot \left( \frac{c_1(\mathcal{E}_{\chi_i^\lambda}) \cdot H^{d-1}}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})} - \frac{c_1(\mathcal{F}) \cdot H^{d-1}}{r} \right)$$

Using  $c_i = \frac{c_1(\mathcal{E}_{\chi_i^\lambda}) \cdot H^{d-1}}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})}$  and  $c = \frac{c_1(\mathcal{F}) \cdot H^{d-1}}{r}$ , we can rewrite (7) as:

$$\nu_{d-1}(\delta) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\sum_{i=1}^r \langle \delta, \chi_i^\lambda \rangle \cdot (c_i - c)}{\sqrt{\sum_{i=1}^r (\langle \delta, \chi_i^\lambda \rangle)^2}}$$

□

For each  $\delta \in X_*(T) \cap \mathcal{C}_\lambda$ , the inclusion  $Z_\delta \hookrightarrow T$  induces an identification

$$X_*(Z_\delta)_\mathbb{R} = \bigoplus_{i=1}^h \mathbb{R}z_i \oplus \bigoplus_{j \in J_{P_\delta}} \mathbb{R}\omega_j^\vee \hookrightarrow X_*(Z_\lambda)_\mathbb{R}$$

where the last inclusion comes from  $Z_\delta \subset Z_\lambda$ . Let us denote by  $\chi_i^\delta$  the restriction of each character  $\chi_i$  to  $X_*(Z_\delta)_\mathbb{R}$ . In coordinates of the basis  $\mathcal{B}^\vee$ , the tuple  $(\chi_i^\delta)_{i=1}^r$  is the matrix  $M_\delta$  obtained from  $M$  by deleting the rows corresponding to  $\Delta \setminus I_{P_\delta}$ . Note that the set of all such possible matrices  $M_\delta$  as we vary  $\delta$  is finite, because there are only finitely many possibilities for the set of indexes  $J_{P_\delta}$ .

As a consequence of Lemma 4.23 we get the following immediate corollary.

**Corollary 4.24.** *Set  $\vec{a} = [c_i]_{i=1}^r$  to be the row vector of size  $r$  with the  $i^{\text{th}}$  element given by  $c_i$ , and let  $\vec{1}$  denote the row vector of size  $r$  such that all entries are 1. Let  $\vec{x} = [x_i]$  be the column vector of the coordinates of  $\delta \in \mathcal{C}_\lambda$  in the basis  $\mathcal{B}$ . Then we can express  $\nu_{d-1}$  in terms of the matrix  $M_\lambda$  as follows.*

$$(8) \quad \nu_{d-1}(\vec{x}) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{(\vec{a} - c\vec{1})M_\lambda\vec{x}}{\sqrt{\vec{x}^T M_\lambda^T M_\lambda \vec{x}}}$$

□

**Lemma 4.25.** *There exists a linear functional  $\psi : X^*(Z_\lambda)_\mathbb{R} \rightarrow \mathbb{R}$  such that  $c_i = \psi(\chi_i^\lambda)$  for all  $i$ .*

*Proof.* Recall that  $j : U \hookrightarrow X_s$  is a big open set and  $X_s$  is smooth. The pushforward  $j_*$  induces an isomorphism of Picard groups  $j_* : \text{Pic}(U) \rightarrow \text{Pic}(X_s)$ . Let  $X^*(L_\lambda)$  be the group of characters of the Levi subgroup  $L_\lambda$ . The  $L_\lambda$ -bundle  $\mathcal{G}_{L_\lambda}$  on  $U$  induces a homomorphism  $X^*(L_\lambda) \rightarrow \text{Pic}(U)$  where any character  $\chi \in X^*(L_\lambda)$  is mapped to the associated line bundle  $\mathcal{G}_{L_\lambda} \times^{L_\lambda} \chi$ . By post-composing with  $j_*$  we get a homomorphism

$$X^*(L_\lambda) \rightarrow \text{Pic}(U) \rightarrow \text{Pic}(X_s)$$

On the other hand, the first Chern class induces a homomorphism

$$\text{Pic}(X_s) \rightarrow \mathbb{Z}, \quad \mathcal{L} \mapsto c_1(\mathcal{L}) \cdot H^{d-1} \in \mathbb{Z}.$$

We compose this with the previous map to get a homomorphism

$$\varphi : X^*(L_\lambda) \rightarrow \mathbb{Z}, \quad \chi \mapsto c_1(j_*(\mathcal{G}_{L_\lambda} \times^{L_\lambda} \chi)) \cdot H^{d-1}$$

Tensor with  $\mathbb{R}$  to obtain a linear functional  $\varphi_\mathbb{R} : X^*(L_\lambda)_\mathbb{R} \rightarrow \mathbb{R}$ .

The inclusion  $Z_\lambda \hookrightarrow L_\lambda$  yields a restriction map on characters  $\text{res} : X^*(L_\lambda)_\mathbb{R} \rightarrow X^*(Z_\lambda)_\mathbb{R}$ . This morphism  $\text{res}$  is an isomorphism since the inclusion induces an isogeny between  $Z_\lambda$  and  $L_\lambda/[L_\lambda, L_\lambda]$ . We set  $\psi$  to be the linear functional

$$\psi = \varphi_\mathbb{R} \circ (\text{res})^{-1} : X^*(Z_\lambda) \rightarrow \mathbb{R}, \quad \bar{\chi} \mapsto \varphi_\mathbb{R}(\chi) = c_1(j_*(\mathcal{G}_{L_\lambda} \times^{L_\lambda} \chi)) \cdot H^{d-1}$$

for a character  $\bar{\chi}$  of  $Z_\lambda$  that comes from the restriction of a character  $\chi$  of  $L_\lambda$ . We are left to prove that  $\psi(\chi_i^\lambda) = c_i$  for all  $i$ .

By definition  $\mathcal{E}_{\chi_i^\lambda}|_U$  is the associated vector bundle  $\mathcal{G}_{L_\lambda} \times^{L_\lambda} V_{\chi_i^\lambda}$ . Hence we have  $\det(\mathcal{E}_{\chi_i^\lambda}|_U) = \mathcal{G}_\lambda \times^{L_\lambda} \det(V_{\chi_i^\lambda})$ . Therefore:

$$c_i = \frac{c_1(\mathcal{E}_{\chi_i^\lambda}) \cdot H^{d-1}}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})} = \frac{c_1(\det(\mathcal{E}_{\chi_i^\lambda})) \cdot H^{d-1}}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})} = \frac{c_1(j_* \det(\mathcal{E}_{\chi_i^\lambda}|_U)) \cdot H^{d-1}}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})} = \frac{\varphi_{\mathbb{R}}(\det(V_{\chi_i^\lambda}))}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})}$$

As a representation of  $Z_\lambda$ , the weight space  $V_{\chi_i^\lambda}$  is a direct sum of  $\text{rk}(\mathcal{E}_{\chi_i^\lambda})$ -many copies of the character  $\chi_i^\lambda$ . Hence  $\det(V_{\chi_i^\lambda}) = \text{rk}(\mathcal{E}_{\chi_i^\lambda}) \cdot \chi_i^\lambda$  as a  $Z_\lambda$ -character. This means

$$c_i = \frac{\varphi_{\mathbb{R}}(\det(V_{\chi_i^\lambda}))}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})} = \frac{\psi(\text{rk}(\mathcal{E}_{\chi_i^\lambda}) \cdot \chi_i^\lambda)}{\text{rk}(\mathcal{E}_{\chi_i^\lambda})} = \psi(\chi_i^\lambda)$$

□

After all of our preliminary work, we are now ready to prove the main proposition of this subsection.

**Proposition 4.26.** *The polynomial numerical invariant  $\nu$  on the stack  $\text{Bun}_\rho(X)$  satisfies the HN boundedness condition.*

*Proof.* We shall prove property (\*) in Lemma 4.18. Fix an algebraically closed extension  $K \supset k$  and a scheme  $S$  of finite type over  $K$ . After replacing  $k$  with the extension  $K$ , we can assume that the ground field  $k$  is algebraically closed. Fix a family  $g : S \rightarrow \text{Bun}_\rho(X)$  represented by a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  on  $X_S$ .

Choose a closed point  $s \in S$ . Let  $f = (\lambda, \mathcal{G}_\lambda)$  be a nondegenerate weighted reduction of  $(\mathcal{F}|_{X_s}, \sigma|_{X_s})$ . Let  $U$  be a big open subset of  $X_s$  where  $\mathcal{F}|_{X_s}$  is locally-free and  $(\lambda, \mathcal{G}_\lambda)$  is defined. Suppose that  $\nu(f) > 0$ . The first step will be to construct another weighted parabolic reduction  $f'$  that satisfies  $\nu(f') \geq \nu(f)$ .

Let  $P_\lambda \subset G$  the parabolic subgroup associated to the cocharacter  $\lambda$ . Note that the discussion and lemmas before the proposition apply to the weighted parabolic reduction  $(\lambda, \mathcal{G}_\lambda)$  of  $(\mathcal{F}|_{X_s}, \sigma|_{X_s})$ . The scale-invariant continuous function  $\nu_{d-1}$  (see (4) in Lemma 4.23) attains a maximal value  $\nu_{d-1}(\delta^*)$  for some  $\delta^*$  on the domain  $\mathcal{C}_\lambda$  (by restricting to the compact intersection of  $\mathcal{C}_\lambda$  with the unit sphere). We can assume without loss of generality that the maximum  $\delta^*$  is in the interior of the cone  $\mathcal{C}_\lambda$ . Otherwise, let  $F$  be the unique facet of  $\mathcal{C}_\lambda$  such that  $\delta^* \in \text{Int}(F)$ . This facet  $F$  is the smaller cone of dominant coweights of a parabolic subgroup  $P_{\delta^*} \supset P_\lambda$ . Then we can replace  $\lambda$  with a  $\delta^*$  and replace the parabolic reduction  $(\lambda, \mathcal{G}_\lambda)$  with  $(\delta^*, \mathcal{G}_{\delta^*})$  for the bigger parabolic.

Since the maximum  $\delta^*$  is in the interior of  $\mathcal{C}_\lambda$ , it must be a critical point of the function  $\nu_{d-1} : \mathcal{C}_\lambda \rightarrow \mathbb{R}$  as in (8) in Corollary 4.24. If  $\nu_{d-1}$  is not identically 0 (equivalently  $(\vec{a} - c\vec{1})M_\lambda \neq 0$  in (8)), the computation of the gradient yields the coordinates  $\vec{z}$  of a critical point, up to scaling:

$$(9) \quad \vec{z} = (M_\lambda^T M_\lambda)^{-1} M_\lambda^T (\vec{a}^T - c\vec{1}^T)$$

We consider three cases to construct a modified weighted parabolic reduction  $f'$ :

(C1) Suppose that  $\nu_{d-1}(\lambda) \neq \nu_{d-1}(\delta^*)$ . Then  $\nu_{d-1}$  is not identically 0. Since all of the matrices and vectors in (9) have rational coordinates, we can scale  $\delta^*$  to have rational coordinates. After further scaling, we can assume that  $\delta^*$  is a cocharacter in  $\mathcal{C}_\lambda \cap X_*(T)$ . It has coordinates

$$\vec{y} = q\vec{z} = q(M_\lambda^T M_\lambda)^{-1} M_\lambda^T (\vec{a}^T - c\vec{1}^T)$$

for  $q \in \mathbb{Q}_{>0}$  (the negative multiples correspond to minimal critical points). Set the modified weighted parabolic reduction to be  $f' = (\delta^*, \mathcal{G}_{\delta^*})$  with  $\nu_{d-1}(\delta^*) > \nu_{d-1}(\lambda)$ . By construction we have  $\nu(f') > \nu(f)$ .

- (C2) Suppose that  $\nu_{d-1}(\lambda) = \nu_{d-1}(\delta^*)$  and the function  $\nu_{d-1}$  is not identically 0. Then  $\lambda$  is a critical point for  $\nu_{d-1}$  in the interior of the cone  $\mathcal{C}_\lambda$  with coordinates  $\vec{y} = q\vec{z}$ . In this case we define the weighted parabolic reduction  $f'$  to be the original  $f = (\lambda, \mathcal{G}_\lambda)$  and set  $\delta^* = \lambda$ .
- (C3) Suppose that  $\nu_{d-1}(\lambda) = \nu_{d-1}(\delta^*)$  and  $\nu_{d-1}$  is identically 0. In this case  $\vec{z} = 0$ . We set  $f' = f$  and denote  $\delta^* = \lambda$ .

Let  $f' = (\delta^*, \mathcal{G}_{\delta^*})$  with maximizing  $\delta^*$  as explained for each case above. Let  $(\mathcal{F}_m^{\delta^*})_{m \in \mathbb{Z}}$  be its corresponding filtration with associated graded  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m^{\delta^*} / \mathcal{F}_{m+1}^{\delta^*}$ . We are left to find a uniform lower bound for the slopes  $\hat{\mu}_{d-1}(\mathcal{F}_m^{\delta^*} / \mathcal{F}_{m+1}^{\delta^*})$ . Since the slope of  $\mathcal{F}_m^{\delta^*} / \mathcal{F}_{m+1}^{\delta^*}$  is the same as the slope of its reflexive hull  $(\mathcal{F}_m^{\delta^*} / \mathcal{F}_{m+1}^{\delta^*})^{\vee\vee}$ , this is equivalent to giving a uniform lower bound for

$$\hat{\mu}_{d-1} \left( (\mathcal{F}_m^{\delta^*} / \mathcal{F}_{m+1}^{\delta^*})^{\vee\vee} \right) = \hat{\mu}_{d-1} \left( \bigoplus_{\{\chi_i^\lambda \mid \langle \delta^*, \chi_i^\lambda \rangle = m\}} \mathcal{E}_{\chi_i^\lambda} \right)$$

(See Equation (5) in the proof of Lemma 4.23). By additivity of degrees and ranks, it suffices to give a uniform lower bound for the slope  $\hat{\mu}(\mathcal{E}_{\chi_i^\lambda})$  of each direct summand above. Hirzebruch-Riemann-Roch implies that

$$\hat{\mu}_{d-1}(\mathcal{E}_{\chi_i^\lambda}) = \frac{1}{A_d} \left( c_i + t_1 \cdot H^{d-1} \right),$$

Therefore it suffices to give a lower bound for the numbers  $c_i$ . Denote the coordinates in  $\mathcal{B}^\vee$  of the linear functional  $\psi$  of Lemma 4.25 by the column vector  $\vec{\psi}$ . Observe that  $\psi(\chi_i^\lambda) = c_i$  translates into  $\vec{a}^T = M_\lambda \vec{\psi}$ . We shall prove uniform bounds for the vector  $\vec{a} = [c_i]$ . In cases (C1) and (C2) the coordinates  $\vec{y}$  of  $\delta^*$  are given by

$$\vec{y} = q\vec{z} = q(M_\lambda^T M_\lambda)^{-1} M_\lambda^T (\vec{a}^T - c\vec{1})$$

We can plug-in  $\vec{a}^T = M_\lambda \vec{\psi}$  to obtain the following:

$$\vec{y} = q(M_\lambda^T M_\lambda)^{-1} M_\lambda^T M_\lambda \vec{\psi} - qc(M_\lambda^T M_\lambda)^{-1} M_\lambda^T \vec{1}^T = q\vec{\psi} - qc(M_\lambda^T M_\lambda)^{-1} M_\lambda^T \vec{1}^T$$

Multiply both sides by  $M_\lambda$  to get

$$(10) \quad \vec{a}^T = M_\lambda \vec{\psi} = \frac{1}{q} M_\lambda \vec{y} + c M_\lambda (M_\lambda^T M_\lambda)^{-1} M_\lambda^T \vec{1}^T.$$

Since the family  $\mathcal{F}$  is bounded, we see that  $c = \frac{c_1(\mathcal{F}|_{X_s}) \cdot H^{d-1}}{r}$  admits uniform upper and lower bounds that do not depend on  $s$ . Also we know that there are finitely many possibilities for the matrix  $M_\lambda$ , one for each possible choice of subset  $I_{P_\lambda}$  of simple roots corresponding to the parabolic  $P_\lambda$ . This gives uniform upper and lower bounds  $C_{max}$  and  $C_{min}$  for the coefficients of matrix of the second term  $c M_\lambda (M_\lambda^T M_\lambda)^{-1} M_\lambda^T \vec{1}^T$  in (10), independent of  $s$  and the weighted parabolic reduction  $f$ .

The  $i^{th}$  coefficient of the row vector  $M_\lambda \vec{y}$  is the weight  $\langle \delta^*, \chi_i^\lambda \rangle$  of the sheaf  $\mathcal{E}_{\chi_i^\lambda}$ . From (10) we get for each index  $i$ :

$$C_{min} + \frac{1}{q} \text{wgt}(\mathcal{E}_{\chi_i^\lambda}) \leq c_i \leq C_{max} + \frac{1}{q} \text{wgt}(\mathcal{E}_{\chi_i^\lambda})$$

We can add  $t_1 \cdot H^{d-1}$  to get

$$(11) \quad C_{min} + \frac{1}{q} \text{wgt}(\mathcal{E}_{\chi_i^\lambda}) + t_1 \cdot H^{d-1} \leq A_d \cdot \hat{\mu}_{d-1}(\mathcal{E}_{\chi_i^\lambda}) \leq C_{max} + \frac{1}{q} \text{wgt}(\mathcal{E}_{\chi_i^\lambda}) + t_1 \cdot H^{d-1}$$



Let  $\text{wgt}(\mathcal{E}_{\chi_{i_{\min}}^\lambda})$  be the smallest weight among all of the  $\mathcal{E}_{\chi_i^\lambda}$ . Since  $q > 0$ , we obtain

$$\frac{1}{A_d} C_{\min} + \frac{1}{qA_d} \text{wgt}(\mathcal{E}_{\chi_{i_{\min}}^\lambda}) + \frac{1}{A_d} t_1 \cdot H^{d-1} \leq \frac{1}{A_d} C_{\min} + \frac{1}{qA_d} \text{wgt}(\mathcal{E}_{\chi_i^\lambda}) + \frac{1}{A_d} t_1 \cdot H^{d-1} \leq \widehat{\mu}_{d-1}(\mathcal{E}_{\chi_i^\lambda})$$

for each  $i$ . So it suffices to give a uniform lower bound for  $\frac{1}{qA_d} \text{wgt}(\mathcal{E}_{\chi_{i_{\min}}^\lambda})$ . Replace  $\mathcal{E}_{\chi_i^\lambda}$  by  $\mathcal{E}_{\chi_{i_{\min}}^\lambda}$  in the upper inequality of (11) to get

$$\widehat{\mu}_{d-1}(\mathcal{E}_{\chi_{i_{\min}}^\lambda}) - \frac{1}{A_d} C_{\max} - \frac{1}{A_d} t_1 \cdot H^{d-1} \leq \frac{1}{qA_d} \text{wgt}(\mathcal{E}_{\chi_{i_{\min}}^\lambda}).$$

Therefore we just need to give a uniform lower bound for  $\widehat{\mu}_{d-1}(\mathcal{E}_{\chi_{i_{\min}}^\lambda})$ .

Set  $m_{\min} := \langle \delta^*, \chi_{i_{\min}}^\lambda \rangle$ . Note that  $\mathcal{E}_{\chi_{i_{\min}}^\lambda}$  is a direct summand of the reflexive hull of the graded piece  $\mathcal{F}_{m_{\min}}^{\delta^*} / \mathcal{F}_{m_{\min}+1}^{\delta^*}$  of the filtration  $(\mathcal{F}_m^{\delta^*})_{m \in \mathbb{Z}}$ . We can think of  $\mathcal{E}_{\chi_{i_{\min}}^\lambda}$  as a quotient of the reflexive hull  $(\mathcal{F}_{m_{\min}}^{\delta^*} / \mathcal{F}_{m_{\min}+1}^{\delta^*})^{\vee\vee}$ , and there is a corresponding quotient sheaf  $\mathcal{F}_{m_{\min}}^{\delta^*} / \mathcal{F}_{m_{\min}+1}^{\delta^*} \twoheadrightarrow \mathcal{Q}$  with the same slope as  $\mathcal{E}_{\chi_{i_{\min}}^\lambda}$ .

Since the weight of  $\mathcal{E}_{\chi_{i_{\min}}^\lambda}$  is minimal, it follows that  $\mathcal{F}_{m_{\min}}^{\delta^*} / \mathcal{F}_{m_{\min}+1}^{\delta^*}$  is the nontrivial graded piece with the smallest weight. This implies that  $\mathcal{F}_{m_{\min}}^{\delta^*} / \mathcal{F}_{m_{\min}+1}^{\delta^*}$  is a quotient of  $\mathcal{F}|_{X_s}$ . Hence  $\mathcal{Q}$  is also a quotient of  $\mathcal{F}|_{X_s}$ . It follows that

$$\widehat{\mu}_{d-1}(\mathcal{E}_{\chi_{i_{\min}}^\lambda}) = \widehat{\mu}_{d-1}(\mathcal{Q}) \geq \widehat{\mu}_{d-1}(\mathcal{F}|_{X_s})_{\min},$$

where  $\widehat{\mu}_{d-1}(\mathcal{F}|_{X_s})_{\min}$  is the smallest slope appearing among the associated graded pieces of the Gieseker-Harder-Narasimhan filtration of  $\mathcal{F}|_{X_s}$  (c.f. [HL97, Def. 1.3.2]). The quantity  $\widehat{\mu}_{d-1}(\mathcal{F}|_{X_s})_{\min}$  is uniformly bounded as we vary  $t \in T$ , by constructibility of the locus of each fixed Harder-Narasimhan type (see e.g. [Nit11]). This concludes the proof of cases (C1) and (C2).

If we are in case (C3) we have  $\vec{z} = 0$  (see (9)). Then:

$$(M_\lambda^T M_\lambda)^{-1} M_\lambda^T \vec{a}^T = c(M_\lambda^T M_\lambda)^{-1} M_\lambda^T \vec{1}^T$$

Using  $\vec{a}^T = M_\lambda \vec{\psi}$  we get  $\vec{a}^T = cM_\lambda (M_\lambda^T M_\lambda)^{-1} M_\lambda^T \vec{1}^T$ , which is the second term in (10) and is bounded by our discussion of the cases before.  $\square$

**4.5.  $\Theta$ -stratification, semistability and leading term filtrations.** After all of our work in this section, we can now construct a  $\Theta$ -stratification on  $\text{Bun}_\rho(X)$ .

**Theorem 4.27.** *The polynomial numerical invariant  $\nu$  defines a  $\Theta$ -stratification on the stack  $\text{Bun}_\rho(X)$ .*

*Proof.* [HLHJ21, Thm. 2.26] (or [HL14, Thm. B]) shows that it suffices to check that  $\nu$  satisfies HN boundedness and is strictly  $\Theta$ -monotone on the stack  $\text{Bun}_\rho(X)$ . This follows by Propositions 4.26 and 4.16.  $\square$

The following is a translation of the notion of  $\Theta$ -semistability from [HL14, §2.1] for our numerical invariant  $\nu$ .

**Definition 4.28.** *Let  $K \supset k$  be an algebraically closed field extension. Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . We say that  $(\mathcal{F}, \sigma)$  is Gieseker semistable if for all weighted parabolic reductions  $(\lambda, \mathcal{G}_\lambda)$  with associated underlying filtrations  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  (as in Definition 3.15) we have*

$$\sum_{m \in \mathbb{Z}} m \cdot (\overline{p}_{\mathcal{F}_m / \mathcal{F}_{m+1}} - \overline{p}_{\mathcal{F}}) \cdot \text{rk}(\mathcal{F}_m / \mathcal{F}_{m+1}) \leq 0$$

In general, if  $K \supset k$  is any field extension, then we say that a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  on  $X_K$  is Gieseker semistable if the pullback  $(\mathcal{F}|_{X_{\overline{K}}}, \sigma|_{X_{\overline{K}}})$  is semistable for any algebraic closure  $\overline{K} \supset K$ . A  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  on  $X_K$  is called unstable if it is not Gieseker semistable.

The summation by parts argument in [HLHJ21, Prop. 4.1] shows that the expression in Definition 4.28 can be rewritten as

$$(12) \quad \sum_{m \in \mathbb{Z}} m \cdot (\overline{p}_{\mathcal{F}_m/\mathcal{F}_{m+1}} - \overline{p}_{\mathcal{F}}) \cdot \text{rk}(\mathcal{F}_m/\mathcal{F}_{m+1}) = \sum_{m \in \mathbb{Z}} (\overline{p}_{\mathcal{F}_m} - \overline{p}_{\mathcal{F}}) \cdot \text{rk}(\mathcal{F}_m)$$

By Theorem 4.27, the polynomial numerical invariant  $\nu$  defines a  $\Theta$ -stratification on  $\text{Bun}_{\rho}(X)$ . In particular the locus of semistable  $\rho$ -sheaves is an open substack, which we will denote by  $\text{Bun}_{\rho}(X)^{ss}$ . Note that  $\text{Bun}_{\rho}(X)^{ss}$  is a disjoint union of open and closed substacks where the Hilbert polynomial of the underlying torsion-free sheaf is constant. For every rational polynomial  $P \in \mathbb{Q}[n]$ , we will denote by  $\text{Bun}_{\rho}(X)^{ss, P}$  the open and closed substack of  $\text{Bun}_{\rho}(X)^{ss}$  parametrizing Gieseker semistable  $\rho$ -sheaves  $(\mathcal{F}, \sigma)$  such that the Hilbert polynomial of  $\mathcal{F}$  is  $P$ .

We will next explain what it means to have a  $\Theta$ -stratification on  $\text{Bun}_{\rho}(X)$  in terms of canonical filtrations.

**Definition 4.29.** *Suppose that  $K$  is algebraically closed. Let  $(\mathcal{F}, \sigma)$  be an unstable  $\rho$ -sheaf on  $X_K$ . A weighted parabolic reduction  $f = (\lambda, \mathcal{G}_{\lambda})$  is called canonical if the polynomial invariant  $\nu(f)$  is maximal among all weighted parabolic reductions of  $(\mathcal{F}, \sigma)$ . We say that the associated underlying filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  to  $f = (\lambda, \mathcal{G}_{\lambda})$  is a canonical filtration of  $(\mathcal{F}, \sigma)$ .*

We already know that a weighted parabolic reduction  $(\lambda, \mathcal{G}_{\lambda})$  determines a unique underlying filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$ . By Proposition 3.9, the converse is true. Namely, the underlying filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  uniquely determines the weighted parabolic reduction  $(\lambda, \mathcal{G}_{\lambda})$  over an open subset  $U$  where  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m/\mathcal{F}_{m+1}$  is locally-free. Hence the data of a canonical weighted parabolic reduction is equivalent to specifying its underlying canonical filtration.

**Definition 4.30.** *Let  $K \supset k$  be an arbitrary field extension. Let  $(\mathcal{F}, \sigma)$  be an unstable  $\rho$ -sheaf on  $X_K$ . A  $\mathbb{Z}$ -weighted filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  of  $\mathcal{F}$  is called a canonical filtration if for any algebraic closure  $\overline{K} \supset K$  we have that  $(\mathcal{F}_m|_{X_{\overline{K}}})_{m \in \mathbb{Z}}$  is a canonical filtration of  $(\mathcal{F}|_{X_{\overline{K}}}, \sigma|_{X_{\overline{K}}})$ .*

The fact that  $\nu$  defines a  $\Theta$ -stratification on  $\text{Bun}_{\rho}(X)$  implies that any field valued point of  $\text{Bun}_{\rho}(X)$  admits a  $\nu$ -maximizing  $\Theta$ -filtration that is unique up to scaling the weights. We can use our concrete descriptions of  $\Theta$ -filtrations for  $\text{Bun}_{\rho}(X)$  to translate this into the following.

**Proposition 4.31.** *Let  $K \supset k$  be an arbitrary field extension. Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . Then,  $(\mathcal{F}, \sigma)$  admits a canonical filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  which is uniquely determined up to scaling all of the indexes by a constant rational number.*

Furthermore, if the group  $G_K$  is split, then there exists a (uniquely determined) canonical weighted parabolic reduction  $(\lambda, \mathcal{G}_{\lambda})$  defined over  $K$  such that its associated underlying filtration is  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$ .  $\square$

**Definition 4.32.** *Let  $(\mathcal{F}, \sigma)$  be field valued point of  $\text{Bun}_{\rho}(X)$ . Suppose that  $(\mathcal{F}, \sigma)$  is unstable. We call the canonical filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  the leading term HN filtration of  $(\mathcal{F}, \sigma)$ . It is uniquely determined up to scaling the weights, by Proposition 4.31.*

On the other hand, if  $(\mathcal{F}, \sigma)$  is Gieseker semistable, then we define the leading term HN filtration to be  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  where

$$\mathcal{F}_m = \begin{cases} \mathcal{F} & \text{if } m \leq 0 \\ 0 & \text{otherwise} \end{cases}$$

**Remark 4.33.** *Suppose that  $f$  is the leading term HN filtration of  $(\mathcal{F}, \sigma)$ . If the polynomial  $\nu(f)$  has degree  $i$ , then we say that  $i$  is the leading degree of  $(\mathcal{F}, \sigma)$ . One should think of the leading term HN filtration as an analogue of the slope filtration. Here the coefficient of degree  $i$  in the reduced Hilbert polynomial plays the role of the slope. This can be made precise in the case of torsion-free sheaves, see Subsection 9.1.*

The existence of a  $\Theta$ -stratification also implies that the notion of leading term HN filtration behaves well in families. In order to explain this precisely, we need to first define what is a filtration of a family of  $\rho$ -sheaves.

Let  $S$  be a  $k$ -scheme and let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_S$ . A filtration of  $\mathcal{F}$  is a  $\mathbb{Z}$ -weighted filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  of  $\mathcal{F}$  satisfying the following properties.

- (1)  $\mathcal{F}_{m+1} \subset \mathcal{F}_m$ .
- (2)  $\mathcal{F}_m = 0$  for  $m \gg 0$  and  $\mathcal{F}_m = \mathcal{F}$  for  $m \ll 0$ .
- (3)  $\mathcal{F}_m/\mathcal{F}_{m+1}$  is a  $S$ -flat family of torsion-free sheaves.

An argument similar to Proposition 3.8 shows that the data of such filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  is equivalent to the data of a morphism of stacks  $\Theta_S \rightarrow \mathrm{Coh}_r^{tf}(X)$  such that the restriction to  $1_S$  recovers  $\mathcal{F}$ . This amounts to a torsion-free sheaf  $\tilde{\mathcal{F}}$  on  $X_{\Theta_S}$ . Let  $U$  be a big open subset of  $X_S$  where  $\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m/\mathcal{F}_{m+1}$  is locally-free. Then, the restriction  $\tilde{\mathcal{F}}|_{U \times_S \Theta_S}$  is a  $\mathrm{GL}(V)$ -bundle on  $U \times_S \Theta_S$  and  $\sigma$  induces a  $G$ -reduction of structure group of the restriction  $\tilde{\mathcal{F}}|_{U \times_S 1_S}$ .

**Definition 4.34.** *We say that  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  as above is a filtration of  $(\mathcal{F}, \sigma)$  if the  $G$ -reduction of structure group (uniquely) extends to  $U \times_S \Theta_S$ .*

The uniqueness of the extension shows that this is a property, not a structure. An argument similar to Proposition 3.12 implies that such a filtration of  $(\mathcal{F}, \sigma)$  is equivalent to a morphism  $\Theta_S \rightarrow \mathrm{Bun}_\rho(X)$  such that the restriction to  $1_S$  recovers  $(\mathcal{F}, \sigma)$ .

**Definition 4.35.** *Let  $S$  be a  $k$ -scheme and let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_S$ . We say that a filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  of  $(\mathcal{F}, \sigma)$  is a relative leading term HN filtration if for all points  $s \in S$  the restriction  $(\mathcal{F}_m|_{X_s})_{m \in \mathbb{Z}}$  is the leading term HN filtration of  $(\mathcal{F}|_{X_s}, \sigma|_{X_s})$ .*

Suppose that  $S$  is a connected scheme of finite type over a field extension  $K \supset k$  and assume that  $G_K$  is split. In this case, every relative filtration comes from a relative weighted parabolic reduction. Indeed, let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_S$ , and let  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  be a relative filtration of  $(\mathcal{F}, \sigma)$ . Let  $\tilde{\mathcal{G}}$  denote the corresponding  $G$ -bundle on  $\Theta_K \times_K U$  for some big open subset  $U \subset X_S$ . We denote by  $\mathcal{G}$  the  $G$ -bundle on  $U$  corresponding to  $(\mathcal{F}, \sigma)$  (i.e. the restriction  $\tilde{\mathcal{G}}|_{1 \times_S}$ ). Since  $X$  is geometrically connected and smooth, the fibers of  $X_S \rightarrow S$  are integral, hence the fibers of  $U \rightarrow S$  are also integral. Since  $S$  is connected, we conclude that  $U$  is a connected scheme of finite type over  $K$ . Proposition 3.14 shows that  $\tilde{\mathcal{G}}$  arises from applying the Rees construction to a weighted parabolic reduction  $(\lambda, \mathcal{G}_\lambda)$  of the  $G$ -bundle  $\mathcal{G}$  on  $U$ . We call  $(\lambda, \mathcal{G}_\lambda)$  a relative weighted parabolic reduction of  $(\mathcal{F}, \sigma)$  and note that the filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  is uniquely determined by  $(\lambda, \mathcal{G}_\lambda)$ . Then we say that the filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  comes from the relative weighted parabolic reduction  $(\lambda, \mathcal{G}_\lambda)$ .

Let  $S$  be a  $k$ -scheme and  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_S$ . Let  $H \rightarrow S$  be a locally closed stratification of  $S$ . This means that  $H$  is the disjoint union of locally closed subschemes of  $S$  and the morphism  $H \rightarrow S$  is surjective.

**Definition 4.36.** *We say that the  $S$ -scheme  $H$  is universal for leading term HN filtrations of  $(\mathcal{F}, \sigma)$  if it represents the functor that classifies pairs  $(T, [(\mathcal{F}_m)_{m \in \mathbb{Z}}])$ , where  $T$  is a  $S$ -scheme and  $[(\mathcal{F}_m)_{m \in \mathbb{Z}}]$  is an equivalence class of relative leading term HN filtrations of  $(\mathcal{F}|_{X_T}, \sigma|_{X_T})$  up to rational scaling the weights.*

The fact that the numerical invariant  $\nu$  induces a  $\Theta$ -stratification implies the following property of the leading term HN filtration in families, called a “schematic stratification” in [Nit11, GN14, GN18].

**Proposition 4.37.** *Relative leading term HN filtrations induce a stratification of  $\mathrm{Bun}_\rho(X)$  by locally closed substacks. Equivalently, for any  $k$ -scheme  $S$  and every  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  on  $X_S$  there is a unique locally closed stratification  $H \rightarrow S$  that is universal for leading term HN filtrations of  $(\mathcal{F}, \sigma)$ .*

*If, in addition,  $S$  is locally of finite type over a field extension  $K \supset k$  such that  $G_K$  is split, then the restriction of the universal relative leading term HN filtration<sup>4</sup> to each connected component of the stratification  $H$  comes from a relative weighted parabolic reduction.  $\square$*

## 5. COMPLETENESS OF THE STACK $\mathrm{Bun}_\rho(X)$

In this section we prove that the stack  $\mathrm{Bun}_\rho(X)$  satisfies the existence part of the valuative criterion for properness [Sta21, Tag 0CLK] in the case of complete discrete valuation rings (see Theorem 5.7).

The ideas that we use in this section (purely ramified extensions, Bruhat-Tits buildings) already appear in [BS02, Section 6] (see [BP03] for positive characteristic). Besides, V. Balaji has also considered extensions of families of principal bundles in higher dimensions, in the context of Donaldson-Uhlenbeck compactification [Bal07]. The case of positive characteristic for curves has also been studied by J. Heinloth [Hei08b, Hei10]. For torsion free sheaves, the valuative criteria was proved by S. Langton [Lan75].

Let  $R$  be a complete discrete valuation  $k$ -algebra. We set  $K = \mathrm{Frac}(R)$ , and choose a uniformizer  $\pi$ . By the Cohen structure theorem,  $R$  is isomorphic to the power series ring  $\kappa[[\pi]]$ , where  $\kappa$  is the residue field. Consider the base change  $X_R$ . Let  $\eta_\kappa$  denote the generic point of the special fiber  $X_\kappa$ . Similarly, let  $\eta_K$  be the generic point of the generic fiber  $X_K$ . Note that the local  $\mathcal{O}_{X_R, \eta_\kappa}$  is a discrete valuation ring with generic point  $\eta_K$  and special point  $\eta_\kappa$ .

The following is a local characterization of when a  $\rho$ -sheaf extends.

**Lemma 5.1.** *Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf over  $X_K$ . Let  $\mathcal{P}$  be the corresponding  $G$ -bundle over the generic fiber  $\eta_K$ . The following statements are equivalent:*

- (1) *There is a  $\rho$ -sheaf on  $X_R$  that restricts to  $(\mathcal{F}, \sigma)$  over  $X_K$ .*
- (2) *There is a  $G$ -bundle on  $\mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa})$  that restricts to  $\mathcal{P}$  over  $\eta_K$ .*

*Proof.* (1)  $\implies$  (2). Suppose that  $(\tilde{\mathcal{F}}, \zeta)$  is a  $\rho$ -sheaf on  $X_R$  that extends  $(\mathcal{F}, \sigma)$ . Let  $U$  be a big open subset of  $X_R$  where  $\tilde{\mathcal{F}}$  is locally-free. Let  $\tilde{\mathcal{P}}$  be the  $G$ -bundle on  $U$  corresponding to the restriction of  $(\tilde{\mathcal{F}}, \zeta)$ . Then  $\tilde{\mathcal{P}}|_{\mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa})}$  is a  $G$ -bundle on  $\mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa})$  that extends  $\mathcal{P}$ .

(2)  $\implies$  (1). Suppose that there is a  $G$ -bundle  $\tilde{\mathcal{P}}$  on  $\mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa})$  that extends  $\mathcal{P}$ . Let  $\mathcal{A}$  be the corresponding  $\mathrm{GL}(V)$ -bundle on  $\mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa})$  obtained via extension of structure group using  $\rho$ .  $\mathcal{A}$  is a vector bundle of rank  $r$  on  $\mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa})$ . By construction we have a canonical identification  $\mathcal{A}|_{\eta_K} \xrightarrow{\sim} \mathcal{F}|_{\eta_K}$ . Now [Lan75, Prop. 6] implies that there is a (unique) family of torsion-free sheaves  $\tilde{\mathcal{F}}$  on  $X_R$  along with isomorphisms  $\tilde{\mathcal{F}}|_{X_K} \xrightarrow{\sim} \mathcal{F}$  and  $\tilde{\mathcal{F}}|_{\mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa})} \xrightarrow{\sim} \mathcal{A}$  that are compatible with the identification  $\mathcal{A}|_{\eta_K} \xrightarrow{\sim} \mathcal{F}|_{\eta_K}$ .

We have a section  $s : \mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa}) \rightarrow \mathrm{Red}_G(\tilde{\mathcal{F}})|_{\mathrm{Spec}(\mathcal{O}_{X_R, \eta_\kappa})}$  corresponding to the  $G$ -reduction  $\tilde{\mathcal{P}}$ . It identifies over the point  $\eta_K$  with the section  $\sigma : X_K \rightarrow \mathrm{Red}_G(\mathcal{F})$ . Since

<sup>4</sup>More precisely, we choose a representative up to scaling.

$\text{Red}_G(\tilde{\mathcal{F}})$  is of finite type over  $X_R$ , we can spread the compatible pair  $s$  and  $\sigma$  to a section

$$\tilde{\sigma} : X_R \setminus Z \rightarrow \text{Red}_G(\tilde{\mathcal{F}})$$

defined away from a proper closed subscheme  $Z \subsetneq X_\kappa$  of the special fiber  $X_\kappa$ . Note that every point  $z \in Z$  has codimension at least 2 in  $X_R$ . Since  $X_R$  is smooth over the regular scheme  $\text{Spec}(R)$ , it is itself regular [Sta21, Tag 07NF]. It follows that for every  $z \in Z$ , the local ring  $\mathcal{O}_{X_R, z}$  has depth at least 2. By the proof of Lemma 2.2 (c), the section  $\tilde{\sigma}$  on  $X_R \setminus Z$  uniquely extends to a section  $\zeta : X_R \rightarrow \text{Red}_G(\tilde{\mathcal{F}})$  defined over the whole  $X_R$ . The  $\rho$ -sheaf  $(\tilde{\mathcal{F}}, \zeta)$  restricts to  $(\mathcal{F}, \sigma)$ , as desired.  $\square$

We would like to show that the extension result in Lemma 5.1 (2) is true after maybe passing to a finite extension  $R' \supset R$  of the discrete valuation ring  $R$ . In order to do so, we prove the following general result which could be of independent interest. <sup>5</sup>

**Proposition 5.2.** *Let  $\kappa$  be a perfect field. Let  $G$  be a quasi-split connected reductive group over  $\kappa$ . Suppose that the two following conditions are satisfied:*

- (1) *The maximal central torus  $Z \subset G$  is split over  $\kappa$ .*
- (2) *Let  $H$  denote the kernel of the simply-connected cover  $\widetilde{G}/Z \rightarrow G/Z$  of the semisimple group  $G/Z$ . The characteristic of  $\kappa$  does not divide the order of  $H$ .*

*Set  $\kappa[[\pi]]$  to be the formal power series ring. Let  $\mathcal{P}$  be a  $G$ -bundle on the fraction field  $\text{Spec}(\kappa((\pi)))$ . Then, there exists some  $m \geq 1$  such that the pullback of  $\mathcal{P}$  to  $\kappa((\pi^{\frac{1}{m}}))$  extends to a  $G$  bundle on  $\text{Spec}(\kappa[[\pi^{\frac{1}{m}}]])$ .*

*Proof.* Define  $K := \kappa((\pi))$  and  $\mathcal{O}_K = \kappa[[\pi]]$ . We fix the choice of a separable closure  $K_s$  of  $K$ . Set  $K_{unr}$  to be the maximal unramified extension of  $K$  inside  $K_s$ . We will denote by  $\mathcal{O}_{K_{unr}}$  the ring of integers of  $K_{unr}$ . Let  $\Gamma_K := \text{Gal}(K_s/K)$  be the absolute Galois group of  $K$ . We denote by  $\Gamma_{unr} := \text{Gal}(K_{unr}/K)$  the unramified Galois group.

We denote by  $H^1(K, G)$  (resp.  $H^1(\mathcal{O}_K, G)$ ) the pointed set of  $G$ -bundles on  $\text{Spec}(K)$  (resp.  $\text{Spec}(\mathcal{O}_K)$ ). We have a natural restriction morphism  $H^1(\mathcal{O}_K, G) \rightarrow H^1(K, G)$ . We are given an element  $\mathcal{P} \in H^1(K, G)$ . Our goal is to show that, after maybe replacing  $K$  with an extension  $K(\pi^{\frac{1}{m}})$ , there exists some  $\tilde{\mathcal{P}} \in H^1(\mathcal{O}_K, G)$  that restricts to  $\mathcal{P}$ .

Since  $G$  is smooth, the base-change  $\mathcal{P}_{K_s}$  is trivializable over  $\text{Spec}(K_s)$ . The descent data for  $\mathcal{P}$  yield a class in the nonabelian Galois cohomology set  $H^1(\Gamma_K, G(K_s))$ . Since  $G$  is connected and  $\kappa$  is perfect, [BT87, 3.14 (3)] shows that  $H^1(\Gamma_K, G(K_s)) = H^1(\Gamma_{unr}, K_{unr})$ . So the torsor  $\mathcal{P}$  comes from an unramified cohomology class  $\bar{c} \in H^1(\Gamma_{unr}, K_{unr})$ . Notice that there is a natural action of  $\Gamma_{unr}$  on  $G(\mathcal{O}_{K_{unr}})$ . There is a map  $H^1(\Gamma_{unr}, G(\mathcal{O}_{K_{unr}})) \rightarrow H^1(\Gamma_{unr}, G(K_{unr}))$  induced by the inclusion of  $\Gamma_{unr}$ -groups  $G(\mathcal{O}_{K_{unr}}) \hookrightarrow G(K_{unr})$ . Our goal is to show that, after maybe replacing  $K$  with  $K(\pi^{\frac{1}{m}})$ , we have that  $\bar{c}$  comes from a class in  $H^1(\Gamma_{unr}, G(\mathcal{O}_{K_{unr}}))$ .

Let us first suppose that  $G$  is semisimple. There is a finite unramified Galois extension  $L \supset K$  contained in  $K_{unr}$  such that  $\mathcal{P}_L$  is trivializable. Let  $\mathcal{O}_L$  denote the ring of integers of  $L$ . Set  $\Gamma_{L/K} := \text{Gal}(L/K)$ . This is a finite quotient of  $\Gamma_{unr}$ . The descent data for  $\mathcal{P}$  yield a class  $\bar{c}$  in the nonabelian Galois cohomology set  $H^1(\Gamma_{L/K}, G(L))$ .

Since  $G_K$  is quasi-split, it follows that  $G_K$  is residually quasi-split [BT87, 2.2]. Therefore, we can use the description of unramified Galois cohomology given in [BT87], we just need to be a bit careful if  $G$  is not simply-connected. We will denote by  $G^{00}$  the subgroup of  $G(K_{unr})$  generated by the  $K_{unr}$ -points of all parahoric subgroups of  $G_{K_{unr}}$ . By [BT87, 1.8],  $G^{00}$  admits the following description. Let  $T$  denote a maximal torus of  $G$  over  $\kappa$ . We can

<sup>5</sup>V. Balaji pointed out to us that Proposition 5.2 could also be deduced from [RR84, Proposition 3.12].

think of  $T_K$  as the centralizer of a maximal split torus of  $G_K$ , since  $G_K$  is quasi-split. Let  $j : \tilde{G} \rightarrow G$  be a simply connected cover of  $G$ . Then, we have  $G^{00} = T(\mathcal{O}_{K_{unr}}) \cdot j(\tilde{G}(K_{unr}))$ .

Choose a cocycle  $c \in Z^1(\Gamma_{L/K}, G(L))$  representing  $\bar{c}$ . We will show that, after maybe replacing  $K$  with some ramified extension  $K(\pi^{\frac{1}{m}})$ , the cocycle  $c$  takes values in the subgroup  $G^{00}$ . Let  $H$  denote the finite kernel of the isogeny  $j : \tilde{G} \rightarrow G$ . Note that  $H$  is a finite product of groups  $\mu_j$  of  $j^{\text{th}}$  roots of unity. By assumption (2), the characteristic of  $\kappa$  does not divide any of the orders of the  $\mu_j$  appearing in  $H$ . After maybe enlarging the unramified extension  $L$ , we can assume that all of these roots of unity are contained in  $L$ . So  $H_L$  is a finite constant étale group scheme. Since  $H$  is smooth, the following sequence of groups is exact.

$$1 \rightarrow H(K_s) \rightarrow \tilde{G}(K_s) \rightarrow G(K_s) \rightarrow 1$$

We can take  $\text{Gal}(K_s/L)$  invariants to get the following exact sequence of pointed sets

$$1 \rightarrow H(L) \rightarrow \tilde{G}(L) \rightarrow G(L) \rightarrow H^1(\text{Gal}(K_s/L), H(K_s))$$

Hence the obstructions to lifting to  $\tilde{G}(L)$  the finitely many elements in the image of the cocycle  $c \in Z^1(\Gamma_{L/K}, G(L))$  consist of finitely many cohomology classes in  $H^1(\text{Gal}(K_s/L), H(K_s))$ . By Kummer theory, these are killed after taking some cyclic extensions of  $L$ . After further enlarging the unramified extension  $L$ , we can assume that these cyclic extensions are totally ramified. Such cyclic extension must be of the form  $L(\pi^{\frac{1}{m}})$  for some uniformizer  $\pi_L$  of  $L$ . Such uniformizer must be of the form  $\pi_L = u\pi$ , where  $u \in \mathcal{O}_L^\times$ . After replacing  $L$  with  $L(u^{\frac{1}{m}})$ , we can assume that the obstructions in  $H^1(\text{Gal}(K_s/L), H(K_s))$  vanish on the extension  $L(\pi^{\frac{1}{m}})$  for some  $m \geq 1$ . So after replacing  $K$  with some sufficiently big ramified extension  $K(\pi^{\frac{1}{m}})$ , we can assume that  $c$  takes values in the group  $j(\tilde{G}(L)) \subset G^{00}$ . Hence  $\bar{c}$  is a class in the Galois cohomology set  $H^1(\Gamma_{L/K}, G^{00} \cap G(L))$ . Since  $G_K$  is residually-quasisplit, the result [BT87, 3.15] and the construction of [BT87, 3.4 Lem. 2] show that  $\bar{c}$  is represented by a cocycle  $z \in Z^1(\Gamma_{L/K}, P(\mathcal{O}_L))$  taking values in the group of  $\mathcal{O}_L$ -points of a parahoric  $P$  of  $G_K$  defined over  $K$ . We can without loss of generality take  $P$  to be in the apartment corresponding to a maximal split torus  $S$  inside  $T_K$ , because  $G(K)$  acts transitively on apartments. The argument in [Lar95, Lem. 2.4] (see also [Ser96, Proposition 8] and [Gil94, Lemma I.1.3.2]) shows that there exists some  $m \geq 1$  and some  $K$ -rational cocharacter  $\lambda : \mathbb{G}_m \rightarrow S$  such that  $\lambda(\pi^{\frac{1}{m}}) \in G(K[\pi^{\frac{1}{m}}])$  conjugates the base-changed parahoric  $P_{\mathcal{O}_{K(\pi^{\frac{1}{m}})}}$  into the hyperspecial parahoric  $G_{\mathcal{O}_{K(\pi^{\frac{1}{m}})}}$ . After replacing  $K$  with  $K(\pi^{\frac{1}{m}})$ , we can use the element  $\lambda(\pi^{\frac{1}{m}})$  to conjugate the cocycle  $z$  into a cocycle  $z' \in Z^1(\Gamma_{L/K}, G(\mathcal{O}_L))$ . This shows that  $\bar{c}$  comes from a cohomology class in  $H^1(\Gamma_{L/K}, G(\mathcal{O}_L))$ , and thus concludes the proof in the case when  $G$  is semisimple.

We now deal with the general case. Let  $G$  be an arbitrary quasi-split connected reductive group over  $\kappa$ . Let  $Z$  denote the maximal central torus of  $G$ . We have a short exact sequence of algebraic groups

$$1 \rightarrow Z \rightarrow G \rightarrow \bar{G} \rightarrow 1$$

Here  $\bar{G} = G/Z$  is a quasi-split semisimple group over  $\kappa$  satisfying condition (2) of the proposition. By assumption  $Z$  is split, and so it is isomorphic to a finite product  $(\mathbb{G}_m)^h$  of copies of  $\mathbb{G}_m$ . We have an exact sequence

$$1 \rightarrow Z(K_{unr}) \rightarrow G(K_{unr}) \rightarrow \bar{G}(K_{unr}) \rightarrow H^1(\Gamma_{K_s/K_{unr}}, Z(K_s))$$

Hilbert's Theorem 90 implies that  $H^1(\Gamma_{K_s/K_{unr}}, Z(K_s)) = 0$ , and so we have a short exact sequence of  $\Gamma_{unr}$ -groups

$$1 \rightarrow Z(K_{unr}) \rightarrow G(K_{unr}) \rightarrow \bar{G}(K_{unr}) \rightarrow 1$$

The same thing holds for  $\mathcal{O}_{K_{unr}}$ -points. After taking  $\Gamma_{unr}$ -invariants and replacing  $Z$  with  $(\mathbb{G}_m)^h$ , we get a diagram of pointed sets

$$\begin{array}{ccccccc}
H^1(\Gamma_{unr}, \mathcal{O}_{K_{unr}}^\times)^{\oplus h} & \rightarrow & H^1(\Gamma_{unr}, G(\mathcal{O}_{K_{unr}})) & \rightarrow & H^1(\Gamma_{unr}, \overline{G}(\mathcal{O}_{K_{unr}})) & \rightarrow & H^2(\Gamma_{unr}, \mathcal{O}_{K_{unr}}^\times)^{\oplus h} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
H^1(\Gamma_{unr}, K_{unr}^\times)^{\oplus h} & \rightarrow & H^1(\Gamma_{unr}, G(K_{unr})) & \rightarrow & H^1(\Gamma_{unr}, \overline{G}(K_{unr})) & \rightarrow & H^2(\Gamma_{unr}, K_{unr}^\times)^{\oplus h}
\end{array}$$

After an application of Hilbert's Theorem 90 to the first column, we get

$$\begin{array}{ccccccc}
0 & \longrightarrow & H^1(\Gamma_{unr}, G(\mathcal{O}_{K_{unr}})) & \longrightarrow & H^1(\Gamma_{unr}, \overline{G}(\mathcal{O}_{K_{unr}})) & \xrightarrow{f_1} & H^2(\Gamma_{unr}, \mathcal{O}_{K_{unr}}^\times)^{\oplus h} \\
& & \downarrow & & \downarrow & & \downarrow g \\
0 & \longrightarrow & H^1(\Gamma_{unr}, G(K_{unr})) & \xrightarrow{f_2} & H^1(\Gamma_{unr}, \overline{G}(K_{unr})) & \xrightarrow{f_3} & H^2(\Gamma_{unr}, K_{unr}^\times)^{\oplus h}
\end{array}$$

where the two horizontal rows are exact.

Let  $\bar{c} \in H^1(\Gamma_{unr}, G(K_{unr}))$  be our unramified Galois cohomology class coming from  $\mathcal{P}$ . Our goal is to lift  $\bar{c}$  to a class in  $H^1(\Gamma_{unr}, G(\mathcal{O}_{K_{unr}}))$ . Consider the image  $f_2(\bar{c}) \in H^1(\Gamma_{unr}, \overline{G}(K_{unr}))$ . By the proof of the semisimple case, after maybe replacing  $K$  with an extension  $K(\pi^{\frac{1}{m}})$  we can lift  $f_2(\bar{c})$  to a cohomology class  $\bar{d} \in H^1(\Gamma_{unr}, \overline{G}(\mathcal{O}_{K_{unr}}))$ . We now want to lift  $\bar{d}$  to a class in  $H^1(\Gamma_{unr}, G(\mathcal{O}_{K_{unr}}))$  whose image in  $H^1(\Gamma_{unr}, G(K_{unr}))$  is the original  $\bar{c}$ . For this it suffices to show that  $f_1(\bar{d}) = 0$ .

We have  $g(f_1(\bar{d})) = f_3(f_2(\bar{c})) = 0$ . Therefore, it suffices to show that  $g$  is injective. The morphism  $g$  is the direct sum of  $h$  copies of the morphism  $H^2(\Gamma_{unr}, \mathcal{O}_{K_{unr}}^\times) \rightarrow H^2(\Gamma_{unr}, K_{unr}^\times)$  coming from the inclusion of  $\Gamma_{unr}$ -groups  $\mathcal{O}_{K_{unr}}^\times \hookrightarrow K_{unr}^\times$ . These groups fit into a short exact sequence

$$1 \rightarrow \mathcal{O}_{K_{unr}}^\times \rightarrow K_{unr}^\times \rightarrow \mathbb{Z} \rightarrow 1$$

Here  $\Gamma_{unr}$  acts trivially on  $\mathbb{Z}$ . We get an exact sequence of abelian groups

$$(13) \quad H^1(\Gamma_{unr}, \mathbb{Z}) \rightarrow H^2(\Gamma_{unr}, \mathcal{O}_{K_{unr}}^\times) \rightarrow H^2(\Gamma_{unr}, K_{unr}^\times)$$

Since  $\Gamma_{unr}$  acts trivially on  $\mathbb{Z}$ , we have that  $H^1(\Gamma_{unr}, \mathbb{Z})$  is the group of continuous homomorphisms from  $\Gamma_{unr}$  to  $\mathbb{Z}$ . Since  $\Gamma_{unr}$  is profinite, the image of such a homomorphism is finite. Therefore it must be 0, because  $\mathbb{Z}$  is torsion-free. We conclude that  $H^1(\Gamma_{unr}, \mathbb{Z}) = 0$ . Hence (13) shows that  $g$  is injective, as desired.  $\square$

**Remark 5.3.** *The argument that allows us to go from semisimple to reductive groups is similar to the argument of  $G$ . Faltings in [Fal93, Theorem II.4].*

**Remark 5.4.** *It follows from the argument that  $m \geq 1$  can be chosen independently of  $\mathcal{P}$ . One can give a uniform upper bound that depends only on the root data of  $G/Z$ .*

**Remark 5.5.** *The argument above works verbatim for a general complete discrete valuation ring  $R$  with perfect residue field  $\kappa$  (i.e. mixed characteristic). In that case we can take  $G$  to be a reductive group over  $R$  with connected fibers such that the generic fiber  $G_K$  is residually quasi-split and the analogous conditions (1) and (2) are satisfied.*

In order to construct our generic  $G$ -bundle extension, we will make use of formal gluing, also known as Beauville-Laszlo gluing [BL95]. Suppose that  $S \rightarrow \widehat{S}$  is a faithfully flat morphism of rings. Let  $f \in S$  be a nonzerodivisor. We denote by  $\text{Glue}_{\text{Aff}}$  the category consisting of triples  $(\widehat{A}, A_f, \varphi)$ , where

- (1)  $\widehat{A}$  is an affine scheme over  $\text{Spec}(\widehat{S})$ .
- (2)  $A_f$  is an affine scheme over  $\text{Spec}(S_f)$ .
- (3)  $\varphi$  is an isomorphism  $\varphi : \widehat{A}|_{\text{Spec}(\widehat{S}_f)} \xrightarrow{\sim} A_f|_{\text{Spec}(\widehat{S}_f)}$ .

A morphism  $(\widehat{A}, A_f, \varphi) \rightarrow (\widehat{B}, B_f, \psi)$  consists of a pair of morphisms  $(g_1 : \widehat{A} \rightarrow \widehat{B}, g_2 : A_f \rightarrow B_f)$  such that their base-changes to  $\mathrm{Spec}(\widehat{S}_f)$  fit into a commutative diagram

$$\begin{array}{ccc} \widehat{A}|_{\mathrm{Spec}(\widehat{S}_f)} & \xrightarrow{g_1|_{\mathrm{Spec}(\widehat{S}_f)}} & A_f|_{\mathrm{Spec}(\widehat{S}_f)} \\ \downarrow \varphi & & \downarrow \psi \\ \widehat{B}|_{\mathrm{Spec}(\widehat{S}_f)} & \xrightarrow{g_2|_{\mathrm{Spec}(\widehat{S}_f)}} & B_f|_{\mathrm{Spec}(\widehat{S}_f)} \end{array}$$

Given an affine scheme  $A$  over  $\mathrm{Spec}(S)$ , we can form the two pullbacks  $A_{\mathrm{Spec}(\widehat{S})}$  and  $A_{\mathrm{Spec}(S_f)}$ . Note that there is a canonical isomorphism  $\varphi_{A,can} : \left(A_{\mathrm{Spec}(\widehat{S})}\right)|_{\mathrm{Spec}(\widehat{S}_f)} \xrightarrow{\sim} \left(A_{\mathrm{Spec}(S_f)}\right)|_{\mathrm{Spec}(\widehat{S}_f)}$ . This defines a functor

$$\mathrm{Aff}_{can} : \mathrm{Aff}_{\mathrm{Spec}(S)} \rightarrow \mathrm{Glue}_{\mathrm{Aff}}, \quad A \mapsto (A_{\mathrm{Spec}(\widehat{S})}, A_{\mathrm{Spec}(S_f)}, \varphi_{A,can})$$

Let  $G$  be an affine flat group scheme over  $\mathrm{Spec}(S)$ . We can similarly define a category  $\mathrm{Glue}_{G\mathrm{Bun}}$  by replacing the tuple of affine schemes with  $G$ -bundles and requiring that  $\varphi$  is a morphism of  $G$ -bundles. Denote by  $BG(S)$  the groupoid of  $G$ -bundles over  $\mathrm{Spec}(S)$ . The functor  $\mathrm{Aff}_{can}$  induces another functor

$$G\mathrm{Bun}_{can} : BG(S) \rightarrow \mathrm{Glue}_{G\mathrm{Bun}}, \quad \mathcal{P} \mapsto (\mathcal{P}|_{\mathrm{Spec}(\widehat{S})}, \mathcal{P}|_{\mathrm{Spec}(S_f)}, \varphi_{\mathcal{P},can})$$

The following lemma is usually known in the literature as Beauville-Laszlo gluing.

**Lemma 5.6.** *Suppose that the faithfully flat morphism  $S \rightarrow \widehat{S}$  induces an isomorphism  $S/(f \cdot S) \xrightarrow{\sim} \widehat{S}/(f \cdot \widehat{S})$ . Then both of the functors  $\mathrm{Aff}_{can}$  and  $G\mathrm{Bun}_{can}$  described above are equivalences of categories.*

*Proof.* We shall use [Sta21, Tag 05ES]. Since the gluing of modules preserves the tensor product structure on modules (cf. [Sta21, Tag 05EU]), the equivalence in [Sta21, Tag 05ES] induces an equivalence of monoid objects (i.e. algebras). By applying the  $\mathrm{Spec}$  functor we deduce that  $\mathrm{Aff}_{can}$  is an equivalence, and that it is compatible with fiber products relative to the base. This implies that  $\mathrm{Aff}_{\widehat{S}}$  induces an equivalence of categories of affine schemes equipped with a right  $G$ -action<sup>6</sup>. If  $\mathcal{P}$  is a scheme equipped with a right  $G$ -action, then by descent for the flat cover  $\mathrm{Spec}(\widehat{S}) \rightarrow \mathrm{Spec}(S)$  we know that  $\mathcal{P}$  is a  $G$ -bundle if and only if  $\mathcal{P}|_{\mathrm{Spec}(\widehat{S})}$  is a  $G$ -bundle. This shows that the equivalence on affine schemes equipped with a right  $G$ -action restricts to an equivalence  $G\mathrm{Bun}_{can}$  on the subcategories of  $G$ -bundles.  $\square$

Now we are ready to prove the main theorem of this section.

**Theorem 5.7.** *The structure morphism  $\mathrm{Bun}_{\rho}(X) \rightarrow \mathrm{Spec}(k)$  satisfies the existence part of the valuative criterion for properness [Sta21, Tag 0CLK] in the case of complete discrete valuation rings. In other words, for all complete discrete valuation rings  $R$  with fraction field  $K$  and all diagrams*

$$\begin{array}{ccc} \mathrm{Spec}(K) & \xrightarrow{f} & \mathrm{Bun}_{\rho}(X) \\ \downarrow & & \downarrow \\ \mathrm{Spec}(R) & \longrightarrow & \mathrm{Spec}(k) \end{array}$$

*there is a finite extension  $R' \supset R$  of complete discrete valuation rings and a morphism  $\mathrm{Spec}(R') \rightarrow \mathrm{Bun}_{\rho}(X)$  fitting into a 2-commutative diagram*

<sup>6</sup>One defines the corresponding gluing category similarly as in the case for  $G$ -bundles above.



$$\begin{array}{ccc}
\mathrm{Spec}(K') & \xrightarrow{f_{K'}} & \mathrm{Bun}_\rho(X) \\
\downarrow & \nearrow & \downarrow \\
\mathrm{Spec}(R') & \longrightarrow & \mathrm{Spec}(k)
\end{array}$$

*Proof.* We will use the notation from Lemma 5.1. By Lemma 5.1, it suffices to find a finite extension  $R' \supset R$  of discrete valuation rings such that the base-change  $\mathcal{P}_{K'}$  on the generic point  $\eta_{K'}$  of  $\mathrm{Spec}(\mathcal{O}_{X_{R'}, \eta_{K'}})$  admits an extension to a  $G$ -bundle  $\widehat{\mathcal{P}}$  on the whole  $\mathrm{Spec}(\mathcal{O}_{X_{R'}, \eta_{K'}})$ .

We can replace  $k$  with  $\kappa$  in order to assume that  $\kappa$  is the ground field. By the Cohen structure theorem, we have  $R = \kappa[[\pi]]$ . After replacing  $\kappa$  with a finite extension  $\kappa'$ , we can assume that  $G$  is split. Set  $S = \mathcal{O}_{X_R, \eta_\kappa}$ . Note that  $S \supset R$  is a discrete valuation ring with uniformizer  $\pi \in R$ . Let  $\widehat{S}$  denote the completion  $\widehat{\mathcal{O}}_{X_R, \eta_\kappa}$ . Let  $\widehat{\mathcal{P}}$  denote the base change of  $\mathcal{P}$  to  $\mathrm{Spec}(\mathrm{Frac}(\widehat{S}))$ . We have that  $\widehat{S} = \kappa_S[[\pi]]$ , where  $\kappa_S \supset \kappa$  is the residue field of  $S$  (equivalently  $\kappa_S$  is the fraction field of the special fiber  $X_\kappa$ ).

By formal gluing (Lemma 5.6 applied to  $S \rightarrow \widehat{S}$  and  $f = \pi$ ), extending  $\mathcal{P}$  to  $\mathrm{Spec}(S)$  is equivalent to extending  $\widehat{\mathcal{P}}$  to  $\mathrm{Spec}(\widehat{S})$ . Indeed, such an extension  $(\widehat{\mathcal{P}})^\sim$  to  $\mathrm{Spec}(\widehat{S})$  yields a triple  $(\mathcal{P}, (\widehat{\mathcal{P}})^\sim, \varphi)$ , which by Lemma 5.6 glues to an extension of  $\mathcal{P}$  over  $\mathrm{Spec}(S)$ . Hence we may replace  $S$  with  $\widehat{S}$ .

Note that we are allowed to replace  $\kappa[[\pi]]$  with a finite totally ramified extension  $\kappa[[\pi^{\frac{1}{m}}]]$  for some  $m \geq 1$ . This replaces  $\widehat{S}$  with  $\kappa_S[[\pi^{\frac{1}{m}}]]$ . The theorem is therefore implied by Proposition 5.2 (by taking  $\kappa = \kappa_S$  in that proposition).  $\square$

Since the semistable locus  $\mathrm{Bun}_\rho(X)^{ss}$  comes from a  $\Theta$ -stratification, we automatically have semistable reduction for  $\rho$ -sheaves by [AHLH19, §6]. We record this result as a separate theorem.

**Theorem 5.8** (Semistable Reduction). *The substack  $\mathrm{Bun}_\rho(X)^{ss}$  of Gieseker semistable points satisfies the existence part of the valuative criterion for properness.*  $\square$

The arguments in this section yield an interesting result in the case of curves. Suppose that  $X$  is a smooth projective curve. Then a  $\rho$ -sheaf is the same thing as a  $G$ -bundle. The arguments above imply the following result about extending  $G$ -bundles.

**Corollary 5.9.** *Let  $X$  be a smooth projective geometrically connected curve over a field  $\kappa$  of arbitrary characteristic. Let  $G$  be a quasi-split connected reductive group over  $\kappa$  satisfying the following two conditions:*

- (1) *The maximal central torus of  $G$  is split over  $\kappa$ .*
- (2) *Let  $H$  denote the kernel of the simply-connected cover  $[\widetilde{G}, \widetilde{G}] \rightarrow [G, G]$  of the derived subgroup of  $G$ . Then, the characteristic of  $\kappa$  does not divide the order of  $H$ .*

*Let  $R$  be a complete discrete valuation ring over  $\kappa$  with fraction field  $K$ . Suppose that the residue field of  $R$  is perfect. Let  $\mathcal{P}$  be a  $G$ -bundle on the generic fiber  $X_K$ . Then, after maybe taking a root of the uniformizer of  $R$ , we can extend  $\mathcal{P}$  to a  $G$ -bundle on  $X_R$ .*

*Proof.* This follows by applying the proof of Lemma 5.1, using formal gluing as in Theorem 5.7 and then applying Proposition 5.2.  $\square$

## 6. MODULI SPACE FOR THE SEMISTABLE LOCUS

**6.1. The degree of a  $\rho$ -sheaf.** Fix the choice of an algebraic closure  $\overline{k} \supset k$ . We denote by  $\Gamma_k$  the absolute Galois group  $\Gamma_k := \mathrm{Gal}(\overline{k}/k)$ . Let  $X^*(G_{\overline{k}})$  be the group of characters of

the base-change  $G_{\bar{k}}$ . Let  $X^*(G_{\bar{k}})^\vee$  denote the dual group  $\text{Hom}(X^*(G_{\bar{k}}), \mathbb{Z})$ . This group admits a natural continuous action of the profinite absolute Galois group  $\Gamma_k$ . We write  $\Lambda(G) := (X^*(G_{\bar{k}})^\vee)_{\Gamma_k}$  to denote the quotient group of  $\Gamma_k$ -coinvariants of  $(X^*(G_{\bar{k}})^\vee)$ . Note that  $\Lambda(G)$  does not depend on the choice of algebraic closure  $\bar{k}$ .

**Example 6.1.** *If the group  $G$  is split over  $k$ , then  $X^*(G_{\bar{k}})$  coincides with the group  $X^*(G)$  of characters defined over  $k$ . The action of the Galois group  $\Gamma_k$  on  $X^*(G)$  is trivial. Therefore  $\Lambda(G) = X^*(G)^\vee$ .*

**Example 6.2.** *If the group  $G$  is semisimple, then  $\Lambda(G) = 0$ .*

Let  $K \supset k$  be an arbitrary algebraically closed field extension and let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . Let  $U$  be any big open subset of  $X_K$  where  $\mathcal{F}$  is locally-free and denote by  $j$  be the open inclusion  $j : U \hookrightarrow X_K$ . If  $\mathcal{G}$  is the corresponding  $G_K$ -bundle on  $U$ , then for any character  $\chi \in X^*(G_K)$  we can form the associated line bundle  $\mathcal{G} \times^{G_K} \chi$  over  $U$ . Since  $X_K$  is smooth, the pushforward  $j_*(\mathcal{G} \times^{G_K} \chi)$  is a line bundle on  $X_K$  whose first Chern class is  $c_1(j_*(\mathcal{G} \times^{G_K} \chi))$ . Let  $H$  be the hyperplane class on  $X$  corresponding to the line bundle  $\mathcal{O}_X(1)$  and define the group homomorphism (c.f Lemma 4.25):

$$\psi : X^*(G_K) \rightarrow \mathbb{Z}, \quad \psi(\chi) := c_1(j_*(\mathcal{G} \times^{G_K} \chi)) \cdot H^{d-1}.$$

If we denote by  $L$  the algebraic closure of  $k$  in  $K$ , then base-change induces a canonical identification  $X^*(G_L) \xrightarrow{\sim} X^*(G_K)$ . Choose an isomorphism  $\bar{k} \xrightarrow{\sim} L$  over  $k$ . This yields an isomorphism of abelian groups  $X^*(G_{\bar{k}}) \xrightarrow{\sim} X^*(G_L)$  which identifies  $\psi$  with an element of  $X^*(G_{\bar{k}})^\vee$ . Since any two choices of isomorphisms  $\bar{k} \xrightarrow{\sim} L$  differ by an element of the Galois group  $\Gamma_k$ ,  $\psi$  yields a well-defined element of the quotient  $\Lambda(G)$  that is independent of the choice of  $\bar{k} \xrightarrow{\sim} L$ .

**Definition 6.3.** *Let  $K \supset k$  be an algebraically closed field extension. Let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . We define the degree  $\text{deg}(\mathcal{F}, \sigma)$  to be the element  $\vartheta \in \Lambda(G)$  constructed above.*

*For an arbitrary field extension  $K \supset k$ , we set  $\text{deg}(\mathcal{F}, \sigma)$  to be  $\text{deg}(\mathcal{F}|_{X_{\bar{K}}}, \sigma|_{X_{\bar{K}}})$  for any choice of an algebraically closed field extension  $\bar{K} \supset K$ .*

**Proposition 6.4.** *Let  $S$  be a  $k$ -scheme and let  $(\mathcal{F}, \sigma)$  be a  $\rho$ -sheaf on  $X_S$ . Let  $\text{deg} : S \rightarrow \Lambda(G)$  denote the function defined by  $\text{deg}(s) := \text{deg}(\mathcal{F}|_{X_s}, \sigma|_{X_s})$ . Then  $\text{deg}$  is locally constant on  $S$ .*

*Proof.* Since the stack  $\text{Bun}_\rho(X)$  is locally of finite type over  $k$ , we can reduce to the case when  $S$  is of finite type over  $k$ . Given that the notion of degree is well-behaved under taking field extensions, we can replace  $k$  with an algebraic closure to assume that the ground field is algebraically closed and, hence  $\Gamma_k$  is trivial.

Let  $U$  be a big open subset of  $X_S$  where  $\mathcal{F}$  is locally-free and denote by  $j : U \hookrightarrow X_S$  the open immersion. For each point  $s \in S$ , let  $j_s$  be the base-change  $j_s : U_s \hookrightarrow X_s$ . Denote by  $\mathcal{G}$  the corresponding  $G$ -bundle on  $U$ . Let  $\chi \in X^*(G)$  be a character of  $G$  and denote by  $\mathcal{L}$  the associated line bundle  $\mathcal{G} \times^G \chi$  on  $U$ . We want to show that the function  $f : S \rightarrow \mathbb{Z}$  given by

$$f(s) := c_1((j_s)_*(\mathcal{L}|_{U_s})) \cdot H^{d-1}$$

is locally constant on  $S$ .

We first claim that  $f$  is constructible on  $S$ . By Noetherian induction we need to prove that if  $S$  is integral then  $f$  is constant on an open subset of  $S$ . Assume that  $S$  is integral. Since  $k$  is perfect, there exists an open dense subset of  $S$  that is smooth over  $k$ . Therefore we can reduce to the case when  $S$  is a smooth variety. For each  $s \in S$ , consider the following Cartesian diagram

$$\begin{array}{ccc}
U_s & \longrightarrow & U \\
\downarrow j_s & & \downarrow j \\
X_s & \longrightarrow & X_S
\end{array}$$

We get an induced base-change morphism of quasicoherent sheaves  $\varphi : j_*(\mathcal{L})|_{X_s} \rightarrow (j_s)_*(\mathcal{L}|_{U_s})$  where the right-hand side is a line bundle. Since  $S$  is smooth over  $k$  and  $X_S \rightarrow S$  is a smooth morphism, we conclude that  $X_S$  is also smooth over  $k$ , so  $X_S$  is a regular scheme. The complement of the open subset  $U$  has codimension  $\geq 2$  inside  $X_S$ , therefore the pushforward  $j_*(\mathcal{L})$  is a line bundle. It follows that  $(j_*(\mathcal{L})|_{X_s})$  is a line bundle, and so  $\varphi$  is a morphism of line bundles on  $X_s$ . Note that the restriction of  $\varphi$  to the big open subset  $U_s \subset X_s$  is an isomorphism by construction. We conclude that  $\varphi$  must be an isomorphism of line bundles. This shows that we can equivalently write  $f$  as

$$f(s) = c_1((j_*\mathcal{L})|_{X_s}) \cdot H^{d-1}$$

Since  $j_*\mathcal{L}$  is a line bundle its first Chern class is locally constant in families, then  $f$  is constant on the variety  $S$ . This concludes the proof that  $f$  is constructible.

In order to show that  $f$  is locally constant, it suffices to check that  $f$  is constant when the base  $S$  is of the form  $\text{Spec}(R)$  for some discrete valuation ring  $R$ . In this case  $S$  is a regular integral scheme. Since  $X_S \rightarrow S$  is smooth, it follows that  $X_S$  is regular [Sta21, Tag 07NF] and we can apply the same argument as in the proof of constructibility to see that  $f$  is constant on  $S$ .  $\square$

As a consequence of Proposition 6.4, we have the following.

**Corollary 6.5.** *For every  $\vartheta \in \Lambda(G)$ , there exists an open and closed substack  $\text{Bun}_\rho(X)_\vartheta \subset \text{Bun}_\rho(X)$  parametrizing  $\rho$ -sheaves with degree  $\vartheta$ . Then*

$$\text{Bun}_\rho(X) = \bigsqcup_{\vartheta \in \Lambda(G)} \text{Bun}_\rho(X)_\vartheta$$

**6.2. Boundedness of the semistable locus.** Let  $P \in \mathbb{Q}[n]$  be a Hilbert polynomial and let  $\vartheta \in \Lambda(G)$ . We define  $\text{Bun}_\rho(X)_\vartheta^{ss,P} := \text{Bun}_\rho(X)^{ss,P} \cap \text{Bun}_\rho(X)_\vartheta$ . By Corollary 6.5 we get a decomposition of the semistable locus

$$\text{Bun}_\rho(X)^{ss,P} = \bigsqcup_{\vartheta \in \Lambda(G)} \text{Bun}_\rho(X)_\vartheta^{ss,P}$$

The next proposition shows that each piece  $\text{Bun}_\rho(X)_\vartheta^{ss,P}$  is bounded.

**Proposition 6.6.** *For all  $\vartheta \in \Lambda(G)$  and all  $P \in \mathbb{Q}[n]$ , the stack  $\text{Bun}_\rho(X)_\vartheta^{ss,P}$  is quasicompact.*

*Proof.* After replacing  $k$  with a field extension, we can assume that  $k$  is algebraically closed. Since the forgetful morphism  $\text{Forget} : \text{Bun}_\rho(X) \rightarrow \text{Coh}_r^{tf}(X)$  is of finite type, it is sufficient to show that the restriction  $\text{Forget}|_{\text{Bun}_\rho(X)_\vartheta^{ss,P}}$  factors through a quasicompact open substack of  $\text{Coh}_r^{tf}(X)$ . We shall show that the following set:

$$\mathfrak{S} = \left\{ \text{Forget}((\mathcal{F}, \sigma)) \mid (\mathcal{F}, \sigma) \in \text{Bun}_\rho(X)_\vartheta^{ss,P}(k) \right\}$$

is a bounded family of torsion-free sheaves.

Let  $(\mathcal{F}, \sigma) \in \text{Bun}_\rho(X)_\vartheta^{ss,P}(k)$ . We claim that there exists a uniform upper bound  $\mu_{max}$  (only depending on  $\vartheta$ ) such that for all torsion-free subsheaves  $\mathcal{V} \subset \mathcal{F}$  we have  $\hat{\mu}_{d-1}(\mathcal{V}) \leq \mu_{max}$ . Since the Hilbert polynomial  $P$  of  $\mathcal{F}$  is fixed, it will follow that the family  $\mathfrak{S}$  is bounded by [HL97, Thm. 3.3.7].

We are left to show the claim. Since  $G$  is a reductive group over  $k$  of characteristic 0,  $G$  is linearly reductive and the representation  $\rho : G \rightarrow \mathrm{GL}(V)$  decomposes into a direct sum of irreducible representations  $V = \bigoplus_{i=1}^q V_i$ . If we denote  $H := \prod_{i=1}^q \mathrm{GL}(V_i)$ , then  $\rho$  factors through a faithful homomorphism  $\varphi : G \rightarrow H$ . Since each representation  $V_i$  is irreducible, it follows that the image of  $\varphi$  is not contained in any proper parabolic subgroup of  $H$ .

Let  $U$  be a big open subset where  $\mathcal{F}$  is locally-free and let  $\mathcal{G}$  be the corresponding  $G$ -bundle on  $U$  which, by Proposition 10.4, is slope semistable in the sense of Ramanathan [Ram79]. By the analogue of [BG03, Thm. 5.1] for rational principal bundles (as explained in [BG03, §7]), the restriction of  $\mathcal{G}$  to a general smooth complete intersection curve  $C \hookrightarrow X$  of degree sufficiently large is a (slope) semistable  $G$ -bundle as in [Ram75]. Let  $\mathcal{H} := \varphi_*(\mathcal{G})$  be the associated  $H$ -bundle on  $U$ . Notice that  $\varphi_*(\mathcal{G}|_C)$  is the restriction  $\mathcal{H}|_C$ . By [BH04, Prop. 5.3]  $\varphi_*(\mathcal{G}|_C)$  is (slope) semistable on  $C$ , so the restriction of  $\mathcal{H}$  to a general smooth complete intersection curve  $C$  of degree sufficiently large is slope semistable in the sense of Ramanathan. By the argument in [BG03, Remark 5.2], it follows that  $\mathcal{H}$  is slope semistable.

This  $H$ -bundle  $\mathcal{H}$  amounts to the data of a  $q$ -tuple  $(\mathcal{E}_{U,i})_{i=1}^q$  of vector bundles on  $U$ . Let  $j$  denote the open inclusion  $j : U \hookrightarrow X$ . For each  $i$ , define  $\mathcal{E}_i := j_*(\mathcal{E}_{U,i})$ . The fact that  $\mathcal{H}$  is semistable implies that each  $\mathcal{E}_i$  is a slope semistable sheaf: otherwise the slope filtration of  $\mathcal{E}_i$  would restrict over  $U$  to a destabilizing filtration of the  $\mathcal{E}_{U,i}$ , thus contradicting the fact that  $\mathcal{H}$  is semistable.

For each  $i$ ,  $\det(V_i)$  is a character of  $G$  and, by the definition of degree,  $\vartheta(\det(V_i)) = c_1(\mathcal{E}_i) \cdot H^{d-1}$ . By Hizebruch-Riemann-Roch, the slope  $\widehat{\mu}_{d-1}(\mathcal{E}_i)$  is given by

$$\widehat{\mu}_{d-1}(\mathcal{E}_i) = \frac{1}{A_d \mathrm{rk}(\mathcal{E}_i)} c_1(\mathcal{E}_i) \cdot H^{d-1} + \frac{1}{A_d} t_1 \cdot H^{d-1} = \frac{1}{A_d \dim(V_i)} \vartheta(\det(V_i)) + \frac{1}{A_d} t_1 \cdot H^{d-1}$$

Therefore  $\widehat{\mu}_{d-1}(\mathcal{E}_i)$  is completely determined by  $\vartheta$ , and does not depend on  $(\mathcal{F}, \sigma)$ . Denote by  $\mu_{max}$  the maximum of the slopes  $\widehat{\mu}_{d-1}(\mathcal{E}_1), \widehat{\mu}_{d-1}(\mathcal{E}_2), \dots, \widehat{\mu}_{d-1}(\mathcal{E}_q)$ .

We have  $\bigoplus_{i=1}^q \mathcal{E}_i = \mathcal{F}^{\vee\vee}$ , where  $\mathcal{F}^{\vee\vee}$  denotes the reflexive hull of  $\mathcal{F}$ , because both sides restrict to the same associated vector bundle  $\rho_*(\mathcal{G})$  on  $U$ . If we set  $\mathcal{F}_j := \left( \bigoplus_{i=1}^j \mathcal{E}_i \right) \cap \mathcal{F} \subset \mathcal{F}^{\vee\vee}$ , this defines a filtration of  $\mathcal{F}$  by saturated subsheaves:

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \dots \subset \mathcal{F}_{q-1} \subset \mathcal{F}_q = \mathcal{F}$$

For each  $j$  we have  $(\mathcal{F}_j/\mathcal{F}_{j-1})|_U = \mathcal{E}_{U,j}$ , and therefore  $(\mathcal{F}_j/\mathcal{F}_{j-1})^{\vee\vee} = \mathcal{E}_j$ . This implies that  $\widehat{\mu}_{d-1}(\mathcal{F}_j/\mathcal{F}_{j-1}) = \widehat{\mu}_{d-1}(\mathcal{E}_j)$ . Since  $\mathcal{E}_j$  is slope semistable, it also follows that  $\mathcal{F}_j/\mathcal{F}_{j-1}$  is a slope semistable sheaf.

We have therefore found a filtration of  $\mathcal{F}$  such that each graded piece  $\mathcal{F}_j/\mathcal{F}_{j-1}$  is a slope semistable sheaf such that  $\widehat{\mu}_{d-1}(\mathcal{F}_j/\mathcal{F}_{j-1}) \leq \mu_{max}$ . This implies that subsheaves  $\mathcal{V} \subset \mathcal{F}$  satisfy  $\widehat{\mu}_{d-1}(\mathcal{V}) \leq \mu_{max}$ , as desired.  $\square$

**Remark 6.7.** *If the representation  $\rho$  is central (as in Definition 10.5), then the degree  $\vartheta$  of a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  is determined by the slope of  $\mathcal{F}$ . In particular, the degree  $\vartheta$  is determined by the Hilbert polynomial  $P$  of  $\mathcal{F}$ . This shows that in this case the notion of degree is superfluous and it suffices to specify the Hilbert polynomial  $P$  alone. Hence, Proposition 6.6 shows that  $\mathrm{Bun}_\rho(X)^{ss,P}$  is bounded for  $\rho$  central.*

**6.3. Moduli space for the semistable locus.** The work done in previous sections allows us to show that the stack of semistable  $\rho$ -sheaves admits a proper good moduli space in the sense of [Alp13].

**Theorem 6.8.** *For each choice of Hilbert polynomial  $P \in \mathbb{Q}[n]$  and degree  $\vartheta \in \Lambda(G)$ , the stack  $\mathrm{Bun}_\rho(X)_{\vartheta}^{ss,P}$  admits a proper good moduli space  $M_\rho(X)_{\vartheta}^{ss,P}$ .*

*Proof.* It suffices to check the hypotheses of [HLHJ21, Thm. 2.26] (see also [HL14, Thm. B]).

- (a) Proposition 4.16 implies that the numerical invariant  $\nu$  is strictly  $\Theta$ -monotone and strictly  $S$ -monotone.
- (b) By Proposition 6.6, the stack  $\text{Bun}_\rho(X)^{ss,P}$  is bounded.
- (c) By Theorem 5.7, the stack  $\text{Bun}_\rho(X)$  satisfies the existence part of the valuative criterion of properness.

□

If the group  $G$  is semisimple, then the representation  $\rho$  factors through the subgroup  $\text{SL}(V) \subset \text{GL}(V)$ . In this case the moduli space of semistable  $\rho$ -sheaves was already constructed in [Sch02, GLSS08] using GIT. For semisimple groups the proof above provides an alternative stack-theoretic approach which is intrinsic, in the sense that it does not appeal to an embedding into an external GIT problem.

On the other hand, suppose that the representation  $\rho$  is central. Then, the degree  $\vartheta$  is determined by the Hilbert polynomial  $P$  and each  $\text{Bun}_\rho(X)^{ss,P}$  admits a proper good moduli space  $M_\rho(X)^{ss,P}$ . Let  $\text{Bun}_G(X)^s$  denote the stack of slope stable  $G$ -bundles on  $X$ . By Corollary 10.9, the stack  $\text{Bun}_\rho(X)^{ss}$  of Gieseker semistable  $\rho$ -sheaves contains  $\text{Bun}_G(X)^s$  as an open substack. Therefore the separated moduli space  $M_\rho(X)^{ss} = \bigsqcup_{P \in \mathbb{Q}[n]} M_\rho(X)^{ss,P}$  contains an open subspace  $M_G(X)^s$  which is a good moduli space for the stack  $\text{Bun}_G(X)^s$ .

## 7. GENERALIZATION: $\rho$ -SHEAVES FOR PRODUCTS OF GENERAL LINEAR GROUPS

In this section we define the more general notion of  $\rho$ -sheaf valued in a product of general linear groups. All of the arguments from previous sections will apply in this more general context.

Example 10.10 shows that the notion of Gieseker stability could be quite restrictive if we do not assume that the representation  $\rho$  is central. Since the dimension of the center of  $\text{GL}(V)$  is 1, the faithful representation  $\rho$  can only be central if the center of  $G$  has dimension at most 1. In order to deal with reductive groups  $G$  with large center, it is convenient to allow the codomain of  $\rho$  to be a product of general linear groups. All of the previous results extend to this more general context.

Fix a tuple  $V^\bullet = (V^i)_{i=1}^b$  of finite-dimensional vector spaces over  $k$  with  $r_i := \dim V^i$ . Define  $\text{GL}(V^\bullet) := \prod_{i=1}^b \text{GL}(V^i)$ . We fix once and for all a faithful representation  $\rho : G \rightarrow \text{GL}(V^\bullet)$ , which induces representations  $\rho_i : G \rightarrow \text{GL}(V^i)$  by composing with the natural projections  $\text{GL}(V^\bullet) \rightarrow \text{GL}(V^i)$ .

Let  $S$  be a  $k$ -scheme. Suppose that we are given a tuple  $\mathcal{F}^\bullet = (\mathcal{F}^i)_{i=1}^b \in \prod_{i=1}^b \text{Coh}_{r_i}^{tf}(X)(S)$  of families of torsion-free sheaves on  $X_S$ . We set

$$E_{V^i}(\mathcal{F}^i) := \text{Sym}^\bullet(\mathcal{F}^i \otimes_k (V^i)^\vee) \otimes_{\mathcal{O}_{X_S}} \text{Sym}^\bullet(\det(\mathcal{F}^i)^\vee \otimes_k \det(V^i))$$

We also define  $H(\mathcal{F}^i, V^i) := \underline{\text{Spec}}_{X_S}(E_{V^i}(\mathcal{F}^i))$  and  $H(\mathcal{F}^\bullet, V^\bullet) := \prod_{i=1}^b H(\mathcal{F}^i, V^i)$ . Each scheme  $H(\mathcal{F}^i, V^i)$  admits a right action of the group  $G$  via the representation  $\rho_i$ , as described in Subsection 2.3. Equip  $H(\mathcal{F}^\bullet, V^\bullet)$  with the corresponding diagonal action of  $G$  and form the relatively affine GIT quotient  $H(\mathcal{F}^\bullet, V^\bullet)//G$  by taking the ring of  $G$ -invariants.

Let  $U$  be a big open subscheme such that  $\mathcal{F}^i|_U$  is locally-free for all  $i$ . The same construction as in Subsection 2.3 applies to yield a divisor  $\delta_U^i - 1$  on  $H(\mathcal{F}^i, V^i)|_U$ . This divisor is stable under the right action of  $G$ . We descend these divisors to get a closed subscheme  $\prod_{i=1}^b (H(\mathcal{F}^i, V^i)|_U)_{\alpha_U^i=0} \hookrightarrow (H(\mathcal{F}^\bullet, V^\bullet)|_U)//G$ . Define the scheme of  $G$ -reductions  $\text{Red}_G(\mathcal{F}^\bullet)$  to be the scheme-theoretic closure of this subset in  $H(\mathcal{F}^\bullet, V^\bullet)//G$ .

**Definition 7.1.** A  $\rho$ -sheaf on  $X_S$  is a pair  $(\mathcal{F}^\bullet, \sigma)$ , where  $\mathcal{F}^\bullet \in \prod_{i=1}^b \text{Coh}_{r_i}^{tf}(X)(S)$  and  $\sigma$  is a section  $\sigma : X_S \rightarrow \text{Red}_G(\mathcal{F}^\bullet)$ .

The tuple  $\mathcal{F}^\bullet|_U$  of locally-free sheaves defines a  $\text{GL}(V^\bullet)$ -bundle  $\mathcal{P}$  on  $U$ . The same argument as in Subsection 2.3 shows that the data of a section  $\sigma : X_S \rightarrow \text{Red}_G(\mathcal{F}^\bullet)$  is equivalent to a  $G$ -reduction of structure group for the  $\text{GL}(V^\bullet)$ -bundle  $\mathcal{P}$ . In particular we obtain a rational  $G$ -bundle  $\mathcal{G}$  defined on  $U$ . This allows to define the notion of degree  $\text{deg}(\mathcal{F}^\bullet, \sigma)$  of a  $\rho$ -sheaf as in Definition 6.3.

We denote by  $\text{Bun}_\rho(X)$  the stack on groupoids that takes an affine  $k$ -scheme  $T$  to the groupoid of  $\rho$ -sheaves on  $X_T$ . There is a natural forgetful morphism  $\text{Forget} : \text{Bun}_\rho(X) \rightarrow \prod_{i=1}^b \text{Coh}_{r_i}^{tf}(X)$  defined by  $\text{Forget}(\mathcal{F}^\bullet, \sigma) = \mathcal{F}^\bullet$ , which is affine and of finite type by the same reasoning as in Proposition 3.6. Therefore  $\text{Bun}_\rho(X)$  is an algebraic stack with affine diagonal and locally of finite type over  $k$ . All of the discussion for filtrations in Subsection 3.3 applies in this more general context, the only difference being that the underlying associated filtration is now a tuple of filtrations  $(\mathcal{F}_m^\bullet)_{m \in \mathbb{Z}}$ , where each  $(\mathcal{F}_m^\bullet)_{m \in \mathbb{Z}}$  is a filtration of the torsion-free sheaf  $\mathcal{F}^i$ .

We define a family of line bundles  $L_n$  on  $\text{Bun}_\rho(X)$  as follows. For each  $i$ , we have a family of line bundles  $L_n^i$  on  $\text{Coh}_{r_i}^{tf}(X)$  defined by the same formulas as in Definition 4.1. Then, we define the box product  $L_n := \boxtimes_{i=1}^b L_n^i$  on the stack  $\prod_{i=1}^b \text{Coh}_{r_i}^{tf}(X)$ , and use the morphism  $\text{Forget}$  to pullback  $L_n$  to a line bundle on  $\text{Bun}_\rho(X)$ , also denoted by  $L_n$ .

More explicitly, for every morphism  $f : T \rightarrow \text{Bun}_\rho(X)$  represented by a  $\rho$ -sheaf  $(\mathcal{F}^\bullet, \sigma)$  on  $X_T$ , we have

$$f^* L_n = \left( \bigotimes_{i=1}^b f^* M_n^i \right) \otimes \left( \bigotimes_{i=1}^b (f^* b_d^i)^{-\otimes \bar{p}_{\mathcal{F}^i}(n)} \right)$$

where  $f^* M_n^i := \det(R\pi_{T*}(\mathcal{F}^i(n)))$  and  $b_d^i := \bigotimes_{j=0}^d (M_j^i)^{\otimes (-1)^j \binom{d}{j}}$ .

We will use this family of line bundles  $L_n$  to define a polynomial numerical invariant on  $\text{Bun}_\rho(X)$ . Let  $(\mathcal{F}^\bullet, \sigma)$  be a  $\rho$ -sheaf on  $X_K$  for some field extension  $K \supset k$ . For every nondegenerate filtration  $f$  with underlying associated filtration  $(\mathcal{F}_m^\bullet)_{m \in \mathbb{Z}}$ , we set  $\nu(f)$  to be the polynomial

$$\nu(f) = \frac{\text{wgt}(L_n|_0)}{\sqrt{\sum_{i=1}^b \sum_{m \in \mathbb{Z}} m^2 \cdot \text{rk}(\mathcal{F}_m^i / \mathcal{F}_{m+1}^i)}}$$

The same computation as in Proposition 4.4 shows that this can be rewritten in terms of the reduced Hilbert polynomials as follows:

$$\nu(f) = \sqrt{A_d} \cdot \frac{\sum_{i=1}^b \sum_{m \in \mathbb{Z}} m \cdot (\bar{p}_{\mathcal{F}_m^i / \mathcal{F}_{m+1}^i} - \bar{p}_{\mathcal{F}^i}) \cdot \text{rk}(\mathcal{F}_m^i / \mathcal{F}_{m+1}^i)}{\sqrt{\sum_{i=1}^b \sum_{m \in \mathbb{Z}} m^2 \cdot \text{rk}(\mathcal{F}_m^i / \mathcal{F}_{m+1}^i)}}$$

One can define a version of the affine Grassmannian in this more general context. Let  $S$  be a quasicompact  $k$ -scheme and let  $\mathcal{F}^\bullet$  be a tuple of torsion-free sheaves in  $\prod_{i=1}^b \text{Coh}_{r_i}^{tf}(X)(S)$ . Let  $D \hookrightarrow X_S$  be an effective Cartier divisor that is flat over  $S$  and set  $Q = X_S \setminus D$ . Assume that we have a section  $\sigma : Q \rightarrow \text{Red}_G(\mathcal{F}^\bullet)$  defined over  $Q$ .

The affine Grassmannian  $\text{Gr}_{X_S, D, \mathcal{F}^\bullet, \sigma}$  is the functor from  $(\text{Aff}_S)^{\text{op}}$  into sets defined as follows. For each affine scheme  $T$  in  $\text{Aff}_S$ ,  $\text{Gr}_{X_S, D, \mathcal{F}^\bullet, \sigma}(T)$  is the set of equivalence classes of the following data

- (1) A  $\rho$ -sheaf  $(\mathcal{E}^\bullet, \zeta)$  on  $X_T$ .
- (2) A morphism  $\theta^\bullet : \mathcal{F}^\bullet|_{X_T} \rightarrow \mathcal{E}^\bullet$  of tuples such that the restriction  $\theta^\bullet|_{Q_T}$  is an isomorphism. Moreover, we require that the following diagram induced by the isomorphism  $\theta^\bullet|_{Q_T}$  commutes.

$$\begin{array}{ccc}
Q_T & \xrightarrow{\zeta|_{Q_T}} & \mathrm{Red}_G(\mathcal{E}^\bullet|_{Q_T}) \\
& \searrow^{\sigma|_{Q_T}} & \uparrow \wr \\
& & \mathrm{Red}_G(\mathcal{F}^\bullet|_{Q_T})
\end{array}$$

There is a forgetful morphism  $\mathrm{Gr}_{X_S, D, \mathcal{F}^\bullet, \sigma} \rightarrow \prod_{i=1}^b \mathrm{Gr}_{X_S, D, \mathcal{F}^i}$  that takes a point  $(\mathcal{E}^\bullet, \zeta, \theta^\bullet)$  to the tuple  $(\mathcal{E}^i, \theta^i)_{i=1}^b$ . The same proof as in Proposition 4.9 shows that this forgetful morphism exhibits  $\mathrm{Gr}_{X_S, D, \mathcal{F}^\bullet, \sigma}$  as a closed ind-subscheme of  $\prod_{i=1}^b \mathrm{Gr}_{X_S, D, \mathcal{F}^i}$ .

Let  $P^\bullet = (P^i)_{i=1}^b$  be a tuple of rational polynomials. For each  $N \geq 0$  we define the subfunctor

$$\mathrm{Gr}_{X_S, D, \mathcal{F}^\bullet, \sigma}^{\leq N, P^\bullet} := \mathrm{Gr}_{X_S, D, \mathcal{F}^\bullet, \sigma} \cap \prod_{i=1}^b \mathrm{Gr}_{X_S, D, \mathcal{F}^i}^{\leq N, P^i}$$

which is a closed subscheme of  $\mathrm{Gr}_{X_S, D, \mathcal{F}^\bullet, \sigma}$  and it is projective over  $S$ . There exists an integer  $m \gg 0$  such that, for all  $n \geq m$ , the restriction of the dual  $L_n^\vee$  is  $S$ -ample on the projective stratum  $\mathrm{Gr}_{X_S, D, \mathcal{F}^\bullet, \sigma}^{\leq N, P^\bullet}$ .

The proof of Proposition 4.16 applies without change in this more general context. The argument of HN boundedness in Proposition 4.26 also applies, one only needs to keep track of one extra index  $1 \leq i \leq b$  in all the formulas. In particular, Theorem 4.27 holds and the numerical invariant  $\nu$  induces a  $\Theta$ -stratification on the stack  $\mathrm{Bun}_\rho(X)$ . All of the arguments in Subsection 4.5 and Section 5 apply without change.

The tuple of rational polynomials  $P^\bullet = (P^i)_{i=1}^b$  defines an open substack  $\mathrm{Bun}_\rho(X)^{ss, P^\bullet} \subset \mathrm{Bun}_\rho(X)$  parametrizing Gieseker semistable  $\rho$ -sheaves  $(\mathcal{F}^\bullet, \sigma)$  such that the Hilbert polynomial of  $\mathcal{F}^i$  is  $P^i$ . If we fix the degree  $\vartheta$  of the  $\rho$ -sheaf, we obtain an open and closed substack  $\mathrm{Bun}_\rho(X)_{\vartheta}^{ss, P^\bullet} \subset \mathrm{Bun}_\rho(X)^{ss, P^\bullet}$ , which is quasicompact by the same proof as in Proposition 6.6. It follows by [HLHJ21, Thm. 2.26] that  $\mathrm{Bun}_\rho(X)_{\vartheta}^{ss, P^\bullet}$  admits a proper good moduli space  $M_\rho(X)_{\vartheta}^{ss, P^\bullet}$ .

**Theorem 7.2.** *The polynomial numerical invariant  $\nu$  induces a  $\Theta$ -stratification on the stack  $\mathrm{Bun}_\rho(X)$ . In particular the leading term HN filtration induces a stratification of  $\mathrm{Bun}_\rho(X)$  by locally closed substacks.*

*For each choice of degree  $\vartheta \in \Lambda(G)$  and tuple of Hilbert polynomials  $P^\bullet$ , the open substack  $\mathrm{Bun}_\rho(X)_{\vartheta}^{ss, P^\bullet}$  admits a proper good moduli space  $M_\rho(X)_{\vartheta}^{ss, P^\bullet}$ .  $\square$*

If the representation  $\rho$  is central, then the degree  $\vartheta$  is determined by the tuple of Hilbert polynomials  $P^\bullet$  and each  $\mathrm{Bun}_\rho(X)_{\vartheta}^{ss, P^\bullet}$  admits a proper good moduli space  $M_\rho(X)_{\vartheta}^{ss, P^\bullet}$ . In this case, the stack  $\mathrm{Bun}_\rho(X)^{ss}$  of Gieseker semistable  $\rho$ -sheaves contains the stack  $\mathrm{Bun}_G(X)^s$  of slope stable  $G$ -bundles as an open substack, by Corollary 10.9. Therefore the separated moduli space  $M_\rho(X)^{ss} = \bigsqcup_{P^\bullet \in (\mathbb{Q}[n])^b} M_\rho(X)_{\vartheta}^{ss, P^\bullet}$  contains an open subspace  $M_G(X)^s$  which is a good moduli space for the stack  $\mathrm{Bun}_G(X)^s$ . Hence  $M_\rho(X)^{ss}$  is a locally proper compactification of the moduli of slope stable  $G$ -bundles on  $X$  (c.f. [BG03, §7]).

Under the assumption that the representation  $\rho$  is central, the  $\Theta$ -stratification induced by the leading term HN filtration refines the slope stratification of  $\mathrm{Bun}_G(X)$  described in [GN18, GN16], by Proposition 10.11.

## 8. GIESEKER-HARDER-NARASIMHAN FILTRATIONS FOR $\rho$ -SHEAVES

The example in Subsection 9.1 shows that the leading term HN filtration of a  $\rho$ -sheaf is analogous to the leading term filtration of a torsion-free sheaf as defined in [HLHJ21, §5.2]. In this subsection we describe a procedure that iterates the leading term HN filtration in

order to construct a finer filtration for  $\rho$ -sheaves, called the Gieseker-Harder-Narasimhan filtration (GHN filtration). We prove that this filtration induces a stratification of  $\text{Bun}_\rho(X)$  by locally closed substacks.

To understand the notation and constructions in this section, it might help the reader to look beforehand at the examples in Subsections 9.1 and 9.2.

**8.1. Lexicographic filtrations.** Fix a positive integer  $q$ . We equip the group  $\mathbb{Z}^q$  with the lexicographic order where two tuples  $\vec{m} = (m_i)_{i=1}^q, \vec{n} = (n_i)_{i=1}^q \in \mathbb{Z}^q$  satisfy  $\vec{m} > \vec{n}$  if and only if there exists an index  $j$  such that  $m_j > n_j$  and  $m_i = n_i$  for all  $i < j$ . The lexicographic order endows  $\mathbb{Z}^q$  with the structure of a totally well-ordered abelian group. For  $\vec{m} = (m_1, m_2, \dots, m_{q-1}, m_q) \in \mathbb{Z}^q$ , we define the successor by  $\vec{m} + 1 := (m_1, m_2, \dots, m_{q-1}, m_q + 1)$ .

Let  $K \supset k$  be a field extension and let  $\mathcal{F}$  be a torsion-free sheaf on  $X_K$ .

**Definition 8.1.** *A lexicographic  $\mathbb{Z}^q$ -filtration of  $\mathcal{F}$  is a sequence of subsheaves  $f = (\mathcal{F}_{\vec{m}})_{\vec{m} \in \mathbb{Z}^q}$  indexed by  $\mathbb{Z}^q$  such that the following are satisfied.*

- (1)  $\mathcal{F}_{\vec{m}} \subset \mathcal{F}_{\vec{n}}$  whenever  $\vec{m} > \vec{n}$ .
- (2) There exists an  $M_{\max} \in \mathbb{Z}^q$  such that  $\mathcal{F}_{\vec{m}} = 0$  for all  $\vec{m} > M_{\max}$ , and there exists an  $M_{\min} \in \mathbb{Z}^q$  such that  $\mathcal{F}_{\vec{m}} = \mathcal{F}$  for all  $\vec{m} < M_{\min}$ .
- (3) For all  $\vec{m}$ , we have that  $\mathcal{F}/\mathcal{F}_{\vec{m}}$  is torsion-free.

The associated graded of a lexicographic  $\mathbb{Z}^q$ -filtration  $f = (\mathcal{F}_{\vec{m}})_{\vec{m} \in \mathbb{Z}^q}$  is defined as  $\text{gr}(f) := \bigoplus_{\vec{m} \in \mathbb{Z}^q} \mathcal{F}_{\vec{m}}/\mathcal{F}_{\vec{m}+1}$ . Conditions (2) and (3) in Definition 8.1 ensure that  $\text{gr}(f)$  is a torsion-free sheaf with the same rank as  $\mathcal{F}$ . Note that  $\text{gr}(f)$  is naturally equipped with a  $\mathbb{Z}^q$ -grading. We say that  $\vec{m} \in \mathbb{Z}^q$  is a jumping point of  $f$  if the quotient  $\mathcal{F}_{\vec{m}}/\mathcal{F}_{\vec{m}+1}$  is not the zero sheaf. There are finitely many jumping points.

**Definition 8.2.** *Let  $f$  be a lexicographic  $\mathbb{Z}^q$ -filtration. Let  $\vec{p} = (p_i)_{i=1}^q$  be a tuple of positive rational numbers. For all coordinates  $i$ , we can scale the weights of  $f$  at the  $i^{\text{th}}$  coordinate by the rational number  $p_i$ . Doing this for all  $i$  yields a lexicographic multi-weighted filtration denoted by  $\vec{p} \cdot f$  that could possibly have rational weights. If  $\vec{p} \cdot f$  has integer weights, then we say that the lexicographic  $\mathbb{Z}^q$ -filtration  $\vec{p} \cdot f$  is obtained by scaling the weights of  $f$  by  $\vec{p}$ .*

The scheme  $\mathbb{A}_k^q = \text{Spec}(k[t_1, t_2, \dots, t_q])$  admits an action of  $\mathbb{G}_m^q$  over  $k$ , where the  $i^{\text{th}}$  coordinate of  $\mathbb{G}_m^q$  acts with weight  $-1$  on  $t_i$  and weight  $0$  on the other coordinates. Set  $\Theta_k^q := [\mathbb{A}_k^q/\mathbb{G}_m^q]$  and denote by  $\Theta_K^q := [\mathbb{A}_K^q/(\mathbb{G}_m^q)_K]$  the base change. A multi-graded version of the Rees construction in Proposition 3.8 can be applied to associate to any lexicographic  $\mathbb{Z}^q$ -filtration a morphism of stacks  $\Theta_K^q \rightarrow \text{Coh}^{tf}(X)$  in the following way.

**Definition 8.3.** *Given a lexicographic  $\mathbb{Z}^q$ -filtration  $f = (\mathcal{F}_{\vec{m}})_{\vec{m} \in \mathbb{Z}^q}$  of  $\mathcal{F}$ , we set multi-graded Rees construction  $\tilde{\mathcal{F}}_f$  to be the  $\mathbb{G}_m^q$ -equivariant  $\mathbb{A}_K^q$ -flat sheaf on  $\mathbb{A}_K^q \times X$  defined by*

$$\tilde{\mathcal{F}}_f = \bigoplus_{\vec{m} \in \mathbb{Z}^q} \left( \mathcal{F}_{\vec{m}} \cdot \prod_{i=1}^q t_i^{m_i} \right)$$

Note that this is a coherent sheaf by condition (2) in Definition 8.1. If  $\vec{1}_K$  denotes the point  $(1, 1, \dots, 1) \in \mathbb{A}_K^q$ , the fiber at  $\vec{1}_K \times X$  is isomorphic to the original sheaf  $\mathcal{F}$  and the fiber at  $0_K \times X$  is the  $\mathbb{Z}^q$ -graded sheaf  $\text{gr}(f)$ . The locus of points in  $\mathbb{A}_K^q$  such that the fiber of  $\tilde{\mathcal{F}}_f$  is torsion-free is open by the slicing criterion for flatness [Sta21, Tag 046Y], and the fiber at the origin  $0_K$  is torsion-free by condition (3) in Definition 8.1. By  $\mathbb{G}_m^q$ -equivariance, it follows that  $\tilde{\mathcal{F}}_f$  is a family of torsion-free sheaves. So indeed  $\tilde{\mathcal{F}}_f$  represents a morphism  $\Theta_K^q \rightarrow \text{Coh}^{tf}(X)$ .



More generally, a lexicographic  $\mathbb{Z}^q$ -filtration  $f$  of a tuple  $\mathcal{F}^\bullet = (\mathcal{F}_i)_{i=1}^b$  of torsion-free sheaves on  $X_K$  is defined to be a tuple  $f = (f_i)_{i=1}^b$  where  $f_i$  is a lexicographic  $\mathbb{Z}^q$ -filtration of  $\mathcal{F}^i$ . Here it also makes sense to scale the weights by a tuple  $\vec{p} = (p_j)_{j=1}^q$  of rational numbers; we just scale the weights of each  $f_i$  by  $\vec{p}$ . If  $\mathcal{F}^\bullet$  is a point in  $\prod_{i=1}^b \text{Coh}_{r_i}^{tf}(X)$ , then the multi-graded Rees construction applied to each component yields a morphism  $\Theta_K^q \rightarrow \prod_{i=1}^b \text{Coh}_{r_i}^{tf}(X)$  represented by a tuple  $\widetilde{\mathcal{F}}^\bullet_f$  such that the fiber at  $\vec{1}_K \times X$  recovers  $\mathcal{F}^\bullet$ .

Fix a tuple  $V^\bullet$  of  $k$ -vector spaces and let  $\rho : G \rightarrow \text{GL}(V^\bullet)$  be faithful representation, as in Section 7. Let  $(\mathcal{F}^\bullet, \sigma)$  be a  $\rho$ -sheaf on  $X_K$  and suppose that we are given a lexicographic  $\mathbb{Z}^q$ -filtration  $f$  of  $\mathcal{F}^\bullet$ . Applying the multi-graded Rees construction, we obtain a homomorphism  $\Theta_K^q \rightarrow \prod_{i=1}^b \text{Coh}_{r_i}^{tf}(X)$  represented by a tuple  $\widetilde{\mathcal{F}}^\bullet_f$ . If  $U$  is a big open subset of  $X_K$  such that  $\text{gr}(f)$  is locally-free, the restriction  $\widetilde{\mathcal{F}}^\bullet_f|_{\Theta_K^q \times U}$  is a tuple of locally-free sheaves, which we can view as a  $\text{GL}(V^\bullet)$ -bundle on  $\Theta_K^q \times_K U$ . The section  $\sigma$  induces a  $G$ -reduction of structure group of the restriction  $\widetilde{\mathcal{F}}^\bullet_f|_{\vec{1}_K \times_K U}$ .

**Definition 8.4.** *We say that  $f$  is a lexicographic  $\mathbb{Z}^q$ -filtration of  $(\mathcal{F}^\bullet, \sigma)$  if the  $G$ -reduction of  $\widetilde{\mathcal{F}}^\bullet_f|_{\vec{1}_K \times_K U}$  induced by  $\sigma$  extends to a  $G$ -reduction defined over  $\Theta_K^q \times U$ .*

Since the  $(\mathbb{G}_m^q)_K$ -orbit of  $\vec{1}_K$  is scheme-theoretically dense in  $\mathbb{A}_K^q$ , if such extension exists then it must be unique. So, again, being a filtration of  $(\mathcal{F}^\bullet, \sigma)$  is a property, not a structure.

**8.2. Multi-weighted parabolic reductions.** For this subsection we fix a field extension  $K \supset k$ .

Let  $H$  be a reductive group over  $K$  and let  $\vec{\lambda} = (\lambda_i)_{i=1}^q$  be a tuple of  $q$  cocharacters of  $H$ . The first cocharacter  $\lambda_1$  defines a parabolic subgroup  $P_{\lambda_1}$  with corresponding Levi subgroup  $L_{\lambda_1}$ . We denote by  $q_{\lambda_1} : P_{\lambda_1} \rightarrow L_{\lambda_1}$  the quotient morphism. Suppose that the cocharacter  $\lambda_2 : \mathbb{G}_m \rightarrow H$  factors through the subgroup  $L_{\lambda_1}$  and let  $\overline{\lambda}_2 : \mathbb{G}_m \rightarrow L_{\lambda_1}$  be the corresponding cocharacter. This induces a parabolic subgroup  $P_{\overline{\lambda}_2} \subset L_{\lambda_1}$  with corresponding Levi subgroup  $L_{\overline{\lambda}_2}$ . The preimage  $P_{(\lambda_1, \lambda_2)} := (q_{\lambda_1})^{-1}(P_{\overline{\lambda}_2})$  is a parabolic subgroup of  $H$  contained in  $P_{\lambda_1}$  (c.f. second paragraph of the proof of [BH04, Prop. 3.1]). We set  $L_{(\lambda_1, \lambda_2)} := L_{\overline{\lambda}_2}$ , the joint centralizer of  $\lambda_1$  and  $\lambda_2$ . By definition,  $L_{(\lambda_1, \lambda_2)}$  is a Levi subgroup of  $P_{(\lambda_1, \lambda_2)}$ ; we denote by  $q_{(\lambda_1, \lambda_2)} : P_{(\lambda_1, \lambda_2)} \rightarrow L_{(\lambda_1, \lambda_2)}$  the quotient morphism. If the cocharacter  $\lambda_3$  factors through the  $L_{(\lambda_1, \lambda_2)}$ , then we can similarly use  $q_{(\lambda_1, \lambda_2)}$  to define a smaller parabolic subgroup  $P_{(\lambda_1, \lambda_2, \lambda_3)}$  with Levi subgroup  $L_{(\lambda_1, \lambda_2, \lambda_3)}$  given by the joint centralizer of the cocharacters  $\lambda_1$ ,  $\lambda_2$  and  $\lambda_3$ , and iterate this process.

**Definition 8.5.** *We say that the tuple  $\vec{\lambda}$  is nested if at the  $i^{\text{th}}$  step of the procedure described above we have that  $\lambda_i$  factors through the subgroup  $L_{(\lambda_1, \lambda_2, \dots, \lambda_{i-1})}$ .*

For a nested tuple  $\vec{\lambda} = (\lambda_1, \dots, \lambda_q)$ , this procedure terminates at the  $q^{\text{th}}$  step and yields a parabolic subgroup  $P_{\vec{\lambda}}$  of  $H$  with Levi subgroup  $L_{\vec{\lambda}}$ . Furthermore these groups fit into nested chains of subgroups:

$$\begin{aligned} P_{\vec{\lambda}} &\subset P_{(\lambda_1, \lambda_2, \dots, \lambda_{q-1})} \subset \dots \subset P_{(\lambda_1, \lambda_2)} \subset P_{\lambda_1} \subset H \\ L_{\vec{\lambda}} &\subset L_{(\lambda_1, \lambda_2, \dots, \lambda_{q-1})} \subset \dots \subset L_{(\lambda_1, \lambda_2)} \subset L_{\lambda_1} \subset H \end{aligned}$$

**Definition 8.6.** *Let  $Y$  be a  $K$ -scheme. Let  $\mathcal{H}$  be a  $H$ -bundle on  $Y$ . A  $q$ -weighted parabolic reduction of  $\mathcal{H}$  consists of a pair  $(\vec{\lambda}, \mathcal{H}_{\vec{\lambda}})$ , where  $\vec{\lambda}$  is a tuple of nested cocharacters of  $H$  and  $\mathcal{H}$  is a reduction of structure group of  $\mathcal{H}$  to the subgroup  $P_{\vec{\lambda}}$ .*

Let  $\theta : H \rightarrow F$  be a homomorphism of reductive groups over  $K$ . Let  $\vec{\lambda} = (\lambda_i)_{i=1}^q$  be a nested tuple of cocharacters of  $H$ . We set  $\theta_*(\vec{\lambda}) = (\theta \circ \lambda_i)_{i=1}^q$ , which can be checked to

be a nested tuple of cocharacters of  $F$ . The homomorphism  $\theta$  restricts to a well-defined homomorphism  $\theta : P_{\vec{\lambda}} \rightarrow P_{\theta_*(\vec{\lambda})}$ . Let  $Y$  be a  $K$ -scheme and let  $\mathcal{H}$  be a  $H$ -bundle on  $Y$ . Given a  $q$ -weighted parabolic reduction  $(\vec{\lambda}, \mathcal{H}_{\vec{\lambda}})$  of  $H$ , we can associate a  $q$ -weighted parabolic reduction  $(\theta_*(\vec{\lambda}), \theta_*\mathcal{H}_{\vec{\lambda}})$  of  $\theta_*\mathcal{H}$ . Here  $\theta_*\mathcal{H}_{\vec{\lambda}}$  is the parabolic reduction obtained from  $\mathcal{H}_{\vec{\lambda}}$  via the restriction  $\theta : P_{\vec{\lambda}} \rightarrow P_{\theta_*(\vec{\lambda})}$ .

We now return to our original group  $G_K$ . Suppose that we are given a  $G_K$ -bundle  $\mathcal{G}$  on  $Y$ . Given a  $q$ -weighted parabolic reduction  $(\vec{\lambda}, \mathcal{G}_{\vec{\lambda}})$  as above, we shall construct a  $G_K$ -bundle on  $\Theta_K^q \times_K Y$  by iterating the Rees construction.

Take the last cocharacter  $\lambda_q$  of  $\vec{\lambda}$  and consider the  $\mathbb{G}_m$ -equivariant  $P_{\vec{\lambda}}$ -bundle  $\tilde{\mathcal{G}}^1$  on  $\mathbb{A}_K^1 \times_K Y$  given by the Rees construction  $\text{Rees}(\mathcal{G}_{\vec{\lambda}}, \lambda_q)$ . Using  $P_{\vec{\lambda}} \subset P_{(\lambda_1, \lambda_2, \dots, \lambda_{q-1})}$ , we extend the structure group of  $\tilde{\mathcal{G}}^1$  to the bigger parabolic  $P_{(\lambda_1, \lambda_2, \dots, \lambda_{q-1})}$ . Given that the subsequent cocharacter  $\lambda_{q-1}$  factors through  $P_{(\lambda_1, \lambda_2, \dots, \lambda_{q-1})}$ , we apply the Rees construction to the  $P_{(\lambda_1, \lambda_2, \dots, \lambda_{q-1})}$ -bundle  $\tilde{\mathcal{G}}^1$  on  $\mathbb{A}_K^1 \times_K Y$  using the cocharacter  $\lambda_{q-1}$ . This way we obtain the  $\mathbb{G}_m^2$ -equivariant  $P_{(\lambda_1, \lambda_2, \dots, \lambda_{q-1})}$ -bundle  $\tilde{\mathcal{G}}^2 := \text{Rees}(\tilde{\mathcal{G}}^1, \lambda_{q-1})$  on  $\mathbb{A}_K^2 \times_K Y$ . We iterate this procedure: at the  $i^{\text{th}}$  step we have a  $P_{(\lambda_1, \lambda_2, \dots, \lambda_{q-i})}$ -bundle  $\tilde{\mathcal{G}}^i$  on  $\mathbb{A}_K^i \times_K Y$  and we apply the Rees construction using the cocharacter  $\lambda_{q-i}$  to get the  $\mathbb{G}_m^{i+1}$ -equivariant  $P_{(\lambda_1, \lambda_2, \dots, \lambda_{q-i-1})}$ -bundle  $\tilde{\mathcal{G}}^{i+1} := \text{Rees}(\tilde{\mathcal{G}}^i, \lambda_{q-i})$  on  $\mathbb{A}_K^{i+1} \times_K Y$ .

**Definition 8.7.** We denote by  $\text{Rees}(\mathcal{G}_{\vec{\lambda}}, \vec{\lambda})$  the  $P_{\lambda_1}$ -bundle on  $\Theta_K^q \times_K Y$  obtained at the  $q^{\text{th}}$  step of the procedure described above.

By construction, the restriction  $\text{Rees}(\mathcal{G}_{\vec{\lambda}}, \vec{\lambda})|_{\tilde{\Gamma}_K \times_K Y}$  is canonically isomorphic to the  $P_{\lambda_1}$ -bundle obtained via  $\mathcal{G}_{\vec{\lambda}}$  using the inclusion  $P_{\vec{\lambda}} \subset P_{\lambda_1}$ . We can extend the structure group from  $P_{\lambda_1}$  to  $G_K$  in order to obtain a  $G_K$ -bundle  $\text{Rees}(\mathcal{G}_{\vec{\lambda}}, \vec{\lambda}) \times^{P_{\lambda_1}} G_K$  on  $\Theta_K^q \times_K Y$ . The restriction to  $\tilde{\Gamma}_K \times_K Y$  is canonically isomorphic to the original  $G_K$ -bundle  $\mathcal{G}$ .

**Example 8.8.** If the group is a general linear group  $\text{GL}(V)$ , this process can be described more concretely as follows. Let  $\mathcal{E}$  be a  $\text{GL}(V)_K$ -bundle on  $Y$ . We can think of  $\mathcal{E}$  as a vector bundle of constant rank. Let  $(\vec{\lambda}, \mathcal{E}_{\vec{\lambda}})$  be a  $q$ -weighted parabolic reduction of  $\mathcal{E}$ . For each index  $i$ , we set  $\mathcal{E}_{(\lambda_1, \lambda_2, \dots, \lambda_i)}$  to be the  $P_{(\lambda_1, \lambda_2, \dots, \lambda_i)}$ -parabolic reduction of  $\mathcal{E}$  obtained from  $\mathcal{E}_{\vec{\lambda}}$  by extending the structure group  $P_{\vec{\lambda}} \subset P_{(\lambda_1, \lambda_2, \dots, \lambda_i)}$ . The pair  $(\lambda_i, \mathcal{E}_{(\lambda_1, \lambda_2, \dots, \lambda_i)})$  amounts to  $\mathbb{Z}$ -weighted filtration  $(\mathcal{E}_m^{(i)})_{m \in \mathbb{Z}}$  of  $\mathcal{E}$ . We use this to define a lexicographic  $\mathbb{Z}^q$ -filtration  $f = (\mathcal{E}_{\vec{m}})_{\vec{m} \in \mathbb{Z}^q}$  of  $\mathcal{E}$  by vector subbundles where for a tuple  $\vec{m} = (m_i)_{i=1}^q$ , we set  $\mathcal{E}_{\vec{m}} = \bigcap_{i=1}^q \mathcal{E}_{m_i}^{(i)}$ . Then we apply the multi-graded Rees construction to  $f$  in order to obtain a vector bundle  $\tilde{\mathcal{E}}_f$  on  $\Theta_K^q \times_K Y$ . The  $\text{GL}(V)_K$ -bundle  $\tilde{\mathcal{E}}_f$  is precisely  $\text{Rees}(\mathcal{G}_{\vec{\lambda}}, \vec{\lambda}) \times^{P_{\lambda_1}} \text{GL}(V)_K$ . A similar description holds for  $\text{GL}(V^\bullet)$ -bundles, by applying this construction to each component of a tuple  $\mathcal{E}^\bullet$ .

Now let  $(\mathcal{F}^\bullet, \sigma)$  be a  $\rho$ -sheaf over  $X_K$  and let  $f = (\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  be a lexicographic  $\mathbb{Z}^q$ -filtration of  $\mathcal{F}^\bullet$ . Let  $U$  be a big open subset of  $X_K$  where the tuple  $\text{gr}(f)$  is locally-free. Then the restriction  $(\tilde{\mathcal{F}}_f^\bullet)|_{\Theta_K^q \times_K U}$  is locally-free, so we can view it as a  $\text{GL}(V^\bullet)$ -bundle. Since  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  is a filtration of the  $\rho$ -sheaf, this  $\text{GL}(V^\bullet)$  bundle comes equipped with a canonical reduction of structure group  $\tilde{\mathcal{G}}$  to  $G_K$ . Let  $\mathcal{G}$  be the  $G_K$ -bundle on  $U$  corresponding to the  $\rho$ -sheaf  $(\mathcal{F}^\bullet, \sigma)$  (equivalently  $\mathcal{G} = \tilde{\mathcal{G}}|_{\tilde{\Gamma}_K \times_K U}$ ).

**Definition 8.9.** We say that the lexicographic filtration  $f = (\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  comes from a  $q$ -weighted parabolic reduction  $(\vec{\lambda}, \mathcal{G}_{\vec{\lambda}})$  of  $\mathcal{G}$  if  $\tilde{\mathcal{G}} = \text{Rees}(\mathcal{G}_{\vec{\lambda}}, \vec{\lambda}) \times^{P_{\lambda_1}} G_K$ .

More explicitly, if  $j : U \hookrightarrow X_K$  is a big open subset where the tuple  $\mathcal{F}^\bullet$  is locally-free, a  $q$ -weighted parabolic reduction  $(\vec{\lambda}, \mathcal{G}_{\vec{\lambda}})$  induces a  $q$ -weighted parabolic reduction

$(\rho_*(\vec{\lambda}), \rho_*\mathcal{G}_{\vec{\lambda}})$  of the  $\mathrm{GL}(V^\bullet)_K$ -bundle  $\mathcal{F}^\bullet|_U := \mathcal{E}^\bullet$ . We define a lexicographic  $\mathbb{Z}^q$ -filtration  $(\mathcal{E}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  of  $\mathcal{E}^\bullet$  associated to the  $q$ -weighted parabolic reduction  $(\rho_*(\vec{\lambda}), \rho_*\mathcal{G}_{\vec{\lambda}})$  and, for each  $\vec{m} \in \mathbb{Z}^q$ , set  $\tilde{\mathcal{E}}_{\vec{m}}^\bullet := j_*(\mathcal{E}_{\vec{m}}^\bullet) \cap \mathcal{F}^\bullet$  (the intersection being taken for each coordinate of the tuple). This defines a lexicographic  $\mathbb{Z}^q$ -filtration of  $\mathcal{F}^\bullet$ , and  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q} = (\tilde{\mathcal{E}}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  exactly when  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  comes from the  $q$ -weighted parabolic reduction  $(\vec{\lambda}, \mathcal{G}_\lambda)$ .

**8.3. The GHN filtration for  $\rho$ -sheaves.** Fix a tuple  $V^\bullet = (V^i)_{i=1}^b$  of vector spaces over  $k$  and let  $\rho : G \rightarrow \mathrm{GL}(V^\bullet)$  be a faithful representation. In this subsection, we recursively apply the leading term HN filtration to the associated graded Levi sheaves in order to obtain a sequence of multi-weighted parabolic reductions of  $(\mathcal{F}^\bullet, \sigma)$ .

Let  $K \supset k$  be a field extension such that  $G_K$  is split and let  $(\mathcal{F}^\bullet, \sigma)$  be an unstable  $\rho$ -sheaf on  $X_K$ . Let us proceed to describe the recursive construction of the GHN filtration of  $(\mathcal{F}^\bullet, \sigma)$ .

First Step: Let  $f_{can,1} := (\mathcal{F}_m^\bullet)_{m \in \mathbb{Z}}$  be the leading term HN filtration of  $(\mathcal{F}^\bullet, \sigma)$  and let  $(\lambda_1, \mathcal{G}_{\lambda_1})$  be the corresponding weighted parabolic reduction. The cocharacter  $\rho \circ \lambda_1$  induces a grading on each vector space  $V_i$ . This induces a nontrivial direct sum decomposition of the tuple  $V^\bullet$ . Indeed, if the decomposition was trivial then  $\rho \circ \lambda_1$  would be a cocharacter of the center of  $\mathrm{GL}(V^\bullet)$  and using the same reasoning as in Proposition 10.6 we would get  $\nu(f|_{can,1}) = 0$ , contradicting the fact that  $f|_{can,1}$  is destabilizing.

We can think of this direct sum decomposition as a finer tuple  $V_{\lambda_1}^\bullet$  of direct summands, where we have increased the number of indexes. The group  $\mathrm{GL}(V_{\lambda_1}^\bullet)$  is the Levi subgroup of  $\mathrm{GL}(V^\bullet)$  determined by the cocharacter  $\rho \circ \lambda_1$  which, by definition, contains the image of the Levi  $L_{\lambda_1} \subset G$  under  $\rho$ . We denote by  $\rho_{\lambda_1} : L_{\lambda_1} \rightarrow \mathrm{GL}(V_{\lambda_1}^\bullet)$  the restriction to  $L_{\lambda_1}$  of the homomorphism  $\rho$ .

The description of the associated graded  $f_{can,1}|_0 = (\bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m^\bullet / \mathcal{F}_{m+1}^\bullet, \mathrm{gr}(\sigma))$  given at the end of Subsection 3.3 shows that we can canonically view  $f_{can,1}|_0$  as a  $\rho_{\lambda_1}$ -sheaf for the Levi  $L_{\lambda_1}$ . We call this the 1<sup>st</sup> associated graded Levi sheaf and denote it by  $\mathrm{gr}_1(\mathcal{F}^\bullet, \sigma) := (\mathrm{gr}_1(\mathcal{F}^\bullet), \mathrm{gr}_1(\sigma))$ .

Second Step: It might happen that the  $\rho_{\lambda_1}$ -sheaf  $\mathrm{gr}_1(\mathcal{F}^\bullet, \sigma)$  is unstable (see Subsection 9.1 below). If that is the case, we can consider the leading term HN filtration  $f_{can,2} = (\mathrm{gr}_1(\mathcal{F}^\bullet)_m)_{m \in \mathbb{Z}}$  of the  $\rho_{\lambda_1}$ -sheaf  $\mathrm{gr}_1(\mathcal{F}^\bullet, \sigma)$ . This is by definition a  $\mathbb{Z}$ -filtration of  $\mathrm{gr}_1(\mathcal{F}^\bullet)$ . Note that  $(L_{\lambda_1})_K$  must also be split, and therefore there exists a weighted parabolic reduction  $(\lambda_2, \mathcal{G}_{\lambda_2})$  of  $\mathrm{gr}_1(\mathcal{F}^\bullet, \sigma)$  that induces the leading term HN filtration  $(\mathrm{gr}_1(\mathcal{F}^\bullet)_m)_{m \in \mathbb{Z}}$ .

By construction we can think of  $(\lambda_1, \lambda_2)$  as a nested pair. We set  $L_{(\lambda_1, \lambda_2)}$  to be the Levi subgroup of  $L_{\lambda_1}$  centralizing  $\lambda_2$  and denote by  $P_{(\lambda_1, \lambda_2)}$  the preimage under  $q_{\lambda_1} : P_{\lambda_1} \rightarrow L_{\lambda_1}$  of  $P_{\lambda_2}$ . By [BH04, Lem. 2.1], the  $P_{\lambda_2}$ -parabolic reduction  $\mathcal{G}_{\lambda_2}$  lifts to a unique  $P_{(\lambda_1, \lambda_2)}$ -parabolic reduction  $\mathcal{G}_{(\lambda_1, \lambda_2)}$  that recovers the  $P_{\lambda_1}$ -reduction  $\mathcal{G}_{\lambda_1}$  under the extension of groups  $P_{(\lambda_1, \lambda_2)} \subset P_{\lambda_1}$ .

Each piece in the  $\mathbb{Z}$ -filtration  $(\mathrm{gr}_1(\mathcal{F}_m^\bullet))_{m \in \mathbb{Z}}$  of the associated graded  $\mathrm{gr}_1(\mathcal{F}^\bullet) = \bigoplus_{m \in \mathbb{Z}} \mathcal{F}_m^\bullet / \mathcal{F}_{m+1}^\bullet$  can be lifted to the original  $\mathcal{F}^\bullet$  by taking the preimages in each subquotient. This way we obtain a lexicographic  $\mathbb{Z}^2$ -filtration  $(\mathcal{F}_{(m_1, m_2)}^\bullet)_{(m_1, m_2) \in \mathbb{Z}^2}$  of the original tuple  $\mathcal{F}^\bullet$ . This is the lexicographic filtration coming from the 2-weighted parabolic reduction  $((\lambda_1, \lambda_2), \mathcal{G}_{(\lambda_1, \lambda_2)})$ . In particular  $(\mathcal{F}_{(m_1, m_2)}^\bullet)_{(m_1, m_2) \in \mathbb{Z}^2}$  is actually a lexicographic  $\mathbb{Z}^2$ -filtration of  $(\mathcal{F}^\bullet, \sigma)$ .

The cocharacter  $\lambda_2$  induces a nontrivial direct sum decomposition of the tuple  $V_{\lambda_1}^\bullet$ . We can think of this decomposition as a (strictly) finer tuple  $V_{(\lambda_1, \lambda_2)}^\bullet$  of direct summands. The restriction of  $\rho$  to the Levi subgroup  $L_{(\lambda_1, \lambda_2)}$  factors through  $\mathrm{GL}(V_{(\lambda_1, \lambda_2)}^\bullet)$ , which is a smaller Levi subgroup of  $\mathrm{GL}(V^\bullet)$  contained in the Levi  $\mathrm{GL}(V_{\lambda_1}^\bullet)$ . We denote this restriction

by  $\rho_{(\lambda_1, \lambda_2)} : L_{(\lambda_1, \lambda_2)} \rightarrow \mathrm{GL}(V_{(\lambda_1, \lambda_2)}^\bullet)$ . The associated graded  $f_{can, 2}|_0$  acquires canonically the structure of a  $\rho_{(\lambda_1, \lambda_2)}$ -sheaf, denoted by  $\mathrm{gr}_2(\mathcal{F}^\bullet, \sigma) := (\mathrm{gr}_2(\mathcal{F}^\bullet), \mathrm{gr}_2(\sigma))$ . We call this the 2<sup>nd</sup> associated graded Levi sheaf. We have that  $\mathrm{gr}_2(\mathcal{F}^\bullet)$  is the associated graded for the lexicographic  $\mathbb{Z}^2$ -filtration  $(\mathcal{F}_{(m_1, m_2)}^\bullet)_{(m_1, m_2) \in \mathbb{Z}^2}$  that we have defined. Moreover, its corresponding  $L_{(\lambda_1, \lambda_2)}$ -rational bundle is the same as the one obtained from the  $P_{(\lambda_1, \lambda_2)}$ -parabolic reduction  $\mathcal{G}_{(\lambda_1, \lambda_2)}$  via the Levi quotient map  $P_{(\lambda_1, \lambda_2)} \rightarrow L_{(\lambda_1, \lambda_2)}$ . So we can view  $\mathrm{gr}_2(\mathcal{F}^\bullet)$  as the associated graded of the 2-weighted parabolic reduction  $((\lambda_1, \lambda_2), \mathcal{G}_{(\lambda_1, \lambda_2)})$ .<sup>7</sup>

**Recursion:** If  $\mathrm{gr}_2(\mathcal{F}^\bullet, \sigma)$  is not Gieseker semistable, then we can iterate this construction and consider the leading term HN filtration  $f_{can, 3}$ , and so on. At the  $j^{\mathrm{th}}$  step of this iteration we have a  $j$ -weighted parabolic reduction  $((\lambda_1, \dots, \lambda_j), \mathcal{G}_{(\lambda_1, \dots, \lambda_j)})$  of  $(\mathcal{F}^\bullet, \sigma)$ . The associated underlying lexicographic  $\mathbb{Z}^j$ -filtration  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^j}$  coming from  $((\lambda_1, \dots, \lambda_j), \mathcal{G}_{(\lambda_1, \dots, \lambda_j)})$  is uniquely determined up to scaling the weights. Recall that  $(\lambda_1, \dots, \lambda_j)$  determines a Levi subgroup  $L_{(\lambda_1, \dots, \lambda_j)}$  that centralizes all the cocharacters in the tuple. We have an induced direct sum decomposition  $V_{(\lambda_1, \dots, \lambda_j)}^\bullet$ , so that the representation  $\rho$  restricted to  $L_{(\lambda_1, \dots, \lambda_j)}$  factors as  $\rho_{(\lambda_1, \dots, \lambda_j)} : L_{(\lambda_1, \dots, \lambda_j)} \rightarrow \mathrm{GL}(V_{(\lambda_1, \dots, \lambda_j)}^\bullet)$ . We also have a  $j^{\mathrm{th}}$  associated graded Levi  $\rho_{(\lambda_1, \dots, \lambda_j)}$ -sheaf  $(\mathrm{gr}_j(\mathcal{F}^\bullet), \mathrm{gr}_j(\sigma))$ .

**Termination:** At each step of the process, the nontrivial direct sum decomposition of  $V^\bullet$  is finer than in the previous step. Since the vector spaces  $V^i$  are finite dimensional, this process must terminate eventually. Suppose that it terminates at the  $q^{\mathrm{th}}$  step to yield a  $q$ -weighted parabolic reduction  $(\vec{\lambda}, \mathcal{G}_{\vec{\lambda}})$ . We call this a Gieseker-Harder-Narasimhan (multi-weighted) parabolic reduction of  $(\mathcal{F}^\bullet, \sigma)$ . The corresponding  $q^{\mathrm{th}}$  associated graded Levi  $\rho_{\vec{\lambda}}$ -sheaf  $(\mathrm{gr}_q(\mathcal{F}^\bullet), \mathrm{gr}_q(\sigma))$  is called the Gieseker-Harder-Narasimhan associated Levi of  $(\mathcal{F}^\bullet, \sigma)$ . By construction,  $(\mathrm{gr}_q(\mathcal{F}^\bullet), \mathrm{gr}_q(\sigma))$  is Gieseker semistable. We call the lexicographic  $\mathbb{Z}^q$ -filtration  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  the (multi-weighted) Gieseker-Harder-Narasimhan filtration of the  $\rho$ -sheaf  $(\mathcal{F}^\bullet, \sigma)$ . This filtration is uniquely determined up to scaling the weights by a rational tuple  $\vec{p}$ , because of the uniqueness of the leading term HN filtration up to scaling at each step of the process.

This construction is recorded in the following definition.

**Definition 8.10.** *Suppose that the field extension  $K \supset k$  is algebraically closed and let  $(\mathcal{F}^\bullet, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . The Gieseker-Harder-Narasimhan filtration of  $(\mathcal{F}^\bullet, \sigma)$  is the unique (up to scaling of the weights) lexicographic  $\mathbb{Z}^q$ -filtration  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  of  $(\mathcal{F}^\bullet, \sigma)$  constructed above. Its Gieseker-Harder-Narasimhan associated Levi  $\rho_{\vec{\lambda}}$ -sheaf, as described above, is Gieseker semistable.*

*If  $K \supset k$  is not algebraically closed, then by uniqueness we can descend the GHN filtration of the base change to any algebraic closure of  $K$ . Therefore we obtain a unique (up to scaling the weights) lexicographic  $\mathbb{Z}^q$ -filtration  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$ , which we call the GHN filtration of  $(\mathcal{F}^\bullet, \sigma)$ .*

**Remark 8.11.** *In particular, when the  $\rho$ -sheaf  $(\mathcal{F}^\bullet, \sigma)$  is Gieseker semistable, the GHN filtration constructed above produces a trivial filtration as in Definition 4.32.*

**Remark 8.12.** *If  $G_K$  is split then, by construction, the GHN filtration  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  comes from a  $q$ -weighted parabolic reduction  $(\vec{\lambda}, \mathcal{G}_{\vec{\lambda}})$ .*

**Remark 8.13.** *Suppose in addition that  $\rho$  is central. Then by Proposition 10.11 the rational  $P_{\vec{\lambda}}$ -parabolic reduction  $\mathcal{G}_{\vec{\lambda}}$  is finer than the canonical slope parabolic reduction of the corresponding rational  $G$ -bundle, as in [AAB02] or [BG03, §7]. More precisely, the parabolic subgroup  $P_{\vec{\lambda}}$  is contained in the canonical slope parabolic subgroup  $P$  and*

<sup>7</sup>We entrust the reader with the precise formulation of what this means.

the rational  $P$ -parabolic reduction  $\mathcal{G}_P$  obtained from  $\mathcal{G}_{\vec{\lambda}}$  via extension of structure group  $P_{\vec{\lambda}} \subset P$  recovers the canonical slope reduction.

In the case of torsion-free sheaves (i.e. when  $G = \mathrm{GL}(V)$  and  $\rho$  is the identity), this filtration recovers the classical Gieseker-Harder-Narasimhan filtration. We prove this in Subsection 9.1, where we describe the procedure above in the case of torsion-free sheaves.

**8.4. The GHN filtration in families.** In this subsection we define a notion of relative GHN filtrations. We prove that relative GHN filtrations induce a stratification of  $\mathrm{Bun}_\rho(X)$  by locally closed substacks.

Let  $S$  be a  $k$ -scheme. Let  $(\mathcal{F}^\bullet, \sigma)$  be a  $\rho$ -sheaf on  $X_S$ .

**Definition 8.14.** A relative lexicographic  $\mathbb{Z}^q$ -filtration of the underlying tuple of sheaves  $\mathcal{F}^\bullet = (\mathcal{F}_i)_{i=1}^b$  is a sequence of tuples of subsheaves  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  satisfying the following conditions

- (1)  $\mathcal{F}_{\vec{m}}^\bullet \subset \mathcal{F}_{\vec{n}}^\bullet$  whenever  $\vec{m} > \vec{n}$ .
- (2) There exists some  $M_{max} \in \mathbb{Z}^q$  such that  $\mathcal{F}_{\vec{m}}^\bullet = 0$  for all  $\vec{m} > M_{max}$ . Similarly, there exists some  $M_{min} \in \mathbb{Z}^q$  such that  $\mathcal{F}_{\vec{m}}^\bullet = \mathcal{F}^\bullet$  for all  $\vec{m} < M_{min}$ .
- (3) For all  $\vec{m}$ , we have that  $\mathcal{F}^\bullet / \mathcal{F}_{\vec{m}}^\bullet$  is a tuple of families of torsion-free sheaves on  $X_S$ .

For any such filtration  $f = (\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$ , we can apply the multi-graded Rees construction in order to obtain a tuple  $\widetilde{\mathcal{F}}_f^\bullet$  of families of torsion-free sheaves on  $\Theta^q \times X_S$ . Let  $U$  be a big open subset of  $X_S$  where the associated graded  $\mathrm{gr}(f)$  is locally-free. One can use  $\mathbb{G}_m^q$ -equivariance to conclude that the restriction  $(\widetilde{\mathcal{F}}_f^\bullet)|_{\Theta^q \times U}$  is locally-free. In particular, it can be viewed as a  $\mathrm{GL}(V^\bullet)$  bundle. The section  $\sigma$  gives a  $G$ -reduction of structure group of  $(\widetilde{\mathcal{F}}_f^\bullet)|_{\widetilde{1} \times U}$ .

**Definition 8.15.** We say that  $f = (\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  is a relative lexicographic  $\mathbb{Z}^q$ -filtration of the  $\rho$ -sheaf  $(\mathcal{F}^\bullet, \sigma)$  if the  $G$ -reduction induced by  $\sigma$  extends (uniquely) to  $\Theta_k^q \times U$ .

We can also define relative  $q$ -weighted reductions for  $(\mathcal{F}^\bullet, \sigma)$  in a similar way as in the case when  $S = \mathrm{Spec}(K)$ . One can say as well when a relative lexicographic  $\mathbb{Z}^q$ -filtration of  $(\mathcal{F}^\bullet, \sigma)$  comes from a  $q$ -weighted parabolic reduction  $(\vec{\lambda}, \mathcal{G}_\lambda)$ , we omit the details.

**Definition 8.16.** Let  $S$  and  $(\mathcal{F}^\bullet, \sigma)$  be as above and let  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  be a lexicographic  $\mathbb{Z}^q$ -filtration of  $(\mathcal{F}^\bullet, \sigma)$ , where we might allow a different choice of  $q$  over each connected component of  $S$ . We say that  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  is a GHN filtration for  $(\mathcal{F}^\bullet, \sigma)$  if for all points  $s \in S$  the restriction  $(\mathcal{F}_{\vec{m}}^\bullet|_{X_s})_{\vec{m} \in \mathbb{Z}^q}$  is the GHN filtration of  $(\mathcal{F}^\bullet|_{X_s}, \sigma|_{X_s})$ .

**Definition 8.17.** Let  $H \rightarrow S$  be a locally closed stratification of  $S$ . We say that the  $S$ -scheme  $H$  is universal for GHN filtrations of  $(\mathcal{F}^\bullet, \sigma)$  if it represents the functor that classifies pairs  $(T, [(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}])$ , where  $T$  is a  $S$ -scheme and  $[(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}]$  is an equivalence class of relative GHN filtrations of  $(\mathcal{F}^\bullet|_{X_T}, \sigma|_{X_T})$ , up to scaling the weights.

**Theorem 8.18.** The relative GHN filtrations induce a stratification of  $\mathrm{Bun}_\rho(X)$  by locally closed substacks. Equivalently, for any  $k$ -scheme  $S$  and every  $\rho$ -sheaf  $(\mathcal{F}^\bullet, \sigma)$  there exists a unique locally closed stratification  $H$  of  $S$  that is universal for GHN filtrations of  $(\mathcal{F}^\bullet, \sigma)$ .

If, in addition,  $S$  is locally of finite type over a field extension  $K \supset k$  such that  $G_K$  is split, then the restriction of the universal relative GHN filtration<sup>8</sup> to each connected component of the stratification comes from a relative multi-weighted parabolic reduction.

*Proof.* Let  $S$  be a  $k$ -scheme and let  $(\mathcal{F}^\bullet, \sigma)$  be a  $\rho$ -sheaf on  $X_S$ . We will denote by  $\mathrm{RelGies}(\mathcal{F}^\bullet, \sigma)$  the functor that parametrizes pairs  $(T, [(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}])$ , where  $T$  is a  $S$ -scheme

<sup>8</sup>More precisely, we choose a representative up to scaling.

and  $[(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}]$  is an equivalence class of relative GHN filtrations of  $(\mathcal{F}^\bullet|_{X_T}, \sigma|_{X_T})$  up to scaling the weights. We would like to show that  $\text{RelGies}(\mathcal{F}^\bullet, \sigma)$  is represented by a locally closed stratification of  $S$ . Since locally closed immersions satisfy étale descent, we can check this after base-changing to a field extension  $K \supset k$  such that  $G_K$  is split. After replacing  $k$  with  $K$ , we can assume without loss of generality that  $G$  is split. Moreover, since  $\text{Bun}_\rho(X)$  is locally of finite type over  $k$ , we can reduce to the case when  $S$  is of finite type over  $k$ . We will construct  $H$  recursively.

First Step: We denote by  $H_1$  the locally closed stratification of  $S$  that is universal for leading term HN filtrations of  $(\mathcal{F}^\bullet, \sigma)$  (Proposition 4.37). There is a natural forgetful functor  $\text{RelGies}(\mathcal{F}^\bullet, \sigma) \rightarrow H_1$  given by sending a relative GHN filtration  $(\mathcal{F}_{\vec{m}}^\bullet)_{\vec{m} \in \mathbb{Z}^q}$  to the relative leading term HN filtration  $(\mathcal{F}_m^\bullet)_{m \in \mathbb{Z}}$  defined by

$$\mathcal{F}_m^\bullet := \bigcup_{(m_2, m_3, \dots, m_q) \in \mathbb{Z}^{q-1}} \mathcal{F}_{(m, m_2, \dots, m_q)}$$

The fact that this is a relative leading term HN filtration follows from the pointwise construction of the GHN filtration and the fact that pulling back to a fiber commutes with taking unions.

Choose a representative  $f_1$  for the universal leading term HN filtration of  $(\mathcal{F}^\bullet|_{X_{H_1}}, \sigma|_{X_{H_1}})$ . By the last statement in Proposition 4.37, the restriction of  $f_1$  to each component of  $H_1$  comes from a relative weighted parabolic reduction  $(\lambda_1, \mathcal{G}_{\lambda_1})$ . We can use this to define, on each component of  $H_1$ , a relative 1<sup>st</sup> associated graded Levi  $\rho_{\lambda_1}$ -sheaf, in exactly the same way as in the construction of the GHN filtration in Subsection 8.3. Putting all of these together for all connected components, we obtain what we can call a relative 1<sup>st</sup> associated graded Levi  $\rho_{\lambda_1}$ -sheaf  $\text{gr}_1 := \text{gr}_1(\mathcal{F}^\bullet|_{X_{H_1}}, \sigma|_{X_{H_1}})$  defined on  $H_1$ , where we allow a different choice of the cocharacter  $\lambda_1$  over each connected component of  $H_1$ .

Second Step: We can apply Proposition 4.37 to the restriction of  $\text{gr}_1$  on each connected component in order to obtain a refinement  $H_2$  of the stratification  $H_1$ . The scheme  $H_2$  is universal for the leading term HN filtration of  $\text{gr}_1$ , hence we can choose a representative  $f_2$  for the universal filtration. By the last statement of Proposition 4.37, the restriction of  $f_2$  to each connected component of  $H_2$  comes from a weighted reduction  $(\lambda_2, \mathcal{G}_{\lambda_2})$  of  $\text{gr}_1$ . The procedure to construct the GHN filtration can be applied verbatim to this relative situation. In particular, we can construct a relative lexicographic  $\mathbb{Z}^2$ -filtration of  $(\mathcal{F}^\bullet|_{X_{H_2}}, \sigma|_{X_{H_2}})$ . This lexicographic filtration comes from a 2-weighted parabolic reduction  $((\lambda_1, \lambda_2), \mathcal{G}_{(\lambda_1, \lambda_2)})$ . By construction,  $H_2$  is universal for lexicographic  $\mathbb{Z}^2$ -filtrations obtained in the second step of the construction of the GHN filtration. We can define a forgetful morphism  $\text{RelGies}(\mathcal{F}^\bullet, \sigma) \rightarrow H_2$  in the same way as in the case of the first step  $H_1$ . In this case we remember the first two coordinates of the  $\mathbb{Z}^q$ -filtration and take the union over the other coordinates.

Recursion: Now we can iterate as in the construction of the (pointwise) GHN filtration. This process must terminate by the same reasoning as in Subsection 8.3, namely, the dimensions of the vector spaces in the tuple  $V^\bullet$  are finite. If it terminates at the  $q^{\text{th}}$  step, then the corresponding locally closed stratification  $H_q$  is universal for GHN filtrations of  $(\mathcal{F}^\bullet, \sigma)$ . Furthermore, on each connected component of  $H_q$  the restriction of (a representative up to scaling of) the universal GHN filtration comes from a relative multi-weighted parabolic reduction.  $\square$

**Remark 8.19.** *It follows from the proof of Theorem 8.18 that this stratification of the stack is obtained by recursively taking intersections of certain  $\Theta$ -stratifications. At each step we are using the leading term HN  $\Theta$ -stratification defined on what is called “the center” of the  $\Theta$ -stratum in the previous step. In this case each center can be thought of as  $\text{Bun}_{\rho_{(\lambda_1, \lambda_2, \dots, \lambda_i)}}(X)$  for some representation  $\rho_{(\lambda_1, \lambda_2, \dots, \lambda_i)}$  of a Levi subgroup. This is what allows the iterative procedure.*

## 9. EXAMPLES

In this section we give some examples of the theory developed.

**9.1. Torsion-free sheaves.** Suppose that  $G = \mathrm{GL}(V)$  and  $\rho$  is the identity. Then  $\mathrm{Bun}_\rho(X)$  is equivalent to the stack  $\mathrm{Coh}_r^{tf}(X)$  of torsion-free sheaves of rank  $r$  on  $X$ . In this case Proposition 3.8 gives a description of a filtration of a torsion-free sheaf  $\mathcal{F}$  in  $\mathrm{Bun}_\rho(X)$  as a decreasing  $\mathbb{Z}$ -filtration of  $\mathcal{F}$  by saturated subsheaves  $\mathcal{F}_m \subset \mathcal{F}$ . The proof of [HLHJ21, Prop. 5.16] shows that a torsion-free sheaf in  $\mathrm{Bun}_\rho(X)$  is Gieseker semistable in our sense if and only if it is Gieseker semistable in the sense of [HL97, Defn. 1.2.4].

The leading term HN filtration for torsion-free sheaves in  $\mathrm{Bun}_\rho(X)$  recovers the weighted leading term filtration described in [HLHJ21, Thm. 5.11], which we reproduce here. Let

$$0 = \mathcal{F}_0^{\mathrm{HN}} \subset \mathcal{F}_1^{\mathrm{HN}} \subset \mathcal{F}_2^{\mathrm{HN}} \subset \dots \subset \mathcal{F}_n^{\mathrm{HN}} = \mathcal{F}$$

denote the Gieseker-Harder-Narasimhan filtration of  $\mathcal{F}$  (c.f. [HL97, Thm. 1.3.4]). Then we have

$$\bar{\rho}_{\mathcal{F}_1^{\mathrm{HN}}} > \bar{\rho}_{\mathcal{F}_2^{\mathrm{HN}}/\mathcal{F}_1^{\mathrm{HN}}} > \dots > \bar{\rho}_{\mathcal{F}_n^{\mathrm{HN}}/\mathcal{F}_{n-1}^{\mathrm{HN}}}$$

Let  $i$  denote the minimal index such that

$$\hat{\mu}_h(\mathcal{F}_1^{\mathrm{HN}}) = \hat{\mu}_h(\mathcal{F}_2^{\mathrm{HN}}/\mathcal{F}_1^{\mathrm{HN}}) = \dots = \hat{\mu}_h(\mathcal{F}_n^{\mathrm{HN}}/\mathcal{F}_{n-1}^{\mathrm{HN}})$$

for all  $h > i$ . In other words,  $i$  is the largest integer such that the  $i^{\mathrm{th}}$  slopes of the graded pieces of the filtration are not all equal. We have

$$\hat{\mu}_i(\mathcal{F}_1^{\mathrm{HN}}) \geq \hat{\mu}_i(\mathcal{F}_2^{\mathrm{HN}}/\mathcal{F}_1^{\mathrm{HN}}) \geq \dots \geq \hat{\mu}_i(\mathcal{F}_n^{\mathrm{HN}}/\mathcal{F}_{n-1}^{\mathrm{HN}})$$

where at least one of the inequalities above is strict.

We define indexes  $h_0 < h_1 < h_2 < \dots < h_j = n$  as follows. We let  $h_0 = 0$ , and let

$$\{h_m\}_{m=1}^j = \{1 \leq l \leq n \mid \hat{\mu}_i(\mathcal{F}_{l+1}^{\mathrm{HN}}/\mathcal{F}_l^{\mathrm{HN}}) < \hat{\mu}_i(\mathcal{F}_l^{\mathrm{HN}}/\mathcal{F}_{l-1}^{\mathrm{HN}})\}$$

This ensures that  $\hat{\mu}_i(\mathcal{F}_{h_k}^{\mathrm{HN}}/\mathcal{F}_{h_{k-1}}^{\mathrm{HN}}) > \hat{\mu}_i(\mathcal{F}_{h_{k+1}}^{\mathrm{HN}}/\mathcal{F}_{h_k}^{\mathrm{HN}})$ . For each  $k$ , we set  $\mathcal{F}_k^{\mathrm{l-term}} = \mathcal{F}_{h_k}^{\mathrm{HN}}$ .

**Definition 9.1.** *The filtration*

$$0 = \mathcal{F}_0^{\mathrm{l-term}} \subset \mathcal{F}_1^{\mathrm{l-term}} \subset \mathcal{F}_2^{\mathrm{l-term}} \subset \dots \subset \mathcal{F}_j^{\mathrm{l-term}} = \mathcal{F}$$

is called the unweighted leading term filtration of  $\mathcal{F}$ . We call  $i$  the leading term index.

We remark that, if  $\mathcal{F}$  is slope unstable, then  $i = d - 1$  (recall that  $\hat{\mu}_{d-1}(\mathcal{F})$  is the slope of  $\mathcal{F}$ ), and the leading term filtration is the slope-Harder-Narasimhan filtration.

**Definition 9.2.** *Let  $\mathcal{F} \in \mathrm{Bun}_\rho(X) = \mathrm{Coh}_r^{tf}(X)$  and assume that  $\mathcal{F}$  is unstable. Suppose that the unweighted leading term filtration for  $\mathcal{F}$  is given by*

$$0 = \mathcal{F}_0^{\mathrm{l-term}} \subset \mathcal{F}_1^{\mathrm{l-term}} \subset \mathcal{F}_2^{\mathrm{l-term}} \subset \dots \subset \mathcal{F}_j^{\mathrm{l-term}} = \mathcal{F}$$

with leading slope  $\hat{\mu}_i$ . Set  $\mu = \hat{\mu}_i(\mathcal{F})$ ,  $\mu_k = \hat{\mu}_i(\mathcal{F}_k^{\mathrm{l-term}}/\mathcal{F}_{k-1}^{\mathrm{l-term}})$ . Choose a positive rational number  $L$  such that  $L \cdot \mu \in \mathbb{Z}$  and  $L \cdot \mu_k \in \mathbb{Z}$  for all  $k$ .

Then, we define the weighted leading term filtration  $(\mathcal{F}^{\mathrm{Can}})_{m \in \mathbb{Z}}$  as follows (up to scaling of the weights).

- (1)  $\mathcal{F}_m^{\mathrm{Can}} = 0$  for  $m > L(\mu_1 - \mu)$ .
- (2) Let  $1 \leq k \leq j - 1$ . For each  $L(\mu_k - \mu) \geq m > L(\mu_{k+1} - \mu)$ , we set  $\mathcal{F}_m^{\mathrm{Can}} = \mathcal{F}_k^{\mathrm{l-term}}$ .
- (3)  $\mathcal{F}_m^{\mathrm{Can}} = \mathcal{F}$  for  $L(\mu_j - \mu) \geq m$ .

In other words, the canonical filtration is given by the leading term filtration and the  $k^{\mathrm{th}}$  jump happens at the integer  $L(\mu_k - \mu)$ .

We end this subsection by explicitly describing the procedure for the construction of the GHN filtration in this case. Let  $0 = \mathcal{F}_0^{\text{l-term}} \subset \mathcal{F}_1^{\text{l-term}} \subset \mathcal{F}_2^{\text{l-term}} \subset \dots \subset \mathcal{F}_{j_1}^{\text{l-term}} = \mathcal{F}$  be the unweighted leading term filtration, with leading term index  $i_1$ . By assigning weights as above, this yields the leading term HN filtration  $({}^{(1)}\mathcal{F}_m)_{m \in \mathbb{Z}}$  of  $\mathcal{F}$ . The 1<sup>st</sup> associated graded Levi  $\rho_{\lambda_1}$ -sheaf consists of the tuple

$$\text{gr}_1(\mathcal{F}) = \left( \mathcal{F}_i^{\text{l-term}} / \mathcal{F}_{i-1}^{\text{l-term}} \right)_{i=1}^{j_1}$$

The leading term HN filtration of  $\text{gr}_1(\mathcal{F})$  is obtained by taking the leading term filtration of each element  $\mathcal{F}_i^{\text{l-term}} / \mathcal{F}_{i-1}^{\text{l-term}}$  in the tuple. Each corresponding leading term index will be strictly smaller than  $i_1$ . The sheaves appearing in the leading term filtration of  $\mathcal{F}_i^{\text{l-term}} / \mathcal{F}_{i-1}^{\text{l-term}}$  are of the form  $\mathcal{F}_h^{\text{HN}} / \mathcal{F}_{i-1}^{\text{l-term}}$  for some index  $1 \leq h \leq n$ . We denote the corresponding lexicographic  $\mathbb{Z}^2$ -filtration of  $\mathcal{F}$  by  $({}^{(2)}\mathcal{F}_{\vec{m}})_{\vec{m} \in \mathbb{Z}^2}$ . Let  $\vec{m}_1 > \vec{m}_2 > \dots > \vec{m}_{j_2}$  denote the jumping points of  $({}^{(2)}\mathcal{F}_{\vec{m}})_{\vec{m} \in \mathbb{Z}^2}$ . The unweighted filtration

$$0 \subset ({}^{(2)}\mathcal{F}_{\vec{m}_1}) \subset ({}^{(2)}\mathcal{F}_{\vec{m}_2}) \subset \dots \subset ({}^{(2)}\mathcal{F}_{\vec{m}_{j_2}}) = \mathcal{F}$$

is coarser than the Gieseker-Harder-Narasimhan filtration, but finer than the leading term filtration. The reduced Hilbert polynomials of the graded pieces are strictly increasing, and each graded piece  $({}^{(2)}\mathcal{F}_{\vec{m}_i}) / ({}^{(2)}\mathcal{F}_{\vec{m}_{i-1}})$  is semistable up to a degree  $i_2$  in the Hilbert polynomial that is strictly smaller than the degree  $i_1$  in the previous step. The 2<sup>nd</sup> associated graded Levi sheaf is  $\text{gr}_2(\mathcal{F}) = \left( ({}^{(2)}\mathcal{F}_{\vec{m}_i}) / ({}^{(2)}\mathcal{F}_{\vec{m}_{i-1}}) \right)_{i=1}^{j_2}$ . We iterate this procedure, at each step we take the leading term filtration of the graded pieces.

Suppose that this process terminates at the  $q^{\text{th}}$  step. Let  $({}^{(q)}\mathcal{F}_{\vec{m}})_{\vec{m} \in \mathbb{Z}^q}$  be the GHN filtration obtained at this  $q^{\text{th}}$  step. Denote by  $\vec{m}_1 > \vec{m}_2 > \dots > \vec{m}_{j_q}$  the jumping points of the GHN filtration. Consider the unweighted filtration

$$(14) \quad 0 \subset ({}^{(q)}\mathcal{F}_{\vec{m}_1}) \subset ({}^{(q)}\mathcal{F}_{\vec{m}_2}) \subset \dots \subset ({}^{(q)}\mathcal{F}_{\vec{m}_{j_q}}) = \mathcal{F}.$$

The reduced Hilbert polynomials of the graded pieces of this filtration are strictly increasing by construction. Moreover, each graded piece must be Gieseker semistable, because the  $q^{\text{th}}$  step is the last one of the procedure. Hence this filtration is the classical Gieseker-Harder-Narasimhan filtration of  $\mathcal{F}$ . The associated  $q$ -weights in the GHN filtration encode information about how high is the degree of the difference of reduced Hilbert polynomials between two consecutive graded pieces.

We have therefore shown the following.

**Proposition 9.3.** *Let  $\mathcal{F}$  be a torsion-free sheaf. Let  $({}^{(q)}\mathcal{F}_{\vec{m}})_{\vec{m} \in \mathbb{Z}^q}$  be the GHN filtration. The corresponding unweighted filtration (14) coincides with the classical Gieseker-Harder-Narasimhan filtration of  $\mathcal{F}$ .  $\square$*

**9.2. Example of Gieseker-Harder-Narasimhan filtration.** We compare the different filtrations that we have considered in an example. Let  $G = \text{SO}(7)$  and let  $\rho : \text{SO}(7) \rightarrow \text{GL}(7)$  be the standard representation. Set  $X = \mathbb{P}^3$ , and let  $L \cong \mathbb{P}^1 \subset \mathbb{P}^3$  be a fixed line. Choose two different points  $Z = \{p, p'\} \subset \mathbb{P}^3$ . We denote by  $\mathcal{I}_L$  the ideal sheaf of  $L$ ,  $\mathcal{I}_p$  the ideal sheaf of  $p$ , and  $\mathcal{I}_Z$  the ideal sheaf of  $Z$ . Set

$$\mathcal{F} = \mathcal{I}_L \oplus \mathcal{I}_Z \oplus \mathcal{I}_p \oplus \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}.$$



Let  $\varphi : \mathcal{F} \otimes \mathcal{F} \rightarrow \mathcal{O}$  be the symmetric bilinear form defined by the matrix

$$\begin{pmatrix} 0 & \cdots & & & & & 1 \\ \vdots & & & & & & \\ & & & & & 1 & \\ & & & & 1 & & \\ & & & 1 & & & \\ & & 1 & & & & \\ & 1 & & & & & \vdots \\ 1 & & & & \cdots & & 0 \end{pmatrix}$$

Set  $U = \mathbb{P}^3 \setminus (L \cup Z)$ . The sheaf  $\mathcal{F}|_U$  is locally free, and in fact trivial. Let  $\varphi'$  be the canonical trivialization of  $\det \mathcal{F}$ . Clearly  $\det \varphi = \varphi'^2$ , hence the pair  $(\varphi, \varphi')$  induces a reduction of the trivial  $\mathrm{GL}(7)$ -bundle to  $\mathrm{SO}(7)$ . The rational  $\mathrm{SO}(7)$ -bundle  $\mathcal{G} \rightarrow U$  which it defines is trivial. Let  $\psi$  denote the canonical isomorphism between  $\rho_*(\mathcal{G})$  and  $\mathcal{F}|_U$ .

The principal  $\rho$ -sheaf  $(\mathcal{G}, \mathcal{F}, \psi)$  is slope semistable (Definition 10.7). The rational principal  $\mathrm{SO}(7)$ -bundle  $\mathcal{G}$  is trivial, hence also slope semistable (Definition 10.3). Hence the Harder-Narasimhan canonical reduction defined in [AAB02] is trivial.

The Hilbert polynomials are

$$\begin{aligned} P_{\mathcal{O}}(n) &= \frac{1}{6}(n^3 + 6n^2 + 10n + 3) \\ P_{\mathcal{I}_p}(n) &= \frac{1}{6}(n^3 + 6n^2 + 10n - 3) \\ P_{\mathcal{I}_Z}(n) &= \frac{1}{6}(n^3 + 6n^2 + 10n - 9) \\ P_{\mathcal{I}_L}(n) &= \frac{1}{6}(n^3 + 6n^2 - 6n - 3) \end{aligned}$$

Note that the three first polynomials only differ in the last term, and the last polynomial differs in the last two terms.

The principal  $\rho$ -sheaf  $(\mathcal{G}, \mathcal{F}, \psi)$  is Gieseker unstable (Definition 10.2). We will now calculate the leading term HN filtration. We work in the standard basis where the bilinear form is given by the matrix above. We will use cocharacters of  $\mathrm{SO}(7)$  of the form

$$\mathrm{diag}(t^a, t^b, t^c, 1, t^{-c}, t^{-b}, t^{-a})$$

with  $a \geq b \geq c \geq 0$ . The associated filtration  $\mathcal{F}_m$  satisfies  $\mathcal{F}_m^\perp = \mathcal{F}_{-m+1}$ . Recall the definition of orthogonal of a subsheaf  $\mathcal{F}_i \subset \mathcal{F}$

$$\mathcal{F}_i^\perp := \ker(\mathcal{F} \xrightarrow{\varphi} \mathcal{F}^\vee \rightarrow \mathcal{F}_i^\vee).$$

The leading term HN filtration is

$$\underbrace{\mathcal{F}}_{\mathcal{F}_{-1}} \supset \overbrace{\mathcal{I}_Z \oplus \mathcal{I}_p \oplus \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}}^{\mathcal{F}_0} \supset \underbrace{\mathcal{O}}_{\mathcal{F}_1}$$

where  $\mathcal{F}_1$  is the last summand in  $\mathcal{F}$ , and  $\mathcal{F}_1^\perp = \mathcal{F}_0$ . By definition of leading term HN filtration, this is the unique filtration, with  $\mathcal{F}_m^\perp = \mathcal{F}_{-m+1}$ , which gives a maximum of the numerical function (3).

This filtration has been obtained by the cocharacter  $\lambda_1 = \mathrm{diag}(t, 1, 1, 1, 1, 1, t^{-1})$ . The associated Levi is  $L_{\lambda_1} = \mathbb{G}_m \times \mathrm{SO}(5)$ , the representation  $\rho_{\lambda_1}$  is

$$\begin{aligned} \rho_{\lambda_1} : \mathbb{G}_m \times \mathrm{SO}(5) &\longrightarrow \mathrm{GL}(1) \times \mathrm{GL}(5) \times \mathrm{GL}(1) \\ (t, g) &\longmapsto (t, g, t^{-1}) \end{aligned}$$

and the associated graded sheaf is

$$\underbrace{\mathcal{I}_L}_{\mathcal{F}^{-1} := \mathcal{F}_{-1}/\mathcal{F}_0} \oplus \overbrace{\mathcal{I}_Z \oplus \mathcal{I}_p \oplus \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}}^{\mathcal{F}^0 := \mathcal{F}_0/\mathcal{F}_{-1}} \oplus \underbrace{\mathcal{O}}_{\mathcal{F}^1 := \mathcal{F}_1}.$$

The  $\mathrm{SO}(5)$ -sheaf defined by  $\mathcal{F}^0$  is not Gieseker stable. Its leading term HN filtration is given by the cocharacter  $\lambda_2 = \mathrm{diag}(t^2, t, 1, t^{-1}, t^{-2})$  and it is

$$\underbrace{\mathcal{F}^0}_{\mathcal{F}^{-2}} \supset \underbrace{\mathcal{I}_p \oplus \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}}_{\mathcal{F}^{-1}} \supset \underbrace{\mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}}_{\mathcal{F}_0^0} \supset \underbrace{\mathcal{O} \oplus \mathcal{O}}_{\mathcal{F}_1^0} \supset \underbrace{\mathcal{O}}_{\mathcal{F}_2^0}$$

Finally, the (multi-weighted) Gieseker-Harder-Narasimhan filtration is

$$\underbrace{\mathcal{F}^{(-1,0)}}_{\mathcal{F}} \supset \underbrace{\mathcal{I}_Z \oplus \mathcal{I}_p \oplus \mathcal{O}^{\oplus 4}}_{\mathcal{F}^{(0,-2)}} \supset \underbrace{\mathcal{I}_p \oplus \mathcal{O}^{\oplus 4}}_{\mathcal{F}^{(0,-1)}} \supset \underbrace{\mathcal{O}^{\oplus 4}}_{\mathcal{F}^{(0,0)}} \supset \underbrace{\mathcal{O}^{\oplus 3}}_{\mathcal{F}^{(0,1)}} \supset \underbrace{\mathcal{O}^{\oplus 2}}_{\mathcal{F}^{(0,2)}} \supset \underbrace{\mathcal{O}}_{\mathcal{F}^{(1,0)}}$$

The process stops here because the associated Levi principal sheaf is Gieseker semistable. Indeed, the Levi is  $L_{\lambda_2} = \mathbb{G}_m \times (\mathbb{G}_m \times \mathbb{G}_m) \subset \mathbb{G}_m \times \mathrm{SO}(5)$ , the representation is

$$\begin{aligned} \rho_{\lambda_2} : \mathbb{G}_m \times (\mathbb{G}_m \times \mathbb{G}_m) &\longrightarrow \mathrm{GL}(1)^{\times 7} \\ (a, b, c) &\mapsto (a, b, c, 1, c^{-1}, b^{-1}, a^{-1}) \end{aligned}$$

and the associated graded is isomorphic to  $\mathcal{F}$ .

## 10. COMPARISON WITH PREVIOUS WORK

In this section we compare our notions of stability and Harder-Narasimhan filtrations of principal  $\rho$ -sheaves with other definitions in the literature.

**10.1. Comparison with previous notions of semistability.** The problem of finding a compactification of the moduli space by introducing singular degenerations of principal bundles has already been considered in the literature, and the objects have been called (honest) singular principal bundles, principal sheaves or principal  $\rho$ -sheaves ([Sch02, GS03, GS05, GLSS08, GLSS10]). All notions fit the same framework: we fix a representation  $\rho$  of  $G$  (sometimes  $\rho$  is not required to be injective) and the objects are triples consisting of a principal  $G$ -bundle  $P$  on a big open subset  $U \subset X$ , a torsion free sheaf  $\mathcal{E}$  on  $X$ , and an isomorphism  $\psi$  on  $U$  between the vector bundle associated to  $P$  using the representation  $\rho$  and the restriction  $\mathcal{E}|_U$ .

Unfortunately, the names are not always the same (for instance, singular principal bundles in [GLSS08] were called honest singular principal bundles in [Sch02]), and the vector bundle that appears in the isomorphism  $\psi$  is sometimes  $P(\rho^\vee)$ , using the dual of the representation  $\rho$  (in [GLSS08, Sch02, GLSS10]) and sometimes  $P(\rho)$ , using directly the representation (in [GS05, GS03]). In particular, we warn the reader that, in [GLSS10, §1.2], a *principal  $\rho$ -sheaf* was defined using the dual representation, and  $\rho$  was not required to be injective. In this subsection we will see that our definition of Gieseker semistability for principal  $\rho$ -sheaves coincides with the definitions that have already appeared.

Let us recall some of the main definitions of this article.

**Definition 10.1.** *Let  $G$  be a connected reductive group. Let*

$$\rho = (\rho_1, \dots, \rho_b) : G \longrightarrow \mathrm{GL}(V^1) \times \dots \times \mathrm{GL}(V^b)$$

*be an injective homomorphism. Let  $S$  be a  $k$ -scheme. A principal  $\rho$ -sheaf on  $X_S$  is a triple  $(\mathcal{G}, \mathcal{F}^\bullet, \psi^\bullet)$  where  $\mathcal{G}$  is a principal  $G$ -bundle on a big open set  $U \subset X_S$ ,  $\mathcal{F}^\bullet = (\mathcal{F}^i)_{i=1}^b$  is a tuple of torsion free sheaves on  $X_S$ , and  $\psi^\bullet = (\psi^i)_{i=1}^b$  is a tuple of isomorphisms  $(\rho_i)_* \mathcal{G} \cong \mathcal{F}^i|_U$ .*

We generalize Definition 4.28 in the following way, using the summation by parts argument as in Equation (12).

**Definition 10.2.** Let  $K \supset k$  be an algebraically closed field. A principal  $\rho$ -sheaf on  $X_K$  is Gieseker semistable if for all cocharacters  $\lambda : (\mathbb{G}_m)_K \rightarrow G_K$  and all weighted parabolic reductions  $(\lambda, \mathcal{G}_\lambda)$  with associated underlying filtrations  $(\mathcal{F}_m^i)_{m \in \mathbb{Z}}$  of  $\mathcal{F}^i$ , (as in Definition 3.15) we have

$$(15) \quad \sum_{i=1}^b \sum_{m \in \mathbb{Z}} (\mathrm{rk}(\mathcal{F}^i) \cdot P_{\mathcal{F}_m^i} - \mathrm{rk}(\mathcal{F}_m^i) \cdot P_{\mathcal{F}^i}) \leq 0$$

We start by comparing our notion of Gieseker semistability with Ramanathan's notion of slope semistability for the underlying rational  $G$ -bundle.

**Definition 10.3** ([Ram79]). A rational  $G$ -bundle on a polarized smooth projective variety  $X$  is a  $G$ -bundle  $\mathcal{G}$  on a big open set  $U \subset X$ . It is called slope stable (respectively, semistable) if for all reductions of structure group  $\mathcal{G}_P \rightarrow U' \subset U$  to a proper parabolic subgroup  $P \subset G$  on a big open subset  $U'$  and all dominant characters  $\chi$  of  $P$ , the associated line bundle  $\chi_*(\mathcal{G}_P)$  satisfies

$$\mathrm{deg} \chi_*(\mathcal{G}_P) < 0 \quad (\text{respectively, } \leq)$$

Here the degree is defined with respect to the fixed polarization. A rational  $G$ -bundle is called slope unstable if it is not slope semistable.

**Proposition 10.4.** Let  $K \supset k$  be a field extension. Let  $(\mathcal{F}^\bullet, \sigma)$  be a  $\rho$ -sheaf on  $X_K$ . Suppose that  $U$  is a big open subset of  $X_K$  such that the tuple  $\mathcal{F}^\bullet|_U$  is locally-free. Let  $\mathcal{G}$  denote the  $G$ -bundle on  $U$  corresponding to  $(\mathcal{F}^\bullet, \sigma)$ . If  $(\mathcal{F}^\bullet, \sigma)$  is Gieseker semistable, then the rational  $G$ -bundle  $\mathcal{G}$  is slope semistable in the sense of [Ram79].

*Proof.* We will freely use the notation from the proof of Proposition 4.26 and assume, after replacing  $K$  with a field extension, that  $K$  is algebraically closed. Suppose that the rational  $G$ -bundle  $\mathcal{G}$  is not slope semistable. Then there exists a proper parabolic subgroup  $P_\lambda \subset G$  and a canonical (slope)  $P_\lambda$ -reduction of structure group  $\mathcal{G}_{P_\lambda}$  defined over a big open subset  $Q$  contained in  $U$ , as in [AAB02]. After replacing  $U$  with the smaller subset  $Q$ , we can assume without loss of generality that  $\mathcal{G}_{P_\lambda}$  is defined over  $U$ .

Let  $\mathcal{G}_{L_\lambda}$  be the  $L_\lambda$ -bundle associated to the  $P_\lambda$ -bundle  $\mathcal{G}_\lambda$  via the quotient  $P_\lambda \twoheadrightarrow L_\lambda$ . Let  $Z_\lambda$  denote the center of  $P_\lambda$ . The rational  $L_\lambda$ -bundle  $\mathcal{G}_{L_\lambda}$  determines a linear functional  $\psi : X^*(Z_\lambda)_\mathbb{R} \rightarrow \mathbb{R}$  as in the proof of Lemma 4.25. Since  $\psi$  actually lies in  $X_*(Z_\lambda)_\mathbb{Q}$ , there is a multiple  $N\psi$  that lies in the lattice of cocharacters  $X_*(Z_\lambda)$ . By the properties of the canonical slope reduction [AAB02],  $\psi$  lies in the interior of the cone of  $P_\lambda$ -dominant coweights. It follows that  $P_\lambda = P_{(N\psi)}$ . Therefore  $f = (N\psi, \mathcal{G}_{P_\lambda})$  is a nondegenerate weighted parabolic reduction of  $(\mathcal{F}, \sigma)$ .

The group  $G$  acts on each vector space  $V^j$  via the homomorphism  $\rho_j$ . For each  $j$ , we denote by  $(\chi_{i,j})_{i=1}^{r_j}$  the tuple of weights of the split maximal torus  $T$  acting on  $V^j$  (with possible repetitions). Recall the function  $\nu_{d-1}$  on the dominant cone  $\mathcal{C}_\lambda$  described in Lemma 4.23 (4). In this case the formula (4) includes an extra summation

$$(16) \quad \nu_{d-1}(\delta) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\sum_{j=1}^b \sum_{i=1}^{r_j} \langle \delta, \chi_{i,j}^\lambda \rangle \cdot (c_{i,j} - c_j)}{\sqrt{\sum_{j=1}^b \sum_{i=1}^{r_j} (\langle \delta, \chi_{i,j}^\lambda \rangle)^2}}$$

where we define  $c_{i,j} = \langle \psi, \chi_{i,j}^\lambda \rangle$  and  $c_j = \frac{1}{r_j} \langle \psi, \sum_{i=1}^{r_j} \chi_{i,j}^\lambda \rangle$ . Expression (16) for  $\delta = N\psi$  yields

$$\nu_{d-1}(N\psi) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\sum_{j=1}^b \sum_{i=1}^{r_j} \langle N\psi, \chi_{i,j}^\lambda \rangle \cdot \left( \langle \psi, \chi_{i,j}^\lambda \rangle - \frac{1}{r_j} \sum_{l=1}^{r_j} \langle \psi, \chi_{i,l}^\lambda \rangle \right)}{\sqrt{\sum_{j=1}^b \sum_{i=1}^{r_j} (\langle N\psi, \chi_{i,j}^\lambda \rangle)^2}} =$$

$$(17) \quad \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\sum_{j=1}^b \left( \sum_{i=1}^{r_j} \left( \langle \psi, \chi_{i,j}^\lambda \rangle \right)^2 \right) - \frac{1}{r_j} \left( \sum_{i=1}^{r_j} \langle \psi, \chi_{i,j}^\lambda \rangle \right) \cdot \left( \sum_{l=1}^{r_j} \langle \psi, \chi_{l,j}^\lambda \rangle \right)}{\sqrt{\sum_{j=1}^b \sum_{i=1}^{r_j} \left( \langle N\psi, \chi_{i,j}^\lambda \rangle \right)^2}}$$

We shall show that the numerator of (17) is positive. This will contradict Gieseker semistability of  $(\mathcal{F}, \sigma)$ . For each index  $j$ , let  $\vec{v}_j := [\langle \psi, \chi_{i,j}^\lambda \rangle]_{i=1}^{r_j} \in \mathbb{R}^{r_j}$  and set  $\vec{1}_j \in \mathbb{R}^{r_j}$  to be the vector with all coordinates equal to 1. The numerator in (17) is

$$\sum_{j=1}^b \left( \sum_{i=1}^{r_j} \left( \langle \psi, \chi_{i,j}^\lambda \rangle \right)^2 \right) - \frac{1}{r_j} \left( \sum_{i=1}^{r_j} \langle \psi, \chi_{i,j}^\lambda \rangle \right)^2 = \sum_{j=1}^b \frac{1}{r_j} \left[ \left( \vec{1}_j \cdot_{\mathbb{R}} \vec{1}_j \right) \left( \vec{v}_j \cdot_{\mathbb{R}} \vec{v}_j \right) - \left( \vec{1}_j \cdot_{\mathbb{R}} \vec{v}_j \right)^2 \right]$$

where  $\cdot_{\mathbb{R}}$  denotes the standard inner product on  $\mathbb{R}^r$ . The Cauchy-Schwarz inequality implies that the quantity above is always nonnegative and, moreover, it is strictly positive if there exists some index  $j$  such that  $\vec{1}_j$  and  $\vec{v}_j$  are not scalar multiples of each other. Therefore,  $\nu_{d-1}(N\psi) > 0$  whenever  $\langle \psi, \chi_{i,j}^\lambda \rangle \neq \langle \psi, \chi_{l,j}^\lambda \rangle$  for some  $j$  and  $i \neq l$ . We are left to show that there are  $j$  and two indexes  $i \neq l$  satisfying this condition.

Let  $\{\zeta_{l,j}\}$  denote the subset of  $\{\chi_{i,j}\}$  consisting of  $B$ -highest weights for an irreducible component of the representation  $\rho_j$ . Each  $\zeta_{l,j}$  can be uniquely expressed as a linear combination  $\sum_{i=1}^h x_{z_i} z_i^\vee + \sum_{\alpha \in \Delta} x_\alpha \alpha$ , where all coefficients  $x_\alpha$  are nonnegative. Since the parabolic  $P_\lambda$  is proper, we can choose some  $\alpha \in I_{P_\lambda}$ . Since  $\bigoplus \rho_j$  is faithful, at least one of the  $\zeta_{l,j}$  must have a nonzero coefficient  $x_\alpha > 0$ ; we fix such  $l, j$ . The restriction  $\chi_{l,j}^\lambda \in X^*(Z_\lambda)$  is a  $P_\lambda$ -dominant character such that the coefficient of  $\alpha$  is not 0. Since  $\psi$  is in the interior of the the  $P_\lambda$ -dominant cone in  $X_*(Z_\lambda)_{\mathbb{R}}$ , it follows that  $\langle \psi, \chi_{l,j}^\lambda \rangle > 0$ .

Let  $W$  denote the Weyl group of  $G$  permutting the set of characters  $\{\chi_{i,j}\}_{i=1}^{r_j}$  of the representation  $\rho_j$ . Let  $W_{P_\lambda}$  denote the subgroup of  $W$  generated by the reflections of the simple roots in  $I_{P_\lambda}$ .  $W_{P_\lambda}$  acts on the characters  $X^*(Z_\lambda)$ , and the action is transitive on the set of Weyl chambers in  $X^*(Z_\lambda)_{\mathbb{R}}$ . The set of restrictions  $\{\chi_{i,j}^\lambda\}_{i=1}^{r_j} \subset X^*(Z_\lambda)$  is preserved by  $W_{P_\lambda}$ . The character  $\chi_{l,j}^\lambda$  is not fixed by the action of  $W_\lambda$ , because the coefficient of  $\alpha$  is not 0. Therefore, there exists some  $w \in W_{P_\lambda}$  such that  $\chi_{i,j}^\lambda := w \cdot \chi_{l,j}^\lambda$  is contained in the the  $P_\lambda$ -anti-dominant cone of  $X^*(Z_\lambda)_{\mathbb{R}}$ . This implies that  $\langle \psi, \chi_{i,j}^\lambda \rangle < 0$ . Since  $\langle \psi, \chi_{i,j}^\lambda \rangle < 0 < \langle \psi, \chi_{l,j}^\lambda \rangle$ , we conclude that  $\langle \psi, \chi_i^\lambda \rangle \neq \langle \psi, \chi_j^\lambda \rangle$ , as desired.  $\square$

Example 10.10 below shows that the converse is not true even if we assume that the underlying rational  $G$ -bundle is stable in the sense of Ramanathan. This seems quite restrictive. In order to obtain a reasonable notion of stability, it will be useful to impose the following condition on our representation  $\rho$ .

**Definition 10.5.** *We say that a homomorphism of groups  $\rho : G_1 \rightarrow G_2$  is central if the image of the identity component of the center of  $G_1$  is in the center of  $G_2$ .*

Unless otherwise mentioned, for the rest of this subsection we will assume that  $\rho$  is central. This will allow us to have a better behaved notion of stability which can be compared to previous notions in the literature. First, we note that under this additional condition we can check semistability only using cocharacters of the derived subgroup of  $G$ .

**Proposition 10.6.** *Let  $K \supset k$  be an algebraically closed field, and suppose that  $\rho$  is central. Then in the semistability condition of Definition 10.2 it is enough to look at cocharacters  $\lambda : (\mathbb{G}_m)_K \rightarrow [G_K, G_K]$  whose image is in the derived subgroup  $[G_K, G_K] \subset G_K$ .*

*Proof.* Let  $Z \subset G_K$  be the center and  $[G_K, G_K] \subset G_K$  be the commutator subgroup. Consider the exact sequence

$$1 \rightarrow F \rightarrow Z \times [G_K, G_K] \rightarrow G_K \rightarrow 1$$

where  $F = Z \cap [G_K, G_K]$  is a finite group (it is the center of the semisimple group  $[G_K, G_K]$ ). Let  $\lambda : (\mathbb{G}_m)_K \rightarrow G_K$  be a cocharacter. Since  $F$  is finite, there is an integer  $N$  such that  $\lambda^N$  lifts to  $Z \times [G_K, G_K]$ . If we replace  $\lambda$  by  $\lambda^N$  then the parabolic subgroup  $\mathcal{G}_\lambda$  associated to  $\lambda$  does not change. The only change in the associated filtration  $(\mathcal{F}_m^i)_{m \in \mathbb{Z}}$  is that the indexes are multiplied by  $N$ , and the expression (15) does not change. Therefore, to check (semi)stability we can assume that the cocharacter  $\lambda$  lifts to the product  $Z \times [G_K, G_K]$ . Assume  $\lambda$  lifts, and write  $\lambda = (\lambda_1, \lambda_2)$  where  $\lambda_1$  is the lift to  $Z$  and  $\lambda_2$  is the lift to  $[G_K, G_K]$ .

We claim that the filtrations associated to  $(\lambda_1, \lambda_2)$  and  $(1, \lambda_2)$  on each  $V^i$  only differ by a constant shift in the indexes. Indeed, since  $\rho$  is central, the image of the composition  $\rho_i \circ \lambda_1$  lies in the center of  $\mathrm{GL}(V^i)$ , so  $(\mathbb{G}_m)_K$  acts on  $V^i$ , via  $\rho_i \circ (\lambda_1, 1)$ , with a fixed weight  $m_i$ . If  $v \in V^i$  is a vector where  $(\mathbb{G}_m)_K$  acts with weight  $m$  via  $\rho_i \circ (\lambda_1, \lambda_2)$ , then the action, via  $\rho_i \circ (1, \lambda_2)$ , has weight  $m - m_i$ . Therefore, the filtrations  $(\mathcal{F}_m^i)_{m \in \mathbb{Z}}$  induced by  $\rho_i \circ (1, \lambda_2)$  and  $\rho_i \circ (\lambda_1, \lambda_2)$  just differ by a shift by  $m_i$ , as claimed.

Finally, since the sum in (15) does not change by a shift by  $m_i$  on each  $V^i$ , we may assume that  $\lambda_1$  is equal to the constant 1. In other words, we may assume that  $\lambda$  factors through  $[G_K, G_K]$ .  $\square$

Proposition 10.6 shows that the following definition is consistent with our earlier notions.

**Definition 10.7.** *Suppose that  $\rho$  is central. Let  $K \supset k$  be an algebraically closed field. A principal  $\rho$ -sheaf on  $X_K$  is Gieseker stable (respectively, Gieseker semistable) if for all nontrivial cocharacters  $\lambda : (\mathbb{G}_m)_K \rightarrow [G_K, G_K]$  and all weighted parabolic reductions  $(\lambda, \mathcal{G}_\lambda)$  with associated underlying filtrations  $(\mathcal{F}_m^i)_{m \in \mathbb{Z}}$  of  $\mathcal{F}^i$ , (as in Definition 3.15) we have*

$$\sum_{i=1}^b \sum_{m \in \mathbb{Z}} (\mathrm{rk}(\mathcal{F}^i) \cdot P_{\mathcal{F}_m^i} - \mathrm{rk}(\mathcal{F}_m^i) \cdot P_{\mathcal{F}_m^i}) < 0 \quad (\text{respectively, } \leq)$$

A principal  $\rho$ -sheaf is slope stable (respectively, semistable) if we replace the numerical condition with

$$\sum_{i=1}^b \sum_{m \in \mathbb{Z}} (\mathrm{rk}(\mathcal{F}^i) \cdot \deg \mathcal{F}_m^i - \mathrm{rk}(\mathcal{F}_m^i) \cdot \deg \mathcal{F}^i) < 0 \quad (\text{respectively, } \leq).$$

We say that a principal  $\rho$ -sheaf is Gieseker unstable (resp. slope unstable) if it is not Gieseker semistable (resp. it is not slope semistable).

We immediately obtain that slope stable implies Gieseker stable, and Gieseker semistable implies slope semistable. We proceed now to compare with other notions of stability in the literature.

In [GLSS10, §1.1, p. 282] it is assumed that the field is algebraically closed,  $G$  is an arbitrary nonconnected reductive group, and the homomorphism  $\rho$  is assumed to be central. Furthermore, [GLSS10, §1.1, p. 282] requires that the image of  $\rho$  lies in the special group of  $V = V^1 \oplus \dots \oplus V^b$  as follows:

$$\rho : G \rightarrow \left( \mathrm{GL}(V^1) \times \dots \times \mathrm{GL}(V^b) \right) \cap \mathrm{SL}(V)$$

But this is not an essential condition. If we are given a representation  $\rho = (\rho_1, \dots, \rho_b)$  we can add one more summand  $V^{b+1} = k$  and the representation

$$\rho' = (\rho_1, \dots, \rho_b, (\det(\rho_1) \cdots \det(\rho_b))^{-1})$$

will automatically land in the special group. A  $\rho'$ -sheaf is  $\rho$ -sheaf together with an isomorphism  $\det(\mathcal{F}^1) \otimes \dots \otimes \det(\mathcal{F}^b) \cong (\mathcal{F}^{b+1})^\vee$ . The  $\rho'$ -sheaf is (semi)stable in the sense of [GLSS10] if and only if the corresponding  $\rho$ -sheaf is (semi)stable.

Therefore, a principal  $\rho$ -sheaf is the same thing as a singular principal  $\rho^\vee$ -bundle in [GLSS10] and our definition of (semi)stability coincides with the definition in [GLSS10, §1.1, p. 282] (recall that they assume that  $\rho$  is central, so we can use Proposition 10.6).

Therefore, all the comparisons that are made in [GLSS10, §6] also hold for us. If we consider orthogonal or symplectic groups with the standard representation, a principal  $\rho$ -sheaf is the same thing as an orthogonal or symplectic sheaf as defined in [GS03], and the conditions of (semi)stability defined in this article coincide with those in [GS03].

[GS05] uses the adjoint representation  $\rho : G \rightarrow \mathrm{GL}(\mathfrak{g}')$  into the semisimple part  $\mathfrak{g}' \subset \mathfrak{g}$  of the Lie algebra (note that the adjoint representation is trivial on the center  $\mathfrak{z}$  of the Lie algebra). If the center of  $G$  is trivial this representation is faithful. Then we can apply the techniques of this article. A principal  $\rho$ -sheaf is the same thing as a principal  $G$ -sheaf as defined in [GS05], and the conditions of stability and semistability coincide with those in [GS05].

**Lemma 10.8** (Lemma 2.5.4 and Remark 2.5.5 in [GLSS10]). *A rational principal bundle  $\mathcal{G}$  defined over  $U$  is slope stable (respectively, slope semistable) if for every big open subset  $U' \subset U$ , every nontrivial cocharacter  $\lambda : (\mathbb{G}_m)_K \rightarrow [G_K, G_K] \subset G_K$  and every reduction  $\mathcal{G}_\lambda \rightarrow U' \subset U$  to the parabolic subgroup  $P_\lambda$  defined by  $\lambda$ , we have*

$$\sum_{i=1}^b \sum_{m \in \mathbb{Z}} (\mathrm{rk}(\mathcal{F}) \cdot \deg \mathcal{F}_m^i - \mathrm{rk}(\mathcal{F}_i) \cdot \deg \mathcal{F}^i) < 0 \quad (\text{respectively, } \leq)$$

where  $\mathcal{F}^i = (\rho_i)_* \mathcal{G}$  and  $(\mathcal{F}_m^i)_{m \in \mathbb{Z}}$  is the filtration associated to the weighted parabolic reduction.  $\square$

The following corollary follows immediately.

**Corollary 10.9.** *If  $\rho$  is central, then a principal  $\rho$ -sheaf is slope (semi)stable if and only if the underlying rational principal bundle is slope (semi)stable in the sense of Ramanathan (Definition 10.3).  $\square$*

If  $\dim X = 1$ , then the only big open subset of  $X$  is  $X$  itself. It follows that a principal  $\rho$ -sheaf on a curve is the same thing as a principal  $G$ -bundle, and the notion of Gieseker (semi)stability coincides with the notion introduced by Ramanathan in [Ram75].

If the representation  $\rho$  is not central, then our notion of Gieseker semistability can be quite restrictive. This is illustrated by the next example.

**Example 10.10.** *We construct a principal  $\rho$ -sheaf, with  $\rho$  not central (Definition 10.5), which is Gieseker unstable, but the underlying rational principal bundle is stable in the sense of Ramanathan (as in Lemma 10.8).*

Let  $W$  be a vector space over  $k$ . Set  $V = W \oplus W$  and  $G = \mathrm{GL}(W) \times \mathrm{GL}(W)$ . We define  $\rho : G \rightarrow \mathrm{GL}(V)$  to be the faithful representation given by taking the direct sum of automorphisms

$$\rho(A, B) = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$$

Note that, in the notation of Definition 10.2, we are taking  $b = 1$ , and then  $\rho$  is not central.

Let  $(\mathcal{E}, \mathcal{E}')$  be a pair of slope stable vector bundles of rank  $\dim W$  on  $X$  and with  $d = \deg \mathcal{E} > \deg \mathcal{E}' = d'$ . This pair defines a principal  $G$ -bundle  $P$  on  $X$ . In particular, it defines a principal  $\rho$ -sheaf. It is stable in the sense of Ramanathan (Definition 10.3 or, equivalently, Lemma 10.8 for the identity representation of  $\mathrm{GL}(W) \times \mathrm{GL}(W)$ ). However it is not Gieseker semistable in the sense of Definition 10.2. Indeed, consider the central cocharacter  $\lambda$  of  $G$  defined by  $\lambda(t) = t^d \mathrm{Id}_W \oplus t^{d'} \mathrm{Id}_W$  (note that the image is not in the subgroup  $[G, G]$ ). The corresponding parabolic  $P_\lambda$  is the whole group  $G$ . We have a weighted

parabolic reduction  $f = (\lambda, \mathcal{G}_\lambda)$ , where  $\mathcal{G}_\lambda$  is the trivial parabolic reduction over  $X$ . The underlying associated filtration  $(\mathcal{F}_m)_{m \in \mathbb{Z}}$  of  $\mathcal{F}$  is given by

$$\mathcal{F}_m = \begin{cases} \mathcal{E} \oplus \mathcal{E}' & \text{if } m \leq d' \\ \mathcal{E} & \text{if } d' < m \leq d \\ 0 & \text{if } d < m \end{cases}$$

We have

$$\mathrm{rk}(\mathcal{E} \oplus \mathcal{E}') \cdot \mathrm{deg}(\mathcal{E}) - \mathrm{rk}(\mathcal{E}) \cdot \mathrm{deg}(\mathcal{E} \oplus \mathcal{E}') = (d - d') \dim W > 0$$

and hence the numerical condition in the Gieseker semistability is not satisfied.

**10.2. Comparison of the leading term filtration with previous notions in the literature.** We end this section by comparing the leading term HN filtration with the canonical (slope) parabolic reductions for  $G$ -bundles described in [AAB02, GN18].

**Proposition 10.11.** *Suppose that the representation  $\rho$  is central. Let  $K \supset k$  be a field extension such that  $G_K$  is split. Let  $(\mathcal{F}^\bullet, \sigma)$  be a  $\rho$ -sheaf on  $X_K$  and choose a big open subset  $U \subset X_K$  such that  $\mathcal{F}|_U$  is locally-free. Let  $\mathcal{G}$  denote the rational  $G$ -bundle on  $U$  corresponding to  $(\mathcal{F}^\bullet, \sigma)$  and suppose that  $\mathcal{G}$  is (slope) unstable in the sense of Ramanathan. If  $f = (\lambda, \mathcal{G}_\lambda)$  is the leading term HN weighted parabolic reduction of  $(\mathcal{F}^\bullet, \sigma)$ , then  $\mathcal{G}_\lambda$  is the canonical (slope) parabolic reduction as defined in [AAB02].*

*Proof.* We will use the setup and notation from Propositions 4.26 and 10.4. The group  $G$  acts on each vector space  $V^j$  via the homomorphism  $\rho_j$ . For each  $j$ , we denote by  $(\chi_{i,j})_{i=1}^{r_j}$  the tuple of weights of the split maximal torus  $T$  acting on  $V^j$  (with possible repetitions). Since the representation  $\rho$  is faithful, the collection of all the weights  $(\chi_{i,j})$  determines a positive definite inner product  $\cdot_\chi$  on  $X_*(T)_\mathbb{R}$  given by

$$\delta \cdot_\chi \gamma := \sum_{j=1}^b \sum_{i=1}^{r_j} \langle \delta, \chi_{i,j} \rangle \cdot \langle \gamma, \chi_{i,j} \rangle$$

which is invariant under the action of the Weyl group  $W$ . We can restrict this inner product to  $X_*(Z_\lambda)_\mathbb{R}$  in order to obtain a  $W_{P_\lambda}$ -invariant positive definite inner product.

Recall that we have

$$(18) \quad \nu_{d-1}(\delta) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\sum_{j=1}^b \sum_{i=1}^{r_j} \langle \delta, \chi_{i,j}^\lambda \rangle \cdot (c_{i,j} - c_j)}{\sqrt{\sum_{j=1}^b \sum_{i=1}^{r_j} \left( \langle \delta, \chi_{i,j}^\lambda \rangle \right)^2}}$$

where we define  $c_{i,j} = \langle \psi, \chi_{i,j}^\lambda \rangle$  and  $c_j = \frac{1}{r_j} \langle \psi, \sum_{i=1}^{r_j} \chi_{i,j}^\lambda \rangle$ . Set  $\mathrm{det}_j = \det(V^j) \circ \rho_j = \sum_{i=1}^{r_j} \chi_{i,j}$ . Rewrite (18) as

$$(19) \quad \nu_{d-1}(\delta) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\sum_{j=1}^b \sum_{i=1}^{r_j} \langle \delta, \chi_{i,j}^\lambda \rangle \cdot \langle \psi, \chi_{i,j}^\lambda \rangle - \frac{1}{r_j} \langle \delta, \mathrm{det}_j \rangle \cdot \langle \psi, \mathrm{det}_j \rangle}{\sqrt{\sum_{j=1}^b \sum_{i=1}^{r_j} \left( \langle \delta, \chi_{i,j}^\lambda \rangle \right)^2}} = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\delta \cdot_\chi \psi - \sum_{j=1}^b \frac{1}{r_j} \langle \delta, \mathrm{det}_j \rangle \cdot \langle \psi, \mathrm{det}_j \rangle}{\sqrt{\delta \cdot_\chi \delta}}$$

Recall that

$$X_*(Z_\lambda)_\mathbb{R} = \bigoplus_{l=1}^h \mathbb{R}z_l \oplus \bigoplus_{n \in J_{P_\lambda}} \mathbb{R}\omega_n^\vee$$

Choose a pair  $z_l$  and  $\omega_n^\vee$ . Let  $s_n$  denote the simple reflection of  $X_*(T)_\mathbb{R}$  determined by the root  $\alpha_n$ . Since  $\cdot_\chi$  is  $W_\lambda$ -invariant, we have

$$z_l \cdot_\chi \omega_n^\vee = (s_n z_l) \cdot_\chi (s_n \omega_n^\vee) = z_l \cdot_\chi (-\omega_n^\vee).$$

This shows that  $z_l \cdot_\chi \omega_n^\vee = 0$ , and so the central cocharacters  $X_*(Z)_\mathbb{R} = \bigoplus_{i=1}^h \mathbb{R}z_i$  are orthogonal to the fundamental coweights. Therefore the projection  $\pi_Z : X_*(Z_\lambda)_\mathbb{R} \rightarrow \bigoplus_{n \in J_{P_\lambda}} \mathbb{R}\omega_n^\vee$  is an orthogonal projection with respect to  $\cdot_\chi$ .

By assumption, the central torus  $Z$  maps to the center of  $\prod_{j=1}^b \mathrm{GL}(V^j)$ . For every index  $l$ , since  $\rho_j \circ z_l$  lands in the center of  $\mathrm{GL}(V^j)$ , we must have  $\langle z_l, \chi_{i,j} \rangle = \langle z_l, \chi_{p,j} \rangle$  for any  $i, p$ . It follows that  $\langle z_l, \det_j \rangle = r_j \langle z_l, \chi_{1,j} \rangle$  for the fixed index  $i = 1$ . This implies that

$$\delta \cdot_\chi \psi - \sum_{j=1}^b \frac{1}{r_j} \langle \delta, \det_j \rangle \cdot \langle \psi, \det_j \rangle = \pi_Z(\delta) \cdot_\chi \pi_Z(\psi)$$

Therefore, (19) turns out to be:

$$(20) \quad \nu_{d-1}(\delta) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \frac{\pi_Z(\delta) \cdot_\chi \pi_Z(\psi)}{\sqrt{\delta \cdot_\chi \delta}}$$

By Corollary 10.9, the  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  is unstable. Using the proof of Proposition 10.6 we can conclude that the image of  $\lambda$  is not contained in the center of  $G$ . Therefore the parabolic  $P_\lambda$  is a proper subgroup of  $G$ . We denote by  $f = (\lambda, \mathcal{G}_\lambda)$  the leading term HN weighted parabolic reduction of  $(\mathcal{F}, \sigma)$ . To prove that  $\mathcal{G}_\lambda$  is the canonical parabolic reduction of the rational  $G$ -bundle  $\mathcal{G}$ , we need to show the following (cf. [AAB02, Thm. 6]):

- (i) The linear functional  $\psi \in X_*(Z_\lambda)_\mathbb{R}$  lies in the interior  $\mathrm{Int}(\mathcal{C}_\lambda)$  of the dominant cone  $\mathcal{C}_\lambda$ .
- (ii) The rational  $L_\lambda$ -bundle  $\mathcal{G}_{L_\lambda}$  is slope semistable.

We proceed with the proof of these statements.

(i) Since  $f = (\lambda, \mathcal{G}_\lambda)$  maximizes  $\nu(f)$  in the interior of  $\mathcal{C}_\lambda$ , we know that  $\lambda$  is a critical point of  $\nu_{d-1}$ . It follows from Cauchy-Schwarz that this critical point is a scalar multiple of  $\pi_Z(\psi)$ . We must have  $\lambda = Q \cdot \pi_Z(\psi)$  for some constant  $Q > 0$ , because  $\nu(f) > 0$  and then  $\nu_{d-1}(\lambda) \geq 0$ . Since  $\lambda \in \mathrm{Int}(\mathcal{C}_\lambda)$  by definition, it follows that  $\pi_Z(\psi) \in \mathrm{Int}(\mathcal{C}_\lambda)$ . Therefore  $\psi \in \mathrm{Int}(\mathcal{C}_\lambda)$ , as desired.

(ii) Suppose for the sake of contradiction that  $\mathcal{G}_\lambda$  is not slope semistable. Let  $\mathcal{G}_{\bar{P}}$  be the canonical slope parabolic reduction of  $\mathcal{G}_{L_\lambda}$  for some proper parabolic subgroup  $\bar{P} \subset L_\lambda$ , as in [AAB02]. This reduction is defined over a big open subset. After shrinking the original open subset  $U$ , we can assume that everything is defined over  $U$ .

Let  $P$  denote the inverse image of  $\bar{P}$  under the canonical quotient map  $P_\lambda \twoheadrightarrow L_\lambda$ . Note that  $P$  is a parabolic subgroup of  $G$  (c.f. second paragraph of the proof of [BH04, Prop. 3.1]). Let  $\iota : P \hookrightarrow P_\lambda$  denote the inclusion and let  $q : P \twoheadrightarrow \bar{P}$  denote the quotient map. By [BH04, Lem. 2.1], there exists a unique  $P$ -parabolic reduction  $\mathcal{G}_P$  of  $\mathcal{G}$  such that  $\iota_*(\mathcal{G}_P) = \mathcal{G}_\lambda$  and  $q_*(\mathcal{G}_P) = \mathcal{G}_{\bar{P}}$ . Let  $L$  be the unique Levi subgroup of  $P$  containing the maximal torus  $T \subset P_\lambda$ , and let  $Z_L$  denote the maximal central torus of  $L$ . The canonical inclusion  $Z_\lambda \subset Z_L$  induces an inclusion  $X_*(Z_\lambda)_\mathbb{R} \subset X_*(Z_L)_\mathbb{R}$ .

Since  $\lambda = Q \cdot \pi_Z(\psi)$  for  $Q > 0$ , (20) is given by

$$\nu_{d-1}(\lambda) = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \sqrt{\pi_Z(\psi) \cdot_\chi \pi_Z(\psi)}.$$

On the other hand, let

$$\psi_L = z + \sum_{n \in J_P} a_n \omega_n^\vee \in X_*(Z_L)_\mathbb{R} = X_*(Z)_\mathbb{R} \oplus \bigoplus_{n \in J_P} \mathbb{R}\omega_n^\vee$$

be the linear functional corresponding to the  $L$ -bundle  $\mathcal{G}_L$  defined as in the proof of Lemma 4.25. Here  $z \in X_*(Z)_\mathbb{R}$  and  $a_m \in \mathbb{R}$ .



By construction,  $\mathcal{G}_{L_\lambda}$  is isomorphic to the  $L_\lambda$ -bundle obtained from  $\mathcal{G}_L$  by extension of structure group via the inclusion  $L \subset L_\lambda$ . This implies that  $\psi = z + \sum_{n \in J_{P_\lambda}} a_n \omega_n^\vee$ , hence

$$(21) \quad \psi_L = \psi + \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee$$

In particular we have  $a_n > 0$  for all  $q \in J_{P_\lambda}$ , because  $\psi \in \text{Int}(\mathcal{C}_\lambda)$ .

We know that the bundle  $q_*(\mathcal{G}_\lambda)$  is the canonical slope parabolic reduction of  $\mathcal{G}_{L_\lambda}$ . This implies that  $a_n > 0$  for all the other indexes  $n \in J_P \setminus J_{P_\lambda}$ . We conclude that  $\psi_L$  is in the interior of the  $P$ -dominant cone  $\mathcal{C}_P$ . This shows that there exists some positive integer  $N > 0$  such that the tuple  $f' = (N \cdot \pi_Z(\psi_L), \mathcal{G}_P)$  is a well-defined weighted parabolic reduction of  $(\mathcal{F}, \sigma)$ . The version of (20) for this parabolic reduction yields

$$\nu_{d-1}(f') = \frac{1}{\sqrt{A_d}(d-1)!} \cdot \sqrt{\pi_Z(\psi_L) \cdot_\chi \pi_Z(\psi_L)}.$$

Equation (21) implies  $\pi_Z(\psi_L) = \pi_Z(\psi) + \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee$ , therefore:

$$(22) \quad \begin{aligned} \pi_Z(\psi_L) \cdot_\chi \pi_Z(\psi_L) &= \pi_Z(\psi) \cdot_\chi \pi_Z(\psi) + 2\pi_Z(\psi) \cdot_\chi \left( \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee \right) + \\ &\quad \left( \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee \right) \cdot_\chi \left( \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee \right) \end{aligned}$$

Note that  $\cdot_\chi$  is a Weyl group invariant positive definite inner product on  $X_*(T)_\mathbb{R}$ . In particular we must have  $\omega_l^\vee \cdot_\chi \omega_n^\vee \geq 0$  for all  $l, n$ . Since  $a_n > 0$  for all  $n \in J_P$ , it follows that

$$\pi_Z(\psi) \cdot_\chi \left( \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee \right) = \left( \sum_{l \in J_{P_\lambda}} a_l \omega_l^\vee \right) \cdot_\chi \left( \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee \right) \geq 0$$

On the other hand, since  $\cdot_\chi$  is positive definite we have

$$\left( \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee \right) \cdot_\chi \left( \sum_{n \in J_P \setminus J_{P_\lambda}} a_n \omega_n^\vee \right) > 0.$$

Plugging the last two inequalities in (22) we get  $\pi_Z(\psi_L) \cdot_\chi \pi_Z(\psi_L) > \pi_Z(\psi) \cdot_\chi \pi_Z(\psi)$ . Therefore  $\nu_{d-1}(f') > \nu_{d-1}(\lambda) = \nu_{d-1}(f)$ . Hence, the leading coefficient of  $\nu(f')$  is strictly bigger than the leading coefficient of  $\nu(f)$ , thus contradicting the maximality of  $\nu(f)$ .  $\square$

**Remark 10.12.** *When the representation  $\rho$  is central, Proposition 10.11 shows that the leading term HN  $\Theta$ -stratification of  $\text{Bun}_\rho(X)$  is a refinement of the slope stratification for the stack of  $G$ -bundles defined in [GN18, GN16] in the split case. This in turn implies that the finer stratification induced by GHN filtrations is also a refinement of [GN18, GN16].*

## APPENDIX A. GIESEKER-HARDER-NARASIMHAN FILTRATIONS IN POSITIVE CHARACTERISTIC

Many of the arguments in this article do not use the hypothesis that the characteristic of  $k$  is 0. In this appendix we explain what needs to be modified in order to define Gieseker-Harder-Narasimhan filtrations in positive characteristic. We can show that  $\nu$  defines a weak  $\Theta$ -stratification on  $\text{Bun}_\rho(X)$  (as in [HLHJ21, Defn. 2.23]). This allows us to define the GHN filtration of a  $\rho$ -sheaf after maybe passing to a purely inseparable extension of the ground field.

Fix an arbitrary field  $k$ . We use the same setup and notation as in the characteristic 0 case ( $X$  is a polarized smooth projective geometrically irreducible  $k$ -variety,  $G$  is a connected reductive group over  $k$ , and  $\rho$  is a faithful representation of  $G$ ). Note that everything in Subsection 2.2 applies to general  $k$ .

The construction of  $\text{Red}_G(\mathcal{F})$  in Subsection 2.3 can be done in a similar way in arbitrary characteristic. The proofs of all the results in Subsection 2.3 remain valid, with the only exception that in Lemma 2.16 it is no longer clear that the natural base-change morphism  $\text{Red}_G(\mathcal{F}) \times_{X_S} X_T \rightarrow \text{Red}_G(\mathcal{F}|_{X_T})$  is necessarily an isomorphism. This is because it is no longer true that formation of  $G$ -invariants commutes with arbitrary base-change. We note, however, that for any big open  $U \subset X_S$  such that  $\mathcal{F}$  is locally free, the restriction  $\text{Red}_G(\mathcal{F}) \times_{X_S} U_T \rightarrow \text{Red}_G(\mathcal{F}|_{X_T})_{U_T}$  is an isomorphism. This is because both sides are naturally identified with the quotient  $\mathcal{P}|_{U_T}/G$  by Proposition 2.17. This is used to show the following lemma.

**Lemma A.1.** *Let  $S$  be a  $k$ -scheme, and let  $\mathcal{F}$  be a relative torsion-free sheaf of rank  $r$  on  $X_S$ . For any  $T \rightarrow S$ , there is a natural bijection between the following two sets*

- (A) *Sections of the structure morphism  $\text{Red}_G(\mathcal{F}|_{X_T}) \rightarrow X_T$ .*
- (B) *Sections of the structure morphism  $\text{Red}_G(\mathcal{F}) \times_{X_S} X_T \rightarrow X_T$ .*

*Proof.* Let  $U \subset X_S$  denote a big open subset such that  $\mathcal{F}|_U$  is locally-free. By Lemma 2.2 (c), sections in the set (A) are in correspondence with sections  $U_T \rightarrow \text{Red}_G(\mathcal{F}|_{X_T})_{U_T}$  defined on the big open  $U_T \subset X_T$ . Since the base-change morphism  $\text{Red}_G(\mathcal{F}) \times_{X_S} X_T \rightarrow \text{Red}_G(\mathcal{F}|_{X_T})$  restricts to an isomorphism over the open set  $U_T$ , these sections are in bijection with sections  $U_T \rightarrow \text{Red}_G(\mathcal{F}) \times_{X_S} U_T$ . Another application of Lemma 2.2 (c) shows that all such sections extend uniquely to a morphism  $X_T \rightarrow \text{Red}_G(\mathcal{F}) \times_{X_S} X_T$  as in (B).  $\square$

In other words, even though the schemes  $\text{Red}_G(\mathcal{F}) \times_{X_S} X_T$  and  $\text{Red}_G(\mathcal{F}|_{X_T})$  are not necessarily isomorphic, they have the same sections over  $X_T$  because they are isomorphic over a big open subset.

The notion of  $\rho$ -sheaf and the stack  $\text{Bun}_\rho(X)$  can be defined in the same way for arbitrary characteristic.

**Definition A.2.** *A  $\rho$ -sheaf is a pair  $(\mathcal{F}, \sigma)$  where*

- (1)  *$\mathcal{F}$  is a family of torsion-free sheaves of rank  $r$  on  $X_S$ .*
- (2)  *$\sigma$  is a section of the structure morphism  $\text{Red}_G(\mathcal{F}) \rightarrow X_S$ .*

*The stack  $\text{Bun}_\rho(X)$  is the pseudofunctor from  $(\text{Aff}_k)^{op}$  into groupoids that sends an affine scheme  $T \in \text{Aff}_k$  to*

$$\text{Bun}_\rho(X)(T) = [ \text{groupoid of } \rho\text{-sheaves } (\mathcal{F}, \sigma) \text{ on } X_S \ ] .$$

Once we know Lemma A.1, everything in Section 3 applies verbatim. Similarly, the definitions and arguments in Subsections 4.1, 4.2, 4.3 and 4.4 apply without a change. Hence we have the following result in arbitrary characteristic

**Proposition A.3.** *The polynomial numerical invariant  $\nu$  on the stack  $\text{Bun}_\rho(X)$  is strictly  $\Theta$ -monotone, strictly  $S$ -monotone, and satisfies the HN boundedness condition.  $\square$*

Proposition A.3 yields the necessary hypotheses to apply [HLHJ21, Thm. 2.26] (or [HL14, Thm. B]). This yields the following theorem (see [HLHJ21, Defn. 2.23] for a definition of weak  $\Theta$ -stratification).

**Theorem A.4.** *The polynomial numerical invariant  $\nu$  defines a weak  $\Theta$ -stratification on the stack  $\text{Bun}_\rho(X)$ .  $\square$*

Note that in this case we only get a weak  $\Theta$ -stratification. In particular, the corresponding notion of leading term HN filtration of a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  is not necessarily defined over the field  $K$  of definition of  $(\mathcal{F}, \sigma)$ ; one might need to allow to pass to a finite purely inseparable extension of  $K$ . This is to be expected for small characteristics, because of the failure of Behrend's conjecture even in the case when  $X$  is a curve [Hei08a]. We note that it is still the case that the leading HN filtration is unique up to scaling, even though it might not be defined over the ground field.

The analogous definition of relative leading HN filtrations (Definition 4.32) applies in arbitrary characteristic. However, since we are working with a weak  $\Theta$ -stratification, the universal stratification  $H \rightarrow S$  is no longer a locally closed immersion. Instead, we get that the universal morphism  $H \rightarrow S$  parametrizing relative leading term HN filtrations is surjective, radicial and of finite type. Each connected component of  $H$  is finite over its image.

As long as the characteristic of  $k$  is not too small, the proofs in Section 5 can still be applied to show Theorem 5.7. More precisely, we just need that condition (2) in Proposition 5.2 holds.

**Theorem A.5.** *Choose an algebraic closure  $\bar{k}$  of  $k$ . Let  $Z \subset G_{\bar{k}}$  denote the maximal central torus of the base-change  $G_{\bar{k}}$ . Suppose that the characteristic of  $k$  does not divide the order of the kernel of the simply-connected cover  $\widetilde{G_{\bar{k}}/Z} \rightarrow G_{\bar{k}}/Z$  of the semisimple group  $G_{\bar{k}}/Z$ . Then the morphism  $\text{Bun}_{\rho}(X) \rightarrow \text{Spec}(k)$  satisfies the existence part of the valuative criterion for properness [Sta21, Tag 0CLK] in the case of complete discrete valuation rings (as in Theorem 5.7).  $\square$*

The definitions and constructions in Sections 7 and 8 apply in a similar way. The only change is that the GHN filtration of a  $\rho$ -sheaf  $(\mathcal{F}, \sigma)$  need not be defined over the field  $K$  of definition, but again one might need to pass to a finite purely inseparable extension. The analogue of Theorem 8.18 still holds; the only thing that needs to be modified is to replace the locally closed stratification  $H \rightarrow S$  with a surjective radicial morphism of finite type.

Finally, we note that the proof of Proposition 10.11 applies to arbitrary characteristic. This means that if the representation  $\rho$  is central, then the stratification of  $\text{Bun}_{\rho}(X)$  by radicial morphisms induced by our Gieseker-Harder-Narasimhan filtrations refines the theory developed in [GN16] in arbitrary characteristic.

**Remark A.6.** *The attentive reader might have noticed that we did not mention Section 6 in this appendix. Unfortunately the results in [HL14, AHLH19] that we are using to build moduli spaces do not apply in arbitrary characteristic.*

**Remark A.7.** *Using the same techniques, one could generalize the results in this appendix to the case when  $X \rightarrow S$  is a smooth projective morphism with geometrically connected fibers over an arbitrary base-scheme  $S$ ,  $G \rightarrow S$  is a reductive group scheme with connected fibers, and  $\rho$  is a faithful representation  $\rho : G \hookrightarrow \prod_i GL(V^i)$  for some vector bundles  $V^i$  on  $S$ . We have chosen not to pursue this level of generality in this paper.*

## REFERENCES

- [AAB02] B. Anchoche, H. Azad, and I. Biswas. Harder-Narasimhan reduction for principal bundles over a compact Kähler manifold. *Math. Ann.*, 323(4):693–712, 2002.
- [AB83] M. F. Atiyah and R. Bott. The Yang-Mills equations over Riemann surfaces. *Philos. Trans. Roy. Soc. London Ser. A*, 308(1505):523–615, 1983.
- [AHLH19] J. Alper, D. Halpern-Leistner, and J. Heinloth. Existence of moduli spaces for algebraic stacks. <https://arxiv.org/abs/1812.01128>, 2019. arXiv preprint.
- [Alp13] J. Alper. Good moduli spaces for Artin stacks. *Ann. Inst. Fourier (Grenoble)*, 63(6):2349–2402, 2013.

- [Bal07] V. Balaji. Principal bundles on projective varieties and the Donaldson-Uhlenbeck compactification. *J. Differential Geom.*, 76(3):351–398, 2007.
- [Beh91] K. A. Behrend. *The Lefschetz trace formula for the moduli stack of principal bundles*. 1991. Thesis (Ph.D.)–University of California, Berkeley.
- [Beh95] K. A. Behrend. Semi-stability of reductive group schemes over curves. *Math. Ann.*, 301(2):281–305, 1995.
- [BG03] I. Biswas and T. L. Gómez. Restriction theorems for principal bundles. *Math. Ann.*, 327(4):773–792, 2003.
- [BH04] I. Biswas and Y. I. Holla. Harder-Narasimhan reduction of a principal bundle. *Nagoya Math. J.*, 174:201–223, 2004.
- [BL95] Arnaud Beauville and Yves Laszlo. Un lemme de descente. *C. R. Acad. Sci. Paris Sér. I Math.*, 320(3):335–340, 1995.
- [BP03] V. Balaji and A. J. Parameswaran. Semistable principal bundles. II. Positive characteristics. *Transform. Groups*, 8(1):3–36, 2003.
- [BS02] V. Balaji and C. S. Seshadri. Semistable principal bundles. I. Characteristic zero. volume 258, pages 321–347. 2002. Special issue in celebration of Claudio Procesi’s 60th birthday.
- [BT87] F. Bruhat and J. Tits. Groupes algébriques sur un corps local. Chapitre III. Compléments et applications à la cohomologie galoisienne. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 34(3):671–698, 1987.
- [BZ16] I. Biswas and A. Zamora. On the Gieseker Harder-Narasimhan filtration for principal bundles. *Bull. Sci. Math.*, 140(4):58–69, 2016.
- [Fal93] G. Faltings. Stable  $G$ -bundles and projective connections. *J. Algebraic Geom.*, 2(3):507–568, 1993.
- [Ful98] W. Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, second edition, 1998.
- [Gil94] Ph. Gille. *Torseurs sur la droite affine et  $R$ -équivalence*. PhD thesis, Université Paris-Sud XI, Orsay, 1994.
- [GLSS08] T. L. Gómez, A. Langer, A. H. W. Schmitt, and I. Sols. Moduli spaces for principal bundles in arbitrary characteristic. *Adv. Math.*, 219(4):1177–1245, 2008.
- [GLSS10] T. L. Gómez, A. Langer, A. H. W. Schmitt, and Ignacio Sols. Moduli spaces for principal bundles in large characteristic. In *Teichmüller theory and moduli problem*, volume 10 of *Ramanujan Math. Soc. Lect. Notes Ser.*, pages 281–371. Ramanujan Math. Soc., Mysore, 2010.
- [GN14] S. Gurjar and N. Nitsure. Schematic Harder-Narasimhan stratification for families of principal bundles and  $\Lambda$ -modules. *Proc. Indian Acad. Sci. Math. Sci.*, 124(3):315–332, 2014.
- [GN16] S. Gurjar and N. Nitsure. Harder-Narasimhan stacks for principal bundles in higher dimensions and arbitrary characteristics. <https://arxiv.org/abs/1605.08997>, 2016. arXiv preprint.
- [GN18] S. Gurjar and N. Nitsure. Schematic Harder-Narasimhan stratification for families of principal bundles in higher dimensions. *Math. Z.*, 289(3-4):1121–1142, 2018.
- [GS03] T. L. Gómez and I. Sols. Stable tensors and moduli space of orthogonal sheaves. <https://arxiv.org/abs/math/0103150>, 2003. arXiv preprint.
- [GS05] T. L. Gómez and I. Sols. Moduli space of principal sheaves over projective varieties. *Ann. of Math. (2)*, 161(2):1037–1092, 2005.
- [GSZ15] T. L. Gómez, Ignacio Sols, and A. Zamora. A GIT interpretation of the Harder-Narasimhan filtration. *Rev. Mat. Complut.*, 28(1):169–190, 2015.
- [Hei08a] J. Heinloth. Bounds for Behrend’s conjecture on the canonical reduction. *Int. Math. Res. Not. IMRN*, (14):Art. ID rnn045, 17, 2008.
- [Hei08b] J. Heinloth. Semistable reduction for  $G$ -bundles on curves. *J. Algebraic Geom.*, 17(1):167–183, 2008.
- [Hei10] J. Heinloth. Addendum to “Semistable reduction of  $G$ -bundles on curves”. *J. Algebraic Geom.*, 19(1):193–197, 2010.
- [Hei17] J. Heinloth. Hilbert-Mumford stability on algebraic stacks and applications to  $\mathcal{G}$ -bundles on curves. *Épjournal Geom. Algébrique*, 1:Art. 11, 37, 2017.
- [HL97] D. Huybrechts and M. Lehn. *The geometry of moduli spaces of sheaves*. Aspects of Mathematics, E31. Friedr. Vieweg & Sohn, Braunschweig, 1997.
- [HL14] Daniel Halpern-Leistner. On the structure of instability in moduli theory. <https://arxiv.org/abs/1411.0627>, 2014. ArXiv version.
- [HLHJ21] D. Halpern-Leistner, A. Fernandez Herrero, and T. Jones. Moduli spaces of sheaves via affine Grassmannians. <https://arxiv.org/abs/2107.02172>, 2021.
- [HN75] G. Harder and M. S. Narasimhan. On the cohomology groups of moduli spaces of vector bundles on curves. *Math. Ann.*, 212:215–248, 1974/75.

- [HR15] Jack Hall and David Rydh. General Hilbert stacks and Quot schemes. *Michigan Math. J.*, 64(2):335–347, 2015.
- [HR19] Jack Hall and David Rydh. Coherent Tannaka duality and algebraicity of Hom-stacks. *Algebra Number Theory*, 13(7):1633–1675, 2019.
- [Kem78] G. R. Kempf. Instability in invariant theory. *Ann. of Math. (2)*, 108(2):299–316, 1978.
- [Kob87] Shoshichi Kobayashi. *Differential geometry of complex vector bundles*, volume 15 of *Publications of the Mathematical Society of Japan*. Princeton University Press, Princeton, NJ; Princeton University Press, Princeton, NJ, 1987. Kanô Memorial Lectures, 5.
- [Lan75] S. G. Langton. Valuative criteria for families of vector bundles on algebraic varieties. *Ann. of Math. (2)*, 101:88–110, 1975.
- [Lar95] M. Larsen. Maximality of Galois actions for compatible systems. *Duke Math. J.*, 80(3):601–630, 1995.
- [LMB00] Gérard Laumon and Laurent Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.
- [MFK94] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric invariant theory*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (2)*. Springer-Verlag, Berlin, third edition, 1994.
- [Nit11] N. Nitsure. Schematic Harder-Narasimhan stratification. *Internat. J. Math.*, 22(10):1365–1373, 2011.
- [Ram75] A. Ramanathan. Stable principal bundles on a compact Riemann surface. *Math. Ann.*, 213:129–152, 1975.
- [Ram79] A. Ramanathan. Moduli for principal bundles. In *Algebraic geometry (Proc. Summer Meeting, Univ. Copenhagen, Copenhagen, 1978)*, volume 732 of *Lecture Notes in Math.*, pages 527–533. Springer, Berlin, 1979.
- [RR84] M. S. Raghunathan and A. Ramanathan. Principal bundles on the affine line. *Proc. Indian Acad. Sci. Math. Sci.*, 93(2-3):137–145, 1984.
- [Sch02] A. H. W. Schmitt. Singular principal bundles over higher-dimensional manifolds and their moduli spaces. *Int. Math. Res. Not.*, (23):1183–1209, 2002.
- [Ser96] J.-P. Serre. Exemples de plongements des groupes  $\mathrm{PSL}_2(\mathbf{F}_p)$  dans des groupes de Lie simples. *Invent. Math.*, 124(1-3):525–562, 1996.
- [Sta21] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2021.
- [Zam13] A. Zamora. GIT characterizations of Harder-Narasimhan filtrations. <https://arxiv.org/abs/1407.4223>, 2013. Ph.D. Thesis, Universidad Complutense de Madrid.
- [Zam16] A. Zamora. Harder-Narasimhan filtration for rank 2 tensors and stable coverings. *Proc. Indian Acad. Sci. Math. Sci.*, 126(3):305–327, 2016.

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, 310 MALOTT HALL, ITHACA, NEW YORK 14853, USA.

*Email address:* [ajf254@cornell.edu](mailto:ajf254@cornell.edu)

INSTITUTO DE CIENCIAS MATEMÁTICAS (CSIC-UAM-UC3M-UCM), NICOLÁS CABRERA 15, CAMPUS CANTOBLANCO UAM, 28049 MADRID, SPAIN

*Email address:* [tomas.gomez@icmat.es](mailto:tomas.gomez@icmat.es)

DEPARTAMENTO DE MATEMÁTICA APLICADA A LAS TIC, ETSI INFORMÁTICOS, UNIVERSIDAD POLITÉCNICA DE MADRID, CAMPUS DE MONTEGANCEDO, 28660 MADRID, SPAIN

*Email address:* [alfonso.zamora@upm.es](mailto:alfonso.zamora@upm.es)