

# AUTOMORPHISMS OF SEMIDIRECT PRODUCTS FIXING THE NON-NORMAL SUBGROUP

RATAN LAL AND VIPUL KAKKAR

ABSTRACT. In this paper, we describe the automorphism group of semidirect product of two groups that fixes the non-normal subgroup of it. We have computed these automorphisms for the non-abelian metacyclic  $p$ -group and non-abelian  $p$ -groups ( $p \geq 5$ ) of order  $p^4$ , where  $p$  is a prime.

**Keywords:** Automorphism Group; Semidirect Product;  $p$ -group.

**2020 Mathematics Subject Classification:** 20D45, 20E36

## 1. INTRODUCTION

Let  $K$  be a non-normal subgroup of a group  $G$ . Let  $S$  be a right transversal to  $K$  in  $G$  with  $1 \in S$ . Then the group operation on  $G$  induces a binary operation on  $S$  with respect to it  $S$  becomes a right loop. Let  $Aut_K(G) = \{\theta \in Aut(G) \mid \theta(K) = K\}$ . In [7, Theorem 2.6, p. 73], R. Lal obtained that  $\theta \in Aut_K(G)$  can be identified with the triple  $(\alpha, \gamma, \delta)$ , where  $\alpha \in Map(S, S)$ ,  $\gamma \in Map(S, K)$  and  $\delta \in Aut(K)$  satisfying the conditions in [7, Definition 2.5, p. 73].

In case, if there is a right transversal  $H$  to  $K$  in  $G$  which is a normal subgroup of  $G$ , then  $G$  is the semidirect product of  $K$  and  $H$ . In this case, the conditions on  $\alpha, \gamma$  and  $\delta$  agrees with the conditions given in [6, Lemma 1.1, p. 1000]. These conditions are given as follows.

- (C1)  $\alpha(hh') = \alpha(h)\alpha(h')^{\gamma(h)}$ ,
- (C2)  $\gamma(h^k) = \gamma(h)^{\delta(k)}$ ,
- (C3)  $\alpha(h^k) = \alpha(h)^{\delta(k)}$ ,
- (C4) For any  $h'k' \in G$ , there exists a unique  $hk \in G$  such that  $\alpha(h) = h'$  and  $\gamma(h)\delta(k) = k'$ .

**Remark 1.1.** In [7], the author put the non-normal subgroup in the left in the factorization of  $G$ . To match the terminology with that in [4], we put the non-normal subgroup  $K$  in the right, that is  $G = HK$ . Through out the paper, we will use the terminology used in [4]. We will identify the internal semidirect product  $G = HK$  with the external semidirect product  $H \rtimes_{\phi} K$ , where  $\phi : K \rightarrow Aut(H)$  is a homomorphism.

M . J. Curran [4] and D. Jill [5] studied the automorphism group of the group  $G$  that fixes the normal subgroup  $H$ . In this paper, we study the automorphism group of the group  $G$  that fixes the non-normal subgroup  $K$ . We have computed the automorphism group,  $Aut_K(G)$

in case of non-abelian metacyclic  $p$ -groups and  $p$ -groups ( $p \geq 5$ ) of order  $p^4$ , where  $p$  is a prime.

## 2. STRUCTURE OF THE AUTOMORPHISM GROUP $Aut_K(G)$

Let  $\alpha \in Map(H, H)$ ,  $\gamma \in Hom(H, K)$  and  $\delta \in Aut(K)$ . Consider a set

$$\hat{\mathcal{M}}_K = \left\{ \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \mid \alpha, \gamma, \delta \text{ satisfy the conditions (C1) - (C4)} \right\}.$$

Then  $\hat{\mathcal{M}}_K$  is a group with the usual multiplication of matrices defined as,

$$\begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} \alpha' & 0 \\ \gamma' & \delta' \end{pmatrix} = \begin{pmatrix} \alpha\alpha' & 0 \\ \gamma\alpha' + \delta\gamma' & \delta\delta' \end{pmatrix},$$

where  $\alpha\alpha'$ ,  $\delta\delta'$  are the usual composition of maps and  $(\gamma\alpha' + \delta\gamma')(h) = \gamma\alpha'(h)\delta\gamma'(h)$ . Clearly,  $Aut_K(G)$  is a subgroup of  $Aut(G)$ .

**Proposition 2.1.** *The group  $Aut_K(G)$  is isomorphic to the group  $\hat{\mathcal{M}}_K$ .*

*Proof.* Define a map  $\psi : Aut_K(G) \rightarrow \hat{\mathcal{M}}_K$  by  $\psi(\theta) = \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix}$ , for all  $\theta \in Aut_K(G)$ , where  $\alpha : H \rightarrow H$ ,  $\gamma : H \rightarrow K$  and  $\delta : K \rightarrow K$  are defined as,  $\theta(h) = \alpha(h)\gamma(h)$  and  $\delta(k) = \theta(k)|_K$  for all  $h \in H$  and  $k \in K$ . By the similar argument as in [4, Theorem 1, p. 206], one can observe that  $\psi$  is an isomorphism.  $\square$

From now on we will identify automorphisms in  $Aut_K(G)$  with the matrices in  $\hat{\mathcal{M}}_K$ . Now, we have the following remark.

**Remark 2.1.** (i)  $\begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} \in Aut_K(G)$  if and only if  $\alpha \in Aut(H)$

and  $\alpha(h^k) = \alpha(h)^k$  for all  $h \in H$  and  $k \in K$ .

(ii)  $\begin{pmatrix} 1 & 0 \\ \gamma & 1 \end{pmatrix} \in Aut_K(G)$  if and only if  $\gamma(H) \subseteq C_K(H)$  and  $\gamma(h^k) = \gamma(h)^k$ , for all  $h \in H$  and  $k \in K$ , where  $C_K(H) = \{k \in K \mid h^k = h, \forall h \in H\}$  is the centralizer of  $H$  in  $K$ .

(iii)  $\begin{pmatrix} 1 & 0 \\ 0 & \delta \end{pmatrix} \in Aut_K(G)$  if and only if  $k^{-1}\delta(k) \in C_K(H)$  for all  $k \in K$ .

Now, let us consider the following subsets of  $Aut(H)$ ,  $Aut(K)$  and  $Aut(H) \times Aut(K)$ ,

$$\begin{aligned} U &= \{\alpha \in Aut(H) \mid \alpha(h^k) = \alpha(h)^k, \forall h \in H, k \in K\}, \\ V &= \{\delta \in Aut(K) \mid k^{-1}\delta(k) \in C_K(H), \forall k \in K\}, \\ W &= \{(\alpha, \delta) \in Aut(H) \times Aut(K) \mid \alpha(h^k) = \alpha(h)^{\delta(k)}, \forall h \in H, k \in K\}. \end{aligned}$$

Clearly,  $U, V$  and  $W$  are the subgroups of  $Aut(H), Aut(K)$ , and  $Aut(H) \times Aut(K)$ , respectively. The corresponding subgroups of the group  $Aut_K(G)$  are

$$A = \left\{ \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} \mid \alpha \in U \right\}, D = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & \delta \end{pmatrix} \mid \delta \in V \right\},$$

and  $E = \left\{ \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \mid (\alpha, \delta) \in W \right\}$ . Note that, if  $\alpha \in U$  and  $\delta \in V$ , then  $(\alpha, \delta) \in W$ . Therefore,  $U \times V \leq W$ .

Clearly,  $E$  is a subgroup of  $Aut_K(G)$ . However, one can check that  $E$  need not be a normal subgroup of  $Aut_K(G)$ . Let

$$C = \left\{ \begin{pmatrix} 1 & 0 \\ \gamma & 1 \end{pmatrix} \in Aut_K(G) \mid \gamma(H) \subseteq C_K(H) \text{ and} \right. \\ \left. \gamma(h^k) = \gamma(h)^k, \text{ for all } h \in H \text{ and } k \in K \right\}.$$

One can easily observe that  $C$  is a normal subgroup of the group  $Aut_K(G)$ . Clearly,  $E \cap C = 1$ . Now, let  $\begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \in Aut_K(G)$ . Then,

$$\begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \gamma\alpha^{-1} & 1 \end{pmatrix} \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix}.$$

Hence,  $Aut_K(G) = CE, C \trianglelefteq Aut_K(G), E \leq Aut_K(G)$ , and  $C \cap E = 1$ . Thus, we have proved the following theorem,

**Theorem 2.1.** *Let  $G = H \rtimes K$  be the semidirect product. Then  $Aut_K(G) \simeq C \rtimes E$ .*

### 3. COMPUTATION OF $Aut_K(G)$ FOR SOME GROUPS

In this section, we will compute the automorphism group  $Aut_K(G)$  for non-abelian metacyclic  $p$ -groups and  $p$ -groups ( $p \geq 5$ ) of order  $p^4$ , where  $p$  is a prime.  $\mathbb{Z}_m$  will denote the cyclic group of order  $m$ .

#### Metacyclic $p$ -groups

First, assume that  $p$  is odd. A non-abelian split metacyclic  $p$ -group  $G$  is of the form  $G = \langle a, b \mid a^{p^m} = 1 = b^{p^n}, a^b = a^{1+p^{m-r}} \rangle$ , where  $m \geq 2, n \geq 1$ , and  $1 \leq r \leq \min\{m-1, n\}$ . Let  $H = \langle a \rangle, K = \langle b \rangle$  and  $\phi : K \rightarrow Aut(H)$  be defined by  $\phi(b)(a) = a^{1+p^{m-r}}$ . Then  $G = H \rtimes_{\phi} K$ .

Note that  $[H, K] = \langle a^{p^{m-r}} \rangle \simeq \mathbb{Z}_{p^r}$ . Since  $K$  is abelian, by [2, Corollary 2.2, p. 490],  $\gamma(h^k) = \gamma(h)$  is equivalent to  $\gamma \in Hom(H/[H, K], K)$ . Define  $\gamma_i : H \rightarrow K$  by  $\gamma_i(a) = b^i, 1 \leq i \leq p^n$  when  $m-r \geq n$  and by  $\gamma_i(a) = b^{ip^{n-m+r}}, 1 \leq i \leq p^{m-r}$  when  $m-r < n$ . Since  $[H, K] \subseteq Ker \gamma_i$ , it will induce a homomorphism from  $H/[H, K]$  to  $K$ . Let  $c = \begin{pmatrix} 1 & 0 \\ \gamma_1 & 1 \end{pmatrix}$ . Then, one can easily observe that  $\gamma_1(H) \subseteq C_K(H)$ . Therefore,  $Hom(H/[H, K], K) \simeq C = \langle c \rangle \simeq \mathbb{Z}_{p^{\min\{m-r, n\}}}$ .

Also,  $C_K(H) = \langle b^{p^r} \rangle \simeq \mathbb{Z}_{p^{n-r}}$  and for  $b \in K$ ,  $b^{-1}\delta(b) \in C_K(H)$ . Therefore, there are  $p^{n-r}$  choices for  $\delta(b)$ . If  $\delta_1(b) = b^{1+p^r}$ , then  $V = \langle \delta_1 \rangle \simeq \mathbb{Z}_{p^{n-r}}$  and so,  $D \simeq \mathbb{Z}_{p^{n-r}}$ . Now, for all  $\alpha \in \text{Aut}(H)$ ,  $\alpha(a^b) = \alpha(a^{1+p^{m-r}}) = \alpha(a)^{1+p^{m-r}} = \alpha(a)^b$ . Therefore,  $U = \text{Aut}(H) \simeq \mathbb{Z}_{p^{m-1}(p-1)}$  and so,  $A \simeq \mathbb{Z}_{p^{m-1}(p-1)}$ . Then, by Theorem [4, Theorem 2, p. 207],  $E = A \times D$ . Now, by Theorem 2.1,  $\text{Aut}_K(G) \simeq \mathbb{Z}_{p^{\min\{m-r, n\}}} \rtimes (\mathbb{Z}_{p^{m-1}(p-1)} \times \mathbb{Z}_{p^{n-r}})$ . Hence,  $\text{Aut}_K(G)$  is a subgroup of index  $p^{\min\{m, n\}}$  in the group  $\text{Aut}(G)$ .

Now, assume  $p = 2$ . Then, as given in [3], the non-abelian split metacyclic 2-group is one of the following three forms,

- (i)  $G = \langle a, b \mid a^{2^m} = 1 = b^{2^n}, a^b = a^{1+2^{m-r}} \rangle, 1 \leq r \leq \min\{m-2, n\}, m \geq 3, n \geq 1$ .
- (ii)  $G = \langle a, b \mid a^{2^m} = 1 = b^{2^n}, a^b = a^{-1+2^{m-r}} \rangle, 1 \leq r \leq \min\{m-2, n\}, m \geq 3, n \geq 1$ .
- (iii)  $G = \langle a, b \mid a^{2^m} = 1 = b^{2^n}, a^b = a^{-1} \rangle, m \geq 2, n \geq 1$ .

Let  $H = \langle a \rangle \simeq \mathbb{Z}_{2^m}$  and  $K = \langle b \rangle \simeq \mathbb{Z}_{2^n}$ . We will compute the automorphism group,  $\text{Aut}_K(G)$  in the above three cases (i) – (iii). Using the similar argument as for odd prime  $p$  above, in the case (i),  $[H, K] = \langle a^{2^{m-r}} \rangle \simeq \mathbb{Z}_{2^r}$  and  $C_K(H) = \langle b^{2^r} \rangle \simeq \mathbb{Z}_{2^{n-r}}$ . Then  $\text{Hom}(H/[H, K], K) \simeq \mathbb{Z}_{2^{\min\{m-r, n\}}}$ . Thus  $C \simeq \mathbb{Z}_{2^{\min\{m-r, n\}}}$ ,  $A \simeq \mathbb{Z}_2 \times \mathbb{Z}_{2^{m-2}}$  and  $D \simeq \mathbb{Z}_2 \times \mathbb{Z}_{2^{n-r-1}}$ . Hence,  $\text{Aut}_K(G) \simeq \mathbb{Z}_{2^{\min\{m-r, n\}}} \rtimes (\mathbb{Z}_2 \times \mathbb{Z}_{2^{m-2}} \times \mathbb{Z}_2 \times \mathbb{Z}_{2^{n-r-1}})$ .

In the case (ii),  $[H, K] = \langle a^2 \rangle \simeq \mathbb{Z}_{2^{m-1}}$  and  $C_K(H) = \langle b^{2^r} \rangle \simeq \mathbb{Z}_{2^{n-r}}$ . Thus,  $C \simeq \mathbb{Z}_2$ ,  $A \simeq \mathbb{Z}_2 \times \mathbb{Z}_{2^{m-2}}$  and  $D \simeq \mathbb{Z}_2 \times \mathbb{Z}_{2^{n-r-1}}$ . Hence,  $\text{Aut}_K(G) \simeq \mathbb{Z}_2 \times (\mathbb{Z}_2 \times \mathbb{Z}_{2^{m-2}} \times \mathbb{Z}_2 \times \mathbb{Z}_{2^{n-r-1}})$ . Similarly, in the case (iii),  $[H, K] = \langle a^2 \rangle \simeq \mathbb{Z}_{2^{m-1}}$ , and  $C_K(H) = \langle b^2 \rangle \simeq \mathbb{Z}_{2^{n-1}}$ . Thus,  $C \simeq \mathbb{Z}_2$ ,  $A \simeq \mathbb{Z}_2 \times \mathbb{Z}_{2^{m-2}}$  and  $D \simeq \mathbb{Z}_2 \times \mathbb{Z}_{2^{n-2}}$ . Hence,  $\text{Aut}_K(G) \simeq \mathbb{Z}_2 \times (\mathbb{Z}_2 \times \mathbb{Z}_{2^{m-2}} \times \mathbb{Z}_2 \times \mathbb{Z}_{2^{n-2}})$ .

### Non-abelian $p$ -groups of order $p^4$ ( $p \geq 5$ )

Burnside in [1] classified  $p$ -groups of order  $p^4$ , where  $p$  is a prime. Below, we list 10 non-abelian  $p$ -groups ( $p \geq 5$ ) of order  $p^4$  up to isomorphism.

- (i)  $G_1 = \langle a, b \mid a^{p^3} = 1 = b^p, a^b = a^{1+p^2} \rangle$ ,
- (ii)  $G_2 = \langle a, b \mid a^{p^2} = 1 = b^{p^2}, a^b = a^{1+p} \rangle$ ,
- (iii)  $G_3 = \langle a, b, c \mid a^{p^2} = 1 = b^p = c^p, cb = a^pbc, ab = ba, ac = ca \rangle$ ,
- (iv)  $G_4 = \langle a, b, c \mid a^{p^2} = 1 = b^p = c^p, ca = a^{1+p}c, ab = ba, cb = bc \rangle$ ,
- (v)  $G_5 = \langle a, b, c \mid a^{p^2} = 1 = b^p = c^p, ca = abc, ab = ba, bc = cb \rangle$ ,
- (vi)  $G_6 = \langle a, b, c \mid a^{p^2} = 1 = b^p = c^p, ba = a^{1+p}b, ca = abc, bc = cb \rangle$ ,
- (vii)  $G_7 = \langle a, b, c \mid a^{p^2} = 1 = b^p = c^p, ba = a^{1+p}b, ca = a^{1+p}bc, cb = a^pbc \rangle$ ,
- (viii)  $G_8 = \langle a, b, c \mid a^{p^2} = 1 = b^p = c^p, ba = a^{1+p}b, ca = a^{1+dp}bc, cb = a^{dp}bc, d \not\equiv 0, 1 \pmod{p} \rangle$ ,
- (ix)  $G_9 = \langle a, b, c, d \mid a^p = b^p = c^p = d^p = 1, dc = acd, bd = db, ad = da, bc = cb, ac = ca, ab = ba \rangle$ ,

$$(x) \ G_{10} = \langle a, b, c, d \mid a^p = b^p = c^p = d^p = 1, dc = bcd, db = abd, ad = da, bc = cb, ac = ca, ab = ba \rangle.$$

Observe that  $G_1$  and  $G_2$  are metacyclic  $p$ -groups.  $Aut_K(G_1)$  and  $Aut_K(G_2)$  (for the corresponding  $K$ ) can be calculated by the above discussion.

**The group  $G_3$ .** Let  $H = \langle a, b \mid a^{p^2} = b^p = 1, ab = ba \rangle$  and  $K = \langle c \mid c^p = 1 \rangle$ . Then  $G_3 \simeq H \rtimes_{\phi} K$ , where  $\phi : K \rightarrow Aut(H)$  is given by  $\phi(c)(a) = a$  and  $\phi(c)(b) = a^p b$ . Note that  $[a^u b^v, c] = (a^u b^v) c (a^u b^v)^{-1} c^{-1} = a^u b^v (a^{u+pv} b^v)^{-1} = a^{-pv}$ . Therefore,  $[H, K] = \langle a^p \rangle \simeq \mathbb{Z}_p$ . Also, if  $c^s \in C_K(H)$ , then  $a^i b^j = c^s a^i b^j c^{-s} = a^{i+ps} b^j$ . Therefore,  $js \equiv 0 \pmod{p}$  for all  $j$  and hence,  $C_K(H) = \{1\}$ . This implies that  $Hom(H/[H, K], K)$  is the trivial group. Since  $K$  is abelian, by [2, Corollary 2.2, p. 490]  $C$  is the trivial group. Note that, each  $\alpha \in Aut(H)$  defined by  $\alpha(a) = a^i b^j$  and  $\alpha(b) = a^{pm} b^l$  can be expressed as a matrix  $\begin{pmatrix} i & j \\ m & l \end{pmatrix}$ , where  $0 \leq i \leq p^2 - 1, \gcd(p, i) = 1, 0 \leq m, j \leq p - 1$  and  $1 \leq l \leq p - 1$ . Also, let  $\delta \in Aut(K) \simeq \mathbb{Z}_{p-1}$  be defined by  $\delta(c) = c^r$ , where  $1 \leq r \leq p - 1$ . Now, if  $(\alpha, \delta) \in W$ , then (i)  $\alpha(a^c) = \alpha(a)^{\delta(c)}$  and (ii)  $\alpha(b^c) = \alpha(b)^{\delta(c)}$ . By (i),  $a^i b^j = \alpha(a) = \alpha(a^c) = \alpha(a)^{\delta(c)} = (a^i b^j)^{c^r} = a^i a^{prj} b^j = a^{i+prj} b^j$ . Thus,  $prj \equiv 0 \pmod{p^2}$  which implies that  $j = 0$ . Now, by (ii),  $a^{pi+pm} b^l = \alpha(a^p b) = \alpha(b^c) = \alpha(b)^{\delta(c)} = (a^{pm} b^l)^{c^r} = a^{pm} (b^l)^{c^r} = a^{pm} a^{prl} b^l = a^{pm+prl} b^l$ . Thus,  $i \equiv rl \pmod{p}$ . Let  $t$  be a primitive root of  $1 \pmod{p}$  and  $x = \left( \begin{pmatrix} t+p & 0 \\ 0 & t \end{pmatrix}, \delta_1 \right), y = \left( \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \delta_1 \right)$  and  $z = \left( \begin{pmatrix} t+p & 0 \\ 0 & 1 \end{pmatrix}, \delta_t \right)$ , where  $\delta_{\rho}(c) = c^{\rho}$ . Then  $W \simeq \langle x, y, z \mid x^{p(p-1)} = 1 = y^p = z^{p(p-1)}, xz = zx, xy = yx, zyz^{-1} = y^{t-1} \rangle$ . Therefore,  $W \simeq (\mathbb{Z}_p \times \mathbb{Z}_{p(p-1)}) \rtimes \mathbb{Z}_{p(p-1)}$  and so,  $E \simeq (\mathbb{Z}_p \times \mathbb{Z}_{p(p-1)}) \rtimes \mathbb{Z}_{p(p-1)}$ . Hence, by Theorem 2.1,  $Aut_K(G_3) \simeq (\mathbb{Z}_p \times \mathbb{Z}_{p(p-1)}) \rtimes \mathbb{Z}_{p(p-1)}$ .

**The group  $G_4$ .** Let  $H = \langle a, b \mid a^{p^2} = b^p = 1, ab = ba \rangle$  and  $K = \langle c \mid c^p = 1 \rangle$ . Then  $G_4 \simeq H \rtimes_{\phi} K$ , where  $\phi : K \rightarrow Aut(H)$  is given by  $\phi(c)(a) = a^{1+p}$  and  $\phi(c)(b) = b$ . Note that  $[H, K] = \langle a^p \rangle \simeq \mathbb{Z}_p$ . By the similar argument as in the case  $G_3$  above,  $C_K(H) = \{1\}$ . Since  $K$  is abelian, by [2, Corollary 2.2, p. 490]  $C$  is the trivial group. Note that, any  $\alpha \in Aut(H)$  defined by  $\alpha(a) = a^i b^j$  and  $\alpha(b) = a^{pm} b^l$  can be expressed as a matrix  $\begin{pmatrix} i & j \\ m & l \end{pmatrix}$ , where  $0 \leq i \leq p^2 - 1, \gcd(p, i) = 1, 0 \leq m, j \leq p - 1$  and  $1 \leq l \leq p - 1$ . Also, let  $\delta \in Aut(K) \simeq \mathbb{Z}_{p-1}$  be defined by  $\delta(c) = c^r$ , where  $1 \leq r \leq p - 1$ . Now, if  $(\alpha, \delta) \in W$ , then (i)  $\alpha(a^c) = \alpha(a)^{\delta(c)}$  and (ii)  $\alpha(b^c) = \alpha(b)^{\delta(c)}$ . Note that  $\alpha(b^c) = \alpha(b) = a^{pm} b^l$  and  $\alpha(b)^{\delta(c)} = (a^{pm} b^l)^{c^r} = (a^{pm})^{c^r} b^l = a^{pm(1+p)^r} b^l = a^{pm} b^l$ . Therefore, each  $\alpha \in Aut(H)$  satisfies (ii). Now, by (i),  $(a^i b^j)^{1+p} = \alpha(a^{1+p}) = \alpha(a^c) = \alpha(a)^{\delta(c)} = (a^i b^j)^{c^r} = (a^i)^{c^r} b^j = a^{i(1+p)^r} b^j$ . Thus,  $i(p+1) \equiv i(1+p)^r \pmod{p^2}$

mod  $p^2$ ) which implies that  $r = 1$ . Therefore,  $W \simeq \text{Aut}(H) \simeq \mathbb{Z}_{p-1} \times (((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_p) \rtimes \mathbb{Z}_{p-1})$ . Hence,  $E \simeq \mathbb{Z}_{p-1} \times (((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_p) \rtimes \mathbb{Z}_{p-1})$ . Thus,  $\text{Aut}_K(G_4) \simeq \mathbb{Z}_{p-1} \times (((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_p) \rtimes \mathbb{Z}_{p-1})$ .

**The group  $G_5$ .** Let  $H = \langle b, c \mid b^p = c^p = 1, bc = cb \rangle$  and  $K = \langle a \mid a^{p^2} = 1 \rangle$ . Then  $G_5 \simeq H \rtimes_{\phi} K$ , where  $\phi : K \rightarrow \text{Aut}(H)$  is given by  $\phi(a)(b) = b$  and  $\phi(a)(c) = b^{-1}c$ . Note that  $[H, K] = \langle b \rangle \simeq \mathbb{Z}_p$ . Also, if  $a^s \in C_K(H)$ , then  $b^i c^j = a^s b^i c^j a^{-s} = b^{i-js} c^j$ . Therefore,  $s \equiv 0 \pmod{p}$  and  $C_K(H) = \langle a^p \rangle$ . Since  $K$  is abelian, by [2, Corollary 2.2, p. 490],  $\gamma(h^k) = \gamma(h)$  is equivalent to  $\gamma \in \text{Hom}(H/[H, K], K)$ . Define  $\gamma_k \in \text{Hom}(H/[H, K], K)$  by  $\gamma_k(b) = 1$  and  $\gamma_k(c) = a^{pk}$  for all  $0 \leq k \leq p-1$ . Since  $[H, K] \subseteq \text{Ker} \gamma_k$ , it will induce a homomorphism from  $H/[H, K]$  to  $K$ . Let  $c = \begin{pmatrix} 1 & 0 \\ \gamma_1 & 1 \end{pmatrix}$ . Then, one can easily observe that  $\gamma_1(H) \subseteq C_K(H)$ . Therefore,  $\text{Hom}(H/[H, K], K) \simeq C = \langle c \rangle \simeq \mathbb{Z}_p$ . Note that, any  $\alpha \in \text{Aut}(H) \simeq \text{GL}(2, p)$  defined as,  $\alpha(b) = b^i c^j$  and  $\alpha(c) = b^l c^m$  can be represented as a matrix,  $\begin{pmatrix} i & j \\ l & m \end{pmatrix}$ , where  $0 \leq l, j \leq p-1$  and  $1 \leq i, m \leq p-1$ . Also, let  $\delta \in \text{Aut}(K) \simeq \mathbb{Z}_{p(p-1)}$  be defined by  $\delta(a) = a^r$ , where  $r \in \mathbb{Z}_{p^2}$ ,  $\text{gcd}(p, r) = 1$ . Now, if  $(\alpha, \delta) \in W$ , then (i)  $\alpha(b^a) = \alpha(b)^{\delta(a)}$  and (ii)  $\alpha(c^a) = \alpha(c)^{\delta(a)}$ . By (i),  $b^i c^j = \alpha(b) = \alpha(b^a) = \alpha(b)^{\delta(a)} = (b^i c^j)^{a^r} = b^i b^{-rj} c^j = b^{i-rj} c^j$ . Thus,  $rj \equiv 0 \pmod{p}$  which implies that  $j = 0$ . Now, by (ii),  $b^{-i+l} c^m = \alpha(b^{-1}c) = \alpha(c^a) = \alpha(c)^{\delta(a)} = (b^l c^m)^{a^r} = b^l (c^m)^{a^r} = b^l b^{-rm} c^m = b^{l-rm} c^m$ . Thus,  $i \equiv rm \pmod{p}$ . Let  $t$  be a primitive root of 1  $\pmod{p}$  and  $x = \left( \begin{pmatrix} t & 0 \\ 1 & t \end{pmatrix}, \delta_1 \right)$ , and  $y = \left( \begin{pmatrix} t+p & 0 \\ 0 & 1 \end{pmatrix}, \delta_t \right)$ , where  $\delta_{\rho}(a) = a^{\rho}$ . Then  $W \simeq \langle x, y \mid x^{p(p-1)} = 1, y^{p(p-1)} = 1, yxy^{-1} = x^{\lambda} \rangle$ , where  $x^{\lambda} = \begin{pmatrix} t & 0 \\ (t+p)^{-1} & t \end{pmatrix}$ . Then  $W \simeq \mathbb{Z}_{p(p-1)} \rtimes \mathbb{Z}_{p(p-1)}$  and so,  $E \simeq \mathbb{Z}_{p(p-1)} \rtimes \mathbb{Z}_{p(p-1)}$ . Hence,  $\text{Aut}_K(G_5) \simeq \mathbb{Z}_p \times (\mathbb{Z}_{p(p-1)} \rtimes \mathbb{Z}_{p(p-1)})$ .

**The group  $G_6$ .** Let  $H = \langle a, b \mid a^{p^2} = b^p = 1, ba = a^{1+pb} \rangle$  and  $K = \langle c \mid c^p = 1 \rangle$ . Then  $G_6 \simeq H \rtimes_{\phi} K$ , where  $\phi : K \rightarrow \text{Aut}(H)$  is given by  $\phi(c)(a) = ab$  and  $\phi(c)(b) = b$ . Note that  $[H, K] = \langle b^{-1}, a^p \rangle \simeq \mathbb{Z}_p \times \mathbb{Z}_p$ . By the similar argument as in the case  $G_3$  above,  $C_K(H) = \{1\}$  and hence  $C$  is the trivial group. Now,  $\alpha \in \text{Aut}(H)$  as given in [2] can be expressed as a matrix  $\begin{pmatrix} \eta & \beta \\ \xi & 1 \end{pmatrix}$ , where  $\eta(a) = a^i, 0 \leq i \leq p^2 - 1, \text{gcd}(p, i) = 1, \beta(b) = a^{pj}, 0 \leq j \leq p-1, \xi(a) = b^k, 0 \leq k \leq p-1$ , and  $1(b) = b$ . Also,  $\delta \in \text{Aut}(K)$  is given by  $\delta(c) = c^r, 1 \leq r \leq p-1$ . Now, if  $(\alpha, \delta) \in W$ , then (i)  $\alpha(a^c) = \alpha(a)^{\delta(c)}$  and (ii)  $\alpha(b^c) = \alpha(b)^{\delta(c)}$ . Note that  $\alpha(b^c) = \alpha(b) = a^{pj}b$  and  $\alpha(b)^{\delta(c)} = (a^{pj}b)^{c^r} = (c^r a c^{-r})^{pj} b = (a b^r)^{pj} b = a^{pj+rp \frac{pj(pj-1)}{2}} b^{pjr+1} = a^{pj} b$ . Therefore, each  $\alpha \in \text{Aut}(H)$  satisfies (ii). Now, by (i),  $a^{i+pj} b^{k+1} = \alpha(ab) = \alpha(a^c) = \alpha(a)^{\delta(c)} =$

$(a^i b^k)^{c^r} = (c^r a c^{-r})^i b^k = (a b^r)^i b^k = a^{i+r p \frac{i(i-1)}{2}} b^{ri+k}$ . Thus,  $ri \equiv 1 \pmod{p}$  which gives that  $i \equiv 2j+1 \pmod{p}$ . Therefore,  $i \in \{(2j+1) + \lambda p \mid \lambda \in \mathbb{Z}_p\}$ . Let  $t$  be a primitive root of  $1 \pmod{p}$  and  $x = \left( \begin{pmatrix} t+p & 0 \\ 0 & 1 \end{pmatrix}, \delta_t \right)$ ,  $y = \left( \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \delta_1 \right)$ , and  $z = \left( \begin{pmatrix} 1+p & 0 \\ 0 & 1 \end{pmatrix}, \delta_1 \right)$ , where  $\delta_\rho(a) = a^\rho$ . Then  $W \simeq \langle x, y, z \mid x^{p(p-1)} = 1 = y^p = z^p, xyx^{-1} = y^e, xz = zx, yz = zy \rangle$ , where  $y^e = \begin{pmatrix} 1 & 0 \\ (t+p)^{-1} & 1 \end{pmatrix}$ . Hence,  $W \simeq \mathbb{Z}_p \times ((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_{p-1})$  and so,  $E \simeq \mathbb{Z}_p \times ((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_{p-1})$ . Thus,  $\text{Aut}_K(G_6) \simeq \mathbb{Z}_p \times ((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_{p-1})$ .

**The group  $G_7$ .** Let  $H = \langle a, b \mid a^{p^2} = b^p = 1, ba = a^{1+pb} \rangle$  and  $K = \langle c \mid c^p = 1 \rangle$ . Then,  $G_7 \simeq H \rtimes_\phi K$ , where  $\phi : K \rightarrow \text{Aut}(H)$  is given by  $\phi(c)(a) = a^{1+pb}$  and  $\phi(c)(b) = a^p b$ . Note that  $[H, K] = \langle b, a^p \rangle \simeq \mathbb{Z}_p \times \mathbb{Z}_p$ . By the similar argument as in the case  $G_3$  above,  $C_K(H) = \{1\}$  and hence  $C$  is the trivial group. Each  $\alpha \in \text{Aut}(H)$  can be expressed as a matrix  $\begin{pmatrix} \eta & \beta \\ \xi & 1 \end{pmatrix}$ , where  $\eta(a) = a^i, 0 \leq i \leq p^2 - 1, \gcd(i, p) = 1$ ,  $\beta(b) = a^{pj}, \xi(a) = b^k, 0 \leq j, k \leq p-1$ , and  $1(b) = b$ . Also,  $\delta \in \text{Aut}(K)$  is given by  $\delta(c) = c^r, 1 \leq r \leq p-1$ . Now, if  $(\alpha, \delta) \in W$ , then (i)  $\alpha(a^c) = \alpha(a)^{\delta(c)}$  and (ii)  $\alpha(b^c) = \alpha(b)^{\delta(c)}$ .

By (ii),  $a^{pi+pb} = \alpha(a^p b) = \alpha(b^c) = \alpha(b)^{\delta(c)} = (a^{pj} b)^{c^r} = (c^r a c^{-r})^{pj} (c^r b c^{-r}) = (a^{1+p \frac{r(r+1)}{2}} b^r)^{pj} (a^{rp} b) = a^{p^2 j \frac{r(r+1)}{2}} (a b^r)^{pj} a^{rp} b = a^{pj+pr} b$ . Thus  $i \equiv r \pmod{p}$ . Now, by (i),  $a^{i(1+p)+pj} b^{k+1} = \alpha(a^{1+pb}) = \alpha(a^c) = \alpha(a)^{\delta(c)} = (a^i b^k)^{c^r} = (c^r a c^{-r})^i (c^r b c^{-r})^k = a^{i+r p i \frac{r+i}{2}} b^{ri} (a^{rp} b)^k = a^{i+r p i \frac{r+i}{2} + r p k} b^{ri+k}$ . Thus,  $ri \equiv 1 \pmod{p}$  and  $ip + pj \equiv p r i \frac{r+i}{2} + r p k \pmod{p^2}$  implies that  $i + j \equiv r + r k \pmod{p}$ . So,  $j \equiv r k \pmod{p}$ . Using  $r \equiv i \pmod{p}$  and  $ri \equiv 1 \pmod{p}$ , we get  $i^2 \equiv 1 \pmod{p^2}$ .

Let  $t$  be a primitive root of  $1 \pmod{p}$  and  $x = \left( \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \delta_1 \right)$ ,  $y = \left( \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \delta_1 \right)$  and  $z = \left( \begin{pmatrix} t+p & 0 \\ 0 & 1 \end{pmatrix}, \delta_t \right)$ , where  $\delta_\rho(a) = a^\rho$ . Then  $W \simeq \langle x, y, z \mid x^p = 1 = y^2 = z^{p(p-1)}, xy = yx^{-1}, xzx^{-1} = x^t, yz = zy \rangle \simeq \mathbb{Z}_{p(p-1)} \times (\mathbb{Z}_p \rtimes \mathbb{Z}_2)$  and so,  $E \simeq \mathbb{Z}_{p(p-1)} \times (\mathbb{Z}_p \rtimes \mathbb{Z}_2)$ . Hence,  $\text{Aut}_K(G_7) \simeq \mathbb{Z}_{p(p-1)} \times (\mathbb{Z}_p \rtimes \mathbb{Z}_2) \simeq D_{2p} \times \mathbb{Z}_{p(p-1)}$ , where  $D_{2p}$  is the dihedral group of order  $2p$ .

**The group  $G_8$ .** Let  $H = \langle a, b \mid a^{p^2} = b^p = 1, ba = a^{1+pb} \rangle$  and  $K = \langle c \mid c^p = 1 \rangle$ . Then  $G_8 \simeq H \rtimes_\phi K$ , where  $\phi : K \rightarrow \text{Aut}(H)$  is given by  $\phi(c)(a) = a^{1+dp}$  and  $\phi(c)(b) = a^{dp} b, d \not\equiv 0, 1 \pmod{p}$ . By the similar argument as for the group  $G_7$ , we get,  $C$  is the trivial group and  $E \simeq \mathbb{Z}_{p(p-1)} \times (\mathbb{Z}_p \rtimes \mathbb{Z}_2)$ . Hence,  $\text{Aut}_K(G_8) \simeq D_{2p} \times \mathbb{Z}_{p(p-1)}$ .

**The group  $G_9$ .** Let  $H = \langle a, b, c \mid a^p = b^p = c^p = 1, ab = ba, bc = cb, ac = ca \rangle$ , and  $K = \langle d \mid d^p = 1 \rangle$ . Then  $G_9 \simeq H \rtimes_{\phi} K$ , where  $\phi : K \rightarrow \text{Aut}(H)$  is given by  $\phi(d)(a) = a$ ,  $\phi(d)(b) = b$ , and  $\phi(d)(c) = ac$ .

Note that  $[H, K] = \langle a \rangle \simeq \mathbb{Z}_p$ . By the similar argument as in the case  $G_3$  above,  $C_K(H) = \{1\}$  and hence  $C$  is the trivial group. Note that,  $\text{Aut}(H) \simeq GL(3, p)$  and  $\text{Aut}(K) \simeq \mathbb{Z}_{p-1}$ . So, any automorphism  $\alpha \in \text{Aut}(H)$  can be identified as an element in  $GL(3, p)$ . Let  $\alpha \in \text{Aut}(H)$  and  $\delta \in \text{Aut}(K)$  be defined as,  $\alpha(a) = a^i b^j c^k$ ,  $\alpha(b) = a^l b^m c^n$ ,  $\alpha(c) = a^{\lambda} b^{\mu} c^{\rho}$ , and  $\delta(d) = d^r$ , where  $1 \leq i, m, \rho, r \leq p-1$  and  $0 \leq j, k, l, n, \lambda, \mu \leq p-1$ . Now, if  $(\alpha, \delta) \in W$ , then (i)  $\alpha(a^d) = \alpha(a)^{\delta(d)}$ , (ii)  $\alpha(b^d) = \alpha(b)^{\delta(d)}$  and (iii)  $\alpha(c^d) = \alpha(c)^{\delta(d)}$ .

Note that,  $d^r c d^{-r} = a^r c$ . By (i),  $a^i b^j c^k = \alpha(a) = \alpha(a^d) = \alpha(a)^{\delta(d)} = (a^i b^j c^k)^{d^r} = a^i b^j (d^r c d^{-r})^k = a^i b^j (a^r c)^k = a^{i+r k} b^j c^k$ . Therefore,  $r k \equiv 0 \pmod{p}$  which implies that  $k = 0$ . Now, by (ii),  $a^l b^m c^n = \alpha(b) = \alpha(b^d) = \alpha(b)^{\delta(d)} = (a^l b^m c^n)^{d^r} = a^{l+r n} b^m c^n$ . Therefore,  $r n \equiv 0 \pmod{p}$  which implies that  $n = 0$ . By (iii),  $a^{i+\lambda} b^{j+\mu} c^{\rho} = \alpha(ac) = \alpha(c^d) = \alpha(c)^{\delta(d)} = (a^{\lambda} b^{\mu} c^{\rho})^{d^r} = a^{\lambda+r \rho} b^{\mu} c^{\rho}$ . Thus,  $i = r \rho$  and  $j = 0$ . So, we have,

$$\alpha = \begin{pmatrix} r\rho & 0 & 0 \\ l & m & 0 \\ \lambda & \mu & \rho \end{pmatrix}. \text{ Let } t \text{ be a primitive root of } 1 \pmod{p} \text{ and } u = \begin{pmatrix} \begin{pmatrix} t & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & t \end{pmatrix}, \delta_1 \\ \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \delta_1 \end{pmatrix}, v = \begin{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & t & 0 \\ 0 & 0 & 1 \end{pmatrix}, \delta_1 \\ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}, \delta_1 \end{pmatrix}, w = \begin{pmatrix} \begin{pmatrix} t & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \delta_t \\ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \delta_1 \end{pmatrix}, x =$$

where  $\delta_s(d) = d^s$ . Then  $W \simeq \langle u, v, w, x, y, z \mid u^{p-1} = 1 = v^{p-1} = w^{p-1} = x^p = y^p = z^p, uv = vu, uw = wu, uy = yu, vw = wv, vy = yv, wz = zw, xy = yx, yz = zy, uxu^{-1} = x^{t-1}, uzu^{-1} = z^t, vxv^{-1} = x^t, vzv^{-1} = z^{t-1}, wxw^{-1} = x^{t-1}, wyw^{-1} = y^{t-1}, zx = xyz \rangle \simeq (((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_p) \times \mathbb{Z}_{p-1}) \rtimes (\mathbb{Z}_{p-1} \times \mathbb{Z}_{p-1})$  and so,  $E \simeq (((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_p) \times \mathbb{Z}_{p-1}) \rtimes (\mathbb{Z}_{p-1} \times \mathbb{Z}_{p-1})$ . Hence,  $\text{Aut}_K(G_9) \simeq (((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_p) \times \mathbb{Z}_{p-1}) \rtimes (\mathbb{Z}_{p-1} \times \mathbb{Z}_{p-1})$ .

**The group  $G_{10}$ .** Let  $H = \langle a, b, c \mid a^p = b^p = c^p = 1, ab = ba, bc = cb, ac = ca \rangle$  and  $K = \langle d \mid d^p = 1 \rangle$ . Then,  $G_{10} \simeq H \rtimes_{\phi} K$ , where  $\phi : K \rightarrow \text{Aut}(H)$  is given by  $\phi(d)(a) = a$ ,  $\phi(d)(b) = ab$ , and  $\phi(d)(c) = bc$ .

Note that  $[H, K] = \langle a, b \rangle \simeq \mathbb{Z}_p \times \mathbb{Z}_p$ . By the similar argument as above  $C$  is the trivial group. Note that,  $\text{Aut}(H) \simeq GL(3, p)$  and  $\text{Aut}(K) \simeq \mathbb{Z}_{p-1}$ . So, any automorphism  $\alpha \in \text{Aut}(H)$  can be identified as an element in  $GL(3, p)$ . Let  $\alpha \in \text{Aut}(H)$  and  $\delta \in \text{Aut}(K)$  be defined as,  $\alpha(a) = a^i b^j c^k$ ,  $\alpha(b) = a^l b^m c^n$ ,  $\alpha(c) = a^{\lambda} b^{\mu} c^{\rho}$ , and  $\delta(d) = d^r$ , where  $1 \leq i, m, \rho, r \leq p-1$  and  $0 \leq j, k, l, n, \lambda, \mu \leq p-1$ . Now, if  $(\alpha, \delta) \in W$ , then (i)  $\alpha(a^d) = \alpha(a)^{\delta(d)}$ , (ii)  $\alpha(b^d) = \alpha(b)^{\delta(d)}$  and (iii)  $\alpha(c^d) = \alpha(c)^{\delta(d)}$ .



Note that,  $d^r b d^{-r} = a^r b$  and  $d^r c d^{-r} = a^{\frac{r(r-1)}{2}} b^r c$ . By (i),  $a^i b^j c^k = \alpha(a) = \alpha(a^d) = \alpha(a)^{\delta(d)} = (a^i b^j c^k)^{d^r} = a^i (d^r b d^{-r})^j (d^r c d^{-r})^k = a^i (a^r b)^j (a^{\frac{r(r-1)}{2}} b^r c)^k = a^{i+rj+k\frac{r(r-1)}{2}} b^{j+r k} c^k$ . Thus  $k = 0$  and  $j = 0$ . Now, by (ii),  $a^{i+l} b^m c^n = \alpha(ab) = \alpha(b^d) = \alpha(b)^{\delta(d)} = (a^l b^m c^n)^{d^r} = a^{l+rm+n\frac{r(r-1)}{2}} b^{m+rn} c^n$ . Thus,  $n = 0$  and  $i = rm$ . By (iii),  $a^{l+\lambda} b^{m+\mu} c^\rho = \alpha(bc) = \alpha(c^d) = \alpha(c)^{\delta(d)} = (a^\lambda b^\mu c^\rho)^{d^r} = a^{\lambda+r\mu+\rho\frac{r(r-1)}{2}} b^{\mu+r\rho} c^\rho$ . Thus,  $m = r\rho$  and  $l = r\mu + \rho\frac{r(r-1)}{2} \pmod{p}$ . So, we have,  $\alpha = \begin{pmatrix} r^2\rho & 0 & 0 \\ l & r\rho & 0 \\ \lambda & \mu & \rho \end{pmatrix}$ , where

$l = r\mu + \rho\frac{r(r-1)}{2} \pmod{p}$ . Let  $t$  be a primitive root of  $1 \pmod{p}$  and  $x = \left( \begin{pmatrix} t & 0 & 0 \\ 0 & t & 0 \\ 0 & 0 & t \end{pmatrix}, \delta_1 \right)$ ,  $y = \left( \begin{pmatrix} t^2 & 0 & 0 \\ 0 & t & 0 \\ 0 & 0 & 1 \end{pmatrix}, \delta_t \right)$  and  $z = \left( \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}, \delta_1 \right)$ ,

where  $\delta_s(d) = d^s$ . Note that,  $\langle z \rangle = \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \right\rangle$  is an abelian group of order  $p^2$ . Therefore,  $W \simeq \langle x, y, z \mid x^{p-1} = y^{p-1} = z^p, xy = yx, xz = zx, yzy^{-1} = z^u \rangle$ , where  $z^u = \begin{pmatrix} 1 & 0 & 0 \\ t^{-1} & 1 & 0 \\ t^{-2} & t^{-1} & 1 \end{pmatrix}$ . Thus

$W \simeq (\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes (\mathbb{Z}_{p-1} \times \mathbb{Z}_{p-1})$  and so,  $E \simeq (\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes (\mathbb{Z}_{p-1} \times \mathbb{Z}_{p-1})$ . Hence,  $\text{Aut}_K(G_{10}) \simeq \mathbb{Z}_{p-1} \times ((\mathbb{Z}_p \times \mathbb{Z}_p) \rtimes \mathbb{Z}_{p-1})$ .

#### ACKNOWLEDGEMENTS

The first author is supported by the Junior Research Fellowship of UGC, India.

#### REFERENCES

- [1] Burnside, W., Theory of groups of finite order, *Cambridge University Press*, 1897.
- [2] Bidwell, J. N. S., and Curran, M. J., The automorphism group of a split metacyclic  $p$ -group, *Arch. Math.*, vol. 87, pp. 488 - 497, 2006.
- [3] Curran, M. J., The automorphism group of a split metacyclic 2 - group, *Arch. Math.*, vol. 89, pp. 10 - 23, 2007, doi:10.1007/s00013-007-2107-5.
- [4] Curran, M. J., Automorphisms of semidirect products, *Math. Proc. R. Ir. Acad.*, 108A, pp. 205 - 210, 2008.
- [5] Dietz, J., Automorphisms of products of groups, in *CM. Campbell, M.R. Quick, E.F. Robertson, G.C. Smith (eds), Groups St Andrews, LMS Lecture Notes Series*, vol. 339, pp. 28 - 305, 2005.
- [6] Hsu N.C., The group of automorphisms of the holomorph of a group, *Pacific J. Math.* 11 (1961), pp. 999-1012.
- [7] Lal, R., Trnasversls in Groups *J. Algebra*, 181, pp. 70 - 81, 1996.

DEPARTMENT OF MATHEMATICS, CENTRAL UNIVERSITY OF RAJASTHAN, RAJASTHAN

Email address: vermarattan789@gmail.com, vplkakk@gmail.com