

UNREFINABLE PARTITIONS INTO DISTINCT PARTS IN A NORMALIZER CHAIN

RICCARDO ARAGONA, ROBERTO CIVINO, NORBERTO GAVIOLI,
AND CARLO MARIA SCOPPOLA

ABSTRACT. In a recent paper on a study of the Sylow 2-subgroups of the symmetric group with 2^n elements it has been show that the growth of the first $(n - 2)$ consecutive indices of a certain normalizer chain is linked to the sequence of partitions of integers into distinct parts. Unrefinable partitions into distinct parts are those in which no part x can be replaced with integers whose sum is x obtaining a new partition into distinct parts. We prove here that the $(n - 1)$ -th index of the previously mentioned chain is related to the number of unrefinable partitions into distinct parts satisfying a condition on the minimal excludant.

1. INTRODUCTION

The sequence (b_j) of the number of partitions of integers into distinct parts has been extensively studied in past and recent years and is a well-understood integer sequence [Eul48, And94] appearing in different areas of mathematics. Recently, triggered by a problem in algebraic cryptography related to translation subgroups in the symmetric group with 2^n elements [CDVS06, ACGS19, CBS19, CCS21], it has been shown that such a sequence is related to the growth of the indices of consecutive terms in a chain of normalizers [ACGS21]. More precisely, let Σ_n be a Sylow 2-subgroup of the symmetric group $\text{Sym}(2^n)$ and T be an elementary abelian regular subgroup of Σ_n . Defining $N_n^0 = N_{\Sigma_n}(T)$ and recursively $N_n^i = N_{\Sigma_n}(N_n^{i-1})$, the authors proved that the number

$$\log_2 |N_n^i : N_n^{i-1}|$$

is independent of n for $1 \leq i \leq n - 2$, and is equal to the $(i + 2)$ -th term of the sequence (a_j) of the partial sums of (b_j) (cfr. Table 1).

The result is obtained by proving that the terms in the chain are *saturated* subgroups, i.e. generated by *rigid commutators*, a family of left-normed commutators involving a special set of generators of Σ_n . We invite the interested reader to refer to Aragona et al. [ACGS21], where rigid commutators and saturated groups are introduced and described in detail.

When $i > n - 2$, the behavior of the chain does not seem to show any recognizable pattern and the study of its combinatorial nature is still open. Nonetheless, further

2010 *Mathematics Subject Classification.* 20B30, 20B35, 20D20, 11P81, 05A17.

Key words and phrases. Symmetric group on 2^n elements; Partitions into distinct parts; Unrefinable partitions; Minimal excludant; Sylow 2-subgroups.

All the authors are members of INDAM-GNSAGA (Italy) and of the group “Crittografia e Codici” of the Italian Mathematical Union. R. Civino is partially funded by the Centre of Excellence EX-EMERGE at University of L’Aquila. N. Gavioli is a member of the Centre of Excellence EX-EMERGE at University of L’Aquila.

j	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
b_j	0	0	0	1	1	2	3	4	5	7	9	11	14	17	21
a_j	0	0	0	1	2	4	7	11	16	23	32	43	57	74	95

TABLE 1. First values of the sequences (b_j) and (a_j)

investigations on experimental evidences led us to notice that the first exception to the rule, i.e. $\log_2 |N_n^{n-1} : N_n^{n-2}|$, is linked to the number of partitions of n into distinct parts which do not admit a *refinement*, i.e. those partitions $n = p_1 + p_2 + \dots + p_t$ where no integer p_i can be replaced by $l \geq 2$ distinct integers q_1, q_2, \dots, q_l , whose sum is a_i and such that the resulting partition

$$n = p_1 + p_2 + \dots + p_{i-1} + (q_1 + q_2 + \dots + q_l) + p_{i+1} + \dots + p_t$$

is still a partition into distinct parts. In our knowledge, *unrefinable* partitions have not been investigated so far.

We prove here (cfr. Theorem 9) that a transversal of N_n^{n-2} in N_n^{n-1} is made of rigid commutators which are in one-to-one correspondence with unrefinable partitions into distinct parts with a condition on their minimal excludants, where the minimal excludant of a partition is the least positive integer which does not appear in the partition.

A brief summary on rigid commutators and saturated subgroups is presented in Sec. 2, which also contains a representation of the Sylow 2-subgroup of the symmetric group and to the precise definition of the normalizer chain under investigation. The proof of Theorem 9 and considerations on unrefinable partitions and rigid commutators can be found in Sec. 3.

2. PRELIMINARIES

Let n be a non-negative integer. Let us define a set of permutations $\{s_i \mid 1 \leq i \leq n\}$ which generates a Sylow 2-subgroup of the symmetric group on 2^n letters.

The Sylow 2-subgroup. Let us consider the set

$$\mathcal{T}_n = \{w_1 \dots w_n \mid w_i \in \{0, 1\}\}$$

of binary words of length n , where \mathcal{T}_0 contains only the empty word. The infinite rooted binary tree \mathcal{T} is defined as the graph whose vertices are $\bigcup_{j \geq 0} \mathcal{T}_j$ and where two vertices, say $w_1 \dots w_n$ and $v_1 \dots v_m$, are connected by an edge if $|m - n| = 1$ and $w_i = v_i$ for $1 \leq i \leq \min(m, n)$. The empty word is the root of the tree and it is connected with both the two words of length 1.

We can define a sequence $\{s_i\}_{i \geq 1}$ of automorphisms of this tree. Each s_i necessarily fixes the root, which is the only vertex of degree 2. The automorphism s_1 changes the value w_1 of the first letter of every non-empty word into $\bar{w}_1 \stackrel{\text{def}}{=} (w_1 + 1) \bmod 2$ and leaves the other letters unchanged. If $i \geq 2$, we define

$$(w_1 \dots w_k) s_i \stackrel{\text{def}}{=} \begin{cases} \text{empty word} & \text{if } n = 0 \\ w_1 \dots \bar{w}_i \dots w_k & \text{if } k \geq i \text{ and } w_1 = \dots = w_{i-1} = 0 \\ w_1 \dots w_k & \text{otherwise.} \end{cases}$$

In general, s_i leaves a word unchanged unless the word has length at least i and the letters preceding the i -th one are all zero, in which case the i -th letter is increased by 1 modulo 2. If $i \leq n$ and the word $w_1 \dots w_n \in \mathcal{T}_n$ is identified with the integer $1 + \sum_{i=1}^n 2^{n-i} w_i \in \{1, \dots, 2^n\}$, then s_i acts on \mathcal{T}_n as the permutation whose cyclic decomposition is

$$\prod_{j=1}^{2^{n-i}} (j, j + 2^{n-i})$$

which has order 2. In particular, the group $\langle s_1, \dots, s_n \rangle$ acts faithfully on the set \mathcal{T}_n , whose cardinality is 2^n , as a Sylow 2-subgroup Σ_n of the symmetric group $\text{Sym}(2^n)$.

Rigid commutators. The *commutator* of two elements h and k in a group G is defined as $[h, k] \stackrel{\text{def}}{=} h^{-1}k^{-1}hk = h^{-1}h^k$. The *left-normed commutator* of the m elements $g_1, \dots, g_m \in G$ is the usual commutator if $m = 2$ and is recursively defined by

$$[g_1, \dots, g_{n-1}, g_n] \stackrel{\text{def}}{=} [[g_1, \dots, g_{n-1}], g_n]$$

if $m \geq 3$. In this paper we will only focus on left-normed commutators in s_1, \dots, s_n , therefore, for the sake of simplicity, we write $[i_1, \dots, i_k]$ to denote the left-normed commutator $[s_{i_1}, \dots, s_{i_k}]$, when $k \geq 2$. We also write $[i]$ to denote the element s_i and we set $[\]$ to be the identity permutation.

Definition 1. A left-normed commutator $r = [i_1, \dots, i_k] \in \Sigma_n$ is called *rigid* when we have $i_1 > i_2 > \dots > i_k$ or $r = [\]$. The set of all the rigid commutators of Σ_n is denoted by \mathcal{R} and we let $\mathcal{R}^* \stackrel{\text{def}}{=} \mathcal{R} \setminus \{[\]\}$.

Definition 2. A subgroup of Σ_n is said to be *saturated* if it is generated by rigid commutators.

Let us define a special set $\{t_1, t_2, \dots, t_n\}$ of rigid commutators where

$$t_i \stackrel{\text{def}}{=} [i, i-1, \dots, 2, 1]. \quad (2.1)$$

Remark 1. The saturated subgroup $T \stackrel{\text{def}}{=} \langle t_1, t_2, \dots, t_n \rangle$ is an elementary abelian regular subgroup of Σ_n .

Theorem 3 ([ACGS21]). *The normalizer N in Σ_n of a saturated subgroup of Σ_n is also saturated, provided that N contains T .*

Let $i_1 > i_2 > \dots > i_k$ and let $X = \{1, \dots, i_1\} \setminus \{i_1, \dots, i_k\}$. For the purposes of this work it is more convenient to use the notation

$$\vee[i_1; X]$$

to denote the rigid commutator $[i_1, \dots, i_k]$. The commutator of two rigid commutators is again a rigid commutator which, in the above notation, can be computed in the following way:

Proposition 4 ([ACGS21]). *Let $1 \leq a, b \leq n$ and let I and J be subsets of $\{1, 2, \dots, a-1\}$ and $\{1, 2, \dots, b-1\}$ respectively. Then*

$$[\vee[a; I], \vee[b; J]] = \begin{cases} \vee[\max(a, b); (I \cup J) \setminus \{\min(a, b)\}] & \text{if } \min(a, b) \in I \cup J \\ 1 & \text{otherwise.} \end{cases}$$

The normalizer chain. The *normalizer chain starting at T* is defined as

$$N_n^i \stackrel{\text{def}}{=} \begin{cases} N_{\Sigma_n}(T) & \text{if } i = 0, \\ N_{\Sigma_n}(N_n^{i-1}) & \text{if } i \geq 1. \end{cases} \quad (2.2)$$

Notice that N_n^0 is the Sylow 2-subgroup U_n of the affine group $\text{AGL}(2, n)$. It is worth noticing here that $N_{\Sigma_n}(N_n^i) = N_{\text{Sym}(2^n)}(N_n^i)$, for all $i \geq 0$ [ACGS20].

In order to show where partitions of integers come into play, let us briefly describe the generators of the first $n - 2$ normalizers of the chain. First, let us determine the permutations in Σ_n normalizing T : for $1 \leq j < i \leq n$ let us define

$$X_{ij} \stackrel{\text{def}}{=} \{1, \dots, i\} \setminus \{j\} \text{ and } u_{ij} \stackrel{\text{def}}{=} [X_{ij}] = \vee[i; \{j\}] \in \mathcal{R}^*, \quad (2.3)$$

and let us set

$$\mathcal{U}_n \stackrel{\text{def}}{=} \{t_1, \dots, t_n, u_{ij} \mid 1 \leq j < i \leq n\} \subseteq \mathcal{R}^*.$$

Next, let us define

$$\mathcal{W}_{ij} \stackrel{\text{def}}{=} \left\{ \vee[i; I] \in \mathcal{R}^* \mid I \subseteq \{1, 2, \dots, i-1\}, |I| \geq 2, \sum_{x \in I} x = j \right\} \quad (2.4)$$

for each $1 \leq i \leq n$ and j , and

$$\mathcal{N}_n^i \stackrel{\text{def}}{=} \begin{cases} \mathcal{U}_n & \text{if } i = 0 \\ \mathcal{N}_n^{i-1} \dot{\cup} \left(\dot{\bigcup}_{j=1}^i \mathcal{W}_{n+j-i, j+2} \right) & \text{for } i > 0. \end{cases} \quad (2.5)$$

Note that, if $j \leq i - 2$, then $|\mathcal{W}_{i,j}| = b_j$, i.e. it corresponds to the number of partitions of j into at least two distinct parts.

The previous elements generate the subgroups in the normalizer chain:

Theorem 5 ([ACGS21]). *For $i \leq n - 2$, the group $\langle \mathcal{N}_n^i \rangle$ is the i -th term N_n^i of the normalizer chain. In particular, the subgroup N_n^i of Σ_n is generated by U_n and the rigid commutators $\vee[a; X]$ such that*

- $|X| \geq 2$,
- $\sum_{x \in X} x \leq i + 2 - (n - a)$.

The following straightforward consequence is derived:

Corollary 6 ([ACGS21]). *For $1 \leq i \leq n - 2$, the number $\log_2 |N_n^i : N_n^{i-1}|$ is independent of n . It equals the $(i + 2)$ -th term of the sequence (a_j) of the partial sums of the sequence (b_j) counting the number of partitions of j into at least two distinct parts.*

3. UNREFINABLE PARTITIONS

Every non-empty finite subset $X \subseteq \mathbb{N} \setminus \{0\}$ represents a partition of the integer

$$\sum X \stackrel{\text{def}}{=} \sum_{x \in X} x$$

into *distinct* parts. Some partitions, e.g. $7 = 1 + 2 + 4$, are not refinable, some others are. For example, in $10 = 1 + 4 + 5$, the part 5 can be replaced by $2 + 3$ obtaining a partition of $10 = 1 + 2 + 3 + 4$ into distinct parts. More precisely:

Definition 7. A finite non-empty subset $X \subseteq \mathbb{N} \setminus \{0\}$ is *refinable* if there exists $x \in X$ and a subset $Y \subseteq \mathbb{N} \setminus (X \cup \{0\})$ such that $x = \sum Y$. If so, X induces a *refinable partition* into distinct parts of $\sum X$. Equivalently, $(X \setminus \{x\}) \dot{\cup} Y$ is again a partition into distinct parts of $\sum X$, called a *refinement* of X . We say that X is an *unrefinable* partition of $\sum X$ into distinct parts if X is not refinable.

Remark 2. Notice that if X is a refinable partition, then there exists Y as in Definition 7, $|Y| = 2$, such that $x = \sum Y$.

Let us now recall the concept of minimal excludant, better known from its use in combinatorial game theory [Spr35, Gru39, FP15] and recently considered in the theory of partitions of integers [AN19, AN20, BM20].

Definition 8. The *minimal excludant* $\text{mex}(X)$ of a set X of positive integers is the least positive integer that is not an element of X .

Rigid commutators and unrefinable partitions are linked by the following consideration depending on Proposition 4. Suppose that $a > 1$ and that $X \subseteq \{1, \dots, a-1\}$ is refinable. If $x \in X$ and Y are as in Definition 7, then

$$[\vee[a; X], \vee[x; Y]] = \vee[a; Z]$$

where $Z \stackrel{\text{def}}{=} (X \setminus \{x\}) \dot{\cup} Y$ is the refinement of X obtained by *replacing* x with Y . Notice that $\sum Z = \sum X$. Since $\sum Y = x$, by Theorem 5 the rigid commutator $\vee[x; Y]$ belongs to N_n^{n-2} for $n \geq x$. Conversely, if

$$[\vee[a; X], \vee[x; Y']] = \vee[a; Z]$$

for some Y' and if $\sum Z = \sum X$, then $x \in X$ and $Z = (X \setminus \{x\}) \dot{\cup} Y$ is a refinement of X , where $Y \stackrel{\text{def}}{=} Y' \setminus X$ is such that $\sum Y = x$.

Bearing this in mind, let us prove the following:

Theorem 9. *Let $a > 1$, $X \subseteq \{1, 2, \dots, a-1\}$ and let $k = \text{mex}(X)$ be the minimal excludant of X . The rigid commutator $\vee[a; X]$ belong to $N_n^{n-1} \setminus N_n^{n-2}$ if and only if*

- (1) $\sum X = a + 1$,
- (2) X is an unrefinable partition,
- (3) $a \leq n < a + k$.

Proof. Suppose that $c \stackrel{\text{def}}{=} \vee[a; X]$ belongs to $N_n^{n-1} \setminus N_n^{n-2}$ and let $b = \sum X$ and $h = \min(X)$. By Theorem 5, since $c \notin N_n^{n-2}$, we have $b \geq a + 1$. Let us define

$$d \stackrel{\text{def}}{=} \begin{cases} u_{h, h-1} & \text{if } h > 1 \\ t_1 & \text{if } h = 1, \end{cases}$$

where $u_{h, h-1}$ and t_1 are respectively defined in Eq. (2.3) and Eq. (2.1). From Proposition 4, the commutator $[c, d]$ is rigid of the form $\vee[a; Y]$, where $\sum Y = \sum X - 1 = b - 1$. Since $d \in N_n^{n-2}$, we have that $[c, d] \in N_n^{n-2}$, and in particular $\sum Y = b - 1 \leq a$. We already know that $a \leq b - 1$, and so we have the equality $\sum X = b = a + 1$, as claimed in (1).

Suppose now that X is refinable. We have already observed that there exist $x \in X$ and a rigid commutator $\vee[x; Y] \in N_n^{n-2}$, i.e. $\sum Y = x$, such that

$$[c, \vee[x; Y]] = \vee[a; Z],$$

with $\sum Z = \sum X = a + 1$. This implies that $\vee[a; Z] \notin N_n^{n-2}$, and so c does not normalize N_n^{n-2} , in contradiction to the assumption. This proves (2).

Finally, we obviously have $k \leq a$. Let us prove that the equality cannot hold. Suppose indeed that $k = a$, then X is the set $\{1, \dots, a-1\}$. By (1), we have the quadratic equation

$$a + 1 = \sum X = (a^2 - a)/2$$

which admits no integer solutions. Now, let $l \stackrel{\text{def}}{=} a + k$ and let us prove that $n < l$. Suppose that $n \geq l$. By definition of l and by Theorem 5 we have that $d \stackrel{\text{def}}{=} \vee[l; \{a, k\}] \in N_n^{n-2} \setminus N_n^{n-3}$. Since $c \in N_n^{n-1}$, then $N_n^{n-2} \ni [c, d] = \vee[l; X \dot{\cup} \{k\}]$. This is possible only if $l \geq \sum X + k = a + 1 + k = l + 1$, a contradiction. The claim (3) is then proved.

Conversely, suppose that (1), (2) and (3) hold and let us consider a generic rigid commutator in N_n^{n-2}

$$d \stackrel{\text{def}}{=} \vee[m; V] \in N_n^{n-2}.$$

We have $\sum V \leq m$ and $m \leq n < a + k$. Let us prove first that $[c, d] \in N_n^{n-2}$. To do so we may assume $[c, d] \neq 1$, otherwise there is nothing to prove. Three cases need to be distinguished.

Assume first that $m < a$. Since $[c, d] \neq 1$, then $[c, d] = \vee[a; Z]$, where $Z = (X \setminus \{m\}) \cup V$. Notice that $\sum Z < \sum X = a + 1$, where the equality is excluded since Z would otherwise represent a refinement of X , which is unrefinable by hypothesis. Hence $\sum Z \leq a$, and so $[c, d] \in N_n^{n-2}$, as claimed. The case $m = a$ has already been discussed since it corresponds to $[c, d] = 1$. To conclude, suppose that $a < m \leq n < a + k$. Since $[c, d] \neq 1$, by Proposition 4 we have $a \in V$, and consequently

$$\sum (V \setminus \{a\}) \leq m - a < k. \quad (3.1)$$

By Eq. (3.1), we have that $V \setminus \{a\}$ contains only elements smaller than k and, since $k = \text{mex}(X)$ is minimum positive integer in the complementary set of X in \mathbb{N} , the set $V \setminus \{a\}$ has to be a subset of X . Thus we have $[c, d] = \vee[m; Z]$, where $Z = X \cup (V \setminus \{a\}) = X$. Hence $\sum Z = \sum X = a + 1 \leq m$ and so $[c, d] \in N_n^{n-2}$.

We proved that $[c, d] \in N_n^{n-2}$ for all $d \in N_n^{n-2} \cap \mathcal{R}^*$. Now, since N_n^{n-2} is a saturated subgroup of Σ_n we have, by Theorem 3, that $c \in N_{\Sigma_n}(N_n^{n-2}) = N_n^{n-1}$. Moreover, $c \notin N_n^{n-2}$ since $\sum X = a + 1 > a$. \square

Using the GAP implementation of the algorithmic version of Theorem 5, available at GITHUB (<https://github.com/ngunivaq/normalizer-chain>), it is easy to compute the first normalizers in the chain of Eq.(2.2). The first values of $\log_2 |N_n^i : N_n^{i-1}|$ are shown in Table 2. Notice that for $i \leq n - 2$ the blue numbers correspond to those of the sequence (a_j) , whereas black bold numbers represent $\log_2 |N_n^{n-1} : N_n^{n-2}|$ and correspond to the number of unrefinable partitions as in Theorem 9. Moreover, notice that the diagonals in the blue area of Table 2 contain the same sequence of numbers. This is not the case when looking at the bold diagonal related to the $(n-1)$ -th normalizer. The reason for this is explained below.

Remark 3. Recall that, if $\sum X \leq a$, then the rigid commutator $c = \vee[a; X]$ belongs to N_n^{n-2} . In particular $c \in N_m^{m-2}$ for all $m \geq n$. A similar property does not hold for rigid commutators in N_n^{n-1} . In this case, we can show that if $c \in N_n^{n-1} \setminus N_n^{n-2}$, then $c \in N_m^{m-1}$ for finitely many m . Indeed, if $a \leq m$, by Theorem 9 we have that

n	$\log_2 N_n^i : N_n^{i-1} $ for $1 \leq i \leq 14$													
3	1	0	0	0	0	0	0	0	0	0	0	0	0	0
4	1	2	1	1	0	0	0	0	0	0	0	0	0	0
5	1	2	4	1	2	2	1	1	1	1	0	0	0	0
6	1	2	4	7	2	4	4	1	1	2	2	2	2	1
7	1	2	4	7	11	4	7	3	4	2	2	4	4	4
8	1	2	4	7	11	16	7	5	6	2	6	6	3	3
9	1	2	4	7	11	16	23	4	9	4	11	4	12	9
10	1	2	4	7	11	16	23	32	4	14	5	20	7	19
11	1	2	4	7	11	16	23	32	43	5	22	7	32	4
12	1	2	4	7	11	16	23	32	43	57	7	32	12	43
13	1	2	4	7	11	16	23	32	43	57	74	12	42	18
14	1	2	4	7	11	16	23	32	43	57	74	95	8	24
15	1	2	4	7	11	16	23	32	43	57	74	95	121	8

TABLE 2. Values of $\log_2|N_n^i : N_n^{i-1}|$ for small i and n

$c \in N_m^{m-1}$ only if m is in the interval $a \leq m < a + \text{mex}(X)$.

It is natural to ask when $k = \text{mex}(X)$ is the largest possible with respect to a . Since $\sum X = a + 1$, it is clear that this happens exactly when X is a triangular partition, i.e. $X = \{1, \dots, k-1\}$. We then have that

$$a = \sum X - 1 = k(k-1)/2 - 1 = (k+1)(k-2)/2$$

is the integer preceding the $(k-1)$ -th triangular number $k(k-1)/2$. Since $a = (k+1)(k-2)/2 > (k-2)^2/2$ we have that $\sqrt{2a} + 2$ is an upper bound for the largest possible value of k . In this case, if $a \geq 3$, then $c \in N_n^{n-1} \setminus N_n^{n-2}$ implies

$$n < a + k \leq a + \sqrt{2a} + 2 \leq (\sqrt{a} + 1)^2,$$

i.e. $a > n - 1 - 2\sqrt{n}$.

Example 10. When $n = 8$ we have computed that $\log_2|N_8^7 : N_8^6| = 7$. Indeed, applying Theorem 9 it can be shown that the normalizer N_8^7 can be generated by the generators of N_8^6 and by the following 7 rigid commutators of the type $\vee[a; X]$

$$\begin{aligned} \vee[5; \{3, 2, 1\}] &= [5, 4], \\ \vee[6; \{4, 2, 1\}] &= [6, 5, 3], \\ \vee[7; \{5, 2, 1\}] &= [7, 6, 4, 3], \\ \vee[7; \{4, 3, 1\}] &= [7, 6, 5, 2], \\ \vee[8; \{6, 2, 1\}] &= [8, 7, 5, 4, 3], \\ \vee[8; \{5, 3, 1\}] &= [8, 7, 6, 4, 2], \\ \vee[8; \{4, 3, 2\}] &= [8, 7, 6, 5, 1]. \end{aligned}$$

It is natural to wonder whether a general closed formula for $\log_2|N_n^{n-1} : N_n^{n-2}|$ may be found, even though at the time of writing this does not seem an easy task. To our knowledge, the problem of determining a closed formula or a generating function for the sequence c_j of the number of unrefinable partitions into distinct parts is open. The first values of c_j are shown here in Table 3, while a list of the first

j	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
c_j	1	1	1	1	1	2	1	1	2	3	1	2	2	3	5

TABLE 3. First values of the sequences (c_j)

1000 integers can be found in the On-Line Encyclopedia of Integer Sequences [OEI, <https://oeis.org/A179009>]. The related problem of determining a closed formula for the number of unrefinable partitions with a given minimal excludant, related to the number of partitions of Theorem 9, seems at the moment out of reach.

REFERENCES

- [ACGS19] R. Aragona, R. Civino, N. Gavioli, and C. M. Scoppola, *Regular subgroups with large intersection*, Ann. Mat. Pura Appl. (4) **198** (2019), no. 6, 2043–2057.
- [ACGS20] R. Aragona, R. Civino, N. Gavioli, and C. M. Scoppola, *A Chain of Normalizers in the Sylow 2-subgroups of the symmetric group on 2^n letters*, Preprint published electronically at <https://arxiv.org/abs/2008.13423>, 2020, To appear in *Indian J. Pure Appl. Math.*
- [ACGS21] R. Aragona, R. Civino, N. Gavioli, and C. M. Scoppola, *Rigid commutators and a normalizer chain*, Monatshefte für Mathematik (2021), 1–25.
- [AN19] G. E. Andrews and D. Newman, *Partitions and the minimal excludant*, Ann. Comb. **23** (2019), no. 2, 249–254.
- [AN20] ———, *The minimal excludant in integer partitions*, J. Integer Seq. **23** (2020), no. 2, Art. 20.2.3, 11.
- [And94] G. E. Andrews, *Number theory*, Dover Publications, Inc., New York, 1994, Corrected reprint of the 1971 original.
- [BM20] C. Ballantine and M. Merca, *The minimal excludant and colored partitions*, Sémin. Lothar. Combin. **84B** (2020), Art. 23, 12.
- [CBS19] R. Civino, C. Blondeau, and M. Sala, *Differential attacks: using alternative operations*, Des. Codes Cryptogr. **87** (2019), no. 2-3, 225–247.
- [CCS21] M. Calderini, R. Civino, and M. Sala, *On properties of translation groups in the affine general linear group with applications to cryptography*, J. Algebra **569** (2021), 658–680.
- [CDVS06] A. Caranti, F. Dalla Volta, and M. Sala, *Abelian regular subgroups of the affine group and radical rings*, Publ. Math. Debrecen **69** (2006), no. 3, 297–308.
- [Eul48] L. Euler, *Introductio in analysin infinitorum*, vol. 1, MM Bousquet, Lausanne, 1748.
- [FP15] A. S. Fraenkel and U. Peled, *Harnessing the unwieldy MEX function*, Games of No Chance **4** (2015), 77–94.
- [Gru39] P. M. Grundy, *Mathematics and Games*, Eureka **2** (1939), 6–9.
- [OEI] *The On-Line Encyclopedia of Integer Sequences*, Published electronically at <https://oeis.org>, Accessed: 2021-07-07.
- [Spr35] R. Sprague, *Über mathematische kampfspiele*, Tohoku Mathematical Journal, First Series **41** (1935), 438–444.

(R. Aragona, R. Civino, N. Gavioli) DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL’INFORMAZIONE E MATEMATICA, UNIVERSITÀ DEGLI STUDI DELL’AQUILA, VIA VETOIO, I-67100 COPPITO (AQ), ITALY

Email address, Riccardo Aragona: riccardo.aragona@univaq.it

Email address, Roberto Civino: roberto.civino@univaq.it

Email address, Norberto Gavioli: norberto.gavioli@univaq.it

Email address, Carlo Maria Scoppola: carlo.scoppola@univaq.it