

ON FANO THREEFOLDS OF DEGREE 22 AFTER CHELTSOV AND SHRAMOV

KENTO FUJITA

ABSTRACT. It has been known that nonsingular Fano threefolds of Picard rank one with the anti-canonical degree 22 admitting faithful actions of the multiplicative group form a one-dimensional family. Cheltsov and Shramov showed that all but two of them admit Kähler-Einstein metrics. In this paper, we show that the remaining Fano threefolds also admit Kähler-Einstein metrics.

CONTENTS

1. Introduction	1
2. Log Fano pairs	3
3. On the volume functions	6
4. On quasi-log schemes	10
5. Proof of Theorem 1.2	12
References	13

1. INTRODUCTION

Let X be a nonsingular Fano threefold over the complex number field with $\text{Pic}(X) = \mathbb{Z}[-K_X]$. Such X with large automorphism groups have been studied by many authors. For example, Mukai and Umemura systematically studied in [MU83] the Fano threefold V^{MU} with the anti-canonical degree 22, so-called *the Mukai-Umemura threefold*, which is obtained by the unique $\text{SL}(2, \mathbb{C})$ -equivariant nonsingular projective compactification of $\text{SL}(2, \mathbb{C})/I_h$ with the Picard rank one, where $I_h \subset \text{SL}(2, \mathbb{C})$ is the icosahedral group. The automorphism group of V^{MU} is equal to $\text{PGL}(2, \mathbb{C})$. On the other hand, Prokhorov showed in [Pro90] that, if $\text{Aut}(X)$ is not finite, then the anti-canonical degree of X must be equal to 22. Moreover, he determined all of such X . For example,

Date: October 13, 2021.

2010 Mathematics Subject Classification. Primary 14J45; Secondary 14L24.

Key words and phrases. Fano varieties, K-stability, Kähler-Einstein metrics.

there is a unique Fano threefold V^a such that $\text{Aut}^0(V^a)$ is equal to the additive group \mathbb{C}^+ .

Nowadays, the structures of such X are well-understood thanks to the works of Kuznetsov, Prokhorov, Shramov [KPS18], and Kuznetsov, Prokhorov [KP18]. The family of nonsingular Fano threefolds X with $\text{Pic}(X) = \mathbb{Z}[-K_X]$, $(-K_X)^3 = 22$, $X \not\cong V^a$ and $\text{Aut}(X)$ infinite are parametrized by $\mathbb{C} \setminus \{0, 1\}$. Let us denote the family by $\{V_u\}_{u \in \mathbb{C} \setminus \{0, 1\}}$. Then $V_{-1/4}$ is isomorphic to V^{MU} , and the automorphism group of V_u is equal to $G := \mathbb{C}^* \rtimes (\mathbb{Z}/2\mathbb{Z})$ unless $u = -1/4$. Moreover, any V_u can be obtained by the two-ray game from the blowup of some 3-dimensional quadric hypersurface along a certain nonsingular sextic rational curve.

In [CS18], Cheltsov and Shramov considered the problem for the existence of Kähler-Einstein metrics for the above V_u . If $u = -1/4$, then the Fano threefold is the Mukai-Umemura threefold V^{MU} . In this case, Donaldson already showed in [Don07] the existence of Kähler-Einstein metrics on V^{MU} by showing that the $\text{PGL}(2, \mathbb{C})$ -invariant α -invariant $\alpha_{\text{PGL}(2, \mathbb{C})}(V^{\text{MU}})$ (see §2) of V^{MU} is equal to $5/6$. In fact, Tian showed in [Tia87] that, for a Fano manifold X and a reductive subgroup $\Gamma \subset \text{Aut}(X)$, if $\alpha_\Gamma(X) > \dim X / (\dim X + 1)$ holds, then X admits Kähler-Einstein metrics. Cheltsov and Shramov considered the remaining cases by evaluating the G -invariant α -invariant of V_u . More precisely, they showed the following:

Theorem 1.1 ([CS18, Theorem 1.5]). *We have*

$$\alpha_G(V_u) = \begin{cases} 4/5 & \text{if } u \neq 3/4 \text{ and } u \neq 2, \\ 3/4 & \text{if } u = 3/4, \\ 2/3 & \text{if } u = 2. \end{cases}$$

Thus, together with Tian's result, if $u \neq 3/4$ and $u \neq 2$, then V_u admits Kähler-Einstein metrics. However, if $u = 3/4$ or $u = 2$, then the existence of Kähler-Einstein metrics of V_u was not known at that time. The purpose of this paper is to show the existence of Kähler-Einstein metrics for $V_{3/4}$ and V_2 . The main result is the following:

Theorem 1.2. *Both $V_{3/4}$ and V_2 admit Kähler-Einstein metrics.*

Thus, together with Matsushima's obstruction [Mat57], we gave a complete answer for the existence of Kähler-Einstein metrics of nonsingular Fano threefolds X with $\text{Pic}(X) = \mathbb{Z}[-K_X]$ and $\text{Aut}(X)$ infinite; such X admits Kähler-Einstein metrics if and only if $X \cong V^a$.

The main technique to prove Theorem 1.2 is the “ G -valuative criterion” (see §2, [ACCFKMGSV] and [Zhu20]), which is a G -equivariant version of [Lil17, Theorem 3.7] and [Fjt19a, Theorem 1.6]. Moreover,

we need deep analyses [CS18] of G -invariant curves on V_u which might be possible destabilizing centers of G -invariant prime divisors over V_u . Technically, the theory of quasi-log schemes [Fjn17] and the careful analysis of the volume functions (see §3) play important roles in order to show Theorem 1.2. Especially, we use a subadjunction-type result for projective qlc strata on quasi-log schemes [Fjn21] (see Theorem 4.1). We expect that these techniques will be applied for many other Fano varieties (cf. [ACCFKMGSSV]).

Acknowledgments. The author thank Carolina Araujo, Ana-Maria Castravet, Ivan Cheltsov, Osamu Fujino, Anne-Sophie Kaloghiros, Akihiro Kanemitsu, Jesus Martinez-Garcia, Constantin Shramov, Hendrick Süß and Nivedita Viswanathan for many discussions. This work started while the author participated the AIM workshop named “K-stability and related topics” on January 2020. The author thanks the staffs for the stimulating environment. This work was supported by JSPS KAKENHI Grant Number 18K13388.

2. LOG FANO PAIRS

In this section, we recall the definition of α -invariant for log Fano pairs and see the relationship between α -invariant and the existence of Kähler-Einstein metrics. For the minimal model program, we refer the readers to [KM98].

Definition 2.1. A *log Fano pair* (X, Δ) is a pair of a complex normal projective variety X and an effective \mathbb{Q} -Weil divisor Δ on X such that the pair (X, Δ) is klt and $-(K_X + \Delta)$ is an ample \mathbb{Q} -divisor. If X is nonsingular and $\Delta = 0$, then X is said to be a *Fano manifold*.

Definition 2.2 (α -invariant (see [Dem08] for example)). Let (X, Δ) be a log Fano pair and let $\Gamma \subset \text{Aut}(X, \Delta)$ be an algebraic subgroup.

- (1) The Γ -invariant α -invariant $\alpha_\Gamma(X, \Delta)$ of (X, Δ) is defined as the supremum of $\alpha \in \mathbb{Q}_{>0}$ such that the pair $(X, \Delta + \frac{\alpha}{m}\mathcal{D})$ is lc for any $m \in \mathbb{Z}_{>0}$ with $-m(K_X + \Delta)$ Cartier and for any nonempty Γ -invariant sub-linear system $\mathcal{D} \subset |-m(K_X + \Delta)|$.
- (2) For any scheme-theoretic point $\eta \in X$, we define $\alpha_{\Gamma, \eta}(X, \Delta)$ as the supremum of $\alpha \in \mathbb{Q}_{>0}$ such that the pair $(X, \Delta + \frac{\alpha}{m}\mathcal{D})$ is lc at η for any $m \in \mathbb{Z}_{>0}$ with $-m(K_X + \Delta)$ Cartier and for any nonempty Γ -invariant sub-linear system $\mathcal{D} \subset |-m(K_X + \Delta)|$.
- (3) If $\Delta = 0$, then we simply write $\alpha_\Gamma(X)$ and $\alpha_{\Gamma, \eta}(X)$ in place of $\alpha_\Gamma(X, 0)$ and $\alpha_{\Gamma, \eta}(X, 0)$, respectively.

Lemma 2.3. *Let (X, Δ) be a log Fano pair, $\Gamma \subset \text{Aut}(X, \Delta)$ be an algebraic subgroup, and $\eta \in X$ be a scheme-theoretic point. Assume that the identity component Γ_0 of Γ is solvable.*

- (1) *The value $\alpha_\Gamma(X, \Delta)$ is equal to the supremum of $\alpha \in \mathbb{Q}_{>0}$ such that the pair $(X, \Delta + \alpha D)$ is lc for any effective Γ -invariant \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -(K_X + \Delta)$.*
- (2) *The value $\alpha_{\Gamma, \eta}(X, \Delta)$ is equal to the supremum of $\alpha \in \mathbb{Q}_{>0}$ such that the pair $(X, \Delta + \alpha D)$ is lc at η for any effective Γ -invariant \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -(K_X + \Delta)$.*

Proof. We only prove (2). By [BS64, Lemme 5.11], there is a finite algebraic subgroup $\Gamma_1 \subset \Gamma$ such that Γ_1 meets every connected component of Γ . Set $d := \#\Gamma_1$. Fix $\alpha \in \mathbb{Q}_{>0}$.

Assume that $(X, \Delta + \frac{\alpha}{m}\mathcal{D})$ is lc at η for any m and $\mathcal{D} \subset |-m(K_X + \Delta)|$. Then, for any effective Γ -invariant \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -(K_X + \Delta)$, since $\{mD\} \subset |-m(K_X + \Delta)|$ is a Γ -invariant sub-linear system for m sufficiently divisible, we know that the pair $(X, \Delta + \frac{\alpha}{m}\{mD\})$ is lc at η , i.e., the pair $(X, \Delta + \alpha D)$ is lc at η .

Conversely, assume that $(X, \Delta + \alpha D)$ is lc at η for any effective Γ -invariant \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -(K_X + \Delta)$. Take any Γ -invariant sub-linear system $\mathcal{D} \subset |-m(K_X + \Delta)|$. Since Γ_0 is connected and solvable, there exists a Γ_0 -invariant divisor $D_0 \in \mathcal{D}$ by the Borel fixed point theorem [Hum75, §21.2]. Let us set

$$\tilde{D} := \sum_{h \in \Gamma_1} h(D_0) \in d\mathcal{D}.$$

Since $h(D_0) \in \mathcal{D}$ is Γ_0 -invariant for any $h \in \Gamma_1$, the divisor $\tilde{D} \in d\mathcal{D}$ is Γ -invariant. Set $D := \frac{1}{md}\tilde{D} \sim_{\mathbb{Q}} -(K_X + \Delta)$. Since $(X, \Delta + \alpha D)$ is lc at η , the pair $(X, \Delta + \frac{\alpha}{md}(d\mathcal{D}))$ is also lc at η . This is equivalent to the pair $(X, \Delta + \frac{\alpha}{m}\mathcal{D})$ being lc at η . \square

We recall the words on *K-stability* of log Fano pairs. The original notion of K-stability was introduced by Tian [Tia97] and Donaldson [Don02] by using the languages of *test configurations*. In this paper, we only treat its simplification due to Li [Li17] and the author [Fjt19a].

Definition 2.4. Let (X, Δ) be an n -dimensional log Fano pair and let F be a prime divisor over X obtained by a log resolution $\pi: \tilde{X} \rightarrow X$ of (X, Δ) (that is, F is a prime divisor on \tilde{X}).

- (1) Let $A_{X, \Delta}(F)$ be the log discrepancy of (X, Δ) along F , that is, 1 plus the coefficient of $K_{\tilde{X}} - \pi^*(K_X + \Delta)$ along F .
- (2) For any effective \mathbb{Q} -Cartier \mathbb{Q} -divisor D , let $\text{ord}_F D \in \mathbb{Q}_{\geq 0}$ be the coefficient of π^*D along F . For any $m \in \mathbb{Z}_{\geq 0}$ with $-m(K_X +$

Δ) Cartier and for any $j \in \mathbb{R}_{\geq 0}$, let

$$H^0(X, -m(K_X + \Delta) - jF) \subset H^0(X, -m(K_X + \Delta))$$

be the sub-vector space corresponds to the sub-linear system $| -m(K_X + \Delta) - jF| \subset | -m(K_X + \Delta)|$ consisting all $D \in | -m(K_X + \Delta)|$ with $\text{ord}_F D \geq j$.

(3) For any $x \in \mathbb{R}_{\geq 0}$, let us set

$$\text{vol}(-(K_X + \Delta) - xF) := \lim_{m \rightarrow \infty} \frac{\dim H^0(X, -m(K_X + \Delta) - mxF)}{m^n/n!},$$

where m runs through all positive integers with $-m(K_X + \Delta)$ Cartier (the limit exists by [Laz04a, Laz04b]). Obviously, $\text{vol}(-(K_X + \Delta) - 0 \cdot F) = (-m(K_X + \Delta))^n$ holds. It follows from the definition that

$$\text{vol}(-(K_X + \Delta) - xF) = \text{vol}_{\tilde{X}}(-\pi^*(K_X + \Delta) - xF).$$

In particular, by [Laz04a, Laz04b], the function $\text{vol}(-(K_X + \Delta) - xF)$ is a non-increasing and continuous function over $x \in [0, \infty)$. Moreover, if $x \gg 0$, then $\text{vol}(-(K_X + \Delta) - xF) = 0$ holds. Let us set

$$\tau_{X,\Delta}(F) := \sup\{\tau \in \mathbb{R}_{>0} \mid \text{vol}(-(K_X + \Delta) - \tau F) > 0\}.$$

(4) Let us set

$$S_{X,\Delta}(F) := \frac{1}{(-m(K_X + \Delta))^n} \int_0^\infty \text{vol}(-(K_X + \Delta) - xF) dx.$$

The above definitions do not depend on the choice of π . We often write $A(F)$, $\tau(F)$, $S(F)$ in place of $A_{X,\Delta}(F)$, $\tau_{X,\Delta}(F)$, $S_{X,\Delta}(F)$, just for simplicity.

The following lemma is well-known:

Lemma 2.5. *Let (X, Δ) be a log Fano pair, let $\Gamma \subset \text{Aut}(X, \Delta)$ be an algebraic subgroup, and let $\eta \in X$ be a scheme-theoretic point. For any Γ -invariant prime divisor F over X with $\eta \in c_X(F)$, we have*

$$\frac{A(F)}{\tau(F)} \geq \alpha_{\Gamma,\eta}(X, \Delta),$$

where $c_X(F)$ is the center of F on X .

Proof. Take any $\tau = \frac{j}{m} \in (0, \tau(F)) \cap \mathbb{Q}$. Then the sub-linear system

$$| -m(K_X + \Delta) - jF| \subset | -m(K_X + \Delta)|$$

is nonempty, Γ -invariant, and vanishes along F at least j times. Thus we have

$$\begin{aligned} \alpha_{\Gamma, \eta}(X, \Delta) &\leq \text{lct}_{\eta} \left(X, \Delta; \frac{1}{m} | -m(K_X + \Delta) - jF | \right) \\ &\leq \frac{A_{X, \Delta}(F)}{\text{ord}_F \left(\frac{1}{m} | -m(K_X + \Delta) - jF | \right)} \leq \frac{A_{X, \Delta}(F)}{\tau}, \end{aligned}$$

where lct_{η} is the log canonical threshold at η . \square

We see a G -invariant version of [Li17, Fjt19a] and [Fjt19c]. See also [Zhu20] for more general frameworks.

Proposition 2.6 (see [ACCFKMGSSV]). *Let X be an n -dimensional Fano manifold and let $\Gamma \subset \text{Aut}(X)$ be a reductive subgroup.*

- (1) (see also [Zhu20, Corollary 4.13]) *Assume that $A(F) > S(F)$ holds for any Γ -invariant prime divisor over X with the \mathbb{C} -algebra*

$$\bigoplus_{m, j \in \mathbb{Z}_{\geq 0}} H^0(X, -mK_X - jF)$$

finitely generated. Then X admits Kähler-Einstein metrics.

- (2) (see also [Tia87] and [Fjt19c, Theorem 1.3]) *If $\alpha_{\Gamma}(X) \geq \frac{n}{n+1}$, then X admits Kähler-Einstein metrics.*

For the complete proof, see [ACCFKMGSSV]. We only give a sketch of the idea. For (1), for any Γ -equivariant special degeneration of X in the sense of [DS16], the corresponding prime divisor F over X in [Fjt19a, Theorem 5.1] obviously satisfies the assumptions in (1). By [Fjt19a, Theorem 5.1], the signatures of the Donaldson-Futaki invariant of the special degeneration and $A(F) - S(F)$ are same. Thus we can apply [DS16, Theorem 1]. For (2), we may assume that there exists a Γ -invariant prime divisor F over X with $A(F) \leq S(F)$ such that the \mathbb{C} -algebra in (1) is finitely generated. By Lemma 2.5 and [Fjt19c, Theorem 4.1], X is isomorphic to \mathbb{P}^n and then we complete the proof.

3. ON THE VOLUME FUNCTIONS

In this section, we generalize [Fjt19b, Proposition 2.1] in order to show that V_2 admits Kähler-Einstein metrics.

Proposition 3.1. *Let (X, Δ) be an n -dimensional log Fano pair, let F be a prime divisor over X , and let $0 < a < b$ be positive real numbers. Assume that*

$$\text{vol}(-(K_X + \Delta) - xF) = \left(\frac{b-x}{b-a} \right)^n \text{vol}(-(K_X + \Delta) - aF)$$

for any $x \in [a, b]$. Then we have

$$S(F) \leq \frac{(n-1)a + b}{n+1}.$$

Proof. The proof looks similar to the argument in the proof of [SZ19, Theorem 1.2]. From the assumption, we have $\tau(F) = b$. Set $V := (-K_X + \Delta)^n$. By [BFJ09, Theorem A], the function $\text{vol}(-(K_X + \Delta) - xF)$ is \mathcal{C}^1 over $x \in [0, b]$. Let us set

$$f(x) := -\frac{1}{n} \frac{d}{dx} \text{vol}(-(K_X + \Delta) - xF)$$

as in [Fjt19b, Proof of Proposition 2.1]. (We note that $f(x)$ is a restricted volume function in the sense of [ELMNP09].) Then, for any $x \in [a, b]$, we have

$$f(x) = \left(\frac{b-x}{b-a}\right)^{n-1} f(a).$$

As in [Fjt19b, Proof of Proposition 2.1], we have

$$\begin{aligned} V &= n \int_0^b f(x) dx, \\ S(F) &= \frac{1}{V} \cdot n \int_0^b x f(x) dx, \\ f(x) &\geq \left(\frac{x}{a}\right)^{n-1} f(a) \text{ for any } x \in [0, a]. \end{aligned}$$

In particular, we get

$$\begin{aligned} V &\geq n \left(\int_0^a \left(\frac{x}{a}\right)^{n-1} f(a) dx + \int_a^b \left(\frac{b-x}{b-a}\right)^{n-1} f(a) dx \right) \\ &= b \cdot f(a). \end{aligned}$$

Set

$$g(x) := \begin{cases} \left(\frac{x}{a}\right)^{n-1} \cdot \frac{V}{b} & \text{for } x \in [0, a], \\ \left(\frac{b-x}{b-a}\right)^{n-1} \cdot \frac{V}{b} & \text{for } x \in [a, b]. \end{cases}$$

Then we have

$$\begin{aligned} V &= n \int_0^b g(x) dx, \\ g(x) &\geq f(x) \text{ for any } x \in [a, b]. \end{aligned}$$

Moreover, the function

$$f(x)^{\frac{1}{n-1}} - g(x)^{\frac{1}{n-1}} = f(x)^{\frac{1}{n-1}} - \frac{x}{a} \cdot \left(\frac{V}{b}\right)^{\frac{1}{n-1}}$$

is \mathcal{C}^0 and concave over $x \in [0, a]$ (by [ELMNP09, Theorem A]). Moreover, we have $f(0)^{\frac{1}{n-1}} - g(0)^{\frac{1}{n-1}} \geq 0$ and $f(a)^{\frac{1}{n-1}} - g(a)^{\frac{1}{n-1}} \leq 0$. Thus there exists $c \in [0, a]$ such that

$$\begin{aligned} f(x) &\geq g(x) \quad \text{for any } x \in [0, c], \\ g(x) &\geq f(x) \quad \text{for any } x \in [c, a]. \end{aligned}$$

Thus we get

$$\begin{aligned} n \int_0^b xf(x)dx - cV &= n \int_0^b (x-c)f(x)dx \\ &\leq n \int_0^b (x-c)g(x)dx = n \int_0^b xg(x)dx - cV. \end{aligned}$$

Since

$$n \int_0^b xg(x)dx = \frac{(n-1)a+b}{n+1}V,$$

we get the assertion. \square

Proposition 3.2. *Let (X, Δ) be an n -dimensional log Fano pair, let $\eta \in X$ be a scheme-theoretic point, and let $0 < t \leq s$ be positive real numbers. Assume that there exists a prime divisor T on X with $T \sim_{\mathbb{Q}} -k(K_X + \Delta)$ for some $k \in \mathbb{Q}_{>0}$ such that*

- *the pair $(X, \Delta + \frac{1}{k}T)$ is lc at η , and*
- *for any effective \mathbb{Q} -divisor $D' \sim_{\mathbb{Q}} -(K_X + \Delta)$ with $T \not\subset \text{Supp } D'$, the pair $(X, \Delta + sD')$ is lc at η .*

Then, for any prime divisor F over X with $\eta \in c_X(F)$, we have the following:

- (1) *If $s^{-1}A(F) \leq \frac{1}{k} \text{ord}_F T$, then we have*

$$\begin{aligned} &\text{vol}(-(K_X + \Delta) - xF) \\ &= \left(\frac{\frac{1}{k} \text{ord}_F T - x}{\frac{1}{k} \text{ord}_F T - s^{-1}A(F)} \right)^n \text{vol}(-(K_X + \Delta) - s^{-1}A(F)F) \end{aligned}$$

for any $x \in [s^{-1}A(F), \frac{1}{k} \text{ord}_F T]$.

- (2) *We have*

$$S(F) \leq \frac{A(F)}{n+1} ((n-1)s^{-1} + t^{-1}).$$

Proof. (1) Let us fix a log resolution $\pi: \tilde{X} \rightarrow X$ of (X, Δ) with $F \subset \tilde{X}$. Take any effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -(K_X + \Delta)$. Then there uniquely exists $e \in [0, 1] \cap \mathbb{Q}$ such that we can write $D = \frac{e}{k}T + (1-e)D'$ with $D' \sim_{\mathbb{Q}} -(K_X + \Delta)$ effective and $T \not\subset \text{Supp } D'$. Since $(X, \Delta + sD')$ is lc at η , we have $\text{ord}_F D' \leq s^{-1}A(F)$. Assume that $x \in (s^{-1}A(F), \frac{1}{k} \text{ord}_F T] \cap$

\mathbb{Q} satisfies that $\text{ord}_F D \geq x$. (In other words, $mD \in |-m(K_X + \Delta) - mxF|$ holds for a sufficiently divisible $m \in \mathbb{Z}_{>0}$.) Then we have

$$e \geq \frac{x - s^{-1}A(F)}{\frac{1}{k} \text{ord}_F T - s^{-1}A(F)},$$

since

$$x \leq \text{ord}_F D \leq e \cdot \frac{1}{k} \text{ord}_F T + (1 - e)s^{-1}A(F).$$

This implies that the linear system $|-m(K_X + \Delta) - mxF|$ has a fixed divisor

$$m \cdot \frac{x - s^{-1}A(F)}{\frac{1}{k} \text{ord}_F T - s^{-1}A(F)} \cdot \frac{1}{k} T$$

for $m \in \mathbb{Z}_{>0}$ sufficiently divisible. Thus we get

$$\begin{aligned} & \text{vol}(-(K_X + \Delta) - xF) \\ &= \text{vol}_{\bar{X}} \left(-\pi^*(K_X + \Delta) - xF - \frac{x - s^{-1}A(F)}{\frac{1}{k} \text{ord}_F T - s^{-1}A(F)} \frac{1}{k} (\pi^*T - (\text{ord}_F T)F) \right) \\ &= \left(\frac{\frac{1}{k} \text{ord}_F T - x}{\frac{1}{k} \text{ord}_F T - s^{-1}A(F)} \right)^n \text{vol}(-(K_X + \Delta) - s^{-1}A(F)F). \end{aligned}$$

(2) Since the pair $(X, \Delta + \frac{t}{k}T)$ is lc at η , we have $\frac{1}{k} \text{ord}_F T \leq t^{-1}A(F)$. If $\frac{1}{k} \text{ord}_F T \leq s^{-1}A(F)$, then we have $\text{ord}_F D \leq s^{-1}A(F)$ for any effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -(K_X + \Delta)$. Thus we get the inequality $s \leq \frac{A(F)}{\tau(F)}$. By [Fjt19b, Proposition 2.1], we get

$$\begin{aligned} S(F) &\leq \frac{n}{n+1} \tau(F) \leq \frac{n}{n+1} s^{-1}A(F) \\ &\leq \frac{A(F)}{n+1} ((n-1)s^{-1} + t^{-1}). \end{aligned}$$

Thus we may assume that $\frac{1}{k} \text{ord}_F T > s^{-1}A(F)$. In this case, we can apply (1). We have

$$\begin{aligned} S(F) &\leq \frac{1}{n+1} \left((n-1)s^{-1}A(F) + \frac{1}{k} \text{ord}_F T \right) \\ &\leq \frac{A(F)}{n+1} ((n-1)s^{-1} + t^{-1}) \end{aligned}$$

by Proposition 3.1. □

4. ON QUASI-LOG SCHEMES

The theory of *quasi-log schemes* plays an important role for the study of birational geometry. In this section, we see a kind of subadjunction theorem for projective qlc strata on quasi-log schemes, which is a direct consequence of the recent work [Fjn21]. For the theory of quasi-log schemes, we refer the readers to [Fjn17].

An \mathbb{R} -Weil divisor D on a normal projective variety is said to be *pseudo-effective* in this paper if $D+A$ is big (i.e., there exists an effective \mathbb{R} -Weil divisor E such that $D+A-E$ is an ample \mathbb{R} -divisor) for any ample \mathbb{R} -divisor A .

Theorem 4.1 (see [Fjn21, Lemma 4.17, Theorems 1.9 and 7.1]). *Let $[X, \omega]$ be a quasi-log scheme, let $C \subset X$ be a qlc stratum of $[X, \omega]$, and let $\nu: \bar{C} \rightarrow C$ be the normalization of C . Assume that C is a projective variety. Then $\nu^*(\omega|_C) - K_{\bar{C}}$ is a pseudo-effective \mathbb{R} -Weil divisor on \bar{C} .*

Proof. By [Fjn21, Lemma 4.19], we may assume that $X = C$. Moreover, by [Fjn21, Theorem 1.9], we may further assume that $X = C$ is normal. By [Fjn21, Theorem 7.1], there exists a projective birational morphism $p: X' \rightarrow X$ from a smooth projective variety X' such that we can write

$$K_{X'} + B_{X'} + M_{X'} = p^*\omega,$$

where $B_{X'}$ is an effective \mathbb{R} -Weil divisor on X' with $B_{X'}^{\leq 0}$ p -exceptional, and $M_{X'}$ is a nef \mathbb{R} -divisor on X' . This immediately implies that

$$\omega - K_X = p_*(B_{X'} + M_{X'})$$

is a pseudo-effective \mathbb{R} -Weil divisor on X . □

As a corollary, we get the following result, which is important for the proof of Theorem 1.2.

Corollary 4.2. *Let (X, Δ) be a log Fano pair, let $D \sim_{\mathbb{Q}} -(K_X + \Delta)$ be an effective \mathbb{Q} -divisor, and let $\alpha \in (0, 1) \cap \mathbb{Q}$. Assume that the pair $(X, \Delta + \alpha D)$ is not klt. Let $\text{Nklt}(X, \Delta + \alpha D)$ be the locus of non-klt points of $(X, \Delta + \alpha D)$.*

- (1) *The locus $\text{Nklt}(X, \Delta + \alpha D)$ is connected.*
- (2) *Take any 1-dimensional irreducible component $B \subset \text{Nklt}(X, \Delta + \alpha D)$ with its reduced scheme structure. Then B is a rational curve with*

$$(-(K_X + \Delta) \cdot B) \leq \frac{2}{1 - \alpha}.$$

Moreover, if any irreducible component of $\text{Nklt}(X, \Delta + \alpha D)$ is of dimension ≤ 1 , then $B \simeq \mathbb{P}^1$ and the restriction homomorphism

$$H^0(X, L) \rightarrow H^0(B, L|_B)$$

is surjective for any nef line bundle L on X .

Proof. By [Fjn17, 6.4.1], the pair $(X, \Delta + \alpha D)$ admits a quasi-log structure $[X, \omega]$ with $\omega = K_X + \Delta + \alpha D$, and $N := \text{Nklt}(X, \Delta + \alpha D)$ has a natural scheme structure with

$$N = \bigcup_C C \cup X_{-\infty},$$

where C are the lc centers of $(X, \Delta + \alpha D)$ and $X_{-\infty}$ is the non-qlc locus of $(X, \Delta + \alpha D)$.

For any nef line bundle L on X , since

$$L - (K_X + \Delta + \alpha D) \sim_{\mathbb{Q}} L + (1 - \alpha)(-K_X - \Delta)$$

is ample, we have

$$H^i(X, L \otimes I_N) = 0$$

for any $i > 0$ by [Fjn17, Theorem 6.3.5 (ii)], where $I_N \subset \mathcal{O}_X$ is the defining ideal sheaf of $N \subset X$.

(1) For $L := \mathcal{O}_X$, we get the surjection

$$H^0(X, \mathcal{O}_X) \rightarrow H^0(N, \mathcal{O}_N).$$

Thus N is connected.

(2) After replacing α with the log canonical threshold of $(X, \Delta; D)$ at the generic point of B , we may assume that B is an lc center of $(X, \Delta + \alpha D)$. Take the normalization $\nu: \bar{B} \rightarrow B$. By Theorem 4.1, the \mathbb{Q} -divisor

$$\nu^*((K_X + \Delta + \alpha D)|_B) - K_{\bar{B}}$$

is pseudo-effective. This implies that $\bar{B} \simeq \mathbb{P}^1$ and

$$\begin{aligned} 0 &\leq \deg_{\bar{B}}(\nu^*((K_X + \Delta + \alpha D)|_B) - K_{\bar{B}}) \\ &= -(1 - \alpha)(-(K_X + \Delta) \cdot B) + 2. \end{aligned}$$

Now we assume that $\dim N \leq 1$. Since $H^1(X, \mathcal{O}_X) = H^2(X, I_N) = 0$, we get $H^1(N, \mathcal{O}_N) = 0$. From the assumption $\dim N \leq 1$, we get $H^1(B, \mathcal{O}_B) = 0$, i.e., $B \simeq \mathbb{P}^1$. Moreover, by [KM98, Lemma 4.13 and the proof of Lemma 4.50], the restriction homomorphism

$$H^0(N, L|_N) \rightarrow H^0(B, L|_B)$$

is surjective for any nef line bundle L on X . Since $H^1(X, L \otimes I_N) = 0$,

$$H^0(X, L) \rightarrow H^0(B, L|_B)$$

is also surjective. Thus we get the assertions. \square

5. PROOF OF THEOREM 1.2

By Theorem 1.1 and Proposition 2.6 (2), we may assume that $u = 2$. Set $X := V_2$. Take any G -invariant prime divisor F over X . (Recall that $G := \mathbb{C}^* \rtimes (\mathbb{Z}/2\mathbb{Z})$.) By Proposition 2.6 (1), it is enough to show the inequality $A(F) > S(F)$. Set $C := c_X(F)$ and let $\eta \in X$ be the generic point of C . By Lemma 2.5 and [Fjt19b, Proposition 2.1], we may assume that $\alpha_{G,\eta}(X) \leq \frac{3}{4}$. Note that C is a G -invariant subvariety on X . Thus C is not a closed point by [CS18, Lemma 2.23]. Moreover, if F is a prime divisor on X , then $A(F) > S(F)$ holds by [Fjt16, Corollary 9.3]. Thus we may assume that C is a G -invariant curve. By Lemma 2.3, there exists a G -invariant effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -K_X$ and there exists $\alpha \in [\alpha_{G,\eta}(X), 4/5) \cap \mathbb{Q}$ such that the pair $(X, \alpha D)$ is lc but not klt at η . We note that the non-klt locus of the pair $(X, \alpha D)$ is of dimension ≤ 1 since $\text{Pic}(X) = \mathbb{Z}[-K_X]$ and $\alpha < 1$.

By Corollary 4.2, C is a smooth rational curve with

$$(-K_X \cdot C) \leq \frac{2}{1-\alpha} < 10.$$

Moreover, for any $m \in \mathbb{Z}_{\geq 0}$, the restriction homomorphism

$$H^0(X, \mathcal{O}_X(-mK_X)) \rightarrow H^0(C, \mathcal{O}_X(-mK_X)|_C)$$

is surjective. Thus $C \subset X \subset \mathbb{P}^{13} = \mathbb{P}^*H^0(X, -K_X)$ is a G -invariant rational normal curve in \mathbb{P}^{13} with $\deg C < 10$.

By [CS18, Proposition 4.12, Lemma 7.7 and Corollary 7.10] and the assumption $\alpha_{G,\eta}(X) \leq \frac{3}{4}$, we may assume that $C = \mathcal{C}_4$, where $\mathcal{C}_4 \subset X$ is the unique G -invariant rational normal curve of anti-canonical degree 4 in X (see [CS18] for the definition of the curve \mathcal{C}_4). By [CS18, Lemma 5.2 and the proof of Lemma 7.14], there exists a prime divisor $T'_{15} \sim -K_X$ on X such that

- the pair $(X, \frac{2}{3}T'_{15})$ is lc at η , and
- the pair (X, D') is lc at η for any effective \mathbb{Q} -divisor $D' \sim_{\mathbb{Q}} -K_X$ with $T'_{15} \not\subset \text{Supp } D'$.

By Proposition 3.2 (2), we get

$$S(F) \leq \frac{A(F)}{4} \left(2 \cdot 1^{-1} + \left(\frac{2}{3} \right)^{-1} \right) = \frac{7}{8}A(F).$$

Thus we get the desired inequality $A(F) > S(F)$ and we complete the proof of Theorem 1.2.

REFERENCES

- [ACCFKMGSSV] C. Araujo, A-M. Castravet, I. Cheltsov, K. Fujita, A-S. Kaloghiros, J. Martinez-Garcia, C. Shramov, H. Süß and N. Viswanathan, *The Calabi problem for Fano threefolds*, MPIM preprint 2021-31.
- [BFJ09] S. Boucksom, C. Favre and M. Jonsson, *Differentiability of volumes of divisors and a problem of Teissier*, J. Algebraic Geom. **18** (2009), no. 2, 279–308.
- [BS64] A. Borel and J.-P. Serre, *Théorèmes de finitude en cohomologie galoisienne*, Comment. Math. Helv. **39** (1964), 111-164.
- [CS18] I. Cheltsov and C. Shramov, *Kähler-Einstein Fano threefolds of degree 22*, arXiv:1803.02774v4.
- [Dem08] J.-P. Demailly, Appendix to I. Cheltsov and C. Shramov’s article. “Log canonical thresholds of smooth Fano threefolds” : *On Tian’s invariant and log canonical thresholds*, Russian Math. Surveys **63** (2008), no. 5, 945–950.
- [Don02] S. Donaldson, *Scalar curvature and stability of toric varieties*, J. Differential Geom. **62** (2002), no. 2, 289–349.
- [Don07] S. Donaldson, *A note on the α -invariant of the Mukai-Umemura 3-fold*, arXiv:0711.4357v1.
- [DS16] V. Datar and G. Székelyhidi, *Kähler-Einstein metrics along the smooth continuity method*, Geom. Funct. Anal. **26**, no. 4, 975–1010.
- [ELMNP09] L. Ein, R. Lazarsfeld, M. Mustață, M. Nakamaye and M. Popa, *Restricted volumes and base loci of linear series*, Amer. J. Math. **131** (2009), no. 3, 607–651.
- [Fjn17] O. Fujino, *Foundations of the minimal model program*, MSJ Memoirs, **35**. Mathematical Society of Japan, Tokyo, 2017.
- [Fjn21] O. Fujino, *Cone theorem and Mori hyperbolicity*, arXiv::2102.11986v1.
- [Fjt16] K. Fujita, *On K -stability and the volume functions of \mathbb{Q} -Fano varieties*, Proc. Lond. Math. Soc. **113** (2016), no. 5, 541–582.
- [Fjt19a] K. Fujita, *A valuative criterion for uniform K -stability of \mathbb{Q} -Fano varieties*, J. Reine Angew. Math. **751** (2019), 309–338.
- [Fjt19b] K. Fujita, *Uniform K -stability and plt blowups of log Fano pairs*, Kyoto J. Math. **59** (2019), no. 2, 399–418.
- [Fjt19c] K. Fujita, *K -stability of Fano manifolds with not small alpha invariants*, J. Inst. Math. Jussieu **18** (2019), no. 3, 519–530.
- [Hum75] J. Humphreys, *Linear algebraic groups*, Graduate Texts in Mathematics, No. **21**. Springer-Verlag, New York-Heidelberg, 1975.
- [KM98] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, With the collaboration of C. H. Clemens and A. Corti. Cambridge Tracts in Math., **134**, Cambridge University Press, Cambridge, 1998.
- [KP18] A. Kuznetsov and Y. Prokhorov, *Prime Fano threefolds of genus 12 with a \mathbb{G}_m -action and their automorphisms*, Épijournal Geom. Algébrique **2** (2018), Art. 3, 14 pp.
- [KPS18] A. Kuznetsov, Y. Prokhorov and C. Shramov, *Hilbert schemes of lines and conics and automorphism groups of Fano threefolds*, Jpn. J. Math. **13** (2018), no. 1, 109–185.
- [Laz04a] R. Lazarsfeld, *Positivity in algebraic geometry, I: Classical setting: line bundles and linear series*, Ergebnisse der Mathematik und ihrer Grenzgebiete. (3) **48**, Springer, Berlin, 2004.

- [Laz04b] R. Lazarsfeld, *Positivity in algebraic geometry, II: Positivity for Vector Bundles, and Multiplier Ideals*, Ergebnisse der Mathematik und ihrer Grenzgebiete. (3) **49**, Springer, Berlin, 2004.
- [Li17] C. Li, *K-semistability is equivariant volume minimization*, Duke Math. J. **166** (2017), no. 16, 3147–3218.
- [Mat57] Y. Matsushima, *Sur la structure du groupe d'homéomorphismes analytiques d'une certaine variété kählérienne*, Nagoya Math. J. **11** (1957), 145–150.
- [MU83] S. Mukai and H. Umemura, *Minimal rational threefolds*, Algebraic geometry (Tokyo/Kyoto, 1982), 490–518, Lecture Notes in Math., **1016**, Springer, Berlin, 1983.
- [Pro90] Y. Prokhorov, *Automorphism groups of Fano 3-folds*, Uspekhi Mat. Nauk **45** (1990), no. 3 (273), 195–196.
- [SZ19] C. Stibitz and Z. Zhuang, *K-stability of birationally superrigid Fano varieties*, Compos. Math. **155** (2019), 1845–1852.
- [Tia87] G. Tian, *On Kähler-Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$* , Invent. Math. **89** (1987), no. 2, 225–246.
- [Tia97] G. Tian, *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **130** (1997), no. 1, 1–37.
- [Zhu20] Z. Zhuang, *Optimal destabilizing centers and equivariant K-stability*, arXiv:2004.09413v1.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA
UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN
Email address: fujita@math.sci.osaka-u.ac.jp