

# Solutions of the Dirac equation in one fixed and one moving wall well

Qiuyu Shan

## Abstract

It is very important which the Hamiltonian of the quantum system is time changing, especially the potential well that its width can change, the schrodinger equation and klein Gordon equation of this kind of circumstance are solved by some studies, but the Dirac equation haven't be solved, so this article discussed the solution of the Dirac equation in this kind of circumstance. We find that the momentum of the particle must be a imaginary number and it will be faster than light in a particular case. Then we discussed a new way to define the simultaneity which is different from Einstein.

Keywords: Time dependent potential well; Dirac equation; Klein Gordon equation; moving boundaries;  
PACS 03.65.Pm – Relativistic wave equations

## 1. Introduction

It is a common situation in quantum mechanics that the Hamiltonian changes with time. One useful example is the case of the one dimensional well which the well wall is moving, particularly the one dimensional infinite well that one wall of the well is still and another is moving with a constant speeds. S. W. Doescher and M. H. Rice<sup>[1]</sup> solved this problem in the nonrelativistic case, then D.N.Pinder<sup>[2]</sup> discussed the approximate method in the moving wall well, S. V. Mousavi<sup>[3]</sup> and Sara Di Martino<sup>1</sup>, Fabio Anz`a, Paolo Facchi, Andrzej Kossakowski, Giuseppe Marmo, Antonino Messina, Benedetto Militello, Saverio Pascazio<sup>[4]</sup> gave us further analyses of the solution of Schrodinger equation and Sebastián Carrasco, José Rogan, Juan Alejandro Valdivia<sup>[5]</sup> discussed how to controlling the Quantum State in this case.

E-mail: shanqiuyu@outlook.com

Michael Koehn<sup>[6-7]</sup> and Iwo Bialynicki-Birula<sup>[8]</sup> solved the Klein Gordon equation with the time varying potential. But as far as I know that we haven't discussed the finite potential well and we also haven't discussed the solution of Dirac equation in this case. So this paper will solve the Dirac equation in this potential well:

$$V(z) = \begin{cases} V_0 & (z \leq 0) & \text{I} \\ 0 & (0 < z < L_0 + vt - vt_0) & \text{II} \\ V_0 & (z \geq L_0 + vt - vt_0) & \text{III} \end{cases} \quad (1)$$

Where  $V_0$  is constant,  $v$  is the speed of the wall,  $L_0$  is the initial width of the well,  $t_0$  is the moment that the movement of wall begins, we can requiring that  $L_0 = vt_0$ , and we will use this assumption in this paper if no specification is stated, in other words make the range of  $V = 0$  is  $[0, vt]$ .

The Dirac equation can be written as

$$i\hbar \frac{\partial \psi}{\partial t} = -i\hbar c \left( \alpha^1 \frac{\partial \psi}{\partial x^1} + \alpha^2 \frac{\partial \psi}{\partial x^2} + \alpha^3 \frac{\partial \psi}{\partial x^3} \right) + \beta mc^2 \psi \quad (2)$$

If we choose a particular Basis, and discussed the particle which is up-spin and propagates in the  $z$  direction, The Dirac equation can be written as

$$i\hbar \frac{\partial \psi}{\partial t} = -i\hbar c \left( \alpha^3 \frac{\partial \psi}{\partial z} \right) + \beta mc^2 \psi \quad (3)$$

And the form of the solution is

$$\psi(z, t) = \begin{bmatrix} \varphi_0(z, t) \\ 0 \\ \varphi_2(z, t) \\ 0 \end{bmatrix} \quad (4)$$

## 2. Dirac equation in the time independent potential well

First, consider the potential well as

$$V(z) = \begin{cases} V_0 & (z \leq 0) & \text{I} \\ 0 & (0 < z < L_0) & \text{II} \\ V_0 & (z \geq L_0) & \text{III} \end{cases} \quad (5)$$

If the eigenvalue of the energy is  $E$ , the part of solution which is the function of space in this three regions can be written as

$$\text{I : } \varphi_{\text{I}}(z) = se^{ik_2z} \begin{pmatrix} 1 \\ 0 \\ \frac{\hbar ck_2}{E - V_0 + mc^2} \\ 0 \end{pmatrix} + be^{-ik_2z} \begin{pmatrix} 1 \\ 0 \\ -\frac{\hbar ck_2}{E - V_0 + mc^2} \\ 0 \end{pmatrix} \quad (6)$$

$$\text{II : } \varphi_{\text{II}}(z) = he^{ik_1z} \begin{pmatrix} 1 \\ 0 \\ \frac{\hbar ck_1}{E + mc^2} \\ 0 \end{pmatrix} + Je^{-ik_1z} \begin{pmatrix} 1 \\ 0 \\ -\frac{\hbar ck_1}{E + mc^2} \\ 0 \end{pmatrix} \quad (7)$$

$$\text{III : } \varphi_{\text{III}}(z) = qe^{ik_2z} \begin{pmatrix} 1 \\ 0 \\ \frac{\hbar ck_2}{E - V_0 + mc^2} \\ 0 \end{pmatrix} + re^{-ik_2z} \begin{pmatrix} 1 \\ 0 \\ -\frac{\hbar ck_2}{E - V_0 + mc^2} \\ 0 \end{pmatrix} \quad (8)$$

The solution in this three regions can be written as  $\psi_{\text{I}}(z,t) = \varphi_{\text{I}}(z)e^{\frac{-i(E-V_0)t}{\hbar}}$ ,  
 $\psi_{\text{II}}(z,t) = \varphi_{\text{II}}(z)e^{\frac{-iEt}{\hbar}}$ ,  $\psi_{\text{III}}(z,t) = \varphi_{\text{III}}(z)e^{\frac{-i(E-V_0)t}{\hbar}}$ .

Where  $k_2 = \frac{\sqrt{(E-V_0)^2 - m^2c^4}}{\hbar c}$ ,  $k_1 = \frac{\sqrt{E^2 - m^2c^4}}{\hbar c}$ , from the boundary conditions and

the eigenvalue of the energy we know that the constant  $a, b, h, J, q, r$  can be written as

$$z = 0 : \begin{cases} s + b = h + J \\ s - b = \frac{k_2(E + mc^2)}{k_1(E - V_0 + mc^2)}(h - J) \end{cases} \quad (9)$$

$$z = L_0 : \begin{cases} qe^{ik_2L_0} + re^{-ik_2L_0} = he^{ik_1L_0} + Je^{-ik_1L_0} \\ qe^{ik_2L_0} - re^{-ik_2L_0} = \frac{k_1(E - V_0 + mc^2)}{k_2(E + mc^2)}(he^{ik_1L_0} - Je^{-ik_1L_0}) \end{cases} \quad (10)$$

In particular, if  $V_0 > 0$  and  $|E - V_0| < mc^2$ ,  $k_2 = i|k_2|$ , The solution in I and III is an exponential function and  $b = q = 0$ .

### 3. Klein Gordon equation in a time-dependent potential well

We will discuss this potential well first,

$$V(z) = \begin{cases} 0 & 0 < z < vt \\ \infty & z \leq 0, z \geq vt \end{cases} \quad (11)$$

If the wave travels in the  $z$  direction, the Klein Gordon equation can be written as

$$\hbar^2 \frac{\partial^2 \varphi(z,t)}{\partial t^2} - \hbar^2 c^2 \frac{\partial^2 \varphi(z,t)}{\partial z^2} + m^2 c^4 \varphi(z,t) = 0 \quad (12)$$

The boundary conditions is  $\varphi(0,t) = \varphi(vt,t) = 0$ .

One can give the substitution

$$x = \sqrt{\frac{ct+z}{ct-z}}, \quad y = \sqrt{c^2 t^2 - z^2} \quad (13)$$

Like the method of Iwo Bialynicki-Birula<sup>[8]</sup> and Koehn M<sup>[6]</sup>, the Klein Gordon equation will be written as

$$\begin{aligned} & -\hbar^2 y^2 \frac{\partial^2 \varphi(x,y)}{\partial y^2} + \hbar^2 x^2 \frac{\partial^2 \varphi(x,y)}{\partial x^2} - \hbar^2 y \frac{\partial \varphi(x,y)}{\partial y} \\ & + \hbar^2 x \frac{\partial \varphi(x,y)}{\partial x} - m^2 c^2 y^2 \varphi(x,y) = 0 \end{aligned} \quad (14)$$

The boundary conditions is  $\varphi(1,y) = \varphi\left(\sqrt{\frac{c+v}{c-v}}, y\right) = 0$

We can assume that  $\varphi(x,y) = f(x)g(y)$  and obtain

$$\begin{aligned} & \frac{\hbar^2 x^2}{f(x)} \frac{d^2 f(x)}{dx^2} + \frac{\hbar^2 x}{f(x)} \frac{df(x)}{dx} \\ & = \frac{\hbar^2 y^2}{g(y)} \frac{d^2 g(y)}{dy^2} + \frac{\hbar^2 y}{g(y)} \frac{dg(y)}{dy} - m^2 c^2 y^2 = a \end{aligned} \quad (15)$$

Where  $a$  is a constant, it is not connected with  $x, y$ .

We can solve the two equations above easily,

$$f(x) = c_1 x^{\frac{\sqrt{a}}{\hbar}} + c_2 x^{-\frac{\sqrt{a}}{\hbar}} \quad (16)$$

$$g(y) = c_3 J_{\frac{\sqrt{a}}{\hbar}}\left(\frac{mcy}{\hbar}\right) + c_4 Y_{\frac{\sqrt{a}}{\hbar}}\left(\frac{mcy}{\hbar}\right) \quad (17)$$

Where  $c_1, c_2, c_3, c_4$  are constants,  $J_{\frac{\sqrt{a}}{\hbar}}\left(\frac{mcy}{\hbar}\right)$  and  $Y_{\frac{\sqrt{a}}{\hbar}}\left(\frac{mcy}{\hbar}\right)$  are Bessel function, we can get the solution of the Klein Gordon equation

$$\varphi(x, y) = \sum_a (c_1 x^{\frac{\sqrt{a}}{\hbar}} + c_2 x^{-\frac{\sqrt{a}}{\hbar}}) [c_3 J_{\frac{\sqrt{a}}{\hbar}}\left(\frac{mcy}{\hbar}\right) + c_4 Y_{\frac{\sqrt{a}}{\hbar}}\left(\frac{mcy}{\hbar}\right)] \quad (18)$$

Using the boundary condition and define  $ik_n = \frac{\sqrt{a}}{\hbar}$ ,  $k_n = \frac{n\pi}{\ln\sqrt{\frac{c+v}{c-v}}}$  we can get the

solution of the Klein Gordon equation in this potential well

$$\varphi(x, y) = \sum_n \sin(k_n \ln x) [c_3 J_{\frac{\sqrt{a}}{\hbar}}\left(\frac{mcy}{\hbar}\right) + c_4 Y_{\frac{\sqrt{a}}{\hbar}}\left(\frac{mcy}{\hbar}\right)] \quad (19)$$

## 4. Dirac equation in a time-dependent potential well

We will discuss the Dirac equation in the potential well as we defined in (1), the form of the Dirac equation in this case is

$$\begin{cases} i\hbar \frac{\partial \varphi_0(z, t)}{\partial t} = -i\hbar c \frac{\partial \varphi_2(z, t)}{\partial z} + mc^2 \varphi_0(z, t) \\ i\hbar \frac{\partial \varphi_2(z, t)}{\partial t} = -i\hbar c \frac{\partial \varphi_0(z, t)}{\partial z} - mc^2 \varphi_2(z, t) \end{cases} \quad (20)$$

If  $U_1(z, t) = \varphi_0(z, t) + \varphi_2(z, t)$ ,  $U_2(z, t) = \varphi_0(z, t) - \varphi_2(z, t)$ , one can obtain

$$\begin{cases} i\hbar \frac{\partial U_1(z, t)}{\partial t} = -i\hbar c \frac{\partial U_1(z, t)}{\partial z} + mc^2 U_2(z, t) \\ i\hbar \frac{\partial U_2(z, t)}{\partial t} = i\hbar c \frac{\partial U_2(z, t)}{\partial z} + mc^2 U_1(z, t) \end{cases} \quad (21)$$

And then

$$\hbar^2 \frac{\partial^2 U_1(z, t)}{\partial t^2} - \hbar^2 c^2 \frac{\partial^2 U_1(z, t)}{\partial z^2} + m^2 c^4 U_1(z, t) = 0 \quad (22)$$

This is a Klein Gordon equation in fact, so we define  $x = \sqrt{\frac{ct+z}{ct-z}}$ ,  $y = \sqrt{c^2 t^2 - z^2}$

and get the solution

$$\begin{aligned}
U_1(x, y) = & \sum_a (c_1 x^{\frac{\sqrt{a}}{h}}) [c_2 J_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h}) + c_3 Y_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h})] \\
& + (c_4 x^{-\frac{\sqrt{a}}{h}}) [c_5 J_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h}) + c_6 Y_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h})]
\end{aligned} \tag{23}$$

We also can calculate  $U_2(x, y)$

$$\begin{aligned}
U_2(x, y) = & \sum_a i(c_1 x^{\frac{\sqrt{a}}{h}-1}) [c_2 J_{\frac{\sqrt{a}}{h}-1}(\frac{mcy}{h}) + c_3 Y_{\frac{\sqrt{a}}{h}-1}(\frac{mcy}{h})] \\
& - i c_4 (x^{-\frac{\sqrt{a}}{h}-1}) [c_5 J_{\frac{\sqrt{a}}{h}+1}(\frac{mcy}{h}) + c_6 Y_{\frac{\sqrt{a}}{h}+1}(\frac{mcy}{h})]
\end{aligned} \tag{24}$$

And so

$$\begin{aligned}
\varphi_0(x, y) = & \frac{1}{2} \sum_a \{ (c_1 x^{\frac{\sqrt{a}}{h}}) [c_2 J_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h}) + c_3 Y_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h})] \\
& + (c_4 x^{-\frac{\sqrt{a}}{h}}) [c_5 J_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h}) + c_6 Y_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h})] \\
& + i(c_1 x^{\frac{\sqrt{a}}{h}-1}) [c_2 J_{\frac{\sqrt{a}}{h}-1}(\frac{mcy}{h}) + c_3 Y_{\frac{\sqrt{a}}{h}-1}(\frac{mcy}{h})] \\
& - i c_4 (x^{-\frac{\sqrt{a}}{h}-1}) [c_5 J_{\frac{\sqrt{a}}{h}+1}(\frac{mcy}{h}) + c_6 Y_{\frac{\sqrt{a}}{h}+1}(\frac{mcy}{h})] \}
\end{aligned} \tag{25}$$

$$\begin{aligned}
\varphi_2(x, y) = & \frac{1}{2} \sum_a \{ (c_1 x^{\frac{\sqrt{a}}{h}}) [c_2 J_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h}) + c_3 Y_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h})] \\
& + (c_4 x^{-\frac{\sqrt{a}}{h}}) [c_5 J_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h}) + c_6 Y_{\frac{\sqrt{a}}{h}}(\frac{mcy}{h})] \\
& - i(c_1 x^{\frac{\sqrt{a}}{h}-1}) [c_2 J_{\frac{\sqrt{a}}{h}-1}(\frac{mcy}{h}) + c_3 Y_{\frac{\sqrt{a}}{h}-1}(\frac{mcy}{h})] \\
& + i c_4 (x^{-\frac{\sqrt{a}}{h}-1}) [c_5 J_{\frac{\sqrt{a}}{h}+1}(\frac{mcy}{h}) + c_6 Y_{\frac{\sqrt{a}}{h}+1}(\frac{mcy}{h})] \}
\end{aligned} \tag{26}$$

We notice that  $L_0$  and  $vt_0$  can be substituted for each other in our calculation, if  $L_0 = vt_0$ ,  $L_1 = vt_1$ , the solution at  $t_1$ ,  $\varphi_0(z, t_1)$  and  $\varphi_2(z, t_1)$  should

satisfy the boundary condition which is given by  $L_0 = vt_1$ , so we can write the solution at one moment in the case that  $V = 0$ ,  $\varphi_0(z, t_1)$  and  $\varphi_2(z, t_1)$  as the sum of plane waves and calculate the solution in the region that  $V = V_0$  by using the method in the second part.

There is a important quantity in the equation of the potential well (1), it is the speed of the movement of the well,  $v$ , However, this velocity is not necessarily the velocity which the potential well wall moves, but may also be the velocity which the potential well wall builds up or collapses. It is interesting to note that the speed of building or collapsing of the wall could exceed the speed of light because it is not the speed of a really object. We will discuss the particle in the region II when  $v < -c$ .

If  $V_0 = \infty$ , we will get that  $U_1(x, y) = 0$  where  $x = 0, x = \sqrt{\frac{c+v}{c-v}}$ , we can know that

$\frac{\sqrt{a}}{\hbar}$  must be a real number and the momentum of the particle in the region II could be an imaginary number, it means that the speed of the particle in the region II would faster than light. We can find an example about it, when  $t = t_0$  and the mass of the particle is  $m$ , the wave function of the particle is

$$\varphi_0(x, y) = \frac{c_1}{2} \left[ x^{\frac{\sqrt{a}}{\hbar}} J_{\frac{\sqrt{a}}{\hbar}} \left( \frac{mcy}{\hbar} \right) + ix^{\frac{\sqrt{a}-1}{\hbar}} J_{\frac{\sqrt{a}-1}{\hbar}} \left( \frac{mcy}{\hbar} \right) \right] \text{ and}$$

$$\varphi_2(x, y) = \frac{c_1}{2} \left[ x^{\frac{\sqrt{a}}{\hbar}} J_{\frac{\sqrt{a}}{\hbar}} \left( \frac{mcy}{\hbar} \right) - ix^{\frac{\sqrt{a}-1}{\hbar}} J_{\frac{\sqrt{a}-1}{\hbar}} \left( \frac{mcy}{\hbar} \right) \right]$$

Where  $x = \sqrt{\frac{ct_0 + z}{ct_0 - z}}$ ,  $y = \sqrt{c^2 t_0^2 - z^2}$ , if  $\frac{\sqrt{a}}{\hbar}$  is real integers, we can get that the

average value of the momentum of the particle in the region II would be a complex

number, if the momentum is  $p = \frac{mv}{\sqrt{1 - \frac{v^2}{c^2}}}$ , we will know that there must be the

particle which the momentum od it is an imaginary number.

Another interesting conclusion is that the wave function of the particle in anywhere will change when the well begin to move or build or collapse. The change of the wave function and the moving of the well is simultaneity. That means that we can find a new way to define the simultaneity which is different from Einstein's method in his famous paper.<sup>[9]</sup>

## 5. Conclusion

It is very difficult for us to solve the exact solution of Schrodinger equation, Klein Gordon equation or Dirac equation if the Hamiltonian of the quantum system is time varying, solving the Dirac equation is even more difficult because it is an equation set, this paper discussed the solution of Dirac equation when the particle is moving in the one dimensional, finite potential well that one wall of the well is still and another is moving with a constant speeds, because the Klein tunneling, the properties of Dirac particle is very different from the solution of Schrodinger equation and Klein Gordon equation. We discussed the motion with uniform velocity only, but an extension of this paper is any movement of the potential well, it will be useful.

We also discussed the character of the particle when  $v < -c$  and find that its momentum must be a imaginary number and it will be faster than light. Then we discussed a new way to define the simultaneity.

## References

- [1] Doescher, and W. S. . "Infinite Square-Well Potential with a Moving Wall." American Journal of Physics 37.12(1969):1246-1249.
- [2] Pinder, and N. D. . "The contracting square quantum well." American Journal of Physics 58.1(1990):54-58.
- [3] Mousavi, S. V. . "Quantum effective force in an expanding infinite square-well potential and Bohmian perspective." Physica Scripta 86.3(2012):471-473.
- [4] Martino, S. D. , et al. "A quantum particle in a box with moving walls." Journal of Physics A Mathematical and Theoretical 46.36(2013).
- [5] Carrasco, S. , J. Rogan , and J. A. Valdivia . "Controlling the Quantum State with a time varying potential." Scientific Reports 7.1(2017).
- [6] Koehn.M. . "Solutions of the Klein-Gordon equation in an infinite square-well



potential with a moving wall." (2013).

[7] Koehn. M. . "Relativistic Wavepackets in Classically Chaotic Quantum Cosmological Billiards." *Physical Review D Particles & Fields* 85.6(2011).

[8] Bialynicki-Birula Iwo. "Solutions of the d'Alembert and Klein-Gordon equations confined to a region with one fixed and one moving wall." *Epl* 101.6(2013):60003.

[9] Einstein, B. A. . "On the Electrodynamics of Moving Bodies." *Journal of Genetics & Molecular Biology* 16.3(2005):160-183.