

VELDKAMP QUADRANGLES AND POLAR SPACES

BERNHARD MÜHLHERR AND RICHARD M. WEISS

ABSTRACT. Veldkamp polygons are certain graphs $\Gamma = (V, E)$ such that for each $v \in V$, Γ_v is endowed with a symmetric anti-reflexive relation \equiv_v . These relations are all trivial if and only if Γ is a thick generalized polygon. A Veldkamp polygon is called flat if no two vertices have the same set of vertices that are opposite in a natural sense. We explore the connection between Veldkamp quadrangles and polar spaces. Using this connection, we give the complete classification of flat Veldkamp quadrangles in which some but not all of the relations \equiv_v are trivial.

1. INTRODUCTION

In 2.8, we introduce the notion of a Veldkamp polygon. A Veldkamp polygon is a graph $\Gamma = (V, E)$ such that for each vertex $v \in V$, the set Γ_v of vertices adjacent to v is endowed with a symmetric, anti-reflexive relation \equiv_v satisfying several axioms. The relations \equiv_v are called the *local opposition relations* of Γ . If (u, v, w) is a path of length 2 and $u \equiv_v w$, we say that the path (u, v, w) is *straight*. An arbitrary path is straight if every subpath of length 2 is straight. Thus every path is straight if and only if the local opposition relations \equiv_v are all trivial (in the sense that $u \equiv_v w$ for all distinct $u, w \in \Gamma_v$). This is the case if and only if Γ is a thick generalized polygon.

The thick generalized polygons that arise in the theory of algebraic groups all satisfy the Moufang condition. This is a transitivity condition on the root groups. There are natural extensions of the notion of a root group and the Moufang condition to Veldkamp polygons. A Veldkamp polygon satisfying this Moufang condition is the same thing as a Tits polygon, a notion introduced in [3].

There is a standard way to construct a Tits polygon from an irreducible spherical building of rank at least 3 and a Tits index that is compatible in a suitable sense. Every irreducible spherical building of rank at least 3 has at least one compatible Tits index. Thus the theory of Tits polygons allows us, in particular, to study an arbitrary spherical building as a rank 2 structure on which the methods in [12] can be brought to bear (as in [3]–[9]). This was our original reason for introducing Tits polygons. In [4], we also showed that there is an equivalence between Tits polygons and a certain class of root graded groups.

In [15, Def. 3.1], Veldkamp introduced the notion of a Veldkamp plane. (Veldkamp called them Barbilian planes.) This notion arose in his work on projective geometry over rings. In the same paper Veldkamp showed the importance of the notion of the stable rank of a ring (a notion he found in K -theory) in ring-geometry,

Date: November 16, 2021.

2000 Mathematics Subject Classification. 20E42, 51E12, 51E24.

Key words and phrases. building, Tits polygon, polar space.

a subject first studied by Corrado Segre early in the 20th century (see [16]). Veldkamp was able to show that many of the basic results of projective geometry, with suitable modifications, hold over rings of stable rank 2. In particular, he showed that the geometries that arise from a free module of rank 3 over an associative ring of stable rank 2 and Veldkamp planes satisfying something slightly stronger than the Moufang condition (see [15, Def. 4.16]) are essentially the same. This result was subsequently strengthened first by Faulkner in [2, Thm 9] and then by ourselves in [5, Thms. 7.2 and 7.3].

A Veldkamp plane as defined by Veldkamp is the same thing as a Veldkamp n -gon as defined in 2.8 with $n = 3$, where the local opposition relations play the role of the relation $\not\approx$ in [15, Def. 3.1]. Thus our notion of a Veldkamp polygon is a natural generalization of the notion of a Veldkamp plane.

We say that a Veldkamp polygon is *green* if some (or all) of its local opposition relations are trivial, and we call a Veldkamp polygon *flat* if no two vertices have the same set of vertices that are opposite as defined in 2.9. All generalized polygons are both green and flat.

In this paper, the focus is on Veldkamp quadrangles (i.e. a Veldkamp n -gon with $n = 4$). We prove four theorems. In our first theorem (see 4.23, 4.26 and 4.30), we show that every green Veldkamp quadrangle Γ has a canonical flat quotient $\bar{\Gamma}$ (which is also a green Veldkamp quadrangle) and a natural surjective homomorphism π from Γ to $\bar{\Gamma}$.

In our second theorem (see 5.18 and 5.22), we show that there is a one-to-one correspondence (up to isomorphism) between flat green Veldkamp quadrangles and non-degenerate polar spaces with thick lines.

The notion of a polar space was also introduced by Veldkamp. Polar spaces of rank 2 are generalized quadrangles. Non-degenerate polar spaces of finite rank at least 3 and characteristic different from 2 were classified by Veldkamp in [14]. This result was extended to arbitrary characteristic by Tits in [11]. Several people contributed to the extension of Tits' result to include polar spaces of infinite rank. We cite [10] for this part of the classification.

In our third theorem (see 6.16), we use the classification of non-degenerate polar spaces of rank at least 3 to show that every flat green Veldkamp quadrangle is either a generalized quadrangle or a Tits quadrangle obtained from a quadratic space or a pseudo-quadratic space (which we interpret to include the symplectic case) or a Tits quadrangle of type D_3 or type E_7 as defined in 6.13 and 6.15. In particular, a flat green Veldkamp quadrangle that is not a generalized quadrangle always satisfies the Moufang condition.

There are, of course, also many generalized quadrangles that do not satisfy the Moufang condition. Generalized quadrangles that do satisfy the Moufang condition were classified in [12, 17.4]. There are six families. Three of them have a natural description in terms of polar spaces, but three of them do not; see 6.17.

In our fourth theorem (see 6.5 and 6.12), we show that to every non-degenerate quadratic space and every non-degenerate pseudo-quadratic space (with infinite Witt index allowed), there does, in fact, exist a corresponding flat green Veldkamp quadrangle.

Conventions 1.1. As in [12], we set

$$[a, b] = a^{-1}b^{-1}ab$$

for elements a, b in a group and we apply permutations from left to right.

ACKNOWLEDGMENT: The work of the first author was partially supported by DFG Grant MU1281/7-1 and the work of the second author was partially supported by Simons Foundation Collaboration Grant 516364.

2. VELDKAMP POLYGONS

In this section we introduce the notion of a Veldkamp polygon. In 2.15, we describe the connection between Veldkamp polygons and generalized polygons.

Definition 2.1. An *opposition relation* on a set X is a relation on X that is symmetric and anti-reflexive. Let \equiv be an opposition relation on X . We say that two elements of X are *opposite* if $x \equiv y$. We say that \equiv is *trivial* if every two distinct elements of X are opposite.

Definition 2.2. Let \equiv be an opposition relation on a set X . For each $k \geq 1$, we say that \equiv is *k-plump* if for every subset S of X such that $|S| \leq k$, there exists an element of X that is opposite every element of S . Note that if \equiv is *k-plump* for some $k \geq 1$, then it is also $(k - 1)$ -plump and $|X| \geq k + 1$.

Notation 2.3. By *graph* we mean a pair (V, E) consisting of a set V and a set E of 2-element subsets of V . The elements of V and E are called *vertices* and *edges*. For each $v \in V$, we let Γ_v denote the set of *neighbors* of v . In other words,

$$\Gamma_v = \{u \in V \mid \{u, v\} \in E\}.$$

Let $\Gamma = (V, E)$ be a graph. A *subgraph* of Γ is a pair (V_1, E_1) , where V_1 is a non-empty subset of V and E_1 is a subset of $\{e \in E \mid e \subset V_1\}$. For each non-empty subset V_1 of V , the subgraph of Γ *spanned by* V_1 is the subgraph (V_1, E_1) , where $E_1 = \{e \in E \mid e \subset V_1\}$. An *s-path* in Γ is a sequence (x_0, x_1, \dots, x_s) of $s + 1$ vertices x_i for some $s \geq 0$ such that $x_{i-1} \in \Gamma_{x_i}$ for all $i \in [1, s]$, and $x_{i-2} \neq x_i$ for all $i \in [2, s]$. (Here $[i, j]$ denotes the closed interval $i, i + 1, \dots, j$ if $i \leq j$ and $[i, j] = \emptyset$ if $i > j$.) A *path* is an *s-path* for some $s \geq 0$; the parameter s is called the *length* of the path. The graph Γ is *connected* if there is a path from any given vertex to any other given vertex. The distance $\text{dist}(x, y)$ between two vertices is the minimal length of a path from x to y (assuming that there is one). For each $v \in V$, let

$$\Gamma_m(v) = \{u \in V \mid \text{dist}(u, v) = m\}.$$

Thus $\Gamma_1(v) = \Gamma_v$ for all $v \in V$.

Notation 2.4. An *s-path* is *closed* if its first and last vertices are the same and $s \geq 3$. An *s-circuit* is the subgraph with vertex set $\{x_1, \dots, x_s\}$ and edge set $\{\{x_{i-1}, x_i\} \mid i \in [1, s]\}$ for some closed *s-path* (x_0, x_1, \dots, x_s) . A *circuit* is an *s-circuit* for some $s \geq 3$.

Notation 2.5. Let $\Gamma = (V, E)$ be a graph endowed with an opposition relation \equiv_v on Γ_v for each $v \in V$. We say that two vertices x, y are *opposite at* v if $x, y \in \Gamma_v$ and $x \equiv_v y$. A path (v_0, v_1, \dots, v_s) is called *straight* if v_{i-1} is opposite v_{i+1} at v_i for all $i \in [1, s - 1]$. A circuit is *straight* if every path it contains is straight. We say

that Γ is k -plump for some $k \geq 1$ if all its local opposition relations are k -plump as defined in 2.2.

Definition 2.6. A *Veldkamp graph* is a graph $\Gamma = (V, E)$ endowed with a 2-plump opposition relation on Γ_v for each $v \in V$.

Remark 2.7. Since every Veldkamp graph is 2-plump (and therefore also 1-plump), it follows that for every $s \geq 0$, a straight s -path in a Veldkamp graph can be extended to a straight $(s + 1)$ -path.

Definition 2.8. Let $n \geq 2$. A *Veldkamp n -gon* is a Veldkamp graph Γ such that the following hold:

- (VP1) Γ is connected and bipartite.
- (VP2) For each $k \in [1, n - 1]$ and each straight k -path $\alpha = (v_0, \dots, v_k)$, α is the unique straight path from v_0 to v_k of length at most k .
- (VP3) Every straight $(n + 1)$ -path is contained in a straight $2n$ -circuit.

Note that by (VP2), the straight $2n$ -circuit in (VP3) is unique. A *Veldkamp polygon* is a Veldkamp n -gon for some $n \geq 2$.

Definition 2.9. Let $\Gamma = (V, E)$ be a Veldkamp n -gon for some $n \geq 2$. A *root* in Γ is a straight n -path. Two vertices x and y are *opposite* if there is a root from x to y and two edges e, f of Γ are *opposite* if there exists a straight $(n + 1)$ -path whose first and last edges are e and f . By (VP2), neither a vertex nor an edge can be opposite itself, so both of these relations are, in fact, opposition relations (on V and on E). The set of vertices opposite a given vertex x is denoted by x^{op} and Γ is called *flat* if $x^{\text{op}} = y^{\text{op}}$ for $x, y \in V$ implies that $x = y$.

Suppose for the rest of this section that $\Gamma = (V, E)$ is a Veldkamp n -gon for some $n \geq 2$.

Proposition 2.10. *Suppose that $x, y \in V$ are opposite and that $z \in \Gamma_y$. Then there exists a unique root (x, \dots, z, y) from x to y containing z .*

Proof. Uniqueness holds by (VP2). Since x and y are opposite, there is a root (x, \dots, w, y) from x to y . Since Γ is 2-plump, we can choose $u \in \Gamma_y$ opposite w and z at y . Thus (x, \dots, w, y, u) is a straight $(n + 1)$ -path. By (VP3), it follows that there exists a root (x, \dots, u, y) from x to y that contains u . Then (x, \dots, u, y, z) is a straight $(n + 1)$ -path and the claim follows by a second application of (VP3). \square

Remark 2.11. Suppose that $u, v \in V$ and $\text{dist}(u, v)$ is even. Since Γ is bipartite, every vertex in u^{op} is at even distance to every vertex in v^{op} .

Proposition 2.12. *For each pair e_1, e_2 of edges of Γ , there exists an edge that is opposite both e_1 and e_2 .*

Proof. Let (x_0, \dots, x_k) be a k -path in Γ for some $k \geq 1$ and let c_i denote the edge $\{x_{i-1}, x_i\}$ for each $i \in [1, k]$. Since Γ is connected, it will suffice to show that there exists an edge opposite both c_1 and c_k . We proceed by induction with respect to k . By 2.7, the claim holds for $k = 1$. Suppose that $k \geq 2$ and that f is an edge that is opposite c_1 and c_{k-1} . Let y be the unique vertex of f that is opposite x_{k-1} . Since Γ is bipartite, y is opposite to a unique vertex on c_1 . By 2.11, this vertex is at even distance from x_{k-1} . Thus y is opposite x_p , where $p = 0$ if k is odd and $p = 1$ if k is even.

Let $\alpha = (x_{k-1}, x_k, \dots, y)$ be the unique root from x_{k-1} to y containing x_k and let u be unique the element of Γ_y contained in α . If k is odd, let $\beta = (x_0, x_1, \dots, y)$ be the unique root from x_0 to y containing x_1 and if k is even, let $\beta = (x_1, x_0, \dots, y)$ be the unique root from x_1 to y containing x_0 . In both cases, let v be unique element of Γ_y contained in β . Since Γ is 2-plump, we can choose a vertex $z \in \Gamma_y$ that is opposite u and v . Let $e = \{y, z\}$. Attaching z at the end of α and at the end of β yields straight $(n+1)$ -paths. Hence e is an edge opposite both c_1 and c_k . \square

Corollary 2.13. *Let u, v be vertices such that $\text{dist}(u, v)$ is even. Then $u^{\text{op}} \cap v^{\text{op}} \neq \emptyset$.*

Proof. Let e be an edge containing u and let f be an edge containing v . By 2.12, there exists an edge g opposite both e and f . Let x be the unique vertex of g that is opposite u and let y be the unique vertex of g that is opposite v . By 2.11, $x = y$ and thus $x \in u^{\text{op}} \cap v^{\text{op}}$. \square

Definition 2.14. We recall that a *generalized n -gon* for some $n \geq 2$ is a bipartite graph Ω such that the following hold:

- (i) $\text{diam}(\Omega) = n$.
- (ii) Ω contains no circuits of length less than $2n$.
- (iii) $|\Omega_v| \geq 2$ for all vertices v .

A generalized n -gon Ω is *thick* if $|\Omega_v| \geq 3$ for all vertices v .

Proposition 2.15. *If the local opposition relations are all trivial, then Γ is a thick generalized n -gon, and every thick generalized n -gon, when endowed with the trivial local opposition relation at every vertex, is a Veldkamp n -gon.*

Proof. Suppose that all local opposition relations of Γ are trivial. Then every path is straight. By (VP1), Γ is connected. By (VP3), therefore, $\text{diam}(\Gamma) \leq n$. By (VP2), Γ contains no circuits of length less than $2n$. This implies that $\text{diam}(\Gamma) \geq n$. Since Γ is 2-plump, we have $|\Gamma_x| \geq 3$ for all vertices x . It follows that Γ is a thick generalized n -gon.

Suppose, conversely, that Ω is a thick generalized n -gon endowed with the trivial local opposition relation at every vertex, so that every path is straight. Since Ω is thick, it is 2-plump. By 2.14(ii), Ω satisfies (VP2) and by 2.14(i), it follows that (VP1) and (VP3) hold. Thus Ω is a Veldkamp n -gon. \square

Proposition 2.16. *Suppose that (x_0, x_1, \dots, x_n) and (y_0, y_1, \dots, y_n) are two roots with $x_0 = y_0$ and $x_n = y_n$. Then x_1 and y_1 are opposite at x_0 if and only if x_{n-1} and y_{n-1} are opposite at x_n .*

Proof. Suppose that x_{n-1} and y_{n-1} are opposite at x_n . Then

$$(x_0, x_1, \dots, x_{n-1}, x_n, y_{n-1})$$

is a straight $(n+1)$ -path. By (VP3), there exists a root $\alpha := (z_0, z_1, \dots, z_n)$ such that $z_0 = x_0$, $z_{n-1} = y_{n-1}$, $z_n = x_n$ and z_1 is opposite x_1 at x_0 . By (VP2), it follows that $\alpha = (y_0, y_1, \dots, y_n)$. Hence x_1 is opposite y_1 at x_0 . The converse holds by symmetry. \square

Proposition 2.17. *Suppose that \equiv_v is trivial for some $v \in V$. Then \equiv_w is trivial for all w at even distance from v and if, in addition, n is odd, then \equiv_w is trivial for all $w \in V$.*

Proof. Let (x_0, \dots, x_n) be a root. By 2.16, \equiv_{x_0} is trivial if and only if \equiv_{x_n} is trivial. Thus if $\text{dist}(v, w)$ is even for some $w \in V$, then by 2.13, \equiv_w is trivial. If n is odd, then by (VP1), every vertex of V is at even distance from x_0 or from x_n , so if one local opposition relation is trivial, then they all are. \square

3. VELDKAMP QUADRANGLES

In this section we focus on the case $n = 4$. Let $\Gamma = (V, E)$ be a Veldkamp quadrangle. Since Γ is connected and bipartite, there is a unique decomposition of V into two subsets such that every edge contains one vertex from each of them. We choose one of these two subsets and call it P and the other we call L . We refer to the elements of P as *points* and to the elements of L as *lines*.

Notation 3.1. We call Γ *green* if the local opposition relations at lines are all trivial.

We assume for the rest of this section that our Veldkamp quadrangle Γ is green.

Proposition 3.2. Γ does not contain circuits of length 4.

Proof. Let (a, x, b, y, c) be a 4-path with $c = a \in P$. Since Γ is green, (a, x, b) and (a, y, b) are distinct straight paths from a to b . By (VP2), this is not allowed. \square

Notation 3.3. Let x, y be vertices such that $\text{dist}(x, y) = 2$. By 3.2, there is a unique vertex in $\Gamma_x \cap \Gamma_y$. We denote this vertex by $x \wedge y$.

Proposition 3.4. A 6-circuit does not contain any straight 2-paths (x, a, y) such that $a \in P$.

Proof. Let (a, x, b, y, c, z, a) be a closed 6-path with $a \in P$. It suffices to show that (x, b, y) is not straight. Suppose that it is. Then (a, x, b, y, c) is a root, so by 2.9, a and c are opposite. By 2.10, there is a root (a, u, d, z, c) from a to c that contains z . Thus (a, z) and (a, u, d, z) are both straight paths from a to z . By (VP2), this is not allowed. \square

Proposition 3.5. If $a, b \in P$ are opposite, then $\text{dist}(a, b) = 4$.

Proof. Suppose that $a, b \in P$ are opposite. By 2.9, there is a root from a to b . In particular, $\text{dist}(a, b) \leq 4$. By 3.2, $a \neq b$ and then by 3.4, $\text{dist}(a, b) \neq 2$. Hence $\text{dist}(a, b) = 4$. \square

Proposition 3.6. $\text{diam}(\Gamma) = 4$.

Proof. By 3.5, it suffices to show that $\text{dist}(a, u) \leq 3$ for all $a \in P$ and $u \in L$. Let $a \in P$ and $u \in L$. Choose $b \in \Gamma_u$. By 2.13, we can choose $c \in a^{\text{op}} \cap b^{\text{op}}$. By 2.10, there is a root (b, x_1, x_2, x_3, c) from b to c such that $x_1 = u$ and a root (a, y_1, y_2, y_3, c) from a to c such that $y_3 = x_3$. If $y_2 = x_2$, then (a, y_1, x_2, x_1) is a 3-path from a to u . Suppose that $y_2 \neq x_2$. Then $(a, y_1, y_2, x_3, x_2, x_1)$ is a 5-path from a to u . Since Γ is green, this path is straight. By (VP3), therefore, $\text{dist}(a, u) \leq 3$ also in this case. \square

Proposition 3.7. Let $a, b \in P$ and suppose that $\text{dist}(a, b) = 4$ but a and b are not opposite. Then $a^{\text{op}} = b^{\text{op}}$ and $\Gamma_2(a) = \Gamma_2(b)$.

Proof. Suppose there exists $c \in \Gamma_2(b) \cap a^{\text{op}}$ and let $x = b \wedge c$. By 2.10, there exists a root (a, y_1, y_2, y_3, c) from a to c with $y_3 = x$. Since Γ is green, it follows that

(a, y_1, y_2, y_3, b) is a root. Since $a \notin b^{\text{op}}$ by hypothesis, we obtain a contradiction. Therefore

$$(3.8) \quad \Gamma_2(b) \cap a^{\text{op}} = \emptyset.$$

Now choose $x \in \Gamma_b$. We claim that

$$(3.9) \quad \Gamma_x \setminus \{b\} \subset \Gamma_2(a).$$

By 2.13, we can choose $c \in a^{\text{op}} \cap b^{\text{op}}$. Let (b, x_1, x_2, x_3, c) be the unique root from b to c such that $x_1 = x$ and let (c, y_1, y_2, y_3, a) be the unique root from c to a such that $y_1 = x_3$. By (3.8), (a, y_3, y_2, y_1, x_2) is not a root. Since Γ is green, it follows that $x_2 = y_2$.

Choose $z \in \Gamma_c$ opposite x_3 at c . Let (c, z_1, z_2, z_3, b) and (c, z'_1, z'_2, z'_3, a) be the unique roots from c to b and from c to a such that $z_1 = z'_1 = z$. The closed 8-path $(b, z_3, z_2, z, c, x_3, x_2, x, b)$ is straight. By 2.16, (x, b, z_3) is also straight.

If $z_2 \neq z'_2$, then $(a, z'_3, z'_2, z'_1, z_2)$ is a root and hence $z_2 \in \Gamma_2(b) \cap a^{\text{op}}$. By (3.8), it follows that $z_2 = z'_2$.

Let $d \in \Gamma_x \setminus \{x_2, b\}$. To prove (3.9), we need to show that $d \in \Gamma_2(a)$. The path (d, x, x_2, x_3, c) is a root. Let (c, w_1, w_2, w_3, d) be the unique root from c to d such that $w_1 = z$. By 3.4, we have $w_2 \neq z_2$ since otherwise the straight 2-path (x, b, z_3) is contained in the closed 6-path $(x, b, z_3, z_2, w_3, d, x)$. Therefore (a, z'_3, z_2, z, w_2) is a root. Hence w_2 and a are opposite. Let (w_2, v_1, v_2, v_3, a) be the unique root from w_2 to a such that $v_1 = w_3$. If $v_2 \neq d$, then (d, w_3, v_2, v_3, a) is a root, but this implies that $d \in \Gamma_2(b) \cap a^{\text{op}}$. By (3.8), we must have $v_2 = d$. Therefore $d \in \Gamma_2(a)$. Thus (3.9) holds as claimed.

By 3.6, (3.8) and symmetry, we have $\Gamma_2(a) = \Gamma_2(b)$. Thus

$$(3.10) \quad \Gamma_2(e) = \Gamma_2(f)$$

for all $e, f \in P$ such that $\text{dist}(e, f) = 4$ and $e \notin f^{\text{op}}$. It remains to show that $a^{\text{op}} = b^{\text{op}}$. Suppose that $c \in a^{\text{op}} \setminus b^{\text{op}}$ and let (a, x_1, x_2, x_3, c) be a root from a to c . Thus $z \in \Gamma_2(c) \cap a^{\text{op}}$ for all $z \in \Gamma_{x_3} \setminus \{x_2, c\}$. Therefore

$$(3.11) \quad \Gamma_2(c) \cap a^{\text{op}} \neq \emptyset.$$

Since a and b are not opposite, we have $c \neq b$. By (3.8), $\text{dist}(b, c) \neq 2$. Thus $\text{dist}(b, c) = 4$ by 3.6. Hence $\Gamma_2(a) = \Gamma_2(b) = \Gamma_2(c)$ by (3.10). It follows that $\Gamma_2(a) \cap a^{\text{op}} \neq \emptyset$ by (3.11). By 3.5, this is impossible. With this contradiction, we conclude that $a^{\text{op}} = b^{\text{op}}$. \square

Corollary 3.12. *Suppose that Γ is flat as defined in 2.9 and let $a, b \in P$. Then a and b are opposite if and only if $\text{dist}(a, b) = 4$.*

Proof. This holds by 2.9, 3.5 and 3.7. \square

4. FLAT QUOTIENTS

Our goal in this section is to show that every green Veldkamp quadrangle has a canonical flat quotient. The precise result is formulated in 4.23 and 4.30.

For the rest of this section, we assume that Γ is a green Veldkamp quadrangle and continue with the notation in the previous section.

Notation 4.1. A *weed* is a non-straight 4-path (a, x, b, y, c) such that $a, b, c \in P$ and $\text{dist}(a, c) = 4$.

Remark 4.2. Note that a root cannot also be a weed since roots are straight and weeds are not.

Remark 4.3. Let $\alpha = (a, x, b, y, c)$ be a weed. The point b is called the *middle point* of α . By 3.2, the middle point of a weed α is uniquely determined by the two lines contained in α .

Notation 4.4. Let $a \simeq b$ for $a, b \in P$ if either $a = b$ or there is a weed that begins at a and ends at b .

Proposition 4.5. *Suppose that $u \in v^{\text{op}}$ for some $u, v \in P \cup L$ and $a \simeq b$ for some $a, b \in P$ such that $a \neq b$. Then the following hold:*

- (i) *Every 4-path from u to v is a root and for every $w \in \Gamma_u$, there is a unique 4-path from u to v that contains w .*
- (ii) *For every $x \in \Gamma_a$, there exists a 4-path from a to b containing x , every 4-path from a to b is a weed and $a \notin b^{\text{op}}$.*

Proof. Let $w \in \Gamma_u$. By 3.5 and 3.6, we have $\text{dist}(w, v) = 3$. Hence there exists a 4-path $\alpha = (u, p, q, w, v)$ from u to v that contains w . By 2.10, there exists a root $\beta = (u, m, n, w, v)$ from u to v that passes through w . By 3.4, $q = n$. By 3.2, therefore, $\alpha = \beta$. Hence α is a root and α is the only 4-path from u to v that contains w . Thus (i) holds.

Let $x \in \Gamma_a$. Since $a \neq b$, there exists a weed α from a to b . Hence $\text{dist}(a, b) = 4$ and by 3.6, there exists a 4-path β from a to b containing x . Applying (i), we observe that since α is not a root (by 4.2), neither is β a root. Hence β is a weed. Thus (ii) holds. \square

Proposition 4.6. *Let $a, b \in P$. Then the following hold:*

- (i) *$a \simeq b$ if and only if $a^{\text{op}} = b^{\text{op}}$.*
- (ii) *If $a \simeq b$, then $\Gamma_2(a) = \Gamma_2(b)$.*

Proof. It suffices to assume that $a \neq b$. Suppose that $a \simeq b$. Then $\text{dist}(a, b) = 4$ but by 4.5(ii), a and b are not opposite. By 3.7, therefore, $a^{\text{op}} = b^{\text{op}}$ and $\Gamma_2(a) = \Gamma_2(b)$. Suppose, conversely, that $a^{\text{op}} = b^{\text{op}}$. If $\text{dist}(a, b) = 2$, then (because Γ is green) there exists a root α starting at a and containing b , but then the last vertex of α is contained in $a^{\text{op}} \setminus b^{\text{op}}$. Hence $\text{dist}(a, b) = 4$ by 3.6. Let β be a 4-path from a to b . Since a is not opposite itself, it is not opposite b . It follows that β is a weed and hence $a \simeq b$. \square

Notation 4.7. By 4.6(i), \simeq is an equivalence relation. We denote the equivalence class of a point a by $[a]$.

Notation 4.8. Let $x \sim y$ for $x, y \in L$ if there is a weed containing both x and y . Thus if $x \sim y$, there exist $a, b, c \in P$ such that (a, x, b, y, c) is a weed. Let \approx denote the transitive-reflexive closure of the relation \sim on L . For each $x \in L$, let $[x]$ denote the equivalence class containing x . (Thus all the equivalence classes are singletons if there are no weeds.)

Proposition 4.9. *Let $x, y, z \in \Gamma_a$ for some $a \in P$. Suppose that x is opposite y at a and that $y \sim z$. Then x is opposite z at a .*

Proof. Since $y \sim z$, there exist $b, c \in P$ such that (c, z, a, y, b) is a weed. By 4.4 and 4.6(ii), we have $b \simeq c$ and $\Gamma_2(b) = \Gamma_2(c)$. Choose $d \in \Gamma_x \setminus \{a\}$. By 3.2, the

points b, c, d are pairwise distinct. Since x and y are opposite at a , (d, x, a, y, b) is a root and thus b and d are opposite. By 3.5, therefore, $d \notin \Gamma_2(b)$. Hence $d \notin \Gamma_2(c)$. By 3.6, we have $\text{dist}(c, d) = 4$. Suppose now that x and z are not opposite at a . Then (c, z, a, x, d) is a weed and hence $c \simeq d$. Since $b \simeq c$, it follows that $b \simeq d$. By 4.5(ii), this is impossible. With this contradiction, we conclude that x is opposite z at a . \square

Proposition 4.10. *Let $x, y \in L$ and suppose that $x \approx y$. Then $x^{\text{op}} = y^{\text{op}}$.*

Proof. It suffices to assume that $x \sim y$ and, by symmetry, to show that $x^{\text{op}} \subset y^{\text{op}}$. Let $a = x \wedge y$ and choose $z \in x^{\text{op}}$. By 4.5(i), there exists a root (x, a, u, c, z) from x to z that contains a . The lines x and u are opposite at a . By 4.9, it follows that u is opposite y at a . Thus (y, a, u, c, z) is also a root. Hence $z \in y^{\text{op}}$. \square

We will need the following result in the proof of 4.12.

Lemma 4.11. *Let $a, b, c, e, f \in P$ and $s, t, u, v, w, x, y, z \in L$ and suppose that $x^{\text{op}} = y^{\text{op}}$, that (b, x, a, y, c, u, b) and $\xi := (b, w, e, t, c, u, b)$ are both closed 6-paths and that the 4-paths (y, c, t, e, z) , (y, c, v, f, s) and (x, b, w, e, z) are all roots. Then $\text{dist}(b, f) = 2$.*

Proof. Note that (a, y, c, v, f) is a root and hence a and f are opposite. By 3.5, $a \notin \Gamma_2(f)$ and hence $b \neq f$. By 3.6, therefore, it suffices to show that $\text{dist}(b, f) \neq 4$. We suppose that $\text{dist}(b, f) = 4$. Since $a \in \Gamma_2(b)$, we have $\Gamma_2(b) \neq \Gamma_2(f)$. By 4.6(ii), it follows that b and f are opposite. By 4.5(i), therefore, $\alpha := (b, u, c, v, f)$ is a root.

Suppose now that $v = t$. Then (u, c, v) lies on the root α and on the closed 6-path ξ . By 3.4, this is impossible. Hence $v \neq t$. Hence by 3.2, we have $e \neq f$. Suppose that there exists $r \in \Gamma_e \cap \Gamma_f$. By 4.5(i), (b, w, e, r, f) is a root. Since (x, b, w, e, z) is a root, it follows that (x, b, w, e, r) is also a root. Hence $r \in x^{\text{op}} = y^{\text{op}}$. By 4.5(i), $\delta := (r, f, v, c, y)$ is a root. By 3.5, $r \neq t$ and $r \neq v$. It follows that $\zeta := (r, f, v, c, t, e, r)$ is a closed 6-path. The path (r, f, v) is contained in both δ and ζ . By 3.4 again, this is impossible. Hence the line r does not exist and thus $\text{dist}(e, f) = 4$ by 3.6.

Since b and f are opposite, there exists a root (b, w, g, p, f) from b to f that contains w . Since $\text{dist}(e, f) = 4$, we have $g \neq e$. Since (x, b, w, e, z) is a root, it follows that (x, b, w, g, p) is a root. Thus $p \in x^{\text{op}} = y^{\text{op}}$. By 4.5(i), therefore, (y, c, v, f, p) is a root. Hence (g, p, f, v, c) is a root. Thus g and c are opposite and therefore the $\beta := (g, w, e, t, c)$ is also a root, again by 4.5(i). The path (w, e, t) thus lies on the root β and on the closed 6-path ξ . By 3.4, this is impossible. With this last contradiction, we conclude that $\text{dist}(b, f) \neq 4$. \square

Proposition 4.12. *Let (x, a, y) be a path for some $a \in P$ and some $x, y \in L$ and suppose that $x^{\text{op}} = y^{\text{op}}$. Then for each $c \in \Gamma_y \setminus \{a\}$, there exists $b \in \Gamma_x \setminus \{a\}$ such that (b, x, a, y, c) is a weed.*

Proof. If (x, a, y) is straight, then there exists a root starting at x and containing a and y , but then the last vertex on this root is in x^{op} but not in y^{op} . Hence (x, a, y) is not straight.

Choose $c \in \Gamma_y \setminus \{a\}$. Let $z \in y^{\text{op}}$ and let (y, c, t, e, z) be the root from y to z that contains c . Then (a, y, c, t, e) is also a root, so a is opposite e . Hence by 3.4, $a \notin \Gamma_2(e)$. Since $x^{\text{op}} = y^{\text{op}}$, there exists a unique root (x, b, w, e, z) from x to z that contains e . Since $b \in \Gamma_2(e)$, we have $a \neq b$. Thus by 3.2, $b \neq c$. We want to

show that (b, x, a, y, c) is a weed. By 3.6, it suffices to show that $\text{dist}(b, c) \neq 2$. We assume, therefore, that there exists $u \in \Gamma_b \cap \Gamma_c$.

By 3.2, $u \neq x$ and $u \neq y$. Hence (a, x, b, u, c, y, a) is a closed 6-path. By 3.4, therefore, (x, b, u) is not straight. Since (x, b, w) is straight, we have $u \neq w$. Since (b, w, e) is contained in a root, it is straight and hence $b \neq e$. We already know that $b \neq c$. It follows by 3.2, $u \neq t$ and $w \neq t$. Thus $\xi = (b, w, e, t, c, u, b)$ is a closed 6-path.

Now suppose that (y, c, v, f, s) is an arbitrary root starting with y and then c . Thus $s \in y^{\text{op}} = x^{\text{op}}$. By 4.11, $\text{dist}(b, f) = 2$. By 4.5(i), therefore, (x, b, r, f, s) is the unique root from x to s that contains f , where $r = b \wedge f$. We conclude that b does not depend on the choice of z . Thus u is also independent of the choice of z . Since Γ is 2-plump, we can assume that z is chosen so that t is opposite u at c . Thus the 2-path (t, c, u) is both straight and contained in the 6-circuit ξ . By 3.4, this is impossible. With this contradiction, we conclude that $\text{dist}(b, c) \neq 2$. \square

Proposition 4.13. *Let $x, y \in L$. Then $x^{\text{op}} = y^{\text{op}}$ if and only if $x \approx y$.*

Proof. Suppose that $x^{\text{op}} = y^{\text{op}}$. We claim that $x \approx y$. By 4.12, we have $x \sim y$ if $\text{dist}(x, y) = 2$. By 3.6, therefore, it suffices to assume that $\text{dist}(x, y) = 4$. Let (x, a, z, b, y) be a 4-path from x to y . By 4.12 again, it will suffice to show that $x^{\text{op}} = z^{\text{op}}$ or $z^{\text{op}} = y^{\text{op}}$. By 4.10, we can assume that there is no weed containing x and z and no weed containing z and y . Since x is not opposite to itself, it is not opposite to y . Since $\text{dist}(x, y) = 4$, it follows that (x, a, z) or (z, b, y) is not straight. By symmetry, it suffices to consider the case that (x, a, z) is not straight. Since there is no weed containing x and z , we thus have

$$(4.14) \quad \Gamma_x \subset \Gamma_2(b).$$

Let $w \in x^{\text{op}}$. There exists a root (x, a, u, c, w) from x to w that contains a and a root (y, b, v, d, w) from y to w that contains b . Since (x, a, u) is straight but (x, a, z) is not, we have $u \neq z$ and hence $u \neq v$ by 3.2. We claim that also $c \neq d$. Suppose, instead, that $c = d$. An element of Γ_d is opposite x (respectively y) if and only if it is opposite u (respectively v) at d . Since $x^{\text{op}} = y^{\text{op}}$, it follows that an element of Γ_d is opposite u at d if and only if it is opposite v at d . Thus if g, t are vertices such that (u, d, w, g, t) is a 4-path, then (u, d, w, g, t) is a root if and only if (v, d, w, g, t) is a root. By 4.5(i), therefore, $u^{\text{op}} = v^{\text{op}}$. Hence by 4.12, there exists $f \in \Gamma_u$ such that (f, u, d, v, b) is a weed. In particular, $\text{dist}(f, b) = 4$, so $f \neq a$. By 4.6(ii), we have $\Gamma_2(f) = \Gamma_2(b)$. By (4.14), therefore, $\Gamma_x \subset \Gamma_2(f)$. There thus exist s, h such that (x, a, u, f, s, h, x) is a closed 6-path. The path (x, a, u) is contained in this path and in the root (x, a, u, c, w) . By 3.4, this is impossible. With this contradiction, we conclude that, in fact, $c \neq d$, as claimed. Thus (c, w, d, v, b) is a root. By 4.5(i), therefore, (c, u, a, z, b) is also a root. Hence (w, c, u, a, z) is a root and thus $w \in z^{\text{op}}$. Therefore $x^{\text{op}} \subset z^{\text{op}}$.

Suppose, conversely, that $w \in z^{\text{op}}$. By 4.5(i), there exist roots (z, a, u, c, w) and (z, b, v, d, w) from z to w , one containing a and the other containing b . By 3.2, $u \neq v$ and hence by 3.4, $c \neq d$. Therefore (c, w, d, v, b) is a root. Thus c and b are opposite and hence there is a root (b, y, e, s, c) from b to c that contains y . Therefore there exists $t \in \Gamma_c \setminus \{s\}$ such that (t, c, s, e, y) is a root from t to y . Thus $t \in y^{\text{op}} = x^{\text{op}}$. By 4.5(i), therefore, (t, c, u, a, x) is a root. Thus (u, a, x) and (u, c, w) are both straight. It follows that (x, a, u, c, w) is a root and thus $w \in x^{\text{op}}$. We conclude that $x^{\text{op}} = z^{\text{op}}$. Therefore $x^{\text{op}} = y^{\text{op}}$ implies $x \approx y$. The converse holds by 4.10. \square

Proposition 4.15. Γ is flat if and only if Γ contains no weeds.

Proof. If $\alpha = (a, x, b, y, c)$ is a weed in Γ , then $x^{\text{op}} = y^{\text{op}}$ by 4.10. Hence if Γ is flat, it contains no weeds. Suppose, instead, that Γ is not flat. If there exist distinct points a, b such that $a^{\text{op}} = b^{\text{op}}$, then by 4.6(i), $a \simeq b$ and hence there is a weed from a to b . Suppose that there exist distinct lines x, y such that $x^{\text{op}} = y^{\text{op}}$. By 4.13, $x \approx y$. Hence there exists a line z such that $x \sim z$. It follows that there is a weed containing x and z . \square

Proposition 4.16. Let x, y be elements of L such that $\text{dist}(x, y) = 2$ and $x^{\text{op}} = y^{\text{op}}$ and let $b = x \wedge y$. Then there exists a unique map φ_{xy} from Γ_x to Γ_y mapping b to itself such that for all $a \in \Gamma_x \setminus \{b\}$, $(a, x, b, y, \varphi_{xy}(a))$ is a weed.

Proof. Let $a \in \Gamma_x \setminus \{b\}$. By 4.12, there exists c such that (a, x, b, y, c) is a weed. If $c' \in \Gamma_y \setminus \{b, c\}$, then $c' \in \Gamma_2(c) = \Gamma_2(a)$ by 4.6(ii) and hence (a, x, b, y, c') is not a weed. \square

Remark 4.17. Let x, y be as in 4.16. Applying 4.16 to x, y and to y, x , we obtain maps φ_{xy} and φ_{yx} . These two maps are inverses of each other and hence both are bijections.

Remark 4.18. By 4.10, the map φ_{xy} exists for all $x, y \in L$ such that $x \sim y$.

Proposition 4.19. Let $a \in P$. Then the restriction of the reflexive closure of \sim to Γ_a is an equivalence relation.

Proof. Suppose that $u \sim v$, $v \sim w$ and $w \neq u$ for $u, v, w \in \Gamma_a$. By 4.18, there exist $b \in \Gamma_u$, $c \in \Gamma_v$ and $d \in \Gamma_w$ such that $b \simeq c$ and $c \simeq d$. By 4.7, it follows that $b \simeq d$. By 3.2, $b \neq d$. Hence by 4.5(ii), (b, u, a, w, d) is a weed. Hence $u \sim w$. \square

Proposition 4.20. Let $x, y \in \Gamma_a$ for some $a \in P$ and suppose that $\text{dist}(x, y) = 2$. Then $x \sim y$ if and only if $x \approx y$.

Proof. Let $u_0, u_1, u_2 \in L$ and suppose that $u_0 \in \Gamma_a$, $u_0 \sim u_1$, $u_1 \sim u_2$ and $u_2 \neq u_0$. By 4.18, we can choose a weed (a, u_0, b, u_1, c) starting at a and containing u_0 and u_1 .

We claim that the following hold:

- (i) If $u_2 \in \Gamma_a$, then $u_0 \sim u_2$.
- (ii) If $u_2 \notin \Gamma_a$, then there exists $u'_2 \in \Gamma_a$ such that $u'_2 \sim u_2$ and either $u'_2 \sim u_0$ or $u'_2 = u_0$.

Choose $h \in \Gamma_{u_0} \setminus \{a, b\}$. Let $i = \varphi_{u_0 u_1}(h)$ and $j = \varphi_{u_1 u_2}(i)$. Then $h \simeq i$ and $i \simeq j$ so by 4.4, $h \simeq j$. Since $\text{dist}(a, h) = 2$, we have $a \not\sim h$ and thus $j \neq a$. We also have $j \in \Gamma_{u_2}$. Suppose now that $u_2 \in \Gamma_a$. Since $u_0 \neq u_2$, (h, u_0, a, u_2, j) is a 4-path and therefore a weed by 4.5(ii). Thus (i) holds. Suppose, instead, that $u_2 \notin \Gamma_a$. In this case, we let $g = \varphi_{u_1 u_2}(c)$. Then $g \simeq c$. Since $a \simeq c$, it follows by 4.7 that $a \simeq g$. By 4.5(ii), therefore, there exists a weed (a, u'_2, f, u_2, g) from a to g containing u_2 . Thus $u'_2 \sim u_2$. Let $k = \varphi_{u_2 u'_2}(j)$. Then $j \simeq k$ and hence $h \simeq k$. Since $a \not\sim h$, we have $k \neq a$. We also have $k \in \Gamma_{u'_2}$. Hence if $u_0 \neq u'_2$, then (h, u_0, a, u'_2, k) is a 4-path and thus a weed, again by 4.5(ii), and therefore $u_0 \sim u'_2$. Thus (ii) holds.

Let Ω be the set of sequences u_0, \dots, u_m of elements of L for arbitrary m such that $u_i \sim u_{i-1}$ for all $i \in [1, m]$ and $u_0, u_m \in \Gamma_a$ but neither $u_0 = u_m$ nor $u_0 \sim u_m$ holds. Suppose that Ω is non-empty. Let u_0, \dots, u_m be a sequence in Ω of minimal length. Then $m \geq 2$. Suppose $u_2 \in \Gamma_a$. The sequence u_2, \dots, u_m is too short to be

in Ω . Hence $u_2 \sim u_m$ or $u_2 = u_m$. By (i) and 4.19, it follows that $u_0 \sim u_m$. This contradicts, however, our choice of u_0, \dots, u_m . Suppose instead that $u_2 \notin \Gamma_a$ and let u'_2 be as in (ii). Thus $u'_2 \sim u_2$, but again the sequence $u'_2, u_2, u_3, \dots, u_m$ is too short to be in Ω . Hence $u'_2 \sim u_m$ or $u'_2 = u_m$. Since either $u_0 = u'_2$ or $u_0 \sim u'_2$, we conclude that $u_0 = u_m$ or $u_0 \sim u_m$ by one more application of 4.19. Again this contradicts our choice of u_0, \dots, u_m . Thus Ω is empty. \square

Notation 4.21. Let \bar{P} be the set of equivalence classes of the equivalence relation \simeq on P and let \bar{L} be the set of equivalence classes of the equivalence relation \approx on L . Let $\bar{\Gamma}$ be the bipartite graph whose vertex set is the disjoint union of \bar{P} and \bar{L} , where $[a] \in \bar{P}$ is adjacent to $[x] \in \bar{L}$ whenever there exists an element of $[a]$ adjacent in Γ to an element of $[x]$. We declare that two elements of \bar{P} adjacent in $\bar{\Gamma}$ to a vertex $[x]$ in \bar{L} are *opposite at $[x]$* if they are distinct and we declare that two elements $[x]$ and $[y]$ of \bar{L} adjacent in $\bar{\Gamma}$ to a vertex $[a]$ in \bar{P} are *opposite at $[a]$* if there is point b in $[a]$ such that all the lines in $\Gamma_b \cap [x]$ are all opposite all the lines in $\Gamma_b \cap [y]$. We can identify $\bar{\Gamma}$ with Γ in the case that the equivalence classes $[x]$ for $x \in L$ and $[a]$ for $a \in P$ are all singletons,

Proposition 4.22. Γ is flat if and only if $\Gamma = \bar{\Gamma}$.

Proof. We have $[x] \sim [y]$ for distinct $x, y \in L$ if and only if there is a weed containing x and y and $[a] \simeq [b]$ for distinct $a, b \in P$ if and only if there is a weed containing a and b . Hence Γ contains no weeds if and only if the equivalence classes comprising the vertex set of $\bar{\Gamma}$ are all singletons. Hence Γ contains no weeds if and only if $\Gamma = \bar{\Gamma}$. The claim holds, therefore, by 4.15. \square

Theorem 4.23. Let $\bar{\Gamma}$ be the graph endowed with local opposition relations defined in 4.21. Then $\bar{\Gamma}$ is a flat green Veldkamp quadrangle.

Proof. We will prove 4.23 in a series of steps.

Proposition 4.24. Let $[a] \in \bar{P}$ and $[x] \in \bar{L}$ be adjacent in $\bar{\Gamma}$. Then $\Gamma_b \cap [x] \neq \emptyset$ for each $b \in [a]$ and $\Gamma_y \cap [a] \neq \emptyset$ for each $y \in [x]$.

Proof. Since $[a]$ and $[x]$ are adjacent in $\bar{\Gamma}$, we can assume that $x \in \Gamma_a$. Let $b \in [a] \setminus \{a\}$. Then $a \simeq b$ but $a \neq b$. By 4.5(ii), there exists a weed (a, x, c, y, b) from a to b that contains x . Then $y \in \Gamma_b \cap [x]$.

Now let $z \in [x] \setminus \{x\}$. We want to show that $\Gamma_z \cap [a] \neq \emptyset$. It will suffice to assume that $z \sim x$ (rather than $z \approx x$) and that $z \notin \Gamma_z$. By 4.10 and 4.12, there is a weed starting at a and containing x and z . Thus $e \in \Gamma_z \cap [a]$, where e is the last vertex of this weed. \square

Proposition 4.25. Let $a, b \in P$, $x, y \in \Gamma_a$ and $u, v \in \Gamma_b$. Suppose that x and y are opposite at a , $[a] = [b]$, $[x] = [u]$ and $[y] = [v]$. Then u and v are opposite at b .

Proof. Suppose that $a = b$. By 4.20, we have $x \sim u$ or $x = u$ and $y \sim v$ or $y = v$. By 4.9, therefore, u and v are opposite at a .

Suppose that $a \neq b$. By 4.5(ii), there exist weeds $\alpha = (a, x, d, w, b)$ and $\beta = (a, y, e, z, b)$ from a to b passing through x and through y . Then $[u] = [x] = [w]$ and $[v] = [y] = [z]$. Since x and y are opposite at a , (d, x, a, y, e) is a root. By 4.5(i), therefore, (d, w, b, z, e) is a root. Hence w and z are opposite at b . By the conclusion of the previous paragraph (with b in place of a), it follows that u and v are opposite at b . \square

Proposition 4.26. *Let π be the natural map from Γ onto $\bar{\Gamma}$ sending a to $[a]$ for all $a \in P$ and x to $[x]$ for all $x \in L$. Then the following hold:*

- (i) π maps edges of Γ to edges of $\bar{\Gamma}$.
- (ii) For each vertex u of Γ , paths in $\bar{\Gamma}$ starting at $[u]$ lift via π to paths in Γ starting at u .
- (iii) Let γ be a path in Γ . Then γ is straight if and only if $\pi(\gamma)$ is a straight path in $\bar{\Gamma}$.

Proof. The assertions follow from 4.21, 4.24 and 4.25. We only need to observe that if (a, x, b) is a 2-path with $x \in L$, then $\text{dist}(a, b) = 2$ and therefore $[a] \neq [b]$ and if (x, a, y) is a straight 2-path with $a \in P$, then there is no weed containing x and y and hence $[x] \neq [y]$ by 4.20. \square

Proposition 4.27. $\bar{\Gamma}$ has no circuits of length 4.

Proof. Suppose that $\beta := ([a], [x], [b], [y], [c])$ is a 4-path in $\bar{\Gamma}$ such that $[a] = [c]$. By 4.26(ii), we can assume that $\alpha := (a, x, b, y, c)$ is a 4-path in Γ . By 3.2, $a \neq c$. By 4.5(ii), therefore, α is a weed. Therefore $[x] = [y]$. This contradicts the assumption that β is a path. \square

Proposition 4.28. *Let $\alpha = ([a], [x], [b], [y])$ be a straight 3-path in $\bar{\Gamma}$ with $a \in P$. Then α is the unique straight path of length at most 3 in $\bar{\Gamma}$ from $[a]$ to $[y]$.*

Proof. By 4.26(ii)-(iii), we can assume that (a, x, b, y) is a straight path in Γ . By 4.27, $[a]$ and $[y]$ are not adjacent. Since $\bar{\Gamma}$ is bipartite, it follows that the distance from $[a]$ to $[y]$ in $\bar{\Gamma}$ is 3. Let $\gamma = ([a], [z], [d], [y])$ be an arbitrary straight 3-path from $[a]$ to $[y]$. It will suffice to show that $\gamma = \alpha$.

By 4.26(ii)-(iii), we can assume that (e, z, d, y) is a straight 3-path for some $e \in \Gamma_z \cap [a]$. By 4.6(ii), $\Gamma_2(e) = \Gamma_2(a)$. Hence $b \in \Gamma_2(e)$, so by 3.5, b and e are not opposite. Therefore (e, z, d, y, b) is not a root. Hence $b = d$. Thus $([a], [z], [d])$ and $([a], [x], [b])$ are both paths in $\bar{\Gamma}$ from $[a]$ to $[d]$. By 4.27, it follows that $[z] = [x]$. Therefore $\gamma = \alpha$. \square

We can now show that $\bar{\Gamma}$ satisfies the axioms of a Veldkamp quadrangle. Let $u \in P \cup L$ and let $[v]$ and $[w]$ be two vertices adjacent to $[u]$ in $\bar{\Gamma}$. By 4.26(ii), we can assume that $v, w \in \Gamma_u$. Since Γ is 2-plump, there exists $z \in \Gamma_u$ opposite u and v at u . By 4.26(iii), $[z]$ is opposite $[u]$ and $[v]$ at $[u]$. Thus $\bar{\Gamma}$ is 2-plump.

Since Γ is connected and every vertex of $\bar{\Gamma}$ is in the image of π , 4.26(i) implies that $\bar{\Gamma}$ is connected. It is bipartite by construction. Thus $\bar{\Gamma}$ satisfies (VP1). By 4.27 and 4.28, $\bar{\Gamma}$ satisfies (VP2) with $n = 4$. By 4.26(ii)-(iii), a straight 5-path in $\bar{\Gamma}$ lifts to a straight 5-path in Γ (via π) and a straight 8-circuit in Γ maps to a straight 8-circuit in $\bar{\Gamma}$. Since Γ satisfies (VP3) with $n = 4$, it follows that $\bar{\Gamma}$ does as well. By 4.21, the local opposition relations at elements of \bar{L} are trivial. Thus $\bar{\Gamma}$ is a green Veldkamp quadrangle.

Suppose that $([a], [x], [b], [y], [c])$ is a weed in $\bar{\Gamma}$. Thus the distance between $[a]$ and $[c]$ in $\bar{\Gamma}$ is 4. By 4.26(ii)-(iii), we can assume that $\alpha := (a, x, b, y, c)$ is a path that is not straight and by 4.26(i), $\text{dist}(a, c) = 4$. Hence α is a weed, but this implies that $[a] = [c]$. With this contradiction, we conclude that $\bar{\Gamma}$ does not contain any weeds. By 4.15, it follows that $\bar{\Gamma}$ is flat. This concludes the proof of 4.23. \square

Proposition 4.29. *If Γ is k -plump for some $k \geq 3$, then $\bar{\Gamma}$ is also k -plump.*

Proof. This holds by the same argument just used to show that $\bar{\Gamma}$ 2-plump. \square

Remark 4.30. Note that the construction of $\bar{\Gamma}$ in 4.21 does not involve any choices and thus $\bar{\Gamma}$ is uniquely determined by Γ . We call it the *flat quotient* of Γ .

Proposition 4.31. *The following hold:*

- (i) π restricts to an bijection from Γ_x to $\bar{\Gamma}_{[x]}$ for all $x \in L$.
- (ii) $\pi(\Gamma_a) = \bar{\Gamma}_{[a]}$ for all $a \in P$, but the restriction of π to Γ_a is injective for all $a \in P$ if and only if Γ is flat.

Proof. By 4.21, $\pi(\Gamma_x) \subset \bar{\Gamma}_{[x]}$ and $\pi(\Gamma_a) \subset \bar{\Gamma}_{[a]}$ for all $x \in L$ and all $a \in P$. By 4.26(ii), both of these inclusions are, in fact, equalities and by 4.26(iii), therefore, (i) holds. Let $a \in P$ and suppose that $[y] = [z]$ for some $y, z \in \Gamma_a$. By 4.20, either $y \sim z$ or $y = z$. It follows that the restriction of π to $\Gamma_{[a]}$ is injective if and only if there no weeds whose middle point is a . Hence the restriction of π to $\Gamma_{[a]}$ is injective for all $a \in P$ if and only if Γ contains no weeds. Thus (ii) holds by 4.15. \square

5. POLAR SPACES

In this section we describe the connection between polar spaces and Veldkamp quadrangles. The main results are 5.18 and 5.22. The proofs require a few basic properties of polar spaces for which we will cite results from [1, Chapter 7].

Definition 5.1. A *line space* is a pair $S = (P, L)$ consisting of a set P of points and a subset L of 2^P such that $|x| \geq 2$ for all $x \in L$. The elements of L are called *lines*. A line x is *thick* if $|x| \geq 3$ and S is called *thick* if all its lines are thick. Two points are called *collinear* if there is an element of L that contains them both

Notation 5.2. Let $S = (P, L)$ be a line space. The associated biparite graph $\Gamma = \Gamma_S$ is the graph with vertex set $P \cup L$ and edge set consisting of all pairs $\{a, x\}$ such that $a \in P$, $x \in L$ and $a \in x$.

Notation 5.3. Let $S = (P, L)$ be a line space. For each $a \in L$, we denote by a^\perp the set of all points collinear with a (including a itself) and we set

$$X^\perp = \bigcap_{a \in X} a^\perp$$

for all subsets X of P . A *subspace* is a subset X of P such that for all $x \in L$, either $x \subset X$ or $|x \cap X| \leq 1$. We will often identify a subspace X with the line space

$$S_X := (X, \{x \in L \mid x \subset X\}).$$

For all subsets X of P , let $\langle X \rangle$ denote the intersection of all the subspaces that contain X . This intersection is itself a subspace. It is called the subspace *generated by X* .

Notation 5.4. A *linear space* is a line space such that for every two points there is a unique line containing them both. A *partially linear space* is a line space such that for every two points there is at most one line containing them both. If a, b are distinct collinear points in a partially linear space, then the unique line containing them both is denoted by ab .

Definition 5.5. A *projective plane* is a linear space $S = (P, L)$ such that $|L| \geq 2$ and $x \cap y \neq \emptyset$ for all $x, y \in P$. A *projective space* is a linear space such that every subspace generated by the union of two lines with non-empty intersection is a projective plane.

Definition 5.6. Let $S = (P, L)$ be a projective space. The *dimension* of S is the maximal length k of a properly ascending chain of subspaces X_i such that

$$\emptyset \neq X_1 \subset X_2 \subset \cdots \subset X_k = P.$$

It is denoted by $\dim(S)$. Thus, in particular, lines are subspaces of dimension 1 and $\dim(S) = \infty$ if there no bound on the length of such chains. A non-singular subspace of dimension 2 is called a non-singular *plane*.

Definition 5.7. A *polar space* is a line space $S = (P, L)$ such that the following holds:

(BS) For all $a \in P$ and $x \in L$, either $|a^\perp \cap x| = 1$ or $x \subset a^\perp$.

Note that every subspace of a polar space is itself a polar space.

Definition 5.8. Let $S = (P, L)$ be a line space. A subspace X of S is called *singular* if every two points of X are collinear. Thus lines are, in particular, singular subspaces.

Definition 5.9. A polar space $S = (P, L)$ is *non-degenerate* if $P \neq a^\perp$ for all $a \in P$.

Proposition 5.10. *Let S be a non-degenerate polar space. Then the following hold:*

- (i) S is a partially linear space.
- (ii) Singular subspaces are projective spaces.

Proof. (i) holds by [1, Thm. 7.4.11] and (ii) by [1, Thm. 7.4.13(iv)]. □

Definition 5.11. In light of 5.10(ii), we can define the *rank* of a non-degenerate polar space $S = (P, L)$ to be one more than the supremum of the set

$$\{\dim(E) \mid E \text{ is a non-singular subspace}\}.$$

The rank is denoted by $\text{rk}(S)$. Thus $\text{rk}(S) = 1$ if and only if $L = \emptyset$ but $P \neq \emptyset$, and $\text{rk}(S) = 2$ if and only if $L \neq \emptyset$ but lines are maximal singular subspaces.

Remark 5.12. A polar space is called *thick* in this paper (see 5.1) if all its lines are thick. We note that the definition of a thick polar space in [11, 7.3] is slightly different.

Proposition 5.13. *Let $S = (P, L)$ be a thick non-degenerate polar space and let $a, b \in P$. Then there exists a point that is collinear with neither a nor b .*

Proof. Suppose first that a and b are collinear and let $x = ab$. Since S is thick, we can choose $c \in x$ distinct from a and b . By [1, 7.4.8(iii)], there exists $d \in c^\perp \setminus x^\perp$. This point d is collinear with neither a nor b . Now suppose that a and b are not collinear. Let x be a line containing b and let c be a point on x distinct from b . By (BS), a is collinear with exactly one point on x . Since S is thick, we can assume, therefore, that a and c are not collinear. By another application of [1, 7.4.8(iii)], there exists $d \in c^\perp \setminus x^\perp$. Let $y = cd$. By (BS), there is a unique point e in $y \cap a^\perp$.

Since S is thick, we can choose f on y distinct from c and e . Thus f is not collinear with a , nor, by the choice of b , is it collinear with b . \square

Proposition 5.14. *Let $S = (P, L)$ be a thick non-degenerate polar space and let a and b be two non-collinear points. Then $a^\perp \cap b^\perp$ is a thick non-degenerate polar space.*

Proof. Let $S_1 = a^\perp \cap b^\perp$. If x is a line containing two points d, e of S_1 , then by (BS), $x \subset S_1$. Thus S_1 is a thick subspace. In particular, S_1 is a polar space. By [1, 7.4.8(ii)], it is non-degenerate. \square

Proposition 5.15. *Let $S = (P, L)$ be a thick non-degenerate polar space and let $x \in L$. Then there exists $y \in L$ such that $x^\perp \cap y^\perp$ is a thick non-degenerate polar space and $x \cap y^\perp = \emptyset$.*

Proof. Let $a \in x$. Since S is non-degenerate, we can choose $e \in P \setminus a^\perp$ and by (BS), we can choose $b \in x$ collinear with e . Let $S_1 = a^\perp \cap e^\perp$. By 5.14, S_1 is a thick non-degenerate polar space. In particular, we can choose $d \in S_1$ not collinear with b . Let $y = de$ and let $S_2 = b^\perp \cap d^\perp \cap S_1$. Thus

$$S_2 = a^\perp \cap b^\perp \cap d^\perp \cap e^\perp.$$

By (BS), $a^\perp \cap b^\perp = x^\perp$ and $d^\perp \cap e^\perp = y^\perp$. Hence $S_2 = x^\perp \cap y^\perp$. By a second application of 5.14, S_2 is a thick non-degenerate polar space.

Now suppose that $c \in x \cap y^\perp$. Then $b, c \in e^\perp$ as well as $c \in d^\perp$, so $c \neq b$. By (BS), therefore, $a \in e^\perp$. This contradicts the choice of e . We conclude that $x \cap y^\perp = \emptyset$. \square

Proposition 5.16. *Let $x, y \in L$ be as in 5.15, let $S_2 = x^\perp \cap y^\perp$ and let E be a singular plane containing x . Then $|E \cap S_2| = 1$.*

Proof. By 5.5 and 5.10(ii), E is a projective plane. Thus every two lines contained in E intersect in a point. In particular, $E \subset x^\perp$. Let $d \in y$. By (BS), d^\perp contains a point on each line of E . Hence $|d^\perp \cap E| \geq 2$. It follows that $E \cap d^\perp$ contains a line. Now choose a second point e in y . Then $E \cap e^\perp$ also contains a line. Therefore $E \cap d^\perp \cap e^\perp \neq \emptyset$. By (BS), we have $d^\perp \cap e^\perp = y^\perp$. Thus $E \cap S_2 = E \cap y^\perp \neq \emptyset$.

Suppose that a, b are two points in $E \cap S_2$. Then a and b are collinear. Let $z = ab$. Then $z \subset E \cap S_2$. It follows that z and x intersect in a point that lies in $x \cap S_2 \subset x \cap y^\perp$. By 5.15, however, $x \cap y^\perp = \emptyset$. With this contradiction, we conclude that $|E \cap S_2| \leq 1$. \square

Proposition 5.17. *Let $x \in L$ and let M be the set of singular planes containing x . Then for all $E, F \in M$ there exists an element G of M such that both $\langle E \cup G \rangle$ and $\langle F \cup G \rangle$ are non-singular.*

Proof. Let $y \in L$ be as in 5.15 and let $S_2 = x^\perp \cap y^\perp$. Thus S_2 is a thick non-degenerate polar space and $x \cap y^\perp = \emptyset$. Choose $E, F \in M$. By 5.16, there exist points a, b such that $E \cap S_2 = \{a\}$ and $F \cap S_2 = \{b\}$. By 5.13, there exists $c \in S_2$ that is collinear with neither a nor b . Let $G = \langle x \cup c \rangle$. Since $x \cap y^\perp = \emptyset$, we have $c \notin x$. On the other hand, $c \in x^\perp$ so by [1, 7.4.7(v)], G is singular. Thus $G \in M$. By the choice of c , neither $\langle E \cup G \rangle$ nor $\langle F \cup G \rangle$ is singular. \square

Proposition 5.18. *Let $S = (P, L)$ be a thick, non-degenerate polar space such that $L \neq \emptyset$ and let $\Gamma = \Gamma_S$ be the associated bipartite graph as defined in 5.2. Let Γ be endowed with the following local opposition relations:*

- (a) *For each $a \in P$ and all $x, y \in \Gamma_a$, x is opposite y at a if the subspace $\langle x \cup y \rangle$ is non-singular.*
- (b) *For each $x \in L$ and all $a, b \in \Gamma_x$, a is opposite b at x if $a \neq b$.*

Then Γ is a flat green Veldkamp quadrangle. It is a generalized quadrangle if and only if $\text{rk}(S) = 2$.

Proof. Let $a \in P$ and $x, y \in \Gamma_a$. Then 5.5 and 5.10(ii), imply that if x and y are not opposite at a , then every point in $x \setminus \{a\}$ is collinear with every point in $y \setminus \{a\}$. By [1, 7.4.7(v)], the converse holds. Thus x and y are opposite at a if and only if some point in $x \setminus \{a\}$ is not collinear with some point in $y \setminus \{a\}$. By (BS), it follows that if x and y are opposite at a , then no point in $x \setminus \{a\}$ is collinear with any point in $y \setminus \{a\}$.

We claim that Γ is 2-plump. Let $a \in P$. Since S is non-degenerate, we can choose a point b not collinear with a . Let $x, y \in \Gamma_a$. By (BS), there is a unique point d on x collinear with b and a unique point e on y collinear with b . By 5.13 and 5.14, there exists a point f in $a^\perp \cap b^\perp$ that is collinear with neither d nor e . Let $z = af$. Then $z \in \Gamma_a$ is locally opposite x and y at a . Since S is thick, $|\Gamma_x| \geq 3$ also for $x \in L$. Thus Γ is 2-plump, as claimed.

By (BS), $\text{dist}(a, x) \leq 3$ for all $a \in P$ and all $x \in L$. Thus

$$(5.19) \quad \text{diam}(\Gamma) \leq 4.$$

In particular, Γ is connected. Thus it satisfies (VP1). It follows from 5.10(i) that Γ does not contain any 4-circuits. Let (a, x, b, y) is a 3-path with $a \in P$. Suppose that $\langle x \cup y \rangle$ is non-singular. Then every point in x is collinear with b but no other point in y . Hence $\text{dist}(a, c) = 4$ for all $c \in y \setminus \{b\}$. In other words, (a, x, b, y) does not lie on a 6-circuit. Thus Γ satisfies (VP2). Now suppose that $\langle x \cup y \rangle$ is singular. Then a is collinear with every point on y . Hence $\Gamma_y \subset \Gamma_2(a)$. Thus we conclude that

$$(5.20) \quad \Gamma_w \subset \Gamma_2(f)$$

for all $w \in L$ and all $f \in P$ such that there is a non-straight 3-path from w to f and hence

$$(5.21) \quad e^{\text{op}} = \Gamma_4(e)$$

for all points e .

Now suppose that (a, x, b, y, c, z) be a straight 5-path. Then $\text{dist}(a, c) = 4$ by (5.21), so by (5.19), we have $\text{dist}(a, z) = 3$. Hence there exist d, u such that $(a, x, b, y, c, z, d, u, a)$ is a closed 8-path. The subspace $\langle z \cup u \rangle$ is non-singular since otherwise $c \in \Gamma_z \subset \Gamma_2(a)$ by (5.20). Since the subspace $\langle y \cup z \rangle$ is also non-singular, we have $\text{dist}(b, d) = 4$ by (5.21). Thus $\langle x \cup u \rangle$ is non-singular since otherwise $d \in \Gamma_u \subset \Gamma_2(b)$ by (5.20). We conclude that Γ satisfies (VP3). Thus Γ is a green Veldkamp quadrangle. Suppose that $a^{\text{op}} = b^{\text{op}}$ for $a, b \in P$. Since b is not opposite itself, it follows by (5.21) that $\text{dist}(a, b) \neq 4$. Suppose that $\text{dist}(a, b) = 2$. Since Γ is green, there exists a root (a, w, b, v, c) starting at a and containing b , but then the last vertex c of this root would be in $a^{\text{op}} \setminus b^{\text{op}}$. By (5.19), therefore, $a = b$. By 4.6(i), it follows that Γ contains no weeds. By 4.15, we conclude that Γ is flat.

If $\text{rk}(S) = 2$, then $\langle x \cup y \rangle$ is non-singular for all $x, y \in L$, hence every path in Γ is straight and thus Γ is a generalized 4-gon. Suppose, conversely, that every path in Γ is straight. Then $\langle x \cup y \rangle$ is non-singular for all $x, y \in L$ such that $x \cap y \neq \emptyset$. By 5.5 and 5.10(ii), $x \cap y \neq \emptyset$ for every two lines contained in a singular plane. Hence there are no singular planes. In other words, $\text{rk}(S) = 2$. \square

Theorem 5.22. *Let $\Gamma = (V, E)$ be a flat green Veldkamp quadrangle with point set P and line set L and let $L_1 = \{\Gamma_x \mid x \in L\}$. Then the following hold:*

- (i) $S_\Gamma := (P, L_1)$ is a thick non-degenerate polar space with $L_1 \neq \emptyset$.
- (ii) The construction $\Gamma \rightsquigarrow S_\Gamma$ and the construction $S \rightsquigarrow \Gamma_S$ in 5.18 are inverses of each other.
- (iii) $\text{rk}(S_\Gamma) = 2$ if and only if Γ is a generalized quadrangle.

Proof. Let $a \in P$ and let $x \in L$. By 3.6, $\text{dist}(a, x) \leq 3$. Suppose that $\text{dist}(a, x) = 3$ but $\Gamma_x \not\subset \Gamma_2(a)$. Then there exists a 4-path $\alpha = (a, u, b, x, c)$ such that $\text{dist}(a, c) = 4$. By 4.15, Γ contains no weeds. It follows that α is a root. In particular, (u, b, x) is a straight. Since Γ is green, (a, u, b, x, d) is a root for all $d \in \Gamma_x \setminus \{b\}$. Thus $\Gamma_x \cap \Gamma_2(a) = \{b\}$. We conclude that either $a \in \Gamma_x$ or $\Gamma_x \subset \Gamma_2(a)$ or $|\Gamma_x \cap \Gamma_2(a)| = 1$. Hence S_Γ is a polar space. Since Γ is 2-plump, S_Γ is thick. By 3.5, S_Γ is non-degenerate. Thus (i) holds. The assertion (ii) holds by definition and (iii) follows from (ii) and 5.18. \square

In the next result, we produce examples of green Veldkamp quadrangles that are not flat.

Proposition 5.23. *Let $S = (P, L)$ be a thick non-degenerate polar space, suppose that $\text{rk}(S) \geq 3$ and let $\Gamma = \Gamma_S$ be as in 5.18. Let $d \in P$, let $P_1 = d^\perp \setminus \{d\}$ and let $L_1 = \{x \in L \mid x \subset P_1\}$. Let Ω be the subgraph of Γ spanned by $P_1 \cup L_1$ endowed with the natural restriction of the local opposition relations of Γ to the neighborhoods of Ω . Then the following hold:*

- (i) Ω is a green Veldkamp quadrangle that is not flat.
- (ii) Let $a, b \in P_1$. Then $a^{\text{op}} = b^{\text{op}}$ in Ω if and only if a, b and d are collinear.
- (iii) Let $e \in d^{\text{op}}$, let $P_2 = e^\perp \cap P_1$ and let L_2 be the set of lines contained in P_2 . Then $L_2 \neq \emptyset$, $S_2 := (P_2, L_2)$ is a thick non-degenerate polar space and Γ_{S_2} is, up to isomorphism, the flat quotient of Ω .

Proof. We first want to show that Ω is 2-plump. Let $a \in P_1$, let $u = ad$ and choose $x, y \in \Omega_a$. Since $\Gamma_x \subset P_1$, it follows from (5.21) that the path (u, a, x) is not straight. Hence $E := \langle u \cup x \rangle$ is a singular plane containing u . Similarly, $F := \langle u \cup y \rangle$ is a singular plane containing u . By 5.17, there exists a singular plane G containing u such that neither $\langle E \cup G \rangle$ nor $\langle F \cup G \rangle$ is singular. Let z be a line in G that intersects u in the point a . Since G is singular, the path (u, a, z) is not straight. By (5.20), we have $\Gamma_z \subset P_1$. In other words, $z \in L_1$. If (x, a, z) were not straight, then $\langle x \cup z \rangle$ would be singular, but by [1, 7.4.7(v)], this would imply that $\langle E \cup G \rangle$ is singular. Hence (x, a, z) is straight. Similarly, (y, a, z) is straight. Thus local opposition at a is 2-plump. Since Γ is thick, so is Ω . Therefore Ω is 2-plump.

Again let $a \in P_1$ and $u = ad$. This time let x be an arbitrary element of $L_1 \setminus \Omega_a$. By 3.6, there are 3-paths in Γ from a to x . Let (a, y, b, x) be one of them. Suppose that $y \neq u$. Since $x \in L_1$, we have $b \in d^\perp$. By 3.5, therefore, d and b are not opposite, so (d, u, a, y, b) is not a root. Hence (d, u, a, y) is not straight and thus $y \in L_1$ by (5.20). Thus the distance from a to x in Ω is 3.

Suppose now that $y = u$. Since $x \in L_1$, the path (d, u, b, x) is not straight by (5.21). Hence (a, u, b, x) is not straight. Therefore $\Gamma_x \subset \Gamma_2(a)$ by (5.20). Let $c \in \Gamma_x \setminus \{b\}$. Then there exists a 3-path (a, v, c, x) in Ω from a to x containing c and $v \neq u$. By 3.4, (u, a, v) is not straight. By (5.20), $v \in L_1$. Therefore the distance from a to x in Ω is 3 also in this case. By 3.2, however, (a, u, b) is the only 2-path in Γ from a to b , so the distance from a to b in Ω is not 2. Since $b \in \Omega_x$, we conclude that the distance from a to b in Ω is 4 and that (a, v, c, x, b) is a non-straight 4-path from a to b in Ω .

We have thus shown that

$$(5.24) \quad \text{diam}(\Omega) = 4.$$

In particular, Ω is connected. Therefore Ω satisfies (VP1). Since Γ satisfies (VP2), so does Ω .

Now suppose that (a, x, b, y, c, z) is a straight 5-path in Ω . By (VP2) and (VP3), there is a unique straight 3-path α from a to z in Γ . By (5.24), the distance from a to z in Ω is also 3. By 3.5, $\text{dist}(a, c) = 4$. Thus $\Gamma_z \not\subset \Gamma_2(a)$. By (5.21), it follows that α lies in Ω . Hence Ω satisfies (VP3). Thus Ω is a green Veldkamp quadrangle.

Next let $a, b \in P_1$, let $u = da$ and suppose that $b \in \Gamma_u \setminus \{a\}$. By 3.2, there is no $x \in L_1$ such that Γ_x contains a and b . By (5.24), it follows that the distance from a to b in Ω is 4. By 3.4, there is no root from a to b . Hence every 4-path from a to b in Ω is a weed in Ω . By 4.6(i), we conclude that $a^{\text{op}} = b^{\text{op}}$ in Ω .

Now let $a, b \in P_1$, let $u = ad$ and suppose that $a \neq b$ and $a^{\text{op}} = b^{\text{op}}$ in Ω . By 4.5(ii) and 4.6(i), the distance from a to b in Ω is 4 and every 4-path from a to b in Ω is a weed in Ω . Let (a, y, c, x, b) be a 4-path from a to b in Ω . Thus (a, y, c, x) is not straight and by 4.12, b is the unique element of Γ_x that is not at distance 2 from a in Ω . By (5.21), $b \in \Gamma_x \subset \Gamma_2(a)$. Hence $b \in \Gamma_u$. We conclude that (i) and (ii) hold.

Finally, we let e and $S_2 = (P_2, L_2)$ be as in (iii). By 5.14, S_2 is a thick non-degenerate polar space. Let $\bar{\Omega}$ denote the flat quotient of Ω and let $S_{\bar{\Omega}}$ be as in 5.22(i). Thus by 4.21, the point set of $S_{\bar{\Omega}}$ is the set of equivalence classes $[a]$ for all $a \in P_1$ as defined in 4.7 (with Ω in place of Γ). By (BS), for each $w \in \Gamma_d$, there exists a unique $a \in \Gamma_w \cap P_2$. By (ii), therefore, there exists a canonical bijection ψ from the point set of $S_{\bar{\Omega}}$ to P_2 mapping the equivalence class $[a]$ to the unique vertex in $\Gamma_{ad} \cap P_2$ for all $a \in P_1$. Let $a, b \in P_1$ and suppose that the points $[a], [b]$ of $S_{\bar{\Omega}}$ are distinct. Let $u = ad$ and $v = bd$ and let $a' = \psi([a])$ and $b' = \psi([b])$. Suppose first that $[a]$ and $[b]$ are collinear in $S_{\bar{\Omega}}$. By 4.26(ii), we can assume that the distance between a and b in Ω is 2. Let $x = a \wedge b$. Since x is a vertex of Ω , we have $x \in \Gamma_3(d)$. Hence (a, u, d, v, b, x, a) is a closed 6-circuit. By 3.4, therefore, (u, d, v) is not straight. Therefore also (a', u, d, v, b') is not straight. By 5.18, Γ is flat. By 4.15, therefore, (a', u, d, v, b') is not a weed. Hence $\text{dist}(a', b') = 2$. It follows that a' and b' are collinear in S_2 . Thus ψ maps collinear points of $S_{\bar{\Omega}}$ to collinear points of S_2 . Conversely, if a' and b' are collinear in S_2 , then (since $L_2 \subset L_1$), $[a'] = [a]$ and $[b'] = [b]$ are collinear in $S_{\bar{\Omega}}$. We conclude that $[a]$ and $[b]$ are collinear in $S_{\bar{\Omega}}$ if and only if a' and b' are collinear in S_2 . By [1, 7.4.12], it follows that ψ is an isomorphism from $S_{\bar{\Omega}}$ to S_2 . By 5.22, it follows that ψ induces an isomorphism from $\bar{\Omega}$ to Γ_{S_2} . Thus (iii) holds. \square

6. TITS QUADRANGLES

In this section, we give the classification of green Veldkamp quadrangles that are not generalized quadrangles in 6.12 and 6.16. In particular, we show that these Veldkamp quadrangles are all, in fact, Tits quadrangles as defined in [3, 1.1.6 and 1.1.8].

Definition 6.1. Let Γ be a Veldkamp n -gon for some $n \geq 3$ and let

$$\alpha = (x_0, x_1, \dots, x_n)$$

be a root. We denote by U_α the pointwise stabilizer in $\text{Aut}(\Gamma)$ of the set

$$\Gamma_{x_1} \cup \Gamma_{x_2} \cup \dots \cup \Gamma_{x_{n-1}}.$$

The subgroup U_α is called the *root group* of Γ associated with α .

Definition 6.2. Let Γ be a Veldkamp n -gon for some $n \geq 3$. Then Γ is *Moufang* if for each root $\alpha = (x_0, x_1, \dots, x_n)$, the root group U_α acts transitively on the set of vertices in Γ_{x_n} that are opposite x_{n-1} at x_n .

Proposition 6.3. *Let Γ be a Veldkamp graph, let (x, y, z) be a straight 2-path, let M be a subgroup of the stabilizer of (x, y) in $\text{Aut}(\Gamma)$ and let g be an element of $\text{Aut}(\Gamma)$ mapping (y, x) to (y, z) . Suppose that M acts transitively on the set of vertices in Γ_y that are opposite x . Then $\langle M, M^g \rangle$ acts transitively on Γ_y .*

Proof. The conjugate M^g fixes z and acts transitively on the set of vertices of Γ_y opposite z . Since Γ is 2-plump, it follows that $\langle M, M^g \rangle$ acts transitively on Γ_y . \square

Remark 6.4. Let Γ be a Veldkamp n -gon for some $n \geq 3$. Then Γ is Moufang if and only if it is a Tits n -gon as defined in [3, 1.1.6 and 1.1.8].

Example 6.5. Let Λ be either a quadratic space

$$(K, L, q)$$

(Case I) or a pseudo-quadratic space

$$\Lambda = (K, K_0, \sigma, L, q)$$

as defined in [12, 11.15–11.17] (Case II). In Case II, L is a right vector space over K and the associated sesquilinear form f on L is assumed to be skew-hermitian. In Case I, we write scalars on the left, we let f denote the bilinear form associated with q and we let σ denote the trivial automorphism of K . In both cases, we assume that Λ is non-degenerate as defined in [11, 8.2.3]. Thus

$$\{a \in L \mid q(a) = f(a, a) = 0\} = \{0\}$$

in Case I and

$$\{a \in L \mid f(a, a) = 0 \text{ and } q(a) \in K_0\} = \{0\}$$

in Case II. In Case II, $L = 0$ is allowed, but in Case I, $L \neq 0$. In Case II we assume that $\text{char}(K) \neq 2$ if $K_0 = K$. We do not make any restriction on the Witt index of q in Case I or in Case II.

It is important for the reader to keep in mind that $\{t + t^\sigma \mid t \in K\} \subset K_0$ in Case II. Many of the claims we make below depend on this observation.

Let $V = K^4 \oplus L$ and let $Q: V \rightarrow K$ be given by

$$Q(t_1, t'_1, t_2, t'_2, a) = t_1^\sigma t'_1 + t_2^\sigma t'_2 + q(a)$$

for all $(t_1, t'_1, t_2, t'_2, a) \in V$. In Case I, let

$$F((s_1, s'_1, s_2, s'_2, b), (t_1, t'_1, t_2, t'_2, a)) = s_1 t'_1 + t_1 s'_1 + s_2 t'_2 + t_2 s'_2 + f(a, b)$$

and in Case II, let

$$F((s_1, s'_1, s_2, s'_2, b), (t_1, t'_1, t_2, t'_2, a)) = s_1^\sigma t'_1 - (s'_1)^\sigma t_1 + s_2^\sigma t'_2 - (s'_2)^\sigma t_2 + f(b, a)$$

for all $(s_1, s'_1, s_2, s'_2, b), (t_1, t'_1, t_2, t'_2, a) \in V$. In Case I, Q is a non-degenerate quadratic form on V and F is the bilinear form associated with Q . In Case II, Q is a non-degenerate pseudo-quadratic form and F is the skew-hermitian form associated with Q . In particular, the image of Q is defined only modulo K_0 and when we say that Q vanishes on a given subspace V_1 of V , we mean that $Q(V_1) \subset K_0$. If we are in Case I or in Case II with $K_0 \neq K$, then by [12, 11.19], the form F is uniquely determined by Q and, in particular, F is identically zero on every subspace on which Q vanishes identically. If we are in Case II with $K_0 = K$, then Q vanishes identically on V , K is commutative, $\sigma = 1$, F is alternating and $\text{char}(K) \neq 2$ (by hypothesis). We call this subcase of Case II the *symplectic* case.

Let T denote the additive group of L in Case I and in Case II, let T denote the group with underlying set

$$\{(a, t) \in L \times K \mid q(a) - t \in K_0\}$$

defined in [12, 11.24]. Thus $T = K_0$ if $L = 0$ and

$$(a, t)(b, u) = (a + b, t + u + f(b, a))$$

and

$$(a, t)^{-1} = (-a, -t^\sigma)$$

for all $(a, t), (b, u) \in T$ otherwise. Note that in the symplectic case, the condition $q(a) - t \in K_0$ is vacuous and the underlying set of T is simply the direct product $L \times K$.

For each i , let W_i be the set of subspaces of V of dimension i on which Q and F vanish identically. For $i \geq 2$, we identify each element of W_i with the set of elements of W_1 it contains. This makes (W_1, W_2) into a thick line space. If $V_1 \in W_1$ and $V_2 \in W_2$, then $\dim_K(V_2 \cap V_1^\perp) = 1$ or 2 . (Here orthogonal means orthogonal with respect to the form F .) Thus the line space (W_1, W_2) is, in fact, a polar space as defined in 5.7. Since Λ is non-degenerate, we have $V_1^\perp \neq V$. Hence the polar space (W_1, W_2) is non-degenerate as defined in 5.9. We denote it by S_Λ . For each i , the i -dimensional singular subspaces of $S = S_\Lambda$ are the subspaces in W_{i+1} . Let $\Gamma = \Gamma_S$ be the corresponding Veldkamp quadrangle as defined in 5.22.

We want to show that Γ is a Tits quadrangle and that its root group sequences are as described in [12, 16.3] in Case I and in [12, 16.2 and 16.5] in Case II, but without the assumption in either case that q is anisotropic.

For each isotropic vector u in V (i.e. for each vector u that spans an element of W_1) we will denote simply by u the element of W_1 spanned by u when it is clear that we are referring vertices of Γ . For each pair of orthogonal isotropic vectors u and v , we will denote by uv the element of W_2 they span. Next we set

$$x_1 = (1, 0, 0, 0, 0), \quad x'_1 = (0, 1, 0, 0, 0)$$

as well as

$$x_2 = (0, 0, 1, 0, 0) \text{ and } x'_2 = (0, 0, 0, 1, 0)$$

in V and we identify L with its image in V under the map $u \mapsto (0, 0, 0, 0, u)$. Let Σ be the straight 8-circuit of Γ with vertex set

$$x_1, x_1x'_2, x'_2, x'_1x'_2, x'_1, x'_1x_2, x_2, x_1x_2$$

and let

$$\alpha = (x'_2, x_1x'_2, x_1, x_1x_2, x_2) \text{ and } \gamma = (x'_1x'_2, x'_2, x_1x'_2, x_1, x_1x_2)$$

as well as

$$\beta = (x'_1, x'_1x'_2, x'_2, x_1x'_2, x_1) \text{ and } \delta = (x'_1x_2, x'_1, x'_1x'_2, x'_2, x_1x'_2).$$

Note that the roots α, \dots, δ all contain the edge $\{x_1x'_2, x'_2\}$.

Notation 6.6. Let $\varepsilon = 1$ in Case I and $\varepsilon = -1$ in Case II. Let ρ denote the linear automorphism of V that interchanges x_1 with $\varepsilon x'_2$ and x'_1 with x_2 and acts trivially on L . Let τ denote the linear automorphism of V that interchanges x_1 with $\varepsilon x'_1$, fixes x_2 and x'_2 and acts trivially on L . Both ρ and τ are isometries of both Q and F and hence both induce automorphisms of Γ . We denote these elements by ρ and τ . Both ρ and τ stabilize Σ , ρ fixes the vertices $x_1x'_2$ and x'_1x_2 , τ fixes the vertices x_2 and x'_2 and $\langle \rho, \tau \rangle$ acts transitively on the edge set of Σ . Hence every root contained in Σ is in the same $\langle \rho, \tau \rangle$ -orbit as either α or δ .

Notation 6.7. Suppose we are in Case I. For all $a, b \in L$, let $\dot{\alpha}_a$ be the linear automorphism of V that fixes x_1, x_2 and x'_2 , maps x'_1 to $x'_1 - q(a)x_1 + a$ and maps an arbitrary element $u \in L$ to $u - f(u, a)x_1$ and let $\dot{\beta}_b$ be the linear automorphism of V that fixes x_1, x'_1 and x'_2 , maps x_2 to $x_2 - q(b)x'_2 + b$ and maps an arbitrary element $u \in L$ to $u - f(u, b)x'_2$. For all $s, t \in K$, let $\dot{\gamma}_t$ be the linear automorphism of V that fixes x_1 and x'_2 , acts trivially on L , maps x'_1 to $x'_1 - tx'_2$ and x_2 to $x_2 + tx_1$ and let $\dot{\delta}_s$ be the linear automorphism of V that fixes x'_1 and x'_2 , acts trivially on L and maps x_1 to $x_1 + sx'_2$ and x_2 to $x_2 - sx'_1$.

Notation 6.8. Suppose that we are in Case II. For all $(a, t), (b, s) \in T$, let $\dot{\alpha}_{(a,t)}$ be the linear automorphism of V that fixes x_1, x_2 and x'_2 , maps x'_1 to $x'_1 + x_1t + a$ and maps an arbitrary element $u \in L$ to $u + x_1f(a, u)$ and let $\dot{\beta}_{(b,s)}$ be the linear automorphism of V that fixes x_1, x'_1 and x'_2 , maps x_2 to $x_2 - x'_2s - b$ and maps an arbitrary element $u \in L$ to $u + x'_2f(b, u)$. For all $s, t \in K$, let $\dot{\gamma}_t$ be the linear automorphism of V that fixes x_1 and x'_2 , acts trivially on L , maps x'_1 to $x'_1 + x'_2t^\sigma$ and x_2 to $x_2 - x_1t$ and let $\dot{\delta}_s$ be the linear automorphism of V that fixes x'_1 and x'_2 , acts trivially on L and maps x_1 to $x_1 - x'_2s^\sigma$ and x_2 to $x_2 - x'_1s$.

For all $v, w \in T$ and all $s, t \in K$, the maps $\dot{\alpha}_v, \dot{\beta}_w, \dot{\gamma}_t$ and $\dot{\delta}_s$ defined in 6.7 and 6.8 are all isometries of Q and F and hence induce automorphisms of Γ . We denote these automorphisms by $\alpha_v, \beta_w, \gamma_t$ and δ_s . Let

$$A = \{\alpha_v \mid v \in T\}, B = \{\beta_w \mid w \in T\}, C = \{\gamma_t \mid t \in K\} \text{ and } D = \{\delta_s \mid s \in K\}.$$

Observations 6.9. Let K^+ denote the additive group of K . With a bit of calculation, it can be checked that the following hold:

- (i) A, B, C and D are subgroups of $\text{Aut}(\Gamma)$.
- (ii) $A \subset U_\alpha, B \subset U_\beta, C \subset U_\gamma$ and $D \subset U_\delta$.
- (iii) $v \mapsto \alpha_v$ and $v \mapsto \beta_v$ are isomorphisms from T to A and from T to B and $t \mapsto \gamma_t$ and $t \mapsto \delta_t$ are isomorphisms from K^+ to C and from K^+ to D .

- (iv) A acts transitively on the set of vertices adjacent to x_2 that are opposite x_1x_2 at x_2 and on the set of vertices adjacent to x'_2 that are opposite $x_1x'_2$ at x'_2 .
- (v) B acts transitively on the set of vertices adjacent to x_1 that are opposite $x_1x'_2$ at x_1 and on the set of vertices adjacent to x'_1 that are opposite $x'_1x'_2$ at x'_1 .
- (vi) C acts transitively on $\Gamma_{x_1x_2} \setminus \{x_1\}$ and on $\Gamma_{x'_1x'_2} \setminus \{x'_2\}$.
- (vii) D acts transitively on $\Gamma_{x_1x'_2} \setminus \{x'_2\}$ and on $\Gamma_{x'_1x_2} \setminus \{x'_1\}$.

Now let M denote the group generated by ρ, τ, A, B, C and D . By 6.9(iv)–(vii), the subgroup $\langle A, B, C, D \rangle$ of M acts transitively both on the set of roots of Γ starting with $(x'_2, x_1x'_2)$ and on the set of roots of Γ starting with $(x_1x'_2, x'_2)$. By 6.3 and 6.9(iv) and (vii), the subgroup $\langle A, \tau \rangle$ acts transitively on $\Gamma_{x'_2}$ and the subgroup $\langle C, \rho \rangle$ acts transitively in $\Gamma_{x_1x'_2}$. Since x'_2 and $x_1x'_2$ are adjacent vertices and Γ is connected, it follows that M acts transitively on the set of edges of Γ . By 6.6, we conclude that every root of Γ is in the same M -orbit as either α or γ . Hence by 6.2 and 6.9(ii), (iv) and (vii), Γ is Moufang. By 6.4, therefore, Γ is a Tits quadrangle. By 6.9(ii) and [3, 1.3.25], finally, we have

$$(6.10) \quad A = U_\alpha, \quad B = U_\beta, \quad C = U_\gamma \quad \text{and} \quad D = U_\delta.$$

Definition 6.11. Let Λ, T and f be as in 6.5 (so Λ is non-degenerate and if $K_0 = K$ in Case II, then $\text{char}(K) \neq 2$) and let K^+ be as in 6.9. Let Δ be a Tits quadrangle, let $(\Sigma, i \mapsto w_i)$ be a coordinate system for Δ and let $i \mapsto U_i$ be the corresponding root group labeling as defined in [7, 2.3]. The root groups U_1, \dots, U_4 satisfy the commutator conditions in [7, 2.5]. In particular, $[U_i, U_{i+1}] = 1$ for all i .

- (i) Δ is called *orthogonal of type Λ* if $\Lambda = (K, L, q)$ is a quadratic space and, after replacing the coordinate system by its opposite (as defined in [7, 2.3]) if necessary, there exist isomorphisms $x_i: K^+ \rightarrow U_i$ for $i = 1$ and 3 and $x_i: T \rightarrow U_i$ for $i = 2$ and 4 such that

$$\begin{aligned} [x_2(a), x_4(b)^{-1}] &= x_3(f(a, b)) \quad \text{and} \\ [x_1(t), x_4(a)^{-1}] &= x_2(ta)x_3(tq(a)) \end{aligned}$$

for all $t \in K$ and all $a, b \in L$ and $[U_1, U_3] = 1$.

- (ii) Δ is called *pseudo-quadratic of type Λ* if $\Lambda = (K, K_0, \sigma, L, q)$ is a pseudo-quadratic space and, after replacing the coordinate system by its opposite if necessary, there exist isomorphisms $x_i: T \rightarrow U_i$ for $i = 1$ and 3 and $x_i: K^+ \rightarrow U_i$ for $i = 2$ and 4 such that

$$\begin{aligned} [x_1(a, t), x_3(b, s)^{-1}] &= x_2(f(a, b)), \\ [x_2(v), x_4(w)^{-1}] &= x_3(0, v^\sigma w + w^\sigma v) \quad \text{and} \\ [x_1(a, t), x_4(v)^{-1}] &= x_2(tv)x_3(av, v^\sigma tv) \end{aligned}$$

for all $(a, t), (b, s) \in T$ and all $v, w \in K$.

Here now is our existence result for orthogonal and pseudo-quadratic Tits quadrangles.

Theorem 6.12. *Suppose that Λ either a non-degenerate quadratic space or that $\Lambda = (K, K_0, \sigma, L, q)$ is a non-degenerate pseudo-quadratic space such that $\text{char}(K) \neq 2$ if $K_0 = K$ and let Γ be as in 6.5. Then Γ an orthogonal Tits quadrangle of type Λ if Λ is a quadratic space and Γ is an pseudo-quadratic Tits quadrangle of type Λ if Λ is a pseudo-quadratic space.*

Proof. We continue with all the notation in 6.5, where we already showed that Γ is a Tits quadrangle. In Case I, let

$$(w_1, w_2, \dots, w_8) = (x'_1 x_2, x'_1, x'_1 x'_2, x'_2, x_1 x'_2, x_1, x_1 x_2, x_2),$$

let $x_1(t) = \delta_t$ and $x_3(t) = \gamma_t$ for all $t \in K$ and let $x_2(a) = \beta_a$ and $x_4(a) = \alpha_a$ for all $a \in L$. In Case II, let

$$(w_1, w_2, \dots, w_8) = (x_2, x_1 x_2, x_1, x_1 x'_2, x'_2, x'_1 x'_2, x'_1, x'_1 x_2),$$

let $x_1(a, t) = \alpha_{(a,t)}$ and $x_3(a, t) = \beta_{(a,t)}$ for all $(a, t) \in T$ and let $x_2(v) = \gamma_v$ and $x_4(v) = \delta_v$ for all $v \in K$. In both Case I and Case II, $(\Sigma, i \mapsto w_i)$ is a coordinate system of Γ . Let $i \mapsto U_i$ be the corresponding root group labeling. By 6.5(iii) and (6.10), x_i is injective and its image is U_i for all $i \in [1, 4]$ both in Case I and in Case II. It now requires only the formulas in 6.7, 6.8 and a bit of patience to check that the commutator relations in (i) and (ii) hold. \square

We now describe two further families of thick non-degenerate polar spaces of rank at least 3.

Definition 6.13. Let S be the thick non-degenerate polar space of rank 3 described in [11, 7.13] and let $\Gamma = \Gamma_S$ denote the corresponding Veldkamp quadrangle. Then Γ is a Tits quadrangle and there exist a skew-field K , a coordinate system $(\gamma, i \mapsto w_i)$ with root group labeling $i \mapsto U_i$ and isomorphisms $x_i: K^+ \rightarrow U_i$ for $i = 1$ and 3 and $x_i: K^+ \oplus K^+ \rightarrow U_i$ for $i = 2$ and 4 such that

$$\begin{aligned} [x_1(u), x_4(s, t)^{-1}] &= x_2(su, ut)x_3(sut) \\ [x_2(s, t), x_4(u, v)^{-1}] &= x_3(ut + sv) \end{aligned}$$


for all $u, v, s, t \in K$ as well as $[U_1, U_3] = [U_1, U_2] = [U_2, U_3] = [U_3, U_4] = 1$. We call Γ the Tits quadrangle of type D_3 over K .

Remark 6.14. Let S, Γ and K be as in 6.13. If K is commutative, then Γ is, in fact, of orthogonal type Λ where Λ is the hyperbolic quadratic space of dimension 2 over K . When K is non-commutative, S is non-embeddable and Γ is not of orthogonal or pseudo-quadratic type.

Definition 6.15. Let Γ be the Tits quadrangle obtained by applying the construction described in [3, 1.2.12 and 1.2.28] to the pair (Δ, T) , where Δ is the exceptional building $C_3(J)$ for some octonion division algebra J obtained by Galois descent from the Tits index



and T is the Tits index



and $\text{rk}(S_\Gamma) = 3$. We call Γ the Tits quadrangle of type E_7 over J . Let K denote the center of J . Then the commutator relations associated with Γ are as described in [7, 8.2] with Ξ set equal to the quadrangular algebra $\mathcal{Q}_2(J, K)$ described in [6, §4].

We can now formulate the classification of flat green Veldkamp quadrangles.

Theorem 6.16. *Let Γ be a flat green Veldkamp quadrangle. Then Γ is either a generalized quadrangle or one of the following holds:*

- (i) Γ is a Tits quadrangle of orthogonal of type Λ as defined in 6.11 for some non-degenerate quadratic space Λ with Witt index at least 3.
- (ii) Γ is a Tits quadrangle of pseudo-quadratic of type Λ as defined in 6.11 for some non-degenerate quadratic space Λ with Witt index at least 3.
- (iii) Γ is a Tits quadrangle of type D_3 or E_7 as defined in 6.13 and 6.15.

In particular, if Γ is not a generalized polygon, it is Moufang as defined in 6.2.

Proof. If Γ is not a generalized quadrangle, then by 5.18, the rank of the polar space S_Γ described in 5.22 is at least 3. By [10, 7.9.7], Γ is as in (i) or (ii) if $\text{rk}(S_\Gamma) \geq 4$ including the case that $\text{rk}(S_\Gamma)$ is infinite. We can thus suppose that $\text{rk}(S_\Gamma) = 3$. In this case, the claim holds by [11, Thms. 8.22 and 9.1]. Alternatively, we can cite [12, 40.17 and 40.25(ii)-(iv)] for the classification of thick buildings of type C_3 . If Γ is not of type D_3 , then by [11, Prop. 7.13], the building of type C_3 in [11, Thm. 7.4] is thick. It follows from the classification of thick buildings of type C_3 that either Γ is as in (i) or (ii) or Γ is of type E_7 . \square

Remark 6.17. Moufang quadrangles (i.e. generalized polygons satisfying the Moufang condition) were classified in [12, Thm. 17.4]. The Moufang quadrangles described in [12, 16.2, 16.3 and 16.5] are precisely the Tits quadrangles of orthogonal and pseudo-quadratic type as defined in 6.11 for Λ is anisotropic, but there are also three further families: the indifferent quadrangles described in [12, 16.4], the exceptional quadrangles of type E_6 , E_7 and E_8 described in [12, 16.6] and the exceptional quadrangles of type F_4 described in [12, 16.7]. There are, of course, also many generalized quadrangles that do not satisfy the Moufang condition; see, for example, [13].

Remark 6.18. Let Γ be a Tits n -gon for some n . As in 6.11, we choose a coordinate system $(\gamma, i \mapsto w_i)$ and let $i \mapsto U_i$ be the corresponding root group labeling. Let U_i^\sharp for all i and μ_γ be as in [7, 2.8 and 2.9] and let

$$H_i = \langle \mu_\gamma(a)\mu_\gamma(b) \mid a, b \in U_i^\sharp \rangle$$

for each i . As in [9, 2.23], we say that Γ is *razor-sharp* if for each i , every nontrivial H_i -invariant normal subgroup of the root group U_i contains elements of U_i^\sharp . In [3, 1.6.14], we showed that if Γ is razor-sharp, then $n = 3, 4, 6$ or 8 . In [3], [5] and [8], razor-sharp Tits triangles, hexagons and octagons were classified. Partial results on the case $n = 4$ can be found in [7] and [9]. In particular, it is shown in [9, 3.11] that every normal 5-plump razor-sharp Tits quadrangle is orthogonal, where “normal” is defined in analogy to the eponymous notion in [12, 21.7]. It is perhaps of interest to observe that many of the difficulties in the proof of this result stem from not knowing that the underlying Veldkamp quadrangle is green and that this fact emerges in the proof of [9, 3.11] only at the very end. (It is shown in [9, 3.10] that, conversely, every orthogonal Tits quadrangle is razor-sharp except for those that are of type D_3 ; see 6.14.)

Remark 6.19. Let Γ , $(\gamma, i \mapsto w_i)$ and $i \mapsto U_i$ be as in 6.13. Then the following observations can be shown to hold: There is an isomorphism $x_0: K^+ \oplus K^+ \rightarrow U_0$ such that

$$\begin{aligned} [x_0(s, t), x_3(u)^{-1}] &= x_1(tus)x_2(us, tu) \\ [x_0(s, t), x_2(u, v)^{-1}] &= x_1(tu + vs) \end{aligned}$$

for all $s, t, u, v \in K$. We have $x_4(s, t) \in U_4^\sharp$ if and only if $st \in K^\times$ and if $st \in K^\times$, then

$$\mu(x_4(s, t)) = x_0(t^{-1}, s^{-1})x_4(s, t)x_0(t^{-1}, s^{-1})$$

and conjugation by $\mu(x_4(s, t))$ induces the maps

$$\begin{aligned} x_0(u, v) &\mapsto x_4(svs, tut) \\ x_1(u) &\mapsto x_3(sut) \\ x_2(u, v) &\mapsto x_2(-svt^{-1}, -s^{-1}ut) \\ x_3(u) &\mapsto x_1(s^{-1}ut^{-1}) \\ x_4(u, v) &\mapsto x_0(t^{-1}vt^{-1}, s^{-1}us^{-1}) \end{aligned}$$

for all $u, v \in K$. Thus $\{x_4(u, 0) \mid u \in K\}$ is a nontrivial H_4 -invariant subgroup of U_4 that is disjoint from U_4^\sharp . Hence the Tits quadrangle Γ is not razor-sharp.

For the sake of completeness, we record also the following: There is an isomorphism $x_5: K^+ \rightarrow U_5$ such that

$$[x_2(s, t), x_5(u)^{-1}] = x_3(sut)x_4(su, ut)$$

for all $s, t, u \in K$. We have $x_1(u) \in U_1^\sharp$ if and only if $u \in K^\times$ and if $u \in K^\times$, then

$$\mu_\gamma(x_1(u)) = x_5(u^{-1})x_1(u)x_5(u^{-1})$$

and conjugation by $\mu_\gamma(x_1(u))$ induces the maps

$$\begin{aligned} x_1(s) &\mapsto x_5(u^{-1}su^{-1}) \\ x_2(s, t) &\mapsto x_4(-su^{-1}, -u^{-1}t) \\ x_3(s) &\mapsto x_3(s) \\ x_4(s, t) &\mapsto x_2(su, ut) \\ x_5(s) &\mapsto x_1(usu) \end{aligned}$$

for all $s, t \in K$.

REFERENCES

- [1] F. Buekenhout and A. M. Cohen, *Diagram Geometry*, Springer, Berlin, New York, Heidelberg, 2013.
- [2] J. R. Faulkner, Coordinatization of Moufang-Veldkamp planes, *Geom. Dedicata* **14** (1983), 189-201.
- [3] B. Mühlherr and R. M. Weiss, Tits Polygons, *Mem. A.M.S.*, to appear.
- [4] B. Mühlherr and R. M. Weiss, Root graded groups of rank 2, *J. Comb. Alg.* **3** (2019), 189-214.
- [5] B. Mühlherr and R. M. Weiss, Tits triangles, *Canad. Math. Bull.* **62** (2019), 583-601.
- [6] B. Mühlherr and R. M. Weiss, Isotropic quadrangular algebras, *J. Math. Soc. Japan* **71** (2019), 1321-1380.
- [7] B. Mühlherr and R. M. Weiss, The exceptional Tits quadrangles, *Transform. Groups* **25** (2020), 1289-1344.
- [8] B. Mühlherr and R. M. Weiss, Dagger-sharp Tits octagons, *J. Korean Math. Soc.* **58** (2021), 173-205.
- [9] B. Mühlherr and R. M. Weiss, Orthogonal Tits quadrangles, submitted.
- [10] E. Shult, *Points and Lines*, Springer, Berlin, New York, Heidelberg, 2011.
- [11] J. Tits, *Buildings of spherical type and finite BN-pairs*, Lecture Notes in Math. **386**, Springer, Berlin, New York, Heidelberg, 1974.
- [12] J. Tits and R. M. Weiss, *Moufang Polygons*, Springer, Berlin, New York, Heidelberg, 2002.
- [13] H. Van Maldeghem, *Generalized Polygons*, Birkhäuser, Basel, 1998.
- [14] F. D. Veldkamp, Polar geometry, I-V, *Proc. Kon. Ned. Akad. Wet.* **A62** (1959), 512-551; **A63**, 207-212 (= *Indag. Math.* **21**, **22**).

- [15] F. D. Veldkamp, Projective planes over rings of stable rank 2, *Geom. Dedicata* **11** (1981), 285-308.
- [16] F. D. Veldkamp, Geometry over rings, in: *Handbook of Incidence Geometry*, F. Buekenhout, ed., North-Holland, Amsterdam, 1995.

MATHEMATISCHES INSTITUT, UNIVERSITÄT GIESSEN, 35392 GIESSEN, GERMANY
Email address: `bernhard.m.muehlherr@math.uni-giessen.de`

DEPARTMENT OF MATHEMATICS, TUFTS UNIVERSITY, MEDFORD, MA 02155, USA
Email address: `rweiss@tufts.edu`