

TENSOR PRODUCT THEOREM FOR PARABOLIC λ -CONNECTIONS

MAO SHENG AND JIANPING WANG

ABSTRACT. We establish a tensor product theorem for slope semistable parabolic λ -connections over smooth projective varieties in arbitrary characteristic.

1. INTRODUCTION

A basic result in the theory of vector bundles is the so-called *tensor product theorem*: the tensor product of two slope semistable vector bundles over a smooth projective variety in char zero is again slope semistable. There are several different approaches to establish the theorem. The aim of the paper is to apply the recent theory of Higgs-de Rham flows [LSZ19] to this type problem. Our result is the following generalization of the tensor product theorem to parabolic vector bundles equipped with (not necessarily integrable) λ -connections.

Theorem 1.1. *Let k be an algebraically closed field. Let X be a smooth projective variety over k and L an ample line bundle over X . Let D be a reduced effective normal crossing divisor in X . Let $\lambda \in k$. Let $(E_{\bullet}^i, \nabla^i), i = 1, 2$ be two μ_L -semistable parabolic λ -connections over (X, D) . If either $\text{char } k = 0$ or $\text{char } k = p > 0$ and $\text{rk}(E^1) + \text{rk}(E^2) \leq p + 1$, then their tensor product is μ_L -semistable.*

Note that torsion freeness has been built into our definitions of parabolic sheaves and their tensor products (see §2).

In char zero, for Higgs bundles over X , the above result is due to C. Simpson ([Si, Corollary 3.8]). He first reduced the theorem to the curve case and then used the existence of Higgs-Yang-Mills connections for stable subquotients in a Jordan-Hölder filtration attached to a semistable Higgs bundle. However, in this approach, one might not be able to weaken the regularity condition on X (compare [HL, Theorem 3.1.4], [M, Theorem 5.16]), since the restriction theorem of Mehta-Ramanathan [MR] requires X be smooth.

Our method is basically to algebraize Simpson's approach. We shall use the reduction modulo p . Recall that in char p , Balaji-Parameswaran ([BP, Theorem 1.1]) proved that the tensor product of two semistable Higgs bundles over a smooth projective curve with trivial determinant and the rank

condition ¹ is semistable. In this paper, we shall give it another proof, using the existence result on semistable Higgs-de Rham flows in char p ([LSZ19, Theorem A.1], [La14, Theorem 5.12]). Our approach has some new features which are absent in the GIT approach: (i) the presence of D , that is, considering logarithmic Higgs fields does not introduce additional difficulty; (ii) the same method treats semistable connections on an equal footing. We remark that, even in the curve case, our result for connections cannot be directly deduced from the theorem of Balaji-Parameswaran. On the other hand, a Metha-Ramanathan type result for parabolic λ -connections should be supplied. In this aspect, one has the result of Simpson ([Si, Lemma 3.7]) for Higgs sheaves and the result of Mochizuki ([Mo, Proposition 3.29]) for parabolic Higgs sheaves. We observe that the integrality of Higgs fields plays no role in the proofs, and therefore the restriction theorem holds also for parabolic 0-connections. However, the argument does not work for connections ². We circumvent this difficulty by exploiting again the idea of Simpson on the existence of a gr-semistable filtration on a semistable connection.

In many ways, the notion of a Higgs-de Rham flow is a good analogue of Higgs-Yang-Mills flow in complex algebraic geometry. We would like to remind our readers of the recent work of Arapura [A], the work of Esnault-Groechenig [EG], the work of Langer [La15]-[La16] and the work of Lan-Sheng-Yang-Zuo [LSYZ].

2. PRELIMINARIES ON PARABOLIC λ -CONNECTIONS

Fix a locally noetherian scheme S . Let X be a smooth S -scheme, and $D = \sum_{i=1}^w D_i \subset X$ be an effective relative reduced normal crossing S -divisor. For brevity, we write the pair (X, D) by X_{\log} . Following [MY], we make the following

Definition 2.1. Let E be a torsion free coherent \mathcal{O}_X -module which is flat over S . A parabolic structure on E with parabolic support in D is an \mathbb{R} -indexed filtration of coherent sheaves E_\bullet over X such that

- (0) E_t is flat over S , for $t \in \mathbb{R}$,
- (1) $E_0 = E$,
- (2) $E_t \subseteq E_s$, for $t \geq s$,
- (3) $\exists \epsilon > 0$ such that $E_{t-\epsilon} = E_t$ holds for all $t \in \mathbb{R}$.
- (4) $E_{t+1} = E_t(-D)$, for $t \in \mathbb{R}$.

We call E together with a parabolic structure E_\bullet a *parabolic sheaf* over X_{\log}/S .

¹The rank condition of Balaji-Parameswaran is same as the one given in Theorem 1.1. It is interesting to note that the condition shows up from a totally different reason. See Step 3-4 in the proof of Theorem 1.1.

²Indeed, for a smooth hypersurface $Y \subset X$, a parabolic λ -connection over (X, D) induces a $\Omega_X(\log D + Y)|_Y$ -valued operator by restriction. It is only an $\Omega_X(\log D)|_Y$ -valued operator when $\lambda = 0$.

For an $1 \leq i \leq w$, by (3) there are only finitely many numbers $\{a_{ij}\} \subset [0, 1)$ such that $\text{Gr}_{a_{ij}}(E_\bullet) := (\frac{E_{a_{ij}}}{E_{a_{ij}+\delta}})|_{D_i}$ is nonzero for small enough $\delta > 0$. The collection $\{a_{ij}\}_j$ is called the set of *parabolic weights* of E_\bullet along D_i . A parabolic weight is said to be *rational* if it is a rational number. In this paper, the *parabolic filtration* associated to E_\bullet means the finite filtration

$$E = E_0 \supsetneq E_1 \supsetneq \cdots \supsetneq E_l \supsetneq E_{l+1} = E(-D),$$

by forgetting the parabolic weights of E_\bullet . One defines a morphism of parabolic sheaves, a parabolic coherent subsheaf (quotient sheaf), and the direct sum of parabolic sheaves in the natural way. However they do not form an abelian category, because of the torsion free assumption.

Example 2.2. (a) Trivial parabolic structure. The trivial parabolic structure on E is defined by $E_t := E(-[t]D)$, where $[\cdot]$ is the ceiling function. The parabolic weights are all zero.
 (b) Parabolic vector bundle. For an invertible sheaf L over X and a w -tuple $\vec{a} = (a_i) \in [0, 1)^w$, one defines a parabolic structure on L by

$$L_t^{\vec{a}} := L\left(\sum_{1 \leq i \leq w} -[t - a_i] D_i\right).$$

The parabolic weight of $L_\bullet^{\vec{a}}$ along D_i is a_i . A parabolic vector bundle over X_{\log} is a vector bundle V over X , equipped with a parabolic structure which Zariski locally is a direct sum of $L_\bullet^{\vec{a}}$ s. This is called a *locally abelian* parabolic bundle in [IS].

Lemma 2.3. For $S = \text{Spec } k$ with k algebraically closed and $\dim X = 1$, any parabolic sheaf over X_{\log}/k is a parabolic vector bundle.

Proof. Let E_\bullet be a parabolic sheaf over X_{\log} . As E is torsion free and X is a smooth curve over a field, E is locally free, viz. a vector bundle over X . Let $\{0 = a_0 < a_1 < \cdots < a_l < 1\}$ be the parabolic weights of E_\bullet at a point $x \in D$. Then the parabolic structure E_\bullet restricts to a flag of vector spaces over $k = k(x)$:

$$E(x) = F_0(E) \supsetneq F_1(E_x) \supsetneq \cdots \supsetneq F_l(E_x) \supsetneq F_{l+1}(E_x) = 0.$$

Set $n_i = \dim_k(F_i(E_x)/F_{i-1}(E_x))$. Choose a basis of $E(x)$ to split the above filtration. Then, by Nakayama lemma, it follows that locally around x , there is an isomorphism of parabolic sheaves:

$$E_\bullet \cong \bigoplus_{0 \leq i \leq l} ((\mathcal{O}_X)_\bullet^{a_i})^{\oplus n_i},$$

where $(\mathcal{O}_X)_\bullet^{a_i}$ is the parabolic line bundle as defined in 2.2(b). \square

To define the tensor product of two parabolic sheaves, we set $j : U = X - D \rightarrow X$ to be the open immersion. For any coherent sheaf E over X , put $E_\infty = \varinjlim E(nD) \subset j_* j^* E$. For a parabolic sheaf E_\bullet over X_{\log} , there is

a natural inclusion $E_t \rightarrow E_\infty$ for any t . Then for two parabolic sheaves E_\bullet and F_\bullet over X_{\log} , define for $t \in \mathbb{R}$

$$(E_\bullet \otimes F_\bullet)_t = \text{Im} \left(\bigoplus_{t_1+t_2=t} E_{t_1} \otimes F_{t_2} \longrightarrow E_\infty \otimes F_\infty / \text{torsion} \right).$$

It is not difficult to verify that $(E_\bullet \otimes F_\bullet)_\bullet$ defines a parabolic structure on $(E_\bullet \otimes F_\bullet)_0$.

Now we fix an element $\lambda \in \Gamma(S, \mathcal{O}_S)$, and for a parabolic sheaf E_\bullet over X_{\log}/S , we make the following

Definition 2.4. An S -relative λ -connection ∇ on E_\bullet is a collection $\{\nabla_t\}_{t \in \mathbb{R}}$ of \mathcal{O}_S -linear maps with $\nabla_t : E_t \rightarrow E_t \otimes \Omega_{X/S}(\log D)$ satisfying

- (a) $\nabla(fs) = \lambda df \otimes s + f\nabla(s)$ for $f \in \mathcal{O}_X$ and $s \in E_t$;
- (b) For any $t \geq s$, the following diagram commutes:

$$\begin{array}{ccc} E_t & \xrightarrow{\nabla_t} & E_t \otimes \Omega_{X/S}(\log D) \\ \downarrow \iota_{t,s} & & \downarrow \iota_{t,s} \otimes id \\ E_s & \xrightarrow{\nabla_s} & E_s \otimes \Omega_{X/S}(\log D) \end{array}$$

where $\iota_{t,s} : E_t \rightarrow E_s$ is the natural inclusion.

Remark 2.5. A λ -connection is said to be integrable if $\nabla \wedge \nabla = 0$. The integrability condition is automatic if X/S is of relative dimension one. For a λ -connection ∇ on E_\bullet , there is a natural λ -connection on E_∞ . Therefore, for two parabolic λ -connections (E_\bullet, ∇_E) and (F_\bullet, ∇_F) , their tensor product is defined to be the $(E_\bullet \otimes F_\bullet)_\bullet$, equipped with the λ -connection induced from the tensor product of λ -connections over $E_\infty \otimes F_\infty$.

In order to handle parabolic structures in this paper, we need to recall the generalized Biswas-Iyer-Simpson (BIS) correspondence for parabolic λ -connections in all characteristics ([Bi],[IS],[KS20, §2]). Let k be an algebraically closed field and let N be a positive integer such that $\text{char } k \nmid N$. Let X be a smooth curve over k and let $\pi : X' \rightarrow X$ be a cyclic cover of order N with branch divisor $D \subset X$. The constant group scheme $G \cong \mathbb{Z}/N\mathbb{Z}$ acts on X' with quotient $X'/G \cong X$. Let $D' = \sum_{i=1}^l D'_i$ be the reduced divisor $(\pi^*D)_{red}$. Set X'_{\log} to be the log pair (X', D') .

Proposition 2.6. [KS20, Proposition 2.14] *Notations as above. The parabolic pushforward $\pi_{\text{par}*}$ and parabolic pullback π_{par}^* induce an equivalence of categories between the category of G -equivariant λ -connections over X'_{\log} and the category of parabolic λ -connections over X_{\log} whose parabolic structure is supported along D and has weights in $\frac{1}{N}\mathbb{Z}$.*

When X/k projective, equipped with an ample line bundle L , one may define the *parabolic degree* of a parabolic sheaf E_\bullet over X_{\log}/k by the following

formula:

$$\text{pardeg}_L(E_\bullet) = \text{deg}_L(E_0) + \sum_{1 \leq i \leq w} \sum_j a_{ij} n_{ij} \text{deg}_L D_i,$$

where $n_{ij} := \text{rkGr}_{a_{ij}}(E_\bullet)$. Clearly, for a parabolic sheaf with the trivial parabolic structure, the parabolic degree and the usual degree coincide.

Definition 2.7. Let k be an algebraically closed field. Let (E_\bullet, ∇) be a λ -connection over X_{\log} . It is said to be μ_L -(semi)stable, if for any ∇ -invariant parabolic coherent subsheaf $F_\bullet \subset E_\bullet$, the following inequality holds:

$$\text{rk}(E) \text{pardeg}_L(F_\bullet) (\leq) < \text{rk}(F) \text{pardeg}_L(E_\bullet).$$

For a nonzero parabolic sheaf E_\bullet over X_{\log} , one defines its parabolic slope (with respect to L) by

$$\mu_L(E_\bullet) = \text{pardeg}_L(E_\bullet) / \text{rk}(E).$$

Lemma 2.8. *Notations as above. Let F_\bullet^i , $i = 1, 2$ be two parabolic sheaves over X_{\log} . Then*

$$\mu_L(F_\bullet^1 \otimes F_\bullet^2) = \mu_L(F_\bullet^1) + \mu_L(F_\bullet^2).$$

Proof. We may assume L to be very ample. By Bertini's theorem ([H, Theorem 8.18]), we may restrict F_\bullet^i to a smooth complete intersection curve in the linear system $|L|$, and $\text{pardeg}(F_\bullet^i)$ equals the parabolic degree of the restricted parabolic sheaf. Therefore, we may assume $\dim X = 1$. By Lemma 2.3, we may assume then F_\bullet^i s are parabolic vector bundles.

Consider first the case that the parabolic structures on F_\bullet^i s are trivial. The truth of this case is a standard fact: one may use the splitting principle to reduce the problem to the case of line bundles that is obviously true. Consider further the case that one of the parabolic structures is nontrivial but satisfies either i) $\text{char } k = 0$, the parabolic weights are rational, or ii) $\text{char } k = p > 0$, the parabolic weights are rational whose denominators are coprime to p . One applies Proposition 2.6 to reduce this case to the previous one. For the general case, we use a density argument as follows: let

$$V^i = \{\vec{a}^i = (a_{jk}^i) | a_{jk}^i \in \mathbb{R}, 0 \leq a_{jk}^i < 1\}, i = 1, 2$$

be the weight space containing the parabolic weights (say \vec{a}_0^i) of F_\bullet^i . For $\vec{a}^i \in V^i$, we denote by F_\bullet^{i, \vec{a}^i} the parabolic bundle whose associated parabolic filtration is the same as that of F_\bullet^i but with parabolic weights \vec{a}^i . In this terminology, $F_\bullet^i = F_\bullet^{i, \vec{a}_0^i}$. Define the function

$$\Psi : V^1 \times V^2 \longrightarrow \mathbb{R}, \quad (\vec{a}^1, \vec{a}^2) \longmapsto \mu(F_\bullet^{1, \vec{a}^1} \otimes F_\bullet^{2, \vec{a}^2}) - \mu(F_\bullet^{1, \vec{a}^1}) - \mu(F_\bullet^{2, \vec{a}^2}).$$

By definition of parabolic slope, it is continuous. Moreover, we notice that the subset $Z \subset V^1 \times V^2$, consisting of the parabolic weights in the second case, is actually dense. Since by the second case Ψ restricts to Z as the zero map, it is identically zero. Thus the general case follows. \square

3. RESTRICTION THEOREM

Mehta and Ramanathan [MR, Theorem 6.1] proved that the restriction of a semistable torsion free coherent sheaf to a general hyperplane section of sufficiently high degree is again semistable. Simpson [Si, Lemma 3.7] generalized it to Higgs sheaves, and later Mochizuki [Mo, Proposition 3.29] generalized it further to parabolic Higgs sheaves. Langer [La15, Theorem 10] has obtained a much stronger restriction theorem, in that the degree of a general hypersurface can be made explicit. To our purpose, it suffices to generalize the Mehta-Ramanathan restriction theorem to the semistable parabolic λ -connections.

We resume the setting of Theorem 1.1. Let (E_\bullet, ∇) be a μ_L -semistable parabolic λ -connection over X_{\log} . By Bertini's theorem, for a general member $Y \in |\mathcal{O}_X(m)|$ for m large enough, E_\bullet restricts to a parabolic sheaf $E_\bullet|_Y$ over $Y_{\log} = (Y, Y \cap D)$, and $Y + D$ is normal crossing so that $D \cap Y$ is a reduced effective normal crossing divisor on Y . By a simple local calculation, one sees that the parabolic subsheaf $E_\bullet \otimes \mathcal{O}_X(-Y) \subset E_\bullet$ is annihilated by the composite map

$$E_\bullet \xrightarrow{\nabla} E_\bullet \otimes \Omega_{X/k}(\log D) \xrightarrow{\text{res}} E_\bullet|_Y \otimes \Omega_{X/k}(\log D)|_Y \xrightarrow{\text{id} \otimes \text{ev}} E_\bullet|_Y \otimes \Omega_{Y/k}(\log D \cap Y).$$

So ∇ on E_\bullet yields the operator $\nabla|_Y$ on $E_\bullet|_Y$, which is easily seen to be a parabolic λ -connection over $(Y, D \cap Y)/k$. The aim of the section is the following theorem.

Theorem 3.1. *Let $(X_{\log}, L, E_\bullet, \nabla)$ be as above. Then for general hyperplane sections Y of sufficiently high degree, $(E_\bullet, \nabla)|_Y := (E_\bullet|_Y, \nabla|_Y)$ is a semistable parabolic λ -connection.*

Let us consider an arbitrary parabolic λ -connection (E_\bullet, ∇) over X_{\log}/k . A level n filtration $Fil = Fil_\bullet^*$ of parabolic subsheaves of E_\bullet

$$E_\bullet = Fil_\bullet^0 \supset Fil_\bullet^1 \supset \cdots \supset Fil_\bullet^n \supset 0$$

is said to be *Griffiths transverse* if

$$\nabla(Fil_\bullet^*) \subset Fil_\bullet^{*-1} \otimes \Omega_{X/k}(\log D)$$

holds. Taking grading yields a collection $\{Gr_{Fil_t}(E_t, \nabla)\}_{t \in \mathbb{R}}$ of logarithmic 0-connections and morphisms $\{\phi_{t,s}\}_{t \geq s}$ of logarithmic 0-connections

$$\phi_{t,s} : Gr_{Fil_t}(E_t, \nabla) \rightarrow Gr_{Fil_s}(E_s, \nabla).$$

We consider a construction of Fil_\bullet such that $\phi_{t,s}$ are all injective.

Lemma 3.2. *Let Fil_0 be a saturated and Griffiths transverse filtration on (E_0, ∇) . For $n \in \mathbb{Z}$, set $Fil_n = Fil_0 \otimes Fil_{tr}$ to be the tensor filtration over $V_n \cong V_0 \otimes \mathcal{O}_X(-nD)$, where Fil_{tr} stands for the trivial filtration. For $n \leq t < n + 1$, set $Fil_t = E_t \cap Fil_n$ to be the restricted filtration over E_t . Then, the filtration $Fil = Fil_\bullet$ defined as above is a Griffiths transverse filtration with $\phi_{t,s}$ injective for any $t \geq s$.*

We call Fil the *induced* filtration from Fil_0 .

Proof. Let $\lambda d : \mathcal{O}_X(D) \rightarrow \mathcal{O}_X(D) \otimes \Omega_{X/k}(\log D)$ be the parabolic λ -connection induced by the exterior differential d . Then for any $t \in \mathbb{R}$,

$$(E_{t-1}, \nabla) \cong (E_t, \nabla) \otimes (\mathcal{O}_X(D), \lambda d).$$

Thus for $t \in \mathbb{Z}$, the Griffiths transversality of Fil_t follows from that of Fil_0 . Then, for any $t \in \mathbb{R} - \mathbb{Z}$, Fil_t is also Griffiths transverse by construction. Again by construction, for $n \leq s \leq t < n+1$, $\phi_{t,s}$ is injective. Note that the composite

$$Gr_{Fil_{n+1}}(V_{n+1}, \nabla) \xrightarrow{\phi_{n+1,t}} Gr_{Fil_t}(V_t, \nabla) \xrightarrow{\phi_{t,n}} Gr_{Fil_n}(V_n, \nabla)$$

is $\phi_{n+1,n}$. As Fil_0 is saturated, it follows that $\phi_{n+1,n}$ is injective. So $\phi_{n+1,t}$ is also injective. Then it follows that $\phi_{t,s}$ is injective for any $t \geq s$, by writing it as a composite of finitely many injective morphisms. \square

Definition 3.3. A filtration Fil over (E_\bullet, ∇) is *gr-semistable* if it is the induced filtration from a saturated and Griffiths transverse filtration Fil_0 of finite level over (E_0, ∇) such that the associated graded parabolic 0-connection $Gr_{Fil}(E_\bullet, \nabla)$ is μ_L -semistable.

For $D = \emptyset$, the above notion reduces back to the one considered in Appendix [LSZ19]. We have a natural generalization of [LSZ19, Theorem A.4] to the parabolic setting.

Proposition 3.4. *Let k be an algebraically closed field, and X a smooth projective variety over k , equipped with a reduced effective normal crossing divisor D . Let L be an ample line bundle over X . Let (E_\bullet, ∇) be a μ_L -semistable parabolic λ -connection over (X, D) . Then there exists a gr-semistable filtration Fil over (E_\bullet, ∇) .*

Proof. We shall adapt the proof of [LSZ19, Theorem A.4] to the current setting. One starts with an arbitrary filtration Fil , induced from some saturated Griffiths transverse filtration Fil_0 of level n over (E_0, ∇_0) . Consider the associated graded parabolic 0-connection

$$(E_\bullet = \bigoplus_{i=0}^n E_\bullet^i, \theta) := Gr_{Fil}(E_\bullet, \nabla).$$

If it is semistable, we are done. Otherwise, let $I_{Fil, \bullet}$ be its maximal destabilizer—it is the maximal θ -invariant parabolic subsheaf of E_\bullet with maximal parabolic slope. It is graded, that is, for $0 \leq i \leq n$ and $0 \leq t < 1$, one has

$$I_{Fil,t} = \bigoplus_{i=0}^n I_{Fil,t}^i, \quad I_{Fil,t}^i = I_{Fil,t} \cap E_t^i.$$

Moreover, $I_{Fil,t} = I_{Fil,0} \cap E_t$ for $0 \leq t < 1$ and hence

$$I_{Fil,t}^i = I_{Fil,t} \cap E_t^i = I_{Fil,0} \cap E_t \cap E_t^i = I_{Fil,0} \cap E_t \cap E_0^i = I_{Fil,0}^i \cap E_t^i.$$

The key is the construction of a new filtration $\xi(Fil)$ (of level $n + 1$), which is defined by

$$\xi(Fil)_\bullet^i := \ker(E_\bullet \rightarrow \frac{E_\bullet/Fil_\bullet^i}{I_{Fil,\bullet}^{i-1}}), \quad 1 \leq i \leq n + 1, \quad \xi(Fil)_\bullet^0 = E_\bullet.$$

Claim: $\xi(Fil)$ is the induced filtration from $\xi(Fil)_0$, which is again a saturated Griffiths transverse filtration over (E_0, ∇) . Note that $\xi(Fil)_0$ is nothing but the $\xi(Fil_0)$ in the terminology of [LSZ19, A.6.2], and therefore saturated and Griffiths transverse. It remains to show the induced property. For that, we may assume $0 \leq t < 1$. Note that there is a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & Fil_t^i & \longrightarrow & \xi(Fil)_t^i & \longrightarrow & I_{Fil,t}^{i-1} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & Fil_0^i \cap E_t & \longrightarrow & \xi(Fil)_0^i \cap E_t & \longrightarrow & I_{Fil,0}^i \cap E_t^i \longrightarrow 0 \end{array}$$

The top and bottom horizontal sequences are exact. The left and right vertical morphism are identity, and the middle vertical morphism is injective. It follows that the middle vertical morphism is also identity. The claim is proved.

By the claim, there is a short exact sequence of graded parabolic 0-connections:

$$(3.1) \quad 0 \longrightarrow Gr_{Fil_\bullet}(E_\bullet)/I_{Fil,\bullet} \longrightarrow Gr_{\xi(Fil)_\bullet}(E_\bullet) \longrightarrow I_{Fil,\bullet}^{[1]} \longrightarrow 0$$

Granted the above short sequence, the arguments for [LSZ19, Lemmata A.7-A.8] work verbatim for the parabolic setting. They imply that, there exists some $k_0 \in \mathbb{N}$, such that $\xi^{k_0}(Fil)$ is a gr-semistable filtration. \square

We proceed to the proof of Theorem 3.1.

Proof. We observe that the proof of [Mo, Proposition 3.29] is characteristic free, and does not require components of D be smooth. As remarked in §1, the integrality of a Higgs field has not been used in the proof of [Si, Lemma 3.7]. Therefore, we have the extension of Mochizuki's restriction theorem to a semistable parabolic 0-connection over X_{\log} .

Now let (E_\bullet, ∇) be a semistable parabolic λ -connection with $\lambda \neq 0$. Take a gr-semistable filtration Fil_\bullet on (E_\bullet, ∇) by Proposition 3.4. By [MR, Proposition 1.5], for a general hyperplane section Y , one has

$$Gr_{Fil_\bullet|_Y}(E_\bullet, \nabla)|_Y \cong (Gr_{Fil_\bullet}(E_\bullet, \nabla))|_Y.$$

Note that $Gr_{Fil_\bullet}(E_\bullet, \nabla)$ is a parabolic 0-connection. By Mochizuki's restriction theorem, $(Gr_{Fil_\bullet}(E_\bullet, \nabla))|_Y$ is semistable for a general hyperplane section Y of arbitrarily high degree. Because taking grading preserves the parabolic degree, the semistability of $Gr_{Fil_\bullet|_Y}(E_\bullet, \nabla)|_Y$ implies the semistability of $(E_\bullet, \nabla)|_Y$, as claimed. \square

4. TENSOR PRODUCT THEOREM

In this section, we shall establish Theorem 1.1. To begin with, we prove three preparatory lemmata. The first lemma deals with the stability with respect to varying parabolic weights.

Lemma 4.1. *Notations as Theorem 1.1. Let (E_\bullet, ∇) be a parabolic λ -connection over X_{\log} with parabolic weights $\vec{a} = (a_{ij})$.*

- (1) *If (E_\bullet, ∇) is μ_L -stable with respect to \vec{a} , then for any $n = 1, 2, \dots$, there exists a vector $\vec{b}^n = (b_{ij}^n)$ in euclidean space with $b_{ij}^n \in \mathbb{Q} \cap [0, 1)$ and $0 < |\vec{b}^n - \vec{a}| < \frac{1}{n}$ such that (E_\bullet, ∇) is μ_L -stable with respect to the parabolic weights \vec{b}^n . Moreover, if $\text{char } k > 0$, then one may choose the denominator of each $\{b_{ij}^n\}$ to be coprime to $\text{char } k$.*
- (2) *If there exists a sequence of rational vectors $\vec{b}^n = (b_{ij}^n)$ as above such that (E_\bullet, ∇) is μ_L -semistable with respect to the parabolic weights \vec{b}^n , then (E_\bullet, ∇) is μ_L -semistable with respect to \vec{a} .*

Proof. For a vector $\vec{b} = (b_{ij})$ with $b_{ij} \in [0, 1)$, (E_\bullet, ∇) being μ_L -stable with respect to \vec{b} means that for any ∇ -invariant parabolic coherent subsheaf $F_\bullet \subset E_\bullet$, one has the following inequality

$$(4.1) \quad \deg_L(F_0) < \frac{\text{rk}(F_0)}{\text{rk}(E_0)} \deg_L(E_0) + \sum_{i,j} \left(\frac{\text{rk}(F_0)}{\text{rk}(E_0)} n_{ij} - n_{ij}^{F_\bullet} \right) \deg_L(D_j) b_{ij}.$$

Suppose (E_\bullet, ∇) to be μ_L -stable with respect to \vec{a} . Set

$$A = \deg_L(E_0) + \sum_{i,j} n_{ij} \deg_L(D_j) a_{ij}.$$

Then the inequality 4.1 implies that there is an upper bound $\deg_L(F_0) \leq A$ for all ∇ -invariant coherent subsheaves $F_\bullet \subset E_\bullet$. On the other hand, it is clear that there exists a constant A' so that once $\deg_L(F_0) < A'$ is satisfied, the inequality 4.1 holds automatically. Hence, the stability condition imposes only finitely many inequalities for \vec{b} . We list them as

$$L_1(\vec{b}) > 0, L_2(\vec{b}) > 0, \dots, L_m(\vec{b}) > 0,$$

where each L_i is linear with *rational* coefficients. In particular, it cuts out an *open* rational cone C in the vector space of \vec{b} . Since C contains a solution \vec{a} , it must be of positive dimension. Therefore, we may find an approximating sequence \vec{b}^n to \vec{a} with rational entries, as requested in (i).

Conversely, assume that (E_\bullet, ∇) is μ_L -semistable with respect to \vec{b}^n . Then by the continuity of L_i , it follows that (E_\bullet, ∇) is μ_L -semistable with respect to \vec{a} . □

Next we consider the necessary device for mod p reduction. In particular, we shall assume $\text{char } k = 0$ in the second lemma. We may assume $\lambda \in \{0, 1\}$,

since $\lambda^{-1}\nabla$ is a connection for $\lambda \neq 0$. For the given tuple $\{X, D, L, E_\bullet, \nabla\}$ over k , there exists an integral (affine) scheme S of finite type over \mathbb{Z} , and a new tuple $\{\mathcal{X}, \mathcal{D}, \mathcal{L}, \mathcal{E}_\bullet, \nabla\}$ defined over S , where

- (a) \mathcal{X} is smooth projective over S ,
- (b) \mathcal{D} is an S -divisor with normal crossing,
- (c) \mathcal{L} is an S -ample line bundle over \mathcal{X} ,
- (d) $(\mathcal{E}_\bullet, \nabla)$ is an S -relative parabolic λ -connection over $(\mathcal{X}, \mathcal{D})$,

together with a k -rational point $s_\infty : \text{Spec } k \rightarrow S$ such that

$$\{\mathcal{X}, \mathcal{D}, \mathcal{L}, \mathcal{E}_\bullet, \nabla\}_{s_\infty} := \{\mathcal{X}, \mathcal{D}, \mathcal{L}, \mathcal{E}_\bullet, \nabla\} \times_{\text{Spec } k} S \cong \{X, D, L, E_\bullet, \nabla\}.$$

Such a tuple is called to be an *arithmetic model* of $\{X, D, L, E_\bullet, \nabla\}$.

Lemma 4.2. *Notations as above. Assume $\dim X = 1$. Then (E_\bullet, ∇) is μ_L -(semi)stable if and only if there exists infinitely many prime numbers $\{p_i\}$ and geometric points $\{s_i \in S\}$ with $\text{char } k(s_i) = p_i$ such that $(\mathcal{E}_\bullet, \nabla)_{s_i}$ is $\mu_{\mathcal{L}_{s_i}}$ -(semi)stable.*

Proof. The if direction follows from the fact that the Harder-Narasimhan function is upper-semicontinuous ([Si, Theorem 3], see also §2[N]). The basic idea is that the maximal destabilizer of (E_\bullet, ∇) spreads over an open subscheme $S' \subset S$ containing s_∞ . For $\lambda = 0$, the only if direction follows from the openness of (semi)stability. In any event, we provide below a proof of the only if direction for an arbitrary λ . Assume the contrary. We may shrink S such that for any geometric point $s \in S$, $(\mathcal{E}_\bullet, \nabla)_s$ is unstable. Then there exists a pair of numbers (d, r) , an infinite sequence of prime numbers $\{p_i\}_i$ and a sequence of geometric points $\{s_i\}_i \subset S$ with $\text{char } k(s_i) = p_i$, such that the degree zero term F_0^i of the minimal quotient $(\mathcal{E}_\bullet, \nabla)_{s_i} \twoheadrightarrow (F_\bullet^i, \nabla_{s_i})$ is of $\deg(F_0^i) = d$ and $\text{rk}(F_0^i) = r$. Let $\text{Quot}_{\mathcal{E}_0/\mathcal{X}/S}^{(d,r)}$ be the Quot-scheme parametrizing quotient coherent sheaves of $(\mathcal{E}_0)_s$ whose degree is d and rank is r (see e.g. [M, Theorem 1.5]). There is a *closed* subscheme $\text{Quot}_{\mathcal{E}_0/\mathcal{X}/S}^{(d,r),\nabla}$ parametrizing ∇ -invariant quotient coherent sheaves. To see this, let $Q = \text{Quot}_{\mathcal{E}_0/\mathcal{X}/S}^{(d,r)}$ and $\pi : \mathcal{E}_Q \rightarrow \mathcal{G}$ be the universal quotient. Let $\mathcal{K} = \ker(\pi)$ and ∇_Q be the induced λ -connection on \mathcal{E}_Q . Consider the composite

$$\Phi : \mathcal{K} \xrightarrow{\nabla_Q} \mathcal{E}_Q \otimes \Omega_{\mathcal{X}_Q/Q}(\log \mathcal{D}_Q) \xrightarrow{\pi \otimes \text{id}} \mathcal{G} \otimes \Omega_{\mathcal{X}_Q/Q}(\log \mathcal{D}_Q).$$

Let $\mathcal{Z} \rightarrow \mathcal{X}_Q$ be the closed subscheme defined by the zero locus of Φ . Its composite with the natural projection $\mathcal{X}_Q \rightarrow Q$ gives the projective morphism $\alpha : \mathcal{Z} \rightarrow Q$. Let $\alpha(\mathcal{Z})$ be the scheme theoretical image of α , which is a closed subscheme of Q . Then $\text{Quot}_{\mathcal{E}_0/\mathcal{X}/S}^{(d,r),\nabla} \rightarrow \alpha(\mathcal{Z})$ is the closed subscheme of $\alpha(\mathcal{Z})$ (hence also a closed subscheme of Q) over which α is of relative dimension one. Because $(\text{Quot}_{\mathcal{E}_0/\mathcal{X}/S}^{(d,r),\nabla})_{s_i} \neq \emptyset$ for each i , it follows from Hilbert's Nullstellensatz that $\text{Quot}_{\mathcal{E}_0/\mathcal{X}/S}^{(d,r),\nabla}(k) \neq \emptyset$. But then it means that there is a ∇ -invariant quotient sheaf $E_0 \twoheadrightarrow F_0$ such that $\mu_L(F_\bullet) < \mu_L(E_\bullet)$ where F_\bullet is

the induced parabolic structure on F_0 . Contradiction. This shows the only if direction for semistability. The only if direction for stability is proved along the same line. \square

Finally, we shall need to establish a supplementary property of the generalized BIS correspondence (Proposition 2.6).

Lemma 4.3. *Notations as above. The parabolic pushforward as well as the parabolic pullback commute with direct sum and tensor product.*

Proof. It suffices to show the parabolic pushforward commutes with direct sum and tensor product by the equivalence of categories. Let $\pi : X' \rightarrow X$ be as above. Recall that for a G -equivariant sheaf V over X' , the parabolic structure on $\pi_{\text{par}*}V$ is given by

$$(\pi_{\text{par}*}V)_t := (\pi_*(V \otimes \mathcal{O}_{X'}(\sum_j [-tN]D'_j)))^G.$$

Since direct sum commutes with tensor product, pushforward π_* and taking G -invariants, it follows that parabolic pushforward commutes with direct sum. Let $(V_i, \nabla_i), i = 1, 2$ be two G -equivariant λ -connections over X'_{\log} . One notices that there is a natural morphism of parabolic λ -connections

$$\varphi : \pi_{\text{par}*}(V_1, \nabla_1) \otimes \pi_{\text{par}*}(V_2, \nabla_2) \rightarrow \pi_{\text{par}*}((V_1, \nabla_1) \otimes (V_2, \nabla_2)).$$

We are going to show that φ is an isomorphism. The problem is local. As $\dim X = 1$, by Lemma 2.3 it follows that, we may assume that $V_i, i = 1, 2$ are direct sums of G -equivariant line bundles (via the BIS correspondence, direct sums of parabolic line bundles over X_{\log} and direct sums of G -equivariant line bundles over X' correspond to each other.). We may assume further that V_i s are simply G -equivariant line bundles.

Etale locally we may assume that

$$X' = \{y^N = x\} \subset \mathbb{A}_k^2 \xrightarrow{pr} X = \{y = 0\} \cong \mathbb{A}^1,$$

and $V_i = \mathcal{O}_{X'}e_i$ is the trivial line bundles with base element e_i . Take a generator $\sigma \in G$. Then the G -action on $\mathcal{O}_{X'}$ and V_i are described respectively by

$$\sigma(y) = \xi y, \quad \sigma(e_i) = \xi^{-n_i} e_i,$$

where ξ is an N -th primitive root of unity, and $0 \leq n_i/N < 1$ is the parabolic weight of $\pi_{\text{par}*}V_i$. Then the parabolic structure of $\pi_{\text{par}*}V_i$ is determined by

$$(\pi_{\text{par}*}V_i)_t = \begin{cases} \mathcal{O}_X y^{n_i} e_i & 0 \leq t \leq \frac{n_i}{N} \\ \mathcal{O}_X x y^{n_i} e_i & \frac{n_i}{N} < t < 1. \end{cases}$$

It follows that the parabolic structure of $\pi_{\text{par}*}V_1 \otimes \pi_{\text{par}*}V_2$ is determined by

$$(\pi_{\text{par}*}V_1 \otimes \pi_{\text{par}*}V_2)_t = \begin{cases} \mathcal{O}_X y^m e_1 \otimes e_2 & 0 \leq t \leq \frac{m}{N} \\ \mathcal{O}_X x y^m e_1 \otimes e_2 & \frac{m}{N} < t < 1. \end{cases}$$

Here $m = n_1 + n_2$ if $n_1 + n_2 < N$, and $m = n_1 + n_2 - N$ if $n_1 + n_2 \geq N$. On the other hand, $V_1 \otimes V_2$ is the trivial bundle with base element $e_1 \otimes e_2$.

It is straightforward to verify that the last expression is nothing but the parabolic structure of $\pi_{\text{par}*}(V_1 \otimes V_2)$ for $0 \leq t < 1$. Therefore φ is indeed an isomorphism. The lemma is proved. \square

Now we proceed with the proof of Theorem 1.1.

Proof. By Theorem 3.1, we may assume $\dim X = 1$ in the following. We divide the whole proof into five steps. In the first four steps, we shall assume $(E_{\bullet}^i, \nabla^i), i = 1, 2$ to be μ_L -stable.

Step 1. Let X be a smooth projective curve defined over k , and $D \subset X$ a reduced effective divisor. By Lemma 4.1, we may assume the parabolic weights of $E_{\bullet}^i, i = 1, 2$ are contained in $\frac{1}{N}\mathbb{Z}$ and $(N, p) = 1$ when $\text{char } k = p > 0$. Fix a closed point $x_0 \in X$ away from D . We may choose another reduced effective divisor B such that the degree of $\tilde{D} = D + B$ is divisible by N . Let $\pi : X' \rightarrow X$ be the cyclic cover with branch divisor \tilde{D} . We claim that the parabolic pullback $\pi_{\text{par}}^*(E_{\bullet}^i, \nabla^i)$, which is a G -equivariant logarithmic λ -connection over $(X', \pi^{-1}\tilde{D})$, is semistable. Indeed, if it were unstable, then the maximal destabilizer would be G -equivariant and invariant under $\pi_{\text{par}}^*\nabla^i$. By the generalized BIS correspondence, it descends to a parabolic subsheaf of E_{\bullet}^i which is ∇^i -invariant. The claim follows. Moreover by Lemma 4.3, we have

$$\pi_{\text{par}}^*((E_{\bullet}^1, \nabla^1) \otimes (E_{\bullet}^2, \nabla^2)) \cong \pi_{\text{par}}^*(E_{\bullet}^1, \nabla^1) \otimes \pi_{\text{par}}^*(E_{\bullet}^2, \nabla^2).$$

Therefore, we may assume the parabolic structure of E_{\bullet}^i to be trivial (Example 2.2 (a)).

Step 2. When $\text{char } k = 0$, we take an arithmetic model of $\{X, D, L, E_{\bullet}^1 \oplus E_{\bullet}^2, \nabla^1 \oplus \nabla^2\}$. By Lemma 4.2, there exists an infinite sequence of geometric points $\{s_i\} \subset S$ such that both $(\mathcal{E}_{\bullet}^1, \nabla)_{s_i}$ and $(\mathcal{E}_{\bullet}^2, \nabla)_{s_i}$ are stable. We remove those (at most finitely many) s_i s satisfying $p_i < \text{rk}(E^1) + \text{rk}(E^2) - 1$ from the sequence. By Lemma 4.2 again, the char zero statement follows from the char p statement.

Step 3. We choose an arbitrary $W_2(k)$ -lifting of the pair (X, D) , and let C^{-1} be inverse Cartier transform of Ogus-Vologodsky with respect to this lifting (see Appendix [LSYZ]). Let $(E^i, \nabla^i), i = 1, 2$ be two stable logarithmic λ -connections over (X, D) . Then by Proposition 3.4, there exists a filtration Fil_{-1}^i over E^i such that $\text{Gr}_{Fil_{-1}^i}(E^i, \nabla^i)$ is a graded semistable logarithmic Higgs bundle over (X, D) . Because of the rank assumption, one has $\text{rkGr}_{Fil_{-1}^i}(E^i, \nabla^i) \leq p$. Therefore, there exists a semistable Higgs-de Rham flow initializing $\text{Gr}_{Fil_{-1}^i}(E^i, \nabla^i)$ for $i = 1, 2$ ([LSZ19, Theorem A.1],

[La14, Theorem 5.12]). That is, we obtain a flow of the following form:

$$\begin{array}{ccccc}
 (E^i, \nabla^i) & & (H_0^i, \nabla_0^i) & & (H_1^i, \nabla_1^i) \\
 \searrow^{Gr_{Fil_{-1}^i}} & & \nearrow^{C^{-1}} & \searrow^{Gr_{Fil_0^i}} & \nearrow^{C^{-1}} & \searrow^{Gr_{Fil_1^i}} \\
 & (E_0^i, \theta_0^i) & & (E_1^i, \theta_1^i) & & \dots,
 \end{array}$$

in which each Higgs term in bottom is semistable. Since a Higgs-de Rham flow preserves the rank, the rank assumption implies that $\text{rk}E_j^1 + \text{rk}E_j^2 \leq p+1$ for each $j \geq 0$. So the nilpotent indices l_j^i s of θ_j^i s satisfy $l_j^1 + l_j^2 \leq p-1$. By Lemma 3.4 [KS20], it follows that

$$C^{-1}((E_j^1, \theta_j^1) \otimes (E_j^2, \theta_j^2)) \cong (H_j^1, \nabla_j^2) \otimes (H_j^2, \nabla_j^2).$$

Thus, we obtain the tensor product of flows:

$$\begin{array}{ccc}
 (E^1, \nabla^1) \otimes (E^2, \nabla^2) & & (H_0^1, \nabla_0^2) \otimes (H_0^2, \nabla_0^2) \\
 \searrow^{Gr_{Fil_{-1}^1 \otimes Fil_{-1}^2}} & & \nearrow^{C^{-1}} & \searrow^{Gr_{Fil_0^1 \otimes Fil_0^2}} \\
 & (E_0^1, \theta_0^1) \otimes (E_0^2, \theta_0^2) & & \dots.
 \end{array}$$

In the next step, we are going to show the above flow is semistable, that is, each Higgs term in bottom is semistable. In particular, $(E_0^1, \theta_0^1) \otimes (E_0^2, \theta_0^2)$ is semistable. It follows that $(E^1, \nabla^1) \otimes (E^2, \nabla^2)$ is semistable, as desired.

Step 4. Set $r_i = \text{rk}(E^i)$ and $d_i = \text{deg}(E^i)$. In case $r_i | d_i$, one may twist (E_0^i, θ_0^i) with some invertible sheaf of degree $-d_i/r_i$. The resulting logarithmic Higgs bundle is still semistable and of degree zero. As $r_1 + r_2 \leq p+1$, we may assume that $p \geq 3$ and $r_i \leq p-1$ for each i . After taking the base change via a certain cyclic cover of degree r_i , one may reduce the problem to the situation that $r_i | d_i$. Therefore, we may assume $\text{deg} E_0^i = 0$ and consequently $\text{deg} E_j^i = 0$ for $i = 1, 2$ and $j \geq 0$. Since the ranks and the degrees of $\{E_j^i\}_j$ are constant for $i = 1, 2$, they form *bounded* families. It follows that $\{E_j^1 \otimes E_j^2\}_j$ is also a bounded family. In particular, the degrees of subsheaves of $\{E_j^1 \otimes E_j^2\}_j$ have an upper bound by a theorem of Grothendieck (see e.g. [M, Proposition 3.12]). But then, if there were a θ -invariant subsheaf $F \subset E_0^1 \otimes E_0^2$ of positive degree, then by the flow it would give rise to a θ -invariant subsheaf of $E_j^1 \otimes E_j^2$, whose degree exceeds the upper bound when j becomes large enough. Contradiction.

Step 5. The final step reduces the general case to the previous one by using a Jordan-Hölder (JH) filtration attached to a semistable parabolic λ -connection. Let

$$0 = (E_{\bullet}^{i,0}, \nabla^i) \subset (E_{\bullet}^{i,1}, \nabla^i) \subset \dots \subset (E_{\bullet}^{i,m_i}, \nabla^i) = (E_{\bullet}^i, \nabla^i),$$

be a JH filtration of $(E_{\bullet}^i, \nabla^i)$, $i = 1, 2$, that is, each subquotient

$$\mathrm{Gr}_{E_{\bullet}^i}^j := (E_{\bullet}^{i,j}/E_{\bullet}^{i,j-1}, \nabla^i), \quad i = 1, 2, \quad 1 \leq j \leq m_i$$

is μ_L -stable of slope $\mu_L(E_{\bullet}^i)$. Then we define a filtration of parabolic λ -connections over the tensor product as follows:

$$0 = (F_{\bullet}^0, \nabla) \subset (F_{\bullet}^1, \nabla) \subset \cdots \subset (F_{\bullet}^{m_1 m_2}, \nabla) = (E_{\bullet}^1, \nabla^1) \otimes (E_{\bullet}^2, \nabla^2),$$

where for $0 \leq i \leq m_1 - 1$ and $1 \leq j \leq m_2$

$$F_{\bullet}^{im_2+j} = E_{\bullet}^{1,i} \otimes E_{\bullet}^2 + E_{\bullet}^{1,i+1} \otimes E_{\bullet}^{2,j}.$$

One verifies that

$$(F_{\bullet}^j/F_{\bullet}^{j-1}, \nabla) \cong \mathrm{Gr}_{E_{\bullet}^1}^{\lfloor \frac{j}{m_2} \rfloor} \otimes \mathrm{Gr}_{E_{\bullet}^2}^{j - (\lfloor \frac{j}{m_2} \rfloor - 1)m_2}.$$

By the previous discussion, $(F_{\bullet}^j/F_{\bullet}^{j-1}, \nabla)$ is μ_L -semistable of slope

$$\begin{aligned} \mu_L \left(\mathrm{Gr}_{E_{\bullet}^1}^{\lfloor \frac{j}{m_2} \rfloor} \otimes \mathrm{Gr}_{E_{\bullet}^2}^{j - (\lfloor \frac{j}{m_2} \rfloor - 1)m_2} \right) &= \mu_L \left(\mathrm{Gr}_{E_{\bullet}^1}^{\lfloor \frac{j}{m_2} \rfloor} \right) + \mu_L \left(\mathrm{Gr}_{E_{\bullet}^2}^{j - (\lfloor \frac{j}{m_2} \rfloor - 1)m_2} \right) \\ &= \mu_L(E_{\bullet}^1) + \mu_L(E_{\bullet}^2), \end{aligned}$$

where the first equality follows from Lemma 2.8. It follows that $(E_{\bullet}^1, \nabla^1) \otimes (E_{\bullet}^2, \nabla^2)$ is μ_L -semistable. \square

Acknowledgements. We are grateful to H el ene Esnault for many helpful comments and valuable suggestions. Especially, she pointed out to us an error in our early argument about the Mehta-Ramanathan type theorem for parabolic λ -connections (see footnote 2) as well as an error in Lemma 3.2.

REFERENCES

- [A] D. Arapura, Kodaira–Saito vanishing via Higgs bundles in positive characteristic, *J. Reine Angew. Math.*, 755, 293–312, 2019.
- [BP] V. Balaji, A.J. Parameswaran, Tensor product theorem for Hitchin pairs, an algebraic approach, *Annales de l’Institut Fourier*, 61(6):2361–2403, 2011.
- [Bi] I. Biswas, Parabolic bundles as orbifold bundles, *Duke Mathematical Journal*, Vol. 88, No. 2. 1997.
- [EG] H. Esnault, M. Groechenig, Rigid connections and F -isocrystals, *Acta Mathematica*, 225(1), 103–158, 2020.
- [H] R. Hartshorne, Algebraic geometry, Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.
- [HL] D. Huybrechts, M. Lehn, The geometry of moduli spaces of sheaves, Cambridge Mathematical Library. Cambridge University Press, 2edition, 2010.
- [IS] J. Iyer, C. Simpson, A relation between the parabolic Chern characters of the de Rham bundles, *Math. Ann.* Vol 338, 347–383, 2007.
- [KS20] R. Krishnamoorthy, M. Sheng, Periodic de Rham bundles over curves, arXiv:2011.03268, 2020.
- [LSZ15] G. Lan, M. Sheng, K. Zuo, Nonabelian Hodge theory in positive characteristic via exponential twisting, *Math. Res. Lett.* 22 (2015), no. 3, 859–879.
- [LSZ19] G. Lan, M. Sheng, K. Zuo, Semistable Higgs bundles, periodic Higgs bundles and representations of algebraic fundamental groups, *Journal of the European Mathematical Society*, vol. 21, 3053–3112, 2019.

- [LSYZ] G. Lan, M. Sheng, Y. Yang, K. Zuo, Uniformization of p-adic curves via Higgs-de Rham flows, *J. Reine Angew. Math.* 747 (2019), 63-108.
- [La14] A. Langer, Semistable modules over Lie algebroids in positive characteristic, *Doc. Math.*, vol 19, 509-540, 2014.
- [La15] A. Langer, Bogomolov's inequality for Higgs sheaves in positive characteristic, *Invent. Math.*, 199:889-920, 2015.
- [La16] A. Langer, The Bogomolov–Miyaoaka–Yau inequality for logarithmic surfaces in positive characteristic, *Duke Mathematical Journal*, 165(14):2737-2769, 2016.
- [N] N. Nitsure, Schematic Harder-Narasimhan stratification. *Internat. J. Math.* 22 (2011), no. 10, 1365-1373.
- [M] M. Maruyama, Moduli spaces of stable sheaves on schemes: restriction theorems, boundedness and the GIT construction, volume MSJ Memoirs 33, Mathematical Society of Japan, Tokyo, 2016.
- [MY] M. Maruyama, K. Yokogawa, Moduli of parabolic stable sheaves, *Mathematische Annalen*, 293, no. 1, 77-99, 1992.
- [MR] V. B. Mehta, A. Ramanathan, Semistable sheaves on projective varieties and their restriction to curves, *Math. Ann.*, 258(3):213-224, 1982.
- [Mo] T. Mochizuki, Kobayashi-Hitchin correspondence for tame harmonic bundles and an application. *Astérisque* No. 309 (2006).
- [Sh] S. Shatz, The decomposition and specialization of algebraic families of vector bundles. *Compositio Math.* 35 (1977), no. 2, 163-187.
- [Si] C. Simpson, Higgs bundles and local systems, *Inst. Hautes Études Sci. Publ. Math.* No. 75 (1992), 5-95.

Email address: msheng@ustc.edu.cn

Email address: jianpw@ustc.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI, 230026, CHINA