

LIMIT VARIETIES OF MONOIDS SATISFYING A CERTAIN IDENTITY

SERGEY V. GUSEV, YU XIAN LI, AND WEN TING ZHANG

ABSTRACT. A *limit variety* is a variety that is minimal with respect to being non-finitely based. Since the turn of the millennium, much attention has been given to the classification of limit varieties of aperiodic monoids. Seven explicit examples have so far been found, and the task of locating other examples has recently been reduced to two subproblems, one of which is concerned with monoids that satisfy the identity $xsxt \approx xsxtx$. In the present article, we provide a complete solution to this subproblem by showing that there are precisely two limit varieties that satisfy this identity. One of them turns out to be the first example having infinitely many subvarieties.

It is also deduced that the variety generated by any monoid of order five or less contains at most countably many subvarieties.

1. INTRODUCTION

A variety of algebras is *finitely based* (FB) if it has a finite basis of identities; otherwise, it is *non-finitely based* (NFB). The *finite basis problem*—to determine which varieties are finitely based—has been the focus of investigation for many years. In the 1930s, Neumann [26] questioned if every variety of groups is FB, and it was not until 1970 when the existence of NFB examples was independently confirmed by Adyan [1], Ol’shanskiĭ [28], and Vaughan-Lee [35]. The finite basis problem for varieties of semigroups and of monoids gained much interest in the 1960s after Perkins [29] exhibited the first examples of NFB varieties generated by a finite semigroup. For more information on the finite basis problem for these varieties, refer to Gupta and Krasilnikov [5] for varieties of groups and to Volkov [36] for varieties of semigroups and of monoids.

A FB variety satisfies the stronger property of being *hereditary finitely based* (HFB) if all its subvarieties are FB. A *limit variety* is a NFB variety whose proper subvarieties are all FB. By Zorn’s lemma, every NFB variety contains some limit subvariety; in other words, a variety is HFB if and only if it excludes all limit varieties. Therefore, classifying HFB varieties in a certain sense reduces to classifying limit varieties. But finding explicit examples of limit varieties turns out to be highly nontrivial. For instance, no explicit example of limit variety of groups is known so far, even though uncountably many of them exist [15]. Locating such an example

2010 *Mathematics Subject Classification.* 20M07.

Key words and phrases. Monoid, variety, hereditary finitely based variety, limit variety, finite basis problem.

The first author is supported by the Ministry of Science and Higher Education of the Russian Federation (project FEUZ-2020-0016). The second and the third authors are partially supported by the National Natural Science Foundation of China (No. 12171213, 11771191) and the Natural Science Foundation of Gansu Province (No. 20JR5RA275).

remains one of the foremost unsolved problems in the theory of varieties of groups; see Gupta and Krasilnikov [5].

One of the main goals of the present paper is to study limit varieties of monoids, considered as algebras of type $\langle 2, 0 \rangle$. A complete classification of all limit varieties of monoids is highly infeasible since that would include a description of all limit varieties of groups. Therefore it is natural to focus on the class \mathbf{A} of *aperiodic* monoids, that is, monoids whose subgroups are all trivial. The first explicit examples of limit subvarieties of \mathbf{A} were exhibited by Jackson [11] in the early 2000s. Since then, limit subvarieties of \mathbf{A} have received much attention and a few more examples have been found; see, for example, Gusev [6, 7], Gusev and Sapir [9], Lee [18, 19], Sapir [32, 33], Zhang [37], and Zhang and Luo [38].

Presently, seven explicit examples of limit subvarieties of \mathbf{A} are known, and it follows from Gusev and Sapir [9, Sorting Lemma 2] that any other limit subvariety of \mathbf{A} is contained in one of the following varieties or their duals:

$$\mathbf{P} = \text{var}\{xsxt \approx xsxtx\} \quad \text{and} \quad \mathbf{Q} = \text{var}\{xtx \approx txt^2, xy^2tx \approx (xy)^2tx\}.$$

Therefore, to completely classify all limit subvarieties of \mathbf{A} , it suffices to consider only subvarieties of \mathbf{P} and of \mathbf{Q} . Sapir [33] made some progress by exhibiting three new examples of limit subvarieties of \mathbf{Q} .

The present paper is concerned with the variety \mathbf{P} , where the main objective is to provide a complete description of all its limit subvarieties (Theorem 5.11). Specifically, we present two new limit varieties of monoids and show that they exhaust all limit subvarieties of \mathbf{P} . One of these limit varieties contains countably infinitely many subvarieties (Corollary 5.10)—a result that is quite surprising given that every limit variety of monoids previously found is *small* in the sense that it contains only finitely many subvarieties.

Our main result also implies an important fundamental result concerning varieties generated by a monoid of small order. Finite members from several classical classes of algebras, such as groups [27] and associative rings [16, 25], generate small HFB varieties. But this result does not hold for finite semigroups or monoids. The smallest monoids currently known to generate a variety with continuum many subvarieties are of order six; the well-known Brandt monoid

$$B_2^1 = \langle a, b, 1 \mid aba = a, bab = b, a^2 = b^2 = 0 \rangle = \{a, b, ab, ba, 0, 1\}$$

is one such example [12, 14]. As for monoids of order five or less, up to isomorphism and anti-isomorphism, every one of them, with the exception of

$$P_2^1 = \langle a, b, 1 \mid a^2 = ab = a, b^2a = b^2 \rangle = \{a, b, ba, b^2, 1\},$$

generates a small HFB variety [24]. The variety \mathbf{P}_2^1 generated by P_2^1 is not small, but not much else is known about it. An answer to the following question is thus desirable.

Question 1 (Jackson and Lee [12, Question 6.1]). Does the variety \mathbf{P}_2^1 contain uncountable many subvarieties?

The monoid P_2^1 satisfies the identity $xsxt \approx xsxtx$ and so \mathbf{P}_2^1 is a subvariety of \mathbf{P} . It follows from our classification of limit subvarieties of \mathbf{P} that \mathbf{P}_2^1 is HFB (Proposition 6.1). Consequently, the variety generated by any monoid of order five or less is HFB and so contains at most countably many subvarieties.

Remark 1.1. It is relevant to note that P_2^1 was originally investigated as a semi-group by Lee, who first questioned if it generates a HFB variety of semigroups; see Edmunds et al. [4, Question 4.2] and Lee [21, Problem 1.4]. The answer to this question, however, remains elusive.

This article consists of six sections. Background information and some basic results are first given in Section 2. Then several important results on the variety \mathbf{P} and its subvarieties are established in Section 3. Results that are crucial to the proof of the main results are established in Section 4. Limit subvarieties of \mathbf{P} are then classified in Section 5, while results concerning subvarieties of \mathbf{P}_2^1 are established in Section 6.

Many identities will be introduced and used throughout this article. For the reader's convenience, these identities are collected in the appendix for quick referencing.

2. PRELIMINARIES

Acquaintance with rudiments of universal algebra is assumed of the reader. Refer to Burris and Sankappanavar [3] for more information.

2.1. Words, identities, and deduction. Let \mathfrak{A}^* denote the free monoid over a countably infinite alphabet \mathfrak{A} . Elements of \mathfrak{A} are called *letters* and elements of \mathfrak{A}^* are called *words*. The empty word, denoted by λ , is the identity element of \mathfrak{A}^* .

An identity is written as $\mathbf{u} \approx \mathbf{v}$, where $\mathbf{u}, \mathbf{v} \in \mathfrak{A}^*$; it is *non-trivial* if $\mathbf{u} \neq \mathbf{v}$. An identity $\mathbf{u} \approx \mathbf{v}$ is *directly deducible* from an identity $\mathbf{s} \approx \mathbf{t}$ if there exist some words $\mathbf{a}, \mathbf{b} \in \mathfrak{A}^*$ and substitution $\phi: \mathfrak{A} \rightarrow \mathfrak{A}^*$ such that $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\phi(\mathbf{s})\mathbf{b}, \mathbf{a}\phi(\mathbf{t})\mathbf{b}\}$. A non-trivial identity $\mathbf{u} \approx \mathbf{v}$ is *deducible* from a set Σ of identities if there exists some finite sequence $\mathbf{u} = \mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_m = \mathbf{v}$ of distinct words such that each identity $\mathbf{w}_i \approx \mathbf{w}_{i+1}$ is directly deducible from some identity in Σ .

The following assertion is a specialization for monoids of a well-known universal-algebraic fact (see Burris and Sankappanavar [3, Theorem II.14.19]).

Proposition 2.1. *Let \mathbf{V} be the variety defined by some set Σ of identities. Then \mathbf{V} satisfies an identity $\mathbf{u} \approx \mathbf{v}$ if and only if $\mathbf{u} \approx \mathbf{v}$ is deducible from Σ . \square*

2.2. k -decomposition of a word. Here we introduce a series of notions such as k -divider, k -block, k -decomposition, and some others, which appeared in Gusev and Vernikov [10, Chapter 3].

The *content* of a word \mathbf{w} , denoted by $\text{con}(\mathbf{w})$, is the set of all letters occurring in \mathbf{w} . A letter is *simple* [respectively, *multiple*] *in a word \mathbf{w}* if it occurs in \mathbf{w} once [respectively, at least twice]. The set of all simple [respectively, multiple] letters in a word \mathbf{w} is denoted by $\text{sim}(\mathbf{w})$ [respectively, $\text{mul}(\mathbf{w})$]. If \mathbf{w} is a word and X is a set of letters then \mathbf{w}_X denotes the word obtained from \mathbf{w} by deleting all letters from X . If $X = \{x\}$, then we write \mathbf{w}_x rather than $\mathbf{w}_{\{x\}}$. For a word \mathbf{w} and letters $x_1, x_2, \dots, x_k \in \text{con}(\mathbf{w})$, let $\mathbf{w}(x_1, x_2, \dots, x_k)$ denote the word obtained from \mathbf{w} by retaining the letters x_1, x_2, \dots, x_k . Equivalently,

$$\mathbf{w}(x_1, x_2, \dots, x_k) = \mathbf{w}_{\text{con}(\mathbf{w}) \setminus \{x_1, x_2, \dots, x_k\}}.$$

Let \mathbf{w} be a word such that $\text{sim}(\mathbf{w}) = \{t_1, t_2, \dots, t_m\}$. We may assume without loss of generality that $\mathbf{w}(t_1, t_2, \dots, t_m) = t_1 t_2 \cdots t_m$. Then

$$(2.1) \quad \mathbf{w} = t_0 \mathbf{w}_0 t_1 \mathbf{w}_1 \cdots t_m \mathbf{w}_m,$$

where $\mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_m \in \mathfrak{A}^*$ and $t_0 = \lambda$. The words $\mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_m$ are *0-blocks* of \mathbf{w} , while t_0, t_1, \dots, t_m are *0-dividers* of \mathbf{w} . The representation of the word \mathbf{w} as a product of alternating 0-dividers and 0-blocks, starting with the 0-divider t_0 and ending with the 0-block \mathbf{w}_m , is called the *0-decomposition* of \mathbf{w} .

For $k \geq 1$, we define the *k-decomposition* of a word \mathbf{w} recursively as follows. Suppose that (2.1) is the $(k-1)$ -decomposition of \mathbf{w} with $(k-1)$ -dividers t_0, t_1, \dots, t_m and $(k-1)$ -blocks $\mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_m$. For any $i = 0, 1, \dots, m$, let $s_{i1}, s_{i2}, \dots, s_{ir_i}$ be all the letters in $\text{sim}(\mathbf{w}_i)$ that do not occur to the left of \mathbf{w}_i . We may assume that $\mathbf{w}_i(s_{i1}, s_{i2}, \dots, s_{ir_i}) = s_{i1}s_{i2} \cdots s_{ir_i}$. Then

$$(2.2) \quad \mathbf{w}_i = \mathbf{v}_{i0}s_{i1}\mathbf{v}_{i1}s_{i2}\mathbf{v}_{i2} \cdots s_{ir_i}\mathbf{v}_{ir_i}$$

for some $\mathbf{v}_{i0}, \mathbf{v}_{i1}, \dots, \mathbf{v}_{ir_i} \in \mathfrak{A}^*$. Put $s_{i0} = t_i$. Then the words $\mathbf{v}_{i0}, \mathbf{v}_{i1}, \dots, \mathbf{v}_{ir_i}$ are *k-blocks* of \mathbf{w} , while the letters $s_{i0}, s_{i1}, \dots, s_{ir_i}$ are *k-dividers* of \mathbf{w} .

As noted in Gusev and Vernikov [10, Remark 3.1], only the first occurrence of a letter in a given word might be a *k-divider* of the word for some k . In view of this observation, we use below an expression like “a letter x is (or is not) a *k-divider* of a word \mathbf{w} ” to mean that the first occurrence of x in \mathbf{w} has the specified property.

Now decompose every $(k-1)$ -block \mathbf{w}_i from (2.1) in the form (2.2). Then the word \mathbf{w} in (2.1) is decomposed as a product of alternating *k-dividers* and *k-blocks*, starting with the *k-divider* $s_{00} = t_0$ and ending with the *k-block* \mathbf{v}_{mr_m} . This is called the *k-decomposition* of \mathbf{w} .

Remark 2.2. Since the length of the word \mathbf{w} is finite, there is a number k such that the *k-decomposition* of \mathbf{w} coincides with its *n-decomposition* for each $n > k$.

We say that the *k-decomposition* of \mathbf{w} is *maximal* if it coincides with the *n-decomposition* of \mathbf{w} for all $n > k$.

Let $\text{occ}_x(\mathbf{w})$ denote the number of occurrences of a letter x in a word \mathbf{w} . For a given word \mathbf{w} , a letter $x \in \text{con}(\mathbf{w})$, a natural number $i \leq \text{occ}_x(\mathbf{w})$ and an integer $k \geq 0$, we denote by $h_i^k(\mathbf{w}, x)$ the right-most *k-divider* of \mathbf{w} that precedes to the *i*th occurrence of x in \mathbf{w} .

For a given word \mathbf{w} and a letter $x \in \text{con}(\mathbf{w})$, the *depth* of x in \mathbf{w} is the number $D(\mathbf{w}, x)$ defined as follows. If $x \in \text{sim}(\mathbf{w})$, then we put $D(\mathbf{w}, x) = 0$. Suppose now that $x \in \text{mul}(\mathbf{w})$. If there is a natural k such that the first and the second occurrences of x in \mathbf{w} lie in different $(k-1)$ -blocks of \mathbf{w} , then the depth of x in \mathbf{w} is equal to the minimal number k with such a property. If, for any natural k , the first and the second occurrences of x in \mathbf{w} lie in the same *k-block* of \mathbf{w} , then we put $D(\mathbf{w}, x) = \infty$. In other words, $D(\mathbf{w}, x) = k$ if and only if $h_1^{k-1}(\mathbf{w}, x) \neq h_2^{k-1}(\mathbf{w}, x)$ and k is the least number with such a property, while $D(\mathbf{w}, x) = \infty$ if and only if $h_1^{k-1}(\mathbf{w}, x) = h_2^{k-1}(\mathbf{w}, x)$ for any k .

Lemma 2.3 (Gusev and Vernikov [10, Lemma 3.7]). *A letter t is a *k-divider* of a word \mathbf{w} if and only if $D(\mathbf{w}, t) \leq k$. \square*

The *i*th occurrence of a letter x in a word \mathbf{w} is denoted by ${}_i\mathbf{w}x$. We write $({}_i\mathbf{w}x) < ({}_j\mathbf{w}y)$ to indicate that within \mathbf{w} , the *i*th occurrence of x precedes the *j*th occurrence of y .

Lemma 2.4 (Gusev and Vernikov [10, Lemma 3.9]). *Let \mathbf{w} be a word, x be a multiple letter in \mathbf{w} with $D(\mathbf{w}, x) = k$ and t be a $(k-1)$ -divider of \mathbf{w} .*

- (i) *If $t = h_2^{k-1}(\mathbf{w}, x)$, then $({}_1\mathbf{w}x) < ({}_1\mathbf{w}t)$.*

- (ii) If $({}_{1\mathbf{w}}x) < ({}_{1\mathbf{w}}t) < ({}_{2\mathbf{w}}x)$, then $D(\mathbf{w}, t) = k - 1$. Further, if $k > 1$, then $({}_{2\mathbf{w}}x) < ({}_{2\mathbf{w}}t)$. \square

Lemma 2.5 (Gusev and Vernikov [10, Lemma 3.13]). *Let \mathbf{w} be a word, $r > 1$ be a number and y be a letter such that $D(\mathbf{w}, y) = r - 2$. If $({}_{1\mathbf{w}}z) < ({}_{1\mathbf{w}}y)$ for some letter z with $D(\mathbf{w}, z) \geq r$, then $({}_{2\mathbf{w}}z) < ({}_{1\mathbf{w}}y)$.* \square

In the following, to facilitate understanding of the form of words, we will sometimes explicitly display the occurrence of a letter, for example,

$$\mathbf{w} = x_1 \overset{(1)}{x_2} \overset{(2)}{x_1} \overset{(1)}{x_3} \overset{(2)}{x_2} \overset{(3)}{x_1}.$$

Lemma 2.6 (Gusev and Vernikov [10, Lemma 3.14]). *Let $\mathbf{u} \approx \mathbf{v}$ be an identity and ℓ be a natural number. Suppose that*

$$(2.3) \quad \text{sim}(\mathbf{u}) = \text{sim}(\mathbf{v}) \text{ and } \text{mul}(\mathbf{u}) = \text{mul}(\mathbf{v}),$$

$$(2.4) \quad h_i^{\ell-1}(\mathbf{u}, x) = h_i^{\ell-1}(\mathbf{v}, x) \text{ for } i = 1, 2 \text{ and all } x \in \text{con}(\mathbf{u}),$$

and there is a letter x_ℓ such that $D(\mathbf{u}, x_\ell) = \ell$. Then there are letters $x_0, x_1, \dots, x_{\ell-1}$ such that $D(\mathbf{u}, x_s) = D(\mathbf{v}, x_s) = s$ for any $0 \leq s < \ell$ and the identity $\mathbf{u} \approx \mathbf{v}$ has the form

$$\begin{aligned} & \mathbf{u}_{2\ell+1} \overset{(1)}{x_\ell} \mathbf{u}_{2\ell} \overset{(1)}{x_{\ell-1}} \mathbf{u}_{2\ell-1} \overset{(2)}{x_\ell} \mathbf{u}_{2\ell-2} \overset{(1)}{x_{\ell-2}} \mathbf{u}_{2\ell-3} \overset{(2)}{x_{\ell-1}} \mathbf{u}_{2\ell-4} \overset{(1)}{x_{\ell-3}} \\ & \cdot \mathbf{u}_{2\ell-5} \overset{(2)}{x_{\ell-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0 \\ \approx & \mathbf{v}_{2\ell+1} \overset{(1)}{x_\ell} \mathbf{v}_{2\ell} \overset{(1)}{x_{\ell-1}} \mathbf{v}_{2\ell-1} \overset{(2)}{x_\ell} \mathbf{v}_{2\ell-2} \overset{(1)}{x_{\ell-2}} \mathbf{v}_{2\ell-3} \overset{(2)}{x_{\ell-1}} \mathbf{v}_{2\ell-4} \overset{(1)}{x_{\ell-3}} \\ & \cdot \mathbf{v}_{2\ell-5} \overset{(2)}{x_{\ell-2}} \cdots \mathbf{v}_4 \overset{(1)}{x_1} \mathbf{v}_3 \overset{(2)}{x_2} \mathbf{v}_2 \overset{(1)}{x_0} \mathbf{v}_1 \overset{(2)}{x_1} \mathbf{v}_0 \end{aligned}$$

for some $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2r+1}, \mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2r+1} \in \mathfrak{A}^*$. \square

Remark 2.7. Analyzing the proof of Lemma 3.14 in Gusev and Vernikov [10], one can notice that, in Lemma 2.6, the letters $x_0, x_1, \dots, x_{\ell-1}$ can be chosen so that \mathbf{u}_{2s+2} and \mathbf{v}_{2s+2} do not contain any s -dividers of \mathbf{u} and \mathbf{v} , respectively, for any $s = 0, 1, \dots, \ell - 1$. \square

Evidently, if $\mathbf{u} = \mathbf{v}$, then (2.3) and (2.4) are true for all ℓ . So, one can apply Lemma 2.6 and Remark 2.7 for the trivial identity and obtain the following corollary.

Corollary 2.8. *Let \mathbf{u} be a word and k be a natural number. Suppose that there is a letter x_k such that $D(\mathbf{u}, x_k) = k$. Then there exist letters x_0, x_1, \dots, x_{k-1} such that the word \mathbf{u} has the form*

$$(2.5) \quad \begin{aligned} & \mathbf{u}_{2k+1} \overset{(1)}{x_k} \mathbf{u}_{2k} \overset{(1)}{x_{k-1}} \mathbf{u}_{2k-1} \overset{(2)}{x_k} \mathbf{u}_{2k-2} \overset{(1)}{x_{k-2}} \mathbf{u}_{2k-3} \overset{(2)}{x_{k-1}} \mathbf{u}_{2k-4} \overset{(1)}{x_{k-3}} \\ & \cdot \mathbf{u}_{2k-5} \overset{(2)}{x_{k-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0 \end{aligned}$$

for some $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2k+1} \in \mathfrak{A}^*$ such that $D(\mathbf{u}, x_s) = s$ and \mathbf{u}_{2s+2} does not contain any s -dividers of \mathbf{u} for any $s = 0, 1, \dots, k - 1$. \square

Lemma 2.9. *Let k, m be natural numbers and \mathbf{u} be a word of the form*

$$\begin{aligned} & \mathbf{u}_{2k+2} \overset{(1)}{y_m} \mathbf{u}_{2k+1} \overset{(1)}{x_k} \mathbf{u}_{2k} \overset{(1)}{x_{k-1}} \mathbf{u}_{2k-1} \overset{(2)}{x_k} \mathbf{u}_{2k-2} \overset{(1)}{x_{k-2}} \mathbf{u}_{2k-3} \overset{(2)}{x_{k-1}} \\ & \cdot \mathbf{u}_{2k-4} \overset{(1)}{x_{k-3}} \mathbf{u}_{2k-5} \overset{(2)}{x_{k-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0, \end{aligned}$$

where $D(\mathbf{u}, y_m) = m$ and $D(\mathbf{u}, x_s) = s$ for any $0 \leq s \leq k$. If \mathbf{u}_{2k+1} does not contain any $(m-1)$ -divider of \mathbf{u} , then ${}_{2\mathbf{u}}y_m$ occurs in one of the subwords \mathbf{u}_{2m} , \mathbf{u}_{2m-1} or \mathbf{u}_{2m-2} of \mathbf{u} .

Proof. Since $D(\mathbf{u}, y_m) = m > 0$, we have $y_m \in \text{mul}(\mathbf{u})$. Let $y_{m-1} = h_2^{m-1}(\mathbf{u}, y_m)$. Since $({}_{1\mathbf{u}}y_m) < ({}_{1\mathbf{u}}y_{m-1}) < ({}_{2\mathbf{u}}y_m)$ by Lemma 2.4(i), the letter ${}_{1\mathbf{u}}y_{m-1}$ does not occur in \mathbf{u}_{2k+2} . By the hypothesis, ${}_{1\mathbf{u}}y_{m-1}$ does not occur in \mathbf{u}_{2k+1} . Since $({}_{1\mathbf{u}}y_{m-1}) < ({}_{2\mathbf{u}}y_m)$, the letter ${}_{2\mathbf{u}}y_m$ occurs in neither \mathbf{u}_{2k+2} nor \mathbf{u}_{2k+1} . Then $({}_{1\mathbf{u}}x_k) < ({}_{2\mathbf{u}}y_m)$. It follows that $m \leq k+1$. If $m = k+1$, then ${}_{2\mathbf{u}}y_m$ occurs in $\mathbf{u}_{2m-2} = \mathbf{u}_{2k}$ because $h_1^k(\mathbf{u}, y_m) \neq h_2^k(\mathbf{u}, y_m)$ otherwise. If $m = k$, then ${}_{2\mathbf{u}}y_m$ occurs in either \mathbf{u}_{2k} or \mathbf{u}_{2k-1} or \mathbf{u}_{2k-2} because $h_1^{k-1}(\mathbf{u}, y_m) \neq h_2^{k-1}(\mathbf{u}, y_m)$ otherwise. If $k > m$, then ${}_{1\mathbf{u}}y_{m-1}$ and so ${}_{2\mathbf{u}}y_m$ do not occur in both \mathbf{u}_{2k} and \mathbf{u}_{2k-1} because $h_1^{m-1}(\mathbf{u}, x_k) \neq h_2^{m-1}(\mathbf{u}, x_k)$ otherwise. Further, if $k > m+1$, then ${}_{1\mathbf{u}}y_{m-1}$ and so ${}_{2\mathbf{u}}y_m$ do not occur in the subwords \mathbf{u}_{2r+1} , \mathbf{u}_{2r} and \mathbf{u}_{2r-1} of \mathbf{u} for any $r = m+1, m+2, \dots, k-1$ because $h_1^{r-2}(\mathbf{u}, x_r) \neq h_2^{r-2}(\mathbf{u}, x_r)$ otherwise. Finally, if $m > 1$, then $({}_{2\mathbf{u}}y_m) < ({}_{1\mathbf{u}}x_{m-2})$ by Lemma 2.5. This means that if $m > 1$, then ${}_{2\mathbf{u}}y_m$ does not occur in \mathbf{u}_{2m-3} , \mathbf{u}_{2m-4} , \dots , \mathbf{u}_0 . Therefore, ${}_{2\mathbf{u}}y_m$ occurs in either \mathbf{u}_{2m} or \mathbf{u}_{2m-1} or \mathbf{u}_{2m-2} . \square

2.3. Some known facts. For an identity system Σ , we denote by $\text{var } \Sigma$ the variety of monoids defined by Σ . It is well known that the variety

$$\mathbf{LRB} = \text{var}\{xy \approx yx\}$$

of left regular bands is generated by the monoid obtained by adjoining an identity element to the left zero semigroup of order two. The *initial part* of a word \mathbf{w} , denoted by $\text{ini}(\mathbf{w})$, is the word obtained from \mathbf{w} by retaining the first occurrence of each letter. The following statement is well known and can be easily verified.

Lemma 2.10. *A non-trivial identity $\mathbf{u} \approx \mathbf{v}$ holds in the variety \mathbf{LRB} if and only if $\text{ini}(\mathbf{u}) = \text{ini}(\mathbf{v})$.* \square

We use the standard symbol \mathbb{N} to denote the set of all natural numbers. For any $s \in \mathbb{N}$ and $1 \leq q \leq s$, put

$$\mathbf{b}_{s,q} = x_{s-1}x_sx_{s-2}x_{s-1} \cdots x_{q-1}x_q.$$

For brevity, we will write \mathbf{b}_s rather than $\mathbf{b}_{s,1}$. We put also $\mathbf{b}_0 = \lambda$ for convenience. We introduce the following four countably infinite series of identities:

$$\begin{aligned} \alpha_k &: x_k y_k x_{k-1} x_k y_k \mathbf{b}_{k-1} \approx y_k x_k x_{k-1} x_k y_k \mathbf{b}_{k-1}, \\ \beta_k &: x x_k x \mathbf{b}_k \approx x_k x^2 \mathbf{b}_k, \\ \gamma_k &: y_1 y_0 x_k y_1 \mathbf{b}_k \approx y_1 y_0 y_1 x_k \mathbf{b}_k, \\ \delta_k^m &: y_{m+1} y_m x_k y_{m+1} \mathbf{b}_{k,m} y_m \mathbf{b}_{m-1} \approx y_{m+1} y_m y_{m+1} x_k \mathbf{b}_{k,m} y_m \mathbf{b}_{m-1}, \end{aligned}$$

where $k \in \mathbb{N}$ and $1 \leq m \leq k$. The subvariety of a variety \mathbf{V} defined by a set Σ of identities is denoted by $\mathbf{V}\Sigma$. Let

$$\begin{aligned} \mathbf{F} &= \mathbf{P}\{x^2 y^2 \approx y^2 x^2\}, \\ \mathbf{F}_k &= \mathbf{F}\{\alpha_k\}, \\ \mathbf{H}_k &= \mathbf{F}\{\beta_k\}, \\ \mathbf{I}_k &= \mathbf{F}\{\gamma_k\}, \\ \text{and } \mathbf{J}_k^m &= \mathbf{F}\{\delta_k^m\}, \end{aligned}$$

where $k \in \mathbb{N}$ and $1 \leq m \leq k$. The trivial variety of monoids is denoted by \mathbf{T} . Let \mathbf{SL} denote the variety of all semilattice monoids. Put

$$\begin{aligned}\mathbf{C} &= \text{var}\{x^2 \approx x^3, xy \approx yx\}, \\ \mathbf{D} &= \text{var}\{x^2 \approx x^3, x^2y \approx xyx \approx yx^2\}, \\ \mathbf{E} &= \text{var}\{x^2 \approx x^3, x^2y \approx xyx, x^2y^2 \approx y^2x^2\}.\end{aligned}$$

The subvariety lattice of a monoid variety \mathbf{V} is denoted by $\mathfrak{L}(\mathbf{V})$.

Lemma 2.11 (Gusev and Vernikov [10, Proposition 6.1]). *The chain*

$$\begin{aligned}\mathbf{T} \subset \mathbf{SL} \subset \mathbf{C} \subset \mathbf{D} \subset \mathbf{E} \subset \mathbf{F}_1 \subset \mathbf{H}_1 \subset \mathbf{I}_1 \subset \mathbf{J}_1^1 \\ \subset \mathbf{F}_2 \subset \mathbf{H}_2 \subset \mathbf{I}_2 \subset \mathbf{J}_2^1 \subset \mathbf{J}_2^2 \\ \vdots \\ \subset \mathbf{F}_k \subset \mathbf{H}_k \subset \mathbf{I}_k \subset \mathbf{J}_k^1 \subset \mathbf{J}_k^2 \subset \cdots \subset \mathbf{J}_k^k \\ \vdots \\ \subset \mathbf{F}\end{aligned}$$

is the lattice $\mathfrak{L}(\mathbf{F})$. □

Lemma 2.12 (Gusev and Vernikov [10, Proposition 6.9(i)]). *A non-trivial identity $\mathbf{u} \approx \mathbf{v}$ holds in the variety \mathbf{F}_k if and only if (2.3) and (2.4) hold with $\ell = k$.*

An identity $\mathbf{u} \approx \mathbf{v}$ is *k-well-balanced* if the words \mathbf{u} and \mathbf{v} have the same set of k -dividers, these k -dividers appear in \mathbf{u} and \mathbf{v} in the same order, and $i_{\mathbf{u}}x$ occurs in some k -block of \mathbf{u} if and only if $i_{\mathbf{v}}x$ occurs in the corresponding k -block of \mathbf{v} for any letter x and $i \in \{1, 2\}$.

Corollary 2.13. *Let $\mathbf{u} \approx \mathbf{v}$ be an identity of \mathbf{F}_k . Then the identity $\mathbf{u} \approx \mathbf{v}$ is $(k-1)$ -well-balanced.*

Proof. Let $t_0\mathbf{u}_0t_1\mathbf{u}_1 \cdots t_m\mathbf{u}_m$ be the $(k-1)$ -decomposition of \mathbf{u} . According to Lemma 2.12, (2.3) and (2.4) hold with $\ell = k$. Then the $(k-1)$ -decomposition of \mathbf{v} has the form $t_0\mathbf{v}_0t_1\mathbf{v}_1 \cdots t_m\mathbf{v}_m$ by Gusev and Vernikov [10, Lemma 3.8]. Suppose that $i_{\mathbf{u}}x$ occurs in the $(k-1)$ -block \mathbf{u}_j of \mathbf{u} , where $i \in \{1, 2\}$ and $j \in \{0, 1, \dots, m\}$. Then $h_i^{k-1}(\mathbf{u}, x) = t_j$. Since (2.4) holds with $\ell = k$, we have $h_i^{k-1}(\mathbf{v}, x) = t_j$ and so $i_{\mathbf{v}}x$ occurs in the $(k-1)$ -block \mathbf{v}_j of \mathbf{v} . Therefore, the identity $\mathbf{u} \approx \mathbf{v}$ is $(k-1)$ -well-balanced. □

3. SOME PROPERTIES OF THE VARIETY \mathbf{P}

Recall that the variety \mathbf{P} is defined by the identity

$$(3.1) \quad xsxt \approx xsxtx.$$

For any word \mathbf{w} , let $\text{ini}_2(\mathbf{w})$ denote the word obtained from \mathbf{w} by retaining the first and second occurrences of each letter.

The following result is evident.

Lemma 3.1. *For any $\mathbf{w} \in \mathfrak{A}^*$, the identity $\mathbf{w} \approx \text{ini}_2(\mathbf{w})$ holds in the variety \mathbf{P} . □*

Let $\phi: \mathfrak{A} \rightarrow \mathfrak{A}^*$ be a substitution, $x_1, x_2, \dots, x_n \in \mathfrak{A}$ and $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n \in \mathfrak{A}^*$. For brevity, we will say that ϕ is the substitution

$$(x_1, x_2, \dots, x_n) \mapsto (\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n)$$

if $\phi(x_i) = \mathbf{w}_i$ for any $i = 1, 2, \dots, n$ and $\phi(a) = a$ otherwise.

Lemma 3.2. *Let \mathbf{V} be a subvariety of \mathbf{P} .*

- (i) *If $\mathbf{LRB} \not\subseteq \mathbf{V}$, then $\mathbf{V} \subseteq \mathbf{F}$.*
- (ii) *If $\mathbf{F}_1 \not\subseteq \mathbf{V}$, then $\mathbf{V} \subseteq \mathbf{LRB} \vee \mathbf{C} = \text{var}\{x^2 \approx x^3, x^2y \approx xyx\}$.*

Proof. (i) It follows from Lee et al. [23, Theorem 5.17] that \mathbf{V} satisfies the identity $x^2(y^2x^2)^2 \approx (y^2x^2)^2$. It is then easily shown that \mathbf{V} satisfies $x^2y^2 \approx y^2x^2$. Thus, $\mathbf{V} \subseteq \mathbf{F}$.

(ii) If $\mathbf{C} \not\subseteq \mathbf{V}$, then by Gusev and Vernikov [10, Corollary 2.6], the variety \mathbf{V} is *completely regular*, that is, it consists of unions of groups. Then \mathbf{V} satisfies the identity $x \approx x^n$ for some $n \geq 2$. Clearly, this identity together with (3.1) imply the identity $xy \approx xyx$, whence $\mathbf{V} \subseteq \mathbf{LRB}$. Therefore, we may assume that $\mathbf{C} \subseteq \mathbf{V}$. There is an identity $\mathbf{u} \approx \mathbf{v}$ of \mathbf{V} that is not satisfied by \mathbf{F}_1 . According to Gusev and Vernikov [10, Proposition 2.2] and Lemma 2.12, (2.3) holds but (2.4) with $\ell = 1$ does not. Then $h_i^0(\mathbf{u}, x) \neq h_i^0(\mathbf{v}, x)$ for some $x \in \text{con}(\mathbf{u})$ and $i \in \{1, 2\}$. If $i = 1$, then we can multiply both sides of the identity $\mathbf{u} \approx \mathbf{v}$ on the left by xt , where $t \notin \text{con}(\mathbf{u})$. So, we may assume that $i = 2$. Put $y = h_2^0(\mathbf{u}, x)$ and $z = h_2^0(\mathbf{v}, x)$. If y and z occur in \mathbf{u} and \mathbf{v} in different order, then \mathbf{V} is commutative and so $\mathbf{V} \subseteq \mathbf{C}$. Therefore, we may assume without loss of generality that $(1_{\mathbf{u}}z) < (1_{\mathbf{u}}y)$ and $(1_{\mathbf{v}}z) < (1_{\mathbf{v}}y)$. Then the identity $\mathbf{u}(x, y) \approx \mathbf{v}(x, y)$ is equivalent modulo (3.1) to $xyx \approx x^2y$. It follows from Lee [20, Lemma 3.3] that $\mathbf{LRB} \vee \mathbf{C} = \text{var}\{x^2 \approx x^3, x^2y \approx xyx\}$. So, $\mathbf{V} \subseteq \mathbf{LRB} \vee \mathbf{C}$. \square

Lemma 3.3. *Let \mathbf{V} be a subvariety of \mathbf{P} that does not contain \mathbf{F}_{k+1} . Then \mathbf{V} satisfies the identity*

$$\kappa_k : xx_kx\mathbf{b}_k \approx x^2x_k\mathbf{b}_k.$$

Proof. If $\mathbf{LRB} \not\subseteq \mathbf{V}$, then $\mathbf{V} \subseteq \mathbf{F}$ by Lemma 3.2(i). Since $\mathbf{F}_{k+1} \not\subseteq \mathbf{V}$, it follows from Lemma 2.11 that $\mathbf{V} \subseteq \mathbf{J}_k^k$. It is verified in Gusev and Vernikov [10, Lemma 6.3] that \mathbf{J}_k^k satisfies κ_k . Then \mathbf{V} satisfies κ_k as well. Therefore, we may assume that $\mathbf{LRB} \subseteq \mathbf{V}$. Further, if $\mathbf{F}_1 \not\subseteq \mathbf{V}$, then \mathbf{V} satisfies $x^2y \approx xyx$ by Lemma 3.2(ii). Clearly, $x^2y \approx xyx$ implies κ_k . Therefore, we may assume that $\mathbf{F}_1 \subseteq \mathbf{V}$.

Let r be the least number such that $\mathbf{F}_r \subseteq \mathbf{V}$. Then $\mathbf{F}_{r+1} \not\subseteq \mathbf{V}$. Evidently, $r \leq k$. There is an identity $\mathbf{u} \approx \mathbf{v}$ of \mathbf{V} that is not satisfied by \mathbf{F}_{r+1} . According to Lemma 2.12, (2.3) and (2.4) hold with $\ell = 1, 2, \dots, r$, while (2.4) with $\ell = r+1$ does not. Then $h_i^r(\mathbf{u}, x) \neq h_i^r(\mathbf{v}, x)$ for some $x \in \text{con}(\mathbf{u})$ and $i \in \{1, 2\}$. If $i = 1$, then we multiply both sides of the identity $\mathbf{u} \approx \mathbf{v}$ on the left by xt , where $t \notin \text{con}(\mathbf{u}) = \text{con}(\mathbf{v})$. So, we may assume that $i = 2$. Put $a = h_2^r(\mathbf{u}, x)$ and $b = h_2^r(\mathbf{v}, x)$. We may assume without loss of generality that $(1_{\mathbf{u}}b) < (1_{\mathbf{u}}a)$. Then $(1_{\mathbf{v}}b) < (1_{\mathbf{v}}a)$ by Lemma 2.10.

In view of Lemma 2.3, $D(\mathbf{u}, a) = s$ for some $s \leq r$. Put $x_s = a$. According to Lemma 2.6, there exist letters x_0, x_1, \dots, x_{s-1} such that $D(\mathbf{u}, x_j) = D(\mathbf{v}, x_j) = j$

for any $0 \leq j < s$ and the identity $\mathbf{u} \approx \mathbf{v}$ is of the form

$$\begin{aligned} & \mathbf{u}_{2s+1} \overset{(1)}{x_s} \mathbf{u}_{2s} \overset{(1)}{x_{s-1}} \mathbf{u}_{2s-1} \overset{(2)}{x_s} \mathbf{u}_{2s-2} \overset{(1)}{x_{s-2}} \mathbf{u}_{2s-3} \overset{(2)}{x_{s-1}} \mathbf{u}_{2s-4} \overset{(1)}{x_{s-3}} \\ & \cdot \mathbf{u}_{2s-5} \overset{(2)}{x_{s-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0 \\ \approx & \mathbf{v}_{2s+1} \overset{(1)}{x_s} \mathbf{v}_{2s} \overset{(1)}{x_{s-1}} \mathbf{v}_{2s-1} \overset{(2)}{x_s} \mathbf{v}_{2s-2} \overset{(1)}{x_{s-2}} \mathbf{v}_{2s-3} \overset{(2)}{x_{s-1}} \mathbf{v}_{2s-4} \overset{(1)}{x_{s-3}} \\ & \cdot \mathbf{v}_{2s-5} \overset{(2)}{x_{s-2}} \cdots \mathbf{v}_4 \overset{(1)}{x_1} \mathbf{v}_3 \overset{(2)}{x_2} \mathbf{v}_2 \overset{(1)}{x_0} \mathbf{v}_1 \overset{(2)}{x_1} \mathbf{v}_0 \end{aligned}$$

for some $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2s+1}, \mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2s+1} \in \mathfrak{A}^*$. Clearly, ${}_{1\mathbf{u}}x$ occurs in the subword \mathbf{u}_{2s+1} of \mathbf{u} , while ${}_{1\mathbf{v}}x$ and ${}_{2\mathbf{v}}x$ occur in the subword \mathbf{v}_{2s+1} of \mathbf{v} . Since $x_s = h_2^r(\mathbf{u}, x)$ and x_{s-1} is an r -divider by Lemma 2.3, the letter ${}_{2\mathbf{u}}x$ occurs in \mathbf{u}_{2s} . Therefore, the identity

$$\mathbf{u}(x, x_1, x_2, \dots, x_s) \approx \mathbf{v}(x, x_1, x_2, \dots, x_s)$$

is equivalent modulo (3.1) to κ_s . Clearly, κ_s implies κ_k . Therefore, κ_k is satisfied by \mathbf{V} . \square

For any word \mathbf{w} , let $\text{ini}^2(\mathbf{w})$ denote the word obtained from $\text{ini}(\mathbf{w})$ by replacing each letter x with x^2 . For example, if $\mathbf{w} = x^3yzyx$, then $\text{ini}(\mathbf{w}) = xyz$ and so $\text{ini}^2(\mathbf{w}) = x^2y^2z^2$. A word \mathbf{w} is said to be *2-limited* if $\text{occ}_x(\mathbf{w}) \leq 2$ for any letter x .

Lemma 3.4. *Let \mathbf{w} be a word that does not contain simple letters. Then the identities (3.1) and*

$$(3.2) \quad (xy)^2 \approx x^2y^2.$$

imply the identity $\mathbf{w} \approx \text{ini}^2(\mathbf{w})$.

Proof. First, we note that $\mathbf{P}\{(3.2)\}$ satisfies the identity

$$(3.3) \quad x_1^2x_2^2 \cdots x_n^2 \approx (x_1x_2 \cdots x_n)^2$$

for any $n \geq 2$ because the identities

$$(x_1x_2 \cdots x_n)^2 \approx x_1^2(x_2x_3 \cdots x_n)^2 \approx x_1^2x_2^2(x_3x_4 \cdots x_n)^2 \approx \cdots \approx x_1^2x_2^2 \cdots x_n^2$$

follow from (3.2).

In view of Lemma 3.1, we may assume that the word \mathbf{w} is 2-limited. We say that a letter x *forms an island* in \mathbf{w} if $\mathbf{w} = \mathbf{w}_1x^2\mathbf{w}_2$ for some words $\mathbf{w}_1, \mathbf{w}_2 \in \mathfrak{A}^*$ such that $x \notin \text{con}(\mathbf{w}_1\mathbf{w}_2)$. Let k be the number of distinct letters that do not form an island in \mathbf{w} . We will use induction on k .

Induction base: $k = 0$. Then $\mathbf{w} = \text{ini}^2(\mathbf{w})$, and we are done.

Induction step: $k > 0$ and $\mathbf{P}\{(3.2)\}$ satisfies the identity $\mathbf{v}' \approx \text{ini}^2(\mathbf{v}')$ for any 2-limited word \mathbf{v}' with at most $k - 1$ letters that do not form an island in \mathbf{v}' . Then, since \mathbf{w} is 2-limited, there is a letter x such that $\mathbf{w} = \mathbf{w}_1xy_1^{k_1}y_2^{k_2} \cdots y_n^{k_n}x\mathbf{w}_2$, where

$k_1, k_2, \dots, k_n \in \{1, 2\}$ and if $k_i = 1$, then $y_i \in \text{con}(\mathbf{w}_1)$. Then the identities

$$\begin{aligned}
\mathbf{w} &= \mathbf{w}_1 x y_1^{k_1} y_2^{k_2} \cdots y_n^{k_n} x \mathbf{w}_2 \\
&\stackrel{(3.1)}{\approx} \mathbf{w}_1 x y_1^2 y_2^2 \cdots y_n^2 x y_1^2 y_2^2 \cdots y_n^2 \mathbf{w}_2 \\
&\stackrel{(3.3)}{\approx} \mathbf{w}_1 x (y_1 y_2 \cdots y_n)^2 x (y_1 y_2 \cdots y_n)^2 \mathbf{w}_2 \\
&\stackrel{(3.2)}{\approx} \mathbf{w}_1 x^2 (y_1 y_2 \cdots y_n)^4 \mathbf{w}_2 \\
&\stackrel{(3.3)}{\approx} \mathbf{w}_1 x^2 y_1^4 y_2^4 \cdots y_n^4 \mathbf{w}_2 \\
&\stackrel{(3.1)}{\approx} \mathbf{w}_1 x^2 y_1^{k_1} y_2^{k_2} \cdots y_n^{k_n} \mathbf{w}_2 = \mathbf{w}'
\end{aligned}$$

are satisfied by $\mathbf{P}\{(3.2)\}$. Clearly, the word \mathbf{w}' has exactly $k - 1$ letters that do not form an island in \mathbf{w}' . By the induction assumption, $\mathbf{w}' \approx \text{ini}^2(\mathbf{w}')$ holds in $\mathbf{P}\{(3.2)\}$. It remains to note that $\text{ini}^2(\mathbf{w}) = \text{ini}^2(\mathbf{w}')$. Therefore, $\mathbf{P}\{(3.2)\}$ satisfies $\mathbf{w} \approx \text{ini}^2(\mathbf{w})$. \square

Corollary 3.5. *Let \mathbf{w} be a word, k be a number such that the k -decomposition of \mathbf{w} is maximal and \mathbf{a} be a k -block of \mathbf{u} . If $\mathbf{w} = \mathbf{w}'\mathbf{a}\mathbf{w}''$, then the variety $\mathbf{P}\{(3.2)\}$ satisfies the identity $\mathbf{w} \approx \mathbf{w}' \text{ini}^2(\mathbf{a}) \mathbf{w}''$.*

Proof. Let x be a letter such that ${}_1\mathbf{w}x$ occurs in the k -block \mathbf{a} of \mathbf{w} . If ${}_{2\mathbf{w}}x$ does not occur in \mathbf{a} , then $h_1^k(\mathbf{w}, x) \neq h_2^k(\mathbf{w}, x)$ and so x is a $(k + 1)$ -divider of \mathbf{w} . This contradicts the fact that the k -decomposition of \mathbf{w} is maximal. Therefore, ${}_{2\mathbf{w}}x$ occurs in \mathbf{a} as well. We see that $\text{con}(\mathbf{a}) \subseteq \text{mul}(\mathbf{w}'\mathbf{a})$. Then we can apply the identity (3.1) and add some occurrences of letters in the k -block \mathbf{a} so that the resulting k -block \mathbf{b} contains multiple letters only and $\text{ini}(\mathbf{a}) = \text{ini}(\mathbf{b})$. Then identity $\mathbf{b} \approx \text{ini}^2(\mathbf{b})$ follows from (3.1) and (3.2) by Lemma 3.4. Evidently, $\text{ini}^2(\mathbf{a}) = \text{ini}^2(\mathbf{b})$. We see that the identities

$$\mathbf{w} = \mathbf{w}'\mathbf{a}\mathbf{w}'' \approx \mathbf{w}'\mathbf{b}\mathbf{w}'' \approx \mathbf{w}' \text{ini}^2(\mathbf{b}) \mathbf{w}'' = \mathbf{w}' \text{ini}^2(\mathbf{a}) \mathbf{w}''$$

follow from the identities (3.1) and (3.2), and we are done. \square

4. CRITICAL PAIRS

An identity $\mathbf{u} \approx \mathbf{v}$ is said to be *balanced* if $\text{occ}_x(\mathbf{u}) = \text{occ}_x(\mathbf{v})$ for any $x \in \mathfrak{A}$. We say that a pair $\{i_x, j_y\}$ of occurrences of letters x and y in a balanced identity $\mathbf{u} \approx \mathbf{v}$ is *critical* if \mathbf{u} contains ${}_{i\mathbf{u}}x j_{\mathbf{u}}y$ as a subword and $j_{\mathbf{v}}y$ precedes $i_{\mathbf{v}}x$ in \mathbf{v} . Let \mathbf{w} denote the word obtained from \mathbf{u} by replacing ${}_{i\mathbf{u}}x j_{\mathbf{u}}y$ with $j_{\mathbf{u}}y i_{\mathbf{u}}x$. Given a set Δ of identities and a balanced identity $\mathbf{u} \approx \mathbf{v}$, we say that the critical pair $\{i_x, j_y\}$ is *Δ -removable in $\mathbf{u} \approx \mathbf{v}$* if Δ implies $\mathbf{u} \approx \mathbf{w}$.

The following special case of Sapir [31, Lemma 3.4], which can be easily verified by induction, describes the standard method of deriving identities by removing critical pairs. This method traces back to Jackson and Sapir [13] and Sapir [30], for instance.

Lemma 4.1. *Let \mathbf{V} be a monoid variety and Δ be a set of identities. Suppose that each critical pair in every balanced identity of \mathbf{V} is Δ -removable. Then every balanced identity of \mathbf{V} can be derived from Δ .* \square

4.1. **Removing critical pairs of the form $\{1x, 2y\}$.** If $\phi: \mathfrak{A} \rightarrow \mathfrak{A}^*$ is a substitution and ε denote an identity $\mathbf{u} \approx \mathbf{v}$, then, for convenience, we will write $\phi(\varepsilon)$ rather than $\phi(\mathbf{u}) \approx \phi(\mathbf{v})$.

The following lemma generalizes Gusev and Vernikov [10, Lemma 6.3].

Lemma 4.2. *For any $k \in \mathbb{N}$, the identities κ_k and δ_k^k are equivalent within \mathbf{P} .*

Proof. Let ϕ be the substitution $(y_k, y_{k+1}) \mapsto (1, x)$. Then the identity $\phi(\delta_k^k)$ is equal to κ_k . Therefore, δ_k^k implies κ_k . Further, δ_k^k follows from (3.1) and κ_k because

$$\begin{aligned}
y_{k+1}y_kx_ky_{k+1}\mathbf{b}_{k,k}y_k\mathbf{b}_{k-1} &= y_{k+1}y_kx_ky_{k+1}x_{k-1}x_ky_k\mathbf{b}_{k-1} \\
&\stackrel{(3.1)}{\approx} y_{k+1}y_kx_ky_{k+1}x_{k-1}x_ky_kx_k\mathbf{c} \\
&\stackrel{\kappa_k}{\approx} y_{k+1}^2y_kx_kx_{k-1}x_ky_kx_k\mathbf{c} \\
&\stackrel{(3.1)}{\approx} y_{k+1}^2y_kx_kx_{k-1}x_ky_k\mathbf{d} \\
&\stackrel{\kappa_k}{\approx} y_{k+1}y_ky_{k+1}x_kx_{k-1}x_ky_k\mathbf{d} \\
&\stackrel{(3.1)}{\approx} y_{k+1}y_ky_{k+1}x_kx_{k-1}x_ky_k\mathbf{b}_{k-1} \\
&= y_{k+1}y_ky_{k+1}x_k\mathbf{b}_{k,k}y_k\mathbf{b}_{k-1},
\end{aligned}$$

where

$$\mathbf{c} = \begin{cases} \lambda & \text{if } k = 1, \\ x_{k-2}x_{k-1}x_k\mathbf{b}_{k-2} & \text{if } k > 1 \end{cases} \quad \text{and} \quad \mathbf{d} = \begin{cases} \lambda & \text{if } k = 1, \\ x_{k-2}x_kx_{k-1}x_k\mathbf{b}_{k-2} & \text{if } k > 1. \end{cases}$$

The lemma is thus proved. \square

We introduce the following countably infinite series of identities:

$$\varepsilon_{k-1} : y_kx_{k-1}xy_kx\mathbf{b}_{k-1} \approx y_kx_{k-1}y_kx^2\mathbf{b}_{k-1},$$

where $k \in \mathbb{N}$.

Lemma 4.3. *The following proper inclusions hold:*

- (i) $\mathbf{P}\{\gamma_1\} \subset \mathbf{P}\{\gamma_2\} \subset \cdots \subset \mathbf{P}\{\gamma_k\} \subset \cdots \subset \mathbf{P}\{\varepsilon_0\}$;
- (ii) $\mathbf{P}\{\gamma_k\} \subset \mathbf{P}\{\delta_k^1\} \subset \mathbf{P}\{\delta_k^2\} \subset \cdots \subset \mathbf{P}\{\delta_k^k\} \subset \mathbf{P}\{\delta_{k+1}^k\} \subset \cdots \subset \mathbf{P}\{\delta_{k+m}^k\} \subset \cdots \subset \mathbf{P}\{\varepsilon_k\}$;
- (iii) $\mathbf{P}\{\varepsilon_0\} \subset \mathbf{P}\{\varepsilon_1\} \subset \cdots \subset \mathbf{P}\{\varepsilon_k\} \subset \cdots \subset \mathbf{P}\{(3.2)\}$.

Proof. (i) Evidently, $\mathbf{P}\{\gamma_1\} \subseteq \mathbf{P}\{\gamma_2\} \subseteq \cdots \subseteq \mathbf{P}\{\gamma_k\} \subseteq \cdots \subseteq \mathbf{P}\{\varepsilon_0\}$. These inclusions are proper because $\mathbf{I}_{k+1} \subseteq \mathbf{P}\{\gamma_{k+1}\} \subseteq \mathbf{P}\{\varepsilon_0\}$ but $\mathbf{I}_{k+1} \not\subseteq \mathbf{P}\{\gamma_k\}$ for any $k \in \mathbb{N}$.

(ii) The inclusion $\mathbf{P}\{\gamma_k\} \subseteq \mathbf{P}\{\delta_k^1\}$ is evident. Let ϕ_1 and ϕ_2 be the substitutions

$$(x_{m-1}, x_m, x_{m+1}, \dots, x_k) \mapsto (x_mx_{m+1}x_{m-1}x_m, x_{m+1}, x_{m+2}, \dots, x_{k+1})$$

$$\text{and } (x_{m-1}, x_m, x_{m+1}, \dots, x_{k-1}, x_k, y_m) \mapsto (xx_{m-1}, 1, 1, \dots, 1, x, x_m),$$

respectively. Then the identities $\phi_1(\delta_k^m)$ and $\phi_2(\delta_k^m)$ are equivalent modulo (3.1) to δ_{k+1}^m and ε_m , respectively. Therefore, $\mathbf{P}\{\delta_k^m\} \subseteq \mathbf{P}\{\delta_{k+1}^m\}$ and $\mathbf{P}\{\delta_k^m\} \subseteq \mathbf{P}\{\varepsilon_m\}$. If $1 \leq m < k$ and ϕ_3 is the substitution

$$(x_{m-1}, y_m, y_{m+1}) \mapsto (y_{m+1}x_{m-1}, y_{m+1}, y_{m+2}),$$

then the identity $\phi_3(\delta_k^m)$ is equivalent modulo (3.1) to δ_k^{m+1} and so $\mathbf{P}\{\delta_k^m\} \subseteq \mathbf{P}\{\delta_k^{m+1}\}$. To show that the inclusions are proper, it suffices to note that for $1 \leq m \leq k$, $\mathbf{J}_k^m \subseteq \mathbf{P}\{\delta_k^m\} \subseteq \mathbf{P}\{\varepsilon_k\}$ but $\mathbf{J}_k^1 \not\subseteq \mathbf{P}\{\gamma_k\}$, $\mathbf{J}_{k+1}^m \not\subseteq \mathbf{P}\{\delta_k^m\}$ and $\mathbf{J}_k^{m+1} \not\subseteq \mathbf{P}\{\delta_k^m\}$ whenever $m < k$.

(iii) Evidently, $\mathbf{P}\{\varepsilon_0\} \subseteq \mathbf{P}\{\varepsilon_1\} \subseteq \cdots \subseteq \mathbf{P}\{\varepsilon_k\} \subseteq \cdots \subseteq \mathbf{P}\{(3.2)\}$. It is routine to verify that $\mathbf{P}\{\varepsilon_k\}$ violates ε_{k-1} for any $k \in \mathbb{N}$. Therefore, the inclusions are proper. \square

The *head* of a non-empty word \mathbf{w} , denoted by $h(\mathbf{w})$, is the first letter of \mathbf{w} . The following lemma generalizes Gusev and Vernikov [10, Lemma 6.6(i),(ii)].

Lemma 4.4. *Let \mathbf{V} be a subvariety of \mathbf{P} and \mathbf{u} be a word. Further, let $\mathbf{u} = \mathbf{u}'\mathbf{c}\mathbf{u}''$ for some $\mathbf{u}', \mathbf{u}'', \mathbf{c} \in \mathfrak{A}^*$ such that $\mathbf{c} \in \{xy, yx\}$, $x \notin \text{con}(\mathbf{u}')$ and $y \in \text{sim}(\mathbf{u}'')$. Suppose that one of the following holds:*

- (i) \mathbf{V} satisfies γ_k , $D(\mathbf{u}, x) = k$ and $D(\mathbf{u}, y) = 1$;
- (ii) \mathbf{V} satisfies δ_k^m , $D(\mathbf{u}, x) = k$ and $D(\mathbf{u}, y) = m + 1$;
- (iii) \mathbf{V} satisfies ε_k , $D(\mathbf{u}, x) = \infty$ and $D(\mathbf{u}, y) = k + 1$;
- (iv) \mathbf{V} satisfies (3.2), $D(\mathbf{u}, x) = D(\mathbf{u}, y) = \infty$.

Then \mathbf{V} satisfies the identity $\mathbf{u} \approx \mathbf{w}$, where \mathbf{w} is obtained from \mathbf{u} by swapping of the first occurrence of x and the second occurrence of y .

Proof. We consider only the case when $\mathbf{c} = xy$ since the case $\mathbf{c} = yx$ is similar. We also omit the proof of Part (i) because it is very similar to (and in fact simpler than) that of Part (ii).

(ii) Put $x_k = x$ and $y_{m+1} = y$. It follows from Corollary 2.8 that there exist letters x_0, x_1, \dots, x_{k-1} such that $D(\mathbf{u}, x_s) = s$ for any $0 \leq s < k$ and

$$\mathbf{u} = \mathbf{v}_{2k+2}y_{m+1}\mathbf{v}_{2k+1}x_ky_{m+1}\mathbf{p},$$

where

$$\mathbf{p} = \mathbf{v}_{2k}x_{k-1}\mathbf{v}_{2k-1}x_k\mathbf{v}_{2k-2}x_{k-2}\mathbf{v}_{2k-3}x_{k-1} \cdots \mathbf{v}_4x_1\mathbf{v}_3x_2\mathbf{v}_2x_0\mathbf{v}_1x_1\mathbf{v}_0$$

for some $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2k+2} \in \mathfrak{A}^*$. Clearly, for any $s = 1, 2, \dots, k-1$, there exist $\mathbf{v}'_{2s}, \mathbf{v}''_{2s} \in \mathfrak{A}^*$ such that $\mathbf{v}_{2s} = \mathbf{v}'_{2s}\mathbf{v}''_{2s}$, where \mathbf{v}'_{2s} does not contain any $(s-1)$ -divisor of \mathbf{u} , while either $\mathbf{v}''_{2s} = \lambda$ or $h(\mathbf{v}''_{2s})$ is an $(s-1)$ -divisor of \mathbf{u} . Put $\mathbf{v}''_{2k} = \mathbf{v}_{2k}$ and $\mathbf{v}'_0 = \mathbf{v}_0$. Let

$$\mathbf{q}_s = \mathbf{v}''_{2s+2}x_s\mathbf{v}_{2s+1}x_{s+1}\mathbf{v}'_{2s}$$

for any $s = 0, 1, \dots, k-1$. Let ϕ_1 be the substitution

$$(x_0, x_1, \dots, x_{k-1}, y_m, y_{m+1}) \mapsto (\mathbf{q}_0, \mathbf{q}_1, \dots, \mathbf{q}_{k-1}, \mathbf{v}_{2k+1}, y_{m+1}).$$

Then the identity $\phi_1(\delta_k^m)$ coincides with the identity

$$(4.1) \quad y_{m+1}\mathbf{v}_{2k+1}x_ky_{m+1}\mathbf{q} \approx \mathbf{v}_{2k+2}y_{m+1}\mathbf{v}_{2k+1}y_{m+1}x_k\mathbf{q}$$

where

$$\mathbf{q} = \mathbf{q}_{k-1}x_k\mathbf{q}_{k-2}\mathbf{q}_{k-1} \cdots \mathbf{q}_{m-1}\mathbf{q}_m\mathbf{v}_{2k+1}\mathbf{q}_{m-2}\mathbf{q}_{m-1} \cdots \mathbf{q}_1\mathbf{q}_2\mathbf{q}_0\mathbf{q}_1.$$

We note that $\mathbf{p} = \mathbf{q}_{k-1}\mathbf{q}_{k-2} \cdots \mathbf{q}_0$. Since $D(\mathbf{u}, x_s) = s$ and $D(\mathbf{u}, x_{s+1}) = s + 1$, Lemma 2.3 implies that x_s and x_{s+1} are not $(s-1)$ -divisors of \mathbf{u} . If one of the subwords \mathbf{v}''_{2s+2} or \mathbf{v}_{2s+1} of \mathbf{u} contains some $(s-1)$ -divisor d of \mathbf{u} , then $(1_{\mathbf{u}}x_{s+1}) < (1_{\mathbf{u}}d) < (2_{\mathbf{u}}x_{s+1})$ and so $D(\mathbf{u}, x_{s+1}) \leq s$ contradicting $D(\mathbf{u}, x_{s+1}) = s + 1$. Therefore, \mathbf{v}''_{2s+2} and \mathbf{v}_{2s+1} do not contain any $(s-1)$ -divisor of \mathbf{u} . Finally, the subword

\mathbf{v}'_{2s} of \mathbf{u} does not contain any $(s-1)$ -divider of \mathbf{u} by its definition. We see that the subword \mathbf{q}_s of \mathbf{u} does not contain $(s-1)$ -dividers of \mathbf{u} for any $s = 1, 2, \dots, k-1$.

Let z be a letter such that ${}_{1\mathbf{u}}z$ occurs in the subword \mathbf{v}_{2k+1} of \mathbf{u} . Since $D(\mathbf{u}, y_{m+1}) = m+1 > 1$, we have $z \in \text{mul}(\mathbf{u})$. We are going to verify that $z \in \text{mul}(\mathbf{v}_{2k+1}\mathbf{q}_{k-1} \cdots \mathbf{q}_m\mathbf{q}_{m-1})$. If $m = 1$, then this claim is evident because $\mathbf{v}_{2k+1}\mathbf{q}_{k-1} \cdots \mathbf{q}_m\mathbf{q}_{m-1}$ is a suffix of \mathbf{u} in this case. So, we may assume that $m > 1$. Let $z' = h(\mathbf{q}_{m-2})$. Since $z' = h(\mathbf{v}''_{2m-2})$ or $z' = x_{m-2}$, the letter z' is an $(m-2)$ -divider of \mathbf{u} . If $({}_{1\mathbf{u}}z') < ({}_{2\mathbf{u}}z)$, then $h_1^{m-2}(\mathbf{u}, z) \neq h_2^{m-2}(\mathbf{u}, z)$ and so $D(\mathbf{u}, z) \leq m-1$, whence z is an $(m-1)$ -divider of \mathbf{u} by Lemma 2.3. But this contradicts the fact that $D(\mathbf{u}, y_{m+1}) = m+1$ because $({}_{1\mathbf{u}}y_{m+1}) < ({}_{1\mathbf{u}}z) < ({}_{2\mathbf{u}}y_{m+1})$. Therefore, $({}_{2\mathbf{u}}z) < ({}_{1\mathbf{u}}z')$. We see that if a letter occurs in \mathbf{v}_{2k+1} , then the subword $\mathbf{v}_{2k+1}x_k y_{m+1}\mathbf{q}_{k-1} \cdots \mathbf{q}_m\mathbf{q}_{m-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} .

Let z be a letter such that ${}_{1\mathbf{u}}z$ occurs in the subword \mathbf{q}_s of \mathbf{u} for some $s \in \{1, 2, \dots, k-1\}$. We are going to verify that $z \in \text{mul}(\mathbf{q}_s\mathbf{q}_{s-1})$. If $s = 1$, then this claim is evident because \mathbf{q}_1 does not contain 0-dividers of \mathbf{u} and $\mathbf{q}_1\mathbf{q}_0$ is a suffix of \mathbf{u} . So, we may assume that $s > 1$. Clearly, either $z = x_s$ or $z \in \text{con}(\mathbf{v}''_{2s+2}\mathbf{v}_{2s+1}\mathbf{v}'_{2s})$. Evidently, if $z = x_s$, then $z \in \text{con}(\mathbf{q}_{s-1})$. Suppose now that $z \in \text{con}(\mathbf{v}''_{2s+2}\mathbf{v}_{2s+1}\mathbf{v}'_{2s})$. Since the subword \mathbf{q}_s of \mathbf{u} does not contain $(s-1)$ -dividers of \mathbf{u} , this subword does not contain 0-dividers of \mathbf{u} as well by Lemma 2.3. Therefore, the letter z is multiple in \mathbf{u} . Let $z' = h(\mathbf{q}_{s-2})$. Since $z' = h(\mathbf{v}''_{2s-2})$ or $z' = x_{s-2}$, the letter z' is an $(s-2)$ -divider of \mathbf{u} . If $({}_{1\mathbf{u}}z') < ({}_{2\mathbf{u}}z)$, then $h_1^{s-2}(\mathbf{u}, z) \neq h_2^{s-2}(\mathbf{u}, z)$ and so $D(\mathbf{u}, z) \leq s-1$, whence z is an $(s-1)$ -divider of \mathbf{u} by Lemma 2.3. But \mathbf{q}_s does not contain $(s-1)$ -dividers of \mathbf{u} . Therefore, $({}_{2\mathbf{u}}z) < ({}_{1\mathbf{u}}z')$. This means that ${}_{2\mathbf{u}}z$ occurs in the subword $\mathbf{q}_s\mathbf{q}_{s-1}$ of \mathbf{u} . We see that if a letter occurs in \mathbf{q}_s , then the subword $\mathbf{q}_s\mathbf{q}_{s-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} .

Finally, we recall that ${}_{2\mathbf{u}}x_k$ occurs in the subword \mathbf{q}_{k-1} of \mathbf{u} .

In view of the previous three paragraphs, the identity (3.1) implies

$$\mathbf{u} = \mathbf{v}_{2k+2}y_{m+1}\mathbf{v}_{2k+1}x_k y_{m+1}\mathbf{p} \approx \mathbf{v}_{2k+2}y_{m+1}\mathbf{v}_{2k+1}x_k y_{m+1}\mathbf{q}.$$

By a similar argument we can show that the identity (3.1) implies

$$\mathbf{v}_{2k+2}y_{m+1}\mathbf{v}_{2k+1}y_{m+1}x_k\mathbf{p} \approx \mathbf{v}_{2k+2}y_{m+1}\mathbf{v}_{2k+1}y_{m+1}x_k\mathbf{q}.$$

These identities together with (4.1) imply the identity $\mathbf{u} \approx \mathbf{u}'y_x\mathbf{u}''$. Thus, it is satisfied by \mathbf{V} .

(iii) There are two cases.

Case 1: $h(\mathbf{u}'') = x$. If $k = 0$, then ε_k is nothing but the identity $y_1x_0xy_1x \approx y_1x_0y_1x^2$. Since $y \in \text{con}(\mathbf{u}')$, this identity implies $\mathbf{u} \approx \mathbf{u}'y_x\mathbf{u}''$. So, we may assume that $k > 0$. Put $y_{k+1} = y$. In view of Corollary 2.8, there are letters x_0, x_1, \dots, x_k such that $D(\mathbf{u}, x_s) = s$ for any $0 \leq s \leq k$ and

$$\mathbf{u} = \mathbf{v}_{2k+3}y_{k+1}\mathbf{v}_{2k+2}x_k\mathbf{v}_{2k+1}x_k y_{k+1}\mathbf{p},$$

where

$$\mathbf{p} = \mathbf{v}_{2k}x_{k-1}\mathbf{v}_{2k-1}x_k \cdots \mathbf{v}_4x_1\mathbf{v}_3x_2\mathbf{v}_2x_0\mathbf{v}_1x_1\mathbf{v}_0$$

for some $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2k+3} \in \mathfrak{A}^*$. Clearly, for any $s = 1, 2, \dots, k-1$, there exist $\mathbf{v}'_{2s}, \mathbf{v}''_{2s} \in \mathfrak{A}^*$ such that $\mathbf{v}_{2s} = \mathbf{v}'_{2s}\mathbf{v}''_{2s}$, where \mathbf{v}'_{2s} does not contain any $(s-1)$ -divider of \mathbf{u} , while either $\mathbf{v}''_{2s} = \lambda$ or $h(\mathbf{v}''_{2s})$ is an $(s-1)$ -divider of \mathbf{u} . Put $\mathbf{v}''_{2k} = \mathbf{v}_{2k}$

and $\mathbf{v}'_0 = \mathbf{v}_0$. Let

$$\mathbf{q}_s = \mathbf{v}''_{2s+2} x_s \mathbf{v}_{2s+1} x_{s+1} \mathbf{v}'_{2s}$$

for any $s = 0, 1, \dots, k-1$. Let ϕ_2 be the substitution

$$(x_0, x_1, \dots, x_{k-1}, x_k) \mapsto (\mathbf{q}_0, \mathbf{q}_1, \dots, \mathbf{q}_{k-1}, \mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1}).$$

Then the identity $\phi_2(\varepsilon_k)$ coincides with the identity

$$(4.2) \quad y_{k+1} \mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1} x y_{k+1} x \mathbf{q} \approx y_{k+1} \mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1} y_{k+1} x^2 \mathbf{q}$$

where

$$\mathbf{q} = \mathbf{q}_{k-1} \mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1} \mathbf{q}_{k-2} \mathbf{q}_{k-1} \cdots \mathbf{q}_1 \mathbf{q}_2 \mathbf{q}_0 \mathbf{q}_1.$$

We note that $\mathbf{p} = \mathbf{q}_{k-1} \mathbf{q}_{k-2} \cdots \mathbf{q}_0$. By the same arguments as in Part (ii) one can show that

- if a letter occurs in the word $\mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1}$, then the subword $\mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1} x y_{k+1} x \mathbf{q}_{k-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} ;
- if a letter occurs in \mathbf{q}_s , then the subword $\mathbf{q}_s \mathbf{q}_{s-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} .

Then the identity (3.1) implies

$$\mathbf{u} = \mathbf{v}_{2k+3} y_{k+1} \mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1} x y_{k+1} x \mathbf{p} \approx \mathbf{v}_{2k+3} y_{k+1} \mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1} x y_{k+1} x \mathbf{q}.$$

By a similar argument one can show that the identity (3.1) implies

$$\mathbf{v}_{2k+3} y_{k+1} \mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1} y_{k+1} x^2 \mathbf{p} \approx \mathbf{v}_{2k+3} y_{k+1} \mathbf{v}_{2k+2} x_k \mathbf{v}_{2k+1} y_{k+1} x^2 \mathbf{q}.$$

These identities together with (4.2) imply the identity $\mathbf{u} \approx \mathbf{u}' y x \mathbf{u}''$. Thus, it is satisfied by \mathbf{V} .

Case 2: $h(\mathbf{u}'') \neq x$. In view of Remark 2.2, there is a number r such that the r -decomposition of \mathbf{u} is maximal. Since $D(\mathbf{u}, x) = \infty$, the subword ${}_{1\mathbf{u}} x {}_{2\mathbf{u}} y$ is contained in some r -block \mathbf{a} of \mathbf{u} . Then $\mathbf{a} = \mathbf{a}_1 x y \mathbf{a}_2$ and $\mathbf{u} = \mathbf{u}_1 \mathbf{a} \mathbf{u}_2$ for some $\mathbf{a}_1, \mathbf{a}_2, \mathbf{u}_1, \mathbf{u}_2 \in \mathfrak{A}^*$.

Let $\mathbf{u}^* = \mathbf{u}_1 \mathbf{a}_1 x y x \mathbf{a}_2 \mathbf{u}_2$. The word \mathbf{u}^* differs from \mathbf{u} by the occurrences of x only. Then, since $D(\mathbf{u}, x) = D(\mathbf{u}^*, x) = \infty$, the identity $\mathbf{u} \approx \mathbf{u}^*$ is s -well-balanced for any s . Since (3.2) is a consequence of ε_k by Lemma 4.3(iii), it follows from Corollary 3.5 that the identities

$$\mathbf{u} \approx \mathbf{u}_1 \text{ini}^2(\mathbf{a}_1 x y \mathbf{a}_2) \mathbf{u}_2 \quad \text{and} \quad \mathbf{u}^* \approx \mathbf{u}_1 \text{ini}^2(\mathbf{a}_1 x y x \mathbf{a}_2) \mathbf{u}_2$$

hold in \mathbf{V} . Clearly, $\text{ini}^2(\mathbf{a}_1 x y \mathbf{a}_2) = \text{ini}^2(\mathbf{a}_1 x y x \mathbf{a}_2)$ because $x \in \text{con}(\mathbf{a}_2) \setminus \text{con}(\mathbf{a}_1)$. This implies that \mathbf{V} satisfies $\mathbf{u} \approx \mathbf{u}^*$. By a similar argument we can show that $\mathbf{u}_1 \mathbf{a}_1 y x \mathbf{a}_2 \mathbf{u}_2 \approx \mathbf{u}_1 \mathbf{a}_1 y x^2 \mathbf{a}_2 \mathbf{u}_2$ is satisfied by \mathbf{V} . Then $D(\mathbf{u}^*, y) = k+1$ because the identity $\mathbf{u} \approx \mathbf{u}^*$ is s -well-balanced for any s . In view of Case 1, the identity $\mathbf{u}^* \approx \mathbf{u}_1 \mathbf{a}_1 y x^2 \mathbf{a}_2 \mathbf{u}_2$ is satisfied by \mathbf{V} . Therefore, $\mathbf{u} \approx \mathbf{u}' y x \mathbf{u}''$ holds in \mathbf{V} , and we are done.

(iv) In view of Remark 2.2, there is a number r such that the r -decomposition of \mathbf{u} is maximal. Since $D(\mathbf{u}, x) = D(\mathbf{u}, y) = \infty$, the subword ${}_{1\mathbf{u}} x {}_{2\mathbf{u}} y$ is contained in some r -block \mathbf{a} of \mathbf{u} and $\mathbf{a} = \mathbf{a}_1 y \mathbf{a}_2 x y \mathbf{a}_3 x \mathbf{a}_4$ for some $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4 \in \mathfrak{A}^*$. Evidently,

$$\text{ini}^2(\mathbf{a}_1 y \mathbf{a}_2 x y \mathbf{a}_3 x \mathbf{a}_4) = \text{ini}^2(\mathbf{a}_1 y \mathbf{a}_2 y x \mathbf{a}_3 x \mathbf{a}_4).$$

This fact and Corollary 3.5 imply that $\mathbf{u} \approx \mathbf{u}' y x \mathbf{u}''$ is satisfied by \mathbf{V} . \square

Corollary 4.5. *Let \mathbf{V} be a subvariety of $\mathbf{P}\{\kappa_k\}$ and \mathbf{u} be a word. Further, let $\mathbf{u} = \mathbf{u}'xy\mathbf{u}''$ for some $\mathbf{u}', \mathbf{u}'' \in \mathfrak{A}^*$ such that $x, y \in \text{sim}(\mathbf{u}')$. Suppose that either $D(\mathbf{u}, x) > k$ or $D(\mathbf{u}, y) > k$. Then \mathbf{V} satisfies the identity $\mathbf{u} \approx \mathbf{u}'y\mathbf{u}''$.*

Proof. We may assume without loss of generality that $(1_{\mathbf{u}}x) < (1_{\mathbf{u}}y)$. Then $\mathbf{u}' = \mathbf{u}_1x\mathbf{u}_2y\mathbf{u}_3$ for some $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3 \in \mathfrak{A}^*$. Evidently, $D(\mathbf{u}, x) \leq D(\mathbf{u}, y)$. It follows that $k < D(\mathbf{u}, y)$. Let ϕ be a substitution defined as follows: $\phi(c) = c^2$ for any $c \in \text{con}(\mathbf{u}_1)$ and $\phi(c) = c$ otherwise. Put $\mathbf{w} = \phi(\mathbf{u}_2)y\phi(\mathbf{u}_3)xyx\phi(\mathbf{u}'')$. We are going to verify that $D(\mathbf{u}, a) \leq D(\mathbf{w}, a)$ for any $a \in \text{con}(\mathbf{w})$. This fact is evident whenever $D(\mathbf{w}, a) = \infty$. So, we may assume that $D(\mathbf{w}, a) = k < \infty$. We will use induction on k .

Induction base: $k = 0$. Then $a \in \text{sim}(\mathbf{w})$. The definition of \mathbf{w} implies that $a \notin \text{sim}(\mathbf{u}_1)$ and so $a \in \text{sim}(\mathbf{u})$. Therefore, $D(\mathbf{u}, a) = 0$, and we are done.

Induction step: $k > 0$. Then $a \in \text{mul}(\mathbf{w})$. Let $b = h_2^{k-1}(\mathbf{w}, a)$. In view of Lemma 2.4, $D(\mathbf{w}, b) = k - 1$ and $(1_{\mathbf{w}}a) < (1_{\mathbf{w}}b) < (2_{\mathbf{w}}a)$. By the induction assumption $D(\mathbf{u}, b) \leq k - 1$. The definition of \mathbf{w} implies that $b \notin \text{sim}(\mathbf{u}_1)$ and $(1_{\mathbf{u}}a) < (1_{\mathbf{u}}b) < (2_{\mathbf{u}}a)$. Therefore, $D(\mathbf{u}, a) \leq k$, and we are done.

We have proved that $D(\mathbf{u}, a) \leq D(\mathbf{w}, a)$ for any $a \in \text{con}(\mathbf{w})$. In particular, $k < D(\mathbf{w}, y)$. According to Lemmas 4.2 and 4.3(ii),(iii), the variety $\mathbf{P}\{\kappa_k\}$ satisfies the identities (3.2) and ε_r for any $r \geq k$. Hence if $k < D(\mathbf{w}, y) < \infty$, then

$$(4.3) \quad \mathbf{w} = \phi(\mathbf{u}_2)y\phi(\mathbf{u}_3)xyx\phi(\mathbf{u}'') \approx \phi(\mathbf{u}_2)y\phi(\mathbf{u}_3)yx^2\phi(\mathbf{u}'')$$

is satisfied by $\mathbf{P}\{\kappa_k\}$ by Lemma 4.4(iii), and if $D(\mathbf{w}, y) = \infty$, then (4.3) holds in $\mathbf{P}\{\kappa_k\}$ by Lemma 4.4(iv). It remains to note that $\mathbf{P}\{\kappa_k\}$ satisfies

$$\mathbf{u} \stackrel{(3.1)}{\approx} \mathbf{u}_1x\mathbf{w} \stackrel{(4.3)}{\approx} \mathbf{u}_1x\phi(\mathbf{u}_2)y\phi(\mathbf{u}_3)yx^2\phi(\mathbf{u}'') \stackrel{(3.1)}{\approx} \mathbf{u}'y\mathbf{u}''$$

and so $\mathbf{u} \approx \mathbf{u}'y\mathbf{u}''$. \square

Lemma 4.6. *Suppose that $k \in \mathbb{N}$ and $\mathbf{a} \in \mathfrak{A}^*$ are such that $x_0, x_1, \dots, x_{k-1} \notin \text{con}(\mathbf{a})$ and $\text{sim}(\mathbf{a}) = \{x_k\}$. Then the variety $\mathbf{P}\{\kappa_k\}$ satisfies the identities*

$$\mathbf{ab}_k \approx \text{ini}(\mathbf{a})\text{ini}(\mathbf{a}_{x_k})\mathbf{b}_k,$$

$$y_m\mathbf{ab}_{k,m+1}x_{m-1}\mathbf{cb}_{m-1} \approx y_m\text{ini}(\mathbf{a})\text{ini}(\mathbf{a}_{x_k})\mathbf{b}_{k,m+1}x_{m-1}\mathbf{cb}_{m-1},$$

where $\mathbf{c} \in \{x_my_m, y_mx_m\}$.

Proof. It is routine to verify that $D(\mathbf{ab}_k, x_s) = s$ for any $0 \leq s \leq k - 1$. Let z be a letter from $\text{con}(\mathbf{a})$ with minimal depth in \mathbf{ab}_k , which we denote by d . Clearly, $d < \infty$ because $h_1^{k-1}(\mathbf{ab}_k, x_k) \neq x_{k-1} = h_2^{k-1}(\mathbf{ab}_k, x_k)$ and so $D(\mathbf{ab}_k, x_k) \leq k < \infty$. Then, since $\text{con}(\mathbf{a}) \subseteq \text{mul}(\mathbf{ab}_k)$, we have $d > 0$ and there is a $(d - 1)$ -divider t between the first and the second occurrences of z in \mathbf{ab}_k . In view of the choice of z , we have $t \notin \text{con}(\mathbf{a})$. It follows that $z \in \text{sim}(\mathbf{a})$. Hence $z = x_k$. According to Lemma 2.3, x_{k-1} is a $(k - 1)$ -divider of the word \mathbf{ab}_k . Now Lemma 2.4(ii) applies and we conclude that $D(\mathbf{ab}_k, z) = k$. In view of the choice of z , we have $D(\mathbf{ab}_k, a) > k$ for any $a \in \text{con}(\mathbf{a}) \setminus \{x_k\}$ and $D(\mathbf{ab}_k, x_k) = k$. Let $\text{con}(\mathbf{a}) = \{a_1, a_2, \dots, a_r\}$. We may assume without loss of generality that

$$\mathbf{a} = \overset{(1)}{a_1} \mathbf{a}_1 \overset{(1)}{a_2} \mathbf{a}_2 \cdots \overset{(1)}{a_r} \mathbf{a}_r$$

for some $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_r \in \mathfrak{A}^*$. The identity (3.1) allows us to remove any third or subsequent occurrence of letters from the subwords $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_r$ of \mathbf{a} . So, we may

assume that these subwords contain second occurrences of letters only. According to Lemmas 4.2 and 4.3(ii),(iii), the identities (3.2), δ_p^q and ε_p are satisfied by $\mathbf{P}\{\kappa_k\}$ for all $p \geq q \geq k$. Then, since $D(\mathbf{ab}_k, x_k) = k$ and $D(\mathbf{ab}_k, a) > k$ for any $a \in \text{con}(\mathbf{a}) \setminus \{x_k\}$, the identities

$$\mathbf{ab}_k = a_1 \mathbf{a}_1 a_2 \mathbf{a}_2 \cdots a_r \mathbf{a}_r \mathbf{b}_k \approx a_1 a_2 \cdots a_r \mathbf{a}_1 \mathbf{a}_2 \cdots \mathbf{a}_r \mathbf{b}_k = \text{ini}(\mathbf{a}) \mathbf{a}_1 \mathbf{a}_2 \cdots \mathbf{a}_r \mathbf{b}_k$$

hold in $\mathbf{P}\{\kappa_k\}$ by Parts (ii)–(iv) of Lemma 4.4. According to Corollary 4.5, $\mathbf{P}\{\kappa_k\}$ satisfies $\text{ini}(\mathbf{a}) \mathbf{a}_1 \mathbf{a}_2 \cdots \mathbf{a}_{m+1} \mathbf{b}_k \approx \text{ini}(\mathbf{a}) \text{ini}(\mathbf{a}_{x_k}) \mathbf{b}_k$ and so $\mathbf{ab}_k \approx \text{ini}(\mathbf{a}) \text{ini}(\mathbf{a}_{x_k}) \mathbf{b}_k$. Substituting $x_{m-1} \mathbf{c}$ for x_{m-1} in the last identity, we obtain the identity

$$(4.4) \quad \mathbf{ab}_{k,m+1} x_{m-1} \mathbf{c} x_m \mathbf{d} \approx \text{ini}(\mathbf{a}) \text{ini}(\mathbf{a}_{x_k}) \mathbf{b}_{k,m+1} x_{m-1} \mathbf{c} x_m \mathbf{d}$$

where

$$\mathbf{d} = \begin{cases} \mathbf{b}_0 & \text{if } m = 1, \\ \mathbf{b}_{m-1,m-1} \mathbf{c} \mathbf{b}_{m-2} & \text{if } m > 1. \end{cases}$$

Then $\mathbf{P}\{\kappa_k\}$ satisfies

$$\begin{aligned} y_m \mathbf{ab}_{k,m+1} x_{m-1} \mathbf{c} \mathbf{b}_{m-1} &\stackrel{(3.1)}{\approx} y_m \mathbf{ab}_{k,m+1} x_{m-1} \mathbf{c} x_m \mathbf{d} \\ &\stackrel{(4.4)}{\approx} y_m \text{ini}(\mathbf{a}) \text{ini}(\mathbf{a}_{x_k}) \mathbf{b}_{k,m+1} x_{m-1} \mathbf{c} x_m \mathbf{d} \\ &\stackrel{(3.1)}{\approx} y_m \text{ini}(\mathbf{a}) \text{ini}(\mathbf{a}_{x_k}) \mathbf{b}_{k,m+1} x_{m-1} \mathbf{c} \mathbf{b}_{m-1}, \end{aligned}$$

and we are done. \square

Let

$$\Phi = \{(3.1), (3.2), \gamma_k, \delta_k^m, \varepsilon_{k-1} \mid k \in \mathbb{N}, 1 \leq m \leq k\}.$$

Given a monoid variety \mathbf{V} and a system of identities Σ we use $\Sigma(\mathbf{V})$ to denote the set of those identities from Σ , which hold in \mathbf{V} .

Proposition 4.7. *Let \mathbf{V} be a monoid variety from the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}]$, $\mathbf{u} \approx \mathbf{v}$ be an identity of \mathbf{V} and $\{i x, j y\}$ be a critical pair in $\mathbf{u} \approx \mathbf{v}$. If $\{i, j\} = \{1, 2\}$, then the critical pair $\{i x, j y\}$ is Δ -removable for some $\Delta \subseteq \Phi(\mathbf{V})$.*

Proof. It suffices to assume that $(i, j) = (1, 2)$ since the case when $(i, j) = (2, 1)$ is similar. Let \mathbf{w} denote the word obtained from \mathbf{u} by replacing $1\mathbf{u}x 2\mathbf{u}y$ with $2\mathbf{u}y 1\mathbf{u}x$. Since $\{1x, 2y\}$ is a critical pair in $\mathbf{u} \approx \mathbf{v}$, the identity $\mathbf{u}(x, y) \approx \mathbf{v}(x, y)$ is equivalent modulo (3.1) to (3.2). Therefore, (3.2) is satisfied by \mathbf{V} .

Suppose that $D(\mathbf{u}, y) = \infty$. Then x is not an r -divider of \mathbf{u} for any $r \geq 0$. Therefore, $D(\mathbf{u}, x) = \infty$ by Lemma 2.3. Then $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), (3.2)\} \subseteq \Phi(\mathbf{V})$ by Lemma 4.4(iv). So, we may assume that $D(\mathbf{u}, y) < \infty$, say $D(\mathbf{u}, y) = s$ for some $s \in \mathbb{N}$. Further, Lemma 3.1 allows us to assume that the identity $\mathbf{u} \approx \mathbf{v}$ is 2-limited.

There are two cases.

Case 1: $\mathbf{F} \subseteq \mathbf{V}$. In view of Remark 2.2, there is a number k such that the k -decompositions of \mathbf{u} and \mathbf{v} are maximal. According to Corollary 2.13 and the inclusion $\mathbf{F}_{k+1} \subset \mathbf{F}$, the identity $\mathbf{u} \approx \mathbf{v}$ is k -well-balanced. It follows that x is not an r -divider of \mathbf{u} for any $r \geq 0$. Therefore, $D(\mathbf{u}, x) = \infty$ by Lemma 2.3.

Now we apply Lemma 2.6 and conclude that there are letters x_0, x_1, \dots, x_{s-1} such that $D(\mathbf{u}, x_r) = D(\mathbf{v}, x_r) = r$ for any $0 \leq r < s$ and the identity $\mathbf{u} \approx \mathbf{v}$ has

the form

$$\begin{aligned}
& \mathbf{u}_{2s+1} \overset{(1)}{y} \mathbf{u}_{2s} \overset{(1)}{x_{s-1}} \mathbf{u}_{2s-1} \overset{(2)}{y} \mathbf{u}_{2s-2} \overset{(1)}{x_{s-2}} \mathbf{u}_{2s-3} \overset{(2)}{x_{s-1}} \mathbf{u}_{2s-4} \overset{(1)}{x_{s-3}} \\
& \cdot \mathbf{u}_{2s-5} \overset{(2)}{x_{s-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0 \\
\approx & \mathbf{v}_{2s+1} \overset{(1)}{y} \mathbf{v}_{2s} \overset{(1)}{x_{s-1}} \mathbf{v}_{2s-1} \overset{(2)}{y} \mathbf{v}_{2s-2} \overset{(1)}{x_{s-2}} \mathbf{v}_{2s-3} \overset{(2)}{x_{s-1}} \mathbf{v}_{2s-4} \overset{(1)}{x_{s-3}} \\
& \cdot \mathbf{v}_{2s-5} \overset{(2)}{x_{s-2}} \cdots \mathbf{v}_4 \overset{(1)}{x_1} \mathbf{v}_3 \overset{(2)}{x_2} \mathbf{v}_2 \overset{(1)}{x_0} \mathbf{v}_1 \overset{(2)}{x_1} \mathbf{v}_0
\end{aligned}$$

for some words $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2s+1}$ and $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2s+1}$. Clearly, ${}_{1\mathbf{u}}x$ is the last letter of \mathbf{u}_{2s-1} . Since $D(\mathbf{u}, x) = \infty$, we have $({}_{2\mathbf{u}}x) < ({}_{1\mathbf{u}}x_{s-2})$. It follows that ${}_{2\mathbf{u}}x$ occurs in \mathbf{u}_{2s-2} . Further, $D(\mathbf{v}, x) = \infty$ by Corollary 2.13. Since $({}_{2\mathbf{v}}y) < ({}_{1\mathbf{v}}x)$ and the identity $\mathbf{u} \approx \mathbf{v}$ is k -well-balanced, both ${}_{1\mathbf{v}}x$ and ${}_{2\mathbf{v}}x$ occur in \mathbf{v}_{2s-2} . Then we substitute y_s for y in the identity

$$\mathbf{u}(x_0, x_1, \dots, x_{s-1}, x, y) \approx \mathbf{v}(x_0, x_1, \dots, x_{s-1}, x, y)$$

and obtain the identity ε_{s-1} . Therefore, ε_{s-1} is satisfied by \mathbf{V} . Now we apply Lemma 4.4(iii) and conclude that the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \varepsilon_{s-1}\} \subseteq \Phi(\mathbf{V})$.

Case 2: $\mathbf{F} \not\subseteq \mathbf{V}$. In view of Lemma 2.11, there is $k \in \mathbb{N}$ such that $\mathbf{F}_{k+1} \not\subseteq \mathbf{V}$. Let k be the least number with such a property. Then $\mathbf{F}_k \subseteq \mathbf{V}$. According to Lemma 3.3, \mathbf{V} satisfies κ_k . Then δ_k^k holds in \mathbf{V} by Lemma 4.2.

Suppose that $k < s$. Then, by Parts (ii) and (iii) of Lemma 4.3, \mathbf{V} satisfies ε_{s-1} and δ_r^s for any $r \geq s$. Clearly, $D(\mathbf{u}, x) \geq k$ because $D(\mathbf{u}, y) \leq k < s$ otherwise. If $D(\mathbf{u}, x) = \infty$, then Lemma 4.4(iii) applies and we conclude that the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \varepsilon_{s-1}\} \subseteq \Phi(\mathbf{V})$. If $D(\mathbf{u}, x) = r$ for some $k \leq r < \infty$, then, by Lemma 4.4(ii), the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \delta_r^s\} \subseteq \Phi(\mathbf{V})$. So, we may assume without loss of generality that $s \leq k$.

If $D(\mathbf{u}, x) = \infty$, then since (2.3) and (2.4) hold for all $\ell = 1, 2, \dots, k$ by Lemma 2.12, one can repeat literally arguments from the second paragraph of Case 1 and obtain that the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \varepsilon_{s-1}\} \subseteq \Phi(\mathbf{V})$. So, we may assume without loss of generality that $D(\mathbf{u}, x) = p$ for some $p \in \mathbb{N}$.

Then by Corollary 2.8, there exist letters x_0, x_1, \dots, x_{p-1} such that

$$\begin{aligned}
\mathbf{u} = & \mathbf{u}_{2p+1} \overset{(1)}{x} \mathbf{u}_{2p} \overset{(1)}{x_{p-1}} \mathbf{u}_{2s-1} \overset{(2)}{x} \mathbf{u}_{2p-2} \overset{(1)}{x_{p-2}} \mathbf{u}_{2p-3} \overset{(2)}{x_{p-1}} \mathbf{u}_{2p-4} \overset{(1)}{x_{p-3}} \\
& \cdot \mathbf{u}_{2p-5} \overset{(2)}{x_{p-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0
\end{aligned}$$

for some $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2p+1} \in \mathfrak{A}^*$, with $D(\mathbf{u}, x_r) = r$ and \mathbf{u}_{2r+2} not containing any r -dividers of \mathbf{u} for any $r = 0, 1, \dots, k-1$.

Clearly, ${}_{1\mathbf{u}}y$ occurs in \mathbf{u}_{2p+1} , while ${}_{2\mathbf{u}}y$ is the first letter of \mathbf{u}_{2p} . In view of Lemma 2.4, there is a letter y_{s-1} such that $D(\mathbf{u}, y_{s-1}) = s-1$ and $({}_{1\mathbf{u}}y) < ({}_{1\mathbf{u}}y_{s-1}) < ({}_{2\mathbf{u}}y)$. Let $X = \{x_0, x_1, \dots, x_{p-1}, x, y_{s-1}, y\}$.

Subcase 2.1: $s = 1$. Then $y_{s-1} = y_0 \in \text{sim}(\mathbf{u}) = \text{sim}(\mathbf{v})$ and

$$\mathbf{u}(X) = yy_0xyx_{p-1}x\mathbf{b}_{p-1}.$$

Let $p \leq k$. Since $({}_{2\mathbf{v}}y) < ({}_{1\mathbf{v}}x)$ and the identity $\mathbf{u} \approx \mathbf{v}$ is $(k-1)$ -well-balanced by Corollary 2.13, we have $\mathbf{v}(X) = yy_0yx_{p-1}x\mathbf{b}_{p-1}$. Let ϕ be the substitution $(x, y) \mapsto (x_p, y_s)$. Then the identity $\phi(\mathbf{u}(X)) \approx \phi(\mathbf{v}(X))$ coincides with γ_p .

Now let $p > k$. Since the identity $\mathbf{u} \approx \mathbf{v}$ is $(k-1)$ -well-balanced by Corollary 2.13, we have $\mathbf{v}(X) = yy_0 \mathbf{a}' \mathbf{b}_k$ for some word \mathbf{a}' with $\text{sim}(\mathbf{a}') = \{x_k, y\}$ and $\text{mul}(\mathbf{a}') = \{x_{k+1}, x_{k+2}, \dots, x_{p-1}, x\}$. Now Lemma 2.10 and the fact that $({}_{2\mathbf{v}}y) < ({}_{1\mathbf{v}}x)$ imply that $\text{ini}(\mathbf{a}') = yxx_{p-1}x_{p-2} \cdots x_k$. It follows that $\mathbf{a}' = ya$ for some a . Since κ_k holds in \mathbf{V} , Lemma 4.6 implies that \mathbf{V} satisfies the identities

$$\begin{aligned} yy_0 y \mathbf{a} \mathbf{b}_k &\approx yy_0 y \text{ini}(\mathbf{a}) \text{ini}(\mathbf{a}_{x_k}) \mathbf{b}_k = yy_0 y x x_{p-1} x_{p-2} \cdots x_k x_{p-1} x_{p-2} \cdots x_{k+1} \mathbf{b}_k \\ &\approx yy_0 y x x_{p-1} x \mathbf{b}_{p-1, k+1} \mathbf{b}_k = yy_0 y x x_{p-1} x \mathbf{b}_{p-1} \end{aligned}$$

and so the identity γ_p .

We see that γ_p is satisfied by \mathbf{V} in either case. Now we apply Lemma 4.4(i) and conclude that the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \gamma_p\} \subseteq \Phi(\mathbf{V})$.

Subcase 2.2: $s > 1$. Clearly, $p \geq s-1$ because $h_1^p(\mathbf{u}, y) \neq h_2^p(\mathbf{u}, y)$ and so $D(\mathbf{u}, y) < s$ otherwise. Suppose that $p = s-1$. Then $p \leq k$. In view of Corollary 2.13, the identity $\mathbf{u} \approx \mathbf{v}$ is $(k-1)$ -well-balanced. Hence

$$\begin{aligned} \mathbf{v} &= \mathbf{v}_{2p+1} \overset{(1)}{x} \mathbf{v}_{2p} \overset{(1)}{x_{p-1}} \mathbf{v}_{2p-1} \overset{(2)}{x} \mathbf{v}_{2p-2} \overset{(1)}{x_{p-2}} \mathbf{v}_{2p-3} \overset{(2)}{x_{p-1}} \mathbf{v}_{2s-4} \overset{(1)}{x_{p-3}} \\ &\quad \cdot \mathbf{v}_{2p-5} \overset{(2)}{x_{p-2}} \cdots \mathbf{v}_4 \overset{(1)}{x_1} \mathbf{v}_3 \overset{(2)}{x_2} \mathbf{v}_2 \overset{(1)}{x_0} \mathbf{v}_1 \overset{(2)}{x_1} \mathbf{v}_0 \end{aligned}$$

for some words $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2p+1}$. Let ϕ be the substitution $(x, y) \mapsto (x_p, x)$. Then the identity

$$\phi(\mathbf{u}(x_0, x_1, \dots, x_{p-1}, x, y)) \approx \phi(\mathbf{v}(x_0, x_1, \dots, x_{p-1}, x, y))$$

is equal to κ_{s-1} . Then \mathbf{V} satisfies δ_{s-1}^{s-1} by Lemma 4.2. Now we apply Lemma 4.4(ii) and conclude that the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \delta_{s-1}^{s-1}\} \subseteq \Phi(\mathbf{V})$. So, we may assume that $p > s-1$. In particular, $x \neq y_{s-1}$.

Since $s > 1$, the letter y_{s-1} is in $\text{mul}(\mathbf{u}) = \text{mul}(\mathbf{v})$. Since $D(\mathbf{u}, y) = s$, there are no $(s-2)$ -dividers of \mathbf{u} between ${}_{1\mathbf{u}}y$ and ${}_{2\mathbf{u}}y$. This fact and Lemma 2.9 imply that ${}_{2\mathbf{u}}y_{s-1}$ occurs in either \mathbf{u}_{2s-2} or \mathbf{u}_{2s-3} or \mathbf{u}_{2s-4} . Let $y_{s-2} = h_2^{s-1}(\mathbf{u}, y_{s-1})$. Using Lemmas 2.4 and 2.5, one can show that ${}_{1\mathbf{u}}y_{s-2}$ follows ${}_{2\mathbf{u}}x_s$. Since \mathbf{u}_{2s-2} does not contain any $(s-2)$ -dividers of \mathbf{u} , ${}_{1\mathbf{u}}y_{s-2}$ does not occur in \mathbf{u}_{2s-2} . Therefore, ${}_{2\mathbf{u}}y_{s-1}$ occurs in either \mathbf{u}_{2s-3} or \mathbf{u}_{2s-4} . Since $s \leq k$ and $\mathbf{u} \approx \mathbf{v}$ is $(k-1)$ -well-balanced by Corollary 2.13, the letter ${}_{2\mathbf{v}}y_{s-1}$ lies between ${}_{2\mathbf{v}}x_{s-2}$ and ${}_{1\mathbf{v}}x_{s-3}$. This means that ${}_{2\mathbf{v}}y_{s-1}$ occurs in either \mathbf{v}_{2s-3} or \mathbf{v}_{2s-4} .

Clearly,

$$\mathbf{u}(X) = yy_{s-1} xy x_{p-1} x \mathbf{b}_{p-1, s} x_{s-2} \mathbf{c} \mathbf{b}_{s-2},$$

where $\mathbf{c} \in \{x_{s-1} y_{s-1}, y_{s-1} x_{s-1}\}$.

Let $p \leq k$. Since the identity $\mathbf{u} \approx \mathbf{v}$ is $(k-1)$ -well-balanced and $({}_{2\mathbf{v}}y) < ({}_{1\mathbf{v}}x)$, we have

$$\mathbf{v}(X) = yy_{s-1} yx x_{p-1} x \mathbf{b}_{p-1, s} x_{s-2} \mathbf{d} \mathbf{b}_{s-2},$$

where $\mathbf{d} \in \{x_{s-1} y_{s-1}, y_{s-1} x_{s-1}\}$. Then the identity $\mathbf{u}(X) \approx \mathbf{v}(X)$ coincides with

$$(4.5) \quad yy_{s-1} xy x_{p-1} x \mathbf{b}_{p-1, s} x_{s-2} \mathbf{c} \mathbf{b}_{s-2} \approx yy_{s-1} yx x_{p-1} x \mathbf{b}_{p-1, s} x_{s-2} \mathbf{d} \mathbf{b}_{s-2}.$$

Now let $p > k$. Since the identity $\mathbf{u} \approx \mathbf{v}$ is $(k-1)$ -well-balanced and $({}_{2\mathbf{v}}y) < ({}_{1\mathbf{v}}x)$, we have

$$\mathbf{v}(X) = yy_{s-1} \mathbf{a}' \mathbf{b}_{k, s} x_{s-2} \mathbf{d} \mathbf{b}_{s-2},$$

where $\mathbf{d} \in \{x_{s-1} y_{s-1}, y_{s-1} x_{s-1}\}$ and \mathbf{a}' is a word with $\text{sim}(\mathbf{a}') = \{x_k, y\}$ and $\text{mul}(\mathbf{a}') = \{x_{k+1}, x_{k+2}, \dots, x_{p-1}, x\}$. Now Lemma 2.10 and the fact that $({}_{2\mathbf{v}}y) <$

($1_{\mathbf{V}}x$) imply that $\text{ini}(\mathbf{a}') = yxx_{p-1}x_{p-2} \cdots x_k$. Then $\mathbf{a}' = y\mathbf{a}$ for some \mathbf{a} . Since \mathbf{V} satisfies κ_k , Lemma 4.6 implies that \mathbf{V} satisfies also

$$\begin{aligned} yy_{s-1}y \mathbf{a} \mathbf{b}_{k,s} x_{s-2} \mathbf{d} \mathbf{b}_{s-2} &\approx yy_{s-1}y \text{ini}(\mathbf{a}) \text{ini}(\mathbf{a}_{x_k}) \mathbf{b}_{k,s} x_{s-2} \mathbf{d} \mathbf{b}_{s-2} \\ &= yy_{s-1}y xx_{p-1} \cdots x_k xx_{p-1} \cdots x_{k+1} \mathbf{b}_{k,s} x_{s-2} \mathbf{d} \mathbf{b}_{s-2} \\ &\approx yy_{s-1}y xx_{p-1}x \mathbf{b}_{p-1,s} x_{s-2} \mathbf{d} \mathbf{b}_{s-2} \end{aligned}$$

and so (4.5).

We see that (4.5) is satisfied by \mathbf{V} in either case. Then \mathbf{V} satisfies the identity δ_p^{s-1} because

$$\begin{aligned} & y_s \overset{(1)}{y_{s-1}} x_p y_s \cdots x_{s-2} \overset{(2)}{x_{s-1}} \overset{(2)}{y_{s-1}} x_{s-3} x_{s-2} \cdots x_1 x_2 x_0 x_1 \\ (3.1) \quad & \approx y_s \overset{(1)}{y_{s-1}} x_p y_s \cdots (x_{s-2} \overset{(2)}{x_{s-1}} \overset{(2)}{y_{s-1}}) \mathbf{c} x_{s-3} (x_{s-2} \overset{(4)}{x_{s-1}} \overset{(4)}{y_{s-1}}) \cdots x_1 x_2 x_0 x_1 \\ (4.5) \quad & \approx y_s \overset{(1)}{y_{s-1}} y_s x_p \cdots (x_{s-2} \overset{(2)}{x_{s-1}} \overset{(2)}{y_{s-1}}) \mathbf{d} x_{s-3} (x_{s-2} \overset{(4)}{x_{s-1}} \overset{(4)}{y_{s-1}}) \cdots x_1 x_2 x_0 x_1 \\ (3.1) \quad & \approx y_s \overset{(1)}{y_{s-1}} y_s x_p \cdots x_{s-2} \overset{(2)}{x_{s-1}} \overset{(2)}{y_{s-1}} x_{s-3} x_{s-2} \cdots x_1 x_2 x_0 x_1. \end{aligned}$$

Now we apply Lemma 4.4(ii) and conclude that the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \delta_p^{s-1}\} \subseteq \Phi(\mathbf{V})$. \square

4.2. Removing critical pairs of the form $\{2x, 2y\}$. We introduce the following countably infinite series of identities:

$$\begin{aligned} \zeta_{k-1} &: xy_0yx_{k-1}xy\mathbf{b}_{k-1} \approx xy_0yx_{k-1}yx\mathbf{b}_{k-1}, \\ \lambda_k^m &: xy_myx_kxy\mathbf{b}_{k,m}y_m\mathbf{b}_{m-1} \approx xy_myx_kyx\mathbf{b}_{k,m}y_m\mathbf{b}_{m-1}, \\ \eta_{k-1} &: xy_1yy_0xyx_{k-1}y_1\mathbf{b}_{k-1} \approx xy_1yy_0yx_{k-1}y_1\mathbf{b}_{k-1}, \\ \mu_k^m &: xy_{m+1}yy_myx_ky_{m+1}\mathbf{b}_{k,m}y_m\mathbf{b}_{m-1} \approx xy_{m+1}yy_myx_ky_{m+1}\mathbf{b}_{k,m}y_m\mathbf{b}_{m-1}, \\ \nu_{k-1} &: xx_{k-1}yzxyz_\infty^2z\mathbf{b}_{k-1} \approx xx_{k-1}yzzyz_\infty^2z\mathbf{b}_{k-1}, \end{aligned}$$

where $k \in \mathbb{N}$ and $1 \leq m \leq k$.

We omit the proof of the following statement because it is very similar to the proof of Lemma 4.3.

Lemma 4.8. *The following proper inclusions hold:*

- (i) $\mathbf{P}\{\zeta_0\} \subset \mathbf{P}\{\zeta_1\} \subset \cdots \subset \mathbf{P}\{\zeta_k\} \subset \cdots \subset \mathbf{P}\{\nu_0\} \subset \mathbf{P}\{\nu_1\} \subset \cdots \subset \mathbf{P}\{\nu_k\} \subset \cdots$;
- (ii) $\mathbf{P}\{\varepsilon_k\} \subset \mathbf{P}\{\zeta_k\} \subset \mathbf{P}\{\lambda_k^1\} \subset \mathbf{P}\{\lambda_k^2\} \subset \cdots \subset \mathbf{P}\{\lambda_k^k\} \subset \mathbf{P}\{\lambda_{k+1}^k\} \subset \cdots \subset \mathbf{P}\{\lambda_{k+m}^k\} \subset \cdots$;
- (iii) $\mathbf{P}\{\eta_0\} \subset \mathbf{P}\{\eta_1\} \subset \cdots \subset \mathbf{P}\{\eta_k\} \subset \cdots$;
- (iv) $\mathbf{P}\{\mu_k^1\} \subset \mathbf{P}\{\mu_k^2\} \subset \cdots \subset \mathbf{P}\{\mu_k^k\} \subset \mathbf{P}\{\mu_{k+1}^k\} \subset \cdots \subset \mathbf{P}\{\mu_{k+m}^k\} \subset \cdots$. \square

Let $\mathbf{w} = \mathbf{w}_1\mathbf{w}_2\mathbf{w}_3$. We say that the *depth* of the subword \mathbf{w}_2 in \mathbf{w} is equal to k if $D(\mathbf{w}, a) = k$ for some $a \in \text{con}(\mathbf{w}_2) \setminus \text{con}(\mathbf{w}_1)$ and k is the least number with such a property. Equivalently, the depth of the subword \mathbf{w}_2 in \mathbf{w} is equal to k if \mathbf{w}_2 contains a k -divider of \mathbf{w} but does not contain any $(k-1)$ -divider of \mathbf{w} (see Lemma 2.3). If, for any $k \geq 0$, \mathbf{w}_2 does not contain any k -divider of \mathbf{w} , then we say that the depth of the subword \mathbf{w}_2 in \mathbf{w} is *infinite*. Equivalently, the depth of the subword \mathbf{w}_2 in \mathbf{w} is infinite if $D(\mathbf{w}, a) = \infty$ for any $a \in \text{con}(\mathbf{w}_2) \setminus \text{con}(\mathbf{w}_1)$ (see Lemma 2.3).

Lemma 4.9. *Let \mathbf{V} be a subvariety of \mathbf{P} and $\mathbf{u} = \mathbf{u}' \overset{(2)}{x} \overset{(2)}{y} \mathbf{u}''$ for some $\mathbf{u}', \mathbf{u}'' \in \mathfrak{A}^*$.*

Further, let $\mathbf{u}' = \mathbf{u}_1 \overset{(1)}{x} \mathbf{u}_2 \overset{(1)}{y} \mathbf{u}_3$, where the depth of the subwords \mathbf{u}_2 and \mathbf{u}_3 in \mathbf{u} is equal to m and k , respectively. Suppose also that one of the following holds:

- (i) $k \geq 0$, $m = 0$ and \mathbf{V} satisfies ζ_k ;
- (ii) $1 \leq m \leq k$ and \mathbf{V} satisfies λ_k^m .

Then \mathbf{V} satisfies the identity $\mathbf{u} \approx \mathbf{u}'y\mathbf{x}\mathbf{u}''$.

Proof. We omit the proof of Part (i) because it is very similar to (and in fact simpler than) that of Part (ii).

(ii) The fact that the depth of \mathbf{u}_3 is equal to k in \mathbf{u} and Lemma 2.3 imply that \mathbf{u}_3 contains a k -divider x_k of \mathbf{u} such that $D(\mathbf{u}, x_k) = k$. It follows from Corollary 2.8 that there exist letters x_0, x_1, \dots, x_{k-1} such that $D(\mathbf{u}, x_s) = s$ for any $0 \leq s < k$ and

$$\mathbf{u}'' = \mathbf{v}_{2k}x_{k-1}\mathbf{v}_{2k-1}x_k\mathbf{v}_{2k-2}x_{k-2}\mathbf{v}_{2k-3}x_{k-1} \cdots \mathbf{v}_4x_1\mathbf{v}_3x_2\mathbf{v}_2x_0\mathbf{v}_1x_1\mathbf{v}_0,$$

for some $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2k} \in \mathfrak{A}^*$. Clearly, for any $s = 1, 2, \dots, k-1$, there exist $\mathbf{v}'_{2s}, \mathbf{v}''_{2s} \in \mathfrak{A}^*$ such that $\mathbf{v}_{2s} = \mathbf{v}'_{2s}\mathbf{v}''_{2s}$, where \mathbf{v}'_{2s} does not contain any $(s-1)$ -divisor of \mathbf{u} , while either $\mathbf{v}''_{2s} = \lambda$ or $h(\mathbf{v}''_{2s})$ is an $(s-1)$ -divisor of \mathbf{u} . Put $\mathbf{v}''_{2k} = \mathbf{v}_{2k}$ and $\mathbf{v}'_0 = \mathbf{v}_0$. Let

$$\mathbf{q}_s = \mathbf{v}''_{2s+2}x_s\mathbf{v}_{2s+1}x_{s+1}\mathbf{v}'_{2s}$$

for any $s = 0, 1, \dots, k-1$.

Let ϕ be the substitution

$$(x_0, \dots, x_{k-1}, x_k, y_m) \mapsto (\mathbf{q}_0, \dots, \mathbf{q}_{k-1}, \mathbf{u}_3, \mathbf{u}_2).$$

Then the identity $\phi(\lambda_k^m)$ coincides with

$$(4.6) \quad x\mathbf{u}_2y\mathbf{u}_3xy\mathbf{q} \approx x\mathbf{u}_2y\mathbf{u}_3yx\mathbf{q}$$

where

$$\mathbf{q} = \mathbf{q}_{k-1}\mathbf{u}_3\mathbf{q}_{k-2}\mathbf{q}_{k-1} \cdots \mathbf{q}_{m-1}\mathbf{q}_m\mathbf{u}_2\mathbf{q}_{m-2}\mathbf{q}_{m-1} \cdots \mathbf{q}_1\mathbf{q}_2\mathbf{q}_0\mathbf{q}_1.$$

We note that $\mathbf{u}'' = \mathbf{q}_{k-1}\mathbf{q}_{k-2} \cdots \mathbf{q}_0$. By the same arguments as in the proof of Lemma 4.4(ii) we can show that

- if a letter occurs in \mathbf{u}_2 , then the subword $\mathbf{u}_2y\mathbf{u}_3xy\mathbf{q}_{k-1} \cdots \mathbf{q}_m\mathbf{q}_{m-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} ;
- if a letter occurs in \mathbf{u}_3 , then the subword $\mathbf{u}_3xy\mathbf{q}_{k-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} ;
- if a letter occurs in \mathbf{q}_s , then the subword $\mathbf{q}_s\mathbf{q}_{s-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} for any $s = 1, 2, \dots, k-1$.

Then the identity (3.1) implies $\mathbf{u} = \mathbf{u}'xy\mathbf{u}'' \approx \mathbf{u}'xy\mathbf{q}$. By a similar argument we can show that $\mathbf{u}'y\mathbf{x}\mathbf{u}'' \approx \mathbf{u}'yx\mathbf{q}$ follows from (3.1). These identities together with (4.6) imply the identity $\mathbf{u} \approx \mathbf{u}'y\mathbf{x}\mathbf{u}''$. Thus, it is satisfied by \mathbf{V} . \square

The proof of the following lemma is very similar to the proof of Lemma 4.9. We provide it for the sake of completeness.

Lemma 4.10. *Let \mathbf{V} be a subvariety of \mathbf{P} and $\mathbf{u} = \mathbf{u}' \overset{(2)}{x} \overset{(2)}{y} \mathbf{u}''$ for some $\mathbf{u}', \mathbf{u}'' \in \mathfrak{A}^*$. Further, let*

$$\mathbf{u}' = \mathbf{u}_1 \overset{(1)}{x} \mathbf{u}_2 \overset{(1)}{y_{m+1}} \mathbf{u}_3 \overset{(1)}{y} \mathbf{u}_4 \quad \text{and} \quad \mathbf{u}'' = \mathbf{u}_5 \overset{(1)}{x_k} \mathbf{u}_6 \overset{(1)}{y_{m+1}} \mathbf{u}_7,$$

where $D(\mathbf{u}, y_{m+1}) = m + 1$, $D(\mathbf{u}, x_k) = k$ and y_{m+1} is a letter such that $(2_{\mathbf{u}}a) \leq (2_{\mathbf{u}}y_{m+1})$ for any $a \in \{c \mid 1_{\mathbf{u}}c \text{ lies in } \mathbf{u}_2y_{m+1}\mathbf{u}_3\}$. Suppose that the depth of the subwords $\mathbf{u}_2y_{m+1}\mathbf{u}_3$, \mathbf{u}_4 and $\mathbf{u}_5x_k\mathbf{u}_6$ is equal to $m + 1$, m and k , respectively. Suppose also that one of the following holds:

- (i) $k \geq 0$, $m = 0$ and \mathbf{V} satisfies η_k ;
- (ii) $1 \leq m \leq k$ and \mathbf{V} satisfies μ_k^m .

Then \mathbf{V} satisfies the identity $\mathbf{u} \approx \mathbf{u}'y\mathbf{x}\mathbf{u}''$.

Proof. We omit the proof of Part (i) because it is very similar to (and in fact simpler than) that of Part (ii).

(ii) It follows from Corollary 2.8 and the fact that the depth of $\mathbf{u}_5x_k\mathbf{u}_6$ is equal to k that there exist letters x_0, x_1, \dots, x_{k-1} such that $D(\mathbf{u}, x_s) = s$ for any $0 \leq s < k$ and

$$\mathbf{u}_7 = \mathbf{v}_{2k}x_{k-1}\mathbf{v}_{2k-1}x_k\mathbf{v}_{2k-2}x_{k-2}\mathbf{v}_{2k-3}x_{k-1} \cdots \mathbf{v}_4x_1\mathbf{v}_3x_2\mathbf{v}_2x_0\mathbf{v}_1x_1\mathbf{v}_0,$$

for some $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2k} \in \mathfrak{A}^*$. Clearly, for any $s = 1, 2, \dots, k - 1$, there exist $\mathbf{v}'_{2s}, \mathbf{v}''_{2s} \in \mathfrak{A}^*$ such that $\mathbf{v}_{2s} = \mathbf{v}'_{2s}\mathbf{v}''_{2s}$, where \mathbf{v}'_{2s} does not contain any $(s - 1)$ -divisor of \mathbf{u} , while either $\mathbf{v}''_{2s} = \lambda$ or $h(\mathbf{v}''_{2s})$ is an $(s - 1)$ -divisor of \mathbf{u} . Put $\mathbf{v}''_{2k} = \mathbf{v}_{2k}$ and $\mathbf{v}'_0 = \mathbf{v}_0$. Let

$$\mathbf{q}_s = \mathbf{v}''_{2s+2}x_s\mathbf{v}_{2s+1}x_{s+1}\mathbf{v}'_{2s}$$

for any $s = 0, 1, \dots, k - 1$.

Let ϕ be the substitution

$$(x_0, \dots, x_{k-1}, x_k, y_m, y_{m+1}) \mapsto (\mathbf{q}_0, \dots, \mathbf{q}_{k-1}, \mathbf{u}_5x_k\mathbf{u}_6, \mathbf{u}_4, \mathbf{u}_2y_{m+1}\mathbf{u}_3).$$

Then the identity $\phi(\mu_k^m)$ coincides with

$$(4.7) \quad \mathbf{x}\mathbf{u}_2y_{m+1}\mathbf{u}_3y\mathbf{u}_4xy\mathbf{u}_5x_k\mathbf{u}_6\mathbf{q} \approx \mathbf{x}\mathbf{u}_2y_{m+1}\mathbf{u}_3y\mathbf{u}_4y\mathbf{x}\mathbf{u}_5x_k\mathbf{u}_6\mathbf{q}$$

where

$$\mathbf{q} = \mathbf{u}_2y_{m+1}\mathbf{u}_3\mathbf{q}_{k-1}\mathbf{u}_5x_k\mathbf{u}_6\mathbf{q}_{k-2}\mathbf{q}_{k-1} \cdots \mathbf{q}_{m-1}\mathbf{q}_m\mathbf{u}_4\mathbf{q}_{m-2}\mathbf{q}_{m-1} \cdots \mathbf{q}_1\mathbf{q}_2\mathbf{q}_0\mathbf{q}_1.$$

We note that $\mathbf{u}_7 = \mathbf{q}_{k-1}\mathbf{q}_{k-2} \cdots \mathbf{q}_0$. The choice of y_{m+1} implies that if z is a letter such that $1_{\mathbf{u}}z$ occurs in the subword $\mathbf{u}_2y_{m+1}\mathbf{u}_3$ of \mathbf{u} , then $(2_{\mathbf{u}}z) \leq (2_{\mathbf{u}}y_{m+1})$. We see that if a letter lies in $\mathbf{u}_2y_{m+1}\mathbf{u}_3$, then the subword $\mathbf{u}_2y_{m+1}\mathbf{u}_3y\mathbf{u}_4xy\mathbf{u}_5x_k\mathbf{u}_6y_{m+1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} . By the same arguments as in the proof of Lemma 4.4(ii) we can show that

- if a letter occurs in \mathbf{u}_4 , then the subword $\mathbf{u}_4xy\mathbf{u}_5x_k\mathbf{u}_6y_{m+1}\mathbf{q}_{k-1} \cdots \mathbf{q}_m\mathbf{q}_{m-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} ;
- if a letter occurs in $\mathbf{u}_5x_k\mathbf{u}_6$, then the subword $\mathbf{u}_5x_k\mathbf{u}_6y_{m+1}\mathbf{q}_{k-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} ;
- if a letter occurs in \mathbf{q}_s , then the subword $\mathbf{q}_s\mathbf{q}_{s-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} for any $s = 1, 2, \dots, k - 1$.

Then the identity (3.1) implies

$$\mathbf{u} = \mathbf{u}'xy\mathbf{u}_5x_k\mathbf{u}_6y_{m+1}\mathbf{u}_7 \approx \mathbf{u}'xy\mathbf{u}_5x_k\mathbf{u}_6y_{m+1}\mathbf{q}.$$

By a similar argument we can show that

$$\mathbf{u}'y\mathbf{x}\mathbf{u}_5x_k\mathbf{u}_6y_{m+1}\mathbf{u}_7 \approx \mathbf{u}'y\mathbf{x}\mathbf{u}_5x_k\mathbf{u}_6y_{m+1}\mathbf{q}$$

follows from (3.1). These identities together with (4.7) imply the identity $\mathbf{u} \approx \mathbf{u}'y\mathbf{x}\mathbf{u}''$. Thus, it is satisfied by \mathbf{V} . \square

Lemma 4.11. *Let \mathbf{V} be a subvariety of \mathbf{P} and $\mathbf{u} = \mathbf{u}' \overset{(2)}{x} \overset{(2)}{y} \mathbf{u}''$ for some $\mathbf{u}', \mathbf{u}'' \in \mathfrak{A}^*$. Further, let $\mathbf{u}' = \mathbf{u}_1 \overset{(1)}{x} \mathbf{u}_2 \overset{(1)}{y} \mathbf{u}_3$, where the depth of the subword \mathbf{u}_2 in \mathbf{u} is equal to k , while the depth of the subword \mathbf{u}_3 in \mathbf{u} is infinite. Suppose that \mathbf{V} satisfies ν_k . Then \mathbf{V} satisfies the identity $\mathbf{u} \approx \mathbf{u}'y\mathbf{x}\mathbf{u}''$.*

Proof. The fact that the depth of \mathbf{u}_2 is equal to k in \mathbf{u} and Lemma 2.3 imply that \mathbf{u}_2 contains a k -divider x_k of \mathbf{u} such that $D(\mathbf{u}, x_k) = k$. It follows from Corollary 2.8 and the fact that the depth of \mathbf{u}_3 is infinite that there exist letters x_0, x_1, \dots, x_{k-1} such that $D(\mathbf{u}, x_s) = s$ for any $0 \leq s < k$ and

$$\mathbf{u}'' = \mathbf{v}_{2k}x_{k-1}\mathbf{v}_{2k-1}x_k\mathbf{v}_{2k-2}x_{k-2}\mathbf{v}_{2k-3}x_{k-1} \cdots \mathbf{v}_4x_1\mathbf{v}_3x_2\mathbf{v}_2x_0\mathbf{v}_1x_1\mathbf{v}_0,$$

for some $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_{2k} \in \mathfrak{A}^*$. Clearly, for any $s = 1, 2, \dots, k-1$, there exist $\mathbf{v}'_{2s}, \mathbf{v}''_{2s} \in \mathfrak{A}^*$ such that $\mathbf{v}_{2s} = \mathbf{v}'_{2s}\mathbf{v}''_{2s}$, where \mathbf{v}'_{2s} does not contain any $(s-1)$ -divider of \mathbf{u} , while either $\mathbf{v}''_{2s} = \lambda$ or $h(\mathbf{v}''_{2s})$ is an $(s-1)$ -divider of \mathbf{u} . Besides that, there are words \mathbf{v}'_{2k} and \mathbf{v}''_{2k} such that $\mathbf{v}_{2k} = \mathbf{v}'_{2k}\mathbf{v}''_{2k}$, where \mathbf{v}'_{2k} does not contain r -dividers of \mathbf{u} for any $r \in \mathbb{N} \cup \{0\}$, while either $\mathbf{v}''_{2s} = \lambda$ or $h(\mathbf{v}''_{2s})$ is an r' -divider of \mathbf{u} for some $r' \in \mathbb{N} \cup \{0\}$. Put $\mathbf{v}'_0 = \mathbf{v}_0$. Let

$$\mathbf{q}_s = \mathbf{v}''_{2s+2}x_s\mathbf{v}_{2s+1}x_{s+1}\mathbf{v}'_{2s}$$

for any $s = 0, 1, \dots, k-1$.

Let ϕ be the substitution

$$(x_0, x_1, \dots, x_{k-1}, x_k, z_\infty, z) \mapsto (\mathbf{q}_0, \mathbf{q}_1, \dots, \mathbf{q}_{k-1}, \mathbf{u}_2, \mathbf{v}'_{2k}, \mathbf{u}_3).$$

Then the identity $\phi(\nu_k)$ coincides with

$$(4.8) \quad \mathbf{x}\mathbf{u}_2\mathbf{y}\mathbf{u}_3\mathbf{x}\mathbf{y}\mathbf{q} \approx \mathbf{x}\mathbf{u}_2\mathbf{y}\mathbf{u}_3\mathbf{y}\mathbf{x}\mathbf{q}$$

where

$$\mathbf{q} = (\mathbf{v}'_{2k})^2\mathbf{u}_3\mathbf{q}_{k-1}\mathbf{u}_2\mathbf{q}_{k-2}\mathbf{q}_{k-1} \cdots \mathbf{q}_1\mathbf{q}_2\mathbf{q}_0\mathbf{q}_1.$$

We note that $\mathbf{u}'' = \mathbf{v}'_{2k}\mathbf{q}_{k-1}\mathbf{q}_{k-2} \cdots \mathbf{q}_0$. Since the subword $\mathbf{u}_3\mathbf{x}\mathbf{y}\mathbf{v}'_{2k}$ of \mathbf{u} does not contain r -dividers of \mathbf{u} for any $r \in \mathbb{N} \cup \{0\}$ and the first letter of \mathbf{q}_{k-1} is an r' -divider of \mathbf{u} for some $r' \in \mathbb{N} \cup \{0\}$, if a letter occurs in the subword $\mathbf{u}_3\mathbf{x}\mathbf{y}\mathbf{v}'_{2k}$ of \mathbf{u} , then this subword contains some non-first occurrence of this letter. By the same arguments as in the proof of Lemma 4.4(ii) we can show that

- if a letter occurs in \mathbf{u}_2 , then the subword $\mathbf{u}_2\mathbf{y}\mathbf{u}_3\mathbf{x}\mathbf{y}\mathbf{v}'_{2k}\mathbf{q}_{k-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} ;
- if a letter occurs in \mathbf{q}_s , then the subword $\mathbf{q}_s\mathbf{q}_{s-1}$ of \mathbf{u} contains some non-first occurrence of this letter in \mathbf{u} for any $s = 1, 2, \dots, k-1$.

Then the identity (3.1) implies $\mathbf{u} = \mathbf{u}'\mathbf{x}\mathbf{y}\mathbf{u}'' \approx \mathbf{u}'\mathbf{x}\mathbf{y}\mathbf{q}$. By a similar argument we can show that $\mathbf{u}'\mathbf{y}\mathbf{x}\mathbf{u}'' \approx \mathbf{u}'\mathbf{y}\mathbf{x}\mathbf{q}$ follows from (3.1). These identities together with (4.8) imply the identity $\mathbf{u} \approx \mathbf{u}'\mathbf{y}\mathbf{x}\mathbf{u}''$. Thus, it is satisfied by \mathbf{V} . \square

Let $\mathbf{w} = \mathbf{w}_1\mathbf{w}_2\mathbf{w}_3$. The depth of \mathbf{w}_2 in \mathbf{w} is *strictly infinite* if

- the depth of \mathbf{w}_2 in \mathbf{w} is infinite, in particular, $\text{con}(\mathbf{w}_2) \subseteq \text{mul}(\mathbf{w})$;
- for any $a \in \text{sim}(\mathbf{w}_2) \setminus \text{con}(\mathbf{w}_1)$, there are no first occurrences of letters between the last letter of \mathbf{w}_2 and the second occurrence of a in \mathbf{w} .

For an arbitrary $n \in \mathbb{N}$, we denote by S_n the full symmetric group on the set $\{1, 2, \dots, n\}$.

Lemma 4.12. *Let $\mathbf{u} = \mathbf{u}_1 \overset{(1)}{x} \mathbf{u}_2 \overset{(1)}{y} \mathbf{u}_3 \overset{(2)}{x} \overset{(2)}{y} \mathbf{u}_4$. If the depth of \mathbf{u}_3 in \mathbf{u} is strictly infinite, then the identity $\mathbf{u} \approx \mathbf{u}_1 \mathbf{x} \mathbf{u}_2 \mathbf{y} \mathbf{u}_3 \mathbf{y} \mathbf{x} \mathbf{u}_4$ follows from the identities (3.1) and*

$$(4.9) \quad xtyxy \approx xty^2x.$$

Proof. First, we note that if $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3$ are words with $\text{sim}(\mathbf{w}_3) \subseteq \text{con}(\mathbf{w}_1 \mathbf{w}_2)$, then the identities (3.1) and (4.9) imply the identity

$$\mathbf{w}_1 \mathbf{x} \mathbf{w}_2 \mathbf{y} \mathbf{w}_3 \mathbf{x} \mathbf{y} \approx \mathbf{w}_1 \mathbf{x} \mathbf{w}_2 \mathbf{y} \mathbf{w}_3 \mathbf{y} \mathbf{x}$$

because

$$\mathbf{w}_1 \mathbf{x} \mathbf{w}_2 \mathbf{y} \mathbf{w}_3 \mathbf{x} \mathbf{y} \overset{(3.1)}{\approx} \mathbf{w}_1 \mathbf{x} \mathbf{w}_2 \mathbf{y} \mathbf{w}_3 \mathbf{x} \mathbf{y} \mathbf{w}_3 \overset{(4.9)}{\approx} \mathbf{w}_1 \mathbf{x} \mathbf{w}_2 (\mathbf{y} \mathbf{w}_3)^2 \mathbf{x} \overset{(3.1)}{\approx} \mathbf{w}_1 \mathbf{x} \mathbf{w}_2 \mathbf{y} \mathbf{w}_3 \mathbf{y} \mathbf{x}.$$

By the hypothesis, $\mathbf{u}_3 = \mathbf{a}_0 z_1 \mathbf{a}_1 z_2 \mathbf{a}_2 \cdots z_n \mathbf{a}_n$ and $\mathbf{u}_4 = \mathbf{b}_1 z_{1\pi} \mathbf{b}_2 z_{2\pi} \cdots \mathbf{b}_n z_{n\pi} \mathbf{b}'$, where $\pi \in S_n$ and $\mathbf{a}_0, \mathbf{a}_1, \mathbf{b}_1, \mathbf{a}_2, \mathbf{b}_2, \dots, \mathbf{a}_n, \mathbf{b}_n$ do not contain first occurrences of letters in \mathbf{u} . In view of the saying in previous paragraph, the identities (3.1) and (4.9) imply the identities

$$\begin{aligned} \mathbf{u} &\approx \mathbf{u}_1 \mathbf{x} \mathbf{u}_2 \mathbf{y} \mathbf{u}_3 z_n \mathbf{x} \mathbf{y} (\mathbf{b}_1 z_{1\pi} \mathbf{b}_2 z_{2\pi} \cdots \mathbf{b}_n z_{n\pi})_{\{z_n\}} \mathbf{b}' \\ &\approx \mathbf{u}_1 \mathbf{x} \mathbf{u}_2 \mathbf{y} \mathbf{u}_3 z_n z_{n-1} \mathbf{x} \mathbf{y} (\mathbf{b}_1 z_{1\pi} \mathbf{b}_2 z_{2\pi} \cdots \mathbf{b}_n z_{n\pi})_{\{z_n, z_{n-1}\}} \mathbf{b}' \\ &\quad \vdots \\ &\approx \mathbf{u}_1 \mathbf{x} \mathbf{u}_2 \mathbf{y} \mathbf{u}_3 z_n z_{n-1} \cdots z_1 \mathbf{x} \mathbf{y} \mathbf{b}_1 \mathbf{b}_2 \cdots \mathbf{b}_n \mathbf{b}' \\ &\approx \mathbf{u}_1 \mathbf{x} \mathbf{u}_2 \mathbf{y} \mathbf{u}_3 z_n z_{n-1} \cdots z_1 \mathbf{y} \mathbf{x} \mathbf{b}_1 \mathbf{b}_2 \cdots \mathbf{b}_n \mathbf{b}'. \end{aligned}$$

By a similar argument we can show that (3.1) and (4.9) imply

$$\mathbf{u}_1 \mathbf{x} \mathbf{u}_2 \mathbf{y} \mathbf{u}_3 \mathbf{y} \mathbf{x} \mathbf{u}_4 \approx \mathbf{u}_1 \mathbf{x} \mathbf{u}_2 \mathbf{y} \mathbf{u}_3 z_n z_{n-1} \cdots z_1 \mathbf{y} \mathbf{x} \mathbf{b}_1 \mathbf{b}_2 \cdots \mathbf{b}_n \mathbf{b}'.$$

We see that $\mathbf{u} \approx \mathbf{u}_1 \mathbf{x} \mathbf{u}_2 \mathbf{y} \mathbf{u}_3 \mathbf{y} \mathbf{x} \mathbf{u}_4$ follows from (3.1) and (4.9). \square

Lemma 4.13. *Let \mathbf{V} be a monoid variety from the interval $[\mathbf{F}_1, \mathbf{P}]$ and $\mathbf{u} \approx \mathbf{v}$ be an identity of \mathbf{V} such that*

$$\mathbf{u} = \mathbf{w}_1 \overset{(1)}{x} \mathbf{w}_2 \overset{(1)}{y} \mathbf{w}_3 \overset{(2)}{x} \overset{(2)}{y} \mathbf{w}_4$$

for some words $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4$ and $(2_{\mathbf{v}}y) < (2_{\mathbf{v}}x)$. Suppose that the depth of the subwords \mathbf{w}_2 and \mathbf{w}_3 of \mathbf{u} is equal to m and k , respectively.

- (i) *If $k \geq 0$ and $m = 0$, then \mathbf{V} satisfies ζ_k .*
- (ii) *If $1 \leq m \leq k$, then \mathbf{V} satisfies λ_k^m .*

Proof. In view of Lemma 3.1, we may assume without loss of generality that the identity $\mathbf{u} \approx \mathbf{v}$ is 2-limited.

Let y_m be an m -divider of \mathbf{u} that occurs in \mathbf{w}_2 and let x_k be a k -divider of \mathbf{u} that occurs in \mathbf{w}_3 . In view of Lemma 2.3, $D(\mathbf{u}, x_k) = k$ and $D(\mathbf{u}, y_m) = m$.

If $k = 0$, then $m = 0$ as well. Then Corollary 2.13 and the inclusion $\mathbf{F}_1 \subset \mathbf{V}$ imply that the identity $\mathbf{u}(x, y, x_0, y_0) \approx \mathbf{v}(x, y, x_0, y_0)$ coincides with the identity ζ_0 , and we are done. So, we may assume that $k > 0$.

Suppose that the identity $\mathbf{u} \approx \mathbf{v}$ is not k -well-balanced. In view of Corollary 2.13 and the inclusion $\mathbf{F}_1 \subset \mathbf{V}$, the identity $\mathbf{u} \approx \mathbf{v}$ is 0-well-balanced. Then there is the smallest p such that $0 < p \leq k$ and the identity $\mathbf{u} \approx \mathbf{v}$ is not p -well-balanced. In view of Corollary 2.13, $\mathbf{F}_{p+1} \not\subseteq \mathbf{V}$. According to Lemma 3.3, \mathbf{V} satisfies κ_p . Then Lemmas 4.2, 4.3(ii) and 4.8(ii) imply that \mathbf{V} satisfies ζ_k and λ_k^r for any

$r = 1, 2, \dots, m$, and we are done. So, we may assume that the identity $\mathbf{u} \approx \mathbf{v}$ is k -well-balanced.

Now we apply Corollary 2.8 to deduce that there exist letters x_0, x_1, \dots, x_{k-1} such that

$$\begin{aligned} \mathbf{u} = & \mathbf{u}_{2k+1} \overset{(1)}{x_k} \mathbf{u}_{2k} \overset{(1)}{x_{k-1}} \mathbf{u}_{2k-1} \overset{(2)}{x_k} \mathbf{u}_{2k-2} \overset{(1)}{x_{k-2}} \mathbf{u}_{2s-3} \overset{(2)}{x_{k-1}} \mathbf{u}_{2k-4} \overset{(1)}{x_{k-3}} \\ & \cdot \mathbf{u}_{2k-5} \overset{(2)}{x_{k-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0 \end{aligned}$$

for some $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2k+1} \in \mathfrak{A}^*$, $D(\mathbf{u}, x_r) = r$ and \mathbf{u}_{2r+2} does not contain any r -dividers of \mathbf{u} for any $r = 0, 1, \dots, k-1$. The choice of x_k implies that both ${}_{2\mathbf{u}}x$ and ${}_{2\mathbf{u}}y$ lie in \mathbf{u}_{2k} . Put

$$X = \{x_0, x_1, \dots, x_k, x, y_m, y\}.$$

Suppose that $m = 0$. Then $y_0 \in \text{sim}(\mathbf{u}) = \text{sim}(\mathbf{v})$ and $\mathbf{u}(X) = xy_0yx_kxy \mathbf{b}_k$. Since the identity $\mathbf{u} \approx \mathbf{v}$ is k -well-balanced, $\mathbf{v}(X) = xy_0yx_kyx \mathbf{b}_k$ and, therefore, the identity $\mathbf{u}(X) \approx \mathbf{v}(X)$ coincides with ζ_k , we are done.

Suppose now that $1 \leq m \leq k$. Then $y_m \in \text{mul}(\mathbf{u}) = \text{mul}(\mathbf{v})$. According to the choice of y_m and x_k , the subwords \mathbf{w}_2 and \mathbf{w}_3 do not contain any $(m-1)$ -divider of \mathbf{u} . It follows from Lemma 2.9 that ${}_{2\mathbf{u}}y_m$ occurs in either \mathbf{u}_{2m} or \mathbf{u}_{2m-1} or \mathbf{u}_{2m-2} . Let $y_{m-1} = h_2^{m-1}(\mathbf{u}, y_m)$. Since \mathbf{u}_{2m} does not contain any $(m-1)$ -dividers of \mathbf{u} , the letter ${}_{1\mathbf{u}}y_{m-1}$ does not occur in \mathbf{u}_{2m} . Therefore, ${}_{2\mathbf{u}}y_m$ occurs in either \mathbf{u}_{2m-1} or \mathbf{u}_{2m-2} . Then

$$\mathbf{u}(X) = xy_myx_kxy \mathbf{b}_{k,m+1} x_{m-1} \mathbf{c} \mathbf{b}_{m-1},$$

where $\mathbf{c} \in \{x_my_m, y_mx_m\}$. Since $\mathbf{u} \approx \mathbf{v}$ is k -well-balanced, ${}_{2\mathbf{v}}y_m$ lies between ${}_{1\mathbf{v}}x_{m-1}$ and ${}_{1\mathbf{v}}x_{m-2}$ and

$$\mathbf{v}(X) = xy_myx_kyx \mathbf{b}_{k,m+1} x_{m-1} \mathbf{d} \mathbf{b}_{m-1},$$

where $\mathbf{d} \in \{x_my_m, y_mx_m\}$. We see that the identity $\mathbf{u}(X) \approx \mathbf{v}(X)$ coincides with

$$(4.10) \quad xy_myx_kxy \mathbf{b}_{k,m+1} x_{m-1} \mathbf{c} \mathbf{b}_{m-1} \approx xy_myx_kyx \mathbf{b}_{k,m+1} x_{m-1} \mathbf{d} \mathbf{b}_{m-1}.$$

Then \mathbf{V} satisfies λ_k^m because

$$\begin{aligned} & x \overset{(1)}{y_m} yx_kxy \cdots x_{m-1} \overset{(2)}{x_m} \overset{(2)}{y_m} x_{m-2} x_{m-1} \cdots x_1 x_2 x_0 x_1 \\ (3.1) \quad & \approx x \overset{(1)}{y_m} yx_kxy \cdots (x_{m-1} \overset{(2)}{x_m} \overset{(2)}{y_m}) \mathbf{c} x_{m-2} (x_{m-1} \overset{(4)}{x_m} \overset{(4)}{y_m}) \cdots x_1 x_2 x_0 x_1 \\ (4.10) \quad & \approx x \overset{(1)}{y_m} yx_kyx \cdots (x_{m-1} \overset{(2)}{x_m} \overset{(2)}{y_m}) \mathbf{d} x_{m-2} (x_{m-1} \overset{(4)}{x_m} \overset{(4)}{y_m}) \cdots x_1 x_2 x_0 x_1 \\ (3.1) \quad & \approx x \overset{(1)}{y_m} yx_kyx \cdots x_{m-1} \overset{(2)}{x_m} \overset{(2)}{y_m} x_{m-2} x_{m-1} \cdots x_1 x_2 x_0 x_1, \end{aligned}$$

and we are done. \square

Lemma 4.14. *Let \mathbf{V} be a monoid variety from the interval $[\mathbf{F}_1, \mathbf{P}]$ and $\mathbf{u} \approx \mathbf{v}$ be an identity of \mathbf{V} such that*

$$\mathbf{u} = \mathbf{w}_1 \overset{(1)}{x} \mathbf{w}_2 \overset{(1)}{y_{m+1}} \mathbf{w}_3 \overset{(1)}{y} \mathbf{w}_4 \overset{(2)}{x} \overset{(2)}{y} \mathbf{w}_5 \overset{(2)}{y_{m+1}} \mathbf{w}_6$$

for some words $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4, \mathbf{w}_5, \mathbf{w}_6$ and $({}_{2\mathbf{v}}x) < ({}_{2\mathbf{v}}y)$. Suppose that $({}_{2\mathbf{u}}a) < ({}_{2\mathbf{u}}y_{m+1})$ for any $a \in \text{con}(\mathbf{w}_2\mathbf{w}_3) \setminus \text{con}(\mathbf{w}_1)$. Suppose also that the depth of the subwords $\mathbf{w}_2y_{m+1}\mathbf{w}_3$, \mathbf{w}_4 and \mathbf{w}_5 of \mathbf{u} is equal to $m+1$, m and k , respectively.

- (i) If $k \geq 0$ and $m = 0$, then \mathbf{V} satisfies η_k .
- (ii) If $1 \leq m \leq k$, then \mathbf{V} satisfies μ_k^m .

Proof. If $\mathbf{LRB} \not\subseteq \mathbf{V}$, then $\mathbf{V} \subseteq \mathbf{F}$ by Lemma 3.2. Evidently, \mathbf{F} satisfies η_k and μ_k^m for any $k \geq 0$ and $m \leq k$. So, we may further assume that $\mathbf{LRB} \subseteq \mathbf{V}$. Further, Lemma 3.1 allows us to assume that the identity $\mathbf{u} \approx \mathbf{v}$ is 2-limited.

Let x_k be a k -divider of \mathbf{u} that occurs in \mathbf{w}_5 . In view of Lemma 2.3, $D(\mathbf{u}, x_k) = k$ and $D(\mathbf{u}, y_{m+1}) = m + 1$. Now by Corollary 2.8, there exist letters x_0, x_1, \dots, x_{k-1} such that $D(\mathbf{u}, x_r) = r$ and

$$\begin{aligned} \mathbf{u} = & \mathbf{u}_{2k+1} \overset{(1)}{x_k} \mathbf{u}_{2k} \overset{(1)}{x_{k-1}} \mathbf{u}_{2k-1} \overset{(2)}{x_k} \mathbf{u}_{2k-2} \overset{(1)}{x_{k-2}} \mathbf{u}_{2k-3} \overset{(2)}{x_{k-1}} \mathbf{u}_{2k-4} \overset{(1)}{x_{k-3}} \\ & \cdot \mathbf{u}_{2k-5} \overset{(2)}{x_{k-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0 \end{aligned}$$

for some $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2k+1} \in \mathfrak{A}^*$ such that \mathbf{u}_{2r+2} does not contain any r -dividers of \mathbf{u} for any $r = 0, 1, \dots, k-1$. Since the depth of \mathbf{w}_5 in \mathbf{u} is equal to k , the letter ${}_{2\mathbf{u}}y_{m+1}$ lies in \mathbf{u}_{2k} . Let y_m be an m -divider of \mathbf{u} that occurs in \mathbf{w}_4 . Clearly,

$$(1\mathbf{u}x) < (1\mathbf{u}y_{m+1}) < (1\mathbf{u}y) < (1\mathbf{u}y_m) < (2\mathbf{u}x) < (2\mathbf{u}y) < (1\mathbf{u}x_k).$$

Put

$$X = \{x_0, x_1, \dots, x_k, x, y_m, y_{m+1}, y\}.$$

There are two cases.

Case 1: the identity $\mathbf{u} \approx \mathbf{v}$ is k -well-balanced.

Suppose that $k \geq 0$ and $m = 0$. Then $y_m = y_0 \in \text{sim}(\mathbf{u}) = \text{sim}(\mathbf{v})$ and $\mathbf{u}(X) = xy_1yy_0xyx_ky_1\mathbf{b}_k$. Since the identity $\mathbf{u} \approx \mathbf{v}$ is k -well-balanced, we have $\mathbf{v}(X) = xy_1yy_0yx_ky_1\mathbf{b}_k$. Therefore, $\mathbf{u}(X) \approx \mathbf{v}(X)$ coincides with η_k , and we are done.

Suppose now that $1 \leq m \leq k$. Then $y_m \in \text{mul}(\mathbf{u}) = \text{mul}(\mathbf{v})$. Clearly, the subwords \mathbf{w}_4 and \mathbf{w}_5 do not contain any $(m-1)$ -divider of \mathbf{u} . It follows from Lemma 2.9 that ${}_{2\mathbf{u}}y_m$ occurs in either \mathbf{u}_{2m} or \mathbf{u}_{2m-1} or \mathbf{u}_{2m-2} . Let $y_{m-1} = h_2^{m-1}(\mathbf{u}, y_m)$. Since \mathbf{u}_{2m} does not contain any $(m-1)$ -dividers of \mathbf{u} , the letter ${}_{1\mathbf{u}}y_{m-1}$ does not occur in \mathbf{u}_{2m} . Therefore, ${}_{2\mathbf{u}}y_m$ occurs in either \mathbf{u}_{2m-1} or \mathbf{u}_{2m-2} . Then

$$\mathbf{u}(X) = xy_{m+1}yy_myx_ky_{m+1}\mathbf{b}_{k,m+1}x_{m-1}\mathbf{c}\mathbf{b}_{m-1},$$

where $\mathbf{c} \in \{x_my_m, y_mx_m\}$. Since $\mathbf{u} \approx \mathbf{v}$ is k -well-balanced, ${}_{2\mathbf{v}}y_m$ lies between ${}_{1\mathbf{v}}x_{m-1}$ and ${}_{1\mathbf{v}}x_{m-2}$ and

$$\mathbf{v}(X) = xy_{m+1}yy_myx_ky_{m+1}\mathbf{b}_{k,m+1}x_{m-1}\mathbf{d}\mathbf{b}_{m-1},$$

where $\mathbf{d} \in \{x_my_m, y_mx_m\}$. We see that the identity $\mathbf{u}(X) \approx \mathbf{v}(X)$ coincides with

$$(4.11) \quad \begin{aligned} & xy_{m+1}yy_myx_ky_{m+1}\mathbf{b}_{k,m+1}x_{m-1}\mathbf{c}\mathbf{b}_{m-1} \\ & \approx xy_{m+1}yy_myx_ky_{m+1}\mathbf{b}_{k,m+1}x_{m-1}\mathbf{d}\mathbf{b}_{m-1}. \end{aligned}$$

Then \mathbf{V} satisfies μ_k^m because

$$\begin{aligned} & x \overset{(1)}{y_{m+1}} y y_m x y x_k y_{m+1} \cdots x_{m-1} \overset{(2)}{x_m} \overset{(2)}{y_m} x_{m-2} x_{m-1} \cdots x_1 x_2 x_0 x_1 \\ (3.1) \quad & \approx x \overset{(1)}{y_{m+1}} y y_m x y x_k y_{m+1} \cdots (x_{m-1} \overset{(2)}{x_m} \overset{(2)}{y_m}) \mathbf{c} x_{m-2} (x_{m-1} \overset{(4)}{x_m} \overset{(4)}{y_m}) \cdots x_1 x_2 x_0 x_1 \\ (4.11) \quad & \approx x \overset{(1)}{y_{m+1}} y y_m y x x_k y_{m+1} \cdots (x_{m-1} \overset{(2)}{x_m} \overset{(2)}{y_m}) \mathbf{d} x_{m-2} (x_{m-1} \overset{(4)}{x_m} \overset{(4)}{y_m}) \cdots x_1 x_2 x_0 x_1 \\ (3.1) \quad & \approx x \overset{(1)}{y_{m+1}} y x_m y x x_k y_{m+1} \cdots x_{m-1} \overset{(2)}{x_m} \overset{(2)}{y_m} x_{m-2} x_{m-1} \cdots x_1 x_2 x_0 x_1, \end{aligned}$$

and we are done.

Case 2: the identity $\mathbf{u} \approx \mathbf{v}$ is not k -well-balanced. In view of Corollary 2.13 and the inclusion $\mathbf{F}_1 \subset \mathbf{V}$, the identity $\mathbf{u} \approx \mathbf{v}$ is 0-well-balanced. Then there is the smallest n such that $0 < n \leq k$ and the identity $\mathbf{u} \approx \mathbf{v}$ is not n -well-balanced. In view of Corollary 2.13, $\mathbf{F}_{n+1} \not\subset \mathbf{V}$. Then $\mathbf{F}_n \subseteq \mathbf{V}$. According to Lemma 3.3, \mathbf{V} satisfies κ_n .

Suppose that $k \geq 0$ and $m = 0$. Then $y_m = y_0 \in \text{sim}(\mathbf{u}) = \text{sim}(\mathbf{v})$ and $\mathbf{u}(X) = xy_1yy_0xyx_ky_1\mathbf{b}_k$. Lemma 2.10 and the inclusion $\mathbf{LRB} \subset \mathbf{V}$ together with the fact that the identity $\mathbf{u} \approx \mathbf{v}$ is $(n-1)$ -well-balanced imply $\mathbf{v}(X) = xy_1yy_0\mathbf{a}\mathbf{b}_n$ for some word \mathbf{a} with $\text{sim}(\mathbf{a}) = \{x_n, x, y_1, y\}$ and $\text{mul}(\mathbf{a}) = \{x_{n+1}, x_{n+2}, \dots, x_k\}$. According to Lemma 2.10, $\text{ini}(\mathbf{a}_{\{x,y\}}) = x_kx_{k-1} \cdots x_n$. Further, $\mathbf{a}(x, y) = yx$ because $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$. Then \mathbf{V} satisfies the identities

$$\begin{aligned} xy_1yy_0\mathbf{a}\mathbf{b}_n &\approx xy_1yy_0\mathbf{a}xyy_1\mathbf{b}_n && \text{by (3.1)} \\ &\approx xy_1yy_0\text{ini}(\mathbf{a}xyy_1)\text{ini}((\mathbf{a}xyy_1)_{x_n})\mathbf{b}_n && \text{by Lemma 4.6} \\ &= xy_1yy_0\text{ini}(\mathbf{a})\text{ini}(\mathbf{a}_{x_n})\mathbf{b}_n. \end{aligned}$$

Since $({}_{1\mathbf{a}}y) < ({}_{1\mathbf{a}}x)$, it follows from Proposition 4.7 that $\Phi(\mathbf{V})$ together with

$$(4.12) \quad xytxy \approx xytyx$$

imply the identity

$$\text{ini}(\mathbf{a})\text{ini}(\mathbf{a}_{x_n})\mathbf{b}_n \approx \mathbf{a}'\mathbf{b}_n,$$

where $\text{sim}(\mathbf{a}') = \{x_n\}$ and $\text{ini}(\mathbf{a}') = yxx_ky_1x_{k-1} \cdots x_n$. Now we apply Lemma 4.6 again and obtain that \mathbf{V} satisfies the identities

$$\begin{aligned} \mathbf{a}'\mathbf{b}_n &\approx \text{ini}(\mathbf{a}')\text{ini}(\mathbf{a}'_{x_n})\mathbf{b}_n = yxx_ky_1x_{k-1} \cdots x_n yxx_ky_1x_{k-1} \cdots x_{n+1}\mathbf{b}_n \\ &= \text{ini}(y^2x^2x_ky_1^2\mathbf{b}_{k,n+1})\text{ini}((y^2x^2x_ky_1^2\mathbf{b}_{k,n+1})_{x_n})\mathbf{b}_n \approx y^2x^2x_ky_1^2\mathbf{b}_{k,n+1}\mathbf{b}_n \\ &= y^2x^2x_ky_1^2\mathbf{b}_k. \end{aligned}$$

Then the identities

$$xy_1yy_0\mathbf{a}'\mathbf{b}_n \approx xy_1yy_0y^2x^2x_ky_1^2\mathbf{b}_k \stackrel{(3.1)}{\approx} xy_1yy_0yxx_ky_1\mathbf{b}_k$$

hold in \mathbf{V} . Therefore, \mathbf{V} satisfies

$$xy_1yy_0\mathbf{a}\mathbf{b}_n \approx xy_1yy_0yxx_ky_1\mathbf{b}_k$$

and so η_k .

Suppose now that $1 \leq m \leq k$. Then $y_m \in \text{mul}(\mathbf{u}) = \text{mul}(\mathbf{v})$. It is routine to verify that the depth of the letter y in both sides of the identity μ_k^m is equal to $m+1$. Thus, if $n \leq m$, then \mathbf{V} satisfies μ_k^m by Corollary 4.5. So, we may assume that $m < n$. Clearly, the subwords \mathbf{w}_4 and \mathbf{w}_5 do not contain any $(m-1)$ -divider of \mathbf{u} . It follows from Lemma 2.9 that ${}_{2\mathbf{u}}y_m$ occurs in either \mathbf{u}_{2m} or \mathbf{u}_{2m-1} or \mathbf{u}_{2m-2} . Let $y_{m-1} = h_2^{m-1}(\mathbf{u}, y_m)$. Since \mathbf{u}_{2m} does not contain any $(m-1)$ -divider of \mathbf{u} , the letter ${}_{1\mathbf{u}}y_{m-1}$ does not occur in \mathbf{u}_{2m} . Therefore, ${}_{2\mathbf{u}}y_m$ occurs in either \mathbf{u}_{2m-1} or \mathbf{u}_{2m-2} . Then

$$\mathbf{u}(X) = xy_{m+1}yy_mxyx_ky_{m+1}\mathbf{b}_{k,m+1}x_{m-1}\mathbf{c}\mathbf{b}_{m-1},$$

where $\mathbf{c} \in \{x_my_m, y_mx_m\}$. Lemma 2.10 and the inclusion $\mathbf{LRB} \subset \mathbf{V}$ together with the facts that $m < n$ and $\mathbf{u} \approx \mathbf{v}$ is $(n-1)$ -well-balanced imply that ${}_{2\mathbf{v}}y_m$ lies between ${}_{1\mathbf{v}}x_{m-1}$ and ${}_{1\mathbf{v}}x_{m-2}$ and

$$\mathbf{v}(X) = xy_{m+1}yy_m\mathbf{a}\mathbf{b}_{n,m+1}x_{m-1}\mathbf{d}\mathbf{b}_{m-1},$$

$\text{sim}(\mathbf{a}) = \{x_n, x, y, y_{m+1}\}$, $\text{mul}(\mathbf{a}) = \{x_{n+1}, x_{n+2}, \dots, x_k\}$ and $\mathbf{d} \in \{x_m y_m, y_m x_m\}$. As in the previous paragraph, one can show that \mathbf{V} satisfies the identity

$$xy_{m+1}yy_m \mathbf{a} \mathbf{b}_{n,m+1} x_{m-1} \mathbf{d} \mathbf{b}_{m-1} \approx xy_{m+1}yy_m y x x_k y_{m+1} \mathbf{b}_{k,m+1} x_{m-1} \mathbf{d} \mathbf{b}_{m-1}$$

and so the identity (4.11). As we have verified in Case 1, the last identity together with (3.1) imply μ_k^m . We see that the identity μ_k^m holds in \mathbf{V} . \square

Lemma 4.15. *Let \mathbf{V} be a variety from the interval $[\mathbf{LRB}, \mathbf{P}]$. Suppose that \mathbf{V} satisfies the identity*

$$(4.13) \quad xx_p y z x y z_\infty^2 z \mathbf{b}_p \approx xx_p \mathbf{c} \mathbf{b}_p,$$

where

$$(4.14) \quad \mathbf{c} = \mathbf{c}_1 y \mathbf{c}_2 y \mathbf{c}_3 x \mathbf{c}_4 z_\infty^2 \mathbf{c}_5 \quad \text{and} \quad \mathbf{c}_1 \mathbf{c}_2 \mathbf{c}_3 \mathbf{c}_4 \mathbf{c}_5 = z^2.$$

Then the identity ν_p holds in \mathbf{V} .

Proof. According to Lemma 2.10 and the inclusion $\mathbf{LRB} \subset \mathbf{V}$, $\text{ini}(\mathbf{c}_x) = yzz_\infty$. Therefore, $\mathbf{c}_1 = \lambda$ and $\mathbf{c}_2 \mathbf{c}_3 \mathbf{c}_4 \neq \lambda$. If $\mathbf{c}_2 = \lambda$, then it follows from Proposition 4.7 that $\Phi(\mathbf{V})$ implies the identity $xx_p \mathbf{c} \mathbf{b}_p \approx xx_p y z \mathbf{d} \mathbf{b}_p$, where \mathbf{d} is a word obtained from $y \mathbf{c}_3 x \mathbf{c}_4 z_\infty^2 \mathbf{c}_5$ by removing the first occurrence of z . So, we may assume that $z \in \text{con}(\mathbf{c}_2)$. If $z \in \text{con}(\mathbf{c}_5)$, then (4.13) is nothing but ν_p . So, it remains to consider the case when $\mathbf{c}_5 = \lambda$. In this case, we substitute $z_\infty z$ for z_∞ in the identity $\mathbf{u}(z, z_\infty) \approx \mathbf{v}(z, z_\infty)$. As a result, we obtain an identity that is equivalent modulo (3.1) to (3.2). Since $\text{ini}^2(\mathbf{c}) = y^2 z^2 x^2 z_\infty^2 = \text{ini}^2(yzyxz_\infty^2 z)$, Corollary 3.5 implies that \mathbf{V} satisfies the identity $xx_p \mathbf{c} \mathbf{b}_p \approx xx_p y z y x z_\infty^2 z \mathbf{b}_p$ and, therefore, the identity ν_p . \square

Let \mathbf{O}_1 denote the variety defined by the identities (3.1), (4.9) and

$$(4.15) \quad xzytxyz_\infty^2 z \approx xzytyxz_\infty^2 z.$$

Put

$$\Psi_1 = \{(3.1), (4.9), (4.15), \zeta_{k-1}, \lambda_k^m, \eta_{k-1}, \mu_k^m, \nu_{k-1} \mid k \in \mathbb{N}, 1 \leq m \leq k\}.$$

Proposition 4.16. *Let \mathbf{V} be a monoid variety from the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{O}_1]$, $\mathbf{u} \approx \mathbf{v}$ be an identity of \mathbf{V} and $\{2x, 2y\}$ be a critical pair in $\mathbf{u} \approx \mathbf{v}$. Then the critical pair $\{2x, 2y\}$ is Δ -removable for some $\Delta \subseteq \Psi_1(\mathbf{V})$.*

Proof. We consider only the case when $(1_{\mathbf{u}}x) < (1_{\mathbf{u}}y)$ since the case when $(1_{\mathbf{u}}y) < (1_{\mathbf{u}}x)$ is similar. Then $\mathbf{u} = \mathbf{w}_1 \overset{(1)}{x} \mathbf{w}_2 \overset{(1)}{y} \mathbf{w}_3 \overset{(2)}{x} \overset{(2)}{y} \mathbf{w}_4$ for some words $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4$. According to the inclusion $\mathbf{LRB} \subset \mathbf{V}$ and Lemma 2.10, $(1_{\mathbf{v}}x) < (1_{\mathbf{v}}y)$. Let \mathbf{w} be the word obtained from \mathbf{u} by replacing $2_{\mathbf{u}}x 2_{\mathbf{u}}y$ with $2_{\mathbf{u}}y 1_{\mathbf{u}}x$. Lemma 3.1 allows us to assume that the identity $\mathbf{u} \approx \mathbf{v}$ is 2-limited.

In view of Remark 2.2, there is a number d such that the d -decomposition of \mathbf{u} is maximal. Let \mathbf{a} denote the d -block of \mathbf{u} , which contains the critical pair $\{2x, 2y\}$. Then $\mathbf{a} = \mathbf{a}' \overset{(2)}{x} \overset{(2)}{y} \mathbf{a}''$ and $\mathbf{u} = \mathbf{u}' \mathbf{a} \mathbf{u}''$ for some $\mathbf{a}', \mathbf{a}', \mathbf{u}', \mathbf{u}'' \in \mathfrak{A}^*$. Clearly, $\text{con}(\mathbf{a}) \subseteq \text{mul}(\mathbf{u}' \mathbf{a})$.

If $\text{con}(\mathbf{w}_2) \subseteq \text{mul}(\mathbf{u}' \mathbf{a})$, then the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), (4.15)\} \subseteq \Psi_1(\mathbf{V})$ because

$$\mathbf{u} \overset{(3.1)}{\approx} \mathbf{w}_1 x \mathbf{w}_2 y \mathbf{w}_3 x y (\mathbf{a}'')^2 \mathbf{w}_2 \mathbf{u}'' \overset{(4.15)}{\approx} \mathbf{w}_1 x \mathbf{w}_2 y \mathbf{w}_3 y x (\mathbf{a}'')^2 \mathbf{w}_2 \mathbf{u}'' \overset{(3.1)}{\approx} \mathbf{w}.$$

So, we may assume that $\text{con}(\mathbf{w}_2) \not\subseteq \text{mul}(\mathbf{u}'\mathbf{a})$. Then there exist a letter x_p and a number $p \in \mathbb{N} \cup \{0\}$ such that the following claims hold:

- ${}_{1\mathbf{u}}x_p$ occurs in the subword \mathbf{w}_2 of \mathbf{u} ;
- $D(\mathbf{u}, x_p) = p$;
- the depth of \mathbf{w}_2 in \mathbf{u} is equal to p ;
- if $p > 0$, then ${}_{2\mathbf{u}}x_p$ occurs in the subword \mathbf{u}'' of \mathbf{u} .

There are two subcases.

Case 1: $D(\mathbf{u}, y) = \infty$. Then the first occurrence of y in \mathbf{u} lies in the d -block \mathbf{a} . This means that $\mathbf{a}' = \mathbf{a}_1 y \mathbf{w}_3$ for some $\mathbf{a}_1 \in \mathfrak{A}^*$ such that $y \notin \text{con}(\mathbf{a}_1)$.

Clearly, the depth of \mathbf{w}_3 in \mathbf{u} is infinite. If the depth of \mathbf{w}_3 in \mathbf{u} is strictly infinite, then the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), (4.9)\} \subseteq \Psi_1(\mathbf{V})$ by Lemma 4.12. So, it remains to consider the case when the depth of \mathbf{w}_3 in \mathbf{u} is not strictly infinite. Then there are letters z and z_∞ such that $({}_{1\mathbf{u}}z) < ({}_{1\mathbf{u}}z_\infty) < ({}_{2\mathbf{u}}z)$ and ${}_{1\mathbf{u}}z$ lies in the subword \mathbf{w}_3 of \mathbf{u} , while ${}_{1\mathbf{u}}z_\infty$ and ${}_{2\mathbf{u}}z$ lie in the subword \mathbf{a}'' of \mathbf{u} .

Now by Corollary 2.8, there exist letters x_0, x_1, \dots, x_{p-1} such that

$$\begin{aligned} \mathbf{u} = & \mathbf{u}_{2p+1} \overset{(1)}{x_p} \mathbf{u}_{2p} \overset{(1)}{x_{p-1}} \mathbf{u}_{2p-1} \overset{(2)}{x_p} \mathbf{u}_{2p-2} \overset{(1)}{x_{p-2}} \mathbf{u}_{2p-3} \overset{(2)}{x_{p-1}} \mathbf{u}_{2p-4} \overset{(1)}{x_{p-3}} \\ & \cdot \mathbf{u}_{2p-5} \overset{(2)}{x_{p-2}} \cdots \mathbf{u}_4 \overset{(1)}{x_1} \mathbf{u}_3 \overset{(2)}{x_2} \mathbf{u}_2 \overset{(1)}{x_0} \mathbf{u}_1 \overset{(2)}{x_1} \mathbf{u}_0 \end{aligned}$$

for some $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{2p+1} \in \mathfrak{A}^*$ with $D(\mathbf{u}, x_r) = r$. Clearly, $({}_{1\mathbf{u}}x_p) < ({}_{1\mathbf{u}}y) < ({}_{2\mathbf{u}}z) < ({}_{1\mathbf{u}}x_{p-1})$.

Corollary 2.13 and the inclusion $\mathbf{F}_1 \subset \mathbf{V}$ imply that the identity $\mathbf{u} \approx \mathbf{v}$ is 0-well-balanced. Then if the identity $\mathbf{u} \approx \mathbf{v}$ is not p -well-balanced, then there is the smallest k such that $0 < k \leq p$ and the identity $\mathbf{u} \approx \mathbf{v}$ is not k -well-balanced. In view of Corollary 2.13, $\mathbf{F}_{k+1} \not\subseteq \mathbf{V}$. According to Lemma 3.3, \mathbf{V} satisfies κ_k . Then Lemmas 4.2, 4.3(ii) and 4.8(i) imply that \mathbf{V} satisfies ν_p . Now we apply Lemma 4.11 and conclude that the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \nu_p\} \subseteq \Psi_1(\mathbf{V})$.

So, we may assume that the identity $\mathbf{u} \approx \mathbf{v}$ is p -well-balanced. Now we substitute yxz_∞^2 for z_∞ in the identity

$$\mathbf{u}(x_0, x_1, \dots, x_p, x, y, z, z_\infty) \approx \mathbf{v}(x_0, x_1, \dots, x_p, x, y, z, z_\infty)$$

Since $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$, we obtain an identity that is equivalent modulo (3.1) to the identity (4.13) such that (4.14) is true. Now we apply Lemma 4.15 and conclude that ν_p is satisfied by \mathbf{V} . Then the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \nu_p\} \subseteq \Psi_1(\mathbf{V})$ by Lemma 4.11.

Case 2: $D(\mathbf{u}, y) < \infty$, say $D(\mathbf{u}, y) = s + 1$ for some $s \in \mathbb{N} \cup \{0\}$. This means that the depth of the subword \mathbf{w}_3 in \mathbf{u} is equal to s . The choice of x_p implies that $p \leq s + 1$.

Suppose that $p \leq s$. If $p = 0$, then \mathbf{V} satisfies ζ_s by Lemma 4.13(i). Then we apply Lemma 4.9(i) and conclude that the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \zeta_s\} \subseteq \Psi_1(\mathbf{V})$. If $p > 0$, then \mathbf{V} satisfies λ_s^p by Lemma 4.13(ii). Then the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \lambda_s^p\} \subseteq \Psi_1(\mathbf{V})$ by Lemma 4.9(ii).

Suppose now that $p = s + 1$. Evidently, we may assume without loss of generality that $({}_{2\mathbf{u}}a) \leq ({}_{2\mathbf{u}}x_p)$ for any $a \in \text{con}(\mathbf{w}_2) \setminus \text{con}(\mathbf{w}_1)$. Let $\mathbf{u}'' = \mathbf{u}_1'' {}_{2\mathbf{u}}x_p \mathbf{u}_2''$. Since the first letter of \mathbf{u}'' is a d -divider of \mathbf{u} , the depth of the subword $\mathbf{a}''\mathbf{u}_1''$ in \mathbf{u} is equal to some number, say k . Clearly, $p - 1 = s \leq k$. If $p = 1$, then \mathbf{V} satisfies η_k by Lemma 4.14(i). Then we apply Lemma 4.10(i) and conclude that the identity $\mathbf{u} \approx \mathbf{w}$

follows from $\{(3.1), \eta_k\} \subseteq \Psi_1(\mathbf{V})$. If $p > 1$, then \mathbf{V} satisfies μ_k^s by Lemma 4.14(ii). Then the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \mu_k^s\} \subseteq \Psi_1(\mathbf{V})$ by Lemma 4.10(ii). \square

Let $\mathbf{O}_2 = \mathbf{P}\{(4.15), \nu_1\}$. Put

$$\Psi_2 = \{(3.1), (4.9), (4.15), \zeta_{k-1}, \lambda_k^m, \eta_{k-1}, \mu_k^m, \nu_0, \nu_1 \mid k \in \mathbb{N}, 1 \leq m \leq k\}.$$

Proposition 4.17. *Let \mathbf{V} be a monoid variety from the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{O}_2]$, $\mathbf{u} \approx \mathbf{v}$ be an identity of \mathbf{V} and $\{2x, 2y\}$ be a critical pair in $\mathbf{u} \approx \mathbf{v}$. Then the critical pair $\{2x, 2y\}$ is Δ -removable for some $\Delta \subseteq \Psi_2(\mathbf{V})$.*

Proof. We consider only the case when $(1_{\mathbf{u}}x) < (1_{\mathbf{u}}y)$ since that case when $(1_{\mathbf{u}}y) < (1_{\mathbf{u}}x)$ is similar. Then $\mathbf{u} = \mathbf{w}_1 \overset{(1)}{x} \mathbf{w}_2 \overset{(1)}{y} \mathbf{w}_3 \overset{(2)}{x} \overset{(2)}{y} \mathbf{w}_4$ for some words $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4$. The inclusion $\mathbf{LRB} \subset \mathbf{V}$ and Lemma 2.10 imply that $(1_{\mathbf{v}}x) < (1_{\mathbf{v}}y)$. Let \mathbf{w} be the word obtained from \mathbf{u} by replacing $2_{\mathbf{u}}x 2_{\mathbf{u}}y$ with $2_{\mathbf{u}}y 1_{\mathbf{u}}x$. Lemma 3.1 allows us to assume that the identity $\mathbf{u} \approx \mathbf{v}$ is 2-limited.

In view of Remark 2.2, there is a number d such that the d -decomposition of \mathbf{u} is maximal. Let \mathbf{a} denote the d -block of \mathbf{u} , which contains the critical pair $\{2x, 2y\}$. Then $\mathbf{a} = \mathbf{a}' 2_{\mathbf{u}}x 2_{\mathbf{u}}y \mathbf{a}''$ and $\mathbf{u} = \mathbf{u}' \mathbf{a} \mathbf{u}''$ for some $\mathbf{a}', \mathbf{a}'', \mathbf{u}', \mathbf{u}'' \in \mathfrak{A}^*$. Clearly, $\text{con}(\mathbf{a}) \subseteq \text{mul}(\mathbf{u}'\mathbf{a})$.

Case 1: $D(\mathbf{u}, y) = \infty$. Then the first occurrence of y in \mathbf{u} lies in the d -block \mathbf{a} . This means that $\mathbf{a}' = \mathbf{a}_1 y \mathbf{w}_3$ for some $\mathbf{a}_1 \in \mathfrak{A}^*$ such that $y \notin \text{con}(\mathbf{a}_1)$.

Suppose that \mathbf{w}_2 contains some simple letter t of \mathbf{u} . In view of the inclusion $\mathbf{F}_1 \subset \mathbf{V}$ and Corollary 2.13, the identity $\mathbf{u} \approx \mathbf{v}$ is 0-well-balanced. Hence $(1_{\mathbf{v}}x) < (1_{\mathbf{v}}t) < (1_{\mathbf{v}}y)$. Then the \mathbf{V} satisfies (4.9) because it coincides with $\mathbf{u}(x, y, t) \approx \mathbf{v}(x, y, t)$. Hence \mathbf{V} is a subvariety of \mathbf{O}_1 . Now Proposition 4.16 applies and we conclude that $\mathbf{u} \approx \mathbf{w}$ follows from $\Psi_1(\mathbf{V})$. In view of Lemma 4.8(i), $\mathbf{P}\{\nu_1\} \subseteq \mathbf{P}\{\nu_r\}$ for any $r \in \mathbb{N}$. Then, since ν_1 holds in \mathbf{V} , the identities in $\Psi_1(\mathbf{V})$ follow from $\Psi_2(\mathbf{V})$. Therefore, $\Psi_2(\mathbf{V})$ implies $\mathbf{u} \approx \mathbf{w}$.

So, it remains to consider the case when \mathbf{w}_2 does not contain simple letters of \mathbf{u} . Let ϕ be the substitution $(x_0, x_1, z_\infty, z) \mapsto (\mathbf{u}'', \mathbf{w}_2, \mathbf{a}'', \mathbf{w}_3)$. Then the identity $\phi(\nu_1)$ coincides with

$$(4.16) \quad x \mathbf{w}_2 y \mathbf{w}_3 x y (\mathbf{a}'')^2 \mathbf{w}_3 \mathbf{u}'' \mathbf{w}_2 \approx x \mathbf{w}_2 y \mathbf{w}_3 y x (\mathbf{a}'')^2 \mathbf{w}_3 \mathbf{u}'' \mathbf{w}_2.$$

Then the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), \nu_1\} \subseteq \Psi_2(\mathbf{V})$ because

$$\mathbf{u} \stackrel{(3.1)}{\approx} \mathbf{w}_1 x \mathbf{w}_2 y \mathbf{w}_3 x y (\mathbf{a}'')^2 \mathbf{w}_3 \mathbf{u}'' \mathbf{w}_2 \stackrel{(4.16)}{\approx} \mathbf{w}_1 x \mathbf{w}_2 y \mathbf{w}_3 y x (\mathbf{a}'')^2 \mathbf{w}_3 \mathbf{u}'' \mathbf{w}_2 \stackrel{(3.1)}{\approx} \mathbf{w},$$

and we are done.

Case 2: $D(\mathbf{u}, y) < \infty$, say $D(\mathbf{u}, y) = s + 1$ for some $s \in \mathbb{N} \cup \{0\}$. This means that the depth of the subword \mathbf{w}_3 in \mathbf{u} is equal to s . If $\text{con}(\mathbf{w}_2) \subseteq \text{mul}(\mathbf{u}'\mathbf{a})$, then the identity $\mathbf{u} \approx \mathbf{w}$ follows from $\{(3.1), (4.15)\} \subseteq \Psi_2(\mathbf{V})$ because

$$\mathbf{u} \stackrel{(3.1)}{\approx} \mathbf{w}_1 x \mathbf{w}_2 y \mathbf{w}_3 x y (\mathbf{a}'')^2 \mathbf{w}_2 \mathbf{u}'' \stackrel{(4.15)}{\approx} \mathbf{w}_1 x \mathbf{w}_2 y \mathbf{w}_3 y x (\mathbf{a}'')^2 \mathbf{w}_2 \mathbf{u}'' \stackrel{(3.1)}{\approx} \mathbf{w}.$$

So, we may assume that $\text{con}(\mathbf{w}_2) \not\subseteq \text{mul}(\mathbf{u}'\mathbf{a})$. Then there exist a letter x_p and a number $p \in \mathbb{N} \cup \{0\}$ such that $D(\mathbf{u}, x_p) = p$; the letters $1_{\mathbf{u}}x_p$ and $2_{\mathbf{u}}x_p$ occur in the subwords \mathbf{w}_2 and \mathbf{u}'' of \mathbf{u} , respectively; and the depth of \mathbf{w}_2 in \mathbf{u} is equal to p . The choice of x_p implies that $p \leq s + 1$. Further, by similar arguments as in the last two paragraphs of the proof of Proposition 4.16, one can show that $\mathbf{u} \approx \mathbf{v}$ follows from $\Phi_2(\mathbf{V})$. \square

Corollary 4.18. *For any $i = 1, 2$, each variety from the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{O}_i]$ can be defined within \mathbf{O}_i by identities from $\Phi \cup \Psi_i$. Consequently, the variety \mathbf{O}_i is HFB.*

Proof. Let \mathbf{V} be a variety from $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{O}_i]$. The inclusion $\mathbf{O}_i \subseteq \mathbf{P}$ and Lemmas 2.12 and 3.1 imply that \mathbf{V} can be defined within \mathbf{P} by a set of 2-limited balanced identities. In view of Lemma 2.10, $\text{ini}(\mathbf{u}) = \text{ini}(\mathbf{v})$ for any identity $\mathbf{u} \approx \mathbf{v}$ of \mathbf{V} . Then Lemma 4.1 and Propositions 4.7, 4.16 and 4.17 imply that the identity system $(\Phi \cup \Psi_i)(\mathbf{V})$ forms an identity basis for \mathbf{V} . It follows from Lemmas 4.3 and 4.8 that every subset of $\Phi \cup \Psi_i$ defines a FB subvariety of \mathbf{P} . Therefore, \mathbf{V} is FB.

So, it remains to show that if \mathbf{U} is a subvariety of \mathbf{O}_i that does not contain $\mathbf{LRB} \vee \mathbf{F}_1$, then \mathbf{U} is FB. Clearly, either $\mathbf{LRB} \not\subseteq \mathbf{U}$ or $\mathbf{F}_1 \not\subseteq \mathbf{U}$. Then by Lemma 3.2, either $\mathbf{U} \subseteq \mathbf{F}$ or $\mathbf{U} \subseteq \mathbf{LRB} \vee \mathbf{C}$. Since \mathbf{F} is HFB by Lemma 2.11 and $\mathbf{LRB} \vee \mathbf{C}$ is also HFB [20, Proposition 4.1], \mathbf{U} is HFB in any case. \square

5. LIMIT SUBVARIETIES OF \mathbf{P}

5.1. The limit variety \mathbf{J}_1 .

Lemma 5.1. *The variety $\mathbf{P}\{\kappa_1\}$ violates the identity (4.12).*

Proof. Let \mathbf{u} be a word such that $\mathbf{P}\{\kappa_1\}$ satisfies $xytxy \approx \mathbf{u}$ and $\text{ini}_2(\mathbf{u}) = xytxy$. In view of Proposition 2.1, it suffices to show that if the identity $\mathbf{u} \approx \mathbf{v}$ is directly deducible from some identity $\mathbf{s} \approx \mathbf{t} \in \{(3.1), \kappa_1\}$, that is, $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\xi(\mathbf{s})\mathbf{b}, \mathbf{a}\xi(\mathbf{t})\mathbf{b}\}$ for some words $\mathbf{a}, \mathbf{b} \in \mathfrak{A}^*$ and some endomorphism ξ of \mathfrak{A}^* , then $\text{ini}_2(\mathbf{v}) = xytxy$. Arguing by contradiction suppose that $\text{ini}_2(\mathbf{v}) \neq xytxy$. According to Lemmas 2.10 and 2.12 and the inclusion $\mathbf{LRB} \vee \mathbf{F}_1 \subset \mathbf{P}\{\kappa_1\}$, $\text{ini}_2(\mathbf{v}) = xytyx$. In particular, $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$.

Clearly, $\mathbf{s} \approx \mathbf{t}$ does not coincide with (3.1) because it does not change the first and the second occurrences of letters in a word. Suppose that $\mathbf{s} \approx \mathbf{t}$ coincides with κ_1 . We consider only the case when $(\mathbf{s}, \mathbf{t}) = (xx_1xx_0x_1, x^2x_1x_0x_1)$ because the other case is similar. Since the left side of κ_1 differs from the right side of κ_1 only in swapping of the first occurrence of x_1 and the second occurrence of x and $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$, there are three possibilities:

- $\xi({}_{1\mathbf{s}}x_1)$ contains ${}_{2\mathbf{u}}x$ and $\xi({}_{2\mathbf{s}}x)$ contains ${}_{2\mathbf{u}}y$;
- ${}_{2\mathbf{u}}x{}_{2\mathbf{u}}y$ is a subword of $\xi({}_{1\mathbf{s}}x_1)$ and $\xi({}_{2\mathbf{s}}x)$ contains some non-first and non-second occurrence of y ;
- $\xi({}_{1\mathbf{s}}x_1)$ contains ${}_{1\mathbf{u}}x$ and ${}_{2\mathbf{u}}x{}_{2\mathbf{u}}y$ is a subword of $\xi({}_{2\mathbf{s}}x)$.

In any case, $\xi({}_{1\mathbf{s}}x)$ contains ${}_{1\mathbf{u}}y$, while $\xi({}_{2\mathbf{s}}x)$ contains some non-first occurrence of y . Then $t \in \text{con}(\xi(x_1x))$ because $({}_{1\mathbf{u}}y) < ({}_{1\mathbf{u}}t) < ({}_{2\mathbf{u}}y)$. But this is impossible because $t \in \text{sim}(\mathbf{u})$, while $x, x_1 \in \text{mul}(\mathbf{s})$. Therefore, $\mathbf{s} \approx \mathbf{t}$ cannot coincide with κ_1 as well. \square

For arbitrary $n \in \mathbb{N}$ and $\pi, \tau \in S_{2n}$, we put

$$\begin{aligned} \mathbf{w}_n[\pi, \tau] &= \left(\prod_{i=1}^n x_i t_i \right) x \left(\prod_{i=1}^{2n} z_i \right) y \left(\prod_{i=n+1}^{2n} t_i x_i \right) txy \left(\prod_{i=1}^{2n} x_{i\pi} z_{i\tau} \right), \\ \mathbf{w}'_n[\pi, \tau] &= \left(\prod_{i=1}^n x_i t_i \right) x \left(\prod_{i=1}^{2n} z_i \right) y \left(\prod_{i=n+1}^{2n} t_i x_i \right) tyx \left(\prod_{i=1}^{2n} x_{i\pi} z_{i\tau} \right). \end{aligned}$$

Let $\mathbf{J}_1 = \text{var } \Omega_1$, where

$$\Omega_1 = \{(3.1), \kappa_1, \mathbf{w}_n[\pi, \tau] \approx \mathbf{w}'_n[\pi, \tau] \mid n \in \mathbb{N}, \pi, \tau \in S_{2n}\}.$$

Lemma 5.2. *The variety \mathbf{J}_1 violates the identity*

$$(5.1) \quad xsytxy \approx xsytyx.$$

Proof. Let \mathbf{u} be a word such that \mathbf{J}_1 satisfies $xsytxy \approx \mathbf{u}$ and $\text{ini}_2(\mathbf{u}) = xsytxy$. In view of Proposition 2.1, to verify that \mathbf{J}_1 violates (5.1), it suffices to show that if the identity $\mathbf{u} \approx \mathbf{v}$ is directly deducible from some identity $\mathbf{s} \approx \mathbf{t} \in \Omega_1$, that is, $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\xi(\mathbf{s})\mathbf{b}, \mathbf{a}\xi(\mathbf{t})\mathbf{b}\}$ for some words $\mathbf{a}, \mathbf{b} \in \mathfrak{A}^*$ and some endomorphism ξ of \mathfrak{A}^* , then $\text{ini}_2(\mathbf{v}) = xsytxy$. Arguing by contradiction suppose that $\text{ini}_2(\mathbf{v}) \neq xsytxy$. According to Lemma 2.12 and the inclusion $\mathbf{F}_1 \subset \mathbf{J}_1$, $\text{ini}_2(\mathbf{v}) = xsytyx$. In particular, $(2_{\mathbf{v}}y) < (2_{\mathbf{v}}x)$.

In view of Lemma 5.1, $\mathbf{s} \approx \mathbf{t}$ cannot coincide with (3.1) or κ_1 . Therefore, $\mathbf{s} \approx \mathbf{t}$ coincides with $\mathbf{w}_k[\pi, \tau] \approx \mathbf{w}'_k[\pi, \tau]$ for some $k \in \mathbb{N}$ and $\pi, \tau \in S_{2k}$. We consider only the case when $(\mathbf{s}, \mathbf{t}) = (\mathbf{w}_k[\pi, \tau], \mathbf{w}'_k[\pi, \tau])$ because the other case is similar. Since $\mathbf{w}_k[\pi, \tau]$ differs from $\mathbf{w}'_k[\pi, \tau]$ only in swapping of the second occurrences of letters x and y and $(2_{\mathbf{v}}y) < (2_{\mathbf{v}}x)$, there are three possibilities:

- (a) $\xi(2_{\mathbf{s}}x)$ contains $2_{\mathbf{u}}x$ and $\xi(2_{\mathbf{s}}y)$ contains $2_{\mathbf{u}}y$;
- (b) $2_{\mathbf{u}}x2_{\mathbf{u}}y$ is a subword of $\xi(2_{\mathbf{s}}x)$ and $\xi(2_{\mathbf{s}}y)$ contains some occurrence of y ;
- (c) $\xi(2_{\mathbf{s}}x)$ contains $1_{\mathbf{u}}x$ and $2_{\mathbf{u}}x2_{\mathbf{u}}y$ is a subword of $\xi(2_{\mathbf{s}}y)$.

Suppose that (a) holds. Then $\xi(1_{\mathbf{s}}x)$ contains $1_{\mathbf{u}}x$ and $\xi(1_{\mathbf{s}}y)$ contains $1_{\mathbf{u}}y$. Therefore $s \in \text{con}(\xi(xz_1z_2 \cdots z_{2k}y))$, contradicting $s \in \text{sim}(\mathbf{u})$ and $\{x, y, z_1, z_2, \dots, z_{2k}\} \subset \text{mul}(\mathbf{s})$. Suppose that (b) holds. Then $\xi(1_{\mathbf{s}}x)$ contains both $1_{\mathbf{u}}x$ and $1_{\mathbf{u}}y$. Hence $s \in \text{con}(\xi(x))$, which contradicts $s \in \text{sim}(\mathbf{u})$ and $x \in \text{mul}(\mathbf{s})$. Finally, (c) is impossible because $\xi(2_{\mathbf{s}}x)$ cannot contain any first occurrence of a letter in \mathbf{u} . We see that $\mathbf{s} \approx \mathbf{t}$ cannot coincide with $\mathbf{w}_k[\pi, \tau] \approx \mathbf{w}'_k[\pi, \tau]$ as well. Therefore, \mathbf{J}_1 violates (5.1). \square

Lemma 5.3. *Let \mathbf{V} be a subvariety of \mathbf{P} . If \mathbf{V} does not contain \mathbf{J}_1 , then \mathbf{V} satisfies the identity (4.15).*

Proof. If \mathbf{V} does not contain \mathbf{LRB} or \mathbf{F}_1 , then \mathbf{V} is contained in either \mathbf{F} or $\mathbf{LRB} \vee \mathbf{C}$ by Lemma 3.2. Evidently, both \mathbf{F} and $\mathbf{LRB} \vee \mathbf{C}$ satisfy (4.15). So, we may assume that \mathbf{V} belongs to the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}]$.

In view of Lemma 2.12 and the inclusion $\mathbf{F}_1 \subset \mathbf{V}$, we have $\text{sim}(\mathbf{a}) = \text{sim}(\mathbf{b})$ and $\text{mul}(\mathbf{a}) = \text{mul}(\mathbf{b})$ for any identity $\mathbf{a} \approx \mathbf{b}$ of \mathbf{V} . This fact and Lemma 3.1 imply that any identity of \mathbf{V} is equivalent to a 2-limited balanced identity. Then Lemma 4.1 implies that there exists a 2-limited balanced identity $\mathbf{u} \approx \mathbf{v}$ of \mathbf{V} with a critical pair $\{ix, jy\}$ that is not Ω_1 -removable. It follows from the inclusion $\mathbf{LRB} \subset \mathbf{V}$ and Lemma 2.10 that $(i, j) \neq (1, 1)$. Then $(i, j) \in \{(1, 2), (2, 1), (2, 2)\}$ because the identity $\mathbf{u} \approx \mathbf{v}$ is 2-limited.

Suppose that $\{i, j\} = \{1, 2\}$. According to Proposition 4.7, the identity $\mathbf{u} \approx \mathbf{v}$ may be chosen from the set

$$\Phi = \{(3.1), (3.2), \gamma_k, \delta_k^m, \varepsilon_{k-1} \mid k \in \mathbb{N}, 1 \leq m \leq k\}.$$

Evidently, $\mathbf{u} \approx \mathbf{v}$ does not coincide with (3.1) because it holds in \mathbf{J}_1 . Since \mathbf{J}_1 satisfies κ_1 , it also satisfies δ_k^m and ε_k for any $k \in \mathbb{N}$ and $1 \leq m \leq k$ by Lemmas 4.2 and 4.3(ii),(iii). Besides that, (3.2) holds in \mathbf{J}_1 because it is a consequence of κ_1 .

It follows that $\mathbf{u} \approx \mathbf{v}$ coincides with either ε_0 or γ_k for some $k \in \mathbb{N}$. It is easy to see that (4.15) is a consequence of ε_0 and so γ_k for any $k \in \mathbb{N}$.

So, it remains to consider the case when $(i, j) = (2, 2)$. By symmetry, we may assume that $(1\mathbf{u}x) < (1\mathbf{u}y)$. Then $(1\mathbf{v}x) < (1\mathbf{v}y)$ by Lemma 2.10 and the inclusion $\mathbf{LRB} \subset \mathbf{V}$. If there are no simple letters between $1\mathbf{u}y$ and $2\mathbf{u}y$, then $D(\mathbf{u}, y) > 1$ and so the critical pair $\{2x, 2y\}$ is $\{(3.1), \kappa_1\}$ -removable in $\mathbf{u} \approx \mathbf{v}$ by Corollary 4.5. So, we may assume that there is a simple letter t between $1\mathbf{u}y$ and $2\mathbf{u}y$.

If, for any letter a with

$$(1\mathbf{u}x) < (1\mathbf{u}a) < (1\mathbf{u}y) < (2\mathbf{u}y) < (2\mathbf{u}a),$$

there are no first occurrences of letters between $2\mathbf{u}y$ and $2\mathbf{u}a$, then it is easy to see that one can choose $n \in \mathbb{N}$ and $\pi, \tau \in S_{2n}$ so that the critical pair $\{2x, 2y\}$ is $\{(3.1), \mathbf{w}_n[\pi, \tau] \approx \mathbf{w}'_n[\pi, \tau]\}$ -removable in $\mathbf{u} \approx \mathbf{v}$. So, we may assume that there are letters z and z_∞ such that

$$(1\mathbf{u}x) < (1\mathbf{u}z) < (1\mathbf{u}y) < (2\mathbf{u}y) < (1\mathbf{u}z_\infty) < (2\mathbf{u}z).$$

Now we substitute yxz_∞^2 for z_∞ in the identity $\mathbf{u}(x, y, t, z, z_\infty) \approx \mathbf{v}(x, y, t, z, z_\infty)$. Since $(2\mathbf{v}y) < (2\mathbf{v}x)$, we obtain an identity that is equivalent modulo (3.1) to

$$xzytxyz_\infty^2 z \approx xzyt\mathbf{w},$$

where $\mathbf{w} \in \{zyxz_\infty^2, yzxz_\infty^2, yxzz_\infty^2, yxz_\infty^2 z\}$. If $\mathbf{w} = yxz_\infty^2 z$, then \mathbf{V} satisfies (4.15). So, it remains to consider the case when

$$\mathbf{w} \in \{zyxz_\infty^2, yzxz_\infty^2, yxzz_\infty^2\}.$$

In this case, the variety \mathbf{V} satisfies the identity

$$xzyt\mathbf{w} \approx xzytyxzz_\infty^2$$

because this identity is a consequence of the identity (4.12), which is evidently satisfied by \mathbf{V} . Now Proposition 4.7 applies and we conclude that the identity $xzytyxzz_\infty^2 \approx xzytyxz_\infty z z_\infty$ follows from $\Phi(\mathbf{V})$. Hence \mathbf{V} satisfies

$$(5.2) \quad xzytxyz_\infty^2 z \approx xzytyxz_\infty z z_\infty.$$

Finally, we substitute z_∞^2 for z_∞ in the identity (5.2). We obtain an identity that is equivalent modulo (3.1) to (4.15). We see that (4.15) is satisfied by \mathbf{V} in either case. \square

Proposition 5.4. *The variety \mathbf{J}_1 is a limit variety.*

Proof. According to Lemmas 4.2, 4.3(ii) and 4.8(i),(ii), ν_1 is satisfied by \mathbf{J}_1 . Then Lemma 5.3 implies that any proper subvariety of \mathbf{J}_1 is contained in \mathbf{O}_2 and so is FB by Corollary 4.18.

So, it remains to establish that \mathbf{J}_1 is NFB. It suffices to verify that, for any $n \in \mathbb{N}$, the set of identities

$$\Delta_n = \{(3.1), \kappa_1, \mathbf{w}_k[\pi, \tau] \approx \mathbf{w}'_k[\pi, \tau] \mid k < n, \pi, \tau \in S_{2k}\}$$

does not imply the identity $\mathbf{w}_n[\varepsilon, \varepsilon] \approx \mathbf{w}'_n[\varepsilon, \varepsilon]$, where ε is the trivial permutation on $\{1, 2, \dots, 2n\}$. Let \mathbf{w} be a word such that \mathbf{J}_1 satisfies $\mathbf{w}_n[\varepsilon, \varepsilon] \approx \mathbf{w}$. According to Lemmas 2.10 and 2.12 and the inclusion $\mathbf{LRB} \vee \mathbf{F}_1 \subseteq \mathbf{J}_1$,

$$\mathbf{w} = \left(\prod_{i=1}^n x_i t_i \right) x \left(\prod_{i=1}^{2n} z_i \right) y \left(\prod_{i=n+1}^{2n} t_i x_i \right) t \mathbf{w}'$$

for some word \mathbf{w}' with $\text{con}(\mathbf{w}') = \{x, y, x_1, z_1, x_2, z_2, \dots, x_{2n}, z_{2n}\}$. If ${}_{1\mathbf{w}'}x$ follows ${}_{1\mathbf{w}'}x_1$ in \mathbf{w}' , then the identity $\mathbf{w}_n[\varepsilon, \varepsilon](x, t, x_1, t_1) \approx \mathbf{w}(x, t, x_1, t_1)$ is equivalent modulo (3.1) to (5.1). But this is impossible by Lemma 5.2. Therefore, ${}_{1\mathbf{w}'}x$ precedes ${}_{1\mathbf{w}'}x_1$ in \mathbf{w}' . By a similar argument we can show that $({}_{1\mathbf{w}'}y) < ({}_{1\mathbf{w}'}x_1)$ and

$$({}_{1\mathbf{w}'}x_1) < ({}_{1\mathbf{w}'}z_1) < ({}_{1\mathbf{w}'}x_2) < ({}_{1\mathbf{w}'}z_2) < \dots < ({}_{1\mathbf{w}'}x_{2n}) < ({}_{1\mathbf{w}'}z_{2n}).$$

This means $\text{ini}(\mathbf{w}') = \mathbf{a}x_1z_1x_2z_2 \cdots x_{2n}z_{2n}$, where $\mathbf{a} \in \{xy, yx\}$. So, we have proved that $\text{ini}_2(\mathbf{w}) \in \{\mathbf{w}_n[\varepsilon, \varepsilon], \mathbf{w}'_n[\varepsilon, \varepsilon]\}$.

Let \mathbf{u} be a word such that \mathbf{J}_1 satisfies $\mathbf{w}_n[\varepsilon, \varepsilon] \approx \mathbf{u}$ and $\text{ini}_2(\mathbf{u}) = \mathbf{w}_n[\varepsilon, \varepsilon]$. According to Proposition 2.1, to verify that $\mathbf{w}_n[\varepsilon, \varepsilon] \approx \mathbf{w}'_n[\varepsilon, \varepsilon]$ does not follow from Δ_n , it suffices to establish that if the identity $\mathbf{u} \approx \mathbf{v}$ is directly deducible from some identity $\mathbf{s} \approx \mathbf{t}$ in Δ_n , that is, $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\xi(\mathbf{s})\mathbf{b}, \mathbf{a}\xi(\mathbf{t})\mathbf{b}\}$ for some words $\mathbf{a}, \mathbf{b} \in \mathfrak{A}^*$ and some endomorphism ξ of \mathfrak{A}^* , then $\text{ini}_2(\mathbf{v}) = \mathbf{w}_n[\varepsilon, \varepsilon]$. Arguing by contradiction suppose that $\text{ini}_2(\mathbf{v}) \neq \mathbf{w}_n[\varepsilon, \varepsilon]$. In view of the previous paragraph, $\text{ini}_2(\mathbf{v}) = \mathbf{w}'_n[\varepsilon, \varepsilon]$. In particular, $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$.

Since $\text{ini}_2(\mathbf{u}(x, y, t)) = xytxy$, Lemma 5.1 implies that $\mathbf{s} \approx \mathbf{t}$ cannot coincide with (3.1) and κ_1 . Suppose that $\mathbf{s} \approx \mathbf{t}$ coincides with $\mathbf{w}_k[\pi, \tau] \approx \mathbf{w}'_k[\pi, \tau]$ for some $k < n$ and $\pi, \tau \in S_{2k}$. We consider only the case when $(\mathbf{s}, \mathbf{t}) = (\mathbf{w}_k[\pi, \tau], \mathbf{w}'_k[\pi, \tau])$ because the other case is similar. Since $\mathbf{w}_k[\pi, \tau]$ differs from $\mathbf{w}'_k[\pi, \tau]$ only in swapping of the second occurrences of letters x and y and $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$, there are three possibilities:

- (a) $\xi({}_{2\mathbf{s}}x)$ contains ${}_{2\mathbf{u}}x$ and $\xi({}_{2\mathbf{s}}y)$ contains ${}_{2\mathbf{u}}y$;
- (b) ${}_{2\mathbf{u}}x{}_{2\mathbf{u}}y$ is a subword of $\xi({}_{2\mathbf{s}}x)$ and $\xi({}_{2\mathbf{s}}y)$ contains some occurrence of y ;
- (c) $\xi({}_{2\mathbf{s}}x)$ contains ${}_{1\mathbf{u}}x$ and ${}_{2\mathbf{u}}x{}_{2\mathbf{u}}y$ is a subword of $\xi({}_{2\mathbf{s}}y)$.

Suppose that (a) holds. Then $\xi({}_{1\mathbf{s}}x)$ contains ${}_{1\mathbf{u}}x$ and $\xi({}_{1\mathbf{s}}y)$ contains ${}_{1\mathbf{u}}y$. Hence $t_n, z_1 \notin \text{con}(\xi(x))$ and $z_{2n}, t_{n+1} \notin \text{con}(\xi(y))$ because $\text{ini}_2(\mathbf{u}) \neq \mathbf{w}_n[\varepsilon, \varepsilon]$ otherwise. Therefore, $\xi(x) = x$ and $\xi(y) = y$. It follows that $\xi(\prod_{i=1}^{2k} z_i) = \prod_{i=1}^{2n} z_i$. Then $\xi({}_{1\mathbf{s}}z_p)$ contains ${}_{1\mathbf{u}}z_q {}_{1\mathbf{u}}z_{q+1}$ for some $p \leq 2k$ and $q < 2n$. Since $\text{ini}_2(\mathbf{u}) = \mathbf{w}_n[\varepsilon, \varepsilon]$, the word $\xi({}_{2\mathbf{s}}z_p)$ does not contain ${}_{2\mathbf{u}}z_q {}_{2\mathbf{u}}z_{q+1}$. Hence ${}_{2\mathbf{u}}z_q$ is contained in the subword $\xi(\prod_{i=1}^{r-1} (x_{i\pi} z_{i\tau}) x_{r\pi})$ of \mathbf{u} , where $r\tau = p$. Then ${}_{1\mathbf{u}}z_q$ must be contained in $\xi({}_{1\mathbf{s}}a)$ for some $a \in \{z_{1\tau}, x_{1\pi}, z_{2\tau}, x_{2\pi}, \dots, z_{(r-1)\tau}, x_{(r-1)\pi}, x_{r\pi}\}$. This contradicts the fact that ${}_{1\mathbf{u}}z_q$ is contained in $\xi({}_{1\mathbf{s}}z_p)$. Therefore, (a) is impossible. Suppose that (b) holds. Then $\xi({}_{1\mathbf{s}}x)$ contains both ${}_{1\mathbf{u}}x$ and ${}_{1\mathbf{u}}y$. This is only possible when $xz_1z_2 \cdots z_{2n}y$ is a subword of $\xi(x)$. But this contradicts the fact that $\text{ini}_2(\mathbf{u}) = \mathbf{w}_n[\varepsilon, \varepsilon]$ and, therefore, (b) is impossible as well. Finally, (c) is impossible because $\xi({}_{2\mathbf{s}}x)$ cannot contain any first occurrence of a letter in \mathbf{u} . We see that $\mathbf{s} \approx \mathbf{t}$ cannot coincide with $\mathbf{w}_k[\pi, \tau] \approx \mathbf{w}'_k[\pi, \tau]$. So, we have proved that Δ_n does not imply $\mathbf{w}_n[\varepsilon, \varepsilon] \approx \mathbf{w}'_n[\varepsilon, \varepsilon]$ and so \mathbf{J}_1 is NFB. \square

5.2. The limit variety \mathbf{J}_2 . For arbitrary $n \in \mathbb{N}$ and $\pi, \tau \in S_{2n}$, define

$$\begin{aligned} \mathbf{z}_n[\pi, \tau] &= \left(\prod_{i=1}^{2n} x_i t_i \right) x \left(\prod_{i=1}^n z_i s_i \right) y \left(\prod_{i=1}^n z_{n+i} \right) xy \left(\prod_{i=1}^{2n} x_{i\pi} z_{i\tau} \right) t \left(\prod_{i=1}^n s_i \right), \\ \mathbf{z}'_n[\pi, \tau] &= \left(\prod_{i=1}^{2n} x_i t_i \right) x \left(\prod_{i=1}^n z_i s_i \right) y \left(\prod_{i=1}^n z_{n+i} \right) yx \left(\prod_{i=1}^{2n} x_{i\pi} z_{i\tau} \right) t \left(\prod_{i=1}^n s_i \right). \end{aligned}$$

Let $\mathbf{J}_2 = \text{var } \Omega_2$, where

$$\Omega_2 = \{(3.1), \eta_1, \mathbf{z}_n[\pi, \tau] \approx \mathbf{z}'_n[\pi, \tau] \mid n \in \mathbb{N}, \pi, \tau \in S_{2n}\}.$$

Lemma 5.5. *Let $\mathbf{u} \approx \mathbf{v}$ be an identity of \mathbf{J}_2 . Then $(1_{\mathbf{u}}x) < (2_{\mathbf{u}}y)$ if and only if $(1_{\mathbf{v}}x) < (2_{\mathbf{v}}y)$ for any letters x and y .*

Proof. In view of Proposition 2.1, it suffices to show that the required conclusion is true for an identity $\mathbf{u} \approx \mathbf{v}$ of \mathbf{J}_2 that is directly deducible from some identity $\mathbf{s} \approx \mathbf{t} \in \Omega_2$, that is, $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\xi(\mathbf{s})\mathbf{b}, \mathbf{a}\xi(\mathbf{t})\mathbf{b}\}$ for some words $\mathbf{a}, \mathbf{b} \in \mathfrak{A}^*$ and some endomorphism ξ of \mathfrak{A}^* .

If $\mathbf{s} \approx \mathbf{t}$ coincides with (3.1), then the required conclusion is true because (3.1) does not change the first and the second occurrences of letters in a word. Suppose that $\mathbf{s} \approx \mathbf{t}$ coincides with some identity from $\Omega_2 \setminus \{(3.1)\}$. Then \mathbf{s} differs from \mathbf{t} only in swapping of the second occurrences of x and y . Clearly, both $\xi(2_{\mathbf{s}}x)$ and $\xi(2_{\mathbf{s}}y)$ cannot contain first occurrences of letters. It follows that if $(1_{\mathbf{u}}x) < (2_{\mathbf{u}}y)$ for some letters x and y , then $(1_{\mathbf{v}}x) < (2_{\mathbf{v}}y)$. \square

Lemma 5.6. *The variety $\mathbf{P}\{\eta_1\}$ violates the identity*

$$(5.3) \quad xx_1yxyx_0x_1 \approx xx_1y^2xx_0x_1.$$

Proof. Let \mathbf{u} be a word such that $\mathbf{P}\{\eta_1\}$ satisfies $xx_1yxyx_0x_1 \approx \mathbf{u}$ and $\text{ini}_2(\mathbf{u}) = xx_1yxyx_0x_1$. In view of Proposition 2.1, it suffices to establish that if the identity $\mathbf{u} \approx \mathbf{v}$ is directly deducible from some identity $\mathbf{s} \approx \mathbf{t} \in \{(3.1), \eta_1\}$, that is $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\xi(\mathbf{s})\mathbf{b}, \mathbf{a}\xi(\mathbf{t})\mathbf{b}\}$ for some words $\mathbf{a}, \mathbf{b} \in \mathfrak{A}^*$ and some endomorphism ξ of \mathfrak{A}^* , then $\text{ini}_2(\mathbf{v}) = xx_1yxyx_0x_1$. Arguing by contradiction suppose that $\text{ini}_2(\mathbf{v}) \neq xx_1yxyx_0x_1$. According to Lemmas 2.10, 2.12 and 5.5 and the inclusions $\mathbf{LRB} \vee \mathbf{F}_1 \subseteq \mathbf{J}_2 \subseteq \mathbf{P}\{\eta_1\}$, we have $\text{ini}_2(\mathbf{v}) = xx_1y^2xx_0x_1$. In particular, $(2_{\mathbf{v}}y) < (2_{\mathbf{v}}x)$.

Clearly, $\mathbf{s} \approx \mathbf{t}$ does not coincide with (3.1) because it does not change the first and the second occurrences of letters in a word. Suppose that $\mathbf{s} \approx \mathbf{t}$ coincides with η_1 . We consider only the case when

$$(\mathbf{s}, \mathbf{t}) = (xy_1yy_0xyx_1y_1x_0x_1, xy_1yy_0yxx_1y_1x_0x_1)$$

because the other case is similar. Since the left side of η_1 differs from the right side of η_1 only in swapping of the second occurrences of x and y and $(2_{\mathbf{v}}y) < (2_{\mathbf{v}}x)$, there are three possibilities:

- (a) $\xi(2_{\mathbf{s}}x)$ contains $2_{\mathbf{u}}x$ and $\xi(2_{\mathbf{s}}y)$ contains $2_{\mathbf{u}}y$;
- (b) $2_{\mathbf{u}}x2_{\mathbf{u}}y$ is a subword of $\xi(2_{\mathbf{s}}x)$ and $\xi(2_{\mathbf{s}}y)$ contains some non-first and non-second occurrence of y ;
- (c) $\xi(2_{\mathbf{s}}x)$ contains $1_{\mathbf{u}}x$ and $2_{\mathbf{u}}x2_{\mathbf{u}}y$ is a subword of $\xi(2_{\mathbf{s}}y)$.

Suppose that (a) holds. Then $\xi(1_{\mathbf{s}}x)$ contains $1_{\mathbf{u}}x$ and $\xi(1_{\mathbf{s}}y)$ contains $1_{\mathbf{u}}y$. It follows that $x_1 \in \text{con}(\xi(xy_1y))$. Clearly, $x_1 \notin \text{con}(xy)$ because otherwise, some occurrence of x_1 lies between $2_{\mathbf{u}}x$ and $2_{\mathbf{u}}y$, contradicting $\text{ini}_2(\mathbf{u}) = xx_1yxyx_0x_1$. Therefore, $1_{\mathbf{u}}x_1$ coincides with $\xi(1_{\mathbf{s}}y_1)$ and so $\xi(x) = x$, $\xi(y) = y$ and $\xi(y_1) = x_1$. It follows that $x_0 \in \text{con}(\xi(x_1))$, contradicting $x_0 \in \text{sim}(\mathbf{u})$ and $x_1 \in \text{mul}(\mathbf{s})$. Suppose now that (b) holds, then $\xi(1_{\mathbf{s}}x)$ contains both $1_{\mathbf{u}}x$ and $1_{\mathbf{u}}y$. It follows that $x_1 \in \text{con}(\xi(x))$. But this is impossible because there are no occurrences of x_1 between $2_{\mathbf{u}}x$ and $2_{\mathbf{u}}y$. Finally, (c) is impossible because $\xi(2_{\mathbf{s}}x)$ cannot contain any first occurrence of a letter in \mathbf{u} . We see that $\mathbf{s} \approx \mathbf{t}$ cannot coincide with η_1 as well. Therefore, $\mathbf{P}\{\eta_1\}$ violates (5.3). \square

Lemma 5.7. *The variety \mathbf{J}_2 violates the identity (4.9).*

Proof. Let \mathbf{u} be a word such that \mathbf{J}_2 satisfies $xyxy \approx \mathbf{u}$ and $\text{ini}_2(\mathbf{u}) = xyxy$. In view of Proposition 2.1, to verify that \mathbf{J}_2 violates (4.9), it suffices to show that if the identity $\mathbf{u} \approx \mathbf{v}$ is directly deducible from some identity $\mathbf{s} \approx \mathbf{t} \in \Omega_2$, that is $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\xi(\mathbf{s})\mathbf{b}, \mathbf{a}\xi(\mathbf{t})\mathbf{b}\}$ for some words $\mathbf{a}, \mathbf{b} \in \mathfrak{A}^*$ and some endomorphism ξ of \mathfrak{A}^* , then $\text{ini}_2(\mathbf{v}) = xyxy$. Arguing by contradiction suppose that $\text{ini}_2(\mathbf{v}) \neq xyxy$. According to Lemmas 2.10, 2.12 and 5.5 and the inclusion $\mathbf{LRB} \vee \mathbf{F}_1 \subset \mathbf{J}_2$, $\text{ini}_2(\mathbf{v}) = xty^2x$. In particular, $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$.

Since $\text{ini}_2(\mathbf{u}) \approx \text{ini}_2(\mathbf{v})$ coincides with (4.9), in view of Lemma 5.6 and the fact that (4.9) implies (5.3), the identity $\mathbf{s} \approx \mathbf{t}$ cannot coincide with (3.1) or η_1 . Therefore, $\mathbf{s} \approx \mathbf{t}$ coincides with $\mathbf{z}_k[\pi, \tau] \approx \mathbf{z}'_k[\pi, \tau]$ for some $k \in \mathbb{N}$ and $\pi, \tau \in S_{2k}$. We consider only the case when $(\mathbf{s}, \mathbf{t}) = (\mathbf{z}_k[\pi, \tau], \mathbf{z}'_k[\pi, \tau])$ because the other case is similar. Since $\mathbf{z}_k[\pi, \tau]$ differs from $\mathbf{z}'_k[\pi, \tau]$ only in swapping of the second occurrences of letters x and y and $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$, there are three possibilities:

- (a) $\xi({}_{2\mathbf{s}}x)$ contains ${}_{2\mathbf{u}}x$ and $\xi({}_{2\mathbf{s}}y)$ contains ${}_{2\mathbf{u}}y$;
- (b) ${}_{2\mathbf{u}}x{}_{2\mathbf{u}}y$ is a subword of $\xi({}_{2\mathbf{s}}x)$ and $\xi({}_{2\mathbf{s}}y)$ contains some occurrence of y ;
- (c) $\xi({}_{2\mathbf{s}}x)$ contains ${}_{1\mathbf{u}}x$ and ${}_{2\mathbf{u}}x{}_{2\mathbf{u}}y$ is a subword of $\xi({}_{2\mathbf{s}}y)$.

Suppose that (a) holds. Then $\xi({}_{1\mathbf{s}}x)$ contains ${}_{1\mathbf{u}}x$ and $\xi({}_{1\mathbf{s}}y)$ contains ${}_{1\mathbf{u}}y$, so that $t \in \text{con}(\xi(xz_1s_1z_2s_2 \cdots z_k s_k y))$, contradicting $\{x, y, z_1, s_1, \dots, z_k, s_k\} \subset \text{mul}(\mathbf{s})$ and $t \in \text{sim}(\mathbf{u})$. Suppose that (b) holds. Then $\xi({}_{1\mathbf{s}}x)$ contains both ${}_{1\mathbf{u}}x$ and ${}_{1\mathbf{u}}y$, so that $t \in \text{con}(\xi(x))$, contradicting $t \in \text{sim}(\mathbf{u})$ and $x \in \text{mul}(\mathbf{s})$. Finally, (c) is impossible because $\xi({}_{2\mathbf{s}}x)$ cannot contain any first occurrence of a letter in \mathbf{u} . We see that $\mathbf{s} \approx \mathbf{t}$ cannot coincide with $\mathbf{z}_k[\pi, \tau] \approx \mathbf{z}'_k[\pi, \tau]$ as well. Therefore, \mathbf{J}_2 violates (4.9). \square

Lemma 5.8. *Let \mathbf{V} be a subvariety of \mathbf{P} . If \mathbf{V} does not contain \mathbf{J}_2 , then \mathbf{V} satisfies one of the identities (4.9) or ν_1 .*

Proof. If \mathbf{V} does not contain \mathbf{LRB} or \mathbf{F} , then \mathbf{V} is contained in either \mathbf{F} or $\mathbf{P}\{\kappa_r\}$ for some $r \in \mathbb{N}$ by Lemmas 3.2(i), 2.11 and 3.3. Evidently, \mathbf{F} satisfies the identity (4.9), while $\mathbf{P}\{\kappa_r\}$ satisfies this identity by Corollary 4.5. So, we may assume that \mathbf{V} belongs to the interval $[\mathbf{LRB} \vee \mathbf{F}, \mathbf{P}]$. In particular, Corollary 2.13 implies that every identity of \mathbf{V} is r -well-balanced for any $r \geq 0$. In view of this fact and Lemma 3.1, any identity of \mathbf{V} is equivalent to a 2-limited balanced identity. Then Lemma 4.1 implies that there exist a 2-limited balanced identity $\mathbf{u} \approx \mathbf{v}$ of \mathbf{V} and a critical pair $\{{}_i x, {}_j y\}$ in $\mathbf{u} \approx \mathbf{v}$ that is not Ω_2 -removable in $\mathbf{u} \approx \mathbf{v}$. It follows from the inclusion $\mathbf{LRB} \subset \mathbf{V}$ and Lemma 2.10 that $(i, j) \neq (1, 1)$. Then $(i, j) \in \{(1, 2), (2, 1), (2, 2)\}$ because the identity $\mathbf{u} \approx \mathbf{v}$ is 2-limited.

Suppose that $\{i, j\} = \{1, 2\}$. According to Proposition 4.7, the identity $\mathbf{u} \approx \mathbf{v}$ may be chosen from the set

$$\Phi = \{(3.1), (3.2), \gamma_k, \delta_k^m, \varepsilon_{k-1} \mid k \in \mathbb{N}, 1 \leq m \leq k\}.$$

Evidently, $\mathbf{u} \approx \mathbf{v}$ does not coincide with (3.1) because it holds in \mathbf{J}_2 . It follows from Lemma 4.3 that every identity from $\Phi \setminus \{(3.1)\}$ together with (3.1) imply the identity (3.2). It is routine to verify that (4.9) follows from $\{(3.1), (3.2)\}$. Therefore, (4.9) is satisfied by \mathbf{V} , and we are done.

So, it remains to consider the case when $(i, j) = (2, 2)$. By symmetry, we may assume that $({}_{1\mathbf{u}}x) < ({}_{1\mathbf{u}}y)$. Then $({}_{1\mathbf{v}}x) < ({}_{1\mathbf{v}}y)$ by Lemma 2.10 and the inclusion $\mathbf{LRB} \subset \mathbf{V}$. If there is a simple letter t between ${}_{1\mathbf{u}}x$ and ${}_{1\mathbf{u}}y$, then the identity $\mathbf{u}(x, y, t) \approx \mathbf{v}(x, y, t)$ coincides with (4.9) because $\mathbf{u} \approx \mathbf{v}$ is 0-well-balanced. So, we

may assume that there are no simple letters between ${}_{1\mathbf{u}}x$ and ${}_{1\mathbf{u}}y$. If, for any letter a with

$$({}_{1\mathbf{u}}x) < ({}_{1\mathbf{u}}a) < ({}_{1\mathbf{u}}y) < ({}_{2\mathbf{u}}y) < ({}_{2\mathbf{u}}a),$$

there are no simple letters between ${}_{2\mathbf{u}}y$ and ${}_{2\mathbf{u}}a$, then it is routine to show that the critical pair $\{{}_{2x}, {}_{2y}\}$ is $\{(3.1), \eta_1\}$ -removable. So, we may assume that there are letters $x_0 \in \text{sim}(\mathbf{u})$ and $x_1 \in \text{mul}(\mathbf{u})$ such that

$$({}_{1\mathbf{u}}x) < ({}_{1\mathbf{u}}x_1) < ({}_{1\mathbf{u}}y) < ({}_{2\mathbf{u}}y) < ({}_{1\mathbf{u}}x_0) < ({}_{2\mathbf{u}}x_1).$$

If there is a simple letter t between ${}_{1\mathbf{u}}y$ and ${}_{2\mathbf{u}}x$, then the identity $\mathbf{u}(x_0, x_1, x, y, t) \approx \mathbf{v}(x_0, x_1, x, y, t)$ coincides with the identity η_0 (up to renaming of letters) because $\mathbf{u} \approx \mathbf{v}$ is 1-well-balanced. It is easy to see that η_0 implies ν_1 , and we are done. So, it remains to consider the case when there are no simple letters between ${}_{1\mathbf{u}}y$ and ${}_{2\mathbf{u}}x$. If, for any letter a with

$$({}_{1\mathbf{u}}y) < ({}_{1\mathbf{u}}a) < ({}_{2\mathbf{u}}x) < ({}_{2\mathbf{u}}y) < ({}_{2\mathbf{u}}a),$$

there are no first occurrences of letters between ${}_{2\mathbf{u}}y$ and ${}_{2\mathbf{u}}a$, then it is easy to see that one can choose $n \in \mathbb{N}$ and $\pi, \tau \in S_{2n}$ so that the critical pair $\{{}_{2x}, {}_{2y}\}$ is $\{(3.1), \mathbf{z}_n[\pi, \tau] \approx \mathbf{z}'_n[\pi, \tau]\}$ -removable in $\mathbf{u} \approx \mathbf{v}$. So, we may assume that there are letters z and z_∞ such that

$$({}_{1\mathbf{u}}y) < ({}_{1\mathbf{u}}z) < ({}_{2\mathbf{u}}x) < ({}_{2\mathbf{u}}y) < ({}_{1\mathbf{u}}z_\infty) < ({}_{2\mathbf{u}}z).$$

We may assume without loss of generality that $({}_{1\mathbf{u}}z_\infty) < ({}_{2\mathbf{u}}z) < ({}_{1\mathbf{u}}x_0)$ because we can substitute $z_\infty z x_0$ for x_0 . Then we substitute $yz z_\infty^2$ for z_∞ in the identity

$$\mathbf{u}(x_0, x_1, x, y, z_\infty, z) \approx \mathbf{v}(x_0, x_1, x, y, z_\infty, z).$$

Since the identity $\mathbf{u} \approx \mathbf{v}$ is 1-well-balanced and $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$, we obtain an identity, which is equivalent modulo (3.1) to

$$xx_1 y z x y z_\infty^2 z x_0 x_1 \approx xx_1 c x_0 x_1$$

such that (4.14) is true. Now Lemma 4.15 applies and we conclude that ν_1 is satisfied by \mathbf{V} . We see that \mathbf{V} satisfies either ν_1 or (4.9) in either case. \square

Proposition 5.9. *The variety \mathbf{J}_2 is a limit variety.*

Proof. It is routine to verify that \mathbf{J}_2 satisfies (4.15). Then Lemma 5.8 implies that any proper subvariety of \mathbf{J}_2 is contained in either \mathbf{O}_1 or \mathbf{O}_2 and so is FB by Corollary 4.18.

So, it remains to establish that \mathbf{J}_2 is NFB. It suffices to verify that, for any $n \in \mathbb{N}$, the set of identities

$$\Delta_n = \{(3.1), \eta_1, \mathbf{z}_k[\pi, \tau] \approx \mathbf{z}'_k[\pi, \tau] \mid k < n, \pi, \tau \in S_{2k}\}$$

does not imply the identity $\mathbf{z}_n[\varepsilon, \varepsilon] \approx \mathbf{z}'_n[\varepsilon, \varepsilon]$, where ε is the trivial permutation on $\{1, 2, \dots, 2n\}$. Let \mathbf{w} be a word such that \mathbf{J}_2 satisfies $\mathbf{z}_n[\varepsilon, \varepsilon] \approx \mathbf{w}$. According to the inclusion $\mathbf{LRB} \subset \mathbf{J}_2$ and Lemmas 2.10 and 5.5,

$$\mathbf{w} = \left(\prod_{i=1}^{2n} x_i t_i \right) x \left(\prod_{i=1}^n z_i s_i \right) y \left(\prod_{i=1}^n z_{n+i} \right) \mathbf{w}' t \mathbf{w}''$$

for some words \mathbf{w}' and \mathbf{w}'' such that $\text{con}(\mathbf{w}') = \{x, y, x_1, z_1, x_2, z_2, \dots, x_{2n}, z_{2n}\}$ and $\text{con}(\mathbf{w}'') = \{s_1, s_2, \dots, s_n\}$. If ${}_{1\mathbf{w}'}x$ follows ${}_{1\mathbf{w}'}x_1$ in \mathbf{w}' , then the identity $\mathbf{w}_n[\varepsilon, \varepsilon](x, x_1, t_1) \approx \mathbf{w}(x, x_1, t_1)$ is equivalent modulo (3.1) to (4.9). But this is

impossible by Lemma 5.7. Therefore, ${}_{1\mathbf{w}'}x$ precedes ${}_{1\mathbf{w}'}x_1$ in \mathbf{w}' . By a similar argument we can show that $({}_{1\mathbf{w}'}y) < ({}_{1\mathbf{w}'}x_1)$ and

$$({}_{1\mathbf{w}'}x_1) < ({}_{1\mathbf{w}'}z_1) < ({}_{1\mathbf{w}'}x_2) < ({}_{1\mathbf{w}'}z_2) < \cdots < ({}_{1\mathbf{w}'}x_{2n}) < ({}_{1\mathbf{w}'}z_{2n}).$$

This means that $\text{ini}(\mathbf{w}') = \mathbf{a}x_1z_1x_2z_2 \cdots x_{2n}z_{2n}$, where $\mathbf{a} \in \{xy, yx\}$. So, we have proved that $\text{ini}_2(\mathbf{w}) = \mathbf{z}_n[\theta]$ for some $\theta \in S_n$, where

$$\mathbf{z}_n[\theta] = \left(\prod_{i=1}^{2n} x_i t_i \right) x \left(\prod_{i=1}^n z_i s_i \right) y \left(\prod_{i=1}^n z_{n+i} \right) \mathbf{a} \left(\prod_{i=1}^{2n} x_i z_i \right) t \left(\prod_{i=1}^n s_i \theta \right).$$

Let \mathbf{u} be a word such that \mathbf{J}_2 satisfies $\mathbf{z}_n[\varepsilon, \varepsilon] \approx \mathbf{u}$ and $({}_{2\mathbf{u}}x) < ({}_{2\mathbf{u}}y)$. In view of the previous paragraph, $\text{ini}_2(\mathbf{u}) = \mathbf{z}_n[\theta_1]$ for some $\theta_1 \in S_n$. According to Proposition 2.1, to verify that $\mathbf{z}_n[\varepsilon, \varepsilon] \approx \mathbf{z}'_n[\varepsilon, \varepsilon]$ does not follow from Δ_n , it suffices to show that if an identity $\mathbf{u} \approx \mathbf{v}$ is directly deducible from some identity $\mathbf{s} \approx \mathbf{t}$ in Δ_n , that is, $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\xi(\mathbf{s})\mathbf{b}, \mathbf{a}\xi(\mathbf{t})\mathbf{b}\}$ for some words $\mathbf{a}, \mathbf{b} \in \mathfrak{A}^*$ and some endomorphism ξ of \mathfrak{A}^* , then $({}_{2\mathbf{v}}x) < ({}_{2\mathbf{v}}y)$. Arguing by contradiction suppose that $({}_{2\mathbf{v}}y) < ({}_{2\mathbf{v}}x)$. Then, in view of the observation in the previous paragraph, $\text{ini}_2(\mathbf{v}) = \mathbf{z}_n[\theta_2]$ for some $\theta_2 \in S_n$.

Further, since $\text{ini}_2(\mathbf{u}(x, y, s_1, t)) = xs_1yxyts_1$ and $\text{ini}_2(\mathbf{v}(x, y, s_1, t)) = xs_1y^2xts_1$, Lemma 5.6 implies that $\mathbf{s} \approx \mathbf{t}$ cannot coincide with (3.1) and η_1 . Suppose that $\mathbf{s} \approx \mathbf{t}$ coincides with $\mathbf{z}_k[\pi, \tau] \approx \mathbf{z}'_k[\pi, \tau]$ for some $k < n$ and $\pi, \tau \in S_{2k}$. We consider only the case when $(\mathbf{s}, \mathbf{t}) = (\mathbf{z}_k[\pi, \tau], \mathbf{z}'_k[\pi, \tau])$ because the other case is similar. Since $\mathbf{z}_k[\pi, \tau]$ differs from $\mathbf{z}'_k[\pi, \tau]$ only in swapping of the second occurrences of letters x and y , there are three possibilities:

- (a) $\xi({}_{2\mathbf{s}}x)$ contains ${}_{2\mathbf{u}}x$ and $\xi({}_{2\mathbf{s}}y)$ contains ${}_{2\mathbf{u}}y$;
- (b) ${}_{2\mathbf{u}}x{}_{2\mathbf{u}}y$ is a subword of $\xi({}_{2\mathbf{s}}x)$ and $\xi({}_{2\mathbf{s}}y)$ contains some occurrence of y ;
- (c) $\xi({}_{2\mathbf{s}}x)$ contains ${}_{1\mathbf{u}}x$ and ${}_{2\mathbf{u}}x{}_{2\mathbf{u}}y$ is a subword of $\xi({}_{2\mathbf{s}}y)$.

Suppose that (a) holds. Then $\xi({}_{1\mathbf{s}}x)$ contains ${}_{1\mathbf{u}}x$ and $\xi({}_{1\mathbf{s}}y)$ contains ${}_{1\mathbf{u}}y$. Hence $t_{2n}, z_1 \notin \text{con}(\xi(x))$ and $s_n, z_{n+1} \notin \text{con}(\xi(y))$ because the equality $\text{ini}_2(\mathbf{u}) = \mathbf{z}_n[\theta_1]$ is false otherwise. Therefore, $\xi(x) = x$ and $\xi(y) = y$, whence $\xi(\prod_{i=1}^k z_{k+i}) = \prod_{i=1}^n z_{n+i}$. Then $\xi({}_{1\mathbf{s}}z_p)$ contains ${}_{1\mathbf{u}}z_q {}_{1\mathbf{u}}z_{q+1}$ for some $k+1 \leq p \leq 2k$ and $n+1 \leq q < 2n$. Since $\text{ini}_2(\mathbf{u}) = \mathbf{z}_n[\theta_1]$, the word $\xi({}_{2\mathbf{s}}z_p)$ cannot contain ${}_{2\mathbf{u}}z_q$. Then ${}_{2\mathbf{u}}z_q$ is contained in the subword $\xi(\prod_{i=1}^{r-1} (x_i \pi z_i \tau) x_{r\pi})$ of \mathbf{u} , where $r\tau = p$. Hence ${}_{1\mathbf{u}}z_q$ must be contained in $\xi({}_{1\mathbf{s}}a)$ for some

$$a \in \{z_{1\tau}, x_{1\pi}, z_{2\tau}, x_{2\pi}, \dots, z_{(r-1)\tau}, x_{(r-1)\pi}, x_{r\pi}\}.$$

This contradicts the fact that ${}_{1\mathbf{u}}z_q$ is contained in $\xi({}_{1\mathbf{s}}z_p)$. Therefore, (a) is impossible. Suppose that (b) holds. Then $\xi({}_{1\mathbf{s}}x)$ contains both ${}_{1\mathbf{u}}x$ and ${}_{1\mathbf{u}}y$. This is only possible when $xz_1s_1z_2s_2 \cdots z_y$ is a subword of $\xi(x)$. But this contradicts the fact that $\text{ini}_2(\mathbf{u}) = \mathbf{z}_n[\theta_1]$ and, therefore, (b) is impossible as well. Finally, (c) is impossible because $\xi({}_{2\mathbf{s}}x)$ cannot contain any first occurrence of a letter in \mathbf{u} . We see that $\mathbf{s} \approx \mathbf{t}$ cannot coincide with $\mathbf{z}_k[\pi, \tau] \approx \mathbf{z}'_k[\pi, \tau]$. So, we have proved that Δ_n does not imply $\mathbf{z}_n[\varepsilon, \varepsilon] \approx \mathbf{z}'_n[\varepsilon, \varepsilon]$ and so \mathbf{J}_2 is NFB. \square

The following fact provides the first example of a limit variety of monoids with infinitely many subvarieties.

Corollary 5.10. *The limit variety \mathbf{J}_2 have countably infinitely many subvarieties.*

Proof. Since \mathbf{J}_2 is a limit variety by Proposition 5.9, it contains at most countably infinitely many subvarieties. It remains to note that \mathbf{J}_2 has infinitely many subvarieties because $\mathbf{F}_1 \subset \mathbf{F}_2 \subset \cdots \subset \mathbf{F}_k \subset \cdots \subset \mathbf{J}_2$ by Lemma 2.12. \square

5.3. Classification of limit subvarieties of \mathbf{P} . The following is the main result of the paper.

Theorem 5.11. *The following statements on any subvariety \mathbf{V} of \mathbf{P} are equivalent:*

- (i) \mathbf{V} is HFB;
- (ii) $\mathbf{V} \subseteq \mathbf{O}_1$ or $\mathbf{V} \subseteq \mathbf{O}_2$;
- (iii) $\mathbf{J}_1, \mathbf{J}_2 \not\subseteq \mathbf{V}$.

Consequently, \mathbf{J}_1 and \mathbf{J}_2 are the only limit subvarieties of \mathbf{P} .

Proof. The implication (ii) \rightarrow (i) holds by Corollary 4.18, while the implication (i) \rightarrow (iii) holds because the varieties \mathbf{J}_1 and \mathbf{J}_2 are NFB by Propositions 5.4 and 5.9, respectively. By Lemmas 5.3 and 5.8, the statement $\mathbf{J}_1, \mathbf{J}_2 \not\subseteq \mathbf{V}$ implies that \mathbf{V} satisfies either $\{(4.15), (4.9)\}$ or $\{(4.15), \nu_1\}$, whence either $\mathbf{V} \subseteq \mathbf{O}_1$ or $\mathbf{V} \subseteq \mathbf{O}_2$. Therefore the implication (iii) \rightarrow (ii) holds. \square

6. THE MONOID P_2^1

6.1. The variety \mathbf{P}_2^1 is HFB.

Proposition 6.1. *The monoid variety \mathbf{P}_2^1 is HFB.*

Proof. Since the identities (3.1) and (5.1) constitute an identity basis for the variety \mathbf{P}_2^1 [22, Corollary 6.6], the inclusion $\mathbf{P}_2^1 \subseteq \mathbf{O}_1$ is easily deduced. Then \mathbf{P}_2^1 is HFB because \mathbf{O}_1 is HFB by Corollary 4.18. \square

Lemma 6.2 (Lee and Zhang [24]). *Up to isomorphism and anti-isomorphism, every monoid of order five or less, with the possible exception of P_2^1 , generates a small HFB variety.*

Corollary 6.3. *The variety generated by any monoid of order five is HFB and so contains at most countably many subvarieties.* \square

In contrast, as we have mentioned in the introduction, there exist monoids of order six that generate varieties with continuum many subvarieties [12, 14].

Remark 6.4. Up to isomorphism and anti-isomorphism, every semigroup of order five or less that is different from P_2^1 generates a HFB variety of semigroups [21]. As observed in Remark 1.1, whether or not P_2^1 generates a HFB variety of semigroups is presently an open question.

6.2. Subvarieties of \mathbf{P}_2^1 . In this subsection, we describe the subvariety lattice of \mathbf{P}_2^1 . Recall from Subsection 4.1 that

$$\Phi = \{(3.1), (3.2), \gamma_k, \delta_k^m, \varepsilon_{k-1} \mid k \in \mathbb{N}, 1 \leq m \leq k\}.$$

Proposition 6.5. *Each variety \mathbf{V} from the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1]$ can be defined within \mathbf{P}_2^1 by identities from Φ .*

Proof. The inclusion $\mathbf{F}_1 \subset \mathbf{V}$ and Lemma 2.12 imply that any 2-limited identity of \mathbf{V} is balanced. In view of this fact and Lemma 3.1, \mathbf{V} can be defined within \mathbf{P}_2^1 by a set of 2-limited balanced identities.

Let $\mathbf{u} \approx \mathbf{v}$ be a non-trivial 2-limited balanced identity of \mathbf{V} . Then there exists a critical pair $\{i x, j y\}$ in $\mathbf{u} \approx \mathbf{v}$. Let \mathbf{w} denote the word obtained from \mathbf{u} by replacing $i \mathbf{u} x j \mathbf{u} y$ with $j \mathbf{u} y i \mathbf{u} x$. It follows from the inclusion $\mathbf{LRB} \subset \mathbf{V}$ and Lemma 2.10 that $(i, j) \neq (1, 1)$. Then $(i, j) \in \{(1, 2), (2, 1), (2, 2)\}$ because the identity $\mathbf{u} \approx \mathbf{v}$ is 2-limited. If $(i, j) = (2, 2)$, then the critical pair $\{i x, j y\}$ is $\{(5.1)\}$ -removable. Now let $\{i, j\} = \{1, 2\}$. Then the critical pair $\{i x, j y\}$ is $\Phi(\mathbf{V})$ -removable by Proposition 4.7. We see that $\Phi(\mathbf{V}) \cup \{(5.1)\}$ implies $\mathbf{u} \approx \mathbf{w}$ in either case. According to Lemma 4.1, $\mathbf{u} \approx \mathbf{v}$ is a consequence of $\Phi(\mathbf{V}) \cup \{(5.1)\}$ and, therefore, \mathbf{V} is defined within \mathbf{P}_2^1 by $\Phi(\mathbf{V})$. \square

Lemma 6.6. *For any $1 \leq m \leq k$, we have $\mathbf{P}\{\delta_k^m, \varepsilon_0\} \subseteq \mathbf{P}\{\gamma_{k+1}\}$.*

Proof. By Lemmas 4.2 and 4.3(ii), the identity κ_k is satisfied by $\mathbf{P}\{\delta_k^m, \varepsilon_0\}$. Then the identity γ_{k+1} holds in $\mathbf{P}\{\delta_k^m, \varepsilon_0\}$ because

$$\begin{aligned}
& y_1 y_0 y_1 x_{k+1} x_k x_{k+1} \mathbf{b}_k \stackrel{\kappa_k}{\approx} y_1 y_0 y_1 x_{k+1}^2 x_k \mathbf{b}_k \stackrel{\varepsilon_0}{\approx} y_1 y_0 x_{k+1} y_1 x_{k+1} x_k \mathbf{b}_k \\
(3.1) \quad & \approx y_1 y_0 x_{k+1} y_1 x_{k+1} x_k \mathbf{b}_{k,m} y_1 \mathbf{b}_{m-1} \stackrel{\delta_k^m}{\approx} y_1 y_0 x_{k+1} y_1 x_k x_{k+1} \mathbf{b}_{k,m} y_1 \mathbf{b}_{m-1} \\
(1.1) \quad & \approx y_1 y_0 x_{k+1} y_1 x_k x_{k+1} \mathbf{b}_k.
\end{aligned}$$

The lemma is thus proved. \square

Lemma 6.7. *For any $1 \leq m \leq \ell \leq k$, we have $\mathbf{P}\{\delta_k^\ell, \varepsilon_m\} \subseteq \mathbf{P}\{\delta_{k+1}^m\}$.*

Proof. For convenience, let $\mathbf{b}_{p-1,p} = \lambda$ for any $p \in \mathbb{N}$. We notice that the identity κ_k is satisfied by $\mathbf{P}\{\delta_k^k, \varepsilon_m\}$ by Lemmas 4.2 and 4.3(ii). Then the identity δ_{k+1}^m holds in $\mathbf{P}\{\delta_k^k, \varepsilon_m\}$ because

$$\begin{aligned}
& y_{m+1} y_m y_{m+1} x_{k+1} x_k x_{k+1} \mathbf{b}_{k,m} y_m \mathbf{b}_{m-1} \\
(3.1) \quad & \approx y_{m+1} y_m y_{m+1} x_{k+1} x_k x_{k+1} \mathbf{b}_{k,m} \mathbf{d}_1 \\
& \stackrel{\kappa_k}{\approx} y_{m+1} y_m y_{m+1} x_{k+1}^2 x_k \mathbf{b}_{k,m} \mathbf{d}_1 \\
(3.1) \quad & \approx y_{m+1} y_m y_{m+1} x_{k+1}^2 (x_k \mathbf{b}_{k,m}) \mathbf{d}_2 \\
& \stackrel{\varepsilon_m}{\approx} y_{m+1} y_m x_{k+1} y_{m+1} x_{k+1} (x_k \mathbf{b}_{k,m}) \mathbf{d}_2 \\
(3.1) \quad & \approx y_{m+1} y_m x_{k+1} y_{m+1} x_{k+1} x_k \mathbf{b}_{k,k} y_{m+1} \mathbf{b}_{k-1,m} \mathbf{d}_1 \\
& \stackrel{\delta_k^k}{\approx} y_{m+1} y_m x_{k+1} y_{m+1} x_k x_{k+1} \mathbf{b}_{k,k} y_{m+1} \mathbf{b}_{k-1,m} \mathbf{d}_1 \\
(3.1) \quad & \approx y_{m+1} y_m x_{k+1} y_{m+1} x_k x_{k+1} \mathbf{b}_{k,m} y_m \mathbf{b}_{m-1},
\end{aligned}$$

where

$$\mathbf{d}_1 = \begin{cases} y_m & \text{if } m = 1, \\ (y_m x_{m-2}) x_{m-1} & \text{if } m = 2, \\ (y_m x_{m-2}) x_{m-1} x_{m-3} (y_m x_{m-2}) \mathbf{b}_{m-3} & \text{if } m \geq 3, \end{cases}$$

$$\mathbf{d}_2 = \begin{cases} y_m & \text{if } m = 1, \\ x_{m-2} (x_k \mathbf{b}_{k,m}) \mathbf{b}_{m-2} & \text{if } m \geq 2. \end{cases}$$

To complete the proof, it remains to notice that $\mathbf{P}\{\delta_k^\ell, \varepsilon_m\} \subseteq \mathbf{P}\{\delta_k^k, \varepsilon_m\}$ by Lemma 4.3(ii). \square




FIGURE 1. The lattice $\mathcal{L}(P_2^1)$

Proposition 6.8. *The lattice $\mathfrak{L}(\mathbf{P}_2^1)$ is as shown in Fig. 1.¹*

Proof. Lemma 3.2(ii) implies that the lattice $\mathfrak{L}(\mathbf{P}_2^1)$ is the disjoint union of the lattice $\mathfrak{L}(\mathbf{LRB} \vee \mathbf{C})$ and the interval $[\mathbf{F}_1, \mathbf{P}_2^1]$. As verified in Lee [19, Fig. 2], the lattice $\mathfrak{L}(\mathbf{LRB} \vee \mathbf{C})$ is as shown in Fig. 1. In view of Lemma 3.2(i), the interval $[\mathbf{F}_1, \mathbf{P}_2^1]$ is a disjoint union of the intervals $[\mathbf{F}_1, \mathbf{F}]$ and $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1]$. According to Lemma 2.11, the interval $[\mathbf{F}_1, \mathbf{F}]$ has the form shown in Fig. 1. Since any identity from $\Phi \setminus \{(3.1)\}$ together with (3.1) imply (3.2) by Lemma 4.3, the variety $\mathbf{P}_2^1\{(3.2)\}$ is the greatest proper subvariety of \mathbf{P}_2^1 . (This result is also deducible from Lee and Li [22, Lemma 6.7].) Let \mathbf{V} be a variety from $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1\{(3.2)\}]$.

Suppose that $\mathbf{P}_2^1\{\delta_1^1\} \not\subseteq \mathbf{V}$. Since $\mathbf{P}_2^1\{\delta_1^1\}$ satisfies the identities (3.2), δ_k^m and ε_k for any $k \in \mathbb{N}$ and $1 \leq m \leq k$ by Parts (ii) and (iii) of Lemma 4.3, Proposition 6.5 implies that \mathbf{V} satisfies ε_0 or γ_ℓ for some $\ell \in \mathbb{N}$. Now we apply Lemma 4.3(i) and obtain that $\mathbf{P}_2^1\{\gamma_\ell\} \subseteq \mathbf{P}_2^1\{\varepsilon_0\}$ for any $\ell \in \mathbb{N}$. Hence \mathbf{V} satisfies ε_0 . Thus, \mathbf{V} belongs to the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1\{\varepsilon_0\}]$.

Suppose now that $\mathbf{P}_2^1\{\delta_k^k\} \subseteq \mathbf{V}$ for all $k \in \mathbb{N}$. Then, since $\mathbf{P}_2^1\{\delta_k^k\}$ violates ε_{k-1} , Lemma 4.3 and Proposition 6.5 imply that $\mathbf{V} = \mathbf{P}_2^1\{(3.2)\}$.

Finally, suppose that $\mathbf{P}_2^1\{\delta_k^k\} \subseteq \mathbf{V}$ but $\mathbf{P}_2^1\{\delta_{k+1}^{k+1}\} \not\subseteq \mathbf{V}$ for some $k \in \mathbb{N}$. Notice that

- (a) $\mathbf{J}_s^1 \not\subseteq \mathbf{P}_2^1\{\gamma_s\}$ but $\mathbf{J}_s^1 \subseteq \mathbf{P}_2^1\{\delta_s^1\}$ for any $s \in \mathbb{N}$;
- (b) $\mathbf{I}_{s+1} \not\subseteq \mathbf{P}_2^1\{\delta_s^s\}$ but $\mathbf{I}_{s+1} \subseteq \mathbf{P}_2^1\{\gamma_{s+1}\}$ for any $s \in \mathbb{N}$;
- (c) $\mathbf{J}_s^{t+1} \not\subseteq \mathbf{P}_2^1\{\delta_s^t\}$ but $\mathbf{J}_s^{t+1} \subseteq \mathbf{P}_2^1\{\delta_s^{t+1}\}$ for any $1 \leq t < s$;
- (d) $\mathbf{J}_{s+1}^t \not\subseteq \mathbf{P}_2^1\{\delta_s^t\}$ but $\mathbf{J}_{s+1}^t \subseteq \mathbf{P}_2^1\{\delta_{s+1}^t\}$ for any $1 \leq t \leq s$.

These facts together with Lemma 4.3(ii),(iii) imply that $\mathbf{P}_2^1\{\delta_k^k\}$ violates the identities γ_p and δ_p^q , where $p \in \mathbb{N}$ and $1 \leq q < k$. Now we apply Lemma 4.3(ii),(iii) again and conclude that $\mathbf{P}_2^1\{\delta_{k+1}^{k+1}\}$ satisfies δ_r^m and ε_r for all r and m such that $k+1 \leq m \leq r$. Then, since $\mathbf{P}_2^1\{\delta_k^k\} \subseteq \mathbf{V}$ and $\mathbf{P}_2^1\{\delta_{k+1}^{k+1}\} \not\subseteq \mathbf{V}$, Proposition 6.5 implies that \mathbf{V} satisfies one of the identities ε_k or δ_p^k for some $p \geq k$. Now we apply Lemma 4.3(ii) again and obtain that $\mathbf{P}_2^1\{\delta_p^k\} \subseteq \mathbf{P}_2^1\{\varepsilon_k\}$. Hence \mathbf{V} satisfies ε_k . Thus, \mathbf{V} belongs to the interval $[\mathbf{P}_2^1\{\delta_k^k\}, \mathbf{P}_2^1\{\varepsilon_k\}]$.

We see that the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1\{(3.2)\}]$ is a disjoint union of the intervals $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1\{\varepsilon_0\}]$, $\{\mathbf{P}_2^1\{(3.2)\}\}$ and $[\mathbf{P}_2^1\{\delta_k^k\}, \mathbf{P}_2^1\{\varepsilon_k\}]$ for all $k \in \mathbb{N}$. So, it remains to verify that these intervals are as shown in Fig. 1.

First, we will show that the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1\{\varepsilon_0\}]$ has the form shown in Fig. 1. In view of Lemma 4.3, $\mathbf{P}_2^1\{\varepsilon_0\}$ satisfies $\{(3.2), \varepsilon_r \mid r \in \mathbb{N}\}$. Then Proposition 6.5 implies that every variety from the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1\{\varepsilon_0\}]$ can be defined within $\mathbf{P}_2^1\{\varepsilon_0\}$ by some possibly empty subset of $\{\gamma_p, \delta_p^q \mid 1 \leq q \leq p\}$; in particular $\mathbf{LRB} \vee \mathbf{F}_1 = \mathbf{P}_2^1\{\gamma_1\}$. It follows from Lemma 6.6 that $\mathbf{P}_2^1\{\delta_p^q, \varepsilon_0\} \subseteq \mathbf{P}_2^1\{\gamma_{p+1}\}$ for any $1 \leq q \leq p$. Besides that, Lemma 4.3(ii) implies that $\mathbf{P}_2^1\{\gamma_p\} \subseteq \mathbf{P}_2^1\{\delta_p^q, \varepsilon_0\}$. Therefore, the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1\{\varepsilon_0\}]$ is a union of the singleton interval $\{\mathbf{P}_2^1\{\varepsilon_0\}\}$ and the intervals of the form $[\mathbf{P}_2^1\{\gamma_p\}, \mathbf{P}_2^1\{\gamma_{p+1}\}]$, where $p \in \mathbb{N}$. In view of Lemma 4.3, $\mathbf{P}_2^1\{\gamma_{p+1}\}$ satisfies γ_s , δ_s^t and ε_r for all s , t and r such that $p+1 \leq s$, $1 \leq t \leq s$ and $r \geq 0$. It follows from (a)–(d) that $\mathbf{P}_2^1\{\gamma_p\}$ violates γ_s and δ_s^t for all s and t such that $1 \leq t \leq s < p$. In view of these facts and Proposition 6.5, every variety from the interval $[\mathbf{P}_2^1\{\gamma_p\}, \mathbf{P}_2^1\{\gamma_{p+1}\}]$ can

¹For convenience, some varieties in Fig. 1 are marked by the identities, which define them within the variety \mathbf{P}_2^1 . For instance, the identity γ_1 marks the variety $\mathbf{P}_2^1\{\gamma_1\}$.

be defined within $\mathbf{P}_2^1\{\gamma_{p+1}\}$ by some possibly empty subset of $\{\gamma_p, \delta_p^1, \delta_p^2, \dots, \delta_p^p\}$. Now we apply Lemma 4.3 again and obtain that the interval $[\mathbf{P}_2^1\{\gamma_p\}, \mathbf{P}_2^1\{\gamma_{p+1}\}]$ forms the chain

$$\mathbf{P}_2^1\{\gamma_p\} \subseteq \mathbf{P}_2^1\{\delta_p^1, \gamma_{p+1}\} \subseteq \mathbf{P}_2^1\{\delta_p^2, \gamma_{p+1}\} \subseteq \dots \subseteq \mathbf{P}_2^1\{\delta_p^p, \gamma_{p+1}\} \subseteq \mathbf{P}_2^1\{\gamma_{p+1}\}.$$

It remains to notice that all these inclusions are strict by (a)–(d). We see that the interval $[\mathbf{P}_2^1\{\gamma_p\}, \mathbf{P}_2^1\{\gamma_{p+1}\}]$ and, therefore, the interval $[\mathbf{LRB} \vee \mathbf{F}_1, \mathbf{P}_2^1\{\varepsilon_0\}]$ are as shown in Fig. 1.

It remains to show that, for any $k \in \mathbb{N}$, the interval $[\mathbf{P}_2^1\{\delta_k^k\}, \mathbf{P}_2^1\{\varepsilon_k\}]$ has the form shown in Fig. 1. The proof of this fact is very similar to the arguments from the previous paragraph but we provide it for the sake of completeness. It follows from (a)–(d) that $\mathbf{P}_2^1\{\delta_k^k\}$ violates γ_r, δ_s^t and ε_t for all r, s and t such that $r \geq 0, s \geq 1, t \leq s$ and $1 \leq t < k$. In view of Lemma 4.3, $\mathbf{P}_2^1\{\varepsilon_k\}$ satisfies $\{(3.2), \varepsilon_r \mid r \geq k\}$. Then Proposition 6.5 implies that every variety from the interval $[\mathbf{P}_2^1\{\delta_k^k\}, \mathbf{P}_2^1\{\varepsilon_k\}]$ can be defined within $\mathbf{P}_2^1\{\varepsilon_k\}$ by some possibly empty subset of $\{\delta_p^q \mid k \leq q \leq p\}$. It follows from Lemma 6.7 that $\mathbf{P}_2^1\{\delta_p^q, \varepsilon_k\} \subseteq \mathbf{P}_2^1\{\delta_{p+1}^k\}$ for any $k \leq q \leq p$. Besides that, Lemma 4.3 implies that $\mathbf{P}_2^1\{\delta_p^k\} \subseteq \mathbf{P}_2^1\{\delta_p^q, \varepsilon_k\}$. Therefore, the interval $[\mathbf{P}_2^1\{\delta_k^k\}, \mathbf{P}_2^1\{\varepsilon_k\}]$ is the union of the singleton interval $\{\mathbf{P}_2^1\{\varepsilon_k\}\}$ and the intervals of the form $[\mathbf{P}_2^1\{\delta_p^k\}, \mathbf{P}_2^1\{\delta_{p+1}^k\}]$, where $p \geq k$. Finally, Lemma 4.3, Proposition 6.5 and (a)–(d) imply that every variety from the interval $[\mathbf{P}_2^1\{\delta_p^k\}, \mathbf{P}_2^1\{\delta_{p+1}^k\}]$ can be defined within $\mathbf{P}_2^1\{\delta_{p+1}^k\}$ by some possibly empty subset of $\{\delta_p^k, \delta_p^{k+1}, \dots, \delta_p^p\}$. Now we apply Lemma 4.3 again and obtain that the interval $[\mathbf{P}_2^1\{\delta_p^k\}, \mathbf{P}_2^1\{\delta_{p+1}^k\}]$ forms the chain

$$\mathbf{P}_2^1\{\delta_p^k\} \subseteq \mathbf{P}_2^1\{\delta_p^{k+1}, \delta_{p+1}^k\} \subseteq \mathbf{P}_2^1\{\delta_p^{k+2}, \delta_{p+1}^k\} \subseteq \dots \subseteq \mathbf{P}_2^1\{\delta_p^p, \delta_{p+1}^k\} \subseteq \mathbf{P}_2^1\{\delta_{p+1}^k\}.$$

It remains to notice that all these inclusions are strict by (a)–(d). We see that the interval $[\mathbf{P}_2^1\{\delta_p^k\}, \mathbf{P}_2^1\{\delta_{p+1}^k\}]$ and so the interval $[\mathbf{P}_2^1\{\delta_k^k\}, \mathbf{P}_2^1\{\varepsilon_k\}]$ are as shown in Fig. 1. \square

6.3. Finitely universal varieties. Following Shevrin et al. [34], a variety of algebras is *finitely universal* if every finite lattice is order-embeddable into its subvariety lattice. The first example of a finitely universal variety of semigroups was presented by Burris and Nelson [2] half a century ago. In contrast, the first example of a finitely universal variety of monoids was just recently found, in particular, there exist finitely universal varieties of monoids that are finitely generated [8, Section 5]. In view of this result, it is natural to find the least order of a monoid that generates a finitely universal variety. The following statement shows that this order is more than five.

Proposition 6.9. *Every monoid of order five or less generates a variety that is not finitely universal.*

Proof. In view of Lemma 6.2, it suffices to show that \mathbf{P}_2^1 is not finitely universal. It is clear from Fig. 1 that every proper subvariety of \mathbf{P}_2^1 has at most two coverings, so that \mathbf{P}_2^1 is indeed not finitely universal. For instance, the modular non-distributive lattice in Fig. 2 is not embeddable in the lattice $\mathfrak{L}(\mathbf{P}_2^1)$. \square

Remark 6.10. In contrast, up to isomorphism and anti-isomorphism, there exist precisely four distinct semigroups of order four that generate a finitely universal




FIGURE 2. A modular non-distributive lattice

variety, while the variety generated by any other semigroup of order four or less is not finitely universal [17].

APPENDIX. TABLES OF IDENTITIES

TABLE 1. Identities labeled by Greek letters

Identity	Page number
α_k	6
β_k	6
γ_k	6
δ_k^m	6
ε_k	11
ζ_k	19
η_k	19
κ_k	8
λ_k^m	19
μ_k^m	19
ν_k	19

TABLE 2. “Non-local” identities labeled by numbers

Identity	Page number
(3.1)	7
(3.2)	9
(4.9)	23
(4.12)	26
(4.15)	27
(5.1)	31

ACKNOWLEDGEMENTS

The authors thank Edmond W. H. Lee for careful reading the previous version and his valuable comments and suggestions that contributed to a significant improvement of the manuscript. The authors are also grateful the anonymous referee for the suggestion to add tables of identities.

REFERENCES

- [1] S. I. Adyan, *Infinite irreducible systems of group identities*, Dokl. Akad. Nauk SSSR **190** (1970), 499–501 [Russian; Engl. translation: Soviet Math. Dokl. **11** (1970), 113–115].
- [2] S. Burris and E. Nelson, *Embedding the dual of Π_m in the lattice of equational classes of commutative semigroups*, Proc. Amer. Math. Soc., **30** (1971), 37–39.
- [3] S. Burris and H. P. Sankappanavar, *A Course in Universal Algebra*, Springer Verlag, New York, 1981.
- [4] C. C. Edmunds, E. W. H. Lee and K. W. K. Lee, *Small semigroups generating varieties with continuum many subvarieties*, Order, **27** (2010), 83–100.
- [5] C. K. Gupta and A. Krasilnikov, *The finite basis question for varieties of groups—some recent results*, Illinois J. Math., **47** (2003), 273–283.
- [6] S. V. Gusev, *A new example of a limit variety of monoids*, Semigroup Forum, **101** (2020), 102–120.
- [7] S. V. Gusev, *Limit varieties of aperiodic monoids with commuting idempotents*, J. Algebra Appl., **20**, No. 9 (2021), 2150160.
- [8] S. V. Gusev and E. W. H. Lee, *Varieties of monoids with complex lattices of subvarieties*, Bull. Lond. Math. Soc., **52** (2020), 762–775.
- [9] S. V. Gusev and O. B. Sapir, *Classification of limit varieties of J -trivial monoids*, Comm. Algebra, accepted; DOI: 10.1080/00927872.2021.2024561.
- [10] S. V. Gusev and B. M. Vernikov, *Chain varieties of monoids*, Dissert. Math., **534** (2018), 1–73.
- [11] M. Jackson, *Finiteness properties of varieties and the restriction to finite algebras*, Semigroup Forum, **70** (2005), 154–187.
- [12] M. Jackson and E. W. H. Lee, *Monoid varieties with extreme properties*, Trans. Amer. Math. Soc., **370** (2018), 4785–4812.
- [13] M. Jackson and O. B. Sapir, *Finitely based, finite sets of words*, Internat. J. Algebra Comput., **10** (2000), 683–708.
- [14] M. Jackson and W. T. Zhang, *From A to B to Z* , Semigroup Forum, **103** (2021), 165–190.
- [15] P. A. Kozhevnikov, *On nonfinitely based varieties of groups of large prime exponent*, Comm. Algebra, **40** (2012), 2628–2644.
- [16] R. L. Kruse, *Identities satisfied by a finite ring*, J. Algebra, **26** (1973), 298–318.
- [17] E. W. H. Lee, *Minimal semigroups generating varieties with complex subvariety lattices*, Int. J. Algebra Comput., **17** (2007), 1553–1572.
- [18] E. W. H. Lee, *Finitely generated limit varieties of aperiodic monoids with central idempotents*, J. Algebra Appl., **8** (2009), 779–796.
- [19] E. W. H. Lee, *Maximal Specht varieties of monoids*, Mosc. Math. J., **12** (2012), 787–802.
- [20] E. W. H. Lee, *Varieties generated by 2-testable monoids*, Studia Sci. Math. Hungar., **49** (2012), 366–389.
- [21] E. W. H. Lee, *Finite basis problem for semigroups of order five or less: generalization and revisitation*, Studia Logica, **101** (2013), 95–115.
- [22] E. W. H. Lee, and J. R. Li, *Minimal non-finitely based monoids*, Dissert. Math., **475** (2011), 1–65.
- [23] E. W. H. Lee, J. Rhodes and B. Steinberg, *Join irreducible semigroups*, Int. J. Algebra Comput., **29** (2019), 1249–1310.
- [24] E. W. H. Lee and W. T. Zhang, *The smallest monoid that generates a non-Cross variety*, Xiamen Daxue Xuebao Ziran Kexue Ban, **53** (2014), 1–4 [Chinese].
- [25] I. V. L’vov, *Varieties of associative rings. I*, Algebra i Logika, **12** (1973), 269–297 [Russian; Engl. translation: Algebra and Logic, **12** (1973), 150–167].
- [26] B. H. Neumann, *Identical relations in groups. I*, Math. Ann. **114** (1937), 506–525.
- [27] S. Oates and M. B. Powell, *Identical relations in finite groups*, J. Algebra, **1** (1964), 11–39.
- [28] A. Yu. Ol’shanskii, *The finite basis problem for identities in groups*, Izv. Akad. Nauk SSSR Ser. Mat. **34** (1970), 376–384 [Russian; Engl. translation: Math. USSR Izv. **4** (1970), 381–389].
- [29] P. Perkins, *Bases for equational theories of semigroups*, J. Algebra **11** (1969), 298–314.
- [30] O. B. Sapir, *Finitely based words*, Internat. J. Algebra Comput., **10** (2000), 457–480.
- [31] O. B. Sapir, *Finitely based monoids*, Semigroup Forum, **90** (2015), 587–614.
- [32] O. B. Sapir, *Limit varieties of J -trivial monoids*, Semigroup Forum, **103** (2021), 236–260.

- [33] O. B. Sapir, *Limit varieties generated by finite non-J-trivial aperiodic monoids*, manuscript; available at: <https://arxiv.org/abs/2012.13598v1>
- [34] L. N. Shevrin, B. M. Vernikov and M. V. Volkov, *Lattices of semigroup varieties*, Izv. VUZ. Matematika, No. 3 (2009), 3–36 [Russian; Engl. translation: Russ. Math. Iz. VUZ, **53**, No. 3 (2009), 1–28].
- [35] M. R. Vaughan-Lee, *Uncountably many varieties of groups*, Bull. London Math. Soc. **2** (1970), 280–286.
- [36] M. V. Volkov, *The finite basis problem for finite semigroups*, Sci. Math. Jpn., **53** (2001), 171–199.
- [37] W. T. Zhang, *Existence of a new limit variety of aperiodic monoids*, Semigroup Forum, **86** (2013), 212–220.
- [38] W. T. Zhang and Y. F. Luo, *A new example of limit variety of aperiodic monoids*, manuscript; available at: <https://arxiv.org/abs/1901.02207>

URAL FEDERAL UNIVERSITY, INSTITUTE OF NATURAL SCIENCES AND MATHEMATICS, LENINA 51, EKATERINBURG 620000, RUSSIA
Email address: sergey.gusb@gmail.com

LANZHOU UNIVERSITY, SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU 730000, PEOPLE'S REPUBLIC OF CHINA
Email address: liyuxian21@lzu.edu.cn

LANZHOU UNIVERSITY, SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU 730000, PEOPLE'S REPUBLIC OF CHINA
Email address: zhangwt@lzu.edu.cn