

AN UNDECIDABLE EXTENSION OF MORLEY'S THEOREM ON THE NUMBER OF COUNTABLE MODELS

CHRISTOPHER J. EAGLE¹, CLOVIS HAMEL², SANDRA MÜLLER³,
AND FRANKLIN D. TALL⁴

ABSTRACT. We show that Morley's theorem on the number of countable models of a countable first-order theory becomes an undecidable statement when extended to second-order logic. More generally, we calculate the number of equivalence classes of σ -projective equivalence relations in several models of set theory. Our methods include random and Cohen forcing, Woodin cardinals and Inner Model Theory.

1. INTRODUCTION

Vaught's Conjecture, which asserts that a countable first-order theory must have either at most countably many or exactly 2^{\aleph_0} many non-isomorphic countable models, is one of the most important problems in Model Theory. While the question itself is model-theoretic, it is known to have deep connections to both Descriptive Set Theory and Topological Dynamics. Since Vaught's original paper [46], Vaught's Conjecture has been verified for a number of classes of theories, such as theories of trees [39], ω -stable theories [37], \aleph_1 -minimal theories [28], and varieties (in the sense of universal algebra) [14], among others.

A strong positive result about Vaught's Conjecture that applies to all first-order theories is a result of the late Michael Morley [29], which states that the number of isomorphism classes of countable models of a countable first-order theory is always either at most \aleph_1 or exactly 2^{\aleph_0} . In this form Morley's Theorem requires no proof at all if we are in a universe of set theory where the continuum hypothesis holds. However, not long after Morley's paper appeared it was noticed that this result can be improved. Under a standard identification between countable models and elements of 2^ω (described in Section 2 below), we have the following strengthening of Morley's result:

Date: February 9, 2022.

2020 Mathematics Subject Classification. Primary 03C85, 03C55, 03E35, 03C52; Secondary 03E45, 03E55, 03E60, 03E15, 03C80.

Key words and phrases. Morley's theorem, countable models, random and Cohen forcing, σ -projective equivalence relations, Woodin cardinals, inner model theory.

¹ Supported by an NSERC Discovery Grant.

² Supported by an NSERC Vanier Scholarship.

³ Supported by L'ORÉAL Austria, in collaboration with the Austrian UNESCO Commission and in cooperation with the Austrian Academy of Sciences - Fellowship *Determinacy and Large Cardinals* and Elise Richter grant number V844 of the Austrian Science Fund (FWF).

⁴ Supported by NSERC grant A-7354.

Theorem (Absolute Morley Theorem). *Let T be a first-order theory (or, more generally, a sentence of $L_{\omega_1, \omega}$) in a countable signature. Then either T has at most \aleph_1 isomorphism classes of countable models, or there is a perfect set of non-isomorphic countable models of T .*

The previous theorem does not appear in Morley’s paper, but many of the ideas are there for its proof. An elegant proof relies on an important Descriptive Set Theory result by Burgess [5]:

Theorem (Burgess). *Let E be a Σ_1^1 equivalence relation on \mathbb{R} . If there is no perfect set of pairwise inequivalent reals, then there are at most \aleph_1 equivalence classes.*

The intersection of Descriptive Set Theory and Model Theory has become a widely studied topic, to such extent that there are versions of Vaught’s Conjecture that completely fall in the domain of Descriptive Set Theory, as they concern equivalence relations more general than isomorphism between countable models (see [10]). The main idea behind the connection between these two disciplines is that countable models of a second-order theory can be *coded* as reals, usually as elements of 2^ω . It is then easy to prove that the isomorphism relation (which can be formulated as the existence of a certain function) is a Σ_1^1 equivalence relation. Finally, the proof of the Absolute Morley Theorem follows from the fact proved by Morley [29] that the set of countable models of a theory in a countable fragment of $L_{\omega_1, \omega}$ is Borel. As is customary in Descriptive Set Theory, we make extensive use of the fact that sets of reals *definable* from a real can also be coded by a real.

In this paper we concern ourselves with logics that are stronger than first-order logic, especially second-order logic and fragments thereof. Vaught’s Conjecture is false for second-order logic if the continuum hypothesis fails; one easy counterexample is that one can express in second-order logic that a linear order is a well-order, and hence there is a second-order theory whose countable models are (up to isomorphism) exactly the countable ordinals. In this paper we investigate versions of the Absolute Morley Theorem for second-order logic. Our main results are the following:

Theorem A. *Force over L by first adding \aleph_2 Cohen reals and then adding \aleph_3 random reals. In the resulting universe of set theory, there is a second-order theory T in a countable signature such that the number of non-isomorphic countable models of T is exactly \aleph_2 , while $2^{\aleph_0} = \aleph_3$.*

Theorem B. *Beginning with a supercompact cardinal, carry out the standard forcing iteration for producing a model of the Proper Forcing Axiom. In the resulting universe of set theory, if T is a second-order theory in a countable signature, then either T has at most \aleph_1 isomorphism classes of countable models, or there is a perfect set of non-isomorphic models of T .*

These two results together show that, modulo the consistency of the existence of a supercompact cardinal, the extension of the absolute version of Morley’s Theorem from first-order logic to second-order logic is undecidable.

Our proof of Theorem B is based on results of Foreman and Magidor [9], and in fact produces a similar conclusion not just for isomorphism classes of countable

models of a second-order theory, but in fact for any equivalence relation in $L(\mathbb{R})$. This suggests the possibility of reducing the large cardinal strength below a supercompact cardinal for obtaining our desired result. Moreover, the full strength of second-order logic is significantly more than is needed for expressing most theories of interest in mathematics. We therefore also wish to consider the required large cardinal strength for fragments of second-order logic with a prescribed maximum quantifier complexity (such as existential second-order logic). We obtain the following:

Theorem C. *If there are infinitely many Woodin cardinals, then there is a model of set theory in which the Absolute Morley Theorem holds for second-order theories in countable signatures.*

Although we believe our use of large cardinals is necessary, we don't have a proof of this.

Problem 1. *Prove that large cardinals are necessary to prove the consistency of the conclusion of Theorem C.*

The remainder of the paper is structured as follows. In Section 2 we briefly review the syntax and semantics of second-order logic, and also the method by which countable structures can be encoded as elements of 2^ω . We also consider the descriptive complexity of the set of models of a second-order theory. Next, in Section 3 we prove Theorem A, and in Section 4 we sketch a proof of a weaker version of Theorem B. In Section 5 we develop finer results concerning the large cardinal strength necessary for the consistency of the Absolute Morley theorem for fragments of second-order logic. In Section 6 we investigate Absolute Morley for certain game and partially ordered quantifiers. In Section 7 we prove Theorems B and C.

Acknowledgement. We are grateful to Professor Magidor for explaining the proof of Theorem 4.1 to the fourth author.

2. PRELIMINARIES

2.1. Coding countable structures as reals. All of the logics we will consider make use of the same notion of *structures* as is used in first-order logic. For simplicity we will consider only relational structures. Structures with function and constant symbols can be easily incorporated into this framework by coding the functions and constants as relations. Thus, for us, a *signature* consists of a collection of relation symbols, each with a specified *arity*. If S is a signature, then the S -*structures* are defined exactly as in first-order logic.

To use tools from Descriptive Set Theory we need to code countable structures as elements of a Polish space. The method for doing so is standard (see for example [10]), but we review it here for the convenience of the reader.

Let $S = \{R_i\}_{i \in I}$ be a signature, where each R_i is a relation symbol of arity n_i , and I is countable. Suppose that \mathcal{M} is a countable S -structure. Up to isomorphism, we may assume that the underlying set of \mathcal{M} is ω . For each $i \in I$ the interpretation

$R_i^{\mathcal{M}}$ of R_i is a subset of ω^{n_i} , and so we identify R_i with an element of $2^{\omega^{n_i}}$. As the structure \mathcal{M} is completely determined by the interpretations of each of the relation symbols, we may identify \mathcal{M} with an element of $\prod_{i \in I} 2^{\omega^{n_i}}$. This identification provides a bijective map from the collection of S -structures with universe ω to $\prod_{i \in I} 2^{\omega^{n_i}}$. We thus view the Cantor set $\prod_{i \in I} 2^{\omega^{n_i}}$ as being the space of countable S -structures. We thus define $\text{Mod}_S = \prod_{i \in I} 2^{\omega^{n_i}}$.

If σ is a sentence of some logic, we define $\text{Mod}_S(\sigma) = \{\mathcal{M} \in \text{Mod}_S : \mathcal{M} \models \sigma\}$ (if S is clear from context we may omit it). In the case where σ is a sentence of $L_{\omega_1, \omega}$, the set $\text{Mod}_S(\sigma)$ is a Borel subset of Mod_S , and moreover every isomorphism-invariant Borel subset of Mod_S is of the form $\text{Mod}_S(\sigma)$ for some $L_{\omega_1, \omega}$ sentence σ [10, Theorem 11.3.6].

We are now prepared to define the main equivalence relations we study in this paper.

Definition 2.1. Let S be a countable signature, and let T be an S -theory of some logic. The equivalence relation of *isomorphism of models of T* is the equivalence relation \cong_T on Mod_S defined by declaring that $\mathcal{M} \cong_T \mathcal{N}$ if and only if either neither of the two structures is a model of T , or $\mathcal{M} \cong \mathcal{N}$.

The equivalence classes of \cong_T are thus one class for each isomorphism class of models of T , together with one additional class containing all elements of $\text{Mod}_S \setminus \text{Mod}_S(T)$. We are only interested in the classes corresponding to isomorphism types of models of T ; the following lemma (which we often use without explicit mention) allows us to move from a perfect set of \cong_T -inequivalent structures to a perfect set of non-isomorphic models of T . Recall that a set of reals A has the *perfect set property* if either A is countable or A includes a non-empty perfect set.

Lemma 2.2. *Let X be a Polish space, and let $A \subseteq X$ be a non-empty perfect set. For every $x \in X$, $A \setminus \{x\}$ includes a non-empty perfect set.*

Proof. Perfect sets are closed; in a Polish space they are therefore G_δ . Therefore the set $A \setminus \{x\}$ is a Borel set in X . Since Borel sets have the perfect set property and $A \setminus \{x\}$ is uncountable, $A \setminus \{x\}$ includes a non-empty perfect set. \square

The following proposition follows immediately from the definition of \cong_T , and will be key for us later.

Proposition 2.3. *Let S be a countable signature, and let T be an S -theory of some logic. The complexity of the equivalence relation \cong_T is the minimum complexity that includes both Σ_1^1 and the complexity of the complement of $\text{Mod}_S(T)$.*

We end this section with a standard definition that will be used many times in what follows.

Definition 2.4. An equivalence relation E on a Polish space X is *thin* if there is no perfect set of pairwise E -inequivalent elements of X .

In particular, many of the results we are interested in are about counting the number of equivalence classes of \cong_T in cases where that relation is thin.

2.2. Second-order logic. The logics we study in this paper are all closely related to second-order logic, so we include a review of that logic here. The reader familiar with the “full” semantics of second-order logic can safely skip this section, while the reader interested in more about second-order logic should consult [45].

On the syntactic side the definitions closely follow the corresponding recursive definitions for first-order logic, but with additional clauses describing the use of second-order variables. Indeed, the first significant difference from first-order logic is that our formulas will allow several kinds of variables, specifically:

- Variables to represent individual elements of structures. (These are the *first-order variables*).
- For each n , variables to represent sets of n -tuples of elements. (These are the *n -ary relation variables*).

The *atomic second-order S -formulas* are defined recursively:

- If x and x' are first-order variables, then $x = x'$ is an atomic S -formula.
- If x_1, \dots, x_n are first-order variables, and R is an n -ary relation symbol in S , then $R(x_1, \dots, x_n)$ is an atomic S -formula.
- If x_1, \dots, x_n are first-order variables, and U is an n -ary relation variable, then $U(x_1, \dots, x_n)$ is an atomic S -formula.

The *second-order S -formulas* are defined recursively:

- Atomic second-order S -formulas are second-order S -formulas.
- The second-order S -formulas are closed under conjunction, disjunction, and negation.
- If ϕ is an S -formula and x is a first-order variable then $(\exists x)\phi$ and $(\forall x)\phi$ are second-order S -formulas.
- If ϕ is an S -formula and U is a relation variable then $(\exists U)\phi$ and $(\forall U)\phi$ are second-order S -formulas.

In cases where it helps with clarity, we sometimes indicate that a quantifier is a second-order quantifier by adding a superscript 1; thus the notation \exists^1 is sometimes used for emphasis when we are using existential quantification over a second-order variable. Analogously, \forall^1 is used to denote a second-order universal quantifier.

Finally, we come to the second-order satisfaction relation. We use this relation with the *full semantics*. These semantics are defined as for first-order logic, with the following additions. Suppose that $\phi(P)$ is a second-order S -formula, where P is a relation variable (ϕ may have other variables that are not displayed).

- If P is an n -ary relation variable and $A \subseteq M^n$, then $\mathcal{M} \models \phi(A)$ if and only if $(\mathcal{M}, A) \models \phi$, where (\mathcal{M}, A) is the expanded structure obtained by interpreting P as A .
- $\mathcal{M} \models (\exists P)\phi(P)$ if and only if there is some $A \subseteq M^n$ such that $\mathcal{M} \models \phi(A)$. The definition for the second-order universal quantifier is similar.

While we will not be directly dealing with the meta-mathematical properties of second-order logic in this paper, we emphasize to the reader that second-order model theory is significantly different from first-order model theory. In particular, both the compactness and Löwenheim-Skolem theorems fail for second-order logic.

In Section 5 we will need to consider restricted classes of second-order formulas. For that purpose the following fact (see [45, Section 4]) is very useful:

Fact 2.5. *Every second-order formula is equivalent to a second-order formula in prenex normal form; that is, to a formula where all quantifiers appear at the beginning of the formula, and all second-order quantifiers precede all first-order quantifiers.*

In light of this fact, when discussing second-order formulas in general we will assume they are already written in prenex normal form. To avoid overlap of terminology, we will say that a second-order formula is \forall_n if it is equivalent to a prenex formula that begins with a second-order universal quantifier and has a total of n blocks of quantifiers, and likewise a formula is \exists_n if it is equivalent to a prenex formula that begins with a second-order existential quantifier and has a total of n blocks of quantifiers. We refer to a second-order theory as a \forall_n theory if it has an axiomatization using sentences that are at most \forall_n , and likewise for \exists_n theories.

2.3. Descriptive complexity of second-order theories. Morley's proof of his result in [29] relies heavily on the fact that $\text{Mod}_S(\sigma)$ is Borel when σ is an $L_{\omega_1, \omega}$ sentence. In the context of a sentence of second-order logic the sets $\text{Mod}_S(\sigma)$ are no longer necessarily Borel. Instead, the models of a single second-order sentence will form a projective set, with the complexity of the set being determined by the quantifier complexity of the sentence.

Lemma 2.6. *Let S be a countable signature, and let σ be a second-order S -sentence. Then $\text{Mod}_S(\sigma)$ is projective. More specifically, if σ is a \forall_n formula then $\text{Mod}_S(\sigma)$ is a Π_n^1 set, and if σ is an \exists_n formula then $\text{Mod}_S(\sigma)$ is a Σ_n^1 set.*

Proof. The proof is by induction on the complexity of σ . It is well-known (see e.g. [10, Lemma 11.3.3]) that if σ is first-order then $\text{Mod}_S(\sigma)$ is Borel. Conjunctions correspond to intersections and negations to complements, so the case of interest is the second-order existential quantifier.

Suppose that \mathcal{M} is an S -structure, and X is a second-order variable, and that $\mathcal{M} \models (\exists X)\varphi(X)$. Then there is an $S \cup \{X\}$ -structure \mathcal{M}' such that $\mathcal{M}' \models \varphi$ and its reduct to S is \mathcal{M} , i.e. $\mathcal{M}' \upharpoonright S = \mathcal{M}$. It is easily verified that the projection $f : \text{Mod}_{S \cup \{X\}} \rightarrow \text{Mod}_S$ given by $f(\mathcal{M}) = \mathcal{M} \upharpoonright S$ is continuous. We have:

$$\begin{aligned} \text{Mod}_S(\exists X\varphi) &= \{\mathcal{M} \in \text{Mod}_S : \mathcal{M} \models (\exists X)\varphi(X)\} \\ &= \{\mathcal{M} \in \text{Mod}_S : (\exists \mathcal{N} \in \text{Mod}_{S \cup \{X\}})\mathcal{N} \models \varphi(X) \text{ and } \mathcal{N} \upharpoonright S = \mathcal{M}\} \\ &= \{\mathcal{M} \in \text{Mod}_S : (\exists \mathcal{N} \in \text{Mod}_{S \cup \{X\}}(\varphi))f(\mathcal{N}) = \mathcal{M}\} \\ &= f[\text{Mod}_{S \cup \{X\}}(\varphi)] \end{aligned}$$

Thus the second-order existential quantifier corresponds to projection, completing the proof. \square

It is easy to see that the relation of isomorphism of countable structures is a Σ_1^1 relation on Mod_S . Combining this with Lemma 2.6 and the fact that the projective classes Π_n^1 and Σ_n^1 are closed under countable intersection (see [20, Proposition 37.1]) we obtain:

Proposition 2.7. *If T is a second-order theory of bounded quantifier complexity, then \cong_T is a projective equivalence relation. More specifically, for $n > 1$, if T is a \forall_n theory then \cong_T is a Σ_n^1 relation, and if T is an \exists_n theory then \cong_T is a Π_n^1 relation. If T is an existential second-order sentence (i.e. \exists_1), then \cong_T is Δ_2^1 .*

If we wish to consider second-order theories of unbounded quantifier complexity then we will need the following definition that generalizes the projective hierarchy:

Definition 2.8. The collection of σ -projective sets is the smallest σ -algebra containing the open subsets (of \mathbb{R}) and closed under projections.

Lemma 2.9. *Let S be a countable signature, and let T be a second-order S -theory. Then $\text{Mod}_S(T)$ is σ -projective; it is in fact a countable intersection of projective sets.*

Proof. $\text{Mod}_S(T) = \bigcap_{\sigma \in T} \text{Mod}_S(\sigma)$, so this follows directly from Lemma 2.6. \square

3. MORLEY'S THEOREM FAILS CONSISTENTLY FOR SECOND-ORDER LOGIC

Our strategy for showing the consistent failure of Morley's Theorem for second-order logic is to force over L to add \aleph_2 Cohen reals, and then force over the resulting model to add \aleph_3 random reals. After that forcing, we have $2^{\aleph_0} = \aleph_3$. In this final model $L[G][H]$ of set theory there are exactly \aleph_2 Cohen reals over L (random reals don't add Cohen reals: [2, Section 7.2]), which will enable us to construct a second-order theory with exactly \aleph_2 isomorphism classes of countable models.

Recall that a real r is *Cohen* over L if and only if r is not in any first category F_σ set F of reals, F having a constructible code (Solovay, see e.g. [2, Theorem 3.1.3]).

We will need the following lemma:

Lemma 3.1. *If M is a transitive set, then the structure (M, \in) does not have non-trivial automorphisms.*

Proof. Suppose there is a non-trivial automorphism τ of the structure (M, \in) . Since \in is well-founded, let a be an \in -minimal element of M on which τ is not the identity. Now, M is transitive and well-founded, and a and $\tau(a)$ have the same elements. Thus $a = \tau(a)$, a contradiction. \square

We are going to use the following theorem from [18]:

Theorem 3.2. *There is a sentence σ_1 in the language of set theory using an unary predicate symbol \dot{A} as a parameter such that for any A and any transitive class N , $(N, \in, A \cap N) \models \sigma_1$ if and only if $N = L[A]$ or $N = L_\delta[A]$ for some limit $\delta > \omega$.*

Theorem 3.3. *It is consistent with ZFC that there exists a countable second-order theory with exactly \aleph_2 non-isomorphic countable models while the continuum is \aleph_3 .*

Proof. Starting from L , we force to add \aleph_2 Cohen reals over L and then add \aleph_3 random reals over the resulting model. This latter model is then of the form $L[G][H]$ where G and H are the generic filters associated to the Cohen and random forcings, respectively.

We consider the signature $S = \{\in\}$ where \in is a binary relation symbol. We shall be considering only structures where $\mathcal{M} = (M, \in^{\mathcal{M}})$ is a model of a fragment of ZF. To avoid confusion, we denote the ω of a given model by $\omega^{\mathcal{M}}$. Moreover, since ω is definable, we are going to use ω on the syntactic side to denote what would be interpreted as $\omega^{\mathcal{M}}$ in any of our models \mathcal{M} . Analogously, given that we are only going to work with models of enough set theory in which L is definable, we are going to use L to denote the syntactic object that will be interpreted as $L^{\mathcal{M}}$ by a model \mathcal{M} . Note that $(M, \in^{\mathcal{M}})$ is not necessarily transitive.

We shall produce a second-order formula to say that there is a Cohen real over L . For that purpose we use that “ r is a Cohen real over L ” is equivalent to the statement “ r is not in any first category F_σ set F of reals, F having a constructible code” (i.e. in L). We treat first category F_σ sets as follows: fix $\{O_n : n < \omega\}$ to be the base for ω^ω corresponding to open intervals with rational endpoints enumerated in a recursive fashion. If $C = \bigcup_{n < \omega} F_n$ is F_σ and each F_n is closed and nowhere dense, C satisfies:

$$(\forall n < \omega)(\forall m < \omega)(\exists B_n \in \mathcal{P}(\omega))[(F_n = \omega^\omega \setminus \bigcup_{k \in B_n} O_k) \wedge (O_m \not\subseteq F_n) \wedge (C = \bigcup_{k < \omega} F_k)].$$

Notice that the quantification “ $\exists B_n \in \mathcal{P}(\omega)$ ” is a first-order quantification because we are considering only models of (sufficiently large fragments of) ZF, and here the quantification ranges over the model’s version of the power set of its version of ω , which are first-order definable sets in the language of set theory. Furthermore, we denote by A_c the set coded by c as in Kanamori [18, p. 137]. Then we can denote the statement that c is a code for A_c and A_c is an F_σ first category set by the first-order formula $F_\sigma(c)$.

Our objective is to write down a theory such that each of its isomorphism classes contains a model of the form $L_\delta[r]$ where $\delta < \omega_1$ and r is Cohen over L_δ . We use the formula σ_1 from Theorem 3.2 under the following considerations: We use second-order quantification to state the existence of a binary relation that will yield the graph of a Cohen real when restricted to $\omega \times \omega$. Thus the unary predicate \dot{A} from Theorem 3.2 is taken to be simply belonging to the set of pairs satisfying the binary relation R and we denote the resulting formula by σ_1^R .

The following sentence ψ asserts the existence of a Cohen real the graph of which satisfies the formula σ_1^R :

$$\begin{aligned}
& (\exists^1 R)(\forall x)(\forall y)(\forall c \in \omega^\omega)(\exists f \in \omega^\omega)(\forall n \in \omega)(\exists! m \in \omega) \\
& R(n, m) \wedge (R(x, y) \implies (x, y) \in \omega \times \omega) \\
& \wedge \\
& (\forall n)(\forall m)(f(n) = m \iff R(n, m)) \wedge (c \in \mathbf{L} \wedge F_\sigma(c) \implies f \notin A_c) \\
& \wedge \\
& \sigma_1^R
\end{aligned}$$

The sentence ψ only has one second-order (existential) quantifier and so the complexity of $\text{Mod}_S(\psi)$ is Σ_1^1 . Notice that R is the only variable under the scope of a second-order quantifier, being a binary relation symbol. As mentioned before, \mathbf{L} refers to the syntactic placeholder for what will be interpreted as $L^\mathcal{M}$ in any model \mathcal{M} that we consider. Thus $c \in \mathbf{L}$ is just a first-order quantification in ψ . For simplicity, we denote the function (in fact, the real) f that appears in ψ by r .

We emphasize that the sentence σ_1^R , which we take from Kanamori, is a sentence in the signature $S \cup \{R\}$. It is for this reason that we need to use second-order quantification to obtain r ; even though we go on to require that R is closely tied to an element f_R of our model, in using the result from Kanamori it is important that R itself is a relation on our model, not an element of the model.

We proceed with a variation of an argument that appears in [18]. The following sentences constitute our second-order theory T :

- (1) ψ
- (2) $(\forall^1 A)(\exists x \in A)(x \cap A = \emptyset)$.
- (3) Enough ZF including extensionality.

Notice that the resulting complexity of $\text{Mod}(T)$ remains Δ_2^1 , and so by Proposition 2.7 the complexity of the isomorphism relation \cong_T is Δ_2^1 .

In order to count the models of T we will use the following claim:

Claim 1. *For every $\mathcal{N} \models T$ there is a unique ordinal $\gamma < \omega_1$ and real r which is Cohen over L_γ such that $\mathcal{N} \cong L_\gamma[r]$.*

We indiscriminately use the notation $L_\gamma[r]$ to denote both the forcing extension which is obtained by adding the real r to L_γ , and the γ -th level of the hierarchy of sets constructible relative to r . The reason is that when r is generic over L_γ the universes of these two models are the same, see [21, Exercise VII B10].

Our strategy is to count equivalence classes of the isomorphism relation on $\text{Mod}(T)$ by looking at the models in each class that are of the form $L_\gamma[r]$ where $\gamma < \omega_1$ and $L_\gamma[r] \models$ “ r is Cohen over L ”, which means that r is Cohen over L_γ . We show that for any such γ in $L[G][H]$, there are only \aleph_2 Cohen reals over L_γ : suppose there is a Cohen over L_γ real in $L[G][H] \setminus L[G]$. It is standard to see (the reader is referred to [4]) that such a real would be added by a countable initial segment of the random forcing and so it is added by one random real s , since the forcing

for adding one random real is isomorphic to that of adding countably many. Now s being random over $L[G]$ is equivalent to that s is not in any Borel null set with a code in $L[G]$ (Solovay, see e.g. [2, Theorem 3.1.3]), so it is clear that it is also random over $L_\gamma[G]$ (and over L_γ) and so it could not have added any Cohen over $L_\gamma[G]$ (or over L_γ) reals. In $L[G]$ there are only \aleph_2 reals and we have just shown that no element of $L[G][H] \setminus L[G]$ can be Cohen over an L_γ and so, for any $\gamma < \omega_1$, there are only \aleph_2 r 's such that $L_\gamma[r] \models$ “ r is Cohen over L ”.

Since there are only \aleph_2 reals that are Cohen over a given L_γ , and only \aleph_1 choices for γ , the proof is complete once we prove Claim 1.

For existence, suppose that $\mathcal{N} = (N, \in^{\mathcal{N}}) \models T$. By clauses (2) and (3) of the definition of T the structure $(N, \in^{\mathcal{N}})$ is extensional and well-founded, and hence by the Mostowski collapse there is a transitive set X such that $(X, \in) \cong (N, \in^{\mathcal{N}})$, where the membership relation on X is the membership relation from V . The key observation is that, by second-order semantics, since \mathcal{N} “thinks” it is well-founded, it really is. Through this isomorphism, we get a model $\mathcal{N}' \models T$ where the underlying set is X and $\in^{\mathcal{N}'} = \in$. By clause (1) of T and Theorem 3.2, $X = L_\gamma[r]$ for some γ , $\gamma < \omega_1$, where r is Cohen over L . Thus, each isomorphism class of T contains a model of the form $L_\gamma[r]$.

Now, suppose that we are given two different models $L_\alpha[r_1]$ and $L_\beta[r_2]$. It is immediate to see that if $\alpha \neq \beta$, then the two models are non-isomorphic. If $\alpha = \beta$, we distinguish two subcases: First, suppose that r_1 and r_2 are mutually generic and that $L_\alpha[r_1] \cong L_\alpha[r_2]$. Then we must have that the isomorphisms $\pi_{1,2} : L_\alpha[r_1] \rightarrow L_\alpha[r_2]$ and $\pi_{2,1} : L_\alpha[r_2] \rightarrow L_\alpha[r_1]$ must be the identity. Otherwise, notice that their compositions $\pi_{1,2} \circ \pi_{2,1} : L_\alpha[r_1] \rightarrow L_\alpha[r_1]$ and $\pi_{2,1} \circ \pi_{1,2} : L_\alpha[r_2] \rightarrow L_\alpha[r_2]$ yield automorphisms of $L_\alpha[r_1]$ and $L_\alpha[r_2]$ (respectively), and so such compositions must be the identity since these models have no non-trivial automorphisms by Lemma 3.1. Now, since such isomorphisms can never be \in -decreasing (by the same \in -induction argument as in Lemma 3.1) and their compositions yield the identity, they must be identity themselves and so $L_\alpha[r_1] = L_\alpha[r_2]$, which contradicts the mutual genericity of r_1 and r_2 . Secondly, when r_1 and r_2 are not mutually generic, we simply have $L_\alpha[r_1] = L_\alpha[r_2]$ since each is definable from the other. Notice that we have \aleph_2 mutually generic Cohen reals over L in $L[G][H]$. Thus each isomorphism class of T contains an $L_\gamma[r]$ for r Cohen over L_γ and $\gamma < \omega_1$, while two mutually generic Cohen reals over the same L_γ yield non-isomorphic models of T , and so there are \aleph_2 isomorphism classes and yet the continuum is \aleph_3 in $L[G][H]$. \square

4. MORLEY’S THEOREM IS CONSISTENTLY TRUE FOR SECOND-ORDER LOGIC

In this section we sketch that, modulo a supercompact cardinal, the extension of Morley’s result to second-order logic is consistently true. This completes the proof of the undecidability of the second-order version of Morley’s Theorem. We will obtain finer results, with smaller large cardinals, in later sections. In fact, the consistency of the Absolute Morley Theorem follows directly from the following result of Foreman and Magidor.

Theorem 4.1 ([9]). *If it is consistent that there is a supercompact cardinal, then it is consistent that $\neg\text{CH}$ and every equivalence relation on the power set of \mathbb{R} that is in $L(\mathbb{R})$ has $\leq \aleph_1$ or a perfect set of inequivalent elements.*

Foreman and Magidor do not specifically state Theorem 4.1 in [9], but it is implicit in their work. In this section we sketch a proof of a weaker version of Theorem 4.1. Since our intended audience includes model theorists who may not be familiar with large cardinals or inner models of set theory, we first need to briefly explicate “supercompact” and “ $L(\mathbb{R})$ ”. But first, for the set theorist reader, we should mention that with the development of Inner Model Theory since [9], it is now clear that the supercompact cardinal above may be reduced to the assumption that there is a proper class of Woodin cardinals (defined in Section 7). We will consider and prove various extensions and improvements of Theorem 4.1 in Section 7 below.

Definition 4.2. A cardinal κ is *supercompact* if for every cardinal $\lambda \geq \kappa$ there exists an elementary embedding j_λ of V into an inner model M (i.e. a proper class model of ZFC included in V) with *critical point* κ (least ordinal j_λ moves) and $\lambda < j_\lambda(\kappa)$, such that M^λ is included in M .

Supercompactness allows us to reflect certain kinds of relatively simple statements about cardinals equal to or beyond the supercompact cardinal back down to cardinals below it. Supercompactness combined with iterated forcing is a useful method in the establishing of the consistency of unbounded universal statements. See [18, Section 22].

In Section 2 we showed that when σ is a second-order sentence, $\text{Mod}(\sigma)$ is projective; it is not generally Borel. If T is a second-order theory, then $\text{Mod}(T)$ is the countable intersection of projective sets, but these projective sets may have unbounded complexity, so $\text{Mod}(T)$ is σ -projective but may not be projective. Since some results from the literature are set in the broader context of sets in $L(\mathbb{R})$, we remind the reader of that setting.

Definition 4.3. $L(\mathbb{R})$, the collection of all sets constructible from \mathbb{R} , is the smallest inner model of V containing \mathbb{R} as a member. It is defined in analogy to L and $L[\mathbb{R}]$ by:

$$\begin{aligned} L_0(\mathbb{R}) &= \text{Trc}(\mathbb{R}) \text{ the transitive closure of } \mathbb{R} \\ L_{\alpha+1}(\mathbb{R}) &= \text{Def}(L_\alpha(\mathbb{R})) \\ L_\delta(\mathbb{R}) &= \bigcup_{\alpha < \delta} L_\alpha(\mathbb{R}) \text{ for limit } \delta > 0 \\ L(\mathbb{R}) &= \bigcup_{\alpha \in ON} L_\alpha(\mathbb{R}) \end{aligned}$$

We are actually interested in $L(\mathbb{R}) \cap \mathcal{P}(\mathbb{R})$, but will usually just write $L(\mathbb{R})$. Notice that each element C of $L(\mathbb{R})$ can be defined by a formula of the language of set theory with a real and finitely many ordinals as parameters. We speak of the real together with the ordinals as a “code” of C . We will speak more about coding later. In particular, by coding, we can easily see that σ -projective sets are in $L(\mathbb{R})$. Weaker large cardinal hypotheses suffice for our σ -projective purposes, as we shall see in Sections 5 and 7, and these are sufficient for our applications to second-order logic. We do not know of any interesting logics that lead us beyond the σ -projective sets toward $L(\mathbb{R})$, but any logic we can define by using a formula of set theory with

finitely many real and ordinal parameters is fair game. The model of set theory that Foreman and Magidor use is actually one familiar to set theorists. It is the usual model for the *Proper Forcing Axiom*.

- Definition 4.4.**
- (1) If κ is a cardinal, $H(\kappa)$ is the set of all sets the transitive closures of which have cardinality strictly less than κ . It is known that if κ is regular uncountable, $(H(\kappa), \in)$ models ZFC except perhaps for the power set axiom.
 - (2) Given a forcing notion P and κ a regular cardinal satisfying $\kappa > 2^{|P|}$, we say that a countable elementary submodel N of $H(\kappa)$ is *suitable for P* if $\mathcal{P}(P) \in N$.
 - (3) Given a forcing notion P , fix κ a regular cardinal satisfying $\kappa > 2^{|P|}$. We say that P is *proper* if whenever $p \in P$, there is a $q \in P$ such that $q \leq p$ and for every suitable countable elementary submodel M of $H(\kappa)$, and every $D \in M$ which is dense in P , $D \cap M$ is predense below q (i.e. every $r \leq q$ is compatible with an element of $D \cap M$).

The Proper Forcing Axiom (PFA) is like MA_{\aleph_1} , but instead of meeting \aleph_1 dense sets for countable chain condition partial orders, one does this for the wider class of proper partial orders. The usual model is obtained by iterating proper posets of size less than a supercompact cardinal κ via countable support iteration κ many times, and then using a reflection argument to argue that all collections of \aleph_1 dense sets in any proper partial order have been dealt with. For details, see e.g. [17]. It is known that PFA implies $2^{\aleph_0} = \aleph_2$ [3, 47].

We will briefly sketch the ideas of the Foreman-Magidor proof in the special case we are interested in, namely equivalence relations - such as σ -projective ones - that are coded by a real. A detailed analysis of the proof will appear in Section 7, where we find weaker hypotheses that still enable us to compute the possible numbers of equivalence classes for thin equivalence relations definable more stringently than just being in $L(\mathbb{R})$. For example, let us consider all sets of reals - e.g. the σ -projective ones - that are constructible from a real. The idea of the Foreman-Magidor proof is to note that a code for such a set (being a real, it is of size less than the supercompact) appears at some initial stage of the iteration; if such an equivalence relation is thin, then they show that the interpretation of the code when the code appears is also thin (“downwards generic absoluteness”), and that the rest of the proper forcing adds no new equivalence classes to thin equivalence relations (“upwards generic absoluteness”). When the code appears at some initial stage, we may without loss of generality assume CH holds at that stage, since it holds cofinally often in the iteration. At that stage then, its interpretation has no more than \aleph_1 equivalence classes. By upwards generic absoluteness, its interpretation at the final stage, which is just the equivalence relation we started with, then has no more than \aleph_1 equivalence classes. It is convenient that we can refer to a well-known model of set theory, but the Foreman-Magidor proof does not actually use most of the properties of that model. For further discussion, see Section 7.

This sketch sounds to a set theorist like a typical supercompact reflection proof, but there are some nuances. In Section 7, with the benefit of research after [9], we will prove sharper results. We need to talk about codes. Think for example of the pair $\langle 0, 1 \rangle$ as the real $2^1 \times 3^2$. This real can be thought of as coding the

open interval $(0, 1)$, which of course has different extensions in different models. σ -projective sets are coded by a real, just as Borel sets are. The reader is probably familiar with the idea of coding a Borel set: one first lists the pairs of rational numbers in some recursive way, then countable sequences of such pairs to code open sets, then countable sequences of these, as well as countable intersections of complements of these, then continue recursively to list countable sequences of what has gone before, etc. For a careful exposition of coding, see [22]. Sets in $L(\mathbb{R})$ are coded by a real plus finitely many ordinals. This makes things more complicated, since those ordinals may be bigger than the large cardinal that we are collapsing to $\aleph_2 = 2^{\aleph_0}$. Since we don't currently have any examples of interesting logics that are in $L(\mathbb{R})$ but are not definable from a real, here we just sketched the Foreman-Magidor proof for logics that are in $L(\mathbb{R})$ and definable from a real. We will prove - indeed improve - their theorem later.

5. FINER ANALYSIS: DESCRIPTIVE SET THEORY

In the previous section we noted that from the consistency of a supercompact cardinal we can obtain the consistency of the Absolute Morley Theorem for second-order logic. In this section we start examining how weakenings of the $L(\mathbb{R})$ results of Foreman-Magidor [9] can be achieved via weaker assumptions. We achieve this by cobbling together results from the literature. In Section 7 we employ the machinery of Inner Model Theory to achieve more precise conclusions.

Most of this section concerns Descriptive Set Theory, and as such when we discuss issues of definability we mean "definable" in the sense used in Descriptive Set Theory. In particular, typical definability notions we consider are "Borel", "analytic", "projective", or being in $L(\mathbb{R})$.

Morley and Burgess use that projections of Borel sets are analytic. To extend their results to second-order logic we will use that countable intersections of projective sets are σ -projective. Morley quotes the classical theorem of Descriptive Set Theory that analytic sets have the perfect set property; we want to make similar assertions for σ -projective sets. However such assertions are no longer theorems of ZFC: Gödel showed that under $V = L$, there is an uncountable co-analytic (Π_1^1) set which does not include a perfect set. See e.g. [18] for a proof. Co-analytic sets do, however, have cardinality $\leq \aleph_1$ or 2^{\aleph_0} .

One way of extending the perfect set property to more complicated sets of reals is via large cardinals, which imply restrictions of the Axiom of Determinacy to various classes of definable sets of reals, which in turn imply that if such sets are uncountable, they include perfect sets. We refer the reader to Section 27 of [18] for an introduction to determinacy and the formal definitions of games and winning strategies. In the standard Descriptive Set Theory abuse of notation, we think of \mathbb{R} as ω^ω . This is harmless since the descriptive structures of the reals, of the Baire space ω^ω and of the Cantor space are isomorphic.

Definition 5.1. Let \mathcal{C} be a collection of subsets of \mathbb{R} (we are interested in \mathcal{C} 's which are composed of sets which are "definable" in some sense). $AD_{\mathcal{C}}$ is the assertion that given any $C \in \mathcal{C}$, in any game where players alternately pick natural numbers,

with Player I trying to get the resulting infinite sequence to be in C and Player II trying to prevent that, one of the players has a winning strategy. *Projective determinacy (PD)* is the assertion $\text{AD}_{\mathcal{P}}$ where \mathcal{P} is the collection of all projective sets. For specific levels of the projective hierarchy, instead of e.g. $\text{AD}_{\Sigma_{17}^1}$, it is common to write $\text{Det}(\Sigma_{17}^1)$.

Letting \mathcal{B} stand for the collection of all Borel sets, $\text{AD}_{\mathcal{B}}$ is a theorem of ZFC ([24], see also [20]); AD for progressively larger definable classes follows from progressively larger cardinals. $L(\mathbb{R})$ cannot satisfy the Axiom of Choice if we suppose that it satisfies the Axiom of Determinacy, but there is much to be said for assuming $\text{AD}_{L(\mathbb{R})}$ within a ZFC environment, since it follows from large cardinals and imposes a pleasant regularity on the constructible sets of reals. See [18, Section 32]. For example, note the following result from [50]:

Theorem 5.2 (Steel-Woodin). *If there is a supercompact cardinal (much weaker cardinals suffice), then $\text{AD}_{L(\mathbb{R})}$ holds.*

The conclusion says that determinacy holds for all constructible sets of reals. For familiar classes \mathcal{C} , $\text{AD}_{\mathcal{C}}$ implies every uncountable member of \mathcal{C} includes a perfect set [6].

We want to build a model of set theory in which the isomorphism relation for countable models of a second-order theory has either at most \aleph_1 classes or has a perfect set of non-isomorphic models. We don't know of any results that deal specifically with this question, but there is a long line of research extending Silver [38] which tries to count the number of equivalence classes of a definable equivalence relation among sets of reals.

5.1. Thin equivalence relations. A great deal is known about equivalence relations on Polish spaces that do, or do not, have a perfect set of pairwise inequivalent elements. Here we remind the reader of some of the standard terminology and some relevant results from the literature. We also prove that Morley's Theorem extends to sentences of universal second-order logic.

Although results about determinacy and the perfect set property are motivating, there does not seem to be a direct link between a projective class having the perfect set property and whether an equivalence relation of that complexity has perfectly many equivalence classes. In fact, Σ_2^1 has the perfect set property because each Σ_2^1 set is the union of \aleph_1 Borel sets (a classical result), but it is undecidable, modulo large cardinals, whether or not thin Σ_2^1 equivalence relations have $\leq \aleph_1$ equivalence classes.

Recall from Section 2 that an equivalence relation E on a Polish space X is *thin* if there is no perfect set of pairwise E -inequivalent elements of X . The theorem of Burgess [5] mentioned in the introduction can thus be stated as follows:

Theorem 5.3 (Burgess). *A Σ_1^1 thin equivalence relation on a Polish space X has at most \aleph_1 equivalence classes.*

As an immediate consequence, we obtain the following generalization of Morley's result.

Theorem 5.4. *Let S be a countable signature, and let σ be a universal second-order S sentence (that is, a sentence for which the second-order quantifiers in the prenex form are all universal). Either there is a perfect set of non-isomorphic models of σ , or there are at most \aleph_1 non-isomorphic models of σ .*

Proof. By Proposition 2.7 the relation \cong_σ is Σ_1^1 , and hence Burgess' theorem applies. \square

Schlicht's Example 5.7 below establishes limits to what we can prove in ZFC plus large cardinals about the number of equivalence classes of thin projective equivalence relations. We shall see below, however, that large cardinals do enable us to extend ZFC results. Of course generic extensions of large cardinal models give us even more power: viz. [9].

Example 5.5. Let $S = \{<\}$, and let σ be the second-order sentence expressing that $<$ is a well-order. That is, σ is the conjunction of the (first-order) axioms for being a linear order with the second-order statement

$$(\forall^1 A) (((\exists z) A(z)) \rightarrow (\exists x) (A(x) \wedge (\forall y)(A(y) \rightarrow x \leq y))).$$

Up to isomorphism the countable models of σ are exactly the countable ordinals, so σ has \aleph_1 pairwise non-isomorphic models, regardless of the size of the continuum. This example shows that Vaught's Conjecture is consistently false (and hence undecidable, since it is true assuming the continuum hypothesis) for second-order logic, and even for single sentences of universal second-order logic.

For this example, σ is a universal second-order statement, so \cong_σ is Σ_1^1 by Proposition 2.7. This example shows that Silver's Dichotomy for Π_1^1 equivalence relations (see [10, Theorem 5.3.5]) does not extend to Σ_1^1 relations. Since isomorphism of countable structures is already Σ_1^1 even without restricting to models of a specific sentence, we see that there is little hope for directly applying Silver's Dichotomy to make progress on Vaught's conjecture, even for first-order theories.

The situation for existential second-order sentences is no better than it is for universal sentences, as shown by the next example, which is due to Kunen (see [39, 1.4.3]).

Example 5.6. Let $S = \{<\}$, and let σ be the second-order sentence obtained by taking the conjunction of the (first-order) axioms for $<$ being a linear order with the second-order statement expressing that for any two elements a and b there is an automorphism of $<$ sending a to b . Kunen shows that every model of σ is of the form Z^α or $Z^\alpha \cdot \eta$ for some countable ordinal α , where $Z = \{0\}$, the product order is lexicographic, and η is the ordertype of \mathbb{Q} . In particular, σ has exactly \aleph_1 countable models (and Steel [39] also points out that σ does not have a perfect set of non-isomorphic countable models).

For this example σ is an existential second-order statement, so \cong_σ is Δ_2^1 by Proposition 2.7.

The third example we wish to consider is due to Schlicht [36]. Some background is required before we can state the example. Recall that a *prewellordering* on a set X is a relation \leq on X which is reflexive, transitive, connected (i.e. for any $x, y \in X$, $x \leq y$ or $y \leq x$), and every nonempty subset of X has a least element. There is a natural equivalence relation associated with a prewellorder: $x \sim y$ if and only if $x \leq y$ and $y \leq x$. It is interesting to note that, as pointed out in [36, Remark 5.25], Projective Determinacy (*PD*) implies every thin projective equivalence relation is induced by a projective prewellorder.

For $n \geq 1$, the n th *projective ordinal* δ_n^1 is the supremum of lengths of Δ_n^1 prewellorders. The statement that “ $x^\#$ exists” is equivalent to there being a non-trivial elementary embedding from $L[x]$ to $L[x]$ - see e.g. [18, Section 9]. If there is a measurable cardinal, or even just a cardinal κ such that $\kappa \rightarrow (\omega)_2^{<\omega}$, then $x^\#$ exists for every real x . *PD* also implies this conclusion. *PD* follows from the existence of infinitely many Woodin cardinals (defined in Section 7); $AD_{L(\mathbb{R})}$ follows from the existence of infinitely many Woodin cardinals with a measurable above all of them, see e.g. [33].

We are now ready to describe Schlicht’s example. For more details, see [36].

Example 5.7. Assume $x^\#$ exists for every real x . Let $\langle \iota_\alpha^x : \alpha \in ORD \rangle$ enumerate the x -indiscernibles and define $u_2^x = \iota_{\omega_1+1}^x$. The prewellorder defined by

$$x \leq y \iff u_2^x \leq u_2^y$$

is Δ_3^1 and has length δ_2^1 . On the other hand, assuming ZF plus PD, Martin proved that $\delta_2^1 \leq \aleph_2$, and that $\delta_1^1 = \aleph_1$. But, also assuming ZF plus PD, Kechris and Moschovakis proved $\delta_1^1 < \delta_2^1$ (the results of Martin and Kechris-Moschovakis are found in [19]). The length of a prewellorder is equal to the number of equivalence classes of the corresponding equivalence relation. Thus, assuming ZF plus PD, we get a Δ_3^1 equivalence relation with exactly \aleph_2 countable models.

This example shows that the result of Foreman and Magidor (used in Section 4) in the usual PFA model does not hold in ZFC plus a sufficiently large cardinal. Indeed, suppose for example that we have a supercompact cardinal, so that PD holds and we have sharps. Add \aleph_3 Cohen reals. The supercompact is still supercompact but Schlicht’s Δ_3^1 example exists and is thin and of size \aleph_2 .

In between Burgess’ dichotomy for Σ_1^1 equivalence relations and Schlicht’s Δ_3^1 example, one might wonder about Σ_2^1 equivalence relations, especially in view of the Shoenfield Absoluteness Theorem. The following result is encouraging:

Lemma 5.8 ([12]). *Assume PD. Thin Σ_2^1 equivalence relations are Δ_2^1 .*

However, Schlicht’s analysis of the number of classes of projective equivalence relations specializes in the Σ_2^1 case to yield:

Theorem 5.9. *In ZFC there is no upper bound below 2^{\aleph_0} for the number of equivalence classes of thin Σ_2^1 equivalence relations.*

Proof. In the proof of Theorem 3.3, rather than adding \aleph_2 Cohen reals and \aleph_3 random reals, we could have added κ many Cohen reals and κ^+ many random

reals, for any regular κ , to get 2^{\aleph_0} to be κ^+ while there is a thin Δ_2^1 equivalence relation with κ many equivalence classes. \square

Theorem 5.9 is actually a special case of Schlicht's Lemma 4.17 in [36], but that is not so easy to see, since his statement and proof involve Woodin cardinals, determinacy, and *premise*. He uses Harrington forcing from [11] instead of random forcing. Also note that our result is somewhat stronger: *there is no upper bound below 2^{\aleph_0} on the cardinality of thin Δ_2^1 equivalence relations.*

Thus, even for the relatively simple case of Δ_2^1 we must either assume large cardinals or, as in the Foreman-Magidor theorem, consider generic extensions of the universe in order to get a Burgess type result. What we can get is:

Theorem 5.10. *Assume PD. Then thin Σ_2^1 equivalence relations have at most \aleph_1 equivalence classes.*

Proof. Harrington-Shelah [13] showed the conclusion for Π_2^1 , assuming there was a Cohen real over L . They then applied Lemma 5.8, getting Σ_2^1 thin equivalence relations are Π_2^1 . But *PD* - indeed $\text{Det}(\Pi_1^1)$ - implies there is a Cohen real over L because it implies $\aleph_1^L (= (2^{\aleph_0})^L)$ is countable, so there are only countably many dense sets to meet. See more on Harrington-Shelah in Sections 5.2 and 6.1. \square

Corollary 5.11. *If there is a supercompact cardinal, then thin Σ_2^1 equivalence relations have at most \aleph_1 equivalence classes.*

Remark 5.12. Theorem 5.10 and Corollary 5.11 do not contradict Theorem 3.3: in a universe in which PD holds, our Cohen plus random forcing over L only creates an equivalence relation with \aleph_2^L , i.e. countably many, equivalence classes.

Remark 5.13. There is another interesting way of looking at Theorem 5.9. Harrington [11] proved:

Lemma 5.14. *There is a model of ZFC in which:*

- (1) 2^{\aleph_0} can be as large as one likes,
- (2) Martin's Axiom holds,
- (3) Every set of reals of size less than 2^{\aleph_0} is Π_2^1 .

Given a set A of reals, consider the trivial equivalence relation $E(A)$ on \mathbb{R} defined by:

$$x \sim y \text{ if and only if } x = y \text{ or } x \notin A \text{ or } y \notin A.$$

Then E has $|A| + 1$ equivalence classes. If A is Π_2^1 , then $E(A)$ is Σ_2^1 . Thus in Harrington's model, for each $\kappa < 2^{\aleph_0}$, there are thin Σ_2^1 equivalence relations with κ -many equivalence classes. Again, this does not contradict Theorem 5.10 because Harrington's model is constructed as a cardinal-preserving extension of a model in which $\aleph_1 = \aleph_1^L$, which contradicts PD.

Remark 5.15. Using newer Inner Model Theory techniques, the assumption of PD in Lemma 5.8 and Theorem 5.10 can be considerably reduced, as can the assumption

of supercompactness in Corollary 5.11. Woodin cardinals are defined in Section 7. We have:

Theorem 5.16. *Assume there is a Woodin cardinal with a measurable cardinal above; then thin Σ_2^1 equivalence relations are Δ_2^1 .*

Proof. See [27] and [16, Lemma 2.5]. □

Corollary 5.17. *Assume there is a Woodin cardinal with a measurable cardinal above. Then thin Σ_2^1 equivalence relations have at most \aleph_1 equivalence classes.*

Note that a measurable cardinal yields Π_1^1 -determinacy [23].

5.2. \aleph_1 -Souslin and Co- \aleph_1 -Souslin Equivalence Relations. In Foreman-Magidor [9] and other works involving determinacy, κ -Souslin sets and their complements play a role. They also appear in the study of thin equivalence relations.

Definition 5.18 ([13]).

- (a) T is a tree on the set Y if: $T \subseteq Y^{<\omega}$, and $(\eta \in T, \tau \subseteq \eta) \implies \tau \in T$.
- (b) For T a tree on Y , $[T] := \{f : \omega \rightarrow Y, (\forall n)(f|_n \in T)\}$.
- (c) For T a tree on $\kappa \times X$, let

$$p[T] := \{g : \omega \rightarrow X, \text{ and for some } h : \omega \rightarrow \kappa, \langle h, g \rangle \in [T]\}.$$

(Here we identify $\langle h, g \rangle$ with the function $f : \omega \rightarrow \kappa \times X$ where $f(n) = \langle h(n), g(n) \rangle$).

- (d) A binary relation R on ω^ω is κ -Souslin (via T) if: T is a tree on $\kappa \times (\omega^2)$ and $R = p[T]$; (R is co- κ -Souslin if: \overline{R} is κ -Souslin (where \overline{R} = complement of R)).

The following theorem follows easily from results in [13].

Theorem 5.19. *The assumption MA_{ω_1} implies thin co- \aleph_1 -Souslin equivalence relations on \mathbb{R} have $\leq \aleph_1$ equivalence classes.*

Proof. Since the tree T for the complement of the co- \aleph_1 -Souslin equivalence relation E has cardinality \aleph_1 , it follows that $L[T] \models \text{CH}$, so $|2^\omega|^{L[T]} = \aleph_1$, which MA_{ω_1} implies is $< 2^{\aleph_0}$. Then MA_{ω_1} easily implies there is a real Cohen-generic over $L[T]$. That, according to [13], implies E remains an equivalence relation after a Cohen real is adjoined to $L[T]$. We now need another result from [13]:

Lemma 5.20. *Suppose E is a thin co- κ -Souslin relation via T , i.e. T is the tree for the complement of E . Assume E is an equivalence relation after adding a Cohen real to $L[T]$. Then E has at most κ equivalence classes.*

We refer to the second proof of this in [13]. The first “proof” assumes “not strongly thick” rather than “thin”, but only defines “strongly thick” for κ -Souslin relations. Furthermore it claims “the method of [] (sic) shows that E is strongly thick.” □

There is a connection between (co)- κ -Souslin and our previous discussion of Σ_2^1 and Π_2^1 . First we quote:

Lemma 5.21 ([25]). *A relation R is \aleph_1 -Souslin if and only if it is the union of \aleph_1 Borel sets.*

The relationship between \aleph_1 -Souslin and Σ_2^1 is quite interesting. It is a classical result that

Lemma 5.22. *Every Σ_2^1 set is the union of \aleph_1 Borel sets, i.e. is \aleph_1 -Souslin.*

Thus Π_2^1 sets are co- \aleph_1 -Souslin, so we have

Corollary 5.23. *The assumption MA_{ω_1} implies that Π_2^1 equivalence relations have $\leq \aleph_1$ equivalence classes.*

Corollary 5.24. *The assumption MA_{ω_1} implies an existential second-order theory has either a perfect set of pairwise non-isomorphic countable models or has $\leq \aleph_1$ of them.*

On the other hand, Martin and Solovay [26] prove:

Lemma 5.25. *Assume MA_{ω_1} . If there is a $t \subseteq \omega$ such that $\omega_1 = \omega_1^{L[t]}$, then the union of \aleph_1 Borel sets is Σ_2^1 .*

They show that their extra hypothesis is necessary. Interestingly, their use of almost disjoint coding to make an \aleph_1 -Souslin set Σ_2^1 is also found in Harrington [11], which Schlicht used instead of random forcing to prove Theorem 5.9.

Problem 2. *Is it consistent that CH fails but that thin \aleph_1 -Souslin equivalence relations have $\leq \aleph_1$ equivalence classes?*

One is tempted to apply Theorem 5.10 plus Lemma 5.25, but their hypotheses are not mutually consistent.

6. OTHER LOGICS

In this section we consider the applicability of our results to logics other than second-order logic.

6.1. Game quantifiers. Moschovakis [30, 31] and others, e.g. [45], have considered well-ordered quantifiers of length ω . The question of the number of equivalence classes for theories involving these fits in nicely with topics we have been discussing. In particular, we have the closed and open game quantifiers discussed in [30]:

Definition 6.1. The expression

$$(\forall x_0 \exists y_0 \forall x_1 \exists y_1 \dots) \bigwedge_{n < \omega} \varphi_n(x_0, y_0, \dots, x_n, y_n)$$

is a *closed game quantifier sentence of length ω* . The truth value of this game expression in a model \mathcal{M} is equivalent to the existence of a winning strategy for

Player II in the following game:

$$\begin{array}{c|ccc} \text{I} & a_0 & a_1 & \dots \\ \hline \text{II} & b_0 & b_1 & \dots \end{array}$$

Here $a_0, b_0, a_1, b_1, \dots$ are elements of M and b_0, b_1, \dots are chosen such that

$$\mathcal{M} \models \varphi_0(a_0, b_0),$$

$$\mathcal{M} \models \varphi_1(a_0, b_0, a_1, b_1),$$

etc. (otherwise Player II loses).

The open game quantifier interchanges the universal and existential quantifiers and exchanges the infinite conjunction for an infinite disjunction. Thus, the negation of a closed game quantifier sentence is an open game quantifier sentence. Explicitly, an *open game quantifier sentence of length ω* has the following form:

$$(\exists x_0 \forall y_0 \exists x_1 \forall y_1 \dots) \bigvee_{n < \omega} \varphi_n(x_0, y_0, \dots, x_n, y_n)$$

The truth value of an open game quantifier sentence in a model \mathcal{M} is equivalent to the existence of at least one play of the game in which Player I wins; equivalently, Player II does not have a winning strategy. Note that, unless there are enough determinacy assumptions, this does not necessarily imply that Player I does have a winning strategy.

Definition 6.2 ([31]). A *pointclass* Γ is a collection of sets such that each element of Γ is a subset of some product space X . A pointclass is *adequate* if it contains all recursive pointsets and is closed under recursive substitution, \wedge , \vee , and bounded existential and universal quantification. For simplicity, all our pointclasses are boldface.

For example, each level Σ_n^1 or Π_n^1 , $n < \omega$, of the projective hierarchy is adequate.

Now we introduce the game operator \wp following Moschovakis. The reader is referred to [31] for a thorough exposition on the subject.

Definition 6.3. (1) Given a set $P \subseteq \mathcal{X} \times \omega^\omega$, we define the set $\wp P$ as follows:
 $x \in \wp P \iff$ Player I wins the game $\{\alpha : P(x, \alpha)\}$, i.e. the game with parameter x where Player I plays each $\alpha(2n)$ and Player II plays each $\alpha(2n+1)$, and Player I wins if and only if $P(x, \alpha)$.
 (2) Given a pointclass Γ , we define $\wp \Gamma = \{\wp P : P \subseteq \mathcal{X} \times \omega^\omega \wedge P \in \Gamma\}$.

The following theorem relates the game operator \wp and the open game quantifier:

Theorem 6.4. *Given a language, fix a theory T and a sequence of formulas $\langle \varphi_n : n < \omega \rangle$ for which there is an adequate pointclass Γ such that for any $\mathcal{M} \in \text{Mod}(T)$ and $a_1, a_2, \dots \in M$,*

$$\text{Mod}(T \wedge \bigvee_{n < \omega} \varphi_n(a_1, \dots, a_{2n})) \in \Gamma$$

(as a set of reals coding the respective models). Then the formula ψ defined by applying an open game quantifier to the sequence $\langle \varphi_n : n < \omega \rangle$ satisfies

$$\text{Mod}(\psi) \in \mathfrak{D}\Gamma.$$

Proof. For simplicity, we assume that our models are enumerated and that their elements are exactly the elements of ω . Define P to be the following set:

$$\{(\mathcal{M}, \alpha) \in \text{Mod}(T) \times \omega^\omega : \mathcal{M} \models T \wedge (\exists 0 < n < \omega) \mathcal{M} \models \varphi_n(\alpha(0), \alpha(1), \dots, \alpha(2n-1))\}.$$

Notice that for a fixed model \mathcal{M} of T , the projection $\{\alpha : (\mathcal{M}, \alpha) \in P\}$ corresponds to the set of plays which result in Player I winning the game corresponding to ψ . Thus $\mathfrak{D}P$ is the set of models for which Player I wins the game and so $\mathfrak{D}P = \text{Mod}(\psi)$.

Finally, notice that $(\mathcal{M}, \alpha) \in P$ if and only if

$$\mathcal{M} \models T \wedge \bigvee_{n < \omega} \varphi_n(\alpha(0), \alpha(1), \dots, \alpha(2n-1))$$

That is $(\mathcal{M}, \alpha) \in P$ if and only if $\mathcal{M} \in \text{Mod}(T \wedge \bigvee_{n < \omega} \varphi_n(\alpha(0), \dots, \alpha(2n-1)))$. Thus $\text{Mod}(\psi) = \mathfrak{D}P \in \mathfrak{D}\Gamma$. \square

Corollary 6.5. *Let $\langle \varphi_n : n < \omega \rangle$ be a sequence of second-order formulas of uniformly bounded complexity, i.e. there is an $m < \omega$ such that every φ_n has complexity at most Σ_m^1 . If ψ is the formula obtained by applying an open game quantifier to $\langle \varphi_n : n < \omega \rangle$, then $\text{Mod}(\psi)$ has complexity at most $\mathfrak{D}\Sigma_m^1$.*

We need the following notation from [31]:

Definition 6.6. Suppose Γ is a pointclass and $P \in \Gamma$, $P \subseteq \mathcal{X} \times \omega^\omega$. We denote by $\exists^1 P$ the set $\{x \in \mathcal{X} : (\exists^1 y)P(x, y)\}$. We define the pointclass $\exists^1 \Gamma$ to be $\{\exists^1 P : P \in \Gamma\}$. The definitions of $\forall^1 P$ and $\forall^1 \Gamma$ are analogous.

Using the notation in the previous definition, we can use the expression $\exists^1 \Gamma \subseteq \Gamma$ to denote that Γ is closed under second-order existential quantification, e.g. $\exists^1 \Sigma_2^1 \subseteq \Sigma_2^1$.

Under further assumptions, classes of the form $\mathfrak{D}\Sigma_m^1$ are well understood. Moschovakis [31] proves:

Lemma 6.7 ([31]). *Suppose Γ is an adequate pointclass and $\text{Det}(\Gamma)$ holds. If $\exists^1 \Gamma \subseteq \Gamma$, then $\mathfrak{D}\Gamma = \forall^1 \Gamma$.*

With further work, the following result is obtained:

Lemma 6.8 ([31]). *If PD holds, then*

$$\mathfrak{D}\Sigma_1^0 = \Pi_1^1, \quad \mathfrak{D}\Pi_1^1 = \Sigma_2^1, \quad \mathfrak{D}\Sigma_2^1 = \Pi_3^1, \dots$$

Thus, assuming PD, the open and closed game quantifiers do not lead us out of the projective sets. Moreover, since the class of σ -projective sets is an adequate pointclass that is closed under second-order existential and universal quantifications by definition, we have:

Corollary 6.9. *If σ -projective determinacy holds, then $\mathfrak{D}A$ is σ -projective for any σ -projective A .*

In particular, we can also obtain a ZFC result on isomorphism relations since Borel determinacy holds in ZFC:

Theorem 6.10. *If a sentence σ consists of an open game quantifier followed by a sequence of $L_{\omega_1, \omega}$ formulas, then $\text{Mod}(\sigma)$ is $\mathbf{\Pi}_1^1$, so the associated equivalence relation is $\mathbf{\Sigma}_1^1$. Thus if the relation \cong_σ is thin, there are $\leq \aleph_1$ equivalence classes.*

Proof. In this situation $\text{Mod}(\sigma)$ is obtained by applying \mathfrak{D} to a Borel set, and hence is $\mathbf{\Pi}_1^1$; it follows that \cong_σ is $\mathbf{\Sigma}_1^1$ by Lemma 2.6. The result on the number of equivalence classes then follows by Burgess's dichotomy theorem (Theorem 5.3). \square

Theorem 6.11. *In the usual model for PFA, if \mathcal{L} is the closure of second-order logic under open and closed game quantifiers and σ is an \mathcal{L} -sentence, then the associated isomorphism relation \cong_σ has either a perfect set of pairwise non-isomorphic elements or $\leq \aleph_1$ of them.*

Proof. The relation \cong_σ is in $L(\mathbb{R})$, indeed it is σ -projective, so the result follows from Foreman-Magidor (Theorem 4.1). \square

We can actually do better than this. See Theorem 7.1 below to see that we can get the conclusion of Theorem 6.11 from a model which only assumes that there are infinitely many Woodin cardinals.

For one application of the closed game quantifier to a first-order formula, we get that the complexity of the equivalence relation is $\mathbf{\Delta}_2^1$, assuming $\text{Det}(\mathbf{\Pi}_1^1)$. We can now prove:

Theorem 6.12. *If $\text{Det}(\mathbf{\Pi}_1^1)$ then for a thin equivalence relation defined by an $L_{\omega_1, \omega}$ formula preceded by a closed game quantifier, the number of equivalence classes is $\leq \aleph_1$.*

Proof. By [18, 11.5, 27.13], $\text{Det}(\mathbf{\Pi}_1^1)$ implies $\aleph_1^{L[a]}$ is countable for all reals a . By [13] that implies $\mathbf{\Pi}_2^1$ thin equivalence relations have $\leq \aleph_1$ equivalence classes. \square

As noted earlier, by [23], $\text{Det}(\mathbf{\Pi}_1^1)$ follows from the existence of a measurable cardinal.

6.2. Partially Ordered Quantifiers. Partially ordered (first-order) quantifiers were introduced by Henkin [15], who noted that the usual first-order logic could not adequately express a situation in which for all x there exists a y , and for all z there is a w such that $R(x, y, z, w)$, but the y does not depend on z and the w does not depend on x . Since then, a number of authors have investigated such quantifiers and even infinitary versions of them [48], [7], [43], etc. A comprehensive treatment was given in A. K. Swett's doctoral dissertation [43]. Swett assumed the partial order was well-founded. He gave an example to show that the semantics could be ill-defined if one dropped that assumption. It was shown by Walkoe [48] and

Enderton [7] that sentences with finite partially ordered quantifiers are equivalent to existential second-order sentences, and hence by Corollary 5.24 we obtain:

Theorem 6.13. *Assuming MA_{ω_1} , thin equivalence relations in first-order logic augmented with finite partially ordered quantifiers have $\leq \aleph_1$ equivalence classes.*

Corollary 6.14. *Assuming MA_{ω_1} , the number of countable models of a countable theory in first-order logic augmented with finite partially ordered quantifiers is $\leq \aleph_1$ or $= 2^{\aleph_0}$.*

We have not investigated the infinite partially ordered cases, except for the open and closed game quantifiers.

7. σ -PROJECTIVE EQUIVALENCE RELATIONS

The set of reals coding the countable models of a second-order *sentence* can be arbitrarily high in the projective hierarchy. The set of reals coding the countable models of a second-order *theory* can thus be the intersection of countably many projective sets, and these projective sets may have unbounded projective complexity. In order to extend our work to include second-order theories we therefore need information about the set of equivalence classes of a σ -projective equivalence relation. In this section we obtain results along these lines. For this section we temporarily set aside our intended applications in second-order logic and focus only on the results we need in Descriptive Set Theory. As such, we use the terminology of Descriptive Set Theory throughout; the only appearance of second-order logic in this section is at the very end, in Theorem 7.12. In particular, we emphasize that in this section “definable” is meant in the sense of Descriptive Set Theory.

At this point we should warn the reader that for this section not only do we assume familiarity with a substantial amount of Set Theory as we have implicitly so far but towards the end of the section we also assume some knowledge of *Inner Model Theory*. For the set theoretic background, i.e., up to and including the proof of Theorem 7.1 modulo Lemmas 7.4 and 7.7, standard textbooks such as [18, 17, 35, 22] suffice. For the rest of this section we suppose familiarity with the basics of Inner Model Theory as for example covered in the first five sections of [42].

Foreman and Magidor [9] used a supercompact cardinal to produce a model of $\neg CH$ in which every equivalence relation in $L(\mathbb{R})$ on the power set of \mathbb{R} has $\leq \aleph_1$ or a perfect set of inequivalent elements. Their model is the standard model for PFA but they could have instead used the forcing we apply in the proof of Theorem 7.1. Then, by results of Woodin, they could have reduced the large cardinal hypothesis in their result from a supercompact cardinal to a proper class of Woodin cardinals or even to infinitely many Woodin cardinals with a measurable cardinal above them all. There are two main consequences of supercompactness that Foreman-Magidor used in their proof. First, they used generic absoluteness of $L(\mathbb{R})$ for forcings below the supercompact cardinal. Second, they used that sets in $L(\mathbb{R})$ are weakly homogeneously Suslin at the supercompact cardinal (see for example [35, Problem 13.3] or [18, Section 32] for the definition of κ -weakly homogeneously Suslin). But for both of these applications, supercompactness is not necessary. Suppose κ is an

inaccessible cardinal and above κ there are infinitely many Woodin cardinals with a measurable cardinal above them all. Then the Foreman-Magidor result follows using the forcing we use in the proof of Theorem 7.1 by appealing to the following results of Woodin. First, generic absoluteness of $L(\mathbb{R})$ for forcings of size $\leq \kappa$ holds by Theorem [49, Theorem 2.31], see Theorem 6.1 and Remark 6.6 in [41] for a proof. Second, every set of reals in $L(\mathbb{R})$ is κ -weakly homogeneously Suslin by [49, Theorem 2.13].

For our application to variants of Morley's Theorem we are interested in equivalence relations that are much simpler than arbitrary equivalence relations in $L(\mathbb{R})$. If we focus on σ -projective equivalence relations we can reduce the large cardinal hypothesis to infinitely many Woodin cardinals (and even below, see the discussion below and Theorem 7.11). This leads to a significantly lower large cardinal hypothesis as the mice we consider do not have inner models with infinitely many Woodin cardinals. Defining mice is beyond the scope of this paper but we refer the interested reader to [35] or [42].

Theorem 7.1. *Suppose there are infinitely many Woodin cardinals. Then there is a model of $\neg\text{CH}$ in which every σ -projective equivalence relation on the power set of \mathbb{R} has $\leq \aleph_1$ or a perfect set of inequivalent elements.*

We recall the definition of Woodin cardinals for the reader's convenience. In contrast to other large cardinal notions such as measurable cardinals or supercompact cardinals, Woodin cardinals are not critical points of strong elementary embeddings. They are limits of such critical points in the following sense.

Definition 7.2. (1) Let $\kappa < \delta$ be ordinals and $A \subseteq V_\delta$. Then κ is called *A-reflecting in δ* if and only if for all $\eta < \delta$ there is an elementary embedding $i: V \rightarrow M$ with critical point κ such that $i(\kappa) > \eta$ and

$$i(A) \cap V_\eta = A \cap V_\eta.$$

(2) A cardinal δ is a *Woodin cardinal* if and only if for all $A \subseteq \delta$ there is some $\kappa < \delta$ that is *A-reflecting in δ* .

Remark. Note that the hypothesis of Theorem 7.1 is not optimal. We will state a sharper theorem below but decided to start with this version as the statement of Theorem 7.1 does not require any knowledge of Inner Model Theory.

The proof of Theorem 7.1 uses the two following inner model theoretic lemmas that we prove below. Again, the large cardinal hypothesis in the statement of the lemmas is not optimal and we will improve it before proving the lemmas. We need one more definition before we can state the lemmas.

Recall that projective sets of reals can be defined in second-order arithmetic with parameters by Σ_n and Π_n formulas, see for example [18, Section 12, The Definability Context]. We extend this hierarchy to formulas in the language $\mathcal{L}_{\omega_1, \omega}$ defining σ -projective sets. As in the projective case, we say a formula φ is $\Sigma_{\alpha+1}$ for some $\alpha < \omega_1$ if it is of the form $(\exists x)\psi$ for some Π_α formula ψ . For limit ordinals $\lambda < \omega_1$ we say a formula φ is Σ_λ if and only if it is of the form $\bigvee_{k < \omega} \psi_k$ for Σ_{λ_k} -formulae ψ_k with $\lambda_k < \lambda$ for all $k < \omega$. Moreover, a formula φ is Π_α for some $\alpha < \omega_1$ if it is $\mathcal{L}_{\omega_1, \omega}$ -equivalent to $\neg\psi$ for some Σ_α formula ψ .

Definition 7.3. A formula $\varphi(v)$ in the language $\mathcal{L}_{\omega_1, \omega}$ is called σ -projective if and only if it is Σ_α for some $\alpha < \omega_1$.

Remark. Note that a set of reals A is σ -projective if and only if there is a Σ_α -formula φ for some $\alpha < \omega_1$ and a parameter $z \in {}^\omega\omega$ such that A is $\Sigma_\alpha(z)$ definable in second-order arithmetic in z , i.e.,

$$A = \{x \in {}^\omega\omega : \mathcal{A}^2(z) \models \varphi(x)\},$$

where $\mathcal{A}^2(z)$ is the two-sorted structure

$$(\omega, {}^\omega\omega, ap, +, \times, exp, <, 0, 1, z)$$

as in [18, Section 12, The Definability Context]. Here $ap: {}^\omega\omega \rightarrow \omega$ denotes the binary operation of *application*, i.e., $ap(x, n) = x(n)$.

Lemma 7.4. *Let M be a model of ZFC with a sequence of Woodin cardinals $(\delta_i : i < \omega)$. Let \mathbb{P} be a partial order in M with $|\mathbb{P}| < \delta_0$ and let g be \mathbb{P} -generic over M . Then for every σ -projective formula $\varphi(v)$ and every $x \in ({}^\omega\omega)^M$,*

$$(\mathcal{A}^2(x))^M \models \varphi(x) \text{ if and only if } (\mathcal{A}^2(x))^{M[g]} \models \varphi(x).$$

It is known that assuming large cardinals weaker than the existence of infinitely many Woodin cardinals (more precisely only the existence of certain inner models with finitely many Woodin cardinals) σ -projective sets are determined, see for example [1]. We argue here that they are $<\eta$ -universally Baire for some ordinal η , as for example defined in [17, Definition 32.21] or [35, Definition 8.6]. We recall the definition here.

Definition 7.5. Let (S, T) be trees on $\omega \times \kappa$ for some ordinal κ and let η be an ordinal. We say (S, T) is η -absolutely complementing if and only if

$$p[S] = {}^\omega\omega \setminus p[T]$$

in every $\text{Col}(\omega, \eta)$ -generic extension of V .

Definition 7.6 (Feng-Magidor-Woodin, [8]). Let A be a set of reals.

- (1) We say A is $<\eta$ -universally Baire ($<\eta$ -uB) if for every ordinal $\nu < \eta$, there are ν -absolutely complementing trees (S, T) with $p[S] = A$.
- (2) We say A is *universally Baire* (uB) if it is $<\eta$ -universally Baire for every ordinal η .

Lemma 7.7. *Let M be a model of ZFC with a sequence of Woodin cardinals $(\delta_i : i < \omega)$. Then all σ -projective sets in M are $<\delta_0$ -universally Baire.*

We now prove Theorem 7.1 modulo Lemmas 7.4 and 7.7.

Proof of Theorem 7.1. Let M be a model of ZFC with a sequence of Woodin cardinals $(\delta_i : i < \omega)$ and let κ be the least inaccessible cardinal in M . In particular, $\kappa < \delta_0$. We consider a generic extension $M[G]$ of M via the following forcing: Consider the countable support iteration \mathbb{P}_κ of length κ of the following partial orders

$$\{\dot{\mathbb{Q}}_\alpha : \alpha < \kappa\}.$$

At an even stage $\alpha < \kappa$ that is not a limit stage of cofinality ω , let $\dot{\mathbb{Q}}_\alpha$ be the usual countably closed collapse of the continuum (of the current stage of the iteration) to ω_1 . At limit stages $\alpha < \kappa$ of cofinality ω , let $\dot{\mathbb{Q}}_\alpha$ be the trivial forcing. At an odd stage $\beta < \kappa$ let $\dot{\mathbb{Q}}_\beta$ add β Cohen reals. Note that each individual forcing $\dot{\mathbb{Q}}_\alpha$ and hence the whole iteration \mathbb{P}_κ is proper and in particular preserves \aleph_1 . The final model $M[G]$ satisfies

$$2^{\aleph_0} = \aleph_2$$

but CH holds at cofinally many initial segments $M[G \upharpoonright \alpha]$ of the iteration. Here $G \upharpoonright \alpha$ denotes the canonical restriction of the generic G to the initial segment \mathbb{P}_α up to α of the iteration.

We claim that in $M[G]$ every σ -projective equivalence relation on the power set of \mathbb{R} has $\leq \aleph_1$ or a perfect set of inequivalent elements. Let E be a σ -projective equivalence relation on the power set of \mathbb{R} and suppose E is thin, i.e., it does not have a perfect set of inequivalent elements. We will argue that it has $\leq \aleph_1$ equivalence classes. Let φ be a σ -projective formula defining E in $M[G]$, i.e.,

$$E = \{(x, y) \in \mathbb{R}^{M[G]} : (\mathcal{A}^2(x, y))^{M[G]} \models \varphi(x, y)\}.$$

Let $M[G \upharpoonright \alpha]$ be an initial segment of the iteration such that CH holds in $M[G \upharpoonright \alpha]$. Let E_α be the equivalence relation defined by

$$E_\alpha = \{(x, y) \in \mathbb{R}^{M[G \upharpoonright \alpha]} : (\mathcal{A}^2(x, y))^{M[G \upharpoonright \alpha]} \models \varphi(x, y)\}.$$

Claim 1. *E_α is thin in $M[G \upharpoonright \alpha]$.*

Proof. Suppose toward a contradiction that E_α is not thin, i.e., it has a perfect set of inequivalent elements. Then the reals representing these classes are still elements of $M[G]$. As the size of the forcing \mathcal{P}_κ is small, $M[G \upharpoonright \alpha]$ still has the sequence of Woodin cardinals $(\delta_i : i < \omega)$. So by Lemma 7.4 the reals representing the perfect set of inequivalent elements under E_α are also pairwise E -inequivalent as E is the equivalence relation given by φ . This implies that E is also not thin, contradicting our choice of E . \square

So E_α has at most \aleph_1 equivalence classes in $M[G \upharpoonright \alpha]$. The following claim finishes the proof of Theorem 7.1 (modulo Lemmas 7.4 and 7.7).

Claim 2. *Every real added by the forcing iteration from $M[G \upharpoonright \alpha]$ to $M[G]$ is E -equivalent to a real in $M[G \upharpoonright \alpha]$. In particular, E also has at most \aleph_1 equivalence classes in $M[G]$.*

Proof. As in the previous claim, $M[G \upharpoonright \alpha]$ still has the sequence of Woodin cardinals $(\delta_i : i < \omega)$. So by Lemma 7.7, E_α is $< \delta_0$ -universally Baire. This allows us to canonically extend E_α to equivalence relations in generic extensions for partial orders of size $< \delta_0$. Here the canonical extension of E_α to $M[G]$ is E .

Note that universal Baireness is all that is used of the hypothesis that the equivalence relation is weakly homogeneously Suslin in the proof of [9, Theorem 3.4]. Hence, we can apply this theorem to E_α and obtain that since E_α does not have a perfect set of inequivalent reals in $M[G]$, no real added by the forcing iteration from $M[G \upharpoonright \alpha]$ to $M[G]$ is E -equivalent to a real in $M[G \upharpoonright \alpha]$. \square

□

Now we turn to the proofs of Lemmas 7.4 and 7.7. We in fact prove strengthenings of them here that will also lead to a strengthening of Theorem 7.1. Our aim is to reduce the large cardinal hypothesis in these results. For notational simplicity we focus on the case of countable intersections of projective sets instead of arbitrary σ -projective sets but a relatively straightforward generalization of the ideas described below yields similar results for all σ -projective sets. Recall that the set of all countable models of a second-order S -theory for a countable signature S is a countable intersection of projective sets.

Before we can state the strengthenings of Lemmas 7.4 and 7.7 we need the following definition that introduces the large cardinal hypothesis we want to work from. Note that here we pass from large cardinals to inner models with large cardinals – a central theme in Inner Model Theory.

Definition 7.8. We say M is a *ladder premouse above η* if and only if M is a proper class fine structural premouse as for example in [42] with cardinals $\eta < \delta_0 < \delta_1 < \dots$ such that

- (1) η is the second inaccessible cardinal in M ,
- (2) for each $n < \omega$,

$$M_n^\#(M|\delta_n) \models \text{“}\delta_n \text{ is Woodin”},$$

and

- (3) for each $n < \omega$,

$$M_n^\#(M|\delta_n) \trianglelefteq M.$$

Here $M_n^\#(M|\delta_n)$ denotes the least ω_1 -iterable premouse constructed above $M|\delta_n$ that is not n -small above δ_n , see for example [42, p. 1660], [40], or the introduction of [32] for the definition and some properties of this model. It is beyond the scope of this paper to formally introduce $M_n^\#(M|\delta_n)$ but we would like to mention that the premouse $M_n^\#(M|\delta_n)$ is a model that has n Woodin cardinals above δ_n . Nevertheless these Woodin cardinals will not remain Woodin in the full model M . So the minimal ladder premouse (if it exists) does not have Woodin cardinals. It has inner models with n Woodin cardinals for every natural number n but it does not have an inner model with infinitely many Woodin cardinals. This in particular explains why the *consistency strength* of the existence of a ladder mouse is below the existence of a model with infinitely many Woodin cardinals as assumed in Theorem 7.1, even though there is no canonical large cardinal hypothesis below the existence of infinitely many Woodin cardinals that implies the existence of a ladder mouse. Note that the existence of a ladder premouse easily follows from the hypothesis of Theorem 7.1, i.e., in a model M of ZFC with a sequence of Woodin cardinals $(\delta_i : i < \omega)$.

Remark. A similar notion of ladder premouse figures prominently in the computation of the sets of reals of initial segments of $L(\mathbb{R})$ as the sets of reals of mice, see for example [34] or the discussion in [44, Section 8.4].

Now we can state Lemmas 7.4 and 7.7 in terms of ladder premice. Note that if M is a ladder mouse above η , then the reals $({}^\omega\omega)^M$ of M are clearly all contained in $M|\eta$, the model up to level η .

Lemma 7.9. *Let M be a ladder premouse above η for some ordinal η . Let \mathbb{P} be a partial order in M with $|\mathbb{P}| < \eta$ and let g be \mathbb{P} -generic over M . Then for every Π_ω formula $\varphi(v)$ and every $x \in ({}^\omega\omega)^M$,*

$$(\mathcal{A}^2(x))^M \models \varphi(x) \text{ if and only if } (\mathcal{A}^2(x))^{M[g]} \models \varphi(x).$$

Lemma 7.10. *Let M be a ladder premouse above η for some ordinal η witnessed by cardinals $\delta_0 < \delta_1 < \dots$. Then all sets in M that are obtained as a countable intersection of projective sets are $<\eta$ -universally Baire.*

For the proofs of Lemmas 7.9 and 7.10 we require that the reader is familiar with the basics of Inner Model Theory as for example covered in the first five sections of [42].

Proof of Lemma 7.9. Let M be a ladder premouse above η for some ordinal η . Then M is clearly closed under the operations $z \mapsto M_n^\#(z)$ for all reals z and natural numbers n . As the size of the forcing \mathbb{P} is below η , it is easy to see that any \mathbb{P} -generic extension $M[G]$ of M is also closed under the operations $z \mapsto M_n^\#(z)$ for all reals z and natural numbers n .

Therefore, the following claim finishes the proof of Lemma 7.9.

Claim 1. *Let φ be a Σ_ω formula, say $\varphi = \bigvee_{k < \omega} \psi_k$, and let N be any model that is closed under the operations $z \mapsto M_n^\#(z)$ for all reals z and natural numbers n . Then for every $x \in ({}^\omega\omega)^N \cap ({}^\omega\omega)^V$,*

$$(\mathcal{A}^2(x))^N \models \varphi(x) \text{ if and only if } (\mathcal{A}^2(x))^V \models \varphi(x).$$

Proof. We start with the right-to-left implication. Let $x \in ({}^\omega\omega)^N \cap ({}^\omega\omega)^V$ and suppose

$$(\mathcal{A}^2(x))^V \models \varphi(x).$$

That means $(\mathcal{A}^2(x))^V \models \psi_k(x)$ for some $k < \omega$. Suppose for notational simplicity that ψ_k is a Σ_k formula. Then, by correctness¹ of the inner model $M_k^\#(x)$,

$$(\mathcal{A}^2(x))^{M_k^\#(x)} \models \psi_k(x).$$

By our assumption, N is closed under the operation $z \mapsto M_k^\#(z)$ for all reals $z \in ({}^\omega\omega)^N$. In particular, $(M_k^\#(x))^N = (M_k^\#(x))^V$ and $(\mathcal{A}^2(x))^{(M_k^\#(x))^N} \models \psi_k(x)$. Therefore, by correctness of $(M_k^\#(x))^N$ in N ,

$$(\mathcal{A}^2(x))^N \models \psi_k(x).$$

Hence, $(\mathcal{A}^2(x))^N \models \varphi(x)$, as desired. The left-to-right implication follows by the same argument. \square

\square

¹This result is due to Woodin, see for example [32, Lemma 1.17] for the precise statement and a proof.

Proof of Lemma 7.10. Let $A = \bigcap_{k < \omega} A_k$ be a countable intersection of projective sets A_k , $k < \omega$. Suppose for notational simplicity that each A_k is a Σ^1_k set. Then by the usual argument for universal Baireness (or even homogeneity) from Woodin cardinals, see for example [35, Problem 13.4],

$$M_k^\#(M|\delta_k) \models A_k \text{ is } <\delta_k\text{-universally Baire.}$$

In particular, A_k is $<\eta$ -universally Baire in $M_k^\#(M|\delta_k)$. Let $\nu < \eta$ and let (S_k^ν, T_k^ν) be a pair of ν -absolutely complementing trees with $p[S_k^\nu] = A_k$ in $M_k^\#(M|\delta_k)$ and in M for each $k < \omega$. By combining the trees (S_k^ν, T_k^ν) we can easily obtain a pair of ν -absolutely complementing trees (S^ν, T^ν) in M with $p[S] = A$. Therefore, A is $<\eta$ -universally Baire in M . \square

The proof of Theorem 7.1 did not use any large cardinal hypothesis beyond an inaccessible cardinal except for the applications of Lemmas 7.4 and 7.7. So by exchanging these with Lemmas 7.9 and 7.10 we obtain the following theorem.

Theorem 7.11. *Let M be a ladder premouse above η for some ordinal η . Then there is a model of \neg CH in which every equivalence relation on the power set of \mathbb{R} that is obtained as a countable intersection of projective sets has $\leq \aleph_1$ or a perfect set of inequivalent elements.*

Finally, we apply the results of this section to second-order logic again.

Theorem 7.12. *If there are infinitely many Woodin cardinals (or there exists a ladder premouse above some η), then there is a model of set theory in which the Absolute Morley Theorem holds for second-order theories in countable signatures.*

Proof. If T is a second-order theory in a countable signature then the relation \cong_T is σ -projective (in fact, it is obtained as an intersection of countably many projective sets), so this follows from Theorem 7.1 (or Theorem 7.11). \square

REFERENCES

1. J. P. Aguilera, S. Müller, and P. Schlicht, *Long games and σ -projective sets*, Annals of Pure and Applied Logic **172** (2021), 102939.
2. T. Bartoszynski and H. Judah, *Set theory, on the structure of the real line*, A. K. Peters Ltd., Wellesley, MA, 1995.
3. M. Bekkali, *Topics in set theory: Lebesgue measurability, large cardinals, forcing axioms, and ρ -functions; notes on lectures by S. Todorcevic*, Springer-Verlag, Berlin, 1991.
4. A. Blass, *Combinatorial cardinal characteristics of the continuum*, Handbook of set theory, vol. 1,2,3, Springer, Dordrecht, 2010, pp. 395–489.
5. J. Burgess, *Infinitary languages and descriptive set theory*, Ph.D. thesis, University of California at Berkeley, Berkeley, California, 1974.
6. M. Davis, *Infinite games of perfect information*, Advances in Game Theory (M. Dresher, L.S. Shapley, and A.W. Tucker, eds.), Annals of Mathematics Studies, vol. 52, Princeton University Press, 1964, pp. 85–101.
7. H.B. Enderton, *A mathematical introduction to logic*, Academic Press, New York-London, 1972.
8. Q. Feng, M. Magidor, and H. Woodin, *Universally Baire Sets of Reals*, Set Theory of the Continuum (New York, NY) (H. Judah, W. Just, and H. Woodin, eds.), Springer US, 1992, pp. 203–242.

9. M. Foreman and M. Magidor, *Large cardinals and definable counterexamples to the continuum hypothesis*, Annals of Pure and Applied Logic **76** (1995), no. 1, 47–97.
10. S. Gao, *Invariant descriptive set theory*, Chapman and Hall/CRC, New York, 2008.
11. L. Harrington, *Long projective wellorderings*, Ann. Math. Logic **12** (1977), 1–24.
12. L. Harrington and R. Sami, *Equivalence relations, projective and beyond*, Logic Colloquium'78, North-Holland, Amsterdam, 1979, pp. 247–264.
13. L. Harrington and S. Shelah, *Counting equivalence classes for co- κ -souslin equivalence relations*, Logic Colloquium'80, Prague, 1980, North-Holland, Amsterdam, 1982, pp. 147–152.
14. B. Hart, S. Starchenko, and M. Valeriote, *Vaught's conjecture for varieties*, Trans. Amer. Math. Soc. **342** (1994), no. 1, 173–196.
15. L. Henkin, *Some remarks on infinitely long formulas*, Infnitistic Methods (Proc. Sympos. Foundations of Math., Warsaw, 1959, Pergamon, Oxford; Państwowe Wydawnictwo Naukowe, Warsaw, 1961, pp. 167–183.
16. G. Hjorth, *Some applications of coarse inner model theory*, J. Symb. Logic **62** (1997), 337–365.
17. T. J. Jech, *Set theory*, Springer Monographs in Mathematics, Springer, 2003.
18. A. Kanamori, *The higher infinite*, 2nd ed., Springer Monographs in Mathematics, Springer, 2009.
19. A.S. Kechris, *AD and projective ordinals*, Cabal seminar 76-77 (Berlin) (A.S. Kechris et. al., ed.), Lecture notes in mathematics, vol. 689, Springer-Verlag, 1978, pp. 91–132.
20. ———, *Classical descriptive set theory*, Springer-Verlag, New York, 1995.
21. K. Kunen, *Set theory: An introduction to independence proofs*, Studies in logic and the foundations of mathematics, vol. 102, Elsevier, Amsterdam, 1980.
22. ———, *Set theory*, College Publications, London, 2011.
23. D.A. Martin, *Measurable cardinals and analytic games*, Fund. Math. **66** (1970), 287–291.
24. ———, *Borel determinacy*, Ann. Math. **102** (1973), 363–371.
25. ———, *Wadge degrees and projective ordinals*, The Cabal Seminar, Volume II (A.S. Kechris, B. Löwe, and J.R. Steel, eds.), Lect. Notes Logic, no. 37, Cambridge University Press, 2011, pp. 484–508.
26. D.A. Martin and R.M. Solovay, *Internal Cohen extensions*, Ann. Math. Logic **2** (1970), 143–178.
27. D.A. Martin and J.R. Steel, *Iteration trees*, J. Amer. Math. Soc. **7** (1994), 1–73.
28. L. Mayer, *Vaught's conjecture for o-minimal theories*, J. Symb. Logic **53** (1988), no. 1, 146–159.
29. M. Morley, *The number of countable models*, J. Symb. Logic **35** (1970), 14–18.
30. Y.N. Moschovakis, *The game quantifier*, Proc. Amer. Math. Soc. **31** (1972), 245–250.
31. ———, *Descriptive set theory*, Mathematical Surveys and Monographs, vol. 155, American Mathematical Society, Providence, R.I., 2009.
32. S. Müller, R. Schindler, and W. H. Woodin, *Mice with finitely many Woodin cardinals from optimal determinacy hypotheses*, Journal of Mathematical Logic **20** (2020).
33. I. Neeman, *Determinacy in $L(\mathbb{R})$* , Handbook of Set Theory (M. Foreman and A. Kanamori, eds.), Springer, 2010, pp. 1877–1950.
34. M. Rudominer, *Mouse sets*, Annals of Pure and Applied Logic **87** (1997), no. 1, 1–100.
35. R. Schindler, *Set theory*, Universitext, Springer-Verlag, 2014.
36. P. Schlicht, *Thin equivalence relations and inner models*, Ann. Pure and Appl. Logic **165** (2014), 1577–1625.
37. S. Shelah, L. Harrington, and M. Makkai, *A proof of Vaught's conjecture for ω -stable theories*, Israel J. Math. **49** (1984), 259–280.
38. J.H. Silver, *Counting the number of equivalence classes of Borel and coanalytic equivalence relations*, Ann. Math. Logic **18** (1980), 1–28.
39. J. Steel, *On Vaught's conjecture*, Cabal seminar 76-77 (Berlin) (A.S. Kechris et. al., ed.), Lecture notes in mathematics, vol. 689, Springer-Verlag, 1978, pp. 193–208.
40. J. R. Steel, *Projectively well-ordered inner models*, Annals of Pure and Applied Logic **74** (1995), 77–104.
41. ———, *The derived model theorem*, Logic Colloquium 2006 (S. B. Cooper, H. Geuvers, A. Pillay, and J. Väänänen, eds.), Lecture Notes in Logic, Cambridge University Press, 2009, p. 280–327.
42. ———, *An Outline of Inner Model Theory*, Handbook of Set Theory (M. Foreman and A. Kanamori, eds.), Springer, 2010.

43. A.K. Swett, *Herbrand's theorem in infinitary logic*, Ph.D. thesis, University of Toronto, 1973.
44. S. Uhlenbrock (now Müller), *Pure and Hybrid Mice with Finitely Many Woodin Cardinals from Levels of Determinacy*, Ph.D. thesis, WWU Münster, 2016.
45. J. Väänänen, *Second-order and higher-order logic*, The Stanford Encyclopedia of Philosophy (E. Zalta, ed.), The Metaphysics Research Lab Center for the Study of Language and Information at Stanford University, Stanford, 2020, URL: <https://plato.stanford.edu/archives/fall2020/entries/logic-higher-order/>.
46. R. Vaught, *Denumerable models of complete theories*, Proc. Sympos. Foundations of Mathematics, Infinitistic Methods (Warsaw), Pergamon Press, 1961, pp. 303–321.
47. B. Velickovic, *Forcing axioms and stationary sets*, Adv. Math. **94** (1992), 256–284.
48. W.J. Walkoe, Jr., *Finite partially-ordered quantification*, J. Symb. Logic **35** (1970), 535–555.
49. W. H. Woodin, *The Axiom of Determinacy, Forcing Axioms, and the Nonstationary Ideal*, De Gruyter series in logic and its applications, vol. 1, De Gruyter, 2010.
50. W.H. Woodin, *Supercompact cardinals, sets of reals, and weakly homogeneous trees*, Proc. Nat. Acad. Sci. U.S.A. **85** (1988), no. 18, 6587–6591.

(C. J. Eagle) UNIVERSITY OF VICTORIA, DEPARTMENT OF MATHEMATICS AND STATISTICS, PO BOX 1700 STN CSC, VICTORIA, BRITISH COLUMBIA, CANADA, V8W 2Y2

Email address: eaglec@uvic.ca

URL: <http://www.math.uvic.ca/~eaglec>

(C. Hamel) UNIVERSITY OF TORONTO, DEPARTMENT OF MATHEMATICS, 40 ST. GEORGE ST., TORONTO, ONTARIO, CANADA M5S 2E4

Email address: chamel@math.toronto.edu

(S. Müller) INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, TU WIEN, WIEDNER HAUPTSTRASSE 8-10/104, 1040 WIEN, AUSTRIA, AND INSTITUT FÜR MATHEMATIK, UNIVERSITÄT WIEN, KOLINGASSE 14-16, 1090 WIEN, AUSTRIA

Email address: sandra.mueller@tuwien.ac.at

URL: <http://www.logic.univie.ac.at/~smueller/>

(F. D. Tall) UNIVERSITY OF TORONTO, DEPARTMENT OF MATHEMATICS, 40 ST. GEORGE ST., TORONTO, ONTARIO, CANADA M5S 2E4

Email address: f.tall@utoronto.ca

URL: <http://www.math.toronto.edu/tall/>