

Double Glueing over Free Exponential: with Measure Theoretic Applications

Masahiro HAMANO

*Miin Wu School of Computing, National Cheng Kung University,
No 1. Academic Road, Tainan City, 70101, TAIWAN*

Abstract

This paper provides a compact method to lift the free exponential construction of Melliès-Tabareau-Tasson over the Hyland-Schalk double glueing for orthogonality categories. A condition "reciprocity of orthogonality" is presented simply enough to lift the free exponential over the double glueing in terms of the orthogonality. Our general method applies to the monoidal category \mathbf{Tsk} of the s-finite transition kernels with countable biproducts. We show (i) \mathbf{Tsk}^{op} has the free exponential, which is shown to be describable in terms of measure theory. (ii) The s-finite transition kernels have an orthogonality between measures and measurable functions in terms of Lebesgue integrals. The orthogonality is reciprocal, hence the free exponential of (i) lifts to the orthogonality category $\mathbf{O}_{\mathcal{I}}(\mathbf{Tsk}^{\text{op}})$, which subsumes Ehrhard et al's probabilistic coherent spaces as the full subcategory of countable measurable spaces. To lift the free exponential, the measure-theoretic uniform convergence theorem commuting Lebesgue integral and limit plays a crucial role. Our measure-theoretic orthogonality is considered as a continuous version of the orthogonality of the probabilistic coherent spaces for linear logic, and in particular provides a two layered decomposition of Crubillé et al's direct free exponential for these spaces.

Contents

1	Lifting Free Exponential of \mathcal{C} to $\mathbf{O}_J(\mathcal{C})$	3
1.1	Melliès-Tabareau-Tasson Free Exponential in a Monoidal \mathcal{C} with Finite Products	3
1.2	Reciprocal Orthogonality in \mathcal{C} and Orthogonality Category $\mathbf{O}_J(\mathcal{C})$	4
1.3	Lifting Free Exponential of \mathcal{C} to $\mathbf{O}_J(\mathcal{C})$	7
2	Monoidal Category \mathbf{Tsk} of s-finite Transition Kernels with Biproducts	9
2.1	Preliminaries from Measure Theory	9
2.2	Transition Kernels	10
2.3	Countable Biproducts in \mathbf{TK}	11
2.4	Monoidal Product and Countable Biproducts in \mathbf{Tsk}	12
3	The Free Exponential in \mathbf{Tsk}^{op}	13
4	The Orthogonality Category $\mathbf{O}_{\mathcal{I}}(\mathbf{Tsk}^{\text{op}})$	18
4.1	Reciprocal Orthogonality between Measures and Measurable Functions in \mathbf{Tsk}^{op}	18
4.2	Push Forward Integral as Reciprocity	18
4.3	Distribution of Monoidal Product over Limit in $\mathbf{O}_{\mathcal{I}}(\mathbf{Tsk}^{\text{op}})$	20
4.4	Characterising Equalsers $\mathbb{A}^{\leq n}$ and Limit $\mathbb{A}^{\leq \infty}$ in $\mathbf{O}_{\mathcal{I}}(\mathbf{Tsk}^{\text{op}})$	22
4.5	Discretisation	27

Email address: hamano@gs.ncku.edu.tw (Masahiro HAMANO)

5	Appendix	29
A.1	Omitted Proofs	29
A.1.1	Proof of Proposition 1.14	29
A.1.2	Proof of Proposition 2.20	29
A.1.3	Demonstration of Eqn.(1) in Def 1.2	30
A.1.4	Proof of Lemma 1.13	30
A.2	Accommodating [6]’s Characterisation $\langle\langle f_1, \dots, f_n \rangle\rangle$ into the Limit L_A of $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{op})$	30
A.3	The Exponential of Pcoh [8]	32

Introduction

This paper is concerned with modelling the exponential connective of linear logic; (i) abstractly for the orthogonality for the double glueing construction and (ii) concretely for a category of s-finite transition kernels using the duality between measures and measurable functions.

The (symmetric) monoidal category provides a minimum categorical counterpart to the tensor connective of linear logic [15]. On top of the monoidality, richer categorical structures are augmented consistently, interpreting other logical components (e.g., the closedness for the linear implication and Barr’s *-autonomy for the duality of linear logic). Understanding categorical properties of the exponential connective ! of linear logic is the difficult part in various works (cf. [3, 19, 21] for surveys) stemming from Seely [25]. Recently Mellès-Tabareau-Tasson [22] formulates the categorical construction to obtain the free commutative comonoid over a symmetric monoidal category with binary products. Their construction interprets the exponential as the limit of the enumerated equalisers for $n!$ -symmetries of n -th powers of the monoidal products between certain rooted objects using the cartesian product.

The most recent application of [22] is done by Crubillé-Ehrhard-Pagani-Tasson [6] to Danos-Ehrhard [8]’s probabilistic coherent spaces, whose exponential is shown to be the free one. The category Pcoh of probabilistic coherent spaces is a probabilistic version of Girard’s denotational semantics Coh of coherent spaces [15]. As the original semantics of linear logic, Coh has the distinctive feature of the linear duality, in terms of the graphical structure on webs, stating that a clique and anti-clique intersect in at most one singleton. Developing the web based method, Ehrhard investigates the linear duality in mathematically richer structures (e.g., Köthe spaces [10]), and his investigation leads Danos-Ehrhard [8] to a probabilistic version of the duality, generalising the web to the non-negative real valued functions on it (i.e., fuzzy web), reminiscent of probability distributions on the web. When each function is identified as a vector enumerating its values, the probabilistic linear duality states that the inner product of the vectors from clique and anti-clique is not greater than 1. Girard earlier addresses this quantitative form of the duality in [16].

The starting point of this paper is our attempt to comprehend [6]’s application to probabilistic semantics more generally, especially free from the web-based method, but extracting the abstract role of the duality more explicitly. We present two new methods respectively to the abstract orthogonality in the category-theory and to probabilistic semantics in the continuous measure-theory: (i) Reciprocal orthogonality enabling us to lift a free exponential to an intricate category (ii) Continuous orthogonality for linear logic between measure and measurable functions.

For (i), we investigate the Hyland-Schalk double glueing construction [19]. Our construction is general enough to yield a probabilistic version of the linear duality. It is well known that the double glueing construction over the category of relations gives rise to the *-autonomy of Coh [3, 19]. This leads to various full completeness theorems, not only for the multiplicative [9] fragment, but also for the multiplicative-additive [2] one. We introduce *reciprocal orthogonality*, characterising the adjunction of the orthogonality. We show that the simple notion of orthogonality interact consistently with equalisers and limit of [22] so to lift the free exponential to the orthogonality categories. Our paper gives a simple insight for the less studied exponential structure inside the double glueing after [19].

For (ii), we investigate the s-finite (i.e., sum of finite) class of the transition kernels [26], which is recently revisited by Staton [27] to analyse factorial commutativity of measure-based denotational semantics for probabilistic programming. The s-finiteness provides a wider extension of the preceding probabilistic semantics using transition kernels, from Kozen’s precursory work [20], then Panangaden’s seminal work of Markov

kernels (a.k.a, stochastic relations) [23, 24] to the recent measure-transformer semantics [4] of finite kernels. Following Staton’s work on the functorial monoidal product, we show another advantage (versus Markov and finiteness): s-finiteness is wide enough to accommodate the free exponential of Melliès-Tabareau-Tasson.

The paper is organised as follows: Section 1 is a categorical study of how to lift a free exponential of \mathcal{C} to a double glueing $\mathbf{O}_J(\mathcal{C})$ with a reciprocal orthogonality. Section 2 is a measure-theoretic study on the s-finite transition kernels \mathbf{Tsk} . Section 3 constructs the free exponential in \mathbf{Tsk}^{op} . Section 4 presents a measure theoretic instance of Section 1 using Section 3, which subsumes the probabilistic coherences as discretisation.

1. Lifting Free Exponential of \mathcal{C} to $\mathbf{O}_J(\mathcal{C})$

This section concerns lifting of Melliès-Tabareau-Tasson free exponential of \mathcal{C} to an orthogonality category $\mathbf{O}_J(\mathcal{C})$ by introducing a notion of reciprocal orthogonality.

1.1. Melliès-Tabareau-Tasson Free Exponential in a Monoidal \mathcal{C} with Finite Products

Definition 1.1 (Melliès-Tabareau-Tasson free exponential [22]). The following four structures uniquely determine the free exponential of a symmetric monoidal category $(\mathcal{C}, \otimes, I)$ with finite cartesian products $\&$:

(\mathbf{E}_A) For any object A of \mathcal{C} and a natural number n , the equaliser exists in \mathcal{C} , denoted by $A^{\leq n}$ with eq_A , for the $n!$ -symmetries on $(A\&I)^{\otimes n}$ as the parallel morphisms:

$$\begin{array}{ccc} A^{\leq n} & \xrightarrow{\text{eq}_A} & (A\&I)^{\otimes n} \xrightarrow{\quad \vdots \quad} (A\&I)^{\otimes n} \\ \uparrow \text{eq}\backslash f & \nearrow f & \downarrow \text{eq} \\ & & (A\&I)^{\otimes n} \xrightarrow{n! \text{ symm.}} (A\&I)^{\otimes n} \end{array}$$

For $f : X \rightarrow (A\&I)^{\otimes n}$ equalising the $n!$ -symmetries, the universal morphisms factoring f is denoted by $\text{eq}\backslash f$ such that $\text{eq} \circ (\text{eq}\backslash f) = f$.

(**distribution of \otimes over \mathbf{E}_A**) The equaliser of \mathbf{E}_A commutes with the tensor product:

$(A^{\leq n} \otimes B, \text{eq} \otimes B)$ becomes the equaliser for the $n!$ -symmetries $\otimes B$ on $(A\&I)^{\otimes n} \otimes B$ for any object B :

$$A^{\leq n} \otimes B \xrightarrow{\text{eq} \otimes B} (A\&I)^{\otimes n} \otimes B \xrightarrow{\quad \vdots \quad} (A\&I)^{\otimes n} \otimes B \\ \text{(n! symm.)} \otimes B$$

(\mathbf{L}_A) For any object A in \mathcal{C} , the following diagram has the the limit $(A^{\leq \infty}, \{p_{\infty, n} : A^{\leq \infty} \rightarrow A^{\leq n}\}_n)$:

$$A^{\leq 0} \xleftarrow{p^{1,0}} A^{\leq 1} \xleftarrow{p^{2,1}} A^{\leq 2} \dots \xleftarrow{p^{n+1,n}} A^{\leq n} \xleftarrow{p^{n+1,n}} A^{\leq n+1} \dots$$

where $p_{n+1, n}$ is the universal morphism guaranteed by \mathbf{E}_A for the composition $((A\&I)^{\otimes n} \otimes p_r) \circ \text{eq}$ equalising $n!$ -symmetries $(A\&I)^{\otimes n}$. p_r is the right projection. See the following:

$$\begin{array}{ccc} A^{\leq n+1} & \xrightarrow{p_{n+1, n}} & A^{\leq n} \\ \downarrow \text{eq} & & \downarrow \text{eq} \\ (A\&I)^{\otimes n+1} & \xrightarrow{(A\&I)^{\otimes n} \otimes p_r} & (A\&I)^{\otimes n} \otimes I \cong (A\&I)^{\otimes n} \end{array}$$

(distribution of \otimes over L_A) The limit of L_A commutes with the monoidal product \otimes :
 $(A^{\leq\infty} \otimes B, \{p_{\infty,n} \otimes B\}_n)$ becomes the limit for the following diagram for any object B :

$$A^{\leq 0} \otimes B \xleftarrow{p_{1,0} \otimes B} \dots \xleftarrow{\dots} A^{\leq n} \otimes B \xleftarrow{p_{n+1,n} \otimes B} A^{\leq n+1} \otimes B \dots$$

The constructions of the equaliser for (E_A) and the limit for (L_A) act not only on objects but also on morphisms functorially preserving the categorical composition.

Definition 1.2 (morphisms $f^{\leq n}$ and $f^{\leq\infty}$). Let $f : A \rightarrow B$.

- There exists the unique morphism for any natural number n by E_A

$$f^{\leq n} := \text{eq}_B \setminus ((f \cdot p_l \& I \cdot p_r)^{\otimes n} \circ \text{eq}_A) : A^{\leq n} \rightarrow B^{\leq n}$$

because the composition $(f \cdot p_l \& I \cdot p_r)^{\otimes n} \circ \text{eq}_A$ equalises the $n!$ -symmetries $(B \& I)^{\otimes n}$. See the following diagram, where p_l and p_r are right and left projections:

$$\begin{array}{ccc} B^{\leq n} & \xrightarrow{\text{eq}_B} & (B \& I)^{\otimes n} \\ \uparrow f^{\leq n} & & \uparrow (f \cdot p_l \& I \cdot p_r)^{\otimes n} \\ A^{\leq n} & \xrightarrow{\text{eq}_A} & (A \& I)^{\otimes n} \end{array}$$

From the universality of eq_B , it holds that $f^{\leq n} = g^{\leq n}$ whenever $(f \cdot p_l \& I \cdot p_r)^{\otimes n} = (g \cdot p_l \& I \cdot p_r)^{\otimes n}$.

- There exists the unique morphism

$$f^{\leq\infty} : A^{\leq\infty} \rightarrow B^{\leq\infty}$$

factoring a cone $\{ A^{\leq\infty} \xrightarrow{p_{\infty,n}} A^{\leq n} \xrightarrow{f^{\leq n}} B^{\leq n} \}_n$ because of the limit $B^{\leq\infty}$ of L_B .
 See the following diagram for $f^{\leq\infty}$ and for the cone.

$$\begin{array}{ccccc} \dots & \longleftarrow & B^{\leq n} & \xleftarrow{p_{n+1,n}} & B^{\leq n+1} & \dots & & B^{\leq\infty} \\ & & \uparrow f^{\leq n} & & \uparrow f^{\leq(n+1)} & & & \uparrow \exists! f^{\leq\infty} \\ \dots & \longleftarrow & A^{\leq n} & \xleftarrow{p_{n+1,n}} & A^{\leq n+1} & \dots & & A^{\leq\infty} \\ & & & \xleftarrow{p_{\infty,n+1}} & & & & \xleftarrow{p_{\infty,n+1}} \end{array}$$

The indexed set indeed is a cone by the equality (1) below, whose demonstration is put in A.1.3.

$$p_{n+1,n} \circ f^{\leq n+1} = f^{\leq n} \circ p_{n+1,n} \tag{1}$$

Obviously the two morphisms in Definition 1.2 are related $f^{\leq n} \circ p_{\infty,n} = p_{\infty,n} \circ f^{\leq\infty}$, which is seen as the limit of (1).

1.2. Reciprocal Orthogonality in \mathcal{C} and Orthogonality Category $\mathbf{O}_J(\mathcal{C})$

Definition 1.3 (orthogonality on \mathcal{C} [19]). An *orthogonality* on a symmetric monoidal category $(\mathcal{C}, \otimes, I)$ is an indexed family of relation \perp_R between the maps $u \in \mathcal{C}(I, R)$ and $x \in \mathcal{C}(R, J)$, where I is the monoidal unit while J is an arbitrary fixed object,

$$I \xrightarrow{u} R \perp_R R \xrightarrow{x} J$$

satisfying the following conditions:

(isomorphism) If $f : R \rightarrow S$ is an isomorphism, then for any $u : I \rightarrow R$ and $x : R \rightarrow J$,

$$u \perp_R x \quad \text{iff} \quad f \circ u \perp_S x \circ f$$

(tensor) Given $u : I \rightarrow R$, $v : I \rightarrow S$, and $h : R \otimes S \rightarrow J$,

$$\begin{aligned} u \perp_R R &\cong R \otimes I \xrightarrow{R \otimes v} R \otimes S \xrightarrow{h} J \quad \text{and} \\ v \perp_S S &\cong I \otimes S \xrightarrow{u \otimes S} R \otimes S \xrightarrow{h} J \quad \text{imply} \quad u \otimes v \perp_{R \otimes S} h. \end{aligned}$$

(identity) For all $u : I \rightarrow R$ and $x : R \rightarrow J$, $u \perp_R x$ implies $\text{Id}_I \perp_I x \circ u$

For $U \subseteq \mathcal{C}(I, R)$, its orthogonal $U^\circ \subseteq \mathcal{C}(R, J)$ is given by

$$U^\circ := \{x : R \rightarrow J \mid \forall u \in U \ u \perp_R x\}$$

This gives a Galois connection so that $U^{\circ\circ} = U^\circ$. The operator $(\)^\circ$ is called the closure operator in the sequel.

Definition 1.4 (reciprocal orthogonality). An orthogonality is *reciprocal* when for every $u : I \rightarrow R$, $x : S \rightarrow J$, and $f : R \rightarrow S$,

$$u \perp_R x \circ f \quad \text{if and only if} \quad f \circ u \perp_S x \quad (2)$$

Definition 1.5. *Precise tensor* is the tensor condition of Definition 1.3 strengthened by replacing “imply” with “iff”

Proposition 1.6 (reciprocity enough for the three conditions of the orthogonality). *The reciprocal orthogonality derives the three conditions (isomorphism), (tensor) and (identity) on Definition 1.4. Moreover (precise tensor) is derived.*

Proof. (isomorphism) $u = f^{-1} \circ f \circ u \perp_R x$ iff $f \circ u \perp_R x \circ f^{-1}$ by reciprocity.

(identity) Similarly but easily by $u = u \circ \text{Id}_I$.

(tensor) Only one premise of the tensor condition implies the conclusion: $u \perp_R h \circ (R \otimes v)$ iff by reciprocity $(R \otimes v) \circ u \perp_R h$, whose left hand is $u \otimes v$.

(precise tensor) $u \otimes v = (R \otimes v) \circ (u \otimes I) = (u \otimes S) \circ (I \otimes v)$, composing $(R \otimes v)$ (resp. $(u \otimes S)$) to the right h in the $\perp_{R \otimes S}$ yields the first (resp. the second) premise of the tensor. \square

The similar condition to (2) in Definition 1.4 is considered in Hyland-Schalk [19] in order to define a certain class of morphisms f , called focused.

Definition 1.7 (focused orthogonality (cf. Example 48 of [19])). An orthogonality is *focused* when it is determined by a subset F of $\mathcal{C}(I, J)$ in the following manner:

$$I \xrightarrow{u} R \perp_R R \xrightarrow{x} J \quad \text{if and only if} \quad x \circ u \in F$$

Proposition 1.8 (reciprocity coincides with focused orthogonality). *An orthogonality is reciprocal if and only if it is focused.*

Proof. (if) The left and the right of (2) are both equivalent to $x \circ f \circ u \in F$.

(only if) Put $F := \{x : I \rightarrow J \mid \text{Id}_I \perp_I x\}$. Then $u \perp_R x$ iff $x \circ u \in F$ by the reciprocity. \square

Definition 1.9 (orthogonality category $\mathbf{O}_J(\mathcal{C})$ [19]). An object of $\mathbf{O}_J(\mathcal{C})$ is a tuple $\mathbb{A} = (A, \mathbb{A}_p, \mathbb{A}_{cp})$ with $\mathbb{A}_p \subseteq \mathcal{C}(I, A)$ and $\mathbb{A}_{cp} \subseteq \mathcal{C}(A, J)$ satisfying;

(mutual orthogonality) $\mathbb{A}_p = (\mathbb{A}_{cp})^\circ$ and $\mathbb{A}_{cp} = (\mathbb{A}_p)^\circ$

Each map from $\mathbb{A} = (A, \mathbb{A}_p, \mathbb{A}_{cp})$ to $\mathbb{B} = (B, \mathbb{B}_p, \mathbb{B}_{cp})$ in $\mathbf{O}_J(\mathcal{C})$ is any \mathcal{C} map $f : A \rightarrow B$ satisfying:

(p) point: $\forall u : I \rightarrow B$ in \mathbb{A}_p , the composition $f \circ u : I \xrightarrow{u} A \xrightarrow{f} B$ belongs to \mathbb{B}_p .

(cp) copoint: $\forall y : B \rightarrow J$ in \mathbb{B}_{cp} , the composition $y \circ f : A \xrightarrow{f} B \xrightarrow{y} J$ belongs to \mathbb{A}_{cp} .

The functor exists $|\cdot| : \mathbf{O}_J(\mathcal{C}) \rightarrow \mathcal{C}$ forgetting the second and the third components of the objects.

Remark 1.10 ($\mathbf{O}_J(\mathcal{C})$ is Hyland Schalk's tight orthogonality category). *The category $\mathbf{O}_J(\mathcal{C})$ is called the tight orthogonality category in Hyland-Schalk [19], whereby it is formulated as a subcategory of the double glueing category over \mathcal{C} (cf. Definition 47 of [19]). Since in the present paper this is the only category concerning with the double glueing construction, we use the simple name.*

Lemma 1.11. *The conditions (p) and (cp) are derivable one another when an orthogonality is reciprocal.*

Proof. By $\mathbb{B}_p = \mathbb{B}_{cp}^\circ$, the condition (p) says $\forall u \in \mathbb{A}_p \forall s \in \mathbb{B}_{cp} f \circ u \perp_B s$. By $\mathbb{A}_{cp} = \mathbb{A}_p^\circ$, the condition (cp) says $\forall s \in \mathbb{B}_{cp} \forall u \in \mathbb{A}_p u \perp_A s \circ f$. The two are equivalent by the reciprocity. \square

By the lemma, when an orthogonality is reciprocal, an alternative definition of the category $\mathbf{O}_J(\mathcal{C})$ is obtained;

Definition 1.12 ($\mathbf{O}_J(\mathcal{C})$ with a reciprocal arthogonality). When an orthogonality on \mathcal{C} is reciprocal, each object of $\mathbf{O}_J(\mathcal{C})$ is represented alternatively by a pair $\mathbb{A} = (A, \mathbb{A}_p)$ satisfying the following instead of the mutual orthogonality:

(double orthogonality): $(\mathbb{A}_p)^{\circ\circ} = \mathbb{A}_p$

Each map between the objects must satisfy the condition (p) only.

A stronger condition suffices in particular when the second components are represented by $\mathbb{A}_p = U^{\circ\circ}$ and $\mathbb{B}_p = V^{\circ\circ}$ with genuine subsets U and V of \mathbb{A}_p and of \mathbb{B}_p respectively:

(\bar{p}) $\forall u \in U$, the composition $f \circ u$ belongs to V .

The sufficiency is because of the monotonicity of the operation $(\)^{\circ\circ}$ and the following lemma.

Lemma 1.13. *For any morphism $f : A \rightarrow B$ and any subset $U \subseteq \mathcal{C}(I, A)$, $f(U^{\circ\circ}) \subseteq f(U)^{\circ\circ}$. (See the proof in A.1.4).*

Both cartesian and monoidal products in \mathcal{C} are lifted to $\mathbf{O}_J(\mathcal{C})$, as formulated in Section 5.3 of [19].

Proposition 1.14 (Product in $\mathbf{O}_J(\mathcal{C})$ [19]). Suppose \mathcal{C} has finite products and an orthogonality is reciprocal. Then $\mathbf{O}_J(\mathcal{C})$ has finite products

$$\mathbb{A} \& \mathbb{B} := (A \& B, \mathbb{A}_p \& \mathbb{B}_p := \{u \& v \mid u \in \mathbb{A}_p \ v \in \mathbb{B}_p\})$$

Note the second component is automatically closed under the double orthogonality. The forgetful $\mathbf{O}_J(\mathcal{C}) \rightarrow \mathcal{C}$ preserves them. The proof is put in A.1.1.

Definition 1.15 (stable tensor (Definition 58 of [19])). An orthogonality on a monoidal category \mathcal{C} stabilises the monoidal product when the following condition holds for all $U \subseteq \mathcal{C}(I, R)$ and $V \subseteq \mathcal{C}(I, S)$:

(stable tensor) $(U^{\circ\circ} \otimes V^{\circ\circ})^\circ = (U^{\circ\circ} \otimes V)^\circ = (U \otimes V^{\circ\circ})^\circ$

The stable tensor is a condition on a representability of certain maps in multicategories when \mathcal{C} has a closed structure on the monoidal product (see Section 5.3 of [19]). However the present paper does not assume the closedness.

The reciprocal orthogonality is strong enough to stabilise the monoidal product in \mathcal{C} :

Lemma 1.16. *Any reciprocal orthogonality stabilises monoidal products.*

Proof. We prove (\supset) of the stable tensor condition as the converse is tautological. Take any $\nu \in RHS$, which means $\forall f \in U^{\circ\circ} \forall g \in V \ f \otimes g = (f \otimes S) \circ (I \otimes g) \perp_{R \otimes S} R \otimes S \xrightarrow{\nu} J$ iff by reciprocity $g \perp_S S \cong I \otimes S \xrightarrow{f \otimes S} R \otimes S \xrightarrow{\nu} J$. But this means $\forall h \in V^{\circ\circ} \ h \perp_S \nu \circ (f \otimes S)$ iff by reciprocity $f \otimes h = (f \otimes S) \circ (I \otimes h) \perp_{R \otimes S} \nu$, which means $\nu \in LHS$. \square

Definition 1.17 (Monoidal product in $\mathbf{O}_J(\mathcal{C})$ [19]). Suppose \mathcal{C} is symmetric monoidal with an orthogonality stabilising \otimes . Then $\mathbf{O}_J(\mathcal{C})$ is symmetric monoidal and the forgetful $\mathbf{O}_J(\mathcal{C}) \rightarrow \mathcal{C}$ preserves the monoidality.

$$\mathbb{A} \otimes \mathbb{B} := (A \otimes B, (\mathbb{A}_p \otimes \mathbb{B}_p)^{\circ\circ})$$

The tensor unit \mathbb{I} is given by $(I, \{\text{Id}_I\}^{\circ\circ})$.

1.3. Lifting Free Exponential of \mathcal{C} to $\mathbf{O}_J(\mathcal{C})$

The equalisers $A^{\leq n}$ s and the limit $A^{\leq \infty}$ of \mathcal{C} are lifted to $\mathbb{A}^{\leq n}$ s and $\mathbb{A}^{\leq \infty}$ in any orthogonality category $\mathbf{O}_J(\mathcal{C})$ using the equaliser and the limit actions on \mathcal{C} -homset (Propositions 1.18 and 1.21). Although the orthogonality automatically guarantees the distribution of the monoidal product over the equaliser $E_{\mathbb{A}}$, it is necessary to impose a certain condition on $p_{\infty, n}$ and on \mathcal{C} -morphisms in order to guarantee the distribution of the monoidal product over the limit $L_{\mathbb{A}}$ (Proposition 1.24).

Proposition 1.18 (equaliser $\mathbb{A}^{\leq n}$ for $E_{\mathbb{A}}$ in $\mathbf{O}_J(\mathcal{C})$). In $\mathbf{O}_J(\mathcal{C})$ for every object $\mathbb{A} = (A, \mathbb{A}_p)$, the following object $\mathbb{A}^{\leq n}$ with $\text{eq}_{\mathbb{A}}$ becomes the equaliser of the $n!$ -symmetries of $(\mathbb{A} \& \mathbb{I})^{\otimes n}$:

$$\begin{aligned} \mathbb{A}^{\leq n} &= (A^{\leq n}, (\mathbb{A}^{\leq n})_p) \quad \text{with} \\ (\mathbb{A}^{\leq n})_p &:= \{\text{eq} \setminus h \mid h \in ((\mathbb{A} \& \mathbb{I})^{\otimes n})_p \text{ equalises the } n! \text{-symmetries of } \mathbb{A}_{\bullet}^{\otimes n}\}^{\circ\circ} \end{aligned} \quad (3)$$

Proof. In the proof, $X_{\bullet} := X \& I$ and $X_{\bullet}^{\otimes n}$ is a short for $(X_{\bullet})^{\otimes n}$. By the definition of the morphisms of the double gluing category, note first: If a morphism h of co-domain $\mathbb{X}^{\otimes n}$ equalises the $n!$ -symmetries of $\mathbb{X}^{\otimes n}$ in $\mathbf{O}_J(\mathcal{C})$, then h does so the $n!$ -symmetries of $X^{\otimes n}$ in \mathcal{C} . The following three conditions need to be checked: (i) The \mathcal{C} -morphism $\text{eq}_{\mathbb{A}} \setminus h$ resides in $\mathbf{O}_J(\mathcal{C})$ for any $h : \mathbb{B} \rightarrow \mathbb{A}_{\bullet}^{\otimes n}$ equalising the $n!$ -symmetries in $\mathbf{O}_J(\mathcal{C})$: For any $b \in \mathbb{B}_p$, $(\text{eq}_{\mathbb{A}} \setminus h) \circ b = \text{eq}_{\mathbb{A}} \setminus (h \circ b)$, which belongs to (inside the scope $^{\circ\circ}$ of) (3) as $h \circ b$ belongs to $(\mathbb{A}_{\bullet}^{\otimes n})_p$ and equalises the $n!$ -symmetries of $\mathbb{A}_{\bullet}^{\otimes n}$ by the first note.

(ii) The \mathcal{C} -morphism $\text{eq}_{\mathbb{A}}$ resides in $\mathbf{O}_J(\mathcal{C})$: (\bar{p}) condition holds directly by the definition (3).

(iii) Any $\mathbf{O}_J(\mathcal{C})$ -morphism $h : \mathbb{B} \rightarrow \mathbb{A}_{\bullet}^{\otimes n}$ equalising the $n!$ -symmetries factors via $\mathbb{A}^{\leq n}$:

By the first note, $h : \mathbb{B} \rightarrow \mathbb{A}_{\bullet}^{\otimes n}$ factors via $A^{\leq n}$ in \mathcal{C} . But by (i), the factorisation is that for $\mathbf{O}_J(\mathcal{C})$. \square

Remark 1.19 (on Proposition 1.18). The homset $(\mathbb{A}^{\leq n})_p$ of (3) in particular contains the following homset (but not vice versa in general)

$$\{ I^{\otimes n} \cong I \xrightarrow{\epsilon_n} I^{\leq n} \xrightarrow{f^{\leq n}} A^{\leq n} \mid I \xrightarrow{f} A \in \mathbb{A}_p \}, \quad \text{where } \epsilon_n \text{ is } I \cong I^{\otimes n} \xrightarrow{\text{eq}_I \setminus (I \& I)^{\otimes n}} I^{\leq n}.$$

Lemma 1.20. The morphism $p_{n+1, n}$ resides in $\mathbf{O}_J(\mathcal{C})$ so that it is a morphism from $\mathbb{A}^{\leq n+1}$ to $\mathbb{A}^{\leq n}$.

Proof. Obviously (\bar{p}) condition holds by the definition (3). \square

By this lemma and Proposition 1.18, $\{p_{n+1, n}\}_n$ becomes a diagram for $L_{\mathbb{A}}$ in $\mathbf{O}_J(\mathcal{C})$. Then

Proposition 1.21 (limit $\mathbb{A}^{\leq \infty}$ for $L_{\mathbb{A}}$ in $\mathbf{O}_J(\mathcal{C})$).

In $\mathbf{O}_J(\mathcal{C})$ for every object $\mathbb{A} = (A, \mathbb{A}_p)$, the following $\mathbb{A}^{\leq \infty}$ with $\{p_{\infty, n} : \mathbb{A}^{\leq \infty} \rightarrow \mathbb{A}^{\leq n}\}_n$ becomes the limit for the sequential diagram $\{p_{\infty, n} : \mathbb{A}^{\leq n+1} \rightarrow \mathbb{A}^{\leq n}\}_n$:

$$\mathbb{A}^{\leq \infty} := (A^{\leq \infty}, (\mathbb{A}^{\leq \infty})_p) \quad \text{with}$$

Proposition 1.23 (distribution of \otimes over E_A in $\mathbf{O}_J(\mathcal{C})$). *For any object \mathbb{B} of $\mathbf{O}_J(\mathcal{C})$, $\mathbb{A}^{\leq n} \otimes \mathbb{B}$ becomes the equaliser of $(n!$ -symmetries) $\otimes \mathbb{B}$ of $(\mathbb{A} \& \mathbb{I})^{\otimes n} \otimes \mathbb{B}$.*

Proof. The assertion is direct by the definition of monoidal product in $\mathbf{O}_J(\mathcal{C})$ as the condition (\bar{p}) holds automatically as follows: $\text{eq} \otimes B : \mathbb{A}^{\leq n} \otimes \mathbb{B} \longrightarrow (\mathbb{A} \& \mathbb{I})^{\otimes n} \otimes \mathbb{B}$. By definition $(\mathbb{A}^{\leq n} \otimes \mathbb{B})_p = ((\mathbb{A}^{\leq n})_p \otimes \mathbb{B}_p)^{\circ\circ}$ and $((\mathbb{A} \& \mathbb{I})^{\otimes n} \otimes \mathbb{B})_p = (((\mathbb{A} \& \mathbb{I})^{\otimes n})_p \otimes \mathbb{B}_p)^{\circ\circ}$. Then for any $x \otimes y \in (\mathbb{A}^{\leq n})_p \otimes \mathbb{B}_p$, $(\text{eq} \otimes B) \circ (x \otimes y) = (\text{eq} \circ x) \otimes y \in ((\mathbb{A} \& \mathbb{I})^{\otimes n})_p \otimes \mathbb{B}_p$. \square

On the other hand, the distribution over the limit $\mathbb{A}^{\leq \infty}$ is not retained in general in $\mathbf{O}_J(\mathcal{C})$. Hence we need to augment the following condition on the orthogonality and the monoidal product:

Proposition 1.24 (condition for monoidal product \otimes to distribute over the limit in $\mathbf{O}_J(\mathcal{C})$). *The following condition in $\mathbf{O}_J(\mathcal{C})$ is necessary and sufficient for the distribution of \otimes over the limit L_A .*

$$((\mathbb{A}^{\leq \infty})_p \otimes \mathbb{B}_p)^{\circ\circ} = \bigcap_{n \in \mathbb{N}} (p_{\infty, n} \otimes B)^{-1} \circ ((\mathbb{A}^{\leq n})_p \otimes \mathbb{B}_p)^{\circ\circ} \quad (5)$$

Notation: $f^{-1} \circ U := \{x : I \longrightarrow X \mid f \circ x \in U\}$ for a morphism $f : X \longrightarrow Y$ in \mathcal{C} and a homset $U \subseteq \mathcal{C}(I, Y)$.

The condition when $\mathbb{B} = \mathbb{I}$ is automatically valid for any $\mathbf{O}_J(\mathcal{C})$. But not necessarily so for a general object \mathbb{B} .

Proof. First note that \subset of (5) is tautological for any \mathcal{C} by definitions (3) and (4). Hence the condition (5) is equivalent to the following (6):

$$\forall n \in \mathbb{N} (p_{\infty, n} \otimes B) \circ u \in ((\mathbb{A}^{\leq n})_p \otimes \mathbb{B}_p)^{\circ\circ} \implies u \in ((\mathbb{A}^{\leq \infty})_p \otimes \mathbb{B}_p)^{\circ\circ} \quad (6)$$

The condition (6) says that x_{∞} becomes the mediating morphism for the cone $\{x_n : \mathbb{I} \longrightarrow \mathbb{A}^{\leq n} \otimes \mathbb{B}\}$ in $\mathbf{O}_J(\mathcal{C})$, hence derives the necessity and sufficiency. In particular, (6) when $\mathbb{B} = \mathbb{I}$ is Proposition 1.21, hence is valid in any $\mathbf{O}_J(\mathcal{C})$. \square

The main theorem of this section is obtained by Propositions 1.23 and 1.24.

Theorem 1.25 (Free Exponential in $\mathbf{O}_J(\mathcal{C})$ with the condition (5)). *Suppose an orthogonality on a monoidal category \mathcal{C} is reciprocal and satisfies the condition (5) of Proposition 1.24. Then the orthogonality category $\mathbf{O}_J(\mathcal{C})$ has the free exponential whenever \mathcal{C} has so that its forgetful image by $\mathbf{O}_J(\mathcal{C}) \longrightarrow \mathcal{C}$ is the free exponential of \mathcal{C} .*

2. Monoidal Category \mathbf{TsK} of s-finite Transition Kernels with Biproducts

This section concerns a measure theoretic study, independent from Section 1. The main sources of the section are Staton [27] and Hamano [18]. We also refer Bauer's book [1] for general measure theory.

2.1. Preliminaries from Measure Theory

This subsection recalls some basic definitions and a theorem from measure theory, necessary in this paper.

(Terminology) \mathbb{N} denotes the set of non negative integers. \mathbb{R}_+ denotes the set of non negative reals. $\overline{\mathbb{R}}_+$ denotes $\mathbb{R}_+ \cup \{\infty\}$. \mathfrak{S}_n denotes the symmetric group over $\{1, \dots, n\}$. For a subset A , χ_A denotes the characteristic function of A . $\delta_{x, y}$ is the Kronecker delta. \uplus denotes the disjoint union of sets.

Definition 2.1 (σ -field \mathcal{X} and measurable space (X, \mathcal{X})).

A σ -field over a set X is a family \mathcal{X} of subsets of X containing \emptyset , closed under the complement and countable union. A pair (X, \mathcal{X}) is called a *measurable space*. The members of \mathcal{X} are called *measurable sets*. The measurable space is often written simply by \mathcal{X} , as X is the largest element in \mathcal{X} . For a measurable set $Y \in \mathcal{X}$, the measurable subspace $\mathcal{X} \cap Y$, called the *restriction on Y* , is defined by $\mathcal{X} \cap Y := \{A \cap Y \mid A \in \mathcal{X}\}$.

Definition 2.2 ($\sigma(\mathcal{F})$ and Borel σ -field \mathcal{B}_+). For a family \mathcal{F} of subsets of X , $\sigma(\mathcal{F})$ denotes the σ -field generated by \mathcal{F} , i.e., the smallest σ -field containing \mathcal{F} . When X is $\overline{\mathbb{R}}_+$ and \mathcal{F} is the family $\mathcal{O}_{\overline{\mathbb{R}}_+}$ of the open sets in $\overline{\mathbb{R}}_+$ (with the topology whose basis consists of the open intervals in \mathbb{R}_+ together with $(a, \infty) := \{x \mid a < x\}$ for all $a \in \mathbb{R}_+$), the σ -field is denoted by \mathcal{B}_+ , whose members are called Borel sets over $\overline{\mathbb{R}}_+$.

Definition 2.3 (measurable function). For measurable spaces (X, \mathcal{X}) and (Y, \mathcal{Y}) , a function $f : X \rightarrow Y$ is $(\mathcal{X}, \mathcal{Y})$ -measurable (often just *measurable*) if $f^{-1}(B) \in \mathcal{X}$ whenever $B \in \mathcal{Y}$. In this paper, a measurable function unless otherwise mentioned is to the Borel set \mathcal{B}_+ over $\overline{\mathbb{R}}_+$ from some measurable space (X, \mathcal{X}) .

Definition 2.4 (measure). A *measure* μ on a measurable space (X, \mathcal{X}) is a function from \mathcal{X} to $\overline{\mathbb{R}}_+$ satisfying (σ -additivity): If $\{A_i \in \mathcal{X} \mid i \in I\}$ is a countable family of pairwise disjoint sets, then $\mu(\bigcup_{i \in I} A_i) = \sum_{i \in I} \mu(A_i)$.

Definition 2.5 (integration). For a measure μ on (X, \mathcal{X}) , and a $(\mathcal{X}, \mathcal{B}_+)$ -measurable function f , the integral of f over X wrt the measure μ is defined by $\int_X f(x) \mu(dx)$, which is simply written $\int_X f d\mu$. It is also written $\int_X d\mu f$.

Theorem 2.6 (monotone convergence). *Let μ be a measure on a measurable space (X, \mathcal{X}) . For an monotonic sequence $\{f_n\}$ of $(\mathcal{X}, \mathcal{B}_+)$ -measurable functions, if $f = \sup_n f_n$, then f is measurable and $\int_X f d\mu = \sup \int_X f_n d\mu$.*

Definition 2.7 (push forward measure $\mu \circ F^{-1}$ along a measurable function F). For a measure μ on (Y, \mathcal{Y}) and a measurable function F from (Y, \mathcal{Y}) to (Y', \mathcal{Y}') , $\mu'(B') := \mu(F^{-1}(B'))$ becomes a measure on (Y', \mathcal{Y}') , called *push forward measure* of μ along F . The push forward measure has a following property, called “*variable change of integral along push forward F* ”:

$$\int_{Y'} g d\mu' = \int_Y (g \circ F) d\mu. \quad \text{That is,} \quad \int_{Y'} g(y') \mu'(dy') = \int_Y g(F(y)) \mu(dy) \quad (7)$$

The push forward measure μ' is often denoted by $\mu \circ F^{-1}$ by abuse of notation.

2.2. Transition Kernels

Definition 2.8 (transition kernel [1]). For measurable spaces (X, \mathcal{X}) and (Y, \mathcal{Y}) , a *transition kernel* from (X, \mathcal{X}) to (Y, \mathcal{Y}) is a function

$$\kappa : X \times \mathcal{Y} \rightarrow \overline{\mathbb{R}}_+ \quad \text{satisfying}$$

- (i) For each $x \in X$, the function $\kappa(x, -) : \mathcal{Y} \rightarrow \overline{\mathbb{R}}_+$ is a measure on (Y, \mathcal{Y}) .
- (ii) For each $B \in \mathcal{Y}$, the function $\kappa(-, B) : X \rightarrow \overline{\mathbb{R}}_+$ is measurable on (X, \mathcal{X}) .

Definition 2.9 (operations κ_* and κ^* of a kernel κ on measures and measurable functions).

Let $\kappa : (X, \mathcal{X}) \rightarrow (Y, \mathcal{Y})$ be a transition kernel.

- For a measure μ on \mathcal{X} ,

$$(\kappa_* \mu)(B) := \int_X \kappa(x, B) \mu(dx)$$

is a measure on \mathcal{Y} , where $B \in \mathcal{Y}$.

- For a measurable function f on \mathcal{Y} ,

$$(\kappa^* f)(x) := \int_Y f(y) \kappa(x, dy)$$

is measurable on \mathcal{X} , where $x \in X$. In particular, for a characteristic function χ_B for any $B \in \mathcal{Y}$,

$$(\kappa^* \chi_B)(x) := \kappa(x, B) \quad (8)$$

It is direct to check, by the monotone convergence theorem 2.6, that κ^*f is measurable.

Definition 2.10 (category TK of transition kernels). TK denotes the category where each object is a measurable space (X, \mathcal{X}) and a morphism is a transition kernel $\kappa(x, B)$ from (X, \mathcal{X}) to (Y, \mathcal{Y}) . The composition is the *convolution* of two kernels $\kappa(x, B) : (X, \mathcal{X}) \rightarrow (Y, \mathcal{Y})$ and $\iota(y, C) : (Y, \mathcal{Y}) \rightarrow (Z, \mathcal{Z})$:

$$\iota \circ \kappa(x, C) = \int_Y \kappa(x, dy) \iota(y, C) \quad (9)$$

$\text{Id}_{(X, \mathcal{X})}$ is *Dirac delta measure* $\delta : (X, \mathcal{X}) \rightarrow (X, \mathcal{X})$, defined by for $x \in X$ and $A \in \mathcal{X}$;

$$\text{if } x \in A \text{ then } \delta(x, A) = 1, \text{ else } \delta(x, A) = 0.$$

Remark 2.11. Measures and measurable functions both reside as morphisms in TK: Let (I, \mathcal{I}) be the singleton measurable space with $I = \{*\}$, hence $\mathcal{I} = \{\emptyset, \{*\}\}$, then

$$\text{TK}(\mathcal{I}, \mathcal{X}) = \{\kappa(*, A) \mid A \in \mathcal{X}\} = \{\text{the measures } \mu \text{ on } (X, \mathcal{X})\}$$

$$\text{TK}(\mathcal{X}, \mathcal{I}) = \{\kappa(x, \{*\}) \mid x \in X\} \cup \{\kappa(x, \emptyset) = 0 : X \rightarrow \overline{\mathbb{R}}_+\} = \{\text{the measurable functions } f \text{ on } (X, \mathcal{X}) \text{ to } \mathcal{B}_+\}$$

The operations κ_* and κ^* of Definition 2.9 are respectively categorical precomposition and composition with κ in TK so that $\kappa_*\mu = \kappa \circ \mu$ and $\kappa^*f = f \circ \kappa$.

In the sequel, when a transition kernel κ has the domain (resp. co-domain) \mathcal{I} in TsK^{op} , then $\kappa(\{*\}, x)$ (resp. $\kappa(X, *)$) is simply written as a measurable function $\kappa(x)$ (resp. measure $\kappa(X)$).

Remark 2.12 (SRel [24, 23]). The category SRel of *stochastic relations* is a wide subcategory of TK strengthening the conditions of Definition 2.8 into (i) $\kappa(x, -)$ is a (*sub*)*probability* measure (i.e., its domain is $[0, 1]$) and (ii) $\kappa(-, B)$ is *bounded* measurable. The morphisms in SRel are called (*sub*)*Markov kernels*.

It is now well known that the composition (9) for Markov kernels comes from Giry's probabilistic monad, resembling the power set monad of the relational composition [17, 24].

2.3. Countable Biproducts in TK

Proposition 2.13 (biproduct \coprod). TK has countable biproducts \coprod .

Proof. Given a countable family $\{(X_i, \mathcal{X}_i)\}_i$ of measurable spaces, we define

$$\coprod_i (X_i, \mathcal{X}_i) := \left(\bigcup_i \{i\} \times X_i, \biguplus_i \mathcal{X}_i \right), \quad (10)$$

where $\biguplus_i \mathcal{X}_i := \{\bigcup_i \{i\} \times A_i \mid A_i \in \mathcal{X}_i\}$ is the σ -field generated by the measurable sets of each summands.

(Coproduct): (10) defines a coproduct for TK. The injection $\text{in}_j : (X_j, \mathcal{X}_j) \rightarrow (\bigcup \{i\} \times X_i, \biguplus \mathcal{X}_i)$ is defined by $\text{in}_j(x_j, \sum_i A_i) := \delta(x_j, A_i)$. The mediating morphism $\oplus_{i \in I} f_i : (\bigcup \{i\} \times X_i, \biguplus \mathcal{X}_i) \rightarrow (Y, \mathcal{Y})$ for given morphisms $f_i : (X_i, \mathcal{X}_i) \rightarrow (Y, \mathcal{Y})$ is defined by $(\oplus_{j \in I} f_j)((i, x), B) = f_i(x, B)$. Note (10) is the same instance as the known coproduct in SRel. However, in the relaxed structure of TK, we have moreover;

(Product): (10) becomes a product for TK. The projection $\text{p}_i : \coprod_i (X_i, \mathcal{X}_i) \rightarrow (X_i, \mathcal{X}_i)$ is given by $\text{p}_i((j, x), A_i) := \delta_{i,j} \cdot \delta(x, A_i)$. The mediating morphism $\&_i g_i$ for given morphisms $g_i : (Y, \mathcal{Y}) \rightarrow (X_i, \mathcal{X}_i)$ is defined to be $(\&_i g_i)(b, \bigcup \{i\} \times A_i) := \sum_i g_i(b, A_i)$. Note the construction for the mediating morphism is not closed in Markov kernels, but is so in transition kernels. This construction shows how values of measurable functions include the infinite real ∞ when I becomes infinite. We check the uniqueness of the mediating morphism, say m :

$$\begin{aligned} (\text{p}_j \circ m)(b, \{j\} \times A_j) &= \int_{\biguplus_i X_i} m(b, d(i, x)) \text{p}_j((i, x), \{j\} \times A_j) = \int_{\biguplus_i X_i} m(b, d(i, x)) \delta_{j,i} \cdot \delta((i, x), \{j\} \times A_j) \\ &= \int_{X_j} m(b, d(i, x)) \delta((i, x), \{j\} \times A_j) = m(b, \{j\} \times A_j) \end{aligned} \quad (11)$$

The required commutativity for m is $p_j \circ m = g_j$, which holds by (11) if and only if $m(b, A_j) = g_j(b, A_j)$ for all j . Since $m(b, -)$ is a measure and $\{A_j\}_{j \in J}$ are disjoint, this yields the definition $\&_i g_i$ of the mediating morphism. \square

The unit of the biproduct is the null measurable space $\mathcal{T} = (\emptyset, \{\emptyset\})$.

2.4. Monoidal Product and Countable Biproducts in \mathbf{TsK}

Definition 2.14 (product of measurable spaces [1, 24]). The product of measurable spaces (X_1, \mathcal{X}_1) and (X_2, \mathcal{X}_2) is the measurable space $(X_1 \times X_2, \mathcal{X}_1 \otimes \mathcal{X}_2)$, where $\mathcal{X}_1 \otimes \mathcal{X}_2$ denotes the σ -field over the cartesian product $X_1 \times X_2$ generated by *measurable rectangles* $A_1 \times A_2$'s such that $A_i \in \mathcal{X}_i$.

In order to accommodate measures into the product of measurable spaces, each measure μ_i on (X_i, \mathcal{X}_i) needs to be extended uniquely to the product. The condition of σ -finiteness ensures this, yielding the unique product measure over the product measurable space:

Definition 2.15 (σ -finiteness [1, 24]). A measure μ on (X, \mathcal{X}) is σ -finite when the set X is written as a countable union of sets of finite measures. That is, $\exists A_1, A_2, \dots \in \mathcal{X}$ such that $\mu(A_i) < \infty$ and $X = \cup_{i=1}^{\infty} A_i$.

Definition 2.16 (product measure [1, 24]). For σ -finite measures μ_i on (X_i, \mathcal{X}_i) with $i = 1, 2$, there exists a unique measure μ on $(X_1 \times X_2, \mathcal{X}_1 \otimes \mathcal{X}_2)$ such that $\mu(A_1 \times A_2) = \mu_1(A_1)\mu_2(A_2)$. μ is written $\mu_1 \otimes \mu_2$ and called the *product measure* of μ_1 and μ_2 .

The product measure derived from σ -finite measures guarantees the basic theorem in measure theory, stating double integration is treated as iterated integrations.

Theorem 2.17 (Fubini-Tonelli [1, 24]). For σ -finite measures μ_i on (X_i, \mathcal{X}_i) with $i = 1, 2$ and a $(\mathcal{X}_1 \otimes \mathcal{X}_2, \mathcal{B}_+)$ -measurable function f ,

$$\int_{X_1 \times X_2} f d(\mu_1 \otimes \mu_2) = \int_{X_2} d\mu_2 \int_{X_1} f d\mu_1 = \int_{X_1} d\mu_1 \int_{X_2} f d\mu_2$$

The Fubini-Tonelli Theorem becomes crucial category theoretically to us for the following two (i) dealing with functoriality of morphisms on the product measurable spaces (cf. Proposition 2.20 below) (ii) giving a new instance of the orthogonality using measure theory (cf. Proposition 4.2 of Section 4).

Since the σ -finiteness retained in the category \mathbf{TK} is not closed under the categorical composition, we define the two classes of transition kernels, *finiteness* and *s-finiteness*, respectively by tightening and loosening the σ -finiteness so that the both classes are preserved under the composition of \mathbf{TK} .

Definition 2.18 (s-finite kernels [26, 27]). Let κ be a transition kernel from (X, \mathcal{X}) to (Y, \mathcal{Y}) .

- κ is called *finite* when $\sup_{x \in X} \kappa(x, Y) < \infty$; i.e., the condition says that up to the scalar $0 < a < \infty$ factor determined by the sup, κ is Markovian.
- κ is called *s-finite* (i.e., sum of finite) when $\kappa = \sum_{i \in \mathbb{N}} \kappa_i$ where each κ_i is a finite kernel from (X, \mathcal{X}) to (Y, \mathcal{Y}) and the sum is defined by $(\sum_{i \in \mathbb{N}} \kappa_i)(x, B) := \sum_{i \in \mathbb{N}} \kappa_i(x, B)$. This is well-defined because any countable sum of kernels from (X, \mathcal{X}) to (Y, \mathcal{Y}) becomes a kernel of the same type.

In the definition of s-finiteness, note that $(\sum_{i \in \mathbb{N}} \kappa_i)^* = \sum_{i \in \mathbb{N}} \kappa_i^*$ and $(\sum_{i \in \mathbb{N}} \kappa_i)_* = \sum_{i \in \mathbb{N}} (\kappa_i)_*$ for the operations of Definition 2.9: That is, the preservation of the operation $(\)^*$ (resp. of $(\)_*$) means the commutativity of integral over countable sum of measures (resp. of measurable functions).

Remark 2.19. The both classes of the finite kernels and of the s-finite kernels are closed under the categorical composition of \mathbf{TK} . This is directly calculated for the finite kernels, to which the s-finite ones are reduced by virtue of the note in the above paragraph. We refer to the proof of Lemma 3 of [27] for the calculation.

The original Fubini-Tonelli (Theorem 2.17) for the σ -finite measures extends to the s-finite measures:

Proposition 2.20 (Fubini-Tonelli extending for s-finite measures (cf. Proposition 5 of Staton [27])). *For the same f as Theorem 2.17 but μ_1 and μ_2 are s-finite measures, it holds;*

$$\int_{X_2} d\mu_2 \int_{X_1} f d\mu_1 = \int_{X_1} d\mu_1 \int_{X_2} f d\mu_2$$

See the proof in A.1.2.

It is derived from Proposition 2.20 that the s-finite transition kernels form a monoidal category.

Definition 2.21 (monoidal subcategories \mathbf{TsK} of s-finite kernels). \mathbf{TsK} is a wide subcategory of \mathbf{TK} , whose morphisms are the *s-finite* transition kernels. \mathbf{TsK} has a symmetric monoidal product \otimes : On objects is by Definition 2.16. Given morphisms $\kappa_1 : (X_1, \mathcal{X}_1) \rightarrow (Y_1, \mathcal{Y}_1)$ and $\kappa_2 : (X_2, \mathcal{X}_2) \rightarrow (Y_2, \mathcal{Y}_2)$, their product is defined explicitly:

$$(\kappa_1 \otimes \kappa_2)((x_1, x_2), C) := \int_{Y_1} \kappa_1(x, dy_1) \int_{Y_2} \kappa_2(x, dy_2) \chi_C((y_1, y_2))$$

Alternatively, thanks to Fubini-Tonelli (Proposition 2.20), the product is implicitly defined as the unique transition kernel $\kappa_2 \otimes \kappa_1 : (X_1 \times X_2, \mathcal{X}_1 \otimes \mathcal{X}_2) \rightarrow (Y_1 \times Y_2, \mathcal{Y}_1 \otimes \mathcal{Y}_2)$ satisfying the following for any rectangle $B_1 \times B_2$ with $B_i \in \mathcal{X}_i$:

$$(\kappa_1 \otimes \kappa_2)((x_1, x_2), B_1 \times B_2) = \kappa_1(x_1, B_1) \kappa_2(x_2, B_2)$$

The unit of the monoidal product is the singleton measurable space (I, \mathcal{I}) .

Proposition 2.22 (The biproducts).

\mathbf{TsK} has countable biproducts which are those in \mathbf{TK} residing inside the subcategory.

Proof. The coproduct construction of Proposition 2.13 all works under the additional constraint of the s-finiteness of kernels. For the product construction, the only construction necessary to be checked is that of the mediating morphism $\&_i g_i$, employing the sum over $i \in I$ for a countable infinite I : If given g_i 's of the product construction in Proposition 2.13 are s-finite, then each is written $g_i = \sum_{j \in \mathbb{N}} g_{(i,j)}$, where each $g_{(i,j)}$ is a finite kernel from (Y, \mathcal{Y}) to (X_i, \mathcal{X}_i) , thus so is g_i . In what follows, the index set I is identified with \mathbb{N} . A transition kernel h_n is defined for each $n \in \mathbb{N}$:

$$h_n := \sum_{i+j=n} \text{in}_i \circ g_{(i,j)} : (Y, \mathcal{Y}) \rightarrow \coprod_{j \in I} (X_j, \mathcal{X}_j)$$

where $\text{in}_i : (X_i, \mathcal{X}_i) \rightarrow \coprod_{j \in I} (X_j, \mathcal{X}_j)$ is the coproduct injection. Note h_n is a finite kernel, as the sum specified by the subscript $i + j = n$ is finite. Then in terms of the finite kernels, the mediating morphism $\&_i g_i$ constructed in Proposition 2.13 is represented as follows to become s-finite:

$$\&_{i \in \mathbb{N}} g_i = \sum_{n \in \mathbb{N}} h_n$$

□

Note: In the sequel, only the product structure of \coprod is employed. That is Sections 3 and 4 do not treat \coprod as biproduct but simpler as product. Accordingly, \coprod is written by $\&$.

3. The Free Exponential in \mathbf{TsK}^{op}

This section constructs the free exponential structure of \mathbf{TsK}^{op} . The opposite setting is chosen accordingly to [18] by virtue of the asymmetry between the first and the second arguments of the transition kernels.

In our measure-theoretic framework, the equaliser for the exponential is given slightly in general, not only for a rooted object $\mathcal{X}_\bullet = \mathcal{X} \& \mathcal{I}$ with the monoidal unit \mathcal{I} , but also for a general object \mathcal{X} of \mathbf{TsK}^{op} :

Definition 3.1 (measurable space $(X^{(n)}, \mathcal{X}^{(n)})$). For any measurable space (X, \mathcal{X}) and any natural number n , a measurable space $(X^{(n)}, \mathcal{X}^{(n)})$ is defined by

$$X^{(n)} := \{x_1 \cdots x_n \mid x_i \in X\} \quad \text{and} \quad \mathcal{X}^{(n)} := \{A \subseteq X^{(n)} \mid F^{-1}(A) \in \mathcal{X}^{\otimes n}\},$$

where $F : X^{\otimes n} \rightarrow X^{(n)} \quad (x_1, \dots, x_n) \mapsto x_1 \cdots x_n$

The members $x_1 \cdots x_n$ in $X^{(n)}$ are formal products whose order of factor is irrelevant. On the other hand, the members of $X^{\otimes n}$ are ordered sequences, hence the map F forgets the order of factor.

Note $\mathcal{X}^{(n)}$ is automatically a σ -field over $X^{(n)}$. That is, $(X^{(n)}, \mathcal{X}^{(n)})$ is the push forward measurable space of the n -th direct product $(X^{\otimes n}, \mathcal{X}^{\otimes n})$ along the map $F : X^{\otimes n} \rightarrow X^{(n)}$ forgetting the order.

Notation: An element $x_1 \cdots x_n \in X^{(n)}$ is abbreviated by \mathbf{x} when n is clear from the context, while an element $(x_1, \dots, x_n) \in X^{\otimes n}$ is abbreviated by $\vec{\mathbf{x}}$.

Proposition 3.2 ($\mathcal{X}^{(n)}$ as equaliser). In $\mathbf{TKer}^{\text{op}}$, the object $\mathcal{X}^{(n)}$ becomes the equaliser of the $n!$ -symmetries of $\mathcal{X}^{\otimes n}$. The transition kernel $\text{eq}_{\mathcal{X}} : \mathcal{X}^{(n)} \rightarrow \mathcal{X}^{\otimes n}$ is specified by

$$\text{eq}(-, (x_1, \dots, x_n)) := \delta(-, x_1 \cdots x_n) \quad (12)$$

For any transition kernel κ to $\mathcal{X}^{\otimes n}$ equalising the $n!$ -symmetries, its unique factorization $\text{eq} \backslash \kappa$ via eq is given by

$$\text{eq} \backslash \kappa(-, x_1 \cdots x_n) := \kappa(-, (x_1, \dots, x_n)) \quad (13)$$

$\text{eq} \backslash \kappa$ is well defined (independently of the ordering $x_1 \cdots x_n$), because $\kappa(-, (x_1, \dots, x_n)) = \kappa(-, (x_{\sigma(1)}, \dots, x_{\sigma(n)}))$ for all $\sigma \in \mathfrak{S}_n$.

Proof. Precompose any transition kernel τ (of the codomain $\mathcal{X}^{(n)}$) to $\text{eq}_{\mathcal{X}}$:

$$\text{eq}_{\mathcal{X}} \circ \tau(-, (x_1, \dots, x_n)) = \int_{X^{(n)}} \tau(-, \mathbf{y}) \text{eq}(d\mathbf{y}, (x_1, \dots, x_n)) = \int_{X^{(n)}} \tau(-, \mathbf{y}) \delta(d\mathbf{y}, x_1 \cdots x_n) = \tau(-, x_1 \cdots x_n)$$

Thus $\text{eq} \backslash \kappa$ gives the unique factorisation of κ . \square

The equaliser of Proposition 3.2 acts also on morphisms for any $\kappa : \mathcal{Y} \rightarrow \mathcal{X}$: The transition kernel $\kappa^{(n)} : \mathcal{Y}^{(n)} \rightarrow \mathcal{X}^{(n)}$ is defined by the unique factorization via $\mathcal{X}^{(n)}$ of $\kappa^{\otimes n} \circ \text{eq}_{\mathcal{Y}}$ equalising the $n!$ -symmetries of $\mathcal{X}^{\otimes n}$: See the diagram;

$$\begin{array}{ccccc} & & \text{eq}_{\mathcal{X}} & & \\ & \swarrow & \text{---} & \searrow & \\ \mathcal{X}^{(n)} & \xleftarrow{\kappa^{(n)}} & \mathcal{Y}^{(n)} & \xrightarrow{\text{eq}_{\mathcal{Y}}} & \mathcal{Y}^{\otimes n} & \xrightarrow{\kappa^{\otimes n}} & \mathcal{X}^{\otimes n} \end{array} \quad (14)$$

Example 3.3. $\mathfrak{p}_l^{(n)} : (\mathcal{X}_1 \& \mathcal{X}_2)^{(n)} \rightarrow (\mathcal{X}_1)^{(n)}$ is

$$\mathfrak{p}_l^{(n)}(-, x_1 \cdots x_n) = \delta(-, ((1, x_1), \dots, (1, x_n)))$$

for the left projection $\mathfrak{p}_l : \mathcal{X}_1 \& \mathcal{X}_2 \rightarrow \mathcal{X}_1$, which is defined $\mathfrak{p}_l(-, x) = \delta(-, (1, x))$. The equality is by the following starting from (LHS via (13)) and ending at (RHS via (12)) in $(\mathcal{X}_1 \& \mathcal{X}_2)^{(n)} \rightarrow \mathcal{X}_1^{(n)}$:

$$\begin{aligned} (\mathfrak{p}_l)^{\otimes n} \circ \text{eq}(-, (x_1, \dots, x_n)) &= \int_{(X \uplus Y)^{\otimes n}} \text{eq}(-, \vec{\mathbf{z}}) (\mathfrak{p}_l)^{\otimes n}(d\vec{\mathbf{z}}, (x_1, \dots, x_n)) \\ &= \int_{(X \uplus Y)^{\otimes n}} \text{eq}(-, \vec{\mathbf{z}}) \delta^{\otimes n}(d\vec{\mathbf{z}}, ((1, x_1), \dots, (1, x_n))) \\ &= \text{eq}(-, ((1, x_1), \dots, (1, x_n))) \end{aligned}$$

Proposition 3.2 directly makes the equaliser $\mathcal{X}^{\leq n}$ for $\text{E}_{\mathcal{X}}$ of Definition 1.1 for \mathbf{TsK}^{op} definable by

$$\mathcal{X}^{\leq n} := (\mathcal{X} \& \mathcal{I})^{(n)}$$

Then, we observe that the canonical $p_{n+1,n} : (\mathcal{X}\&\mathcal{I})^{(n+1)} \longrightarrow (\mathcal{X}\&\mathcal{I})^{(n)}$ is described in \mathbf{TsK}^{op} by

$$p_{n+1,n}(-, z_1 \cdots z_n) = \delta(-, z_1 \cdots z_n(2, *)) \quad (15)$$

because of the following equalities starting from (LHS via (13)) and ending at (RHS via (12)), where $\mathcal{X}_{\bullet}^{\otimes n}$ abbreviates $(\mathcal{X}\&\mathcal{I})^{\otimes n}$:

$$\begin{aligned} & (\mathcal{X}_{\bullet}^{\otimes n} \otimes \mathbf{p}_r) \circ \mathbf{eq}(d(y_1, \dots, y_n, y_{n+1}), (z_1, \dots, z_n)) \\ &= \int_{(X \uplus I)^{\otimes (n+1)}} \mathbf{eq}(-, (\vec{y}, y_{n+1})) (\mathcal{X}_{\bullet}^{\otimes n} \otimes \mathbf{p}_r)(d(\vec{y}, y_{n+1}), (z_1, \dots, z_n)) \\ & \hspace{15em} \text{where } (\vec{y}, y_{n+1}) = (y_1, \dots, y_n, y_{n+1}) \\ & \stackrel{FT}{=} \int_{(X \uplus I)^{\otimes n}} \int_{X \uplus I} \mathbf{eq}(-, (\vec{y}, y_{n+1})) (\mathcal{X}_{\bullet}^{\otimes n} \otimes \mathbf{p}_r)(d(\vec{y}, y_{n+1}), (z_1, \dots, z_n)) \\ &= \int_{(X \uplus I)^{\otimes n}} \mathbf{eq}(-, (\vec{y}, (2, *))) \mathcal{X}_{\bullet}^{\otimes n}(\vec{y}, (z_1, \dots, z_n)) \\ &= \mathbf{eq}(-, (z_1, \dots, z_n, (2, *))) \end{aligned}$$

Intuitively, the transition kernel $p_{n+1,n}$ may be seen to forget the rooted element $(2, *)$. This leads us to define that the limit of the sequential diagram of the (rooted) equalisers $\mathcal{X}^{\leq n} = (\mathcal{X}\&\mathcal{I})^{(n)}$ is given by the countable infinite products of the (rootless) equalisers $\mathcal{X}^{(n)}$.

Definition 3.4 (measurable space $!\mathcal{X}$). For any measurable space \mathcal{X} , the following measurable space of the countable infinite products of $\mathcal{X}^{(k)}$ s is denoted by

$$!\mathcal{X} := \left(\bigsqcup_{k \in \mathbb{N}} X^{(k)}, \&\mathcal{X}^{(k)} \right)$$

Theorem 3.5 (limit $L_{\mathcal{X}}$ for \mathbf{TsK}^{op}). For any object \mathcal{X} in \mathbf{TsK}^{op} , $!\mathcal{X}$ of Definition 3.4 and the following morphisms $\{p_{\infty,n}\}_n$ become the limit of $L_{\mathcal{X}}$:

$$\begin{aligned} \mathcal{X}^{\leq \infty} &= !\mathcal{X} \quad \text{and} \quad p_{\infty,n} : \&\mathcal{X}^{(k)} \longrightarrow \mathcal{X}^{\leq n} = (\mathcal{X}\&\mathcal{I})^{(n)} \\ p_{\infty,n}(-, (1, x_1) \cdots (1, x_k)(2, *) \cdots (2, *)) &:= \delta(-, (k, x_1 \cdots x_k)) \end{aligned} \quad (16)$$

In (16), each element of the first set of the tuple $(\mathcal{X}\&\mathcal{I})^{(n)}$ is written $(1, x_1) \cdots (1, x_k) \underbrace{(2, *) \cdots (2, *)}_{n-k}$ with a certain $k \leq n$ such that $x_i \in X$ and $i = 1, \dots, k$.

Proof. The proof consists of the following two claims. Let $\{\tau_n\}_n$ denote a family of $\tau_n : \mathcal{Y} \longrightarrow (\mathcal{X}\&\mathcal{I})^{(n)}$. (Claim 1) A family $\{\tau_n\}_n$ becomes a cone to the sequential diagram $\{p_{n+1,n} : \mathcal{X}^{\leq n+1} \rightarrow \mathcal{X}^{\leq n}\}_n$ if and only if the following holds for all $z_1 \cdots z_n \in (X \uplus I)^{(n)}$ and $n \in \mathbb{N}$:

$$\tau_n(-, z_1 \cdots z_n) = \tau_{n+1}(-, z_1 \cdots z_n(2, *))$$

Thus, in particular, $p_{\infty,n}$ of (16) indexed by n provides a cone.

Proof of Claim 1:

$$\begin{aligned} p_{n+1,n} \circ \tau_{n+1}(-, z_1 \cdots z_n) &= \int_{(X \uplus I)^{(n+1)}} \tau_{n+1}(-, \mathcal{Y}) p_{n+1,n}(d\mathcal{Y}, z_1 \cdots z_n) \\ &= \int_{(X \uplus I)^{(n+1)}} \tau_{n+1}(-, \mathcal{Y}) \delta(d\mathcal{Y}, z_1 \cdots z_n(2, *)) \quad \text{by (15)} \\ &= \tau_{n+1}(-, z_1 \cdots z_n(2, *)) \end{aligned}$$

(Claim 2) Any cone $\{\tau_n\}_n$ factors through the cone $\{p_{\infty,n}\}_n$ uniquely by the following countable infinite product morphism τ whose codomain is $\&\mathcal{X}^{(n)}$;

$$\tau := \&\left(\mathbf{p}_l^{(k)} \circ \tau_k \right), \text{ where } \mathbf{p}_l^{(k)} : (\mathcal{X}\&\mathcal{I})^{(k)} \longrightarrow \mathcal{X}^{(k)} \quad (17)$$

Proof of Claim 2: It suffices to prove the following for any morphism h of the codomain $\&_{n \in \mathbb{N}} \mathcal{X}^{(n)}$ and any natural number n , where \mathfrak{p}_n is the n -th projection;

$$\mathfrak{p}_n \circ h = \mathfrak{p}_l^{(n)} \circ p_{\infty, n} \circ h$$

The sufficiency is because any n -th projection of the mediating τ (LHS put $h := \tau$) turns definable by $\mathfrak{p}_l^{(n)} \circ \tau_n$ (RHS put $\tau_n = p_{\infty, n} \circ \tau$).

For the proof, precompose any h to $p_{\infty, n}$;

$$\begin{aligned} & p_{\infty, n} \circ h(-, (1, x_1) \cdots (1, x_k)(2, *) \cdots (2, *)) && \text{with } (1, x_1) \cdots (1, x_k)(2, *) \cdots (2, *) \in (X \uplus I)^n \\ &= \int_{\uplus_{n \in \mathbb{N}} X^{(n)}} h(-, y) p_{\infty, n}(dy, (1, x_1) \cdots (1, x_k)(2, *) \cdots (2, *)) \\ &= \int_{\uplus_{n \in \mathbb{N}} X^{(n)}} h(-, y) \delta(dy, (k, x_1 \cdots x_k)) && \text{by (16)} \\ &= \int_{X^{(k)}} h(-, y) \delta(dy, (k, x_1 \cdots x_k)) \\ &= h(-, (k, x_1 \cdots x_k)) && \text{with } (k, x_1 \cdots x_k) \in \uplus_{n \in \mathbb{N}} X^{(n)} \end{aligned}$$

Taking $k = n$ in the above, the first line equates using $\mathfrak{p}_l^{(n)}$: $p_{\infty, n} \circ h(-, (1, x_1) \cdots (1, x_n)) = \mathfrak{p}_l^{(n)} \circ p_{\infty, n} \circ h(-, x_1 \cdots x_n)$. Hence we have the following for all $x_1 \cdots x_n \in X^{(n)}$, which is the assertion:

$$\begin{aligned} \mathfrak{p}_l^{(n)} \circ p_{\infty, n} \circ h(-, x_1 \cdots x_n) &= h(-, (n, x_1 \cdots x_n)) \\ &= p_n \circ h(-, x_1 \cdots x_n) \quad \text{by } n\text{-th projection} \end{aligned}$$

See the diagram for the construction of the mediating τ :

$$\begin{array}{ccc} \mathcal{X}^{(n)} & \xleftarrow{\mathfrak{p}_n} & \mathcal{X}^{(n+1)} \dots \\ \uparrow \mathfrak{p}_l^{(n)} & & \uparrow \mathfrak{p}_l^{(n+1)} \\ (\mathcal{X} \& \mathcal{I})^{(n)} & \xleftarrow{p^{n+1, n}} & (\mathcal{X} \& \mathcal{I})^{(n+1)} \dots \\ & & \uparrow \mathfrak{p}_l^{(k)} \\ & & \&_{k \in \mathbb{N}} \mathcal{X}^{(k)} \end{array}$$

$\xleftarrow{\tau_n} \quad \xleftarrow{\tau_{n+1}} \quad \uparrow \tau = \&_{k \in \mathbb{N}} (\mathfrak{p}_l^{(k)} \circ \tau_k)$

Directly by the definition (17), $\mathfrak{p}_l^{(n)} \circ \tau_n = \mathfrak{p}_n \circ \tau$. Hence by the universality of the limit,

$$\mathfrak{p}_l^{(n)} \circ p_{\infty, n} = \mathfrak{p}_n \tag{18}$$

□

The action of the limit of Definition 1.2 is described concretely in TsK^{op} :

Proposition 3.6 (morphisms $\kappa^{(n)}$ and $\kappa^{\leq \infty}$). *Let $\kappa : \mathcal{Y} \rightarrow \mathcal{X}$ in TsK^{op} .*

(i) *The morphism $\kappa^{(n)} : \mathcal{Y}^{(n)} \rightarrow \mathcal{X}^{(n)}$ of (14) is described explicitly as follows for $x_1 \cdots x_n \in X^{(n)}$:*

$$\kappa^{(n)}(-, x_1 \cdots x_n) := \kappa^{\otimes n}(F^{-1}(-), (x_1, \dots, x_n))$$

This in particular stipulates $(\kappa \& \mathcal{I})^{(n)} = \kappa^{\leq n}$, remind Definition 1.2.

(ii) *The unique morphisms $\kappa^{\leq \infty} : \mathcal{Y}^{\leq \infty} \rightarrow \mathcal{X}^{\leq \infty}$ is explicitly described as follows, where \mathfrak{p}_n is the n -th projection of the product:*

$$\kappa^{\leq \infty} = \&_{n \in \mathbb{N}} (\kappa^{(n)} \circ \mathfrak{p}_n)$$

Note $\kappa^{(n)} \circ \mathbf{p}_n : \mathcal{Y}^{\leq \infty} := \&_{k \in \mathbb{N}} \mathcal{Y}^{(k)} \xrightarrow{\mathbf{p}_n} \mathcal{Y}^{(n)} \xrightarrow{\kappa^{(n)}} \mathcal{X}^{(n)}$

Proof. (i) The explicit definition is derived via (13);

$$\begin{aligned} \kappa^{\otimes n} \circ \mathbf{eq}_{\mathcal{Y}}(-, (x_1, \dots, x_n)) &= \int_{Y^{\otimes n}} \mathbf{eq}_{\mathcal{Y}}(-, (y_1, \dots, y_n)) \kappa^{\otimes n}(d(y_1, \dots, y_n), (x_1, \dots, x_n)) \\ &= \int_{Y^{\otimes n}} \delta(-, y_1 \cdots y_n) \kappa^{\otimes n}(d(y_1, \dots, y_n), (x_1, \dots, x_n)) \\ &= \int_{Y^{\otimes n}} \delta(F^{-1}(-), (y_1, \dots, y_n)) \kappa^{\otimes n}(d(y_1, \dots, y_n), (x_1, \dots, x_n)) \\ &= \kappa^{\otimes n}(F^{-1}(-), (x_1, \dots, x_n)) \end{aligned}$$

In the 3rd line, $\delta(F^{-1}(-), (y_1, \dots, y_n))$ replaces $\delta(-, y_1 \cdots y_n)$ equivalently.

(ii) $\kappa^{\leq \infty}$ by Definition 1.2 is the unique factorisation of the cone $\{(\kappa \cdot \mathbf{p}_l \& \mathcal{I} \cdot \mathbf{p}_r)^{(n)} \circ \mathbf{p}_n\}_n$. Hence by (17), $\kappa^{\leq \infty}$ is

$$\&_{n \in \mathbb{N}} (\mathbf{p}_l^{(n)} \circ ((\kappa \cdot \mathbf{p}_l \& \mathcal{I} \cdot \mathbf{p}_r)^{(n)} \circ \mathbf{p}_n)) = \&_{n \in \mathbb{N}} (\kappa^{(n)} \circ \mathbf{p}_n)$$

The equality is by the commutativity of the following diagram, in which the right square commutes by the functoriality of $(-)^{(n)}$ and by $\mathbf{p}_l \circ (\kappa \cdot \mathbf{p}_l \& \mathcal{I} \cdot \mathbf{p}_r) = \kappa \circ \mathbf{p}_l$, and the left triangle does by (18).

$$\begin{array}{ccc} & \mathcal{Y}^{(n)} & \xrightarrow{\kappa^{(n)}} & \mathcal{X}^{(n)} \\ & \nearrow \mathbf{p}_n & & \uparrow \mathbf{p}_l^{(n)} \\ \mathcal{Y} & \xrightarrow{p_{\infty, n}} & (\mathcal{Y} \& \mathcal{I})^{(n)} & \xrightarrow{(\kappa \cdot \mathbf{p}_l \& \mathcal{I} \cdot \mathbf{p}_r)^{(n)}} & (\mathcal{X} \& \mathcal{I})^{(n)} \end{array}$$

□

In TsK^{op} , Fubini-Tobelli Theorem guarantees the distribution properties for the free exponential.

Proposition 3.7 (distribution of \otimes over the equalisers and the limits in TsK^{op}). *For any measurable space \mathcal{Z} ;*

- $\mathcal{X}^{(n)} \otimes \mathcal{Z}$ becomes the equaliser of $(n!$ -symmetries) $\otimes \mathcal{Y}$ of $\mathcal{X}^{\otimes n} \otimes \mathcal{Z}$. Thus in particular, the monoidal product distributes over the equaliser $E_{\mathcal{X}}$.
- The monoidal product distributes over the limit $L_{\mathcal{X}}$.

Proof. The direct generalisation of the respective proofs of Proposition 3.2 and of Theorem 3.5 by replacing $\int_{(-)}$ with $\int_{(-) \otimes \mathcal{Z}} \stackrel{FT}{=} \int_{(-)} \int_{\mathcal{Z}}$.

(first assertion) The generalisation yields

$$(\mathbf{eq}_{\mathcal{X}} \otimes \mathcal{Z}) \circ \tau(-, (k, ((x_1, \dots, x_k), z))) = \tau(-, (k, (x_1 \cdots x_k, z))).$$

(second assertion)

(Claim 1) A family $\{\tau_n : \mathcal{Y} \rightarrow (\mathcal{X} \& \mathcal{I})^{(n)} \otimes \mathcal{Z}\}_n$ becomes a cone to the diagram $\{p_{n, n+1} \otimes \mathcal{Z} : \mathcal{X}^{\leq n+1} \otimes \mathcal{Z} \rightarrow \mathcal{X}^{\leq n} \otimes \mathcal{Z}\}$ if and only if the following holds for all $z_1 \cdots z_n \in (X \uplus I)^{(n)}$ and $z \in \mathcal{Z}$ with any natural number n :

$$\tau_n(-, (z_1 \cdots z_n, z)) = \tau_{n+1}(-, (z_1 \cdots z_n (2, *), z))$$

(Claim 2) First, by the generalisation, any cone $\{\tau_n\}_n$ to the sequential diagram $\{p_{n, n+1} \otimes \mathcal{Z}\}_n$ determines the following universal morphism:

$$\&_{k \in \mathbb{N}} (\mathbf{p}_l^{(k)} \circ \tau_k) : \mathcal{Y} \rightarrow \&_{k \in \mathbb{N}} (\mathcal{X}^{(k)} \otimes \mathcal{Z}) \quad (19)$$

Second, generally in the presence of the tensor factor \mathcal{Z} , the following two measurable spaces are isomorphic; $\&_{k \in \mathbb{N}} (\mathcal{X}^{(k)} \otimes \mathcal{Z}) \cong (\&_{k \in \mathbb{N}} \mathcal{X}^{(k)}) \otimes \mathcal{Z}$ under mapping $(k, (x_1 \cdots x_k, z))$ to $((k, (x_1 \cdots x_k)), z)$ for any natural numbers k . Hence, (19) gives the meditating morphism of the cone. □

By Propositions 3.2 and 3.7 and Theorem 3.5;

Theorem 3.8. *The monoidal category \mathbf{TsK}^{op} with products has the free exponential.*

4. The Orthogonality Category $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$

This section presents a reciprocal orthogonality in \mathbf{TsK}^{op} in terms of Lebesgue integral. Accordingly, a general categorical construction of Section 1 applies to the free exponential of Section 3 in order to obtain the free exponential in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$. In Section 4.3, the condition of Section 1.3 on distribution of monoidal product over the equaliser is shown to hold using the uniform convergence theorem for the integral. Section 4.4 concretely characterises the equaliser and their limit in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$. Section 4.5 shows that our free exponential is considered a continuous extension of the exponential of \mathbf{Pcoh} . In the sequel, the object J for the orthogonality (Definition 1.3) is the monoidal unit \mathcal{I} .

4.1. Reciprocal Orthogonality between Measures and Measurable Functions in \mathbf{TsK}^{op}

Definition 4.1 (inner product). For a measure $\mu \in \mathbf{TKer}^{\text{op}}(\mathcal{X}, \mathcal{I})$ and a measurable function $f \in \mathbf{TKer}^{\text{op}}(\mathcal{I}, \mathcal{X})$, we define

$$\langle f | \mu \rangle_{\mathcal{X}} := \int_{\mathcal{X}} f d\mu$$

Then the two operators in Definition 2.9 become characterised as follows:

Proposition 4.2 (reciprocity between κ^* and κ_*). *In $\mathbf{TsKer}^{\text{op}}$, for any measure $\mu : \mathcal{X} \rightarrow \mathcal{I}$, any measurable function $f : \mathcal{I} \rightarrow \mathcal{Y}$ and any transition kernel $\kappa : \mathcal{Y} \rightarrow \mathcal{X}$,*

$$\langle f | \kappa_* \mu \rangle_{\mathcal{Y}} = \langle \kappa^* f | \mu \rangle_{\mathcal{X}}.$$

$$\text{Equivalently, } \langle f | \mu \circ \kappa \rangle_{\mathcal{Y}} = \langle \kappa \circ f | \mu \rangle_{\mathcal{X}}$$

Proof. The following starts from LHS and ends with RHS of the assertion, using Fubini-Tonelli (Proposition 2.20): $\int_{\mathcal{Y}} f(y) (\kappa_* \mu)(dy) = \int_{\mathcal{Y}} f(y) \int_{\mathcal{X}} \kappa(dy, x) \mu(dx) = \int_{\mathcal{X}} \mu(dx) \int_{\mathcal{Y}} f(y) \kappa(dy, x) = \int_{\mathcal{X}} (\kappa^* f)(x) \mu(dx)$ \square

Proposition 4.3 (orthogonality in terms of integral). *For a measurable function $f \in \mathbf{TsKer}^{\text{op}}(\mathcal{I}, \mathcal{X})$ and a measure $\mu \in \mathbf{TsKer}^{\text{op}}(\mathcal{X}, \mathcal{I})$, the relation $\perp_{\mathcal{X}} \subset \mathbf{TsKer}^{\text{op}}(\mathcal{I}, \mathcal{X}) \times \mathbf{TsKer}^{\text{op}}(\mathcal{X}, \mathcal{I})$ is defined*

$$f \perp_{\mathcal{X}} \mu \quad \text{if and only if} \quad \langle f | \mu \rangle_{\mathcal{X}} \leq 1$$

Then this gives a reciprocal orthogonality.

Proof. By Proposition 4.2. \square

4.2. Push Forward Integral as Reciprocity

The reciprocity of orthogonality also emerges explicitly when category theoretically reformulated the measure-theoretic variable change along the push forward measure (Definition 2.7). This subsection shows that the reciprocity characterises integrals over the push forward measures along the two morphisms eq and $p_{\infty, n}$. The two are the main ingredient for the equaliser and the limit construction in Definition 1.1.

Remark 4.4 (variable change along the push forward $\mathcal{X}^{\otimes n} \rightarrow \mathcal{X}^{(n)}$ as reciprocity for eq). The variable change property along the pushforward $F : \mathcal{X}^{\otimes n} \rightarrow \mathcal{X}^{(n)}$ in Definition 3.4 is reformulated using the categorical composition and precomposition of eq in \mathbf{TsK}^{op} :

$$\int_{\mathcal{X}^{(n)}} f d(\mu \circ \text{eq}) = \int_{\mathcal{X}^{\otimes n}} (\text{eq} \circ f) d\mu \quad (20)$$

The reformulation is obtained because $\mu \circ F^{-1}$ (resp. $f \circ F$) in (7) in Definition 2.7 is $\mu \circ \text{eq}$ (resp. $\text{eq} \circ f$) when any measure μ (resp. measurable function f) is seen as a morphism $\mathcal{X}^{\otimes n} \rightarrow \mathcal{I}$ (resp. $\mathcal{I} \rightarrow \mathcal{X}^{\otimes n}$) in

TsK^{op} .

(20) is obviously a reciprocity of the orthogonality for TsK^{op} because it stipulates

$$\langle f | \mu \circ \text{eq} \rangle_{X^{(n)}} = \langle \text{eq} \circ f | \mu \rangle_{X^{\otimes n}}$$

Alternatively putting $f = \text{eq} \setminus g$ with a measurable function g on $\mathcal{X}^{\otimes n}$,

$$\int_{X^{(n)}} (\text{eq} \setminus g) d(\mu \circ \text{eq}) = \int_{X^{\otimes n}} g d\mu \quad (21)$$

For the morphism $p_{\infty, n}$, we start with characterising composition and precomposition of $p_{\infty, n}$ in TsK^{op} in terms of a measurable function $G_{n, \infty}$:

Lemma 4.5 (composition and precomposition of $p_{\infty, n}$ in terms of a measurable function $G_{n, \infty}$).
A function $G_{n, \infty} : \mathcal{A}^{\leq n} \rightarrow \mathcal{A}^{\leq \infty}$ is defined by

$$\begin{aligned} G_{n, \infty} : \mathcal{A}^{\leq n} &\rightarrow \bigsqcup_k A^{(k)} \\ (1, a_1) \cdots (1, a_k)(2, *) \cdots (2, *) &\mapsto (k, a_1 \cdots a_k) \end{aligned} \quad (22)$$

The function $G_{n, \infty}$ is one to one (but not surjective) and $(\mathcal{A}^{\leq n}, \mathcal{A}^{\leq \infty})$ -measurable.

Then $G_{n, \infty}$ characterises the composition (resp. precomposition) of $p_{\infty, n}$ in TsK^{op} to a measurable function (resp. to a measure) as follows:

(i) For any measurable function $f : \mathcal{I} \rightarrow \mathcal{A}^{\leq \infty} \otimes \mathcal{B}$ in TsK^{op} ,

$$((p_{\infty, n} \otimes \mathcal{B}) \circ f)(I, \mathbb{Y}) = f(I, (G_{n, \infty} \times B)(\mathbb{Y})) \quad \text{for } \mathbb{Y} \in A^{\leq n} \times B$$

(ii) (precomposition of $p_{\infty, n}$ as a pushforward)

For any measure $\mu : \mathcal{A}^{\leq n} \otimes \mathcal{B} \rightarrow \mathcal{I}$ in TsK^{op} ,

$$\mu \circ (p_{\infty, n} \otimes \mathcal{B}) = \mu \circ (G_{n, \infty} \times B)^{-1}$$

That is, the composition $\mu \circ (p_{\infty, n} \otimes \mathcal{B})$ is the push forward measure of μ along the measurable function $G_{n, \infty} \times B$.

Proof. (i) Direct by the definition (16) because

$$p_{\infty, n}(-, \mathbb{Y}) = \delta(-, G_{n, \infty}(\mathbb{Y})).$$

$$\begin{aligned} (ii) \quad &(\mu \circ (p_{\infty, n} \otimes \mathcal{B}))(-, *) \\ &= \int_{A^{\leq n} \times B} (p_{\infty, n} \otimes \mathcal{B})(-, (\mathbb{x}, y)) \mu(d(\mathbb{x}, y), *) \\ &= \int_{A^{\leq n} \times B} \delta(-, (G_{n, \infty} \times B)(\mathbb{x}, y)) \mu(d(\mathbb{x}, y), *) \\ &= \int_{A^{\leq n} \times B} \delta(-, (G_{n, \infty}(\mathbb{x}), y)) \mu(d(\mathbb{x}, y), *) \\ &= \int_{A^{\leq \infty} \times B} \delta(-, (\mathbb{x}', y)) \mu((G_{n, \infty} \times B)^{-1}(d(\mathbb{x}', y)), *) \quad \text{by variable change along } G_{n, \infty} \times B \\ &= \mu((G_{n, \infty} \times B)^{-1}(-), *) \end{aligned}$$

□

Lemma 4.5 relates a reciprocity for $p_{\infty, n}$ with a variable change property:

Corollary 4.6 (variable change along $\mathcal{A}^{\leq n} \otimes \mathcal{B} \rightarrow \mathcal{A}^{\leq \infty} \otimes \mathcal{B}$ as reciprocity for $p_{\infty, n} \otimes \mathcal{B}$). For any measurable function $f : \mathcal{I} \rightarrow \mathcal{A}^{\leq \infty} \otimes \mathcal{B}$ and any measure $\mu : \mathcal{A}^{\leq n} \otimes \mathcal{B} \rightarrow \mathcal{I}$ in \mathbf{Tsk}^{op} ,

$$\int_{\mathcal{A}^{\leq \infty} \times B} f d(\mu \circ (p_{\infty, n} \otimes \mathcal{B})) = \int_{\mathcal{A}^{\leq n} \times B} ((p_{\infty, n} \otimes \mathcal{B}) \circ f) d\mu$$

Proof. The assertion is the variable change property (Definition 2.7) along the push forward $G_{n, \infty} \times B$ but instantiated with (i) and (ii) of Lemma 4.5:

$$\int_{\mathcal{A}^{\leq \infty} \times B} f(\mathfrak{y}) (\mu \circ (G_{n, \infty} \times B)^{-1})(d\mathfrak{y}) = \int_{\mathcal{A}^{\leq n} \times B} f((G_{n, \infty} \times B)(\mathfrak{y}')) \mu(d\mathfrak{y}')$$

□

4.3. Distribution of Monoidal Product over Limit in $\mathbf{O}_{\mathcal{I}}(\mathbf{Tsk}^{\text{op}})$

This subsection is concerned with showing the distribution of the tensor product over the limit $L_{\mathbb{A}}$ in $\mathbf{O}_{\mathcal{I}}(\mathcal{C})$ when $\mathcal{C} = \mathbf{Tsk}^{\text{op}}$. Using the monotone convergence theorem, Lebesgue integrals over the limit measurable space are shown approximated by sequence of integrals over the push forward measurable spaces along $p_{\infty, n} \otimes \mathcal{B}$ s (Proposition 4.8). For the approximation, the reciprocity of $p_{\infty, n}$ in Corollary 4.6 above plays a crucial role. The approximation leads to the satisfaction of the distribution condition (Theorem 4.9).

Definition 4.7 (sequence of measures ν_n on $\mathcal{A}^{\leq n} \otimes \mathcal{B}$ for a measure ν on $\mathcal{A}^{\leq \infty} \otimes \mathcal{B}$). For any measure $\nu : \mathcal{A}^{\leq \infty} \otimes \mathcal{B} \rightarrow \mathcal{I}$ in \mathbf{Tsk}^{op} and a natural number n , a measure ν_n on $\mathcal{A}^{\leq n} \otimes \mathcal{B}$ is defined using $G_{n, \infty}$ of (22);

$$\nu_n : \mathcal{A}^{\leq n} \otimes \mathcal{B} \rightarrow \mathcal{I} \quad X \times Y \mapsto \nu(G_{n, \infty}(X) \times Y)$$

for every rectangle $X \times Y$ with measurable sets $X \in \mathcal{A}^{\leq n}$ and $Y \in \mathcal{B}$.

Note the measure $\nu_n \circ (p_{\infty, n} \otimes \mathcal{B})$, which by Lemma 4.5 (ii), is the push forward measure $\nu_n \circ (G_{n, \infty} \times B)^{-1}$ of ν_n along $G_{n, \infty} \times B$. Thus $\nu_n \circ (p_{\infty, n} \otimes \mathcal{B})$ gives the restriction of the measure ν to the sub measurable space $sub_n(\mathcal{A}^{\leq \infty}) \otimes \mathcal{B}$ of $\mathcal{A}^{\leq \infty} \otimes \mathcal{B}$, where

$$sub_n(\mathcal{A}^{\leq \infty}) := \left(\biguplus_{k \in \mathbb{N}} A^{(k)}, \left(\bigotimes_{k \leq n} A^{(k)} \right) \& \left(\bigotimes_{n < k} \{\emptyset, A^{(k)}\} \right) \right)$$

That is, sub_n collapses the k -th component measurable space $A^{(k)}$ to the rudest one for $n < k$.

Proposition 4.8 (Approximation of ν on $\mathcal{A}^{\leq \infty} \otimes \mathcal{B}$ in terms of the sequence $\{\nu_n \circ (p_{\infty, n} \otimes \mathcal{B})\}_{n \in \mathbb{N}}$ with $n \rightarrow \infty$). For any measurable function $f : \mathcal{I} \rightarrow \mathcal{A}^{\leq \infty} \otimes \mathcal{B}$ in \mathbf{Tsk}^{op} ,

$$\begin{aligned} \int_{\mathcal{A}^{\leq \infty} \times B} f d\nu &= \lim_{n \rightarrow \infty} \int_{\mathcal{A}^{\leq \infty} \times B} f d(\nu_n \circ (p_{\infty, n} \otimes \mathcal{B})) \\ &\geq \int_{\mathcal{A}^{\leq \infty} \times B} f d(\nu_n \circ (p_{\infty, n} \otimes \mathcal{B})) \end{aligned} \quad (23)$$

Proof. We prove Equation (23) using the monotone convergence theorem (as the inequality is direct for any n by the definition ν_n). Given f , we define an increasing sequence $0 \leq f_0 \leq f_1 \leq \dots \leq f_n \leq \dots$ of measurable functions $f_n : \mathcal{I} \rightarrow \mathcal{A}^{\leq \infty} \otimes \mathcal{B}$ by

$$f_n(((k, a_1 \dots a_k), b)) := \begin{cases} 0 & \text{for } k > n \\ f_n(((k, a_1 \dots a_k), b)) & \text{for } k \leq n \end{cases}$$

This yields

$$(p_{\infty, n} \otimes \mathcal{B}) \circ f = (p_{\infty, n} \otimes \mathcal{B}) \circ f_n \quad \text{for any } n \quad (24)$$

Obviously from the definition,

$$\lim_{n \rightarrow \infty} f_n = f$$

The following first equation is by the definition ν_m of Definition 4.7,

$$\begin{aligned} \int_{A \leq \infty \times B} f_n d\nu &= \int_{A \leq \infty \times B} f_n d(\nu_n \circ (p_{\infty, n} \otimes \mathcal{B})) \\ &= \int_{A \leq n \times B} (p_{\infty, n} \otimes \mathcal{B}) \circ f_n d\nu_n && \text{by Cor 4.6} \\ &= \int_{A \leq n \times B} (p_{\infty, n} \otimes \mathcal{B}) \circ f d\nu_n && \text{by (24)} \\ &= \int_{A \leq \infty \times B} f d(\nu_n \circ (p_{\infty, n} \otimes \mathcal{B})) && \text{by Cor 4.6} \end{aligned}$$

Taking the limit $n \rightarrow \infty$ yields the assertion with a bypass of the monotone convergence theorem commuting the limit and the integral:

$$\int_{A \leq \infty \times B} f d\nu = \lim_{n \rightarrow \infty} \int_{A \leq \infty \times B} f_n d\nu = \lim_{n \rightarrow \infty} \int_{A \leq \infty \times B} f d(\nu_n \circ (p_{\infty, n} \otimes \mathcal{B}))$$

□

Finally with Proposition 4.8, we are ready to prove the goal of this subsection.

Theorem 4.9 (\otimes distributes over $L_{\mathbb{A}}$ in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$). *The condition (6) holds for all objects \mathbb{A} and \mathbb{B} in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$. That is, the distribution of the monoidal product over $L_{\mathbb{A}}$ is retained in the orthogonal category.*

Proof. By Proposition 1.24, it suffices to show the condition (6). Take an arbitrary $\nu \in ((\mathbb{A}^{\leq \infty} \otimes \mathbb{B})_p)^\circ$ and consider ν_n of Definition 4.7. Then $\nu_n \in ((\mathbb{A}^{\leq n})_p \otimes \mathbb{B}_p)^\circ$ for all n by the followings implication (25).

$$\begin{aligned} \nu \in ((\mathbb{A}^{\leq \infty} \otimes \mathbb{B})_p)^\circ &= ((\mathbb{A}^{\leq \infty})_p \otimes \mathbb{B}_p)^{\circ \circ \circ} = ((\mathbb{A}^{\leq \infty})_p \otimes \mathbb{B}_p)^\circ \\ \implies \forall n \in \mathbb{N} \quad \nu_n \in ((\mathbb{A}^{\leq n} \otimes \mathbb{B})_p)^\circ &= ((\mathbb{A}^{\leq n})_p \otimes \mathbb{B}_p)^{\circ \circ \circ} = ((\mathbb{A}^{\leq n})_p \otimes \mathbb{B}_p)^\circ \end{aligned} \quad (25)$$

(demonstration of (25)) By the definition of $(\mathbb{A}^{\leq \infty})_p$, an arbitrary element of $(\mathbb{A}^{\leq n})_p \otimes \mathbb{B}_p$ is written by $(p_{\infty, n} \circ x) \otimes y = (p_{\infty, n} \otimes B)(x, y)$ with $x \in (\mathbb{A}^{\leq \infty})_p$ and $y \in \mathbb{B}_p$. Then by Ineqn (23), $\nu_n \circ (p_{\infty, n} \otimes \mathcal{B}) \in ((\mathbb{A}^{\leq \infty})_p \otimes \mathbb{B}_p)^\circ$ so that the orthogonality holds $(x, y) \perp_{\mathcal{A} \leq \infty \otimes \mathcal{B}} \nu_n \circ (p_{\infty, n} \otimes \mathcal{B})$, which is equivalent by reciprocity to $(p_{\infty, n} \otimes \mathcal{B}) \circ (x, y) \perp_{\mathcal{A} \leq n \otimes \mathcal{B}} \nu_n$.

(end of demonstration of (25))

Thus the premise of the condition (6) with $u = f$ yields the following inequality,

$$\begin{aligned} 1 &\geq \int_{A \leq n \times B} (p_{\infty, n} \otimes \mathcal{B}) \circ f d\nu_n \\ &= \int_{A \leq \infty \times B} f d(\nu_n \circ (p_{\infty, n} \otimes \mathcal{B})) && \text{by Cor 4.6} \end{aligned}$$

Making n tend to ∞ , we have by Eqn. (23) of Prop 4.8,

$$1 \geq \int_{A \leq \infty \times B} f d\nu,$$

which is the conclusion of the condition (6). □

As a corollary of Theorem 4.9,

Corollary 4.10. $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$ has the free exponential whose forgetful image by $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}}) \rightarrow \mathbf{TsK}^{\text{op}}$ is the free exponential of Theorem 3.8.

Proof. By Theorem 1.25. □

4.4. Characterising Equalisers $\mathbb{A}^{\leq n}$ and Limit $\mathbb{A}^{\leq \infty}$ in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$

This subsection is concerned with a concrete representation of the free exponential of $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$ guaranteed abstractly in Corollary 4.10 above. This subsection starts with characterising the equaliser $\mathbb{A}^{\leq n}$ in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$ in terms of generators for double orthogonal (Proposition 4.16). The barycenter construction by Crubillé et al [6] is directly applied to our categorical framework \mathbf{TsK}^{op} . Using the characterisation globally over natural numbers n , a characterisation of the limit $\mathbb{A}^{\leq \infty}$ in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$ is obtained in Theorem 4.19. The limit characterisation directly leads to the coincidence of the free exponential with the exponential of the linear comonad in our preceding study [18].

Definition 4.11 (barycenter $s_n(g)$ as composing an endomorphism s_n on $\mathcal{A}^{\otimes n}$). In the category \mathbf{TsK}^{op} , the n -th barycenter s_n is defined as the following endomorphism on $\mathcal{A}^{\otimes n}$:

$$s_n(-, (a_1, \dots, a_n)) = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \delta(-, (a_{\sigma(1)}, \dots, a_{\sigma(n)}))$$

For a measurable function $g : \mathcal{I} \rightarrow \mathcal{A}^{\otimes n}$ in \mathbf{TsK}^{op} , its barycenter $s_n(g) : \mathcal{I} \rightarrow \mathcal{A}^{\otimes n}$ is defined to be the categorical composition $s_n \circ g$ in \mathbf{TsK}^{op} :

$$s_n(g) := \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \sigma(g) \quad \text{where } \sigma(g)(a_1, \dots, a_n) = g(a_{\sigma(1)}, \dots, a_{\sigma(n)})$$

In particular when putting $g = f_1 \otimes \dots \otimes f_n$ of the domain $\mathcal{I}^{\otimes n} \cong \mathcal{I}$ with $f_i : \mathcal{I} \rightarrow \mathcal{A}$ for $i = 1, \dots, n$;

$$s_n(f_1 \otimes \dots \otimes f_n) = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} f_{\sigma(1)} \otimes \dots \otimes f_{\sigma(n)}$$

It is direct that $(s_n(g))(a_1, \dots, a_n) = s_n \circ g(I, (a_1, \dots, a_n)) = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} g(I, (a_{\sigma(1)}, \dots, a_{\sigma(n)}))$. The barycenters characterise the invariant morphisms: $s_n(g) = g$ if and only if g equalises the $n!$ -symmetries of $\mathcal{A}^{\otimes n}$.

Lemma 4.12. For $\mathcal{C} = \mathbf{TsK}^{\text{op}}$, the following equality holds between \mathcal{C} -homsets:

$$\{s_n(g) \mid g \in (\mathbb{A}_p^{\otimes n})^{\circ\circ}\}^{\circ\circ} = \{s_n(g) \mid g \in \mathbb{A}_p^{\otimes n}\}^{\circ\circ},$$

in which $\mathbb{A}_p^{\otimes n}$ is a short for $(\mathbb{A}_p)^{\otimes n}$.

Proof. We prove (\subset) as the converse is tautological. Note LHS of the assertion is $(s_n \circ (\mathbb{A}_p^{\otimes n})^{\circ\circ})^{\circ\circ}$, while RHS is $(s_n \circ \mathbb{A}_p^{\otimes n})^{\circ\circ}$, where the composition to a set is element-wise. It suffices to show

$$s_n \circ (\mathbb{A}_p^{\otimes n})^{\circ\circ} \subset (s_n \circ \mathbb{A}_p^{\otimes n})^{\circ\circ}$$

Take an arbitrary $\nu \in (s_n \circ \mathbb{A}_p^{\otimes n})^{\circ}$, which means $\forall x \in \mathbb{A}_p^{\otimes n} \ s_n \circ x \perp_{\mathcal{A}^{\otimes n}} \nu$. By reciprocity, $x \perp_{\mathcal{A}^{\otimes n}} \nu \circ s_n$, which means $\nu \circ s_n \in (\mathbb{A}_p^{\otimes n})^{\circ}$. Hence $\forall g \in (\mathbb{A}_p^{\otimes n})^{\circ\circ} \ g \perp_{\mathcal{A}^{\otimes n}} \nu \circ s_n$, hence by reciprocity $s_n \circ g \perp_{\mathcal{A}^{\otimes n}} \nu$, which means $s_n \circ g$ belongs to $(s_n \circ \mathbb{A}_p^{\otimes n})^{\circ\circ}$. □

Thanks to Lemma 4.12, we have Proposition 4.16 characterising equalisers in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$. The proposition is obtained with the help of the following Lemmas 4.14 and 4.15, both on the double closure on homsets.

Let us remark here that the category $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$ retains the general equalisers $\mathbb{A}^{(n)}$ lifting those $A^{(n)}$ in $\mathcal{C} = \mathbf{TsK}^{\text{op}}$ of Proposition 3.2. The lifting below in Definition 4.13 is general enough to subsume the definition (3) of Proposition 1.18 when \mathbb{A} is instantiated in particular by $\mathbb{A} \& \mathbb{I}$.

Definition 4.13 (equaliser $\mathbb{A}^{(n)}$ in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$). In $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$ for every object \mathbb{A} , the following object $\mathbb{A}^{(n)}$ with eq of Proposition 3.2 becomes the equaliser of the $n!$ -symmetries of $\mathbb{A}^{\otimes n}$:

$$\begin{aligned} \mathbb{A}^{(n)} &= (A^{(n)}, (\mathbb{A}^{(n)})_p), \quad \text{where} \\ (\mathbb{A}^{(n)})_p &:= \{\text{eq} \setminus h \mid h \in (\mathbb{A}^{\otimes n})_p \text{ equalises the } n! \text{-symmetries of } \mathbb{A}^{\otimes n}\}^{\circ\circ} \end{aligned} \quad (26)$$

Lemma 4.14 ($\text{eq} \setminus$ preserves the double orthogonal $()^{\circ\circ}$ of measurable functions in \mathbf{TsK}^{op}). For any hom-set $V \subseteq \{h : \mathcal{I} \rightarrow \mathcal{A}^{\otimes n} \text{ equalises the } n! \text{-symmetries of } \mathcal{A}^{\otimes n}\}$ in \mathbf{TsK}^{op} , the morphism $\text{eq} : A^{(n)} \rightarrow \mathbb{A}^{\otimes n}$ has the following property:

$$\text{eq} \setminus V^{\circ\circ} = (\text{eq} \setminus V)^{\circ\circ}$$

Proof. Note first that the variable change for the push forward (20) says that $\mu \circ \text{eq} \in (\text{eq} \setminus V)^{\circ}$ if and only if $\mu \in V^{\circ}$.

(\subset) Take any $\text{eq} \setminus g \in LHS$ with $g \in V^{\circ\circ}$. Note any measure in $(\text{eq} \setminus V)^{\circ}$ is a push forward $\mu \circ \text{eq}$, and moreover by the first note $\mu \in V^{\circ}$. Then $\langle \text{eq} \setminus g \mid \mu \circ \text{eq} \rangle_{A^{(n)}} \stackrel{\text{by(21)}}{=} \langle g \mid \mu \rangle_{A^{\otimes n}} \leq 1$, which means $\text{eq} \setminus g \in RHS$.
(\supset) It suffices to prove that LHS of the assertion is double orthogonal. Take any $f \in LHS^{\circ\circ}$, which means $1 \geq \langle f \mid \nu \circ \text{eq} \rangle_{A^{(n)}} \stackrel{\text{by(20)}}{=} \langle \text{eq} \circ f \mid \nu \rangle_{A^{\otimes n}}$ for all $\nu \circ \text{eq} \in LHS^{\circ}$ iff $\nu \in V^{\circ\circ}$ by the first note. But $V^{\circ\circ} = V^{\circ}$, hence $\text{eq} \circ f \in V^{\circ}$, thus $f \in LHS$. \square

Lemma 4.15 (downward closedness of double orthogonal sets consisting of measurable functions). Let $\mathcal{C} = \mathbf{TsK}^{\text{op}}$, and $V^{\circ\circ} = V \subset \mathcal{C}(I, \mathcal{X})$. For any $g \in \mathcal{C}(I, \mathcal{X})$ (i.e., a measurable function g on a measurable space \mathcal{X}),

$$g \leq \exists g' \in V \implies g \in V,$$

in which the order \leq is pointwise order between measurable functions.

Proof. Obvious: $\forall \nu \in V^{\circ} \int_X g \, d\nu \leq \int_X g' \, d\nu \leq 1$. Thus $g \in V^{\circ\circ}$. \square

In what follows, when h has a domain $I^{\otimes n}$, the domain $I^{(k)}$ of $\text{eq} \setminus h$ is identified with $I \cong I^{(k)}$.

Proposition 4.16 (barycentric genetarors for $(\mathbb{A}^{(n)})_p$ and for $(\mathbb{A}^{\leq n})_p$). In $\mathbf{O}_{\mathcal{I}}(\mathcal{C})$ with $\mathcal{C} = \mathbf{TsK}^{\text{op}}$,

(i) (generators for the equaliser $\mathbb{A}^{(n)}$)

$$\begin{aligned} (\mathbb{A}^{(n)})_p &= \{\text{eq} \setminus s_n(g) \mid g \in (\mathbb{A}_p^{\otimes n})^{\circ\circ}\}^{\circ\circ} \\ &= \{\text{eq} \setminus s_n(f_1 \otimes \cdots \otimes f_n) \mid \forall i \, f_i \in \mathbb{A}_p\}^{\circ\circ}, \end{aligned}$$

where $\text{eq} : A^{(n)} \rightarrow A^{\otimes n}$.

(ii) (generators for the equaliser $\mathbb{A}^{\leq n}$)

$$\begin{aligned} (\mathbb{A}^{\leq n})_p &= \{\text{eq} \setminus s_n((f_1 \& \iota_1) \otimes \cdots \otimes (f_n \& \iota_n)) \mid \forall i \, (f_i \in \mathbb{A}_p \text{ and } \iota_i \in \{\text{Id}_I\}^{\circ\circ})\}^{\circ\circ} \\ &= \{\text{eq} \setminus s_n((f_1 \& \text{Id}_I) \otimes \cdots \otimes (f_n \& \text{Id}_I)) \mid \forall i \, f_i \in \mathbb{A}_p\}^{\circ\circ}, \end{aligned}$$

where $\text{eq} : A^{\leq n} \rightarrow (A \& I)^{\otimes n}$.

Proof. (i) By Lemmas 4.12 and 4.14.

(ii) We prove the second line as the first one is by the second equation of (i). First note that $\iota_i \leq \text{Id}_I$ as $\iota_i \in \{\text{Id}_I\}^{\circ\circ}$ is a function on $\{*\} = I$ to $[0, 1]$ (while Id_I maps $*$ to 1). Thus $f_i \otimes \iota_i \leq f_i \otimes \text{id}_I$ for all $i = 1, \dots, n$, hence $s_n((f_1 \& \iota_1) \otimes \cdots \otimes (f_n \& \iota_n)) \leq s_n((f_1 \& \text{Id}_I) \otimes \cdots \otimes (f_n \& \text{Id}_I))$. Hence by the downward closed Lemma 4.15, $\text{eq} \setminus s_n((f_1 \& \iota_1) \otimes \cdots \otimes (f_n \& \iota_n))$ belongs to RHS of the first equation of (ii), which has shown the assertion. \square

Consider Proposition 1.21 applied to $\mathcal{C} = \mathbf{TsK}^{\text{op}}$: The set $(\mathbb{A}^{\leq \infty})_p$ is specified using the description (17) of the mediating morphism x_∞ for \mathbf{TsK}^{op} ;

$$(\mathbb{A}^{\leq \infty})_p = \left\{ \begin{array}{l} x_\infty = \bigotimes_{k \in \mathbb{N}} (p_l^{(k)} \circ x_k) \mid \{x_n : \mathbb{I} \longrightarrow \mathbb{A}^{\leq n}\}_n \text{ is} \\ \text{a cone to the diagram } \{p_{n+1,n}\}_n \text{ in } \mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}}) \end{array} \right\}^{\circ\circ} \quad (27)$$

More specifically, in what follows in Theorem 4.19, $(\mathbb{A}^{\leq \infty})_p$ is characterised in direct terms of \mathbb{A}_p .

We prepare two lemmas for the theorem. In the sequel of the subsection, \bar{f} denotes $f \& \text{Id}_I$.

Lemma 4.17. *For any natural numbers $k \leq n$ and measurable functions $\bar{f}_i : \mathcal{I} \longrightarrow \mathcal{A}$ with $i = 1, \dots, n$, the following equality holds in $\mathcal{I} \longrightarrow \mathcal{A}^{\leq k}$*

$$p_{n,k} \circ \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) = \frac{(n-k)!}{n!} \text{eq} \setminus \sum_{\iota: [k] \hookrightarrow [n]} \bar{f}_{\iota(1)} \otimes \cdots \otimes \bar{f}_{\iota(k)},$$

where $p_{n,k} := p_{k+1,k} \circ \cdots \circ p_{n,n-1}$ and ι is an inclusion function with $[m]$ denoting the set $\{1, \dots, m\}$.

Proof. The following starts from *LHS* and ends with *RHS* under the instantiating at $(I, x_1 \cdots x_k)$ with $x_1 \cdots x_k \in \mathcal{A}^{\leq k}$.

$$\begin{aligned} p_{n,k} \circ \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i)(I, x_1 \cdots x_k) &= \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i)(I, x_1 \cdots x_k \overbrace{(2, *) \cdots (2, *)}^{n-k}) && \text{by (15)} \\ &= \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \bar{f}_{\sigma(1)}(I, x_1) \cdots \bar{f}_{\sigma(k)}(I, x_k) \text{Id}_I(I, *) \cdots \text{Id}_I(I, *) \\ &= \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \prod_{i=1}^k \bar{f}_{\sigma(i)}(I, x_i) = \frac{(n-k)!}{n!} \sum_{\iota: [k] \hookrightarrow [n]} \prod_{i=1}^k \bar{f}_{\iota(i)}(I, x_i) \\ &= \frac{(n-k)!}{n!} \text{eq} \setminus \sum_{\iota: [k] \hookrightarrow [n]} \bar{f}_{\iota(1)} \otimes \cdots \otimes \bar{f}_{\iota(k)}(I, x_1 \cdots x_k) \end{aligned}$$

□

Lemma 4.18 (On $p_{\infty,m}$ image of $(\mathbb{A}^{\leq \infty})_p$). *The following inclusions hold among subsets of $(\mathbb{A}^{\leq m})_p$ for any natural number m .*

$$\begin{aligned} p_{\infty,m} \circ (\mathbb{A}^{\leq \infty})_p &\subset \bigcap_{m \leq n \in \mathbb{N}} \{p_{n,m} \circ \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) \mid \forall i f_i \in \mathbb{A}_p\}^{\circ\circ} \\ &\subset \left(\bigcap_{m \leq n \in \mathbb{N}} \{p_{n,m} \circ \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) \mid \forall i f_i \in \mathbb{A}_p\} \right)^{\circ\circ}, \end{aligned}$$

where $p_{n,m} := p_{m+1,m} \circ \cdots \circ p_{n,n-1}$.

Proof. As the second inclusion is direct by $X^\circ \cap Y^\circ \subset (X \cap Y)^\circ$, we prove the first inclusion, which is derivable from the following inclusion for any natural number $n \geq m$

$$\text{LHS} \subset p_{n,m} \circ (\mathbb{A}^{\leq n})_p$$

together with the characterisation of $(\mathbb{A}^{\leq n})_p$ in Proposition 4.16 (ii) and

$$p_{n+1,n} \circ X^{\circ\circ} \subset (p_{n+1,n} \circ X)^{\circ\circ} \quad (28)$$

(demonstration of (28))

Take any $\nu \in (p_{n+1,n} \circ X)^\circ$; that is, $\forall x \in X$ $1 \geq \langle p_{n+1,n} \circ x | \nu \rangle_{A^{\leq n}} \stackrel{recipro.}{=} \langle x | \nu \circ p_{n+1,n} \rangle_{A^{\leq n+1}}$, which means $\nu \circ p_{n+1,n} \in X^\circ$. Thus $\forall y \in X^{\circ\circ}$ $1 \geq \langle y | \nu \circ p_{n+1,n} \rangle_{A^{\leq n+1}} \stackrel{recipro.}{=} \langle p_{n+1,n} \circ y | \nu \rangle_{A^{\leq n}}$. This has shown that $p_{n+1,n} \circ y$ belongs to RHS of (28). (end of demonstration (28)) \square

We are ready to prove the main technical result of this subsection.

Theorem 4.19 (characterising the limit $\mathbb{A}^{\leq \infty}$ in $\mathbf{O}_{\mathcal{I}}(\mathbf{Tsk}^{\text{op}})$). *The following three subsets of $\mathbf{Tsk}^{\text{op}}(\mathcal{I}, \mathbb{A}^{\leq \infty})$ all coincide:*

- (i) $(\mathbb{A}^{\leq \infty})_p$
- (ii) $\left(\bigcup_{n \in \mathbb{N}} \left\{ \&\mathcal{L}_{k \in \mathbb{N}} \left(\frac{f_1 + \dots + f_n}{n} \right)^{(k)} \mid \forall i f_i \in \mathbb{A}_p \right\} \right)^{\circ\circ}$
- (iii) $\left\{ \&\mathcal{L}_{k \in \mathbb{N}} g^{(k)} \mid g \in \mathbb{A}_p \right\}^{\circ\circ}$

Proof. Circular inclusions (i) \subset (ii) \subset (iii) \subset (i) are shown in the proof.

((ii) \subset (iii)) Obvious: $\frac{f_1 + \dots + f_n}{n} \in \mathbb{A}_p$ as $\forall i \langle f_i | \nu \rangle \leq 1 \Rightarrow \langle \frac{f_1 + \dots + f_n}{n} | \nu \rangle \leq 1$.

((iii) \subset (i)) Straightforward because the generators for (i) contain those for (iii). That is, $\{g^{\leq n} : \mathbb{I} \rightarrow \mathbb{A}^{\leq n}\}_n$ forms a cone whose mediating morphism belonging to (27) is described by the specific form as follows:

$$\&\mathcal{L}_{k \in \mathbb{N}} (\mathbf{p}_l^{(k)} \circ g^{\leq k}) = \&\mathcal{L}_{k \in \mathbb{N}} (\mathbf{p}_l^{(k)} \circ (g \&\mathcal{I})^{(k)}) = \&\mathcal{L}_{k \in \mathbb{N}} (\mathbf{p}_l \circ (g \&\mathcal{I}))^{(k)} = \&\mathcal{L}_{k \in \mathbb{N}} g^{(k)}.$$

((i) \subset (ii)) Since both (i) and (ii) are double orthogonal homsets in \mathbf{Tsk}^{op} , it suffices to prove

$$((\mathbb{A}^{\leq \infty})_p)^\circ \supset \left(\bigcup_{n \in \mathbb{N}} \left\{ \&\mathcal{L}_{k \in \mathbb{N}} \left(\frac{f_1 + \dots + f_n}{n} \right)^{(k)} \mid \forall i f_i \in \mathbb{A}_p \right\} \right)^\circ \quad (29)$$

Take an arbitrary measure $\nu : \mathcal{A}^{\leq \infty} \rightarrow \mathcal{I}$ belonging to RHS of (29), and let ν_m be the measure $\mathcal{A}^{\leq m} \rightarrow \mathcal{I}$ in Definition 4.7, definable from ν .

$$\begin{aligned} 1 &\geq \langle \&\mathcal{L}_{k \in \mathbb{N}} \left(\frac{f_1 + \dots + f_n}{n} \right)^{(k)} | \nu \rangle_{A^{\leq \infty}} && (30) \\ &= \lim_{m \rightarrow \infty} \langle \&\mathcal{L}_{k \in \mathbb{N}} \left(\frac{f_1 + \dots + f_n}{n} \right)^{(k)} | \nu_m \circ p_{\infty, m} \rangle_{A^{\leq \infty}} && \text{by Prop 4.8} \\ &\geq \langle \&\mathcal{L}_{k \in \mathbb{N}} \left(\frac{f_1 + \dots + f_n}{n} \right)^{(k)} | \nu_m \circ p_{\infty, m} \rangle_{A^{\leq \infty}} && \text{by Prop 4.8} \\ &= \langle p_{\infty, m} \circ \&\mathcal{L}_{k \in \mathbb{N}} \left(\frac{f_1 + \dots + f_n}{n} \right)^{(k)} | \nu_m \rangle_{A^{\leq m}} && \text{reciprocity} \\ &= \langle \left(\frac{\bar{f}_1 + \dots + \bar{f}_n}{n} \right)^{(m)} | \nu_m \rangle_{A^{\leq m}} && (31) \end{aligned}$$

The last equality uses

$$p_{\infty, m} \circ \&\mathcal{L}_{k \in \mathbb{N}} \left(\frac{f_1 + \dots + f_n}{n} \right)^{(k)} = \left(\frac{f_1 + \dots + f_n}{n} \right)^{\leq m} = \left(\frac{\bar{f}_1 + \dots + \bar{f}_n}{n} \right)^{(m)} = \left(\frac{\bar{f}_1 + \dots + \bar{f}_n}{n} \right)^{(m)}$$

Take any generator $x_\infty \in (\mathbb{A}^{\leq \infty})_p$ of (4) of Lemma 1.21 and consider the following;

$$\begin{aligned} \langle x_\infty | \nu \rangle_{A^{\leq \infty}} &= \lim_{m \rightarrow \infty} \langle x_\infty | \nu_m \circ p_{\infty, m} \rangle_{A^{\leq \infty}} && \text{by Prop 4.8} \\ &= \lim_{m \rightarrow \infty} \langle p_{\infty, m} \circ x_\infty | \nu_m \rangle_{A^{\leq m}} && \text{by reciproc.} \end{aligned}$$

This means that $\forall m \in \mathbb{N}$ $\nu_m \in (p_{\infty, m} \circ (\mathbb{A}^{\leq \infty})_p)^\circ$ implies $\nu \in$ LHS of of (29). Thus for the assertion, it

suffices to show the following for any natural number m thanks to (the contravariant of) Lemma 4.18:

$$\nu_m \in \left(\bigcap_{m \leq n \in \mathbb{N}} \{p_{n,m} \circ \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) \mid \forall i f_i \in \mathbb{A}_p\} \right)^\circ \quad (32)$$

(demonstration of (32))

Noting the following inequality for measurable functions $\mathcal{I} \rightarrow (\mathcal{A} \& I)^{\otimes m}$ in TsK^{op} ;

$$(\bar{f}_1 + \cdots + \bar{f}_n)^{\otimes m} \geq \sum_{i: [m] \hookrightarrow [n]} \bar{f}_{i(1)} \otimes \cdots \otimes \bar{f}_{i(m)}, \quad (33)$$

we have the following inequality in $\mathcal{I} \rightarrow \mathcal{A}^{\leq m}$;

$$\begin{aligned} \frac{(n-m)!n^m}{n!} &\geq \frac{(n-m)!}{n!} \langle \text{eq} \setminus (\bar{f}_1 + \cdots + \bar{f}_n)^{\otimes m} \mid \nu_m \rangle_{\mathcal{A}^{\leq m}} && \text{by Ineqn. (31)} \\ &\geq \frac{(n-m)!}{n!} \langle \text{eq} \setminus \sum_{i: [m] \hookrightarrow [n]} \bar{f}_{i(1)} \otimes \cdots \otimes \bar{f}_{i(m)} \mid \nu_m \rangle_{\mathcal{A}^{\leq m}} && \text{by Ineqn. (33)} \\ &= \langle p_{n,m} \circ \text{eq} \setminus s_n(\bar{f}_1 \otimes \cdots \otimes \bar{f}_n) \mid \nu_m \rangle_{\mathcal{A}^{\leq m}} && \text{by Lem 4.17} \end{aligned}$$

Making n tend to ∞ ,

$$\lim_{n \rightarrow \infty} \frac{(n-m)!n^m}{n!} = \lim_{n \rightarrow \infty} \frac{n^m}{n(n-1)\cdots(n-(m-1))} = \lim_{n \rightarrow \infty} \frac{1}{1(1-\frac{1}{n})\cdots(1-\frac{m-1}{n})} = 1$$

This means (32) holds.

(end of demonstration of (32))

□

Remark 4.20 (On Theorem 4.19). Given that each generator of (iii) is a measurable function on the measurable space $\mathcal{A}^{\leq \infty} = \bigcup_{n \in \mathbb{N}} \mathcal{A}^{(n)}$, its instantiation at each point $(n, x_1 \cdots x_n)$ from the n -th componential space with $x_1 \cdots x_n \in \mathcal{A}^{(n)}$ is explicitly calculated as follows:

$$\begin{aligned} \left(\bigcup_{n \in \mathbb{N}} g^{(k)} \right) (I, (n, x_1 \cdots x_n)) &= g^{(n)}(I, x_1 \cdots x_n) \\ &= g^{\otimes n}(F^{-1}(I^{(n)}), (x_1, \dots, x_n)) \\ &= g^{\otimes n}(I^{\otimes n}, (x_1, \dots, x_n)) && \text{by } F^{-1}(I^{(n)}) = I^{\otimes n} \\ &= \prod_{i=1}^n g(I, x_i) \end{aligned}$$

The exponential constructed in the present paper coincides with that for linear exponential command independently studied in our preceding [18] (free from Melliès-Tabareau-Tasson construction).

Corollary 4.21. *The free exponential of $\mathbf{O}_{\mathcal{I}}(\text{TsK}^{\text{op}})$ coincides with the exponential structure for the linear exponential comonad of the double glueing $\mathbf{G}(\text{TsK}^{\text{op}})$ in [18].*

Proof. The instantiation of Remark 4.20 coincides with the natural transformation $k : \text{TsK}^{\text{op}}(\mathcal{I}, (-)) \rightarrow \text{TsK}^{\text{op}}(\mathcal{I}, (-)_e)$ defined in Definition 4.4 of [18] making $\text{TsK}^{\text{op}}(\mathcal{I}, (-))$ linear distributive, where $(-)_e$ is the exponential action directly formulated in [18] using the exponential measurable spaces (cf. Section 3.1 of [18]). See Section 4.1 of [18] how the natural transformation k (Definition 4.4 of [18]) yields exponential structure of the double glueing. □

Remark 4.22 (Comparison to [6]'s characterisation of the limit for Pcoh). As seen in Corollary 4.21, our limit characterisation in this subsection directly coincides with the linear exponential comonad in our preceding [18]. In contrast in [6] for probabilistic coherent spaces, Crubillé et al first employed a different characterisation (from our (ii) and (iii) of Theorem 4.19) of the free exponential, and then separately showed

its equivalence to the original exponential of [8]. Accordingly in $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$, our methodology of Theorem 4.19 can accommodate their characterisation $\langle\langle f_1, \dots, f_n \rangle\rangle$ giving an inverse image of $p_{n,m} \circ \text{eq} \setminus s_n(\bar{f}_1 \otimes \dots \otimes \bar{f}_n)$ under $p_{\infty,n}$. See Appendix A.2 how to accommodate their characterisation into our Theorem 4.19.

4.5. Discretisation

Definition 4.23. The discretisation \mathbf{TsK}_{ω} of \mathbf{TsK} is the full subcategory of the countable measurable spaces whose σ -fields are generated by the singleton subsets. In the discretisation, the composition in terms of the convolution (9) collapses to the sum of products of matrices.

$$\iota \circ \kappa(x, C) = \sum_{y \in Y} \kappa(x, \{y\}) \iota(y, C) \quad (34)$$

Obviously \mathbf{TsK}_{ω} is closed under \otimes and the product.

Definition 4.24 (\mathbf{Pcoh} , \otimes , $\&$).

(inner product) $\langle x, x' \rangle := \sum_{a \in A} x_a x'_a$, for $x, x' \subseteq \mathbb{R}_+^A$ with a countable set A .

(polar) $P^{\perp} := \{x' \in \mathbb{R}_+^A \mid \forall x \in P \langle x, x' \rangle \leq 1\}$ for $P \subseteq \mathbb{R}_+^A$.

The category \mathbf{Pcoh} of the probabilistic coherent spaces is defined using the inner product and the polar: (object) $X = (|X|, \mathbf{PX})$, where $|X|$ is a countable set, $\mathbf{PX} \subseteq \mathbb{R}_+^{|X|}$ such that $\mathbf{PX}^{\perp\perp} \subseteq \mathbf{PX}$, and $0 < \sup\{x_a \mid x \in \mathbf{PX}\} < \infty$ for all $a \in |X|$.

(morphism) A morphism from X to Y is an element $u \in \mathbf{P}(X \otimes Y^{\perp})^{\perp}$, which can be seen as a matrix $(u)_{a \in |X|, b \in |Y|}$ of columns from $|X|$ and of rows from $|Y|$. Composition is the product of two matrices such that $(uv)_{a,c} = \sum_{b \in |Y|} u_{a,b} v_{b,c}$ for $u : X \rightarrow Y$ and $v : Y \rightarrow Z$.

\mathbf{Pcoh} has a monoidal product \otimes and a cartesian product $\&$:

(monoidal product \otimes)

$X \otimes Y = (|X| \times |Y|, \{x \otimes y \mid x \in \mathbf{PX} \ y \in \mathbf{PY}\}^{\perp\perp})$.

For $u \in \mathbf{Pcoh}(X_1, Y_1)$ and $v \in \mathbf{Pcoh}(X_2, Y_2)$, $u \otimes v \in \mathbf{Pcoh}(X_1 \otimes X_2, Y_1 \otimes Y_2)$ is $(u \otimes v)_{(a_1, a_2), (b_1, b_2)} = u_{a_1, b_1} v_{a_2, b_2}$.

(product $\&$)

$X_1 \& X_2 = (|X_1| \uplus |X_2|, \{x \in \mathbb{R}_+^{\uplus |X_i|} \mid \forall i \ \pi_i(x) \in \mathbf{P}(X_i)\})$,

where $\pi_i(x)_a$ is $x_{(i,a)}$. $\mathbf{P}(X_1 \& X_2)$ becomes automatically closed under the bipolar.

Proposition 4.25. *The two monoidal categories with finite products are isomorphic: $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}_{\omega}^{\text{op}}) \cong \mathbf{Pcoh}$.*

Proof. In the discretisation the symmetry arises $\text{Hom}(\mathcal{I}, \mathcal{X}) = \text{Hom}(\mathcal{X}, \mathcal{I})$, which means that measures and measurable functions become indistinguishable. Thus the both inner products are the same, hence so are the objects in the both categories. The morphisms of the two categories are identical as the matrices and their products. The definitions of monoidal and (finite) products in the both categories are the same. \square

Moreover, $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}_{\omega}^{\text{op}})$ has the free exponential which is the restriction of that for the whole category $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}^{\text{op}})$ in Corollary 4.10. The free exponential construction of \mathbf{TsK}^{op} in Section 3 and its lifting in Section 4 are a continuous generalisation of Crubillé et al's free exponential for \mathbf{Pcoh} in [6]: To be precise, directly from Proposition 4.25;

Corollary 4.26. *The exponentials of $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}_{\omega}^{\text{op}})$ and of \mathbf{Pcoh} are isomorphic.*

Proof. By the universality of the free exponential, the free exponential of the former category is isomorphic to that of the latter constructed in [6]. \square

We end this section with making the isomorphism of Corollary 4.26 explicit: The following is the exponential of $\mathbf{O}_{\mathcal{I}}(\mathbf{TsK}_{\omega}^{\text{op}})$.

(exponential on objects)

$! \mathbb{X} = (! \mathcal{X} = \&_{k \in \mathbb{N}} \mathcal{X}^{(k)}, (\mathbb{X}^{\leq \infty})_p)$. Every generator $f^{\leq \infty} \circ \epsilon_{\infty} \in (\mathbb{X}^{\leq \infty})_p$ is by Remark 4.20,

$$(f^{\leq \infty} \circ \epsilon_{\infty})(I, (n, x_1 \cdots x_n)) = \prod_{i=1}^n f(I, x_i)$$

(exponential on morphisms) For any $\kappa : \mathbb{Y} \longrightarrow \mathbb{X}$ (hence $\kappa : \mathcal{Y} \longrightarrow \mathcal{X}$), every $! \kappa : ! \mathbb{Y} \longrightarrow ! \mathbb{X}$ is by Proposition 3.6,

$$\begin{aligned}
\kappa^{\leq \infty}(\{y_1 \cdots y_n\}, x_1 \cdots x_n) &= \kappa^{(n)}(\{y_1 \cdots y_n\}, x_1 \cdots x_n) \\
&= \kappa^{\otimes n}(F^{-1}(\{y_1, \dots, y_n\}), (x_1, \dots, x_n)) \\
&= \kappa^{\otimes n}(\bigsqcup_{\sigma \in \mathfrak{S}_n / S_{\vec{x}}} \{(y_{\sigma(1)}, \dots, y_{\sigma(n)})\}, (x_1, \dots, x_n)) \\
&= \sum_{\sigma \in \mathfrak{S}_n / S_{\vec{x}}} \kappa^{\otimes n}(\{(y_{\sigma(1)}, \dots, y_{\sigma(n)})\}, (x_1, \dots, x_n)) \\
&= \sum_{\sigma \in \mathfrak{S}_n / S_{\vec{x}}} \prod_{i=1}^n \kappa(y_{\sigma(i)}, x_i)
\end{aligned}$$

where $S_{\vec{x}} := \{\sigma \in \mathfrak{S}_n \mid (x_1, \dots, x_n) = (x_{\sigma(1)}, \dots, x_{\sigma(n)})\}$ is the stabiliser subgroup fixing $\vec{x} = (x_1, \dots, x_n)$. Note that the actions σ ranging over \mathfrak{S}_n modulo $S_{\vec{x}}$ are independent of the ordering \vec{x} of \mathfrak{x} .

It is direct to see that the above exponential action coincides with that in Pcoh (see A.3) via the following isomorphisms: For any set X , each multiset of cardinality n maps to the unique element of $X^{(n)}$, which gives the isomorphisms:

$$\mathcal{M}_{\text{fin}}(X) \cong \bigotimes_{k \in \mathbb{N}} X^{(k)} \quad [x_1, \dots, x_n] \mapsto (n, x_1 \cdots x_n)$$

Future Directions

We remain it a future work how to construct a certain monoidal closed structure inside TsK , in comparison with the recent development of the higher order probabilistic programming [28], in particular with [13, 12] modelling probabilistic PCF. We will pursue to relate our transition kernels to the continuous denotational semantics of measurable cones and measurable stable functions [14], whose cartesian closed structure is shown in [5] to subsume that of Pcoh.

From a different perspective, we intend to accommodate some probabilistic feed back (or iteration) in the monoidal structure of s-finiteness, as addressed in [27] and analysed in GoI semantics for Bayesian programming [7].

References

References

- [1] H. Bauer, Probability Theory, *De Gruyter Studies in Mathematics 23*, Berlin (1996)
- [2] R. Blute, M. Hamano, P. Scott, Softness of Hypercoherences and MALL Full Completeness, *Ann. Pure and Applied Logic*, 131 (2005), pp. 1-63.
- [3] R. Blute, P. Scott, Category Theory for Linear Logicians, in [11], 1–52 (2004).
- [4] J. Borgström, A.D. Gordon, M. Greenberg, J. Margetson, J. Van Gael, Measure Transformer Semantics for Bayesian Machine Learning, *Logical Methods in Computer Science*, Vol. 9, No. 3, 2013.
- [5] R. Crubillé, Probabilistic Stable Functions on Discrete Cones are Power Series, In *33rd Annual ACM/IEEE Symposium on LICS*, 275-284 (2018)
- [6] R. Crubillé, T. Ehrhard, M. Pagani, C. Tasson, The Free Exponential Modality of Probabilistic Coherence Spaces, In *20th International Conference, Foundations of Software Science and Computation Structures*, LNCS (10203). 20-35 (2017)
- [7] U. Dal Lago, N. Hoshino, The Geometry of Bayesian Programming, in *34th Annual ACM/IEEE Symposium of Logic in Computer Science*, (2019). Long version arXiv:1904.07425 (2019)
- [8] V. Danos, T. Ehrhard, Probabilistic Coherence Spaces as a Model of Higher-order Probabilistic Computation, *Information and Computation*, 209(6), 966-991 (2011)
- [9] H. Devarajan, D. Hughes, G. Plotkin, and V. Pratt, Full completeness of the multiplicative linear logic of Chu spaces, in *14th Annual IEEE Symposium on Logic in Computer Science*, pp. 234-243 (1999)
- [10] T. Ehrhard, On Köthe Sequence Spaces and Linear Logic, *Mathematical Structures in Computer Science* 12(5): 579-623 (2002)
- [11] T. Ehrhard, J-Y. Girard, P. Ruet, P. Scott, (eds). Linear Logic in Computer Science, (*London Mathematical Society Lecture Note Series 316*), (2004)
- [12] T. Ehrhard, M. Pagani, C. Tasson, The computational meaning of probabilistic coherent spaces, in *Twenty-Sixth Annual IEEE Symposium on Logic in Computer Science*, 87-96 (2011)

- [13] T. Ehrhard, M. Pagani, C. Tasson, Making Full Abstraction for Probabilistic PCF, *Journal of the ACM*, 65(4): 23:1-23:44 (2018)
- [14] T. Ehrhard, M. Pagani, C. Tasson, Measurable cones and stable, measurable functions: a model for probabilistic higher-order programming. In *Proceedings of the ACM on Programming Languages*, No. 59 (2018)
- [15] J-Y. Girard, Linear Logic. *Theor. Comput. Sci.* 50: 1-102 (1987)
- [16] J-Y Girard, Between logic and quantics : a tract, in [11], 316-346. (2004)
- [17] M. Giry, A categorical approach to probability theory, Categorical aspects of topology and analysis, 68-85, *Lecture Notes in Math.* 915 Springer (1982)
- [18] M. Hamano, A linear Exponential Comonad in s-finite Transition Kernels and Probabilistic Coherent Spaces, submitted, arXiv:1909.07589 (2019)
- [19] M. Hyland, A. Schalk, Glueing and Orthogonality for Models of Linear Logic, *Theoretical Computer Science* 294 (1-2), 183-231 (2003)
- [20] D. Kozen, Semantics of Probabilistic Programs. *J. Comput. Syst. Sci.* 22(3): 328-350 (1981)
- [21] P-A. Melliès, Categorical semantics of linear logic, *Panoramas et Synthèses* 27, Société Mathématique de France 1-196 (2009)
- [22] P-A. Melliès, N. Tabareau and C. Tasson, An explicit formula for the free exponential modality of linear logic, *Mathematical Structures in Computer Science*, 28 (7), 1253-1286 (2018). The conference version of the same titled paper, in *36th ICALP*, LNCS (5556), 247-260 (2009)
- [23] P. Panangaden, The Category of Markov Kernels, in *First International Workshop on Probabilistic Methods in Verification*, *Electr. Notes Theor. Comput. Sci.* 22: 171-187 (1999)
- [24] P. Panangaden, Labelled Markov Processes, *World Scientific* (2009)
- [25] R. Seely, Linear logic, *-autonomous categories and cofree coalgebras, *AMS Contemporary Mathematics*, 92: 371-382, (1989)
- [26] M. Sharpe, General Theory of Markov Processes, *Pure and Applied Mathematics* Volume 133, Academic Press, 1988
- [27] S. Staton, Commutative Semantics for Probabilistic Programming, In *26th European Symposium on Programming*, LNCS (10201) 855-879 (2017)
- [28] S. Staton, H. Yang, C. Heunen, O. Kammar and F. Wood, Semantics for probabilistic programming: higher-order functions, continuous distributions, and soft constraints. In *31st Annual ACM/IEEE Symposium on LICS* 525-534, (2016)

5. Appendix

A.1. Omitted Proofs

A.1.1. Proof of Proposition 1.14

Proof. First, to check the double orthogonality, it suffices to show that $\mathbb{A}_p \& \mathbb{B}_p = U^\circ$ for certain subset $U \subseteq \mathcal{C}(A \& B, J)$. The U is given by $(\mathbb{A}_p)^\circ \mathfrak{p}_l \cup (\mathbb{B}_p)^\circ \mathfrak{p}_r$. In the following the reciprocity of \perp is used wrt the left and right projections \mathfrak{p}_l and \mathfrak{p}_r , respectively.

(\subseteq) Take any $u \& v$ from (LHS) with $u \in \mathbb{A}_p$ and $v \in \mathbb{B}_p$. Then $\forall a \in \mathbb{A}_p^\circ (\mathfrak{p}_l(u \& v) = u \perp_A a \Leftrightarrow u \& v \perp_{A \& B} a \cdot \mathfrak{p}_l)$ and $\forall b \in \mathbb{B}_p^\circ (\mathfrak{p}_r(u \& v) = u \perp_B b \Leftrightarrow u \& v \perp_{A \& B} b \cdot \mathfrak{p}_r)$.

(\supseteq) Take any ψ from (RHS). Then $\forall a \in \mathbb{A}_{cp} (\psi \perp_{A \& B} a \cdot \mathfrak{p}_l \Leftrightarrow \mathfrak{p}_l \cdot \psi \perp_A a)$, and $\forall b \in \mathbb{B}_{cp} (\psi \perp_{A \& B} b \cdot \mathfrak{p}_r \Leftrightarrow \mathfrak{p}_r \cdot \psi \perp_B b)$. Thus $\psi = \mathfrak{p}_l \cdot \psi \& \mathfrak{p}_r \cdot \psi$ belongs to (LHS).

Second, it is direct that the mediating morphisms for the product resides in $\mathbf{O}_J(\mathcal{C})$: Given $f : \mathbb{C} \rightarrow \mathbb{A}$ and $g : \mathbb{C} \rightarrow \mathbb{B}$, for any $x \in \mathbb{C}_p$ $(f \& g) \cdot x = f \cdot x \& g \cdot x \in \mathbb{A}_p \& \mathbb{B}_p$. \square

A.1.2. Proof of Proposition 2.20

Proof. Write $\mu_1 = \sum_{i \in \mathbb{N}} \mu_1^i$ and $\mu_2 = \sum_{j \in \mathbb{N}} \mu_2^j$ with finite kernels μ_1^i s and μ_2^j s, then the following is from (LHS) to (RHS):

$$\begin{aligned}
& \int_{X_2} \sum_{j \in \mathbb{N}} \mu_2^j(dx_2) \int_{X_1} f \sum_{i \in \mathbb{N}} \mu_1^i(dx_1) \\
&= \sum_{j \in \mathbb{N}} \int_{X_2} \mu_2^j(dx_2) \sum_{i \in \mathbb{N}} \int_{X_1} f \mu_1^i(dx_1) \\
&= \sum_{j \in \mathbb{N}} \sum_{i \in \mathbb{N}} \int_{X_2} \mu_2^j(dx_2) \int_{X_1} f \mu_1^i(dx_1) \\
&= \sum_{j \in \mathbb{N}} \sum_{i \in \mathbb{N}} \int_{X_1} \mu_1^i(dx_1) \int_{X_2} f \mu_2^j(dx_2) && \text{by Thm 2.17} \\
&= \sum_{i \in \mathbb{N}} \int_{X_1} \mu_1^i(dx_1) \sum_{j \in \mathbb{N}} \int_{X_2} f \mu_2^j(dx_2)
\end{aligned}$$

$$= \int_{X_1} \sum_{i \in \mathbb{N}} \mu_1^i(dx_1) \int_{X_2} \sum_{j \in \mathbb{N}} f \mu_2^j(dx_2)$$

The first and the last (resp. the second and the second last) equations are by the commutativity of integral over countable sum of measurable functions (resp. of measures) (cf. the note in Definition 2.18). \square

A.1.3. Demonstration of Eqn.(1) in Def 1.2

Proof. By definition, the left and the right squares and the outer most rectangle (with the two bent horizontal arrows) commute. This implies that any composition in the diagram starting from $A^{\leq n+1}$ and ending at $(B\&I)^{\otimes n}$ defines the same map equalising the $n!$ -symmetries $(B\&I)^{\otimes n}$. Thus there exists the unique morphism from $A^{\leq n+1}$ to $B^{\leq n}$ factoring the same map. The uniqueness implies the commutativity of the middle square, which is the assertion.

$$\begin{array}{ccccc}
& & (B\&I)^{\otimes n} \otimes p_r & & \\
& \swarrow & \xrightarrow{\text{eq}} & \searrow & \\
(B\&I)^{\otimes n} & \xleftarrow{\text{eq}} & B^{\leq n} \xleftarrow{p_{n+1}} B^{\leq n+1} & \xrightarrow{\text{eq}} & (B\&I)^{\otimes n+1} \\
\uparrow (f \cdot p_l \& I \cdot p_r)^{\otimes n} & & \uparrow f^{\leq n} \quad \exists! \quad \uparrow f^{\leq n+1} & & \uparrow (f \cdot p_l \& I \cdot p_r)^{\otimes (n+1)} \\
(A\&I)^{\otimes n} & \xleftarrow{\text{eq}} & A^{\leq n} \xleftarrow{p_{n+1, n}} A^{\leq n+1} & \xrightarrow{\text{eq}} & (A\&I)^{\otimes n+1} \\
& \swarrow & \xrightarrow{\text{eq}} & \searrow & \\
& & (A\&I)^{\otimes n} \otimes p_r & &
\end{array}$$

\square

A.1.4. Proof of Lemma 1.13

Proof. For any $h \in U^{\circ\circ}$, we shall show that $f \circ h \in f(U)^{\circ\circ}$: For the assertion, take any $v \in f(U)^{\circ}$. Then for all $u \in U$, $v \perp_B f \circ u$ iff $v \circ f \perp_B u$ by the reciprocity. This means $v \circ f \in U^{\circ}$, hence $h \perp_A v \circ f$. Thus by the reciprocity $f \circ h \perp_B v$, which is the assertion. \square

A.2. Accommodating [6]'s Characterisation $\langle\langle f_1, \dots, f_n \rangle\rangle$ into the Limit $L_{\mathbb{A}}$ of $\mathbf{O}_{\mathcal{I}}(\text{TsK}^{\text{op}})$

Definition A.1 ($\langle\langle f_1, \dots, f_n \rangle\rangle$). For an object $\mathbb{A} = (\mathcal{A}, \mathbb{A}_p)$ in $\mathbf{O}_{\mathcal{I}}(\text{TsK}^{\text{op}})$ and $f_i \in \mathbb{A}_p$ with $i = 1, \dots, n$, the morphism in TsK^{op}

$$\langle\langle f_1, \dots, f_n \rangle\rangle : \mathcal{I} \longrightarrow \bigotimes_{k \in \mathbb{N}} \mathcal{A}^{(k)} = \mathcal{A}^{\leq \infty}$$

is defined by the following instantiation at each $(k, a_1 \cdots a_k) \in \bigotimes_{k \in \mathbb{N}} \mathcal{A}^{(k)}$.

(a) For $k \leq n$

$$\begin{aligned}
\langle\langle f_1, \dots, f_n \rangle\rangle(I, (k, a_1 \cdots a_k)) &:= \frac{1}{n^k} \sum_{\iota: [k] \hookrightarrow [n]} \prod_{i=1}^k f_{\iota(i)}(I, a_i) & (35) \\
&= \frac{1}{n^k} \sum_{\iota: [k] \hookrightarrow [n]} \prod_{i=1}^k \bar{f}_{\iota(i)}(I, (1, a_i)) & \text{with } \bar{f}_i = f_i \& \mathcal{I} \\
&= \frac{1}{n^k} \sum_{\iota: [k] \hookrightarrow [n]} \text{eq} \setminus \bar{f}_{\iota(1)} \otimes \cdots \otimes \bar{f}_{\iota(k)}(I, (1, a_1) \cdots (1, a_k)) \\
&= \frac{n!}{n^k (n-k)!} p_{n,k} \circ \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i)(I, (1, a_1) \cdots (1, a_k)) & \text{by Lem 4.17}
\end{aligned}$$

That is, in particular letting the above k be n , by the definition of $p_{\infty, n}$ of (16),

$$p_{\infty, n} \circ \langle\langle f_1, \dots, f_n \rangle\rangle = \frac{n!}{n^k (n-k)!} \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) : \mathcal{I} \longrightarrow \mathcal{A}^{\leq n} \quad (36)$$

Hence for general k by composing $p_{n,k}$ to (36), as $p_{n,k} \circ p_{\infty,n} = p_{\infty,k}$,

$$p_{\infty,k} \circ \langle\langle f_1, \dots, f_n \rangle\rangle = \frac{n!}{n^k(n-k)!} p_{n,k} \circ \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) : \mathcal{I} \longrightarrow \mathcal{A}^{\leq n}$$

(b) For $k > n$, $\langle\langle f_1, \dots, f_n \rangle\rangle(I, (k, a_1 \cdots a_k)) := 0$. That is,

$$p_k \circ \langle\langle f_1, \dots, f_n \rangle\rangle := 0 : \mathcal{I} \longrightarrow \mathcal{A}^{(k)}.$$

To sum up (a) and (b), the above instantiations at every k -th projected component $\mathcal{A}^{(k)}$ yields the unique morphism from \mathcal{I} to the product $\bigotimes_{k \in \mathbb{N}} \mathcal{A}^{(k)}$;

$$\langle\langle f_1, \dots, f_n \rangle\rangle := \bigotimes_{k \leq n} \frac{n!}{n^k(n-k)!} p_i^{(k)} \circ p_{n,k} \circ \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) \ \& \ \bigotimes_{k > n} 0,$$

where p_i is the left projection $\mathcal{A} \& \mathcal{I} \longrightarrow \mathcal{A}$ and $\bar{f}_i = f_i \& \mathcal{I}$.

See the following commutative diagram for $\langle\langle f_1, \dots, f_n \rangle\rangle$ composed with the k -th projection p_k with $k \leq n$:

$$\begin{array}{ccccc}
 & & \mathcal{A}^{(k)} & \xlongequal{\quad} & \mathcal{A}^{(k)} \\
 & & \uparrow & & \uparrow \\
 & & (p_i)^{(k)} & & p_k \\
 & & \mathcal{A}^{\leq k} & \xleftarrow{p_{n,k}} & \mathcal{A}^{\leq n} & \xleftarrow{p_{\infty,n}} & \bigotimes_{k \in \mathbb{N}} \mathcal{A}^{(k)} \\
 & & & & \swarrow & & \uparrow \\
 & & & & \frac{n!}{n^k(n-k)!} \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) & & \langle\langle f_1, \dots, f_n \rangle\rangle \\
 & & & & \mathcal{I} & &
 \end{array}$$

Then $\langle\langle f_1, \dots, f_n \rangle\rangle$ in Definition A.1 is shown to provide generators for the homset $(\mathbb{A}^{\leq \infty})_p$:

Proposition A.2. *The double orthogonal homset*

$$\left(\bigcup_{n \in \mathbb{N}} \{ \langle\langle f_1, \dots, f_n \rangle\rangle \mid \forall i f_i \in \mathbb{A}_p \} \right)^{\circ\circ} \quad (37)$$

coincides with the equal subsets in Theorem 4.19.

Proof. ((37) \subset (iii)) Straightforward by Lemma 4.15 with the following inequality for measurable functions $\mathcal{I} \longrightarrow \mathcal{A}^{\leq \infty}$:

$$\langle\langle f_1, \dots, f_n \rangle\rangle \leq \bigotimes_{k \in \mathbb{N}} g^{(k)} \quad \text{with } g = \frac{f_1 + \dots + f_n}{n} \quad (38)$$

(demonstration of (38) As the inequality is point wise, we consider an instantiation at $(k, a_1 \cdots a_k)$ for any $k \leq n$; otherwise LHS=0, whereby the inequality is direct.

$$\begin{aligned}
 (\bigotimes_{k \in \mathbb{N}} g^{(k)})(I, (k, a_1 \cdots a_k)) &= g^{(k)}(I, a_1 \cdots a_k) \\
 &= g(I, a_1) \cdots g(I, a_k) = \left(\frac{1}{n}\right)^k \sum_{j=1}^n f_j(I, a_1) \cdots \sum_{j=1}^n f_j(I, a_k) \geq \text{Eqn. (35)}
 \end{aligned}$$

(end of demonstration of (38))

((i) \subset (37)) We show

$$\left(\bigcup_{n \in \mathbb{N}} \{ \langle \langle f_1, \dots, f_n \rangle \mid \forall f_i \in \mathbb{A}_p \} \right)^\circ \subset ((\mathbb{A}^{\leq \infty})_p)^\circ$$

For any measure $\nu : \mathcal{A}^{\leq \infty} \rightarrow \mathcal{I}$, let $\nu_m : \mathcal{A}^{\leq m} \rightarrow \mathcal{I}$ be the measure in Definition 4.7, definable from ν .

$$\begin{aligned} \langle \langle f_1, \dots, f_n \rangle \mid \nu \rangle_{\mathcal{A}^{\leq \infty}} &= \lim_{m \rightarrow \infty} \langle \langle f_1, \dots, f_n \rangle \mid \nu_m \circ p_{\infty, m} \rangle_{\mathcal{A}^{\leq \infty}} && \text{by Prop 4.8} \\ &\geq \langle \langle f_1, \dots, f_n \rangle \mid \nu_n \circ p_{\infty, n} \rangle_{\mathcal{A}^{\leq \infty}} && \text{by Ineqn (23)} \\ &= \langle p_{\infty, n} \circ \langle \langle f_1, \dots, f_n \rangle \mid \nu_n \rangle_{\mathcal{A}^{\leq n}} \rangle_{\mathcal{A}^{\leq \infty}} && \text{by recipro.} \\ &= \langle \frac{n!}{n^k(n-k)!} \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) \mid \nu_n \rangle_{\mathcal{A}^{\leq n}} && \text{by (36)} \\ &= \langle \text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i) \mid \frac{n!}{n^k(n-k)!} \nu_n \rangle_{\mathcal{A}^{\leq n}} \end{aligned}$$

Thus, for ν belonging to LHS, as $\text{eq} \setminus s_n(\otimes_{i=1}^n \bar{f}_i)$ s form the generators of $(\mathbb{A}^{\leq n})_p$ by Proposition 4.16 (ii),

$$\frac{n!}{n^k(n-k)!} \nu_n \in ((\mathbb{A}^{\leq n})_p)^\circ$$

Take an arbitrary $x_\infty \in \mathbb{A}^{\leq \infty}$, then we know $x_n = p_{\infty, n} \circ x_\infty \in \mathbb{A}^{\leq n}$ for any natural number n , hence

$$1 \geq \langle x_n \mid \frac{n!}{n^k(n-k)!} \nu_n \rangle_{\mathcal{A}^{\leq n}}$$

Equivalently

$$\frac{n^k(n-k)!}{n!} \geq \langle x_n \mid \nu_n \rangle_{\mathcal{A}^{\leq n}} \stackrel{\text{recipro.}}{=} \langle x_\infty \mid \nu_n \circ p_{\infty, n} \rangle_{\mathcal{A}^{\leq \infty}}$$

Making $n \rightarrow \infty$

$$\begin{aligned} 1 &= \lim_{n \rightarrow \infty} \frac{n^k(n-k)!}{n!} \geq \lim_{n \rightarrow \infty} \langle x_\infty \mid \nu_n \circ p_{\infty, n} \rangle_{\mathcal{A}^{\leq \infty}} \\ &= \langle x_\infty \mid \nu \rangle_{\mathcal{A}^{\leq \infty}}, \end{aligned} \quad \text{by Prop 4.8}$$

which means $\nu \in ((\mathbb{A}^{\leq \infty})_p)^\circ$. □

A.3. The Exponential of Pcoh [8]

On objects:

$!X = (M_{\text{fin}}(|X|), P(!X))$ is defined for $X = (|X|, \text{PX})$;

$$P(!X) := \{ u^\dagger \mid u \in P(X) \}^{\perp\perp}, \quad \text{where } u^\dagger([a_1, \dots, a_n]) = \prod_{i=1}^n u_{a_i}$$

On morphisms:

$!t \in \mathbb{R}^{M_{\text{fin}}(I) \times M_{\text{fin}}(J)}$ is defined for given $t \in \mathbb{R}^{I \times J}$:

$$(!t)_{\mu, \nu} := \sum_{\rho \in L(\mu, \nu)} \frac{\nu!}{\rho!} t^\rho = \sum_{\sigma \in \mathfrak{S}_n / S_{\bar{\mu}}} \prod_{i=1}^n t_{b_{\sigma(i)}, a_i} \quad (39)$$

In the first equation of (39), for $\mu \in M_{\text{fin}}(I)$ and $\nu \in M_{\text{fin}}(J)$,

$$L(\mu, \nu) := \left\{ r \in M_{\text{fin}}(I \times J) \mid \begin{array}{l} \forall i \in I \sum_{j \in J} r((i, j)) = \mu(i) \\ \forall j \in J \sum_{i \in I} r((i, j)) = \nu(j) \end{array} \right\}$$

and $\nu! := \prod_{j \in J} \nu(j)!$ and $\rho! := \prod_{(i, j) \in I \times J} \rho((i, j))!$

In the second equation of (39), the multisets are given explicitly by $\mu = [b_1, \dots, b_n]$ and $\nu = [a_1, \dots, a_n]$. $\vec{\mu} := (b_1, \dots, b_n)$ so that the actions σ ranging on the quotient $\mathfrak{S}_n/S_{\vec{\mu}}$ do not depend on the ordering $\vec{\mu}$ of μ .

Remark A.3. Note the action on objects of Pcoh may be seen to be subsumed in that on morphisms, because the definition $u^!$ is alternatively given as follows in terms of the morphism action (39) when $u \in \text{PX}$ is identified uniquely as the matrix in $\mathbb{R}^{\{*\} \times |X|}$.

$$u^!(\mu) := (!u)_{[* \dots *], \mu} = \prod_{i=1}^n u_{*, a_i}$$

In the first equation, $[*, \dots, *]$ is the multiset of $|1| = \{*\}$ whose cardinality is the same as that of μ . The second equation holds because the stabiliser subgroup $S_{(* \dots *)} = \mathfrak{S}_n$, when putting $\mu = [a_1, \dots, a_n]$.