

# ON THE DERIVED CATEGORY OF THE ADJOINT GRASSMANNIAN OF TYPE F

MAXIM N. SMIRNOV

ABSTRACT. We construct a full rectangular Lefschetz collection in the derived category of the adjoint Grassmannian in type  $F_4$ . This gives the first example of a full exceptional collection on this variety and also completes the proof of a conjecture due to Alexander Kuznetsov and the author that relates the structure of the derived category of coherent sheaves to the small quantum cohomology in the case of adjoint varieties for non-simply laced groups.

## 1. INTRODUCTION

Exceptional collections in derived categories of rational homogeneous spaces  $G/P$  have a long history going back to the pioneering works of Beilinson [2] and Kapranov [13]. It is conjectured that the derived category  $\mathbf{D}^b(G/P)$  always has a full exceptional collection and many cases of the conjecture are known by now. For an overview of the state of the art from a few years ago we refer to [17, §1.1]. The main advances since then are [4, 9, 10, 18], where exceptional collections were constructed on some homogeneous varieties of symplectic and orthogonal groups, and the coadjoint variety of type  $F_4$ .

For homogeneous spaces of classical types  $A_n, B_n, C_n, D_n$  and in the exceptional type of small rank  $G_2$  the story is relatively well studied (see [17] and [14, §6.4]). On the contrary, for the exceptional types  $F_4, E_6, E_7, E_8$  there were only two known instances of the conjecture: the cominuscule variety of type  $E_6$  (also called the **Cayley plane**) [8, 19] and the coadjoint variety of type  $F_4$  (this is a general hyperplane section of the Cayley plane) [4].

In this paper we deal with one more case in type  $F_4$  by constructing a full exceptional Lefschetz collection (see [15]) in the bounded derived category of coherent sheaves on the adjoint variety of type  $F_4$ . To state our result we introduce a few facts on the geometry of this variety.

Let  $G$  be a simple algebraic group of Dynkin type  $F_4$  and let  $P = P_1 \subset G$  be the maximal parabolic subgroup associated with the first vertex of its Dynkin diagram

$$\begin{array}{ccccccc} & 1 & & 2 & & 3 & & 4 \\ & \bullet & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \tag{1.1}$$

The homogeneous variety  $X = G/P$  has a natural  $G$ -equivariant projective embedding

$$X \subset \mathbb{P}(\mathfrak{g}), \tag{1.2}$$

where  $G$  acts on its Lie algebra  $\mathfrak{g}$  via the adjoint action. In this embedding  $X$  gets identified with the  $G$ -orbit of the highest root. Hence,  $X$  is called the **adjoint Grassmannian** in type  $F_4$ .

---

M.S. was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) — Projektnummer 448537907.

The variety  $X$  is a smooth projective Fano variety of Picard rank 1 with

$$\dim X = 15 \quad \text{and} \quad \omega_X = \mathcal{O}_X(-8), \quad (1.3)$$

where  $\mathcal{O}_X(1)$  is the ample generator of the Picard group of  $X$ .

Let  $\mathcal{U}^{\omega_4}$  be the irreducible  $G$ -equivariant vector bundle on  $X$  associated with the fourth vertex of the Dynkin diagram (1.1) (see Section 2 for more details). It has the following properties

$$\mathrm{rk}(\mathcal{U}^{\omega_4}) = 6 \quad \text{and} \quad \det(\mathcal{U}^{\omega_4}) = \mathcal{O}(3).$$

This bundle turns out to be exceptional and is used, together with its twists, in our exceptional collection (see Theorem 1.1 below).

Let us now consider the tangent bundle  $T_X$ . It is a  $G$ -equivariant vector bundle on  $X$ . Note, however, that  $T_X$  is not irreducible. More importantly, the vector bundle  $T_X$  is not exceptional. However, a direct computation shows that we have

$$\mathrm{Ext}^\bullet(T_X, \mathcal{O}_X) = \mathbb{k}[-1].$$

Thus, there exists a unique non-trivial  $G$ -equivariant extension

$$0 \rightarrow \mathcal{O}_X \rightarrow \tilde{T}_X \rightarrow T_X \rightarrow 0. \quad (1.4)$$

From (1.3) and (1.4) we immediately obtain

$$\mathrm{rk}(\tilde{T}_X) = 16 \quad \text{and} \quad \det(\tilde{T}_X) = \mathcal{O}_X(8).$$

The vector bundle  $\tilde{T}_X$  is exceptional and is used, together with its twists, in our exceptional collection (see Theorem 1.1 below). Let us also note here that  $\tilde{T}_X$  is isomorphic to the sheaf  $\mathrm{Diff}_X^{\leq 1}$  of differential operators on  $X$  of order less or equal to 1.

Now we are ready to state our main result.

**Theorem 1.1.** *Let  $X$  be the adjoint Grassmannian of Dynkin type  $F_4$ . There exists a full exceptional collection of length 24*

$$\mathbf{D}^b(X) = \langle \mathcal{O}_X, \mathcal{U}^{\omega_4}, \tilde{T}_X, \mathcal{O}_X(1), \mathcal{U}^{\omega_4}(1), \tilde{T}_X(1), \dots, \mathcal{O}_X(7), \mathcal{U}^{\omega_4}(7), \tilde{T}_X(7) \rangle \quad (1.5)$$

*consisting of  $G$ -equivariant vector bundles. This collection is an  $\mathrm{Aut}(X)$ -invariant rectangular Lefschetz collection.*

The concept of adjoint varieties is meaningful in any Dynkin type and their derived categories share some common features. In a joint work with Alexander Kuznetsov [18], drawing motivation from the structure of the small quantum cohomology of these varieties, we have proposed some conjectures on the structure of their derived categories. In the particular case of adjoint varieties in non-simply laced Dynkin types (i.e., types  $B_n$ ,  $C_n$ ,  $F_4$ , or  $G_2$ ) our conjecture [18, Conjecture 1.7] predicts the existence of a full  $\mathrm{Aut}$ -invariant rectangular Lefschetz collection. Therefore, since (1.5) has all these properties, we obtain the following.

**Corollary 1.2.** *The conjecture [18, Conjecture 1.7] holds for the adjoint Grassmannian in type  $F_4$ .*

Since in types  $B_n$ ,  $C_n$ , and  $G_2$  the conjecture [18, Conjecture 1.7] is already known to hold, Corollary 1.2 finishes its proof by treating the only remaining case.

*Remark 1.3.* By [6, Proposition 6.3] the small quantum cohomology ring of  $X$  is known to be semisimple. Hence, Theorem 1.1 provides yet another instance where the first part of Dubrovin's conjecture (see [7]) is known to hold.

The proof of Theorem 1.1 naturally splits into two steps. First, by a direct computation we show that the collection (1.5) is exceptional. Second, we prove that (1.5) is full by restricting it to a family of subvarieties of  $X$  that sweep the whole  $X$  and are all isomorphic to the even orthogonal Grassmannian  $\text{OG}(2, 8)$ . It is essential that for  $\text{OG}(2, 8)$  a suitable exceptional collection is already known by [18]. In this respect the proof is similar to the proofs of fullness given in [16, 18]. The extension  $\tilde{T}_{\text{OG}(2,8)}$  of the tangent bundle  $T_{\text{OG}(2,8)}$  by the structure sheaf as in (1.4) also appears in the aforementioned collection on  $\text{OG}(2, 8)$ .

**Observation 1.4.** Let us discuss how the use of the extension  $\tilde{T}_X$  of the tangent bundle  $T_X$  by the structure sheaf  $\mathcal{O}_X$  to construct exceptional collections can be extended to other homogenous spaces. We restrict our attention to homogenous spaces  $G/P$  with  $P$  maximal parabolic and we label them by pairs  $(D, k)$ , where  $D$  is the Dynkin diagram of  $G$  and  $k$  is the vertex in  $D$  that corresponds to the parabolic  $P$ . We use the Bourbaki numbering of vertices in Dynkin diagrams [5].

For any  $G/P$  as above we have

$$\text{Ext}^\bullet(T_{G/P}, \mathcal{O}_{G/P}) = H^\bullet(G/P, \Omega_{G/P}) = \mathbb{k}[-1].$$

Hence, there always exists a unique non-trivial  $G$ -equivariant extension  $\tilde{T}_{G/P}$  of the tangent bundle  $T_{G/P}$  by the structure sheaf  $\mathcal{O}_{G/P}$ . SageMath [21] computations suggest the following patterns:

(1) For any  $G/P$  we have

$$\text{Ext}^i(T_{G/P}, T_{G/P}) = 0 \quad \text{for } i \geq 2. \tag{1.6}$$

(2) The extension  $\tilde{T}_{G/P}$  is an exceptional object for any  $G/P$  with the following exceptions:

$$\begin{aligned} & (A_n, 1) \text{ and } (A_n, n); \\ & (B_n, n-1) \text{ for } n \geq 3 \text{ and also } (B_2, 2); \\ & (C_n, 1) \text{ and } (C_n, 2) \text{ for } n \geq 3 \text{ and also } (C_2, 1); \\ & (F_4, 4) \text{ and } (G_2, 2). \end{aligned} \tag{1.7}$$

These exceptions seem to be closely related to the failure of the Yoneda map

$$\text{Ext}^1(T_{G/P}, \mathcal{O}_{G/P}) \times \text{Ext}^0(\mathcal{O}_{G/P}, T_{G/P}) \rightarrow \text{Ext}^1(T_{G/P}, T_{G/P}) \tag{1.8}$$

to be an isomorphism. In the cases  $(A_n, 1)$ ,  $(A_n, n)$ ,  $(B_2, 2)$  and  $(C_n, 1)$  the variety  $G/P$  is a projective space and it is well-known that in this case already  $T_{G/P}$  itself is an exceptional object. Hence,  $\text{Ext}^1(T_{G/P}, T_{G/P})$  vanishes and (1.8) is not injective. For all other exceptions the map (1.8) is not surjective.

(3) It is well-known (for example, see [1, Section IV.8, Theorem 2]) that for any  $G/P$  except for  $(B_n, n)$ ,  $(C_n, 1)$  and  $(G_2, 1)$  we have an isomorphism

$$H^0(G/P, T_{G/P}) = \mathfrak{g}.$$

Hence, the above discussion on the Yoneda map (1.8) suggests that for any  $G/P$  with the exceptions given by (1.7), and also  $(B_n, n)$  and  $(G_2, 1)$ , we have an isomorphism

$$\mathrm{Ext}^1(T_{G/P}, T_{G/P}) = \mathfrak{g} \quad (1.9)$$

and all the higher Ext-groups vanish as in (1.6).

(4) The sequence of vector bundles of the form

$$\mathcal{O}_{G/P}, \tilde{T}_{G/P}, \mathcal{O}_{G/P}(1), \tilde{T}_{G/P}(1), \dots, \mathcal{O}_{G/P}(s-1), \tilde{T}_{G/P}(s-1), \\ \mathcal{O}_{G/P}(s), \dots, \mathcal{O}_{G/P}(r-1) \quad (1.10)$$

where  $r$  is the index of  $G/P$  and  $s \in [1, r]$  is the maximal integer defined by the condition

$$\mathrm{Ext}^\bullet(\tilde{T}_{G/P}(i), \tilde{T}_{G/P}) = 0 \quad \text{for all } 0 < i < s,$$

is a Lefschetz collection for any  $G/P$  with the following exceptions:

$$\text{all cases from (1.7) and all quadrics.} \quad (1.11)$$

Note that all quadrics are given by the infinite series  $(B_n, 1)$  for  $n \geq 2$  and  $(D_n, 1)$  for  $n \geq 4$ , but also small rank sporadic cases  $(A_3, 2)$ ,  $(B_3, 3)$ ,  $(C_2, 2)$ ,  $(D_4, 3)$ ,  $(D_4, 4)$ , and  $(G_2, 1)$ .

Moreover, the cases

$$\begin{aligned} & (B_n, 2) \text{ and } (D_n, 2) \text{ for } n \geq 4; \\ & (A_5, k) \text{ for } k \in [2, 4]; \\ & (C_3, 3) \end{aligned} \quad (1.12)$$

are the only cases, where the collection (1.10) is a non-rectangular Lefschetz collection, i.e.  $s \neq r$ .

At this point we do not have rigorous proofs of the above claims. We hope to be able to say more on this in a future work.

*Remark 1.5.* Asking for the Yoneda map (1.8) to be an isomorphism is very close to saying that the pair of vector bundles  $\{\mathcal{O}_{G/P}, T_{G/P}\}$  forms an **exceptional block** in the sense of Kuznetsov and Polishchuk [17, Definition 3.1]. The only difference is that in [17] exceptional blocks always consist of irreducible vector bundles, but the tangent bundle  $T_{G/P}$  is irreducible only for cominuscule  $G/P$ .

The paper is organised as follows. In §2 we recall the necessary general background on  $G$ -equivariant vector bundles on  $G/P$  and collect some auxiliary facts for our particular setting. In §3 we construct and prove exceptionality of (1.5). Finally, in §4 we show that (1.5) is full.

**Acknowledgements.** First and foremost I would like to thank Alexander Kuznetsov for his continuing support and mathematical generosity. Our innumerable discussions have greatly influenced this work. Further, I am very grateful to Michel Brion and Nicolas Perrin for explaining to me the proof of Lemma 4.2. Finally, I would like to thank HIM and MPIM in Bonn for the great working conditions during the preparation of this paper.

## 2. PRELIMINARIES

We work over a fixed algebraically closed field  $\mathbb{k}$  of characteristic zero.

**2.1. Generalities on equivariant vector bundles.** Let  $G$  be a connected simply connected simple algebraic group and  $P$  its parabolic subgroup. Let us also fix a maximal torus  $T$  and a Borel subgroup  $B$  in  $G$  such that we have inclusions

$$T \subset B \subset P \subset G$$

Finally, let  $L \subset P$  be the Levi subgroup of  $P$  containing the maximal torus  $T$ . Note that the intersection  $B \cap L$  is a Borel subgroup in  $L$ .

Recall that there is a monoidal equivalence of categories between the category of  $G$ -equivariant coherent sheaves on  $G/P$  and the category of finite dimensional representations of  $P$

$$\mathrm{Coh}^G(G/P) \xrightarrow{\cong} \mathrm{Rep}(P) \tag{2.1}$$

given by sending a  $G$ -equivariant sheaf  $F$  to its fiber  $F_{[P]}$  at the point  $[P] \in G/P$ . Since the group  $P$  is not reductive, the category  $\mathrm{Rep}(P)$  is not semisimple and working with it can be quite hard. However, any irreducible representation of  $P$  is completely determined by its restriction to  $L$ , as the unipotent radical of  $P$  acts trivially on it. Hence, we can identify irreducible representations of  $P$  and  $L$ . Since  $L$  is a reductive group, we have the highest weight theory describing its finite dimensional irreducible representations. This makes working with irreducible representations of  $P$  much easier.

We denote by  $\mathbf{P}_L = \mathbf{P}_G$  the weight lattices for  $L$  and  $G$  with their natural identification and by

$$\mathbf{P}_G^+ \subset \mathbf{P}_L^+$$

the cones of dominant weights. For each dominant weight  $\lambda \in \mathbf{P}_L^+$  we denote by  $V_L^\lambda$  the corresponding irreducible representation of  $L$  and by  $\mathcal{U}^\lambda$  the  $G$ -equivariant vector bundle on  $G/P$  corresponding to it via the equivalence (2.1). Similarly, for  $\lambda \in \mathbf{P}_G^+$  we denote by  $V_G^\lambda$  the corresponding irreducible representation of  $G$ .

We denote by  $W$  the Weyl group of  $G$  and by  $W^L \subset W$  the Weyl group of  $L$ . We denote by  $w_0 \in W$  and  $w_0^L \in W^L$  the longest elements.

We have the following lemma.

**Lemma 2.1.**

- (1) We have  $(\mathcal{U}^\lambda)^\vee \cong \mathcal{U}^{-w_0^L \lambda}$ .
- (2) If  $V_L^\lambda \otimes V_L^\mu = \bigoplus V_L^\nu$ , then  $\mathcal{U}^\lambda \otimes \mathcal{U}^\mu = \bigoplus \mathcal{U}^\nu$ .

*Proof.* This is an immediate consequence of the equivalence (2.1) and [17, Formula (6)].  $\square$

For equivariant vector bundles of the form  $\mathcal{U}^\lambda$  we also have a very efficient way to compute their cohomology. Let  $\ell: W \rightarrow \mathbb{Z}$  be the length function and let  $\rho \in \mathbf{P}_G^+$  be the sum of fundamental weights of  $G$ .

**Theorem 2.2** (Borel–Weil–Bott). *Consider a vector bundle  $\mathcal{U}^\lambda$  with  $\lambda \in \mathbf{P}_L^+$ . If the weight  $\lambda + \rho$  lies on a wall of a Weyl chamber for the  $W$ -action, then*

$$H^\bullet(G/P, \mathcal{U}^\lambda) = 0.$$

*Otherwise, if  $w \in W$  is the unique element such that the weight  $w(\lambda + \rho)$  is dominant, then*

$$H^\bullet(G/P, \mathcal{U}^\lambda) = V_G^{w(\lambda + \rho) - \rho}[-\ell(w)].$$

**2.2. Specifics of our case.** As in Section 1 we let  $G$  be a connected simple algebraic group of Dynkin type  $F_4$  and  $P = P_1 \subset G$  be a maximal parabolic subgroup associated with the first vertex of its Dynkin diagram (1.1). We then consider the homogenous space

$$X = G/P,$$

which is the adjoint Grassmannian in type  $F_4$ .

The Levi subgroup  $L$  of  $P$  is a reductive group, whose semisimple part is of Dynkin type  $C_3$ . A more detailed description of  $L$  is given below in Section 2.3.

We realize the weight lattice  $\mathbf{P}_G$  of  $G$  inside  $\mathbb{Q}^4$  as in [5, p.211–213]. Let  $e_1, e_2, e_3, e_4$  be the standard basis of  $\mathbb{Q}^4$ ; it is orthonormal with respect to the usual scalar product. The weight lattice  $\mathbf{P}_G$  is generated inside  $\mathbb{Q}^4$  by the fundamental weights  $\omega_1, \dots, \omega_4$  or by the simple roots  $\alpha_1, \dots, \alpha_4$ , which are given by

$$\begin{aligned} \omega_1 &= e_1 + e_2 & \alpha_1 &= e_2 - e_3 \\ \omega_2 &= 2e_1 + e_2 + e_3 & \alpha_2 &= e_3 - e_4 \\ \omega_3 &= \frac{1}{2}(3e_1 + e_2 + e_3 + e_4) & \alpha_3 &= e_4 \\ \omega_4 &= e_1 & \alpha_4 &= \frac{1}{2}(e_1 - e_2 - e_3 - e_4) \end{aligned} \tag{2.2}$$

From (2.2) we see that a weight  $\lambda = (\lambda_1, \lambda_2, \lambda_3, \lambda_4)$  is  $G$ -dominant if and only if

$$\lambda_2 \geq \lambda_3 \geq \lambda_4 \geq 0 \quad \text{and} \quad \lambda_1 \geq \lambda_2 + \lambda_3 + \lambda_4. \tag{2.3}$$

The sum of the fundamental weights appearing in the Borel–Weil–Bott theorem is

$$\rho = \frac{11}{2}e_1 + \frac{5}{2}e_2 + \frac{3}{2}e_3 + \frac{1}{2}e_4. \tag{2.4}$$

All roots of  $G$  are given by

$$\pm e_i, \quad \pm e_i \pm e_j, \quad \frac{1}{2}(\pm e_1 \pm e_2 \pm e_3 \pm e_4) \tag{2.5}$$

and, therefore, the walls of the Weyl chambers are given by the hyperplanes

$$\begin{aligned} \lambda_i &= 0, \quad 1 \leq i \leq 4, \\ \lambda_i &= \pm \lambda_j, \quad 1 \leq i \neq j \leq 4, \\ \sum_{i \in I} \lambda_i &= \sum_{j \in J} \lambda_j, \quad I \sqcup J = \{1, 2, 3, 4\}. \end{aligned} \tag{2.6}$$

Note that in the last condition  $I$  and  $J$  can be empty.

The longest element  $w_0 \in W$  acts as

$$w_0(\lambda_1, \lambda_2, \lambda_3, \lambda_4) = (-\lambda_1, -\lambda_2, -\lambda_3, -\lambda_4). \tag{2.7}$$

Hence, we see that any representation of  $G$  is self-dual. The longest element  $w_0^L \in W^L$  acts by

$$w_0^L(\lambda_1, \lambda_2, \lambda_3, \lambda_4) = (\lambda_2, \lambda_1, -\lambda_3, -\lambda_4). \tag{2.8}$$

The ample generator of the Picard group is denoted by  $\mathcal{O}(1)$  and in terms of the equivalence (2.1) we have

$$\mathcal{O}(t) = \mathcal{U}^{t\omega_1} \quad \text{and} \quad \mathcal{U}^{\lambda+t\omega_1} = \mathcal{U}^\lambda(t) \quad \text{for} \quad t \in \mathbb{Z}. \tag{2.9}$$

Now we can apply Lemma 2.1(1) to compute the necessary duals.

**Corollary 2.3.** *We have*

$$\left(\mathcal{U}^{a_2\omega_2+a_3\omega_3+a_4\omega_4}\right)^\vee = \mathcal{U}^{a_2\omega_2+a_3\omega_3+a_4\omega_4}(-3a_2 - 2a_3 - a_4).$$

*Proof.* Using (2.8) and (2.2) we get

$$\begin{aligned} -w_0^L(a_2\omega_2 + a_3\omega_3 + a_4\omega_4) &= \left(-a_2 - \frac{1}{2}a_3, -2a_2 - \frac{3}{2}a_3 - a_4, a_2 + \frac{1}{2}a_3, \frac{1}{2}a_3\right) = \\ &= (-3a_2 - 2a_3 - a_4)\omega_1 + a_2\omega_2 + a_3\omega_3 + a_4\omega_4, \end{aligned}$$

and the claim now follows from Lemma 2.1(1) and (2.9).  $\square$

**2.3. Tensor products and exterior powers.** To compute tensor products of equivariant vector bundles on a homogeneous space  $G/P$  one can use Lemma 2.1(2), which turns it into the problem of computing tensor products of representations of the Levi subgroup  $L$ . We explain here how to further reduce such computations to the known results on tensor products for the semisimple part of  $L$ . Such results exist in the literature in the form of tables (e.g. [23]) and in the form of software (e.g. [21]). Essentially this is a consistent way to control the twist by the center of  $L$ .

Let us temporarily go back into the general setting of Section 2.1. We denote by  $\alpha_1, \dots, \alpha_n$  and  $\omega_1, \dots, \omega_n$  the simple roots and the fundamental weights of an arbitrary semisimple group  $G$ . We also make use of the  $W$ -invariant scalar product on the ambient space of the root system of  $G$ , which we identify with  $(\mathbf{P}_G)_{\mathbb{Q}}$ ; we denote it by  $(\lambda, \mu)$ . To simplify the notation, let us also assume that the parabolic  $P$  is maximal and corresponds to the  $k$ -th vertex of the Dynkin diagram of  $G$ . Denote by  $R(L)$  the roots of  $L$ , i.e. those roots of  $G$ , whose expression as a linear combination of the simple roots  $\alpha_1, \dots, \alpha_n$  does not involve the simple root  $\alpha_k$ . In other words,  $R(L)$  consists of those roots of  $G$  that are orthogonal to  $\omega_k$  with respect to the scalar product.

Consider two closed algebraic subgroups of  $L$ : the derived group  $L' \subset L$  and the center  $Z(L) \subset L$ . The derived group  $L'$  is a connected simply connected semisimple algebraic group, whose Dynkin diagram is obtained from the Dynkin diagram of  $G$  by removing the  $k$ -th vertex. For the center we have an isomorphism  $Z(L) \simeq \mathbb{G}_m$ . The group law in  $L$  gives a surjective homomorphism of algebraic groups

$$L' \times Z(L) \rightarrow L \tag{2.10}$$

with finite kernel  $L' \cap Z(L)$ . The weight lattices of  $L'$  and  $Z(L)$  are the quotients of  $\mathbf{P}_L = \mathbf{P}_G$

$$\begin{array}{ccc} & \mathbf{P}_L & \\ \varphi_{L'} \swarrow & & \searrow \varphi_Z \\ \mathbf{P}_{L'} = \mathbf{P}_L / (\mathbb{Z}\omega_k) & & \mathbf{P}_{Z(L)} = \mathbf{P}_L / \left( \sum_{\alpha \in R(L)} \mathbb{Q}\alpha \cap \mathbf{P}_L \right) \end{array} \tag{2.11}$$

and the map

$$\mathbf{P}_L \xrightarrow{(\varphi_{L'}, \varphi_Z)} \mathbf{P}_{L'} \oplus \mathbf{P}_{Z(L)} \tag{2.12}$$

is an embedding as a finite index subgroup. Moreover, under  $\varphi_{L'}$  the fundamental weights  $\omega_1, \dots, \omega_n$  of  $G$  get mapped to the fundamental weights  $\omega'_1, \dots, \omega'_{k-1}, \omega'_{k+1}, \dots, \omega'_n$  of  $L'$  (with

the labelling of simple roots inherited from  $G$ ). More precisely, we have

$$\varphi_{L'}(\omega_i) = \begin{cases} \omega'_i & i \neq k, \\ 0 & i = k. \end{cases} \quad (2.13)$$

A more detailed discussion of this setup can be found in [12, II.1.18].

Over  $\mathbb{Q}$  we have an orthogonal direct sum decomposition

$$(\mathbf{P}_L)_{\mathbb{Q}} \cong \left( \sum_{\alpha \in R(L)} \mathbb{Q}\alpha \right) \oplus (\mathbb{Q}\omega_k),$$

that allows us to identify

$$(\mathbf{P}_{L'})_{\mathbb{Q}} \cong \sum_{\alpha \in R(L)} \mathbb{Q}\alpha \quad \text{and} \quad (\mathbf{P}_{Z(L)})_{\mathbb{Q}} \cong \mathbb{Q}\omega_k.$$

Thus, after tensoring with  $\mathbb{Q}$  the map (2.12) becomes an isomorphism

$$(\mathbf{P}_L)_{\mathbb{Q}} \xrightarrow{(\varphi_{L'}, \varphi_Z)} (\mathbf{P}_{L'})_{\mathbb{Q}} \oplus \mathbb{Q}\omega_k \quad (2.14)$$

such that for a  $\lambda = \sum_i a_i \omega_i \in (\mathbf{P}_L)_{\mathbb{Q}}$  we have

$$\begin{aligned} \varphi_{L'}(\lambda) &= \lambda', \quad \text{where} \quad \lambda' = \sum_{i \neq k} a_i \omega'_i, \\ \varphi_Z(\lambda) &= r_\lambda \omega_k, \quad \text{where} \quad r_\lambda = \frac{(\lambda, \omega_k)}{(\omega_k, \omega_k)}. \end{aligned} \quad (2.15)$$

Note that the map

$$\begin{aligned} \varphi_{L'}: \mathbf{P}_L &\rightarrow \mathbf{P}_{L'} \\ \lambda &\mapsto \lambda' \end{aligned}$$

is completely determined by (2.13), as  $\omega_1, \dots, \omega_n$  form a basis of  $\mathbf{P}_L$ . We can also consider the map

$$\begin{aligned} \mathbf{P}_{L'} &\rightarrow \mathbf{P}_L \\ \sum_{i \neq k} a_i \omega'_i &\mapsto \sum_{i \neq k} a_i \omega_i \end{aligned} \quad (2.16)$$

to which we will refer to as a **lifting** from  $\mathbf{P}_{L'}$  to  $\mathbf{P}_L$ .

Now we are ready to discuss tensor products. By (2.10) a representation of  $L$  corresponds to a pair of compatible representations of  $L'$  and  $Z(L)$ . We are going to exploit this to compute tensor operations for representations of  $L$ . For a highest weight irreducible representation  $V_L^\lambda$  its restrictions  $\text{Res}_{L'}^L(V_L^\lambda)$  and  $\text{Res}_{Z(L)}^L(V_L^\lambda)$  are again irreducible. Hence, we have

$$\text{Res}_{L'}^L(V_L^\lambda) = V_{L'}^{\lambda'},$$

where  $\lambda'$  is the image of  $\lambda$  in  $\mathbf{P}_{L'}$  under (2.13), and  $\text{Res}_{Z(L)}^L(V_L^\lambda)$  is given by a character of  $Z(L) \simeq \mathbb{G}_m$ .

**Lemma 2.4.** *Let  $V_L^\lambda, V_L^\mu$  be irreducible representations of  $L$  with highest weights  $\lambda, \mu \in \mathbf{P}_L$ . Let  $V_{L'}^{\lambda'}, V_{L'}^{\mu'}$  be the induced irreducible representations of  $L'$  as above.*



(1) If

$$V_{L'}^{\lambda'} \otimes V_{L'}^{\mu'} = \bigoplus_{\nu' \in \Sigma} \left( V_{L'}^{\nu'} \right)^{m(\lambda', \mu', \nu')}$$

then

$$V_L^\lambda \otimes V_L^\mu = \bigoplus_{\nu' \in \Sigma} \left( V_L^{\nu + (r_\lambda + r_\mu - r_{\nu'}) \omega_k} \right)^{m(\lambda', \mu', \nu')}$$

where for a weight  $\nu' \in \Sigma$  we denote by  $\nu$  its lifting (2.16).

(2) If

$$\Lambda^q V_{L'}^{\lambda'} = \bigoplus_{\nu' \in \Sigma} \left( V_{L'}^{\nu'} \right)^{m(\lambda', \nu')}$$

then

$$\Lambda^q V_L^\lambda = \bigoplus_{\nu' \in \Sigma} \left( V_L^{\nu + (qr_\lambda - r_{\nu'}) \omega_k} \right)^{m(\lambda', \nu')}$$

where for a weight  $\nu' \in \Sigma$  we denote by  $\nu$  its lifting (2.16).

(3) If

$$S^q V_{L'}^{\lambda'} = \bigoplus_{\nu' \in \Sigma} \left( V_{L'}^{\nu'} \right)^{m(\lambda', \nu')}$$

then

$$S^q V_L^\lambda = \bigoplus_{\nu' \in \Sigma} \left( V_L^{\nu + (qr_\lambda - r_{\nu'}) \omega_k} \right)^{m(\lambda', \nu')}$$

where for a weight  $\nu' \in \Sigma$  we denote by  $\nu$  its lifting (2.16).

*Proof.* Let us only prove the first statement, as the others are proved in a very similar way.

As discussed just before the statement of the lemma, to compute the tensor product  $V_L^\lambda \otimes V_L^\mu$  it is enough to determine how  $L'$  and  $Z(L)$  act on it. Restricting to  $L'$  we need to compute the tensor product  $V_{L'}^{\lambda'} \otimes V_{L'}^{\mu'}$ , which decomposes as

$$V_{L'}^{\lambda'} \otimes V_{L'}^{\mu'} = \bigoplus_{\nu' \in \Sigma} \left( V_{L'}^{\nu'} \right)^{m(\lambda', \mu', \nu')}$$

by our assumptions. Therefore,  $V_L^\lambda \otimes V_L^\mu$  has a similar direct sum decomposition

$$V_L^\lambda \otimes V_L^\mu = \bigoplus_{\nu' \in \Sigma} \left( V_L^{\nu + t_{(\lambda, \mu, \nu')} \omega_k} \right)^{m(\lambda', \mu', \nu')},$$

where for a  $\nu' \in \Sigma$  we denote by  $\nu$  its lifting (2.16) and  $t_{(\lambda, \mu, \nu)}$ 's are some integers. The integers  $t_{(\lambda, \mu, \nu)}$  are determined by looking at the action of the center  $Z(L)$ . Since on each  $V_L^{\nu + t_{(\lambda, \mu, \nu)} \omega_k}$  the torus  $Z(L)$  acts by a character, and since characters simply add under the tensor product, we get the relation

$$r_{\nu + t_{(\lambda, \mu, \nu)} \omega_k} = r_\lambda + r_\mu,$$

which implies the desired  $t_{(\lambda, \mu, \nu)} = r_\lambda + r_\mu - r_{\nu'}$ . □

From now on we return back to the case of our adjoint variety in type  $F_4$ . In Example 2.5 we illustrate how to use Lemma 2.4 in practice and then in Lemmas 2.6 and 2.7 we collect all tensor products and exterior/symmetric powers required for this paper.

Using (2.2) and (2.15) we get

$$r_{\omega_1} = 1, \quad r_{\omega_2} = \frac{3}{2}, \quad r_{\omega_3} = 1, \quad r_{\omega_4} = \frac{1}{2}.$$

**Example 2.5.** Let us compute  $\mathcal{U}^{\omega_4}(-1) \otimes \mathcal{U}^{\omega_4}(-1)$  using Lemma 2.4, i.e. we need to compute  $V_L^{\omega_4 - \omega_1} \otimes V_L^{\omega_4 - \omega_1}$ . In the notation of Lemma 2.4 we have

$$\lambda = \mu = \omega_4 - \omega_1 \quad \text{and} \quad \lambda' = \mu' = \omega_4' \quad \text{and} \quad r_\lambda = r_\mu = -\frac{1}{2}.$$

Using the table in [23, p.302] we get

$$V_{L'}^{\omega_4'} \otimes V_{L'}^{\omega_4'} = \mathbb{k} \oplus V_{L'}^{\omega_3'} \oplus V_{L'}^{2\omega_4'}.$$

Thus, we have  $\Sigma = \{0, \omega_3', 2\omega_4'\}$  and all multiplicities  $m(\lambda', \mu', \nu')$  are equal to 1. Therefore, applying Lemma 2.4(1) we get

$$V_L^{\omega_4 - \omega_1} \otimes V_L^{\omega_4 - \omega_1} = V_L^{-\omega_1} \oplus V_L^{\omega_3 - 2\omega_1} \oplus V_L^{2\omega_4 - 2\omega_1},$$

and, finally,

$$\mathcal{U}^{\omega_4}(-1) \otimes \mathcal{U}^{\omega_4}(-1) = \mathcal{O}(-1) \oplus \mathcal{U}^{\omega_3}(-2) \oplus \mathcal{U}^{2\omega_4}(-2).$$

**Lemma 2.6.** *We have*

$$\begin{aligned} \mathcal{U}^{\omega_4} \otimes \mathcal{U}^{\omega_4} &= \mathcal{U}^{2\omega_4} \oplus \mathcal{U}^{\omega_3} \oplus \mathcal{O}(1) \\ \mathcal{U}^{\omega_4} \otimes \mathcal{U}^{\omega_3} &= \mathcal{U}^{\omega_3 + \omega_4} \oplus \mathcal{U}^{\omega_2} \oplus \mathcal{U}^{\omega_4}(1) \\ \mathcal{U}^{\omega_4} \otimes \mathcal{U}^{\omega_2} &= \mathcal{U}^{\omega_2 + \omega_4} \oplus \mathcal{U}^{\omega_3}(1) \\ \mathcal{U}^{\omega_2} \otimes \mathcal{U}^{\omega_3} &= \mathcal{U}^{\omega_2 + \omega_3} \oplus \mathcal{U}^{\omega_3 + \omega_4}(1) \oplus \mathcal{U}^{\omega_4}(2) \\ \mathcal{U}^{\omega_2} \otimes \mathcal{U}^{\omega_2} &= \mathcal{U}^{2\omega_2} \oplus \mathcal{U}^{2\omega_3}(1) \oplus \mathcal{U}^{2\omega_4}(2) \oplus \mathcal{O}(3) \\ \mathcal{U}^{2\omega_4} \otimes \mathcal{U}^{\omega_4} &= \mathcal{U}^{3\omega_4} \oplus \mathcal{U}^{\omega_3 + \omega_4} \oplus \mathcal{U}^{\omega_4}(1) \\ \mathcal{U}^{2\omega_4} \otimes \mathcal{U}^{\omega_3} &= \mathcal{U}^{\omega_3 + 2\omega_4} \oplus \mathcal{U}^{\omega_2 + \omega_4} \oplus \mathcal{U}^{2\omega_4}(1) \oplus \mathcal{U}^{\omega_3}(1) \\ \mathcal{U}^{2\omega_4} \otimes \mathcal{U}^{\omega_2} &= \mathcal{U}^{\omega_2 + 2\omega_4} \oplus \mathcal{U}^{\omega_3 + \omega_4}(1) \oplus \mathcal{U}^{\omega_2}(1) \end{aligned}$$

*Proof.* The proof is a routine calculation similar to Example 2.5 using [23, p.302] and Lemma 2.4.  $\square$

**Lemma 2.7.** *We have*

$$S^k \mathcal{U}^{\omega_4} = \mathcal{U}^{k\omega_4} \quad \forall k \quad \text{and} \quad \Lambda^k \mathcal{U}^{\omega_4} = \begin{cases} \mathcal{O} & k = 0 \\ \mathcal{U}^{\omega_4} & k = 1 \\ \mathcal{U}^{\omega_3} \oplus \mathcal{O}(1) & k = 2 \\ \mathcal{U}^{\omega_2} \oplus \mathcal{U}^{\omega_4}(1) & k = 3 \\ \mathcal{U}^{\omega_3}(1) \oplus \mathcal{O}(2) & k = 4 \\ \mathcal{U}^{\omega_4}(2) & k = 5 \\ \mathcal{O}(3) & k = 6 \end{cases}$$

$$\Lambda^2 \mathcal{U}^{\omega_3} = \mathcal{U}^{\omega_2 + \omega_4} \oplus \mathcal{U}^{2\omega_4}(1)$$

$$\Lambda^3 \mathcal{U}^{\omega_3} = \mathcal{U}^{2\omega_2} \oplus \mathcal{U}^{\omega_3+2\omega_4}(1) \oplus \mathcal{U}^{\omega_2+\omega_4}(1) \oplus \mathcal{U}^{2\omega_4}(2)$$

*Proof.* The proof is a routine calculation similar to Example 2.5 using [23, p.302] and Lemma 2.4.  $\square$

## 2.4. Cohomology computations.

**Lemma 2.8.** *We have*

$$H^\bullet(X, \mathcal{O}) = \mathbb{k}[0], \quad (2.17)$$

$$H^\bullet(X, \mathcal{U}^{\omega_2}(-2)) = \mathbb{k}[-1], \quad (2.18)$$

$$H^\bullet(X, \mathcal{U}^{2\omega_2}(-3)) = V^{\omega_1}[-1]. \quad (2.19)$$

*Proof.* The equality (2.17) follows from the fact that  $X$  is a smooth connected Fano variety together with the Kodaira vanishing theorem.

To prove (2.18) and (2.19) we apply the Borel–Weil–Bott theorem. Indeed, in the first case we have  $\lambda = \omega_2 - 2\omega_1$  and it is easy to see using (2.2)–(2.4) that we have  $s_{\alpha_1}(\lambda + \rho) = \rho$ , hence (2.18) holds. Similarly, in the second case we have  $\lambda = 2\omega_2 - 3\omega_1$  and  $s_{\alpha_1}(\lambda + \rho) = \omega_1 + \rho$ , and so (2.19) follows.  $\square$

By the Borel–Weil–Bott theorem a vector bundle  $\mathcal{U}^\lambda$  is acyclic if and only if the weight  $\lambda + \rho$  lies on a wall of a Weyl chamber for the  $W$ -action; such weights are called **singular**. A routine computation using formulas (2.6) for the walls of the Weyl chambers gives the following.

**Corollary 2.9.** *The following vector bundles are acyclic*

$$\begin{aligned} \mathcal{O}(t) & \text{ for } t \in [-7, -1] \\ \mathcal{U}^{\omega_2}(t) & \text{ for } t \in [-10, -1] \setminus \{-9, -2\} \\ \mathcal{U}^{\omega_3}(t) & \text{ for } t \in [-9, -1] \\ \mathcal{U}^{\omega_4}(t) & \text{ for } t \in [-8, -1] \\ \mathcal{U}^{2\omega_2}(t) & \text{ for } t \in [-10, -1] \setminus \{-3, -2\} \\ \mathcal{U}^{2\omega_3}(t) & \text{ for } t \in [-11, -1] \setminus \{-9, -3\} \\ \mathcal{U}^{2\omega_4}(t) & \text{ for } t \in [-9, -1] \\ \mathcal{U}^{\omega_2+\omega_4}(t) & \text{ for } t \in [-11, -1] \setminus \{-10, -2\} \\ \mathcal{U}^{\omega_3+\omega_4}(t) & \text{ for } t \in [-10, -1] \\ \mathcal{U}^{\omega_2+2\omega_4}(t) & \text{ for } t \in [-12, -1] \setminus \{-11, -2\} \\ \mathcal{U}^{3\omega_4}(t) & \text{ for } t \in [-10, -1] \end{aligned}$$

*Remark 2.10.* Note that for any  $L$ -dominant weight

$$\lambda = \sum_i a_i \omega_i \quad \text{with } a_1 = -1$$

the vector bundle  $\mathcal{U}^\lambda$  is always acyclic. Indeed, this follows from the Borel–Weil–Bott theorem, as  $\lambda + \rho$  is fixed by the simple reflection  $s_{\alpha_1}$  and is, therefore, a singular weight.

### 3. CONSTRUCTION OF THE EXCEPTIONAL COLLECTION

Recall that  $X$  is the adjoint Grassmannian of type  $F_4$ . In this section we show that the sequence of vector bundles

$$\mathcal{O}, \mathcal{U}^{\omega_4}, \tilde{T}_X, \mathcal{O}(1), \mathcal{U}^{\omega_4}(1), \tilde{T}_X(1), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7), \tilde{T}_X(7) \quad (3.1)$$

(where  $\tilde{T}_X$  is an extension of the tangent bundle  $T_X$  defined below in (3.3)) is an exceptional collection on  $X$ . As already mentioned in the Introduction, this collection is a rectangular Lefschetz exceptional collection.

Let us first consider the subcollection of (3.1) generated by  $\mathcal{O}$  and  $\mathcal{U}^{\omega_4}$ .

**Lemma 3.1.** *The collection of vector bundles*

$$\mathcal{O}, \mathcal{U}^{\omega_4}, \mathcal{O}(1), \mathcal{U}^{\omega_4}(1), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7)$$

*is exceptional.*

*Proof.* We first show that subcollections  $\mathcal{O}, \mathcal{O}(1), \dots, \mathcal{O}(7)$  and  $\mathcal{U}^{\omega_4}, \mathcal{U}^{\omega_4}(1), \dots, \mathcal{U}^{\omega_4}(7)$  are exceptional, and then show that they can be merged together as in the statement of the lemma.

By (2.17) any line bundle  $\mathcal{O}(t)$  is exceptional and from Corollary 2.9 it follows that the subcollection  $\mathcal{O}, \mathcal{O}(1), \dots, \mathcal{O}(7)$  is exceptional. Thus, we only need to deal with the subcollection  $\mathcal{U}^{\omega_4}, \mathcal{U}^{\omega_4}(1), \dots, \mathcal{U}^{\omega_4}(7)$ . It is enough to show

$$\mathrm{Ext}^\bullet(\mathcal{U}^{\omega_4}(t), \mathcal{U}^{\omega_4}) = \begin{cases} \mathbb{k}[0] & \text{if } t = 0, \\ 0 & \text{if } 1 \leq t \leq 7. \end{cases}$$

By Corollary 2.3 and Lemma 2.6 we rewrite

$$\begin{aligned} \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_4}(t), \mathcal{U}^{\omega_4}) &= H^\bullet(X, \mathcal{U}^{\omega_4} \otimes \mathcal{U}^{\omega_4}(-t-1)) = \\ &= H^\bullet(X, \mathcal{U}^{2\omega_4}(-t-1)) \oplus H^\bullet(X, \mathcal{U}^{\omega_3}(-t-1)) \oplus H^\bullet(X, \mathcal{O}(-t)). \end{aligned}$$

Finally, applying (2.17) and Corollary 2.9 we get the desired result.

To merge the two subcollections it is enough to show

$$\begin{aligned} \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_4}(t), \mathcal{O}) &= 0 \quad \text{for } 0 \leq t \leq 7, \\ \mathrm{Ext}^\bullet(\mathcal{O}(t), \mathcal{U}^{\omega_4}) &= 0 \quad \text{for } 1 \leq t \leq 7. \end{aligned}$$

Rewriting both Ext-groups in terms of cohomology, and using Corollary 2.3 to dualize  $\mathcal{U}^{\omega_4}(t)$ , we see that the above conditions are equivalent to

$$H^\bullet(X, \mathcal{U}^{\omega_4}(-t)) = 0 \quad \text{for } 1 \leq t \leq 8,$$

which holds by Corollary 2.9. □

Now we start our discussion of the tangent bundle. Recall that for any rational homogeneous space  $G/P$  we have the following description of the tangent bundle. Let  $\mathfrak{g}$  and  $\mathfrak{p}$  be the Lie algebras of  $G$  and  $P$  respectively. The group  $G$  (resp.  $P$ ) acts on the Lie algebra  $\mathfrak{g}$  (resp.  $\mathfrak{p}$ ) via the adjoint representation. It is well-known that under the equivalence (2.1) the tangent bundle  $T_{G/P}$  corresponds to the  $P$ -representation  $\mathfrak{g}/\mathfrak{p}$ . Using the Killing form, we get an isomorphism of  $P$ -representations  $\mathfrak{g}/\mathfrak{p} = (\mathfrak{n})^\vee$ , where  $\mathfrak{n}$  is the nilradical of  $\mathfrak{p}$ . This way we also see that the cotangent bundle  $\Omega_{G/P}$  corresponds to  $\mathfrak{n}$  under (2.1). Finally, we note that  $T_{G/P}$  and  $\Omega_{G/P}$  are irreducible if and only if the homogeneous space  $G/P$  is cominuscule.

Now we are ready to add the (extension of the) tangent bundle to the collection. For a  $G$ -equivariant vector bundle  $E$  we denote by  $\text{ss}(E)$  its semisimplification, i.e. the direct sum of composition factors of  $E$ .

**Lemma 3.2.** *The following statements hold.*

(1) *On  $X$  there exist unique  $G$ -equivariant non-split short exact sequences*

$$0 \rightarrow \mathcal{U}^{\omega_2}(-1) \rightarrow T_X \rightarrow \mathcal{O}(1) \rightarrow 0 \quad (3.2)$$

and

$$0 \rightarrow \mathcal{O} \rightarrow \tilde{T}_X \rightarrow T_X \rightarrow 0, \quad (3.3)$$

where  $T_X$  is the tangent bundle of  $X$  and  $\tilde{T}_X$  is some  $G$ -equivariant vector bundle of rank 16.

(2) *We have*

$$\text{ss}(\tilde{T}_X) = \mathcal{O} \oplus \mathcal{U}^{\omega_2}(-1) \oplus \mathcal{O}(1). \quad (3.4)$$

(3) *We have*

$$T_X = \mathbf{L}_{\langle \mathcal{O}(1) \rangle}(\mathcal{U}^{\omega_2}(-1)) \quad \text{and} \quad \tilde{T}_X = \mathbf{R}_{\langle \mathcal{O} \rangle}(T_X), \quad (3.5)$$

where  $\mathbf{L}$  and  $\mathbf{R}$  stand for the left and right mutation functors.

(4) *We have a  $G$ -equivariant isomorphism*

$$\tilde{T}_X = (\tilde{T}_X)^\vee(1). \quad (3.6)$$

*Proof.* Consider the short exact sequence of  $P$ -representations

$$0 \rightarrow [\mathfrak{n}, \mathfrak{n}] \rightarrow \mathfrak{n} \rightarrow \mathfrak{n}/[\mathfrak{n}, \mathfrak{n}] \rightarrow 0.$$

Since  $X$  is an adjoint variety, by [3, Lemma 41] we have that  $[\mathfrak{n}, \mathfrak{n}]$  is one-dimensional and  $\mathfrak{n}/[\mathfrak{n}, \mathfrak{n}]$  is irreducible. Using explicit formulas (2.5) for the roots of the  $F_4$  root system, it is not difficult to check that we have

$$[\mathfrak{n}, \mathfrak{n}] = \mathfrak{g}_{-\omega_1} = V_L^{-\omega_1},$$

where  $\omega_1 = \Theta$  is the highest root of  $F_4$ , and that the highest weight of the  $L$ -representation  $\mathfrak{n}/[\mathfrak{n}, \mathfrak{n}]$  is  $\omega_2 - 2\omega_1 = -\alpha_1$ . Indeed, examining the weight spaces of  $\mathfrak{n}$  we see that among them the only  $L$ -dominant weights are  $-\omega_1, \omega_4 - \omega_1, \omega_2 - 2\omega_1$ , hence the claim. Therefore, for the cotangent bundle we have a short exact sequence

$$0 \rightarrow \mathcal{O}(-1) \rightarrow \Omega_X \rightarrow \mathcal{U}^{\omega_2}(-2) \rightarrow 0. \quad (3.7)$$

Dualizing (3.7) and using Corollary 2.3 we get (3.2). Since by (2.18) we have

$$\text{Ext}^\bullet(\mathcal{O}(1), \mathcal{U}^{\omega_2}(-1)) = \mathbb{k}[-1], \quad (3.8)$$

there exists a unique non-trivial  $G$ -equivariant extension of the form (3.2).

To prove (3.3) we note

$$\text{Ext}^\bullet(T_X, \mathcal{O}) = H^\bullet(X, \Omega_X) = H^\bullet(X, \mathcal{U}^{\omega_2}(-2)) = \mathbb{k}[-1], \quad (3.9)$$

where we used (3.7), Corollary 2.9, and (2.18). Thus, there exists a unique non-trivial  $G$ -equivariant extension  $\tilde{T}_X$  as in (3.3).

For (3.4) we just use (3.2) and (3.3).

For (3.5) it is enough to show that (3.2) and (3.3) are the corresponding mutation triangles. For the former it follows from (3.8) and for the latter from (3.9).

To show (3.6) we proceed as follows. The exact sequences (3.2)–(3.3) imply that  $\tilde{T}_X[1]$  is the left convolution of the 3-term complex

$$\mathcal{O}(1)[-1] \rightarrow \mathcal{U}^{\omega_2}(-1) \rightarrow \mathcal{O}[1], \quad (3.10)$$

in the triangulated category  $\mathbf{D}^b(X)$ ; the composition is zero due to the vanishing

$$\mathrm{Ext}^\bullet(\mathcal{O}(1), \mathcal{O}) = H^\bullet(X, \mathcal{O}(-1)) = 0 \quad (3.11)$$

that holds by Corollary 2.9 and hence (3.10) is indeed a complex. By (3.11) convolutions of (3.10) are unique. Dualizing (3.10) and then twisting the result by  $\mathcal{O}(1)$  we get a new 3-term complex, whose right convolution is  $(\tilde{T}_X)^\vee(1)$ . However, using Corollary 2.3 we see that the new complex is nothing else but the old complex (3.10) and, therefore, its left convolution must be again  $\tilde{T}_X[1]$ . Finally, since the left convolution always differs from the right convolution just by a shift by  $[1]$ , we obtain the desired claim.  $\square$

*Remark 3.3.* The isomorphism (3.6) gives rise to a  $G$ -equivariant non-degenerate bilinear pairing. Moreover, one can show that this pairing is skew-symmetric, i.e. we have  $G$ -equivariant pairing

$$\Lambda^2 \tilde{T}_X \rightarrow \mathcal{O}(1).$$

This pairing comes from the skew-symmetric  $G$ -equivariant pairing on  $\mathcal{U}^{\omega_2}(-1)$ . As we won't need this fact, we omit the proof.

Using (3.2)–(3.3) we compute global sections

$$H^0(X, T_X) = V^{\omega_1} = \mathfrak{g} \quad \text{and} \quad H^0(X, \tilde{T}_X) = V^{\omega_1} \oplus \mathbb{C}.$$

Since  $X$  is homogeneous, the tangent bundle  $T_X$  is globally generated and we have a surjection

$$V^{\omega_1} \otimes \mathcal{O} \rightarrow T_X.$$

Therefore,  $\tilde{T}_X$  is also globally generated and we have a surjection

$$(V^{\omega_1} \oplus \mathbb{C}) \otimes \mathcal{O} \xrightarrow{(\alpha, \beta)} \tilde{T}_X. \quad (3.12)$$

Recall that a 3-term complex sitting in degrees  $-1, 0, 1$  and exact in the first and last terms is called a **monad**.

**Lemma 3.4.** *There is a monad*

$$\tilde{T}_X(-1) \rightarrow (V^{\omega_1} \oplus \mathbb{C}) \otimes \mathcal{O} \xrightarrow{(\alpha, \beta)} \tilde{T}_X, \quad (3.13)$$

whose middle cohomology is isomorphic to  $\mathcal{U}^{2\omega_4}(-1)$ . In particular, the kernel of (3.12) is an extension of  $\tilde{T}_X(-1)$  and  $\mathcal{U}^{2\omega_4}(-1)$ .

*Proof.* Let us consider the diagram

$$\begin{array}{ccc} \tilde{T}_X(-1) & \xrightarrow{\alpha^t} & V^{\omega_1} \otimes \mathcal{O} \\ \beta^t \downarrow & & \downarrow \alpha \\ \mathcal{O} & \xrightarrow{\beta} & \tilde{T}_X \end{array} \quad (3.14)$$

Here  $\alpha$  is the restriction of (3.12) to  $V^{\omega_1} \otimes \mathcal{O}$ ,  $\beta$  is provided by (3.3) and  $\alpha^t, \beta^t$  are their duals; we also used (3.6) and the self-duality of  $V^{\omega_1} \otimes \mathcal{O}$ . Since all the maps in (3.14) are

G-equivariant, their compositions  $\beta \circ \beta^t$  and  $\alpha \circ \alpha^t$  are also G-equivariant. Note however that the space of equivariant morphisms from  $\tilde{T}_X(-1)$  to  $\tilde{T}_X$  is one-dimensional and is generated by the natural morphism  $\tilde{T}_X(-1) = \tilde{T}_X^\vee \rightarrow \mathcal{O} \rightarrow \tilde{T}_X$  provided by (3.3) and (3.6). Thus, perhaps up to rescaling, this square is commutative.

Taking the total complex of (3.14) we get

$$\tilde{T}_X(-1) \xrightarrow{(\alpha^t, -\beta^t)} (V^{\omega_1} \oplus \mathbb{C}) \otimes \mathcal{O} \xrightarrow{(\alpha, \beta)} \tilde{T}_X$$

Since  $(\alpha, \beta)$  is surjective, the map  $(\alpha^t, -\beta^t)$  is injective. Thus, we have a monad. Taking the alternating sum of ranks of its terms we see that its cohomology is G-equivariant vector bundle of rank 21. Since we have  $V^{\omega_1} = \mathfrak{g}$ , the weights of  $V^{\omega_1}$  are the roots (2.5). Using (2.2) we see that the root  $(1, -1, 0, 0) = 2\omega_4 - \omega_1$  is L-dominant, and since the bundle  $\mathcal{U}^{2\omega_4}(-1)$  is of the required rank 21, the cohomology of the monad has to be  $\mathcal{U}^{2\omega_4}(-1)$ .  $\square$

Applying Lemma 2.6 and Corollaries 2.3 and 2.9 to compute the necessary Ext-groups we obtain the inclusion

$$\mathcal{U}^{\omega_2}(-1) \in \langle \mathcal{U}^{\omega_4}(1), \mathcal{O}(2), \mathcal{U}^{\omega_4}(2), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7) \rangle^\perp. \quad (3.15)$$

Therefore, the left mutation of  $\mathcal{U}^{\omega_2}(-1)$  with respect to  $\langle \mathcal{U}^{\omega_4}(1), \mathcal{O}(2), \mathcal{U}^{\omega_4}(2), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7) \rangle$  is  $\mathcal{U}^{\omega_2}(-1)$  itself. Combining it with Lemma 3.2(3) we get

$$\mathbf{L}_{\langle \mathcal{O}(1), \mathcal{U}^{\omega_4}(1), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7) \rangle}(\mathcal{U}^{\omega_2}(-1)) = T_X,$$

and, therefore, we have

$$T_X \in \langle \mathcal{O}(1), \mathcal{U}^{\omega_4}(1), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7) \rangle^\perp. \quad (3.16)$$

Further, by Lemma 3.2(3) we have  $\mathbf{R}_{\langle \mathcal{O} \rangle}(T_X) = \tilde{T}_X$  and by Lemma 3.1 we have

$$\mathcal{O} \in \langle \mathcal{O}(1), \mathcal{U}^{\omega_4}(1), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7) \rangle^\perp.$$

Hence, we obtain

$$\tilde{T}_X \in {}^\perp\langle \mathcal{O} \rangle \cap \langle \mathcal{O}(1), \mathcal{U}^{\omega_4}(1), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7) \rangle^\perp. \quad (3.17)$$

**Lemma 3.5.** *The sequence of vector bundles  $\tilde{T}_X, \tilde{T}_X(1), \dots, \tilde{T}_X(7)$  is exceptional.*

*Proof.* It is enough to prove

$$\mathrm{Ext}^\bullet(\tilde{T}_X, \tilde{T}_X) = \mathbb{C}[0] \quad (3.18)$$

and

$$\mathrm{Ext}^\bullet(\tilde{T}_X(t), \tilde{T}_X) = 0 \quad \text{for } 1 \leq t \leq 7. \quad (3.19)$$

We split the proof into 3 cases.

*Identity (3.19) for  $2 \leq t \leq 6$ .* Exact sequences (3.2)–(3.3) imply that we have an inclusion

$$\tilde{T}_X \in \mathcal{C} := \langle \mathcal{O}, \mathcal{O}(1), \mathcal{U}^{\omega_2}(-1) \rangle$$

and, therefore, it is enough to show that we have

$$\mathrm{Hom}(\mathcal{C}(t), \mathcal{C}) = 0 \quad \text{for } 2 \leq t \leq 6,$$

which in its turn follows from

$$\begin{aligned}\mathrm{Ext}^\bullet(\mathcal{O}(t), \mathcal{O}) &= H^\bullet(X, \mathcal{O}(-t)) = 0 \quad \text{for } 1 \leq t \leq 7, \\ \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_2}(t-1), \mathcal{O}) &= H^\bullet(X, \mathcal{U}^{\omega_2}(-t-2)) = 0 \quad \text{for } 1 \leq t \leq 6, \\ \mathrm{Ext}^\bullet(\mathcal{O}(t), \mathcal{U}^{\omega_2}(-1)) &= H^\bullet(X, \mathcal{U}^{\omega_2}(-t-1)) = 0 \quad \text{for } 2 \leq t \leq 7, \\ \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_2}(t-1), \mathcal{U}^{\omega_2}(-1)) &= 0 \quad \text{for } 2 \leq t \leq 6.\end{aligned}$$

The first three vanishings hold immediately by Corollary 2.9. To show the last one, using Lemma 2.6 and Corollary 2.3, we rewrite

$$\mathrm{Ext}^\bullet(\mathcal{U}^{\omega_2}(t-1), \mathcal{U}^{\omega_2}(-1)) = H^\bullet(X, \mathcal{U}^{2\omega_2}(-3-t) \oplus \mathcal{U}^{2\omega_3}(-2-t) \oplus \mathcal{U}^{2\omega_4}(-1-t) \oplus \mathcal{O}(-t))$$

and again apply Corollary 2.9.

*Identity (3.19) for  $t = 1$  and  $t = 7$ .* Note that the case  $t = 7$  is equivalent to  $t = 1$  by Serre duality. Thus, we only need to consider the case  $t = 1$ . By Lemma 3.4 it is enough to show vanishings

$$\mathrm{Ext}^\bullet(\tilde{T}_X(1), \mathcal{O}) = 0, \quad \mathrm{Ext}^\bullet(\tilde{T}_X(1), \tilde{T}_X(-1)) = 0, \quad \mathrm{Ext}^\bullet(\tilde{T}_X(1), \mathcal{U}^{2\omega_4}(-1)) = 0.$$

The first one and the last one follow from Lemma 2.6 and Corollaries 2.3 and 2.9 by replacing  $\tilde{T}_X(1)$  with its semisimplification (3.4). The second one follows from (3.19) with  $t = 2$ , which we already know.

*Identity (3.18).* By Lemma 3.4 it is enough to prove

$$\mathrm{Ext}^\bullet(\tilde{T}_X, \mathcal{O}) = 0, \quad \mathrm{Ext}^\bullet(\tilde{T}_X, \tilde{T}_X(-1)) = 0, \quad \mathrm{Ext}^\bullet(\tilde{T}_X, \mathcal{U}^{2\omega_4}(-1)) = \mathbb{k}[-1].$$

The first one holds by (3.17). The second one follows from (3.19) with  $t = 1$ , which we already know. To see the last one, according to (3.4), it is enough to show

$$\begin{aligned}\mathrm{Ext}^\bullet(\mathcal{O}(t), \mathcal{U}^{2\omega_4}(-1)) &= 0, \quad \text{for } 0 \leq t \leq 1, \\ \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_2}(-1), \mathcal{U}^{2\omega_4}(-1)) &= \mathbb{k}[-1],\end{aligned}$$

which follow easily from Lemma 2.6, Corollaries 2.3 and 2.9, and (2.18).  $\square$

**Lemma 3.6.** *Collection (3.1) is exceptional.*

*Proof.* In view of the previous lemmas it is enough to show

$$\mathrm{Ext}^\bullet(\tilde{T}_X(t), \mathcal{O}) = 0 \quad \text{for } 0 \leq t \leq 7$$

and

$$\mathrm{Ext}^\bullet(\tilde{T}_X(t), \mathcal{U}^{\omega_4}) = 0 \quad \text{for } 0 \leq t \leq 7.$$

We treat each of these cases separately.

- (1) For  $t = 0$  this follows from (3.17). For  $1 \leq t \leq 6$  this follows from Corollaries 2.3 and 2.9 by replacing  $\tilde{T}_X$  with its semisimplification (3.4). For  $t = 7$  this follows from the case  $t = 0$  by Serre duality, dualization of factors, and (3.6).
- (2) For  $1 \leq t \leq 7$  this follows from Lemma 2.6 and Corollaries 2.3, 2.9 by replacing  $\tilde{T}_X$  with its semisimplification (3.4).

Let us now assume  $t = 0$ . First we note that dualizing each factor and using Corollary 2.3 and (3.6) we have

$$\mathrm{Ext}^\bullet(\tilde{T}_X, \mathcal{U}^{\omega_4}) \cong \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_4}, \tilde{T}_X).$$



Thus, by Lemma 3.4 it is enough to prove

$$\mathrm{Ext}^\bullet(\mathcal{U}^{\omega_4}, \mathcal{O}) = 0, \quad \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_4}, \tilde{T}_X(-1)) = 0, \quad \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_4}, \mathcal{U}^{2\omega_4}(-1)) = 0.$$

The first one follows from Lemma 3.1. The second one follows from the case  $t = 1$  we have just proved. Finally, the last one follows from Lemma 2.6 and Corollaries 2.3 and 2.9.  $\square$

*Remark 3.7.* We have seen in the proof of the above lemma that the bundles  $\mathcal{U}^{\omega_4}$  and  $\tilde{T}_X$  are completely orthogonal. This also hold for their twists, i.e. we have

$$\mathrm{Ext}^\bullet(\tilde{T}_X(t), \mathcal{U}^{\omega_4}(t)) \cong \mathrm{Ext}^\bullet(\mathcal{U}^{\omega_4}(t), \tilde{T}_X(t)) = 0.$$

#### 4. PROOF OF FULLNESS

Here we prove that the collection (3.1) is full. The proof goes via restriction to some closed subvarieties of  $X$ , where a full exceptional collection is already known. In this respect the proof is similar to the proofs of fullness given in [16, 18].

Let  $\mathcal{D}$  be the full triangulated subcategory of  $\mathbf{D}^b(X)$  generated by the exceptional collection (3.1), i.e. we have

$$\mathcal{D} = \langle \mathcal{O}, \mathcal{U}^{\omega_4}, \tilde{T}_X, \mathcal{O}(1), \mathcal{U}^{\omega_4}(1), \tilde{T}_X(1), \dots, \mathcal{O}(7), \mathcal{U}^{\omega_4}(7), \tilde{T}_X(7) \rangle \subset \mathbf{D}^b(X). \quad (4.1)$$

Since  $\mathcal{D}$  is generated by an exceptional collection, it is admissible. Therefore, there exists a semiorthogonal decomposition

$$\mathbf{D}^b(X) = \langle \mathcal{D}^\perp, \mathcal{D} \rangle.$$

To prove fullness of (3.1) we need to prove  $\mathcal{D}^\perp = 0$ .

This section is structured as follows. In Section 4.1 we begin with some geometric facts that eventually allow us to cover  $X$  by copies of the even orthogonal Grassmannian  $\mathrm{OG}(2, 8)$  (see Lemma 4.4). These facts are also interesting in their own right. In Section 4.2 we collect some auxiliary complexes that allow us to conclude that certain additional  $G$ -equivariant vector bundles are already contained in the subcategory  $\mathcal{D}$  (see Corollary 4.6). Finally, in Section 4.3 we prove fullness by showing in Proposition 4.11 that  $\mathcal{D}^\perp = 0$ .

**4.1. Some geometry.** Since  $X$  is the adjoint variety of a simple group  $G$ , there is a closed embedding

$$X \subset \mathbb{P}(\mathfrak{g}),$$

identifying  $X$  with the orbit of the highest root  $[\Theta] \in \mathbb{P}(\mathfrak{g})$ , where  $G$  acts on its Lie algebra  $\mathfrak{g}$  via the adjoint action. Moreover, since we have an isomorphism of  $G$ -representations  $\mathfrak{g} \cong V^{\omega_1}$ , the above embedding becomes

$$X \subset \mathbb{P}(V^{\omega_1}).$$

Recall from [5, p.202–213] that we have the following formulas for roots of the root systems  $F_4$ ,  $B_4$  and  $D_4$

$$\begin{aligned} F_4: & \quad \pm e_i, \quad \pm e_i \pm e_j, \quad \frac{1}{2}(\pm e_1 \pm e_2 \pm e_3 \pm e_4) \\ B_4: & \quad \pm e_i, \quad \pm e_i \pm e_j \\ D_4: & \quad \pm e_i \pm e_j \end{aligned}$$

The roots for  $F_4$  already appeared in (2.5). From these formulas we clearly see the inclusions of root systems  $D_4 \subset B_4 \subset F_4$ , that give rise to embeddings of the corresponding connected simply connected simple algebraic groups of the corresponding Dynkin types

$$\bar{G} \subset \tilde{G} \subset G.$$

Note that for all three groups we have a common maximal torus

$$T \subset \bar{G} \subset \tilde{G} \subset G. \quad (4.2)$$

We also have embeddings of homogeneous spaces

$$\bar{G}/\bar{P} \subset \tilde{G}/\tilde{P} \subset G/P,$$


where

$$\tilde{P} = P \cap \tilde{G} \quad \text{and} \quad \bar{P} = P \cap \bar{G}.$$

Since the highest root of  $F_4$  is  $\Theta = e_1 + e_2 = \omega_1$  and since its restriction to  $B_4$  (resp.  $D_4$ ) is the second fundamental weight of  $B_4$  (resp.  $D_4$ ), we have

$$\tilde{P} = \tilde{P}_2 \quad \text{and} \quad \bar{P} = \bar{P}_2,$$

where we use the Bourbaki labelling of vertices in Dynkin diagrams  $B_4$  and  $D_4$



and the nodes depicted in solid black correspond to the parabolics  $\tilde{P}$  and  $\bar{P}$  respectively.

For the remainder of this subsection we denote by  $V_{\tilde{G}}^\lambda$  (resp.  $V_{\bar{G}}^\lambda$ ) the irreducible representation of the group  $\tilde{G}$  (resp.  $\bar{G}$ ) with the highest weight  $\lambda$ . Similarly, we denote by  $\tilde{U}^\lambda$  (resp.  $\bar{U}^\lambda$ ) the  $\tilde{G}$ -equivariant (resp.  $\bar{G}$ -equivariant) vector bundle on  $\tilde{G}/\tilde{P}$  (resp.  $\bar{G}/\bar{P}$ ) defined by the irreducible representation with the highest weight  $\lambda$  of the Levi subgroup  $\tilde{L}$  (resp.  $\bar{L}$ ) of the parabolic  $\tilde{P}$  (resp.  $\bar{P}$ ).

A simple computation with root systems (e.g. using [5, p.202–213]) shows that the inclusions  $\bar{G} \subset \tilde{G} \subset G$  induce the following maps on the fundamental weights

$$\begin{array}{lll} \omega_1 \mapsto \tilde{\omega}_2 & \omega_1 \mapsto \bar{\omega}_2 & \tilde{\omega}_1 \mapsto \bar{\omega}_1 \\ \omega_2 \mapsto \tilde{\omega}_1 + \tilde{\omega}_3 & \omega_2 \mapsto \bar{\omega}_1 + \bar{\omega}_3 + \bar{\omega}_4 & \tilde{\omega}_2 \mapsto \bar{\omega}_2 \\ \omega_3 \mapsto \tilde{\omega}_1 + \tilde{\omega}_4 & \omega_3 \mapsto \bar{\omega}_1 + \bar{\omega}_4 & \tilde{\omega}_3 \mapsto \bar{\omega}_3 + \bar{\omega}_4 \\ \omega_4 \mapsto \tilde{\omega}_1 & \omega_4 \mapsto \bar{\omega}_1 & \tilde{\omega}_4 \mapsto \bar{\omega}_4 \end{array} \quad \text{and} \quad (4.3)$$

where we denote by  $\tilde{\omega}_i$  (resp.  $\bar{\omega}_i$ ) the fundamental weights of  $\tilde{G}$  (resp.  $\bar{G}$ ).

Let  $V^\lambda$  be an irreducible highest weight representation of  $G$ . Then its restriction  $V^\lambda|_{\tilde{G}}$  decomposes into a direct sum of irreducible representations of  $\tilde{G}$ . Moreover, the image of  $\lambda$  under (4.3) is the highest weight of one of the summands. The same of course holds

for restrictions from  $G$  to  $\bar{G}$  and from  $\tilde{G}$  to  $\bar{G}$ . For example, this way immediately obtain inclusions

$$V_{\bar{G}}^{\tilde{\omega}_2} \subset V_{\tilde{G}}^{\tilde{\omega}_2} \subset V^{\omega_1}. \quad (4.4)$$

We now give a precise description of the restrictions of  $V^{\omega_1}$  and  $V^{\omega_4}$ .

**Lemma 4.1.** *We have*

$$\begin{aligned} V^{\omega_1}|_{\bar{G}} &= V_{\bar{G}}^{\tilde{\omega}_2} \oplus V_{\bar{G}}^{\tilde{\omega}_4} & V^{\omega_1}|_{\tilde{G}} &= V_{\tilde{G}}^{\tilde{\omega}_2} \oplus V_{\tilde{G}}^{\tilde{\omega}_1} \oplus V_{\tilde{G}}^{\tilde{\omega}_3} \oplus V_{\tilde{G}}^{\tilde{\omega}_4} \\ V^{\omega_4}|_{\bar{G}} &= V_{\bar{G}}^{\tilde{\omega}_1} \oplus V_{\bar{G}}^{\tilde{\omega}_4} \oplus \mathbb{k} & V^{\omega_4}|_{\tilde{G}} &= V_{\tilde{G}}^{\tilde{\omega}_1} \oplus V_{\tilde{G}}^{\tilde{\omega}_3} \oplus V_{\tilde{G}}^{\tilde{\omega}_4} \oplus \mathbb{k}^2 \end{aligned}$$

*Proof.* Let us consider  $V^{\omega_1}|_{\tilde{G}}$ . Note that under (4.3) the weight  $\omega_1$  maps to  $\tilde{\omega}_2$ . Since we have  $\dim V^{\omega_1} = 52$  and  $\dim V_{\tilde{G}}^{\tilde{\omega}_2} = 36$ , the restriction  $V^{\omega_1}|_{\tilde{G}}$  has to be a direct sum of  $V_{\tilde{G}}^{\tilde{\omega}_2}$  and some other irreducible representations of  $\tilde{G}$  of dimension 16. Note, however, that the only irreducible representations of  $\tilde{G}$  of dimension smaller or equal to 30 are  $\mathbb{k}, V_{\tilde{G}}^{\tilde{\omega}_1}, V_{\tilde{G}}^{\tilde{\omega}_4}$ , whose dimensions are 1, 9, 16 respectively. Since the multiplicity of the zero weight in  $V^{\omega_1} = \mathfrak{g}$  is equal to 4, we see that the only possibility is  $V_{\tilde{G}}^{\tilde{\omega}_4}$ .

Similarly one can prove the claim for  $V^{\omega_4}|_{\tilde{G}}$ . Restrictions  $V^{\omega_1}|_{\bar{G}}$  and  $V^{\omega_4}|_{\bar{G}}$  are left as an exercise.  $\square$

**Lemma 4.2.** *The subvarieties  $\tilde{G}/\tilde{P} \subset G/P$  and  $\bar{G}/\bar{P} \subset \tilde{G}/\tilde{P}$  are linear sections in their natural projective embeddings  $G/P \subset \mathbb{P}(V^{\omega_1})$  and  $\tilde{G}/\tilde{P} \subset \mathbb{P}(V_{\tilde{G}}^{\tilde{\omega}_2})$ . More precisely, we have*

$$\tilde{G}/\tilde{P} = G/P \times_{\mathbb{P}(V^{\omega_1})} \mathbb{P}(V_{\tilde{G}}^{\tilde{\omega}_2}) \quad \text{and} \quad \bar{G}/\bar{P} = \tilde{G}/\tilde{P} \times_{\mathbb{P}(V_{\tilde{G}}^{\tilde{\omega}_2})} \mathbb{P}(V_{\tilde{G}}^{\tilde{\omega}_2}),$$

where the inclusions  $V_{\tilde{G}}^{\tilde{\omega}_2} \subset V^{\omega_1}$  and  $V_{\bar{G}}^{\tilde{\omega}_2} \subset V_{\tilde{G}}^{\tilde{\omega}_2}$  are given by (4.4).

*Proof.* Let us first consider the case  $\tilde{G}/\tilde{P} \subset G/P$ . From the classification of symmetric spaces (see for example [22, p. 152]) we see that  $G/\tilde{G}$  is a symmetric space. Hence,  $\tilde{G}$  is a fixed-point subgroup of an involution on  $G$ . Its differential induces a direct sum decomposition of the Lie algebra of  $G$  into eigenspaces

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \quad \text{with} \quad \mathfrak{g}_0 = \text{Lie}(\tilde{G}). \quad (4.5)$$

Since the Dynkin diagram of type  $F_4$  has no automorphisms, the group  $G$  has no outer automorphisms (see [11, Theorem 27.4 and (A.8)]). Hence, this involution is a conjugation by an element  $\tau \in G$  of order 2. Moreover, since the involution fixes  $\tilde{G}$ , it also fixes the maximal torus  $T \subset \tilde{G} \subset G$  (see (4.2)). Since  $C_G(T) = T$ , we obtain  $\tau \in T$ .

Since  $T \subset P$ , the multiplications by  $\tau$  and  $\tau^{-1}$  preserve the parabolic  $P$  and, therefore, the induced action on  $G/P$  is given by left multiplication with  $\tau$ . Applying [20, Theorem A] we see that  $\tilde{G}/\tilde{P} \subset G/P$  is a connected component of the fixed locus  $(G/P)^\tau$ .

The action of the maximal torus  $T$  on  $G/P$  preserves  $(G/P)^\tau$ , since multiplication by  $\tau$  commutes with multiplication by any element of  $T$ . Hence, we have inclusions

$$\tilde{G}/\tilde{P} \subset (G/P)^\tau \subset G/P$$

and the maximal torus  $T$  acts by left multiplication on these spaces in a compatible way. Hence, we also have inclusions on the sets of  $T$ -fixed points

$$\left(\tilde{G}/\tilde{P}\right)^T \subset \left((G/P)^\tau\right)^T \subset \left(G/P\right)^T.$$

The cardinalities of  $\left(\tilde{G}/\tilde{P}\right)^T$  and  $\left(G/P\right)^T$  are equal to the ranks of  $K_0$  for these varieties, which are well well-known to be both equal to 24. Thus, we have equalities

$$\left(\tilde{G}/\tilde{P}\right)^T = \left((G/P)^\tau\right)^T = \left(G/P\right)^T.$$

Therefore, since we already know that  $\tilde{G}/\tilde{P}$  is a connected component of  $(G/P)^\tau$ , we obtain that  $(G/P)^\tau$  is connected and we have the equality

$$\tilde{G}/\tilde{P} = (G/P)^\tau.$$

Consider now the  $G$ -equivariant embedding

$$G/P \subset \mathbb{P}(\mathfrak{g})$$

and note that we have the equality

$$(G/P)^\tau = G/P \times_{\mathbb{P}(\mathfrak{g})} \mathbb{P}(\mathfrak{g})^\tau$$

as a scheme-theoretic intersection. Using (4.5) we see that there is a disjoint union decomposition

$$\mathbb{P}(\mathfrak{g})^\tau = \mathbb{P}(\mathfrak{g}_0) \sqcup \mathbb{P}(\mathfrak{g}_1).$$

Hence, we obtain

$$\tilde{G}/\tilde{P} = G/P \times_{\mathbb{P}(\mathfrak{g})} \mathbb{P}(\mathfrak{g}_0) \quad \text{and} \quad G/P \times_{\mathbb{P}(\mathfrak{g})} \mathbb{P}(\mathfrak{g}_1) = \emptyset.$$

Since we have  $\mathfrak{g}_0 = V_{\tilde{G}}^{\tilde{\omega}_2}$ , the first claim of the lemma is proved.

A very similar argument allows to prove the second claim, since  $\tilde{G}/\tilde{G} \cong \mathrm{SO}_9 / \mathrm{SO}_8$  is again a symmetric space (see [22, p. 152]). We omit the details.  $\square$

**Corollary 4.3.** *We have*

$$\bar{G}/\bar{P} = G/P \times_{\mathbb{P}(V^{\omega_1})} \mathbb{P}(V_{\tilde{G}}^{\tilde{\omega}_2}),$$

where the inclusion  $V_{\tilde{G}}^{\tilde{\omega}_2} \subset V^{\omega_1}$  is given by (4.4).

Our next goal is to prove that  $\bar{G}/\bar{P} \subset G/P$  is the zero locus of a regular section of  $\mathcal{U}^{\omega_4}$ .

**Lemma 4.4.**

- (i) *Up to a scalar multiple there exists a unique non-zero global section  $s_0 \in V^{\omega_4}$  of the vector bundle  $\mathcal{U}^{\omega_4}$  such that*

$$Z(s_0) = \tilde{G}/\tilde{P}.$$

- (ii) *There exist a one parameter family of global sections  $s_t \in V^{\omega_4}$  of the vector bundle  $\mathcal{U}^{\omega_4}$  such that*

$$Z(s_t) = \bar{G}/\bar{P}.$$

*In particular, these global sections are regular.*

*Proof.* To show (i) we proceed as follows. By Lemma 4.1 we know that  $V^{\omega_4}|_{\tilde{G}}$  has a one-dimensional trivial representation as a direct summand. Let  $s_0 \in V^{\omega_4}$  be any non-zero element in this subspace. Our goal is to show  $Z(s_0) = \tilde{G}/\tilde{P}$ . Consider the exact sequence

$$(\mathcal{U}^{\omega_4})^\vee \xrightarrow{s_0^\vee} \mathcal{O}_{G/P} \rightarrow \mathcal{O}_{Z(s_0)} \rightarrow 0.$$

The image of  $s_0^\vee$  is the ideal sheaf  $I_{Z(s_0)}$  of  $Z(s_0)$ . By Corollary 2.3 we have  $(\mathcal{U}^{\omega_4})^\vee(1) = \mathcal{U}^{\omega_4}$  and, therefore, both  $(\mathcal{U}^{\omega_4})^\vee(1)$  and  $I_{Z(s_0)}(1)$  are globally generated. Hence,  $Z(s_0)$  inside  $G/P$  is a linear section with the subspace  $H^0(G/P, I_{Z(s_0)}(1))^\perp$ . Since  $s_0$  is fixed by  $\tilde{G}$ , the map

$$H^0(G/P, \mathcal{U}^{\omega_4}) = V^{\omega_4} \xrightarrow{s_0^\vee(1)} H^0(G/P, \mathcal{O}(1)) = V^{\omega_1},$$

is  $\tilde{G}$ -equivariant and non-zero. Hence, by Lemma 4.1 its image is  $V_{\tilde{G}}^{\tilde{\omega}_4}$  and we get the desired

$$H^0(G/P, I_{Z(s_0)}(1))^\perp = V_{\tilde{G}}^{\tilde{\omega}_2}.$$

Finally, applying Lemma 4.2 we get the claim.

The proof of (ii) is similar but the argument becomes a bit more subtle. By Lemma 4.1 we know that  $V^{\omega_4}|_{\tilde{G}}$  has a two-dimensional trivial representation as a direct summand. Let us denote this subspace  $H \subset V^{\omega_4}$ ; the section  $s_0$  considered in part (i) is contained in  $H$ . As in the proof of (i), any non-zero element  $s \in H$  defines a non-zero  $\tilde{G}$ -equivariant map

$$H^0(G/P, \mathcal{U}^{\omega_4}) = V^{\omega_4} \xrightarrow{s^\vee(1)} H^0(G/P, \mathcal{O}(1)) = V^{\omega_1}, \quad (4.6)$$

and to prove the claim of the lemma, we need to determine the dimension of its image for varying sections  $s$ . Since by Lemma 4.1 the representations  $V^{\omega_4}|_{\tilde{G}}$  and  $V^{\omega_1}|_{\tilde{G}}$  have three irreducible direct summands in common, we have to be more careful than in part (i).

Let us consider the normalizer  $N_G(\tilde{G})$ . From the fact that  $G$  and  $\tilde{G}$  have a common maximal torus (4.2) and from the identification of the Weyl group of  $F_4$  with the automorphism group of  $D_4$  (see [5, p. 213]) it follows that  $N_G(\tilde{G})$  is a semidirect product of  $\tilde{G}$  and the symmetric group  $S_3$ . The group  $S_3$  acts on  $\tilde{G}$  by the outer automorphisms that correspond to permutations of the nodes 1, 3, 4 of the Dynkin diagram of type  $D_4$ .

Let us now consider the unique  $G$ -equivariant map

$$\varphi: V^{\omega_1} \otimes V^{\omega_4} \rightarrow V^{\omega_4} \quad (4.7)$$

given by the action of the Lie algebra  $\mathfrak{g} = V^{\omega_1}$  on its representation  $V^{\omega_4}$ . For any non-zero  $h \in H$  we can consider the specialisations

$$\begin{aligned} \varphi_h: V^{\omega_1} &\longrightarrow V^{\omega_4} \\ v &\mapsto \varphi(v \otimes h) \end{aligned} \quad (4.8)$$

which are nothing else but the duals of the maps (4.6). Thus, to answer our original question about (4.6) we can study kernels of the maps (4.8).

From part (i) we know that there exist  $s_0 \in H$  such that  $\dim(\ker \varphi_{s_0}) = 36$ . More precisely, by Lemma 4.1 we have  $\ker \varphi_{s_0} = V^{\tilde{\omega}_2} \oplus V^{\tilde{\omega}_i}$  for some  $i \in \{1, 3, 4\}$ . Acting on  $s_0$  by  $S_3 = N_G(\tilde{G})/\tilde{G}$  and using the  $G$ -equivariance of the map (4.7) we obtain that on the projective line  $\mathbb{P}(H)$  there exist three points  $[h] \in \mathbb{P}(H)$  such that  $\dim(\ker \varphi_h) = 36$  and for these points we have  $\ker \varphi_h = V^{\tilde{\omega}_2} \oplus V^{\tilde{\omega}_i}$  with distinct  $i \in \{1, 3, 4\}$ . Finally, since (4.8) depends linearly on  $h$ , we conclude that these three points are the only points on  $\mathbb{P}(H)$  with  $\dim(\ker \varphi_h) = 36$  and for all the other points we have  $\ker \varphi_h = V^{\tilde{\omega}_2}$  and  $\dim(\ker \varphi_h) = 28$ .

Going back to the maps (4.6) we see that outside of the three points on  $\mathbb{P}(H)$  constructed above we have the desired

$$H^0(G/P, I_{Z(s)}(1))^\perp = V_{\bar{G}}^{\omega_2}.$$

Finally, applying Lemma 4.2 we get the claim.  $\square$

## 4.2. Auxiliary complexes.

### Lemma 4.5.

(1) *There is a G-equivariant monad*

$$\mathcal{U}^{\omega_4}(-1) \rightarrow V^{\omega_4} \otimes \mathcal{O} \rightarrow \mathcal{U}^{\omega_4}, \quad (4.9)$$

*whose middle cohomology is isomorphic to  $\mathcal{U}^{\omega_3}(-1)$ .*

(2) *There is a G-equivariant monad*

$$\mathcal{U}^{2\omega_4}(-1) \oplus \mathcal{U}^{\omega_3}(-1) \oplus \mathcal{O} \rightarrow V^{\omega_4} \otimes \mathcal{U}^{\omega_4} \rightarrow \mathcal{U}^{2\omega_4} \oplus \mathcal{U}^{\omega_3} \oplus \mathcal{O}(1), \quad (4.10)$$

*whose middle cohomology is isomorphic to  $\mathcal{U}^{\omega_3+\omega_4}(-1) \oplus \mathcal{U}^{\omega_2}(-1) \oplus \mathcal{U}^{\omega_4}$ .*

(3) *There is a G-equivariant monad*

$$\mathcal{U}^{\omega_2+\omega_4}(-1) \oplus \mathcal{U}^{\omega_3} \rightarrow V^{\omega_4} \otimes \mathcal{U}^{\omega_2} \rightarrow \mathcal{U}^{\omega_2+\omega_4} \oplus \mathcal{U}^{\omega_3}(1), \quad (4.11)$$

*whose middle cohomology is isomorphic to  $\mathcal{U}^{\omega_2+\omega_3}(-1) \oplus \mathcal{U}^{\omega_3+\omega_4} \oplus \mathcal{U}^{\omega_4}(1)$ .*

(4) *There is a G-equivariant monad*

$$\mathcal{U}^{\omega_3+\omega_4}(-1) \oplus \mathcal{U}^{\omega_2}(-1) \oplus \mathcal{U}^{\omega_4} \rightarrow V^{\omega_4} \otimes \mathcal{U}^{\omega_3} \rightarrow \mathcal{U}^{\omega_3+\omega_4} \oplus \mathcal{U}^{\omega_2} \oplus \mathcal{U}^{\omega_4}(1), \quad (4.12)$$

*whose middle cohomology is isomorphic to  $\mathcal{U}^{2\omega_3}(-1) \oplus \mathcal{U}^{\omega_2+\omega_4}(-1) \oplus \mathcal{U}^{2\omega_4} \oplus \mathcal{U}^{\omega_3} \oplus \mathcal{O}(1)$ .*

(5) *There is a G-equivariant complex*

$$\mathcal{U}^{2\omega_4}(-2) \rightarrow V^{\omega_4} \otimes \mathcal{U}^{\omega_4}(-1) \rightarrow (\Lambda^2 V^{\omega_4} \otimes \mathcal{O}) \oplus (\mathcal{U}^{2\omega_4}(-1) \oplus \mathcal{U}^{\omega_3}(-1) \oplus \mathcal{O}) \rightarrow V^{\omega_4} \otimes \mathcal{U}^{\omega_4} \rightarrow \mathcal{U}^{2\omega_4} \quad (4.13)$$

*whose only (middle) cohomology is isomorphic to  $\mathcal{U}^{2\omega_4}(-1) \oplus \mathcal{U}^{\omega_2+\omega_4}(-2)$ .*

(6) *There is a G-equivariant complex*

$$\begin{aligned} \mathcal{U}^{3\omega_4}(-3) \rightarrow V^{\omega_4} \otimes \mathcal{U}^{2\omega_4}(-2) \rightarrow (\mathcal{U}^{3\omega_4}(-2) \oplus \mathcal{U}^{\omega_3+\omega_4}(-2) \oplus \mathcal{U}^{\omega_4}(-1)) \oplus (\Lambda^2 V^{\omega_4} \otimes \mathcal{U}^{\omega_4}(-1)) \rightarrow \\ (\Lambda^3 V^{\omega_4} \otimes \mathcal{O}) \oplus (V^{\omega_4} \otimes (\mathcal{U}^{2\omega_4}(-1) \oplus \mathcal{U}^{\omega_3}(-1) \oplus \mathcal{O})) \rightarrow \\ (\mathcal{U}^{3\omega_4}(-1) \oplus \mathcal{U}^{\omega_3+\omega_4}(-1) \oplus \mathcal{U}^{\omega_4}) \oplus (\Lambda^2 V^{\omega_4} \otimes \mathcal{U}^{\omega_4}) \rightarrow V^{\omega_4} \otimes \mathcal{U}^{2\omega_4} \rightarrow \mathcal{U}^{3\omega_4} \end{aligned} \quad (4.14)$$

*whose only (middle) cohomology is isomorphic to*

$$\mathcal{U}^{2\omega_2}(-3) \oplus \mathcal{U}^{\omega_3+2\omega_4}(-2) \oplus \mathcal{U}^{\omega_2+\omega_4}(-2) \oplus \mathcal{U}^{2\omega_4}(-1).$$

(7) *There is a G-equivariant complex*

$$\begin{aligned} \mathcal{U}^{3\omega_4}(-2) \oplus \mathcal{U}^{\omega_3+\omega_4}(-2) \oplus \mathcal{U}^{\omega_4}(-1) \rightarrow V^{\omega_4} \otimes (\mathcal{U}^{2\omega_4}(-1) \oplus \mathcal{U}^{\omega_3}(-1) \oplus \mathcal{O}) \rightarrow \\ \rightarrow (\Lambda^2 V^{\omega_4} \otimes \mathcal{U}^{\omega_4}) \oplus (\mathcal{U}^{3\omega_4}(-1) \oplus \mathcal{U}^{\omega_3+\omega_4}(-1) \oplus \mathcal{U}^{\omega_4}) \oplus (\mathcal{U}^{\omega_3+\omega_4}(-1) \oplus \mathcal{U}^{\omega_2}(-1) \oplus \mathcal{U}^{\omega_4}) \oplus \mathcal{U}^{\omega_4} \rightarrow \\ \rightarrow V^{\omega_4} \otimes (\mathcal{U}^{2\omega_4} \oplus \mathcal{U}^{\omega_3} \oplus \mathcal{O}(1)) \rightarrow \mathcal{U}^{3\omega_4} \oplus \mathcal{U}^{\omega_3+\omega_4} \oplus \mathcal{U}^{\omega_4}(1) \end{aligned} \quad (4.15)$$

whose only (middle) cohomology is isomorphic to

$$\begin{aligned} & (\mathcal{U}^{3\omega_4}(-1) \oplus \mathcal{U}^{\omega_3+\omega_4}(-1) \oplus \mathcal{U}^{\omega_4}) \oplus \\ & \oplus \mathcal{U}^{\omega_2+2\omega_4}(-2) \oplus \mathcal{U}^{\omega_2+\omega_3}(-2) \oplus \mathcal{U}^{\omega_3+\omega_4}(-1) \oplus \mathcal{U}^{\omega_2}(-1) \end{aligned}$$

*Proof.* Let us first consider (4.9). Since the vector bundle  $\mathcal{U}^{\omega_4}$  is globally generated by  $H^0(X, \mathcal{U}^{\omega_4}) = V^{\omega_4}$ , the canonical G-equivariant morphism  $V^{\omega_4} \otimes \mathcal{O} \rightarrow \mathcal{U}^{\omega_4}$  is surjective. Taking its dual, using Corollary 2.3 and the selfduality of the G-representation  $V^{\omega_4}$ , we get the G-equivariant embedding  $\mathcal{U}^{\omega_4}(-1) \rightarrow V^{\omega_4} \otimes \mathcal{O}$ . The composition

$$\mathcal{U}^{\omega_4}(-1) \rightarrow V^{\omega_4} \otimes \mathcal{O} \rightarrow \mathcal{U}^{\omega_4}$$

vanishes, since it is a G-equivariant morphism between two non-isomorphic irreducible G-equivariant vector bundles. Thus, we see that (4.9) is indeed a monad.

A direct computation with weights of  $V^{\omega_4}$  shows that among all 26 weights only

$$\omega_4, \omega_4 - \omega_1, \omega_3 - \omega_1, 0, 0$$

are L-dominant. Since the rank of the cohomology of the monad has to be  $26 - 6 - 6 = 14$ , we see that  $\mathcal{U}^{\omega_3}(-1)$  is the only possibility.

The remaining monads are derived from (4.9) by linear algebra operations:

- (4.10) is obtained from (4.9) by tensoring with  $\mathcal{U}^{\omega_4}$ ;
- (4.11) is obtained from (4.9) by tensoring with  $\mathcal{U}^{\omega_2}$ ;
- (4.12) is obtained from (4.9) by tensoring with  $\mathcal{U}^{\omega_3}$ ;
- (4.13) is the exterior square of (4.9);
- (4.14) is the exterior cube of (4.9);
- (4.15) is obtained from (4.13) by tensoring with  $\mathcal{U}^{\omega_4}$ .

Tensor products, exterior and symmetric powers required for the above computations can be found in Lemmas 2.6 and 2.7.  $\square$

From the above lemma we deduce the following corollary.

**Corollary 4.6.** *We have containments*

$$\mathcal{U}^{\omega_3}(m) \in \mathcal{D} \quad \text{for } m \in [0, 6] \tag{4.16}$$

$$\mathcal{U}^{\omega_2}(m) \in \mathcal{D} \quad \text{for } m \in [-1, 5] \tag{4.17}$$

$$\mathcal{U}^{2\omega_4}(m) \in \mathcal{D} \quad \text{for } m \in [0, 6] \tag{4.18}$$

$$\mathcal{U}^{\omega_2+\omega_4}(m) \in \mathcal{D} \quad \text{for } m \in [0, 4] \tag{4.19}$$

$$\mathcal{U}^{\omega_3+\omega_4}(m) \in \mathcal{D} \quad \text{for } m \in [0, 5] \tag{4.20}$$

$$\mathcal{U}^{\omega_2+\omega_3}(m) \in \mathcal{D} \quad \text{for } m \in [0, 3] \tag{4.21}$$

$$\mathcal{U}^{2\omega_3}(m) \in \mathcal{D} \quad \text{for } m \in [0, 4] \tag{4.22}$$

$$\mathcal{U}^{3\omega_4}(m) \in \mathcal{D} \quad \text{for } m \in [1, 4] \tag{4.23}$$

$$\mathcal{U}^{\omega_3+2\omega_4}(m) \in \mathcal{D} \quad \text{for } m \in [1, 3] \tag{4.24}$$

$$\mathcal{U}^{2\omega_2}(1) \in \mathcal{D} \tag{4.25}$$

$$\mathcal{U}^{\omega_2+2\omega_4}(m) \in \mathcal{D} \quad \text{for } m \in [1, 2] \tag{4.26}$$

*Proof.* We treat each case separately:

- (4.16) follows from (4.9) twisted by  $\mathcal{O}(t)$  for  $1 \leq t \leq 7$  and (4.1).
- (4.17) follows from (3.2), (3.3) twisted by  $\mathcal{O}(t)$  for  $0 \leq t \leq 6$  and (4.1).
- (4.18) follows from (3.13) twisted by  $\mathcal{O}(t)$  for  $1 \leq t \leq 7$  and (4.1).
- (4.19) follows from (4.13) twisted by  $\mathcal{O}(t)$  for  $2 \leq t \leq 6$ , (4.18), (4.16), and (4.1).
- (4.20) follows from (4.10) twisted by  $\mathcal{O}(t)$  for  $1 \leq t \leq 6$ , (4.18), (4.16), and (4.1).
- (4.21) follows from (4.11) twisted by  $\mathcal{O}(t)$  for  $1 \leq t \leq 4$ , (4.16), (4.17), and (4.19).
- (4.22) follows from (4.12) twisted by  $\mathcal{O}(t)$  for  $1 \leq t \leq 5$ , (4.16), (4.17), (4.20), (4.1).
- To show (4.23) we proceed as follows. Tensoring (3.13) with  $\mathcal{U}^{\omega_4}$  we get a monad

$$\tilde{T}_X \otimes \mathcal{U}^{\omega_4}(-1) \rightarrow (V^{\omega_1} \oplus \mathbb{C}) \otimes \mathcal{U}^{\omega_4} \rightarrow \tilde{T}_X \otimes \mathcal{U}^{\omega_4},$$

with cohomology

$$\mathcal{U}^{3\omega_4}(-1) \oplus \mathcal{U}^{\omega_3+\omega_4}(-1) \oplus \mathcal{U}^{\omega_4}.$$

Note also that by (3.2)–(3.3) and Lemma 2.6 we have

$$\text{ss}(\tilde{T}_X \otimes \mathcal{U}^{\omega_4}) = \mathcal{U}^{\omega_4} \oplus \mathcal{U}^{\omega_4}(1) \oplus \mathcal{U}^{\omega_2+\omega_4}(-1) \oplus \mathcal{U}^{\omega_3}.$$

Finally, twisting the above monad by  $\mathcal{O}(t)$  for  $2 \leq t \leq 5$  and using (4.16), (4.19), and (4.1), we get (4.23).

- To show (4.24) we proceed as follows. Tensoring (3.13) with  $\mathcal{U}^{\omega_3}$  we get a monad

$$\tilde{T}_X \otimes \mathcal{U}^{\omega_3}(-1) \rightarrow (V^{\omega_1} \oplus \mathbb{C}) \otimes \mathcal{U}^{\omega_3} \rightarrow \tilde{T}_X \otimes \mathcal{U}^{\omega_3},$$

with cohomology

$$\mathcal{U}^{\omega_3+2\omega_4}(-1) \oplus \mathcal{U}^{\omega_2+\omega_4}(-1) \oplus \mathcal{U}^{2\omega_4} \oplus \mathcal{U}^{\omega_3}$$

Note also that by (3.2)–(3.3) and Lemma 2.6 we have

$$\text{ss}(\tilde{T}_X \otimes \mathcal{U}^{\omega_3}) = \mathcal{U}^{\omega_3} \oplus \mathcal{U}^{\omega_3}(1) \oplus \mathcal{U}^{\omega_2+\omega_3}(-1) \oplus \mathcal{U}^{\omega_3+\omega_4} \oplus \mathcal{U}^{\omega_4}(1).$$

Finally, twisting the above monad by  $\mathcal{O}(t)$  for  $2 \leq t \leq 4$  and using (4.16), (4.20), (4.21), and (4.1), we get (4.24).

- (4.25) follows from (4.14) twisted by  $\mathcal{O}(4)$ , (4.16), (4.18), (4.20) (4.23), and (4.1).
- (4.26) follows from (4.15) twisted by  $\mathcal{O}(t)$  for  $3 \leq t \leq 4$ , (4.16)–(4.18), (4.20), (4.23), and (4.1).

□

**4.3. Proof of fullness.** In this section we finally prove that the collection (3.1) is full by restricting it to a family of subvarieties of  $X$  of the form  $\text{OG}(2, 8)$  for which a convenient full exceptional collection is already known by [18].

Let  $Z = \bar{G}/\bar{P} \cong \text{OG}(2, 8)$  and recall that by Lemma 4.4(2) we have a natural embedding

$$i_\varphi: Z \hookrightarrow X,$$

as the zero locus of a regular section  $\varphi \in H^0(X, \mathcal{U}^{\omega_4}) = V^{\omega_4}$ . Acting by an element  $g \in G$  we get a shifted embedding

$$i_{\varphi_g}: gZ \hookrightarrow X$$

as the zero locus of the section  $\varphi_g = g\varphi$ , where  $gZ \cong Z$ . Since the action of  $G$  on  $X$  is transitive, we have the following simple fact.

**Lemma 4.7.** *Varying  $g \in G$  we can sweep out the whole  $X$  by  $gZ \cong \bar{G}/\bar{P}$ .*

As a corollary we now get the following lemma.



**Lemma 4.8** ([16, Lemma 4.5]). *If for an object  $F \in \mathbf{D}^b(X)$  the restrictions  $i_{\varphi_g}^* F$  vanish for all  $\varphi_g \in H^0(X, \mathcal{U}^{\omega_4})$  as above, then  $F = 0$ .*

Let us now recall a known exceptional collection for  $Z = \text{OG}(2, 8)$ . Consider full triangulated subcategories

$$\mathcal{A} = \langle \mathcal{O}, \mathcal{U}^{\bar{\omega}_1}, \mathcal{U}^{\bar{\omega}_3}, \mathcal{U}^{\bar{\omega}_4} \rangle \subset \mathbf{D}^b(Z)$$

and

$$\mathcal{R} = \langle \mathcal{U}^{2\bar{\omega}_1}(-1), \mathcal{U}^{2\bar{\omega}_3}(-1), \mathcal{U}^{2\bar{\omega}_4}(-1), \tilde{T}_Z(-1) \rangle \subset \mathbf{D}^b(Z),$$

where  $\tilde{T}_Z$  is the extension of the tangent bundle  $T_Z$  by  $\mathcal{O}_Z$  analogous to (3.3). Then, by [18, Remark 3.23] there exists an full exceptional collection of the form

$$\mathbf{D}^b(Z) = \langle \mathcal{R}, \mathcal{A}, \mathcal{A}(1), \mathcal{A}(2), \mathcal{A}(3), \mathcal{A}(4) \rangle.$$

The left mutation of the subcategory  $\langle \mathcal{A}(2), \mathcal{A}(3), \mathcal{A}(4) \rangle$  via  $\langle \mathcal{R}, \mathcal{A}, \mathcal{A}(1) \rangle$  is given by the twist by  $\mathcal{O}(-5)$  and we obtain the decomposition

$$\mathbf{D}^b(Z) = \langle \mathcal{A}(-3), \mathcal{A}(-2), \mathcal{A}(-1), \mathcal{R}, \mathcal{A}(1), \mathcal{A}(2) \rangle.$$

Finally, twisting by  $\mathcal{O}(3)$  we arrive at

$$\mathbf{D}^b(Z) = \langle \mathcal{A}, \mathcal{A}(1), \mathcal{A}(2), \mathcal{R}(3), \mathcal{A}(3), \mathcal{A}(4) \rangle. \quad (4.27)$$

This form of the collection will be used in our proof of fullness below.

We also need the following lemma that computes restrictions of (3.1) to  $Z$ .

**Lemma 4.9.** *There exist isomorphisms*

$$i_{\varphi_g}^* \mathcal{U}^{\omega_4} = \bar{\mathcal{U}}^{\bar{\omega}_1} \oplus \bar{\mathcal{U}}^{\bar{\omega}_3} \oplus \bar{\mathcal{U}}^{\bar{\omega}_4} \quad (4.28)$$

$$i_{\varphi_g}^* \mathcal{U}^{2\omega_4} = \bar{\mathcal{U}}^{2\bar{\omega}_1} \oplus \bar{\mathcal{U}}^{2\bar{\omega}_3} \oplus \bar{\mathcal{U}}^{2\bar{\omega}_4} \oplus \bar{\mathcal{U}}^{\bar{\omega}_1+\bar{\omega}_3} \oplus \bar{\mathcal{U}}^{\bar{\omega}_1+\bar{\omega}_4} \oplus \bar{\mathcal{U}}^{\bar{\omega}_3+\bar{\omega}_4} \quad (4.29)$$

$$i_{\varphi_g}^* \tilde{T}_X = \tilde{T}_{gZ} \oplus i_{\varphi_g}^* \mathcal{U}^{\omega_4} \quad (4.30)$$

*Proof.* It is enough to prove the claims for  $g = 1$ , i.e. we are considering the natural embedding  $i: \bar{G}/\bar{P} \subset G/P$ .

To prove (4.28) we proceed as follows. Since  $\mathcal{U}^{\omega_4}$  is irreducible, the unipotent radical of the parabolic  $P$  acts trivially on the  $P$ -representation  $E$  corresponding to  $\mathcal{U}^{\omega_4}$  under the equivalence (2.1). The  $\bar{P}$ -representation corresponding to  $i^*\mathcal{U}^{\omega_4}$  under (2.1) is simply the restriction  $\text{Res}_{\bar{P}}^P(E)$  of the  $P$ -representation  $E$  to  $\bar{P}$ , and, since the unipotent radical of  $\bar{P}$  is contained in the unipotent radical of  $P$ , the latter also acts trivially on  $E$ . Therefore,  $\text{Res}_{\bar{P}}^P(E)$  is completely reducible and we obtain the complete reducibility of  $i^*\mathcal{U}^{\omega_4}$ . Finally, since  $\mathcal{U}^{\omega_4}$  is generated by its global sections  $H^0(X, \mathcal{U}^{\omega_4}) = V^{\omega_4}$ , the restriction  $i^*\mathcal{U}^{\omega_4}$  is also globally generated. Using the description of  $V^{\omega_4}|_{\bar{G}}$  given in Lemma 4.1 and the Borel–Bott–Weil theorem we get the desired claim.

To prove (4.29) we note that according Lemma 2.7 we have  $\mathcal{U}^{2\omega_4} = S^2\mathcal{U}^{\omega_4}$ . Hence, we have  $i^*\mathcal{U}^{2\omega_4} = S^2(i^*\mathcal{U}^{\omega_4})$ . We leave the rest as an exercise.

To prove (4.30) we proceed as follows. First, we claim that all the statements of Lemma 3.2 hold verbatim for  $T_Z$  and  $\tilde{T}_Z$  if we replace  $\mathcal{U}^{\omega_2}(-1)$  with  $\bar{\mathcal{U}}^{\bar{\omega}_1+\bar{\omega}_3+\bar{\omega}_4}(-1)$ ; the proof is completely analogous. In particular, we have  $\text{Ext}^\bullet(T_Z, \mathcal{O}) = \mathbb{k}[-1]$  and  $\left(\tilde{T}_Z\right)^\vee = \tilde{T}_Z(-1)$ .

From (4.27) and (4.28) we have

$$\mathrm{Ext}^\bullet(\tilde{T}_Z, i^*\mathcal{U}^{\omega_4}) = 0.$$

Dualizing both sides and using the isomorphisms  $(\tilde{T}_Z)^\vee = \tilde{T}_Z(-1)$  and  $(i^*\mathcal{U}^{\omega_4})^\vee = i^*\mathcal{U}^{\omega_4}(-1)$  we get

$$\mathrm{Ext}^\bullet(i^*\mathcal{U}^{\omega_4}, \tilde{T}_Z) = 0.$$

Since by (4.27)–(4.28) we also have  $\mathrm{Ext}^\bullet(i^*\mathcal{U}^{\omega_4}, \mathcal{O}) = 0$ , we obtain  $\mathrm{Ext}^\bullet(i^*\mathcal{U}^{\omega_4}, T_Z) = 0$  and the normal bundle exact sequence

$$0 \rightarrow T_Z \rightarrow i^*T_X \rightarrow i^*\mathcal{U}^{\omega_4} \rightarrow 0 \quad (4.31)$$

splits. Moreover, since we also have  $\mathrm{Ext}^\bullet(i^*T_X, \mathcal{O}) = \mathrm{Ext}^\bullet(T_Z, \mathcal{O}) = \mathbb{k}[-1]$ , we get another split short exact sequence

$$0 \rightarrow \tilde{T}_Z \rightarrow i^*\tilde{T}_X \rightarrow i^*\mathcal{U}^{\omega_4} \rightarrow 0. \quad (4.32)$$

This finishes the proof.  $\square$

We will deduce fullness from the following key fact.

**Lemma 4.10.** *Let  $\Sigma$  be the set of vector bundles defined as*

$$\Sigma := \{\mathcal{O}(t), \mathcal{U}^{\omega_4}(t), \mathcal{U}^{2\omega_4}(2), \tilde{T}_X(2) \mid 0 \leq t \leq 4\}.$$

*Then for any vector bundle  $E \in \Sigma$  we have inclusions*

$$E \otimes \Lambda^k \mathcal{U}^{\omega_4} \in \mathcal{D} \quad \forall k, \quad (4.33)$$

*where the subcategory  $\mathcal{D}$  is defined in (4.1).*

*Proof.* We consider four cases:

- (1) **Case**  $E \in \{\mathcal{O}(t) \mid 0 \leq t \leq 4\}$ . Using Lemma 2.7 we see that the irreducible summands of  $\Lambda^k \mathcal{U}^{\omega_4}(t)$  for  $0 \leq t \leq 4$  are

$$\begin{aligned} \mathcal{O}(m) & \text{ for } m \in [0, 7], \\ \mathcal{U}^{\omega_4}(m) & \text{ for } m \in [0, 6], \\ \mathcal{U}^{\omega_3}(m) & \text{ for } m \in [0, 5], \\ \mathcal{U}^{\omega_2}(m) & \text{ for } m \in [0, 4]. \end{aligned}$$

Hence, the desired inclusion (4.33) follows from (4.16)–(4.17).

- (2) **Case**  $E \in \{\mathcal{U}^{\omega_4}(t) \mid 0 \leq t \leq 4\}$ . Using Lemmas 2.6 and 2.7 we see that the irreducible summands of  $\mathcal{U}^{\omega_4}(t) \otimes \Lambda^k \mathcal{U}^{\omega_4}$  for  $0 \leq t \leq 4$  are

$$\begin{aligned} \mathcal{O}(m) & \text{ for } m \in [1, 7], \\ \mathcal{U}^{\omega_4}(m) & \text{ for } m \in [0, 7], \\ \mathcal{U}^{\omega_3}(m) & \text{ for } m \in [0, 6], \\ \mathcal{U}^{\omega_2}(m) & \text{ for } m \in [0, 5], \\ \mathcal{U}^{2\omega_4}(m) & \text{ for } m \in [0, 6], \\ \mathcal{U}^{\omega_3+\omega_4}(m) & \text{ for } m \in [0, 5], \\ \mathcal{U}^{\omega_2+\omega_4}(m) & \text{ for } m \in [0, 4]. \end{aligned}$$

Hence, the desired inclusions (4.33) hold by Corollary 4.6.

- (3) **Case**  $E = \mathcal{U}^{2\omega_4}(2)$ . Using Lemmas 2.6 and 2.7 we see that the irreducible summands of  $\mathcal{U}^{2\omega_4}(2) \otimes \Lambda^k \mathcal{U}^{\omega_4}$  are

$$\begin{aligned} & \mathcal{U}^{2\omega_4}(m) \quad \text{for } m \in [2, 5], \\ & \mathcal{U}^{3\omega_4}(m) \quad \text{for } m \in [2, 4], \\ & \mathcal{U}^{\omega_3+\omega_4}(m) \quad \text{for } m \in [2, 4], \\ & \mathcal{U}^{\omega_4}(m) \quad \text{for } m \in [3, 5], \\ & \mathcal{U}^{\omega_3+2\omega_4}(m) \quad \text{for } m \in [2, 3], \\ & \mathcal{U}^{\omega_2+\omega_4}(m) \quad \text{for } m \in [2, 3], \\ & \mathcal{U}^{\omega_3}(m) \quad \text{for } m \in [3, 4], \\ & \mathcal{U}^{\omega_2+2\omega_4}(2), \\ & \mathcal{U}^{\omega_2}(3). \end{aligned}$$

Hence, the desired inclusions (4.33) hold by Corollary 4.6.

- (4) **Case**  $E = \tilde{T}_X(2)$ . To show the desired inclusions (4.33) it suffices to work with the direct summands of the semisimplification  $\text{ss}(\tilde{T}_X)(2)$ . By (3.2)–(3.3) we have

$$\text{ss}(\tilde{T}_X)(2) = \mathcal{U}^{\omega_2}(1) \oplus \mathcal{O}(2) \oplus \mathcal{O}(3).$$

Using Lemmas 2.6 and 2.7 we see that the irreducible summands of  $\text{ss}(\tilde{T}_X)(2) \otimes \Lambda^k \mathcal{U}^{\omega_4}$  are

$$\begin{aligned} & \mathcal{O}(m) \quad \text{for } m \in [2, 6], \\ & \mathcal{U}^{\omega_4}(m) \quad \text{for } m \in [2, 5], \\ & \mathcal{U}^{\omega_3}(m) \quad \text{for } m \in [2, 4], \\ & \mathcal{U}^{\omega_2}(m) \quad \text{for } m \in [1, 4], \\ & \mathcal{U}^{\omega_2+\omega_4}(m) \quad \text{for } m \in [1, 3], \\ & \mathcal{U}^{\omega_2+\omega_3}(m) \quad \text{for } m \in [1, 2], \\ & \mathcal{U}^{\omega_3+\omega_4}(m) \quad \text{for } m \in [2, 3], \\ & \mathcal{U}^{2\omega_2}(1), \\ & \mathcal{U}^{2\omega_3}(2), \\ & \mathcal{U}^{2\omega_4}(3). \end{aligned}$$

Hence, the desired inclusions (4.33) hold by Corollary 4.6. □

Finally, we are now ready to prove fullness of (3.1).

**Proposition 4.11.** *If  $F \in \mathbf{D}^b(X)$  is right orthogonal to all vector bundles  $E$  in the exceptional collection (3.1), i.e.  $\text{Ext}^\bullet(E, F) = 0$ , then  $F = 0$ .*

*Proof.* Recall the definition of the set  $\Sigma$  from Lemma 4.10. By our assumptions we have that for any vector bundle  $E \otimes \Lambda^k \mathcal{U}^{\omega_4}$  with  $E \in \Sigma$  there are vanishings

$$\mathrm{Ext}^\bullet(E \otimes \Lambda^k \mathcal{U}^{\omega_4}, F) = H^\bullet(X, (E \otimes \Lambda^k \mathcal{U}^{\omega_4})^\vee \otimes F) = 0. \quad (4.34)$$

Consider the embedding  $i_{\varphi_g}: gZ \hookrightarrow X$ . Since the sections  $\varphi_g$  are regular, we have the Koszul resolution

$$0 \rightarrow \mathcal{O}(-3) \rightarrow \cdots \rightarrow (\Lambda^2 \mathcal{U}^{\omega_4})^\vee \rightarrow (\mathcal{U}^{\omega_4})^\vee \rightarrow \mathcal{O}_X \rightarrow i_{(\varphi_g)*} \mathcal{O}_{gZ} \rightarrow 0. \quad (4.35)$$

Let us now tensor the Koszul complex (4.35) by  $E^\vee \otimes F$  with  $E \in \Sigma$

$$\cdots \rightarrow (\Lambda^2 \mathcal{U}^{\omega_4})^\vee \otimes (E^\vee \otimes F) \rightarrow (\mathcal{U}^{\omega_4})^\vee \otimes (E^\vee \otimes F) \rightarrow E^\vee \otimes F \rightarrow i_{(\varphi_g)*} i_{\varphi_g}^*(E^\vee \otimes F) \rightarrow 0.$$

By (4.34) we have that the cohomology groups of  $(\Lambda^k \mathcal{U}^{\omega_4})^\vee \otimes (E^\vee \otimes F)$  vanish. Hence, the cohomology groups of  $i_{(\varphi_g)*} i_{\varphi_g}^*(E^\vee \otimes F)$  also vanish and we have

$$0 = H^\bullet(X, i_{(\varphi_g)*} i_{\varphi_g}^*(E^\vee \otimes F)) = H^\bullet(\mathrm{OG}(2, 8), i_{\varphi_g}^*(E^\vee \otimes F)) = \mathrm{Ext}^\bullet(i_{\varphi_g}^* E, i_{\varphi_g}^* F) \quad (4.36)$$

Hence,  $i_{\varphi_g}^* F$  is right orthogonal to the objects  $i_{\varphi_g}^* E$  with  $E \in \Sigma$ .

Finally, by Lemma 4.9 and (4.36) we get that  $i_{\varphi_g}^* F$  is right orthogonal to all objects in the collection (4.27). Hence, we obtain  $i_{\varphi_g}^* F = 0$ . Since this holds for any section  $\varphi_g$ , we get  $F = 0$  by Lemma 4.8.  $\square$

## REFERENCES

- [1] Dmitri N. Akhiezer, *Lie group actions in complex analysis*, Aspects of Mathematics, E27, Friedr. Vieweg & Sohn, Braunschweig, 1995.
- [2] A. A. Beilinson, *Coherent sheaves on  $\mathbf{P}^n$  and problems in linear algebra*, Funktsional. Anal. i Prilozhen. **12** (1978), no. 3, 68–69.
- [3] P. Belmans and M. Smirnov, *The Hochschild cohomology of generalised Grassmannians*. Available at [arXiv:1911.09414](https://arxiv.org/abs/1911.09414).
- [4] Pieter Belmans, Alexander Kuznetsov, and Maxim Smirnov, *Derived categories of the Cayley plane and the coadjoint Grassmannian of type F*, Transformation Groups (2021), available at <https://doi.org/10.1007/s00031-021-09657-w>. See also [arXiv:2005.01989](https://arxiv.org/abs/2005.01989).
- [5] N. Bourbaki, *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V: Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines*, Actualités Scientifiques et Industrielles [Current Scientific and Industrial Topics], No. 1337, Hermann, Paris, 1968.
- [6] Pierre-Emmanuel Chaput and Nicolas Perrin, *On the quantum cohomology of adjoint varieties*, Proc. Lond. Math. Soc. (3) **103** (2011), no. 2, 294–330. See also [arXiv:0904.4824](https://arxiv.org/abs/0904.4824).
- [7] Boris Dubrovin, *Geometry and analytic theory of Frobenius manifolds*, Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998), 1998, pp. 315–326.
- [8] Daniele Faenzi and Laurent Manivel, *On the derived category of the Cayley plane II*, Proc. Amer. Math. Soc. **143** (2015), no. 3, 1057–1074. See also [arXiv:1201.6327](https://arxiv.org/abs/1201.6327).
- [9] Anton Fonarev, *Full exceptional collections on Lagrangian Grassmannians*, Int. Math. Res. Not. (2020). Available online ahead of print as [doi:10.1093/imrn/rnaa098](https://doi.org/10.1093/imrn/rnaa098), see also [arXiv:1911.08968](https://arxiv.org/abs/1911.08968).
- [10] Lyalya Guseva, *On the derived category of  $\mathrm{IGr}(3; 8)$* , Mat. Sb. **211** (2020), no. 7, 24–59. See also [arXiv:1810.07777](https://arxiv.org/abs/1810.07777).
- [11] James E. Humphreys, *Linear algebraic groups*, Graduate Texts in Mathematics, No. 21, Springer-Verlag, New York-Heidelberg, 1975.
- [12] Jens Carsten Jantzen, *Representations of algebraic groups*, Pure and Applied Mathematics, vol. 131, Academic Press, Inc., Boston, MA, 1987.
- [13] M. M. Kapranov, *On the derived categories of coherent sheaves on some homogeneous spaces*, Inventiones Mathematicae **92** (1988), no. 3, 479–508.

- [14] Alexander Kuznetsov, *Hyperplane sections and derived categories*, Izv. Ross. Akad. Nauk Ser. Mat. **70** (2006), no. 3, 23–128. See also [arXiv:math/0503700](https://arxiv.org/abs/math/0503700).
- [15] ———, *Homological projective duality*, Publ. Math. Inst. Hautes Études Sci. **105** (2007), 157–220. See also [arXiv:math/0507282](https://arxiv.org/abs/math/0507282).
- [16] ———, *Exceptional collections for Grassmannians of isotropic lines*, Proceedings of the London Mathematical Society. Third Series **97** (2008), no. 1, 155–182. See also [arXiv:math/0512013](https://arxiv.org/abs/math/0512013).
- [17] Alexander Kuznetsov and Alexander Polishchuk, *Exceptional collections on isotropic Grassmannians*, J. Eur. Math. Soc. (JEMS) **18** (2016), no. 3, 507–574. See also [arXiv:1110.5607](https://arxiv.org/abs/1110.5607).
- [18] Alexander Kuznetsov and Maxim Smirnov, *Residual categories for (co)adjoint Grassmannians in classical types*, Compositio Mathematica **157** (2021), no. 6, 1172–1206. See also [arXiv:2001.04148](https://arxiv.org/abs/2001.04148).
- [19] Laurent Manivel, *On the derived category of the Cayley plane*, J. Algebra **330** (2011), 177–187. See also [arXiv:0907.2784](https://arxiv.org/abs/0907.2784).
- [20] R. W. Richardson, *On orbits of algebraic groups and Lie groups*, Bull. Austral. Math. Soc. **25** (1982), no. 1, 1–28.
- [21] The Sage Developers, *Sagemath, the Sage Mathematics Software System (Version 9.0)*, 2020. <https://www.sagemath.org>.
- [22] Dmitry A. Timashev, *Homogeneous spaces and equivariant embeddings*, Encyclopaedia of Mathematical Sciences, vol. 138, Springer, Heidelberg, 2011. Invariant Theory and Algebraic Transformation Groups, 8.
- [23] È. B. Vinberg and A. L. Onishchik, *Seminar po gruppam  $Li$  i algebraicheskim gruppam*, Second, URSS, Moscow, 1995.

UNIVERSITÄT AUGSBURG, INSTITUT FÜR MATHEMATIK, UNIVERSITÄTSSTR. 14, 86159 AUGSBURG,  
GERMANY  
*Email address:* [maxim.smirnov@math.uni-augsburg.de](mailto:maxim.smirnov@math.uni-augsburg.de)  
*Email address:* [maxim.n.smirnov@gmail.com](mailto:maxim.n.smirnov@gmail.com)