

# Universal criticality of thermodynamic geometry for boundary conformal field theories in gauge/gravity duality

---

Morteza Rafiee <sup>a1</sup>, Seyed Ali Hosseini Mansoori <sup>a2</sup>, Shao-Wen Wei <sup>b,c3</sup>, Robert B. Mann <sup>d,e4</sup>,

<sup>a</sup>*Faculty of Physics, Shahrood University of Technology, P.O. Box 3619995161 Shahrood, Iran*

<sup>b</sup>*Institute of Theoretical and Physics Research Center of Gravitation, Lanzhou University, Lanzhou 730000, People's Republic of China*

<sup>c</sup>*Institute of Theoretical Physics and Research Center of Gravitation, Lanzhou University, Lanzhou 730000, China*

<sup>d</sup>*Perimeter Institute, 31 Caroline St. N. Waterloo, Ontario, N2L 2Y5, Canada*

<sup>e</sup>*Department of Physics and Astronomy, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada*

ABSTRACT: According to more recent AdS/CFT interpretation [1], in which varying cosmological constant  $\Lambda$  in the bulk corresponds to varying the curvature radius governing the space on which the field theory resides, we study the criticality of thermodynamic curvatures for thermal boundary conformal field theories (CFT) that are dual to  $d$ -dimensional charged anti-de Sitter (AdS) black holes, embedded in  $D$ -dimensional M-theory/superstring inspired models having  $AdS_d \times S^{d+k}$  spacetime with  $D = 2d + k$ . Analogous with criticality features acquired for charged AdS black holes in the bulk [2], the normalized intrinsic curvature  $R_N$  and extrinsic curvature  $K_N$  of the boundary CFT has critical exponents 2 and 1, respectively. In this respect, the universal amplitude of  $R_N t^2$  is  $\frac{1}{2}$  and  $K_N t$  is  $-\frac{1}{2}$  when  $t \rightarrow 0^-$ , whereas  $R_N t^2 \approx \frac{1}{8}$  and  $K_N t \approx \frac{1}{4}$  in the limit  $t \rightarrow 0^+$  in which  $t = T/T_c - 1$  is the temperature parameter with the critical temperature,  $T_c$ . Interestingly, these critical amplitudes are independent of the number of thermal CFT dimensions and are remarkably similar to one given for higher dimensional charged AdS black holes in the bulk.

---

<sup>1</sup>m.rafee@shahroodut.ac.ir

<sup>2</sup>shosseini@shahroodut.ac.ir

<sup>3</sup>weishw@lzu.edu.cn

<sup>4</sup>rbmann@uwaterloo.ca

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The critical behavior of the boundary conformal field theory</b>	<b>3</b>
<b>3</b>	<b>Criticality of Thermodynamic curvatures</b>	<b>12</b>
<b>4</b>	<b>Conclusions</b>	<b>15</b>

---

## 1 Introduction

The study of thermodynamic properties of black holes in anti-de Sitter (AdS) space has gained significant importance due to gauge/gravity duality [3]. This duality establishes a relationship between a gravitational theory in the bulk of AdS spacetime and a quantum field theory that lives on the boundary of that AdS spacetime. In the spirit of this correspondence, the Hawking-Page phase transition between the stable large Schwarzschild black hole and thermal gas in the AdS space [4] can be interpreted as a confinement/deconfinement phase transition in the dual strongly coupled gauge theory [5].

Quite some time ago, the existence of small-large black hole phase transitions of charged and rotating AdS black holes, along with their similarity with the liquid-gas phase transition of Van der Waals (VdW) type were pointed out [6, 7]. A complete identification between the thermodynamic parameters of a charged black hole and a Van der Waals system was made some years later in the framework of extended black hole thermodynamics, in which the cosmological constant was treated as thermodynamic pressure and its conjugate quantity as a thermodynamic volume of the black hole [8]. This in turn led to a broad range of theoretical developments in the study of thermodynamics and phase transitions of AdS black holes in the extended phase space [9–15].

In the framework of gauge/gravity duality, it is therefore of interest to ask whether the interpretation of the dynamical cosmological constant as pressure is applicable to the CFT side? Early attempts to address this question [16–18] proposed that varying pressure, or cosmological constant  $\Lambda$ , in the bulk corresponds to varying the number of colors,  $N$ , in the boundary Yang–Mills theory, with thermodynamic volume interpreted in the boundary field theory as an associated chemical potential  $\mu$  for color. For example, chemical potentials corresponding to neutral and charged black holes in  $AdS_5 \times S^5$  have been explicitly obtained by  $N^2$  (instead of  $N$ ) as the number of degrees of freedom of the  $\mathcal{N} = 4$   $SU(N)$  Yang–Mills theory in the large  $N$  limit [19–21]. In this approach it is worth emphasizing that varying  $\Lambda$  in the AdS bulk is equivalent to changing the boundary field theory.

An alternative approach followed here is based on a different interpretation, which proposes that  $N$  should be kept fixed, so that field theory remains the same. In this approach, varying  $\Lambda$  in the bulk is equivalent to varying the curvature radius governing the space on which the field theory resides [1]. In other words, in order to keep  $N$  fixed, we retain the standard holographic relation

$$N^p \sim \frac{L^{d-2}}{G_d} \quad (1.1)$$

between  $N$ , the AdS radius  $L$ , and the  $d$ -dimensional gravitational constant  $G_d$ . To keep the number of colors  $N$  fixed we must vary  $G_d$  and  $L$  whilst keeping the combination  $L^{d-2}/G_d$  fixed. Note that the power  $p$  is theory dependent. For example, for a gauge theory like  $\mathcal{N} = 4$  SuperYang-Mills,  $p = 2$ , whereas for M-theory in  $d = 11$  compactified on  $AdS_4 \times S^7$  with a three-dimensional CFT,  $p = 3/2$  [3]. In other words, this interpretation proposes that varying the cosmological constant (or equivalently the AdS length scale) in the bulk corresponds to varying the spatial volume in the dual field theory, which is completely determined by the radius of the sphere on which the field theory resides [1]. From a thermodynamic point of view, the conjugate variable associated with this volume is pressure, and one is thus led to study the extended thermodynamics of the CFT [22]. From this latter interpretation, a generalized Smarr relation can be calculated by taking extended black hole thermodynamics of the boundary field theory [23]. Furthermore a precise boundary description of extended black hole thermodynamics, in which varying  $\Lambda$  only corresponds to varying central charge ( $C$ ) (instead of varying the volume of the boundary field theory) was recently proposed [24]. In addition, by including the variation of the gravitational constant  $G$  in the first law, a new definition for thermodynamic black hole volume can be defined in the bulk, whilst the CFT remains unchanged (fixed  $C$ ) on the boundary [25].

Another perspective on the thermodynamic properties of a system that has grown in interest in recent year entails geometric methods [26–32]. A metric is constructed on the thermodynamic phase space, and its Riemann scalar curvature can reveal some information about black hole phase transitions and criticality. In this regard, it is important to construct an appropriate thermodynamic geometry that yields a one-to-one correspondence between phase transition points and singularities of the corresponding scalar curvature. However, in some contradictory examples [30, 33], the well-known Ruppeiner geometry, which was defined as a Hessian matrix of entropy, was not successful in achieving such a correspondence.

Nevertheless, Ruppeiner geometry provides us with powerful tools for understanding microstructures of thermodynamic systems [27, 34–36]. To tackle the issue of the Ruppeiner geometry mentioned above, a new formulation of such a geometry was developed, based on considerations about thermodynamic potentials related to mass (instead of the entropy) by Legendre transformations [37–39]. This new formalism of thermodynamic geometry (NTG) equips us with a one-to-one despondence between critical points where heat capacity diverges and curvature singularities. It has been shown that the extrinsic curvature of a certain kind of hypersurface immersed in thermodynamic space constructed by NTG metric not only is singular at the phase transition point, but also has the same sign as the heat capacity around

the critical point [39]. However it fails to explain the thermal stability of a thermodynamic system around a critical point.

In addition to these useful features of NTG, intrinsic (scalar) and extrinsic curvatures exhibit universal behavior around the critical point for charged AdS black holes, which have behaviour qualitatively similar to a Van der Waals fluid [8]. Indeed, near the critical point the (normalized) intrinsic and extrinsic NTG curvature have critical exponents 2 and 1 respectively [2]. Moreover their universal critical amplitudes are

$$R_N t^2 \approx \begin{cases} -\frac{1}{2} & \text{for } t > 0, \\ -\frac{1}{8} & \text{for } t < 0, \end{cases} \quad \text{and} \quad K_N t \approx \begin{cases} -\frac{1}{2} & \text{for } t > 0, \\ -\frac{1}{4} & \text{for } t < 0. \end{cases} \quad (1.2)$$

in which the temperature parameter  $t = T/T_c - 1$ , where  $T_c$  is the critical temperature. For  $t > 0$  we get closer to the critical point along the isochore line ( $v = v_c$  where  $v$  is special volume), whereas for  $t < 0$ , we approach the critical point along the coexistence line between small and large phases in the  $P - v$  diagram [2].

In this paper, we examine this universality of thermodynamic geometry for boundary field theories, which are dual to charged AdS black holes in the bulk, according to the AdS/CFT correspondence. Indeed, such an AdS black hole is embedded in  $D$ -dimensional M-theory/superstring inspired models having  $AdS_d \times \mathbb{S}^{d+k}$  space-time with  $D = 2d + k$ .

Our paper is organized as follows. In Section 2, we investigate the behaviour and criticality of some thermodynamic quantities such as heat capacities and compressibilities of the  $(d - 1)$  dimensional boundary CFT that is dual to a  $d$ - dimensional AdS black hole embedded in  $D$ -dimensional superstring/M-theory inspired models. In Section 3, the NTG geometry is employed in investigating phase transitions and critical behavior of boundary field theories. Remarkably, the intrinsic and extrinsic NTG curvatures reveal universal behaviour around the critical point for boundary CFTs. Our conclusions are presented in Section 4.

## 2 The critical behavior of the boundary conformal field theory

We are interested in studying the behaviour and criticality of some thermodynamic quantities of a  $(d - 1)$  dimensional boundary CFT that is dual to a  $d$ - dimensional AdS black hole embedded in  $D$ -dimensional superstring/M-theory inspired models.

To be more precise, a  $d$ -dimensional AdS black hole can be embedded in  $D$ -dimensional superstring/M-theory inspired models having  $AdS_d \times \mathbb{S}^{d+k}$  space-time, where  $D = 2d + k$ . These black hole solutions can be related to  $N$  coincident  $(d - 2)$ -branes that are assumed to move in such higher dimensional models and are labeled by a triplet  $(D; d; k)$ , where  $k$  is associated with the internal space, i.e., the  $\mathbb{S}^{d+k}$  sphere [6]. For example, the triplet  $(D, d, k) = (11, 7, -3)$  is associated with the compactification of M-theory on the sphere  $\mathbb{S}^4$  with  $M5$ -branes (see [40] for more detail). One also can consider eleven-dimensional M-theory, with  $AdS_4 \times \mathbb{S}^7$  space-time in the presence of  $N$  coincident  $M2$ -branes, which is characterised by the triplet  $(D, d, k) = (11, 4, 3)$  [41]. In addition, the ten-dimensional type IIB superstring

theory with the  $AdS_5 \times S^5$  space-time in the presence of  $D3$ -branes is realized by the triplet  $(D, d, k) = (10, 5, 0)$  [6, 42].

Furthermore, the gravitational constant  $G_d$  in such a  $d$ -dimensional AdS black hole is associated to one corresponding to  $D$ -dimensional superstring/M-theory inspired models [43]. Since the  $d$ -dimensional AdS black hole is obtained from compactification of the  $D$ -dimensional theory on the  $S^{d+k}$  sphere of radius  $L$ , the  $d$ -dimensional Newton constant  $G_d$  is descended from its  $D$ -dimensional counterpart  $G_{D=2d+k}$  via

$$G_d = \frac{G_{2d+k}}{\text{Vol}(S^{d+k})} = \frac{G_{2d+k}}{\Omega_{d+k} L^{d+k}}, \quad (2.1)$$

where  $\Omega_{d+k} = 2\pi^{(d+k+1)/2}/\Gamma(\frac{d+k+1}{2})$  is the volume of a unit  $(d+k)$  sphere. The AdS radius  $L$  is also related to the brane number  $N$  via

$$L^{2(d-1)+k} = 2^{-\left(\frac{d(4-d)+3}{2}\right)} \pi^{7(k+2(d-5))-4} N^{\frac{d-1}{2}} \ell_p^{2(d-1)+k}, \quad (2.2)$$

where the  $D$ -dimensional Planck length  $l_p$  is associated with Newton's constant  $G_{2d+k}$  via  $l_p^{2(d-1)+k} = \hbar G_{2d+k}$ . Clearly, varying the AdS radius  $L$  at fixed  $G$  leads to the variation of the color number  $N$  on the field theory side [16–18]. On the other hand, combining the two preceding relations, we arrive at

$$\frac{1}{4\pi\hbar G_d} = \gamma \frac{N^{\frac{d-1}{2}}}{\mathcal{V}} \quad (2.3)$$

which is consistent with Eq. (1.1), where  $\gamma = 2^{\frac{1}{2}((d-4)d-7)} \pi^{14d+7k-75} \Omega_{d+k} \Omega_{d-2}$  is a constant number and  $\mathcal{V} = \Omega_{d-2} L^{d-2}$  can be interpreted as the spatial volume of the boundary conformal field theory by  $(d-1)$  dimensions. It follows that varying the volume of the CFT, keeping  $N^{d-1/2}$  fixed, is completely equivalent to varying  $G_d$ , or the  $d$ -dimensional Planck length [1]. From this latter perspective we study the thermodynamics of the boundary conformal field theory. Indeed, we assume that  $N^{d-1/2}$  should be kept fixed, so that varying  $\Lambda$  in the bulk has the more natural consequence of changing the volume of the boundary conformal field theory [1].

The static and spherically symmetric  $d$ -dimensional charged AdS black hole solution in Einstein-Maxwell-AdS gravity is given by [6]

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d^2\Omega_{d-2}, \quad (2.4)$$

$$f(r) = 1 - \frac{m}{r^{D-3}} + \frac{q^2}{r^{2(D-3)}} + \frac{r^2}{L^2},$$

where

$$M = \frac{(D-2)\Omega_{d-2}m}{16\pi G_D} \quad Q = \sqrt{\frac{(d-2)(d-3)}{2}} \Omega_{d-2}q \quad (2.5)$$

respectively relate the parameters  $m$  and  $q$  to the black hole mass  $M$  and charge  $Q$  [6]. Imposing the condition  $f(r_h) = 0$  at the event horizon  $r_h$  yields

$$M = \frac{(d-2)\Omega_{d-2}}{16\pi G_d} \left( \frac{r_h^{d-1}}{L^2} + r_h^{d-3} + \frac{q^2}{r_h^{d-3}} \right), \quad (2.6)$$

expressing  $M$  in terms of  $r_h$  and  $q$ . Moreover, the Bekenstein-Hawking entropy is

$$S = \frac{A_h}{4\hbar G_d} = \frac{\Omega_{d-2} r_h^{d-2}}{4 \hbar G_d}. \quad (2.7)$$

In the context of the AdS/CFT dictionary, black hole thermodynamics can be understood in terms of the fundamental degrees of freedom of a thermal quantum field theory and vice versa. Therefore, one can map the black hole mass  $M$  to the internal energy  $U$  of the large  $N$  Yang-Mills theory at the boundary, and the temperature  $T$  and entropy  $S$  of the black hole to those of the boundary field theory [6]. In addition, the  $U(1)$  charge of a black hole in the bulk corresponds to  $R$ -charge in the Yang-Mills theory at the boundary [6, 44, 45]. Therefore, by defining dimensionless variables,  $x \equiv r_h/L$  and  $y \equiv q/L^{d-3}$ , and inserting  $G_d$  from Eq. (2.3) into Eq. (2.6), the internal energy and entropy of the boundary CFT can be written as

$$U = M = \frac{(d-2)\hbar\gamma N^{\frac{d-1}{2}}}{4L} \left( y^2 x^{3-d} + x^{d-3} + x^{d-1} \right), \quad S = \pi\gamma N^{\frac{d-1}{2}} x^{d-2} \quad (2.8)$$

The internal energy of the CFT is a quantum mechanical quantity that vanishes as  $\hbar \rightarrow 0$  by keeping  $x, y$  fixed, whereas the mass on the AdS side is classical. It is also convenient to define the dimensionless charge corresponding to  $R$ -charge in the boundary field theory

$$\tilde{Q} = \frac{QL}{4\pi\hbar G_d} = \sqrt{\frac{(d-2)(d-3)}{2}} \gamma N^{\frac{d-1}{2}} y \quad (2.9)$$

using Eq. (2.3). In accord with the first law of thermodynamics,  $dU = TdS + \Phi d\tilde{Q} - Pd\mathcal{V}$ , Eq. (2.8) implies that other thermodynamic variables such as temperature, chemical potential, and pressure are

$$T = \left( \frac{\partial U}{\partial S} \right)_{\tilde{Q}, \mathcal{V}} = \frac{\{U, \tilde{Q}, \mathcal{V}\}_{x,y,L}}{\{S, \tilde{Q}, \mathcal{V}\}_{x,y,L}} = \frac{\hbar}{4\pi L} \left( (d-1)x + \frac{d-3}{x} - (d-3) \frac{y^2}{x^{2d-5}} \right), \quad (2.10)$$

$$\Phi = \left( \frac{\partial U}{\partial \tilde{Q}} \right)_{S, \mathcal{V}} = \frac{\{U, S, \mathcal{V}\}_{x,y,L}}{\{\tilde{Q}, S, \mathcal{V}\}_{x,y,L}} = \frac{(d-2)\hbar y x^{3-d}}{\sqrt{2(d-3)(d-2)}L}, \quad (2.11)$$

$$P = - \left( \frac{\partial U}{\partial \mathcal{V}} \right)_{S, \tilde{Q}} = - \frac{\{U, S, \tilde{Q}\}_{x,y,L}}{\{\mathcal{V}, S, \tilde{Q}\}_{x,y,L}} = \frac{\hbar\gamma N^{\frac{d-1}{2}} (y^2 x^{3-d} + x^{d-3} + x^{d-1})}{4L^{d-1}\Omega_{d-2}} \quad (2.12)$$

In the above equations, we have used Nambu bracket notation [38]. In the context of gauge/gravity duality,  $T$  can be interpreted as the temperature of the quark-gluon plasma [6, 42]. The thermodynamically-conjugate variable to the volume of the CFT, i.e.  $\mathcal{V}$ , is the pressure, which leads to the  $P-\mathcal{V}$  criticality of a boundary CFT [22]. Moreover,  $\Phi$  corresponds to the chemical potential associated with  $R$  current in the dual supersymmetric Yang-Mills theory.

In order to study the phase structure of the boundary CFT, let us first consider isocharge in a  $T-S$  phase diagram, as shown in the left diagram of Fig. 1. We see that there is a critical charge  $y_c$ , at which there is an inflection point

$$\left( \frac{\partial T}{\partial S} \right)_{\tilde{Q}, \mathcal{V}} = \left( \frac{\partial^2 T}{\partial S^2} \right)_{\tilde{Q}, \mathcal{V}} = 0 \quad (2.13)$$

yielding

$$x_c = \frac{d-3}{\sqrt{(d-1)(d-2)}} \quad y_c = \frac{(d-3)^{d-3}}{(d-2)^{\frac{d-2}{2}}(d-1)^{\frac{d-3}{2}}\sqrt{2d-5}}. \quad (2.14)$$

The critical temperature, pressure, chemical potential, and charge are therefore

$$\begin{aligned} T_c &= \frac{(d-3)\sqrt{(d-1)(d-2)}\hbar}{\pi(2d-5)L}, & P_c &= \frac{\gamma(d-3)^{d-3}((d-5)d+7)\hbar\gamma N^{\frac{d-1}{2}}}{(d-2)^{\frac{d-3}{2}}(d-1)^{\frac{d-1}{2}}(2d-5)\Omega_{d-2}L^{d-1}} \\ \Phi_c &= \frac{\hbar}{\sqrt{2(d-3)(2d-5)}L} & \tilde{Q}_c &= \frac{(d-3)^{\frac{2d-5}{2}}\gamma N^{\frac{d-1}{2}}}{(d-2)^{\frac{d-3}{2}}(d-1)^{\frac{d-3}{2}}\sqrt{4d-10}}. \end{aligned} \quad (2.15)$$

This implies that the relevant (dimensionless) parameter that determines the phase transition is  $T\mathcal{V}^{\frac{1}{d-2}}$  (or  $TL$ ) and  $\mathcal{V}$  and  $T$  are not fixed separately [6, 22]. In the other words, since the boundary field theory is conformal,  $\mathcal{V}$  and  $T$  are not independent. It is therefore convenient to consider charge  $\tilde{Q}$  ( $y$ ) rather than  $T$  as the control parameter in discussing the thermodynamics of the system. This latter view is more in keeping with the notion of the quantum phase transition rather than the thermal phase transition [22].

Furthermore the compressibility factor  $Z$  is


$$Z = \frac{\mathcal{V}_c P_c}{T_c} = \frac{\pi(d-3)^{d-4}((d-5)d+7)\gamma N^{\frac{d-1}{2}}}{(d-2)^{\frac{d-2}{2}}(d-1)^{d/2}} \quad (2.16)$$

which has a constant value for each dimension, as we keep  $N$  fixed. For future convenience, we work with dimensionless parameters such as  $\tilde{T} = TL$  and  $\tilde{\Phi} = \Phi L$  in the rest of the paper.

At the critical point, the phase transition becomes second-order and the heat capacity diverges. The heat capacity at constant charge for the boundary theory is

$$\begin{aligned} C_{\tilde{Q}} &= \tilde{T} \left( \frac{\partial S}{\partial \tilde{T}} \right)_{\tilde{Q}} = \tilde{T} \frac{\{S, \tilde{Q}\}_{x,y}}{\{\tilde{T}, \tilde{Q}\}_{x,y}} \\ &= \frac{\pi(d-2)\gamma N^{\frac{d-1}{2}} x^{d-2} (x^{2d} ((d-1)x^2 + d-3) - (d-3)x^6 y^2)}{(2d-5)(d-3)x^6 y^2 + ((d-1)x^2 - d+3)x^{2d}}. \end{aligned} \quad (2.17)$$

Clearly,  $C_{\tilde{Q}}$  diverges at the critical point in the CFT, whereas heat capacity at fixed charge and volume for the black hole in the bulk is finite at the critical point [8]. The behavior of the heat capacity  $C_{\tilde{Q}}$  with respect to  $S$  is illustrated in the right diagram in Fig. 1. For small charge  $y < y_c$ , we clearly see that there are two divergent points. The first and the third regions at small and large values of the entropy have positive heat capacity whereas the second region in the middle has negative heat capacity. Positive/negative values of the heat capacity imply thermodynamic stability/instability. By increasing  $y$ , the region of negative heat capacity shrinks, and as  $y \rightarrow y_c$ , the negative region disappears. There then exists only one divergent point at which the heat capacity goes to positive infinity. Further increasing  $y$ , we see that the divergent behavior of the heat capacity  $C_{\tilde{Q}}$  completely disappears, shown as the black dashed line.



**Figure 1.** **Left:** Diagram of the temperature  $\hat{T}$  as a function of the entropy  $SN^{-3/2}/\gamma$ . **Right:** Diagram of the heat capacity  $C_{\hat{Q}}/\gamma N^{3/2}(\times 10^{-1})$  versus entropy  $SN^{-3/2}/\gamma$ . Both plots have been plotted for a triplet  $(D; d; k) = (11; 4; 3)$  or 3D CFT while  $\hbar = 1$ .


Using the pressure as defined above, the  $P - \mathcal{V}$  criticality of a boundary CFT was investigated explicitly [22]. However there are a number of differences between the  $P - \mathcal{V}$  behavior of the boundary field theory and that of the black hole in the bulk. First, in the boundary theory there is a single phase at temperatures below the critical temperature and two phases above it. Second, it is pressure rather than volume, that jumps across the two-phase regime<sup>1</sup>. Finally, as one keeps  $\mathcal{V}$  and  $\tilde{Q}$  fixed, the critical exponents are not mean field in spite of the fact that the equation of state of the CFT is similar to the Van der Waals equation of state. However, with pressure as the order parameter in the  $P - \mathcal{V}$  plane, the critical exponents are mean field and the phase transition is similar to that for a Van der Waals black hole like in the bulk [22]. However, if  $\tilde{\Phi}$  instead of  $\tilde{Q}$  is taken as the order parameter in the  $\tilde{\Phi} - \tilde{Q}$  diagram, the critical exponents become mean field.

We have already noted a constraint on the thermodynamic variables that arises because the boundary field theory is conformal, namely that only the combination  $T\mathcal{V}^{\frac{1}{d-2}}$  is relevant to the phase structure, not  $\mathcal{V}$  and  $T$  separately. Consequently, in order to understand phase behaviour in terms of  $P$  and  $\mathcal{V}$ , one needs to work with the dimensionless variables  $P/N^{\frac{d-1}{2}}T^{d-1}$  and  $\mathcal{V}T^{d-2}$ , and use charge rather than the temperature as the control parameter [22]. We shall instead extract mean field critical exponents by considering phase behaviour in the  $\tilde{\Phi} - \tilde{Q}$  plane, since these variables do not scale directly with the temperature parameter.

By eliminating  $x$  and  $y$  in favour of  $\hat{Q} = \frac{Q}{Q_c} = \frac{\tilde{Q}}{\tilde{Q}_c}$  and  $\hat{\Phi} = \frac{\Phi}{\Phi_c} = \frac{\tilde{\Phi}}{\tilde{\Phi}_c}$ , upon inserting Eqs.

<sup>1</sup>This jump may be viewed as a change in the number of degrees of freedom. Thus, any change in the number of effective degrees of freedom can contribute to a change in pressure across the phase transition.





**Figure 2.** **Left:** Behavior of the  $\hat{Q}$  as a function of  $\hat{\Phi}$  for different temperatures  $\hat{T} = 0.999$ ,  $\hat{T} = 1$  and  $\hat{T} = 1.001$ . Maxwell equal area rule can be applied for  $\hat{T} > 1$ . **Right:** Behavior of the  $\hat{Q}$  as a function of  $\hat{\Phi}$  for  $\hat{T} = 1.001$  and Maxwell equal area rule has been shown. Both sides have been obtained from a triplet  $(D; d; k) = (11; 4; 3)$  or 3D CFT with  $\hbar = 1$ .

(2.9) and (2.11) into Eq. (2.10) we obtain

$$\tilde{T} = \frac{(d-3)^{\frac{1}{3-d}} \hbar^{\frac{d-2}{3-d}} \gamma^{\frac{1}{3-d}} N^{\frac{d-2}{d-3}} \tilde{Q}^{\frac{1}{d-3}} \tilde{\Phi}^{\frac{1}{3-d}}}{4\pi(d-2)} \quad (d-2)(d-1)h^{\frac{d-2}{d-3}} N^{\frac{d}{3-d}} + (d-3)^{\frac{d-1}{d-3}} \gamma^{\frac{2}{d-3}} N^{\frac{1}{3-d}} \tilde{Q}^{\frac{2}{3-d}} \tilde{\Phi}^{\frac{2}{d-3}} \left( (d-2)h^2 - 2(d-3)\tilde{\Phi}^2 \right) \quad (2.18)$$


for the equation of state. We depict the resultant phase behaviour in Fig. 2, whose left diagram illustrates isotherm curves in the  $\hat{\Phi} - \hat{Q}$  plane. The behaviour is reminiscent of a reverse Van der Waals transition<sup>2</sup> [46, 47], with an inflection point at  $\hat{T} = 1$  and two distinct phases at larger values of  $\hat{T}$ . As a consequence, the critical point derived in Eq. (2.15) is also obtained from

$$\left( \frac{\partial \hat{Q}}{\partial \hat{\Phi}} \right)_{\hat{T}} = \left( \frac{\partial^2 \hat{Q}}{\partial^2 \hat{\Phi}} \right)_{\hat{T}} = 0. \quad (2.19)$$

From the above similarities, formally identifying the variables  $(\hat{Q}, \hat{\Phi})$  of the boundary field theory with  $(P - V)$  of a van der Waals liquid gas system, the phase structure of the boundary field theory mimics qualitatively certain remarkable properties to that of a van der Waals liquid gas system. Note that it is a reverse VdW transition, with one phase at cold temperatures and two phases at hot temperatures. An order parameter in the boundary field theory that measures the phase change across the critical point can be defined in terms of the Maxwell equal-area law as shown in the right diagram of Fig. 2. Analogous to a Van der Waals system, we therefore define

$$\eta = |\hat{\Phi}_2 - \hat{\Phi}_1| \quad (2.20)$$

<sup>2</sup>It is worthwhile noted that there is the Maxwell construction for  $\hat{T} < 1$  in charged AdS black holes.



**Figure 3.** Diagram of  $\ln(|\hat{\Phi}_1 - \hat{\Phi}_2|)$  versus  $\ln(t)$ . The fitted straight line for the data points (red dot) is given by  $\Delta\hat{\Phi} = b(t^\beta)$  with  $b = 1.58872$  and  $\beta = 0.499974$  for a triplet  $(D; d; k) = (11; 4; 3)$  or 3D CFT while  $\hbar = 1$ .

as the order parameter to characterize the phase change of the boundary CFT near the critical point. For  $\hat{T} > 1$  there is a two-phase regime in the boundary system. To describe this phase transition one can replace the oscillating part of the isotherm in Fig. 2 by an isocharge from the equal area relation

$$\int_{\hat{\Phi}_1}^{\hat{\Phi}_2} \hat{Q} d\hat{\Phi} = \hat{Q}^*(\hat{\Phi}_2) - \hat{Q}^*(\hat{\Phi}_1) \quad (2.21)$$

where  $\hat{Q}$  is regarded as a function of  $(\hat{T}, \hat{\Phi})$  from (2.18), i.e.,

$$\hat{Q} = \frac{\left(4(d-2)\hat{T}\hat{\Phi}^{\frac{5}{d-3}} - \frac{2A}{\sqrt{d-3}}\right)^{d-3}}{(4d-10)^{d-3}\hat{\Phi}^4}, \quad (2.22)$$

$$A = \sqrt{(d-2)\left(4(d-5)d(\hat{T}^2-1) + 24\hat{T}^2 - 25\right)\hat{\Phi}^{\frac{10}{d-3}} + (2d-5)\hat{\Phi}^{\frac{2(d+2)}{d-3}}} \quad (2.23)$$

We define

$$t = \hat{T} - 1, \quad \phi = \hat{\Phi} - 1. \quad (2.24)$$

in order to characterize critical exponents describing the behaviour of physical quantities near the critical point.

The critical exponent  $\beta$ , which characterizes the behavior of the chemical potential difference  $\eta$ , along the  $\hat{Q} - \hat{T}$  coexistence curve, is defined as

$$\eta = |\hat{\Phi}_1 - \hat{\Phi}_2| = bt^\beta \quad \text{for } t > 0 \quad (2.25)$$

whose behaviour we illustrate in Fig. 3, plotting  $\ln(|\hat{\Phi}_1 - \hat{\Phi}_2|)$  as a function of  $\ln(t)$ . The fitted straight line for the data points is generated by solving Eq. (2.21) numerically for  $t > 10^{-4}$ . We find  $b = 1.58872$  and  $\beta = 0.499974$ .

In the reduced parameter space, the isothermal compressibility  $\kappa_{\tilde{T}}$ , and the adiabatic compressibility  $\kappa_S$ , are respectively given by

$$\begin{aligned}\kappa_{\tilde{T}} &= -\frac{1}{\tilde{\Phi}} \left( \frac{\partial \tilde{\Phi}}{\partial \tilde{Q}} \right)_{\tilde{T}} = -\frac{1}{\hat{Q}_c \hat{\Phi}} \left( \frac{\partial \hat{\Phi}}{\partial \hat{Q}} \right)_{\hat{T}} \\ \kappa_S &= -\frac{1}{\tilde{\Phi}} \left( \frac{\partial \tilde{\Phi}}{\partial \tilde{Q}} \right)_S = -\frac{1}{\hat{Q}_c \hat{\Phi}} \left( \frac{\partial \hat{\Phi}}{\partial \hat{Q}} \right)_{\hat{S}}\end{aligned}\quad (2.26)$$

By defining the heat capacity at constant chemical potential,  $C_{\tilde{\Phi}} = \tilde{T} \left( \partial S / \partial \tilde{T} \right)_{\tilde{\Phi}}$ , we find a fascinating relation

$$\kappa_{\tilde{T}} C_{\tilde{\Phi}} \left( \kappa_S C_{\tilde{Q}} \right)^{-1} = 1. \quad (2.27)$$

between heat capacities and compressibilities<sup>3</sup>. Generally, near the critical point heat capacity and compressibility exhibit the critical behaviour

$$C_I \approx \begin{cases} t^{-\alpha_I} & t > 0 \\ (-t)^{-\alpha'_I} & t < 0 \end{cases} \quad \text{and} \quad \kappa_I \approx \begin{cases} t^{-\gamma_I} & t > 0 \\ (-t)^{-\gamma'_I} & t < 0 \end{cases} \quad (2.29)$$

along the  $\hat{Q} - \hat{\Phi}$  coexistence curve and isochemical potential  $\hat{\Phi} = 1$  line, respectively.

Along the isochemical potential curve  $\hat{\Phi} = 1$ , substituting  $\hat{T} = 1 + (-t)$  into the heat capacity and compressibility functions and then expanding them to lowest order in  $(-t)$  yields

$$C_{\tilde{Q}} \approx \frac{\pi(d-3)^{d-2} \gamma N^{\frac{d-1}{2}}}{(d-2)^{\frac{d-4}{2}} (d-1)^{\frac{d-2}{2}} (2d-5)(-t)} \quad (2.30)$$

$$C_{\tilde{\Phi}}^c = -\frac{2\pi(d-3)^{d-2} \gamma N^{\frac{d-1}{2}}}{(d-2)^{\frac{d-6}{2}} (d-1)^{\frac{d-2}{2}}} \quad (2.31)$$

$$\kappa_{\tilde{T}} \approx \frac{(d-2)^{\frac{d-5}{2}} (d-1)^{\frac{d-3}{2}}}{\sqrt{4d-10} (d-3)^{\frac{2d-5}{2}} \gamma N^{\frac{d-1}{2}} (-t)} \quad (2.32)$$


$$\kappa_S^c = -\frac{((d-2)(d-1))^{\frac{d-3}{2}} \sqrt{4d-10}}{(d-3)^{\frac{2d-5}{2}} \gamma N^{\frac{d-1}{2}}} \quad (2.33)$$

which determine  $\alpha'_{\tilde{Q}} = \gamma'_{\tilde{T}} = 1$  and  $\alpha'_{\tilde{\Phi}} = \gamma'_S = 0$  for  $t < 0$ . As indicated above, the adiabatic compressibility  $\kappa_S$  and the heat capacity  $C_{\tilde{\Phi}}$  are negative at the critical point, reflecting the instability of the system. Similar instabilities have been observed for charged AdS black holes [7] and rotating AdS black holes with constant angular momentum [48, 49].

<sup>3</sup>This can be shown using the bracket notation. We have

$$\kappa_{\tilde{T}} C_{\tilde{\Phi}} \left( \kappa_S C_{\tilde{Q}} \right)^{-1} = \left( \frac{\partial \tilde{\Phi}}{\partial \tilde{Q}} \right)_{\tilde{T}} \left( \frac{\partial S}{\partial \tilde{T}} \right)_{\tilde{\Phi}} \left( \frac{\partial \tilde{Q}}{\partial \tilde{\Phi}} \right)_S \left( \frac{\partial \tilde{T}}{\partial S} \right)_{\tilde{Q}} = \frac{\{\tilde{\Phi}, \tilde{T}\} \{S, \tilde{\Phi}\} \{\tilde{Q}, S\} \{\tilde{T}, \tilde{Q}\}}{\{\tilde{Q}, \tilde{T}\} \{\tilde{T}, \tilde{\Phi}\} \{\tilde{\Phi}, S\} \{S, \tilde{Q}\}} = 1 \quad (2.28)$$

Note that all brackets are written in the coordinates  $(\tilde{T}, \tilde{\Phi})$ .



**Figure 4.** Diagram of  $\ln(\kappa_{\tilde{T}}\gamma N^{3/2})$  versus  $\ln(t)$  for a triplet  $(D; d; k) = (11; 4; 3)$  or 3D CFT while  $\hbar = 1$ . The fitted straight line for the data points (red dot) is given by  $\kappa_{\tilde{T}}\gamma N^{3/2} = r(t^{-\gamma_{\tilde{T}}})$ . **Left**, for region (I),  $r = -1.4565$  and  $\gamma_{\tilde{T}} = 1.00485$ . **Right**, for region (II)  $r = -1.31674$  and  $\gamma_{\tilde{T}} = 0.995182$ .

We illustrate in Fig. 4 numerical plots of isothermal compressibility  $\kappa_{\tilde{T}}$  versus  $t$  for regions (I) (left) and (II) (right) in Fig. 2 along the  $\hat{Q} - \hat{\Phi}$  coexistence curve for the triplet  $(11; 4; 3)$ . We obtain the predicted asymptotic critical behavior  $\kappa_{\tilde{T}} \approx (t)^{\gamma_{\tilde{T}}}$  with exponent  $\gamma_{\tilde{T}} = 1$ . The coexistence saturated regions (I) and (II) may respectively be interpreted as a gas of particles in quark-gluon plasma phase and hadrons in the confinement phase. Similar results are calculated for other triplets, which we collect in table 1. Note that if pressure is taken as the order parameter (whose value jumps across the line of first order phase transitions in the  $P - \mathcal{V}$  plane) and  $\tilde{Q}$  as the control variable the exponents are always mean field in the large  $N$  limit [22].

Critical exponent	$\beta$	$\alpha_{\tilde{Q}}$	$\gamma_{\tilde{T}}$
(11;4;3); 3D CFT	0.49997	0.999892	1.00002
(10;5;0); 4D CFT	0.49996	0.999919	1.00012
(11;7;-3); 6D CFT	0.49999	1.00003	1.00055

**Table 1.** Critical exponent values of  $\eta$ ,  $C_{\tilde{Q}}/\gamma N^{\frac{d-1}{2}}$  and  $\kappa_{\tilde{T}}\gamma N^{\frac{d-1}{2}}$  for  $t > 0$ . These values have been averaged between the values obtained from region (I) and (II).

### 3 Criticality of Thermodynamic curvatures

Now allow us to apply new formalism of the thermodynamic geometry (NTG) introduced in Ref. [38] for the boundary CFT, which is defined by

$$dl_{NTG}^2 = \frac{1}{T} \left( \eta_i^j \frac{\partial^2 \Xi}{\partial X^j \partial X^i} dX^i dX^j \right) \quad (3.1)$$

where  $\eta_i^j = \text{diag}(-1, 1, \dots, 1)$ ,  $\Xi$  is the thermodynamic potential, and the  $X^i$  can be intensive or extensive variables [38]. Interestingly, this formalism is able to explain a one-to-one correspondence between phase transitions and curvature singularities. For example, by choosing  $\Xi = \tilde{G}(\tilde{T}, \tilde{\Phi})$  where  $\tilde{G} = \tilde{U} - \tilde{T}S - \tilde{\Phi}\tilde{Q}$  is the Gibbs energy and  $X^i = (\tilde{T}, \tilde{\Phi})$  in NTG metric Eq. (3.1), we have

$$g_{\tilde{G}}^{NTG} = \frac{1}{\tilde{T}} \text{diag} \left( -\frac{\partial^2 \tilde{G}}{\partial \tilde{T}^2}, \frac{\partial^2 \tilde{G}}{\partial \tilde{\Phi}^2} \right) = C_{\tilde{\Phi}} \text{diag} \left( \tilde{T}^{-2}, -\mathcal{F} \right) = C_{\tilde{\Phi}} \hat{g}, \quad \text{with } \mathcal{F} = \frac{\left( \frac{\partial \tilde{Q}}{\partial \tilde{\Phi}} \right) \tilde{T}}{\tilde{T} C_{\tilde{\Phi}}} \quad (3.2)$$

in which we have used the first law  $d\tilde{G} = -Sd\tilde{T} - \tilde{Q}d\tilde{\Phi}$  for the Gibbs free energy. According to the new metric  $\hat{g}$ , which is conformally equivalent with the previous one, the Ricci scalar can be written as [38, 50]

$$R = C_{\tilde{\Phi}}^{-1} \left[ \hat{R} - \square \ln C_{\tilde{\Phi}} \right] \quad (3.3)$$

where  $\square = \partial^\mu \partial_\mu$  is d'Alembert operator associated with (3.2) and

$$\hat{R} = \tilde{T} \frac{\partial \ln(\mathcal{F})}{\partial \tilde{T}} \left[ \frac{1}{2} \tilde{T} \frac{\partial \ln(\mathcal{F})}{\partial \tilde{T}} - 1 - \tilde{T} \frac{\partial}{\partial \tilde{T}} \left( \ln \left( \frac{\partial \mathcal{F}}{\partial \tilde{T}} \right) \right) \right] \quad (3.4)$$


As illustrated in Fig. 5, the singularities of the Ricci scalar  $R^{NTG}$  occur exactly at phase transitions of the heat capacity  $C_{\tilde{Q}}$  at constant charge with no other additional roots<sup>4</sup>. As a consequence of the NTG method, there is a one-to-one correspondence between curvature singularities and phase transitions of the heat capacity.

Although NTG curvature indicates the location of phase transition points, it is not able to explain thermodynamic stability of a system. As shown in Fig. 5, the sign of the curvature does not always correspond to the sign of the heat capacity. However it has been shown [39] that the extrinsic curvature of a certain kind of hypersurface immersed in thermodynamic phase space contains unexpected information about stability of a system. More precisely, its extrinsic curvature not only diverges at phase transition points, but also everywhere has the same sign as the heat capacity.

To find this hypersurface, we consider  $\tilde{T}$  constant hypersurfaces in the thermodynamic manifold constructed from the metric elements in (3.2). For such a hypersurface, the unit normal vector is defined as  $n_i = (1, 0)$  and  $n^i = g^{ij} n_j = (\sqrt{|C_{\tilde{\Phi}}|}/\tilde{T}, 0)$ . Therefore, the extrinsic curvature can be defined by  $K = \nabla_i n^i$  which yields

$$K = \frac{\tilde{T}}{2\sqrt{|C_{\tilde{\Phi}}|}} \frac{\partial}{\partial \tilde{T}} \left( \ln \left( |C_{\tilde{\Phi}}| \mathcal{F} \right) \right) \quad (3.5)$$

<sup>4</sup>The conjugate potential  $\tilde{\Xi} = \tilde{U}$ , which obeys  $\Xi + \tilde{\Xi} = 2\tilde{U} - \tilde{T}S - \tilde{\Phi}\tilde{Q}$ , yields the same result [51].



**Figure 5.** Behavior of Ricci scalar, heat capacity  $C_{\tilde{Q}}/\gamma N^{\frac{3}{2}}$  and extrinsic curvature with respect to  $\tilde{\Phi}$ , **Top(Left)**: for  $t < 0$ ,  $R\gamma N^{\frac{3}{2}} (\times 10^{-3})$ ,  $C_{\tilde{Q}}/\gamma N^{\frac{3}{2}} (\times 10^{-1})$  and  $K\gamma^{\frac{1}{2}} N^{\frac{3}{4}} (\times 10^{-1})$  have been plotted. **Top(Right)**: for  $t = 0$ ,  $R(\times 10^{-7})\gamma N^{\frac{3}{2}}$ ,  $C_{\tilde{Q}}/\gamma N^{\frac{3}{2}} (\times 10^{-5})$  and  $K\gamma^{\frac{1}{2}} N^{\frac{3}{4}} (\times 10^{-4})$  have been plotted. **Bottom**: for  $t > 0$ ,  $R\gamma N^{\frac{3}{2}} (\times 10^{-7})$ ,  $C_{\tilde{Q}}/\gamma N^{\frac{3}{2}} (\times 10^{-3})$  and  $K\gamma^{\frac{1}{2}} N^{\frac{3}{4}} (\times 10^{-3})$  have been depicted.

Fig. 5 shows that the extrinsic curvature of the constant  $\tilde{T}$  hypersurface has everywhere the same sign as the heat capacity, in contrast to the scalar curvature.

Let us now examine the intrinsic and extrinsic curvatures obtained in Eqs. (3.3) and (3.8) respectively as the critical point is approached along the isochemical potential line  $\hat{\Phi} = 1$  (or  $\tilde{\Phi} = \tilde{\Phi}_c$ ). Near the critical point we expand  $\hat{Q} = \sum_{i,j} a_{ij} t^i \phi^j$ , where  $a_{00} = 1$ . Since the respective critical behaviour of  $\hat{\Phi}$  along the coexistence and isochemical potential lines is  $\hat{\Phi} = 1$  and  $\hat{\Phi} \approx t^\beta$  (with  $\beta > 1$ ), it is sufficient to consider the expansion of  $\hat{Q}$  to linear order in  $t$ . Therefore, it is easy to see that the last derivative term in Eq. (3.4) can be completely ignored in comparison with other terms near the critical point. Taking advantage of Eqs. (2.26) and (2.29) we obtain

$$\mathcal{F} = \frac{\left(\frac{\partial \hat{Q}}{\partial \hat{\Phi}}\right)_{\tilde{T}}}{\tilde{T} C_{\tilde{\Phi}}} = -(\tilde{T} \tilde{\Phi} C_{\tilde{\Phi}} \kappa_{\tilde{T}})^{-1} \approx -\frac{(1 + (-t))(-t)^{\gamma'_{\tilde{T}}}}{C_{\tilde{\Phi}}^c} \approx -\frac{(-t)^{\gamma'_{\tilde{T}}}}{C_{\tilde{\Phi}}^c} \quad \text{for } t < 0 \quad (3.6)$$

Because  $C_{\tilde{\Phi}}$  is finite at the critical point, the last term in Eq. (3.3) vanishes, and so

$$R_N \equiv RC_{\tilde{\Phi}}^c \approx \hat{R} \approx (1+t') \frac{\partial \ln(\mathcal{F})}{\partial t'} \left[ \frac{1}{2} (1+t') \frac{\partial \ln(\mathcal{F})}{\partial t'} - 1 \right] \approx \frac{\gamma'_{\tilde{T}}{}^2}{2t'^2} \quad (3.7)$$

We also find

$$K_N = K \sqrt{|C_{\tilde{\Phi}}^c|} = \frac{1+t'}{2} \frac{\partial}{\partial t'} \left( \ln(\mathcal{F} C_{\tilde{\Phi}}^c) \right) \approx \frac{\gamma'_{\tilde{T}}}{2t'} \quad (3.8)$$

near the critical point, where we have replaced  $T$  by  $t' + 1$ , with  $t' = -t$ , to lowest order in  $t'$ . In summary, we find

$$\begin{aligned} R_N \equiv RC_{\tilde{\Phi}}^c &\approx \frac{\gamma'_{\tilde{T}}{}^2}{2t'^2} & \text{for } t < 0 \\ K_N \equiv K \sqrt{|C_{\tilde{\Phi}}^c|} &\approx -\frac{\gamma'_{\tilde{T}}}{2t} & \text{for } t < 0 \end{aligned} \quad (3.9)$$

As for VdW AdS black holes [2, 35] we can define normalized intrinsic and extrinsic curvatures. Since the critical exponent  $\gamma'_{\tilde{T}} = 1$  in all cases, we see that  $R_N t^2 = 1/2$  and  $K_N t = -1/2$  for  $t < 0$ , commensurate with the result [2] for d-dimensional charged AdS black holes in the bulk for  $t > 0$ . However, there is a sign difference between our result (3.9) for  $R_N$  and that for charged AdS black holes [2], which comes from  $C_{\tilde{\Phi}} < 0$  at the critical point for the boundary system<sup>5</sup>. This finding demonstrates that there is universal behaviour of thermodynamics curvature near the critical point, namely, they do not depend on the details of the boundary system.

To check this universality property, we approach the critical point along the two-phase regime coexistence line. Using Eqs. (2.29) and (2.25), we find, one obtains

$$\mathcal{F} = -(\tilde{T} \tilde{\Phi} C_{\tilde{\Phi}}^c \kappa_{\tilde{T}})^{-1} \approx -\frac{(1+t)t^{\gamma_{\tilde{T}}-\beta}}{C_{\tilde{\Phi}}^c} \approx -\frac{t^{\gamma_{\tilde{T}}-\beta}}{C_{\tilde{\Phi}}^c} \quad \text{for } t > 0 \quad (3.10)$$

to lowest order in  $t$ . From this we straightforwardly find

$$\begin{aligned} R_N \equiv RC_{\tilde{\Phi}}^c &\approx (1+t) \frac{\partial \ln(\mathcal{F})}{\partial t} \left[ \frac{1}{2} (1+t) \frac{\partial \ln(\mathcal{F})}{\partial t} - 1 \right] \approx \frac{(\beta - \gamma_{\tilde{T}})^2}{2t^2} \\ K_N = K \sqrt{|C_{\tilde{\Phi}}^c|} &= \frac{1+t}{2} \frac{\partial}{\partial t} \left( \ln(\mathcal{F} C_{\tilde{\Phi}}^c) \right) \approx \frac{\gamma_{\tilde{T}} - \beta}{2t} \end{aligned} \quad (3.11)$$

According to the data reported in table 1, the values of the critical exponents  $\beta$  and  $\gamma_{\tilde{T}}$  are essentially the same for all CFT dimensions; we can take  $\gamma_{\tilde{T}} \approx 1$  and  $\beta \approx 1/2$ . From Eqs. (3.9) and (3.11), as  $t \rightarrow 0^\pm$  we get

$$R_N t^2 \approx \begin{cases} \frac{1}{2} & \text{for } t < 0, \\ \frac{1}{8} & \text{for } t > 0, \end{cases} \quad \text{and} \quad K_N t \approx \begin{cases} -\frac{1}{2} & \text{for } t < 0, \\ \frac{1}{4} & \text{for } t > 0. \end{cases} \quad (3.12)$$

<sup>5</sup>The same issue has been reported for charge AdS black holes in a cavity [52].

We emphasize that we approach the critical point along different paths. For  $t < 0$  we approach it along the isochemical potential line ( $\hat{\Phi} = 1$ ), whereas for  $t > 0$  we approach it along the coexistence lines between regions I and II. Due to our choice of various trajectories in the  $\hat{Q} - \hat{\Phi}$  diagram, the discontinuity in the value of thermodynamic curvatures in Eq. (3.12) is unavoidable. Numerically checking this critical behaviour by plotting  $R_N$  and  $K_N$  versus  $t$  (in a log-log plot), we summarize the results for various cases in table 2.

Quantity	Coefficient	(11;4;3)	(10;5;0)	(11;7;-3)
		3D CFT	4D CFT	6D CFT
$\ln  R_N $ (Region (I))	$c_R$	2.00388	2.00567	2.0096
	$d_R$	3.56833	4.54961	5.70467
$\ln  R_N $ (Region (II))	$c_R$	1.99583	1.99386	1.99015
	$d_R$	3.45195	4.37827	5.42222
$\ln  K_N $ (Region (I))	$c_K$	0.993519	0.990341	1.01594
	$d_K$	2.00937	2.44133	3.41461
$\ln  K_N $ (Region (II))	$c_K$	1.00561	1.00776	1.01143
	$d_K$	2.18417	2.2.69318	3.29687

**Table 2.** The coefficients of the numerical fitting of  $\ln |B_N| = -c_B \ln(t) - d_B$  ( $B = R, K$ ), for coexistence saturated Region (I) and coexistence saturated Region (II).

Taking numerical error into account, the slopes of the lines of the curves  $\ln |B_N| = -c_B \ln(t) - d_B$  ( $B = R, K$ ) in table 2 imply that the critical exponents for  $R_N$  and  $K_N$  are roughly  $c_R = 2$  and  $c_K = 1$ , respectively, consistent with those of charged AdS black holes in the bulk. From the values of  $d_R$  and  $d_K$  we collect the amplitudes related to these critical exponents in table 3. We conclude that the universal behavior of both thermodynamic curvatures is independent of the number of CFT dimensions. Indeed, they confirm universal amplitudes presented in Eq. (3.12). On the gravity side, we saw previously that there is the same criticality behavior, i.e. Eq. (1.2) for higher dimensional charged AdS black holes in such a way that it is independent of the spacetime dimension  $d$  [2]. Therefore, we can conclude that such universality holds for both bulk and boundary theories.

## 4 Conclusions

The purpose of the current study was to apply the NTG geometry to determining the critical behavior and phase structure of the thermal boundary conformal field theories near the critical



Quantity	(11;4;3)	(10;5;0)	(11;7;-3)
	3D CFT	4D CFT	6D CFT
$R_N t^2$	0.125214	0.125336	0.125153
$\frac{(\beta - \gamma_{\hat{T}})^2}{2}$	0.125025	0.12508	0.12528
$K_N t$	0.251437	0.253177	0.25903
$\frac{(\gamma_{\hat{T}} - \beta)}{2}$	0.250025	0.25008	0.25028

**Table 3.** Amplitudes of the critical behavior of normalized thermodynamic curvatures near the critical point.

point. According to the AdS/CFT correspondence, such a  $(d-1)$  dimensional boundary CFT is dual to a  $d$ - dimensional AdS black hole embedded in  $D$ -dimensional superstring/M-theory inspired models.

In the framework of extended black hole thermodynamics [8, 53, 54], the cosmological constant is associated with the pressure of the gravitational system,  $P = -\Lambda/8\pi G$ , whereas its conjugate quantity is interpreted as the thermodynamic volume  $V$ . The effect of including these two variables in thermodynamic phase space has led to the discovery of a broad range of new phenomena associated with black holes [46, 55–57]. Nonetheless, it is natural to ask what the interpretation is of the bulk pressure/volume on the dual boundary conformal field theory (CFT) once the cosmological constant is treated as a thermodynamic variable. To this end, there are two interpretations. As argued in [16–18], varying pressure, or  $\Lambda$ , is equivalent to varying the number of colors,  $N$ , in the boundary field theory so that the thermodynamic conjugate of pressure, i.e. the thermodynamic volume, can then be interpreted in the boundary field theory as an associated chemical potential,  $\mu$  for color. Alternatively, the number of colors  $N$  can be kept fixed, so that we are always referring to the same field theory, in which case varying  $\Lambda$  in the bulk has the more natural consequence of varying the volume of the space on which the field theory resides [1].

From this latter interpretation, we obtained critical exponents for heat capacities and compressibilities in the  $\hat{Q} - \hat{\Phi}$  plane with  $\hat{\Phi}$  as the order parameter. The exponents are mean field, and the phase transition is similar to that of a Van der Waals gas, albeit one undergoing a reverse transition. Note that mean field exponents also characterize the phase transition in the  $P - \mathcal{V}$  plane when  $P$  would be the order parameter.

In addition, we studied the critical behavior of the normalized intrinsic and extrinsic thermodynamic curvatures near the critical point. More precisely, the criticality of such a curvature has been analytically and numerically checked form two trajectories, i.e., along the isochemical potential line and the coexistence curve in the  $\hat{Q} - \hat{\Phi}$  diagram on approaching to

the critical point. In analogy with charged AdS black hole cases in the bulk [2], our finding reveals that the critical exponent of the intrinsic and extrinsic curvature is 2 and 1, respectively. Furthermore, the analytical result shows a relation between the amplitude of thermodynamic curvatures and mean field critical exponents as shown in Eqs. (3.9) and (3.11). Interestingly, critical amplitudes do not depend on the number of thermal CFT dimensions. Furthermore, we calculated numerically these amplitudes in Eq. (3.12), which are in consistent with analytical results. Manifestly, there is always a universal feature of thermodynamic curvatures for both the bulk and boundary theories in the context of gauge/gravity duality.

## Acknowledgements

We are grateful to Yu-Xiao Liu and Brian P. Dolan for reading a preliminary version of the draft. This work was supported in part by the Natural Sciences and Engineering Research Council of Canada. This work was also supported by the National Natural Science Foundation of China (Grant No. 12075103).

## References

- [1] A. Karch and B. Robinson, *Holographic Black Hole Chemistry*, *JHEP* **12** (2015) 073, [1510.02472].
- [2] S. A. Hosseini Mansoori, M. Rafiee and S.-W. Wei, *Universal criticality of thermodynamic curvatures for charged AdS black holes*, *Phys. Rev. D* **102** (2020) 124066, [2007.03255].
- [3] J. M. Maldacena, *The Large  $N$  limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231–252, [hep-th/9711200].
- [4] S. W. Hawking and D. N. Page, *Thermodynamics of Black Holes in anti-De Sitter Space*, *Commun. Math. Phys.* **87** (1983) 577.
- [5] E. Witten, *Anti-de Sitter space, thermal phase transition, and confinement in gauge theories*, *Adv. Theor. Math. Phys.* **2** (1998) 505–532, [hep-th/9803131].
- [6] A. Chamblin, R. Emparan, C. V. Johnson and R. C. Myers, *Charged AdS black holes and catastrophic holography*, *Phys. Rev. D* **60** (1999) 064018, [hep-th/9902170].
- [7] A. Chamblin, R. Emparan, C. V. Johnson and R. C. Myers, *Holography, thermodynamics and fluctuations of charged AdS black holes*, *Phys. Rev. D* **60** (1999) 104026, [hep-th/9904197].
- [8] D. Kubiznak and R. B. Mann,  *$P$ - $V$  criticality of charged AdS black holes*, *JHEP* **07** (2012) 033, [1205.0559].
- [9] N. Altamirano, D. Kubiznak and R. B. Mann, *Reentrant phase transitions in rotating anti-de Sitter black holes*, *Phys. Rev. D* **88** (2013) 101502, [1306.5756].
- [10] N. Altamirano, D. Kubizňák, R. B. Mann and Z. Sherkatghanad, *Kerr-AdS analogue of triple point and solid/liquid/gas phase transition*, *Class. Quant. Grav.* **31** (2014) 042001, [1308.2672].

- [11] N. Altamirano, D. Kubiznak, R. B. Mann and Z. Sherkatghanad, *Thermodynamics of rotating black holes and black rings: phase transitions and thermodynamic volume*, *Galaxies* **2** (2014) 89–159, [1401.2586].
- [12] S.-W. Wei and Y.-X. Liu, *Critical phenomena and thermodynamic geometry of charged Gauss-Bonnet AdS black holes*, *Phys. Rev. D* **87** (2013) 044014, [1209.1707].
- [13] R.-G. Cai, L.-M. Cao, L. Li and R.-Q. Yang, *P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space*, *JHEP* **09** (2013) 005, [1306.6233].
- [14] Z. Sherkatghanad, B. Mirza, Z. Mirzaiyan and S. A. Hosseini Mansoori, *Critical behaviors and phase transitions of black holes in higher order gravities and extended phase spaces*, *Int. J. Mod. Phys. D* **26** (2016) 1750017, [1412.5028].
- [15] B. P. Dolan, A. Kostouki, D. Kubiznak and R. B. Mann, *Isolated critical point from Lovelock gravity*, *Class. Quant. Grav.* **31** (2014) 242001, [1407.4783].
- [16] C. V. Johnson, *Holographic Heat Engines*, *Class. Quant. Grav.* **31** (2014) 205002, [1404.5982].
- [17] D. Kastor, S. Ray and J. Traschen, *Chemical Potential in the First Law for Holographic Entanglement Entropy*, *JHEP* **11** (2014) 120, [1409.3521].
- [18] B. P. Dolan, *Bose condensation and branes*, *JHEP* **10** (2014) 179, [1406.7267].
- [19] J.-L. Zhang, R.-G. Cai and H. Yu, *Phase transition and thermodynamical geometry for Schwarzschild AdS black hole in  $AdS_5 \times S^5$  spacetime*, *JHEP* **02** (2015) 143, [1409.5305].
- [20] J.-L. Zhang, R.-G. Cai and H. Yu, *Phase transition and thermodynamical geometry of Reissner-Nordström-AdS black holes in extended phase space*, *Phys. Rev. D* **91** (2015) 044028, [1502.01428].
- [21] R. Maity, P. Roy and T. Sarkar, *Black Hole Phase Transitions and the Chemical Potential*, *Phys. Lett. B* **765** (2017) 386–394, [1512.05541].
- [22] B. P. Dolan, *Pressure and compressibility of conformal field theories from the AdS/CFT correspondence*, *Entropy* **18** (2016) 169, [1603.06279].
- [23] M. Sinamuli and R. B. Mann, *Higher Order Corrections to Holographic Black Hole Chemistry*, *Phys. Rev. D* **96** (2017) 086008, [1706.04259].
- [24] M. R. Visser, *Holographic Thermodynamics Requires a Chemical Potential for Color*, 2101.04145.
- [25] W. Cong, D. Kubiznak and R. B. Mann, *Thermodynamics of AdS Black Holes: Central Charge Criticality*, 2105.02223.
- [26] F. Weinhold, *Metric geometry of equilibrium thermodynamics*, *The Journal of Chemical Physics* **63** (1975) 2479–2483.
- [27] G. Ruppeiner, *Thermodynamics: A riemannian geometric model*, *Physical Review A* **20** (1979) 1608.
- [28] O. Hiroshi, O. Tsunehiro and H. Hiroaki, *Riemann scalar curvature of ideal quantum gases obeying gentile’s statistics*, .
- [29] G. Ruppeiner, *Riemannian geometry in thermodynamic fluctuation theory*, *Reviews of Modern Physics* **67** (1995) 605.

- [30] J. E. Āman, I. Bengtsson and N. Pidokrajt, *Geometry of black hole thermodynamics*, *General Relativity and Gravitation* **35** (2003) 1733–1743.
- [31] H. Quevedo, *Geometrothermodynamics*, *Journal of Mathematical Physics* **48** (2007) 013506.
- [32] B. Mirza and M. Zamaninasab, *Ruppeiner geometry of  $rn$  black holes: flat or curved?*, *Journal of High Energy Physics* **2007** (2007) 059.
- [33] T. Sarkar, G. Sengupta and B. Nath Tiwari, *On the thermodynamic geometry of BTZ black holes*, *JHEP* **11** (2006) 015, [[hep-th/0606084](#)].
- [34] S.-W. Wei and Y.-X. Liu, *Insight into the Microscopic Structure of an AdS Black Hole from a Thermodynamical Phase Transition*, *Phys. Rev. Lett.* **115** (2015) 111302, [[1502.00386](#)].
- [35] S.-W. Wei, Y.-X. Liu and R. B. Mann, *Repulsive Interactions and Universal Properties of Charged Anti-de Sitter Black Hole Microstructures*, *Phys. Rev. Lett.* **123** (2019) 071103, [[1906.10840](#)].
- [36] S.-W. Wei, Y.-X. Liu and R. B. Mann, *Ruppeiner Geometry, Phase Transitions, and the Microstructure of Charged AdS Black Holes*, *Phys. Rev. D* **100** (2019) 124033, [[1909.03887](#)].
- [37] S. A. H. Mansoori and B. Mirza, *Correspondence of phase transition points and singularities of thermodynamic geometry of black holes*, *Eur. Phys. J. C* **74** (2014) 2681, [[1308.1543](#)].
- [38] S. A. Hosseini Mansoori and B. Mirza, *Geometrothermodynamics as a singular conformal thermodynamic geometry*, *Phys. Lett. B* **799** (2019) 135040, [[1905.01733](#)].
- [39] S. A. H. Mansoori, B. Mirza and E. Sharifian, *Extrinsic and intrinsic curvatures in thermodynamic geometry*, *Phys. Lett. B* **759** (2016) 298–305, [[1602.03066](#)].
- [40] A. Belhaj, A. El Balali, W. El Hadri, Y. Hassouni and E. Torrente-Lujan, *Phase transition and shadow behaviors of quintessential black holes in M-theory/superstring inspired models*, *Int. J. Mod. Phys. A* **36** (2021) 2150057, [[2004.10647](#)].
- [41] A. Dabholkar, N. Drukker and J. Gomes, *Localization in supergravity and quantum  $AdS_4/CFT_3$  holography*, *JHEP* **10** (2014) 090, [[1406.0505](#)].
- [42] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253–291, [[hep-th/9802150](#)].
- [43] S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, *Coupling constant dependence in the thermodynamics of  $N=4$  supersymmetric Yang-Mills theory*, *Nucl. Phys. B* **534** (1998) 202–222, [[hep-th/9805156](#)].
- [44] K. Behrndt, M. Cvetič and W. A. Sabra, *Nonextreme black holes of five-dimensional  $N=2$  AdS supergravity*, *Nucl. Phys. B* **553** (1999) 317–332, [[hep-th/9810227](#)].
- [45] K. Behrndt, A. H. Chamseddine and W. A. Sabra, *BPS black holes in  $N=2$  five-dimensional AdS supergravity*, *Phys. Lett. B* **442** (1998) 97–101, [[hep-th/9807187](#)].
- [46] D. Kubizňák, R. B. Mann and M. Teo, *Black hole chemistry: thermodynamics with lambda*, *Classical and Quantum Gravity* **34** (2017) 063001.
- [47] A. M. Frassino, D. Kubiznak, R. B. Mann and F. Simovic, *Multiple Reentrant Phase Transitions and Triple Points in Lovelock Thermodynamics*, *JHEP* **09** (2014) 080, [[1406.7015](#)].

- [48] M. M. Caldarelli, G. Cognola and D. Klemm, *Thermodynamics of Kerr-Newman-AdS black holes and conformal field theories*, *Class. Quant. Grav.* **17** (2000) 399–420, [[hep-th/9908022](#)].
- [49] B. P. Dolan, *Thermodynamic stability of asymptotically anti-de Sitter rotating black holes in higher dimensions*, *Class. Quant. Grav.* **31** (2014) 165011, [[1403.1507](#)].
- [50] S. M. Carroll, *Spacetime and geometry*. Cambridge University Press, 2019.
- [51] S. A. Hosseini Mansoori, *Thermodynamic geometry of the novel 4-D Gauss–Bonnet AdS black hole*, *Phys. Dark Univ.* **31** (2021) 100776, [[2003.13382](#)].
- [52] P. Wang, H. Wu and H. Yang, *Thermodynamic Geometry of AdS Black Holes and Black Holes in a Cavity*, *Eur. Phys. J. C* **80** (2020) 216, [[1910.07874](#)].
- [53] D. Kastor, S. Ray and J. Traschen, *Enthalpy and the Mechanics of AdS Black Holes*, *Class. Quant. Grav.* **26** (2009) 195011, [[0904.2765](#)].
- [54] B. P. Dolan, *Pressure and volume in the first law of black hole thermodynamics*, *Class. Quant. Grav.* **28** (2011) 235017, [[1106.6260](#)].
- [55] S. Gunasekaran, D. Kubizňák and R. B. Mann, *Extended phase space thermodynamics for charged and rotating black holes and born-infeld vacuum polarization*, *Journal of High Energy Physics* **2012** (2012) 110.
- [56] N. Altamirano, D. Kubizňák and R. B. Mann, *Reentrant phase transitions in rotating anti-de sitter black holes*, *Physical Review D* **88** (2013) 101502.
- [57] R. A. Hennigar, R. B. Mann and E. Tjoa, *Superfluid black holes*, *Physical Review Letters* **118** (2017) 021301.