

Amplitudes on abelian categories

Barbara Giunti^{*}, John S. Nolan[†], Nina Otter[‡], Lukas Waas[§]

Abstract

The use of persistent homology in applications is justified by the validity of certain stability results. At the core of such results is a notion of distance between the invariants that one associates to data sets. While such distances are well-understood in the one-parameter case, the situation for multiparameter persistence modules is more challenging, since there exists no generalisation of the barcode.

Here we introduce a general framework to study stability questions in multiparameter persistence. We introduce amplitudes — invariants that arise from assigning a non-negative real number to each persistence module, and which are monotone and subadditive in an appropriate sense — and then study different ways to associate distances to such invariants. Our framework is very comprehensive, as many different invariants that have been introduced in the Topological Data Analysis literature are examples of amplitudes, and furthermore many known distances for multiparameter persistence can be shown to be distances from amplitudes. Finally, we show how our framework can be used to prove new stability results.

Contents

1	Introduction	2
1.1	Related work	5
2	Persistence modules and finiteness conditions	5
2.1	Finiteness conditions	6
3	Amplitudes	10
3.1	Amplitudes in TDA	12
3.2	L^p -amplitudes	16
3.3	Additive amplitudes	20
3.4	The c_p -rank in the amplitude framework	22
4	Distances from amplitudes	25
4.1	The interleaving distance as a distance from an amplitude	28
4.2	Wasserstein distance of p -norm amplitudes	29
5	Continuity and stability	30
5.1	Amplitude-bounding functors	30
5.2	Continuity of amplitude-bounding functors	31
5.3	Stability results for amplitudes	32

^{*}Institute of Geometry, Graz Institute of Technology, Graz, Austria, *email*: bgiunti@tugraz.at

[†]Department of Mathematics, UC Berkeley, Berkeley, CA, US, *email*: john_nolan@berkeley.edu

[‡]Department of Mathematics, UCLA, Los Angeles, CA, US, *email*: n.otter@qmul.ac.uk

[§]Department of Mathematics, Heidelberg University, Germany, *email*: lwaas@mathi.uni-heidelberg.de

5.3.1	Stability under change of amplitudes	33
5.3.2	Interleaving and L^1 -path metric	35
5.3.3	Pulling back along monotone functions	35
5.4	Stability of quotient restrictions	36
5.5	Stability of the 0-th local cohomology	38
5.6	Stability of c_r -rank	39
6	Conclusions	40
A	Noise systems and amplitudes	41
A.1	Noise systems and amplitudes	41
A.2	Distances from noise systems vs. amplitudes	42
B	Additional proofs	44
B.1	Proof of Proposition 3.14	44
B.2	Proof of Theorem 3.38	45
B.3	Proof of Proposition 3.43	46

1 Introduction

Persistent homology is, arguably, the most successful method in Topological Data Analysis (TDA), since it has been used in a variety of applications, ranging from nanoporous materials [LBD⁺17] to pulmonary diseases [BPC⁺18], with new preprints being released on a weekly basis¹. Such applications rest on stability results, which, roughly, guarantee that the pipeline from the input data to the barcodes is Lipschitz, with respect to appropriate notions of distances on the input data and barcodes. Such distances are well-understood in the one-parameter case, since they rely on matchings between intervals in barcodes, and they differ in how the cost between matchings is computed. In what is called the “bottleneck distance”, one considers the cost of matching the largest intervals, while in what are generally called “ p -Wasserstein distances” one takes the weighted sum of the costs of the matching over all the intervals. Of these distances, the bottleneck distance has been most widely used in applications [BW20, HKNU17] and has been studied the most from a theoretical point of view. However, for many applications the bottleneck distance is too coarse, and there have recently been renewed efforts in studying stability results for p -Wasserstein distances [ST20].

Since there is no generalisation of the barcode for multifiltrations of spaces [CZ09], it is thus not surprising that the study of distances in the multiparameter case is much more complex, and this is currently one of the most active areas of research in TDA. The interleaving distance generalises the bottleneck distance for multiparameter persistence, see [Les15] and references therein to other distances for multiparameter persistence. At the same time, it admits a sheaf theoretical interpretation under the identification of observable persistence modules with γ -sheaves [BP21]. More recently, Wasserstein-type distances and ℓ^p -distances have been studied in [BSS20] and [BBL21], respectively.

Invariants and distances are deeply intertwined, as defining a distance in the multiparameter setting entails first choosing an appropriate invariant. Choosing an appropriate invariant, in turn, requires defining what it means to be “persistent” in the multiparameter setting, so as to be able to distinguish between topological signal and noise. To do this, one could first focus either on finding an appropriate notion of signal, or of noise. Several different approaches,

¹A database of applications of methods in TDA, including persistent homology, is being maintained at [GL20], and at the time of writing it includes more than two hundred applications of persistent homology. The entries in the database can be filtered by application area, data format and topological methods.

focused on finding an appropriate notion of signal, have been taken in the literature. The first invariant that was suggested is the “rank invariant” [CZ09], which encodes the information of the ranks of structure morphisms in the module, and captures first-order information in the module (i.e. information about generators and relations between these). In their paper, Carlsson and Zomorodian also showed for the first time in the TDA literature that in an appropriate sense there is no invariant generalising the barcode. Following their publication, research in this area stalled, only to gain momentum again with the introduction of new techniques to the field, stemming from representation theory [BE18, BBLO20, BBCB20], and commutative algebra [HOST19, Tho19, Mil20a, BBOO21]. In a complementary line of work, the authors of [SCL⁺17] proposed an approach to study notions of “noise” for multiparameter persistence modules, with the idea of stabilising invariants by minimising them on disks, rather than proving a stability result for each of them. Note that the notion of minimising invariants in this fashion was also taken, in the 1-parameter case, in [BMSW17].

In our work, we take a different perspective: we focus on classes of invariants that capture a certain notion of “size” or “measure” of persistence, and on ways of associating distances with such invariants. Specifically, given an abelian category \mathcal{A} , we are interested in studying real-valued maps $\alpha: \text{ob } \mathcal{A} \rightarrow [0, \infty]$ that mimic the properties that we would expect a “size” function to satisfy. Namely, we would like the map to be monotone with respect to subobjects and quotients, and to be additive with respect to short exact sequences. However, the latter condition is too restrictive when \mathcal{A} is a category of persistence modules. Indeed, as we show in Section 3.3, any such invariant can be realised by integrating the Hilbert function, thus forgetting all the information carried by the structure morphisms.

This provides the motivation for introducing what we call an “amplitude”: this is a real-valued map that is monotone and subadditive. As it is apparent from the wealth of examples that we provide in Section 3, this framework is very rich. There are several interesting examples of amplitudes in TDA, including some tropical coordinates [Kal19], persistent magnitude [GH21], p -norms [ST20], shift along a cone [SCL⁺17], and support amplitude (Example 3.11). We give a summary of some examples of amplitudes from TDA in Table 1. Thus, an amplitude is a “size” or “measure” of persistence, and we show that this way of thinking about invariants gives a perspective that is very powerful.

In Section 4, we study different ways to associate distances with an amplitude. In particular, we show that many different distances studied in the literature can be expressed as distances arising from amplitudes. For instance, the interleaving distance [Les15], the L^p -distance from [HOST19, LW15], and the metrics from [BSS20] are all either alternatively given by path-metrics

Amplitude α	Location in the paper	Relationship of d_α to known distances
p -norms [ST20]	Example 3.8	equivalent to p -Wasserstein Cor. 4.22
shift amplitude [SCL ⁺ 17]	Example 3.9	equivalent to interleaving Prop. 4.18
tropical coordinates [Kal19]	Example 3.12	bounded by p -Wasserstein and bottleneck
total persistence [CSEHM10]	Example 3.15	bounded by 1-Wasserstein
persistent magnitude [GH21]	Example 3.16	bounded by 1-Wasserstein

Table 1: A few examples of amplitudes from TDA, together with a summary of how the associated path metrics d_α (see Definition 4.4) are related to known distances in the literature.

of amplitudes or equivalent to distances obtained from amplitudes.

When using persistent homology to study data, one often seeks the best way to capture persistence, depending on the application problem. Therefore one is interested in several amplitudes at once and in studying how to relate these different amplitudes in a robust manner. In the one-parameter setting, there is a standard choice of distances on persistence modules, namely the bottleneck and Wasserstein distances [CSEHM10]. When studying the robustness of amplitudes in the one-parameter setting one therefore asks for their distances to be Lipschitz with respect to the bottleneck or Wasserstein distance. To our knowledge, most stability theorems for 1-parameter persistence use the absolute-value distance (Definition 4.1) associated to an amplitude (see, for instance, the stability results in [CSEHM10, Kal19]). This distance has the drawback that it forgets the underlying categorical structure. On the contrary, several of the other distances from amplitudes discussed in Section 4 capture some of this structure and thus give more refined information. In Section 5, we show how these distances can make it easier to prove new stability results, whereas proofs of stability results involving the absolute-value distance are often computationally involved. We thus believe that one of the main contributions of our work might be to highlight the need for considering distances from amplitudes that are more informative than the ones usually used in the literature.

In the multi-parameter case the situation is different, since there no longer is a canonical choice of distance; thus, here one is interested in studying changes of amplitude that are Lipschitz with respect to distances associated to these. We provide such a result in Section 5.3, and we give a high-level schematics of the types of stability results that we study in our framework in Fig. 1.

Finally, we also show how our framework can be used to study stability of invariants that fail to be amplitudes. For example, this is the case for the c_p -rank, which is not an amplitude as it fails to be monotone, but whose stability can nevertheless be studied in our framework (see Section 5.6).

To conclude, we believe that our work gives a perspective on the problem of studying distances and stability in multiparameter persistence that is both rich, as is shown by the wealth of examples of invariants and distances that can be studied in this framework, as well as powerful, as it allows to obtain new stability results, even for invariants that are not amplitudes.




Figure 1: Schematics of types of stability results we are interested in studying in our framework: a choice of invariant determines a distance on persistence modules, and we thus are interested in studying amplitude-bounding functors G that are continuous in an appropriate sense. Given such a continuous functor G , if one has a stability result guaranteeing that the pipeline from data to the choice of invariant α is stable, one can infer a similar stability result also for the invariant β . We discuss such inferred stability results in Section 5.3.1, while we discuss continuity of functors in Section 5.2.

1.1 Related work

As already mentioned above, some of the ideas that we study in our work have been explored in various different formulations, among others, in [SCL⁺17, BSS20, ST20]. Here we summarise briefly the relationship between our work and the already existing literature, and then we discuss in detail this relationship in the main body.

In [SCL⁺17], the authors introduced the concept of a “noise system” as a way to study different choices of what it means to be noise in a persistence module. They also gave a way to define a distance on persistence modules based on a specific choice of noise. As we show in Appendix A, noise systems and amplitudes as two different perspective for the same theory. This in particular allows us to use some of the results from [SCL⁺17] when studying specific distances obtained from amplitudes in Section 4. Furthermore, among our examples of amplitudes, we present two examples (Example 3.9 and Example 3.10) that were originally given in [SCL⁺17].

Notions related to noise systems also appear in recent work [ST20] in the definition of p -norms. Here, we note that certain characteristics of p -norms make them into amplitudes (see Example 3.8). Moreover, in Section 4.2, we show that Wasserstein-type distances defined from p -norms in [ST20] are equivalent to the distances that we can define from the amplitudes corresponding to p -norms.

Finally, in [BSS20], the authors take a more general approach by considering what they call “weights”, which are arbitrary real-valued maps on an abelian category sending the zero object to zero. They use weights to define Wasserstein-type distances on multiparameter persistence modules. We note that the main example of weight that the authors consider is an amplitude.

2 Persistence modules and finiteness conditions

In this section, we first cover some basic notions about multiparameter persistence modules, and then discuss (novel) finiteness conditions for these.

Notation 2.1. We denote by Vect the (abelian) category of finite dimensional vector spaces over a (fixed) field \mathbb{F} . We use \mathcal{Q} to denote an arbitrary poset, and by an abuse of notation we denote by \mathcal{Q} also the posetal category associated to \mathcal{Q} .

Definition 2.2. A functor $\mathcal{Q} \rightarrow \text{Vect}$ is a **persistence module**. When $\mathcal{Q} = S^n$ for S a totally ordered set and $n \geq 1$ a natural number, we also call such a functor an **n -parameter persistence module**. Furthermore, when $n \geq 2$, we alternatively call such a module a **multiparameter persistence module**.

Persistence modules are one of the main objects of study in TDA: by computing simplicial homology with coefficients in a field of an n -parameter family of simplicial complexes, one obtains for each homology degree an n -parameter persistence module.

Example 2.3. Let $X \subset \mathbb{R}^m$ be a finite set of points. Let $p: X \rightarrow \mathbb{R}_{\geq 0}$ be a density estimation of the points, for instance a kernel density estimation. For real numbers $\delta \geq 0$ and $\epsilon \geq 0$ let $V_{\delta, \epsilon}$ be the Vietoris–Rips simplicial complex at scale ϵ built on the subset of points x of X for which $p(x) \geq \delta$. We then obtain a double-nested sequence of simplicial complexes $\{V_{\delta, \epsilon}\}_{\delta, \epsilon \geq 0}$. By computing simplicial homology in degree k with coefficients in a field of these simplicial complexes we obtain a 2-parameter persistence module.

Persistence modules were first introduced for $n = 1$ in the Topological Data Analysis literature in [ZC05]. Any such module that satisfies certain finiteness conditions can be completely characterised up to isomorphism by a finite collection of intervals in the totally ordered set S , see [CK18] and references therein. This collection of intervals is collectively denoted as **barcode**, and describes

the lifetimes (“persistence”) of features in the filtration. Alternatively, such a collection of intervals is interpreted as a collection of points in the plane, called **persistence diagram**, and denoted by the symbol Dgm , whose coordinates are given by the endpoints of the bars in the interval decomposition of the module [CSEH07].

The question of whether barcodes can be generalised to the setting of multiple parameters was answered in the negative by Carlsson and Zomorodian in 2009 [CZ09]. Specifically, they showed that there is no generalisation of the barcodes for multiparameter persistence modules, as there do not exist isomorphism invariants that are discrete and do not depend on the field of coefficients that one chooses to compute homology. It is an ongoing object of research to study invariants of multiparameter persistence modules that are suitable for applications; one usually wishes such invariants to capture what it means to be persistent, to be robust in an appropriate sense, and to be computable. Our work is motivated by the study of robustness of such invariants, which relies on a definition of distance between such modules.

2.1 Finiteness conditions

Several finiteness conditions for persistence modules have been introduced in the literature, and we review some of them in this section. We then introduce cubical encodings, which are a new finiteness condition. Being cubically encoded is a property that is more general than being finitely presented, but at the same time, it is a property that is restrictive enough to guarantee that the category of cubically encoded persistence modules is abelian. This is important because the types of invariants that we study in our work are real-valued functions defined on the class of objects of an abelian category. At the same time, we need to consider a more general condition than being finitely presented when discussing additive amplitudes (see Section 3.3).

We begin by recalling briefly the main finiteness conditions, what the relationship between them is, and illustrate the different concepts with examples in Figure 2:

1. **Tame functors** were introduced in [SCL⁺17]. Roughly, such functors are extensions of functors defined on \mathbb{N}^n to functors defined on \mathbb{Q}^n that are constant on right-open cubes of \mathbb{Q}^n with integer side length.
2. In [LW15], the authors introduce **continuous extensions**, which are more general than tame functors as they are required to be constant on right-open (hyper-)rectangles with real side length, and show that any finitely presented \mathbb{R}^n -graded module can be realised as a continuous extension of a discrete module. We note that if a module is tame in the sense of [SCL⁺17], then it can be realised as a continuous extension in the sense of [LW15].
3. Finally, **finitely encoded modules** were introduced in (a previous version of) [Mil20b]. Note that now the author call them tame. The basic idea is that a finitely encoded module is a module that can be described using a finite poset (see the following for a more precise description). We observe that if a module can be realised as a continuous extension in the sense of [LW15], then it is finitely encoded in the sense of [Mil20b]. Being finitely encoded is a property that is too general for our purposes, because the category of such modules is not abelian [Mil20b, Ex. 4.25]. For this reason we will devote the rest of this section to discussing a specific class of finitely encoded modules.

In the case of 1-parameter persistence modules, all the previous coincide up to a condition at the endpoints of the intervals: for tame and continuous extensions, the intervals are closed on the left and open on the right, while for finitely encoded there are no restrictions.




Figure 2: Examples of different finiteness conditions. Each depicted module is \mathbb{F} in the blue region and 0 otherwise, with the identity as structure morphism whenever possible. The thick line denotes a closed boundary, and the lack of it an open boundary.

For the purposes of our work, we consider a finiteness condition that lies in-between continuous extensions and arbitrary finitely encoded modules, namely we consider the full subcategory of $\text{Vect}^{\mathbb{R}^n}$ given by the persistence modules that are finitely encoded by cubes.

Recall that an **algebra** on a set S is a subset of the power set of S that contains the empty set and is closed under complements and finite unions. We are going to use algebras to specify what kinds of encodings we allow for. For the scope of this paper, we mostly focus on the case of cubes, which provides a finiteness condition for persistence modules slightly more general than being finitely presented. We chose this level of generality because it is a good balance between the types of modules that are most studied in applications and the need to find the correct level of abstraction.

Definition 2.4. An **upset** of a poset \mathcal{Q} is a subset $U \subseteq \mathcal{Q}$ such that for any $u \in U$ and $q \in \mathcal{Q}$ whenever $u \leq q$ then $q \in U$. Similarly, a **downset** of a poset \mathcal{Q} is a subset $D \subseteq \mathcal{Q}$ such that for any $d \in D$ and $q \in \mathcal{Q}$ whenever $d \geq q$ then $q \in D$. An **interval** in \mathcal{Q} is the intersection of an upset and a downset of \mathcal{Q} .

Definition 2.5. A **cube** in \mathbb{R}^n , or in \mathbb{Z}^n , is a subset of the form $I_1 \times \cdots \times I_n$, where the I_j are intervals in \mathbb{R} , resp. \mathbb{Z} . The **algebra of cubes** in \mathbb{R}^n is given by

$$\mathfrak{C} := \{S \subset \mathbb{R}^n \mid S \text{ is a finite union of cubes}\}.$$

We use the same language and notation for the algebra of cubes in \mathbb{Z}^n , mutatis mutandis.

Observe that, in the above definitions, we allow for intervals to be infinite and open or closed at either end.

Definition 2.6. An **encoding** of an object $M \in \text{ob Vect}^{\mathcal{Q}}$ by a poset \mathcal{Q}' is a morphism $e: \mathcal{Q} \rightarrow \mathcal{Q}'$ together with an object $M' \in \text{ob Vect}^{\mathcal{Q}'}$ and an isomorphism $\eta: M \xrightarrow{\sim} e^*M'$ between the object and the pullback along e of M' . Moreover, the encoding is **finite** if \mathcal{Q} is finite and M' is pointwise finite dimensional. Let \mathfrak{X} be an algebra on \mathcal{Q} and $e: \mathcal{Q} \rightarrow \mathcal{Q}'$, $\eta: M \xrightarrow{\sim} e^*M'$ a finite encoding of M . If e is such that all of its **fibers** $e^{-1}(q')$ are elements of \mathfrak{X} , then we say that the encoding is **of class \mathfrak{X}** or an **\mathfrak{X} -encoding**.

Definition 2.7. An object M in $\text{ob Vect}^{\mathcal{Q}}$ is **finitely encoded** if it admits a finite encoding. Moreover, M is **\mathfrak{X} -encoded**, or **encoded by \mathfrak{X}** , if it admits a (finite) \mathfrak{X} -encoding. In the case where $\mathcal{Q} = \mathbb{R}^n$, or $\mathcal{Q} = \mathbb{Z}^n$, and $\mathfrak{X} = \mathfrak{C}$, the algebra of cubes, we say that M is **finitely encoded by cubes** or simply **cubically encoded**.

Remark 2.8. Note that for the definition of a finitely or \mathfrak{X} -encoded module, it is not necessary to work only with functors into the category of vector spaces. Hence, we will sometimes make use of the same language for elements of $\mathcal{A}^{\mathcal{Q}}$, where \mathcal{A} is any abelian category. All of the definitions translate, *mutatis mutandis*.

Notation 2.9. We denote by $\text{PerM}(\mathcal{Q})$ the full subcategory of $\text{Vect}^{\mathcal{Q}}$ given by finitely encoded persistence modules, and by $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ the full subcategory given by the objects that are finitely \mathfrak{X} -encoded. Note that, when $n = 1$, the two categories $\text{PerM}(\mathbb{R}^n)$ and $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ coincide. Finally, we denote by $\text{PerM}_{fp}(\mathbb{R}^n)$ the full (abelian) subcategory of $\text{Vect}^{\mathbb{R}^n}$ given by finitely presented persistence modules.




Figure 3: Two finitely encoded modules.

To clarify the concept of finitely encoded modules, we present a running example. Consider the modules M, N over \mathbb{R}^2 in Fig. 3, which are zero in the blank part of \mathbb{R}^2 and given by \mathbb{F} respectively on the blue and on the green infinite interval (with structure morphisms the identity, whenever possible, and the zero morphism, whenever necessary). These two modules are finitely presented, so they could be described without encoding, but here we use them as a simple example of it. By definition, we need to find two finite posets, two encoding maps and two isomorphisms. For the posets, we take two copies of the poset with two elements and one relation. To avoid confusion, we keep them separate calling them $\mathcal{P}_M = \{m_1 \leq m_2\}$ and $\mathcal{P}_N = \{n_1 \leq n_2\}$. The encoding of M and N are given by $e_M: \mathbb{R}^2 \rightarrow \mathcal{P}_M$ and $e_N: \mathbb{R}^2 \rightarrow \mathcal{P}_N$, defined as:

$$e_M(q) := \begin{cases} m_2 & \text{if } q \geq (2, 1) \\ m_1 & \text{otherwise} \end{cases}, \quad e_N(q) := \begin{cases} n_2 & \text{if } q \geq (1, 3) \\ n_1 & \text{otherwise} \end{cases}.$$

Let M' and N' be the modules over \mathcal{P}_M and \mathcal{P}_N , respectively, which are the trivial vector space when evaluated m_1 and n_1 , the one dimensional vector space when evaluated m_2 and n_2 and the map $0 \rightarrow \mathbb{F}$ for both the structure morphisms $m_1 \rightarrow m_2$ and $n_1 \rightarrow n_2$. The encoding isomorphisms are given by the identity.

Example 2.10. The previous example generalizes to any interval $I \subset \mathcal{Q}$ given by the intersection of an upset and a downset in an algebra \mathfrak{X} . We define $\mathbb{F}[I]$, the **interval module on I** , as the module obtained as follows. Consider the finite poset $\{0 \leq 1\}^2$ and the map of partially ordered sets $e: \mathcal{Q} \rightarrow \{0 \leq 1\}^2$ induced by the two maps $e_D: \mathcal{Q} \rightarrow \{0 \leq 1\}$, $e_U: \mathcal{Q} \rightarrow \{0 \leq 1\}$ defined as:

$$e_D(q) := \begin{cases} 0 & \text{if } q \in D \\ 1 & \text{otherwise} \end{cases}, \quad e_U(q) := \begin{cases} 1 & \text{if } q \in U \\ 0 & \text{otherwise} \end{cases}.$$

Let $M' \in \text{Vect}^{\{0 \leq 1\}^2}$ be the unique persistence module with $M'_{(1,1)} = \mathbb{F}$ and 0 otherwise. Finally, define $\mathbb{F}[I] = e^*M'$. By construction, $\mathbb{F}[I]$ is \mathfrak{X} -encoded. Explicitly, $\mathbb{F}[I]$ is given by \mathbb{F} on I , 0

everywhere else and with identity as structure morphism for all relations in I . In case I is an interval in \mathbb{R} , we often use the more common notation $\mathbb{I}[I]$ instead. \triangleleft

A useful property of finitely encoded modules is that there always exists a common finite encoding.

Lemma 2.11. *Let M, N be two finitely encoded persistence modules over \mathcal{Q} . Then there exist a finite common encoding $e: \mathcal{Q} \rightarrow \mathcal{P}$ and two modules M'' and N'' with two isomorphisms $M \cong e^*M''$ and $N \cong e^*N''$. Furthermore, if both encodings are of class \mathfrak{X} , for some algebra \mathfrak{X} on \mathcal{Q} , then e can also be taken of class \mathfrak{X} .*

Proof. Let $e_M: \mathcal{Q} \rightarrow \mathcal{P}_M$ and $e_N: \mathcal{Q} \rightarrow \mathcal{P}_N$ be finite encodings of M and N , respectively, with modules M', N' . We define a map

$$\begin{aligned} e: \mathcal{Q} &\longrightarrow \mathcal{P}_M \times \mathcal{P}_N \\ q &\mapsto (e_M(q), e_N(q)). \end{aligned}$$

Define two modules M'' and N'' as the pullback along the canonical projections $\pi_M: \mathcal{P}_M \times \mathcal{P}_M \rightarrow \mathcal{P}_M$, $\pi_N: \mathcal{P}_M \times \mathcal{P}_N \rightarrow \mathcal{P}_N$ of M' and N' , respectively. It follows that there exists two isomorphisms $M \cong e^*M''$ and $N \cong e^*N''$, proving the claim. Furthermore, note that the fibers of e are, by construction, given by the pairwise intersection of the fibers of e_M and e_N . Hence, if the latter two are of class \mathfrak{X} , so is e . \square

Coming back to our running example of Fig. 3, we have that M and N can be encoded on the same posets, i.e. on two isomorphic posets. However, the encoding map is different, thus to find a common encoding we use the argument in the above proof. Consider the product poset $\mathcal{P}_M \times \mathcal{P}_N = \{(m_1, n_1), (m_1, n_2), (m_2, n_1), (m_2, n_2)\}$ with structure morphisms $(m_1, n_1) \rightarrow (m_2, n_1)$, $(m_1, n_1) \rightarrow (m_1, n_2)$, $(m_2, n_1) \rightarrow (m_2, n_2)$, and $(m_1, n_2) \rightarrow (m_2, n_2)$. Then the encoding is given by $e: \mathbb{R}^2 \rightarrow \mathcal{P}_M \times \mathcal{P}_N$ defined by:

$$e(q) := \begin{cases} (m_1, n_1) & \text{if } q \not\geq (2, 1) \text{ and } q \not\geq (1, 3) \\ (m_1, n_2) & \text{if } q \not\geq (2, 1) \text{ and } q \geq (1, 3) \\ (m_2, n_1) & \text{if } q \geq (2, 1) \text{ and } q \not\geq (1, 3) \\ (m_2, n_2) & \text{if } q \geq (2, 1) \text{ and } q \geq (1, 3) \end{cases}.$$

Finally, the modules M'' and N'' are given by the trivial vector space on (m_1, n_1) and (m_1, n_2) , and on (m_1, n_1) and (m_2, n_1) , respectively, by the one-dimensional vector space on (m_2, n_1) and (m_2, n_2) , and on (m_1, n_2) and (m_2, n_2) , respectively, and the appropriate maps induced by the structure morphisms in $\mathcal{P}_M \times \mathcal{P}_N$.

As an immediate consequence of Lemma 2.11 and the additivity of pullback functors, we obtain that the sum of two finitely encoded persistence modules (of a certain class \mathfrak{X}) is again finitely (\mathfrak{X} -)encoded. In other words, we have:

Corollary 2.12. *For any algebra \mathfrak{X} on \mathcal{Q} , the category $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ is an additive subcategory of $\text{Vect}^{\mathcal{Q}}$.*

For our purposes, we are particularly interested in the cases where $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ is a full abelian subcategory of $\text{Vect}^{\mathcal{Q}}$. Observe that this is not true in general (see [Mil20b, Ex. 4.25]). However, under certain connectivity assumptions, which are fulfilled for the algebra of cubes and certain subalgebras of the PL and semialgebraic sets, this turns out to be the case. We provide now a brief introduction to the topic, and refer to [Waa21] for more details and the full proofs.

Definition 2.13. We say a poset \mathcal{Q} is \leq -**connected** if it is connected as a category. In other words, if, for every $q, q' \in \mathcal{Q}$, there exists a zigzag

$$q \leq q_1 \leq \cdots \leq q_k \leq q'$$

of inequalities between the two elements, where \leq denotes either \leq or \geq . The maximal \leq -connected subsets of \mathcal{Q} are called the \leq -**components** of \mathcal{Q} .

Remark 2.14. Every cube in $C \subset \mathbb{R}^n$ is \leq -connected. To see this, note that a zigzag between any two elements $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ of the cube is given by the elements

$$x \leq (y_1, x_2, \dots, x_n) \leq (y_1, y_2, x_3, \dots, x_n) \leq \cdots \leq (y_1, \dots, y_{n-1}, x_n) \leq y$$

As a cube is a product of subset of \mathbb{R} , all of these elements lie again in C .

More generally, every half open interval in the piecewise linear or semi-algebraic setup has finitely many \leq -components in \mathbb{R}^n .

Definition 2.15. An algebra \mathfrak{X} on a poset \mathcal{Q} is **connective** if every interval I in \mathfrak{X} has only finitely many \leq -components and all of these components lie in \mathfrak{X} .

Remark 2.16. The algebra of cubes \mathfrak{C} is connective. Indeed, let $I \subset \mathbb{R}^n$ be an interval given by a finite union of cubes. First, note that we can write I as a finite disjoint union of cubes C_i . By Remark 2.14, each of these cubes is \leq -connected. Hence, also all of the \leq -components of I can be written as a finite disjoint union of C_i . As the components are pairwise disjoint and there are only finitely many C_i , there can only be finitely many \leq -components.

Remark 2.17. It turns out (Theorem 2.18) that for connective algebras the category $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ is a full abelian subcategory of $\text{Vect}^{\mathcal{Q}}$. However, the analogous argument as in Remark 2.16 does not work in the piecewise linear or semialgebraic setting, for example. In fact, the antidiagonal $\{(x, y) \mid x = -y\}$ already is a PL interval with infinitely many \leq -connected components. For many applications it can be of interest to allow for PL or semialgebraic encodings (compare [Mil20b]) while still retaining abelianity of the underlying category. Under minor additional topological assumptions, one can generally obtain connectiveness as long as one has control over the topological path components. This will be discussed in greater details in [Waa21].

Finally, the following theorem allows us to freely work in the framework of abelian categories, as long as we restrict to sufficiently nice classes of encodings, such as the cubical one.

Theorem 2.18. ([Waa21, Thm. 3.7]) Let \mathfrak{X} be a connective algebra on the poset \mathcal{Q} . Then the category $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ is a full abelian subcategory of $\text{Vect}^{\mathcal{Q}}$.

Corollary 2.19. The category $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ is a full abelian subcategory of $\text{Vect}^{\mathbb{R}^n}$.

3 Amplitudes

As explained in the Introduction, one of the main goals of our work is to relate different ways to generalise what it means for an element or submodule of a module to be “persistent”. We thus begin with the definition of amplitudes, which are real-valued maps that behave well with respect to the structure of abelian categories, and then proceed to provide many examples of amplitudes.

Definition 3.1. For an abelian category \mathcal{A} , a class function $\alpha: \text{ob } \mathcal{A} \rightarrow [0, \infty]$ is called an **amplitude** if $\alpha(0) = 0$ and for every short exact sequence

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

in \mathcal{A} the following inequalities hold:

$$\begin{aligned} \text{(Monotonicity)} \quad & \alpha(A) \leq \alpha(B) \\ & \alpha(C) \leq \alpha(B) \quad , \end{aligned} \tag{1}$$

$$\text{(Subadditivity)} \quad \alpha(B) \leq \alpha(A) + \alpha(C) \quad . \tag{2}$$

Moreover, if (2) is an equality, the amplitude is said to be **additive**, and if $\alpha(A) < \infty$ for all A in $\text{ob } \mathcal{A}$, then the amplitude is said to be **finite**.

First, we note that an amplitude is an isomorphism invariant:

Lemma 3.2. *Let \mathcal{A} be an abelian category together with an amplitude α . Let $A, B \in \text{ob } \mathcal{A}$ and assume that there is an isomorphism $A \rightarrow B$. Then $\alpha(A) = \alpha(B)$.*

Proof. This is a direct consequence of property (1) of an amplitude. \square

Given any abelian category \mathcal{A} , we always have a trivial amplitude α by setting $\alpha(A) = 0$ for all A in $\text{ob } \mathcal{A}$. We now give some examples of more meaningful amplitudes.

Example 3.3. (Rank amplitude) Let $R\text{-Mod}$ be the abelian category of modules over a fixed commutative ring R . By standard results in algebra, the map $M \mapsto \text{rank}_R M$ gives an additive amplitude on $R\text{-Mod}$. In particular, the map $V \rightarrow \dim V$ gives an additive amplitude on Vect . \triangleleft

Example 3.4. ((\mathbb{Z}/p) -rank amplitude) Let Ab^{fg} be the abelian category of finitely generated abelian groups and group homomorphisms. Fix a prime number p . We define the (\mathbb{Z}/p) -rank of a finitely generated abelian group G as $r_p(G) = \text{rank}_{\mathbb{Z}/p}(G/pG)$. Using the fundamental theorem of finitely generated modules over a principal ideal domain, one can show that, for any prime p , the (\mathbb{Z}/p) -rank r_p gives an amplitude on Ab^{fg} . \triangleleft

Example 3.5. (Length of a module) Recall that the length $\text{Len}_R(M)$ of a module M over a ring R is defined as

$$\text{Len}_R(M) := \sup\{k \mid \exists \text{ a sequence of } R\text{-submodules } 0 = M_0 \subsetneq \cdots \subsetneq M_k = M\}.$$

Here we allow for the possibility $\text{Len}_R(M) = \infty$. Note that $\text{Len}_R(0) = 0$.

Let \mathcal{A} be the abelian category of finitely generated \mathbb{N} -graded modules over $\mathbb{F}[x_1, \dots, x_n]$. Let \mathfrak{p} be a homogeneous prime ideal of $\mathbb{F}[x_1, \dots, x_n]$, and define an amplitude $\mu_{\mathfrak{p}}$ on \mathcal{A} by setting

$$\mu_{\mathfrak{p}}(M) = \text{Len}_{\mathbb{F}[x_1, \dots, x_n]_{\mathfrak{p}}}(M_{\mathfrak{p}}),$$

where the subscript $(-)_{\mathfrak{p}}$ indicates the localisation at \mathfrak{p} . Then $\mu_{\mathfrak{p}}$ is an additive amplitude. In fact, suppose we have a short exact sequence

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0.$$

Then the localized sequence

$$0 \rightarrow A_{\mathfrak{p}} \rightarrow B_{\mathfrak{p}} \rightarrow C_{\mathfrak{p}} \rightarrow 0$$

is still exact. If $A_{\mathfrak{p}}$ and $C_{\mathfrak{p}}$ have finite length, then additivity of $\mu_{\mathfrak{p}}$ follows from the Jordan-Hölder theorem [Lan02, §III.8]. If either (or both) of $A_{\mathfrak{p}}$ or $C_{\mathfrak{p}}$ has infinite length, the same must be true for $B_{\mathfrak{p}}$, so additivity holds in these cases as well. \triangleleft

Example 3.6. (Max degree) Consider the category Ch of bounded, nonnegatively-graded chain complexes over Vect , and chain maps between them. Define an amplitude δ on Ch as

$$\delta(C) := \max\{n \in \mathbb{N} \mid C_n \neq 0\}$$

for all C in ob Ch . It is clear that $\delta(0) = 0$. Since a short sequence in Ch is exact if and only if it is exact degreewise, we also have the properties of monotonicity (1) and subadditivity (2). Moreover, δ is finite. However, it is immediate to see that, in general, (2) is a strict inequality for this amplitude, and thus δ is not additive. More generally, δ is also an amplitude over the category of non-negative chain complexes and chain maps over any abelian category. \triangleleft

Example 3.7. (New amplitudes from old ones) Let α and β be amplitudes on an abelian category \mathcal{A} . Then, for all $t \in [0, \infty)$, the map $t\alpha$ is also an amplitude on \mathcal{A} , as a direct computation shows. It is also straightforward to check that the assignments $A \mapsto \max(\alpha(A), \beta(A))$ and $A \mapsto \alpha(A) + \beta(A)$ are both amplitudes on \mathcal{A} .

However, the assignment $A \mapsto \min(\alpha(A), \beta(A))$ may not be an amplitude on \mathcal{A} . For a simple example where this fails, consider the abelian category 2-Vect of pairs of finite-dimensional \mathbb{F} -vector spaces, where morphisms $(V, W) \rightarrow (V', W')$ are pairs of linear maps $(\varphi: V \rightarrow V', \psi: W \rightarrow W')$. Define amplitudes α and β by $\alpha(V, W) = \dim V$ and $\beta(V, W) = \dim W$. Let $\gamma(V, W) = \min(\alpha(V, W), \beta(V, W))$ for all $(V, W) \in \mathcal{A}$. Then γ is not an amplitude. In fact, consider the s.e.s.

$$0 \rightarrow (\mathbb{F}, 0) \rightarrow (\mathbb{F}, \mathbb{F}) \rightarrow (0, \mathbb{F}) \rightarrow 0.$$

Here $\gamma(\mathbb{F}, \mathbb{F}) = 1$, but $\gamma(\mathbb{F}, 0)$ and $\gamma(0, \mathbb{F})$ are both zero, so the subadditivity axiom fails. \triangleleft

3.1 Amplitudes in TDA

In this section, we describe several examples of invariants from TDA that satisfy the axioms of an amplitude, and three that do not.

Example 3.8. (p -norm) In [ST20], the authors consider p -norms of finitely presented 1-parameter persistence modules and use them to define distances isometric to the p -Wasserstein distances. Here, we use the p -norms to define amplitudes. Let M be a finitely encoded persistence module in $\text{PerM}(\mathbb{R})$ with barcode decomposition given by the collection of interval modules $\{\mathbb{I}\langle b_j, d_j \rangle\}_{j \in J}$, where we used angular brackets to denote that the interval can be open or closed on each endpoint. Then the p -norm of M , for $p \in [1, \infty]$, is

$$\rho_p(M) := \begin{cases} \left(\sum_{j \in J} |d_j - b_j|^p \right)^{\frac{1}{p}} & \text{if } p < \infty \\ \max_{j \in J} |d_j - b_j| & \text{if } p = \infty \end{cases}.$$

Moreover, using the results of [BL15], the authors of [ST20] show that such p -norms satisfy the monotonicity and subadditivity conditions (1)-(2). Thus, in the language of our work, the p -norms define amplitudes on $\text{PerM}(\mathbb{R})$. In particular, for $p = 1$, the amplitude of the 1-norm is additive. This amplitude is in general not finite since infinite interval modules have infinite p -norm. \triangleleft

As we will see in Appendix A.1, noise systems and amplitudes are closely related, and, as we discuss here, the former provide examples for the latter. The example that we describe here was originally presented in [SCL⁺17].

Example 3.9. (Shift amplitude) Let C be a cone in \mathbb{R}^n , i.e. the set of all linear combinations with non-negative coefficients of a finite collection of non-negative elements in \mathbb{R}^n , and $\|\cdot\|$ a norm on \mathbb{R}^n . Then a shift amplitude ζ on $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$ is defined as follows: for any $M \in \text{ob PerM}_{\mathcal{C}}(\mathbb{R}^n)$,

$$\zeta(M) := \inf\{\|c\| \mid c \in C \text{ and } M(t \leq t+c) = 0 \text{ for all } t \in \mathbb{R}^n\}.$$

The fact that this indeed is an amplitude follows from the properties of a noise system and the definition of standard noise in the direction of a cone in [SCL⁺17, Par. 6.2].

If the cone is generated by one element, then we get a simpler description of ζ , which also motivates the name. Let $c \in C$ be the only vector in C with $\|c\| = 1$. Then, for any $M \in \text{ob PerM}_{\mathcal{C}}(\mathbb{R}^n)$,

$$\zeta(M) := \inf\{\varepsilon \in \mathbb{R} \mid M(t \leq t + \varepsilon c) = 0\}.$$

Note that the shift amplitude ζ does not need to be finite, in general. To see this it is enough to consider a module that persists forever in the direction of the cone. In what follows, we often refer to the shift amplitude along a cone generated by one element as **shift along a vector** v . \triangleleft

Example 3.10. We discuss a further example of amplitude from noise systems, originally presented in [SCL⁺17]. We define the following amplitude on $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$

$$\alpha(M) := \max_{x \in \mathbb{R}^n} \dim M(x)$$

for any $M \in \text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$. This is an amplitude by the rank-nullity theorem in Vect and because a sequence is exact in $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$ if it is exact objectwise. In particular, α is a finite amplitude on $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$ but it is not additive. \triangleleft

Example 3.11. (Support amplitude) Let λ^* be the outer measure on \mathbb{R}^n obtained by extending Lebesgue measure to all subsets of \mathbb{R}^n . We can define a non-additive amplitude α on $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$ by

$$\alpha(M) := \lambda^* (\{x \in \mathbb{R}^n \mid M(x) \neq 0\})$$

for any M in $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$. This is an amplitude because of the monotonicity and subadditivity properties of outer measure [Bog07, Section 1.11]. \triangleleft

In the following examples, we describe different vectorisations of the barcode, namely tropical coordinates [Kal19], total persistence [CSEHM10] and persistent magnitude [GH21]. These vectorisations are useful, for example, in machine-learning techniques [GGM19, MOW20]. Each of them aims at capturing different information encapsulated in the barcode decomposition of persistence modules.

Example 3.12. (Tropical coordinates) Tropical coordinates were introduced in [Kal19] as a stable version of the vectorisation of barcodes suggested in [ACC13]. Tropical coordinates are expressions obtained by adding, taking maxima, or subtracting functions of the form

$$E_{m,(0,1)^k(1,0)^l(1,1)^p} = \sigma_{(0,1)^k(1,0)^l(1,1)^p}(\min\{b_1, ml_1\}, \ell_1, \dots, \min\{b_n, ml_n\}, \ell_n) \quad (3)$$

where we are considering a barcode with n intervals, and b_i is the birth-time of interval i , while ℓ_i is its length. The symbols $\sigma_{(0,1)^k(1,0)^l(1,1)^p}$ are the 2-symmetric max-plus polynomials described in [Kal19, Definition 2.10]. See also the discussion before Definition 2.10 in [Kal19] for how to associate a max-plus polynomial to a 2-symmetric polynomial.

We have that $\sigma_{(0,1)^k}(\min\{b_1, ml_1\}, \ell_1, \dots, \min\{b_n, ml_n\}, \ell_n)$ is the sum of the length of the k longest intervals. In this paper, we denote this function by $\ell\text{-trop}^k$, and we will show in Proposition 3.14 that it is an amplitude, for all $k \in \mathbb{N}$. The remaining two expressions are a little

more complicated. We have that $\sigma_{(1,0)^l}(\min\{b_1, m\ell_1\}, \ell_1, \dots, \min\{b_n, m\ell_n\}, \ell_n)$ is the sum of l largest expressions of the form $\min\{b_i, m\ell_i\}$, and the evaluation of $\sigma_{(1,1)^p}$ is the sum of the p largest expressions of the form $\min\{b_i + \ell_i, (m+1)\ell_i\}$. As we will show in Proposition 3.14, the former is not an amplitude. We reserve the study of the latter for future work.

Arbitrary tropical coordinates are defined by taking sums, maxima, or differences of expressions as in Eq. (3). We note that, when restricting to maxima and sums (see Example 3.7), to obtain an amplitude, by Proposition 3.14, it is sufficient to take $l = 0$ and $p = 0$ in Eq. (3). However, the difference of two amplitudes does in general not give an amplitude. We reserve for future work the study of which other tropical coordinates are amplitudes. \triangleleft

Definition 3.13. Let $M \in \text{ob PerM}(\mathbb{R})$ be an interval-decomposable module given by $M = \bigoplus_{i=1}^n \mathbb{I}\langle b_i, b_i + \ell_i \rangle$, where the each interval can be closed or open at endpoints. Then $\ell\text{-trop}^k(M)$ is defined to be the sum of the length of the k -longest intervals in M .

Proposition 3.14. *On the category $\text{PerM}(\mathbb{R})$:*

- i) *The tropical coordinate $\ell\text{-trop}^k$ is an amplitude, for all $k \in \mathbb{N}$;*
- ii) *The tropical coordinate $\sigma_{(1,0)^l}(\min\{b_1, m\ell_1\}, \ell_1, \dots, \min\{b_n, m\ell_n\}, \ell_n)$ is not an amplitude for any $m, l \in \mathbb{N}$.*

For the proof, we refer to Appendix B, precisely Proposition B.1.

The next vectorisation extracts the information of how long, in total, an homological class can be found in the data, and it is appropriately called total persistence.

Example 3.15. (Total persistence) Consider a persistence module $M \in \text{ob PerM}(\mathbb{R})$ with a decomposition $M \cong \bigoplus_{i=1}^m \mathbb{I}\langle b_i, d_i \rangle$. Its **total persistence** [CSEHM10] is given by $\text{totpers} = \sum_{i=1}^m (d_i - b_i)$. Since this is, in particular, equal to the tropical coordinate $\ell\text{-trop}^m$, and since the latter is an amplitude by Proposition 3.14, by passing to a sufficiently large m , we have that totpers is an amplitude. \triangleleft

Example 3.16. (Persistent magnitude) In [GH21], the authors introduced a new invariant for 1-parameter persistence modules, called persistent magnitude. Let M be finitely presented with an interval decomposition $M \cong \bigoplus_{i=1}^m \mathbb{I}[b_i, d_i]$. Then the **persistent magnitude** of M is the non-negative real number

$$|M| := \sum_{i=1}^m e^{-b_i} - e^{-d_i}.$$

The persistent magnitude is additive with respect to short exact sequences [GH21, Proposition 5.3], and thus $|\cdot|$ is an additive amplitude on the full subcategory of finitely presented persistence modules.

Govc and Hepworth introduced this invariant motivated by the following asymptotic relationship between the magnitude of a metric space A and its blurred magnitude homology:

$$|tA| = \sum_{k=0}^{\infty} (-1)^k \sum_{j=1}^{m_k} \left(e^{-ta_{k_j}} - e^{-tb_{k_j}} \right) \quad \text{for all } t > 0 \text{ large enough,}$$

where tA is the metric space obtained by A by rescaling the distances by t , and $\{[a_{k_j}, b_{k_j}]\}_{j=1}^{m_k}$ is the barcode of the k -th blurred magnitude homology of A , introduced in [Ott18].

We note that the authors in [GH21] also define persistent magnitude for \mathbb{N} -graded persistence modules $M \in \text{ob PerM}_{fp}(\mathbb{R}^n)$, motivated by the fact that persistent homology yields such modules, one in each homology degree. In the present work, when considering TDA applications, we focus

on invariants associated to a single homology degree. In future work, we intend to study also invariants associated to all homology degrees at once, and how these are related to the invariants studied in the present manuscript. \triangleleft

We conclude this section with some non-examples.

Example 3.17. (Number of indecomposables) Let \mathcal{A} be an abelian, Krull-Schmidt category. We can define a real-valued function w on the objects of \mathcal{A} as follows: $w: \text{ob } \mathcal{A} \rightarrow \mathbb{N}$, $M \mapsto n$, where n is the number of indecomposable of M .

Note that such a function is not an amplitude. To see this, take the object M to be the following commutative ladder (see [BE18]) in the abelian category of tame multi-parameter persistent modules (see Section 2):

$$\begin{array}{ccccccccc}
 \mathbb{F} & \xrightarrow{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} & \mathbb{F}^2 & \xrightarrow{1} & \mathbb{F}^2 & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & \mathbb{F} & \longrightarrow & \mathbf{0} \\
 \uparrow & & \uparrow \begin{bmatrix} 1 \\ 0 \end{bmatrix} & & \uparrow \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} & & \uparrow \begin{bmatrix} 1 & 1 \end{bmatrix} & & \uparrow \\
 \mathbf{0} & \longrightarrow & \mathbb{F} & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & \mathbb{F}^2 & \xrightarrow{1} & \mathbb{F}^2 & \xrightarrow{\begin{bmatrix} 0 & 1 \end{bmatrix}} & \mathbb{F}
 \end{array} .$$

Define M' as the subobject of M which is equal to M in the upper row and zero otherwise. Then $w(M) = 1$ and $w(M') = 2$, but $M' \hookrightarrow M$. Thus, w is not monotone with respect to subobjects and thus is not an amplitude. \triangleleft

Example 3.18. (Rank and multirank) The **rank invariant** is a function $\text{rank}_M: \mathcal{Q} \times \mathcal{Q} \rightarrow \mathbb{N}$ for a \mathcal{Q} -module M that takes a pair of points in the parameter space to the rank of the structure morphism

$$\text{rank}_M: (s, q) \mapsto \text{rank}(M(s) \rightarrow M(q))$$

where the rank of the morphism is 0 if the points are not comparable or s does not precede q . The rank invariant was first introduced in [CSZ10] for the poset $\mathcal{Q} = \mathbb{N}^n$, and has since been studied for arbitrary posets, for instance in the recent work [Tho19].

By properties of vector spaces, for every fixed pair of points $(s, q) \in \mathcal{Q} \times \mathcal{Q}$ and for any s.e.s. $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$, we have

$$\text{rank}_{M'}(s, q), \text{rank}_{M''}(s, q) \leq \text{rank}_M(s, q) \quad (4)$$

However, the rank is not subadditive. Set for example $M \cong \mathbb{I}[0, 2)$, $M' \cong \mathbb{I}[1, 2)$, and $M'' \cong \mathbb{I}[0, 1)$, with the obvious inclusion and surjection. Then, taking $(s, q) = (0, 2)$, we obtain a contradiction.

Similar arguments also hold for the m-rank in [Tho19]. \triangleleft

Even if the previous invariant is not an amplitude, we stress that it possible to describe it as a composition of a functor with an amplitude. Indeed, for every \mathcal{Q} -module M , $\text{rank}_M(s, q)$ is obtained by the composition of the non-exact functor $\text{im}(M(p \leq q))$ with the amplitude dim . This is not the only non-amplitude invariant that can be described using an amplitude, as we will show in Section 3.4 for the next example. This description is quite fruitful: as we will discuss in Section 5.6, it allows us to study the stability of non-amplitude invariants via the stability of amplitudes.

Example 3.19. Here we recall the definition of $c_{\mathfrak{p}}$ -rank proposed in [HOST19, Section 4.4]. Consider the abelian category \mathcal{A} of finitely generated \mathbb{N}^n -graded modules M over the polynomial ring $\mathbb{F}[x_1, \dots, x_n]$ on n variables and with coefficients in a field \mathbb{F} . Let \mathfrak{p} be an associated prime ideal of M , namely, \mathfrak{p} is a prime ideal such that

$$\mathfrak{p} = \left\{ s \in \mathbb{F}[x_1, \dots, x_n] \mid s \cdot m = 0 \text{ for some } m \in M \right\}$$

One has that any associated prime ideal \mathfrak{p} in $\mathbb{F}[x_1, \dots, x_n]$ is generated by a subset of the variables. For simplicity, assume that $\mathfrak{p} = \langle x_1, \dots, x_t \rangle$. The 0-th local cohomology of M with respect to \mathfrak{p} is the module over $\mathbb{F}[x_1, \dots, x_n]$ given by:

$$H_{\mathfrak{p}}^0(M) = \{m \in M \mid \mathfrak{p}^k \cdot m = 0 \text{ for all } k \text{ large enough}\}.$$

When M is finitely generated as a module over $\mathbb{F}[x_1, \dots, x_n]$, then so is $H_{\mathfrak{p}}^0(M)$. Furthermore, one also has that $H_{\mathfrak{p}}^0(M)$ is finitely generated over the ring $\mathbb{F}[x_{t+1}, \dots, x_n]$. The $c_{\mathfrak{p}}$ -rank of M is defined to be the rank of $H_{\mathfrak{p}}^0(M)$ as a module over $\mathbb{F}[x_{t+1}, \dots, x_n]$, see [HOST19, Definition 4.19].

Intuitively, the $c_{\mathfrak{p}}$ -rank gives information about the number of ways in which a module goes to infinity in the directions orthogonal to \mathfrak{p} , see [HOST19, Proposition 4.20] for a precise statement. This definition of $c_{\mathfrak{p}}$ -rank depends on a specific module M . Since we aim to interpret it as a function on the collection of finitely generated modules, we give a more general definition, and for ease of reference we state it in Definition 3.20. \triangleleft

To give the more general definition of $c_{\mathfrak{p}}$ -rank, we note that every homogeneous prime ideal of $\mathbb{F}[x_1, \dots, x_n]$ is generated by a subset of the variables [HOST19, Lemma 4.6], and thus we may again assume for simplicity that such a prime is of the form $\mathfrak{p} = \langle x_1, \dots, x_t \rangle$. Furthermore, we have that, for any finitely generated module M , the module $H_{\mathfrak{p}}^0(M)$ is finitely generated over $\mathbb{F}[x_{t+1}, \dots, x_n]$, as the statement in Lemma 4.18 in [HOST19] holds more generally for any homogeneous prime ideal \mathfrak{p} . We thus can give the following:

Definition 3.20. Consider the abelian category \mathcal{A} of finitely generated \mathbb{N}^n -graded modules M over the polynomial ring $\mathbb{F}[x_1, \dots, x_n]$ with coefficients in a field \mathbb{F} . Let \mathfrak{p} be any homogeneous prime ideal in $\mathbb{F}[x_1, \dots, x_n]$, and for simplicity assume that $\mathfrak{p} = \langle x_1, \dots, x_t \rangle$. We define the **$c_{\mathfrak{p}}$ -rank** $c_{\mathfrak{p}}(M)$ of any module $M \in \text{ob } \mathcal{A}$ to be the rank of $H_{\mathfrak{p}}^0(M)$ as a module over $\mathbb{F}[x_{t+1}, \dots, x_n]$.

We can interpret the $c_{\mathfrak{p}}$ -rank of a module as measuring the size of its submodule generated by elements living along the coordinate directions orthogonal to those spanned by \mathfrak{p} , see [HOST19, Proposition 4.20]. We note that this interpretation is valid for general homogeneous primes \mathfrak{p} , and not just associated primes; if \mathfrak{p} is a homogeneous prime that is not an associated prime, then the $c_{\mathfrak{p}}$ -rank is zero, and this reflects the fact that there are no elements of the module that live only along the coordinate directions orthogonal to those spanned by the generators of \mathfrak{p} .

Further, we note that this function is not an amplitude, since it fails to be monotone, see Example 3.21. However, we will discuss in Section 3.4 how the continuous version of this function can be constructed as the composition of an amplitude together with two functors, and we will show how one can use our framework to study its stability in Section 5.6.

Example 3.21. Consider the \mathbb{N}^2 -graded modules $A = \mathbb{F}[x_1, x_2]$, $B = x_1^2 \mathbb{F}[x_1, x_2]$ and $C = x_1^2 \mathbb{F}[x_1, x_2] / \langle x_1^2 \rangle$ over $\mathbb{F}[x_1, x_2]$. Furthermore, consider the short exact sequence

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0,$$

where $A \rightarrow B$ is given by multiplication by x_1^2 and $B \rightarrow C$ is the canonical projection. Then $c_{\langle x_1 \rangle}(A) = c_{\langle x_1 \rangle}(B) = 0$ but $c_{\langle x_1 \rangle}(C) = 2$. \triangleleft

3.2 L^p -amplitudes

Here, we define a useful construction of amplitudes on (sub)categories of persistence modules. The idea is to define an amplitude on a functor category by turning an amplitude on the target category into a function, and then taking the L^p -norm of the resulting function with respect to a measure on the source poset category. A guiding example is the integration of the Hilbert

function for finitely encoded persistence modules. We remark that this technique, per se, is not completely new as it was already introduced in a more different setting, for example, in [BSS20]. There the author works with more restrictive finiteness conditions but using the more general notion of **weight function**, with are non-negative real valued functions whose only requirement is to weight the zero object zero.

In this subsection, let \mathcal{A} be any abelian category, equipped with an amplitude α . Further, let \mathcal{Q} be a poset, equipped with a σ -algebra Σ and a measure μ on the latter.

Definition 3.22. For A in $\text{ob } \mathcal{A}^{\mathcal{Q}}$, let $\alpha_A: \mathcal{Q} \rightarrow [0, \infty]$ be the function that assigns to each element of \mathcal{Q} its evaluation on A : $q \mapsto \alpha(A(q))$. Suppose that α_A is measurable for all A in $\text{ob } \mathcal{A}^{\mathcal{Q}}$. Then, given a measure μ on Σ and $p \in [1, \infty)$, the **L^p -amplitude** $\| - \|_p^\alpha$ is defined by

$$\|A\|_p^\alpha := \left(\int_{\mathcal{Q}} \alpha_A^p d\mu \right)^{\frac{1}{p}}.$$

Proposition 3.23. *The map $\| - \|_p^\alpha$ is an amplitude.*

Note that, by linearity of the integral, if α is additive then $\| - \|_1^\alpha$ is also additive.

Proof. Since clearly $\|0\|_p^\alpha = 0$, to check that $\| - \|_p^\alpha$ is indeed an amplitude we need to show that for any s.e.s.

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

in $\mathcal{A}^{\mathcal{Q}}$, we have:

$$\begin{aligned} \|A\|_p^\alpha, \|C\|_p^\alpha &\leq \|B\|_p^\alpha \\ \|B\|_p^\alpha &\leq \|A\|_p^\alpha + \|C\|_p^\alpha \end{aligned}$$

The first statement follows from the fact that α is nonnegative. The second statement follows from the Minkowski inequality [Bog07, Theorem 2.11.9]. \square

For instance, when \mathcal{Q} is \mathbb{N}^n or any finite poset, we can equip it with the discrete σ -algebra and consider the category of \mathcal{Q} -indexed persistence modules in Vect . Then, by definition, every Hilbert function is measurable with respect to any measure on \mathcal{Q} , and we can use it to define an L^p -amplitude using the Hilbert function. Note that this is also the main example of weight function in [BSS20].

If the σ -algebra Σ on \mathcal{Q} is not the whole power set of \mathcal{Q} , the construction of Definition 3.22 faces an obvious technical difficulty: there is no guarantee that, in general, the function α_A will be measurable. Hence, it does not always make sense to integrate such functions. To solve this issue, we introduce the notion of α -measurable subcategories of the functor category $\mathcal{A}^{\mathcal{Q}}$, for which we need the definition of σ -algebra (or more generally algebra) generated by the upsets of \mathcal{Q} (see Definition 2.4).

Definition 3.24. Let S be a set of subsets of a poset \mathcal{Q} . The (σ -)algebra **generated** by S is the smallest (σ -)algebra containing all the elements of S .

Example 3.25. Given an algebra \mathfrak{X} on \mathcal{Q} , we denote by \mathfrak{X}_{Up} the algebra generated by the upsets of \mathfrak{X} . Note that this algebra is explicitly described as the algebra of finite unions of intervals in \mathcal{Q} , which are intersections of upsets and downsets in \mathfrak{X} . For a \mathfrak{X} -encoding map $\mathcal{Q} \rightarrow \mathcal{Q}'$ these intervals are the only possible fibers. \triangleleft

Definition 3.26. Let \mathcal{C} be a subcategory of $\mathcal{A}^{\mathcal{Q}}$ and \mathfrak{A} an algebra on \mathcal{Q} . \mathcal{C} is **α -measurable** (with respect to \mathfrak{A}) if, for every $C \in \text{ob } \mathcal{C}$, the function α_C is \mathfrak{A} -measurable.

Even for pointwise finite modules, to is easy it find non-measurable modules.

Example 3.27. Consider the persistence module M given by \mathbb{F} on $V \subset \mathbb{R}$, where V is the Vitali set and by 0 on $\mathbb{R} \setminus V$, with 0 as all the nontrivial structure morphisms. Then M is pointwise finite dimensional. However its Hilbert function is not measurable. \triangleleft

The definition of L^p -amplitudes extends straightforwardly to such α -measurable categories.

Example 3.28. (Generalized L^p -amplitudes) Let \mathcal{C} be an α -measurable abelian subcategory of $\mathcal{A}^{\mathcal{Q}}$ with respect to an algebra \mathfrak{A} . In particular, $\mathcal{C} \hookrightarrow \mathcal{A}^{\mathcal{Q}}$ is exact. Let μ be a measure on \mathfrak{A} . Then, similarly to Definition 3.22, we also have a family of amplitudes, denoted with the same symbol $\| - \|_p^\alpha$, for $p \in [1, \infty)$, on the subcategory \mathcal{C} :

$$\|A\|_p^\alpha := \left(\int_{\mathcal{Q}} \alpha_A^p d\mu \right)^{\frac{1}{p}}.$$

The fact that also in this case $\| - \|_p^\alpha$, for all $p \in [1, \infty)$, is an amplitude on \mathcal{C} follows from the same arguments used in Definition 3.22, together with the exactness assumptions on $\mathcal{C} \hookrightarrow \mathcal{A}$. \triangleleft

In the finitely encoded setting, however, the σ -additivity condition of a measure is really not necessary to obtain a well defined notion of integration. For this, recall the definition of a content:

Definition 3.29. Given an algebra \mathfrak{A} on a set S , a function $\nu: \mathfrak{A} \rightarrow [0, \infty]$ is called a **content** if

1. $\nu(\emptyset) = 0$
2. $\nu(X \cup Y) = \nu(X) + \nu(Y)$ for all $X, Y \in \mathfrak{A}$ with $X \cap Y = \emptyset$.

In other words, contents are to algebras what measures are to σ -algebras, by replacing the σ -additivity condition by additivity.

Example 3.30. Measurable functions with only finitely many values $f = \sum_{i=0}^n \lambda_i 1_{X_i}$ $X_i \in \mathfrak{A}$, $\lambda_i \in [0, \infty]$ admit a well defined notion of integration with respect to contents, given by:

$$\int f d\nu = \sum_{i=0}^n \lambda_i \nu(X_i).$$

As a consequence, even in the framework of a content ν on an algebra \mathfrak{A} an α -measurable abelian subcategory $\mathcal{C} \subset \mathcal{A}^{\mathcal{Q}}$ admits a notion of L^p -amplitude. To see this, note that when $A \in \mathcal{A}^{\mathcal{Q}}$ is finitely encoded then α_A^p only takes finitely many different values. Hence, under this assumption the construction of the L^p -amplitude, for $p \in [1, \infty)$, is given by

$$\|A\|_p^\alpha := \left(\int_{\mathcal{Q}} \alpha_A^p d\nu \right)^{\frac{1}{p}}$$

also in the setting of algebras and contents. \triangleleft

The following lemma characterizes the precise relationship between the allowable class of encodings \mathfrak{X} and the measurability of $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$.

Lemma 3.31. *Let \mathfrak{A} and \mathfrak{X} be two algebras on \mathcal{Q} . Then the following statements are equivalent:*

1. $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ is dim-measurable with respect to \mathfrak{A} .
2. Every upset $U \subset \mathcal{Q}$, $U \in \mathfrak{X}$ is \mathfrak{A} -measurable or equivalently, $\mathfrak{X}_{\text{Up}} \subset \mathfrak{A}$.

Moreover, the implication $2. \implies 1.$ is still valid if, instead of $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$, we consider the class of \mathfrak{X} -encoded objects in any functor category $\mathcal{A}^{\mathcal{Q}}$ and, instead of the Hilbert function, we take the function α_A for any amplitude α on \mathcal{A} .

Proof. ($1. \implies 2.$) Given an upset $U \in \mathfrak{X}$, the indicator module $\mathbb{F}[U]$ defines a \mathfrak{X} -encoded persistence module with Hilbert function the characteristic function of U . Therefore, since by assumption the Hilbert function of any encoded module is \mathfrak{A} -measurable, we have that U is \mathfrak{A} -measurable.

($2. \implies 1.$) For the converse implication, assume that every upset is measurable. Let M be a persistence module given by a finite encoding $M \cong f^*N$, for $f: \mathcal{Q} \rightarrow \mathcal{Q}'$ monotone increasing and $N \in \text{Vect}^{\mathcal{P}}$, with \mathcal{P} finite. Note that, as f is monotone increasing, the inverse image under f of any point is an interval. In particular, f is \mathfrak{A} -measurable with respect to the algebras \mathfrak{X}_{Up} on \mathcal{Q} and the discrete one on \mathcal{Q}' . As $\mathfrak{X}_{\text{Up}} \subset \mathfrak{A}$, this also means f is \mathfrak{A} -measurable. Then $\dim(N): \mathcal{Q}' \rightarrow \mathbb{N}$ is measurable as the algebra on the source is discrete. Hence, $\dim(M) = \dim(N) \circ f$ is \mathfrak{A} measurable as a composition of measurable functions, proving the claim.

The proof of the second implication in the general setting is identical, using the fact that an amplitude is an isomorphism invariant. \square

Corollary 3.32. *The category of cubically encoded persistence modules $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ is dim-measurable with respect to the Lebesgue algebra on \mathbb{R}^n .*

A proof of this corollary can be also obtained, more straightforwardly, from the fact that any cube in \mathbb{R}^n is Lebesgue measurable. However, Lemma 3.31 allows for the following, more general statement, which in turn allows for algebras \mathfrak{X} more complicated than the cubical one.

Corollary 3.33. *Let \mathfrak{X} be any algebra on \mathbb{R}^n . Then $\text{PerM}_{\mathfrak{X}}(\mathbb{R}^n)$ is dim-measurable with respect to the Lebesgue σ -algebra.*

Proof. This follows from the fact that any upset in \mathbb{R}^n is Lebesgue measurable, together with Lemma 3.31. See [GG06, Thm. 4.4]² for the case of $[0, 1]^n$. The proof there easily generalizes to \mathbb{R}^n . \square

Example 3.34. By Example 3.28, we obtain L^p -amplitudes for $\text{PerM}_{\mathfrak{X}}(\mathbb{R}^n)$ with respect to any content (in particular the Lebesgue measure) on the Lebesgue algebra on \mathbb{R}^n , whenever \mathfrak{X} is such that $\text{PerM}_{\mathfrak{X}}(\mathbb{R}^n)$ defines an abelian category (in particular by Theorem 2.18 in the connective case). In what follows, we are primarily interested in the case $\mathcal{A} = \text{Vect}$, $\alpha = \dim$, and $\mathfrak{X} = \mathfrak{C}$, i.e. in the functor category of cubically encoded persistence modules. Unless otherwise specified, we always consider the p -Hilbert amplitude with respect to the Lebesgue measure on \mathbb{R}^n . \triangleleft

While in the previous example we would not need the generality of α -measurability, we presented the constructions in this generality to indicate that several other target categories and amplitudes are possible.

Remark 3.35. *The category $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n \times \mathbb{R}^m)$ is equivalently given by the category of cubically encoded objects in the functor category $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)^{\mathbb{R}^m}$. This allows one to start with an amplitude on $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ and turn it into amplitude on $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n \times \mathbb{R}^m)$ by taking the induced L^p -amplitude with respect to some measure on \mathbb{R}^m . In other words, one obtains an amplitude for cubically encoded $\mathbb{R}^n \times \mathbb{R}^m$ modules by using an amplitude of n -parameter persistence modules and integrating on \mathbb{R}^m . This approach seems most promising in the case of $n = 1$, where there is a rich reserve of computable amplitudes available.*

²The paper [GG06] also provides an example of an upper set in $[0, 1]^2$ which is not Borel measurable, so the conditions do not hold if the Lebesgue measure is replaced by the Borel measure.

Example 3.36. Consider the abelian category Ch , with the max degree amplitude δ of Example 3.6. Then the category of tame parametrised chain complexes $\text{tame}([0, \infty), \text{Ch})$ is δ -measurable with respect to the Borel algebra. Indeed, from the definition of a tame chain complex [CGL21], we have that every X in $\text{tame}([0, \infty), \text{Ch})$ admits a cubical finite encoding. Thus, we can define a family of amplitudes $\| - \|_p^\delta$, for $p \in [1, \infty)$, on $\text{tame}([0, \infty), \text{Ch})$. Note that, in general, these L^p -amplitudes are not additive. \triangleleft

3.3 Additive amplitudes

In the previous section, we described how to construct an amplitude for cubically (or more generally \mathfrak{X} -)encoded persistence modules by taking the L^p -norm of the Hilbert function with respect to a measure (more generally a content) on the underlying poset \mathcal{Q} . In the case where $p = 1$, this amplitude is additive. As we will see next, even if additivity may seem a useful property, it is a rather restrictive condition. In fact, for subcategories of finitely encoded persistence modules, any additive amplitude can be obtained by integrating the Hilbert function with respect to a content on the indexing poset. This may be understood as a Riesz-(Markov)-Representation type theorem for additive amplitudes (see [Rud87, Thm. 2.14] for a reference of the latter). In particular, additive amplitudes do not distinguish between persistence modules that have the same Hilbert function but very different structure morphisms.

We begin with the case where \mathcal{Q} is a finite poset, and then derive the general case from this. Here, we can work with the discrete σ -algebra and the notions of content and measure agree. We want to construct a map that associates to any measure on \mathcal{Q} an additive amplitude on the category $\text{Vect}^{\mathcal{Q}}$ of persistence modules over \mathcal{Q} . Let $\text{Meas}(\mathcal{Q})$ be the set of measures μ over the discrete σ -algebra $\Sigma_{\mathcal{Q}}$. By Definition 3.22, we have a well-defined map:

$$\begin{aligned} \psi: \text{Meas}(\mathcal{Q}) &\longrightarrow \text{AdAmp}(\text{Vect}^{\mathcal{Q}}) \\ \mu &\mapsto \int_{\mathcal{Q}} \dim(-) d\mu \end{aligned} \tag{5}$$

where $\text{AdAmp}(\text{Vect}^{\mathcal{Q}})$ is the set of additive amplitudes on $\text{Vect}^{\mathcal{Q}}$. Even if, a priori, $\text{AdAmp}(\text{Vect}^{\mathcal{Q}})$ could be a class, it will turn out to be of set size.

Proposition 3.37. *The map ψ in Eq. (5) is a bijection.*

Namely, if α is an additive amplitude, then there exists a unique measure μ on the discrete σ -algebra of \mathcal{Q} such that $\alpha(M) = \int_{\mathcal{Q}} \dim(M) d\mu$.

Proof. To prove injectivity, assume that we have $\int_{\mathcal{Q}} \dim(-) d\mu = \int_{\mathcal{Q}} \dim(-) d\mu'$, for two measures μ and μ' over the discrete σ -algebra of \mathcal{Q} . For each $S \subseteq \mathcal{Q}$ consider the persistence module N_S that has \mathbb{F} on S and 0 otherwise, and such that the (non-identity) structure morphisms are all zero. By assumption, we get $\mu(S) = \mu'(S)$ for each subset S , and therefore the two measures are equal.

To prove surjectivity, let $\alpha: \text{ob Vect}^{\mathcal{Q}} \rightarrow [0, \infty]$ be an additive amplitude. We define a measure μ as follows: for any $S \subseteq \mathcal{Q}$ we let $\mu(S) = \alpha(N_S)$. We then have that $\mu(\emptyset) = 0$. Given two disjoint subsets S, T of \mathcal{Q} consider the s.e.s.

$$0 \longrightarrow N_S \longrightarrow N_{S \cup T} \longrightarrow N_T \longrightarrow 0$$

where the first morphism is the inclusion and the second the projection. Since α is additive, we obtain that $\mu(S \cup T) = \mu(S) + \mu(T)$. Therefore for any persistence module M of the form $\bigoplus_{i=1}^n N_{S_i}$, where $S_i \subseteq \mathcal{Q}$, we have that $\alpha(M) = \sum_{i=1}^n \mu(S_i) = \int_{\mathcal{Q}} \dim(M) d\mu$.

Next, we show that given any persistence module $M \in \text{ob Vect}^{\mathcal{Q}}$, if we let \hat{M} be the persistence module obtained from it by replacing all the (nontrivial) structure morphisms with zero morphisms, we have that $\alpha(M) = \alpha(\hat{M})$. As every such module with trivial structure morphisms is of the shape $M \cong \bigoplus_{q \in \mathcal{Q}} \bigoplus_{i=1}^{\dim(M_q)} N_{\{u\}}$, this concludes the proof. With this goal, we say that M is U -supported, for U an upset in \mathcal{Q} , if $M_q = 0$ for all $q \notin U$. We now prove $\alpha(M) = \alpha(\hat{M})$ for all $M \in \text{Vect}^{\mathcal{Q}}$ by induction on the cardinality k of U , where U is an upset such that M is U -supported. In particular, for $U = \mathcal{Q}$ we obtain the required result. For $k = 0$ the statement is obvious. Now for an upset of cardinality $k + 1$, let $u \in U$ be a minimal element of U and M be an U -supported module. Let N be the module given by M_q for $q \in U \setminus \{u\}$ and 0 otherwise, with the structure morphisms inherited by that of M . Since u is minimal, $U \setminus \{u\}$ is an upset and thus there is a natural monomorphism $N \hookrightarrow M$. But now the cokernel of this morphism is a module of the form $\bigoplus_{i=1}^{n_u} N_{\{u\}}$ and since the induction hypothesis holds for N , by the additivity of α the claim holds for M . \square

Proposition 3.37 generalises to $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ for any connective algebra \mathfrak{X} and in particular to the category of cubically encoded persistence modules $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$. However, one needs to be careful, because what arises from an additive amplitude is generally not a measure, but rather a content. This is not surprising insofar, as we made no requirements as to how countable colimits should behave with respect to an amplitude. Nevertheless, the idea is the same: all the information given by an additive amplitude on $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ can be retrieved by integrating the Hilbert function.

Now let \mathcal{Q} be an arbitrary poset and \mathfrak{X} be a connective algebra (or more generally any algebra such that $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ is an abelian category). Further, let $\text{Con}(\mathcal{Q}, \mathfrak{X}_{\text{Up}})$ be the set of contents ν over the algebra \mathfrak{X}_{Up} .

Theorem 3.38. *The following map is a bijection:*

$$\begin{aligned} \varphi: \text{Con}(\mathcal{Q}, \mathfrak{X}_{\text{Up}}) &\longrightarrow \text{AdAmp}(\text{PerM}_{\mathfrak{X}}(\mathcal{Q})) \\ \nu &\mapsto \int_{\mathcal{Q}} \dim(-) d\nu . \end{aligned} \tag{6}$$

Namely, any additive amplitude on $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ is uniquely given by integrating the Hilbert functions with respect to a content.

Proof. Since this proof is quite technical, we display it in Appendix B.2. \square

Note that, in the case of $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$, this leads to an even stronger result. In fact, one only needs to specify an additive function on the cubes, not on all finite unions of cubes.

Corollary 3.39. *Let \mathfrak{C} be the set (not the algebra) of cubes in \mathbb{R}^n . Every additive amplitude on $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ is uniquely given through Theorem 3.38 by extending a function $\nu: \mathfrak{C} \rightarrow [0, \infty]$ which fulfills the following two properties to a content.*

1. $\nu(\emptyset) = 0$;
2. For any partition $C = \bigsqcup_{i=1, \dots, n} C_i$ of a cube into disjoint smaller cubes we have

$$\nu \left(\bigsqcup_{i=1, \dots, n} C_i \right) = \sum_{i=1, \dots, n} \nu(C_i).$$

Proof. This follows from the fact that any such function ν extends uniquely to a content on the algebra of cubes \mathfrak{C} , and by Theorem 3.38. \square

Example 3.40. As we mentioned in Example 3.8, the 1-norm ρ_1 of [ST20] is an additive amplitude. Thus, by Theorem 3.38, ρ_1 can be realised by integrating the Hilbert function with respect to some measure (or content). For interval modules of the form $\mathbb{I}\langle b, d \rangle$, with $b, d \in [-\infty, \infty]$, where the angle indicates the interval can either be open or closed, we have $\rho_1(\mathbb{I}\langle b, d \rangle) = d - b$. Hence, the corresponding content has to agree with the Lebesgue measure for all intervals. This already uniquely determines the content as being given by the Lebesgue measure on the algebra of cubes, as the latter algebra is given by finite unions of intervals. Then, using the results of Section 4.2, the 1-Wasserstein distance between two 1-parameter persistence modules M and N can be obtained by minimizing the following integrals

$$\int \dim(\ker \varphi \oplus \operatorname{coker} \varphi) d\lambda + \int \dim(\ker \psi \oplus \operatorname{coker} \psi) d\lambda$$

with respect to the span $M \xleftarrow{\varphi} C \xrightarrow{\psi} N$. Thus, we find, not too surprisingly, a direct link between the 1-Wasserstein distance and the Hilbert function. Moreover, it is now possible to compute, or at least approximate, the 1-Wasserstein distance using techniques from variational methods and numerical analysis. \triangleleft

3.4 The c_p -rank in the amplitude framework

In Section 3.1, we have seen that while many important real-valued invariants from TDA fulfill the axioms of an amplitude, not all invariants are amplitudes. Moreover, we remarked that there exist non-amplitude invariants that can be described using an amplitude. In this section, we discuss one such invariant: the c_p -rank (see Definition 3.20), showing how it is given by the composition of two functors with an amplitude. As a consequence, this invariant can also be studied using the amplitude framework, for instance when it comes to its stability, as we discuss in Section 5. We first generalise the c_p -rank to the continuous (i.e., the \mathbb{R}^n -indexed) setting, which will allow us to investigate its stability in Section 5.6. To do so, we discuss two constructions — the continuous 0-th local cohomology and the quotient restriction — which will be of independent interest in Section 5. We suggest to the reader that is especially interested in continuity and stability results, but not particularly in the c_p -rank invariant, to read Definition 3.45 and Definition 3.47, and otherwise skip this section.

Remark 3.41. *Throughout this section, we will constantly make use of the fact that persistence modules over $\mathcal{Q} = \mathbb{Z}^n, \mathbb{R}^n$, or more generally any abelian partially ordered group, admit a category theoretical, as well as a commutative algebra perspective. They can be equivalently thought of as functors in $\operatorname{Vect}^{\mathcal{Q}}$ as well as as \mathcal{Q} graded modules over the monomial ring $\mathbb{F}[\mathcal{Q}_{\geq 0}]$. Using this perspective, we will frequently denote the application of $M(u \leq v)$ to an element $m \in M_u$ in the form*

$$x^{u-v} m := M(u \leq v) m,$$

for $M \in \operatorname{Vect}^{\mathcal{Q}}$, $u \leq v \in \mathcal{Q}$.

We begin by interpreting the (discrete) c_p -rank as a composition of two functors and an amplitude. Let \mathfrak{p} be a homogeneous prime ideal of $\mathbb{F}[x_1, \dots, x_n]$. Under a change of coordinates, we may assume that \mathfrak{p} is of the shape $\langle x_1, \dots, x_t \rangle$. Now, given a persistence module M over \mathbb{Z}^n , we can express its 0-th local cohomology with respect to \mathfrak{p} as

$$H_{\mathfrak{p}}^0(M) = \ker \left(M \rightarrow \prod_{0 \leq i \leq r} M_{x_i} \right).$$

Here, M_{x_i} is the persistence module obtained by localizing x_i . An immediate application of [Mil20a, Thm. 5.2] shows that this construction indeed preserves cubical encodings and hence

descends to a functor $\text{PerM}_{\mathcal{C}}(\mathbb{Z}^n)$

$$H_{\mathfrak{p}}^0: \text{PerM}_{\mathcal{C}}(\mathbb{Z}^n) \rightarrow \text{PerM}_{\mathcal{C}}(\mathbb{Z}^n).$$

Next, consider the functor

$$(-)/\mathfrak{J}_{\mathfrak{p}}: \text{PerM}_{\mathcal{C}}(\mathbb{Z}^n) \rightarrow \text{PerM}_{\mathcal{C}}(\mathbb{Z}^r)$$

obtained by modding out by the ideal $\mathfrak{J}_{\mathfrak{p}} = \langle x_{t+1} - 1, \dots, x_n - 1 \rangle$. Finally, equip \mathbb{Z}^r with the counting measure. We can now give the following definition:

Definition 3.42. The $c_{\mathfrak{p}}$ -rank for cubically encoded \mathbb{Z}^n -modules is defined as the composition:

$$\text{ob}(\text{PerM}_{\mathcal{C}}(\mathbb{Z}^n)) \xrightarrow{H_{\mathfrak{p}}^0} \text{ob}(\text{PerM}_{\mathcal{C}}(\mathbb{Z}^n)) \xrightarrow{(-)/\mathfrak{J}_{\mathfrak{p}}} \text{ob}(\text{PerM}_{\mathcal{C}}(\mathbb{Z}^r)) \xrightarrow{\|\cdot\|_1^{\text{dim}}} [0, \infty].$$

In other words, the $c_{\mathfrak{p}}$ -rank is given by the 1-Hilbert amplitude of $(H_{\mathfrak{p}}^0(M))/\mathfrak{J}_{\mathfrak{p}}$, for M in $\text{ob} \text{PerM}_{\mathcal{C}}(\mathbb{Z}^n)$.

A priori, the definition of $c_{\mathfrak{p}}$ -rank given in Definition 3.42 might conflict with the one for finitely presented persistent modules given in Definition 3.20. However, as the following proposition shows, these definitions are equivalent. At the same time, this latter definition suggests a natural generalisation of the $c_{\mathfrak{p}}$ -rank to \mathbb{R}^n modules, which we give in Definition 3.49.

Proposition 3.43. *Let M be a cubically encoded persistence module over \mathbb{Z}^n and \mathfrak{p} a homogeneous prime of $\mathbb{F}[x_1, \dots, x_n]$. Then the two definitions of $c_{\mathfrak{p}}$ -rank of M given in Definition 3.20 and Definition 3.42 coincide.*

Proof. The proof is provided in Appendix B.3. \square

To investigate whether the $c_{\mathfrak{p}}$ -rank is continuous in an appropriate sense, one needs a notion of $c_{\mathfrak{p}}$ -rank for persistence modules over \mathbb{R}^n . This also requires a corresponding definition of 0-th local cohomology, which is of interest independently of its application to the $c_{\mathfrak{p}}$ -rank (see Section 5.5). The role of homogeneous prime ideals for \mathbb{Z}^n -modules is taken by the faces of the positive cone for \mathbb{R}^n -modules:

Definition 3.44. A **face** of the positive cone \mathbb{R}^n is a set of the form

$$\tau = \left\{ \sum_i \alpha_i e_{j_i} \mid \alpha_i \geq 0 \right\},$$

where $\{i_0, \dots, i_k\}$ is a subset of $\{1, \dots, n\}$ and e_j denotes the j -th standard unit vector in \mathbb{R}^n . Moreover, τ^c denotes the face **orthogonal** to τ , i.e. it is the cone spanned by the standard basis vectors in \mathbb{R}^n which are not in τ .

Definition 3.45. Let τ be a face of the positive cone in \mathbb{R}^n and $M \in \text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$. We define the following persistence module

$$H_{\tau}^0(M) := \ker(M \rightarrow \prod_{0 \neq \rho \subset \tau} M_{\rho}),$$

where the product is taken over the nontrivial faces of τ , and M_{τ} denotes the localisation of M at τ as in [Mil20a, Def. 2.17]. Note that localization in this sense sends the indicator modules of cubes into indicator modules of cubes and is exact [Mil20a, Lem. 2.24]. Hence, as an application of [Mil20b, Thm. 6.12], localization preserves cubical encodings. In particular, since $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$ is abelian, and the product is finite H_{τ}^0 is again cubically encoded. This construction therefore induces a functor

$$H_{\tau}^0: \text{PerM}_{\mathcal{C}}(\mathbb{R}^n) \rightarrow \text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$$

which we call the **0-th local cohomology with respect to τ** .

Remark 3.46. Note that, in the definition of H_τ^0 in Definition 3.45, we might have equivalently just taken the product over the 1-dimensional faces of τ . Indeed, let $\{v_0, \dots, v_k\} \subset \tau_i$ be any set of generators for a face τ_i of τ . Then $m \in M$ being mapped to 0 in $M_{\mathbb{R}_{\geq 0} v_j}$, for all $0 \leq j \leq k$, already implies that m is mapped to 0 in M_τ .

The role of modding out the ideal $\mathfrak{I}_\mathfrak{p}$ in Definition 3.42 is taken by the quotient restriction functor from [Mil20a, Def. 2.22].

Definition 3.47. Let τ be a face of the positive cone in \mathbb{R}^n of dimension t . Given a persistence module $M \in \text{Vect}^{\mathbb{R}^n}$, the **quotient restriction** M/τ of M along τ is defined as the $\mathbb{R}^n/\mathbb{R}\tau$ -module given by

$$(M/\tau)_{v+\mathbb{R}\tau} := \varinjlim_{u \in v+\tau} M_u$$

with the induced structure maps.

In other words, M can be thought as the quotient by equivalence relation $m \sim x^v m$, for $m \in M$, $v \in \tau$, of an \mathbb{R}^n -graded module.

Lemma 3.48. The quotient restriction preserves cubical encodings. Further, if we identify $\mathbb{R}^n/\mathbb{R}\tau$ with $\mathbb{R}\tau^c$, which in turns is naturally identified with \mathbb{R}^{n-t} via the basis of unit vectors for \mathbb{R}^n , this construction induces a functor

$$(-)/\tau: \text{PerM}_{\mathcal{C}}(\mathbb{R}^n) \rightarrow \text{PerM}_{\mathcal{C}}(\mathbb{R}^{n-t})$$

which is exact.

Proof. Quotient restriction is equivalently defined as first localizing at τ and then restricting along $\mathbb{R}\tau^c$ (see [Mil20b, Rem. 2.23]). As both the localisation as well as the restriction to a subposet are exact, the functor $\text{Vect}^{\mathbb{R}^n} \rightarrow \text{Vect}^{\mathbb{R}^n}$ defined in this fashion is exact. Furthermore, it sends interval modules of cubes to interval modules of cubes. More specifically, if $\mathbb{F}[C]$ is the interval module of an upset-cube C , then $\mathbb{F}[C]_\tau = \mathbb{F}[C + \mathbb{R}\tau]$. The restriction to $\mathbb{R}\tau$ acts as $\mathbb{F}[C + \mathbb{R}\tau] \mapsto \mathbb{F}[(C + \mathbb{R}\tau) \cap \mathbb{R}\tau^c]$. The latter is again the indicator module of a cube. By an application of [Mil20b, Thm. 6.12] and using exactness, we thus obtain that quotient restriction indeed sends cubically encoded modules into cubically encoded modules. \square

We now have all it is needed to define the continuous c_τ -rank.

Definition 3.49. Let τ be a face of the positive cone in \mathbb{R}^n of dimension t . The **c_τ -rank** $c_\tau: \text{PerM}_{\mathcal{C}}(\mathbb{R}^n) \rightarrow [0, \infty]$ is defined as the composition

$$\text{ob}(\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)) \xrightarrow{H_\tau^0} \text{ob}(\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)) \xrightarrow{(-)/\tau^c} \text{ob}(\text{PerM}_{\mathcal{C}}(\mathbb{R}^{n-t})) \xrightarrow{\|\cdot\|_1^{\dim}} [0, \infty].$$

This definition is compatible with the discrete one in the following sense:

Proposition 3.50. Let $f_n: \mathbb{R}^n \rightarrow \mathbb{Z}^n$ the map of posets given by taking the floor map (i.e. the map that sends a real to the largest integer smaller than it) in each component. Let \mathfrak{p} be a homogeneous prime of $\mathbb{Z}[x_1, \dots, x_n]$ with t generators and τ the corresponding face of the positive cone of dimension t . Then, for each $M \in \text{PerM}_{\mathcal{C}}(\mathbb{Z}^n)$, we have

$$c_{\mathfrak{p}}(M) = c_\tau(f_n^* M).$$

Proof. We use the notation introduced before Definition 3.42. A straightforward verification shows that $H_\tau^0 \circ f_n^* \cong f_n^* \circ H_{\mathfrak{p}}^0$ and $(-)/\tau^c \circ f_n^* \cong f_n^* \circ (-)/\mathfrak{J}$. Next, note that the map f_{n-t} is measure preserving with respect to the Lebesgue measure and the counting measure, as the inverse of a point is given by a cube of volume one, with parts of its boundary removed. In particular, $\|f_{n-t}^* N\|_1^{\dim} = \|N\|_1^{\dim}$ for all $N \in \text{PerM}_{\mathcal{C}}(\mathbb{Z}^{n-t})$. Piecing these equalities together gives the result. \square

4 Distances from amplitudes

There are several ways to define a distance from an amplitude. We describe some of them here, with special attention to the ones already suggested in the literature. Throughout this section, where not otherwise specified, we let \mathcal{A} be an abelian category and α an amplitude on \mathcal{A} .

We start by giving a naive definition of distance:

Definition 4.1. The **absolute-value distance** of an amplitude α is defined as

$$d_{|\alpha|}(A, B) := |\alpha(A) - \alpha(B)|$$

for any pair of objects A, B in $\text{ob } \mathcal{A}$.

The fact that this function is indeed a distance follows by the properties of the absolute-value function. If there are at least two non-isomorphic objects A, B , with $\alpha(A) = \alpha(B)$, then $d_{|\alpha|}(A, B) = 0$ and α is a pseudometric. One would have that $d_{|\alpha|}$ would be a metric if α were such that $\alpha(A) \neq \alpha(B)$ for all $A \not\cong B$ in $\text{ob } \mathcal{A}$.

The distance $d_{|\alpha|}$ is straightforward, but it may not be able to capture all the information we want, as it forgets the underlying categorical structure. Nevertheless, such distances are often used in applications, for instance, $d_{|\text{totpers}|}$ is used in one of the first stability results in persistent homology [CSEHM10], while $d_{|\ell\text{-trop}^k|}$ is used in the stability results for tropical coordinates in [Kal19]. We thus believe that one of the main contributions of our work might be to highlight the need for considering distances from amplitudes that are more informative than the ones usually used in the literature.

Example 4.2. Consider as amplitude the p -norm amplitude ρ_p of Example 3.8, for $p \in [1, \infty)$, and the two interval persistence modules $\mathbb{I}[a, a+r)$ and $\mathbb{I}[b, b+r)$, for $r > 0 \in \mathbb{R}$ and $a, b \in \mathbb{R}$ with $a+r < b$. Then $d_{|\rho_p|}(\mathbb{I}[a, a+r), \mathbb{I}[b, b+r)) = 0$. However, we often would like to say that the distance between them is non-zero, since their endpoints differ. \triangleleft

We may then try to define a more distinguishing distance along the lines of the L^p -distances in [Tho19], at least for functor categories.

Example 4.3. Given a poset \mathcal{Q} equipped with a σ -algebra Σ and a measure μ on Σ , we can define a distance on $\mathcal{A}^{\mathcal{Q}}$ using the amplitude α on \mathcal{A} . Let α be such that α_A is measurable for all $A \in \text{ob } \mathcal{A}^{\mathcal{Q}}$. Then, for $1 \leq p < \infty$, we can define the following distance:

$$d_{L^p}(A, B) := \left(\int_{\mathcal{Q}} |\alpha_A - \alpha_B|^p d\mu \right)^{\frac{1}{p}}$$

for any pair of objects A, B in $\mathcal{A}^{\mathcal{Q}}$. \triangleleft

As in Example 3.28, one could generalise the above distance to the case in which α_{\bullet} is not measurable on the whole $\mathcal{A}^{\mathcal{Q}}$, but only on an abelian subcategory \mathcal{C} of it. For example, when considering $\mathcal{C} = \text{PerM}_{\mathcal{E}}(\mathbb{R}^n)$ with the Hilbert function, we retrieve the definition of the L^p -distance given in [LW15, HOST19]. However, since this distance does not consider the structure morphisms of a persistent object, it also fails to distinguish objects that we would like to tell apart. For example, take $\alpha = \dim$ on Vect and $\mathcal{Q} = [0, \infty)$ with the Lebesgue measure. Let M and N be the two 1-parameter persistence modules given by $\mathbb{I}[0, 1) \oplus \mathbb{I}[1, 2)$ and $\mathbb{I}[0, 2)$, respectively. Then $d_{L^p}(M, N) = 0$ for all $p \in [1, \infty)$. Note that a similar distance is also discussed in [Tho19], where, however, the main focus of the author is the L^p -distance given by the (multi-)rank function, which we do not consider as it is not an amplitude (see Example 3.18).

To remedy this shortcoming of the previous example, we provide other distances, using morphisms to compare objects.

Definition 4.4. Let A, B be two object in \mathcal{A} . A **zigzag** between A and B is a sequence γ of morphisms between A and B : $A \xleftarrow{\gamma_1} C_1 \xrightarrow{\gamma_2} \cdots \xleftarrow{\gamma_{n-1}} C_n \xrightarrow{\gamma_n} B$. The **cost** of a zigzag γ (with respect to the amplitude α) is:

$$c_\alpha(\gamma) := \sum_{i=1}^n \alpha(\ker \gamma_i) + \alpha(\operatorname{coker} \gamma_i).$$

For any pair of objects A, B in \mathcal{A} , the **path metric** d_α between A and B is given by

$$d_\alpha(A, B) := \inf_{\gamma} c_\alpha(\gamma),$$

with the convention $\inf \emptyset = \infty$.

Remark 4.5. We use the name path metric, which was used in a preliminary version of [BSS20]. In the definition of cost in the preliminary version, the authors of [BSS20] restrict the domain of the cost function to a subset S of the objects in \mathcal{A} . Since an amplitude is defined on all the objects, we avoided the mention of S in the definition.

The path metric is an extended pseudometric on $\operatorname{ob} \mathcal{A}$. Indeed, a zigzag of isomorphisms has cost zero, thus all isomorphic objects have distance zero from each other. Symmetry follows directly from the definition of a zigzag, and the triangle inequality follows by juxtaposing two zigzags. Moreover, if α is finite, then the path metric is a pseudometric on $\operatorname{ob} \mathcal{A}$. For the sake of readability, in the rest of the paper we often drop the specification and simply say metric.

Example 4.6. Recall the interval modules $\mathbb{I}[a, a+r)$ and $\mathbb{I}[b, b+r)$ of Example 4.2. We have $d_{\rho_p}(\mathbb{I}[a, a+r), \mathbb{I}[b, b+r)) = 2r$ and $d_{m_{\rho_p}}(\mathbb{I}[a, a+r), \mathbb{I}[b, b+r)) = r$. \triangleleft

The absolute-value amplitude distance can be used to obtain a lower bound for the path metric.

Lemma 4.7. For any \mathcal{A} be an abelian category with amplitude α . Then, for any $A, B \in \operatorname{ob} \mathcal{A}$, the following inequality holds:

$$d_{|\alpha|}(A, B) \leq d_\alpha(A, B).$$

In other words, α is 1-Lipschitz with respect to d_α .

Proof. By the triangle inequality and by definition of $d_{|\alpha|}$ and of d_α , it suffices to show that whenever there is a morphism $f: A \rightarrow B$ with $c_\alpha(f) \leq \varepsilon$, then

$$|\alpha(A) - \alpha(B)| \leq \varepsilon.$$

Again by the triangle inequality and epi-mono factorization, we may restrict to the case where f is either an epi- or monomorphisms. If f is a mono, we then have a short exact sequence $0 \rightarrow A \rightarrow B \rightarrow \operatorname{coker} f \rightarrow 0$ with $\alpha(\operatorname{coker}(f)) \leq \varepsilon$. Hence, by monotonicity and subadditivity of amplitudes follows that,

$$\alpha(A) \leq \alpha(B) \leq \alpha(A) + \varepsilon,$$

showing the required inequality. The case of f an epimorphism follows dually. \square

Remark 4.8. Consider the case of $\alpha = \|\cdot\|_{\dim}^p$, the p -Hilbert amplitude with respect to the Lebesgue measure (see Example 3.30). Both the absolute-value distance as well as the distance of Example 4.3 are significantly less discriminative than the path metric with respect to α . Indeed, of these three extended pseudo metrics only the path metric takes the transition morphism into account. Two persistence modules M and N (with measurable Hilbert function) fulfill $d_{|\alpha|} = 0$

(or $d_{L^p}(M, N) = 0$) if and only if their Hilbert functions agree almost everywhere. On the other hand, as shown in [Waa21], when \mathfrak{X} is a connective algebra generated by closed upsets, then the path metric associated to α defined on $\text{PerM}_{\mathfrak{X}}(\mathbb{R}^n)$ fulfills $d_{\alpha}(M, N) = 0 \iff M \cong N$. More generally, this is the case on this category for any amplitude β on $\text{PerM}_{\mathfrak{X}}(\mathbb{R}^n)$ fulfilling $\beta(M) = 0 \implies M \cong 0$.

Following the insight of [SCL⁺17], we aim to give an alternative definition for the path metric. The proofs of this alternative definition have been formulated in [SCL⁺17] for the specific setting of tame multi-parameter persistence modules (see Section 2) and additive cost, but they hold more generally for any abelian category. Therefore, we will provide precise references without displaying all the details.

Our goal is to obtain a metric from the shortest possible zigzag between two objects that allows for different costs. Since we cannot hope to extract a metric using a zigzag of length 1, as the symmetry axiom will not be satisfied, we investigate zigzags of length 2, i.e. (co)spans, which will provide the wanted alternative definition.

We recall that a subadditive real function on $S \subset \mathbb{R}^n$ is a function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ such that $f(v + w) \leq f(v) + f(w)$ for all $v, w \in S$.

Definition 4.9. A **cost function** is a function $f: [0, \infty]^4 \rightarrow [0, \infty]$ that is monotone, subadditive and such that $f(0) = 0$ and $f(x_1, x_2, x_3, x_4) = f(x_3, x_2, x_1, x_4) = f(x_1, x_4, x_3, x_2)$. Given a (co)span $\mathbf{c}: A \xleftarrow{\varphi} C \xrightarrow{\psi} B$ in \mathcal{A} and a cost function f , the **f -cost of \mathbf{c}** is

$$f_{\alpha}(\mathbf{c}) := f(\alpha(\ker \varphi), \alpha(\text{coker } \varphi), \alpha(\ker \psi), \alpha(\text{coker } \psi)).$$

Remark 4.10. For every cost function f , we have that, given a cospan $A \xleftarrow{\varphi} C \xrightarrow{\psi} B$, there exists a span $A \xrightarrow{\psi'} C \xleftarrow{\varphi'} B$ with smaller or equal f -cost, and vice-versa. The proof is in [SCL⁺17, Cor. 8.3] for $f = \|\cdot\|_1$, but it is immediately generalised in our setting.

We are now ready to prove that the f -cost induces a metric:

Proposition 4.11. Let \mathcal{A} be an abelian category and α an amplitude over \mathcal{A} and f a cost function. Then, for any pair of objects A and B in $\text{ob } \mathcal{A}$, the following map define a (pseudo, extended) metric:

$$d_{f_{\alpha}}(A, B) = \inf \left\{ f_{\alpha}(\mathbf{c}) \mid \mathbf{c}: A \xleftarrow{\varphi} C \xrightarrow{\psi} B \right\},$$

with the convention that $\inf \emptyset = \infty$.

Proof. It is clear that $d(A, A) = 0$ and that $d(A, B) = d(B, A)$ for every A, B in $\text{ob } \mathcal{A}$. For the triangle inequality, consider two cospans $\mathbf{c}_1: A \xleftarrow{\varphi_1} H_1 \xrightarrow{\psi_1} B$ and $\mathbf{c}_2: B \xleftarrow{\varphi_2} H_2 \xrightarrow{\psi_2} C$. By Remark 4.10, we have $\mathbf{c}_3: A \xleftarrow{\varphi_1} H_1 \xleftarrow{\varphi_3} H_3 \xrightarrow{\psi_3} H_2 \xrightarrow{\psi_2} C$. Thus, it is left to show that $f_{\alpha}(\mathbf{c}_3) \leq f_{\alpha}(\mathbf{c}_1) + f_{\alpha}(\mathbf{c}_2)$. This follows by the argument presented in [SCL⁺17, Prop. 8.2] and the properties of f . \square

In particular, we obtain the wanted characterization of the path metric:

Corollary 4.12. If f is given by the 1-norm, then $d_{\alpha} = d_{f_{\alpha}}$.

Since the path metric is the most used in our work, we keep the notation d_{α} for denoting it.

This result is allowed by the axioms of amplitudes, and it is not true in the more general case of weight functions [BSS20]. Indeed, one of the motivation for the introduction of the axioms of noise systems [SCL⁺17] was to allow for studying distances using only (co)spans instead of arbitrary zigzags.

Remark 4.13. *Proposition 4.11 strengthens the arguments why it is better to restrict to zigzag of length 2. Indeed, in this case use more general f -costs. For example, we can choose as f any norm, and, by the equivalence of norms in \mathbb{R}^n , we can deduce equivalences of f -path metrics (see Corollary 4.22).*

Remark 4.14. *As shown in Appendix A, every amplitude α is related to a noise system $\mathcal{S}(\alpha)$. As explained in [SCL⁺17], every noise system defines a pseudometric, via ε -closeness (see Definition 4.9). We have that the pseudometric given by α and $\mathcal{S}(\alpha)$ coincide (see Appendix A).*

Corollary 4.12 allows us to recover an amplitude from its path metric.

Lemma 4.15. *Let \mathcal{A} be an abelian category and α an amplitude on \mathcal{A} . Then*

$$\alpha(A) = d_\alpha(A, 0)$$

for each $A \in \mathcal{A}$.

Proof. We observe that $\alpha(A) = c_\alpha(A \rightarrow 0)$, and thus $d_\alpha(A, 0) \leq \alpha(A)$.

For the converse inequality consider a cospan $A \xrightarrow{\varphi} B \leftarrow 0$. It suffices to show that $c_\alpha(A \rightarrow 0) \leq c_\alpha(\varphi) + c_\alpha(0 \rightarrow B)$ to obtain the missing inequality by passing to the infimum. Since $0 = 0 \circ \varphi$, we have

$$c_\alpha(A \rightarrow 0) \leq c_\alpha(\varphi) + c_\alpha(B \rightarrow 0) = c_\alpha(\varphi) + c_\alpha(0 \rightarrow B). \quad \square$$

Note that the above it not true in general for the f -path metric. Indeed, consider for example $f = \max$, and the rank amplitude over Vect , i.e. \dim (see Example 3.3). Let V in Vect with $\dim(V) = 2$. A direct computation shows that $d_{m_{\dim}}(V, 0) = 1$.

Remark 4.16. *In the literature, the path metric appears more often than the f -path metric given by the maximum. The choice on which to use is usually motivated by the applications at hand. However, this choice is less arbitrary than what may seem at first, since, as a direct computation shows, the metrics are equivalent. Namely, let A, B be two objects in \mathcal{A} . Then*

$$d_{m_\alpha}(A, B) \leq d_\alpha(A, B) \leq 2d_{m_\alpha}(A, B).$$

Moreover, by Example 4.6, the second inequality is sharp.

4.1 The interleaving distance as a distance from an amplitude

In this section, we study the relation of the path metrics with the interleaving distance [Les15]. In particular, we are interested in determining whether there exists an amplitude α that gives rise to the interleaving distance. By Remark 4.16, we can restrict to the path metric. From Lemma 4.15, we have that there is only one possible candidate for the amplitude giving the interleaving distance on \mathbb{R}^n indexed persistence modules, which is identified by the distance of objects to 0. In particular, the only possible amplitude is the shift amplitude along a vector v in the positive cone. For $M \in \text{Vect}^{\mathbb{R}^n}$, we denote by $M^{+\varepsilon}$ the shifted persistence module given by pulling back along $\mathbb{R}^n \rightarrow \mathbb{R}^n$, $x \mapsto x + \varepsilon v$.

Let \mathcal{A} be an abelian category and $\varsigma: \mathcal{A}^{\mathbb{R}^n} \rightarrow [0, \infty]$ the shift amplitude along a vector v (see Example 3.9).

Definition 4.17. An ε -interleaving between A and B in $\mathcal{A}^{\mathbb{R}^n}$ is given by morphisms $\varphi: A \rightarrow B^{+\varepsilon}$ and $\psi: B \rightarrow A^{+\varepsilon}$ in $\mathcal{A}^{\mathbb{R}^n}$, such that $\varphi^t \circ \psi^{t+\varepsilon v} = A^{t \leq t+2\varepsilon v}$ and $\psi^t \circ \varphi^{t+\varepsilon v} = B^{t \leq t+2\varepsilon v}$ for all $t \in [0, \infty)$. The **interleaving distance** between two objects A and B in $\mathcal{A}^{\mathbb{R}^n}$ is given by

$$d_I(A, B) := \inf\{\varepsilon > 0 \mid \exists \text{ an } \varepsilon\text{-interleaving between } A \text{ and } B\}$$

with the convention $\inf \emptyset = \infty$.

We have that the interleaving distance on $\mathcal{A}^{\mathbb{R}^n}$ is equivalent to d_ζ .

Proposition 4.18. *For any $A, B \in \mathcal{A}^{\mathbb{R}^n}$, we have:*

$$d_I(A, B) \leq d_\zeta(A, B) \leq 6d_I(A, B).$$

Proof. For $\mathcal{A} = \text{Vect}$ and tame functors, the proof is given in [GC17, Prop. 12.2]. The argument given there carries through for any abelian category. \square

Remark 4.19. *The previous statement, being formulated for the whole functor category, is quite general. However, the result still holds if one restricts to any full (closed under shifts) abelian subcategory of $\mathcal{A}^{\mathbb{R}^n}$, such as the full subcategory of cubically encoded modules.*

4.2 Wasserstein distance of p -norm amplitudes

In this subsection, we show that for 1-parameter persistence modules the path metric d_{ρ_p} and the Wasserstein distance are equivalent. We begin by recalling the standard definition of p -Wasserstein distance.

Definition 4.20. Let M, N be objects in $\text{PerM}_{\mathcal{C}}(\mathbb{R})$ and $p \in [1, \infty)$. The **p -Wasserstein distance** between M and N is given by

$$W_p(\text{Dgm}(M), \text{Dgm}(N)) = \inf_L \left(\sum_{x \in \text{Dgm}(M)} \|x - L(x)\|_p \right)^{\frac{1}{p}}$$

where, possibly adding points on the diagonal, $L: \text{Dgm}(M) \rightarrow \text{Dgm}(N)$ is a bijection.

We now translate in our setting an alternative definition of the Wasserstein distance presented in [ST20], i.e. as a f -path metric with f given by $\| - \|_p$ and $\alpha = \rho_p$.

Definition 4.21. [ST20] Let M and N be two objects in $\text{PerM}(\mathbb{R})$, and $p \in [1, \infty]$. The algebraic p -Wasserstein distance between M and N is given by

$$W_p^{alg}(M, N) = d_{f_\alpha}(M, N) = \inf_{(C, \varphi, \psi)} \|(\rho_p(\ker \varphi), \rho_p(\text{coker } \varphi), \rho_p(\ker \psi), \rho_p(\text{coker } \psi))\|_p$$

where the infimum is taken over the spans $M \xleftarrow{\varphi} C \xrightarrow{\psi} N$, for $C \in \text{ob PerM}_{\mathcal{C}}(\mathbb{R})$ and f is induced by the $\| - \|_p$.

In [ST20, Th 8.25 and 8.26], the authors prove that Definition 4.21 is indeed an alternative definition for the Wasserstein distance of persistence diagrams. We use the result here, with slightly stronger hypothesis: we require the persistence modules to be finitely encoded and not pointwise finite dimensional with bounded p -energy. The result then holds with simpler proof since now it relies almost entirely on the theorem about the structure of persistence submodules and quotients of [BL15].

Using the equivalences of norms in \mathbb{R}^4 and the fact that $W_p^{alg} = d_{f_{\rho_p}}$, where f is given by the p -norm on \mathbb{R}^4 , we obtain the following equivalence:

Corollary 4.22. *For $p \in [1, \infty]$ and $M, N \in \text{PerM}_{\mathcal{C}}(\mathbb{R})$ the following equivalence of metrics holds:*

$$W_p^{alg}(M, N) \leq d_{\rho_p}(M, N) \leq 4^{1-\frac{1}{p}} W_p^{alg}(M, N).$$

In particular, for $p = 1$, we obtain $W_1^{alg} = d_{\rho_1}$.

In [BSS20, Th. 5.16], a version of the case $p = 1$ of Corollary 4.22 is proven under slightly different assumptions.

5 Continuity and stability

As explained in [BdSS15], in TDA there are different types of stability that one might consider. The typical TDA workflow goes from data to invariants passing through parametrised (typically, filtered) simplicial complexes and persistent modules, and one can study the stability for each of these passages, or for some composition of them. Here we present a new possibility, naturally suggested by our approach. Namely, we consider functors from the category of cubically persistence modules to itself, possibly changing the source poset and the amplitudes. Then, studying the stability of these functors, we can infer new stability results from old ones.

With this goal in mind, we will define a category of amplitudes (Section 5.1) whose morphisms are what we called amplitude-bounding functors. To obtain continuity and stability results, one can now consider an amplitude-bounding functor

$$F: (\text{PerM}_{\mathcal{C}}(\mathbb{R}^n), \alpha) \rightarrow (\text{PerM}_{\mathcal{C}}(\mathbb{R}^n), \alpha')$$

with the additional property of being Lipschitz with respect to the distances induced by the amplitudes:

$$d_{\alpha'}(F(M), F(N)) \leq K d_{\alpha}(M, N)$$

for all persistence modules M, N and for some $K > 0$. Then any stability result of the form

$$d_{\alpha}(H_k(D), H_k(E)) \leq K' d(D, E)$$

where D, E and d are appropriate input data sets and distances between these for the persistent homology pipeline, such as finite metric spaces together with the Gromow-Hausdorff distance, or Lipschitz functions on a topological space together with the sup-norm, would give us a stability result for α' :

$$d_{\alpha'}(H_k(D), H_k(E)) \leq K d_{\alpha}(H_k(D), H_k(E)) \leq K K' d(D, E).$$

The proof of new stability results usually requires substantial work, and we therefore believe this to be a very powerful application of the change in perspective that our framework gives.

5.1 Amplitude-bounding functors

We now introduce amplitude-bounding functors.

Definition 5.1. Let Amp , the **category of amplitudes**, be the category where:

1. Objects are pairs (\mathcal{A}, α) consisting of an abelian category \mathcal{A} together with an amplitude α on \mathcal{A} .
2. Morphisms $(\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$ are **amplitude-bounding** functors, i.e. additive functors $F: \mathcal{A} \rightarrow \mathcal{A}'$ such that there exists a non-negative real number K so that $\alpha'(FA) \leq K\alpha(A)$ for all objects A . Given such a K , we say that F has **constant K** .

One could also generalise Definition 5.1 by choosing a bound not given by a linear function, but more generally by any continuous function. In other words, one could consider functors $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$ such that $\alpha'(F(A)) \leq f(\alpha(A))$ for a continuous monotonous function $f: [0, \infty] \rightarrow [0, \infty]$. We explore this approach a little in Proposition 5.5, however it is too general for our main scope.

Examples of amplitude-bounding functors used in TDA applications include the identity functor (see also the discussion at the beginning of Section 5.3), as well as functors induced by common transformations of the indexing poset, such as Example 5.2. An example of a non-exact amplitude-bounding functor is given by 0-th local cohomology (see Proposition 5.28).

Example 5.2. (Rescaling of metric spaces) If (X, d_X) is a metric space, and t is any positive real number, one defines the rescaled space $tX := (X, td_X)$ where $td_X: X \times X \rightarrow \mathbb{R}_{\geq 0}: d_X(x, y) \mapsto td_X(x, y)$. As was observed in [GH21, Sec. 5.2], the rescaling of a metric space induces a rescaling of the corresponding persistence modules obtained by computing persistent homology with the Čech or Vietoris–Rips complex. More precisely, given $M \in \text{ob PerM}(\mathbb{R})$ and $t \in (0, \infty)$, consider the functor

$$s_t: \mathbb{R} \rightarrow \mathbb{R}$$

that sends u to u/t . Then the rescaled persistence module tM is the composition $M \circ s_t$. Therefore this induces a functor

$$F_t: \text{PerM}(\mathbb{R}) \rightarrow \text{PerM}(\mathbb{R})$$

sending a persistence module M to its rescaling tM . Since M is 1-parameter and finitely encoded, the barcode of tM is obtained from that of M by rescaling the intervals of the barcode of M by t . Thus, we have an amplitude-bounding functor

$$F_t: (\text{PerM}(\mathbb{R}), \rho_p) \rightarrow (\text{PerM}(\mathbb{R}), \rho_p),$$

where ρ_p is the amplitude from [ST20] (see Example 3.8), with $\rho_p(F_t(M)) = t\rho_p(M)$. Similarly, we can consider the amplitude $|\cdot|$ given by the persistent magnitude (see Example 3.16), and thus

$$F_t: (\text{PerM}(\mathbb{R}), |\cdot|) \rightarrow (\text{PerM}(\mathbb{R}), |\cdot|)$$

is an amplitude-bounding functor with constant $K = e^t$. ◁

5.2 Continuity of amplitude-bounding functors

As mentioned, in many applications the preferred distance is the absolute-value distance $d_{|\alpha|}$. However, we saw that this distance is not discriminative enough. In this section, we provide other arguments in this regard (Section 5.3.1). Moreover, while with the more discriminative distance d_α , as we show next, one can find general properties to infer new stability results using the notion of amplitude-bounding functor, one cannot, in general, derive continuity or stability with respect to the absolute-value distance.

We begin by showing that an additive functor that is K -Lipschitz is amplitude-bounding, with constant K .

Proposition 5.3. *Let $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$ be an additive functor for which there exists $K \geq 0$ such that $d_{\alpha'}(F(A), F(B)) \leq Kd_\alpha(A, B)$ for all $A, B \in \text{ob } \mathcal{A}$. Then $\alpha'(F(A)) \leq K\alpha(A)$ for all $A \in \text{ob } \mathcal{A}$.*

Proof. By Lemma 4.15 we know that $\alpha'(F(A)) = d_{\alpha'}(F(A), 0)$. Since F is additive, we have that $F0 \cong 0$. Thus, we obtain $\alpha'(F(A)) = d_{\alpha'}(F(A), 0) \leq Kd_\alpha(A, 0) = \alpha(A)$. □

Conversely, we are interested in studying under which additional assumptions a functor that is amplitude bounding is continuous in an appropriate sense.

Proposition 5.4. *Let $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$ be an exact functor that satisfies $\alpha'(F(A)) \leq K\alpha(A)$ for all $A \in \text{ob } \mathcal{A}$. Then we have $d_{\alpha'}(F(A), F(B)) \leq Kd_\alpha(A, B)$ for all $A, B \in \text{ob } \mathcal{A}$.*

The proof follows straightforwardly from the definition of amplitude-bounding and exact functors.

Recall the generalisation of the definition of amplitude-bounding functor given by taking a continuous monotonous function $f: [0, \infty] \rightarrow [0, \infty]$ such that $\alpha'(F(A)) \leq f(\alpha(A))$. A direct application of Corollary 4.12 then gives the following:

Proposition 5.5. *Let $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$ be an exact functor that satisfies $\alpha'(F(A)) \leq f(\alpha(A))$ for all $A \in \text{ob } \mathcal{A}$, for a monotonous function $f: [0, \infty] \rightarrow [0, \infty]$. Then we have*

$$d_{\alpha'}(F(A), F(B)) \leq 4f(d_{\alpha}(A, B))$$

for all $A, B \in \text{ob } \mathcal{A}$. In particular, if f is continuous in 0, F induces a continuous function with respect to the topologies induced by d_{α} and $d_{\alpha'}$.

For an amplitude bounding functor to be Lipschitz, it really suffices to consider a notion of exactness up to objects with amplitude 0. We call a sequence $\cdots \rightarrow A_k \rightarrow A_{k+1} \rightarrow \cdots$ in (\mathcal{A}, α) **α -exact** if its homology at each k has amplitude 0. In other words, if the sequence is exact after modding out by the full subcategory of objects with amplitude 0 (see [Sta18, Tag 02MN] to make this rigorous).

Proposition 5.6. *Let $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$ be an additive amplitude-bounding functor with constant K , with the following two properties for any short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ in \mathcal{A} :*

1. *If A has finite amplitude, then the sequence $F(A) \rightarrow F(B) \rightarrow F(C) \rightarrow 0$ is α' -exact;*
2. *If C has finite amplitude, then the sequence $0 \rightarrow F(A) \rightarrow F(B) \rightarrow F(C)$ is α' -exact.*

Then F is K -Lipschitz with respect to the induced amplitude metrics.

Proof. Without loss of generality (by passing to the appropriate quotient categories if necessary), we may assume that the sequences above are actually exact, and not just α -exact. Next, observe that, for the computation of the metric, only the morphisms that have finite cost are relevant. It suffices to show that for any such morphism f we have $c'_{\alpha'}(F(f)) \leq c_{\alpha}(f)$. Similarly as before, we can reduce to the case where f is a monomorphism. Hence, we have a s.e.s. $0 \rightarrow A \xrightarrow{f} B \rightarrow \text{coker } f \rightarrow 0$. By assumption, F sends this s.e.s. into an exact sequence $0 \rightarrow F(A) \xrightarrow{F(f)} F(B) \rightarrow F(\text{coker } f)$. In particular, $\text{coker}(F(f))$ is a subobject of $F(\text{coker } f)$. Hence,

$$c_{\alpha'}(F(f)) = \alpha'(\text{coker}(F(f))) \leq \alpha'(F(\text{coker } f)) \leq K\alpha(\text{coker } f) = c_{\alpha}(f)$$

which finishes the proof. □

Remark 5.7. *Note that the conditions of Proposition 5.6 are not far from requiring exactness since, after quotienting out, they amount in practice to requiring exactness on the full abelian subcategory of objects with finite amplitude. Indeed, if the target category \mathcal{A}' is such that every α' -exact sequence is exact, these conditions imply that the respective functor induces an exact functor $\mathcal{B} \rightarrow \mathcal{A}'$, where \mathcal{B} is the full abelian subcategory of finite amplitude objects in \mathcal{A} . Nevertheless, this (slightly greater) degree of generality can turn out useful (see the proof of Proposition 5.28).*

5.3 Stability results for amplitudes

Often, in applications of persistent homology, one is interested in using different vectorisations of barcodes, or different distances between barcodes, and in fine-tuning the choice of vectorisations or distances using heuristic or theoretical observations. Our framework allows us to compare different ways of measuring persistence.

5.3.1 Stability under change of amplitudes

When working with data, one may want to study what information different types of vectorisations and distances are carrying. In our setting, this amounts to considering the identity functor on the category of persistence modules while changing the amplitudes. In particular, we now present three stability results: stability of the persistent magnitude, of the ρ_p -path metric, and of the ℓ -trop^k-path metric.

With this goal in mind, we begin by studying when the identity functor is amplitude bounding for the above amplitudes. Then, using Proposition 5.4 (since the identity functor is trivially exact) and [ST20, Theorem 4.7], we obtain the wanted stability results.

Since persistent magnitude is defined only for finitely presented modules, while ρ_p and ℓ -trop^k are more general, we first discuss the former alone and then the two latter together.

Lemma 5.8. *Let $\text{id}: (\text{PerM}_{fp}(\mathbb{R}), \text{totpers}) \rightarrow (\text{PerM}_{fp}(\mathbb{R}), |\cdot|)$, where $|\cdot|$ is the persistent magnitude and $\text{PerM}_{fp}(\mathbb{R})$ is the full subcategory of $\text{Vect}^{\mathbb{R}}$ given by finitely presented persistence modules. Then id is amplitude-bounding.*

Proof. Let $0 \leq b \leq d$ be arbitrary real numbers. Then we have

$$e^{-b} - e^{-d} \leq e^b(e^{-b} - e^{-d}) = 1 - e^{-(d-b)} \leq d - b$$

where the last inequality holds since for all real numbers x we have $1 + x \leq e^x$. Therefore, given any interval $[b, d)$, we have that the difference $e^{-b} - e^{-d}$ is bounded from above by the length of the interval. Thus, given a module $M \in \text{ob PerM}_{fp}(\mathbb{R})$ with $M \cong \bigoplus_{i=1}^m \mathbb{I}[b_i, d_i)$, we have that $|M| = \sum_{i=1}^m e^{-b_i} - e^{-d_i} \leq \sum_{i=1}^m (d_i - b_i) = \text{totpers}(M)$. \square

Proposition 5.9. *Let $\text{id}: (\text{PerM}_{fp}(\mathbb{R}), \text{totpers}) \rightarrow (\text{PerM}_{fp}(\mathbb{R}), |\cdot|)$. Then $d_{|\cdot|}(M, N) \leq d_{\text{totpers}}(M, N)$ for any $M, N \in \text{ob PerM}(\mathbb{R})$.*

Proof. This follows from Lemma 5.8 and Proposition 5.4. \square

Corollary 5.10. *Given any persistence modules $M, N \in \text{ob PerM}_{fp}(\mathbb{R})$, we have that*

$$d_{|\cdot|}(M, N) \leq W_1(\text{Dgm}(M), \text{Dgm}(N)).$$

Furthermore, let K be any finite CW complex, and $f, g: K \rightarrow \mathbb{R}$ monotone maps. Let $H_k(f)$ and $H_k(g)$ be the k -th persistent homology computed with respect to the corresponding sublevel set filtrations. Then we have:

$$d_{|\cdot|}(H_k(f), H_k(g)) \leq \|f - g\|.$$

Proof. By Lemma 5.8 we know that given any module $M \in \text{ob PerM}_{\mathcal{C}}(\mathbb{R})$,

$$|M| \leq \text{totpers}(M) = \rho_1(M).$$

Thus, by Proposition 5.4 we have that

$$d_{|\cdot|}(M, N) \leq d_{\rho_1}(\text{Dgm}(M), \text{Dgm}(N))$$

for all persistence modules $M, N \in \text{ob PerM}_{\mathcal{C}}(\mathbb{R})$. Now, the first inequality follows by Corollary 4.22. The second inequality follows from the first, and [ST20, Theorem 4.7]. \square

We now focus on ρ_p and ℓ -trop^k. For the former, we will make use of the following standard result:

Lemma 5.11. *Let $1 \leq p \leq \infty$ and let $\| - \|_p$ be the usual p -norm on Euclidean spaces. If $1 \leq p \leq q \leq \infty$ then $\| - \|_q \leq \| - \|_p$ and $\| - \|_p \leq c \| - \|_q$, where $c = n^{\frac{1}{p} - \frac{1}{q}}$, for n the number of summands.*

Proposition 5.12. *Let $p, q \in [1, \infty]$ and let $\text{id}: (\text{PerM}(\mathbb{R}), \rho_p) \rightarrow (\text{PerM}(\mathbb{R}), \rho_q)$ be the identity functor. Then id is amplitude bounding with constant $K = 1$ if and only if $p \leq q$. Moreover, if $p > q$, then there does not exist a K such that the identity functor is amplitude bounding.*

Proof. If $p \leq q$, then the sufficient condition follows by Lemma 5.11. For the other implication, we see from the second inequality of Lemma 5.11 that the bounding constant K depends on the number of intervals in the decomposition of the persistence module at hand. By adding a finite number of additional interval, it follows that there cannot be a fixed constant K , and the claim is proved. \square

Proposition 5.13. *Let $p \leq q \in [1, \infty)$ and let $\text{id}: (\text{PerM}(\mathbb{R}), \rho_p) \rightarrow (\text{PerM}(\mathbb{R}), \rho_q)$ be the identity functor. Then $d_{\rho_q}(M, N) \leq d_{\rho_p}(M, N)$ for any $M, N \in \text{ob PerM}(\mathbb{R})$.*

Proof. This follows from Proposition 5.12 and Proposition 5.4. \square

We now prove the analogous results in for the amplitude ℓ -trop^k.

Lemma 5.14. *Let $k, l \in \mathbb{N}$ and let $\text{id}: (\text{PerM}(\mathbb{R}), \ell\text{-trop}^k) \rightarrow (\text{PerM}(\mathbb{R}), \ell\text{-trop}^l)$ be the identity. If $k \geq l$, id is amplitude bounding with constant $K = 1$. If $1 \leq k \leq l$, then id is amplitude bounding with $K = l/k$.*

Proof. If $k \geq l$, then id is amplitude-bounding with constant $K = 1$, by definition of the tropical coordinates (Definition 3.13). The second claim is immediate from the elementary fact that, for a family of positive numbers $(x_i)_{1 \leq i \leq N}$,

$$k \max_{S \subset \{1, \dots, N\}, |S|=l} \sum x_i \leq l \max_{S \subset \{1, \dots, N\}, |S|=k} \sum x_i. \quad \square$$

Proposition 5.15. *Let $k, l \geq 1$ and let $\text{id}: (\text{PerM}(\mathbb{R}), \ell\text{-trop}^k) \rightarrow (\text{PerM}(\mathbb{R}), \ell\text{-trop}^l)$ be given by the identity. Then $d_{\ell\text{-trop}^l}(M, N) \leq \max\{1, \frac{l}{k}\} d_{\ell\text{-trop}^k}(M, N)$ for any $M, N \in \text{ob PerM}(\mathbb{R})$.*

Proof. This follows from Lemma 5.14 and Proposition 5.4. \square

Remark 5.16. *Proposition 5.15 allows for a different perspective on the distances $d_{\ell\text{-trop}^k}$: they monotonously interpolate between the interleaving (i.e. shift amplitude) and 1-Wasserstein distances. The distance $\ell\text{-trop}^1$ agrees with the shift amplitude ζ with respect to the positive unit vector in \mathbb{R} . In particular, by Proposition 4.18, $d_{\ell\text{-trop}^1}$ is equivalent to d_I up to a factor of 6. At the same time, for any $M \in \text{PerM}(\mathbb{R})$, we have $d_{\ell\text{-trop}^k}(M) = \rho_1(M)$, for k sufficiently large. As a consequence of this result and Corollary 4.22, for any $M, N \in \text{PerM}(\mathbb{R})$, it also follows that $d_{\ell\text{-trop}^k}(M, N) = W^1(M, N)$ for k sufficiently large. Hence, we have*

$$d_I(M, N) \leq d_{\ell\text{-trop}^k}(M, N) \xrightarrow{k \rightarrow \infty} W^1(M, N).$$

Finally, we obtain the wanted stability results:

Corollary 5.17. *(Stability of ρ_p and $\ell\text{-trop}^k$) Denote by T either ρ_p or $\ell\text{-trop}^k$. Let K be any finite CW complex, and $f, g: K \rightarrow \mathbb{R}$ monotone maps. Let $H_k(f)$ and $H_k(g)$ be the k -th persistent homology computed with respect to the corresponding sublevel set filtrations. Then, for all $p \in [1, \infty]$ and $k \in \mathbb{N}$, we have:*

$$d_T(H_k(f), H_k(g)) \leq \|f - g\|.$$

In [Kal19], the author proves a stability result for the absolute-value distance of ℓ -trop^k with respect to the p -Wasserstein distances as well as the bottleneck distance. This double stability property is also a consequence of the fact the absolute-value distance does not discriminate much. On the other hand, the path metrics are better discriminator, at the price of not be as stable as the absolute-value distance.

Example 5.18. Consider the two interval modules $M \cong \mathbb{I}[0, 3)$ and $N \cong \mathbb{I}[4, 5)$. Their bottleneck distance is 1.5. We now compute their path metric $d_{\ell\text{-trop}^k}$, for $k \in \mathbb{N}$. By definition, to calculate $d_{\ell\text{-trop}^k}(M, N)$ we need to minimize the cost of the cospan $M \leftarrow C \rightarrow N$. However, since M and N do not overlap, by matching theorem [BL15, Theorem 4.2], we have $d_{\ell\text{-trop}^k}(M, N) = 4$. \triangleleft

5.3.2 Interleaving and L^1 -path metric

The result in Proposition 5.5 allows for a simple proof of the following stability result between interleaving distance and the measure distances defined in [BSS20]. Recall that what is defined as the measure distance with respect to the Lebesgue measure in the latter source is alternatively given by $d_{\|\cdot\|_1^{\dim}}$, with $\|\cdot\|_1^{\dim}$ the L^1 -path metric with respect to the Hilbert amplitude (see Example 3.34).

Theorem 5.19. *Let d_I be the interleaving distance on $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$, with respect to the vector $v = (1, \dots, 1) \in \mathbb{R}^n$, and $d_{\|\cdot\|_1^{\dim}}$ the path distance on $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ induced by the Hilbert amplitude. Then we have*

$$d_I(M, N) \leq 4 \left(d_{\|\cdot\|_1^{\dim}}(M, N) \right)^{\frac{1}{n}}$$

for all $M, N \in \text{ob PerM}_{\mathfrak{C}}(\mathbb{R}^n)$.

Proof. We have seen in Proposition 4.18 that $d_I \leq d_{\zeta}$ where ζ is the shift amplitude induced by v , with respect to the supremum norm. Hence, by Proposition 5.5, it suffices to show that $\sqrt[n]{\cdot}$ defines a function f as in Proposition 5.5 with respect to the identity functor and $d_{\|\cdot\|_1^{\dim}}$ and ζ . So, let $M \in \text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ and, for any $u \in \mathbb{R}^n$ and $m \in M_u$, consider the unique morphism $\mathbb{F}[u + \mathbb{R}_{\geq 0}^n] \rightarrow M$ such that $1 \mapsto m$. Then its image is of the form $\mathbb{F}[S]$ where $S = u + \mathbb{R}_{\geq 0}^n \cap D$ for D a downset in \mathbb{R}^n . By monotonicity of $\|\cdot\|_1^{\dim}$, we have that $\|M\|_1^{\dim} \geq \lambda(S)$. Note that if $x^{\varepsilon} m \neq 0$ then S contains at least the cube of volume ε with lower left corner at u . The latter has volume ε^n . Hence,

$$\|M\|_1^{\dim} \geq \lambda(S) \geq \sup\{\varepsilon \mid x^{\varepsilon} m \neq 0 \text{ for some } m \in M_w, w \in \mathbb{R}^n\}^n = \zeta(M)^n.$$

Taking the n -th square root of the resulting inequality concludes the proof. \square

Remark 5.20. *Theorem 5.19 holds more generally for any connective algebra \mathfrak{X} that is closed under shifts.*

5.3.3 Pulling back along monotone functions

In Example 5.2, we have presented an example of amplitude-bounding functor on the category of persistence modules that arises from transformations between data sets in the form of metric spaces. More generally, one can study for which type of maps of the indexing poset one obtains an amplitude-bounding functor.

Throughout this section, we fix an abelian category \mathcal{A} equipped with an amplitude α . We suppose \mathcal{Q} and \mathcal{Q}' are measure spaces equipped with partial orders, with $\mathcal{C} \subseteq \mathcal{A}^{\mathcal{Q}}$ and $\mathcal{D} \subseteq \mathcal{A}^{\mathcal{Q}'}$ both α -measurable (see Definition 3.26).

Example 5.21. Let $\varphi: \mathcal{Q}' \rightarrow \mathcal{Q}$ be a monotone increasing function. Then pullback along φ gives a functor $\varphi^*: \mathcal{A}^{\mathcal{Q}} \rightarrow \mathcal{A}^{\mathcal{Q}'}$. Suppose that

1. The functor φ^* restricts to a functor $\varphi^*: \mathcal{C} \rightarrow \mathcal{D}$;
2. For any measurable set $S \subseteq \mathcal{Q}$, the set $\varphi^{-1}(S) \subseteq \mathcal{Q}'$ is measurable and satisfies $\mu'(\varphi^{-1}(S)) \leq K\mu(S)$ for some $K \in \mathbb{R}$ independent of S .

Then, as we show next, φ^* is amplitude bounding and exact. ◁

Proposition 5.22. *Let $p \in [1, \infty)$. The functor φ^* of Example 5.21 is an exact amplitude-bounding functor*

$$\varphi^*: (\mathcal{C}, \|\cdot\|_p^\alpha) \rightarrow (\mathcal{D}, \|\cdot\|_p^\alpha),$$

with constant $K^{1/p}$, where $\|\cdot\|_p^\alpha$ is defined in Definition 3.22.

Proof. Follows by classical techniques of measure theory and functional analysis. □

Remark 5.23. *The claim of Proposition 5.22 is also true for $p = \infty$ if we take $K^{1/\infty} = 1$. The argument is straightforward.*

Given persistence modules with a given domain \mathcal{Q} , one may restrict one's attention to a subposet of \mathcal{Q} with a compatible measure in a stable fashion. For example, one could look at the $\mathbb{R}_{\geq 0}$ -indexed portion of a \mathbb{R} -indexed persistence module, and this process would not increase the amplitude.

Example 5.24. Let \mathcal{Q} be a poset with measure μ , $\mathcal{Q}' \subseteq \mathcal{Q}$ a measurable subposet equipped with the induced measure, and $i: \mathcal{Q}' \rightarrow \mathcal{Q}$ the standard inclusion. Let $\mathcal{C} \subseteq \mathcal{A}^{\mathcal{Q}}$ and $\mathcal{D} \subseteq \mathcal{A}^{\mathcal{Q}'}$ be two abelian α -measurable subcategories with $i^*(\mathcal{C}) \subseteq \mathcal{D}$. Then the restriction functor $i^*: (\mathcal{C}, \|\cdot\|_p^\alpha) \rightarrow (\mathcal{D}, \|\cdot\|_p^\alpha)$ is amplitude-bounding with constant 1. ◁

The upshot of the preceding is that transformations of the poset underlying a category of persistence modules often induce stable, predictable changes in the persistence modules. Indeed, since the functor is exact, we can apply Proposition 5.4, for example, to Example 5.21, to carry stability results over the new poset.

5.4 Stability of quotient restrictions

In this section, we prove several stability results for the quotient-restriction functor introduced in Definition 3.47. We will discuss an application of these results in Section 5.6.

Proposition 5.25. *Let τ be a face of the positive cone of \mathbb{R}^n of dimension t , where $1 \leq t \leq n-1$. Without loss of generality, we take $\tau = \mathbb{R}_{\geq 0}\langle e_1, \dots, e_t \rangle$ and identify $\mathbb{R}^n/\mathbb{R}\tau$ with \mathbb{R}^{n-t} . Then the quotient restriction functor $(-)/\tau: \text{PerM}_{\mathcal{C}}(\mathbb{R}^n) \rightarrow \text{PerM}_{\mathcal{C}}(\mathbb{R}^{n-t})$ fulfills the following properties:*

1. For all $M, N \in \text{ob PerM}_{\mathcal{C}}(\mathbb{R}^n)$:

$$\begin{aligned} d_{\zeta'}(M/\tau, N/\tau) &\leq d_{\zeta}(M, N), \\ d_I(M/\tau, N/\tau) &\leq d_I(M, N) \end{aligned}$$

where ζ and ζ' are the shift amplitudes with respect to any fixed vector $v \in \mathbb{R}_{\geq 0}^n$ and its image v' under the projection to the first $n-t$ components $\mathbb{R}^n \rightarrow \mathbb{R}^{n-t}$, respectively, and d_I, d_I' denote the respective interleaving distances with respect to these vectors;

2. For all $M, N \in \text{ob PerM}_{\mathfrak{C}}(\mathbb{R}^n)$, for all $K > 0$ and for all $1 \leq p < \infty$:

$$d_{\|\cdot\|_p^{\text{dim}}} (M/\tau, N/\tau) \leq K d_{\|\cdot\|_p^{\text{dim}}} (M, N);$$

3. $(-)/\tau$ is not continuous with respect to the path metric induced by $\|\cdot\|_p^{\text{dim}}$ on target and source, with respect to any choice of finite measure μ on the Lebesgue σ -algebra on \mathbb{R}^n and any nonzero measure μ' on the one on \mathbb{R}^{n-t} , for $1 \leq p < \infty$.

Remark 5.26. Before we move on to the proof of the proposition, one should note that 2. should be understood more in terms of a general incompatibility between the L^p -amplitudes and the changes of dimension of the underlying poset than in terms of a stability result. To be more precise, topologically, Lemma 5.27 translates into the statement that $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ (as a space) is the topologically disjoint union of the fibers of quotient restriction. From this perspective, while quotient restriction is continuous with respect to the respective L^p -distances, this is because it is in fact locally constant. Therefore, we can interpret Proposition 5.25 as saying that while quotient restriction interacts well with shift amplitudes, it is essentially incompatible with any of the L^p -amplitudes.

Proof. As the quotient restriction is an exact functor, by Proposition 5.3 to prove the amplitude parts of statements 1.-2. it suffices to show that quotient restriction is amplitude bounding in the appropriate sense.

1. For the statement on the shift amplitudes, note that for each $m \in M/\tau$ and $\varepsilon > 0$ we have

$$x^{\varepsilon v'} m = x^{\varepsilon v'} (x^{\varepsilon(v-v')} m) = x^{\varepsilon v} m.$$

using the notation from Remark 3.41. Thus, if multiplying by $x^{\varepsilon v}$ annihilates M , multiplying by $x^{\varepsilon v'}$ annihilates M/τ . From this, by Remark 4.19, the statement on interleaving distances already follows up to a factor of 6. For the stricter inequality, note that the quotient restriction sends ε -interleavings into ε -interleavings.

2. This statement is a consequence of the following Lemma 5.27.

3. Finally, consider any cubical upset $U \subset \mathbb{R}^{n-t}$ with $\mu'(U) \neq 0$, which, if $\mu' \neq 0$, exists by continuity from below of μ' . Further, define

$$U_n = n e_1 + \mathbb{R}_{\geq 0}^t \times U$$

for any $n \in \mathbb{N}$. Now, consider $\mathbb{F}[U_n]$. By our assumption on τ , we have that $\mathbb{R}_{\geq 0}^t = \tau$, and therefore $\mathbb{F}[U_n]/\tau \cong \mathbb{F}[U]$. Therefore, we have that $\|\mathbb{F}[U_n]/\tau\|_p^{\text{dim}} = c > 0$ independently of n . Furthermore, we have $\bigcap_{n \in \mathbb{N}} U_n = \emptyset$. As finite measures are continuous from above, we have that $\|\mathbb{F}[U_n]\|_p^{\text{dim}} = \mu(U_n)^p$ converges to 0. Hence, $\mathbb{F}[U_n]$ converges to 0 in the path metric, showing that $(-)/\tau$ is not continuous. \square

Lemma 5.27. Let τ be a face of the positive cone in \mathbb{R}^n of dimension t , with $0 < t \leq n - 1$. Let $1 \leq q, p < \infty$ and consider \mathbb{R}^n and \mathbb{R}^{n-t} equipped with the Lebesgue measure. Let $M \in \text{ob PerM}_{\mathfrak{C}}(\mathbb{R}^n)$. Then

$$\|M\|_p^{\text{dim}} < \infty \implies \|M/\tau\|_q^{\text{dim}} = 0.$$

Proof. Similarly as in the statement of Proposition 5.25, we assume that τ is spanned by the first t unit vectors, and we identify \mathbb{R}^n/τ with \mathbb{R}^{n-t} . For $M \in \text{ob PerM}_{\mathfrak{C}}(\mathbb{R}^n)$, assume that $\|M\|_q^{\text{dim}} > 0$. Hence we have that $\|M/\tau\|_1^{\text{dim}} > 0$. Choose a cubical encoding of M/τ given by a map $e: \mathbb{R}^{n-t} \rightarrow \mathcal{Q}$. By assumption, at least one of the fibers F of this map has non-zero measure and the dimension of M/τ over this fiber is nonzero. As $\lambda(F) > 0$, there exists a

closed cube $Q \subset F$, with positive volume $\lambda(Q) > 0$ and minimal element $u \in Q$, which is fully contained in F . As all structure morphisms of M/τ which are going from points in F to points in F are isomorphisms, we have that $x^v \hat{m} \neq 0$, for $v \in -u + Q$. Now, let $\hat{m} \in M/\tau(u)$. For any representative $m \in M$ mapping to \hat{m} , we have that $x^v m \neq 0$ for $v \in \tau \times (-u + Q)$. In particular, $\dim(M) \geq 1$ holds on the latter product, which has Lebesgue measure $\infty \cdot \lambda(Q) = \infty$. Consequently we have $\|M\|_q^{\dim} > \infty$, for all $1 \leq p < \infty$. \square

5.5 Stability of the 0-th local cohomology

In Section 3.4, we described an extension of 0-th local cohomology to the setting of \mathbb{R}^n -modules. The following proposition summarizes its stability properties. The discussion is intended as an example of how the framework of amplitudes allows for the investigation of stability with respect to varying notions of size of a persistence module.

Proposition 5.28. *Let τ be some face of the positive cone in \mathbb{R}^n with positive dimension. The 0-th local cohomology $H_\tau^0 : \text{PerM}_{\mathcal{C}}(\mathbb{R}^n) \rightarrow \text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$, with respect to τ , is*

1. *Amplitude bounding with constant $K = 1$ with respect to any amplitude on $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$;*
2. *1-Lipschitz continuous with respect to the interleaving distance d_I ;*
3. *Not continuous with respect to $d_{\|\cdot\|_p^{\dim}}$, with respect to any finite, nonzero measure on the Lebesgue σ -algebra on \mathbb{R}^n , for $1 \leq p < \infty$;*
4. *1-Lipschitz continuous with respect to $d_{\|\cdot\|_p^{\dim}}$ with the Lebesgue measure on \mathbb{R}^n , for $1 \leq p < \infty$.*

Note that the 0-th local cohomology is not an exact functor, and thus we cannot use Proposition 5.4.

Proof. 1. This is an immediate consequence of the monotonicity of amplitudes, since $H_\tau^0(M)$ is given by a subobject of M .

2. The second claim follows from the fact that kernels, products, and localisations all commute with shifts and therefore, so does H_τ^0 . In particular, H_τ^0 carries interleavings into interleavings.

3. For the third statement, consider closed cubes $Q_k := [0, k]^n \subset \mathbb{R}^n$ with side-length k , for k in \mathbb{N} . Then $M^k := \mathbb{F}[Q_k]$ defines a sequence of finitely presented modules converging to $\mathbb{F}[\mathbb{R}_{\geq 0}^n]$ in $d_{\|\cdot\|_p^{\dim}}$ with respect to any finite measure. As $\tau \neq 0$, we have that $H_\tau^0(\mathbb{F}[\mathbb{R}_{\geq 0}^n]) = 0$. At the same time, it holds that $H_\tau^0(M^k) = M^k$, as Q_k is bounded in all directions. The latter sequence does not converge to 0, and the claim follows.

4. Finally, for the last statement we are going to show that H_τ^0 fulfils the conditions of Proposition 5.6. The second condition is immediate from left exactness of H_τ^0 . For the first condition, let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be an exact sequence in $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$, with M' with finite amplitude. Consider the following diagram with exact horizontal rows:

$$\begin{array}{ccccccccc}
0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \prod_{0 \neq \rho \subset \tau} M'_\rho & \longrightarrow & \prod_{0 \neq \rho \subset \tau} M_\rho & \longrightarrow & \prod_{0 \neq \rho \subset \tau} M''_\rho & \longrightarrow & 0
\end{array}$$

Assume that we have shown that $\alpha(\prod_{0 \neq \rho \subset \tau} M'_\rho) = 0$. Then, by the snake lemma, we have a short exact sequence $0 \rightarrow H_\tau^0(M') \rightarrow H_\tau^0(M) \rightarrow H_\tau^0(M'') \rightarrow N$, where N is a quotient of $\prod_{\rho \subset \tau} M'_\rho$. In particular, it fulfils $\alpha(N) = 0$, i.e. $N \cong 0$ in $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)/0_{\|\cdot\|_p^{\dim}}$. Hence, the former sequence

with N replaced by 0 is $\| - \|_p^{\dim}$ -exact. As $\prod_{0 \neq \rho \subset \tau} M'_\rho$ is a finite product, it suffices to show that M'_ρ has amplitude 0, for $\rho \subset \tau$ a face of positive dimension. Without loss of generality, we assume that $\rho = \mathbb{R}_{\geq 0} \langle e_1, \dots, e_t \rangle$ for some t smaller than the dimension of τ . Note that, for a cubically encoded module N to have L^p -amplitude 0, it suffices that each submodule $N' \subset N$ generated by a homogeneous element $x \in N$ has L^p -amplitude 0. Indeed, this condition already implies that the fibers of the encoding over which the module is nontrivial are all 0-sets. So, we assume that $m \in M_\rho$ is supported on a set S with nonzero measure. As all transition morphisms in direction ρ are isomorphisms, we may assume m to come from some homogeneous $m' \in M'$ with $x^v m' \neq 0$ for all $v \in \rho$. By shifting, we may assume $\deg m' = 0$. By assumption, there exists some $\varepsilon > 0$ such that $x^u m \neq 0$ for all u in the complement ρ^c of ρ whose norm is lesser or equal to ε . Thus, also $x^{u+v} m' \neq 0$ for $v \in \rho$ and $u \in \rho^c$ with $\|u\| \leq \varepsilon$. But then, m' is supported on $(\mathbb{R}^t \times [0, \varepsilon)^{n-t}) \cap \mathbb{R}_{\geq 0}^n$. Hence, m' is supported on a set with infinite Lebesgue measure, in contradiction to the amplitude finiteness of M' . \square

5.6 Stability of c_τ -rank

We conclude by showing some stability result for the c_τ -rank. This result are a consequence of the fact that the c_τ -rank can be expressed as a composition of two functors with an amplitude (Definition 3.49), even if it is not an amplitude itself (Example 3.21).

Proposition 5.29. *The c_τ -rank as a function of pseudo metric spaces $c_\tau(\text{ob PerM}_{\mathcal{C}}(\mathbb{R}^n), \text{d}) \rightarrow [0, \infty]$, where d is any of the metrics below, is:*

1. *Not continuous when $\text{d} = \text{d}_I$, the interleaving distance along any diagonal vector;*
2. *Not continuous when $\text{d} = \text{d}_{\| - \|_p^{\dim}}$, where $\| - \|_p^{\dim}$ is with respect to any finite, nonzero measure on the Lebesgue σ -algebra on \mathbb{R}^n , for $1 \leq p < \infty$;*
3. *K -Lipschitz continuous when $\text{d} = \text{d}_{\| - \|_1^{\dim}}$, where $\| - \|_1^{\dim}$ is with respect to the Lebesgue measure, for any $K > 0$ for $1 \leq p < \infty$.*

Proof. Without loss of generality, let $\tau = \langle e_1, \dots, e_t \rangle$.

1. To show c_τ is not continuous with respect to the interleaving distance, consider the sequence of persistence modules

$$M_k := \mathbb{F} \left[\left[0, \frac{1}{k}\right] \times [0, 1]^{t-1} \times \mathbb{R}_{\geq 0}^{n-t} \right]^k$$

for $k \in \mathbb{N}$. For $k \rightarrow \infty$, M_k converges to 0 in the interleaving distance. At the same time, we have

$$H_\tau^0(M_k)/\tau^c = \mathbb{F} \left[\left[0, \frac{1}{k}\right] \times [0, 1]^{t-1} \right]^k.$$

Thus:

$$c_\tau(M_k) = \|\mathbb{F} \left[\left[0, \frac{1}{k}\right] \times [0, 1]^{t-1} \right]^k\|_1^{\dim} = k \lambda \left(\left[0, \frac{1}{k}\right] \times [0, 1]^{t-1} \right) = 1$$

showing that c_τ -rank is not continuous in 0 with respect to the interleaving distance.

2. In the case $\text{d} = \text{d}_{\| - \|_p^{\dim}}$, with respect to some finite measure, again consider the sequence above, but this time with M_k replaced by $M_k := \mathbb{F}[ke_{k+1} + [0, 1]^t \times \mathbb{R}_{\geq 0}^{n-t}]$. By an analogous computation, we have $c_\tau\text{-rank}(M_k) = 1$. At the same time, M_k converges to 0, by the continuity from above of finite measures.

3. Finally, when $\text{d} = \text{d}_{\| - \|_1^{\dim}}$ with respect to the Lebesgue measure, observe how, by Proposition 5.25 and Proposition 5.28, $(-)/\tau^c \circ H_\tau^0$ is amplitude bounding, for any constant $K > 0$, for the amplitude $\| - \|_1^{\dim}$ with respect to the Lebesgue measure both on target and source. Finally, any amplitude is 1-Lipschitz with respect to the path metric induced by it (see Lemma 4.7), giving the result by composition of Lipschitz functions. \square

Note that the Lipschitz-continuity with respect to any K and L^p -amplitudes essentially translates to the fact that the fibers of the c_τ -rank all have infinite distance from each other. Hence, with respect to these distances the c_τ -rank is actually locally constant. Summarizing, the c_τ -rank does not fulfill a meaningful continuity result with respect to any of the amplitude distances of Proposition 5.29.

There are two possible ways to amend the difficulties we described in the previous paragraph. The first one is to consider metrics specifically depending on τ . In application, different parameter directions might often carry very different information. Hence, one could for example consider $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$ as $\text{PerM}_{\mathcal{C}}(\mathbb{R}^{n-t})$ valued persistence modules over \mathbb{R}^t and then study L^p -distance with respect to different amplitudes on $\text{PerM}_{\mathcal{C}}(\mathbb{R}^n)$. Alternatively, in the case where τ is a face of codimension 1, one may consider the barcode of $H_\tau^0(M)/\tau^c$ as an invariant instead. This contains more information than the c_τ -rank. At the same time, by the isometry theorem in [Les15, Thm. 3.4], together with Proposition 5.25 and Proposition 5.28, this gives an invariant that is Lipschitz continuous with respect to the interleaving distance.

6 Conclusions

Motivated by the study of distances between multiparameter persistence modules, in our work we introduce a general framework that allows to define distances based on a specific choice of an invariant. Our framework is very rich, as it encompasses many examples of invariants that have already been studied in the TDA literature, and powerful, as it allows us to derive new stability results. We believe that our framework opens a new avenue of research in TDA, with many interesting directions to be explored. Here, we discuss two directions that we intend to study as next steps.

Discriminating power of distances from amplitudes. As we have discussed in Section 4, the distance from amplitudes that is most widely used in applications of 1-parameter persistence is the absolute-value distance, which is very coarse, as it forgets all the categorical structure. Furthermore, for specific types of applications of 1-parameter persistent homology distances different from the Wasserstein-type distances might be better suited, such as distances obtained from certain vectorisations of barcodes. Thus, we intend to gain better insight into the discriminating power of several distances from amplitudes, both in the 1-parameter as well as in the multiparameter case, using several data sets of viral diseases.

Computability of distances from amplitudes. It is well known that several distances in the TDA literature are hard to compute, and a lot of effort is being done in computing approximations instead of exact distances, see, for instance, [Gäf16, KMN17]. There are several considerations that we can make for the distances that we introduce in our work, and which might help in reducing the computational complexity. Some distances, even if not particularly discriminating, can be calculated directly from the amplitudes, and thus the problem of calculating such distances is equivalent to the one of computing the amplitude themselves. This is not the case for path metrics, for which the problem is more complex. Path metrics associated with additive amplitudes may be approximated using techniques from variational methods and numerical analysis, since, as we show in Section 3.3, this distance is given by optimising an integral. For path metrics associated with amplitudes that are not additive, more refined techniques will be needed to compute approximations, and we intend to develop such techniques in future work.

Acknowledgments

We would like to thank the organisers of the 2020 Applied Category Theory Adjoint School, for bringing us together in this collaboration. Part of this work was initiated during a postdoctoral fellowship of NO and an internship of LW at the Max Planck Institute for Mathematics in the Sciences, and NO and LW would like to thank the Max Planck Institute for their generous support. We thank Håvard B. Bjerkevik, Wojciech Chachólski, Adrian Clough, Nicolò De Ponti Oliver Gäfvert, and Ezra Miller for useful discussions. BG was supported by the grant FAR2019-UniMORE and by the Austrian Science Fund (FWF) grant number P 29984-N35.

A Noise systems and amplitudes

We recall the main notions about noise systems from [SCL⁺17]. Whereas in [SCL⁺17] these notions were defined for tame persistence modules $\mathbb{Q}^n \rightarrow \text{Vect}$, we note that everything generalises to the setting of arbitrary abelian categories, and therefore we adopt this level of generality here.

Definition A.1. Let \mathcal{A} be an abelian category and $[0, \infty)$ the poset of non-negative reals. A **noise system** in \mathcal{A} is a collection $\mathcal{S} = \{\mathcal{S}_\varepsilon\}_{\varepsilon \in [0, \infty)}$ of collections of objects of \mathcal{A} , indexed by non-negative reals ε , such that:

- (i) the zero object belongs to \mathcal{S}_ε for any ε ;
- (ii) if $0 \leq \tau < \varepsilon$, then $\mathcal{S}_\tau \subseteq \mathcal{S}_\varepsilon$;
- (iii) if $0 \rightarrow F \rightarrow G \rightarrow H \rightarrow 0$ is an exact sequence in \mathcal{A} , then
 - if G is in \mathcal{S}_ε , then so are F and H (subobject property);
 - if F is in \mathcal{S}_ε and H is in \mathcal{S}_τ , then G is in $\mathcal{S}_{\varepsilon+\tau}$ (additivity property).

In [SCL⁺17], the authors provide several examples of noise systems. Among these examples, they discuss the property for a noise system of being closed under direct sum. Given a noise system \mathcal{S} on an abelian category \mathcal{A} , \mathcal{S} is closed under direct sum if, for all $\varepsilon \in \mathbb{R}$ and for any pair of objects M, N in \mathcal{S}_ε , then $M \oplus N$ is also in \mathcal{S}_ε . Later works [GC17, RC18] describe a machinery called **contour** that generates noise systems that are in particular closed under direct sums. Since being closed under direct sums is not standard for an amplitude, in this work we decided not to focus on this property. However, since every noise system can be seen as an amplitude (see Proposition A.3), the theory described in [GC17, RC18] can be used to generate amplitudes that on direct sums take as value the maximum of the evaluations on single elements.

A.1 Noise systems and amplitudes

We now study in detail the relation between amplitudes and noise systems. In particular, we will show that noise systems and amplitudes are the same by way of an adjunction. With this goal, we begin by defining the category of noise systems.

Definition A.2. Let Ns , the **category of noise systems**, be the category defined as follows.³ Objects of Ns are pairs $(\mathcal{A}, \mathcal{S})$ consisting of an abelian category \mathcal{A} and a noise system $\mathcal{S} = \{\mathcal{S}_\varepsilon\}_{\varepsilon \geq 0}$. Morphisms $(\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$ in Ns are additive functors $F: \mathcal{A} \rightarrow \mathcal{A}'$ such that there exists a non-negative real number $K \geq 0$ such that, for any $A \in \text{ob } \mathcal{A}$, if $A \in \mathcal{S}_\varepsilon$, then $F(A) \in \mathcal{S}'_{K\varepsilon}$. We call any such functor **noise bounding** with constant K .

³As we are dealing with categories of large categories here, we do not expect Ns to be locally small.

We now establish the relationship between noise systems and amplitudes. Essentially, this relationship comes down to characterizing a norm on a vector space by the set of ε -balls around the origin.

Proposition A.3. *Given an abelian category \mathcal{A} with a noise system $\mathcal{S} = \{\mathcal{S}_\varepsilon\}_{\varepsilon \geq 0}$, the function $\alpha_{\mathcal{S}}$ on $\text{ob } \mathcal{A}$ defined by*

$$\alpha_{\mathcal{S}}(A) := \inf \{ \varepsilon \mid A \in \mathcal{S}_\varepsilon \},$$

with the convention that $\inf \emptyset = \infty$, is an amplitude on \mathcal{A} . This construction gives rise to a functor $\text{NA}: \text{Ns} \rightarrow \text{Amp}$ defined by sending:

- *An object $(\mathcal{A}, \mathcal{S})$ of Ns to the object $(\mathcal{A}, \alpha_{\mathcal{S}})$ of Amp , and*
- *A noise-bounding functor $F: (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$ to an amplitude-bounding functor $F: (\mathcal{A}, \alpha_{\mathcal{S}}) \rightarrow (\mathcal{A}', \alpha_{\mathcal{S}'})$.*

Proof. All of the verifications here are straightforward. □

Having constructed amplitudes from noise systems in Proposition A.3, we are led to consider if the converse is true.

Definition A.4. Let α be an amplitude on an abelian category \mathcal{A} . The **noise system associated to α** , $\mathcal{S}(\alpha)$, is defined by

$$A \in \mathcal{S}(\alpha)_\varepsilon \iff \alpha(A) \leq \varepsilon,$$

for all $\varepsilon \in \mathbb{R}_{\geq 0}$. As in Proposition A.3, this construction induces a functor $\text{AN}: \text{Amp} \rightarrow \text{Ns}$ in the obvious way.

Proposition A.5. *The functors $\text{AN}: \text{Amp} \leftrightarrow \text{Ns}: \text{NA}$ induce an equivalence of categories. The natural isomorphisms of the equivalence are all given by the identity on the underlying categories, and are noise (amplitude) bounding for any $K > 1$.*

Proof. Both natural isomorphisms $\text{NA} \circ \text{AN} \cong 1_{\text{Amp}}$ and $\text{AN} \circ \text{NA} = 1_{\text{Ns}}$ are induced by the identity. The former is easily seen to fulfill $\text{NA}(\text{AN}(\alpha)) = \alpha$ directly. For the latter, note first that $\mathcal{S}_\varepsilon \subset \text{AN}(\text{NA}(\mathcal{S}))_\varepsilon$. Hence, the identity in one direction is noise bounding with constant 1. Conversely, for any $A \in \text{AN}(\text{NA}(\mathcal{S}))_\varepsilon$, we have $A \in \mathcal{S}_{K\varepsilon}$ for any $K > 1$. Hence, the converse direction is noise bounding with constant $K > 1$. □

A.2 Distances from noise systems vs. amplitudes

In this section, we relate distances obtained from noise systems and amplitudes, respectively. Noise systems and amplitudes induce metrics on the underlying abelian categories (see Definition 4.4), and we show that passing from a noise system to its corresponding amplitude does not affect the resulting metric. This all assembles into a commutative diagram of functors between various categories of noise systems, amplitudes and pseudometric spaces.

We begin by defining the relevant categories.

Definition A.6. Let **Pseudom** be the **category of pseudometric spaces** and Lipschitz maps, i.e. K -Lipschitz maps for some $K \geq 0$.

Here, we use the usual definition of “pseudometric space”, except that due to size considerations, we do not require the underlying objects of our pseudometric spaces to be sets.

To ensure that the functors of interest between abelian categories induce Lipschitz maps between the corresponding pseudometric spaces, we restrict our attention to “exact” functors, as in the following definition.

Definition A.7. Let Ns^{ex} be the subcategory of Ns where morphisms are *exact* noise-bounding functors. Moreover, let Amp^{ex} be the subcategory of Amp where morphisms are *exact* amplitude-bounding functors.

We are now ready to investigate the functoriality of the standard metric constructions.

Proposition A.8. *There exists a functor $\text{Ns}^{ex} \rightarrow \text{Pseudom}$*

(i) *Sending a pair $(\mathcal{A}, \mathcal{S})$ to the pseudometric space $(\text{ob } \mathcal{A}, d_{\mathcal{S}})$, where $d_{\mathcal{S}}$ is the metric defined by*

$$d_{\mathcal{S}}(A, B) = \inf\{\varepsilon \mid A \text{ and } B \text{ are } \varepsilon\text{-close}\},$$

where as usual $\inf \emptyset = \infty$, and the definition of ε -closeness is given in [SCL⁺17, Def. 8.4];

(ii) *Sending an exact noise-bounding functor $F : (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$ to the map on objects $F : (\text{ob } \mathcal{A}, d_{\mathcal{S}}) \rightarrow (\text{ob } \mathcal{A}', d_{\mathcal{S}'})$.*

Furthermore, if $F : (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$ is exact and noise-bounding with constant K , then the map on objects is K -Lipschitz.

Proof. The fact that $d_{\mathcal{S}}$ is a pseudometric is proved, in a special case, in [SCL⁺17, Prop. 8.7] and their arguments generalize immediately to the case we are considering. Hence, the only nontrivial part of the proposition is the last claim, since this implies the functor $\text{Ns}^{ex} \rightarrow \text{Pseudom}$ is well-defined on morphisms. So, suppose we are given an exact functor $F : (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$ which is noise-bounding with constant K . Furthermore, suppose $d_{\mathcal{S}}(A, B) < \varepsilon$ for some $A, B \in \text{ob } \mathcal{A}$. Then we can choose a span $A \xleftarrow{f_0} C \xrightarrow{f_1} B$ with $\ker f_0 \in \mathcal{S}_{\varepsilon_1}$, $\ker f_1 \in \mathcal{S}_{\varepsilon_2}$, $\text{coker } f_0 \in \mathcal{S}_{\varepsilon_3}$, and $\text{coker } f_1 \in \mathcal{S}_{\varepsilon_4}$, for some ε_i satisfying $\sum_{i=1}^4 \varepsilon_i < \varepsilon$. Since F is noise bounding and exact, the span $FA \xleftarrow{Ff_0} FC \xrightarrow{Ff_1} FB$ serves as a witness to the inequality $d_{\mathcal{S}'}(FA, FB) < K\varepsilon$. Because this holds for any $\varepsilon > d_{\mathcal{S}}(A, B)$, we see that $d_{\mathcal{S}'}(FA, FB) \leq Kd_{\mathcal{S}}(A, B)$, i.e. F is K -Lipschitz. \square

Corollary A.9. *There exists a functor $\text{Amp}^{ex} \rightarrow \text{Pseudom}$*

(i) *Sending an object (\mathcal{A}, α) to the pseudometric space $(\text{ob } \mathcal{A}, d_{\alpha})$, where d_{α} is as defined in Definition 4.4;*

(ii) *Sending an exact amplitude-bounding functor $F : (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$ to the map on objects $F : (\text{ob } \mathcal{A}, d_{\alpha}) \rightarrow (\text{ob } \mathcal{A}', d_{\alpha'})$.*

Furthermore, if $F : (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$ is exact and amplitude bounding with constant K , then $F : (\text{ob } \mathcal{A}, d_{\alpha}) \rightarrow (\text{ob } \mathcal{A}', d_{\alpha'})$ is K -Lipschitz.

Proof. By Corollary 4.12, such a functor is uniquely given by the composition $\text{Amp}^{ex} \xrightarrow{\text{AN}} \text{Ns}^{ex} \rightarrow \text{Pseudom}$ \square

Applying Proposition A.5, we obtain from this:

Corollary A.10. *The functors of Proposition A.8, Corollary A.9, Proposition A.3, and Definition A.4 assemble into an on the nose commutative triangles*

$$\begin{array}{ccc} \text{Amp}^{ex} & \xrightarrow{\text{AN}} & \text{Ns}^{ex} \\ & \searrow & \swarrow \\ & \text{Pseudom} & \end{array}, \quad \begin{array}{ccc} \text{Ns}^{ex} & \xrightarrow{\text{NA}} & \text{Amp}^{ex} \\ & \searrow & \swarrow \\ & \text{Pseudom} & \end{array}.$$

B Additional proofs

B.1 Proof of Proposition 3.14

We recall the statement:

Proposition B.1. *On the category $\text{PerM}(\mathbb{R})$:*

- i) *The tropical coordinate $\ell\text{-trop}^k$ is an amplitude, for all $k \in \mathbb{N}$;*
- ii) *The tropical coordinate $\sigma_{(1,0)^t}(\min\{b_1, m\ell_1\}, \ell_1, \dots, \min\{b_n, m\ell_n\}, \ell_n)$ is not an amplitude for any $m, l \in \mathbb{N}$.*

In the proof we will make use of the matching theorem in [BL15, Theorem 4.2], and for ease of reference, we now briefly recall the main results that we will use.

Remark B.2. *For an interval decomposable persistence module M , we denote by $\mathcal{D}_M = \{(b_i^M, d_i^M) \mid i = 1, \dots, n\}$ the multiset of its barcode. Furthermore, we denote by b^M the set of left endpoints of intervals in the decomposition and similarly by d^M the set of right endpoints. Suppose we are given a short exact sequence of interval decomposables $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$. Then, by possibly adding empty intervals to the barcode, we can find a common indexing of A , B and C , such that: $d_i^A = d_i^B$, $b_i^B = b_i^C$ and $b_i^A \geq b_i^B$, $d_i^B \geq d_i^C$.*

Proof. i) Recall that $\ell\text{-trop}^k$ is the sum of the k longest intervals. Clearly $\ell\text{-trop}^k(0) = 0$. Further, note that monotonicity is an immediate consequence of Remark B.2, as every bar in a submodule (resp. quotient) of a module B corresponds injectively to a (potentially) longer bar in the latter. We are left with showing subadditivity. As ephemeral modules (i.e. modules such that are supported on points) do not add to $\ell\text{-trop}^k$, we may as well assume all intervals involved to be left closed right open, to simplify the proof. Given a short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ of interval decomposable modules, let us first assume that B has lesser or equal to k interval components. Then, by Remark B.2, so do A and C . Hence, in this case $\ell\text{-trop}^k$ agrees with $\|\cdot\|_1^{\text{dim}}$ (see Example 3.30), that is it is given by the sum of the lengths of all intervals, which is additive. For the general case, let B' denote the submodule of B given by the k -longest bars (in case there are multiple such modules, choose any). We then have a commutative diagram with vertical monomorphisms and exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A' & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & 0 \end{array}$$

where A' is the intersection of B' with the kernel of $B \rightarrow C$, and C' is the image of B' in C . Note that then, again by Remark B.2, the upper exact sequence fulfills additivity for $\ell\text{-trop}^k$, as all modules involved have less or equal to k interval summands. Hence, we conclude

$$\ell\text{-trop}^k B = \ell\text{-trop}^k B' = \ell\text{-trop}^k A' + \ell\text{-trop}^k C' \leq \ell\text{-trop}^k A + \ell\text{-trop}^k C$$

since we already proved monotonicity.

ii) Consider two persistence modules A and B such that they are both constituted by only one interval module, with $b^\bullet < \ell^\bullet$, and such that $A \xrightarrow{\neq} B$. Then $\sigma_{(1,0)^t}(\min\{b^A, m\ell^A\}, \ell^A) > \sigma_{(1,0)^t}(\min\{b^B, m\ell^B\}, \ell^B)$. Thus, $\sigma_{(1,0)^t}$ does not satisfy the monotonicity condition (1) and is not an amplitude. \square

B.2 Proof of Theorem 3.38

As the definition of an amplitude does not involve any requirements with respect to countable families of objects, we can not expect measures to correspond to additive amplitudes in an infinite setting. For this reason, following the perspective of Example 3.30, it turns out that additive amplitudes arise from contents. For the remainder of this section, let \mathfrak{X} denote an algebra with respect to which $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ is an abelian subcategory of $\text{Vect}^{\mathcal{Q}}$.

The proof of Theorem 3.38 is partitioned into lemmas. We begin by showing that additive amplitudes in fact ignore the structure morphisms in the finitely encoded setting.

Lemma B.3. *Let α be an additive amplitude on $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$. For any $M, N \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$, if they have the same Hilbert function then $\alpha(M) = \alpha(N)$.*

Proof. We reduce the problem to the case where \mathcal{Q} is finite. In this case, the result is an immediate consequence of Proposition 3.37. By Lemma 2.11, we have that M and N admit a common \mathfrak{X} -encoding over the same poset \mathcal{Q}' by the same encoding map $e: \mathcal{Q} \rightarrow \mathcal{Q}'$. Furthermore, by restricting to its image if necessary, we can assume e to be surjective. Let M' and N' be the respective encoding modules in $\text{Vect}^{\mathcal{Q}'}$. By assumption we have

$$\dim(M') \circ e = \dim(M) = \dim(N) = \dim(N') \circ e$$

and, thus, by surjectivity of e

$$\dim(M') = \dim(N'). \quad (7)$$

As e^* is exact, composing α with e^* again defines an additive amplitude α' on $\text{Vect}^{\mathcal{Q}'}$. By definition, it holds that $\alpha(M) = \alpha'(M')$ and $\alpha(N) = \alpha'(N')$. Hence, by Eq. (7) it suffices to show that $\alpha'(M') = \alpha'(N')$, which is true by Proposition 3.37. \square

Next, we need to construct modules taking the role of indicator functions for elements of \mathfrak{X}_{Up} . Note that, as we are not working over a finite poset, we can not simply take the nontrivial structure morphisms to be 0, or we would not obtain a finitely encoded module.

Lemma B.4. *For each $S \in \mathfrak{X}_{\text{Up}}$ there exists a \mathfrak{X} -encoded module that has Hilbert function equal to the indicator function χ_S of S .*

Proof. Every $S \in \mathfrak{X}_{\text{Up}}$ can be written as $S = (U_1 \cap D_1) \cup \dots \cup (U_n \cap D_n)$, where $\{U_i\}_i$ is a family of upsets and $\{D_i\}_i$ a family of downsets in \mathfrak{X} . Denote by \mathcal{U} and \mathcal{D} the powerset of $\{U_i\}_i$ and $\{D_i\}_i$ respectively. These powersets can be seen as partially ordered sets, ordered in \mathcal{U} via \supset and in \mathcal{D} via \subset . Further, denote by \mathcal{U}_q (resp. \mathcal{D}_q) the collection of elements of \mathcal{U} (resp. \mathcal{D}) that contain q . The map of posets

$$\begin{aligned} e: \mathcal{Q} &\longrightarrow \mathcal{U} \times \mathcal{D} \\ q &\longmapsto (\mathcal{U}_q, \mathcal{D}_q) \end{aligned}$$

is a map of posets by construction. Each fiber is given by the intersection of a set of upsets and downsets in \mathfrak{X} . In particular, e defines a \mathfrak{X} -encoding map. Let S' be the subset of $\mathcal{U} \times \mathcal{D}$ given by all pairs $(\{U_{i_1}, \dots, U_{i_k}\}, \{D_{j_1}, \dots, D_{j_l}\})$ such that the intersection $\{i_1, \dots, i_k\} \cap \{j_1, \dots, j_l\}$ is nonempty, where the sets of indices $\{i_1, \dots, i_k\}$ and $\{j_1, \dots, j_l\}$ are subsets of $\{1, \dots, n\}$. Then

$$S = e^{-1}(S'). \quad (8)$$

Let M be the persistence module on $\mathcal{U} \times \mathcal{D}$ whose Hilbert function is 1 on S' and 0 otherwise, and such that its structure morphisms are all zero. Then $e^*(M)$ is \mathfrak{X} -encoded by definition and has χ_S as its Hilbert function, by Eq. (8). \square

We now have all the needed tools for the proof of Theorem 3.38.

Proof of Theorem 3.38. The proof of injectivity is identical to the one for Proposition 3.37, after replacing the indicator modules used there by the ones given by Lemma B.4 thanks to Lemma B.3.

For surjectivity, we use again the same technique of the proof of Proposition 3.37, however, we need to be careful on the details. Consider $\alpha \in \text{AdAmp}(\text{PerM}_{\mathfrak{X}}(\mathcal{Q}))$. First, for each $X \in \mathfrak{X}_{\text{Up}}$, choose a \mathfrak{X} -encoded module N_X such that its Hilbert function is the indicator function χ_X , as in Lemma B.4 and Lemma B.3.

Then set $\nu(X) := \alpha(N_X)$. By Lemma B.3, this construction does not depend on the choice of module N_X . For any two disjoint measurable sets X, X' , by Lemma B.3 we obtain that

$$\alpha(N_{X \cup X'}) = \alpha(N_X \oplus N_{X'}).$$

From this equation and the additivity of α , together with the fact that $\alpha(0) = 0$, we obtain that ν defines a content. It remains to be seen that $\|M\|_{\text{dim}}^1 = \alpha(M)$, for all $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$. Note that, by Lemma B.3, we may replace M in this equation by $\bigoplus_{n \in \mathbb{N}} (N_{\dim(M)^{-1}(n)})^n$, which is a finite sum. Hence, by the definition of ν and the additivity of α , we have:

$$\begin{aligned} \alpha(M) &= \alpha\left(\bigoplus_{n \in \mathbb{N}} (N_{\dim(M)^{-1}(n)})^n\right) = \sum_{n \in \mathbb{N}} n \alpha(N_{\dim(M)^{-1}(n)}) \\ &= \sum_{n \in \mathbb{N}} n \nu(\dim(M)^{-1}(n)) = \int_{\mathcal{Q}} \dim(M) d\nu. \end{aligned} \quad \square$$

B.3 Proof of Proposition 3.43

We recall the statement:

Proposition B.5. *Let M be a cubically encoded persistence module over \mathbb{Z}^n and \mathfrak{p} a homogeneous prime of $\mathbb{F}[x_1, \dots, x_n]$. Then the two definition of $c_{\mathfrak{p}}$ -rank of M given in Example 3.19 and Definition 3.42 coincide.*

Proof. Without loss of generality, let $\mathfrak{p} = \langle x_1, \dots, x_t \rangle$. As both constructions of $c_{\mathfrak{p}}$ -rank require to apply first the 0-th local cohomology functor, we may assume that $M = H_{\mathfrak{p}}^0(M)$ to simplify notation. Note that $\|M/\mathfrak{J}M\|_1^{\text{dim}} = \dim_{\mathbb{F}}(M/\mathfrak{J}M)$, where by $\dim_{\mathbb{F}}$ we mean the dimension of the \mathbb{Z}^n -graded objects considered as an \mathbb{F} vector space. As in Definition 3.42, $M/\mathfrak{J}M$ can be computed by first localizing at the elements x_{t+1}, \dots, x_n and then restricting the resulting module to $\mathbb{Z}^t \times \{0\} \subset \mathbb{Z}^n$ (compare [Mil20a, Rem. 2.23] for the analogous real cases). The localized module M_{x_{t+1}, \dots, x_n} is a \mathbb{Z}^n graded module over $\mathbb{F}[x_1, \dots, x_t, x_{t+1}^{-1}, \dots, x_n^{-1}]$. The computation of $\dim_{\mathbb{F}}(M/\mathfrak{J}M)$ is independent from the multiplicative structure in direction of x_1, \dots, x_t . Hence, we may just consider M_{x_{t+1}, \dots, x_n} as a \mathbb{Z}^{n-t} -graded $R = \mathbb{F}[x_{t+1}^{-1}, \dots, x_n^{-1}]$ -module, denoted by M' . Then M' is a free R -free module of rank $\dim_{\mathbb{F}}(M'_0)$. Since the restriction to $\mathbb{Z}^t \times \{0\} \subset \mathbb{Z}^n$ corresponds to evaluating at 0 in this setting, we have

$$\|M/\mathfrak{J}M\|_1^{\text{dim}} = \dim_{\mathbb{F}}(M/\mathfrak{J}M) = \dim_{\mathbb{F}}(M_0) = \text{rank}_R(M').$$

By definition, $c_{\mathfrak{p}}$ -rank in the sense of Example 3.19 is given by the $\text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n])$ -dimension of $M \otimes_{\mathbb{F}[x_{t+1}, \dots, x_n]} \text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n])$. By standard commutative algebra, the latter admits the alternative description:

$$\begin{aligned} M \otimes_{\mathbb{F}[x_{t+1}, \dots, x_n]} \text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n]) &\cong (M \otimes_{\mathbb{F}[x_{t+1}, \dots, x_n]} R) \otimes_R \text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n]) \\ &\cong M' \otimes_R \text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n]). \end{aligned}$$

As M' is, by construction, a free module of rank $\dim_{\mathbb{F}}(M'_0)$, the dimension of the right hand side expression is $\text{rank}_R(M')$. This finishes the proof. \square

References

- [ACC13] Aaron Adcock, Erik Carlsson, and Gunnar Carlsson. The ring of algebraic functions on persistence bar codes. *Homology, Homotopy and Applications*, 18(1):381–402, 04 2013.
- [BBCB20] Magnus Bakke Botnan and William Crawley-Boevey. Decomposition of persistence modules. *Proceedings of the AMS*, 148:4581–4596, 2020.
- [BBL21] Håvard Bakke Bjerkevik and Michael Lesnick. ℓ^p -distances on multiparameter persistence modules, 2021. Preprint, available at arXiv:2106.13589.
- [BBLO20] Magnus Bakke Botnan, Vadim Lebovici, and Steve Oudot. On rectangle-decomposable 2-parameter persistence modules. In *36th International Symposium on Computational Geometry, SoCG 2020, June 23-26, 2020, Zürich, Switzerland*, volume 164 of *LIPICs*, pages 22:1–22:16, 2020.
- [BBOO21] Magnus Bakke Botnan, Steffen Oppermann, and Steve Oudot. Signed barcodes for multi-parameter persistence via rank decompositions and rank-exact resolutions, 2021. Preprint, available at arXiv:2107.06800.
- [BdSS15] Peter Bubenik, Vin de Silva, and Jonathan Scott. Metrics for generalized persistence modules. *Foundations of Computational Mathematics*, 15(6):1501–1531, Dec 2015.
- [BE18] Mickaël Buchet and Emerson G. Escolar. Realizations of indecomposable persistence modules of arbitrarily large dimension. In *34th International Symposium on Computational Geometry (SoCG 2018)*, volume 99 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 15:1–15:13, Dagstuhl, Germany, 2018.
- [BL15] Ulrich Bauer and Michael Lesnick. Induced matchings of barcodes and the algebraic stability of persistence. *Journal of Computational Geometry*, 6(2):162–191, 2015.
- [BMSW17] Ulrich Bauer, Axel Munk, Hannes Sieling, and Max Wardetzky. Persistence barcodes versus Kolmogorov signatures: Detecting modes of one-dimensional signals. *Foundations of Computational Mathematics*, 17(1):1–33, Feb 2017.
- [Bog07] Vladimir I. Bogachev. *Measure theory*, volume 1. Springer Science & Business Media, 2007.
- [BP21] Nicolas Berkouk and François Petit. Ephemeral persistence modules and distance comparison. *Algebraic & Geometric Topology*, 21(1):247–277, Feb 2021.
- [BPC⁺18] Francisco Belchi, Mariam Pirashvili, Joy Conway, Michael Bennett, Ratko Djukanovic, and Jacek Brodzki. Lung topology characteristics in patients with chronic obstructive pulmonary disease. *Scientific Reports*, 8(1):5341, 2018.
- [BSS20] Peter Bubenik, John A. Scott, and Don Stanley. An algebraic wasserstein distance for generalized persistence modules, 2020. Preprint, available at arXiv:1809.09654v2.
- [BW20] David Bramer and Guo-Wei Wei. Atom-specific persistent homology and its application to protein flexibility analysis. *Computational and Mathematical Biophysics*, 8(1):1–35, 2020.
- [CGL21] Wojciech Chachólski, Barbara Giunti, and Claudia Landi. Invariants for tame parametrised chain complexes. *Homology, Homotopy and Applications*, 23(2):183–213, 2021.

- [CK18] René Corbet and Michael Kerber. The representation theorem of persistence revisited and generalized. *Journal of Applied and Computational Topology*, 2:1–31, 2018.
- [CSEH07] David Cohen-Steiner, Herbert Edelsbrunner, and John Harer. Stability of persistence diagrams. *Discrete & Computational Geometry*, 37:103–120, 2007.
- [CSEHM10] David Cohen-Steiner, Herbert Edelsbrunner, John Harer, and Yuriy Mileyko. Lipschitz functions have l_p -stable persistence. *Foundations of Computational Mathematics*, 10(2):127–139, 2010.
- [CSZ10] Gunnar Carlsson, Gurjeet Singh, and Afra J. Zomorodian. Computing multidimensional persistence. *Journal of Computational Geometry*, 1(1):72–100, 2010.
- [CZ09] Gunnar Carlsson and Afra J. Zomorodian. The theory of multidimensional persistence. *Discrete & Computational Geometry*, 42(1):71–93, Jul 2009.
- [Gäf16] Oliver Gäfvert. Algorithms for multidimensional persistence, 2016. Master dissertation, KTH Royal Institute of Technology.
- [GC17] Oliver Gäfvert and Wojciech Chachólski. Stable invariants for multidimensional persistence, 2017. Preprint, available at arXiv:1703.03632.
- [GG06] Benjamin T. Graham and Geoffrey R. Grimmett. Influence and sharp-threshold theorems for monotonic measures. *The Annals of Probability*, 34(5):1726–1745, 2006.
- [GGM19] Noah Giansiracusa, Robert Giansiracusa, and Chul Moon. Persistent homology machine learning for fingerprint classification. In *2019 18th IEEE International Conference On Machine Learning And Applications (ICMLA)*, pages 1219–1226, 2019.
- [GH21] Dejan Govc and Richard Hepworth. Persistent magnitude. *Journal of Pure and Applied Algebra*, 225, 2021.
- [GL20] Barbara Giunti and Jānis Lazovskis. Database of TDA real-world applications, 2020. Available at www.zotero.org/groups/2425412/tda-applications.
- [HKNU17] Christoph Hofer, Roland Kwitt, Marc Niethammer, and Andreas Uhl. Deep learning with topological signatures. *Advances in Neural Information Processing Systems*, 30:1633–1643, 2017.
- [HOST19] Heather A. Harrington, Nina Otter, Hal Schenck, and Ulrike Tillmann. Stratifying multiparameter persistent homology. *SIAM Journal on Applied Algebra and Geometry*, 3(3):439–471, 2019.
- [Kal19] Sara Kališnik. Tropical coordinates on the space of persistence barcodes. *Foundations of Computational Mathematics*, 19(1):101–129, 2019.
- [KMN17] Michael Kerber, Dmitriy Morozov, and Arnur Nigmatov. Geometry helps to compare persistence diagrams. *ACM J. Exp. Algorithmics*, 22:1–20, 2017.
- [Lan02] Serge Lang. *Algebra*. Springer New York, third edition, 2002.
- [LBD⁺17] Yongjin Lee, Senja D. Barthel, Paweł Dłotko, S. Mohamad Moosavi, Kathryn Hess, and Berend Smit. Quantifying similarity of pore-geometry in nanoporous materials. *Nature communications*, 8:15396, 2017.

- [Les15] Michael Lesnick. The theory of the interleaving distance on multidimensional persistence modules. *Foundations of Computational Mathematics*, 15(3):613–650, 2015.
- [LW15] Michael Lesnick and Matthew Wright. Interactive visualization of 2-d persistence modules, 2015. Preprint, available at arXiv:1512.00180.
- [Mil20a] Ezra Miller. Essential graded algebra over polynomial rings with real exponents, 2020. Preprint, available at arXiv:2008.03819.
- [Mil20b] Ezra Miller. Homological algebra of modules over posets, 2020. Preprint, available at arXiv:2008.00063.
- [MOW20] Guido Montúfar, Nina Otter, and Yuguang Wang. Can neural networks learn persistent homology features?, 2020. Topological Data Analysis and Beyond Workshop at the *34th Conference on Neural Information Processing Systems (NeurIPS 2020)*, Vancouver, Canada, available at arXiv:2011.14688.
- [Ott18] Nina Otter. Magnitude meets persistence. Homology theories for filtered simplicial sets. *arXiv:1807.01540*, 2018. To appear in *Homology, Homotopy and Applications*.
- [RC18] Henri Riihimäki and Wojciech Chachólski. Generalized persistence analysis based on stable rank invariant, 2018. Preprint, available at arXiv:1807.01217.
- [Rud87] Walter Rudin. *Real and Complex Analysis, 3rd Ed.* McGraw-Hill, Inc., USA, 1987.
- [SCL⁺17] Martina Scolamiero, Wojciech Chachólski, Anders Lundman, Ryan Ramanujam, and Sebastian Öberg. Multidimensional persistence and noise. *Foundations of Computational Mathematics*, 17:1367–1406, 2017.
- [ST20] Primoz Skraba and Katharine Turner. Wasserstein stability for persistence diagrams, 2020. Preprint, available at arXiv:2006.16824.
- [Sta18] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2018.
- [Tho19] Ashleigh L. Thomas. Invariants and metrics for multiparameter persistent homology, 2019. PhD thesis, Duke University.
- [Waa21] Lukas Waas. Notes on finitely encoded modules, 2021. Available at www.mathi.uni-heidelberg.de/~lwaas/work/pers_notes_01.
- [ZC05] A. Zomorodian and G. Carlsson. Computing persistent homology. *Discrete & Computational Geometry*, 33:249–274, 2005.