

THE ISOMORPHISM PROBLEM OF PROJECTIVE SCHEMES AND RELATED ALGORITHMIC PROBLEMS

TAKEHIKO YASUDA

ABSTRACT. We discuss the isomorphism problem of projective schemes; given two projective schemes, can we algorithmically decide whether they are isomorphic? We give affirmative answers in the case of one-dimensional projective schemes, the case of smooth irreducible varieties with a big canonical sheaf or a big anti-canonical sheaf, and the case of K3 surfaces with a finite automorphism group. As related algorithmic problems, we also discuss decidability of positivity properties of invertible sheaves, and approximation of the nef cone and the pseudo-effective cone.

CONTENTS

1. Introduction	2
Acknowledgments	4
2. Preliminaries	4
3. Semi-decidability of the isomorphism problem of projective schemes	5
4. Hilbert schemes	7
4.1. The Hilbert scheme of a projective space	7
4.2. The universal family	9
4.3. The Hilbert scheme of a general projective scheme	9
4.4. The action of GL_r	11
5. Projective equivalence	12
6. Hom schemes and Iso schemes	12
7. One-dimensional schemes	15
8. Varieties with a big canonical sheaf or a big anti-canonical sheaf	18
9. Computing intersection numbers	20
10. Positivity of invertible sheaves	21
10.1. Global generation	21
10.2. Very ampleness	22
10.3. Ampleness	22
10.4. Nefness, bigness and pseudo-effectivity	23
10.5. Approximating nef and pseudo-effective cones	25
11. K3 surfaces	26
References	28

This work was supported by JSPS KAKENHI Grant Number JP18H01112.

1. INTRODUCTION

The main purpose of this paper is to discuss the isomorphism problem of projective schemes over the field $\overline{\mathbb{Q}}$ of algebraic numbers. Is it algorithmically decidable whether two given projective $\overline{\mathbb{Q}}$ -schemes are isomorphic? What if we restrict ourselves to some class of projective schemes, for example, the class of smooth projective varieties having a prescribed invariant. Poonen [Poo11] writes that Totaro asked him about this problem in 2007. The case of smooth irreducible curves was treated earlier in the 2005 paper [BGJGP05] by Baker, González-Jiménez, González, and Poonen. The same problem was asked also by Arapura on MathOverflow¹ in 2010.

To the best of the author's knowledge, the decidability of the isomorphism problem has been proved in the following two cases:

- (1) Smooth irreducible curves ([BGJGP05, Lem. 5.1] for the case of genus $\neq 1$ and Poonen's comment in the MathOverflow thread mentioned above for the case of genus one).
- (2) Varieties of general type (see [Poo14, Rem. 12.3] for the proof due to Totaro).

In the same MathOverflow thread as above, there is also discussion about the cases of K3 surfaces and abelian surfaces, which has not reached a definite conclusion.

The main result of the paper is to prove that the isomorphism problem is decidable in the following cases:

- (1) One-dimensional projective schemes (Theorem 7.3).
- (2) Smooth projective varieties with a big canonical sheaf or a big anti-canonical sheaf (Theorem 8.1).
- (3) K3 surfaces with a finite automorphism group (Theorem 11.3).

The first two cases slightly generalize ones mentioned above. As an application of the first case, we show the decidability also in the case of one-dimensional reduced *quasi-projective* schemes (Theorem 7.4). In birational geometry, varieties of Kodaira dimensions $-\infty$ and 0 as well as ones of general type have special importance, as they are considered as building blocks of all varieties. With case (2) above being solved, varieties of Kodaira dimension 0 would be the remaining most important case. Besides K3 surfaces, the isomorphism problem for abelian varieties should be important, but we do not discuss it in this paper.

Remark 1.1. It appears difficult to apply the global Torelli theorem for K3 surfaces [BHPV04, p. 332] to solve the isomorphism problem affirmatively. We can approximate the Hodge structure on cohomology groups with arbitrary precision [Sim08]. But the moduli space of (marked) K3 surfaces is not Hausdorff [BHPV04, p. 334]. This suggests that we cannot detect non-isomorphism of K3 surfaces by approximation.

Our strategy to prove these results is to compute the Iso schemes $\underline{\text{Iso}}_{P_i}(X, Y)$ for the given projective schemes X and Y and for finitely many polynomials P_i . The entire Iso scheme $\underline{\text{Iso}}(X, Y)$ is the moduli scheme of isomorphisms $X \xrightarrow{\sim} Y$ and can be embedded into the Hilbert scheme $\text{Hilb}(X \times Y)$ by sending an isomorphism $f: X \rightarrow Y$

¹<https://mathoverflow.net/questions/21883/isomorphism-problem-for-commutative-algebras-and-schemes>

to its graph $\Gamma_f \subset X \times Y$. The Hilbert scheme is decomposed as $\text{Hilb}(X \times Y) = \coprod_P \text{Hilb}_P(X \times Y)$, where P runs over countably many polynomials. This induces a decomposition $\underline{\text{Iso}}(X, Y) = \coprod_P \underline{\text{Iso}}_P(X, Y)$ of the Iso scheme. For each polynomial P , $\underline{\text{Iso}}_P(X, Y)$ is of finite type, but the entire $\underline{\text{Iso}}(X, Y)$ is not generally so. We explain how to algorithmically compute $\underline{\text{Iso}}_P(X, Y)$ for each P . Having the method of computing Iso schemes, we then construct an algorithm for each of the classes of projective schemes mentioned above that produces finitely many polynomials P_1, \dots, P_n from the given projective schemes X and Y . These polynomials satisfy the condition that X and Y are isomorphic if and only if $\underline{\text{Iso}}_{P_i}(X, Y) \neq \emptyset$ for some i . Then, whether X and Y are isomorphic or not is checked by computing these Iso schemes. In construction of finitely many polynomials as above, we use the Kodaira vanishing as a key ingredient in case (2) and use computation of the nef cone in case (3).

We also discuss several algorithmic problems related to the isomorphism problem. Firstly, we explicitly describe an algorithm to check whether two given projective schemes embedded in the same projective space are projectively equivalent (Section 5). If two projective schemes are projectively equivalent, then they are isomorphic, but the converse does not generally hold. Secondly, partly using computation of intersection numbers, we discuss positivity properties of invertible sheaves from the algorithmic viewpoint. We see that global generation of a coherent sheaf and very ampleness of an invertible sheaf on a projective scheme is decidable. Using the Nakai-Moishezon criterion for ampleness and computation of intersection numbers, we see that, if the scheme is smooth and irreducible, then ampleness of an invertible sheaf is also decidable (Proposition 10.4). We do not know whether other positivities, bigness, nefness and pseudo-effectivity, are decidable. However, if we can compute the Picard number, then we can approximate the nef cone and the pseudo-effective cone with arbitrary precision (Proposition 10.14). Note that Poonen, Testa and van Luijk [PTL15] proved that the Picard number of a smooth irreducible projective variety is computable, if the Tate conjecture is true. This is the case for K3 surfaces. For a K3 surface with a finite automorphism group, we can compute its nef cone (not approximately but exactly), which is used to show the decidability in case (3) above.

To end this introduction, we mention a few more related works. Truong [Tru18] proved the decidability of the bounded birationality problem. Namely, he proved that for projective varieties $X \subset \mathbb{P}^m$ and $Y \subset \mathbb{P}^n$ and for a positive integer d , we can decide whether there exists a rational map $\mathbb{P}^m \dashrightarrow \mathbb{P}^n$ of degree $\leq d$ that restricts to a birational map $X \dashrightarrow Y$. He also proved the decidability of the *bounded* isomorphism problem in the case where one of the two given varieties is smooth. To prove these results, he showed computability of a variety parametrizing rational maps $\mathbb{P}^m \dashrightarrow \mathbb{P}^n$ with this property, which is similar to our computability result regarding Iso schemes.

The isomorphism problem that we consider in this paper is a special case of the problem regarding the existence of a morphism $X \rightarrow Y$ of k -schemes possibly imposed with some condition for a more general field or ring k . For example, the famous negative solution by Davis, Matiyasevich, Putnam, and Robinson to Hilbert's tenth problem says that the existence of a morphism $\text{Spec } \mathbb{Z} \rightarrow Y$ with Y an affine scheme of finite type over \mathbb{Z} is undecidable. One of the other undecidability results in this direction is the

one of Kanel'-Belov and Chilikov [KBC19] (see also [Kol20]) that the existence of an embedding $X \hookrightarrow Y$ of varieties over \mathbb{R} or $\overline{\mathbb{Q}}$ is undecidable.

The rest of the paper is organized as follows. In Section 2, we set up our basic convention. In particular, we clarify what we mean by saying that some object (for example, a scheme or an invertible sheaf) is given. In Section 3, we show that the isomorphism problem of projective schemes is semi-decidable. In Section 4, we explain how to compute the Hilbert scheme for each polynomial. In Section 5, we apply computation of the Hilbert scheme to show that it is decidable whether two projective schemes embedded in the same projective space is projectively equivalent. Although each result in sections 3 to 5 would be known to specialists, we include them for the sake of reader's convenience. The reader who knows these materials well may skip these sections. In Section 6, we explain how to compute the Hom scheme and the Iso scheme for each polynomial. In Section 7, we show the decidability of the isomorphism problem for one-dimensional projective schemes and the one for one-dimensional quasi-projective reduced schemes. In Section 8, we do the same for the case of smooth irreducible varieties with a big canonical sheaf or a big anti-canonical sheaf. In Section 9, we explain how to compute intersection numbers on a smooth irreducible projective variety. In Section 10, we discuss decidability of positivity properties of invertible sheaves. In particular, we show that ampleness of an invertible sheaf on a smooth variety is decidable and that the nef cone and the pseudo-effective cone are approximated by rational polyhedral cones with arbitrary precision. In Section 11, we discuss the isomorphism problem for K3 surfaces as well as smooth varieties with a rational polyhedral nef cone.

Acknowledgments. The author is grateful to Bjorn Poonen and Burt Totaro for pointing out references and for valuable suggestions. He would also like to thank Ichiro Shimada, Sho Ejiri, and an anonymous referee for helpful comments.

2. PRELIMINARIES

Throughout the paper, we work over the field of complex algebraic numbers, $\overline{\mathbb{Q}} \subset \mathbb{C}$, which is denoted by k . As explained in [Sim08, Section 2.1], elements of this field are expressed by finite data and four basic arithmetic operations on them, addition, subtraction, multiplication and division, are algorithmically computable. We can also algorithmically decide whether or not two expressions give the same number. It follows that we can also express polynomials with coefficients in k by finite data and algorithmically compute their addition, subtraction and multiplication. We can also compute the Gröbner basis of an ideal in a polynomial ring $k[x_1, \dots, x_m]$. Thus we can also make various computation based on the Gröbner basis. For example, we can algorithmically check whether or not an ideal is contained in another ideal in the same polynomial ring (this is an application of the ideal membership test; see [Eis95, 15.10.1]). We also note that the elements of k as well as the elements of $k[x_1, \dots, x_m]$ are enumerable.

When we say that a projective scheme X is given, we mean that we are given finitely many homogeneous polynomials $f_1, \dots, f_l \in k[x_1, \dots, x_m]$ such that X is the closed subscheme of $\mathbb{P}^{m-1} = \text{Proj } k[x_1, \dots, x_m]$ defined by the ideal (f_1, \dots, f_l) . In particular, we are given an embedding $\iota: X \hookrightarrow \mathbb{P}^{m-1}$ into a projective space, the induced very ample invertible sheaf $\iota^* \mathcal{O}_{\mathbb{P}^{m-1}}(1)$ and the homogeneous coordinate ring

$R_X := k[x_1, \dots, x_m]/(f_1, \dots, f_l)$, which is also denoted by R omitting the subscript X . From these data, we can compute the standard affine charts $X_i := X \cap \{x_i \neq 0\}$ for $1 \leq i \leq m$, which cover X . Indeed, if $k[x_1, \dots, \check{x}_i, \dots, x_m]$ is the polynomial ring with x_i removed and if $f_j^{(i)}$ denotes the polynomial obtained from f_j by substituting 1 for x_i , then X_i is the closed subscheme of $\mathbb{A}^{m-1} = \text{Spec } k[x_1, \dots, \check{x}_i, \dots, x_m]$ defined by $f_1^{(i)}, \dots, f_l^{(i)}$.

For a projective scheme $X \subset \mathbb{P}^{m-1}$ defined by f_1, \dots, f_l , we suppose that every coherent sheaf of X (in particular, an invertible sheaf) is represented by a finitely generated module over $R = R_X$. In turn, every finitely generated R -module is represented by a matrix $A \in M_{r \times s}(R)$ which defines a free presentation of M ,

$$\bigoplus_{j=1}^s R(b_j) \xrightarrow{A} \bigoplus_{i=1}^r R(a_i) \rightarrow M \rightarrow 0.$$

Here maps are supposed to be degree-preserving and $R(a)$ denotes the graded free R -module of rank one defined by $R(a)_c = R_{a+c}$.

When two projective schemes $X \subset \mathbb{P}^{m-1}$ and $Y \subset \mathbb{P}^{n-1}$ are given, we can embed the product $X \times Y$ into \mathbb{P}^{mn-1} via the Segre embedding $\mathbb{P}^{m-1} \times \mathbb{P}^{n-1} \hookrightarrow \mathbb{P}^{mn-1}$. When we say that a morphism $f: X \rightarrow Y$ is given, we mean that its graph $\Gamma_f \subset X \times Y$ is given as a closed subscheme of \mathbb{P}^{mn-1} .

3. SEMI-DECIDABILITY OF THE ISOMORPHISM PROBLEM OF PROJECTIVE SCHEMES

In this section, we show the probably well-known fact that the isomorphism problem of projective schemes is semi-decidable; there exists an algorithm such that, when two projective schemes are given as an input, then the algorithm stops after finitely many steps if and only if these schemes are isomorphic. The algorithm given in this section is a very naive one and would be very inefficient. An approach via Iso schemes would give a more efficient algorithm (see Remark 6.7).

Remark 3.1. Poonen pointed out to the author that the isomorphism problem of finite-type k -schemes is also semi-decidable and it appears well-known. Roughly, the proof is by checking whether the given schemes have the “same” affine open coverings. Our proof below for projective schemes is more along our basic strategy in terms of graphs. Arguments in it will be repeated in computation of Iso schemes in Section 6.

Let $X \subset \mathbb{P}^{m-1}$ and $Y \subset \mathbb{P}^{n-1}$ be projective schemes. We first enumerate all the closed subschemes of $X \times Y$. To do so, we enumerate all the finite sequences f_1, \dots, f_l of homogeneous polynomials in $k[w_{ij} \mid 1 \leq i \leq m, 1 \leq j \leq n]$. For each positive integer i , let I_i be the ideal generated by the i -th sequence and let $Z_i \subset \mathbb{P}^{nm-1}$ be the closed subscheme corresponding to I_i . Thus we obtain the sequence $Z_i, i > 0$ of closed subschemes such that for every closed subscheme $Z \subset \mathbb{P}^{nm-1}$, there exists $i > 0$ such that $Z = Z_i$. For each i , we can check whether or not Z_i is included in $X \times Y$. Removing the ones not included in $X \times Y$, we can algorithmically produce every closed subscheme of $X \times Y$ one by one. If we prefer, we may remove redundancies to get a sequence where every closed subscheme of $X \times Y$ appears exactly once. We let $Z_i, i > 0$ be thus obtained sequence of closed subschemes of $X \times Y$.

Proposition 3.2. *The isomorphism problem of projective schemes is semi-decidable*

Proof. For each integer $i > 0$, from Lemma 3.3 below, we can algorithmically check whether Z_i is the graph of an isomorphism $X \xrightarrow{\sim} Y$. As soon as one finds that this is the case, we stop this algorithm. \square

Lemma 3.3. *We can algorithmically check whether or not a given closed subscheme $\Gamma \subset X \times Y$ is the graph of an isomorphism $X \xrightarrow{\sim} Y$.*

Proof. We need to check whether the two projections $\Gamma \rightarrow X$ and $\Gamma \rightarrow Y$ are both isomorphisms. We discuss only the former projection, denoting it by f . Let $X = \bigcup X_j$ be the standard affine open covering and let R_j be the coordinate ring of X_j . Let Γ_j be the preimage of X_j by the morphism $\Gamma \rightarrow X$, which is a closed subscheme of $\mathbb{P}_{R_j}^{n-1} = \text{Proj } R_j[y_1, \dots, y_n]$. The morphism $f: \Gamma \rightarrow X$ is an isomorphism if and only if every

$$f_j := f|_{\Gamma_j}: \Gamma_j \rightarrow X_j$$

is an isomorphism. We can compute the coherent sheaf Ω_{Γ_j/X_j} of differentials (see Remark 3.4) and check whether or not it is the zero sheaf. Thus we can algorithmically check whether or not f_j is unramified. If it is not unramified, then it is not an isomorphism. Suppose that f_j is unramified. Then it is also a finite morphism. We compute an R_j -module M_j such that $\widetilde{M}_j := (f_j)_* \mathcal{O}_{\Gamma_j}$ (see Remark 3.5). Consider the following three conditions:

- (1) $\text{Supp}(M_j) = X_j$.
- (2) $V(\text{Fitt}_1(M_j)) = \emptyset$, where $\text{Fitt}_1(M_j)$ denote the first Fitting ideal of M_j .
- (3) $\text{Tor}^{R_j}((R_j)_{\text{red}}, M_j \otimes (R_j)_{\text{red}}) = 0$.

The second condition means that for every point $x \in X_j$, the stalk $(\widetilde{M}_j)_x$ is generated by one element as an $\mathcal{O}_{X_j, x}$ -module. From [Har77, II, Exercise 5.8], the first two conditions together are equivalent to that $M_j \otimes (R_j)_{\text{red}}$ is a flat $(R_j)_{\text{red}}$ -module of constant rank one. Under these conditions, the third condition means that M_j is a flat R_j -module (see [Mat89, Th. 22.3] or [Sta21, tag 051C]). We conclude that these three conditions all hold if and only if the finite unramified morphism $f_j: \Gamma_j \rightarrow X_j$ is surjective and flat of constant rank one, that is, an isomorphism. \square

Remark 3.4 (cf. [Sti, Prop. 5.7]). Let R be a commutative ring and let

$$X = \text{Proj } R[x_1, \dots, x_r]/(f_1, \dots, f_l)$$

be a projective scheme over R , where f_1, \dots, f_l are homogeneous polynomials. Then the cotangent sheaf $\Omega_{X/R}$ is associated to the homology module of the sequence

$$\bigoplus_{i=1}^l S(-\deg f_i) \xrightarrow{(\partial f_i / \partial x_j)_{i,j}} S(-1)^{\oplus r} \xrightarrow{(x_1 \cdots x_r)} S.$$

Stillman's notes cited above treat the case where R is a field, but this lemma holds for an arbitrary R . This is a straightforward consequence of [Har77, Prop. 8.12 and Th. 8.13].

Remark 3.5. Let $X \subset \mathbb{P}_R^n$ be a projective scheme over a commutative ring R , let $\pi: X \rightarrow \text{Spec } R$ be the structure morphism and let \mathcal{M} be a coherent sheaf on X . Algorithms computing the pushforward $\pi_*\mathcal{M}$ (and more generally, higher direct images $R^i\pi_*\mathcal{M}$) are explained in [Smi98, ES08].

4. HILBERT SCHEMES

In this section, we discuss how to compute Hilbert schemes of general projective schemes and their universal families. Their computability, Proposition 4.4, has been already proved in [PTL15, Lem. 8.23]. As this result is the core of our approach, we explain it in more details below.

4.1. The Hilbert scheme of a projective space. Bayer [Bay82] explained how to compute equations defining the Hilbert scheme $\text{Hilb}_P(\mathbb{P}^{r-1})$ for each Hilbert polynomial P as a closed subset of a Grassmannian variety. It turned out that his equations also give the right scheme structure of the Hilbert scheme. We recall this description of the Hilbert scheme $\text{Hilb}_P(\mathbb{P}^{r-1})$, closely following the presentation by Iarrobino and Kleiman in [IK99, Appendix C] but with emphasis on algorithmic aspects.

Throughout this section, we fix a positive integer $r > 0$. Let $R := k[x_1, \dots, x_r] = \bigoplus_{i \geq 0} R_i$ with R_i denoting the degree- i part and let $\mathbb{P}^{r-1} = \text{Proj } R$, the $(r-1)$ -dimensional projective space. For a closed subscheme $Z \subset \mathbb{P}^{r-1}$ defined by a homogeneous ideal $I \subset R$, the *Hilbert polynomial* of Z is a polynomial $P \in \mathbb{Q}[t]$ such that $P(i) = \dim_k(R/I)_i$ for $i \gg 0$, where $(R/I)_i$ denotes the degree i part of the graded ring R/I . A polynomial is said to be a *Hilbert polynomial* if it is the Hilbert polynomial of some closed subscheme $Z \subset \mathbb{P}^{r-1}$. For each Hilbert polynomial P , the *Hilbert scheme* $\text{Hilb}_P(\mathbb{P}^{r-1})$ for P is the moduli scheme of closed subschemes $Z \subset \mathbb{P}^{r-1}$ with the Hilbert polynomial P .

For a Hilbert polynomial P , there exists a unique sequence of positive integers, $0 < a_0 \leq \dots \leq a_k$ such that $0 \leq k \leq r-2$ and

$$\binom{r+t-1}{r-1} - P(t) = \binom{t-a_0+r-1}{r-1} + \dots + \binom{t-a_k+r-1-k}{r-1-k}.$$

We can algorithmically compute these integers a_0, \dots, a_k from the polynomial P . The *Gotzmann number* $\varphi(P)$ of P is defined to be a_k .

We now fix a Hilbert polynomial P and an integer $d \geq \varphi(P)$. Let $r_d := \dim_k R_d$, $p := P(d)$ and $p^\vee := r_d - p$. Let $\text{Grass}_{p^\vee}(R_d)$ be the Grassmannian parameterizing p^\vee -dimensional subspaces of the r_d -dimensional vector space R_d . There exists a closed embedding

$$\begin{aligned} \text{Hilb}_P(\mathbb{P}^{r-1}) &\hookrightarrow \text{Grass}_{p^\vee}(R_d), \\ [Z] &\mapsto [(I_Z)_d] \end{aligned}$$

where $I_Z \subset R$ is the saturated ideal of Z and $(I_Z)_d$ is its degree- d part. In particular, the closed subscheme Z is recovered from the subspace $(I_Z)_d \subset R_d$. Indeed Z is defined by the ideal $((I_Z)_d) \subset R$ generated by $(I_Z)_d$.

Let M_d be the set of the monomials of degree d , which is a basis of R_d . The Grassmannian $\text{Grass}_{p^\vee}(R_d)$ has the standard affine open covering

$$\text{Grass}_{p^\vee}(R_d) = \bigcup_{\substack{K \subset M_d \\ \#K=p}} U_K.$$

Each affine chart U_K is isomorphic to the $p \cdot p^\vee$ -dimensional affine space $\mathbb{A}^{p \cdot p^\vee}$. In what follows, we identify M_d with $\{1, 2, \dots, r_d\}$ say by the lex order. Then, the affine chart $U_K = \mathbb{A}^{p \cdot p^\vee}$ is the space of r_d -by- p^\vee matrices

$$(4.1) \quad A = (a_{i,j})_{i \in M_d, 1 \leq j \leq p^\vee}$$

such that the p^\vee -by- p^\vee submatrix

$$(a_{i,j})_{i \in M_d \setminus K, 1 \leq j \leq p^\vee}$$

is the identity matrix. Note that the j -th column of A corresponds to the homogeneous polynomial $\sum_{i=1}^{r_d} a_{i,j} m_i$, where m_i denotes the i -th monomial in M_d . Thus we can write the coordinate ring of U_K as

$$k[U_K] = k[u_{i,j} \mid i \in K, 1 \leq j \leq p^\vee],$$

where $u_{i,j}$ are indeterminates corresponding to entries $a_{i,j}$ above, respectively. For example, if K consists of the last p monomials in M_d , then a matrix A as above is of the form:

$$A = \begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \\ a_{p^\vee+1,1} & \cdots & a_{p^\vee+1,p^\vee} \\ \vdots & \ddots & \vdots \\ a_{r_d,1} & \cdots & a_{r_d,p^\vee} \end{pmatrix}$$

The $p \cdot p^\vee$ free entries in the last p rows serve as coordinates of the affine space U_K . For general K , if we write $A = (\mathbf{a}_1 \cdots \mathbf{a}_{p^\vee})$ with column vectors \mathbf{a}_i , then the matrix A corresponds to the subspace $\langle \mathbf{a}_1, \dots, \mathbf{a}_{p^\vee} \rangle \subset R_d$. A matrix as above is also regarded as a linear map

$$k^{p^\vee} \rightarrow k^{r_d} = R_d.$$

For each i with $1 \leq i \leq r$, let B_i be the r_{d+1} -by- p^\vee matrix corresponding to the composite map

$$k^{p^\vee} \xrightarrow{A} R_d \xrightarrow{\times x_i} R_{d+1} = k^{r_{d+1}}.$$

These matrices are easily computed from A . Indeed their nonzero entries are the ones of A suitably arranged. Finally we define the r_{d+1} -by- $r p^\vee$ matrix

$$B := (B_1 \mid \cdots \mid B_r)$$

by lining B_i 's horizontally. If $V_A \subset R_d$ is the subspace corresponding to A , then the image of B regarded as the map $k^{r p^\vee} \rightarrow R_{d+1}$ is nothing but $R_1 \cdot V_A \subset R_{d+1}$. We have that the point $[V_A] \in \text{Grass}_{p^\vee}(R_d)$ lies in $\text{Hilb}_P(\mathbb{P}^{r-1})$ if and only if the inequality

$$\dim_k R_1 \cdot V_A \leq r_{d+1} - P(d+1) =: q^\vee$$

holds. Note that the inequality is equivalent to the equality, since the opposite inequality \geq always holds. Thus, on the affine chart $U_K \subset \text{Grass}_{p^\vee}(R_d)$, the Hilbert scheme $\text{Hilb}_P(\mathbb{P}^{r-1})$ is cut out, at least set-theoretically, by the $(q^\vee + 1)$ -by- $(q^\vee + 1)$ minors of the matrix B ; each such minor is a polynomial in coordinates $a_{i,j}$ of $U_K = \mathbb{A}^{p^\vee}$. It turns out that this is the case also scheme-theoretically. Precisely:

Proposition 4.1. *The closed subscheme $\text{Hilb}_P(\mathbb{P}^{r-1}) \cap U_K$ of $U_K = \mathbb{A}^{p^\vee}$ is defined by the $(q^\vee + 1)$ -by- $(q^\vee + 1)$ minors of the matrix B .*

4.2. The universal family. We can also compute the universal family

$$\mathcal{Z}_K \subset (\text{Hilb}_P(\mathbb{P}^{r-1}) \cap U_K) \times \mathbb{P}^{r-1}$$

over $\text{Hilb}_P(\mathbb{P}^{r-1}) \cap U_K$ as follows. Recall that $U_K = \mathbb{A}^{p^\vee}$ has coordinates $u_{i,j}$ ($i \in K$, $1 \leq j \leq p^\vee$) and consider the universal r_d -by- p^\vee matrix

$$(4.2) \quad \mathcal{A} = \mathcal{A}_K := (u_{i,j})_{i \in M_d, 1 \leq j \leq p^\vee}.$$

Here, for $i \in K$, the entry $u_{i,j}$ is the indeterminate $u_{i,j} \in k[U_K]$ and, for $i \in M_d \setminus K$, $u_{i,j}$ is defined to be either 1 or 0 so that the p^\vee -by- p^\vee matrix

$$(u_{i,j})_{i \in M_d \setminus K, 1 \leq j \leq p^\vee}$$

is the identity matrix. At each point $(a_{i,j}) \in U_K$, the universal matrix to the matrix A in (4.1). The j -th column of \mathcal{A} defines the universal j -th polynomial

$$h_{K,j} := \sum_{i=1}^{r_d} u_{i,j} m_i = n_i + \sum_{i \in K} u_{i,j} m_i \in k[U_K][x_1, \dots, x_r],$$

where m_i is the i -th monomial in M_d as before and n_i is the i -th monomial in $M_d \setminus K$. They are homogeneous of degree d with respect to variables x_1, \dots, x_r .

Proposition 4.2. *Let $I_K \subset k[U_K]$ be the defining ideal of $\text{Hilb}_P(\mathbb{P}^{r-1}) \cap U_K \subset U_K$, which can be computed as in Proposition 4.1. Then the universal family \mathcal{Z}_K is written as*

$$\mathcal{Z}_K = \text{Proj} \frac{(k[U_K]/I_K)[x_1, \dots, x_r]}{(h_{K,1}, \dots, h_{K,p^\vee})}.$$

4.3. The Hilbert scheme of a general projective scheme. Next we consider the Hilbert scheme $\text{Hilb}_P(X)$ of a projective scheme $X \subset \mathbb{P}_k^{r-1}$. This is the moduli scheme of those closed subschemes $Z \subset X$ that have the Hilbert polynomial P as a closed subscheme of \mathbb{P}_k^{r-1} . Let Q be the Hilbert polynomial of X . If $d \geq \varphi(Q)$, then I_X is d -regular. In particular, the truncated ideal $(I_X)_{\geq d}$, which is the part of the saturated ideal I_X with degree $\geq d$, is generated by the degree- d part $(I_X)_d$. Now we choose the integer d to be $\max\{\varphi(P), \varphi(Q)\}$. For a closed subscheme $Z \subset \mathbb{P}_k^{r-1}$ with the Hilbert polynomial P , we have $Z \subset X$ if and only if $(I_Z)_d \supset (I_X)_d$. Let

$$f_1 = \begin{pmatrix} f_{1,1} \\ \vdots \\ f_{r_d,1} \end{pmatrix}, \dots, f_l = \begin{pmatrix} f_{1,l} \\ \vdots \\ f_{r_d,l} \end{pmatrix} \in k^{r_d} = R_d$$

be a basis of $(I_X)_d$. Note that we can explicitly construct such a basis from the given finitely many defining polynomials of X . Firstly, there is an algorithm to compute the saturation $(I : (x_1, \dots, x_r)^\infty)$ of a homogeneous ideal $I \subset k[x_1, \dots, x_r]$, see [Eis95, page 360]. If g_1, \dots, g_m are thus computed generators of the saturated ideal I_X , which we may assume have degree $\leq d$ (since those of degree $> d$ are redundant), then $(I_X)_d$ is generated by elements of the form xg_i , where x is a monomial of degree $d - \deg g_i$. We can choose a basis from them in a standard linear algebra procedure. As before, a point of $U_K = \mathbb{A}^{p \cdot p^\vee}$ is identified with a r_d -by- p^\vee matrix A . To such a matrix, we associate the r_d -by- $(p^\vee + l)$ matrix $C_A = (A | f_1 | \dots | f_l)$. For example, if K is the first p monomials in M_d , then

$$C_A = \begin{pmatrix} 1 & & 0 & f_{1,1} & \cdots & f_{1,l} \\ & \ddots & & & & \\ 0 & & 1 & \vdots & \ddots & \vdots \\ a_{p^\vee+1,1} & \cdots & a_{p^\vee+1,p^\vee} & \vdots & & \vdots \\ \vdots & \ddots & \vdots & & & \\ a_{r_d,1} & \cdots & a_{r_d,p^\vee} & f_{r_d,1} & \cdots & f_{r_d,l} \end{pmatrix}.$$

For a closed subscheme $Z \subset \mathbb{P}^{r-1}$ with $[Z] \in U_K$, the following conditions are equivalent:

- (1) $(I_Z)_d \supset (I_X)_d$.
- (2) $\text{rank } C_A = p^\vee$.
- (3) all the $(p^\vee + 1)$ -by- $(p^\vee + 1)$ minors of C_A vanish.

The minors in the last condition are polynomials in the coordinates of U_K .

Proposition 4.3. *The closed subscheme*

$$(4.3) \quad \text{Hilb}_P(X)_K := \text{Hilb}_P(X) \cap U_K$$

of $U_K = \mathbb{A}^{p \cdot p^\vee}$ is defined by the defining polynomials of $\text{Hilb}_P(\mathbb{P}^{r-1})$ given in Proposition 4.1 and the above minors of C_A .

We can compute the universal family $\mathcal{Z}_{X,K} \subset \text{Hilb}_P(X)_K \times X$ over $\text{Hilb}_P(X)_K$ in a similar way as in the case of \mathcal{Z}_K .

If $I_{X,K} \subset k[U_K]$ denotes the defining ideal of $\text{Hilb}_P(X) \cap U_K$, then

$$(4.4) \quad \begin{aligned} \mathcal{Z}_{X,K} &= \text{Proj} \frac{(k[U_K]/I_{X,K})[x_1, \dots, x_r]}{(f_1, \dots, f_l, h_{K,1}, \dots, h_{K,p^\vee})} \\ &\subset \text{Proj} \frac{(k[U_K]/I_{X,K})[x_1, \dots, x_r]}{(f_1, \dots, f_l)} = \text{Hilb}_P(X)_K \times X. \end{aligned}$$

As a conclusion of the above computation, we have:

Proposition 4.4 ([PTL15, Lem. 8.23]). *There is an algorithm such that given a projective variety $X \subset \mathbb{P}^{r-1}$ and a Hilbert polynomial P , then it outputs a positive integer d and computes the closed subscheme $\text{Hilb}_P(X) \subset \text{Grass}_{p^\vee}(R_d)$ in terms of defining equations on each affine chart U_K together with defining equations of the universal families $\mathcal{Z}_{X,K} \subset \text{Hilb}_P(X)_K \times X$.*

4.4. **The action of GL_r .** Since it will be used to prove the decidability of projective equivalence in Section 5, we describe the natural action of the general linear group GL_r on the Hilbert scheme $\mathrm{Hilb}_P(\mathbb{P}^{r-1})$. The action of GL_r on \mathbb{P}^{r-1} induces an action of GL_r on $\mathrm{Hilb}_P(\mathbb{P}^{r-1})$. The Hilbert scheme has the open covering $\mathrm{Hilb}_P(\mathbb{P}^{r-1}) = \bigcup_K U_K$ and we have an explicit presentation of U_K for each K .

Definition 4.5. We define $V_{K,K'}$ to be the preimage of $U_{K'}$ by the morphism

$$\mu: \mathrm{GL}_r \times U_K \rightarrow \mathrm{Hilb}_P(\mathbb{P}^{r-1}).$$

We explain how to compute an explicit presentation of the affine scheme $V_{K,K'}$ as well as the morphism

$$(4.5) \quad \mu_{K,K'} := \mu|_{V_{K,K'}}: V_{K,K'} \rightarrow U_{K'}.$$

Let S_r be the coordinate ring of GL_r , that is, the localization $k[\underline{s}]_D$ of the polynomial ring $k[\underline{s}] = k[s_{i,j} \mid 1 \leq i, j \leq r]$ by the determinant $D = \det(s_{i,j})$. We have the universal matrix

$$\begin{pmatrix} s_{1,1} & \cdots & s_{1,r} \\ \vdots & \ddots & \vdots \\ s_{r,1} & \cdots & s_{r,r} \end{pmatrix} \in \mathrm{GL}_r(S_r).$$

The action $\mathrm{GL}_r \times \mathbb{A}^r \rightarrow \mathbb{A}^r$ is given by the following ring map:

$$\begin{aligned} k[x_1, \dots, x_r] &\rightarrow S_r[x_1, \dots, x_r] \\ x_i &\mapsto \sum_{j=1}^r s_{i,j} x_j \end{aligned}$$

As before, we fix $d \geq \varphi(P)$. We write a monomial in $k[x_1, \dots, x_r]$ as $x^e = x_1^{e_1} \cdots x_r^{e_r}$ with multi-index notation. The last map sends a monomial x^e of degree d to the polynomial,

$$\prod_{i=1}^d \left(\sum_{j=1}^r s_{i,j} x_j \right)^{e_i} =: \sum_{e'} \eta_{e,e'} x^{e'},$$

which is homogeneous of degree d both in x_1, \dots, x_r and in $s_{i,j}$. Here $\eta_{e',e}$ is a homogeneous polynomial of degree d in $k[\underline{s}]$. For each e , we can compute $\eta_{e,e'}$'s explicitly. We get the map

$$\begin{aligned} \mathrm{GL}_r &\rightarrow \mathrm{GL}_{r_d} \\ (a_{i,j})_{1 \leq i, j \leq r} &\mapsto (\eta_{e,e'}(\underline{a}))_{e, e' \in M_d}, \end{aligned}$$

which induces an action of GL_r on \mathbb{A}^{r_d} . In terms of coordinate rings, this is given by

$$S_{r_d} \rightarrow S_r, \quad s_{e,e'} \mapsto \eta_{e,e'}(s_{1,1}, \dots, s_{r,r}).$$

Let $K, K' \subset M_d$ be two subsets with $\sharp K = \sharp K' = p$. Let $\mathcal{A}_K = (u_{i,j})$ be the universal r_d -by- p^\vee matrix for K (see (4.2)). Compute the matrix product

$$\mathcal{B} := (\eta_{e,e'}) \cdot \mathcal{A}_K \in \mathrm{M}_{r_d \times p^\vee}(k(\underline{s})[\underline{u}])$$

with $k(\underline{s})$ the fraction field of $k[\underline{s}]$ and $k(\underline{s})[\underline{u}] = k(\underline{s})[u_{i,j} \mid i \in K, 1 \leq j \leq p^\vee]$. By Gaussian elimination in the field $k(\underline{s}, \underline{u})$, we can algorithmically and uniquely transform

\mathcal{B} to a matrix belonging to $U_{K'}(k(\underline{s}, \underline{u}))$ by applying elementary column operations finitely many times; let $\theta = (\theta_{i,j}) \in M_{r_d \times p^\vee}(k(\underline{s}, \underline{u}))$ be the resulting matrix. Here the p^\vee -by- p^\vee matrix $(\theta_{i,j})_{i \in M_d \setminus K', 1 \leq j \leq r_d}$ is the identity matrix. We get $p \cdot p^\vee$ rational functions $\theta_{i,j} \in k(\underline{s}, \underline{u})$ for $i \in K'$ and $1 \leq j \leq p^\vee$. We see that the rational map $\mathrm{GL}_r \times U_K \dashrightarrow U_{K'}$ is given by:

$$\begin{aligned} k[\underline{u}] &\rightarrow k(\underline{s}, \underline{u}) \\ u_{i,j} &\mapsto \theta_{i,j} \end{aligned}$$

Let $V_{K,K'} \subset \mathrm{GL}_r \times U_K$ be the domain of this rational map, that is, the preimage of $U_{K'}$ by the map $\mathrm{GL}_r \times U_K \rightarrow \mathrm{Hilb}_P(\mathbb{P}^{r-1})$. This is an affine scheme, since $\mathrm{GL}_r \times U_K$ and $U_{K'}$ are affine and $\mathrm{Hilb}_P(\mathbb{P}^{r-1})$ is separated. The coordinate ring $T_{K,K'}$ of $V_{K,K'}$ is obtained by adjoining $\theta_{i,j}$'s to $S_r[\underline{u}]$.

5. PROJECTIVE EQUIVALENCE

Two projective schemes X and Y embedded in the same projective space \mathbb{P}^{r-1} are said to be *projectively equivalent* if there exists an invertible matrix $g \in \mathrm{GL}_r(k)$ such that $g(X) = Y$.

Proposition 5.1. *Let X and Y be projective schemes embedded in the same projective space \mathbb{P}^{r-1} . Then we can algorithmically check whether X and Y are projectively equivalent.*

Proof. We follow the notation of Section 4. We can compute the Hilbert polynomials of X and Y (see [Eis95, Sections 15.1.1 and 15.10.2]) and check whether they are the same. If they are different, then there is no $g \in \mathrm{GL}_r(k)$ as in the proposition. Suppose that they are the same and denote it by P . Let $d := \varphi(P)$, the Gotzmann number of P , let I_X and I_Y be the saturated ideals of X and Y and let $(I_X)_d$ and $(I_Y)_d$ be their degree- d parts. We compute bases of $(I_X)_d$ and $(I_Y)_d$, which are represented by r_d -by- p^\vee matrices A_X and A_Y respectively. We then compute their reduced column echelon forms and denote them by B_X and B_Y . Let K and K' be the set of indices such that the corresponding rows of B_X and B_Y have pivots. Then B_X and B_Y define the points $[X] \in U_K$ and $[Y] \in U_{K'}$ respectively. Consider the morphism $\mu_{K,K'}: V_{K,K'} \rightarrow U_{K'}$ (see (4.5)) and the morphism

$$p: V_{K,K'} \hookrightarrow \mathrm{GL}_r \times U_K \xrightarrow{\text{projection}} U_K$$

of affine schemes. We compute the closed subset

$$(\mu_{K,K'})^{-1}([Y]) \cap p^{-1}([X]) \subset V_{K,K'},$$

which is the set of pairs $(g, [X])$ such that $g \in \mathrm{GL}_r(k)$ and $g(X) = Y$. Thus, there exists $g \in \mathrm{GL}_r(k)$ as in the proposition if and only if this closed subset is not empty. \square

6. HOM SCHEMES AND ISO SCHEMES

For projective schemes X and Y , the *Hom scheme*, denoted by $\underline{\mathrm{Hom}}(X, Y)$, and *Iso scheme*, denoted by $\underline{\mathrm{Iso}}(X, Y)$, are the moduli schemes of morphisms $X \rightarrow Y$ and

isomorphisms $X \rightarrow Y$ respectively. From [Kol96, p. 16], we have an open immersion

$$\begin{aligned} \underline{\text{Hom}}(X, Y) &\rightarrow \text{Hilb}(X \times Y), \\ [f: X \rightarrow Y] &\mapsto [\Gamma_f] \end{aligned}$$

where $\Gamma_f \subset X \times Y$ is the graph of f . Namely, we can identify $\underline{\text{Hom}}(X, Y)$ with the locus of points $[Z] \in \text{Hilb}(X \times Y)$ such that the first projection $Z \rightarrow X$ is an isomorphism. Similarly we can identify $\underline{\text{Iso}}(X, Y)$ with the locus of points $[Z]$ where both the projections $Z \rightarrow X$ and $Z \rightarrow Y$ are isomorphisms. Thus, if we embed also $\underline{\text{Hom}}(Y, X)$ into $\text{Hilb}(X \times Y)$ via the obvious isomorphism $\text{Hilb}(X \times Y) \cong \text{Hilb}(Y \times X)$, then we have

$$\underline{\text{Iso}}(X, Y) = \underline{\text{Hom}}(X, Y) \cap \underline{\text{Hom}}(Y, X).$$

We now fix embeddings $X \subset \mathbb{P}^{m-1}$ and $Y \subset \mathbb{P}^{n-1}$ and embed the product $X \times Y$ into \mathbb{P}^{mn-1} by the Segre embedding. Then the Hilbert scheme $\text{Hilb}(X \times Y)$ decomposes into the disjoint union of countably many open and closed subschemes as

$$\text{Hilb}(X \times Y) = \coprod_P \text{Hilb}_P(X \times Y).$$

Here P runs over Hilbert polynomials. Recall that $\text{Hilb}_P(X \times Y)_K$ denotes $\text{Hilb}_P(X \times Y) \cap U_K$, see (4.3).

Definition 6.1. We define

$$\begin{aligned} \underline{\text{Hom}}_P(X, Y) &:= \underline{\text{Hom}}(X, Y) \cap \text{Hilb}_P(X \times Y), \\ \underline{\text{Hom}}_P(X, Y)_K &:= \underline{\text{Hom}}(X, Y) \cap \text{Hilb}_P(X \times Y)_K, \\ \underline{\text{Iso}}_P(X, Y) &:= \underline{\text{Iso}}(X, Y) \cap \text{Hilb}_P(X \times Y), \\ \underline{\text{Iso}}_P(X, Y)_K &:= \underline{\text{Iso}}(X, Y) \cap \text{Hilb}_P(X \times Y)_K. \end{aligned}$$

We have the open coverings,

$$\begin{aligned} \underline{\text{Hom}}_P(X, Y) &= \bigcup_K \underline{\text{Hom}}_P(X, Y)_K, \\ \underline{\text{Iso}}_P(X, Y) &= \bigcup_K \underline{\text{Iso}}_P(X, Y)_K. \end{aligned}$$

Definition 6.2. We define the *Hilbert polynomial* of an isomorphism $f: X \rightarrow Y$ to be the Hilbert polynomial of its graph $\Gamma_f \subset X \times Y$ as a closed subscheme of \mathbb{P}^{mn-1} . Namely the Hilbert polynomial of f is the polynomial P such that $[f] \in \underline{\text{Iso}}_P(X, Y)$.

When we show the decidability of the isomorphism problem for several classes of projective schemes, our strategy will be to construct finitely many polynomials P_1, \dots, P_l from given projective schemes X and Y that satisfy the following property; if X and Y are isomorphic, then there exists an isomorphism $f: X \rightarrow Y$ having the Hilbert polynomial P_i for some $i \in \{1, \dots, l\}$. Namely X and Y are isomorphic if and only if $\bigcup_{i=1}^l \underline{\text{Iso}}_{P_i}(X, Y) \neq \emptyset$. Then what remains to do is to compute $\underline{\text{Iso}}_{P_i}(X, Y)$ for every $i \in \{1, \dots, l\}$.

For each P and K , we have

$$\underline{\text{Iso}}_P(X, Y)_K = \underline{\text{Hom}}_P(X, Y)_K \cap \underline{\text{Hom}}_P(Y, X)_K.$$

Thus, computation of $\underline{\text{Iso}}_P(X, Y)_K$ is reduced to the one of $\underline{\text{Hom}}_P(X, Y)_K$. We focus on the latter computation in what follows. Replacing X in (4.4) with $X \times Y$, we get explicit presentation of the universal family

$$\mathcal{Z}_{X \times Y, K} \subset \text{Hilb}_P(X \times Y)_K \times X \times Y.$$

From discussion above, the open subscheme $\underline{\text{Hom}}_P(X, Y)_K$ of $\text{Hilb}_P(X \times Y)_K$ is the largest open subscheme U such that the composite morphism

$$g: \mathcal{Z}_{X \times Y, K} \hookrightarrow \text{Hilb}_P(X \times Y)_K \times X \times Y \xrightarrow{\text{projection}} \text{Hilb}_P(X \times Y)_K \times X$$

is an isomorphism over $U \times X$. Since we are given the embedding $X \subset \mathbb{P}^{m-1}$, we have the standard affine open covering $X = \bigcup_{i=1}^m X_i$. Let

$$g_i: g^{-1}(X_i) \rightarrow \text{Hilb}_P(X \times Y)_K \times X_i$$

be restriction of g . This is a projective morphism with the target being affine. Let $U_i \subset \text{Hilb}_P(X \times Y)_K$ be the largest open subset such that g_i is an isomorphism over $U_i \times X_i$. Then

$$U = \underline{\text{Hom}}_P(X, Y)_K = \bigcap_{i=1}^m U_i.$$

There exists an algorithm to compute each U_i :

Proposition 6.3. *Let H and X be affine schemes and let $f: Z \rightarrow H \times X$ be a projective morphism. Then there exists an algorithm to compute the largest open subset $U \subset H$ such that f is an isomorphism over $U \times X$.*

Proof. We first compute the closed subset $C_1 := \text{Supp}(f_*\Omega_{Z/H \times X})$ of $H \times X$ (see Remarks 3.4 and 3.5). Next we compute the closed subset $C_2 := (H \times X) \setminus V$, where V is the invertible locus of $f_*\mathcal{O}_Z$ (see Lemma 6.4). The desired open subset $U \subset H$ is $H \setminus p_H(C_1 \cup C_2)$, where p_H denotes the projection $H \times X \rightarrow H$. Indeed, obviously $H \setminus p_H(C_1 \cup C_2)$ is contained in the desired subset. On the other hand, f is unramified (in particular, finite and affine) over $H \setminus p_H(C_1)$. That $f_*\mathcal{O}_Z$ is invertible on $H \setminus p_H(C_1 \cup C_2)$ means that f is an isomorphism over this open subset. \square

Lemma 6.4. *There is an algorithm to compute the invertible locus of a coherent sheaf on an affine scheme, that is, the largest open subset on which the sheaf is invertible.*

Proof. Let X be an affine scheme and let \mathcal{M} be a coherent sheaf on X . We first compute the closed subset

$$C_1 := \overline{X \setminus \text{Supp}(\mathcal{M})} \subset X$$

(see Remark 6.5). Its complement $X \setminus C_1$ is the largest open subset of X that is included in $\text{Supp}(\mathcal{M})$. We then put $\mathcal{M}' := \mathcal{M}|_{X_{\text{red}}}$ and compute the closed subset

$$C_2 := V(\text{Fitt}_1(\mathcal{M}')),$$

where Fitt_1 denotes the first Fitting ideal. Its complement $X \setminus C_2$ is the locus of points $x \in X$ where \mathcal{M}'_x is generated by one element as an $\mathcal{O}_{X, x}$ -module. Finally we compute

$$C_3 := \text{Supp}(\mathcal{T}or^{\mathcal{O}_X}(\mathcal{O}_{X_{\text{red}}}, \mathcal{M})).$$

From [Mat89, Th. 22.3], its complement $X \setminus C_3$ is the locus where $\mathcal{M}|_X$ is flat. Now the open subset

$$U := X \setminus (C_1 \cup C_2 \cup C_3)$$

is the largest open subset such that

- $\text{Supp}(\mathcal{M}|_U) = U$,
- for every $x \in U$, \mathcal{M}_x is generated by one element as an $\mathcal{O}_{X,x}$ -module,
- $\mathcal{M}|_U$ is a flat \mathcal{O}_U -module.

Therefore, U is the desired open subset. We can compute C_1 , C_2 and C_3 by some algorithms, for example, ones implemented to Macaulay2 [GS]. \square

Remark 6.5. Let $X = \text{Spec } R$ and let $C = V(f_1, \dots, f_n) \subset X$ be a closed subset. Then the closed subset $\overline{X \setminus C}$ is defined by the ideal

$$\bigcap_{i=1}^n \text{Ker}(R \rightarrow R_{f_i}).$$

Indeed, $X \setminus C$ is covered by the affine open subsets $U_i = \{f_i \neq 0\}$ and we have $\overline{X \setminus C} = \bigcup \overline{U_i}$. Each $\overline{U_i}$ is defined by the ideal $\text{Ker}(R \rightarrow R_{f_i})$.

We conclude:

Proposition 6.6. *For each Hilbert polynomial P , we can explicitly compute open subschemes $\underline{\text{Hom}}_P(X, Y)$ and $\underline{\text{Iso}}_P(X, Y)$ of $\text{Hilb}_P(X \times Y)$ by means of explicit presentation of open subsets $\underline{\text{Hom}}_P(X, Y)_K$ and $\underline{\text{Iso}}_P(X, Y)_K$ of $\text{Hilb}_P(X \times Y)_K$ for each K .*

Remark 6.7. Using Iso schemes, we can give an alternative proof of the semi-decidability of the isomorphism, which was proved in Section 3. We enumerate all the Hilbert polynomials as P_i , $i \in \mathbb{Z}_{>0}$. For each $i > 0$, we compute the Iso scheme $\underline{\text{Iso}}_{P_i}(X, Y)$. We stop if we get a non-empty Iso scheme.

7. ONE-DIMENSIONAL SCHEMES

In this section, we show that the isomorphism problem for one-dimensional projective schemes and the one for one-dimensional reduced quasi-projective schemes are decidable. This generalizes the known case of smooth irreducible curves ([BGJGP05, Lem. 5.1] for the case of genus $\neq 1$ and the MathOverflow thread mentioned in Introduction for elliptic curves). We need the following version of the Riemann-Roch formula for one-dimensional projective schemes.

Proposition 7.1 ([Vak15, Exercise 18.4.S]). *Let X be a one-dimensional projective scheme and let X_i , $1 \leq i \leq l$, be its one-dimensional irreducible components given with reduced structure and let η_i be the generic point of X_i . Let \mathcal{L} be an invertible sheaf on X and let \mathcal{F} be a coherent sheaf on X . Then*

$$(7.1) \quad \chi(\mathcal{F} \otimes \mathcal{L}) - \chi(\mathcal{F}) = \sum_{i=1}^l \text{length}(\mathcal{F}_{\eta_i}) \deg(\mathcal{L}|_{X_i}).$$

Here $\text{length}(\mathcal{F}_{\eta_i})$ is the length of \mathcal{F}_{η_i} as an \mathcal{O}_{X,η_i} -module. In particular,

$$(7.2) \quad \chi(\mathcal{L}) - \chi(\mathcal{O}) = \sum_{i=1}^l \text{length}(\mathcal{O}_{X,\eta_i}) \deg(\mathcal{L}|_{X_i}).$$

Proof. The outline of the proof is written in [Vak15]. For the sake of completeness, we write it down in more details. We first observe that zero-dimensional connected components of X do not contribute to either side of (7.1). Therefore we may suppose that X has only one-dimensional irreducible components, that is, $X = \bigcup_{i=1}^l X_i$. Note also that both sides of (7.1) are also additive for short exact sequences; if $v(\mathcal{F})$ denotes either side of the equality, for a short exact sequence of coherent \mathcal{O}_X -modules,

$$0 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_2 \rightarrow \mathcal{F}_3 \rightarrow 0,$$

we have $v(\mathcal{F}_2) = v(\mathcal{F}_1) + v(\mathcal{F}_3)$. This implies that for a filtration of coherent sheaves,

$$\mathcal{F} = \mathcal{F}_0 \supset \mathcal{F}_1 \supset \cdots \supset \mathcal{F}_n = 0,$$

we have

$$(7.3) \quad v(\mathcal{F}) = \sum_i v(\mathcal{F}_i/\mathcal{F}_{i+1}).$$

Let $\mathcal{I} \subset \mathcal{O}_X$ be the defining ideal sheaf of the associated reduced scheme X_{red} of X , which is necessarily nilpotent. We apply equality (7.3) to the filtration

$$\mathcal{F} \supset \mathcal{I}\mathcal{F} \supset \mathcal{I}^2\mathcal{F} \supset \cdots \supset \mathcal{I}^n\mathcal{F} = 0.$$

Thus it suffices to show (7.1) for sheaves $\mathcal{I}^i\mathcal{F}/\mathcal{I}^{i+1}\mathcal{F}$. Since they are $\mathcal{O}_{X_{\text{red}}}$ -modules, in turn, it suffices to show the proposition in the case where X is reduced. Let us now write $\mathcal{L} = \mathcal{O}_X(\sum_{j=1}^m n_j p_j)$, where n_i are integers and p_i are closed points of X at which X is smooth and \mathcal{F} is locally free. We prove (7.1) in this situation by induction on $n := \sum_{j=1}^m |n_j|$. If $n = 0$, this is obvious. If $n = 1$, then \mathcal{L} is either $\mathcal{O}_X(p)$ or $\mathcal{O}_X(-p)$. In the latter case, from the exact sequence

$$0 \rightarrow \mathcal{F} \otimes \mathcal{L} \rightarrow \mathcal{F} \rightarrow \mathcal{F}|_p \rightarrow 0,$$

we have

$$\chi(\mathcal{F} \otimes \mathcal{L}) - \chi(\mathcal{F}) = -\chi(\mathcal{F}|_p) = -\text{length}(\mathcal{F}_{\eta_i}),$$

where i is such that $p \in X_i$. If $\mathcal{L} = \mathcal{O}_X(p)$ and if we put $\mathcal{F}' := \mathcal{F} \otimes \mathcal{L}$, then

$$\chi(\mathcal{F} \otimes \mathcal{L}) - \chi(\mathcal{F}) = -(\chi(\mathcal{F}' \otimes \mathcal{O}_X(-p)) - \chi(\mathcal{F}')) = \text{length}(\mathcal{F}_{\eta_i}).$$

Thus (7.1) holds when $n = 1$. For general $n \geq 1$, if we write $\mathcal{L} = \mathcal{L}' \otimes \mathcal{L}''$ with \mathcal{L}' and \mathcal{L}'' having smaller n , we have

$$\begin{aligned} \chi(\mathcal{F} \otimes \mathcal{L}) - \chi(\mathcal{F}) &= (\chi((\mathcal{F} \otimes \mathcal{L}') \otimes \mathcal{L}'') - \chi(\mathcal{F} \otimes \mathcal{L}')) + (\chi(\mathcal{F} \otimes \mathcal{L}') - \chi(\mathcal{F})) \\ &= \sum_{i=1}^l \text{length}(\mathcal{F}_{\eta_i}) \deg(\mathcal{L}'|_{X_i}) + \sum_{i=1}^l \text{length}(\mathcal{F}_{\eta_i}) \deg(\mathcal{L}''|_{X_i}) \\ &= \sum_{i=1}^l \text{length}(\mathcal{F}_{\eta_i}) \deg(\mathcal{L}|_{X_i}). \end{aligned}$$

□

Consider two one-dimensional projective schemes $X \subset \mathbb{P}^{m-1}$ and $Y \subset \mathbb{P}^{n-1}$, which have very ample invertible sheaves \mathcal{L} and \mathcal{M} corresponding to the given embeddings to projective spaces respectively. Suppose that there exists an isomorphism $f: X \rightarrow Y$. We will bound possibilities for the Euler characteristic of $\mathcal{L} \otimes f^*\mathcal{M}$ without using data of f .

Corollary 7.2. *Let X_i , $1 \leq i \leq l$ and Y_j , $1 \leq j \leq m$ be the one-dimensional irreducible components of X and Y respectively. We give them with reduced structure. Let η_i be the generic point of X_i . Let $d := \sum_{j=1}^m \deg(\mathcal{M}|_{Y_j})$. Then there exists a partition of d into positive integers, $d = \sum_{i=1}^l d_i$, such that*

$$\chi(\mathcal{L} \otimes f^*\mathcal{M}) = \chi(\mathcal{O}_X) + \sum_{i=1}^l \text{length}(\mathcal{O}_{X,\eta_i}) (\deg(\mathcal{L}|_{X_i}) + d_i).$$

Proof. We put $d_i := \deg(f^*\mathcal{M}|_{X_i})$ and apply the second equality in Proposition 7.1 with $\mathcal{L} \otimes f^*\mathcal{M}$ in place of \mathcal{L} . □

The Hilbert polynomial of f (see Definition 6.2) is equal to the Hilbert polynomial of X with respect to the very ample sheaf $\mathcal{L} \otimes f^*\mathcal{M}$. It is the polynomial $P(t)$ of degree at most one such that

$$P(0) = \chi(\mathcal{O}_X) \text{ and } P(1) = \chi(\mathcal{L} \otimes f^*\mathcal{M}).$$

Theorem 7.3. *The isomorphism problem for one-dimensional projective schemes is decidable.*

Proof. Let $X \subset \mathbb{P}^{m-1}$ and $Y \subset \mathbb{P}^{n-1}$ be one-dimensional projective schemes. We compute their one-dimensional irreducible components with reduced structure X_i , $1 \leq i \leq l$ and Y_j , $1 \leq j \leq m$ respectively; there exist algorithms to compute associated reduced schemes and (geometric) irreducible components (see [EHV92, Chi86]). Then we compute $d = \sum_{j=1}^m \deg(\mathcal{M}|_{Y_j})$.

For each partition

$$\lambda: d = d_1 + \cdots + d_l$$

of d into l positive integers, we compute

$$e_\lambda := \chi(\mathcal{O}_X) + \sum_{i=1}^l \text{length}(\mathcal{O}_{X,\eta_i}) (\deg(\mathcal{L}|_{X_i}) + d_i)$$

and define the polynomial

$$P_\lambda(t) := (e_\lambda - \chi(\mathcal{O}_X))t + \chi(\mathcal{O}_X).$$

The polynomials P_λ are the only potential Hilbert polynomials for an isomorphism $f: X \rightarrow Y$ if any. For each λ , we compute $\underline{\text{Iso}}_{P_\lambda}(X, Y)$. If one of them is non-empty, then X and Y are isomorphic. Otherwise, they are not isomorphic. □

Theorem 7.4. *The isomorphism problem for one-dimensional reduced quasi-projective schemes is decidable. Here we suppose that each quasi-projective scheme X is given an embedding $X \hookrightarrow \mathbb{P}^{m-1}$ and represented by two projective schemes $\overline{X} \subset \mathbb{P}^{m-1}$, the closure of X in \mathbb{P}^{m-1} , and $\overline{X} \setminus X \subset \mathbb{P}^{m-1}$.*

Proof. Let $X \subset \mathbb{P}^{m-1}$ and $Y \subset \mathbb{P}^{n-1}$ be quasi-projective one-dimensional reduced schemes and let $\overline{X} \subset \mathbb{P}^{m-1}$ and $\overline{Y} \subset \mathbb{P}^{n-1}$ be their closures respectively. If \overline{X} is singular at some point of $\overline{X} \setminus X$, then we resolve this singularity by repeating blowups. Note that a blowup of \mathbb{P}^{m-1} at a point is a closed subvariety of $\mathbb{P}^{m-1} \times \mathbb{P}^{m-2}$ and hence one of $\mathbb{P}^{m(m-1)-1}$ by the Segre embedding. Thus a blowup of \overline{X} at a point has an embedding into $\mathbb{P}^{m(m-1)-1}$, which can be explicitly computed. Therefore we can replace the embedding $X \subset \mathbb{P}^{m-1}$ so that \overline{X} becomes smooth at every point of $\overline{X} \setminus X$. Similarly for Y . If $\overline{X} \setminus X$ and $\overline{Y} \setminus Y$ have different numbers of points, then X and Y are not isomorphic. Thus, we may suppose that they have the same number of points. Now X and Y are isomorphic if and only if there exists an isomorphism $f: \overline{X} \rightarrow \overline{Y}$ such that $\overline{Y} \setminus Y \subset f(\overline{X} \setminus X)$. From the assumption which we just put, $\overline{Y} \setminus Y \subset f(\overline{X} \setminus X)$ implies $\overline{Y} \setminus Y = f(\overline{X} \setminus X)$. Following the algorithm described in the proof of Theorem 7.3, we can compute the Iso scheme

$$\underline{\text{Iso}}(\overline{X}, \overline{Y}) \left(= \bigcup_{\lambda} \underline{\text{Iso}}_{P_{\lambda}}(\overline{X}, \overline{Y}) \right),$$

where λ runs over partitions of a positive integer d as in the proof of Theorem 7.3. In particular, since there are only finitely many partitions, we can algorithmically compute this Iso scheme. Recall that the Iso scheme is by definition a subscheme of the Hilbert scheme $\text{Hilb}(\overline{X} \times \overline{Y})$. Let

$$\mathcal{U} \subset \underline{\text{Iso}}(\overline{X}, \overline{Y}) \times \overline{X} \times \overline{Y}$$

be the universal family. Let $A \subset \mathcal{U}$ be the preimage of $\overline{X} \setminus X$ by the projection $\mathcal{U} \rightarrow \overline{X}$. Let us write $\overline{Y} \setminus Y = \{y_1, \dots, y_n\}$ and let $B_1, \dots, B_n \subset \mathcal{U}$ be the preimages of y_1, \dots, y_n by the projection $\mathcal{U} \rightarrow \overline{Y}$ respectively. Let $\pi: \mathcal{U} \rightarrow \underline{\text{Iso}}(\overline{X}, \overline{Y})$ be the projection. For each i , we claim that

$$\pi(A \cap B_i) = \{[f] \in \underline{\text{Iso}}(\overline{X}, \overline{Y}) \mid y_i \in f(\overline{X} \setminus X)\}.$$

Indeed, $\pi^{-1}([f])$ is identical to the graph $\Gamma_f \subset \overline{X} \times \overline{Y}$ and we have

$$\pi^{-1}([f]) \cap A \cap B_i = \{(x, y_i) \in \overline{X} \times \overline{Y} \mid x \in \overline{X} \setminus X\}.$$

The last set is non-empty if and only if $y_i \in f(\overline{X} \setminus X)$. This shows the above claim.

Thus $\bigcap_{i=1}^n \pi(A \cap B_i)$ is exactly the locus of isomorphisms $f: \overline{X} \rightarrow \overline{Y}$ with $\overline{Y} \setminus Y \subset f(\overline{X} \setminus X)$. We compute this closed subset $\bigcap_{i=1}^n \pi(A \cap B_i)$ and check whether this is empty. The given quasi-projective schemes X and Y are isomorphic if and only if this is not empty. \square

8. VARIETIES WITH A BIG CANONICAL SHEAF OR A BIG ANTI-CANONICAL SHEAF

In this section, we show the decidability of the isomorphism problem for varieties as in the title, generalizing the case of general type solved by Totaro (see [Poo14, Rem.

12.3]). The key ingredients are computation of Iso schemes and the Kodaira vanishing theorem.

Theorem 8.1. *Let X and Y be smooth irreducible projective varieties. Suppose that either ω_X or ω_X^{-1} is big. Then we can algorithmically decide whether X and Y are isomorphic.*

Proof. It is enough to consider the case where X and Y have the same dimension $d > 0$. We may also suppose that either both ω_X and ω_Y are big or both ω_X^{-1} and ω_Y^{-1} are big. For, otherwise, X and Y are not isomorphic. We denote these big sheaves by \mathcal{B}_X and \mathcal{B}_Y respectively. Note that we can algorithmically decide which of ω_X and ω_X^{-1} is big by checking the birationality of maps $\Phi_{\omega_X^{\otimes n}}$, $n \in \mathbb{Z}$ in turn (see Section 10.4). Let \mathcal{L} and \mathcal{M} be the very ample invertible sheaves on X and Y respectively corresponding to the given embeddings into projective spaces. We compute the least positive integer e such that $\mathcal{L}^{\otimes e} \otimes \omega_X^{-1}$ is ample (see Section 10.3). We replace \mathcal{L} with $\mathcal{L}^{\otimes e}$, which amounts to replacing the embedding $X \hookrightarrow \mathbb{P}^{m-1}$ with the one obtained by the e -uple Veronese embedding $\mathbb{P}^{m-1} \hookrightarrow \mathbb{P}^{\binom{m-1+e}{m-1}-1}$. Now $\mathcal{L} \otimes \omega_X^{-1}$ is ample. In this situation, we will algorithmically output finitely many polynomials Q_1, \dots, Q_c such that the Hilbert polynomial of every isomorphism $f: X \rightarrow Y$ (if any) is one of them. To do so, we first note that for every positive integer l , $(\mathcal{L} \otimes f^*\mathcal{M})^{\otimes l} \otimes \omega_X^{-1}$ is ample. From the Kodaira vanishing, we have

$$H^i(X, (\mathcal{L} \otimes f^*\mathcal{M})^{\otimes l}) = 0 \quad (i > 0).$$

Hence the Hilbert polynomial P_f of f satisfies

$$P_f(l) = \chi(X, (\mathcal{L} \otimes f^*\mathcal{M})^{\otimes l}) = h^0(X, (\mathcal{L} \otimes f^*\mathcal{M})^{\otimes l}).$$

Here h^0 means the dimension of H^0 . Then we compute the least positive integer q such that $\mathcal{B}_X^{\otimes q} \otimes \mathcal{L}^{-1}$ and $\mathcal{B}_Y^{\otimes q} \otimes \mathcal{M}^{-1}$ are both effective. Then, there exists an injection

$$0 \rightarrow (\mathcal{L} \otimes f^*\mathcal{M})^{\otimes l} \rightarrow \mathcal{B}_X^{\otimes 2ql},$$

which implies

$$P_f(l) = h^0(X, (\mathcal{L} \otimes f^*\mathcal{M})^{\otimes l}) \leq h^0(X, \mathcal{B}_X^{\otimes 2ql}).$$

Note that the obtained upper bound $h^0(X, \mathcal{B}_X^{\otimes 2ql})$ of $P_f(l)$ is independent of the isomorphism $f: Y \rightarrow X$. In general, a polynomial

$$h(t) = a_d t^d + \dots + a_0$$

of degree d is determined by its values at $d + 1$ distinct points t_0, \dots, t_d , by Lagrange interpolation. We compute $h^0(X, \mathcal{B}_X^{\otimes 2ql})$ for $1 \leq l \leq d + 1$. For each tuple $\lambda = (\lambda_1, \dots, \lambda_{d+1})$ of nonnegative integers with $\lambda_i \leq h^0(X, \mathcal{B}_X^{\otimes 2q_i})$, we compute the polynomial $Q_\lambda(t)$ such that $Q_\lambda(i) = \lambda_i$. Thus obtained finitely many polynomials Q_λ are the desired ones. For each λ , we compute $\text{Iso}_{Q_\lambda}(X, Y)$ and check whether it is empty or not. If one of them is not empty, then X and Y are isomorphic. Otherwise, they are not isomorphic. \square

9. COMPUTING INTERSECTION NUMBERS

The aim of this section is to prove the following proposition.

Proposition 9.1 ([PTL15, Lem. 8.7]). *For a smooth irreducible projective variety X , an irreducible closed subset $Z \subset X$ of codimension c and an invertible sheaf \mathcal{L} on X , we can algorithmically compute the intersection number $Z \cdot \mathcal{L}^{\dim Z} \in \mathbb{Z}$.*

This will be used in Section 10 to discuss decidability of various positivity properties of invertible sheaves. The proof in [PTL15] uses étale cohomology (in fact, its authors considered, more generally, the intersection number of cycles of complementary dimensions). We give an alternative proof using Simpson's algorithm [Sim08, Section 2.5] to compute singular cohomology. We only consider the case where the ambient variety X is smooth, as Simpson's algorithm is valid only for smooth varieties. According to his algorithm, for a smooth projective variety $X \subset \mathbb{P}^{r-1}$, we can compute a finite simplicial complex $\mathcal{H}(X)$ in $\mathbb{P}^{r-1}(\mathbb{C})$ which is homotopy equivalent to $X(\mathbb{C})$. In particular, we can compute the singular homology $H_i(X(\mathbb{C}), \mathbb{Z})$ and cohomology $H^i(X(\mathbb{C}), \mathbb{Z})$ using $\mathcal{H}(X)$. Their elements are represented by simplicial i -chains and i -cochains on $\mathcal{H}(X)$ respectively.

When we have a morphism $f: Z \rightarrow X$ of smooth projective varieties, denoting its graph by Γ_f , we can compute maps of simplicial complexes

$$\mathcal{H}(Z) \leftarrow \mathcal{H}(\Gamma_f) \rightarrow \mathcal{H}(X)$$

and the induced map

$$H_i(Z(\mathbb{C}), \mathbb{Z}) \cong H_i(\Gamma_f(\mathbb{C}), \mathbb{Z}) \xrightarrow{f_*} H_i(X(\mathbb{C}), \mathbb{Z}).$$

If X and Z are irreducible and have dimensions d and p respectively and if f is generically finite onto the image, then the cycle class $[f] \in H^{2d-2p}(X(\mathbb{C}), \mathbb{Z})$ of f is computed to be the element corresponding to $f_*([Z]) \in H_{2p}(X(\mathbb{C}), \mathbb{Z})$ via the Poincaré duality

$$H^{2d-2p}(X(\mathbb{C}), \mathbb{Z}) \cong H_{2p}(X(\mathbb{C}), \mathbb{Z}).$$

Here $[Z] \in H_{2p}(Z(\mathbb{C}), \mathbb{Z})$ denotes the fundamental class of Z . When $Z \subset X$ is a (possibly singular) irreducible closed subvariety of dimension p , then we can algorithmically construct a resolution of singularities $f: \tilde{Z} \rightarrow Z \subset X$ (see [Vil89, VU92, BM91, Bie97, BS00]) and define the cycle class $[Z] \in H^{2d-2p}(X(\mathbb{C}), \mathbb{Z})$ to be $[f]$.

For an invertible sheaf \mathcal{L} on X , we can compute a divisor D on X such that $\mathcal{L} \cong \mathcal{O}_X(D)$, for example, by an algorithm given in [SY18, Section 3]. If we write $D = \sum_{i=1}^n a_i D_i$ with D_i prime divisors and a_i integers, then the cohomology class $[\mathcal{L}]$ of \mathcal{L} is defined to be $[D] = \sum_{i=1}^n a_i [D_i]$.

We can also compute the cup product

$$H^i(X(\mathbb{C}), \mathbb{Z}) \times H^j(X(\mathbb{C}), \mathbb{Z}) \rightarrow H^{i+j}(X(\mathbb{C}), \mathbb{Z})$$

again by using the representation of $H^i(X(\mathbb{C}), \mathbb{Z})$ in terms of the simplicial complex $\mathcal{H}(X)$. In summary, in the situation of Proposition 9.1, we can algorithmically compute elements $[Z], [\mathcal{L}] \in H^{2d-2c}(X(\mathbb{C}), \mathbb{Z})$ as represented by explicit cochains on $\mathcal{H}(X)$ and compute the product $[Z][\mathcal{L}]^{\dim Z} \in H^{2d}(X(\mathbb{C}), \mathbb{Z})$ with respect to the cup product. The desired intersection number $Z \cdot \mathcal{L}^{\dim Z} \in \mathbb{Z}$ is then computed as the integer n such that

$[Z][\mathcal{L}]^{\dim Z} = n[\mathbf{pt}]$, where $[\mathbf{pt}]$ is the cycle class of a point of $X(\mathbb{C})$. This completes the proof of Proposition 9.1.

10. POSITIVITY OF INVERTIBLE SHEAVES

Positivity properties of invertible sheaves, such as ample, big, and nef, are closely related to the isomorphism problem. In this section, we discuss the decidability problem of these properties. We also show that, for a smooth variety whose Picard number can be computable, we can approximate its nef cone and pseudo-effective cone with arbitrary precision.

10.1. Global generation.

Proposition 10.1. *For a projective variety X and a coherent sheaf \mathcal{L} on it, we can algorithmically check whether it is globally generated. (We do not assume that \mathcal{L} is invertible, although it is the case of our main interest.)*

Proof. Let R be the homogeneous coordinate ring of X and let L be the given finitely generated graded R -module, which defines \mathcal{L} . We can compute the graded R -module $L' := \bigoplus_{v \geq 0} H^0(X, \tilde{L}(v))$ as the Hom module $\text{Hom}_R(R_{\geq r}, L)_{\geq 0}$ for some sufficiently large r , see Theorem 8.2 of Chapter 8 by Eisenbud in the book [Vas98]. Let

$$F_1 \rightarrow F_0 \rightarrow L' \rightarrow 0$$

be the obtained minimal free presentation of L' . Here the arrows are degree-preserving R -linear maps and the free module F_0 is written as

$$F_0 = \bigoplus_{i=1}^c R y_i$$

with homogeneous generators y_i with

$$0 \leq \deg(y_1) \leq \dots \leq \deg(y_c).$$

Let y_1, \dots, y_n ($n \leq c$) be the ones of degree 0, which are regarded as a basis of $H^0(X, \tilde{L})$, and let

$$F'_0 := \bigoplus_{i=1}^n R y_i.$$

The derived map $F'_0 \rightarrow L'$ induces the map of sheaves,

$$\mathcal{O}_X \otimes H^0(X, \tilde{L}) \rightarrow \tilde{L}.$$

The sheaf \tilde{L} is globally generated if and only if this map is surjective. We can check the latter condition, for example, by computing the support of the \mathcal{O}_X -module corresponding to the graded R -module $\text{Coker}(F'_0 \rightarrow L')$ and see whether it is empty. \square

10.2. Very ampleness.

Proposition 10.2. *For a projective variety X and an invertible sheaf \mathcal{L} on it, we can algorithmically check whether it is very ample.*

Proof. We first check the global generation of \mathcal{L} . If \mathcal{L} is not globally generated, then \mathcal{L} is not very ample. Suppose that \mathcal{L} is globally generated. We then compute the morphism $\Phi_{\mathcal{L}}: X \rightarrow \mathbb{P}^{n-1}$ associated to \mathcal{L} , where $n = \dim H^0(X, \mathcal{L})$. Let L be the given graded R -module defining \mathcal{L} and we construct a map of R -modules, $F'_0 \rightarrow L'$, as in the proof of Proposition 10.1. Let M be the image of this map, which defines the same sheaf on X as L' and L do. We have a free presentation

$$F'_1 \rightarrow F'_0 \rightarrow M \rightarrow 0.$$

From [Eis95, Prop. A2.2], this induces the exact sequence

$$\mathrm{Sym}_R(F'_0) \otimes_R F'_1 \rightarrow \mathrm{Sym}_R(F'_0) \rightarrow \mathrm{Sym}_R(M) \rightarrow 0.$$

If the basis of F'_1 maps to

$$g_i = \sum_{j=1}^n g_{ij} y_j \quad (i = 1, \dots, r)$$

with r denoting the rank of F'_1 , then

$$\begin{aligned} \mathrm{Sym}_R(M) &= R[y_1, \dots, y_n]/(g_1, \dots, g_r) \\ &= k[x_1, \dots, x_m, y_1, \dots, y_n]/(f_1, \dots, f_t, g_1, \dots, g_r), \end{aligned}$$

which is bi-graded. This defines a closed subscheme $\Gamma \subset \mathbb{P}^{m-1} \times \mathbb{P}^{n-1}$. The projection $\Gamma \rightarrow \mathbb{P}^{m-1}$ is an isomorphism onto X . In other words, Γ is the graph of a morphism $X \rightarrow \mathbb{P}^{n-1}$. The last morphism is the morphism $\Phi_{\tilde{\mathcal{L}}}$ associated to the globally generated invertible sheaf $\tilde{\mathcal{L}}$. Now we can compute the image $Y := \Phi_{\tilde{\mathcal{L}}}(X)$ of $\Phi_{\tilde{\mathcal{L}}}$ by projective elimination (see Remark 10.3). From Lemma 3.3, we can check whether the morphism $X \rightarrow Y$, which corresponds to $\Gamma \subset \mathbb{P}^{m-1} \times \mathbb{P}^{n-1}$, is an isomorphism. Our invertible sheaf \mathcal{L} is very ample if and only if the last morphism is an isomorphism. \square

Remark 10.3 (Projective elimination). Suppose that a closed subscheme $V \subset \mathbb{P}^{m-1} \times \mathbb{P}^{n-1}$ is defined by a bi-homogeneous ideal $I \subset k[x_1, \dots, x_m, y_1, \dots, y_n]$. Then the scheme-theoretic image $p_2(V)$ by the second projection is defined by the ideal

$$(I : (x_1, \dots, x_m)^\infty) \cap k[y_1, \dots, y_n].$$

At the set-theoretic level, this is written [GP08, page 503]; the closed subset $p_2(V)$ is the zero set of the last homogeneous ideal. The scheme-theoretic version follows from [GP08, Lemma A.7.9].

10.3. Ampleness.

Proposition 10.4. *We can algorithmically decide whether an invertible sheaf on a smooth irreducible projective variety is ample.*

Proof. Let X be a projective scheme and let \mathcal{L} be an invertible sheaf on X . From an effective version of Matsusaka's Big Theorem [Siu93], there exists a positive integer $m(\mathcal{L})$ explicitly determined by $\mathcal{L}^{\dim X}$, $\omega_X \cdot \mathcal{L}^{\dim X - 1}$, and $\dim X$ such that \mathcal{L} is ample if and only if $\mathcal{L}^{\otimes m(\mathcal{L})}$ is very ample. Thus, we only need to compute the number $m(\mathcal{L})$ and check whether $\mathcal{L}^{\otimes m(\mathcal{L})}$ is very ample. \square

10.4. Nefness, bigness and pseudo-effectivity. These properties in the title of invertible sheaves are all positivity properties in some sense, which are weaker than ampleness, and play important roles in birational geometry. In what follows, we restrict ourselves to the case where the ambient scheme is irreducible and smooth, unless otherwise noted.

Definition 10.5. Let X be a smooth irreducible projective variety and let \mathcal{L} be an invertible sheaf on X . The \mathcal{L} is *big* if for some integer $n > 0$, the rational map associated to \mathcal{L}^n ,

$$\Phi_{\mathcal{L}^n} : X \dashrightarrow \mathbb{P}^m,$$

is birational onto the image. The \mathcal{L} is *nef* if for every irreducible curve $C \subset X$, we have $C \cdot \mathcal{L} \geq 0$. The \mathcal{L} is *pseudo-effective* if its class in $\text{NS}(X) \otimes \mathbb{R}$ is the limit of classes of effective divisors.

Note that it is easy to check whether \mathcal{L} is effective (that is, isomorphic to $\mathcal{O}_X(D)$ for some effective divisor D) by computing the cohomology group $H^0(X, \mathcal{L})$. For each n , we can algorithmically check whether the map $\Phi_{\mathcal{L}^n}$ is birational, see [Sim04, DHS12]. See also [BHSS19] for implementation of such an algorithm. Thus, bigness of an invertible sheaf is semi-decidable. On the other hand, *not* being nef is a semi-decidable property. Indeed, we enumerate irreducible curves in X as C_1, C_2, \dots and for each i , we compute the intersection number $C_i \cdot \mathcal{L}$, until we get a negative intersection number. As for pseudo-effectivity, we have the following theorem [BDPP13, 0.2 Theorem]: \mathcal{L} is pseudo-effective if and only if $\mathcal{L} \cdot C \geq 0$ for every irreducible curve $C \subset X$ which moves in a family covering X . Using this, we can prove:

Proposition 10.6. *Not being pseudo-effective is a semi-decidable property.*

Proof. We enumerate all irreducible curves on X as C_1, C_2, \dots . Consider the following algorithm:

- (1) Put $n = 1$.
- (2) Check whether C_n moves in a family covering X as follows. We first compute the Hilbert polynomial P_n of C_n and then compute the connected component W of $\text{Hilb}_{P_n}(X)$ containing $[C_n]$. We then check whether the universal family \mathcal{U}_W on W maps onto X ; C_n is a movable curve if and only if this is the case. When C_n is movable, we compute the intersection number $C_n \cdot D$ and stop the algorithm if $C_n \cdot D < 0$.
- (3) Put $n = n + 1$ and go back to (2).

From [BDPP13, 0.2 Theorem], this algorithm stops after finitely many steps if and only if D is not pseudo-effective. The proposition follows. \square

In summary, the following properties of invertible sheaves on a smooth irreducible projective variety are semi-decidable:

- (1) Being big.
- (2) Not being nef.
- (3) Not being pseudo-effective.

It is now quite natural to ask:

Problem 10.7. Are the three properties, big, nef and pseudo-effective, decidable?

Remark 10.8. Note that when the given invertible sheaf is known to be nef, then we only need to compute the intersection number $\mathcal{L}^{\dim X}$ to check whether \mathcal{L} is big (see [Laz04, Theorem 2.2.16]).

In the case of the canonical sheaf ω_X , the most important invertible sheaf, we may take advantage of the following conjectures:

Conjecture 10.9 (The abundance conjecture). *For a smooth projective variety X , the canonical sheaf ω_X is nef if and only if it is semi-ample (that is, \mathcal{L}^n is globally generated for some $n > 0$).*

Conjecture 10.10 (The non-vanishing conjecture). *For a smooth projective variety X , the canonical sheaf ω_X is pseudo-effective if and only if it is \mathbb{Q} -linearly equivalent to an effective \mathbb{Q} -divisor.*

The abundance conjecture is recognized as one of the most important conjectures in the minimal model program. The importance of the non-vanishing conjecture was pinned down by Birkar [Bir11]. The above form of the non-vanishing conjecture is slightly different from the one considered by Birkar. However Hashizume [Has18] proved that they are equivalent.

Proposition 10.11. *Let X be a smooth irreducible projective variety.*

- (1) *If the abundance conjecture holds for X , then the nefness of ω_X is decidable.*
- (2) *If the weak nonvanishing conjecture holds for X , then the pseudo-effectivity of ω_X is decidable.*

Proof. (1) We first note that we can compute the canonical sheaf [Sti, Section 5.6]. We enumerate all irreducible curves in X as C_1, C_2, \dots . Consider the following algorithm:

- (1) Put $n = 1$.
- (2) We check whether $\omega_X^{\otimes n}$ is globally generated. If this is the case, then stop the algorithm and output True.
- (3) We check whether $C_n \cdot \omega_X < 0$. If this is the case, then stop the algorithm and output False.
- (4) Put $n = n + 1$ and go to (2).

If the abundance conjecture holds, then this algorithm always stops after finitely many steps and outputs True if ω_X is nef and False if ω_X is not nef.

(2) Effective \mathbb{Q} -divisors on X are enumerable. For each of them, we can compute its class in $H^2(X(\mathbb{C}), \mathbb{Q})$ by the method explained in Section 9 and check whether it coincides with the class of ω_X . From the non-vanishing conjecture, we see that the

pseudo-effectivity of ω_X is semi-decidable. Combining this with Proposition 10.6 shows the assertion. \square

Remark 10.12. If $\dim X \leq 3$, then the bigness of ω_X is also decidable. Indeed, from [HM06, Tak06, Tsu06], for each dimension d , there exists a positive integer n_d such that for every smooth variety X of general type and of dimension d and for every integer $n \geq n_d$, the rational map $\Phi_{\omega_X^{\otimes n}}$ is birational onto the image. Moreover, for $d \leq 3$, we can take $n_1 = 3$, $n_2 = 5$, $n_3 = 126$ (see [Bom73, CC10]); we only need to check whether $\Phi_{\omega_X^{\otimes n_d}}$ is birational onto the image. To generalize this argument to dimensions ≥ 4 , we need to compute n_d .

If ω_X is nef, then we can check its bigness in any dimension, see Remark 10.8. If ω_X is not nef, then we may run the minimal model program. As an output of the program, we would get a Mori fiber space or a minimal model birational to the given variety X . In the former case, ω_X is not pseudo-effective, in particular, not big. In the latter case, we can check the bigness of ω_X by computing the intersection number $(\omega_X)^{\dim X}$. This strategy provides motivation for studying the following problem, which would be important also on its own right:

Problem 10.13. Describe each step of the minimal model program as a strict algorithm, starting from algorithmically finding a ω_X -negative ray of the cone of curves.

10.5. Approximating nef and pseudo-effective cones. Let $\text{NS}(X)_{\mathbb{R}} := \text{NS}(X) \otimes \mathbb{R}$ denote the Néron-Severi group tensored with \mathbb{R} . This is a finite-dimensional \mathbb{R} -vector space and its dimension $\rho(X)$ is called the Picard number of X . The *nef cone* of X , denoted by $\text{Nef}(X)$, is the smallest closed convex cone in $\text{NS}(X)_{\mathbb{R}}$ such that, for an invertible sheaf \mathcal{L} , the class $[\mathcal{L}]$ belongs to it if and only if \mathcal{L} is nef. The *pseudo-effective cone* $\text{PEff}(X)$ is similarly defined. The *ample cone* and the *big cone* are the interiors of the nef cone and the pseudo-effective cone respectively.

As we do not have an algorithm to decide whether a given invertible sheaf is big/nef, we can not compute the cones $\text{PEff}(X)$ and $\text{Nef}(X)$ at least for now. However, if we know the value of the Picard number $\rho(X)$, then we can approximate these cones with arbitrary precision. Note that, if we know the value of $\rho(X)$, then we can compute the subspace $\text{NS}(X)_{\mathbb{R}} \subset H^2(X(\mathbb{C}), \mathbb{R})$ by giving a basis of it. To do so, we only need to compute classes $[D] \in H^2(X(\mathbb{C}), \mathbb{R})$ of divisors $D \subset X$, until we have enough to span a subspace of dimension $\rho(X)$. Poonen, Testa and van Luijk [PTL15] gave an algorithm to compute $\rho(X)$, assuming the Tate conjecture. In particular, we can compute $\rho(X)$ if X is a K3 surface.

Proposition 10.14. *Let X be a smooth irreducible projective variety. Suppose that we know the value of $\rho(X)$. We fix a metric on $\text{NS}(X)_{\mathbb{R}}$. Let $S \subset \text{NS}(X)_{\mathbb{R}}$ be the unit sphere with center at the origin. Then, for any positive real number $\epsilon > 0$, we can algorithmically construct rational polyhedral convex cones A_{ϵ} and B_{ϵ} such that $A_{\epsilon} \subset \text{Nef}(X) \subset B_{\epsilon}$ and $B_{\epsilon} \cap S$ is contained in the ϵ -neighborhood of $A_{\epsilon} \cap S$. Similarly for $\text{PEff}(X)$.*

Proof. Let ρ denote the Picard number of X . From Proposition 10.4, we can enumerate all the ample divisors on X as D_1, D_2, \dots . Let $A_n = \sum_{i=1}^n \mathbb{R}_{\geq 0}[D_i]$ be the convex

cone generated by $[D_1], \dots, [D_n]$. The closure of $\bigcup_{n \geq 0} A_n$ is the nef cone $\text{Nef}(X)$. In particular, each A_n is a rational convex polyhedral cone contained in $\text{Nef}(X)$. We can also enumerate the irreducible curves in X as C_1, C_2, \dots . Let

$$B_n^c := \bigcup_{i=1}^n \{x \in \text{NS}(X)_{\mathbb{R}} \mid x \cdot C_i < 0\} \text{ and}$$

$$B_n := \text{NS}(X)_{\mathbb{R}} \setminus B_n^c = \bigcap_{i=1}^n \{x \in \text{NS}(X)_{\mathbb{R}} \mid x \cdot C_i \geq 0\}.$$

We see that $\bigcup_{n \geq 0} B_n^c = \text{NS}(X)_{\mathbb{R}} \setminus \text{Nef}(X)$. Thus each B_n is a rational convex polyhedral cone containing $\text{Nef}(X)$. It is also strongly convex (that is, it has a vertex at the origin) for $n \gg 0$. We have got two sequences $(A_n)_n$ and $(B_n)_n$ of rational convex polyhedral cones approximating $\text{Nef}(X)$ from inside and outside respectively. Therefore, for $n \gg 0$, A_n and B_n satisfy the desired condition. To see for which value of n this is the case, we first check whether B_n is strongly convex. If this is the case, then for each vertex $w \in B_n \cap S$, we check whether every vertex $w \in B_n \cap S$ is contained in the ϵ -neighborhood of A_n . If this is the case, B_n is contained in the ϵ -neighborhood of A_n . This completes the proof for the nef cone $\text{Nef}(X)$.

As for the pseudo-effective cone $\text{PEff}(X)$, we only need to replace ample divisors with big divisors and irreducible curves with movable irreducible curves. To enumerate movable irreducible curves, we can use the algorithm in the proof of 10.6. To enumerate big divisors, we can use the following algorithm: We first enumerate all the divisors on X as D_1, D_2, \dots .

- (1) Put $n = 1$ and put $b = ()$, the empty ordered tuple.
- (2) For each $i, j \leq n$, if Φ_{i, D_j} is a birational map onto the image and if $D_j \notin b$, then append D_j to b .
- (3) Put $n = n + 1$ and go to (2).

For every big divisor D on X , the above algorithm appends D to b after finitely many steps. Thus, for every positive integer n , we can algorithmically construct the n -th big divisor. (Thus, big divisors on X are listable. But this does not mean that the bigness of each divisor is decidable.) \square

11. K3 SURFACES

In this section, we discuss the isomorphism problem for K3 surfaces, which would be natural as the next case to study after the one-dimensional case and the case with ω_X or ω_X^{-1} big were treated in Sections 7 and 8 respectively. The main result of this section is the decidability of the isomorphism problem for K3 surfaces with an automorphism group finite.

Proposition 11.1. *Let X be a K3 surface. If $\text{Aut}(X)$ is finite, then we can compute the nef cone $\text{Nef}(X)$ by giving finitely many effective divisors D_1, \dots, D_n such that $\text{Nef}(X) = \sum_{i=1}^n \mathbb{R}_{\geq 0}[D_i]$.*

Proof. From [PTL15], we can compute the Picard number $\rho(X)$ and compute $\text{NS}(X)_{\mathbb{R}}$ as explained in Section 10.5. For an effective divisor D on X , we can check whether it

is nef; we check whether $C \cdot D \geq 0$ for every prime divisor C contained in the support of D . Therefore we can enumerate all the effective divisors as D_1, D_2, \dots and all the nef and effective divisors as N_1, N_2, \dots . For each n , let

$$A_n := \sum_{i=1}^n \mathbb{R}_{\geq 0}[N_i],$$

$$B_n := \bigcap_{i=1}^n \{x \in \text{NS}(X)_{\mathbb{R}} \mid x \cdot D_i \geq 0\}.$$

These are rational polyhedral convex cones satisfying

$$(11.1) \quad A_n \subset \text{Nef}(X) \subset B_n.$$

If $\text{Aut}(X)$ is finite, then $\text{PEff}(X)$ is a rational polyhedral cone spanned by effective classes [Kov94]. It follows that every point of $\text{NS}(X)_{\mathbb{Q}} \cap \text{PEff}(X)$ is represented by an effective \mathbb{Q} -divisor. Since $\text{Nef}(X)$ is the dual cone of $\text{PEff}(X)$, it is also rational polyhedral and spanned by finitely many points of $\text{NS}(X)_{\mathbb{Q}}$. Since $\text{Nef}(X) \subset \text{PEff}(X)$, these points are represented by nef and effective divisors after multiplied with some positive integer. We conclude that $\text{Nef}(X)$ is a rational polyhedral cone spanned by nef and effective classes. Therefore, for $n \gg 0$, inclusions on this page are equalities. For each n , we compute A_n and B_n and check whether $A_n = B_n$. If this equality holds, then the cone $A_n = B_n$ is the nef cone. \square

Proposition 11.2. *For a K3 surface X , we can algorithmically decide whether $\text{Aut}(X)$ is finite.*

Proof. We define cones A_n and B_n in $\text{NS}(X)_{\mathbb{R}}$ as in the proof of Proposition 11.1. From [Kov94], $A_n = B_n$ for $n \gg 0$ if and only if $\text{Aut}(X)$ is finite. Therefore we have an algorithm which stops after finitely many steps exactly when $\text{Aut}(X)$ is finite. We denote this algorithm by Θ .

For an automorphism $f: X \rightarrow X$, the tangent space of $\underline{\text{Aut}}(X) = \underline{\text{Iso}}(X, X)$ at $[f]$ is isomorphic to $H^0(X, \mathcal{T}_X)$ with \mathcal{T}_X denoting the tangent sheaf. Since

$$H^0(X, \mathcal{T}_X)^\vee = H^2(X, \omega_X \otimes \Omega_X) = H^2(X, \Omega_X) = 0,$$

the $\underline{\text{Aut}}$ scheme $\underline{\text{Aut}}(X)$ has only isolated points. From [Kon99], any finite subgroup of $\text{Aut}(X)$ has order at most 3840. The following algorithm stops after finitely many steps exactly when $\text{Aut}(X)$ is infinite: We enumerate all the Hilbert polynomials as P_1, P_2, \dots

- (1) Put $n = 1$ and $\text{numAuts} = 0$.
- (2) Put $\text{numAuts} = \text{numAuts} + \#\underline{\text{Iso}}_{P_n}(X, X)$.
- (3) If $\text{numAuts} > 3840$, then stop.
- (4) Put $n = n + 1$ and go to (2).

We denote this algorithm by Θ' . Now the following algorithm is the desired one:

- (1) Put $n = 1$.
- (2) If Θ stops after n steps, then stop and output Finite.
- (3) If Θ' stops after n steps, then stop and output Infinite.
- (4) Put $n = n + 1$ and go to (2).

□

Theorem 11.3. *For K3 surfaces X and Y with finite automorphism groups, we can algorithmically decide whether they are isomorphic.*

Proof. We compute the nef cones $\text{Nef}(X)$ and $\text{Nef}(Y)$. There exist at most finitely many isomorphisms $g: \text{NS}(Y) \rightarrow \text{NS}(X)$ such that $g(\text{Nef}(Y)) = \text{Nef}(X)$. If there is no such isomorphism, then X and Y are not isomorphic. Suppose that this is not the case and let g_1, \dots, g_n be all the isomorphisms with this property. Let \mathcal{L} and \mathcal{M} be the given very ample sheaves on X and Y . We compute the Hilbert polynomial for each $[\mathcal{L}] + g_i[\mathcal{M}]$ and call it by P_i . Note that the Hilbert polynomial of an ample invertible sheaf depends only on its numerical class. Indeed, the numerical class $[\mathcal{N}]$ of an ample invertible sheaf determines the Euler characteristics $\chi(\mathcal{N})$ and $\chi(\mathcal{N}^2)$ from the Riemann-Roch formula for surfaces. These values together with the one of $\chi(\mathcal{O}_X)$ determines the Hilbert polynomial of \mathcal{N} . If there is an isomorphism $f: X \rightarrow Y$, then the induced isomorphism $\text{NS}(Y) \rightarrow \text{NS}(X)$ is one of the g_i 's and the Hilbert polynomial of f is one of the P_i 's. Thus $[f]$ is a point of $\bigcup_{i=1}^n \text{Iso}_{P_i}(X, Y)$. Thus, X and Y are isomorphic if and only if $\bigcup_{i=1}^n \text{Iso}_{P_i}(X, Y) \neq \emptyset$. From Section 6, the last condition can be algorithmically checked. □

Remark 11.4. For a general K3 surface X , there are only finitely many very ample class x with x^2 being the prescribed number modulo the action of $\text{Aut}(X)$ [Ste85, 2.6]. Therefore, if we replace the given very ample sheaf of X by a suitable automorphism of X , we can find an isomorphism $X \rightarrow Y$ (if any) with the Hilbert polynomial in a finite set of potential candidates. But there is a priori no way to know which automorphism does this job.

Remark 11.5. Let X and Y be K3 surfaces and let \mathcal{L} be the given very ample sheaf of Y . There is no intrinsic invariants of X and Y to determine the place of $f^*[\mathcal{L}]$ in the ample cone of X for a potential isomorphism $f: X \rightarrow Y$. Indeed, when $\rho = 2$ and they have infinite automorphisms, then a very ample class l with $l^2 = d$ is sent to infinitely many distinct lattice points on the curve $x^2 = d$. For two lattice points l_1 and l_2 on the curve, the sum $l_1 + l_2$ can have arbitrarily large Euler characteristic.

Remark 11.6. Discussion in this section indicates that the isomorphism problem is closely related to complexity of the automorphism group. Recently, Lesieutre [Les18] showed that there exists a projective variety X whose automorphism group is discrete, but not finitely generated (see also [DO19]). This result may be considered to suggest that the isomorphism problem for general projective schemes is not decidable.

REFERENCES

- [Bay82] D. Bayer. *The division algorithm and the hilbert scheme*. PhD thesis, Harvard University, 1982.
- [BDPP13] Sébastien Boucksom, Jean-Pierre Demailly, Mihai Păun, and Thomas Peternell. The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. *Journal of Algebraic Geometry*, 22(2):201–248, 2013.
- [BGJGP05] Matthew H. Baker, Enrique González-Jiménez, Josep González, and Bjorn Poonen. Finiteness Results for Modular Curves of Genus at Least 2. *American Journal of Mathematics*, 127(6):1325–1387, 2005.

- [BHPV04] W. Barth, K. Hulek, Chris Peters, and A. van de Ven. *Compact Complex Surfaces*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics. Springer-Verlag, Berlin Heidelberg, 2nd edition, 2004.
- [BHSS19] C. J. Bott, S. Hamid Hassanzadeh, Karl Schwede, and Daniel Smolkin. RationalMaps, a package for Macaulay2. *arXiv:1908.04337*, August 2019.
- [Bie97] Edward Bierstone. Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant. *Inventiones mathematicae*, 128(2):207–302, April 1997.
- [Bir11] Caucher Birkar. On existence of log minimal models II. 2011(658):99–113, September 2011. Publisher: De Gruyter Section: Journal für die reine und angewandte Mathematik.
- [BM91] Edward Bierstone and Pierre D. Milman. A simple constructive proof of Canonical Resolution of Singularities. In Teo Mora and Carlo Traverso, editors, *Effective Methods in Algebraic Geometry*, Progress in Mathematics, pages 11–30. Birkhäuser, Boston, MA, 1991.
- [Bom73] Enrico Bombieri. Canonical models of surfaces of general type. *Inst. Hautes Études Sci. Publ. Math.*, pages 171–219, 1973.
- [BS00] Gábor Bodnár and Josef Schicho. A Computer Program for the Resolution of Singularities. In Herwig Hauser, Joseph Lipman, Frans Oort, and Adolfo Quirós, editors, *Resolution of Singularities: A research textbook in tribute to Oscar Zariski Based on the courses given at the Working Week in Obergurgl, Austria, September 7–14, 1997*, Progress in Mathematics, pages 231–238. Birkhäuser, Basel, 2000.
- [CC10] Jungkai A. Chen and Meng Chen. Explicit birational geometry of threefolds of general type, I. *Annales scientifiques de l'École Normale Supérieure*, 43(3):365–394, 2010.
- [Chi86] A. L. Chistov. Algorithm of polynomial complexity for factoring polynomials and finding the components of varieties in subexponential time. *Journal of Soviet Mathematics*, 34(4):1838–1882, August 1986.
- [DHS12] A. V. Doria, S. H. Hassanzadeh, and A. Simis. A characteristic-free criterion of birationality. *Advances in Mathematics*, 230(1):390–413, May 2012.
- [DO19] Tien-Cuong Dinh and Keiji Oguiso. A surface with discrete and nonfinitely generated automorphism group. *Duke Mathematical Journal*, 168(6):941–966, April 2019.
- [EHV92] David Eisenbud, Craig Huneke, and Wolmer Vasconcelos. Direct methods for primary decomposition. *Inventiones mathematicae*, 110(1):207–235, December 1992.
- [Eis95] David Eisenbud. *Commutative Algebra: with a View Toward Algebraic Geometry*. Graduate Texts in Mathematics. Springer-Verlag, New York, 1995.
- [ES08] David Eisenbud and Frank-Olaf Schreyer. Relative Beilinson monad and direct image for families of coherent sheaves. *Transactions of the American Mathematical Society*, 360(10):5367–5396, April 2008.
- [GP08] G.-M. Greuel and Gerhard Pfister. *A Singular Introduction to Commutative Algebra*. Springer-Verlag, Berlin Heidelberg, 2nd edition, 2008.
- [GS] Daniel R. Grayson and Michael E. Stillman. *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [Har77] Robin Hartshorne. *Algebraic Geometry*. Graduate Texts in Mathematics. Springer-Verlag, New York, 1977.
- [Has18] Kenta Hashizume. On the Non-vanishing Conjecture and Existence of Log Minimal Models. *Publications of the Research Institute for Mathematical Sciences*, 54(1):89–104, January 2018.
- [HM06] Christopher D. Hacon and James McKernan. Boundedness of pluricanonical maps of varieties of general type. *Inventiones mathematicae*, 166(1):1–25, October 2006.
- [IK99] Anthony Iarrobino and Vassil Kanev. Sums of powers of linear forms, and gorenstein algebras. In Anthony Iarrobino and Vassil Kanev, editors, *Power Sums, Gorenstein Algebras, and Determinantal Loci*, Lecture Notes in Mathematics, pages 57–72. Springer, Berlin, Heidelberg, 1999.

- [KBC19] A. Ya. Kanel'-Belov and A. A. Chilikov. On the Algorithmic Undecidability of the Embeddability Problem for Algebraic Varieties over a Field of Characteristic Zero. *Mathematical Notes*, 106(1):299–302, July 2019.
- [Kol96] János Kollár. *Rational Curves on Algebraic Varieties*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics. Springer-Verlag, Berlin Heidelberg, 1996.
- [Kol20] J. Kollár. Pell surfaces. *Acta Mathematica Hungarica*, 160(2):478–518, April 2020.
- [Kon99] Shigeyuki Kondō. The Maximum Order of Finite Groups of Automorphisms of K3 Surfaces. *American Journal of Mathematics*, 121(6):1245–1252, 1999.
- [Kov94] Sándor J. Kovács. The cone of curves of a K3 surface. *Mathematische Annalen*, 300(1):681–691, September 1994.
- [Laz04] R. K. Lazarsfeld. *Positivity in Algebraic Geometry I: Classical Setting: Line Bundles and Linear Series*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics, Positivity in Algebraic Geometry. Springer-Verlag, Berlin Heidelberg, 2004.
- [Les18] John Lesieutre. A projective variety with discrete, non-finitely generated automorphism group. *Inventiones mathematicae*, 212(1):189–211, April 2018.
- [Mat89] Hideyuki Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1989.
- [Poo11] Bjorn Poonen. Automorphisms mapping a point into a subvariety. *Journal of Algebraic Geometry*, 20(4):785–794, 2011.
- [Poo14] Bjorn Poonen. Undecidable problems: a sampler. In *Interpreting Gödel: Critical Essays*. Cambridge University Press, August 2014.
- [PTL15] Bjorn Poonen, Damiano Testa, and Ronald van Luijk. Computing Néron–Severi groups and cycle class groups. *Compositio Mathematica*, 151(4):713–734, April 2015.
- [Sim04] Aron Simis. Cremona transformations and some related algebras. *Journal of Algebra*, 280(1):162–179, October 2004.
- [Sim08] Carlos Simpson. Algebraic cycles from a computational point of view. *Theoretical Computer Science*, 392(1-3):128–140, February 2008.
- [Siu93] Yum-Tong Siu. An effective Matsusaka big theorem. *Annales de l'Institut Fourier*, 43(5):1387–1405, 1993.
- [Smi98] Gregory G. Smith. Computing Global Extension Modules for Coherent Sheaves on a Projective Scheme. July 1998.
- [Sta21] The Stacks Project Authors. The Stacks project, 2021. <https://stacks.math.columbia.edu>.
- [Ste85] Hans Sterk. Finiteness results for algebraic K3 surfaces. *Mathematische Zeitschrift*, 189(4):507–513, December 1985.
- [Sti] Michael Stillman. Computing with sheaves and sheaf cohomology in algebraic geometry: preliminary version. page 41.
- [SY18] Karl Schwede and Zhaoning Yang. Divisor Package for Macaulay2. *Journal of Software for Algebra and Geometry*, 8(1):87–94, September 2018.
- [Tak06] Shigeharu Takayama. Pluricanonical systems on algebraic varieties of general type. *Inventiones mathematicae*, 165(3):551, April 2006.
- [Tru18] Tuyen Trung Truong. Bounded birationality and isomorphism problems are computable. *arXiv:1801.00901*, July 2018.
- [Tsu06] Hajime Tsuji. Pluricanonical systems of projective varieties of general type I. *Osaka Journal of Mathematics*, 43(4):967–995, December 2006.
- [Vak15] Ravi Vakil. Foundations of Algebraic Geometry (april 29, 2015 draft), 2015. <http://math.stanford.edu/~vakil/216blog/FOAGapr2915public.pdf>.
- [Vas98] Wolmer Vasconcelos. *Computational Methods in Commutative Algebra and Algebraic Geometry*. Algorithms and Computation in Mathematics. Springer-Verlag, Berlin Heidelberg, 1998.

- [Vil89] Orlando Villamayor. Constructiveness of Hironaka's resolution. *Annales scientifiques de l'École Normale Supérieure*, 22(1):1–32, 1989.
- [VU92] O. E. Villamayor U. Patching local uniformizations. *Annales scientifiques de l'École normale supérieure*, 25(6):629–677, 1992.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY TOYONAKA, OSAKA 560-0043, JAPAN

Email address: yasuda.takehiko.sci@osaka-u.ac.jp