

# GREATEST RICCI LOWER BOUNDS OF PROJECTIVE HOROSPHERICAL MANIFOLDS OF PICARD NUMBER ONE

DONGSEON HWANG, SHIN-YOUNG KIM, AND KYEONG-DONG PARK

**ABSTRACT.** A horospherical variety is a normal  $G$ -variety such that a connected reductive algebraic group  $G$  acts with an open orbit isomorphic to a torus bundle over a rational homogeneous manifold. The projective horospherical manifolds of Picard number one are classified by Pasquier, and it turned out that the automorphism groups of all nonhomogeneous ones are non-reductive, which implies that they admit no Kähler–Einstein metrics. As a numerical measure of the extent to which a Fano manifold is close to be Kähler–Einstein, we compute the greatest Ricci lower bounds of projective horospherical manifolds of Picard number one using the barycenter of each moment polytope with respect to the Duistermaat–Heckman measure based on a recent work of Delcroix and Hultgren. In particular, the greatest Ricci lower bound of the odd symplectic Grassmannian  $SGr(n, 2n + 1)$  can be arbitrarily close to zero as  $n$  grows.

## 1. INTRODUCTION

One of the most important problems in Kähler geometry is to find a Kähler–Einstein metric. Whereas Calabi–Yau manifolds and Kähler manifolds of general type always admit Kähler–Einstein metrics by the works of Aubin [Aub78] and Yau [Yau78], Fano manifolds do not necessarily admit Kähler–Einstein metrics in general. By the Yau–Tian–Donaldson conjecture, now completely solved for anti-canonically polarized Fano varieties ([CDS15a, CDS15b, CDS15c], [Tia15] and [LXZ21]), a Fano variety admits a Kähler–Einstein metric if and only if it is K-polystable.

The *greatest Ricci lower bound*  $R(X)$  of a Fano manifold  $X$  is defined as

$$R(X) := \sup\{0 \leq t \leq 1 : \text{there exists a Kähler form } \omega \in c_1(X) \text{ with } \text{Ric}(\omega) \geq t\omega\}.$$

This invariant was first studied by Tian [Tia92], and was explicitly defined by Rubinstein [Rub08, Rub09], where it was called Tian’s  $\beta$ -invariant. It was further studied by Székelyhidi [Szé11]; Song and Wang [SW16]. Note that if  $X$  admits a Kähler–Einstein metric then the greatest Ricci lower bound  $R(X)$  is equal to one. Thus  $R(X)$  can be regarded as a measure of the extent to which a Fano manifold is close to be Kähler–Einstein. More precisely,  $R(X)$  is shown to be the same as the maximum existence time  $t \in [0, 1]$  of Aubin and Yau’s continuity path: for a Kähler form  $\omega \in c_1(X)$ , we want to find a Kähler form  $\omega_t$  satisfying the equation

$$\text{Ric}(\omega_t) = t\omega_t + (1 - t)\omega$$

depending on a parameter  $t \in [0, 1]$ , which for  $t = 1$  gives the Einstein equation.

The greatest Ricci lower bounds have been considered in certain subclasses of Fano varieties. For any toric Fano manifold  $X$ , Li [Li11] found an explicit formula for the greatest Ricci lower bound  $R(X)$  purely in terms of the moment polytope associated to  $X$ . This result is extended to Fano manifolds with torus actions of complexity one by Cable [Cab19] and homogeneous toric bundles by Yao [Yao17]. When  $X$  is a smooth Fano equivariant compactifications of complex Lie groups, Delcroix [Del17] obtained a formula for the greatest Ricci lower bound, where the barycenter of the moment polytope with respect to the Lebesgue measure is replaced by the barycenter with respect to the Duistermaat–Heckman measure. Furthermore, the recent result of Delcroix and Hultgren [DH20] extend the formula to the case of *horosymmetric* manifolds introduced in [Del20b], which is a class of spherical varieties including horospherical manifolds and smooth projective symmetric varieties. Since any  $\mathbb{Q}$ -Fano spherical variety admits a special test configuration with horospherical central fiber by [Del20a, Corollary 3.31], for the Donaldson–Futaki invariants of  $\mathbb{Q}$ -Fano spherical varieties, it is enough to calculate them in the case of horospherical varieties.

---

2010 *Mathematics Subject Classification.* Primary: 14M27, 32Q26, Secondary: 32M12, 32Q20, 53C55.

*Key words and phrases.* greatest Ricci lower bounds, horospherical varieties, algebraic moment polytopes, Kähler–Einstein metrics, odd symplectic Grassmannians.

The purpose of this paper is to compute the greatest Ricci lower bounds of projective horospherical manifolds of Picard number one. Recall that the value is always one for rational homogeneous manifolds because they are Kähler–Einstein (see [Mat72, Section 5]). The horospherical manifolds are the simplest examples of spherical varieties including both toric manifolds and rational homogeneous manifolds. All nonhomogeneous projective horospherical manifolds of Picard number one are completely classified by Pasquier ([Pas09, Theorem 0.1]). These are two orbit varieties, an open orbit and a closed orbit, under the action of their automorphism groups which are non-reductive. See Theorem 2.10 for the precise statements and notation following [PP10]. We also write down some known interesting geometry of these nonhomogeneous horospherical manifolds in Subsection 2.3. Contrary to smooth projective *symmetric* varieties of Picard number one (see [LPY21]), all nonhomogeneous projective horospherical manifolds of Picard number one admit no Kähler–Einstein metrics by a theorem of Matsushima in [Mat57] since their automorphism groups are not reductive.

**Theorem 1.1.** *For each nonhomogeneous projective horospherical manifold  $X$  of Picard number one, its greatest Ricci lower bound  $R(X)$  is as in Table 1, in which  $n \geq 3$  for  $X^1(n)$  and  $n \geq k \geq 2$  for  $X^3(n, k)$ .*

$X$	$\dim X$	$R(X)$
$X^1(n)$	$\frac{n(n+3)}{2}$	$\frac{n \int_{-n}^2 (2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt}{\int_{-n}^2 (2-t)(n+t)^n(t+2n+2)^{\frac{n(n-1)}{2}} dt}$
$X^2$	9	$\frac{20}{21} \approx 0.952$
$X^3(n, n)$	$\frac{n(n+3)}{2}$	$\frac{2 \times (2n+1)!}{(n+2)(2^n \times n!)^2}$
$X^3(n, k)$	$\frac{k(4n-3k+3)}{2}$	$\frac{(2n-2k+2) \int_{-k}^{2n-2k+2} (k+t)^{k-1}(2n-2k+2-t)^{2n-2k+1}(4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1}(2n-2k+2-t)^{2n-2k+2}(4n-3k+4-t)^{k-1} dt}$
$X^4$	23	$\frac{178992099}{243545402} \approx 0.734$
$X^5$	7	$\frac{56}{67} \approx 0.8358$

TABLE 1. Dimensions and the greatest Ricci lower bounds of nonhomogeneous projective horospherical manifolds of Picard number one.

In the case of  $X^1(n)$  and  $X^3(n, k)$ , it is possible to compute the exact number  $R(X)$  once the integers  $n$  and  $k$  are given. For instance, see the following Table 2 for the approximate values of  $R(X^1(n))$ ,  $R(X^3(n, 2))$ ,  $R(X^3(n, 3))$  and  $R(X^3(n, 4))$  for some  $n$ .

$n$	3	4	5	6	7	10	20	30	50	70
$R(X^1(n))$	0.8955	0.8755	0.8686	0.8685	0.8715	0.8863	0.9251	0.9451	0.9644	0.9737
$R(X^3(n, 2))$	0.972	0.984	0.99	0.993	0.995	0.9975	0.9994	0.9997	0.9999	0.99995
$R(X^3(n, 3))$	0.875	0.9375	0.9625	0.975	0.982	0.9917	0.9980	0.9991	0.9997	0.99984
$R(X^3(n, 4))$	–	0.820	0.902	0.938	0.958	0.9813	0.9958	0.9982	0.9994	0.99968

TABLE 2. Approximate values of  $R(X^1(n))$ ,  $R(X^3(n, 2))$ ,  $R(X^3(n, 3))$  and  $R(X^3(n, 4))$ .

We do not know how to further simplify the integrals even though we have found a nice expression in the case of  $X^3(n, n)$ . However, this complicated expression is still useful to find the following asymptotic behavior of the greatest Ricci lower bounds.

- Corollary 1.2.** (1)  $R(X^1(n))$  converges to 1 as  $n$  increases, that is,  $\lim_{n \rightarrow \infty} R(X^1(n)) = 1$ .  
 (2) For a fixed integer  $k \geq 2$ ,  $R(X^3(n, k))$  converges to 1 as  $n$  increases, that is,  $\lim_{n \rightarrow \infty} R(X^3(n, k)) = 1$ .  
 (3)  $R(X^3(n, n))$  converges to zero as  $n$  increases, that is,  $\lim_{n \rightarrow \infty} R(X^3(n, n)) = 0$ .

The greatest Ricci lower bound  $R(X)$  of a Fano manifold  $X$  is closely related with Tian's  $\alpha$ -invariant and the  $\delta$ -invariant defined by Fujita and Odaka [FO18], which have been played an important role of verifying the existence of Kähler–Einstein metrics for explicitly given Fano varieties. In particular, if  $X$  does not admit a Kähler–Einstein metric, then we have  $R(X) = \delta(X, -K_X)$  by [CRZ19, Theorem 5.7] and [BBJ21, Corollary 7.6] (see also [Gol20, Section 3.3]), and  $R(X) \geq \alpha(X) \cdot \frac{\dim X + 1}{\dim X}$  by [Tia87, Theorem 2.1]. It is interesting to note that the greatest Ricci lower bound, the alpha-invariant and the delta-invariant can be arbitrarily close to 0 for the odd symplectic Grassmannians  $X^3(n, n) = \text{SGr}(n, 2n + 1)$  by Corollary 1.2 (3).

**Corollary 1.3** (=Corollary 3.15). *For every sufficiently small number  $\epsilon > 0$ , there exists a natural number  $n$  such that  $R(\text{SGr}(n, 2n + 1)) < \epsilon$ ,  $\alpha(\text{SGr}(n, 2n + 1)) < \epsilon$  and  $\delta(\text{SGr}(n, 2n + 1)) < \epsilon$ .*

The paper is organized as follows. In Section 2, we recall the theory of spherical varieties by focusing on horospherical varieties, and we state an explicit expression of the greatest Ricci lower bounds for horospherical manifolds in terms of moment polytopes obtained by Delcroix and Hultgren. In Section 3, we obtain the greatest Ricci lower bound of a projective horospherical manifold of Picard number one by computing the barycenter of its moment polytope with respect to the Duistermaat–Heckman measure.

**Acknowledgments.** K.-D. Park would like to express his gratitude to Jihun Park for helpful discussions on K-stability of Fano manifolds. The authors would like to thank Yanir Rubinstein for valuable comments.

D. Hwang was supported by the Samsung Science and Technology Foundation under Project SSTF-BA1602-03. S.-Y. Kim was supported by the Institute for Basic Science (IBS-R003-D1). K.-D. Park was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2019R1A2C3010487).

## 2. SPHERICAL VARIETIES AND HOROSPHERICAL VARIETIES

Throughout the paper, we denote by  $G$  a connected reductive algebraic group over  $\mathbb{C}$  and by  $X$  an irreducible normal variety over  $\mathbb{C}$ .

**2.1. Spherical varieties and colors.** We review general notions and results about spherical varieties for our use. See [Kno91], [Tim11] and [Gan18] for the reference.

**Definition 2.1.** A normal variety  $X$  equipped with an action of  $G$  is called *spherical* if a Borel subgroup  $B$  of  $G$  acts on  $X$  with an open dense orbit.

Let  $X$  be a spherical variety,  $B$  be a Borel subgroup of  $G$ , and  $T$  be a maximal torus of  $B$ . Let  $G/H$  be an open dense  $G$ -orbit of  $X$ . For a character  $\chi \in \mathfrak{X}(B)$ , let  $\mathbb{C}(G/H)_\chi^{(B)} = \{f \in \mathbb{C}(G/H) : b \cdot f = \chi(b)f \text{ for all } b \in B\}$  be the set of  $B$ -semi-invariant functions in  $\mathbb{C}(G/H)$  associated to  $\chi$ , where  $\mathbb{C}(G/H)$  denotes the function field of  $G/H$ . Recall that  $\mathfrak{X}(B) = \mathfrak{X}(T)$  and  $\mathbb{C}(G/H) = \mathbb{C}(X)$  in our case.

The *spherical weight lattice*  $\mathcal{M}$  of  $G/H$  is defined as a subgroup of the character group  $\mathfrak{X}(T)$  such that each element  $\chi \in \mathcal{M}$  has the non-zero set of  $B$ -semi-invariant functions, that is,

$$\mathcal{M} = \{\chi \in \mathfrak{X}(T) : \mathbb{C}(G/H)_\chi^{(B)} \neq 0\}.$$

Note that every function  $f_\chi$  in  $\mathbb{C}(G/H)_\chi^{(B)}$  is determined by its weight  $\chi$  up to constant. This is because any  $B$ -invariant rational function on  $X$  is constant, that is,  $\mathbb{C}(G/H)^B = \mathbb{C}$ . The spherical weight lattice  $\mathcal{M}$  is a free abelian group of finite rank. We define the *rank* of  $G/H$  as the rank of the lattice  $\mathcal{M}$ . Let  $\mathcal{N} = \text{Hom}(\mathcal{M}, \mathbb{Z})$  denote its dual lattice together with the natural pairing  $\langle \cdot, \cdot \rangle : \mathcal{N} \times \mathcal{M} \rightarrow \mathbb{Z}$ .

It is known that every open  $B$ -orbit of a spherical variety is an affine variety and so its complement has pure codimension one and is a finite union of  $B$ -stable prime divisors.

**Definition 2.2.** For a spherical variety  $X$ , a  $B$ -stable but not  $G$ -stable prime divisor in  $X$  is called a *color* of  $X$ . We denote by  $\mathfrak{D} = \{D_1, \dots, D_k\}$  the set of colors of  $X$ .

We note that a color of  $X$  corresponds to a color of the open  $G$ -orbit  $G/H$  of  $X$ . As a  $B$ -semi-invariant function  $f_\chi$  in  $\mathbb{C}(G/H)_\chi^{(B)}$  is unique up to constant, we define the *color map*  $\rho: \mathfrak{D} \rightarrow \mathcal{N}$  by  $\langle \rho(D), \chi \rangle = \nu_D(f_\chi)$  for  $\chi \in \mathcal{M}$ , where  $\nu_D$  is the discrete valuation associated to a divisor  $D$ , that is,  $\nu_D(f)$  is the vanishing order of  $f$  along  $D$ . Unfortunately, the color map is not injective in general.

Every discrete  $\mathbb{Q}$ -valued valuation  $\nu$  of the function field  $\mathbb{C}(G/H)$  induces a homomorphism  $\hat{\rho}(\nu): \mathcal{M} \rightarrow \mathbb{Q}$  defined by  $\langle \hat{\rho}(\nu), \chi \rangle = \nu(f_\chi)$ . Hence, we get a map

$$\hat{\rho}: \{\text{discrete } \mathbb{Q}\text{-valued valuations on } G/H\} \rightarrow \mathcal{N} \otimes \mathbb{Q}.$$

Luna and Vust [LV83] showed that the restriction of  $\hat{\rho}$  to the set of all  $G$ -invariant discrete valuations on  $G/H$  is injective. Let  $\mathcal{V}$  be the set of  $G$ -invariant discrete  $\mathbb{Q}$ -valued valuations on  $G/H$ . Since the map  $\hat{\rho}$  is injective, we may consider  $\mathcal{V}$  as a subset of  $\mathcal{N} \otimes \mathbb{Q}$ . It is known that  $\mathcal{V}$  is a full-dimensional (co)simplicial cone of  $\mathcal{N} \otimes \mathbb{Q}$ , which is called the *valuation cone* of  $G/H$ . For a  $G$ -stable divisor  $E$  in  $X$ , we simplify the notation  $\hat{\rho}(\nu_E)$  as  $\hat{\rho}(E)$  which is in  $\mathcal{V}$ .

Since  $B$  stabilizes the open  $B$ -orbit, the stabilizer  $P$  of the open  $B$ -orbit under the  $G$ -action is a parabolic subgroup of  $G$ . Then, we have the algebraic Levi decomposition  $P = L \ltimes P^u$ , where  $P^u$  is the unipotent radical of  $P$  and  $L$  is the Levi factor of  $P$ . Let  $\Phi^+$  be the set of positive root of  $G$ ,  $\Phi_L$  be the set of roots of  $L$  and  $\Phi_{P^u}$  be the set of roots of the unipotent radical  $P^u$ . Let  $\rho_G$  be the sum of all fundamental weights of the root system for  $G$ ,  $\rho_L$  be the sum of all fundamental weights of the root system for the Levi factor  $L$  and let  $\rho_P$  be the half of the sum of roots of  $P$ . We formulate the following lemma for later use in Section 3.

**Lemma 2.3.** *In the above notation, we have the following.*

$$2\rho_P = \sum_{\alpha \in \Phi_{P^u}} \alpha = \sum_{\alpha \in \Phi^+ \setminus \Phi_L} \alpha = 2\rho_G - 2\rho_L.$$

**2.2. Horospherical varieties.** A normal  $G$ -variety  $X$  is *horospherical* if  $G$  acts with an open orbit  $G/H$  isomorphic to a torus bundle over a rational homogeneous manifold  $G/P$ , or equivalently, if the isotropy subgroup  $H$  of a general point contains the unipotent radical  $U$  of a Borel subgroup  $B$ . In this case,  $G/H$  above is called a *horospherical homogeneous space* and  $X$  is also called an *embedding of a horospherical homogeneous space*  $G/H$ . The dimension of the torus fiber is called the *rank* of a horospherical variety. A smooth horospherical variety is called a *horospherical manifold*. Note that every horospherical variety is spherical. Indeed, as the full flag variety  $G/B$  contains an open  $U$ -orbit by the Bruhat decomposition, if  $U \subset H$  then  $G/B$  has an open  $H$ -orbit; hence  $G/H$  contains an open  $B$ -orbit.

The spherical data of a projective horospherical variety can be described as follows.

**Proposition 2.4** ([Tim11, Section 28]). *For a projective horospherical variety  $X$  with an open orbit  $G/H$ , we denote by  $\pi: X \rightarrow G/P$  the torus fibration map, where a parabolic subgroup  $P = N_G(H)$  is the normalizer of  $H$  in  $G$ . Then,*

- the spherical weight lattice  $\mathcal{M}$  is the lattice  $\mathfrak{X}(T/T \cap H)$ , which implies that the rank of  $X$  is equal to the dimension of the torus fiber of  $\pi$ ,
- the valuation cone  $\mathcal{V}$  is  $\mathcal{N} \otimes \mathbb{Q}$ ,
- $\mathfrak{D}(X) = \{\pi^{-1}(D_\alpha) : D_\alpha \text{ is a Schubert divisor associated to a simple root } \alpha \text{ of } G/P\}$ , and
- the image  $\rho(\pi^{-1}(D_\alpha))$  under the color map  $\rho$  is the restriction  $\alpha^\vee|_{\mathcal{M}}$  of the simple coroot associated to a Schubert divisor  $D_\alpha$  of  $G/P$ .

Moreover, the colors of  $X$  are the closures  $\overline{Bw_0s_\alpha P/H}$  in  $X$  of  $B$ -stable irreducible divisors of  $G/H$ , where  $w_0$  is the longest element in the Weyl group of  $(G, T)$  and  $s_\alpha$  is the simple reflection associated to a simple root  $\alpha$  not in  $P$ .

The anticanonical divisor  $-K_X$  of a spherical variety  $X$  is explicitly described in [Bri97, Theorem 4.2], [Lun97, Section 3.6]. We use the following special case for projective horospherical varieties.

**Proposition 2.5** ([Pas08, Proposition 3.1]). *Let  $X$  be a projective horospherical variety with an open orbit  $G/H$ . Denote by  $-K_X$  the anticanonical divisor. Then we have the following linear equivalence.*

$$-K_X \sim \sum_{i=1}^k \langle \alpha_i^\vee, 2\rho_P \rangle D_i + \sum_{j=1}^{\ell} E_j$$

for colors  $D_i \in \mathfrak{D}(X)$  and  $G$ -stable divisors  $E_j$  in  $X$  where  $\alpha_i^\vee$  is the coroot of the simple root  $\alpha_i$  of  $G$ . Here, the coefficient  $\langle \alpha_i^\vee, 2\rho_P \rangle$  of each color is positive.

The algebraic moment polytope  $\Delta(X, L)$  for a polarized (spherical)  $G$ -variety  $(X, L)$  was introduced by Brion in [Bri87] as a purely algebraic version of the Kirwan polytope. This is indeed the convex hull of finitely many points in  $\mathfrak{X}(T) \otimes \mathbb{R}$  (see [Bri87]). Based on the works of Brion [Bri89, Bri97], Pasquier [Pas08, Chapter 3] described the algebraic moment polytope of anticanonical line bundle on a Fano horospherical variety, so called  $G/H$ -reflexive polytope.

**Proposition 2.6.** *Let  $X$  be a Gorenstein Fano horospherical variety. Then, assuming the notation in Proposition 2.5, the moment polytope  $\Delta(X, K_X^{-1})$  is  $G/H$ -reflexive and*

$$\Delta(X, K_X^{-1}) = 2\rho_P + Q_X^*,$$

where the polytope  $Q_X$  is the convex hull of the set

$$\left\{ \frac{\rho(D_i)}{\langle \alpha_i^\vee, 2\rho_P \rangle} : i = 1, \dots, k \right\} \cup \{ \hat{\rho}(E_j) : j = 1, \dots, \ell \}$$

in  $\mathcal{N} \otimes \mathbb{R}$  and its dual polytope  $Q_X^*$  is defined as  $\{m \in \mathcal{M} \otimes \mathbb{R} : \langle n, m \rangle \geq -1 \text{ for every } n \in Q_X\}$ .

To study Fano horospherical manifold, it is sufficient to consider the  $G/H$ -reflexive polytopes thanks to the following theorem.

**Theorem 2.7** ([Pas08, Proposition 3.4]). *Let  $G/H$  be a homogeneous horospherical space. There is a bijection between the set of Gorenstein Fano embeddings of  $G/H$  and the set of  $G/H$ -reflexive polytopes in  $\mathcal{N} \otimes \mathbb{R}$  up to suitable isomorphisms.*

With the description of the moment polytope, we can compute the greatest Ricci lower bound of a Fano horospherical manifold.

**Proposition 2.8** ([DH20, Corollary 1.3]). *The greatest Ricci lower bound  $R(X)$  of a Fano horospherical manifold  $X$  is equal to*

$$\sup \left\{ t \in (0, 1) : 2\rho_P + \frac{t}{1-t} (2\rho_P - \mathbf{bar}_{DH}(\Delta)) \in \text{Relint}(\Delta) \right\},$$

where  $\text{Relint}(\Delta)$  means the relative interior of the moment polytope and  $\mathbf{bar}_{DH}(\Delta)$  is the barycenter of the moment polytope  $\Delta(X, K_X^{-1})$  with respect to the Duistermaat–Heckman measure

$$\prod_{\alpha \in \Phi_{Pu}} \kappa(\alpha, p) dp.$$

Here,  $\kappa$  denotes the Killing form on the Lie algebra  $\mathfrak{g}$  of  $G$  and  $p \in \mathfrak{X}(T) \otimes \mathbb{R}$ .

By Proposition 2.8, we immediately get the following elementary geometric expression for  $R(X)$ .

**Corollary 2.9.** *Let  $X$  be a Fano horospherical manifold without a Kähler–Einstein metric. Denote by  $R(X)$  the greatest Ricci lower bound of  $X$ . Let  $A$  be the point corresponding to  $2\rho_P \in \mathfrak{X}(T)$  and  $B$  the barycenter  $\mathbf{bar}_{DH}(\Delta)$  of the moment polytope  $\Delta(X, K_X^{-1})$  with respect to the Duistermaat–Heckman measure. If  $Q$  is the point at which the half-line starting from the barycenter  $B$  in the direction of  $A$  intersects the boundary of the moment polytope  $\Delta$ , then we have  $R(X) = \frac{|A\overrightarrow{Q}|}{|B\overrightarrow{Q}|}$ .*

**2.3. Projective horospherical manifolds of Picard number one.** In this subsection,  $X$  is a projective horospherical manifold of Picard number one with an open  $G$ -orbit  $G/H$  and  $P$  a parabolic subgroup of  $G$ . Then  $X$  is Fano since it is of Picard number one and every spherical variety is rational (see [Per14, Corollary 2.1.3]).

Let  $\{\alpha_1, \dots, \alpha_n\}$  be a system of simple roots of  $G$  following the standard numbering (e.g. [Hum72]) and  $\varpi_i$  be the  $i$ -th fundamental weight. Let  $P^{\alpha_i}$  denotes the maximal parabolic subgroup associated to the simple root  $\alpha_i$  for each  $i$  and  $V(\varpi_i)$  be the irreducible  $G$ -representation with the highest weight  $\varpi_i$ .

For a highest weight vector  $v_i$  of  $V(\varpi_i)$ , the  $G$ -orbit of  $[v_i]$  in  $\mathbb{P}(V(\varpi_i))$  is closed and isomorphic to  $G/P^{\alpha_i}$ , which is a rational homogeneous manifold and denoted by  $(G, \alpha_i)$ . If  $v_i$  and  $v_j$  are highest weight vectors of  $V(\varpi_i)$  and  $V(\varpi_j)$  such that  $i \neq j$ , the open  $G$ -orbit of  $[v_i + v_j] \in \mathbb{P}(V(\varpi_i) \oplus V(\varpi_j))$  is isomorphic to a  $\mathbb{C}^*$ -bundle over a rational homogeneous manifold  $G/P$  where  $P = P^{\alpha_i} \cap P^{\alpha_j}$ . Since the closure of the open  $G$ -orbit is a normal variety according to [Hon16, Proposition 2.1], it is a horospherical  $G$ -variety and we denote it by  $(G, \alpha_i, \alpha_j)$ .

The projective horospherical manifolds of Picard number one are classified by Pasquier [Pas09] using the fact that any nonlinear and  $G$ -nonhomogeneous ones are of the form  $(G, \alpha_i, \alpha_j)$ . In particular, if  $G$ -nonhomogeneous  $X$  is homogeneous under its automorphism group, it is either projective space, quadric  $Q^{2n}$ , Grassmannian  $\text{Gr}(i, n)$  with  $2 \leq i < n$  or spinor variety  $\mathbb{S}_n = \text{Spin}(2n)/P^{\alpha_n}$ , all which are rational homogeneous manifolds.

**Theorem 2.10** ([Pas09, Theorem 0.1]). *Let  $X$  be a projective horospherical  $G$ -manifold of Picard number one. If  $X$  is nonhomogeneous, it is of rank one and its automorphism group is a connected non-reductive linear algebraic group. Moreover,  $X$  is uniquely determined by its two closed  $G$ -orbits  $Y$  and  $Z$ , isomorphic to rational homogeneous manifolds  $G/P^{\alpha_i}$  and  $G/P^{\alpha_j}$ , respectively, where  $(G, \alpha_i, \alpha_j)$  is one of the following triples:*

- (1)  $X^1(n) := (B_n, \alpha_{n-1}, \alpha_n)$  with  $n \geq 3$ ;
- (2)  $X^2 := (B_3, \alpha_1, \alpha_3)$ ;
- (3)  $X^3(n, k) := (C_n, \alpha_k, \alpha_{k-1})$  with  $n \geq k \geq 2$ ;
- (4)  $X^4 := (F_4, \alpha_2, \alpha_3)$ ;
- (5)  $X^5 := (G_2, \alpha_2, \alpha_1)$ .

Even for nonhomogeneous case, Hong [Hon16, Proposition 5.1] proved that a smooth projective horospherical variety  $X$  of Picard number one can be embedded as a linear section into a rational homogeneous manifold of Picard number one except when it is  $X^1(n)$  for  $n \geq 7$ . However, for  $n \geq 7$  it remains open whether  $X^1(n)$  is a linear section of a rational homogeneous manifold or not. Furthermore, Manivel and Pasquier [MP20] realized the blow-up of  $X$  along the closed orbit  $Z$  as the zero locus of a general section of a vector bundle over some homogeneous space. For numerical invariants such as dimensions and the first Chern classes of  $X$ , see [GPPS21, Section 1.6].

Each of these nonhomogeneous horospherical variety has many interesting geometry. For  $n \geq k \geq 2$ ,  $X^3(n, k)$  is isomorphic to the so-called an *odd symplectic Grassmannian*  $\text{SGr}(k, 2n+1)$  studied in [Mih07], which is defined as a variety parametrizing  $k$ -dimensional isotropic subspaces in a  $(2n+1)$ -dimensional complex vector space equipped with a skew-symmetric bilinear form of maximal rank. They appears as smooth Schubert varieties in symplectic Grassmannians ([Hon15]) of which standard embeddings are characterized by means of varieties of minimal rational tangents in [KP18]. It is also known that odd symplectic Grassmannians are rigid under global Kähler deformation. This is proved in [Par16] for odd Lagrangian Grassmannians  $X^3(n, n)$  and in [HL19] for general odd symplectic Grassmannians  $X^3(n, k)$ . Furthermore, Hwang and Li [HL19] characterized odd symplectic Grassmannians  $X^3(n, k)$ , among Fano manifolds of Picard number one, by their varieties of minimal rational tangents at a general point. In addition, Kim [Kim17, Section 3] proved the vanishing of the first Lie algebra cohomology of  $X$  for their characterizations. Recently, Kanemitsu [Kan21] proved that the tangent bundles of  $X^4$  and  $X^1(n)$  for  $n \geq 4$  are unstable in the sense of Mumford–Takemoto which disproves a conjecture on stability of tangent bundles of Fano manifolds of Picard number one. We know that  $X^2$  is isomorphic to a general hyperplane section of the 10-dimensional spinor variety  $\mathbb{S}_5 \subset \mathbb{P}^{15}$ . It was studied, for instance, in [Muk89] and [HM05] as a Fano manifold of coindex 3 and as the variety of minimal rational tangents of a 15-dimensional  $F_4$ -homogeneous variety  $F_4/P^{\alpha_4}$ , respectively. About  $X^5$ , there is a result by Pasquier and Perrin [PP10] that construct explicitly a smooth flat Fano deformation from the orthogonal Grassmannian  $(B_3, \alpha_2) = \text{OGr}(2, 7)$  to  $X^5$  using complexified octonions.

Combining Propositions 2.4, 2.5, 2.6, Theorem 2.10 and [Pas09, Theorem 0.2], we can easily compute the moment polytopes of the anticanonical line bundles on nonhomogeneous horospherical manifolds  $(G, \alpha_i, \alpha_j)$  of Picard number one.

**Lemma 2.11.** *Let  $X$  be a nonhomogeneous projective horospherical manifold of Picard number one. If  $X$  is of type  $(G, \alpha_i, \alpha_j)$ , then the spherical weight lattice  $\mathcal{M}$  is generated by  $\varpi_i - \varpi_j$  and the moment polytope  $\Delta(X, K_X^{-1})$  is the line segment  $\{2\rho_P + t(\varpi_i - \varpi_j) : -\langle \alpha_i^\vee, 2\rho_P \rangle \leq t \leq \langle \alpha_j^\vee, 2\rho_P \rangle\}$  in  $2\rho_P + \mathcal{M} \otimes \mathbb{R}$ , where  $2\rho_P$  is the sum of all roots in the unipotent radical of a submaximal parabolic subgroup  $P^{\alpha_i} \cap P^{\alpha_j}$ .*

*Proof.* By Propositions 2.4 and Theorem 2.10, the spherical weight lattice  $\mathcal{M}$  is generated by  $\varpi_i - \varpi_j$ . By [Pas09, Theorem 0.2] and Theorem 2.10, one can see that  $X$  has three  $G$ -orbits and the codimensions of the two closed orbits are at least two. By Proposition 2.5,  $-K_X = \langle \alpha_i^\vee, 2\rho_P \rangle D_i + \langle \alpha_j^\vee, 2\rho_P \rangle D_j$  is a  $B$ -stable divisor representing the anticanonical line bundle  $K_X^{-1}$ . Because  $\rho(D_i) = \alpha_i^\vee$  and  $\rho(D_j) = \alpha_j^\vee$ , Proposition 2.6 gives us the inequalities

$$\left\langle \frac{\alpha_i^\vee}{\langle \alpha_i^\vee, 2\rho_P \rangle}, t(\varpi_i - \varpi_j) \right\rangle \geq -1 \iff t \geq -\langle \alpha_i^\vee, 2\rho_P \rangle$$

and

$$\left\langle \frac{\alpha_j^\vee}{\langle \alpha_j^\vee, 2\rho_P \rangle}, t(\varpi_i - \varpi_j) \right\rangle \geq -1 \iff t \leq \langle \alpha_j^\vee, 2\rho_P \rangle.$$

Therefore, the result follows.  $\square$

### 3. GREATEST RICCI LOWER BOUNDS OF PROJECTIVE HOROSPHERICAL MANIFOLDS OF PICARD NUMBER ONE

In this section we prove Theorem 1.1. In the course of the proof, we explicitly present the algebraic moment polytope using Lemma 2.11 and then compute the greatest Ricci lower bound for every nonhomogeneous projective horospherical manifold of Picard number one using Corollary 2.9. The most tricky part of the proof is to compute the barycenter of the moment polytope with respect to the Duistermaat–Heckman measure.

**3.1. Greatest Ricci lower bound of  $X^5$ .** Let  $X^5$  be the projective horospherical manifold of type  $(G_2, \alpha_2, \alpha_1)$  in Theorem 2.10. The spherical weight lattice  $\mathcal{M}$  is generated by  $\varpi_1 - \varpi_2$  for the fundamental weights  $\varpi_1, \varpi_2 \in \mathfrak{X}(T)$  of a maximal torus  $T \subset G_2$ , so that the dual lattice  $\mathcal{N}$  is generated by  $\frac{1}{2}\alpha_1^\vee - \frac{1}{2}\alpha_2^\vee$  for the coroots  $\alpha_1^\vee$  and  $\alpha_2^\vee$ .

Choose a realization of the root system  $G_2$  in the Euclidean plane  $\mathbb{R}^2$  with  $\alpha_1 = (1, 0)$  and  $\alpha_2 = \left(-\frac{3}{2}, \frac{\sqrt{3}}{2}\right)$ . Then, the complex Lie group  $G_2$  has the six positive roots:

$$\begin{aligned} \Phi^+ &= \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2, 3\alpha_1 + \alpha_2, 3\alpha_1 + 2\alpha_2\} \\ &= \left\{ (1, 0), \left(-\frac{3}{2}, \frac{\sqrt{3}}{2}\right), \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \left(\frac{3}{2}, \frac{\sqrt{3}}{2}\right), (0, \sqrt{3}) \right\}. \end{aligned}$$

From the relation  $(\alpha_i^\vee, \varpi_j) = \delta_{ij}$ , the fundamental weights corresponding to the system of simple roots are  $\varpi_1 = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$  and  $\varpi_2 = (0, \sqrt{3})$ .


**Proposition 3.1.** *The moment polytope  $\Delta_5 = \Delta(X^5, K_{X^5}^{-1})$  is the line segment connecting two points  $4\varpi_1$  and  $4\varpi_2$  in  $\mathfrak{X}(T) \otimes \mathbb{R}$ , that is,*

$$\Delta(X^5, K_{X^5}^{-1}) = \{(2\varpi_1 + 2\varpi_2) + t(\varpi_1 - \varpi_2) : -2 \leq t \leq 2\}.$$

*Proof.* Since  $P = P^{\alpha_1} \cap P^{\alpha_2} = B$ , we have  $2\rho_P = 2\rho_{G_2} = 2\varpi_1 + 2\varpi_2$  and the assertion follows from Lemma 2.11 (see Figure 1).  $\square$

**Remark 3.2.** Considering the minimal embedding  $X^5 \hookrightarrow \mathbb{P}(V_{G_2}(\varpi_1) \oplus V_{G_2}(\varpi_2))$  and  $K_{X^5}^{-1} = \mathcal{O}(4)$ , motivated by the original definition of Brion, we have

$$\begin{aligned} H^0(X^5, \mathcal{O}(1)) &= V_{G_2}(\varpi_1) \oplus V_{G_2}(\varpi_2) \quad \text{and} \\ H^0(X^5, K_{X^5}^{-1}) &= V_{G_2}(4\varpi_1) \oplus V_{G_2}(3\varpi_1 + \varpi_2) \oplus V_{G_2}(2\varpi_1 + 2\varpi_2) \oplus V_{G_2}(\varpi_1 + 3\varpi_2) \oplus V_{G_2}(4\varpi_2). \end{aligned}$$

FIGURE 1. Moment polytope  $\Delta_5 = \Delta(X^5, K_{X^5}^{-1})$ .

**Proposition 3.3.** *The greatest Ricci lower bound of  $X^5$  is  $R(X^5) = \frac{56}{67} \approx 0.8358$ .*

*Proof.* For  $p = (x, y)$ , the Duistermaat–Heckman measure on  $\mathfrak{X}(T) \otimes \mathbb{R}$  is given as

$$\prod_{\alpha \in \Phi^+} \kappa(\alpha, p) dp = x \left( -\frac{3}{2}x + \frac{\sqrt{3}}{2}y \right) \left( -\frac{1}{2}x + \frac{\sqrt{3}}{2}y \right) \left( \frac{1}{2}x + \frac{\sqrt{3}}{2}y \right) \left( \frac{3}{2}x + \frac{\sqrt{3}}{2}y \right) (\sqrt{3}y) dx dy =: P_{DH}(x, y) dx dy.$$

From Proposition 3.1, the moment polytope  $\Delta_5$  is parametrized as

$$\gamma_5(t) = \left( 1 + \frac{t}{2}, \frac{\sqrt{3}}{2}(2+t) + \sqrt{3}(2-t) \right)$$

for  $-2 \leq t \leq 2$ . Then, we can compute the volume (or length)

$$\text{Vol}_{DH}(\Delta_5) = \int_{-2}^2 P_{DH}(\gamma_5(t)) dt = 9216,$$

and the barycenter

$$\mathbf{bar}_{DH}(\Delta_5) = (\bar{x}, \bar{y}) = \frac{1}{\text{Vol}_{DH}(\Delta_5)} \int_{\Delta_5} p \prod_{\alpha \in \Phi^+} \kappa(\alpha, p) dp = \left( \frac{45}{56}, \frac{179\sqrt{3}}{56} \right) \approx (0.804, 3.196 \times \sqrt{3})$$

of the moment polytope  $\Delta_5$  with respect to the Duistermaat–Heckman measure. Since  $\mathbf{bar}_{DH}(\Delta_5) = \gamma_5(-\frac{11}{28})$ , we get the greatest Ricci lower bound  $R(X^5) = \frac{2}{2 + \frac{11}{28}} = \frac{56}{67}$  by Corollary 2.9.  $\square$

Using the basis of the weight lattice instead of the orthogonal coordinate system is more effective for calculations of the remaining cases. From the relations

$$\kappa(\alpha_1, \varpi_1) = \frac{1}{2}, \quad \kappa(\alpha_2, \varpi_1) = 0, \quad \kappa(\alpha_1, \varpi_2) = 0, \quad \kappa(\alpha_2, \varpi_2) = \frac{3}{2},$$



we have the Duistermaat–Heckman polynomial evaluated on the moment polytope  $\Delta_5$

$$\begin{aligned} P_{DH}(\gamma_5(t)) &= \prod_{\alpha \in \Phi^+} \kappa(\alpha, (2+t)\varpi_1 + (2-t)\varpi_2) \\ &= \frac{1}{2}(2+t) \cdot \frac{3}{2}(2-t) \cdot \left\{ \frac{1}{2}(2+t) + \frac{3}{2}(2-t) \right\} \cdot \left\{ 2 \cdot \frac{1}{2}(2+t) + \frac{3}{2}(2-t) \right\} \\ &\quad \times \left\{ 3 \cdot \frac{1}{2}(2+t) + \frac{3}{2}(2-t) \right\} \cdot \left\{ 3 \cdot \frac{1}{2}(2+t) + 2 \cdot \frac{3}{2}(2-t) \right\}. \end{aligned}$$

As  $\frac{1}{\text{Vol}_{DH}(\Delta_5)} \int_{-2}^2 (2+t) \cdot P_{DH}(\gamma_5(t)) dt = \frac{45}{28}$  and  $\frac{1}{\text{Vol}_{DH}(\Delta_5)} \int_{-2}^2 (2-t) \cdot P_{DH}(\gamma_5(t)) dt = \frac{67}{28}$ , we get the same result  $\mathbf{bar}_{DH}(\Delta_5) = \frac{45}{28}\varpi_1 + \frac{67}{28}\varpi_2 = \gamma_5\left(-\frac{11}{28}\right)$ .

**3.2. Greatest Ricci lower bound of  $X^2$ .** Let  $X^2$  be the projective horospherical manifold of type  $(B_3, \alpha_1, \alpha_3)$  in Theorem 2.10.

**Proposition 3.4.** *The moment polytope  $\Delta_2 = \Delta(X^2, K_{X^2}^{-1})$  is the line segment connecting two points  $7\varpi_1$  and  $7\varpi_3$  in  $\mathfrak{X}(T) \otimes \mathbb{R}$ , that is,*

$$\Delta(X^2, K_{X^2}^{-1}) = \{(3\varpi_1 + 4\varpi_3) + t(\varpi_1 - \varpi_3) : -3 \leq t \leq 4\}.$$

*Proof.* Since the stabilizer of the open  $B$ -orbit in  $X^2$  is  $P$ , we know that  $\Phi_{P^u} = \Phi_{B_3}^+ \setminus \{\alpha_2\}$ . Recall that the fundamental weights  $\varpi_1, \varpi_2, \varpi_3$  and the second simple root  $\alpha_2$  of  $B_3$  can be expressed as  $\varpi_1 = L_1, \varpi_2 = L_1 + L_2, \varpi_3 = \frac{L_1 + L_2 + L_3}{2}$  and  $\alpha_2 = L_2 - L_3$  in terms of an orthonormal basis  $\{L_1, L_2, L_3\}$  of the dual Cartan subalgebra. Then, as  $\alpha_2 = -\varpi_1 + 2\varpi_2 - 2\varpi_3$  we have

$$2\rho_P = 2\rho_{B_3} - 2\rho_L = 2(\varpi_1 + \varpi_2 + \varpi_3) - \alpha_2 = 2(\varpi_1 + \varpi_2 + \varpi_3) - (-\varpi_1 + 2\varpi_2 - 2\varpi_3) = 3\varpi_1 + 4\varpi_3.$$

The result follows from Lemma 2.11.  $\square$

**Proposition 3.5.** *The greatest Ricci lower bound of  $X^2$  is  $R(X^2) = \frac{20}{21} \approx 0.952$ .*

*Proof.* By Proposition 3.4, the moment polytope  $\Delta_2$  is parametrized as

$$\gamma_2(t) = (3+t)\varpi_1 + (4-t)\varpi_3$$

for  $-3 \leq t \leq 4$ . From the relations

$$\kappa(\alpha_1, \varpi_1) = 1, \kappa(\alpha_2, \varpi_1) = 0, \kappa(\alpha_3, \varpi_1) = 0, \quad \kappa(\alpha_1, \varpi_3) = 0, \kappa(\alpha_2, \varpi_3) = 0, \kappa(\alpha_3, \varpi_3) = \frac{1}{2}$$

and  $\Phi_{P^u} = \Phi_{B_3}^+ \setminus \{\alpha_2\} = \{\alpha_1, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, \alpha_2 + 2\alpha_3, \alpha_1 + \alpha_2 + 2\alpha_3, \alpha_1 + 2\alpha_2 + 2\alpha_3\}$ , we have the Duistermaat–Heckman polynomial evaluated on the moment polytope  $\Delta_2$

$$\begin{aligned} P_{DH}(\gamma_2(t)) &= \prod_{\alpha \in \Phi_{P^u}} \kappa(\alpha, (3+t)\varpi_1 + (4-t)\varpi_3) \\ &= (3+t) \cdot \frac{1}{2}(4-t) \cdot (3+t) \cdot \frac{1}{2}(4-t) \cdot \left\{ (3+t) + \frac{1}{2}(4-t) \right\} \cdot \left\{ 2 \cdot \frac{1}{2}(4-t) \right\} \\ &\quad \times \left\{ (3+t) + 2 \cdot \frac{1}{2}(4-t) \right\} \cdot \left\{ (3+t) + 2 \cdot \frac{1}{2}(4-t) \right\} \\ &= \frac{49}{4}(3+t)^2(4-t)^3 \left( 5 + \frac{t}{2} \right). \end{aligned}$$

As  $\frac{1}{\text{Vol}_{DH}(\Delta_2)} \int_{-3}^4 (3+t) \cdot P_{DH}(\gamma_2(t)) dt = \frac{63}{20}$  and  $\frac{1}{\text{Vol}_{DH}(\Delta_2)} \int_{-3}^4 (4-t) \cdot P_{DH}(\gamma_2(t)) dt = \frac{77}{20}$ , we get the barycenter  $\mathbf{bar}_{DH}(\Delta_2) = \frac{63}{20}\varpi_1 + \frac{77}{20}\varpi_3 = \gamma_2\left(\frac{3}{20}\right)$ . Therefore, the greatest Ricci lower bound of  $X^2$  is  $R(X^2) = \frac{3}{3 + \frac{3}{20}} = \frac{20}{21}$  by Corollary 2.9.  $\square$

**3.3. Greatest Ricci lower bound of  $X^4$ .** Let  $X^4$  be the projective horospherical manifold of type  $(F_4, \alpha_2, \alpha_3)$  in Theorem 2.10.

**Proposition 3.6.** *The moment polytope  $\Delta_4 = \Delta(X^4, K_{X^4}^{-1})$  is the line segment connecting two points  $6\varpi_2$  and  $6\varpi_3$  in  $\mathfrak{X}(T) \otimes \mathbb{R}$ , that is,*

$$\Delta(X^4, K_{X^4}^{-1}) = \{(3\varpi_2 + 3\varpi_3) + t(\varpi_2 - \varpi_3) : -3 \leq t \leq 3\}.$$

*Proof.* Since  $P = P^{\alpha_2} \cap P^{\alpha_3}$ , we know that  $\Phi_{P^u} = \Phi_{F_4}^+ \setminus \{\alpha_1, \alpha_4\}$  and  $2\rho_P = 2\rho_{F_4} - \alpha_1 - \alpha_4 = 2(\varpi_1 + \varpi_2 + \varpi_3 + \varpi_4) - (2\varpi_1 - \varpi_2 - \varpi_3 + 2\varpi_4) = 3\varpi_2 + 3\varpi_3$ . The result follows from Lemma 2.11.  $\square$

**Proposition 3.7.** *The greatest Ricci lower bound of  $X^4$  is  $R(X^4) = \frac{178992099}{243545402} \approx 0.734$ .*

*Proof.* By Proposition 3.6, the moment polytope  $\Delta_4$  is parametrized as

$$\gamma_4(t) = (3+t)\varpi_2 + (3-t)\varpi_3$$

for  $-3 \leq t \leq 3$ . From the relations

$$\begin{aligned} \kappa(\alpha_1, \varpi_2) &= 0, \kappa(\alpha_2, \varpi_2) = 1, \kappa(\alpha_3, \varpi_2) = 0, \kappa(\alpha_4, \varpi_2) = 0, \\ \kappa(\alpha_1, \varpi_3) &= 0, \kappa(\alpha_2, \varpi_3) = 0, \kappa(\alpha_3, \varpi_3) = \frac{1}{2}, \kappa(\alpha_4, \varpi_3) = 0 \end{aligned}$$

and  $\Phi_{P^u} = \Phi_{F_4}^+ \setminus \{\alpha_1, \alpha_4\} = \{\alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_3 + \alpha_4, \alpha_1 + \alpha_2 + \alpha_3, \alpha_2 + 2\alpha_3, \alpha_2 + \alpha_3 + \alpha_4, \alpha_1 + \alpha_2 + 2\alpha_3, \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4, \alpha_2 + 2\alpha_3 + \alpha_4, \alpha_1 + 2\alpha_2 + 2\alpha_3, \alpha_1 + \alpha_2 + 2\alpha_3 + \alpha_4, \alpha_2 + 2\alpha_3 + 2\alpha_4, \alpha_1 + 2\alpha_2 + 2\alpha_3 + \alpha_4, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4, \alpha_1 + 2\alpha_2 + 3\alpha_3 + \alpha_4, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4, \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4, \alpha_1 + 2\alpha_2 + 4\alpha_3 + 2\alpha_4, \alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4, 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4\}$ , we have the Duistermaat–Heckman polynomial evaluated on the moment polytope  $\Delta_4$

$$\begin{aligned} P_{DH}(\gamma_4(t)) &= \prod_{\alpha \in \Phi_{P^u}} \kappa(\alpha, (3+t)\varpi_2 + (3-t)\varpi_3) \\ &= (3+t)^2 \left\{ \frac{1}{2}(3-t) \right\}^2 \left\{ (3+t) + \frac{1}{2}(3-t) \right\}^4 \left\{ (3+t) + 2 \cdot \frac{1}{2}(3-t) \right\}^6 \left\{ 2(3+t) + 2 \cdot \frac{1}{2}(3-t) \right\}^3 \\ &\quad \times \left\{ 2(3+t) + 3 \cdot \frac{1}{2}(3-t) \right\}^2 \left\{ 2(3+t) + 4 \cdot \frac{1}{2}(3-t) \right\} \left\{ 3(3+t) + 4 \cdot \frac{1}{2}(3-t) \right\}^2. \end{aligned}$$

$$\text{As } \frac{1}{\text{Vol}_{DH}(\Delta_4)} \int_{-3}^3 (3+t) \cdot P_{DH}(\gamma_4(t)) dt = \frac{243545402}{59664033} \text{ and } \frac{1}{\text{Vol}_{DH}(\Delta_4)} \int_{-3}^3 (3-t) \cdot P_{DH}(\gamma_4(t)) dt = \frac{114438796}{59664033},$$

we get the barycenter  $\mathbf{bar}_{DH}(\Delta_4) = \frac{243545402}{59664033}\varpi_2 + \frac{114438796}{59664033}\varpi_3 = \gamma_4\left(\frac{64553303}{59664033}\right)$ . Therefore, the greatest Ricci lower bound of  $X^4$  is  $R(X^4) = \frac{3}{3 + \frac{64553303}{59664033}} = \frac{178992099}{243545402}$  by Corollary 2.9.  $\square$

**3.4. Greatest Ricci lower bounds of  $X^1(n)$ .** For  $n \geq 3$ , let  $X^1(n)$  be the projective horospherical manifold of type  $(B_n, \alpha_{n-1}, \alpha_n)$  in Theorem 2.10.

**Proposition 3.8.** *The moment polytope  $\Delta_1 = \Delta(X^1(n), K_{X^1(n)}^{-1})$  is the line segment connecting two points  $(n+2)\varpi_{n-1}$  and  $(n+2)\varpi_n$  in  $\mathfrak{X}(T) \otimes \mathbb{R}$ , that is,*

$$\Delta(X^1(n), K_{X^1(n)}^{-1}) = \{(n\varpi_{n-1} + 2\varpi_n) + t(\varpi_{n-1} - \varpi_n) : -n \leq t \leq 2\}.$$

*Proof.* Let  $L_1, L_2, \dots, L_n$  be the orthonormal basis of  $\mathfrak{X}(T) \otimes \mathbb{Q}$ . Then the set of all positive roots of  $B_n$  consists of the following: for  $1 \leq i < j \leq n$ ,

$$L_i - L_j, \quad L_i + L_j, \quad L_i.$$

Hence,  $2\rho_G = \sum_{i < j} (L_i - L_j) + \sum_{i < j} (L_i + L_j) + \sum_{i=1}^n L_i = \sum_{i=1}^n (2n - 2i + 1)L_i$ . On the other hand, since the reductive part  $L$  of  $P = P^{\alpha_{n-1}} \cap P^{\alpha_n}$  is of type  $A_{n-2}$ , the positive roots of  $L$  are  $L_i - L_j$  for  $1 \leq i < j \leq n-1$  and  $2\rho_L = \sum_{1 \leq i < j \leq n-1} (L_i - L_j) = \sum_{i=1}^{n-1} (n-2i)L_i$ . Therefore,

$$2\rho_G - 2\rho_L = \sum_{i=1}^n (2n - 2i + 1)L_i - \sum_{i=1}^{n-1} (n-2i)L_i = n \sum_{i=1}^{n-1} L_i + \sum_{i=1}^n L_i = n\varpi_{n-1} + 2\varpi_n.$$

The result follows from Lemma 2.11.  $\square$

**Proposition 3.9.** *The greatest Ricci lower bound of  $X^1(n)$  is*

$$R(X^1(n)) = \frac{n \int_{-n}^2 (2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt}{\int_{-n}^2 (2-t)(n+t)^n(t+2n+2)^{\frac{n(n-1)}{2}} dt}.$$

*Proof.* By Proposition 3.8, the moment polytope  $\Delta_1$  is parametrized as

$$\gamma_1(t) = (n+t)\varpi_{n-1} + (2-t)\varpi_n$$

for  $-n \leq t \leq 2$ . In order to obtain the Duistermaat–Heckman polynomial evaluated on the moment polytope  $\Delta_1$ , the number of roots in  $\Phi_{Pu}$  for each value of the coefficients of  $\alpha_{n-1}, \alpha_n$  must be counted specifically:

- (i) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 1, c_n = 0\}$  is  $n-1$ ,
- (ii) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 0, c_n = 1\}$  is 1,
- (iii) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 1, c_n = 1\}$  is  $n-1$ ,
- (iv) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 1, c_n = 2\}$  is  $n-1$ , and
- (v) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 2, c_n = 2\}$  is  $\frac{(n-1)(n-2)}{2}$ .

Then, from  $\kappa(\alpha_{n-1}, \varpi_{n-1}) = 1$  and  $\kappa(\alpha_n, \varpi_n) = \frac{1}{2}$  we have

$$\begin{aligned} P_{DH}(\gamma_1(t)) &= \prod_{\alpha \in \Phi_{Pu}} \kappa(\alpha, (n+t)\varpi_{n-1} + (2-t)\varpi_n) \\ &= (n+t)^{n-1} \cdot \frac{1}{2}(2-t) \left\{ (n+t) + \frac{1}{2}(2-t) \right\}^{n-1} \\ &\quad \times \left\{ (n+t) + 2 \cdot \frac{1}{2}(2-t) \right\}^{n-1} \left\{ 2(n+t) + 2 \cdot \frac{1}{2}(2-t) \right\}^{\frac{(n-1)(n-2)}{2}} \\ &= (n+t)^{n-1} \cdot \frac{1}{2}(2-t) \left( \frac{1}{2}t + n + 1 \right)^{n-1} (n+2)^{n-1} (t+2n+2)^{\frac{(n-1)(n-2)}{2}} \\ &= \frac{(n+2)^{n-1}}{2^n} (2-t)(n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}}. \end{aligned}$$

As  $\frac{1}{\text{Vol}_{DH}(\Delta_1)} \int_{-n}^2 (2-t) \cdot P_{DH}(\gamma_1(t)) dt = \frac{\int_{-n}^2 (2-t)^2 (n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt}{\int_{-n}^2 (2-t)(n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt}$ , we get the barycenter

$\text{bar}_{DH}(\Delta_1) = \gamma_1 \left( 2 - \frac{\int_{-n}^2 (2-t)^2 (n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt}{\int_{-n}^2 (2-t)(n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt} \right)$  of the moment polytope  $\Delta_1$  with

respect to the Duistermaat–Heckman measure. Since  $2 - \frac{\int_{-n}^2 (2-t)^2 (n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt}{\int_{-n}^2 (2-t)(n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt} > 0$  by

Lemma 3.10, the greatest Ricci lower bound of  $X^1(n)$  is

$$\begin{aligned} R(X^1(n)) &= \frac{n}{\left( 2 - \frac{\int_{-n}^2 (2-t)^2 (n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt}{\int_{-n}^2 (2-t)(n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt} \right) - (-n)} \\ &= \frac{n \int_{-n}^2 (2-t)(n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt}{(n+2) \int_{-n}^2 (2-t)(n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt - \int_{-n}^2 (2-t)^2 (n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt} \\ &= \frac{n \int_{-n}^2 (2-t)(n+t)^{n-1} (t+2n+2)^{\frac{n(n-1)}{2}} dt}{\int_{-n}^2 (2-t)(n+t)^n (t+2n+2)^{\frac{n(n-1)}{2}} dt} \end{aligned}$$

by Corollary 2.9.  $\square$

**Lemma 3.10.** *For any  $n \geq 3$ , the inequality*

$$\int_{-n}^2 (2-t)^2(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt < 2 \int_{-n}^2 (2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt$$

holds.

*Proof.* It suffices to show that  $\int_{-n}^2 t(2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt > 0$  for  $n \geq 3$ . For a fixed  $n$ , since  $t+2n+2$  is an increasing function,  $t(2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} > t(2-t)(n+t)^{n-1}(2n+2)^{\frac{n(n-1)}{2}}$  for  $0 < t < 2$ . On the other hand, when  $-n < t < 0$  we also have the same inequality  $t(2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} > t(2-t)(n+t)^{n-1}(2n+2)^{\frac{n(n-1)}{2}}$  since  $t < 0$ . Thus,

$$\begin{aligned} & \int_{-n}^2 t(2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt \\ &= \int_{-n}^0 t(2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt + \int_0^2 t(2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt \\ &> \int_{-n}^0 t(2-t)(n+t)^{n-1}(2n+2)^{\frac{n(n-1)}{2}} dt + \int_0^2 t(2-t)(n+t)^{n-1}(2n+2)^{\frac{n(n-1)}{2}} dt \\ &= (2n+2)^{\frac{n(n-1)}{2}} \int_{-n}^2 t(2-t)(n+t)^{n-1} dt \\ &= (2n+2)^{\frac{n(n-1)}{2}} \int_0^{n+2} (s-n)(n+2-s)s^{n-1} ds \quad (\text{let } s := n+t) \\ &= (2n+2)^{\frac{n(n-1)}{2}} \int_0^{n+2} \{-s^{n+1} + 2(n+1)s^n - n(n+2)s^{n-1}\} ds \\ &= (2n+2)^{\frac{n(n-1)}{2}} \left[ -\frac{s^{n+2}}{n+2} + 2(n+1) \cdot \frac{s^{n+1}}{n+1} - n(n+2) \cdot \frac{s^n}{n} \right]_0^{n+2} = 0 \end{aligned}$$

as claimed. The proof is complete.  $\square$

The approximate values of the greatest Ricci lower bounds  $R(X^1(n))$  for some  $n$  are summarized as shown in the following (the first row of Table 2).

$n$	3	4	5	6	7	10	20	30	50	70
$R(X^1(n))$	0.8955	0.8755	0.8686	0.8685	0.8715	0.8863	0.9251	0.9451	0.9644	0.9737

Interestingly, we observe that the value  $R(X^1(n))$  decreases from  $n = 3$  to 6 and then increases again from  $n = 7$ . In addition,  $R(X^1(n))$  becomes arbitrarily close to 1 as  $n$  becomes larger and larger.

**Corollary 3.11.** *The greatest Ricci lower bound  $R(X^1(n))$  of  $X^1(n)$  converges to 1 as  $n$  increases, that is,  $\lim_{n \rightarrow \infty} R(X^1(n)) = 1$ .*

*Proof.* From Proposition 3.9, we have  $R(X^1(n)) = \frac{n}{n+2 - \frac{\int_{-n}^2 (2-t)^2(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt}{\int_{-n}^2 (2-t)(n+t)^{n-1}(t+2n+2)^{\frac{n(n-1)}{2}} dt}}$ . Since all

functions  $2-t$ ,  $n+t$  and  $t+2n+2$  in the integrands are positive for  $-n < t < 2$ , we get an inequality  $R(X^1(n)) > \frac{n}{n+2}$ ; hence  $\lim_{n \rightarrow \infty} R(X^1(n)) \geq \lim_{n \rightarrow \infty} \frac{n}{n+2} = 1$ . As  $R(X^1(n)) \leq 1$  by definition, we conclude that  $\lim_{n \rightarrow \infty} R(X^1(n)) = 1$ .  $\square$

**3.5. Greatest Ricci lower bounds of  $X^3(n, k)$ .** For  $n \geq k \geq 2$ , let  $X^3(n, k)$  be the projective horospherical manifold of type  $(C_n, \alpha_k, \alpha_{k-1})$  in Theorem 2.10.

**Proposition 3.12.** *The moment polytope  $\Delta_3 = \Delta(X^3(n, k), K_{X^3(n, k)}^{-1})$  is the line segment connecting two points  $(2n-k+2)\varpi_{k-1}$  and  $(2n-k+2)\varpi_k$  in  $\mathfrak{X}(T) \otimes \mathbb{R}$ , that is,*

$$\Delta(X^3(n, k), K_{X^3(n, k)}^{-1}) = \{(k\varpi_{k-1} + (2n-2k+2)\varpi_k) + t(\varpi_{k-1} - \varpi_k) : -k \leq t \leq 2n-2k+2\}.$$

*Proof.* Let  $L_1, L_2, \dots, L_n$  be the orthonormal basis of  $\mathfrak{X}(T) \otimes \mathbb{Q}$ . Then the set of all positive roots of  $C_n$  consists of the following: for  $1 \leq i < j \leq n$ ,

$$L_i - L_j, \quad L_i + L_j, \quad 2L_i.$$

Hence,  $2\rho_G = \sum_{i < j} (L_i - L_j) + \sum_{i < j} (L_i + L_j) + 2 \sum_{i=1}^n L_i = \sum_{i=1}^n (2n - 2i + 2)L_i$ . On the other hand, since the reductive part  $L$  of  $P = P^{\alpha_{k-1}} \cap P^{\alpha_k}$  is of type  $A_{k-2} \times C_{n-k}$ , the positive roots of  $L$  are  $L_i - L_j$  for  $1 \leq i < j \leq k-1$ ,  $L_i \pm L_j$  for  $k+1 \leq i < j \leq n$ , and  $2L_i$  for  $i = k+1, \dots, n$ . Therefore, we have  $2\rho_L = \sum_{i=1}^{k-1} (k-2i)L_i + \sum_{i=k+1}^n (2n-2i+2)L_i$  and

$$\begin{aligned} 2\rho_G - 2\rho_L &= \sum_{i=1}^k (2n-2i+2)L_i - \sum_{i=1}^{k-1} (k-2i)L_i \\ &= (2n-2k+2) \sum_{i=1}^k L_i + k \sum_{i=1}^{k-1} L_i = (2n-2k+2)\varpi_k + k\varpi_{k-1}. \end{aligned}$$

The result follows from Lemma 2.11.  $\square$

As  $\alpha_n$  is a unique long simple root of the symplectic group  $C_n = \mathrm{Sp}(2n, \mathbb{C})$  and the other simple roots are short, it is necessary to separate the calculations in the two cases: (Case I)  $k = n$ , (Case II)  $2 \leq k \leq n-1$ .

**Proposition 3.13.** *For  $n \geq 2$ , the greatest Ricci lower bound of an odd Lagrangian Grassmannian  $X^3(n, n)$  is*

$$R(X^3(n, n)) = \frac{2}{(n+2) \int_0^1 (1-t^2)^n dt} = \frac{2 \times (2n+1)!}{(n+2)(2^n \times n!)^2}.$$

*Proof.* By Proposition 3.12, the moment polytope  $\Delta_3$  of  $X^3(n, n)$  is parametrized as

$$\gamma_3(t) = (n+t)\varpi_{n-1} + (2-t)\varpi_n$$

for  $-n \leq t \leq 2$ . Let's count the number of roots in  $\Phi_{P^u}$  for each value of the coefficients of  $\alpha_{n-1}, \alpha_n$ :

- (i) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 1, c_n = 0\}$  is  $n-1$ ,
- (ii) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 0, c_n = 1\}$  is 1,
- (iii) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 1, c_n = 1\}$  is  $n-1$ , and
- (iv) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{n-1} = 2, c_n = 1\}$  is  $\frac{n(n-1)}{2}$ .

Then, from  $\kappa(\alpha_{n-1}, \varpi_{n-1}) = 1$  and  $\kappa(\alpha_n, \varpi_n) = 2$  we have

$$\begin{aligned} P_{DH}(\gamma_3(t)) &= \prod_{\alpha \in \Phi_{P^u}} \kappa(\alpha, (n+t)\varpi_{n-1} + (2-t)\varpi_n) \\ &= (n+t)^{n-1} \cdot 2(2-t) \{(n+t) + 2(2-t)\}^{n-1} \{2(n+t) + 2(2-t)\}^{\frac{n(n-1)}{2}} \\ &= 2(2n+4)^{\frac{n(n-1)}{2}} \cdot (n+t)^{n-1} (2-t)(n+4-t)^{n-1}. \end{aligned}$$

Thus, we get the volume of the moment polytope  $\Delta_3$  with respect to the Duistermaat–Heckman measure

$$\begin{aligned} \mathrm{Vol}_{DH}(\Delta_3) &= \int_{-n}^2 P_{DH}(\gamma_3(t)) dt = \int_{-n}^2 2(2n+4)^{\frac{n(n-1)}{2}} \cdot (n+t)^{n-1} (2-t)(n+4-t)^{n-1} dt \\ &= 2(2n+4)^{\frac{n(n-1)}{2}} \int_{-n}^2 (2-t) \{(n+t)(n+4-t)\}^{n-1} dt \\ &= 2(2n+4)^{\frac{n(n-1)}{2}} \int_{-n}^2 (2-t) \{-(t-2)^2 + (n+2)^2\}^{n-1} dt \\ &= 2(2n+4)^{\frac{n(n-1)}{2}} \left[ \frac{1}{2n} \{-(t-2)^2 + (n+2)^2\}^n \right]_{-n}^2 \\ &= 2(2n+4)^{\frac{n(n-1)}{2}} \cdot \frac{(n+2)^{2n}}{2n} \end{aligned}$$

As we know that  $\int (2-t)\{-(t-2)^2 + (n+2)^2\}^{n-1} dt = \frac{1}{2n}\{-(t-2)^2 + (n+2)^2\}^n + C$  from the previous calculation result, we can use integration by parts

$$\begin{aligned} \int_{-n}^2 (2-t) \cdot \frac{P_{DH}(\gamma_3(t))}{2(2n+4)^{\frac{n(n-1)}{2}}} dt &= \int_{-n}^2 (2-t) \cdot (2-t)\{-(t-2)^2 + (n+2)^2\}^{n-1} dt \\ &= \left[ (2-t) \cdot \frac{1}{2n}\{-(t-2)^2 + (n+2)^2\}^n \right]_{-n}^2 - \int_{-n}^2 (-1) \cdot \frac{1}{2n}\{-(t-2)^2 + (n+2)^2\}^n dt \\ &= \frac{1}{2n} \int_{-n}^2 \{-(t-2)^2 + (n+2)^2\}^n dt = \frac{(n+2)^{2n}}{2n} \int_{-n}^2 \left\{ 1 - \left( \frac{2-t}{n+2} \right)^2 \right\}^n dt \\ &= \frac{(n+2)^{2n}}{2n} \int_1^0 (1-s^2)^n (-n+2) ds = \frac{(n+2)^{2n}}{2n} \cdot (n+2) \int_0^1 (1-s^2)^n ds \end{aligned}$$

Here, we use the substitution  $s = \frac{2-t}{n+2}$  for the last line of equations. Thus, we have

$$\frac{1}{\text{Vol}_{DH}(\Delta_3)} \int_{-n}^2 (2-t) \cdot P_{DH}(\gamma_3(t)) dt = (n+2) \int_0^1 (1-t^2)^n dt,$$

from which we get the barycenter  $\mathbf{bar}_{DH}(\Delta_3) = \gamma_3 \left( 2 - (n+2) \int_0^1 (1-t^2)^n dt \right)$  of the moment polytope  $\Delta_3$  with respect to the Duistermaat–Heckman measure. Since  $2 - (n+2) \int_0^1 (1-t^2)^n dt < 0$  by the first statement of Lemma 3.14, the greatest Ricci lower bound of  $X^3(n)$  is

$$R(X^3(n, n)) = \frac{2}{2 - \{2 - (n+2) \int_0^1 (1-t^2)^n dt\}} = \frac{2}{(n+2) \int_0^1 (1-t^2)^n dt} = \frac{2 \times (2n+1)!}{(n+2)(2^n \times n!)^2}$$

by Corollary 2.9 and the second statement of Lemma 3.14.  $\square$

**Lemma 3.14.** *For any  $n \geq 2$ , the inequality  $(n+2) \int_0^1 (1-t^2)^n dt > 2$  holds. Furthermore, we have  $(n+2) \int_0^1 (1-t^2)^n dt = \frac{(n+2)(2^n \times n!)^2}{(2n+1)!}$  for any  $n \geq 0$ .*

*Proof.* Putting a sequence  $a_n := (n+2) \int_0^1 (1-t^2)^n dt$  for  $n \geq 0$ , we will find a recurrence relation for  $a_n$ . Using integration by parts, we have

$$\begin{aligned} \int_0^1 (1-t^2)^{n+1} dt &= \left[ t \cdot (1-t^2)^{n+1} \right]_0^1 - \int_0^1 t \cdot (n+1)(1-t^2)^n (-2t) dt \\ &= 2(n+1) \int_0^1 t^2 (1-t^2)^n dt \\ &= \int_0^1 (1-t^2)^n dt - \int_0^1 t^2 (1-t^2)^n dt \end{aligned}$$

from which we deduce that  $\int_0^1 (1-t^2)^n dt = (2n+3) \int_0^1 t^2 (1-t^2)^n dt$ . Then,

$$\begin{aligned} a_{n+1} &= \{(n+1)+2\} \int_0^1 (1-t^2)^{n+1} dt = (n+3) \int_0^1 \{(1-t^2)^n - t^2(1-t^2)^n\} dt \\ &= (n+3) \left( 1 - \frac{1}{2n+3} \right) \int_0^1 (1-t^2)^n dt = \frac{n+3}{n+2} \cdot \frac{2n+2}{2n+3} a_n. \end{aligned}$$

From the recurrence relation, we obtain that

$$\begin{aligned} a_{n+2} &= \frac{n+4}{n+3} \cdot \frac{2n+4}{2n+5} a_{n+1} = \frac{n+4}{n+2} \cdot \frac{(2n+2)(2n+4)}{(2n+3)(2n+5)} a_n, \\ a_{n+3} &= \frac{n+5}{n+4} \cdot \frac{2n+6}{2n+7} a_{n+2} = \frac{n+5}{n+2} \cdot \frac{(2n+2)(2n+4)(2n+6)}{(2n+3)(2n+5)(2n+7)} a_n. \end{aligned}$$

More generally, we see that  $a_{n+p} = \frac{n+p+2}{n+2} \cdot \frac{(2n+2)(2n+4)(2n+6) \times \cdots \times (2n+2p)}{(2n+3)(2n+5)(2n+7) \times \cdots \times (2n+2p+1)} a_n$  for any integer  $p \geq 0$ . Substituting  $n = 0$ , we get  $a_p = \frac{p+2}{2} \cdot \frac{2 \times 4 \times 6 \times \cdots \times 2p}{3 \times 5 \times 7 \times \cdots \times (2p+1)} a_0 = (p+2) \cdot \frac{2^p \times p!}{(2p+1)!} \cdot \frac{1}{2^p \times p!}$ .

Thus, the general term of the sequence  $a_n$  is obtained:  $a_n = (n+2) \cdot \frac{(2^n \times n!)^2}{(2n+1)!}$ .

Since  $a_{n+1} = \frac{n+3}{n+2} \cdot \frac{2n+2}{2n+3} a_n = \left(1 + \frac{n}{2n^2+7n+6}\right) a_n$  and  $a_2 = 4 \int_0^1 (1-t^2)^2 dt = \frac{32}{15} > 2$ , the inequality  $a_n > 2$  holds for every natural number  $n \geq 2$ .  $\square$

From the proof of Lemma 3.14, we see that the sequence  $(n+2) \int_0^1 (1-t^2)^n dt$  is increasing as  $n$  increases; hence the greatest Ricci lower bounds  $R(X^3(n, n))$  decreases. As specific examples, the approximate values of  $R(X^3(n, n))$  for small  $n$  are summarized as shown in the following table.

$n$	2	3	4	5	6	7
$R(X^3(n, n))$	$\frac{15}{16} = 0.9375$	$\frac{7}{8} = 0.875$	$\frac{105}{128} \approx 0.820$	$\frac{99}{128} \approx 0.773$	$\frac{3003}{4096} \approx 0.733$	$\frac{715}{1024} \approx 0.698$

TABLE 3. Approximate values of  $R(X^3(n, n))$ .

**Corollary 3.15.** *The greatest Ricci lower bound  $R(X^3(n, n))$  of  $X^3(n, n)$  converges to 0 as  $n$  increases, that is,  $\lim_{n \rightarrow \infty} R(X^3(n, n)) = 0$ . In addition, Tian's alpha-invariant  $\alpha(X^3(n, n))$  also converges to 0 as  $n$  increases.*

*Proof.* Since the formula  $R(X^3(n, n)) = \frac{2 \times (2n+1)!}{(n+2)(2^n \times n!)^2}$  in Proposition 3.13 is expressed as factorials, it is useful to use an approximation for them. Stirling's approximation gives bounds of  $n!$  valid for all positive integers  $n$ :  $\sqrt{2\pi n} \left(\frac{n}{e}\right)^n \leq n! \leq e\sqrt{n} \left(\frac{n}{e}\right)^n$ . Thus, we get an inequality

$$\begin{aligned} R(X^3(n, n)) &= \frac{2 \times (2n+1)!}{(n+2)(2^n \times n!)^2} \leq \frac{2 \times e\sqrt{2n+1} \left(\frac{2n+1}{e}\right)^{2n+1}}{(n+2) \left\{2^n \times \sqrt{2\pi n} \left(\frac{n}{e}\right)^n\right\}^2} \\ &= \frac{2 \times \sqrt{2n+1} (2n+1)^{2n+1}}{(n+2) \times 2^{2n} \times 2\pi n \times n^{2n}} = \frac{2\sqrt{2n+1}}{\pi(n+2)} \cdot \left(1 + \frac{1}{2n}\right)^{2n+1}. \end{aligned}$$

As  $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{2n}\right)^{2n+1} = e$  and  $\lim_{n \rightarrow \infty} \frac{\sqrt{2n+1}}{n+2} = 0$ , we conclude that  $\lim_{n \rightarrow \infty} R(X^3(n, n)) = 0$ .

On the other hand, Tian [Tia87] showed that a lower bound of  $R(X)$  in terms of the alpha-invariant  $\alpha(X)$  for  $X$  which does not admit a Kähler–Einstein metric:  $R(X) \geq \alpha(X) \cdot \frac{\dim X + 1}{\dim X}$ . Hence,  $\dim X^3(n, n) = \frac{n(n+3)}{2}$  implies that  $\lim_{n \rightarrow \infty} \alpha(X^3(n, n)) = 0$ .  $\square$

Now, let's move on to the next case for  $2 \leq k \leq n-1$ .

**Proposition 3.16.** *For  $n > k \geq 2$ , the greatest Ricci lower bound of  $X^3(n, k)$  is*

$$R(X^3(n, k)) = \frac{(2n - 2k + 2) \int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+2} (4n-3k+4-t)^{k-1} dt}.$$

*Proof.* By Proposition 3.12, the moment polytope  $\Delta_3$  of  $X^3(n, k)$  is parametrized as

$$\gamma_3(t) = (k+t)\varpi_{k-1} + (2n-2k+2-t)\varpi_k$$

for  $-k \leq t \leq 2n-2k+2$ . Let's count the number of roots in  $\Phi_{P^u}$  for each value of the coefficients of  $\alpha_{k-1}, \alpha_k$ :

- (i) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{k-1} = 1, c_k = 0\}$  is  $k-1$ ,
- (ii) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{k-1} = 0, c_k = 1\}$  is  $2(n-k)$ ,
- (iii) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{k-1} = 1, c_k = 1\}$  is  $2(k-1)(n-k)$ ,
- (iv) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{k-1} = 0, c_k = 2\}$  is 1,
- (v) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{k-1} = 1, c_k = 2\}$  is  $k-1$ , and
- (vi) the cardinality of the set  $\{\sum_{i=1}^n c_i \alpha_i \in \Phi^+ : c_{k-1} = 2, c_k = 2\}$  is  $\frac{k(k-1)}{2}$ .

Then, from  $\kappa(\alpha_{k-1}, \varpi_{k-1}) = 1$  and  $\kappa(\alpha_k, \varpi_k) = 1$  we have

$$\begin{aligned} P_{DH}(\gamma_3(t)) &= \prod_{\alpha \in \Phi_{P^u}} \kappa(\alpha, (k+t)\varpi_{k-1} + (2n-2k+2-t)\varpi_k) \\ &= (k+t)^{k-1} (2n-2k+2-t)^{2(n-k)} \{(k+t) + (2n-2k+2-t)\}^{2(k-1)(n-k)} \\ &\quad \times \{2(2n-2k+2-t)\} \{(k+t) + 2(2n-2k+2-t)\}^{k-1} \{2(k+t) + 2(2n-2k+2-t)\}^{\frac{k(k-1)}{2}} \\ &= 2(2n-k+2)^{2(k-1)(n-k)} \{2(2n-k+2)\}^{\frac{k(k-1)}{2}} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1}. \end{aligned}$$

$$\text{As } \frac{1}{\text{Vol}_{DH}(\Delta_3)} \int_{-k}^{2n-2k+2} (k+t) \cdot P_{DH}(\gamma_3(t)) dt = \frac{\int_{-k}^{2n-2k+2} (k+t)^k (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt},$$

$$\text{we get the barycenter } \mathbf{bar}_{DH}(\Delta_3) = \gamma_3 \left( \frac{\int_{-k}^{2n-2k+2} (k+t)^k (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt} - k \right)$$

of the moment polytope  $\Delta_3$  with respect to the Duistermaat–Heckman measure.

Since  $\frac{\int_{-k}^{2n-2k+2} (k+t)^k (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt} - k < 0$  by Lemma 3.17, the greatest Ricci lower bound of  $X^3(n, k)$  is

$$\begin{aligned} R(X^3(n, k)) &= \frac{2n-2k+2}{(2n-2k+2) - \left( \frac{\int_{-k}^{2n-2k+2} (k+t)^k (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt} - k \right)} \\ &= \frac{2n-2k+2}{(2n-k+2) - \frac{\int_{-k}^{2n-2k+2} (k+t)^k (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}} \\ &= \frac{(2n-2k+2) \int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+2} (4n-3k+4-t)^{k-1} dt} \end{aligned}$$

by Corollary 2.9. □



**Lemma 3.17.** *For any  $n > k \geq 2$ , the inequality*

$$\frac{\int_{-k}^{2n-2k+2} (k+t)^k (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt} < k$$

holds.

*Proof.* It suffices to show that  $\int_{-k}^{2n-2k+2} t(k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt < 0$  for  $n > k \geq 2$ . Putting the left side of this inequality as  $I(n, k)$ , we have

$$\begin{aligned} I(n, k) &= \int_{-k}^{2n-2k+2} t(2n-2k+2-t)^{2n-2k+1} \{(k+t)(4n-3k+4-t)\}^{k-1} dt \\ &= \int_{-k}^{2n-2k+2} t(2n-2k+2-t)^{2n-2k} \cdot (2n-2k+2-t) \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^{k-1} dt. \end{aligned}$$

Since  $\int (2n-2k+2-t) \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^{k-1} dt = \frac{1}{2k} \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^k + C$ , we can use integration by parts repetitively:

$$\begin{aligned} I(n, k) &= \left[ t(2n-2k+2-t)^{2n-2k} \cdot \frac{1}{2k} \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^k \right]_{-k}^{2n-2k+2} \\ &\quad - \int_{-k}^{2n-2k+2} (2n-2k+2-t)^{2n-2k-1} \{(2n-2k+2) - (2n-2k+1)t\} \\ &\quad \times \frac{1}{2k} \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^k dt \\ &= \frac{1}{2k} \int_{-k}^{2n-2k+2} (2n-2k+2-t)^{2n-2k-2} \{(2n-2k+1)t - (2n-2k+2)\} \\ &\quad \times (2n-2k+2-t) \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^k dt \\ &= \frac{1}{2k} \left[ \{(2n-2k+1)t - (2n-2k+2)\} (2n-2k+2-t)^{2n-2k-2} \right. \\ &\quad \times \left. \frac{1}{2(k+1)} \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^{k+1} \right]_{-k}^{2n-2k+2} - \frac{1}{2k} \int_{-k}^{2n-2k+2} (2n-2k+2-t)^{2n-2k-3} \\ &\quad \times [-(2n-2k+1)(2n-2k-1)t + (2n-2k+2)\{1 + (2n-2k) + (2n-2k-2)\}] \\ &\quad \times \frac{1}{2(k+1)} \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^{k+1} dt. \end{aligned}$$

If  $n-k=2$ , then we have

$$\begin{aligned} I(n, k) &= \frac{1}{2k} \cdot \frac{1}{2(k+1)} \int_{-k}^{2n-2k+2} [(2n-2k+1)(2n-2k-1)t - (2n-2k+2)\{1 + (2n-2k) + (2n-2k-2)\}] \\ &\quad \times (2n-2k+2-t) \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^{k+1} dt \\ &= \frac{1}{2k} \cdot \frac{1}{2(k+1)} \cdot \frac{1}{2(k+2)} \left[ (2n-2k+2)(2n-2k)(2n-2k-2)(2n-k+2)^{2(k+2)} \right. \\ &\quad \left. - (2n-2k+1)(2n-2k-1) \int_{-k}^{2n-2k+2} \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^{k+2} dt \right]. \end{aligned}$$

As a result, we obtain that

$$\begin{aligned} I(n, k) &= \frac{1}{2k} \cdot \frac{1}{2(k+1)} \times \cdots \times \frac{1}{2n} \left[ (2n-2k+2)(2n-2k)(2n-2k-2) \times \cdots \times 4 \times 2 \times (2n-k+2)^{2n} \right. \\ &\quad \left. - (2n-2k+1)(2n-2k-1)(2n-2k-3) \times \cdots \times 3 \times \int_{-k}^{2n-2k+2} \{-(2n-2k+2-t)^2 + (2n-k+2)^2\}^n dt \right]. \end{aligned}$$

For a fixed  $\ell := n - k$ , if we show the following inequality, then we get the conclusion we want:

$$(n + \ell + 2) \int_0^1 (1 - s^2)^n ds > \frac{(2\ell + 2)(2\ell)(2\ell - 2) \times \cdots \times 4 \times 2}{(2\ell + 1)(2\ell - 1)(2\ell - 3) \times \cdots \times 3} \quad \text{for } n \geq \ell + 2.$$

We give a proof by induction on  $n$ .

(i) First, the statement holds for  $n = \ell + 2$  because  $(2\ell + 4) \int_0^1 (1 - s^2)^{\ell + 2} ds > \frac{(2\ell + 2)(2\ell)(2\ell - 2) \times \cdots \times 4 \times 2}{(2\ell + 1)(2\ell - 1)(2\ell - 3) \times \cdots \times 3}$

for any  $\ell \geq 1$ . Indeed, we can show that the ratio  $f(\ell) := \frac{(2\ell + 4) \int_0^1 (1 - s^2)^{\ell + 2} ds}{(2\ell + 2)(2\ell)(2\ell - 2) \times \cdots \times 4 \times 2}$  is greater than  $\frac{(2\ell + 4) \int_0^1 (1 - s^2)^{\ell + 2} ds}{(2\ell + 1)(2\ell - 1)(2\ell - 3) \times \cdots \times 3}$

one. Since  $\int_0^1 (1 - s^2)^{\ell + 3} ds = \frac{2\ell + 6}{2\ell + 7} \int_0^1 (1 - s^2)^{\ell + 2} ds$  from the proof of Lemma 3.14, we can express each element of the sequence  $f(\ell)$  as a function of the preceding ones:

$$\begin{aligned} f(\ell + 1) &= \frac{(2\ell + 6) \int_0^1 (1 - s^2)^{\ell + 3} ds}{(2\ell + 4)(2\ell + 2)(2\ell) \times \cdots \times 4 \times 2} = \frac{(2\ell + 6) \cdot \frac{2\ell + 6}{2\ell + 7} \int_0^1 (1 - s^2)^{\ell + 2} ds}{\frac{2\ell + 4}{2\ell + 3} \cdot (2\ell + 2)(2\ell) \times \cdots \times 4 \times 2} \\ &= \frac{(\ell + 3)^2(2\ell + 3)}{(\ell + 2)^2(2\ell + 7)} f(\ell). \end{aligned}$$

From the recurrence relation, we obtain that

$$\begin{aligned} f(\ell + 2) &= \frac{(\ell + 4)^2(2\ell + 5)}{(\ell + 3)^2(2\ell + 9)} f(\ell + 1) = \frac{(\ell + 4)^2(2\ell + 5)}{(\ell + 3)^2(2\ell + 9)} \cdot \frac{(\ell + 3)^2(2\ell + 3)}{(\ell + 2)^2(2\ell + 7)} f(\ell) \\ &= \frac{(\ell + 4)^2(2\ell + 3)(2\ell + 5)}{(\ell + 2)^2(2\ell + 7)(2\ell + 9)} f(\ell), \\ f(\ell + 3) &= \frac{(\ell + 5)^2(2\ell + 7)}{(\ell + 4)^2(2\ell + 11)} f(\ell + 2) = \frac{(\ell + 5)^2(2\ell + 7)}{(\ell + 4)^2(2\ell + 11)} \cdot \frac{(\ell + 4)^2(2\ell + 3)(2\ell + 5)}{(\ell + 2)^2(2\ell + 7)(2\ell + 9)} f(\ell) \\ &= \frac{(\ell + 5)^2(2\ell + 3)(2\ell + 5)}{(\ell + 2)^2(2\ell + 9)(2\ell + 11)} f(\ell). \end{aligned}$$

More generally, we see that  $f(\ell + p) = \frac{(\ell + p + 2)^2(2\ell + 3)(2\ell + 5)}{(\ell + 2)^2(2\ell + 2p + 3)(2\ell + 2p + 5)} f(\ell)$  for any natural number  $p$ . Thus,

$$f(p + 1) = \frac{(p + 3)^2 \times 5 \times 7}{3^2(2p + 5)(2p + 7)} f(1) = \frac{35(p + 3)^2}{9(2p + 5)(2p + 7)} \times \frac{36}{35} = \frac{4(p + 3)^2}{(2p + 5)(2p + 7)} = 1 + \frac{1}{4p^2 + 24p + 35} > 1.$$

(ii) Assume the induction hypothesis that  $(m + \ell + 2) \int_0^1 (1 - s^2)^m ds > \frac{(2\ell + 2)(2\ell)(2\ell - 2) \times \cdots \times 4 \times 2}{(2\ell + 1)(2\ell - 1)(2\ell - 3) \times \cdots \times 3}$

for a particular  $m \geq \ell + 2$ . Since we know that  $\int_0^1 (1 - s^2)^m ds = (2m + 3) \int_0^1 s^2(1 - s^2)^m ds$  from the proof of Lemma 3.14,

$$\begin{aligned} ((m + 1) + \ell + 2) \int_0^1 (1 - s^2)^{m+1} ds &= (m + \ell + 3) \cdot \frac{2m + 2}{2m + 3} \int_0^1 (1 - s^2)^m ds \\ &= \frac{m + \ell + 3}{m + \ell + 2} \cdot \frac{2m + 2}{2m + 3} \cdot (m + \ell + 2) \int_0^1 (1 - s^2)^m ds \\ &= \left\{ 1 + \frac{m - \ell}{2m^2 + (2\ell + 7)m + 3(\ell + 2)} \right\} \cdot (m + \ell + 2) \int_0^1 (1 - s^2)^m ds. \end{aligned}$$

Therefore, if  $n - \ell \geq 2$  then we have

$$\begin{aligned} ((m+1) + \ell + 2) \int_0^1 (1-s^2)^{m+1} ds &> (m + \ell + 2) \int_0^1 (1-s^2)^m ds \\ &> \frac{(2\ell+2)(2\ell)(2\ell-2) \times \cdots \times 4 \times 2}{(2\ell+1)(2\ell-1)(2\ell-3) \times \cdots \times 3}. \end{aligned}$$

Since both the base case and the inductive step have been proved as true, by mathematical induction the inequality holds for every natural number  $n \geq \ell + 2$ .  $\square$

**Remark 3.18.** The formula for the greatest Ricci lower bound of  $X^3(n, n)$  in Proposition 3.13 is consistent with the formula in Proposition 3.16 when  $k = n$  is substituted in it.

Interestingly, we observe that for a fixed integer  $k$  the value  $R(X^3(n, k))$  always approaches to 1 asymptotically as  $n$  increases.

**Corollary 3.19.** *For a fixed integer  $k \geq 2$ , the greatest Ricci lower bound  $R(X^3(n, k))$  of  $X^3(n, k)$  converges to 1 as  $n$  increases, that is,  $\lim_{n \rightarrow \infty} R(X^3(n, k)) = 1$ .*

*Proof.* Recall that  $R(X^3(n, k)) = \frac{2n - 2k + 2}{2n - k + 2 - \frac{\int_{-k}^{2n-2k+2} (k+t)^k (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}{\int_{-k}^{2n-2k+2} (k+t)^{k-1} (2n-2k+2-t)^{2n-2k+1} (4n-3k+4-t)^{k-1} dt}}$  from Proposition 3.16. Since all functions  $k+t$ ,  $2n-2k+2-t$  and  $4n-3k+4-t$  in the integrands are positive for  $-k < t < 2n-2k+2$ , we get an inequality  $R(X^3(n, k)) > \frac{2n-2k+2}{2n-k+2}$ ; hence for a fixed  $k$  we have  $\lim_{n \rightarrow \infty} R(X^3(n, k)) \geq \lim_{n \rightarrow \infty} \frac{2n-2k+2}{2n-k+2} = 1$ . Therefore, we conclude that  $\lim_{n \rightarrow \infty} R(X^3(n, k)) = 1$ .  $\square$

## REFERENCES

- [Aub78] Thierry Aubin, *Équations du type Monge–Ampère sur les variétés kählériennes compactes*, Bull. Sci. Math.(2) **102** (1978), 63–95.
- [BBJ21] Robert Berman, Sébastien Boucksom, and Mattias Jonsson, *A variational approach to the Yau–Tian–Donaldson conjecture*, J. Amer. Math. Soc. (2021), in press. <https://doi.org/10.1090/jams/964>
- [Bri87] Michel Brion, *Sur l'image de l'application moment*, Séminaire d'algèbre Paul Dubreil et Marie-Paule Malliavin (Paris, 1986), Lecture Notes in Math., vol. 1296, Springer, Berlin, 1987, pp. 177–192.
- [Bri89] ———, *Groupe de Picard et nombres caractéristiques des variétés sphériques*, Duke Math. J. **58** (1989), no. 2, 397–424.
- [Bri97] ———, *Curves and divisors in spherical varieties*, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., vol. 9, Cambridge Univ. Press, Cambridge, 1997, pp. 21–34.
- [Cab19] Jacob Cable, *Greatest lower bounds on Ricci curvature for Fano T-manifolds of complexity one*, Bull. Lond. Math. Soc. **51** (2019), no. 1, 34–42.
- [CDS15a] Xiuxiong Chen, Simon Donaldson, and Song Sun, *Kähler–Einstein metrics on Fano manifolds. I: Approximation of metrics with cone singularities*, J. Amer. Math. Soc. **28** (2015), no. 1, 183–197.
- [CDS15b] ———, *Kähler–Einstein metrics on Fano manifolds. II: Limits with cone angle less than  $2\pi$* , J. Amer. Math. Soc. **28** (2015), no. 1, 199–234.
- [CDS15c] ———, *Kähler–Einstein metrics on Fano manifolds. III: Limits as cone angle approaches  $2\pi$  and completion of the main proof*, J. Amer. Math. Soc. **28** (2015), no. 1, 235–278.
- [CRZ19] Ivan Cheltsov, Yanir Rubinstein, and Kewei Zhang, *Basis log canonical thresholds, local intersection estimates, and asymptotically log del Pezzo surfaces*, Selecta Math. **25** (2019), no. 2, Article number 34.
- [Del17] Thibaut Delcroix, *Kähler–Einstein metrics on group compactifications*, Geom. Funct. Anal. **27** (2017), no. 1, 78–129.
- [Del20a] ———, *K-stability of Fano spherical varieties*, Ann. Sci. Éc. Norm. Supér. (4) **53** (2020), no. 3, 615–662.
- [Del20b] ———, *Kähler geometry of horosymmetric varieties, and application to Mabuchi's K-energy functional*, J. Reine Angew. Math. **763** (2020) 129–199.
- [DH20] Thibaut Delcroix and Jakob Hultgren, *Coupled complex Monge–Ampère equations on Fano horosymmetric manifolds*, J. Math. Pures Appl. (2020), in press. <https://doi.org/10.1016/j.matpur.2020.12.002>
- [FO18] Kento Fujita and Yuji Odaka, *On the K-stability of Fano varieties and anticanonical divisors*, Tohoku Math. J. **70** (2018), no. 4, 511–521.
- [Gan18] Jacopo Gandini, *Embeddings of spherical homogeneous spaces*, Acta Math. Sin. (Engl. Ser.) **34** (2018), no. 3, 299–340.
- [Gol20] Aleksei Golota, *Delta-invariants for Fano varieties with large automorphism groups*, Internat. J. Math. **31** (2020), no. 10, 2050077.
- [GPPS21] Richard Gonzales, Clélia Pech, Nicolas Perrin, and Alexander Samokhin, *Geometry of horospherical varieties of Picard rank one*, Int. Math. Res. Not (2021), in press. <https://doi.org/10.1093/imrn/rnaa331>

- [Hon15] Jaehyun Hong, *Classification of smooth Schubert varieties in the symplectic Grassmannians*, J. Korean Math. Soc. **52** (2015), no. 5, 1109–1122.
- [Hon16] ———, *Smooth horospherical varieties of Picard number one as linear sections of rational homogeneous varieties*, J. Korean Math. Soc. **53** (2016), no. 5, 433–446.
- [Hum72] James E. Humphreys, *Introduction to Lie algebras and representation theory*, Graduate Texts in Mathematics 9, Springer-Verlag, New York, 1972.
- [HL19] Jun-Muk Hwang and Qifeng Li, *Characterizing symplectic Grassmannians by varieties of minimal rational tangents*, preprint (2019), arXiv:1901.00357v1.
- [HM05] Jun-Muk Hwang and Ngaiming Mok, *Prolongations of infinitesimal linear automorphisms of projective varieties and rigidity of rational homogeneous spaces of Picard number 1 under Kähler deformation*, Invent. Math. **160** (2005), no. 3, 591–645.
- [Kan21] Akihiro Kanemitsu, *Fano manifolds and stability of tangent bundles*, J. reine angew. Math. **774** (2021), 163–183.
- [Kim17] Shin-Young Kim, *Geometric structures modeled on smooth projective horospherical varieties of Picard number one*, Transform. Groups **22** (2017), no. 2, 361–386.
- [Kno91] Friedrich Knop, *The Luna–Vust theory of spherical embeddings*, Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989), Manoj Prakashan, Madras, 1991, pp. 225–249.
- [KP18] Shin-Young Kim and Kyeong-Dong Park, *Standard embeddings of smooth Schubert varieties in rational homogeneous manifolds of Picard number 1*, Acta. Math. Sin. (Engl. Ser.) **34** (2018), no. 3, 466–487.
- [LPY21] Jae-Hyoun Lee, Kyeong-Dong Park, and Sungmin Yoo, *Kähler–Einstein metrics on smooth Fano symmetric varieties with Picard number one*, Mathematics **9** (2021), no. 1, 102. <https://doi.org/10.3390/math9010102>
- [Li11] Chi Li, *Greatest lower bounds on Ricci curvature for toric Fano manifolds*, Adv. Math. **226** (2011), no. 6, 4921–4932.
- [Lun97] Domingo Luna, *Grosses cellules pour les variétés sphériques*, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., vol. 9, Cambridge Univ. Press, Cambridge, 1997, pp. 267–280.
- [LV83] Domingo Luna and Thierry Vust, *Plongements d’espaces homogènes*, Comment. Math. Helv. **58** (1983), no. 2, 186–245.
- [LXZ21] Yuchen Liu, Chenyang Xu, and Ziquan Zhuang, *Finite generation for valuations computing stability thresholds and applications to K-stability*, preprint (2021), arXiv:2102.09405v2.
- [Mat57] Yozo Matsushima, *Sur la structure du groupe d’homéomorphismes analytiques d’une certaine variété kählérienne*, Nagoya Math. J. **11** (1957), 145–150.
- [Mat72] ———, *Remarks on Kähler–Einstein manifolds*, Nagoya Math. J. **46** (1972), 161–173.
- [Mih07] Ion Alexandru Mihai, *Odd symplectic flag manifolds*, Transform. Groups **12** (2007), no. 3, 573–599.
- [MP20] Boris Pasquier and Laurent Manivel, *Horospherical two-orbit varieties as zero loci*, preprint (2020), arXiv:2012.05513v1.
- [Muk89] Shigeru Mukai, *Biregular classification of Fano 3-folds and Fano manifolds of coindex 3*, Proc. Nat. Acad. Sci. U.S.A. **86** (1989), no. 9, 3000–3002.
- [Par16] Kyeong-Dong Park, *Deformation rigidity of odd Lagrangian Grassmannians*, J. Korean Math. Soc. **53** (2016), no. 3, 489–501.
- [Pas08] Boris Pasquier, *Variétés horosphériques de Fano*, Bull. Soc. Math. France **136** (2008), no. 2, 195–225.
- [Pas09] Boris Pasquier, *On some smooth projective two-orbit varieties of Picard number 1*, Math. Ann. **344** (2009), no. 4, 963–987.
- [Per14] Nicolas Perrin, *On the geometry of spherical varieties*, Transform. Groups **19** (2014), no. 1, 171–223.
- [PP10] Boris Pasquier and Nicolas Perrin, *Local rigidity of quasi-regular varieties*, Math. Z. **265** (2010), no. 3, 589–600.
- [Rub08] Yanir A. Rubinstein, *Some discretizations of geometric evolution equations and the Ricci iteration on the space of Kähler metrics*, Adv. Math. **218** (2008), no. 5, 1526–1565.
- [Rub09] ———, *On the construction of Nadel multiplier ideal sheaves and the limiting behavior of the Ricci flow*, Trans. Amer. Math. Soc. **361** (2009), no. 11, 5839–5850.
- [SW16] Jian Song and Xiaowei Wang, *The greatest Ricci lower bound, conical Einstein metrics and Chern number inequality*, Geom. Topol. **20** (2016), no. 1, 49–102.
- [Szé11] Gábor Székelyhidi, *Greatest lower bounds on the Ricci curvature of Fano manifolds*, Compos. Math. **147**, no. 1, 319–331.
- [Tia87] Gang Tian, *On Kähler–Einstein metrics on certain Kähler manifolds with  $c_1(M) > 0$* , Invent. Math. **89** (1987), no. 2, 225–246.
- [Tia92] ———, *On stability of the tangent bundles of Fano varieties*, Internat. J. Math. **3** (1992), no. 3, 401–413.
- [Tia15] ———, *K-stability and Kähler–Einstein metrics*, Commun. Pure Appl. Math. **68** (2015), no. 7, 1085–1156. [Corrigendum: *K-stability and Kähler–Einstein metrics*, **68** (2015), no. 11, 2082–2083.]
- [Tim11] Dmitry A. Timashev, *Homogeneous spaces and equivariant embeddings*, Encyclopaedia of Mathematical Sciences, Vol. 138, Subseries *Invariant Theory and Algebraic Transformation Groups*, Vol. 8, Springer, Heidelberg, 2011.
- [Yao17] Yi Yao, *Greatest lower bounds on Ricci curvature of homogeneous toric bundles*, Internat. J. Math. **28** (2017), no. 4, 1750024.
- [Yau78] Shing-Tung Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equation, I*, Comm. Pure Appl. Math. **31** (1978), no. 3, 339–411.

DEPARTMENT OF MATHEMATICS, AJOU UNIVERSITY, SUWON 16499, REPUBLIC OF KOREA

*Email address:* [dshwang@ajou.ac.kr](mailto:dshwang@ajou.ac.kr)

CENTER FOR GEOMETRY AND PHYSICS, INSTITUTE FOR BASIC SCIENCE (IBS), POHANG 37673, REPUBLIC OF KOREA

*Email address:* [shinyoungkim@ibs.re.kr](mailto:shinyoungkim@ibs.re.kr)

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY (KIAS), DONGDAEMUN-GU, SEOUL 02455, REPUBLIC OF KOREA

*Email address:* [kdpark@kias.re.kr](mailto:kdpark@kias.re.kr)