

# Gravitational wave echoes from interacting quark stars

Chen Zhang<sup>1,\*</sup>

<sup>1</sup>*Department of Physics, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1*

We show that interacting quark stars (IQSs) composed of interacting quark matter (IQM), including the strong interaction effects such as perturbative QCD corrections and color superconductivity, can be compact enough to feature a photon sphere that is essential to the signature of gravitational wave echoes. We utilize an IQM equation of state unifying all interacting phases by a simple reparametrization and rescaling, through which we manage to maximally reduce the number of degrees of freedom into one dimensionless parameter  $\bar{\lambda}$  that characterizes the relative size of strong interaction effects. It turns out that gravitational wave echoes are possible for IQSs with  $\bar{\lambda} \gtrsim 10$  at large center pressure. Rescaling the dimension back, we illustrate its implication on the dimensional parameter space of effective bag constant  $B_{\text{eff}}$  and the superconducting gap  $\Delta$  with variations of the perturbative QCD parameter  $a_4$  and the strange quark mass  $m_s$ . We calculate the rescaled GW echo frequencies  $\bar{f}_{\text{echo}}$  associated with IQSs, from which we obtain a simple scaling relation for the minimal echo frequency  $f_{\text{echo}}^{\text{min}} \approx 5.76 \sqrt{B_{\text{eff}}/(100 \text{ MeV})^4}$  kHz at the large  $\bar{\lambda}$  limit.

## INTRODUCTION

The recent observations of gravitational wave (GW) signals from compact binary mergers by the LIGO-Virgo collaborations [1–7] have greatly moved our understanding of black holes and compact stars forward. The detected binary black hole merger events inspired many studies on black hole mimickers termed exotic compact objects (ECOs), whose defining feature is their large compactness: their radius is very close to that of a black hole with the same mass while lacking an event horizon. While some non-GW probes of ECOs have been studied [8], most studies are on the distinctive signatures from gravitational wave echoes in the postmerger signals [9–25], in which a wave that falls inside the gravitational potential barrier travels to a reflecting boundary before returning to the barrier at the photon sphere after some time delay.

Considering the detected binary neutron star merger events, we want to explore the possibility of GW echoes also being signature of realistic compact stars. Generating GW echoes requires the star object to feature a photon sphere at  $R_P = 3M$ , where  $M$  is the object’s mass. For compact stars, the minimum radius should be above the Buchdahl’s limit  $R_B = 9/4M$  [26]. Therefore, GW echo signals are possible if  $R_B < R < R_P$ . This compactness criterion excludes the realistic neutron stars [27, 28]. This motivates the exploration of other more compact star objects such as quark stars composed

of quark matter.

Bodmer [29], Witten [30] and Terazawa [31] proposed that quark matter with comparable numbers of  $u$ ,  $d$ ,  $s$  quarks, also called strange quark matter (SQM), might be the ground state of baryonic matter at the zero temperature and pressure. A recent study [32] demonstrated that  $u, d$  quark matter ( $ud$ QM) is, in general, more stable than SQM, and it can be more stable than the ordinary nuclear matter at a sufficiently large baryon number beyond the periodic table. The SQM hypothesis and  $ud$ QM hypothesis, as mentioned above, allow the possibility of bare quark stars, such as strange quark stars (SQSs) [33, 34] that consist of SQM or up-down quark stars ( $ud$ QSs) [35, 36] that consist of  $ud$ QM. In the context of recent LIGO events, there are a lot of studies on the related astrophysical implications of SQSs [37–42] and  $ud$ QSs [35, 43–46], many of which involve interacting quark matter (IQM) that includes the interquark effects such as the perturbative QCD (pQCD) corrections and the color superconductivity. pQCD corrections are due to the gluon-mediated interaction [47–49]. Color superconductivity is the superconductivity in quark matter, arising from the spin-0 Cooper-pair condensation anti-symmetric in color-flavor space [50–52]. This can result in two-flavor color superconductivity, where  $u$  quarks pair with  $d$  quarks [conventionally termed “2SC” (“2SC+s”) without (with) strange quarks], or in a color-flavor locking (CFL) phase, where  $u, d, s$  quarks pair with each

other antisymmetrically.

In order to achieve a large compactness for stars to generate GW echoes, people commonly assumed ad-hoc exotic equations of state (EOS) [28, 56, 57] or special semi-classical treatment of gravity [58]. Here we demonstrate that the physically motivated interacting quark stars (IQSs) composed of IQM can have GW echo signatures within classical Einstein gravity framework. It has been shown [45] that IQSs can meet the various constraints from observed large pulsar masses [59–61], analysis of the NICER X-ray spectral-timing event data [62, 63], and the recent LIGO events [4–7].

Referring to our previous paper [45], we first rewrite the free energy  $\Omega$  of the superconducting quark matter [53] into a general form with the pQCD correction included:

$$\Omega = -\frac{\xi_4}{4\pi^2}\mu^4 + \frac{\xi_4(1-a_4)}{4\pi^2}\mu^4 - \frac{\xi_{2a}\Delta^2 - \xi_{2b}m_s^2}{\pi^2}\mu^2 - \frac{\mu_e^4}{12\pi^2} + B_{\text{eff}}, \quad (1)$$

where  $\mu$  and  $\mu_e$  are the respective average quark and electron chemical potentials. The first term represents the unpaired free quark gas contribution. The second term with  $(1-a_4)$  represents the pQCD contribution from one-gluon exchange for gluon interaction to  $O(\alpha_s^2)$  order. To phenomenologically account for higher-order contributions, we can vary  $a_4$  from  $a_4 = 1$ , corresponding to a vanishing pQCD correction, to very small values where these corrections become large [48, 54, 55]. The term with  $m_s$  accounts for the correction from the finite strange quark mass if applicable, while the term with the gap parameter  $\Delta$  represents the contribution from color superconductivity.

$$(\xi_4, \xi_{2a}, \xi_{2b}) = \begin{cases} ((\frac{1}{3})^{\frac{4}{3}} + (\frac{2}{3})^{\frac{4}{3}})^{-3}, 1, 0 & \text{2SC phase} \\ (3, 1, 3/4) & \text{2SC+s phase} \\ (3, 3, 3/4) & \text{CFL phase} \end{cases}$$

The corresponding equation of state is [45]:

$$p = \frac{1}{3}(\rho - 4B_{\text{eff}}) + \frac{4\lambda^2}{9\pi^2} \left( -1 + \text{sgn}(\lambda) \sqrt{1 + 3\pi^2 \frac{(\rho - B_{\text{eff}})}{\lambda^2}} \right), \quad (2)$$

where

$$\lambda = \frac{\xi_{2a}\Delta^2 - \xi_{2b}m_s^2}{\sqrt{\xi_4 a_4}}. \quad (3)$$

Note that  $\text{sgn}(\lambda)$  represents the sign of  $\lambda$ . One can easily see that a larger  $\lambda$  (i.e., smaller  $B_{\text{eff}}$ ,  $a_4$ ,  $m_s$  or larger  $\Delta$ ) leads to a stiffer EOS which results in a more compact stellar structure that is more likely to have GW echoes. Thus, for this study, we only need to explore positive  $\lambda$  space.

As shown in Ref. [45], one can further remove the  $B_{\text{eff}}$  parameter by doing the dimensionless rescaling:

$$\bar{\rho} = \frac{\rho}{4B_{\text{eff}}}, \quad \bar{p} = \frac{p}{4B_{\text{eff}}}, \quad (4)$$

and

$$\bar{\lambda} = \frac{\lambda^2}{4B_{\text{eff}}} = \frac{(\xi_{2a}\Delta^2 - \xi_{2b}m_s^2)^2}{4B_{\text{eff}}\xi_4 a_4}, \quad (5)$$

so that the EOS Eq. (2) reduces to the dimensionless form

$$\bar{p} = \frac{1}{3}(\bar{\rho} - 1) + \frac{4}{9\pi^2} \bar{\lambda} \left( -1 + \sqrt{1 + \frac{3\pi^2}{\bar{\lambda}} (\bar{\rho} - \frac{1}{4})} \right). \quad (6)$$

As  $\bar{\lambda} \rightarrow 0$ , Eq. (6) reduces to the conventional non-interacting rescaled quark matter EOS  $\bar{p} = (\bar{\rho} - 1)/3$ . When  $\bar{\lambda}$  goes extremely large, Eq. (6) approaches the special form

$$\bar{p}|_{\bar{\lambda} \rightarrow \infty} = \bar{\rho} - \frac{1}{2}, \quad (7)$$

or equivalently  $p = \rho - 2B_{\text{eff}}$  using Eq. (4). We see that strong interaction effects can reduce the surface mass density of a quark star from  $\rho_0 = 4B_{\text{eff}}$  down to  $\rho_0 = 2B_{\text{eff}}$ , and increase the quark matter sound speed  $c_s^2 = \partial p / \partial \rho$  from 1/3 up to 1 (the light speed) maximally.

## GW ECHOES FROM IQS

To study the stellar structure of IQSs, we first rescale the mass and radius into dimensionless form in geometric units ( $G = c = 1$ )<sup>1</sup>

$$\bar{m} = m\sqrt{4B_{\text{eff}}}, \quad \bar{r} = r\sqrt{4B_{\text{eff}}}, \quad (8)$$

so that the Tolman-Oppenheimer-Volkov (TOV) equation [64, 65]

$$\begin{aligned} \frac{dm}{dr} &= 4\pi\rho r^2, \\ \frac{dp}{dr} &= (\rho + p) \frac{m + 4\pi p r^3}{2mr - r^2}, \end{aligned} \quad (9)$$

<sup>1</sup>Note that  $B_{\text{eff}}$ , which is in units of  $\text{MeV}^4$  or  $\text{MeV}/\text{fm}^3$  in natural units, is in dimension of  $[L^{-2}]$  in geometric units here.

can be converted into the dimensionless form (simply replace nonbarred symbols with barred ones). Solving the dimensionless TOV equation, we obtain the results for the rescaled  $\bar{M} - \bar{R}$  shown in Fig. 1. Note that beyond the maximum mass point (red dot), the object begins to be unstable against radial perturbations. It turns out that all  $\bar{M} - \bar{R}$  configurations meet the Buchdahl's limit, and those with  $\bar{\lambda} \gtrsim 10$  can cross the photon sphere line, satisfying the necessary condition to generate GW echoes. Interestingly, referring to Fig 3 of Ref. [45], this  $\bar{\lambda} \gtrsim 10$  range well saturates the joint constraints set by the GW170817 and GW190814 analyses assuming related objects are IQSs.




Figure 1.  $\bar{M}-\bar{R}$  of IQSs for given  $\bar{\lambda}$ , sampling (0, 0.1, 5, 10, 20, 50, 100) from the lighter black line to darker black line respectively. The red line corresponds to  $\bar{\lambda} \rightarrow \infty$ , with the corresponding EOS Eq. (7). The solid dots denote the maximum mass configurations for given  $\bar{\lambda}$ . GW echoes require the star to have  $\bar{M}-\bar{R}$  configurations above the photon sphere line.

From Eq. (5), we see that the echo criterion  $\bar{\lambda} \gtrsim 10$  maps to the constraint on dimensional parameters

$$(\xi_{2a}\Delta^2 - \xi_{2b}m_s^2)^2 \gtrsim 40\xi_4 a_4 B_{\text{eff}}, \quad (10)$$

which can be satisfied for a large strong interaction effect (i.e., large  $\Delta$  or small  $a_4$ ) or a small effective bag constant. Considering  $m_s$  has been constrained in a result of  $95 \pm 5$  MeV [66], we fix  $m_s = (90, 100)$  MeV and obtain Fig. (2) for CFL phase from Eq. (10) for illustration. We can see that the strange quark mass variation 90 – 100 MeV has a negligible effect on the saturation of Eq. (10), while a larger pQCD correction (smaller  $a_4$ ) results in a smaller  $\Delta$  for a given bag constant to meet Eq. (10).




Figure 2.  $B^{1/4}-\Delta$  of IQSs in CFL phase that gives  $\bar{\lambda} = 10$  for  $m_s = 90$  MeV (red solid), and  $m_s = 100$  MeV (black dashed), with  $a_4 = (1, 0.75, 0.5, 0.25)$  from the lighter black line to darker black line respectively. The region below each line represents the corresponding parameter space that has IQSs compact enough to generate GW echoes.

The characteristic echo time is the light time from the star center to the photon sphere [10–12],

$$\tau_{\text{echo}} = \int_0^{3M} \frac{dr}{\sqrt{e^{2\Phi(r)} \left(1 - \frac{2m(r)}{r}\right)}}, \quad (11)$$

where

$$\frac{d\Phi}{dr} = -\frac{1}{\rho + p} \frac{dp}{dr}. \quad (12)$$

We can also do the dimensionless rescaling

$$\bar{\tau}_{\text{echo}} = \tau_{\text{echo}} \sqrt{4 B_{\text{eff}}}, \quad (13)$$

so that Eq.(12) can also be solved in a dimensionless approach. After obtaining the echo time, we directly obtain the rescaled GW echo frequency from the relation [10–12]

$$f_{\text{echo}} = \frac{\pi}{\tau_{\text{echo}}}, \quad (14)$$

and similarly, we rescale it into dimensionless form  $\bar{f}_{\text{echo}}$  via relation

$$\bar{f}_{\text{echo}} = \frac{f_{\text{echo}}}{\sqrt{4 B_{\text{eff}}}}. \quad (15)$$

In Fig. 3, we show the results of rescaled GW echo frequencies  $\bar{f}_{\text{echo}}$  versus the rescaled center pressure  $\bar{p}_c$  for the stellar configurations of Fig. 1 that can generate echoes (i.e.,  $\bar{\lambda} \gtrsim 10$ ). Note that each curve's left and right ends are truncated at the point where  $\bar{R} = 3\bar{M}$ , and at the point of maximum mass, respectively. The gray

dot at  $(\bar{f}_{\text{echo}}, \bar{p}_c) \approx (3.05, 0.90)$  denotes the configuration with  $\bar{\lambda} = 10$ , in which the  $\bar{R} = 3\bar{M}$  point overlaps with the maximum mass point. For the lines of different  $\bar{\lambda}$ , the left end of each curve maps to a similar  $\bar{f}_{\text{echo}} \sim 3.0$  value due to the same compactness there ( $\bar{R} = 3\bar{M}$ ). As the center pressure increases,  $\bar{f}_{\text{echo}}$  decreases due to the increasing compactness until the maximum mass point is reached. The  $\bar{\lambda} \rightarrow \infty$  case has the smallest  $\bar{f}_{\text{echo}}$  at its maximum mass configuration since it maps to the largest compactness. After rescaling back with Eq. (15), we obtain a simple relation between the minimal echo frequency and the effective bag constant<sup>2</sup>

$$f_{\text{echo}}^{\text{min}} \approx 3.16 \sqrt{B_{\text{eff}} / (10 \text{ MeV}/\text{fm}^3)} \text{ kHz}, \quad (16)$$

where  $B_{\text{eff}}$  is in units of  $\text{MeV}/\text{fm}^3$ , or equivalently

$$f_{\text{echo}}^{\text{min}} \approx 5.76 \sqrt{B_{\text{eff}} / (100 \text{ MeV})^4} \text{ kHz}, \quad (17)$$

where  $B_{\text{eff}}$  is in units of  $\text{MeV}^4$ . Thus, we see the minimal echo frequency is on the order of a few kHz when the effective bag constant is on its conventional order of magnitude  $B_{\text{eff}} \sim (100 \text{ MeV})^4$ .




Figure 3.  $\bar{f}_{\text{echo}} - \bar{p}_c$  of IQSs for different  $\bar{\lambda}$ , sampling  $\bar{\lambda} = 10$  (gray dot) and  $\bar{\lambda} = (20, 50, 100)$  from the lighter black line to darker black line respectively. The red line denotes  $\bar{\lambda} \rightarrow \infty$  case, with the corresponding EOS Eq. (7). The left and right ends of each line are truncated at the point where  $\bar{R} = 3\bar{M}$  and at the point of maximum star mass, respectively.

<sup>2</sup>Note that for the ad-hoc EOS  $p = \rho - 4B$  used in Ref. [56], we derived the corresponding scaling relation of  $f_{\text{echo}}$  in a form of simply multiplying the right side of Eq. (16) or (17) by a  $\sqrt{2}$  factor. This can successfully reproduce their results for their bag constant choices.

## SUMMARY

Interacting quark stars composed of interacting quark matter, including the interquark effects such as the pQCD corrections and the color-superconductivity, can have large compactness with large  $\bar{\lambda}$ , which characterizes the size of strong interactions in a dimensionless rescaling approach that can maximally reduce the number of degrees of freedom. We have shown that interacting quark stars with  $\bar{\lambda} \gtrsim 10$  can meet the compactness condition for generating GW echoes, i.e., featuring a photon sphere within the Buchdahl's limit. Taking the CFL phase for illustration, after rescaling the results back into the dimensional form, we explicitly constructed the corresponding dimensional parameter space of  $B_{\text{eff}}$  and  $\Delta$  with variations of  $a_4$  and  $m_s$  in their empirical range. Furthermore, we showed that a smaller echo frequency is achieved for a larger center pressure and a larger  $\bar{\lambda}$ , from which we obtain a general scaling relation for the minimal echo frequency  $f_{\text{echo}}^{\text{min}} \approx 5.76 \sqrt{B_{\text{eff}} / (100 \text{ MeV})^4} \text{ kHz}$ . Therefore, the echo frequencies for IQSs are on the order of a few kilohertz when the effective bag constant is on its conventional order  $B_{\text{eff}} \sim (100 \text{ MeV})^4$ . This study opens up the possibility of gravitational wave echoes being generated from physical compact stars in the conventional Einstein gravity framework.

**Acknowledgments.** We thank Bob Holdom for helpful discussions. This research is supported in part by the Natural Sciences and Engineering Research Council of Canada.

\* [zhangvchen@gmail.com](mailto:zhangvchen@gmail.com)

- [1] B. P. Abbott *et al.* [LIGO Scientific and Virgo], Phys. Rev. Lett. **116**, no.6, 061102 (2016) [arXiv:1602.03837 [gr-qc]].
- [2] B. P. Abbott *et al.* [LIGO Scientific and VIRGO], Phys. Rev. Lett. **118**, no.22, 221101 (2017) [erratum: Phys. Rev. Lett. **121**, no.12, 129901 (2018)] [arXiv:1706.01812 [gr-qc]].
- [3] B. P. Abbott *et al.* [LIGO Scientific and Virgo], Phys. Rev. X **9**, no.3, 031040 (2019) [arXiv:1811.12907 [astro-ph.HE]].
- [4] B. P. Abbott *et al.* [LIGO Scientific and Virgo], Phys. Rev. Lett. **119**, no.16, 161101 (2017) [arXiv:1710.05832]

- [gr-qc].
- [5] B. P. Abbott *et al.* [LIGO Scientific and Virgo Collaborations], *Phys. Rev. X* **9**, 011001 (2019) [arXiv:1805.11579 [gr-qc]].
- [6] B. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), *Astrophys. J. Lett.* **892**, L3 (2020) [arXiv:2001.01761 [astro-ph.HE]].
- [7] R. Abbott *et al.* [LIGO Scientific and Virgo Collaborations], *Astrophys. J. Lett.* **896**, L44 (2020) [arXiv:2006.12611 [astro-ph.HE]].
- [8] B. Holdom, R. B. Mann and C. Zhang, *Phys. Rev. D* **103**, no.12, 124046 (2021) [arXiv:2011.10179 [gr-qc]].
- [9] Y. G. Ignatev and A. V. Zakharov, *Phys. Lett. A* **66**, 3-4 (1978)
- [10] V. Cardoso, E. Franzin and P. Pani, *Phys. Rev. Lett.* **116**, no.17, 171101 (2016) [erratum: *Phys. Rev. Lett.* **117**, no.8, 089902 (2016)] [arXiv:1602.07309 [gr-qc]].
- [11] V. Cardoso, S. Hopper, C. F. B. Macedo, C. Palenzuela and P. Pani, *Phys. Rev. D* **94**, no.8, 084031 (2016) [arXiv:1608.08637 [gr-qc]].
- [12] V. Cardoso and P. Pani, [arXiv:1707.03021 [gr-qc]].
- [13] J. Abedi, H. Dykaar and N. Afshordi, *Phys. Rev. D* **96**, no.8, 082004 (2017) [arXiv:1612.00266 [gr-qc]].
- [14] Z. Mark, A. Zimmerman, S. M. Du and Y. Chen, “A recipe for echoes from exotic compact objects,” *Phys. Rev. D* **96**, no.8, 084002 (2017) [arXiv:1706.06155 [gr-qc]].
- [15] R. S. Conklin, B. Holdom and J. Ren, *Phys. Rev. D* **98**, no.4, 044021 (2018) [arXiv:1712.06517 [gr-qc]].
- [16] R. S. Conklin and B. Holdom, “Gravitational wave echo spectra,” *Phys. Rev. D* **100**, no.12, 124030 (2019) [arXiv:1905.09370 [gr-qc]].
- [17] R. S. Conklin, “Gravitational Wave Perturbations on a Kerr Background and Applications for Echoes,” *Phys. Rev. D* **101**, no.4, 044045 (2020) [arXiv:1911.07122 [gr-qc]].
- [18] J. Abedi and N. Afshordi, *JCAP* **11**, 010 (2019) [arXiv:1803.10454 [gr-qc]].
- [19] V. Cardoso and P. Pani, *Living Rev. Rel.* **22**, no.1, 4 (2019) [arXiv:1904.05363 [gr-qc]].
- [20] J. Abedi and N. Afshordi, [arXiv:2001.00821 [gr-qc]].
- [21] V. Ferrari and K. D. Kokkotas, *Phys. Rev. D* **62**, 107504 (2000) [arXiv:gr-qc/0008057 [gr-qc]].
- [22] B. Holdom and J. Ren, *Phys. Rev. D* **95**, no.8, 084034 (2017) [arXiv:1612.04889 [gr-qc]].
- [23] B. Holdom, [arXiv:1905.08849 [gr-qc]].
- [24] J. Ren, *Phys. Rev. D* **100**, no.12, 124012 (2019) [arXiv:1905.09973 [gr-qc]].
- [25] B. Holdom, *Phys. Rev. D* **101**, no.6, 064063 (2020) [arXiv:1909.11801 [gr-qc]].
- [26] H. A. Buchdahl, *Phys. Rev.* **116**, 1027 (1959)
- [27] Chandrasekhar, S. and Ferrari, V., 1991. Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences, 434(1891), pp.449-457.
- [28] P. Pani and V. Ferrari, *Class. Quant. Grav.* **35**, no.15, 15LT01 (2018) [arXiv:1804.01444 [gr-qc]].
- [29] A. R. Bodmer, *Phys. Rev. D* **4**, 1601 (1971).
- [30] E. Witten, *Phys. Rev. D* **30**, 272 (1984).
- [31] H. Terazawa, INS-Report-336 (INS, University of Tokyo, Tokyo) May, 1979.
- [32] B. Holdom, J. Ren and C. Zhang, *Phys. Rev. Lett.* **120**, no.22, 222001 (2018) [arXiv:1707.06610 [hep-ph]].
- [33] P. Haensel, J. L. Zdunik and R. Schaeffer, *Astron. Astrophys.* **160**, 121-128 (1986)
- [34] C. Alcock, E. Farhi and A. Olinto, *Astrophys. J.* **310**, 261-272 (1986)
- [35] C. Zhang, *Phys. Rev. D* **101**, no.4, 043003 (2020) [arXiv:1908.10355 [astro-ph.HE]].
- [36] Q. Wang, C. Shi and H. S. Zong, *Phys. Rev. D* **100**, no.12, 123003 (2019) Erratum: [*Phys. Rev. D* **100**, no.12, 129903 (2019)] [arXiv:1908.06558 [hep-ph]].
- [37] E. P. Zhou, X. Zhou and A. Li, *Phys. Rev. D* **97**, 083015 (2018) [arXiv:1711.04312 [astro-ph.HE]].
- [38] G. F. Burgio, A. Drago, G. Pagliara, H. J. Schulze and J. B. Wei, *Astrophys. J.* **860**, 139 (2018) [arXiv:1803.09696 [astro-ph.HE]].
- [39] Z. Roupas, G. Panotopoulos and I. Lopes, *Phys. Rev. D* **103**, no.8, 083015 (2021) [arXiv:2010.11020 [astro-ph.HE]].
- [40] J. E. Horvath and P. H. R. S. Moraes, *Int. J. Mod. Phys. D* **30**, no.03, 2150016 (2021) [arXiv:2012.00917 [astro-ph.HE]].
- [41] A. Kanakis-Pegios, P. S. Koliogiannis and C. C. Moustakidis, *Symmetry* **13**, no.2, 183 (2021) [arXiv:2012.09580 [astro-ph.HE]].
- [42] T. Harko, K. S. Cheng and Z. Kovacs, *Mon. Not. Roy. Astron. Soc.* **400**, 1632 (2009) [arXiv:0908.2672 [astro-ph.HE]].
- [43] T. Zhao, W. Zheng, F. Wang, C. M. Li, Y. Yan, Y. F. Huang and H. S. Zong, *Phys. Rev. D* **100**, no.4, 043018 (2019) [arXiv:1904.09744 [nucl-th]].
- [44] J. Ren and C. Zhang, *Phys. Rev. D* **102**, no.8, 083003 (2020) [arXiv:2006.09604 [hep-ph]].
- [45] C. Zhang and R. B. Mann, [arXiv:2009.07182 [astro-ph.HE]].
- [46] Z. Cao, L. W. Chen, P. C. Chu and Y. Zhou, [arXiv:2009.00942 [astro-ph.HE]].
- [47] E. Farhi and R. L. Jaffe, *Phys. Rev. D* **30**, 2379 (1984)
- [48] E. S. Fraga, R. D. Pisarski and J. Schaffner-Bielich, *Phys. Rev. D* **63**, 121702 (2001) [arXiv:hep-ph/0101143 [hep-ph]].

- [49] E. S. Fraga, A. Kurkela and A. Vuorinen, *Astrophys. J. Lett.* **781**, no.2, L25 (2014) [arXiv:1311.5154 [nucl-th]].
- [50] M. G. Alford, K. Rajagopal and F. Wilczek, *Nucl. Phys. B* **537**, 443-458 (1999) [arXiv:hep-ph/9804403 [hep-ph]].
- [51] K. Rajagopal and F. Wilczek, *Phys. Rev. Lett.* **86**, 3492-3495 (2001) [arXiv:hep-ph/0012039 [hep-ph]].
- [52] G. Lugones and J. E. Horvath, *Phys. Rev. D* **66**, 074017 (2002) [arXiv:hep-ph/0211070 [hep-ph]].
- [53] M. Alford and K. Rajagopal, *JHEP* **0206**, 031 (2002) [hep-ph/0204001].
- [54] M. Alford, M. Braby, M. W. Paris and S. Reddy, *Astrophys. J.* **629**, 969-978 (2005) [arXiv:nucl-th/0411016 [nucl-th]].
- [55] S. Weissenborn, I. Sagert, G. Pagliara, M. Hempel and J. Schaffner-Bielich, *Astrophys. J. Lett.* **740**, L14 (2011) [arXiv:1102.2869 [astro-ph.HE]].
- [56] M. Mannarelli and F. Tonelli, *Phys. Rev. D* **97**, no.12, 123010 (2018) [arXiv:1805.02278 [gr-qc]].
- [57] J. Bora and U. D. Goswami, *Mon. Not. Roy. Astron. Soc.* **502**, no.2, 1557-1568 (2021) [arXiv:2007.06553 [gr-qc]].
- [58] G. L. Volkmer and D. Hadjimichef, [arXiv:2107.06052 [hep-ph]].
- [59] P. Demorest, T. Pennucci, S. Ransom, M. Roberts and J. Hessels, *Nature* **467**, 1081 (2010) [arXiv:1010.5788 [astro-ph.HE]].
- [60] J. Antoniadis *et al.*, *Science* **340**, 1233232 (2013) [arXiv:1304.6875 [astro-ph.HE]].
- [61] H. T. Cromartie *et al.* [NANOGrav], *Nature Astron.* **4**, 72 (2020) [arXiv:1904.06759 [astro-ph.HE]].
- [62] T. E. Riley *et al.*, *Astrophys. J. Lett.* **887**, L21 (2019) [arXiv:1912.05702 [astro-ph.HE]].
- [63] M. Miller *et al.*, *Astrophys. J. Lett.* **887**, L24 (2019) [arXiv:1912.05705 [astro-ph.HE]].
- [64] R. C. Tolman, *Phys. Rev.* **55**, 364 (1939).
- [65] J. R. Oppenheimer and G. M. Volkoff, *Phys. Rev.* **55**, 374 (1939).
- [66] K. A. Olive and Particle Data Group, *Chinese Physics C* **38**, 090001 (2014).