

Behaviour of trajectories near a two-cycle heteroclinic network

Olga Podvigina

Institute of Earthquake Prediction Theory
and Mathematical Geophysics

84/32 Profsoyuznaya St, 117997 Moscow, Russian Federation

November 23, 2021

Abstract

We study behaviour of trajectories near a type Z heteroclinic network which is a union of two cycles. Analytical and numerical studies indicate that attractiveness of this network can be associated with various kinds of dynamics in its vicinity, one or both of these cycle being fragmentarily asymptotically stable, or both being completely unstable. In the latter case trajectories can switch irregularly between the cycles, or they can make a certain number of turns around one of them before switching to the other one. Regular behaviour of trajectories near a heteroclinic network can be described using the notion of an omnicycle, which we introduce in this paper. In particular, we use it to prove that the network can be fragmentarily asymptotically stable even if both cycles are completely unstable.

1 Introduction

There is a number of reasons for chaotic behaviour of trajectories in a vicinity of heteroclinic network. This can be due to the complexity of the network as, e.g., when a network involves nodes which are chaotic sets [5, 12] or is a union of periodic orbits connected by an infinite number of heteroclinic and homoclinic orbits [33]. Chaotic behaviour can be caused by chaotic forcing, even when the amplitude of the added noise is arbitrarily small [4]. When eigenvalues of linearisations near some of the equilibria in the network are complex, the return map near a heteroclinic cycle may possess a horseshoe, thus resulting in emergence of a strange attractor [3, 2]. Chaos due to the presence of complex eigenvalues of linearisations was found in [17] and [32], where either a horseshoe does not exist, or its existence has not been investigated. Strange attractors may also exist in periodically-perturbed dynamical systems with an attracting heteroclinic network [21, 20].

Examples of more primitive heteroclinic networks comprised of a finite number of equilibria and heteroclinic connections feature only real eigenvalues of linearisation near equilibria;

in their vicinity irregular switching between connections takes place, indicating chaotic behaviour of nearby trajectories [1, 30, 31]. Complexity of a network can be measured by the number of splitting nodes (for the definition see, e.g., [7]). For the networks studied in [1, 30, 31], it is the number of equilibria having more than one outgoing connections (six, nine and five, respectively). In the present paper we consider such a primitive network which has only a single splitting node. Our numerical simulations indicate a possibility of irregular switching of nearby trajectories.

A heteroclinic cycle is a union of nodes and connecting trajectories, where all nodes are required to be distinct. A heteroclinic network is a connected union of heteroclinic cycles. In this paper we introduce the notion of an omnicycle, an ordered sequence of nodes and heteroclinic connections, where repeating nodes are allowed. Regarded as a subset of the phase space, an omnicycle is either a heteroclinic cycle or a network, however in the latter case the stability properties of these objects can be very different. Omnicycles were implicitly employed in [30] to describe the behaviour of trajectories near a heteroclinic network, where existence of an attracting omnicycle was indicated by so-called “regular cycling”. However, the terminology introduced *ibid* was closely related to the specific structure of the considered network, while the definition introduced in the present paper is of a general type.

Omnicycles, similarly to heteroclinic cycles and network, do not exist in a generic dynamical systems, because a connection between saddles can be destroyed by a small perturbation. However, they may exist and be robust in a system where some restrictions are imposed, e.g., symmetries [18, 19], natural constraints, as in population dynamics or evolutionary game theory [1, 14, 15] or due to prescribed patterns of interaction, as in a system of coupled cells or oscillators [9, 6].

An invariant set of a dynamical system is called asymptotically stable if it attracts all trajectories from its small neighbourhood. A non-asymptotically stable set can be stable in a weaker sense and attract a positive measure set of initial conditions from its small neighbourhood. In [22] this property was called fragmentary asymptotic stability. Conditions for asymptotic stability or fragmentary asymptotic stability for certain types of heteroclinic cycles are known for a long time, see, e.g., Podvigina [22] or [29] and references therein. Typically, conditions for stability of a cycle are derived by following nearby trajectories and constructing the respective transition maps. Since it is irrelevant how many times each particular node or connection is present in the sequence, the existing conditions for stability of cycles can be straightforwardly employed for stability of omnicycles. The study of stability of a heteroclinic network is more difficult, because a trajectory can follow different sequences of connections along the network.

If a fragmentarily asymptotically stable (f.a.s.) cycle is a subset of a heteroclinic network then the whole network also is f.a.s. For all networks in \mathbb{R}^4 that were studied in literature the converse is also true: if a heteroclinic network is f.a.s. then at least one of its subcycles is f.a.s. [8, 16, 10, 11, 25, 26, 27]. The same property holds true for some networks in higher-dimensional subspaces, namely stability of a network implies stability of at least one of its subcycles [1, 28]. The authors of [30, 31] presented examples of f.a.s. networks, where all subcycles are completely unstable. To prove stability the authors have shown that certain




Figure 1: The network studied in sections 4 and 5. The crosssections H_L and H_R employed in the derivation of conditions of stability are introduced in section 3.

omnicycles, that are subsets of the network, are f.a.s. (To be more precise, in the proofs of [30] a different technique was employed, which in fact is similar to our theorem 2).

In this paper we study stability of a network in \mathbb{R}^6 which is a union of two cycles, $\mathcal{X} = \mathcal{C}_L \cup \mathcal{C}_R \subset \mathbb{R}^6$, admitted by a \mathbb{Z}_2^6 -equivariant dynamical system

$$\dot{\mathbf{x}} = f(\mathbf{x}), \quad f : \mathbb{R}^6 \rightarrow \mathbb{R}^6, \quad (1)$$

where \mathbb{Z}_2^6 is generated by symmetries reversing the sign of one of the spatial coordinates. Six equilibria involved in the network belong to coordinate axes and heteroclinic connection belongs to coordinate planes. The cycles are:

$$\mathcal{C}_L : (\xi_1 \rightarrow \xi_2 \rightarrow \xi_3 \rightarrow \xi_4 \rightarrow) \text{ and } \mathcal{C}_R : (\xi_1 \rightarrow \xi_2 \rightarrow \xi_5 \rightarrow \xi_6 \rightarrow).$$

In terminology of [22], \mathcal{X} is a *type Z* heteroclinic network and \mathcal{C}_L and \mathcal{C}_R are *type Z* heteroclinic cycles. Generalisation of this classification to omnicycles implies that any omnicycle, which is a subset of \mathcal{X} , is a *type Z* omnicycle. Theorems in [22] that give necessary and sufficient conditions for a *type Z* heteroclinic cycle to be f.a.s. are also applicable to study stability of *type Z* heteroclinic omnicycles.

The conditions are inequalities that involve eigenvalues and eigenvectors of transition matrices, whose entries depend on eigenvalues of Jacobians $df(\xi_j)$. For \mathcal{C}_L and \mathcal{C}_R the respective 4×4 transitions matrices are sparse, hence allowing to calculate eigenvalues and eigenvectors. Transition matrix of the omnicycle \mathcal{C}_{LR} , where

$$\mathcal{C}_{LR} : (\xi_1 \rightarrow \xi_2 \rightarrow \xi_3 \rightarrow \xi_4 \rightarrow \xi_1 \rightarrow \xi_2 \rightarrow \xi_5 \rightarrow \xi_6 \rightarrow),$$

is full, therefore eigenvalues and eigenvectors can not be found analytically in a general case. In order to be able to calculate them, we assume that some eigenvalues of $df(\xi_j)$ are much larger then the others. For vanishing small eigenvalues we calculate eigenvalues and

eigenvectors of the respective sparse transition matrix. Because of continuous dependence of eigenvalues and eigenvectors of a matrix on its entries, the stability or instability conditions are satisfied true for small non-vanishing eigenvalues of $df(\xi_j)$ as well.

The paper is organised as follows: In Section 2 we recall basic definitions of robust heteroclinic cycles, networks and asymptotic stability and in Section 3 we discuss asymptotic stability of omnicycles. In sections 4 and 5 we study analytically and numerically asymptotic stability of network shown in fig. 1. Namely, in section 4 we derive conditions of stability of the cycles \mathcal{C}_L and \mathcal{C}_R and show that the network can be f.a.s. while both cycles are completely unstable. In section 5 we present results of numerical simulations of solutions of system (1), in particular, in the cases when all \mathcal{C}_L , \mathcal{C}_R and \mathcal{C}_{LR} are completely unstable. By varying the r.h.s. of (1) we obtain examples of f.a.s. omnicycles, different from \mathcal{C}_{LR} , or of apparently chaotic switching between \mathcal{C}_L and \mathcal{C}_R .

2 Background and definitions

In this section we introduce the notions of omnicycles and trail-stability, and recall definitions that will be used in the following sections.

2.1 Robust heteroclinic cycles, omnicycles and networks

Consider a smooth Γ -equivariant dynamical systems

$$\dot{x} = f(x), \quad f(\gamma \mathbf{x}) = \gamma f(\mathbf{x}), \quad \text{for all } \gamma \in \Gamma \text{ and } x \in \mathbb{R}^n, \quad (2)$$

where we assume that $\Gamma \subset \mathbf{O}(n)$ is finite. For a group Γ acting on \mathbb{R}^n the *isotropy group* of the point $x \in \mathbb{R}^n$ is the subgroup $\Sigma_x = \{\gamma \in \Gamma : \gamma x = x\}$, and a *fixed-point subspace* of a subgroup $\Sigma \subset \Gamma$ is the subspace $\text{Fix}(\Sigma) = \{\mathbf{x} \in \mathbb{R}^n : \sigma \mathbf{x} = \mathbf{x} \text{ for all } \sigma \in \Sigma\}$. (For more details about equivariant dynamical systems see, e.g., Golubitsky and Stewart [13].)

Let ξ_1, \dots, ξ_m be hyperbolic equilibria of (2), where $\xi_j \neq \xi_k$ for any $1 \leq j, k \leq m$, $j \neq k$, and $\kappa_{j,j+1} \neq \emptyset$ be a set of trajectories from ξ_j to ξ_{j+1} , where $\xi_{m+1} = \xi_1$ is assumed. A *heteroclinic cycle* is an invariant set $X \subset \mathbb{R}^n$ which is a union of equilibria $\{\xi_1, \dots, \xi_m\}$ and heteroclinic connections $\{\kappa_{12}, \dots, \kappa_{m1}\}$. A *heteroclinic network* is a connected union of a finite number of heteroclinic cycles.

Definition 1 *An omnicycle is a union of equilibria $\{\xi_1, \dots, \xi_m\}$ and heteroclinic connections, $\{\kappa_{12}, \dots, \kappa_{m1}\}$, where $\kappa_{ij} \subset W^u(\xi_i) \cap W^s(\xi_j)$ and $\kappa_{ij} \neq \emptyset$.*

I.e., the difference between a heteroclinic cycle and an omnicycle is that for the omnicycle repeating equilibria are allowed.

By analogy with a building block of a heteroclinic cycle [25], we introduce building block of an omnicycle.

Definition 2 *Given an omnicycle $\mathcal{X} = \{\xi_1, \dots, \xi_m; \kappa_{12}, \dots, \kappa_{m1}\}$, we say that its subset $\{\xi_1, \dots, \xi_j; \kappa_{12}, \dots, \kappa_{j,j+1}\}$ and the symmetry $\gamma \in \Gamma$ is a building block of \mathcal{X} if $\gamma \xi_i = \xi_{i+j}$ and $\gamma \kappa_{i,i+1} = \kappa_{i+j,i+j+1}$ for any $\xi_i \in \mathcal{X}$ and $\kappa_{i,i+1} \subset \mathcal{X}$.*

A heteroclinic cycle (or an omnicycle, or a network) \mathcal{X} is called *robust*, if any connection $\kappa_{ij} \subset \mathcal{X}$ belongs to a flow-invariant subspace P_{ij} . In system (2) the invariance of P_{ij} typically follows from its equivariance, namely that $P_{ij} = \text{Fix}(\Sigma_{ij})$ for some $\Sigma_{ij} \subset \Gamma$. By Δ_j we denote the isotropy subgroup of ξ_j and $L_j = \text{Fix}(\Delta_j)$. Denote by P_{ij}^\perp and by L_j^\perp the orthogonal complement to P_{ij} and L_j , respectively, in \mathbb{R}^n .

Given an equilibrium $\xi_j \in L_j = P_{j-1,j} \cap P_{j,j+1}$ that belongs to a robust heteroclinic cycle $\mathcal{C} = (\xi_1 \rightarrow \xi_2 \rightarrow \dots \rightarrow \xi_m \rightarrow)$, the eigenvalues of the Jacobian $df(\xi_j)$ can be divided into *radial* (the associated eigenvectors belong to L_j), *contracting* (the eigenvectors belong to $P_{j-1,j} \ominus L_j$), *expanding* (the eigenvectors belong to $P_{j,j+1} \ominus L_j$) and *transverse* (the remaining ones), where $P \ominus L$ denotes the orthogonal complement to L in P .

Definition 3 (Adapted from [22]) *A heteroclinic network \mathcal{X} (cycle or omnicycle) is called of type Z if*

- for any $\xi_j \in \mathcal{X}$ we have that $\dim P_{i,j} = \dim P_{j,k}$ for any incoming connection $\kappa_{i,j}$ and any outgoing connection $\kappa_{j,k}$;
- for any $\kappa_{i,j} \subset \mathcal{X}$ the isotropy subgroup $\Sigma_{i,j}$ decomposes $P_{i,j}^\perp$ into one-dimensional isotypic components.

2.2 Stability

Denote by $\Phi(\mathbf{x}, \tau)$ a trajectory of system (2) through the point $\mathbf{x} \in \mathbb{R}^n$ and by $\Phi(\mathbf{x}, (\tau_1, \tau_2)) = \cup_{\tau \in (\tau_1, \tau_2)} \Phi(\mathbf{x}, \tau)$ the subset of this trajectory corresponding to the time interval (τ_1, τ_2) . For a set $X \subset \mathbb{R}^n$ and $\varepsilon > 0$, the ε -neighbourhood of X is

$$B_\varepsilon(X) = \{\mathbf{x} \in \mathbb{R}^n : d(\mathbf{x}, X) < \varepsilon\}. \quad (3)$$

Given a compact invariant set X of (2), its δ -local basin of attraction defined as

$$\mathcal{B}_\delta(X) = \{\mathbf{x} \in \mathbb{R}^n : d(\Phi(\mathbf{x}, \tau), X) < \delta \text{ for any } \tau \geq 0 \text{ and } \lim_{\tau \rightarrow \infty} d(\Phi(\mathbf{x}, \tau), X) = 0\}. \quad (4)$$

Definition 4 *An invariant set X is called asymptotically stable, if for any $\delta > 0$ there exists an $\varepsilon > 0$ such that*

$$B_\varepsilon(X) \subset \mathcal{B}_\delta(X).$$

Definition 5 *An invariant set X is called fragmentarily asymptotically stable, if for any $\delta > 0$*

$$\mu(\mathcal{B}_\delta(X)) > 0.$$

(Here μ is the Lebesgue measure of a set in \mathbb{R}^n .)

Definition 6 *An invariant set X is called completely unstable, if there exists $\delta > 0$ such that $\mu(\mathcal{B}_\delta(X)) = 0$.*

Definition 7 Given an omnicycle $\mathcal{X} = \{\xi_1, \dots, \xi_m; \kappa_{12}, \dots, \kappa_{m1}\}$, we say that a trajectory $\Phi(\mathbf{x}, (\tau^{in}, \tau^{fin}))$ ε -follows this omnicycle if

$$\Phi(\mathbf{x}, (\tau^{in}, \tau^{fin})) \subset N_\varepsilon(\mathcal{X})$$

and there exist a number $1 \leq j_0 \leq m$ and time instances $\{\tau'_0, \tau''_0, \tau'_1, \tau''_1, \dots, \tau'_S, \tau''_S\}$, where $\tau^{in} \leq \tau'_0 < \tau''_0 < \tau'_1 < \dots < \tau'_S < \tau''_S \leq \tau^{fin}$, such that

$$\begin{aligned} & \Phi(\mathbf{x}, (\tau'_s, \tau''_s)) \subset B_\varepsilon(\xi_j), \text{ where } j = j_0 + s(\text{mod } m), \text{ for all } 0 \leq s \leq S, \text{ and} \\ & \left(\left(\bigcup_{0 \leq s < S} \Phi(\mathbf{x}, (\tau''_s, \tau'_{s+1})) \right) \cup \Phi(\mathbf{x}, (\tau^{in}, \tau_0)) \cup \Phi(\mathbf{x}, (\tau_S, \tau^{fin})) \right) \cap \left(\bigcup_{1 \leq j \leq m} B_\varepsilon(\xi_j) \right) = \emptyset. \end{aligned}$$

Definition 8 Given an omnicycle \mathcal{X} of (2), its δ -local basin of attraction is defined as

$$\mathcal{B}_\delta^{\text{omni}}(\mathcal{X}) = \{\mathbf{x} \in \mathbb{R}^n : \Phi(\mathbf{x}, (0, \tau)) \delta\text{-follows } \mathcal{X} \text{ for any } \tau > 0 \text{ and } \lim_{\tau \rightarrow \infty} d(\Phi(\mathbf{x}, \tau), \mathcal{X}) = 0\}. \quad (5)$$

Definition 9 An omnicycle \mathcal{X} is called trail-stable, if for any $\delta > 0$

$$\mu(\mathcal{B}_\delta^{\text{omni}}(\mathcal{X})) > 0.$$

Remark 1 For an omnicycle which is a heteroclinic cycle the notions of f.a.s. and trail-stability are equivalent, because trajectories near the cycle visit the equilibria in the prescribed order specified in the definition of the cycle. An omnicycle, which is not a cycle, can be f.a.s. according to definition 5 without being trail-stable.

Remark 2 Suppose \mathcal{X} is a robust type Z omnicycle, which is not a heteroclinic cycle, in a Γ -equivariant system. Then \mathcal{X} is not asymptotically stable. This can be shown by arguments similar to the ones employed in the proof of theorem 3.1 in [28].

3 Stability of omnicycles

In this section we state conditions for trail-stability of type Z omnicycles. Due to similarity between heteroclinic cycles and omnicycles, the theorem of [22], which gives conditions for fragmentary asymptotic stability of heteroclinic cycles of type Z, is also applicable to omnicycles of that type. The conditions involve eigenvalues and eigenvectors of transition matrices, which are constructed from local and global maps defined below.

3.1 Local and global maps

To study stability of an omnicycle, similarly to heteroclinic cycles and networks, the behaviour of nearby trajectories is approximated by local and global maps. By $N_{\tilde{\delta}}(\xi_j)$ we denote a neighbourhood of ξ_j bounded by $|a_s| = \tilde{\delta}$, where a_s are coordinates in the basis

comprised of eigenvectors of $df(\xi_j)$. We assume $\tilde{\delta}$ to be fixed and small, so that in $N_{\tilde{\delta}}(\xi_j)$ the equation (2) can be approximated by its linearisation $df(\xi_j)$, and $\tilde{\delta} \gg \varepsilon$ and δ , where ε and δ are those employed in the definitions of stability in subsection 2.2

Given an omnicycle $\mathcal{C} : (\xi_1 \rightarrow \dots \rightarrow \xi_m \rightarrow)$, let $(\mathbf{u}_j, v_j, w_j, \mathbf{z}_j)$ be local coordinates near ξ_j in the basis, where radial eigenvectors come the first (the respective coordinates are \mathbf{u}_j), followed by the contracting and the expanding eigenvectors, the transverse eigenvectors being the last. (If repeating equilibria are present in the sequence, i.e. $\xi_i = \xi_j$ for $i \neq j$, then such decompositions for eigenvectors of $df(\xi_i)$ and $df(\xi_j)$ can be different.) Denote by $H_{j-1,j}^{(in)}$ the face of $N_{\tilde{\delta}}(\xi_j)$ that intersects with the connection $\kappa_{j-1,j}$ and by $H_{j,j+1}^{(out)}$ the face intersected by $\kappa_{j,j+1}$. In literature, $H_{j-1,j}^{(in)}$ and $H_{j,j+1}^{(out)}$ are often called crosssections or Poincaré sections. For a trajectory close to the cycle the local map relates coordinates of its intersection with $H_{j-1,j}^{(in)}$ to the coordinates of its intersection with $H_{j,j+1}^{(out)}$.

Let the superscripts in and out denote the coordinate in $H_{j-1,j}^{(in)}$ and $H_{j,j+1}^{(out)}$, respectively. The local map $\phi_j : H_{j-1,j}^{(in)} \rightarrow H_{j,j+1}^{(out)}$ (see, e.g., [18] or [22]) is

$$v^{out} = K_{1j}(w^{in})^{-c_j/e_j} \text{ and } \mathbf{z}^{out} = \mathbf{K}_j \mathbf{z}^{in}(w^{in})^{-\mathbf{t}_j/e_j}, \quad (6)$$

where K_{1j} and \mathbf{K}_j are constants (for a fixed $\tilde{\delta}$), and c_j, e_j, \mathbf{t}_j are the contracting, expanding and transverse eigenvalues of $df(\xi_j)$, respectively. The coordinates $\mathbf{u}^{in}, v^{in}, \mathbf{u}^{out}$ and w^{out} are ignored, because they are irrelevant in the study of stability.

Near the connection $\kappa_{j,j+1}$ the system (2) can be approximated by a global map (also called a *connecting diffeomorphism*) $\psi_{j,j+1} : H_{j,j+1}^{(out)} \rightarrow H_{j,j+1}^{(in)}$, which relates coordinates of the point where a trajectory exits $N_{\tilde{\delta}}(\xi_j)$ to coordinates of the point where it enters $N_{\tilde{\delta}}(\xi_{j+1})$. The global map is predominantly linear

$$(w^{j+1,in}, \mathbf{z}^{j+1,in}) = A'_j(v^{j,out}, \mathbf{z}^{j,out}), \quad (7)$$

where A'_j is an $(n_t + 1) \times (n_t + 1)$ matrix. For type Z omnicycles the matrix A'_j is a product of a diagonal matrix A''_j and a permutation matrix A_j .

Denote by g_j the superpositions of the local and global maps:

$$g_j = \phi_j \circ \psi_{j-1,j} : H_j^{(out)} \rightarrow H_{j+1}^{(out)}.$$

We call the set of maps $\{g_{1:m}\} = \{g_1, \dots, g_m\}$, $g_j : \mathbb{R}^{n_t+1} \rightarrow \mathbb{R}^{n_t+1}$, *the collection of maps associated with the omnicycle* $\mathcal{C} = \{\xi_1, \dots, \xi_m\}$. For a given a collection of maps $\{g_1^m\}$ we define superpositions

$$G_j = g_{j-1} \circ \dots \circ g_1 \circ g_m \circ \dots \circ g_{j+1} \circ g_j, \quad G_j : H_j^{(out)} \rightarrow H_j^{(out)}, \quad (8)$$

which approximate the behaviour of trajectories near the cycle and are called *return maps*.

3.2 Transition maps and transition matrices

Let us introduce new coordinates:

$$\eta = (\ln |w|, \ln |z_1|, \dots, \ln |z_{n_t}|). \quad (9)$$

Due to the smallness of ε , in these coordinates the maps g_j become approximately linear, $\eta^{j+1} = M_j \eta^j$, where the transition matrix M_j is a product of permutation matrix A_j that relates local coordinates near ξ_{j+1} to the ones near ξ_j and the matrix

$$B_j := \begin{pmatrix} b_{j,1} & 0 & 0 & \dots & 0 \\ b_{j,2} & 1 & 0 & \dots & 0 \\ b_{j,3} & 0 & 1 & \dots & 0 \\ \cdot & \cdot & \cdot & \dots & \cdot \\ b_{j,N} & 0 & 0 & \dots & 0 \end{pmatrix}. \quad (10)$$

The entries $b_{j,l}$ of the matrix B_j depend on the eigenvalues of the linearisation $df(\xi_j)$ of (2) near ξ_j as follows

$$b_{j,1} = c_j/e_j \text{ and } b_{j,l+1} = -t_{j,l}/e_j, \quad 1 \leq l \leq n_t, \quad 1 \leq j \leq m. \quad (11)$$

Transition matrices of the superposition G_j defined in (8) are the products

$$\mathcal{M}^{(j)} = M_{j-1} \dots M_1 M_m \dots M_{j+1} M_j.$$

Consider the matrix $M := \mathcal{M}^{(1)} = M_m \dots M_1 : \mathbb{R}^N \rightarrow \mathbb{R}^N$; it is a product of the basic transition matrices of the form (10). We separate the coordinate vectors \mathbf{e}_l , $1 \leq l \leq N$, into two groups. The first group is comprised of the vectors \mathbf{e}_l for which there exist such k and j that $(A^{(j)})^k A_{j-1,1} \mathbf{e}_l = \mathbf{e}_1$ (recall that A_j are permutation matrices), the second one incorporates the remaining vectors. Denote by V^{sig} and V^{ins} the subspaces spanned by vectors from the first and second group, respectively (the superscripts “ins” and “sig” stand for significant and insignificant). In the basis where significant vectors come the first, matrix M takes the form

$$M = \left(\begin{array}{c|cccc} M^{\text{sig}} & & & & 0 \\ \hline & 1 & 0 & \dots & 0 \\ M^{\text{ins}} & 0 & 1 & \dots & 0 \\ & \cdot & \cdot & \dots & \cdot \\ & 0 & 0 & \dots & 1 \end{array} \right). \quad (12)$$

Theorem 1 (Adapted from [22].) *Let V^{sig} and V^{ins} be the subspaces defined above.*

- (a) *The subspace V^{ins} is M -invariant and all eigenvalues associated with the eigenvectors from this subspace are one.*
- (b) *Generically all components of eigenvectors that do not belong to V^{ins} are non-zero.*

Remark 3 *In the study of stability instead of whole heteroclinic cycle or omnicycle one can consider its building block, see remark 2. In such a case the transition matrix take the form $M^{\text{block}} = AM$, where A is the permutation matrix relating bases $\{\mathbf{e}_l\}$ and $\{\gamma \mathbf{e}_l\}$ and M has the form (12). The statement (a) therefore becomes [22] “The subspace V^{ins} is M -invariant and the absolute value of all eigenvalues associated with the eigenvectors from this subspace is one.”*

We call *insignificant* the eigenvalues associated with eigenvectors from V^{ins} , and *significant* the rest ones. Generically the absolute values of all significant eigenvalues differ from one. Below $\lambda_{\text{max}} \neq 1$ denotes the largest significant eigenvalue of a transition matrix $\mathcal{M}^{(j)}$.

The construction of transition matrices is identical for heteroclinic cycles and omnicycles. Whether or not a set of trajectories stays close to a given sequence of connections is independent of presence or absence of repeating equilibria or connections in this sequence. Therefore the theorem of [22] that gives necessary and sufficient conditions for a heteroclinic cycle to be f.a.s. can be extended to an omnicycle as follows.

Theorem 2 (Adapted from [22].) *Let M_j be basic transition matrices of a collection of maps $\{g_1^m\}$ associated with an omnicycle of type Z . (For type Z omnicycles the matrices are of the form (10).) Denote by $j = j_1, \dots, j_L$ the indices, for which M_j involves negative entries; all entries are non-negative for all remaining j . Denote by λ_{max} the dominant eigenvalue of $\mathcal{M}^{(j)}$ and by \mathbf{w}_{max} the associated eigenvector.*

(a) *If for at least one $j = j_l + 1$ the matrix $\mathcal{M}^{(j)}$ does not satisfy the following conditions*

- (i) λ_{max} is real;
- (ii) $\lambda_{\text{max}} > 1$;
- (iii) $w_l^{\text{max}} w_q^{\text{max}} > 0$ for all l and q , $1 \leq l, q \leq N$.

then the omnicycle is trail-unstable.

(b) *If the matrices $\mathcal{M}^{(j)}$ satisfy the above conditions (i)-(iii) for all j such that $j = j_l + 1$, then the omnicycle is trail-stable.*

4 Fragmentary asymptotic stability: analytical results

In this section we apply theorem 2 to study dynamics near the network $\mathcal{X} = \mathcal{C}_L \cup \mathcal{C}_R$ shown in fig. 1. The network exists in a Γ -equivariant system (2) in \mathbb{R}^6 , where the group $\Gamma \cong \mathbb{Z}_2^6$ is generated by symmetries inverting the sign of one of coordinates. The equilibria in the network belong to coordinate axes, the connections are one-dimensional and belong to coordinate planes, and the equilibria are stable in directions transverse to the network.

According to theorem 2, stability of a cycle or an omnicycle depends on the dominant eigenvalue and the associated eigenvectors of the respective transition matrices. Transition matrices of the cycles \mathcal{C}_L and \mathcal{C}_R have two-dimensional significant subspace, therefore we can calculate the dominant eigenvalues and the associated eigenvectors.

The significant subspace of the transition matrix of the omnicycle

$$\mathcal{C}_{LR} = (\xi_1 \rightarrow \xi_2 \rightarrow \xi_3 \rightarrow \xi_4 \rightarrow \xi_1 \rightarrow \xi_2 \rightarrow \xi_5 \rightarrow \xi_6 \rightarrow)$$

has dimension four, therefore in a general case its eigenvalues can not be found analytically. To show that the omnicycle can be trail-stable, we calculate the dominant eigenvalue and

the associated eigenvector in the case when some eigenvalues of $df(\xi_j)$ vanish. Since the conditions for stability are non-strict inequalities, the inequalities remains true for small non-vanishing eigenvalues as well. Under our assumption that the chosen eigenvalues of $df(\xi_j)$ vanish, the conditions for stability of \mathcal{C}_{LR} are not compatible with conditions for stability of \mathcal{C}_L and \mathcal{C}_R . So, we prove that the network $\mathcal{X} = \mathcal{C}_L \cup \mathcal{C}_R$ can be f.a.s. while both subcycles are completely unstable.

4.1 Stability of cycles \mathcal{C}_L and \mathcal{C}_R

Since the system under consideration is \mathbb{Z}_2^6 -symmetric, the eigenvalues of $d(\xi_j)$ coincide with the Cartesian basis vectors \mathbf{e}_j , $1 \leq j \leq 6$. In crosssections to the connections comprising the cycle \mathcal{C}_L we take the following local bases:

$$H_{12}^{out} : \{\mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5, \mathbf{e}_6\}, H_{23}^{out} : \{\mathbf{e}_4, \mathbf{e}_1, \mathbf{e}_5, \mathbf{e}_6\}, H_{34}^{out} : \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_5, \mathbf{e}_6\}, H_{41}^{out} : \{\mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_5, \mathbf{e}_6\}. \quad (13)$$

In agreement with (10) the basic transition matrices are:

$$M_{412} = \begin{pmatrix} b_{132} & 1 & 0 & 0 \\ b_{142} & 0 & 0 & 0 \\ b_{152} & 0 & 1 & 0 \\ b_{162} & 0 & 0 & 1 \end{pmatrix}, \quad M_{123} = \begin{pmatrix} b_{243} & 1 & 0 & 0 \\ b_{213} & 0 & 0 & 0 \\ b_{253} & 0 & 1 & 0 \\ b_{263} & 0 & 0 & 1 \end{pmatrix} \quad (14)$$

$$M_{234} = \begin{pmatrix} b_{314} & 1 & 0 & 0 \\ b_{324} & 0 & 0 & 0 \\ b_{354} & 0 & 1 & 0 \\ b_{364} & 0 & 0 & 1 \end{pmatrix}, \quad M_{341} = \begin{pmatrix} b_{421} & 1 & 0 & 0 \\ b_{431} & 0 & 0 & 0 \\ b_{451} & 0 & 1 & 0 \\ b_{461} & 0 & 0 & 1 \end{pmatrix},$$

where

$$b_{ijk} = -\mu_{ij}/\mu_{ik} \quad (15)$$

and μ_{ij} is the eigenvalue of $df(\xi_i)$ associated with the eigenvector \mathbf{e}_j .

The cycle \mathcal{C}_L has only one equilibrium, ξ_2 , where a transverse eigenvalue of linearisation is positive. Therefore, theorem 2 implies that the cycle is f.a.s. whenever the matrix $M_L = M_{123}M_{412}M_{341}M_{234}$ satisfies the conditions (i)-(iii). Multiplying the matrices (14) we obtain that

$$M_L = \begin{pmatrix} a_{11} & a_{12} & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 \\ a_{31} & a_{32} & 1 & 0 \\ a_{41} & a_{42} & 0 & 1 \end{pmatrix}, \quad (16)$$

where

$$\begin{aligned}
a_{11} &= (b_{314}b_{421} + b_{324})(b_{132}b_{243} + b_{142}) + b_{314}b_{431}b_{243} \\
a_{21} &= (b_{314}b_{421} + b_{324})b_{132}b_{213} + b_{314}b_{431}b_{213} \\
a_{31} &= (b_{314}b_{421} + b_{324})(b_{132}b_{253} + b_{152}) + b_{314}b_{431}b_{253} \\
a_{41} &= (b_{314}b_{421} + b_{324})(b_{132}b_{263} + b_{162}) + b_{314}b_{431}b_{263} \\
a_{21} &= b_{421}(b_{132}b_{243} + b_{142}) + b_{431}b_{243}, \quad a_{22} = b_{421}b_{132}b_{213} + b_{431}b_{213} \\
a_{32} &= b_{421}(b_{132}b_{253} + b_{152}) + b_{431}b_{253}, \quad a_{42} = b_{421}(b_{132}b_{263} + b_{162}) + b_{431}b_{263}.
\end{aligned} \tag{17}$$

Due to the assumption that the network \mathcal{X} is stable in the transverse directions, in (17) the ratios b_{ijk} are positive, except for b_{253} . Therefore, the entries of matrix (16) are positive, except possibly for a_{31} and a_{32} .

The dominant eigenvalue of M_L , λ_{\max} , is the largest in absolute value eigenvalue of its upper left 2×2 submatrix which we denote by M'_L . The entries of M'_L are positive, implying that the discriminant of its characteristic polynomial $F(\lambda)$ is positive. Hence, the eigenvalues of M'_L are real, which implies that λ_{\max} satisfies condition (i) of theorem 2. Since $F(a_{11}) = F(a_{22}) = -a_{12}a_{21} < 0$, the dominant eigenvalue satisfies $\lambda_{\max} > \max(a_{11}, a_{22}) > 0$. Condition (ii) is satisfied whenever [24]

$$\max\left(\frac{a_{11} + a_{22}}{2}, a_{11} + a_{22} - a_{11}a_{22} + a_{12}a_{21}\right) > 1. \tag{18}$$

Let $\mathbf{v} = (v_1, v_2, v_3, v_4)$ be the eigenvector of A associated with λ_{\max} and $v_1 = 1$. The component v_2 is positive because

$$v_2 = a_{21}v_1/(\lambda_{\max} - a_{22}). \tag{19}$$

From (16) the latter two components satisfy

$$v_3 = \frac{a_{31}v_1 + a_{32}v_2}{\lambda_{\max} - 1}, \quad v_4 = \frac{a_{41}v_1 + a_{42}v_2}{\lambda_{\max} - 1}. \tag{20}$$

Since $a_{41} > 0$ and $a_{42} > 0$, the latter equality implies that $v_4 > 0$. Substituting (19) into the first inequality in (20) we obtain that (iii) is equivalent to

$$a_{31}(\lambda_{\max} - a_{22}) + a_{32}a_{21} > 0. \tag{21}$$

Therefore, the conditions for fragmentary asymptotic stability of the cycle \mathcal{C}_L are (18) and (21), where λ_{\max} is the largest eigenvalue of the upper left 2×2 submatrix of (16) and the dependence of its entries on μ_{ij} is given by (11) and (17). The permutation of coordinates $(x_1, x_2, x_3, x_4, x_5, x_6) \rightarrow (x_1, x_2, x_5, x_6, x_3, x_4)$ maps \mathcal{C}_L to \mathcal{C}_R . Hence, the condition for stability of \mathcal{C}_R are obtained from those for \mathcal{C}_L by the permutation of subscripts $(123456) \rightarrow (125634)$ of μ_{ij} in (11).

4.2 Trail-stability of \mathcal{C}_{LR} .

In this subsection we show that the omnicycle \mathcal{C}_{LR} can be trail-stable. Stability of the cycle depends on eigenvalues and eigenvectors of its 4×4 transition matrix. In order to be able to satisfy the conditions for stability we make two assumptions. Namely, we assume that the equation (1) has an additional symmetry

$$(x_1, x_2, x_3, x_4, x_5, x_6) \rightarrow (x_1, x_2, x_5, x_6, x_3, x_4) \quad (22)$$

and that the eigenvalues

$$\mu_{24}, \mu_{26}, \mu_{31}, \mu_{35}, \mu_{36}, \mu_{42}, \mu_{45}, \mu_{46}, \mu_{52}, \mu_{53}, \mu_{54}, \mu_{61}, \mu_{63}, \mu_{64} \quad (23)$$

are much smaller than other eigenvalues of the Jacobians $df(\xi_j)$. To prove existence of a system where \mathcal{C}_{LR} is trail-stable, we first prove that for vanishing eigenvalues (23) it is possible to identify other eigenvalues μ_{ij} such that the transition matrix satisfies conditions (i)-(iii) of theorem 2. The conditions are not strict and eigenvalues and eigenvectors of a matrix depend continuously on its entries, therefore (i)-(iii) remain true for sufficiently small eigenvalues in (23) as well. For a given set of eigenvalues of $df(\xi_j)$, a system that has such eigenvalues can be constructed by the procedure discussed in subsection 5.1.

Existence of the symmetry (22) implies that the omnicycle is comprised of two building blocks (see definition 2), therefore to study its trail-stability it is sufficient to consider stability of the return map $H_{23}^{out} \rightarrow H_{25}^{out}$ with respective transition matrix $M_{LR} = M_{125}M_{412}M_{341}M_{234}$. Let the bases in H_{12}^{out} , H_{34}^{out} and H_{41}^{out} be given by (13) and the basis in H_{25}^{out} be $\{\mathbf{e}_6, \mathbf{e}_1, \mathbf{e}_3, \mathbf{e}_4\}$. The basic transition matrices of the cycle \mathcal{C}_{LR} in a system with the symmetry (22) and vanishing eigenvalues (23) are

$$\begin{aligned} M_{412} &= \begin{pmatrix} b_{132} & 1 & 0 & 0 \\ b_{142} & 0 & 0 & 0 \\ b_{152} & 0 & 1 & 0 \\ b_{162} & 0 & 0 & 1 \end{pmatrix}, & M_{234} &:= \begin{pmatrix} 0 & 1 & 0 & 0 \\ b_{324} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\ M_{341} &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ b_{431} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, & M_{125} &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & b_{215} & 0 \\ 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \end{aligned} \quad (24)$$

where b_{ijk} are defined in (11).

Multiplication of the matrices implies that

$$M_{LR} = \begin{pmatrix} \tilde{a}_{11} & 0 & 0 & 1 \\ \tilde{a}_{21} & 0 & \tilde{a}_{23} & 0 \\ 0 & \tilde{a}_{32} & -1 & 0 \\ \tilde{a}_{41} & 0 & 0 & 0 \end{pmatrix}, \quad (25)$$

where

$$\tilde{a}_{11} = b_{162}b_{324}, \quad \tilde{a}_{21} = b_{152}b_{324}b_{213}, \quad \tilde{a}_{23} = b_{213}, \quad \tilde{a}_{32} = b_{413}, \quad \tilde{a}_{41} = b_{142}b_{324}. \quad (26)$$

Upon the permutation of bases in H_{25}^{out} : $\{\mathbf{e}_6, \mathbf{e}_1, \mathbf{e}_3, \mathbf{e}_4\} \rightarrow \{\mathbf{e}_6, \mathbf{e}_4, \mathbf{e}_1, \mathbf{e}_3\}$, matrix (25) takes the form

$$M_{LR} = \begin{pmatrix} \tilde{a}_{11} & 1 & 0 & 0 \\ \tilde{a}_{41} & 0 & 0 & 0 \\ \tilde{a}_{21} & 0 & 0 & \tilde{a}_{23} \\ 0 & 0 & \tilde{a}_{32} & \tilde{a}_{33} \end{pmatrix}. \quad (27)$$

The eigenvalues of this matrix are

$$\lambda_{1,2} = \frac{\tilde{a}_{11} \pm (\tilde{a}_{11}^2 + 4\tilde{a}_{11})^{1/2}}{2}, \quad \lambda_{3,4} = \frac{-1 \pm (1 + 4\tilde{a}_{23}\tilde{a}_{32})^{1/2}}{2}, \quad (28)$$

where the indices 1 and 3 correspond to positive signs in front of the square roots. Therefore, the conditions (i) and (ii) of theorem 2 for the matrix M_{LR} take the form

$$\lambda_1 > \max(1, |\lambda_4|). \quad (29)$$

From (27) the components of the eigenvector $\mathbf{v} = (v_1, v_2, v_3, v_4)$ of M_{LR} associated with λ_1 satisfy

$$v_2 = \frac{v_1 \tilde{a}_{41}}{\lambda_1}, \quad v_3 = \frac{v_4 \tilde{a}_{23}}{\lambda_1}, \quad v_4 = v_1 \tilde{a}_{31} \left(\lambda_1 + 1 - \frac{\tilde{a}_{32} \tilde{a}_{23}}{\lambda_1} \right)^{-1}.$$

Therefore, the condition (iii) of theorem 2 holds true whenever

$$(\lambda_1 + 1)\lambda_1 - \tilde{a}_{32}\tilde{a}_{23} > 0. \quad (30)$$

Finally, we show that conditions (29),(30) imply that the cycle \mathcal{C}_L is completely unstable. (Therefore, by the same arguments as applied to show trail-stability of \mathcal{C}_{LR} , it remains unstable for a slightly perturbed non-vanishing eigenvalues listed in (23).) The transition matrix (16) under the assumptions of symmetry (22) and for vanishing eigenvalues (23) takes the form

$$M_L = \begin{pmatrix} a_{11} & 0 & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 \\ 0 & a_{32} & 1 & 0 \\ a_{41} & 0 & 0 & 1 \end{pmatrix}, \quad (31)$$

where $a_{32} = b_{235}b_{413} < 0$. The components of the dominant eigenvector satisfy $v_3 = v_1 a_{32} / (\lambda_{\max} - 1)$. Hence, $\lambda_{\max} > 1$ implies that $v_1 v_3 < 0$, i.e., for the matrix M_L the conditions (ii) and (iii) of theorem 2 can not be satisfied simultaneously.

Remark 4 To show that the cycle \mathcal{C}_L can be stable in symmetric system (22) assume that the eigenvalues

$$\mu_{13}, \mu_{16}, \mu_{24}, \mu_{26}, \mu_{3j}, j = 1, 3, 5; \mu_{41}, \mu_{46} \quad (32)$$

are much smaller than the other ones. Under the assumption of vanishing eigenvalues (32) the matrix (16) takes the form

$$M_L = \begin{pmatrix} b_{324}b_{142} & 0 & 0 & 0 \\ 0 & b_{213}b_{431} & 0 & 0 \\ 0 & b_{451} - b_{431} & 1 & 0 \\ b_{324}b_{142} & 0 & 0 & 1 \end{pmatrix}. \quad (33)$$

The conditions (i)-(iii) are satisfied if

$$\max(b_{324}b_{142}, b_{213}b_{431}) > 1 \text{ and } b_{451} - b_{431} > 0.$$

These conditions are also satisfied for sufficiently small eigenvalues (32).

5 Numerical examples

In this section we study numerically behaviour of trajectories near the heteroclinic networks $\mathcal{X} = \mathcal{C}_L \cup \mathcal{C}_R$ shown in fig. 1. In subsection 5.1 we discuss construction of a \mathbb{Z}_2^6 -equivariant dynamical system, possessing such a network with prescribed eigenvalues of Jacobians $df(\xi_j)$, $1 \leq j \leq 6$. Then, using results of section 4, we give numerical examples of attracting \mathcal{C}_L and \mathcal{C}_{LR} . Our simulations indicate that when none of \mathcal{C}_L , \mathcal{C}_R or \mathcal{C}_{LR} is attracting, nearby trajectories nevertheless can be attracted to the network displaying regular or irregular switching between \mathcal{C}_L and \mathcal{C}_R .

5.1 Construction of the system

Consider the system

$$\dot{x}_i = x_i(-1 + \sum_{1 \leq j \leq 6} \beta_{ij}x_j^2), \quad \text{where } \beta_{ii} = 1, 0 \leq i \leq 6. \quad (34)$$

By construction (34) is \mathbb{Z}_2^6 -equivariant with the symmetry group generated by the inversions of Cartesian coordinates. Any coordinate axis, plane or a hyperplane is an invariant subspace of (34). It has equilibria $\xi_i = \pm 1$ on each of the coordinate axes, which are attractive in the radial direction.

The restriction of (34) to the coordinate plane $\langle \mathbf{e}_l, \mathbf{e}_k \rangle$ is

$$\dot{x}_l = x_l(-1 + x_l^2 + \beta_{lk}x_k^2), \quad \dot{x}_k = x_k(-1 + x_k^2 + \beta_{kl}x_l^2). \quad (35)$$

It is known (see, e.g., [23] and references therein) that if both ξ_l or ξ_k are stable in the radial direction, one of them is unstable in the orthogonal direction and the other equilibrium is stable, then there exists a robust heteroclinic trajectory connecting these equilibria.

The eigenvalues of $df(\xi_l)$ and $df(\xi_k)$ in the directions of \mathbf{e}_k and \mathbf{e}_k , respectively, are

$$\mu_{lk} = -1 + \beta_{kl}, \quad \mu_{kl} = -1 + \beta_{lk}. \quad (36)$$

Each of β_{ij} enters into only one expression for μ_{kl} , therefore we can design system (34) for any prescribed μ_{kl} compatibles with the connections shown in fig. 1. (Except for radial eigenvalues, which do not enter into the expressions for stability. These eigenvalues can be altered by modifying b_{ii} .)

5.2 Numerical results: symmetric system

Here we present results of numerical simulations (see fig. 2) of solutions to equations (34) in the case when the system has additional symmetry (22). In section 3 we derived conditions for stability of heteroclinic cycles \mathcal{C}_L and \mathcal{C}_R , and omnicycle \mathcal{C}_{LR} , which involve ratios $b_{ijk} = -\mu_{ij}/\mu_{jk}$. Therefore, we can choose μ_{lk} to make the cycles is stable or unstable. The values of μ_{lk} employed in simulations shown in the figure are given in table 1.

plate	eigenvalues
(a)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.2$ $\mu_{21} = -1, \mu_{23} = \mu_{25} = 1/1.4, \mu_{32} = \mu_{52} = -1.5, \mu_{43} = \mu_{65} = -1.1, \mu_{45} = \mu_{63} = -1.7$
(b)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.2$ $\mu_{21} = -1, \mu_{23} = \mu_{25} = 1/1.4, \mu_{32} = \mu_{52} = -1.5, \mu_{35} = \mu_{53} = -0.4, \mu_{43} = \mu_{65} = -1.1$
(c)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.2$ $\mu_{21} = -1, \mu_{23} = \mu_{25} = 1/1.4, \mu_{32} = \mu_{52} = -1.5, \mu_{35} = \mu_{53} = -0.4, \mu_{43} = \mu_{65} = -1.6$
(d)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.2$ $\mu_{21} = -1, \mu_{23} = \mu_{25} = 1/1.4, \mu_{32} = \mu_{52} = -1.5, \mu_{35} = \mu_{53} = -0.4, \mu_{43} = \mu_{65} = -1.4$
(e)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.2$ $\mu_{21} = -1, \mu_{23} = \mu_{25} = 1/1.4, \mu_{32} = \mu_{52} = -1, \mu_{15} = \mu_{13} = -0.4, \mu_{43} = \mu_{65} = -2$

Table 1: Eigenvalues of linearisations employed in numerical simulations shown in fig. 2. We do not list $\mu_{ii} = -1$ or eigenvalues that are equal to -0.01.

We start from stable \mathcal{C}_L and \mathcal{C}_{LR} (plates (a) and (b)) where the values of μ_{lk} given in table 1 are taken to satisfy conditions (18),(21) and (26),(28)-(30). For all other considered variants, where μ_{lk} are taken so that both \mathcal{C}_L and \mathcal{C}_{LR} are unstable, the trajectories staying near the network display regular or irregular behaviour: they can switch between \mathcal{C}_L and \mathcal{C}_R making a fixed number of turns (this number depends on parameters of (34)) around one cycle before switching to the other one, or this number can vary. Examples of such regular or irregular switching are shown in plates (c)-(e).

For $\tau \rightarrow \infty$ the trajectories are attracted by the heteroclinic network, which is indicated by exponentially increasing subsequent time intervals between visiting ξ_2 ¹ (i.e., time intervals

¹On the figures only the coordinates x_3 and x_5 are displayed: x_3 close to one indicates that after ξ_2 a trajectory goes to \mathcal{C}_L , while x_5 close to one corresponds to the trajectory going to \mathcal{C}_R .

needed to make a turn around \mathcal{C}_L or \mathcal{C}_R). On the plots the horizontal axis (time) is scales exponentially, implying that temporal behaviour is visually similar to periodic.

Remark 5 *In simulations the values of β_{lk} are taken such that the eigenvalues of transition matrices M_L and M_R satisfy $\lambda_L^{\max} > 1$ and $\lambda_R^{\max} > 1$. We can not prove that these inequalities imply fragmentary asymptotic stability of the network, however numerical results indicate that this might be the case, because for all performed runs trajectories were attracted by the network.*

5.3 Numerical results: non-symmetric system

In this section we present results of numerical integration of system (34) when additional symmetry (22) is not imposed. Typical behaviour of trajectories near the network is shown in fig.3 and the respective eigenvalues of $df(\xi_j)$ are given in table 2. The eigenvalues are chosen such that none of \mathcal{C}_L or \mathcal{C}_R is f.a.s.

plate	eigenvalues
(a)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.2, \mu_{21} = -1, \mu_{23} = 1/1.4$ $\mu_{25} = 1/1.5, \mu_{32} = \mu_{52} = -1.5, \mu_{43} = 1.8, \mu_{65} = -1.4, \mu_{35} = -0.4, \mu_{53} = -0.5$
(b)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.4, \mu_{21} = -1$ $\mu_{23} = \mu_{25} = 1/1.4, \mu_{32} = \mu_{52} = -1.5, \mu_{15} = -0.4, \mu_{43} = 2.9, \mu_{65} = -1.6$
(c)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.4$ $\mu_{21} = -1, \mu_{23} = \mu_{25} = 1/1.4, \mu_{32} = \mu_{52} = -1.5, \mu_{43} = 2.9, \mu_{65} = -1.6$
(d)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.2, \mu_{21} = -1, \mu_{23} = 1/1.4$ $\mu_{25} = 1/1.7, \mu_{32} = \mu_{52} = -1.5, \mu_{35} = \mu_{53} = -0.4, \mu_{43} = -1.6, \mu_{65} = -1.1$
(e)	$\mu_{12} = \mu_{34} = \mu_{41} = \mu_{56} = \mu_{61} = 1, \mu_{14} = \mu_{16} = -1.2, \mu_{21} = -1, \mu_{23} = 1/1.4$ $\mu_{25} = 1/1.5, \mu_{32} = \mu_{52} = -1.5, \mu_{35} = \mu_{53} = -0.4, \mu_{43} = -1.7, \mu_{65} = -1.5$

Table 2: Eigenvalues of linearisations employed in numerical simulations shown in fig. 3. We do not list $\mu_{ii} = -1$ or eigenvalues that are equal to -0.01.

A regular behaviour (examples are shown in plates (a) and (d)) can be described as a repeating pattern comprised of n_L iterations along \mathcal{C}_L followed by n_R iterations along \mathcal{C}_R . Apparently, this indicates that the respective omnicycle, which can be labelled as (n_L, n_R) , is f.a.s. Often we observe less regular behaviour (as shown in plates (b), (c) and (e)) which either indicate chaos or a long omnicycle. Due to the presence of numerical errors, it makes no sense to investigate which kind of behaviour, chaotic or periodic, takes place: what we see in simulations may be different from what really takes place in the system.

6 Conclusion

In this paper we studied behaviour of trajectories near a network in \mathbb{R}^6 comprised of two heteroclinic cycles. We derived condition for f.a.s. of these heteroclinic cycles, proved that




Figure 2: The dependence on time of x_3 (solid line) and x_5 (dashed line).




Figure 3: The dependence on time of x_3 (solid line) and x_5 (dashed line).

the network can be f.a.s. while both cycles are completely unstable and presented results of numerical simulations indicating that for a positive measure set of initial conditions the behaviour of nearby trajectories is apparently chaotic.

To prove stability of the network we introduce the notions of an omnicycle and trail-stability, which were implicitly used in earlier studies of behaviour of trajectories near a heteroclinic network. The definition of omnicycle is similar to the one of heteroclinic cycle, except that the equilibria and heteroclinic connections are not required to be distinct. Hence, the conditions for trail-stability of an omnicycle are identical to the condition of f.a.s. of a heteroclinic cycle, for type Z objects they were proven in [22].

To prove possibility of the existence of an attracting omnicycle we have assumed that some eigenvalues of linearisations are much smaller than the others, which allows us to calculate eigenvalues and eigenvectors of transition matrices. The assumption of smallness of some of the eigenvalues of linearisations can be useful in other studies of stability of cycles, omnicycles or heteroclinic networks in high-dimensional spaces, where direct calculation of eigenvalues is not possible due to large dimensions of transition matrices.

Acknowledgements

I am grateful to Alexander Lohse for a discussion of definition 7.

References

- [1] M.A.D. Aguiar and S.B.S.D. Castro. Chaotic switching in a two-person game. *Physica D* **239**, 1598–1609 (2010).
- [2] M.A.D. Aguiar, S.B.S.D. Castro and I.S. Labouriau. Dynamics near a heteroclinic network. *Nonlinearity* **18**, 391–414 (2005)
- [3] M.A.D. Aguiar, S.B.S.D. Castro and I.S. Labouriau. Simple Vector Fields with Complex Behavior. *Int. J. Bif. and Chaos* **16**, 369–381 (2006).
- [4] D. Armbruster, E. Stone and V. Kirk. Noisy heteroclinic networks. *Chaos* **13**, 71-79 (2003).
- [5] P. Ashwin, M. Field, A.M. Rucklidge and R. Sturman. Phase resetting effects for robust cycles between chaotic sets, *Chaos* **13**, 973–81, (2003).
- [6] P. Ashwin, G. Orosz and J. Borresen Heteroclinic switching in coupled oscillator networks: dynamics on odd graphs. in *M. Thiel et al. (eds.), Nonlinear Dynamics and Chaos: Advances and Perspectives, Understanding Complex Systems*. Springer (2010).
- [7] P.B. Ashwin, S.B.S.D. Castro and A. Lohse. Almost complete and equable heteroclinic networks, *Journal of Nonlinear Science* **30**, 1–22 (2020).

- [8] W. Brannath. Heteroclinic networks on the tetrahedron. *Nonlinearity* **7**, 1367–1384 (1994).
- [9] P.-L. Buono, M. Golubitsky and A. Palacios. Heteroclinic cycles in rings of coupled cells. *Physica D* **143**, 74–108 (2000).
- [10] S.B.S.D. Castro and A. Lohse Stability in simple heteroclinic networks in \mathbb{R}^4 , *Dynamical Systems*, **29** (4), 451–481 (2014).
- [11] S.B.S.D. Castro and A. Lohse Construction of heteroclinic networks in \mathbb{R}^4 , *Nonlinearity* **26**, 3677 (2016).
- [12] M. Dellnitz, M. Field, M. Golubitsky, J. Ma and A. Hohmann. Cycling chaos. *International journal of bifurcations and chaos* **5**, 1243–1247 (1995).
- [13] M. Golubitsky, I. Stewart and D. Schaeffer. *Singularities and Groups in Bifurcation Theory*. Springer (1988).
- [14] J. Hofbauer and J.W.-H. So. Multiple limit cycles for three-dimensional Lotka-Volterra equations. *Appl. Math. Lett.* **7**, 65–70 (1994).
- [15] J. Hofbauer and K. Sigmund *Evolutionary games and population dynamics*. CUP, Cambridge (1998).
- [16] V. Kirk, M. Silber. A competition between heteroclinic cycles. *Nonlinearity* **7**, 1605–1621 (1994).
- [17] V. Kirk, E. Lane, C.M. Postlethwaite, A.M. Rucklidge and M. Silber. A mechanism for switching near a heteroclinic network. *Dynamical Systems Int. J.* **25**, 323–349 (2010).
- [18] M. Krupa and I. Melbourne. Asymptotic stability of heteroclinic cycles in systems with symmetry. *Ergodic Theory Dyn. Syst.* **15**, 121 – 148 (1995).
- [19] M. Krupa and I. Melbourne. Asymptotic stability of heteroclinic cycles in systems with symmetry. II. *Proc. Roy. Soc. Edinburgh* **134A**, 1177 – 1197 (2004).
- [20] I. Labouriau and A.A.P. Rodrigues. Bifurcations from an attracting heteroclinic cycle under periodic forcing. *J. Differential Equations* **269**, 4137–4174 (2020).
- [21] I. Labouriau and A.A.P. Rodrigues. Periodic forcing of a heteroclinic network (2021), arXiv:2105.10757v1 [math.DS].
- [22] O.M. Podvigina. Stability and bifurcations of heteroclinic cycles of type Z. *Nonlinearity* **25**, 1887 – 1917 (2012).
- [23] O.M. Podvigina. Stability of rolls in rotating magnetoconvection in a layer with no-slip electrically insulating boundaries. *Phys. Rev. E* **81**, 056322 (2010).

- [24] O.M. Podvigina and P. Ashwin. On local attraction properties and a stability index for heteroclinic connections. *Nonlinearity* **24**, 887 – 929 (2011).
- [25] O. Podvigina and P. Chossat. Simple Heteroclinic Cycles in R^4 , *Nonlinearity*, **28**, 901–926 (2015).
- [26] O. Podvigina and P. Chossat. Asymptotic stability of pseudo-simple heteroclinic cycles in R^4 , *J. Nonlinear Sci.*, **27**, 343–375 (2017)
- [27] O. Podvigina and A. Lohse. Simple heteroclinic networks in R^4 , *Nonlinearity*, **32**, 3269–3293 (2019).
- [28] O. Podvigina, S.B.S.D. Castro and I.S. Labouriau. Stability of a heteroclinic network and its cycles: a case study from Boussinesq convection, *Dynamical Systems*, **34** (1), 157–193 (2019)
- [29] O. Podvigina, S.B.S.D. Castro and I.S. Labouriau. Asymptotic stability of robust heteroclinic networks. *Nonlinearity*, **33**, 1757–1788 (2020).
- [30] C.M. Postlethwaite and J.H.P. Dawes. Regular and irregular cycling near a heteroclinic network. *Nonlinearity* **18**, 1477 – 1509, (2005)
- [31] C.M. Postlethwaite and A.M. Rucklidge. Stability of cycling behaviour near a heteroclinic network model of Rock-Paper-Scissors-Lizard-Spock (2021) arXiv:2010.10666 [math.DS].
- [32] A.A.P. Rodrigues. Is there switching without suspended horseshoes? *Boletim da SPM* **74**, 61–79 (2016).
- [33] A.A.P. Rodrigues, I.S. Labouriau and M.A.D. Aguiar. Chaotic double cycling *Dynamical systems* **26**, 199–233 (2011).