

APPROXIMATING APPROXIMATE REASONING: FUZZY SETS AND THE ERSHOV HIERARCHY

NIKOLAY BAZHENOV, MANAT MUSTAFA, SERGEI OSPICHEV,
AND LUCA SAN MAURO

ABSTRACT. Computability theorists have introduced multiple hierarchies to measure the complexity of sets of natural numbers. The Kleene Hierarchy classifies sets according to the first-order complexity of their defining formulas. The Ershov Hierarchy classifies Δ_2^0 sets with respect to the number of mistakes that are needed to approximate them. Biacino and Gerla extended the Kleene Hierarchy to the realm of fuzzy sets, whose membership functions range in a complete lattice L (e.g., the real interval $[0; 1]_{\mathbb{R}}$). In this paper, we combine the Ershov Hierarchy and fuzzy set theory, by introducing and investigating the Fuzzy Ershov Hierarchy. In particular, we focus on the fuzzy n -c.e. sets which form the finite levels of this hierarchy. Intuitively, a fuzzy set is n -c.e. if its membership function can be approximated by changing monotonicity at most $n-1$ times. We prove that the Fuzzy Ershov Hierarchy does not collapse; that, in analogy with the classical case, each fuzzy n -c.e. set can be represented as a Boolean combination of fuzzy c.e. sets; but that, contrary to the classical case, the Fuzzy Ershov Hierarchy does not exhaust the class of all Δ_2^0 fuzzy sets.

1. INTRODUCTION

Crisp properties on a given domain \mathcal{D} — i.e., properties whose membership functions range in the set $\{0, 1\}$ — can be naturally identified with subsets of \mathcal{D} . By adopting this perspective, one may regard classical computability theory as the study of the complexity of crisp properties on the set ω of the natural numbers: e.g., “being even” and “being the code of a Turing machine which halts on a blank tape” are examples of, respectively, a decidable crisp property and an undecidable one.

Computability theorists have introduced multiple hierarchies to measure the complexity of crisp objects. Two such hierarchies will be relevant for the present paper. The Kleene Hierarchy classifies subsets of ω according to the first-order complexity of their defining formulas within arithmetic. The Ershov Hierarchy concentrates on an important initial segment of the Kleene Hierarchy, that of Δ_2^0 sets (which coincide with the sets that are computable in the limit), by classifying such sets with respect to the number of mistakes that are needed to approximate them.

Fuzzy sets, introduced by Zadeh [18] and later developed into a broad area of research, allow to mathematically study *graded properties*, such as those properties with blurry boundaries, and to extend the scope of logic to *approximate reasoning*.

Key words and phrases. Fuzzy set, computability theory, n -computably enumerable set, Ershov hierarchy.

It is natural to ask how to introduce computability theory within fuzzy mathematics. A first approach is to define fuzzy algorithms (as in, e.g., [19, 14, 17, 1]), and then rebuild computability theory by permitting fuzzy computations. A parallel approach is to maintain ordinary Turing machines and just adopt them to calibrate the complexity of fuzzy sets. After all, well-established computability-theoretic hierarchies could be extended to the realm of fuzzy objects. This is the case for the Kleene Hierarchy, which has been extended to fuzzy sets by Biacino and Gerla [2], see also [7, 8, 9].

In this paper, we introduce and investigate the Fuzzy Ershov Hierarchy. That is, we focus on the complexity of approximating fuzzy objects, which belong to the class Δ_2^0 . The key idea for evaluating this complexity is that of a *mind change*, which is borrowed from Ershov [4, 5, 6]. In the classical setting, an approximation to a set A changes its mind on a given input x by switching its guess on whether x belongs to A or not. Moving to fuzzy sets, mind changes will be formalized by changes in the monotonicity of approximating functions. We will prove that, by allowing more and more mind changes, we will be able to capture larger and larger sub-classes of Δ_2^0 fuzzy sets. In particular, it will follow that there are fuzzy sets which cannot be approximated only from above or below, but they require approximations which oscillate “up and down” on the membership degree of x , for some inputs x .

The paper is arranged as follows. In Section 2, we recall preliminaries concerning fuzzy sets, effective reals, and the Ershov Hierarchy. In Section 3, we introduce the Fuzzy Ershov Hierarchy, and we prove the main results of the paper. First, the hierarchy does not collapse (Proposition 3.2). Second, in analogy with the classical case, sets lying at the so-called finite levels of the Fuzzy Ershov Hierarchy can be represented as Boolean combinations of fuzzy sets belonging to the first level, i.e. fuzzy c.e. sets (Theorem 3.1). Third, contrary to the classical case, the Fuzzy Ershov Hierarchy does not exhaust the class of all Δ_2^0 fuzzy sets (Proposition 3.3). In the last section, we conclude by briefly discussing two natural ways of broadening the Fuzzy Ershov Hierarchy.

2. PRELIMINARIES

We assume that the reader is familiar with the basic notions of computability theory. For the background, we refer to the monographs [13, 15]. The preliminaries on fuzzy sets mainly follow [7].

As usual, one fixes an effective bijection $\nu: \mathbb{Q} \rightarrow \omega$. This convention allows to transfer familiar computability-theoretic notions to the rationals: for example, a crisp set $X \subseteq \mathbb{Q}$ is computable iff its image $\nu(X)$ is a computable subset of ω .

Note that, as is custom in computability theory, this paper uses the term *computably enumerable* (or *c.e.*) in place of *recursively enumerable*. For a set X , by $|X|$ we denote the cardinality of X .

2.1. Fuzzy Subsets. Let \mathcal{L} be a complete lattice. A *fuzzy subset* (or an \mathcal{L} -subset) of ω is an arbitrary function $A: \omega \rightarrow \mathcal{L}$. In this paper, for the sake of simplicity, we consider the case when \mathcal{L} is equal to the real interval $[0; 1]_{\mathbb{R}}$. A fuzzy subset A is *crisp* if $A(x) \in \{0, 1\}$ for all $x \in \omega$.

As mentioned in the introduction, a fundamental tool for classifying the complexity of crisp subsets of ω is provided by the Kleene Arithmetical Hierarchy [10] (see, e.g., Chapter 4 in [15] for a detailed discussion). Biacino and Gerla [2] extended the

Kleene Hierarchy to fuzzy subsets. In our paper, we work only with fuzzy subsets belonging to the levels Σ_1^0 , Π_1^0 , and Δ_2^0 of the Kleene Hierarchy. Hence, we give formal definitions only for these levels. For more details, the reader is referred to [2] and § 11.5 in [7]. By $[0; 1]_{\mathbb{Q}}$ we denote the set of all rational numbers q such that $0 \leq q \leq 1$.

Definition 2.1 ([2], see also § 11.2 in [7]). A fuzzy set A is *computably enumerable* (or belongs to the class Σ_1^0) if there is a computable function $f: \omega \times \omega \rightarrow [0, 1]_{\mathbb{Q}}$ such that, for all $x \in \omega$, we have:

- (a) $\lim_{s \rightarrow \infty} f(x, s) = A(x)$;
- (b) $(\forall s)(f(x, s+1) \geq f(x, s))$.

We say that such function f is a Σ_1^0 -*approximation* of the fuzzy set A .

Note that without loss of generality, one can always assume that in the definition above, $f(x, 0)$ equals 0. Hence, fuzzy c.e. sets may intuitively be regarded as fuzzy sets which can be approximated “from below”, in the sense that approximations to fuzzy c.e. sets can only increase over time.

If A and B are fuzzy sets, then one can define set-theoretic operations on them:

- Union: $(A \cup B)(x) = \max\{A(x), B(x)\}$.
- Intersection: $(A \cap B)(x) = \min\{A(x), B(x)\}$.
- Complement: $\overline{A}(x) = 1 - A(x)$.

A fuzzy set A is *co-computably enumerable* (or belongs to the class Π_1^0) if its complement \overline{A} is c.e. Equivalently (see Theorem 5.2 in [7, Chap. 11]), A is co-c.e. if and only if there is a computable function $f: \omega^2 \rightarrow [0; 1]_{\mathbb{Q}}$ such that, for all $x \in \omega$, we have:

- (a) $\lim_s f(x, s) = A(x)$;
- (b') $(\forall s)(f(x, s+1) \leq f(x, s))$.

In the Π_1^0 case, we may assume that $f(x, 0) = 1$, for all x . So, fuzzy co-c.e. sets may be regarded as fuzzy sets which can be approximated “from above”.

Finally, the main object of study of this paper are Δ_2^0 fuzzy sets. A fuzzy set A belongs to the class Δ_2^0 if A lies in both classes Σ_2^0 and Π_2^0 of the Kleene Hierarchy. In this paper, we adopt the following equivalent definition (see Proposition 5.4 in [7, Chap. 11]).

Definition 2.2. A fuzzy set A is Δ_2^0 if and only if there is a computable function $f: \omega^2 \rightarrow [0; 1]_{\mathbb{Q}}$ such that $\lim_s f(x, s) = A(x)$, for all $x \in \omega$. We call such function $f(x, s)$ a Δ_2^0 -*approximation* of the fuzzy set A .

2.2. Effective Reals. Here we briefly discuss some simple results which connect fuzzy subsets of ω with effectively approximable reals. We refer to Chapter 5 in [3] for the detailed background.

We consider reals $\alpha \in [0; 1]_{\mathbb{R}}$. A real α is *left-c.e.* if the set $\{q \in \mathbb{Q} : q < \alpha\}$ is c.e. A real α is *right-c.e.* if the set $\{q \in \mathbb{Q} : q > \alpha\}$ is c.e. By working with the definitions, it is not hard to prove the following result.

Proposition 2.1. *Let A be a fuzzy subset of ω .*

- (1) *A is c.e. if and only if the reals $A(k)$, $k \in \omega$, are uniformly left-c.e., i.e. the set $\{(k, q) \in \omega \times \mathbb{Q} : q < \alpha_k\}$ is c.e.*
- (2) *A is co-c.e. if and only if the reals $A(k)$, $k \in \omega$, are uniformly right-c.e., i.e. the set $\{(k, q) \in \omega \times \mathbb{Q} : q > \alpha_k\}$ is c.e.*

A real α is Δ_2^0 if there is a computable sequence $(q_s)_{s \in \omega}$ of rationals such that $\alpha = \lim_s q_s$ (see, e.g., Theorem 5.1.3 in [3]). We also observe the following, which is an immediate consequence of the definitions:

Proposition 2.2. *A fuzzy set A is Δ_2^0 if and only if the reals $A(k)$, $k \in \omega$, are uniformly Δ_2^0 , i.e., there is a computable sequence $(q_{k,s})_{k,s \in \omega}$ of rationals such that $A(k) = \lim_s q_{k,s}$, for all k .*

2.3. The Classical Ershov Hierarchy. We give few preliminaries on the Ershov Hierarchy [4, 5, 6]; to distinguish it from the fuzzy analogue introduced below, we refer to this hierarchy as the Classical Ershov Hierarchy. For the sake of simplicity, here we discuss only the finite levels of the hierarchy (these finite levels are also called *Difference Hierarchy* in the literature). In this section, all subsets of ω are crisp.

Definition 2.3. Let $n \geq 1$. A set $A \subseteq \omega$ is *n-computably enumerable* (or *n-c.e.*, or belongs to the class Σ_n^{-1}) if there is a computable function $f: \omega \times \omega \rightarrow \{0, 1\}$ such that for all $x \in \omega$, we have

- $\lim_s f(x, s) = A(x)$;
- $f(x, 0) = 0$;
- $|\{s : f(x, s) \neq f(x, s+1)\}| \leq n$.

A set A is *co-n-computably enumerable* (or *co-n-c.e.*, or belongs to the class Π_n^{-1}) if its complement \bar{A} is *n-c.e.*

Historically, the notion of *n-c.e.* sets was introduced by Putnam [12] and Gold [11]. Note that

$$\Sigma_n^{-1} \cup \Pi_n^{-1} \subseteq \Sigma_{n+1}^{-1} \cap \Pi_{n+1}^{-1}.$$

Ershov [4] proved that for each $n \geq 1$, there exists an *n-c.e.* set S_n such that every *n-c.e.* set A is many-one reducible to S_n . In addition, S_n does not belong to Π_n^{-1} . In particular, this implies that the Classical Ershov Hierarchy does not collapse.

Sets from the class Σ_n^{-1} can be represented as Boolean combinations of *c.e.* sets:

Theorem 2.1 ([4]). *Let k be a natural number. A set $A \subseteq \omega$ is $(2k+1)$ -c.e. if and only if there are c.e. sets $W_1, W_2, \dots, W_{2k-1}, W_{2k}, W_{2k+1}$ such that*

$$A = (W_1 \setminus W_2) \cup (W_3 \setminus W_4) \cup \dots \cup (W_{2k-1} \setminus W_{2k}) \cup W_{2k+1}.$$

A set A is $(2k+2)$ -c.e. if and only if there are c.e. sets $W_1, W_2, \dots, W_{2k+1}, W_{2k+2}$ such that

$$A = (W_1 \setminus W_2) \cup (W_3 \setminus W_4) \cup \dots \cup (W_{2k-1} \setminus W_{2k}) \cup (W_{2k+1} \setminus W_{2k+2}).$$

We refer the reader to the survey [16] for more details on the Classical Ershov Hierarchy.

3. FUZZY ERSHOV HIERARCHY

In this section, we extend the classical Difference Hierarchy to the class of fuzzy subsets of ω (see Definition 3.2 below). We establish some initial properties of this hierarchy: the hierarchy does not collapse (Subsection 3.1); it is connected to the Boolean combinations of fuzzy *c.e.* sets (Subsection 3.2); the introduced levels of the hierarchy do not exhaust all fuzzy Δ_2^0 sets (Subsection 3.3).

We begin by illustrating the intuition behind Definition 3.2 with the following example. A fuzzy Δ_2^0 set A is called *3-computably enumerable* if it possesses a

Δ_2^0 -approximation $f(x, s)$, which “changes its mind” *at most two times*. So, for an element $x \in \omega$, the worst case behavior looks like this:

- First, our approximation (non-strictly) increases — i.e. there is a stage s_1 such that $f(x, s) \leq f(x, s+1)$, for all $s < s_1$.
- Second, the approximation starts to decrease until some stage $s_2 > s_1$: $f(x, s_1) > f(x, s_1+1)$ and $f(x, s) \geq f(x, s+1)$, for $s_1 < s < s_2$.
- Then the final change of mind happens: the approximation will forever increase — $f(x, s_2) < f(x, s_2+1)$ and $f(x, s) \leq f(x, s+1)$ for all $s > s_2$.

In order to make this idea formal, we introduce *mind change functions*, which “track down” the described mind changes.

Definition 3.1. Let f be a total function from $\omega \times \omega$ to $[0; 1]_{\mathbb{Q}}$. Its Σ -*mind change function* $m_{\Sigma}^f: \omega \times \omega \rightarrow \{-1, 1\}$ is defined as follows.

- (1) $m_{\Sigma}^f(x, 0) = 1$.
- (2) Suppose that $m_{\Sigma}^f(x, s) = 1$.
 - If $f(x, s) \leq f(x, s+1)$, then $m_{\Sigma}^f(x, s+1) = 1$.
 - If $f(x, s) > f(x, s+1)$, then $m_{\Sigma}^f(x, s+1) = -1$.
- (3) Suppose that $m_{\Sigma}^f(x, s) = -1$.
 - If $f(x, s) \geq f(x, s+1)$, then $m_{\Sigma}^f(x, s+1) = -1$.
 - If $f(x, s) < f(x, s+1)$, then $m_{\Sigma}^f(x, s+1) = 1$.

The Π -*mind change function* $m_{\Pi}^f(x, s)$ is defined similarly to m_{Σ}^f , with the following key modification: we put $m_{\Pi}^f(x, 0) = -1$.

Notice the following: if a function f is computable, then both m_{Σ}^f and m_{Π}^f are also computable. Now we are ready to give the main definition.

Definition 3.2. Let n be a non-zero natural number. A fuzzy set A is *n-computably enumerable* if there is a computable function $f: \omega \times \omega \rightarrow [0, 1]_{\mathbb{Q}}$ such that for all $x \in \omega$, we have:

- $\lim_s f(x, s) = A(x)$;
- $f(x, 0) = 0$;
- $|\{s \in \omega : m_{\Sigma}^f(x, s+1) \neq m_{\Sigma}^f(x, s)\}| \leq n - 1$.

A fuzzy set A is *co-n-computably enumerable* if its complement \bar{A} is *n-c.e.*

Note that 1-c.e. fuzzy sets are precisely the c.e. sets from Definition 2.1. In addition, the following fact is immediate.

Proposition 3.1. *Let $n \geq 1$. A fuzzy set A is co-n-c.e. if and only if there is a computable function $f: \omega \times \omega \rightarrow [0, 1]_{\mathbb{Q}}$ such that for all $x \in \omega$, we have:*

- $\lim_s f(x, s) = A(x)$;
- $f(x, 0) = 1$;
- $|\{s \in \omega : m_{\Pi}^f(x, s+1) \neq m_{\Pi}^f(x, s)\}| \leq n - 1$.

3.1. The Hierarchy Does Not Collapse. In order to show the non-collapse of the hierarchy, it is sufficient to prove the following:

Proposition 3.2. *Let A be a crisp subset of ω . Then A is n-c.e. in the Classical Ershov Hierarchy if and only if A is n-c.e. in the Fuzzy Ershov Hierarchy. A similar fact is true for co-n-c.e. sets.*

Indeed, since the Classical Difference Hierarchy does not collapse, Proposition 3.2 implies that our hierarchy is also non-collapsing.

of Proposition 3.2. (\Rightarrow). Suppose that A is n -c.e. in the classical sense. We fix a computable function $f: \omega^2 \rightarrow \{0, 1\}$ satisfying the conditions from Definition 2.3. It is clear that $f(x, s)$ is a Δ_2^0 -approximation of A , treated as a fuzzy Δ_2^0 set.

For an element $x \in \omega$, consider all stages $s_1 < s_2 < \dots < s_k$ (note that $k \leq n$) such that $f(x, s_i) \neq f(x, s_i + 1)$. A straightforward analysis shows the following:

- If $s \leq s_1$, then $f(x, s) = 0$ and $m_\Sigma^f(x, s) = 1$.
- If $s_{2\ell+1} + 1 \leq s \leq s_{2\ell+2}$, then $f(x, s) = 1$ and $m_\Sigma^f(x, s) = 1$.
- If $s_{2\ell+2} + 1 \leq s \leq s_{2\ell+3}$, then $f(x, s) = 0$ and $m_\Sigma^f(x, s) = -1$.

This implies that $|\{s : m_\Sigma^f(x, s+1) \neq m_\Sigma^f(x, s)\}| \leq k - 1 \leq n - 1$. We deduce that the approximation f witnesses that the set A is fuzzy n -c.e.

(\Leftarrow). Suppose that a crisp A is fuzzy n -c.e. Fix a Δ_2^0 -approximation $f: \omega^2 \rightarrow [0; 1]_{\mathbb{Q}}$ satisfying Definition 3.2. We define a new approximation

$$g(x, s) = \begin{cases} 1, & \text{if } f(x, s) > 1/2, \\ 0, & \text{if } f(x, s) \leq 1/2. \end{cases}$$

Since A is crisp, it is clear that $A(x) = \lim_s g(x, s)$. Notice that $g(x, 0) = f(x, 0) = 0$.

For an element $x \in \omega$, consider all stages $s_1 < s_2 < \dots < s_k$ such that $g(x, s_i) \neq g(x, s_i + 1)$. For $i \leq k$, one can show the following:

- If $i = 2\ell + 1$, then $f(x, s_i) \leq 1/2$, $f(x, s_i + 1) > 1/2$, and $m_\Sigma^f(x, s_i + 1) = 1$.
- If $i = 2\ell + 2$, then $f(x, s_i) > 1/2$, $f(x, s_i + 1) \leq 1/2$, and $m_\Sigma^f(x, s_i + 1) = -1$.

In turn, this implies $k - 1 \leq |\{s : m_\Sigma^f(x, s+1) \neq m_\Sigma^f(x, s)\}| \leq n - 1$. Hence, $k \leq n$, and the function $g(x, s)$ witnesses that the set A is n -c.e. in the classical sense. \square \square

3.2. Boolean Combinations of Fuzzy C.E. Sets. We show that similarly to the Classical Ershov Hierarchy (Theorem 2.1), n -c.e. fuzzy sets admit natural presentations via Boolean combinations of c.e. sets.

Theorem 3.1. *Let $n \in \{2k + 1, 2k + 2\}$. A fuzzy set C is n -c.e. if and only if there are fuzzy c.e. sets $A_1, B_1, A_2, B_2, \dots, A_{k+1}, B_{k+1}$ such that:*

- $C = (A_1 \cap \overline{B_1}) \cup (A_2 \cap \overline{B_2}) \cup \dots \cup (A_{k+1} \cap \overline{B_{k+1}})$;
- if $n = 2k + 1$, then $B_{k+1} = \emptyset$.

Proof. (\Rightarrow). Let $f(x, s)$ be a Δ_2^0 -approximation which witnesses the fact that C is n -c.e. We define the desired fuzzy c.e. sets A_i and B_i via their Σ_1^0 -approximations h_{A_i} and h_{B_i} (in the sense of Definition 2.1), respectively.

The intuition behind these c.e. sets is as follows. For an element $x \in \omega$, we split ω into disjoint intervals: $[0; a_0)$, $[a_0; b_0)$, $[b_0; a_1)$, $[a_1; b_1)$, etc. Our function $f(x, \cdot)$ (non-strictly) increases on the intervals $[0; a_0)$, $[b_0; a_1)$, $[b_1; a_2)$, etc. The function decreases on the rest of the intervals.

- The approximation h_{A_1} of the set A_1 looks like this: it copies $f(x, \cdot)$ on the interval $[0; a_0)$, and then stabilizes, i.e. $h_{A_1}(x, s) = h_{A_1}(x, a_0 - 1)$ for all $s \geq a_0$.

- The function $h_{B_1}(x, \cdot)$ equals zero on $[0; a_0)$. Then it copies $1 - f(x, \cdot)$ on the interval $[a_0; b_0)$. After that, $h_{B_1}(x, \cdot)$ equals *one*.
- The function h_{A_2} equals zero on $[0; b_0)$. Then it copies $f(x, \cdot)$ on the interval $[b_0; a_1)$; after that h_{A_2} stabilizes. Et cetera.

Formally speaking, for a non-zero $i \leq k + 1$, we define:

$$h_{A_i}(x, s) = \begin{cases} 0, & \text{if } |\{t \leq s : m_{\Sigma}^f(x, t+1) \neq m_{\Sigma}^f(x, t)\}| < 2i - 2, \\ f(x, s), & \text{if } |\{t \leq s : m_{\Sigma}^f(x, t+1) \neq m_{\Sigma}^f(x, t)\}| = 2i - 2, \\ h_{A_i}(x, s-1), & \text{otherwise;} \end{cases}$$

$$h_{B_i}(x, s) = \begin{cases} 0, & \text{if } |\{t \leq s : m_{\Sigma}^f(x, t+1) \neq m_{\Sigma}^f(x, t)\}| < 2i - 1, \\ 1 - f(x, s), & \text{if } |\{t \leq s : m_{\Sigma}^f(x, t+1) \neq m_{\Sigma}^f(x, t)\}| = 2i - 1, \\ 1, & \text{otherwise.} \end{cases}$$

It is not hard to see that these approximations induce fuzzy c.e. sets. In addition, if $n = 2k + 1$, then $B_{k+1}(x) = 0$ for all x .

Let D be the fuzzy set $(A_1 \cap \overline{B_1}) \cup \dots \cup (A_{k+1} \cap \overline{B_{k+1}})$. We consider its natural Δ_2^0 -approximation

$$(1) \quad h_D(x, s) = \max\{\min\{h_{A_i}(x, s), 1 - h_{B_i}(x, s)\} : 1 \leq i \leq k + 1\}.$$

Consider the value $v^* = |\{t \in \omega : m_{\Sigma}^f(x, t+1) \neq m_{\Sigma}^f(x, t)\}|$.

If $v^* = 2i - 2$, then there is a stage s^* such that for all $s \geq s^*$, we have $h_D(x, s) = h_{A_i}(x, s) = f(x, s)$. This implies that $D(x) = C(x)$.

If $v^* = 2i - 1$, then consider the highest index s_0 such that $h_{A_i}(x, s_0) = f(x, s_0)$. Then for every $s \geq s_0 + 1$, we have $h_{A_i}(x, s) = f(x, s_0)$, $h_{B_i}(x, s) = 1 - f(x, s)$, and

$$h_D(x, s) = \min(h_{A_i}(x, s), 1 - h_{B_i}(x, s)) = \min(f(x, s_0), f(x, s)) = f(x, s).$$

Again, $D(x) = C(x)$. We deduce that the fuzzy sets C and D are equal.

(\Leftarrow). Let D be a fuzzy Δ_2^0 set defined via the approximation h_D from (1). We prove that *this* approximation h_D witnesses the fact that D is n -c.e.

First, we note the following easy observation (it follows from computable enumerability of fuzzy sets A_i and B_i):

- (*) If $1 - h_{B_i}(x, s_0) < h_{A_i}(x, s_0)$ for some s_0 , then we have $1 - h_{B_i}(x, s) < h_{A_i}(x, s)$ for all $s \geq s_0$.

An informal intuition concerning further proof is as follows. Every (approximation of the) real $(A_i \cap \overline{B_i})(x)$ can be treated as a ‘‘hill’’: first we go up, copying the function $h_{A_i}(x, \cdot)$. When we see the inequality $1 - h_{B_i}(x, s_0) < h_{A_i}(x, s_0)$, we can only go down. Coming back to the whole picture of h_D : whenever the mind-change function $m_{\Sigma}^{h_D}(x, \cdot)$ changes from $+1$ to -1 , it happens because we encountered the *top* of one of the ‘‘hills’’.

At a stage s , consider the following sets: $X_s = \{i : 1 - h_{B_i}(x, s) < h_{A_i}(x, s)\}$ and $Y_s = \{1, 2, \dots, k + 1\} \setminus X_s$. Observation (*) implies that $X_s \subseteq X_{s+1}$ for every s . In addition, $X_0 = \emptyset$.

It is not hard to deduce the following equation:

$$h_D(x, s) = \max\{\max\{1 - h_{B_i}(x, s) : i \in X_s\}, \max\{h_{A_i}(x, s) : i \in Y_s\}\}.$$

Note that for a fixed non-empty set Z , the function $\max\{1 - h_{B_i}(x, s) : i \in Z\}$ is non-increasing, and $\max\{h_{A_i}(x, s) : i \in Z\}$ is non-decreasing.

Suppose that $m_{\Sigma}^{h_D}(x, s) = 1$ and $m_{\Sigma}^{h_D}(x, s+1) = -1$. Choose the greatest $s' < s$ such that either $s' = 0$, or $s' > 0$ and $m_{\Sigma}^{h_D}(x, s') = -1$. Towards a contradiction, assume that $X_{s+1} = X_{s'}$.

Then on one hand, we have

$$h_D(x, s) = \max\{1 - h_{B_i}(x, s) : i \in X_{s'}\} > \max\{h_{A_i}(x, s) : i \in Y_{s'}\}.$$

Indeed, if $h_D(x, s)$ equals $\max\{h_{A_i}(x, s) : i \in Y_{s'}\}$, then we would have

$$h_D(x, s+1) = \max\{h_{A_i}(x, s+1) : i \in Y_{s'}\} \geq h_D(x, s),$$

which contradicts the fact that $m_{\Sigma}^{h_D}(x, s+1) = -1$.

On the other hand, every t such that $s' < t \leq s$ satisfies

$$h_D(x, t) = \max\{h_{A_i}(x, t) : i \in Y_{s'}\}.$$

We obtain a contradiction. Therefore, $X_{s+1} \neq X_{s'}$.

We deduce that for each stage s with $m_{\Sigma}^{h_D}(x, s) = 1$ and $m_{\Sigma}^{h_D}(x, s+1) = -1$, at least one new element is added to the growing set $X = \bigcup_{t \in \omega} X_t$.

Suppose that $n = 2k + 2$. Then one can show that $|X| \leq k + 1$. We notice the following: if $|X|$ is less than $k + 1$, then the number of monotonicity breaks (of the function $m_{\Sigma}^{h_D}(x, \cdot)$) will be strictly less than the corresponding number for the case $|X| = k + 1$. Hence, one can consider only the case when $|X| = k + 1$.

If $|X| = k + 1$, then there is a stage s^* such that for all $s \geq s^*$, we have $h_D(x, s) = \max\{1 - h_{B_i}(x, s) : i \in X\}$, and this function can only decrease. A not difficult combinatorial argument shows that $|\{s \in \omega : m_{\Sigma}^{h_D}(x, s+1) \neq m_{\Sigma}^{h_D}(x, s)\}| \leq 2k + 1$.

If $n = 2k + 1$, then $|X| \leq k$. An argument similar to the one above shows that one can consider only the case when $|X|$ equals k .

If $|X| = k$, then there is a stage s^* such that for $s \geq s^*$, we have $h_D(x, s) = \max\{\max\{1 - h_{B_i}(x, s) : i \in X\}, h_{A_{k+1}}(x, s)\}$. One can show that in this case, $|\{s : m_{\Sigma}^{h_D}(x, s+1) \neq m_{\Sigma}^{h_D}(x, s)\}| \leq 2k$. Theorem 3.1 is proved. \square \square

Corollary 3.1. *Every finite Boolean combination of fuzzy c.e. sets is an n -c.e. set, for some $n \geq 1$.*

3.3. The Introduced Hierarchy Is Not Enough. Here we show that the introduced levels of the Fuzzy Ershov Hierarchy do not exhaust the class of all Δ_2^0 fuzzy subsets of ω .

Proposition 3.3. *There exists a Δ_2^0 fuzzy set A such that for any Δ_2^0 -approximation $f(x, s)$ of A , the sequence $(m_{\Sigma}^f(0, s))_{s \in \omega}$ diverges when s tends to infinity. In particular, A is not n -c.e., for all $n \geq 1$.*

Proof. Choose an arbitrary Δ_2^0 real α , which is not left-c.e. and not right-c.e. (see, e.g., Theorem 5.1.10 in [3] for an example of such real). The desired fuzzy set A is defined as follows: put $A(k) = \alpha$, for all $k \in \omega$. Since α is Δ_2^0 , Proposition 2.2 implies that the set A is Δ_2^0 .

Towards a contradiction, assume that $f(x, s)$ is a Δ_2^0 -approximation of A such that the sequence $(m_{\Sigma}^f(0, s))_{s \in \omega}$ converges. There are two possible cases.

Case 1. Suppose that $\lim_s m_{\Sigma}^f(0, s) = 1$. Then choose a stage s^* such that $m_{\Sigma}^f(0, s) = 1$ for all $s \geq s^*$. It is not hard to show that the set $\{q \in \mathbb{Q} : q < \alpha\}$ is equal to $\{q : (\exists s \geq s^*)[q < f(0, s)]\}$, and hence, this set is c.e. Then the real α is left-c.e., which gives a contradiction.

Case 2. Otherwise, $\lim_s m_\Sigma^f(0, s) = -1$. Then a similar argument shows that the real α is right-c.e. — again, a contradiction.

We deduce that our fuzzy set A has all desired properties. \square \square

4. CONCLUDING REMARKS: BROADENING THE FUZZY ERSHOV HIERARCHY

By way of conclusion, we briefly discuss two natural options for, first, *refining* the Fuzzy Ershov Hierarchy, and, secondly, *extending* the finite levels of the hierarchy. We leave many formal details to future work. In particular, even though we obtained a number of results concerning the new hierarchies discussed below, due to space constraints, we defer these results to an extended version of the present paper.

4.1. Counting Updates. We say that a Δ_2^0 -approximation of a fuzzy set A “has an update” if $f(x, s+1) \neq f(x, s)$, for some $x, s \in \omega$. Observe that our notion of mind change, as in Definition 3.2, keeps track only of those updates which determine a change of monotonicity in the approximating function: e.g., if $f(x, s)$ is a Δ_2^0 -approximation of a fuzzy set A , $m_\Sigma^f(x, s) = 1$, and $f(x, s+1) > f(x, s)$, then $m_\Sigma^f(x, s+1)$ remains equal to 1. So, one may explore what happens if one keeps track of *all* updates for a given Δ_2^0 -approximation. To further motivate such approach, consider the example by Harkleroad [8]:

Example 4.1. As usual, K denotes the Halting problem. Define

$$H(x) = \begin{cases} 1, & \text{if } x \in K, \\ 1/2, & \text{otherwise.} \end{cases}$$

It is easy to see that H is a fuzzy c.e. set. But note that, for any c.e. approximation h of H (recall that one assumes $h(x, 0) = 0$ for all x), there must be an infinite crisp set $Z \subseteq \omega$ such that h requires at least *two* updates to approximate each $x \in Z$ (as otherwise, $K = \{x : (\exists s)(h(x, s) = 1)\}$ would be computable).

So, to distinguish H from fuzzy c.e. sets which can be approximated with at most one update, we propose the following new hierarchy:

Definition 4.1. A fuzzy set A is $[n]_1$ -c.e. if there is a computable function $f: \omega \times \omega \rightarrow [0, 1]_{\mathbb{Q}}$ such that for all $x \in \omega$, we have:

- $f(x, 0) = 0$ and $\lim_s f(x, s) = A(x)$;
- $(\forall s)(f(x, s+1) \geq f(x, s))$;
- $|\{s : f(x, s+1) \neq f(x, s)\}| \leq n$.

For simplicity, the definition above is limited to the first level of the Fuzzy Ershov Hierarchy. But clearly, one could similarly stratify each class of fuzzy n -c.e. sets as follows. Intuitively, a set A is fuzzy $[n_1, \dots, n_m]_m$ -c.e. if there is a Δ_2^0 -approximation f to A that, for each x , can go up at most n_1 times, and then down at most n_2 times, etc. — for m -many ups and downs.

In a future work, we plan to carefully study all such refinements of the Fuzzy Ershov Hierarchy, together with their interplay.

4.2. Going Transfinite. In this paper, we talked only about the finite levels of the Fuzzy Ershov Hierarchy. Similarly to the classical case, we could introduce transfinite levels. These levels are labelled by the notations of constructive ordinals, taken from Kleene’s \mathcal{O} (see § 11.7 in [13]).

Definition 4.2. Let a be an element of \mathcal{O} . We say that a fuzzy Δ_2^0 set A belongs to the class Σ_a^{-1} if there exist a Δ_2^0 -approximation $f(x, s)$ and a computable “counting” function $h: \omega^2 \rightarrow \{b \in \mathcal{O} : b <_{\mathcal{O}} a\}$ such that for all x and s :

- $\lim_s f(x, s) = A(x)$ and $f(x, 0) = 0$;
- $h(x, s + 1) \leq_{\mathcal{O}} h(x, s)$; and
- if $m_{\Sigma}^f(x, s + 1) \neq m_{\Sigma}^f(x, s)$, then $h(x, s + 1) \neq h(x, s)$.

We note the following: if a is the notation of a finite ordinal $n \geq 1$, then the Σ_a^{-1} sets are *precisely* the fuzzy n -c.e. sets.

In an extended version of this work, we will prove that, similarly to Proposition 3.3, the classes Σ_a^{-1} , $a \in \mathcal{O}$, still *do not exhaust* all fuzzy Δ_2^0 sets. This provides a *major difference* with the classical case: every crisp Δ_2^0 set belongs to Σ_a^{-1} for some $a \in \mathcal{O}$ (Theorem 6 in [5], see also Theorem 4.3 in [16]).

Acknowledgements. The authors are grateful to Professor Murat Ramazanov for his hospitality during their visit to Qaraghandy, Kazakhstan, in June 2019. Part of the research contained in the paper was carried out while Bazhenov, San Mauro, and Ospichev were visiting the Department of Mathematics of Nazarbayev University, Nur-Sultan.

REFERENCES

- [1] Benjamin Rene Callejas Bedregal and Santiago Figueira. Classical computability and fuzzy Turing machines. In J. R. Correa, A. Hevia, and M. Kiwi, editors, *LATIN 2006: Theoretical Informatics*, volume 3887 of *LNCS*, pages 154–165, Berlin, 2006. Springer.
- [2] L. Biacino and G. Gerla. Decidability, recursive enumerability and Kleene hierarchy for L -subsets. *Z. Math. Logik Grundlagen Math.*, 35(1):49–62, 1989.
- [3] R. G. Downey and D. R. Hirschfeldt. *Algorithmic randomness and complexity*. Springer, New York, 2010.
- [4] Yu. L. Ershov. A hierarchy of sets. I. *Algebra Logic*, 7(1):25–43, 1968.
- [5] Yu. L. Ershov. On a hierarchy of sets, II. *Algebra Logic*, 7(4):212–232, 1968.
- [6] Yu. L. Ershov. On a hierarchy of sets. III. *Algebra Logic*, 9(1):20–31, 1970.
- [7] G. Gerla. *Fuzzy logic. Mathematical tools for approximate reasoning*, volume 11 of *Trends in Logic*. Kluwer Academic Publishers, Dordrecht, 2001.
- [8] L. Harkleroad. Fuzzy recursion, RET’s and isols. *Z. Math. Logik Grundlagen Math.*, 30(26–29):425–436, 1984.
- [9] L. Harkleroad. Fuzzy regressivity and retraceability. *Z. Math. Logik Grundlagen Math.*, 34(6):523–529, 1988.
- [10] S. C. Kleene. Recursive predicates and quantifiers. *Trans. Am. Math. Soc.*, 53(1):41–73, 1943.
- [11] E. Mark Gold. Limiting recursion. *J. Symb. Log.*, 30(1):28–48, 1965.
- [12] H. Putnam. Trial and error predicates and the solution to a problem of Mostowski. *J. Symb. Log.*, 30(1):49–57, 1965.
- [13] H. Rogers. *Theory of recursive functions and effective computability*. McGraw-Hill, New York, 1967.
- [14] E. S. Santos. Fuzzy algorithms. *Inf. Control*, 17(4):326–339, 1970.
- [15] R. I. Soare. *Turing computability. Theory and applications*. Springer, Berlin, 2016.
- [16] F. Stephan, Y. Yang, and L. Yu. Turing degrees and the Ershov hierarchy. In T. Arai, J. Brendle, H. Kikyo, C. T. Chong, R. Downey, Q. Feng, and H. Ono, editors, *Proceedings of the 10th Asian Logic Conference*, pages 300–321. World Scientific, Singapore, 2009.
- [17] Jiri Wiedermann. Fuzzy Turing machines revised. *Comput. Inform.*, 21(3):251–264, 2002.
- [18] L. A. Zadeh. Fuzzy sets. *Inf. Control*, 8(3):338–353, 1965.
- [19] L. A. Zadeh. Fuzzy algorithms. *Inf. Control*, 12(2):94–102, 1968.

SOBOLEV INSTITUTE OF MATHEMATICS, 4 ACAD. KOPTYUG AVE., NOVOSIBIRSK, 630090, RUSSIA
Email address: `bazhenov@math.nsc.ru`

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCES AND HUMANITIES, NAZARBAYEV UNIVERSITY, 53 QABANBAYBATYR AVENUE, NUR-SULTAN, 010000, KAZAKHSTAN
Email address: `manat.mustafa@nu.edu.kz`

SOBOLEV INSTITUTE OF MATHEMATICS, 4 ACAD. KOPTYUG AVE., NOVOSIBIRSK, 630090, RUSSIA
Email address: `ospichev@math.nsc.ru`

INSTITUTE OF DISCRETE MATHEMATICS AND GEOMETRY, VIENNA UNIVERSITY OF TECHNOLOGY
Email address: `luca.sanmauro@gmail.com`