

Contracting T^2 symmetry local existence for Einstein-Vlasov-Scalar field system

A. T. Lassiye¹ and D. Tegankong²

Department of Mathematics, Advanced Teacher Training College,
University of Yaounde 1, PO Box 47, Yaounde Cameroon
e-mail : ¹ tchuanifils@yahoo.fr, ² david.tegankong@univ-yaounde1.cm

Abstract

The evolution of self-gravitating collision-less matter and scalar waves within the general relativity context is described by Einstein and Vlasov equations. The sources of Einstein equations are generated by a distribution function and a scalar field, respectively subject to the Vlasov and wave equations. We prove in contracting T^2 symmetry case, a local in time existence of solutions.

Keywords: Einstein; Vlasov; Scalar field; T^2 symmetry; conformal coordinates.

MSC2020: 83C05, 83C20, 35A01, 35A02, 35L40, 35L45, 35Q83.

1 Introduction

The question of global existence solutions of Einstein equations with matter or not is very important in the mathematical study of general relativity. An essential tool in an investigation of this type is first of all a local in time existence theorem. In this paper a theorem of this kind is proved for one particular choice of matter model such as collision-less matter described by the Vlasov equation and a scalar field for the cosmological case.

In [10] local in time cosmological solutions of the Einstein Vlasov system with massless scalar field in surface symmetry written in areal coordinates is obtained. The method used is adapted here in the case of T^2 symmetry with nonlinear scalar field in the passed direction in conformal coordinates. There are several reasons why it is of interest to look at the case of a scalar field (cf [10] and references therein). Spacetimes with T^2 symmetry have received much attention by different authors for last years. For example in [2], [9] and [11], global existence result was proved in the case of Einstein-Vlasov system using fundamental local existence in time result of Choquet in [4]. There are three types of time coordinates which have been studied in the inhomogeneous

Einstein-Vlasov system case : constant mean curvature, areal and conformal. A constant mean curvature time coordinate t is one where each hypersurface of constant time has constant mean curvature and on each hypersurface of this kind the value of t is the mean curvature of that slice. In the case of areal coordinates the time coordinate is a function of (or is taken to be proportional to) the area of the surfaces of symmetry. In the case of conformal coordinates the metric is conformally flat on a two-dimensional Lorentzian manifold which is the quotient of spacetime by the symmetry group. The time coordinate $R = t$ is the area of the symmetry orbits. Conformal coordinates has advantages to simplify estimations of solutions. By a long chain of geometrical arguments as in [3], ones deduce the local foliations in the past time direction of the spacetime by areal coordinates system.

The paper proceeds as follows. In section 2, we present in T^2 symmetry the system in conformal coordinates. Section 3 is devoted to a priori estimations of unknowns functions and their derivatives. Section 4 deals on local in time existence of solution based on Picard's iterations.

Let us now recall the formulation of the Einstein-Vlasov-scalar field system. The spacetime is a four-dimensional manifold M , with local coordinates $(x^\lambda) = (t, x^i)$ on which $x^0 = t$ denotes the time and (x^i) the space coordinates. Greek indices always run from 0 to 3, and Latin ones from 1 to 3. On M , a Lorentzian metric g is given with signature $(-, +, +, +)$. We consider a self-gravitating collision-less gas and restrict ourselves to the case where all particles have the same rest mass $m = 1$, and move forward in time. We denote by (p^λ) the momenta of the particles. The conservation of the quantity $g_{\lambda\beta}p^\lambda p^\beta$ requires that the phase space of the particle is the seven-dimensional sub-manifold

$$PM = \{g_{\mu\eta}p^\mu p^\eta = -1; p^0 > 0\}$$

of TM which is coordinatized by (t, x^i, p^i) . The energy-momentum tensor is given by

$$T_{\mu\nu} = T_{\mu\nu}^f + T_{\mu\nu}^\phi \quad (1.1)$$

with:

$$T_{\alpha\beta}^f = - \int_{\mathbb{R}^3} f p_\alpha p_\beta |g|^{1/2} \frac{dp^1 dp^2 dp^3}{p_0} \quad (1.2)$$

$$T_{\alpha\beta}^\phi = \nabla_\alpha \phi \nabla_\beta \phi - \frac{1}{2} g_{\alpha\beta} (\nabla_\sigma \phi \nabla^\sigma \phi + 2V(\phi)) \quad (1.3)$$

where the distribution function of the particles is a non-negative real-valued function denoted by f defined on PM . $p_\lambda = g_{\lambda\beta}p^\beta$, $|g|$ denotes the modulus of determinant of the metric g . A scalar field ϕ is a real-valued C^∞ function on M with nonlinear potential $V \in C^\infty(\mathbb{R}_+)$ which satisfies $V(0) = V_0 > 0$, $V'(0) = 0$ and $V''(0) > 0$ (see [8]). The Einstein field equations are given by

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = 8\pi T_{\mu\nu} \quad (1.4)$$

where $R_{\mu\nu}$ and R are respectively the Ricci tensor and scalar curvature of metric g . The Einstein field equations are coupled to the Vlasov equation (matter

equation for f) and to the wave equation (matter equation for ϕ), which are respectively

$$p^\mu \frac{\partial f}{\partial x^\mu} - \Gamma_{\nu\gamma}^i p^\nu p^\gamma \frac{\partial f}{\partial p^i} = 0 \quad (1.5)$$

$$\nabla^\lambda \nabla_\lambda \phi = V'(\phi). \quad (1.6)$$

Equation (1.6) is a consequence of the divergence-free of the energy-momentum tensor due to the Bianchi identities and since the contribution of f to the energy-momentum tensor is divergence-free.

2 Equations

We refer to [1] for details on the notion of T^2 symmetry. There are several choices of spacetime manifolds compatible with T^2 symmetry. We restrict our attention to the T^3 case. Space-times admitting a T^2 isometry group acting on T^3 space-like surfaces are more general than the T^2 space-times: both families admit two commuting Killing vectors. We consider the case where all unknowns are invariant under the T^2 symmetry with the twists different from zero ([9], [11]). The dynamics of the matter is governed by the Vlasov and the non-linear wave equations. The Vlasov equation models a collision-less system of particles which follow the geodesics of spacetime.

We now consider a solution of the Einstein-Vlasov-scalar field system where all unknowns are invariant under this symmetry. We write the system in conformal coordinates. The circumstances under which coordinates of this type exist are discussed in [1] and references therein. In such coordinates, the metric g takes the form (cf [1], [9], [7])

$$ds^2 = e^{2(\tau-\mu)}(-dt^2 + d\theta^2) + e^{2\mu}[dx + Ady + (G + AH)d\theta]^2 + e^{-2\mu}R^2[dy + Hd\theta]^2 \quad (2.1)$$

where μ , τ , R , G , H and A are unknown real functions of t and θ variables. R is periodic in θ with period 2π . The timelike coordinate t locally labels spatial hypersurfaces of the spacetime. The scalar field is a real function of t and θ . On PM we have

$$p^0 = \sqrt{-g^{00}} \sqrt{1 + g_{ab}p^a p^b}$$

Let us introduce new quantities

$$J = -Re^{-2\tau-4\mu}(G_t + AH_t), \quad K = AJ - R^3 e^{-2\tau} H_t, \quad \Gamma = G_t + AH_t. \quad (2.2)$$

Using the results of [7] and [9], the complete EVSFS can be written in the following form :

The Vlasov equation

$$\begin{aligned}
& \left[(\tau_\theta - \mu_\theta)v^0 + (\tau_t - \mu_t)v^1 + \mu_\theta \frac{(v^2)^2}{v^0} + \left(\mu_\theta - \frac{R_\theta}{R} \right) \frac{(v^3)^2}{v^0} - \frac{A_\theta}{R} e^{2\mu} \frac{v^2 v^3}{v^0} \right] \frac{\partial f}{\partial v^1} \\
& + e^{-\tau} (e^{2\tau} \Gamma v^2 + R H_t v^3) \frac{\partial f}{\partial v^1} - \frac{v^1}{v^0} \frac{\partial f}{\partial \theta} + \left[\mu_t v^2 + \mu_\theta \frac{v^1 v^2}{v^0} \right] \frac{\partial f}{\partial v^2} - \frac{\partial f}{\partial t} \\
& + \left[\left(\frac{R_t}{R} - \mu_t \right) v^3 + \left(\mu_\theta - \frac{R_\theta}{R} \right) \frac{v^1 v^3}{v^0} + \frac{e^{2\mu} v^2}{R} (A_t + A_\theta \frac{v^1}{v^0}) \right] \frac{\partial f}{\partial v^3} = 0 \quad (2.3)
\end{aligned}$$

The Einstein-matter constraint equations

$$\begin{aligned}
\mu_t^2 + \mu_\theta^2 + \frac{e^{4\mu}}{4R^2} (A_t^2 + A_\theta^2) + \frac{R_{\theta\theta}}{R} - \frac{\tau_t R_t}{R} - \frac{\tau_\theta R_\theta}{R} \\
= -\frac{e^{-2\tau+4\mu}}{4} \Gamma^2 - \frac{e^{-2\tau}}{4} H_t^2 - e^{2(\tau-\mu)} \rho \quad (2.4)
\end{aligned}$$

$$2\mu_t \mu_\theta + \frac{e^{4\mu}}{2R^2} A_t A_\theta + \frac{R_{t\theta}}{R} - \frac{\tau_t R_\theta}{R} - \frac{\tau_\theta R_t}{R} = e^{2(\tau-\mu)} J_1 \quad (2.5)$$

The Einstein-matter evolution equations

$$\begin{aligned}
\mu_{tt} - \mu_{\theta\theta} = \frac{\mu_\theta R_\theta}{R} - \frac{\mu_t R_t}{R} + \frac{e^{4\mu}}{2R^2} (A_t^2 - A_\theta^2) + \frac{e^{4\mu-2\tau}}{2} \Gamma^2 \\
+ \frac{e^{4\mu-2\tau}}{2} (\rho - P_1 + P_2 - P_3) \quad (2.6)
\end{aligned}$$

$$\begin{aligned}
A_{tt} - A_{\theta\theta} = \frac{A_t R_t}{R} - \frac{A_\theta R_\theta}{R} + 4(A_\theta \mu_\theta - A_t \mu_t) + R^2 e^{-2\tau} \Gamma H_t \\
+ 2R e^{2(\tau-2\mu)} S_{23} \quad (2.7)
\end{aligned}$$

$$\begin{aligned}
\tau_{tt} - \tau_{\theta\theta} = \mu_\theta^2 - \mu_t^2 + \frac{e^{4\mu}}{4R^2} (A_t^2 - A_\theta^2) - \frac{e^{-2\tau+4\mu}}{4} \Gamma^2 - \frac{3R^2 e^{-2\tau}}{4} H_t^2 \\
- e^{2(\tau-\mu)} P_3 - \frac{2A}{R} e^\tau S_{23} \quad (2.8)
\end{aligned}$$

$$R_{tt} - R_{\theta\theta} = R e^{2(\tau-\mu)} (\rho - P_1) + \frac{R e^{-2\tau+4\mu}}{2} \Gamma^2 + \frac{R^3 e^{-2\tau}}{2} H_t^2 \quad (2.9)$$

$$\begin{aligned}
\phi_{tt} - \phi_{\theta\theta} = \phi_\theta [-4\mu_\theta H^2 R^2 e^{-2\tau} + \frac{R_\theta}{R} + 2A_\theta H (G - R) \\
- \mu_\theta (G + AH)^2 e^{-2(\tau-2\mu)}] - \frac{\phi_t R_t}{R} - e^{-2(\tau-\mu)} V'(\phi) \quad (2.10)
\end{aligned}$$

The Auxiliary equations

$$\partial_\theta (R e^{-2\tau+4\mu} \Gamma) = -2R e^\tau J_2 \quad (2.11)$$

$$\partial_t (R e^{-2\tau+4\mu} \Gamma) = 2R e^\tau S_{12} \quad (2.12)$$

$$\partial_\theta (R^3 e^{-2\tau} H_t) + R e^{-2\tau+4\mu} A_\theta \Gamma = -2R^2 e^{\tau-2\mu} J_3 \quad (2.13)$$

$$\partial_t (R^3 e^{-2\tau} H_t) + R e^{-2\tau+4\mu} A_t \Gamma = 2R^2 e^{\tau-2\mu} S_{13}. \quad (2.14)$$

Since all the particles have proper mass 1, the new variables v^λ are related to the canonical momentum variables p^λ by relations :

$$\begin{aligned} (v^0)^2 &= e^{2(\tau-\mu)}(p^0)^2; & (v^1)^2 &= e^{2(\tau-\mu)}(p^1)^2; \\ (v^2)^2 &= e^{2\mu}[(G+AH)p^1 + p^2 + Ap^3]^2; & (v^3)^2 &= R^2 e^{-2\mu}(Hp^1 + p^3)^2; \end{aligned} \quad \text{so that}$$

$$v^0 = \sqrt{1 + (v^1)^2 + (v^2)^2 + (v^3)^2} > 0$$

The matter terms are then defined by:

$$\rho(t, \theta) = -g^{00}T_{00} = \int_{\mathbb{R}^3} v^0 f(t, \theta, v) dv + \frac{1}{2} e^{-2(\tau-\mu)}(\phi_t^2 + \phi_\theta^2) + V(\phi) \quad (2.15)$$

$$J_1(t, \theta) = -g^{11}T_{01} = \int_{\mathbb{R}^3} v^1 f(t, \theta, v) dv - e^{-2(\tau-\mu)}\phi_t\phi_\theta \quad (2.16)$$

$$J_k(t, \theta) = \int_{\mathbb{R}^3} v^k f(t, \theta, v) dv, \quad k \in \{2, 3\} \quad (2.17)$$

$$P_1(t, \theta) = g^{11}T_{11} = \int_{\mathbb{R}^3} \frac{(v^1)^2}{v^0} f(t, \theta, v) dv + \frac{1}{2} e^{-2(\tau-\mu)}(\phi_t^2 + \phi_\theta^2) - V(\phi) \quad (2.18)$$

$$P_2(t, \theta) = e^{-2\mu}T_{22} = \int_{\mathbb{R}^3} \frac{(v^2)^2}{v^0} f(t, \theta, v) dv + \frac{1}{2} e^{-2(\tau-\mu)}(\phi_t^2 - \phi_\theta^2) - V(\phi) \quad (2.19)$$

$$P_3(t, \theta) = \int_{\mathbb{R}^3} \frac{(v^3)^2}{v^0} f(t, \theta, v) dv + \frac{1}{2} e^{-2(\tau-\mu)}(\phi_t^2 - \phi_\theta^2) - V(\phi) \quad (2.20)$$

$$S_{jk}(t, \theta) = \int_{\mathbb{R}^3} \frac{v^j v^k}{v^0} f(t, \theta, v) dv, \quad j \neq k. \quad (2.21)$$

and

$$T_{33} = A^2 e^{2\mu} P_2 + 2ARS_{23} + R^2 e^{-2\mu} P_3$$

We prescribe initial data at time $t = t_0 > 0$ by

$$(f, R, \tau, \mu, A, H, G, \phi)(t_0) = (\overset{\circ}{f}, \overset{\circ}{R}, \overset{\circ}{\tau}, \overset{\circ}{\mu}, \overset{\circ}{A}, \overset{\circ}{H}, \overset{\circ}{G}, \overset{\circ}{\phi}) \text{ and } (\dot{R}, \dot{\tau}, \dot{\mu}, \dot{A}, \dot{H}, \dot{G}, \dot{\phi})(t_0) = (\bar{R}, \bar{\tau}, \bar{\mu}, \bar{A}, \bar{H}, \bar{G}, \bar{\psi}), \text{ where } \bar{R} = R_t, \dots$$

Let us now remind some regularity definitions which are necessary in the next sections.

Definition 2.1 *Let $I \subseteq]0; +\infty[$ be an interval and $(t, \theta) \in I \times S^1$.*

1. $f \in C^1(I \times S^1 \times \mathbb{R}^3)$ is regular if $f(t, \theta + 2\pi, v) = f(t, \theta, v)$, $f \geq 0$ and $\text{supp}f(t, \theta, \dots, \dots)$ is uniformly compact in θ and locally uniformly compact in t .
2. $\phi \in C^2(I \times S^1)$ is regular if $\phi(t, \theta + 2\pi) = \phi(t, \theta)$.
3. Each component $\chi \in C^2(I \times S^1)$ of metric is regular if $\chi(t, \theta + 2\pi) = \chi(t, \theta)$ and $\partial_t \chi, \partial_\theta \chi \in C^1(I \times S^1)$.
4. ρ (or P_k, J_k, S_{jk}) $\in C^1(I \times S^1)$ is regular if $\rho(t, \theta + 2\pi) = \rho(t, \theta)$.

3 Estimations

In order to obtain local existence in time solution, it is necessary to obtain a priori estimations and uniform bounds on the field components, the distribution function, the scalar field and all their derivatives on a finite time interval $[t_1, t_2]$ on which the local solution can exist. The method and results obtained here are similar to those in [7] and [1].

Let us introduce the null vector fields

$$\partial_\sigma := \frac{1}{\sqrt{2}}(\partial_t + \partial_\theta) \quad \text{and} \quad \partial_\lambda := \frac{1}{\sqrt{2}}(\partial_t - \partial_\theta).$$

For any function F of variables t and θ , set

$$F_\sigma := \partial_\sigma F := \frac{1}{\sqrt{2}}(\partial_t F + \partial_\theta F) := \hat{F} \quad \text{and} \quad F_\lambda := \partial_\lambda F := \frac{1}{\sqrt{2}}(\partial_t F - \partial_\theta F) := \check{F}.$$

Step 1 Monotonicity of R and bounds on its first derivatives.

After some calculation, the constraints equations (2.4) and (2.5) give respectively

$$\begin{aligned} \sqrt{2}\partial_\theta R_\sigma &= 2\tau_\sigma R_\sigma - 2R\mu_\sigma^2 - \frac{e^{4\mu}}{2R^2}A_\sigma^2 - Re^{2(\tau-\mu)}(\rho - J_1) \\ &\quad - \frac{R}{4}e^{-2\tau+4\mu}\Gamma^2 - \frac{R^3e^{-2\tau}}{4}H_t^2 \end{aligned} \quad (3.1)$$

$$\begin{aligned} \sqrt{2}\partial_\theta R_\lambda &= -2\tau_\lambda R_\lambda + 2R\mu_\lambda^2 + \frac{e^{4\mu}}{2R^2}A_\lambda^2 + Re^{2(\tau-\mu)}(\rho + J_1) \\ &\quad + \frac{R}{4}e^{-2\tau+4\mu}\Gamma^2 + \frac{R^3}{4}e^{-2\tau}H_t^2 \end{aligned} \quad (3.2)$$

It follows from (2.15) and (2.16), that $\rho \geq |J_1|$ (in fact $V(\phi) \geq 0$) and also $R > 0$, so

$$\partial_\theta R_\sigma = \sqrt{2}\tau_\sigma R_\sigma - \frac{\sqrt{2}}{2}\mathfrak{h}_1 < \sqrt{2}\tau_\sigma R_\sigma \quad (3.3)$$

$$\partial_\theta R_\lambda = -\sqrt{2}\tau_\lambda R_\lambda + \mathfrak{h}_2 > -\sqrt{2}\tau_\lambda R_\lambda \quad (3.4)$$

where

$$\mathfrak{h}_1 = \frac{e^{4\mu}}{2R}A_\sigma^2 + R\mu_\sigma^2 + R \left[e^{2(\tau-\mu)}(\rho - J_1) + \frac{e^{-2\tau+4\mu}}{4}\Gamma^2 + \frac{R^2e^{-2\tau}}{4}H_t^2 \right] \quad (3.5)$$

$$\mathfrak{h}_2 = R\mu_\lambda^2 + \frac{e^{4\mu}}{2R}A_\lambda^2 + R \left[e^{2(\tau-\mu)}(\rho + J_1) + \frac{e^{-2\tau+4\mu}}{4}\Gamma^2 + \frac{R^2e^{-2\tau}}{2\sqrt{2}}H_t^2 \right] \quad (3.6)$$

As in [7], we obtain after integration,

$$\sqrt{2}R_t(t, \theta) \leq R_\sigma(t_0, \theta + t - t_0) + R_\lambda(t_0, \theta - t + t_0) \leq \sup_{\theta \in S^1} (R_\sigma + R_\lambda)(t_0, \theta). \quad (3.7)$$

and deduce that R_t is bounded into the past and $|R_\theta|$ is also bounded. Consequently, R is uniformly bounded to the past of the initial surface.

Step 2 Bounds on μ , A , τ , ϕ and their first derivatives.

We use the light-cone argument and Gronwall's lemma in this step. The functions involved in this case are quadratic and defined by

$$X = \frac{1}{2}R(\mu_t^2 + \mu_\theta^2) + \frac{e^{4\mu}}{8R}(A_t^2 + A_\theta^2) + \frac{1}{2}[R(\phi_t^2 + \phi_\theta^2) + \phi^2] \quad (3.8)$$

$$Y = R\mu_t\mu_\theta + \frac{e^{4\mu}}{4R}A_tA_\theta + R\phi_t\phi_\theta \quad (3.9)$$

Using (2.6) – (2.7) and (2.10) we find respectively after a lengthy calculation that

$$\begin{aligned} \partial_\lambda(X + Y) &= -\frac{1}{2}R_\sigma \left(\mu_t^2 - \mu_\theta^2 + \frac{e^{4\mu}}{4R^2}(-A_t^2 + A_\theta^2) + \phi_t^2 - \phi_\theta^2 \right) \\ &\quad - Re^{2(\tau-\mu)}\phi_\sigma V'(\phi) + \frac{R}{2}\mu_\sigma \left(e^{-2\tau+4\mu}\Gamma^2 + e^{2(\tau-\mu)}(\rho - P_1 + P_2 - P_3) \right) \\ &\quad + \frac{e^{4\mu}}{2R}A_\sigma(R^2 e^{2(\mu-\tau)}\Gamma H_t + 2Re^{2(\tau-\mu)}S_{23}) + \phi\phi_\lambda \quad . \end{aligned} \quad (3.10)$$

and

$$\begin{aligned} \partial_\sigma(X - Y) &= -\frac{1}{2}R_\lambda \left(\mu_t^2 - \mu_\theta^2 + \frac{e^{4\mu}}{4R^2}(-A_t^2 + A_\theta^2) + \phi_t^2 - \phi_\theta^2 \right) \\ &\quad - Re^{2(\tau-\mu)}\phi_\lambda V'(\phi) + \frac{R}{2}\mu_\lambda \left(e^{-2\tau+4\mu}\Gamma^2 + e^{2(\tau-\mu)}(\rho - P_1 + P_2 - P_3) \right) \\ &\quad + \frac{e^{4\mu}}{2R}A_\lambda(R^2 e^{2(\mu-\tau)}\Gamma H_t + 2Re^{2(\tau-\mu)}S_{23}) + \phi\phi_\sigma \quad . \end{aligned} \quad (3.11)$$

Integrating each of the above equations along null paths $t \mapsto (t, \theta - t_0 + t)$ and $t \mapsto (t, \theta + t_0 - t)$ starting at (t_1, θ) and ending at the initial t_0 -surface, and adding the results we obtain

$$\begin{aligned} X(t_1, \theta) &= \frac{1}{2}(X + Y)(t_0, \theta - (t_0 - t_1)) + \frac{1}{2}(X - Y)(t_0, \theta + t_0 - t_1) \\ &\quad - \frac{1}{2} \int_{t_1}^{t_0} [\chi_1(s, \theta - (s - t_1)) + \chi_2(s, \theta + s - t_1)] ds \\ &\quad - \frac{1}{2} \int_{t_1}^{t_0} \left[\left(\frac{e^{4\mu}}{2R} A_\sigma \zeta_2 \right)(s, \theta - (s - t_1)) + \left(\frac{e^{4\mu}}{2R} A_\lambda \zeta_2 \right)(s, \theta + s - t_1) \right] ds \\ &\quad - \frac{1}{2} \int_{t_1}^{t_0} [(\mu_\sigma \zeta_1)(s, \theta - (s - t_1)) + (\mu_\lambda \zeta_1)(s, \theta + s - t_1)] ds \quad (3.12) \end{aligned}$$

where

$$\begin{aligned}\chi_1 &= -\frac{1}{2}R_\sigma \left(\mu_t^2 - \mu_\theta^2 + \frac{e^{4\mu}}{4R^2}(-A_t^2 + A_\theta^2) + \phi_t^2 - \phi_\theta^2 \right) \\ &\quad - \frac{R_t}{\sqrt{2}}\phi_t^2 + \frac{R_\theta}{\sqrt{2}}\phi_\theta^2 + \phi\phi_\lambda - R\phi_\sigma e^{2(\tau-\mu)}V'(\phi)\end{aligned}\quad (3.13)$$

$$\begin{aligned}\chi_2 &= -\frac{1}{2}R_\lambda \left(\mu_t^2 - \mu_\theta^2 + \frac{e^{4\mu}}{4R^2}(-A_t^2 + A_\theta^2) + \phi_t^2 - \phi_\theta^2 \right) \\ &\quad - \frac{R_t}{\sqrt{2}}\phi_t^2 + \frac{R_\theta}{\sqrt{2}}\phi_\theta^2 + \phi\phi_\sigma - R\phi_\lambda e^{2(\tau-\mu)}V'(\phi)\end{aligned}\quad (3.14)$$

$$\zeta_1 = \frac{R}{2} \left(e^{-2\tau+4\mu}\Gamma^2 + e^{2(\tau-\mu)}(\rho - P_1 + P_2 - P_3) \right) \quad (3.15)$$

$$\zeta_2 = R^2 e^{2(\mu-\tau)}\Gamma H_t + 2R e^{2(\tau-\mu)}S_{23} \quad (3.16)$$

It follows from Step 1 and Step 2 of [7], the uniform bound of

$$\int_t^{t_0} \zeta_1(s, \theta \pm (t_0 - s))ds \quad \text{and} \quad \int_t^{t_0} |\zeta_2|(s, \theta \pm (t_0 - s))ds \quad (3.17)$$

Using Step 8 of [7], we conclude that $\sup_{S^1} X$ is uniformly bounded on (t_1, t_0) , leading to bounds on μ , A , ϕ , $V(\phi)$ and their first derivatives. The bounds of τ and its first derivatives are obtained in a similar way since (2.8) can be written as

$$2\partial_\sigma\tau_\lambda = 2\partial_\lambda\tau_\sigma = \mu_\theta^2 - \mu_t^2 + \frac{e^{4\mu}}{4R^2}(A_t^2 - A_\theta^2) - \frac{e^{-2\tau+4\mu}}{4}\Gamma^2 - \frac{3R^2 e^{-2\tau}}{4}H_t^2 - e^{2(\tau-\mu)}P_3 \quad (3.18)$$

The integrals along the null paths for the matter terms in the right hand side of (3.18) is bounded since A , μ , ϕ and their first derivatives are bounded. We obtain that τ_λ and τ_σ are bounded, and therefore

$$\tau_t = \frac{1}{\sqrt{2}}(\tau_\sigma + \tau_\lambda), \quad \tau_\theta = \frac{1}{\sqrt{2}}(\tau_\sigma - \tau_\lambda) \quad \text{and} \quad \tau \text{ are bounded.}$$

Step 3 Bounds on G , H their derivatives and the support of the momentum. A solution f to the Vlasov equation is given by

$$f(t, \theta, v) = f_0(\Theta(t_0, t, \theta, v), V(t_0, t, \theta, v)) := \overset{\circ}{f}(\Theta(t_0, t, \theta, v), V(t_0, t, \theta, v)),$$

where Θ and V are solutions to the characteristic system

$$\begin{cases} \frac{d\Theta}{ds} = \frac{V^1}{V^0} \\ \frac{dV^1}{ds} = -(\tau_\theta - \mu_\theta)V^0 - (\tau_s - \mu_s)V^1 - \mu_\theta \frac{(V^2)^2}{V^0} + (\mu_\theta - \frac{R_\theta}{R}) \frac{(V^3)^2}{V^0} \\ \quad + \frac{A_\theta}{R} e^{2\mu} \frac{V^2 V^3}{V^0} - e^{-\tau} (e^{2\tau} \Gamma V^2 + R H_s V^3) \\ \frac{dV^2}{ds} = -\mu_s V^2 - \mu_\theta \frac{V^1 V^2}{V^0} \\ \frac{dV^3}{ds} = (\frac{R_s}{R} - \mu_s) V^3 + (\mu_\theta - \frac{R_\theta}{R}) \frac{V^1 V^3}{V^0} + \frac{e^{2\mu} V^2}{R} (A_s + A_\theta \frac{V^1}{V^0}) \end{cases}$$

with $\Theta(t_0, t, \theta, v) = \theta$ and $V(t_0, t, \theta, v) = v$. Since $\|f\|_\infty \leq \|\overset{\circ}{f}\|_\infty$, the control of

$$Q(t) := \sup\{|v| : \exists(s, \theta) \in [t, t_0] \times S^1 ; f(s, \theta, v) \neq 0\} \quad (3.19)$$

gives directly using equations (2.15) – (2.21), bounds of ρ , J_k , P_k et S_{jk} , $j, k \in \{1, 2, 3\}$ $j \neq k$.

Besides

$$\frac{dV^2}{V^2} = \left(-\mu_s - \frac{V^1}{V^0}\mu_\theta\right)ds \quad (3.20)$$

After integration on $[t_-; t]$, (3.20) gives

$$V^2(t) \leq |V^2(t_-)| \exp\left(\int_{t_-}^t \left|\mu_s + \frac{V^1}{V^0}\mu_\theta\right| ds\right). \quad (3.21)$$

The last equation of the previous system gives

$$V^3(t) \leq C(t) \exp\left(\int_{t_-}^t \left|\frac{R_s}{R} - \mu_s + \left(-\mu_\theta + \frac{R_\theta}{R}\right) \frac{V^1}{V^0}\right| ds\right) \quad (3.22)$$

where $C(t) = \sup_{\theta \in S^1} \left|\frac{e^{2\mu} V^2}{R}(A_t + A_\theta \frac{V^1}{V^0})\right|(t, \theta)$, $|\frac{V^1}{V^0}|$, $|\mu_t|$ and $|\mu_\theta|$ are all bounded.

Let us define also

$$Q^j(t) := \sup\{|v^j| : \exists(s, \theta) \in [t, t_0] \times S^1; f(s, \theta, v^k) \neq 0, k \neq j\}. \quad (3.23)$$

Consequently V^2 , V^3 , Q^2 , Q^3 are bounded. Now, for $j = 1$ in (3.23) one obtains from the second equation of previous system :

$$|V^1(t)| \leq |V^1(t_0)| + C(t) \int_t^{t_0} [Q^1(s) + \sup_\theta \Gamma(s, \theta) + \sup_\theta H_t(s, \theta)] ds \quad (3.24)$$

Add and subtract respectively auxiliary equations (2.11) – (2.12) to obtain

$$\partial_\lambda(Re^{-2\tau+4\mu}\Gamma) = \sqrt{2}Re^\tau(S_{12} - J_2) \quad (3.25)$$

$$\partial_\sigma(Re^{-2\tau+4\mu}\Gamma) = \sqrt{2}Re^\tau(S_{12} + J_2) \quad (3.26)$$

Integrating along null paths and using previous steps give

$$\sup_\theta \Gamma(t, \theta) \leq \Gamma(t_0, \theta) + C \int_t^{t_0} [1 + Q^1(s)] ds \quad t_1 < t < t_0 \quad (3.27)$$

Analogously, (2.13) and (2.14) give

$$\sup_\theta H_t(t, \theta) \leq |H_t|(t_0, \theta) + C \int_t^{t_0} [1 + Q^1(s) + \sup_\theta |\Gamma|(s, \theta)] ds \quad t_1 < t < t_0 \quad (3.28)$$

Adding (3.24), (3.27), (3.28) and applying Gronwall's lemma give uniform bounds on H_t , Γ and $Q^1(t)$. Since $Q^1(t)$ and V^k are bounded, the integral terms of the matter quantities are also bounded. We deduce from (2.2) that G_t , and K are bounded. Therefore H and G are bounded. From (2.13), we deduce bounds on $H_{t\theta}$ and then on H_t . Consequently (2.11) gives bounds on $G_{\theta t}$ and then on G_θ . We deduce respectively from (2.14) and (2.12) the bounds of H_{tt} and G_{tt} .

4 Local Existence

In this section, we prove using an iteration the local existence and uniqueness of solutions of the EVSFS.

For all component χ of the metric g , we define the iterates $(\chi_n)_{n \in \mathbb{N}}$, $(\phi_n)_{n \in \mathbb{N}}$, $(f_n)_{n \in \mathbb{N}}$ with first terms

$$\chi_0(t, \theta) \equiv \mathring{\chi}(\theta), \quad \phi_0(t, \theta) \equiv \mathring{\phi}(\theta), \quad f_0(t, \theta) \equiv \mathring{f}(\theta)$$

and

$$\chi_n(t_0, \theta) \equiv \mathring{\chi}(\theta), \quad \phi_n(t_0, \theta) \equiv \mathring{\phi}(\theta), \quad f_n(t_0, \theta) \equiv \mathring{f}(\theta)$$

Consider the previous characteristic system where $R, R_\theta, \tau_\theta, \mu_\theta, A_\theta$ are respectively replaced by

$R_{n-1}, \underline{R}_{n-1}, \underline{\tau}_{n-1}, \underline{\mu}_{n-1}, \underline{A}_{n-1}$. In what follows, $\check{R}_{n-1}, \check{\tau}_{n-1}, \check{\mu}_{n-1}, \check{A}_{n-1}, \check{H}_{n-1}, \dots$ replace respectively $(R_{n-1})_t, (\tau_{n-1})_t, (\mu_{n-1})_t, (A_{n-1})_t, (H_{n-1})_t, \dots$ Let

$$\begin{aligned} \Xi_n &= \left[\frac{V^1}{V^0}; \Xi_n^1; \Xi_n^2; \Xi_n^3 \right] \quad \text{where} \\ \Xi_{n-1}^1 &= -(\underline{\tau}_{n-1} - \underline{\mu}_{n-1})V^0 - (\check{\tau}_{n-1} - \check{\mu}_{n-1})V^1 - e^{-\tau_{n-1}}(e^{2\mu_{n-1}}\Gamma_{n-1}V^2 + R_{n-1}\check{H}_{n-1}V^3) \\ &\quad + \frac{e^{2\mu_{n-1}}\underline{A}_{n-1}}{R_{n-1}} \frac{V^2V^3}{V^0} - \left(\underline{\mu}_{n-1} - \frac{\underline{R}_{n-1}}{R_{n-1}} \right) \frac{(V^3)^2}{V^0} + \underline{\mu}_{n-1} \frac{(V^2)^2}{V^0}; \\ \Xi_{n-1}^2 &= -\check{\mu}_{n-1}V^2 - \underline{\mu}_{n-1} \frac{V^1V^2}{V^0}; \\ \Xi_{n-1}^3 &= \left(\frac{\check{R}_{n-1}}{R_{n-1}} - \check{\mu}_{n-1} \right) V^3 + \left(\underline{\mu}_{n-1} - \frac{\underline{R}_{n-1}}{R_{n-1}} \right) \frac{V^1V^3}{V^0} + \frac{e^{2\mu_{n-1}}}{R_{n-1}} \left(\check{A}_{n-1} + \underline{A}_{n-1} \frac{V^1}{V^0} \right) V^2 \end{aligned}$$

Define $(\Theta_n, V_n)(s, t, \theta, V^1, V^2, V^3)$ with $(t \leq t_0)$ a solution of the characteristic system

$$\frac{d}{ds} ((\Theta, V)_n) = \Xi_{n-1}(s, t, \Theta, V) \quad (4.1)$$

with initial data $(\Theta_n, V_n)(t, t, \theta, V^1, V^2, V^3) = (\theta, v)$ and

$$f_n(t, \theta, V^1, V^2, V^3) = f^\circ((\Theta_n, V_n)(t_0, \theta, V^1, V^2, V^3)) \quad (4.2)$$

where f_n is the solution of

$$\begin{aligned} &\left[(\underline{\tau}_{n-1} - \underline{\mu}_{n-1})v^0 + (\check{\tau}_{n-1} - \check{\mu}_{n-1})v^1 + \underline{\mu}_{n-1} \frac{(v^2)^2}{v^0} + \left(\underline{\mu}_{n-1} - \frac{\underline{R}_{n-1}}{R_{n-1}} \right) \frac{(v^3)^2}{v^0} - \frac{\underline{A}_{n-1}}{R_{n-1}} e^{2\mu_{n-1}} \frac{v^2v^3}{v^0} \right] \frac{\partial f_n}{\partial v^1} \\ &\quad - \left[e^{-\tau_{n-1}}(e^{2\tau_{n-1}}\Gamma_{n-1}v^2 + R_{n-1}\check{H}_{n-1}v^3) \right] \frac{\partial f_n}{\partial v^1} - \frac{\partial f_n}{\partial t} - \frac{v^1}{v^0} \frac{\partial f_n}{\partial \theta} + \left[\check{\mu}_{n-1}v^2 + \underline{\mu}_{n-1} \frac{v^1v^2}{v^0} \right] \frac{\partial f_n}{\partial v^2} \\ &\quad - \left[\left(\frac{\check{R}_{n-1}}{R_{n-1}} - \check{\mu}_{n-1} \right) v^3 + \left(\underline{\mu}_{n-1} - \frac{\underline{R}_{n-1}}{R_{n-1}} \right) \frac{v^1v^3}{v^0} + \frac{e^{2\mu_{n-1}}v^2}{R_{n-1}} \left(\check{A}_{n-1} + \underline{A}_{n-1} \frac{v^1}{v^0} \right) \right] \frac{\partial f_n}{\partial v^3} = 0 \end{aligned}$$

Also, using (3.1) – (3.2):

$$\begin{aligned}\partial_\theta \hat{R}_n &= \hat{\tau}_{n-1} \hat{R}_{n-1} - R_{n-1} \hat{\mu}_{n-1}^2 - \frac{e^{4\mu_{n-1}}}{4R_{n-1}^2} \hat{A}_{n-1}^2 \\ &\quad - R_{n-1} e^{2(\tau_{n-1} - \mu_{n-1})} (\rho_n - J_{n,1}) \\ &\quad - \frac{e^{-2\tau_{n-1} + 4\mu_{n-1}}}{4} \Gamma_{n-1}^2 - \frac{R_{n-1}^2 e^{-2\tau_{n-1}}}{4} \check{H}_{n-1}^2\end{aligned}\quad (4.3)$$

$$\begin{aligned}\partial_\theta \check{R}_n &= -\check{\tau}_{n-1} \check{R}_{n-1} + R_{n-1} \check{\mu}_{n-1}^2 + \\ &\quad \frac{e^{4\mu_{n-1}}}{4R_{n-1}^2} \check{A}_{n-1}^2 + R_{n-1} e^{2(\tau_{n-1} - \mu_{n-1})} (\rho_n + J_{n,1}) \\ &\quad + \frac{e^{-2\tau_{n-1} + 4\mu_{n-1}}}{4} \Gamma_{n-1}^2 + \frac{R_{n-1}^2 e^{-2\tau_{n-1}}}{4} \check{H}_{n-1}^2\end{aligned}\quad (4.4)$$

Define also $\rho_n, \rho_{k,n}, P_{k,n}, k \in \{1; 2; 3\}$ in the same way as ρ, ρ_k, P_k , where $f, R, \phi_t, \phi_\theta, A_t, \mu, \tau, \mu_t, \tau_t$ are respectively replaced by $f_n, R_{n-1}, \check{\phi}_{n-1}, \underline{\phi}_{n-1}, \check{A}_{n-1}, \underline{\mu}_{n-1}, \tau_{n-1}, \check{\mu}_{n-1}, \check{\tau}_{n-1}$.

We define X_n and Y_n like (3.8) – (3.9) by:

$$X_n = \frac{1}{2} R_n (\check{\mu}_n^2 + \underline{\mu}_n^2) + \frac{e^{4\mu_n}}{8R_n} (\check{A}_n^2 + \underline{A}_n^2) + \frac{1}{2} [R_n (\check{\phi}_n^2 + \underline{\phi}_n^2) + \phi_n^2] \quad (4.5)$$

$$Y_n = R_n \check{\mu}_n \underline{\mu}_n + \frac{e^{4\mu_n}}{4R_n} \check{A}_n \underline{A}_n + R_n \check{\phi}_n \underline{\phi}_n \quad (4.6)$$

Let $Z_n = X_n + Y_n$, and $\tilde{Z}_n = X_n - Y_n$, therefore

$$Z_n = R_n \hat{\mu}_n^2 + \frac{e^{4\mu_n}}{4R_n} \hat{A}_n^2 + R_n \hat{\phi}_n^2 + \frac{1}{2} \phi_n^2 \quad (4.7)$$

$$\tilde{Z}_n = R_n \check{\mu}_n^2 + \frac{e^{4\mu_n}}{4R_n} \check{A}_n^2 + R_n \check{\phi}_n^2 + \frac{1}{2} \phi_n^2. \quad (4.8)$$

Now, define

$$\mathfrak{h}_n, \check{\mathfrak{h}}_n, \chi_n, \check{\chi}_n, \zeta_n, \check{\zeta}_n, \hat{\tilde{Z}}_n, \check{\tilde{Z}}_n \quad (4.9)$$

using (3.5), (3.6), (3.13), (3.14), (3.15), (3.16), (3.11) and (3.10) with

$$\mathfrak{h}_1, \mathfrak{h}_2, \chi_1, \chi_2, \zeta_1, \zeta_2, \partial_\sigma Z, \partial_\lambda \tilde{Z} \text{ and}$$

$$R, \phi_t, \phi_\theta, A_t, A_\theta, \mu, \tau, \mu_t, \tau_t, \mu', \tau_\theta, \rho, P_k, (k \in \{1; 2; 3\})$$

respectively replaced by $\mathfrak{h}_n, \check{\mathfrak{h}}_n, \chi_n, \check{\chi}_n, \zeta_n, \check{\zeta}_n, \hat{\tilde{Z}}_n, \check{\tilde{Z}}_n, R_{n-1}, \check{\phi}_{n-1}, \underline{\phi}_{n-1}, \check{A}_{n-1}, \underline{A}_{n-1}, \mu_{n-1}, \tau_{n-1}, \check{\mu}_{n-1}, \check{\tau}_{n-1}, \underline{\mu}_{n-1}, \tau_{n-1}, \rho_n, P_{k,n}, (k \in \{1; 2; 3\})$.

From (4.5) – (4.8) we deduce

$$\partial_\lambda Z_n := \tilde{Z}_n = \chi_{n-1} + \check{\mu}_{n-1} \zeta_{n-1} + \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \check{A}_{n-1} \check{\zeta}_{n-1} \quad (4.10)$$

$$\partial_\sigma \tilde{Z}_n := \hat{\tilde{Z}}_n = \check{\chi}_{n-1} + \hat{\mu}_{n-1} \zeta_{n-1} + \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \hat{A}_{n-1} \check{\zeta}_{n-1} \quad (4.11)$$

From auxiliary equations (2.11) – (2.14) we obtain

$$\partial_\lambda(R_n e^{-2\tau_n + 4\mu_n} \Gamma_n) = -\sqrt{2} R_{n-1} e^{\tau_{n-1}} (J_{2,n} - S_{12,n}) \quad (4.12)$$

$$\partial_\sigma(R_n e^{-2\tau_n + 4\mu_n} \Gamma_n) = -\sqrt{2} R_{n-1} e^{\tau_{n-1}} (J_{2,n} + S_{12,n}) \quad (4.13)$$

$$\partial_\lambda(R_n^3 e^{-2\tau_n} \check{H}_n) + (\partial_\lambda A_n) R_n e^{-2\tau_n + 4\mu_n} \Gamma_n = -\sqrt{2} R_{n-1}^2 e^{\tau_{n-1} - 2\mu_{n-1}} (J_{3,n} - S_{13,n}) \quad (4.14)$$

$$\partial_\sigma(R_n^3 e^{-2\tau_n} \check{H}_n) + (\partial_\sigma A_n) R_n e^{-2\tau_n + 4\mu_n} \Gamma_n = -\sqrt{2} R_{n-1}^2 e^{\tau_{n-1} - 2\mu_{n-1}} (J_{3,n} + S_{13,n}) \quad (4.15)$$

Now we proceed for (4.3)–(4.4) the same way as we did for step 1 to establish the inequality (3.7) and obtain the following analogous inequality:

$$\sqrt{2} \check{R}_n(t, \theta) \leq \sup_{\theta \in S^1} (\hat{R}_{n-1} + \check{R}_{n-1})(t_0, \theta). \quad (4.16)$$

Using this last inequality (4.16), we deduce that $R_n, \check{R}_n, \underline{R}_n$ are C^1 -bounded. Proceeding as in step 2 we have

$$\zeta_n \leq \frac{R_{n-1}}{2} [2e^{2(\tau_{n-1} - \mu_{n-1})} (\rho_n - P_{1;n}) + e^{-2\tau_{n-1} + 4\mu_{n-1}} \Gamma_{n-1}^2] \quad (4.17)$$

$$|\check{\zeta}_n| \leq R_{n-1} e^{2(\tau_{n-1} - \mu_{n-1})} |S_{23,n-1}| + R_{n-1}^2 e^{2(\mu_{n-1} - \tau_{n-1})} |\Gamma_{n-1} \check{H}_{n-1}|$$

i.e

$$|\check{\zeta}_n| \leq R_{n-1} e^{2(\tau_{n-1} - \mu_{n-1})} |S_{23,n-1}| + \frac{1}{2} R_{n-1}^2 e^{-2\tau_{n-1}} (e^{4\mu_{n-1}} \Gamma_{n-1}^2 + \check{H}_{n-1}^2) \quad (4.18)$$

$$e^{2(\tau_n - \mu_n)} P_{2;n} \leq \int_{\mathbb{R}^3} f_n \frac{e^{2(\tau_{n-1} - \mu_{n-1})} (v^2)^2}{v^0} dv + \frac{1}{2} \check{\phi}_{n-1}^2 \leq e^{2(\tau_{n-1} - \mu_{n-1})} (\rho_n - P_{1;n}) + \frac{1}{2} \check{\phi}_{n-1}^2. \quad (4.19)$$

Without lost of generality, we choose $V(\phi_n) = V_0 \exp(\phi_n^2)$ such that

$$\int_t^{t_0} \hat{\phi}_n e^{2(\tau_n - \mu_n)} \frac{dV(\phi_n)}{d\phi_n} ds \leq C \sup_{t \in [t_1, t_0]} X_n(t, \theta), \quad (4.20)$$

$$\int_t^{t_0} \check{\phi}_n e^{2(\tau_n - \mu_n)} \frac{dV(\phi_n)}{d\phi_n} ds \leq C \sup_{t \in [t_1, t_0]} X_n(t, \theta), \quad (4.21)$$

Using bounds of R_n and notation (4.9), we have respectively:

$$\int_{t_1}^{t_0} (\hat{\mu}_n \zeta_n)(s, \theta + s - t) ds \leq \sup_{t \in [t_1, t_0]} \sqrt{X_n(t, \theta)} \quad (4.22)$$

$$\int_{t_1}^{t_0} (\check{\mu}_n \zeta_n)(s, \theta - s + t) ds \leq \sup_{t \in [t_1, t_0]} \sqrt{X_n(t, \theta)} \quad (4.23)$$

$$\int_{t_1}^{t_0} \left(\frac{e^{4\mu_n}}{2R_n} |\hat{A}_n \check{\zeta}_n| \right) (s, \theta + s - t) ds \leq C \sup_{t \in [t_1, t_0]} \sqrt{X_n(t, \theta)} \quad (4.24)$$

$$\int_{t_1}^{t_0} \left(\frac{e^{4\mu_n}}{2R_n} |\check{A}_n \check{\zeta}_n| \right) (s, \theta - s + t) ds \leq C \sup_{t \in [t_1, t_0]} \sqrt{X_n(t, \theta)} \quad (4.25)$$

with $t_- \leq t_1 \leq t_0$.

From (4.20) and (4.21), we obtain

$$\int_{t_1}^{t_0} \chi_n(s, \theta + s - t) ds \leq C \int_{t_1}^{t_0} X_n(s, \theta) ds + C \sup_{\theta} \sqrt{X_n(t, \theta)} \quad (4.26)$$

$$\int_{t_1}^{t_0} \tilde{\chi}_n(s, \theta - s + t) ds \leq C \int_{t_1}^{t_0} X_n(s, \theta) ds + C \sup_{\theta} \sqrt{X_n(t, \theta)} \quad (4.27)$$

Therefore

$$\begin{aligned} \sup_{\theta \in S^1} X_n(t_1, \theta) &\leq C \sup_{(\theta, t) \in S^1 \times [t_-, t_0]} X_n(t, \theta) + C \int_{t_1}^{t_0} \sup_{\theta \in S^1} X_n(s, \theta) ds \\ &\quad + 4 \sup_{\theta \in S^1} X_n(t_0, \theta) + C \sup_{(\theta, t) \in S^1 \times [t_-, t_0] \times S^1} \sqrt{X_n(t, \theta)} \end{aligned} \quad (4.28)$$

and by Gronwall's inequality (4.28) gives bounds of X_n . Consequently from (4.5), $\check{\mu}_n$, $\underline{\mu}_n$, \check{A}_n , \underline{A}_n , $\check{\phi}_n$, $\underline{\phi}_n$ are bounded.

Now from (3.18), we deduce that

$$\begin{aligned} \hat{\tau}_n = \check{\tau}_n &= \hat{\mu}_{n-1}^2 - \check{\mu}_{n-1}^2 + \frac{e^{4\mu_{n-1}}}{4R_{n-1}^2} \hat{A}_{n-1}^2 - \check{A}_{n-1}^2 - \frac{e^{-2\tau_{n-1} + 4\mu_{n-1}}}{4} \Gamma_{n-1}^2 \\ &\quad - \frac{3R_{n-1}^2 e^{-2\tau_{n-1}}}{4} \check{H}_{n-1}^2 - e^{2(\tau_{n-1} - \mu_{n-1})} P_{3,n} \end{aligned} \quad (4.29)$$

After integrating (4.29) along null paths, using (4.18), (4.17), (4.20) and also the fact that A_n , ϕ_n , μ_n are C^1 bounded, we deduce that $\check{\tau}_n$ and $\hat{\tau}_n$ are bounded. Consequently τ_n is C^1 -bounded.

To bound ρ_n , $P_{k;n}$, we use the same approach like step 3. Let us define first

$$Q_n(t) := \sup\{|v| : \exists(s, \theta) \in [t, t_0] \times S^1; f_n(s, \theta, v) \neq 0\} \quad (4.30)$$

where $\Theta_n(t, t, x, v) = \theta$ and $V_n(t, t, x, v) = v$ are solutions of the characteristic system associated to the Vlasov equation (4.2). Let Q_n^j , $j \in \{1, 2, 3\}$ defined by

$$Q_n^j(t) := \sup\{|v^j| : \exists(s, \theta) \in [t, t_0] \times S^1 \text{ such that } f_n(s, \theta, v^k) \neq 0, k \neq j\} \quad (4.31)$$

Using Gronwall's inequality like in 3.22 we have

$$V^3(t) \leq C(t) \exp\left(\int_{t_0}^t \left| \frac{\check{R}_{n-1}}{R_{n-1}} - \check{\mu}_{n-1} + \left(-\underline{\mu}_{n-1} + \frac{\underline{R}_{n-1}}{R_{n-1}}\right) \frac{V^1}{V^0} \right| ds\right) \quad (4.32)$$

where $C(t) = \sup_{\theta \in S^1} \left| \frac{e^{2\mu_{n-1}} V^2}{R_{n-1}} \left(\check{A}_{n-1} + \underline{A}_{n-1} \frac{V^1}{V^0} \right) (t, \theta) \right|$, $|\frac{V^1}{V^0}|$, $|\mu_{n-1}|$ and $|\underline{\mu}_{n-1}|$ are bounded.

We can deduce from $\frac{dV^2}{ds}$ and $\frac{dV^3}{ds}$ that

$$e^{\mu_{n-1}} V^2, \quad A_{n-1} e^{\mu_{n-1}} V^2 + R_{n-1} e^{-\mu_{n-1}} V^3$$

are bounded and also Q_n^2, Q_n^3 . Using the similar approach like (3.27) – (3.28), substituting Q^1 by Q_n^1 , one obtains

$$\sup |V^1(t)| \leq |Q_n^1(t_0)| + C(t) \int_t^{t_0} [Q_n^1(s) + \sup_{\theta} \Gamma_n(s, \theta) + \sup_{\theta} \dot{H}_n(s, \theta)] ds, \quad t < t_0 \quad (4.33)$$

$$\sup_{\theta} \Gamma_n(t, \theta) \leq \Gamma_n(t_0, \theta) + C \int_t^{t_0} [1 + Q_n^1(s)] ds, \quad t < t_0 \quad (4.34)$$

$$\sup_{\theta} \dot{H}_n(t, \theta) \leq |\dot{H}_n|(t_0, \theta) + C \int_t^{t_0} [1 + Q_n^1(s) + \sup_{\theta} |\Gamma_n|(s, \theta)] ds \quad t < t_0 \quad (4.35)$$

These give the uniform bound of Q_n^1, Γ_n and \dot{H}_n . Since $|V^k| \leq V^0, k = 1, 2, 3$, we deduce that $\rho_n, P_{k;n}$ are bounded and therefore $|S_{12;n}|$ and $|S_{13;n}|$ are bounded.

Using previous steps, we deduce the existence of $C(t) > 0$ such that

$$\begin{cases} \|\mu_n\|, \|R_n\|, \|A_n\|, \|\phi_n\|, \|\tau_n\|, \|\check{\mu}_n\|, \|\underline{\mu}_n\|, \|p_{1,n}\|, \|p_{2,n}\|, \|p_{3,n}\|, \|\rho_n\|, \\ \|\check{A}_n\|, \|\check{R}_n\|, \|\check{A}_n\|, \|\check{\phi}_n\|, \|\check{\tau}_n\|, \|\check{\phi}_n\|, \|\check{\tau}_n\|, \|S_{21,n}\|, \|S_{23,n}\|, \|S_{31,n}\| \leq C(t) \end{cases}$$

Proposition 4.1 *Let $[t_2; t_0] \subset]0; t_0]$, be an arbitrary compact subset on which the previous estimates hold. Then on such an interval, the iterates and their derivatives converge uniformly for L^∞ -norm.*

Proof 4.1 *Let $t \in [t_2; t_0]$, we use (4.5) – (4.6) to obtain respectively*

$$\begin{aligned} X_{n+1} &- X_n = \frac{1}{2} (R_{n+1} - R_n) (\check{\mu}_{n+1}^2 + \underline{\mu}_{n+1}^2 + \check{\phi}_{n+1}^2 + \underline{\phi}_{n+1}^2) \\ &+ \frac{1}{2} R_n [(\check{\mu}_{n+1} - \check{\mu}_n)(\check{\mu}_{n+1} + \check{\mu}_n) + (\check{\phi}_{n+1} - \check{\phi}_n)(\check{\phi}_{n+1} + \check{\phi}_n)] \\ &+ \frac{1}{2} R_n [(\underline{\mu}_{n+1} - \underline{\mu}_n)(\underline{\mu}_{n+1} + \underline{\mu}_n) + (\underline{\phi}_{n+1} - \underline{\phi}_n)(\underline{\phi}_{n+1} + \underline{\phi}_n)] \\ &+ \frac{e^{4\mu_{n+1}}}{8R_{n+1}} [(\check{A}_{n+1} - \check{A}_n)(\check{A}_{n+1} + \check{A}_n) + (\underline{A}_{n+1} - \underline{A}_n)(\underline{A}_{n+1} + \underline{A}_n)] \\ &+ \left(\frac{e^{4\mu_{n+1}}}{8R_{n+1}} - \frac{e^{4\mu_n}}{8R_n} \right) (\check{A}_n^2 + \underline{A}_n^2) + \frac{1}{2} (\phi_{n+1} - \phi_n)(\phi_{n+1} + \phi_n) \quad (4.36) \end{aligned}$$

and since $\partial_\tau = \frac{1}{\sqrt{2}}(\partial_t + \partial_\theta)$ and $\partial_\lambda = \frac{1}{\sqrt{2}}(\partial_t - \partial_\theta)$,

$$\begin{aligned} X_{n+1} &- X_n = (R_{n+1} - R_n) (\hat{\mu}_{n+1}^2 + \check{\mu}_{n+1}^2 + \hat{\phi}_{n+1}^2 + \check{\phi}_{n+1}^2) \\ &+ R_n [(\hat{\mu}_{n+1} - \hat{\mu}_n)(\hat{\mu}_{n+1} + \hat{\mu}_n) + (\hat{\phi}_{n+1} - \hat{\phi}_n)(\hat{\phi}_{n+1} + \hat{\phi}_n)] \\ &+ \frac{1}{2} R_n [(\check{\mu}_{n+1} - \check{\mu}_n)(\check{\mu}_{n+1} + \check{\mu}_n) + (\check{\phi}_{n+1} - \check{\phi}_n)(\check{\phi}_{n+1} + \check{\phi}_n)] \\ &+ \frac{e^{4\mu_{n+1}}}{4R_{n+1}} [(\hat{A}_{n+1} - \hat{A}_n)(\hat{A}_{n+1} + \hat{A}_n) + (\check{A}_{n+1} - \check{A}_n)(\check{A}_{n+1} + \check{A}_n)] \\ &+ \left(\frac{e^{4\mu_{n+1}}}{8R_{n+1}} - \frac{e^{4\mu_n}}{4R_n} \right) (\hat{A}_n^2 + \check{A}_n^2) + \frac{1}{2} (\phi_{n+1} - \phi_n)(\phi_{n+1} + \phi_n) \quad (4.37) \end{aligned}$$

Let now define $W_{n+1}(t) = (X_{n+1} - X_n)(t) + (Y_{n+1} - Y_n)(t)$ and $\tilde{W}_{n+1}(t) = (X_{n+1} - X_n)(t) - (Y_{n+1} - Y_n)(t)$. We deduce from (4.37) that

$$\begin{aligned} \partial_\lambda W_{n+1} &= \partial_\lambda[(X_{n+1} - X_n) + (Y_{n+1} - Y_n)] = \chi_n - \chi_{n-1} - (\check{\mu}_n - \check{\mu}_{n-1})(\zeta_n - \zeta_{n-1}) \\ &+ \left(\frac{e^{4\mu_n}}{2R_n} - \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \right) (\check{A}_n - \check{A}_{n-1})(\zeta_n - \zeta_{n-1}) + F_n \end{aligned} \quad (4.38)$$

$$\begin{aligned} \partial_\sigma \tilde{W}_{n+1} &= \partial_\sigma[(X_{n+1} - X_n) - (Y_{n+1} - Y_n)] = \tilde{\chi}_n - \tilde{\chi}_{n-1} - (\hat{\mu}_n - \hat{\mu}_{n-1})(\tilde{\zeta}_n - \tilde{\zeta}_{n-1}) \\ &+ \left(\frac{e^{4\mu_n}}{2R_n} - \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \right) (\hat{A}_n - \hat{A}_{n-1})(\tilde{\zeta}_n - \tilde{\zeta}_{n-1}) + \tilde{F}_n. \end{aligned} \quad (4.39)$$

where

$$\begin{aligned} F_n &= -\hat{\mu}_n \zeta_{n-1} - \hat{\mu}_{n-1} \zeta_n - 2\hat{\mu}_{n-1} \zeta_{n-1} + \frac{e^{4\mu_n}}{2R_n} \hat{A}_{n-1} \tilde{\zeta}_n \\ &+ \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \hat{A}_n \tilde{\zeta}_n + \frac{e^{4\mu_n}}{2R_n} \hat{A}_{n-1} \tilde{\zeta}_{n-1} \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \hat{A}_n \tilde{\zeta}_n + \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \hat{A}_n \tilde{\zeta}_{n-1} \end{aligned} \quad (4.40)$$

$$\begin{aligned} \tilde{F}_n &= -\check{\mu}_n \zeta_{n-1} - \check{\mu}_{n-1} \zeta_n - 2\check{\mu}_{n-1} \zeta_{n-1} + \frac{e^{4\mu_n}}{2R_n} \check{A}_{n-1} \tilde{\zeta}_n \\ &+ \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \check{A}_n \tilde{\zeta}_n + \frac{e^{4\mu_n}}{2R_n} \check{A}_{n-1} \tilde{\zeta}_{n-1} \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \check{A}_n \tilde{\zeta}_n + \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \check{A}_n \tilde{\zeta}_{n-1} \end{aligned} \quad (4.41)$$

Integrating (4.38) – (4.39) along null paths between t_1 and t_0 on the characteristics map $t \mapsto (t; \gamma^\pm(t))$ where $\gamma^\pm(t) = \theta \pm (t_0 - t)$ for all $(t; \theta) \in [t_1; t_0] \times S^1$, and after adding, we obtain

$$\begin{aligned} (X_{n+1} - X_n)(t_1, \theta) &= \frac{1}{4} [W_{n+1}(t_0, \gamma^-(t_1)) + \tilde{W}_{n+1}(t_0, \gamma^+(t_1))] \\ &- \frac{1}{4} \int_{t_1}^{t_0} [(\chi_n - \chi_{n-1})(s, \gamma^-(s)) + (\tilde{\chi}_n - \tilde{\chi}_{n-1})(s, \gamma^+(s))] ds \\ &- \frac{1}{4} \int_{t_1}^{t_0} [(\hat{\mu}_n - \hat{\mu}_{n-1})(\zeta_n - \zeta_{n-1})(t_0, \gamma^-(s))] ds \\ &- \frac{1}{4} \int_{t_1}^{t_0} [(\check{\mu}_n - \check{\mu}_{n-1})(\zeta_n - \zeta_{n-1})(t_0, \gamma^+(s))] ds \\ &- \frac{1}{4} \int_{t_1}^{t_0} \left[\left(\frac{e^{4\mu_n}}{2R_n} - \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \right) (\hat{A}_n - \hat{A}_{n-1})(\tilde{\zeta}_n - \tilde{\zeta}_{n-1})(t_0, \gamma^-(s)) \right] ds \\ &- \frac{1}{4} \int_{t_1}^{t_0} \left[\left(\frac{e^{4\mu_n}}{2R_n} - \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \right) (\check{A}_n - \check{A}_{n-1})(\tilde{\zeta}_n - \tilde{\zeta}_{n-1})(t_0, \gamma^+(s)) \right] ds \\ &+ \frac{1}{4} \int_{t_1}^{t_0} [F_n(t_0, \gamma^+(s)) + F_n(t_0, \gamma^-(s))] ds \end{aligned} \quad (4.42)$$

From (4.1) we have

$$\begin{aligned}
\frac{d}{ds}((\Theta, V)_{n+1} - (\Theta, V)_n)(s, t, \theta, v) &= (\Xi_n - \Xi_{n-1})(s, t, \theta, v) \\
&\leq C |X_n - X_{n-1}|(s, t, \theta) + (|F_n| + |\tilde{F}_n|)(s, \theta) \\
|(\Theta, V)_{n+1} - (\Theta, V)_n|(t_0, t, \theta, v) &\leq |(\Theta, V)_{n+1} - (\Theta, V)_n|(t, t, \theta, v) \\
&+ C \int_t^{t_0} (|X_n - X_{n-1}| + |F_n| + |\tilde{F}_n|)(s) ds
\end{aligned}$$

and from (4.2) we have

$$\begin{aligned}
|(f_{n+1} - f_n)(t)| &\leq \|f_0\| |(\Theta, V)_{n+1} - (\Theta, V)_n|(t_0, t, \theta, v) \\
&\leq C \|f_0\| \left[\int_t^{t_0} |\Xi_n - \Xi_{n-1}|(s) ds + \int_t^{t_0} (|F_n| + |\tilde{F}_n|)(s) ds \right]
\end{aligned}$$

This implies

$$\|\rho_{n+1} - \rho_n\| \leq C |W_n| + C \int_t^{t_0} |X_n - X_{n-1}| ds \quad (4.43)$$

Since $|P_{k;n}|, |S_{k;n}|, |J_{k;n}| \leq |\rho_n|$, we deduce from (4.43) that :

$$\|P_{k;n+1} - P_{k;n}\|; \|J_{k;n+1} - J_{k;n}\|; \|S_{j_k;n+1} - S_{j_k;n}\| \leq C \left[|W_n| + \int_t^{t_0} |X_n - X_{n-1}| ds \right] \quad (4.44)$$

We proceed in the same way as we did for (3.17) and (4.37) to obtain respectively

$$\left| \int_{t_1}^{t_0} (\hat{\mu}_n - \hat{\mu}_{n-1})(\zeta_n - \zeta_{n-1})(s, \gamma^+(s)) ds \right| \leq C \sup_{t \in [t_1, t_0]} \sqrt{|X_n - X_{n-1}|(t, \theta)} \quad (4.45)$$

$$\left| \int_{t_1}^{t_0} (\check{\mu}_n - \check{\mu}_{n-1})(\zeta_n - \zeta_{n-1})(s, \gamma^-(s)) ds \right| \leq C \sup_{t \in [t_1, t_0]} \sqrt{|X_n - X_{n-1}|(t, \theta)} \quad (4.46)$$

$$\begin{aligned}
\int_{t_1}^{t_0} \left[\frac{e^{4\mu_n}}{2R_n} - \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \right] |\hat{A}_n - \hat{A}_{n-1}| |\tilde{\zeta}_n - \tilde{\zeta}_{n-1}|(s, \gamma^+(s)) ds &\leq \\
C \sup_{t \in [t_1, t_0]} |X_n - X_{n-1}|(t, \theta) &\quad (4.47)
\end{aligned}$$

$$\begin{aligned}
\int_{t_1}^{t_0} \left[\frac{e^{4\mu_n}}{2R_n} - \frac{e^{4\mu_{n-1}}}{2R_{n-1}} \right] |\check{A}_n - \check{A}_{n-1}| |\tilde{\zeta}_n - \tilde{\zeta}_{n-1}|(s, \gamma^-(s)) ds &\leq \\
C \sup_{t \in [t_1, t_0]} |X_n - X_{n-1}|(t, \theta) &\quad (4.48)
\end{aligned}$$

Using (4.12) – (4.13), we have respectively

$$\begin{aligned}
\partial_\lambda [R_n e^{-2\tau_n + 4\mu_n} (\Gamma_n - \Gamma_{n-1}) + \Gamma_{n-1} (R_n e^{-2\tau_n + 4\mu_n} - R_{n-1} e^{-2\tau_{n-1} + 4\mu_{n-1}})] &= \\
-2R_{n-1} e^{\tau_{n-1}} (J_{2,n} - J_{2,n-1} + S_{12,n-1} - S_{12,n}) - (J_{2,n-1} - S_{12,n-1}) (R_n e^{\tau_n} - R_{n-1} e^{\tau_{n-1}}) &\quad (4.49)
\end{aligned}$$

$$\begin{aligned}
\partial_\sigma [R_n e^{-2\tau_n + 4\mu_n} (\Gamma_n - \Gamma_{n-1}) + \Gamma_{n-1} (R_n e^{-2\tau_n + 4\mu_n} - R_{n-1} e^{-2\tau_{n-1} + 4\mu_{n-1}})] &= \\
-2R_{n-1} e^{\tau_{n-1}} (J_{2,n} - J_{2,n-1} + S_{12,n} - S_{12,n-1}) - (J_{2,n-1} + S_{12,n-1}) (R_n e^{\tau_n} - R_{n-1} e^{\tau_{n-1}}) &\quad (4.50)
\end{aligned}$$

Add (4.49) – (4.50) to obtain

$$\begin{aligned} & \partial_t [R_n e^{-2\tau_n + 4\mu_n} (\Gamma_n - \Gamma_{n-1}) + \Gamma_{n-1} (R_n e^{-2\tau_n + 4\mu_n} - R_{n-1} e^{-2\tau_{n-1} + 4\mu_{n-1}})] = \\ & -2R_{n-1} e^{\tau_{n-1}} (J_{2,n} - J_{2,n-1} + S_{12,n} - S_{12,n-1}) - J_{2,n-1} (R_n e^{\tau_n} - R_{n-1} e^{\tau_{n-1}}) \quad (4.51) \end{aligned}$$

Integrating with respect to t and obtain

$$\begin{aligned} & |R_n e^{-2\tau_n + 4\mu_n} (\Gamma_n - \Gamma_{n-1}) + \Gamma_{n-1} (R_n e^{-2\tau_n + 4\mu_n} - R_{n-1} e^{-2\tau_{n-1} + 4\mu_{n-1}})|(t, \theta) \leq \\ & \int_t^{t_0} [2R_{n-1} e^{\tau_{n-1}} (|J_{2,n} - J_{2,n-1}| + |S_{12,n} - S_{12,n-1}|) + J_{2,n-1} |R_n e^{\tau_n} - R_{n-1} e^{\tau_{n-1}}|] (s, \theta) ds \quad (4.52) \end{aligned}$$

We deduce from (4.44) that (4.52) gives

$$|\Gamma_n - \Gamma_{n-1}| \leq C \int_{t_1}^{t_0} |X_n - X_{n-1}|(s) ds \quad (4.53)$$

From (4.26) – (4.27), we obtain

$$\left| \int_{t_1}^{t_0} [\chi_n - \chi_{n-1}](s, \theta + s - t) ds \right| \leq C \int_{t_1}^{t_0} |X_n - X_{n-1}|(s, \theta) ds \quad (4.54)$$

$$\left| \int_{t_1}^{t_0} [\tilde{\chi}_n - \tilde{\chi}_{n-1}](s, \theta + s - t) ds \right| \leq C \int_{t_1}^{t_0} |X_n - X_{n-1}|(s, \theta) ds \quad (4.55)$$

Replacing (4.45) – (4.46) – (4.47) – (4.48) – (4.54) – (4.55), in (4.42) gives

$$\sup_{\theta \in S^1} |X_{n+1} - X_n|(t_1, \theta) \leq C \int_{t_1}^{t_0} \sup_{\theta \in S^1} |X_n - X_{n-1}|(s, \theta) ds \quad (4.56)$$

Therefore

$$\sup_{\theta \in S^1} |X_{n+1} - X_n|(t_1, \theta) \leq \left| C^{n+2} \frac{(t_0 - t_1)^{n+1}}{(n+1)!} \right| \text{ for } n \in \mathbb{N}, t \in [t_1; t_0] \quad (4.57)$$

Since the series $\left\{ \frac{(t_0 - t_1)^{n+1}}{(n+1)!} \right\}_n$ converges, we deduce the convergence of $\left(\frac{(t_0 - t_1)^{n+1}}{(n+1)!} \right)_{n \in \mathbb{N}}$

to zero. This implies that $(X_n)_{n \in \mathbb{N}}$ is a Cauchy sequence.

Otherwise, (3.15) and (3.16) imply respectively

$$\begin{aligned} \zeta_{n+1} - \zeta_n &= e^{2(\tau_{n+1} - \mu_{n+1})} [\rho_{n+1} - \rho_n - (P_{1,n+1} - P_{1,n}) + P_{2,n+1} - P_{2,n} - (P_{3,n+1} - P_{3,n})] \\ &+ [(e^{\tau_{n+1}} - e^{\tau_n})(e^{\tau_{n+1}} + e^{\tau_n})e^{-2\mu_{n+1}} + e^{2\tau_n}(e^{-2\mu_{n+1}} - e^{-2\mu_n})] [\rho_n - P_{1,n} + P_{2,n} - P_{3,n}] \\ &+ \frac{R_{n+1}}{2} e^{-2\tau_{n+1} + 4\mu_{n+1}} \Gamma_{n+1}^2 - \frac{R_n}{2} e^{-2\tau_n + 4\mu_n} \Gamma_n^2 \quad (4.58) \end{aligned}$$

$$\begin{aligned} \tilde{\zeta}_{n+1} - \tilde{\zeta}_n &= R_{n+1}^2 e^{-2(\tau_{n+1} + \mu_{n+1})} \{ \Gamma_{n+1} (\check{H}_{n+1} - \check{H}_n) + \check{H}_n (\Gamma_{n+1} - \Gamma_n) \} \\ &+ 2R_{n+1}^2 e^{-2(\tau_{n+1} + \mu_{n+1})} (S_{23,n+1} - S_{23,n}) + 2S_{23,n} (R_{n+1}^2 e^{-2(\tau_{n+1} + \mu_{n+1})} \\ &- R_n^2 e^{-2(\tau_n + \mu_n)}) + \check{H}_n \Gamma_n (R_{n+1}^2 e^{-2(\tau_{n+1} + \mu_{n+1})} - R_n^2 e^{-2(\tau_n + \mu_n)}) \quad (4.59) \end{aligned}$$

$$\text{with } \Gamma_{n+1} - \Gamma_n = (\check{G}_{n+1} - \check{G}_n) + (A_{n+1}\check{H}_{n+1} - A_n\check{H}_n) \quad (4.60)$$

Using (4.53), (4.60) gives

$$\|\underline{G}_{n+1} - \underline{G}_n\|, \|\underline{H}_{n+1} - \underline{H}_n\| \leq C\|X_{n+1} - X_n\| \quad (4.61)$$

From (4.36) we deduce that

$$\|R_{n+1} - R_n\|, \|\check{\mu}_{n+1} - \check{\mu}_n\|, \|\check{A}_{n+1} - \check{A}_n\|, \|\phi_{n+1} - \phi_n\| \leq C\|X_{n+1} - X_n\| \quad (4.62)$$

From (4.58), (4.59), (4.60), (4.43), (4.44) we have

$$\|S_{23,n+1} - S_{23,n}\|, \|\rho_{n+1} - \rho_n\|, \|P_{1,n+1} - P_{1,n}\|, \|P_{2,n+1} - P_{2,n}\|, \|P_{3,n+1} - P_{3,n}\| \leq C\|X_{n+1} - X_n\| \quad (4.63)$$

Relation (4.29) imply

$$\begin{aligned} |\hat{\tau}_{n+1} - \hat{\tau}_n| &= |\hat{\mu}_n\check{\mu}_n - \hat{\mu}_{n-1}\check{\mu}_{n-1} + \frac{e^{4\mu_n}}{4R_n^2}\hat{A}_n\check{A}_n - \frac{e^{4\mu_{n-1}}}{4R_{n-1}^2}\hat{A}_{n-1}\check{A}_{n-1} \\ &+ \frac{R_{n-1}}{4}e^{-2\tau_{n-1}+4\mu_{n-1}}\Gamma_{n-1}^2 - \frac{R_n}{4}e^{-2\tau_n+4\mu_n}\Gamma_n^2 + \frac{3R_{n-1}^2e^{-2\tau_{n-1}}}{4}\check{H}_{n-1}^2 \\ &- \frac{3R_n^2e^{-2\tau_n}}{4}\check{H}_n^2 + e^{2(\tau_{n-1}-\mu_{n-1})}P_{3,n} - e^{2(\tau_n-\mu_n)}P_{3,n+1}| \end{aligned}$$

and we deduce from (4.37) and (4.44)-(4.48) that

$$|\hat{\tau}_{n+1} - \hat{\tau}_n| \leq C(\|X_{n+1} - X_n\| + |P_{3,n+1} - P_{3,n}|) \quad (4.64)$$

Integrating along null paths starting at (t_1, θ) and ending at the initial t_0 -surface, we obtain from (4.62) – (4.63) that

$$\|\tau_{n+1} - \tau_n\| \leq C\|X_{n+1} - X_n\| \quad (4.65)$$

We have (where ν is respectively replaced by μ or A):

$$\nu_{n+1}(t, \theta) - \nu_n(t, \theta) = \int_t^{t_0} (\underline{\nu}_{n+1}(s, \theta) - \underline{\nu}_n(s, \theta)) ds \quad (4.66)$$

Using (4.36), we have:

$$|\nu_{n+1}(t, \theta) - \nu_n(t, \theta)| = \left| \int_t^{t_0} (\underline{\nu}_{n+1}(s, \theta) - \underline{\nu}_n(s, \theta)) ds \right| \leq C \int_t^{t_0} |X_{n+1}(s, \theta) - X_n(s, \theta)| ds$$

This implies that

$$|A_{n+1}(t, \theta) - A_n(t, \theta)|, |\mu_{n+1}(t, \theta) - \mu_n(t, \theta)| \leq C \int_t^{t_0} |X_{n+1}(s, \theta) - X_n(s, \theta)| ds \quad (4.67)$$

Otherwise, from (4.61), we have respectively:

$$|G_{n+1}(t, \theta) - G_n(t, \theta)| = \left| \int_t^{t_0} (\underline{G}_{n+1} - \underline{G}_n)(s, \theta) ds \right| \leq C \left| \int_t^{t_0} (X_{n+1}(s, \theta) - X_n(s, \theta)) ds \right| \quad (4.68)$$

$$|H_{n+1}(t, \theta) - H_n(t, \theta)| = \left| \int_t^{t_0} (\underline{H}_{n+1}(s, \theta) - \underline{H}_n(s, \theta)) ds \right| \leq C \left| \int_t^{t_0} (X_{n+1}(s, \theta) - X_n(s, \theta)) ds \right|. \quad (4.69)$$

Using (4.57), (4.63), (4.65), (4.62), (4.67), (4.68), (4.69), we deduce the uniform convergence for L^∞ norm of the iterates. Since all the iterates are respectively bounded with their derivatives, we deduce that there exist functions $\mu, R, A, \phi, \tau, H, G$ and f such that

$$\mu_n \rightarrow \mu, R_n \rightarrow R, A_n \rightarrow A, \phi_n \rightarrow \phi, \tau_n \rightarrow \tau, H_n \rightarrow H, G_n \rightarrow G, f_n \rightarrow f \quad (4.70)$$

and

$$\mu'_n \rightarrow \mu', R'_n \rightarrow R', A'_n \rightarrow A', \phi'_n \rightarrow \phi', \tau'_n \rightarrow \tau', H'_n \rightarrow H', G'_n \rightarrow G'. \quad (4.71)$$

$$\dot{\mu}_n \rightarrow \dot{\mu}, \dot{R}_n \rightarrow \dot{R}, \dot{A}_n \rightarrow \dot{A}, \dot{\phi}_n \rightarrow \dot{\phi}, \dot{\tau}_n \rightarrow \dot{\tau}, \dot{H}_n \rightarrow \dot{H}, \dot{G}_n \rightarrow \dot{G}; \quad (4.72)$$

where "dot" and "prime" represent respectively partial derivative with respect to t and θ . Therefore $(f, R, \tau, \mu, A, H, G, \phi)$ satisfy the complete Cauchy problem.

Let us now prove the uniqueness of the solution.

Let $\chi_k = (f_k; R_k; \mu_k; A_k; \tau_k; G_k; H_k; \phi_k)_{k=1,2}$ be two regular solutions of the Cauchy problem (2.3) – (2.4) – (2.5), (2.6) – (2.10) for the same initial data $(\overset{\circ}{f}, \overset{\circ}{R}, \overset{\circ}{\tau}, \overset{\circ}{\mu}, \overset{\circ}{A}, \overset{\circ}{H}, \overset{\circ}{G}, \overset{\circ}{\phi}, \overset{\circ}{R}, \overset{\circ}{\tau}, \overset{\circ}{\mu}, \overset{\circ}{A}, \overset{\circ}{H}, \overset{\circ}{G}, \psi)$ at $t = t_0$. Using the fact that χ_k is a solution of the complete system, one proceeds similarly to prove the convergence of the iterates to obtain

$$\beta(t) \leq C \int_t^{t_0} \beta(s) ds$$

where

$$\beta(t) = \sup\{\|f_2(s) - f_1(s)\| + \|R_2(s) - R_1(s)\| + \|\mu_2(s) - \mu_1(s)\| + \|A_2(s) - A_1(s)\| + \|\tau_2(s) - \tau_1(s)\| + \|G_2(s) - G_1(s)\| + \|H_2(s) - H_1(s)\| + \|\phi_2(s) - \phi_1(s)\|; s \in [t, t_0]\}.$$

We deduce that $\beta(t) = 0$ for $t \in]0, t_0]$. This implies that $f_2 = f_1; R_2 = R_1; \mu_2 = \mu_1; A_2 = A_1; \tau_2 = \tau_1; G_2 = G_1; H_2 = H_1$ and $\phi_2 = \phi_1$. We have proved the following result:

Theorem 4.1 (local existence) *Let $\overset{\circ}{f} \in C^1(S^1 \times \mathbb{R}^3)$ with $\overset{\circ}{f} \geq 0$, $\overset{\circ}{f}(\theta + 2\pi, v) = \overset{\circ}{f}(\theta, v)$ for $(\theta, v) \in S^1 \times \mathbb{R}^2$ and*

$$W_0 := \sup\{v | (\theta, v) \in \text{supp } \overset{\circ}{f}\} < \infty$$

Let given regular functions $\bar{R}, \bar{\tau}, \bar{\mu}, \bar{A}, \bar{H}, \bar{G}, \psi \in C^1(\mathbb{R})$ and $\overset{\circ}{R}, \overset{\circ}{\tau}, \overset{\circ}{\mu}, \overset{\circ}{A}, \overset{\circ}{H}, \overset{\circ}{G}, \overset{\circ}{\phi} \in C^2(S^1)$. Then there exists a unique left maximal regular solution $(f, R, \tau, \mu, A, H, G, \phi)$ of the complete EVSFS with $(f, R, \tau, \mu, A, H, G, \phi)(t_0) = (\overset{\circ}{f}, \overset{\circ}{R}, \overset{\circ}{\tau}, \overset{\circ}{\mu}, \overset{\circ}{A}, \overset{\circ}{H}, \overset{\circ}{G}, \overset{\circ}{\phi})$ and $(\dot{R}, \dot{\tau}, \dot{\mu}, \dot{A}, \dot{H}, \dot{G}, \dot{\phi})(t_0) = (\bar{R}, \bar{\tau}, \bar{\mu}, \bar{A}, \bar{H}, \bar{G}, \psi)$ on a time interval $]T, t_0]$ with $T \in [0, t_0[$.

Remark 4.1 *The result we have obtained here proves particularly the local in time existence of solution in Gowdy case with scalar field or not in passed time direction. It generalizes also the T^2 symmetry case without scalar field. It remains using global existence result obtained in [7], looking for stability and asymptotic behaviour of that solution ([8], [5]). Another preoccupation will be focused on what happens in future direction (cf [6]).*

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- [1] H. Andreasson, Global foliations of matter spacetimes with Gowdy symmetry. *Comm. Math. Phys.* 206 (1999), pp.337-365.
- [2] H. Andreasson, A. D. Rendall, and M. Weaver, Existence of CMC and constant areal time foliations in T^2 symmetric spacetimes with Vlasov matter, *Comm. Partial Differential Equations*, 29, N^0 1-2, pp. 237-262 (2004).
- [3] B.K. Berger, P. Chrusciel, J. Isenberg and V. Moncrief, Global foliations of vacuum spacetimes with T^2 isometry, *Ann. Phys.* 260, 117-148 (1997).
- [4] Y. Choquet-Bruhat, Problème de Cauchy pour le système intégral différentiel d'Einstein-Liouville, *Ann. Inst. Fourier*, tome 21, N^0 3 pp. 181-201 (1971).
- [5] M. Dafermos and A.D. Rendall, Strong cosmic censorship for surface-symmetric cosmological spacetimes with collisionless matter, arXiv e-print, (2007). [gr-qc/0701034].
- [6] A.T. Lassiye, Equations d'Einstein-Vlasov en symétrie T^2 avec champ scalaire non linéaire. Thèse de Doctorat/Ph.D en Mathématiques, Université de Yaounde 1 (2020).
- [7] A.T. Lassiye, D. Tegankong, N. Noutchequeme, The Einstein-Vlasov-Scalar field system with Gowdy or T^2 symmetry in contracting direction. *Acta Math Vietnam* (2018) Vol 44 pp. 501-518.
- [8] H. Ringström, Future stability of the Einstein non-linear scalar field system, *Invent. Math.* 173, 123-208, (2008).

- [9] J. Smulevici, On the area of the symmetry orbits of cosmological spacetimes with toroidal or hyperbolic symmetry. *Analysis & PDE*. Vol 4, N^0 2, 191-245 (2011).
- [10] D. Tegankong, N. Noutchequeme and A.D. Rendall , Local existence and continuation criteria for solutions of the Einstein-Vlasov-scalar field system with surface symmetry *Journ. Hyperb. Diff. Eq.* Vol 1, No.4, (2004) 691-624.
- [11] M. Weaver, On the area of the symmetry orbits in T^2 symmetric spacetimes with Vlasov matter, *Classical Quantum Gravity*, 21 (2004), pp.1079-1097.