

Energy-momentum squared symmetric Teleparallel gravity: $f(Q, T_{\mu\nu}T^{\mu\nu})$ gravity

Prabir Rudra^a

^a*Department of Mathematics, Asutosh College, Kolkata-700 026, India*

E-mail: prudra.math@gmail.com, rudra@associates.iucaa.in

ABSTRACT: In this work we propose the $f(Q, T_{\mu\nu}T^{\mu\nu})$ gravity as a further extension of the $f(Q)$ and $f(Q, T)$ gravity theories. The action involves an arbitrary function of the non-metricity Q and $T_{\mu\nu}T^{\mu\nu}$ in the gravity Lagrangian. The field equations for the theory are derived in the metric formalism by varying the action with respect to the metric. The theory involves a non-minimal coupling between the geometric and the matter sectors, and hence the covariant divergence of the energy momentum tensor is non-zero, thus implying the non-conservation of the same. The vacuum solutions of the theory are investigated and it is found that the theory perfectly admits a de-Sitter like evolution of the universe. The cosmological equations are derived and it is found that there are two correction terms arising as modification of the gravity. Two specific toy models of the form $Q + \eta (\mathbf{T}^2)^n$ and $f_0 Q^m (\mathbf{T}^2)^n$ are explored to gain further insights into the dynamics of the theory. It is seen that the field equations of both the models have terms similar to those arising from the quantum gravity effects and are thus responsible for the avoidance of the singularity. One striking feature of the model is that the non-linear correction terms dominate in the early universe and gradually fade away at later times giving standard FLRW universe. Solutions for the FLRW equations are found wherever possible and the evolution of the scale factor and the matter energy density are plotted. Other cosmological parameters like the equation of state and deceleration parameters are also studied. Finally the energy conditions are explored in the background of the theory. Using these conditions and some observational data the theory is considerably constrained. $f(Q, T_{\mu\nu}T^{\mu\nu})$ is a theory that can perfectly explain the cosmological dynamics of both the early and the late universe without resorting to any dark energy.

KEYWORDS: non-metricity, energy-momentum, symmetric teleparallel gravity; cosmology, energy conditions.

Contents

1	Introduction	1
2	Field equations of $f(Q, T_{\mu\nu}T^{\mu\nu})$ Gravity	4
2.1	Conservation of matter energy-momentum tensor	6
2.2	Matter as perfect fluid	7
3	Cosmology of EMSSTG	8
3.1	The de-Sitter solution	9
4	Some specific Toy-models	10
4.1	Model: 1	10
4.2	Model: 2	11
5	Energy Conditions	14
5.1	Model: 1	15
5.2	Model: 2	16
6	Conclusion	17
7	Appendix	18
7.1	Expressions for ζ and ϑ in section 5.2	18

1 Introduction

Currently the most comprehensive theory of gravity available to us is the theory of General Relativity (GR) [1] proposed by Albert Einstein in 1916. After a century of extensive research GR has been able to survive quite well with many of its predictions coming true. Some of them being the perihelion precision of mercury, deflection of light by sun [2], gravitational redshift [3], detection of gravitational waves [4] from the mergers of black holes and neutron stars, imaging a black hole shadow in the M87 galaxy by the Event horizon telescope [5], etc. Extensive reviews in experimental and observational tests of GR can be found in [6, 7]. Nevertheless some observations did not go in favour of GR, quite expectedly. The most important one being the discovery of the late cosmic acceleration [8, 9] at turn of the last century. With this observation to light, GR became incompatible at cosmological distances. One more important observation was that GR was not able to explain the gravitational interaction at the quantum level. Moreover standard cosmology is plagued by problems like the singularity problem, cosmological constant problem, cosmic coincidence problem, etc. So it is clear that we are far from the theory that we need to comprehensively explain our observations. This provided motivation for extensive theoretical research in cosmology over the past century. With the development of quantum mechanics throughout the last century, scientists including Einstein took up the challenge of developing a proper theory of quantum gravity. As a result theories like string theory, loop quantum theory, etc. were proposed, but till date they are far from being comprehensive. To resolve the incompatibility of GR at cosmological scales mainstream research have travelled in two different ways. The first path is the theory of dark energy (DE) where the matter content of the universe is modelled by an exotic fluid with negative pressure driving the

acceleration. Extensive reviews on DE can be found in [10]. The alternative path is the theory of modified gravity, where the gravitational framework of GR has been modified to incorporate the cosmic acceleration. See [11–13] for extensive reviews on modified gravity. In this work we will concentrate on this alternative path of modified gravity theory.

The field equations of GR is basically formulated from the Einstein-Hilbert (EH) action $\mathcal{S} = \int (R + \mathcal{L}_m) \sqrt{-g} d^4x$, where R is the Ricci scalar, g is the determinant of the metric tensor ($g_{\mu\nu}$) that represents the gravitational field and \mathcal{L}_m is the matter Lagrangian. The most fundamental form of modification is imposed by replacing the gravitational Lagrangian R of EH action by an arbitrary function $f(R)$, thus considering a generalized action. This gave rise to $f(R)$ theories [14]. Substantive development in $f(R)$ gravity can be found in [15–20]. Comprehensive reviews on this theory can be found in Refs.[21, 22]. In similar fashion further extensions to the EH action was brought about by replacing the arbitrary function $f(R)$ by $f(R, \mathcal{L}_m)$ [23–26]. In ref.[27] the authors modified the EH action by including an arbitrary function $f(R, T)$, where T is the trace of the energy-momentum tensor (EMT). Here a coupling between the matter sector and the gravity sector was considered via the function $f(R, T)$. It was seen that the covariant divergence of the EMT is non-zero for this theory, which means non-conservation of EMT leading to non-geodesic motion of the massive test particles. The reason for this being the coupling effects between matter and geometry which induces extra acceleration on the particles. Further developments in $f(R, T)$ gravity can be found in [28–34]. Katirci and Kavuk in [35] proposed $f(R, \mathbf{T}^2)$ theory, where $\mathbf{T}^2 = T_{\mu\nu}T^{\mu\nu}$ and $T_{\mu\nu}$ is the EMT. Roshan and Shojai in [36] further developed the theory by investigating the properties of a specific form $R + \eta\mathbf{T}^2$ which was termed as energy-momentum squared gravity (EMSG) in the literature. From the field equations of EMSG it was seen that the equations have a flavour of the quantum geometry effects of loop quantum gravity [37], and allowed a maximum energy density ρ_{max} and a minimum length a_{min} in the early universe. As a result, in a homogeneous and isotropic spacetime this theory admits a cosmological bounce and avoids the existence of the early time singularity. Cosmological models in EMSG was studied by the authors in [38, 39]. An extensive dynamical system analysis in EMSG was presented by the authors of [40]. Observational constraints on EMSG using cosmic chronometers and supernova observations can be found in [41]. Thermodynamics of the apparent horizon in the generalized energymomentum-squared cosmology was studied in [42]. Other substantial developments in EMSG can be found in [43–48].

All the above extensions of GR have one feature common between them, which is the underlying Riemannian geometry [49] (formulated in the Riemann metrical space) lying at the heart of all these classical theories including GR. With the incompatibility of these theories at some scales, naturally a thought arises that if the underlying geometry can be replaced by a far more general geometric structure, then we may be able to remove some of the inconsistencies that has plagued these classical theories over the years. Such a novel attempt was undertaken by Weyl [50], where the main objective was the geometrical unification of gravity and electromagnetism. We know that in Riemannian geometry where the Levi-Civita connection is compatible to the metric and is the basic tool for length comparison between vectors. In Weyl’s theory there is a complete different mechanism where two connections are used, one bearing the information of the length of a vector and the other responsible for the direction of a vector during parallel transport. Physically the length connection is identified with the electromagnetic potential. The most striking feature of the theory being the non-zero covariant divergence of the metric tensor and this property induces a new geometrical quantity known as the *non-metricity*. Weyl’s geometry is a mathematical masterpiece with corresponding rich physical structure.

Since gravity is identified as the manifestation of the geometrical properties of spacetime, search for other developments in geometry continued. Based the works [51–54] Cartan proposed an extension of GR known as the Einstein-Cartan theory. A review on these theories may be found in [55]. The striking feature of Cartan’s geometry was the introduction of the *torsion field*, which

is interpreted as the spin density from the physical point of view [55]. This torsion field may be introduced in Weyl's geometric structure to give a natural extension Weyl-Cartan geometry, which can be an interesting mathematical structure with associated physical implications [56–59]. An extensive review on Riemann-Cartan and Weyl-Cartan geometries can be found in [60]. Another elegant geometrical formalism was given by Weitzenbock known as the Weitzenbock spaces [61], where the manifold is equipped with the properties $R^\alpha_{\beta\gamma\delta} = 0$, $T^\alpha_{\beta\gamma} \neq 0$, $\nabla_\mu g_{\alpha\beta} = 0$, where $R^\alpha_{\beta\gamma\delta}$ is the curvature tensor, $T^\alpha_{\beta\gamma}$ is the torsion tensor and $g_{\alpha\beta}$ is the metric tensor of the associated manifold. If the torsion tensor vanishes the Weitzenbock manifold reduces to the Euclidean manifold. Since the Weitzenbock space is curvature-less, the geometry derived from such spaces possess an important property of absolute parallelism or teleparallelism. This property of Weitzenbock spaces attracted Einstein, who applied this theory in physics by proposing a unified teleparallel theory of gravity and electromagnetism [62]. In teleparallel gravity the basic idea is to replace the metric $g_{\alpha\beta}$ by the tetrad vectors e^j_μ , which generates the torsion that is responsible for the gravitational effects. Since in this theory the concept of torsion exactly replaces the concept of curvature, it is termed as the teleparallel equivalent of general relativity (TEGR) [63–65]. One important thing to note in TEGR is that the spacetime in these theories are flat, due to the absence of curvature. Just like GR was extended to $f(R)$ theories, similarly TEGR was extended to $f(\tau)$ theories [66, 67], where τ is the scalar torsion. One significant advantage of $f(\tau)$ theories over $f(R)$ theories lies in the fact that the field equations for the former is of the second order in contrast to the fourth order field equations of the latter. More importantly the $f(\tau)$ theories have been successful in explaining various astrophysical processes and specifically the late cosmic acceleration without depending on any forms of exotic matter or dark energy [68–76]. Further extensions of the teleparallel framework was done via the Weyl-Cartan-Weitzenbock (WCW) theory [77, 78], where the extra condition of the exact cancellation of curvature by the torsion is introduced in the background of the Weyl-Cartan spacetime.

From the above discussions we have seen that there are two alternative formulations of GR: the first one with $R \neq 0, \tau = 0$ (curvature formulation), and the second one with $R = 0, \tau \neq 0$ (teleparallel formulation). However in both these formulations the non-metricity Q vanish. Geometrically Q represents the variation of length of a vector in parallel transport. Now in a third equivalent formalism of GR, a non-vanishing non-metricity Q was considered as the basic geometrical variable responsible for all forms of gravitational interactions. This theory was termed as the *symmetric teleparallel gravity* (STG) [79]. Here the Einstein pseudotensor plays the role of the energy-momentum density and in the geometric representation it finally becomes a true tensor. Further research saw the STG being extended to $f(Q)$ gravity [80], which is also known as the *coincident general relativity* and *non-metric gravity*. Cosmology of $f(Q)$ gravity and its observational constraints was studied in [81, 82]. Over the past decades there have been various developments in the framework of STG [83–87]. In ref.[87] the authors have proposed an extension of the $f(Q)$ gravity based on the non-minimal coupling between the non-metricity Q and the matter Lagrangian \mathcal{L}_m . Quite expectedly the non-minimal coupling between geometry and matter sectors results in the non-conservation of the energy-momentum tensor and appearance of an extra force in the geodesic equation of motion. Xu et. al in [88] developed another extension of the theory namely the $f(Q, T)$ gravity where the gravity Lagrangian is basically an arbitrary function of Q and the trace of the energy-momentum tensor T . The field equations were derived and the cosmological evolution of the model was studied. It was seen that in all the considered cases the theory supported an accelerated expansion of the universe ending with a de-Sitter type evolution.

In this work we are motivated to explore another extension of the symmetric teleparallel theory. Our basic aim is to extend $f(Q, T)$ gravity to $f(Q, \mathbf{T}^2)$ gravity, where $\mathbf{T}^2 = T_{\mu\nu}T^{\mu\nu}$, following the extension of $f(R, T)$ gravity to $f(R, \mathbf{T}^2)$ gravity. We will call $f(Q, \mathbf{T}^2)$ gravity as the Energy-momentum squared symmetric teleparallel gravity (EMSSTG). The gravitational action will be

constituted by an arbitrary function $f(Q, \mathbf{T}^2)$ of Q and \mathbf{T}^2 . Then varying the action with respect to the metric we can frame the field equations in a metric formalism. Using those equations we can study the cosmological evolution of the theory in detail. Some specific toy models may help in properly understanding the dynamics of the theory. We also intend to study the energy conditions in the background of the theory to put constraints on the model. The paper is organized as follows: The basic field equations of $f(Q, T_{\mu\nu}T^{\mu\nu})$ gravity is derived in section 2. Cosmological evolution of EMSSTG is explored in section 3. Section 4 is dedicated in studying the dynamics of some specific toy models. In section 5 we investigate the energy conditions in the background of EMSSTG thus constraining the theory. Finally the paper ends with some concluding remarks in section 6. In section 7, we provide an appendix showing some expressions that is not included in the main body of the paper.

2 Field equations of $f(Q, T_{\mu\nu}T^{\mu\nu})$ Gravity

In this section we will discuss the various geometrical preliminaries required to frame a relativistic theory of gravitation and go on to derive the field equations of $f(Q, T_{\mu\nu}T^{\mu\nu})$ gravity. We start by considering the action for $f(Q, T_{\mu\nu}T^{\mu\nu})$ gravity as,

$$\mathcal{S} = \int \left[\frac{1}{16\pi} f(Q, \mathbf{T}^2) + \mathcal{L}_m \right] \sqrt{-g} d^4x \quad (2.1)$$

Here $g \equiv \det(g_{\mu\nu})$ and $\mathbf{T}^2 = T_{\mu\nu}T^{\mu\nu}$, where $T_{\mu\nu}$ is the matter energy-momentum tensor. Moreover $f(Q, \mathbf{T}^2)$ is a function of the non-metricity Q and \mathbf{T}^2 and \mathcal{L}_m is the matter Lagrangian. The non-metricity tensor $Q_{\lambda\mu\nu}$ is defined as the covariant derivative of the metric tensor with respect to the Weyl-Cartan connection $\tilde{\Gamma}^\lambda_{\mu\nu}$ [55],

$$\nabla_\lambda g_{\mu\nu} = Q_{\lambda\mu\nu} = -\frac{\partial g_{\mu\nu}}{\partial x^\lambda} + g_{\nu\sigma} \tilde{\Gamma}^\sigma_{\mu\lambda} + g_{\sigma\mu} \tilde{\Gamma}^\sigma_{\nu\lambda} \quad (2.2)$$

The Weyl-Cartan connection $\tilde{\Gamma}^\lambda_{\mu\nu}$ can be written as a combination of three entities: the Christoffel symbol $\Gamma^\lambda_{\mu\nu}$, the contortion tensor $C^\lambda_{\mu\nu}$ and the disformation tensor $L^\lambda_{\mu\nu}$, in the following way,

$$\tilde{\Gamma}^\lambda_{\mu\nu} = \Gamma^\lambda_{\mu\nu} + C^\lambda_{\mu\nu} + L^\lambda_{\mu\nu} \quad (2.3)$$

In the above expression the first term, i.e. the Christoffel symbol is basically the Levi-Civita connection of the metric $g_{\mu\nu}$ given by,

$$\Gamma^\lambda_{\mu\nu} = \frac{1}{2} g^{\lambda\sigma} \left(\frac{\partial g_{\sigma\nu}}{\partial x^\mu} + \frac{\partial g_{\sigma\mu}}{\partial x^\nu} - \frac{\partial g_{\mu\nu}}{\partial x^\sigma} \right) \quad (2.4)$$

The contortion tensor $C^\lambda_{\mu\nu}$ is obtained from the torsion tensor $\tilde{\Gamma}^\lambda_{[\mu\nu]}$, which in turn is defined as,

$$\tilde{\Gamma}^\lambda_{[\mu\nu]} = \frac{1}{2} \left(\tilde{\Gamma}^\lambda_{\mu\nu} - \tilde{\Gamma}^\lambda_{\nu\mu} \right) \quad (2.5)$$

Using the above definition the contortion tensor is defined as,

$$C^\lambda_{\mu\nu} = \tilde{\Gamma}^\lambda_{[\mu\nu]} + g^{\lambda\sigma} g_{\mu\kappa} \tilde{\Gamma}^\kappa_{[\nu\sigma]} + g^{\lambda\sigma} g_{\nu\kappa} \tilde{\Gamma}^\kappa_{[\mu\sigma]} \quad (2.6)$$

The disformation tensor is given by,

$$L^\alpha_{\beta\gamma} = -\frac{1}{2} g^{\alpha\lambda} (\nabla_\gamma g_{\beta\lambda} + \nabla_\beta g_{\lambda\gamma} - \nabla_\lambda g_{\beta\gamma}) = -\frac{1}{2} g^{\alpha\lambda} (Q_{\gamma\beta\lambda} + Q_{\beta\lambda\gamma} - Q_{\lambda\beta\gamma}) \quad (2.7)$$

Using this the non-metricity is given as,

$$Q = -g^{\mu\nu} (L_{\beta\mu}^\alpha L_{\nu\alpha}^\beta - L_{\beta\alpha}^\alpha L_{\mu\nu}^\beta) \quad (2.8)$$

Moreover the Weyl-Cartan torsion tensor $\tau_{\mu\nu}^\lambda$ is defined as,

$$\tau_{\mu\nu}^\lambda = \frac{1}{2} (\tilde{\Gamma}_{\mu\nu}^\lambda - \tilde{\Gamma}_{\nu\mu}^\lambda) \quad (2.9)$$

and the Weyl-Cartan curvature tensor may be defined by,

$$\tilde{R}_{\mu\nu\sigma}^\lambda = \tilde{\Gamma}_{\mu\sigma,\nu}^\lambda - \tilde{\Gamma}_{\mu\nu,\sigma}^\lambda + \tilde{\Gamma}_{\mu\sigma}^\alpha \tilde{\Gamma}_{\alpha\nu}^\lambda - \tilde{\Gamma}_{\mu\nu}^\alpha \tilde{\Gamma}_{\alpha\sigma}^\lambda \quad (2.10)$$

We know that the trace of the energy-momentum tensor $T_{\mu\nu}$ is given as,

$$T = g^{\mu\nu} T_{\mu\nu} = g_{\mu\nu} T^{\mu\nu} \quad (2.11)$$

The trace of the non-metricity tensor is given by,

$$Q_\beta = Q_\beta{}^\lambda{}_\lambda, \quad \tilde{Q}_\beta = Q^\lambda{}_\beta{}_\lambda \quad (2.12)$$

We will bring in the superpotential of the model defined as,

$$\begin{aligned} P^\alpha{}_{\mu\nu} &= \frac{1}{4} \left[-Q^\alpha{}_{\mu\nu} + 2 Q_{(\mu}{}^\alpha{}_{\nu)} + Q^\alpha g_{\mu\nu} - \tilde{Q}^\alpha g_{\mu\nu} - \delta^\alpha_{(\mu} Q_{\nu)} \right] \\ &= -\frac{1}{2} L_{\mu\nu}^\alpha + \frac{1}{4} (Q^\alpha - \tilde{Q}^\alpha) g_{\mu\nu} - \frac{1}{4} \delta^\alpha_{(\mu} Q_{\nu)} \end{aligned} \quad (2.13)$$

Now we can use the non-metricity tensor with the superpotential to obtain the following relation for the non-metricity

$$Q = -Q_{\lambda\alpha\beta} P^{\lambda\alpha\beta} = -\frac{1}{4} \left(-Q^{\lambda\beta\rho} Q_{\lambda\beta\rho} + 2 Q^{\lambda\beta\rho} Q_{\rho\lambda\beta} - 2 Q^\rho \tilde{Q}_\rho + Q^\rho Q_\rho \right) \quad (2.14)$$

Now we proceed to derive the field equations. For this, we take the variation of the action \mathcal{S} in eqn.(2.1) with respect to the metric tensor and obtain,

$$\delta\mathcal{S} = \int \frac{1}{16\pi} \delta [f(Q, \mathbf{T}^2) \sqrt{-g}] d^4x + \int \delta [\mathcal{L}_m \sqrt{-g}] d^4x \quad (2.15)$$

Simplifying the above relation we get,

$$\delta\mathcal{S} = \int \frac{1}{16\pi} \left[-\frac{1}{2} g_{\mu\nu} \delta g^{\mu\nu} f(Q, \mathbf{T}^2) + f_Q \delta Q + f_{\mathbf{T}^2} \delta \mathbf{T}^2 - 8\pi T_{\mu\nu} \delta g^{\mu\nu} \right] \sqrt{-g} d^4x \quad (2.16)$$

where $f_Q = \frac{\partial f}{\partial Q}$ and $f_{\mathbf{T}^2} = \frac{\partial f}{\partial \mathbf{T}^2}$. Now in the above expression we have made use of the relations,

$$\delta(\sqrt{-g}) = -\frac{1}{2} \sqrt{-g} g_{\mu\nu} \delta g^{\mu\nu} \quad (2.17)$$

and

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g} \mathcal{L}_m)}{\delta g^{\mu\nu}} \quad (2.18)$$

Considering that \mathcal{L}_m depends only on the metric tensor and not on its derivatives, we get the relation,

$$T_{\mu\nu} = g_{\mu\nu} \mathcal{L}_m - 2 \frac{\partial \mathcal{L}_m}{\partial g^{\mu\nu}} \quad (2.19)$$

Now we see from eqn.(2.16) that we have to calculate the quantities δQ and $\delta \mathbf{T}^2$. After some rigorous calculations δQ is obtained as,

$$\delta Q = 2P_{\alpha\nu\rho}\nabla^\alpha\delta g^{\nu\rho} - (P_{\mu\alpha\beta}Q_\nu{}^{\alpha\beta} - 2Q_\mu{}^{\alpha\beta}P_{\alpha\beta\nu})\delta g^{\mu\nu} \quad (2.20)$$

For an explicit calculation of δQ see ref.[88]. Now let us define a quantity $\theta_{\mu\nu}$ as,

$$\theta_{\mu\nu} = \frac{\delta(T_{\alpha\beta}T^{\alpha\beta})}{\delta g^{\mu\nu}} \quad (2.21)$$

From the above relation we can write $\delta \mathbf{T}^2 = \delta(T_{\alpha\beta}T^{\alpha\beta}) = \theta_{\mu\nu}\delta g^{\mu\nu}$, where an explicit relation for $\theta_{\mu\nu}$ is calculated as,

$$\theta_{\mu\nu} = 2T_\mu^\alpha T_{\nu\alpha} - 2\mathcal{L}_m \left(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T \right) - TT_{\mu\nu} - 4T^{\alpha\beta} \frac{\delta^2 \mathcal{L}_m}{\delta g^{\mu\nu} \delta g^{\alpha\beta}} \quad (2.22)$$

Using all these in eqn.(2.16) we get the variation of the action as,

$$\begin{aligned} \delta \mathcal{S} = \int \frac{1}{16\pi} \left[-\frac{1}{2}g_{\mu\nu}\delta g^{\mu\nu} f(Q, \mathbf{T}^2) + f_Q \{ 2P_{\alpha\mu\nu}\nabla^\alpha\delta g^{\mu\nu} - (P_{\mu\alpha\beta}Q_\nu{}^{\alpha\beta} - 2Q_\mu{}^{\alpha\beta}P_{\alpha\beta\nu})\delta g^{\mu\nu} \} \right. \\ \left. + f_{\mathbf{T}^2}\theta_{\mu\nu}\delta g^{\mu\nu} - 8\pi T_{\mu\nu}\delta g^{\mu\nu} \right] \sqrt{-g}d^4x \end{aligned} \quad (2.23)$$

Now the term $2f_Q\sqrt{-g}P_{\alpha\mu\nu}\nabla^\alpha\delta g^{\mu\nu}$ on integration and with the use of boundary condition gives $-2\nabla^\alpha(f_Q\sqrt{-g}P_{\alpha\mu\nu})\delta g^{\mu\nu}$. Using this and equating the variation of the action to zero we finally obtain the field equations of $f(Q, \mathbf{T}^2)$ gravity as,

$$-\frac{2}{\sqrt{-g}}\nabla_\alpha(f_Q\sqrt{-g}P_{\mu\nu}^\alpha) - \frac{1}{2}f(Q, \mathbf{T}^2)g_{\mu\nu} + f_{\mathbf{T}^2}\theta_{\mu\nu} - f_Q(P_{\mu\alpha\beta}Q_\nu{}^{\alpha\beta} - 2Q_\mu{}^{\alpha\beta}P_{\alpha\beta\nu}) = 8\pi T_{\mu\nu} \quad (2.24)$$

For the special case $f(Q, \mathbf{T}^2) = f(Q)$, these field equations reduce to those of $f(Q)$ gravity. Moreover for the choice $f(Q, \mathbf{T}^2) = Q$ the above field equations reduce to the field equations of the symmetric teleparallel equivalent of general relativity [83–86, 89]. The (1, 1)- form of the field equations are given by,

$$f_{\mathbf{T}^2}\theta^\mu{}_\nu - 8\pi T^\mu{}_\nu = \frac{2}{\sqrt{-g}}\nabla_\alpha(f_Q\sqrt{-g}P^{\alpha\mu}{}_\nu) + \frac{1}{2}f(Q, \mathbf{T}^2)\delta^\mu{}_\nu + f_Q(P^\mu{}_{\alpha\beta}Q_\nu{}^{\alpha\beta}) \quad (2.25)$$

Moreover we also define the hypermomentum tensor density as [88],

$$H_\rho{}^{\alpha\beta} \equiv \frac{\sqrt{-g}}{16\pi} f_{\mathbf{T}^2} \frac{\delta \mathbf{T}^2}{\delta \hat{\Gamma}^\rho{}_{\alpha\beta}} + \frac{\delta \sqrt{-g} \mathcal{L}_m}{\delta \hat{\Gamma}^\rho{}_{\alpha\beta}} \quad (2.26)$$

Now taking the variation of the gravitational action with respect to the connection we get [88],

$$\nabla_\mu \nabla_\nu (\sqrt{-g} f_Q P^{\mu\nu}{}_\alpha + 4\pi H_\alpha{}^{\mu\nu}) = 0 \quad (2.27)$$

We will be using the above quantities in the next section where we investigate the conservation of the matter energy-momentum tensor.

2.1 Conservation of matter energy-momentum tensor

The covariant derivative of a (1, 1)-form tensor $w^\mu{}_\nu$ may be expressed as,

$$\nabla_\nu w^\mu{}_\nu = D_\mu w^\mu{}_\nu - \frac{1}{2}Q_\lambda w^\lambda{}_\nu - L^\lambda{}_{\mu\nu} w^\mu{}_\lambda \quad (2.28)$$

where D_μ represents the covariant derivative with respect to the Levi-Civita connection ($\Gamma_{\mu\nu}^\alpha$) defined as $\Gamma_{\mu\nu}^\alpha = \hat{\Gamma}_{\mu\nu}^\alpha - L_{\mu\nu}^\alpha$.

Using the above relation we take the covariant derivative of the field equations (2.25) in (1,1)-form and get,

$$D_\mu [f_{\mathbf{T}^2} \theta_\nu^\mu - 8\pi T_\nu^\mu] + \frac{8\pi}{\sqrt{-g}} \nabla_\alpha \nabla_\mu H_\nu^{\alpha\mu} = \frac{1}{2} f_{\mathbf{T}^2} \delta_\nu \mathbf{T}^2 + \frac{1}{\sqrt{-g}} Q_\mu \nabla_\alpha (f_Q \sqrt{-g} P^{\alpha\mu}_\nu) \quad (2.29)$$

From the above relation we see that,

$$D_\mu T_\nu^\mu = \frac{1}{8\pi} \left[D_\mu (f_{\mathbf{T}^2} \theta_\nu^\mu) + \frac{8\pi}{\sqrt{-g}} \nabla_\alpha \nabla_\mu H_\nu^{\alpha\mu} - \frac{1}{\sqrt{-g}} Q_\mu \nabla_\alpha (f_Q \sqrt{-g} P^{\alpha\mu}_\nu) - \frac{1}{2} f_{\mathbf{T}^2} \delta_\nu \mathbf{T}^2 \right] \neq 0 \quad (2.30)$$

So we can see that the matter energy-momentum tensor is not conserved in this case. Now we may introduce a tensor A^ν_α to solve eqn.(2.27) such that [88],

$$\nabla_\mu (\sqrt{-g} f_Q P^{\mu\nu}_\alpha + 4\pi H_\alpha^{\mu\nu}) = \sqrt{-g} A^\nu_\alpha \quad (2.31)$$

After some rigorous calculations we reach the following expression for the covariant derivative of the energy-momentum tensor,

$$D_\mu T_\nu^\mu = \frac{1}{8\pi} \left[D_\mu (f_{\mathbf{T}^2} \theta_\nu^\mu) + \frac{16\pi}{\sqrt{-g}} \nabla_\alpha \nabla_\mu H_\nu^{\alpha\mu} - 8\pi \nabla_\mu \left(\frac{1}{\sqrt{-g}} \nabla_\alpha H_\nu^{\alpha\mu} \right) + 2\nabla_\mu A^\mu_\nu - \frac{1}{2} f_{\mathbf{T}^2} \delta_\nu \mathbf{T}^2 \right] = B_\nu \neq 0 \quad (2.32)$$

As mentioned above the energy-momentum tensor is not conserved for this theory and we get $D_\mu T_\nu^\mu = B_\nu \neq 0$. The non-conservation vector B_ν is a function of the dynamical variables Q , \mathbf{T}^2 and other thermodynamic parameters like energy density and pressure. If the matter content is described by a perfect fluid with density ρ and pressure p and the energy-momentum tensor given by $T^\mu_\nu = (\rho + p) u_\nu u^\mu + p \delta^\mu_\nu$, then following [90] we obtain the energy balance equation as,

$$\dot{\rho} + 3H(\rho + p) = B_\alpha u^\alpha \quad (2.33)$$

It can be clearly seen that this is not the standard continuity equation, but contains correction terms on the RHS. Here $B_\alpha u^\alpha$ is the source term that correlates with the creation or annihilation of energy. If $B_\alpha u^\alpha = 0$, then we have an energy conserved gravitational system, otherwise transfer of energy prevails and particle production takes place in the system. The momentum conservation equation representing the motion of massive particles is given by,

$$\frac{d^2 x^\mu}{ds^2} + \Gamma_{\lambda\sigma}^\mu u^\lambda u^\sigma = \frac{\Upsilon^{\mu\nu}}{\rho + p} (B_\nu - D_\nu p) \quad (2.34)$$

where $\Upsilon^{\mu\nu}$ is the projection operator given by $\Upsilon^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$. The term $\frac{\Upsilon^{\mu\nu}}{\rho + p} (B_\nu - D_\nu p)$ on the RHS is the correction term representing addition force coming from the coupling effects of Q and \mathbf{T}^2 . As a result the dynamical evolution of the massive particles is non-geodesic in nature.

2.2 Matter as perfect fluid

Now we need to consider a specific type of matter component to further simplify the field equations. We assume that the universe is filled with a perfect fluid described by the energy-momentum tensor,

$$T_{\mu\nu} = (\rho + p) u_\mu u_\nu + p g_{\mu\nu} \quad (2.35)$$

where ρ is the energy density and p is the pressure. Moreover u_μ is the four-velocity of the fluid, which is normalized as $u_\mu u^\mu = -1$. Using the above expression we get,

$$T_{\mu\nu}T^{\mu\nu} = \mathbf{T}^2 = \rho^2 + 3p^2 \quad (2.36)$$

We consider the matter Lagrangian $\mathcal{L}_m = p$. From this assumption we see that the final term of $\theta_{\mu\nu}$ in eqn.(2.22) vanishes because pressure p does not depend on the metric tensor. Now using eqn.(2.35) in eqn.(2.22) we get,

$$\theta_{\mu\nu} = -(\rho^2 + 4p\rho + 3p^2) u_\mu u_\nu \quad (2.37)$$

Using the above expressions the field equations takes the following form,

$$\begin{aligned} & -\frac{2}{\sqrt{-g}}\nabla_\alpha (f_Q\sqrt{-g}P_{\mu\nu}^\alpha) - \frac{1}{2}f(Q, \mathbf{T}^2)g_{\mu\nu} - f_Q (P_{\mu\alpha\beta}Q_\nu^{\alpha\beta} - 2Q_{\mu}^{\alpha\beta}P_{\alpha\beta\nu}) \\ & = [8\pi(\rho + p) + f_{\mathbf{T}^2}(\rho^2 + 4p\rho + 3p^2)] u_\mu u_\nu + 8\pi p g_{\mu\nu} \end{aligned} \quad (2.38)$$

3 Cosmology of EMSSTG

In this section we intend to explore the cosmological applications of the above derived theory. We will assume that the universe is described by the homogeneous, isotropic and spatially flat Friedmann-Lemaitre-Robertson-Walker (FLRW) metric given by,

$$ds^2 = -dt^2 + a(t)^2 (dx^2 + dy^2 + dz^2) \quad (3.1)$$

where $a(t)$ is the cosmological scale factor representing the expansion of the universe. The expansion rate may be given by the Hubble parameter H defined as,

$$H \equiv \frac{\dot{a}}{a} \quad (3.2)$$

where dot (.) represents derivative with respect to time. From some rigorous calculations we see that for the above FLRW metric the non-metricity Q may be given in terms of Hubble parameter as,

$$Q = 6H^2 \quad (3.3)$$

Using the metric (3.1) we get the modified Friedmann equations for the $f(Q, \mathbf{T}^2)$ gravity as,

$$6f_Q H^2 - \frac{1}{2}f(Q, \mathbf{T}^2) = 8\pi\rho + f_{\mathbf{T}^2}(\rho^2 + 4p\rho + 3p^2) \quad (3.4)$$

$$6f_Q H^2 - \frac{1}{2}f(Q, \mathbf{T}^2) - 2\left(\dot{f}_Q H + f_Q \dot{H}\right) = -8\pi p \quad (3.5)$$

where as mentioned earlier dot (.) represents derivative with respect to time. It should be noted that we have derived the above FLRW equations for a universe filled with perfect fluid with the energy-momentum tensor given by eqn.(2.35). In the first FLRW equation we see that there are two correction terms $1/2f$ and $f_{\mathbf{T}^2}(\rho^2 + 4p\rho + 3p^2)$ when compared to standard equations of GR. The first term is present in the other extensions of symmetric teleparallel gravity like the $f(Q)$ and $f(Q, T)$ gravity. But the second term is the one to look out for, because this is the term that is characteristic of $f(Q, \mathbf{T}^2)$ theory. Moreover we see that the equations follow perfect correspondence and reduce to those of STG, $f(Q)$ or $f(Q, T)$ under suitable limits. We can write the FLRW equations in the standard form for general relativity as below,

$$3H^2 = 8\pi\rho_{eff} \quad (3.6)$$

$$2\dot{H} + 3H^2 = -8\pi p_{eff} \quad (3.7)$$

where

$$\rho_{eff} = \frac{1}{16\pi f_Q} \left[8\pi\rho + \frac{1}{2}f(Q, \mathbf{T}^2) + f_{\mathbf{T}^2}(\rho^2 + 4p\rho + 3p^2) \right] \quad (3.8)$$

and

$$p_{eff} = \frac{1}{8\pi f_Q} \left[-8\pi p + \frac{1}{2}f(Q, \mathbf{T}^2) - 9f_Q H^2 + 2\dot{f}_Q H \right] \quad (3.9)$$

Now the effective energy density and pressure will follow the standard continuity equation given as,

$$\dot{\rho}_{eff} + 3H(\rho_{eff} + p_{eff}) = 0 \quad (3.10)$$

The effective equation of state ω_{eff} of the dark energy derived from the theory may be given by,

$$\omega_{eff} = \frac{p_{eff}}{\rho_{eff}} = \frac{2 \left[-8\pi p + \frac{1}{2}f(Q, \mathbf{T}^2) - 9f_Q H^2 + 2\dot{f}_Q H \right]}{\left[8\pi\rho + \frac{1}{2}f(Q, \mathbf{T}^2) + f_{\mathbf{T}^2}(\rho^2 + 4p\rho + 3p^2) \right]} \quad (3.11)$$

To study the late cosmic acceleration of universe we have devised a parameter known as the deceleration parameter q , which is defined as,

$$q = -1 - \frac{\dot{H}}{H^2} = \frac{1}{2}(1 + 3\omega_{eff}) \quad (3.12)$$

Using eqn.(3.11) in eqn.(3.12) we can write the deceleration parameter for this theory as,

$$q = \frac{1}{2} \left[1 + \frac{6 \left[-8\pi p + \frac{1}{2}f(Q, \mathbf{T}^2) - 9f_Q H^2 + 2\dot{f}_Q H \right]}{\left[8\pi\rho + \frac{1}{2}f(Q, \mathbf{T}^2) + f_{\mathbf{T}^2}(\rho^2 + 4p\rho + 3p^2) \right]} \right] \quad (3.13)$$

A negative value of this parameter corresponds to the accelerated expansion of the universe. We will study this parameter in detail for the specific models.

3.1 The de-Sitter solution

Now let us explore the vacuum solutions, and correspondingly check that whether our theory admits any de-Sitter like solution or not. To perform that we need to constrain our field equations by considering $\rho = p = 0$ and $H = H_0$ (a constant). Using these in eqns.(3.4) and (3.5) we respectively obtain,

$$6f_Q H_0^2 - \frac{1}{2}f = 0 \quad (3.14)$$

and

$$6f_Q H_0^2 - \frac{1}{2}f - 2H_0 \dot{f}_Q = 0 \quad (3.15)$$

Using the above two equations we get $f_Q = f_0$ (a constant), which on integration gives $f = f_0 Q + \epsilon$ where ϵ is the constant of integration. Finally using this result for f in the eqn.(3.14) we get,

$$H_0 = \sqrt{\frac{\epsilon}{6f_0}} \quad (3.16)$$

which is similar to the results obtained in [87, 88]. For $f_0 = 1$, we get $H_0 = \sqrt{\frac{\epsilon}{6}}$, which is equivalent to the result obtained for general relativity. So it is clear that our EMSSTG theory admits de-Sitter like evolution of the universe when subjected to constraints corresponding to vacuum.

4 Some specific Toy-models

We basically consider two specific toy-models in this section. The models are chosen depending on the nature of coupling between Q and \mathbf{T}^2 . In the first models we will consider an additive form and in the second model we will consider a product form. Both the models should have well-defined STG limits, which is important for correspondence. We will derive the FLRW equations and explore the cosmological evolution of the models.

4.1 Model: 1

Now we need to specify a particular form of $f(Q, \mathbf{T}^2)$ to further explore the field equations. We will assume the general form

$$f(Q, T_{\mu\nu}T^{\mu\nu}) = f(Q, \mathbf{T}^2) = Q + \eta (T_{\mu\nu}T^{\mu\nu})^n = Q + \eta (\mathbf{T}^2)^n \quad (4.1)$$

For $\eta \rightarrow 0$ we recover the STG limit of the theory. Using eqns.(2.35), (2.36), (2.37) and (4.1) in eqn.(2.24) we get simplified field equations as,

$$\begin{aligned} & -\frac{2}{\sqrt{-g}}\nabla_\alpha(\sqrt{-g}P^\alpha{}_{\mu\nu}) - \frac{1}{2}Qg_{\mu\nu} - (P_{\mu\alpha\beta}Q_\nu{}^{\alpha\beta} - 2Q^{\alpha\beta}{}_\mu P_{\alpha\beta\nu}) \\ & = \left[8\pi(\rho + p) + n\eta(\rho^2 + 3p^2)^{n-1}(\rho^2 + 4p\rho + 3p^2)\right]u_\mu u_\nu + \left[8\pi p + \frac{\eta}{2}(\rho^2 + 3p^2)^n\right]g_{\mu\nu} \end{aligned} \quad (4.2)$$

The corresponding Friedmann equations become,

$$3H^2 = 8\pi\rho + \eta(\rho^2 + 3p^2)^{n-1}\left[\left(n + \frac{1}{2}\right)(\rho^2 + 3p^2) + 4np\rho\right] \quad (4.3)$$

$$3H^2 - 2\dot{H} = \frac{\eta}{2}(\rho^2 + 3p^2)^n - 8\pi p \quad (4.4)$$

From the above equations we see that the correction terms are higher order terms in the density ρ which should dominate in the high energy density regime as $\rho \rightarrow \infty$, i.e. early universe. In the later universe these terms gradually fade away giving the effects of the standard FLRW model. Looking at the first Friedmann equation we see that the correction terms are somewhat similar in form to those arising from quantum gravity effects in Loop quantum gravity [91]. These equations are also comparable to those arising from the braneworld models [92]. So we are interested in searching for solutions of bouncing cosmology. We consider the following barotropic equation of state for the fluid

$$p = k\rho \quad (4.5)$$

where k is the barotropic parameter. Using this in eqn.(4.3) we see that the correction terms are quadratic in energy density ρ . At small energy density regime $\rho^2 \approx 0$ and we recover the standard Friedmann equation for $n \geq 1$. At high energy densities (early universe) we see that for $n = 1$, we have two critical points i.e. $H = 0$ at,

$$\rho_{c_1} = 0, \quad \rho_{c_2} = -\frac{16\pi}{\eta(9k^2 + 8k + 3)} \quad (4.6)$$

The first critical point is quite expected and corresponds to an empty universe, where the expansion is supposed to be driven by vacuum energy. The second critical density is more interesting and makes sense only if $\eta(9k^2 + 8k + 3) < 0 \Rightarrow \eta < 0$. So for negative values of η we have a cosmological bounce at a finite non-zero value of energy density. For an early radiation dominated universe i.e. for $k = 1/3$, we see that the bounce occurs at $\rho_c = -\frac{12\pi}{5\eta}$. At this point one can easily check that $\dot{H} = -\frac{176\pi^2}{25\eta} > 0$ for $\eta < 0$. This indicates that the early universe does not start

from an initial singularity but there is a possibility of a cut-off energy density (ρ_c) at which the universe undergoes bounce. So for this model the cosmological singularity problem can be solved by considering suitable initial conditions. We fortunately get an explicit solution of the eqns.(4.3) and (4.4) for $n = 1$ and $k = -1$ as below,

$$a(t) = e^{Ct}, \quad \rho(t) = \frac{\sqrt{2(3C^2\eta + 8\pi^2)} - 4\pi}{2\eta} \quad (4.7)$$

where C is the integration constant. This solution resembles with the solution of the de-Sitter scenario but in the presence of a non-vanishing matter energy-momentum tensor. Note that here we have made use of the result $a(0) = 1$. Here the energy density obtained is constant which can mimic an empty universe in some suitable limits of the parameters (say $\eta \rightarrow \infty$). For such a scenario the deceleration parameter will have a constant negative value throughout the evolution of the universe. So this solution does not generate much cosmological interest.

Now from the eqn.(3.10) using eqns.(3.8), (3.9) and (4.1) we get the continuity equation for this model as,

$$\dot{\rho} = -\frac{3H \left[-18H^2 + 4\dot{H} + 16(1-2k)\pi\rho + \eta(1+3k^2)^{n-1}(3+2n+k(9k+8n+6kn))\rho^{2n} \right]}{2 \left[8\pi + \eta(1+3k^2)^{n-1}n(1+2n+8kn+k^2(3+6n))\rho^{2n-1} \right]} \quad (4.8)$$

Further, for $n = 1$, the above equation reduces to,

$$\dot{\rho} + 3H(\rho + p) = \frac{3H \left[18H^2 - 4\dot{H} + \rho(48k\pi + \eta(1+k(14+k(19+18k))))\rho \right]}{16\pi + 2\eta(3+k(8+9k))\rho} \quad (4.9)$$

The non-conservation term can clearly be seen on the RHS of the above equation. For matter energy-momentum tensor conservation we should have

$$\rho[48k\pi + \eta(1+k(14+k(19+18k)))]\rho = 4\dot{H} - 18H^2 \quad (4.10)$$

It is obvious looking at the above conservation equations (4.8) and (4.9), it is not straightforward to integrate them and get a general solution. But for $k = -1$ and $n = 1$ we can integrate the continuity equations and get expressions similar to eqn.(4.7). Using $k = -1$ and the energy density given in eqn.(4.7) in the above condition for energy momentum conservation (4.10) we get the following solution for the scale factor,

$$a(t) = \frac{B}{\cosh \left[C \left(\frac{9t}{2} - A \right) \right]^{2/9}} \quad (4.11)$$

where A and B are new integration constants and the above expression suggests that if the evolution of the universe follows the above scale factor then the matter energy-momentum tensor is conserved for the model. Obviously it should be kept in mind that this solution is valid only around the Λ CDM regime. In order to get greater insight into the above obtained form of scale factor we generate a plot of $a(t)$ vs t in figure (1). From the figure we see that the evolution of the scale factor obtained from this relation (4.11) complies with our observation of an expanding universe. So the conservation of matter energy momentum may be a possibility for this model subject to proper initial conditions.

4.2 Model: 2

Here we consider a second model in the product form between the two scalar invariants Q and T^2 as below,

$$f(Q, T_{\mu\nu}T^{\mu\nu}) = f(Q, \mathbf{T}^2) = f_0 Q^m (T_{\mu\nu}T^{\mu\nu})^n = f_0 Q^m (\mathbf{T}^2)^n, \quad f_0 \neq 0 \quad (4.12)$$




Fig.1

Fig.1 shows the variation of the scale factor $a(t)$ against t for different initial conditions. We have considered $a(0) \approx 1$.

For $f_0 \rightarrow 1$, $n \rightarrow 0$ and $m \rightarrow 1$ we recover the STG limit of the theory. The modified FLRW equations for this model are given by,

$$3H^2 = \left[\frac{2^{4-m} \pi \rho (\rho^2 + 3p^2)^{1-n}}{f_0 \{(6m - 6n - 3)p^2 + (2m - 2n - 1)\rho^2 - 8np\rho\}} \right]^{1/m} \quad (4.13)$$

$$\dot{H} = \frac{1}{2f_0 m (6^m H^{2m} + 2^{n+1} \times 3^n H^{2n} (n-1))} \times \left[H (\rho^2 + 3p^2)^{-n-1} (3H (\rho^2 + 3p^2)) \right.$$

$$\left. \left(16\pi p + 6^m f_0 H^{2m} (2m-1) (\rho^2 + 3p^2)^n \right) - 2^{n+2} \times 3^n f_0 H^{2n} m n (\rho^2 + 3p^2)^n (\rho\dot{\rho} + 3p\dot{p}) \right] \quad (4.14)$$

We further simplify these equations by considering $n = m = 1$ as given below,

$$3H^2 = -\frac{8\pi\rho}{f_0 (\rho^2 + 8p\rho + 3p^2)} \quad (4.15)$$

$$2f_0 (\rho^2 + 3p^2) \dot{H} - 3f_0 (\rho^2 + 3p^2) H^2 + 4f_0 (\rho\dot{\rho} + 3p\dot{p}) H = 8\pi p \quad (4.16)$$

From the first FLRW equation we directly see that:

$$f_0 (\rho^2 + 8p\rho + 3p^2) < 0 \Rightarrow [f_0 < 0] \text{ OR } [\rho^2 + 8p\rho + 3p^2 < 0 \Rightarrow -\frac{1}{3}(4 + \sqrt{13})\rho < p < \frac{1}{3}(-4 + \sqrt{13})\rho]$$

So we readily get a range of the equation of states for the matter component. This is a very interesting result. We may put the above equations in the standard form as,

$$3H^2 = 8\pi\rho_{eff} \quad (4.17)$$

$$2\dot{H} + 3H^2 = -8\pi p_{eff} \quad (4.18)$$




Fig.2




Fig.3

Fig.2 and 3 shows the variation of the scale factor $a(t)$ and matter energy density $\rho(t)$ against t for model-2 for different initial conditions respectively. For Fig.2 we have taken $C_1 = 5, C_2 = -4$. For Fig.3 we have taken $C_1 = -0.5, f_0 = 0.1, 8\pi = 1$. In the figures we have considered $a(0) \approx 1$.

where

$$\rho_{eff} = -\frac{\rho}{f_0(\rho^2 + 8p\rho + 3p^2)} \quad (4.19)$$

$$p_{eff} = -\frac{1}{8\pi f_0(\rho^2 + 3p^2)} [8\pi p - 4f_0(\rho\dot{\rho} + 3p\dot{p})H + 6f_0(\rho^2 + 3p^2)H^2] \quad (4.20)$$

From eqn.(4.15) we see that $\rho_{c3} = 0$ is a critical point of the model, which corresponds to the empty universe. Unlike the previous model here we do not get a bouncing scenario from a non-zero finite energy density. One more interesting feature of this model is that there are two singularities corresponding to $p = \frac{1}{3}(-4 \pm \sqrt{13})\rho$. Considering barotropic equation of state given in eqn.(4.5) the above conditions reduce to $k = \frac{1}{3}(-4 \pm \sqrt{13})$ for a non-empty universe. In this case we are able to get an explicit general solution for the eqns.(4.15) and (4.16) as given below,

$$a(t) = C_2 \left[(k+1)^2(3k-1)t + 2C_1(3k^2+1) \right]^{-2(3k^2+1)/(k+1)^2(3k-1)} \quad (4.21)$$

$$\rho(t) = \frac{8\pi \left[(k+1)^2(3k-1)t + 2C_1(3k^2+1) \right]^2}{6f_0(3k^2+1)^2(3k^2+8k+1)} \quad (4.22)$$

where C_1 and C_2 are integration constant. It is seen that the scale factor is undefined at $k = -1$ (Λ CDM era) and $k = 1/3$ (radiation era). We have generated plots for both scale factor and matter energy density in figures (2) and (3) respectively. It is seen that the trajectories are quite satisfactory and complies with the observations. We have also plotted the corresponding trajectories for the standard Λ CDM model to get a comparative idea about the parameters of the model.

For the model the continuity equation turns out to be,

$$\dot{\rho} + 3H(\rho + p) = 3H \left[(1+k)\rho + \frac{\zeta + f_0\pi\rho^4\xi(4k\pi - 3f_0H\xi\rho)}{\zeta/\rho + 6\pi f_0^2\xi^2H\rho^4} \right] \quad (4.23)$$

where $\zeta = \frac{4}{f_0(1+8k+3k^2)}$ and $\xi = 1 + 3k^2$. The term on the RHS is the non-conservation term. Setting the non-conservation term equal to zero and solving the corresponding differential equation for the scale factor we have the two following evolution of the universe,

$$a(t) = C_3, \quad a(t) = C_3 e^{\frac{\xi\sigma \left[-9f_0^2(\sigma+1)\xi^7\sigma^3 - 16k\pi^6(2C_1\xi + \sigma^2(3k-1)t) \right]^8}{8\pi^6\sigma^2(6k^2+k-1)[2C_1\xi + \sigma^2(3k-1)t]^9}} \quad (4.24)$$




Fig.4

Fig.4 shows the variation of the deceleration parameter $q(t)$ against t for model-2 for different initial conditions. We have taken $C_1 = -0.5$, $C_2 = 0.1$, $f_0 = 0.2$, $8\pi = 1$.

where $\sigma = 1 + 8k + 3k^2$, $\xi = 1 + 3k^2$, $\varrho = k + 1$ and C_3 is the integration constant. The first value suggests that the scale factor is a constant suggesting a static evolution of the universe. Since the scale factor does not grow with time the universe does not expand. This is contrary to our observations. So we ignore this result. The other expression gives an exponential type evolution of the universe. It is quite certain that this will correspond to a de-Sitter like evolution of the universe for suitable initial conditions. So there exists conditions under which this model may satisfy the standard energy momentum conservation relation. The deceleration parameter for this model is plotted in fig.4. From the figure we see that there is a transition of q from positive level to negative level at some finite value of t . These values of t for each trajectory correspond to the redshift value $z \approx 0.6$, where the universe enters into the accelerating phase from a decelerating one. This is cosmologically viable with the observations.

5 Energy Conditions

Energy conditions are tools to establish the positiveness of the energy-momentum tensor in the presence of matter. These conditions actually describe the attractive nature of gravity and also takes care of the causal and geodesic structure of the spacetime [93]. In this section we explore the energy conditions that the thermodynamic parameters of the $f(Q, \mathbf{T}^2)$ theory need to satisfy and thus put some constraints on the model parameters. We will use a perfect fluid matter distribution. It should be mentioned here that the late cosmic acceleration demands the violation of the strong energy condition, since it requires anti-gravitational effect to play its role. The matter component responsible for this violation may be dark energy. There are basically four energy conditions that can be derived from standard general theory of relativity. Considering isotropic cosmology they are:

$$(I) \text{ Weak Energy Condition (WEC)} \Rightarrow \rho_{eff} \geq 0, \quad \rho_{eff} + p_{eff} \geq 0$$

$$(II) \text{ Null Energy Condition (NEC)} \Rightarrow \quad \rho_{eff} + p_{eff} \geq 0$$

$$(III) \text{ Dominant Energy condition (DEC)} \Rightarrow \quad \rho_{eff} \geq 0, \quad \rho_{eff} \geq |p_{eff}|$$

$$(IV) \text{ Strong Energy condition (SEC)} \Rightarrow \quad \rho_{eff} + 3p_{eff} \geq 0$$

Now using eqns.(3.8) and (3.9) in the WEC we get the following inequalities considering $8\pi = 1$,

$$\frac{1}{f_Q} \left[\rho + \frac{1}{2}f(Q, \mathbf{T}^2) + f_{\mathbf{T}^2} (\rho^2 + 4p\rho + 3p^2) \right] \geq 0 \quad (5.1)$$

$$\frac{1}{f_Q} \left[3f(Q, \mathbf{T}^2) + 4H \left(2\dot{f}_Q - 9Hf_Q \right) + 2(\rho - 2p) + 2f_{\mathbf{T}^2} (\rho^2 + 4p\rho + 3p^2) \right] \geq 0 \quad (5.2)$$

The expression (5.2) is the required condition for the satisfaction of NEC. The dominant energy condition $\rho_{eff} \geq |p_{eff}|$ may be modified as $\rho_{eff} \pm p_{eff} \geq 0$. So along with the conditions given in (5.1) and (5.2) we have another condition for DEC given below,

$$\frac{1}{f_Q} \left[f(Q, \mathbf{T}^2) + 4H \left(2\dot{f}_Q - 9Hf_Q \right) - 2(\rho + 2p) - 2f_{\mathbf{T}^2} (\rho^2 + 4p\rho + 3p^2) \right] \leq 0 \quad (5.3)$$

Finally from the SEC we get,

$$\frac{1}{f_Q} \left[7f(Q, \mathbf{T}^2) + 12H \left(2\dot{f}_Q - 9Hf_Q \right) + 2(\rho - 6p) + 2f_{\mathbf{T}^2} (\rho^2 + 4p\rho + 3p^2) \right] \geq 0 \quad (5.4)$$

So all the energy conditions finally yield four inequalities given by (5.1), (5.2), (5.3) and (5.4), which can be used to constrain the theory. Now we may use our toy-models discussed above to check how viable and effective these energy conditions are in constraining cosmological models. We discuss them one by one below.

5.1 Model: 1

Here we will use the model given in eqn.(4.1) which has two free parameters n and η . We will also consider the barotropic equation of state $p = k\rho$. Using the above relations we give the energy conditions for this model as,

$$\mathbf{WEC:} \quad 3H_0^2 + \rho_0 + \frac{1}{2} (1 + 3k^2)^{n-1} [1 + 2n + k(3k + 8n + 6kn)] \eta \rho_0^{2n} \geq 0$$

and

$$-18H_0^2 + 2(1 - 2k)\rho_0 + (1 + 3k^2)^{n-1} [3 + 2n + k(9k + 8n + 6kn)] \eta \rho_0^{2n} \geq 0 \quad (5.5)$$

$$\mathbf{NEC:} \quad -18H_0^2 + 2(1 - 2k)\rho_0 + (1 + 3k^2)^{n-1} [3 + 2n + k(9k + 8n + 6kn)] \eta \rho_0^{2n} \geq 0 \quad (5.6)$$

$$\mathbf{DEC:} \quad 3H_0^2 + \rho_0 + \frac{1}{2} (1 + 3k^2)^{n-1} [1 + 2n + k(3k + 8n + 6kn)] \eta \rho_0^{2n} \geq 0$$

,

$$-18H_0^2 + 2(1 - 2k)\rho_0 + (1 + 3k^2)^{n-1} [3 + 2n + k(9k + 8n + 6kn)] \eta \rho_0^{2n} \geq 0$$

and

$$(1 + 3k^2)^{n-1} [2n + 8kn + 3k^2(2n - 1) - 1] \eta \rho_0^{2n} + 2(15H_0^2 + \rho_0 + 2k\rho_0) \geq 0 \quad (5.7)$$

$$\text{SEC: } 2(1 - 6k)\rho_0 + (1 + 3k^2)^{n-1} [7 + 2n + k(21k + 8n + 6kn)] \eta \rho_0^{2n} - 66H_0^2 \geq 0 \quad (5.8)$$

Since we are interested in constraining the models we have used the present values of Hubble parameter H_0 and matter energy density ρ_0 in the above energy conditions. The present value of the Hubble parameter is estimated to be $H_0 = 67.9 \text{ kmsec}^{-1} \text{ Mpc}^{-1}$ [94, 95] and that of ρ_0 is $\rho_0 = 9.9 \times 10^{-30} \text{ gm cm}^{-3}$ [96]. Now we may consider various cosmological era for matter by changing the value of k such as $k = 1, 1/3, 0, -1/3, -1$. After putting all these values we will get inequalities connecting only η and n , from where it will be straightforward to put constraints on these two model parameters.

5.2 Model: 2

In this model we have three free parameters f_0 , m and n . Since for this model we have \dot{H} present in f_Q terms, we have used the eqn.(3.12) to define $\dot{H} = -H^2(1 + q)$. Below we present the energy conditions for this model.

WEC:

$$\frac{8kn + (3k^2 + 1)(1 + 2n)}{m} + \frac{2 \times (6H_0^2)^{-m} \rho_0^{1-2n}}{f_0 m (1 + 3k^2)^{n-1}} \geq 0,$$

and

$$\frac{1}{m} \left[3 + 2\varrho(3\varrho - 2)\xi^{-1}n + 2f_0^{-1}H_0^{-2m}\xi^{-n}(3 - 2\varrho)6^{-m}\rho_0^{1-2n} + \frac{2}{3}m \left\{ 2(6H_0^2)^{(n-m)} \left(2(1-n)q_0 + \frac{\xi}{\vartheta} \right) - 9 \right\} \right] \geq 0 \quad (5.9)$$

NEC:

$$\frac{1}{m} \left[3 + 2\varrho(3\varrho - 2)\xi^{-1}n + 2f_0^{-1}H_0^{-2m}\xi^{-n}(3 - 2\varrho)6^{-m}\rho_0^{1-2n} + \frac{2}{3}m \left\{ 2(6H_0^2)^{(n-m)} \left(2(1-n)q_0 + \frac{\xi}{\vartheta} \right) - 9 \right\} \right] \geq 0 \quad (5.10)$$

DEC:

$$\frac{8kn + (3k^2 + 1)(1 + 2n)}{m} + \frac{2 \times (6H_0^2)^{-m} \rho_0^{1-2n}}{f_0 m (1 + 3k^2)^{n-1}} \geq 0,$$

$$\frac{1}{m} \left[7 + 2\varrho(3\varrho - 2)\xi^{-1}n + 2f_0^{-1}H_0^{-2m}\xi^{-n}(3 - 2\varrho)6^{-m}\rho_0^{1-2n} + \frac{2}{3}m \left\{ 2(6H_0^2)^{(n-m)} \left(2(1-n)q_0 + \frac{\xi}{\vartheta} \right) - 9 \right\} \right] \geq 0, \quad (5.11)$$

and

$$\frac{1}{m} \left[1 - 2\varrho(3\varrho - 2)\xi^{-1}n - 2f_0^{-1}H_0^{-2m}\xi^{-n}(3 - 2\varrho)6^{-m}\rho_0^{1-2n} + \frac{2}{3}m \left\{ 2(6H_0^2)^{(n-m)} \left(2(1-n)q_0 + \frac{\xi}{\vartheta} \right) - 9 \right\} \right] \leq 0 \quad (5.12)$$

SEC:

$$\frac{1}{m} \left[3 + 2\varrho(3\varrho - 2)\xi^{-1}n + 2f_0^{-1}H_0^{-2m}\xi^{-n}(7 - 6\varrho)6^{-m}\rho_0^{1-2n} + 2m \left\{ 2(6H_0^2)^{(n-m)} \left(2(1-n)q_0 + \frac{\varsigma}{\vartheta} \right) - 9 \right\} \right] \geq 0 \quad (5.13)$$

where σ , ξ and ϱ have been defined earlier in eqn.(4.24) and the expressions for ς and ϑ are given in the appendix. Just like the previous model, in addition to the current values H_0 , ρ_0 we also have to use $q_0 = -0.503$ [94, 95] for this case, to put constraints on the parameters f_0 , n and m from the above inequalities (which is pretty straightforward). So the above energy conditions can be used as constraint relations to constrain the free parameters of the theory. These constraints may be checked or even developed by a full fledged observational data analysis on the model. That will be a good future project to undertake, but for the time being these constraints give a perfect shape to the theory by closing the model.

6 Conclusion

In the present work we have proposed yet another extension of the symmetric teleparallel gravity by generalizing the gravity Lagrangian with an arbitrary function $f(Q, T_{\mu\nu}T^{\mu\nu})$. The field equations were derived in a metric formalism. The correction terms introduced by the modified gravity were noted. As expected for any gravity theory involving non-minimal coupling between geometry and matter sectors, the covariant divergence of the energy-momentum tensor was non-zero thus implying the non-conservation of the same. The non-conservation term was derived using the field equations. The momentum conservation equation showed the presence of correction terms implying extra force on the massive particles thus making the motion non-geodesic. The field equations were further simplified by considering perfect fluid as the matter component. Using these field equations we resorted to study the cosmological evolution of the theory. The FLRW equations for a flat homogeneous and isotropic spacetime were derived. It was noted that there were two additional modification terms introduced in the equations in contrast to those of standard GR. The two additional terms were of the form $f/2$ and $f_{\mathbf{T}^2}(\rho^2 + 4p\rho + 3p^2)$. The first one came from the coupling between the geometric and matter sector and the second one is completely a source term. These higher order terms dominate in the early universe and gradually fade away at late times giving the effects of standard FLRW universe. These corrections are totally intrinsic and uniquely describes the modified gravity. Expressions for some cosmological parameters like the equation of state and deceleration parameter were derived. We investigated the vacuum solution of the theory and saw that EMSSTG admits a de-Sitter like solution in its framework.

To get more insights into the cosmological framework of the theory we studied two specific toy-models $Q + \eta(\mathbf{T}^2)^n$ and $f_0Q^m(\mathbf{T}^2)^n$. We saw that both the models had STG as a limiting case and hence we can recover the parent theory from the equations. After deriving the FLRW equations for the first model we saw that the equations had a flavour of the quantum gravity effects of the loop quantum gravity. So solutions for bouncing cosmology was investigated and it was found that the model indeed supported a cosmological bounce at a finite time, thus avoiding the singularity. Various constraints were imposed on the model from these relations. Although we did not get a general solution of the model, but for $k = -1$ we obtained a solution which resembled the de-Sitter solution. Then we derived the continuity equation for the model and studied the non-conservation term. Using it we were able to trace the evolution of the scale factor for which the non-conservation term will vanish. Similar studies were undertaken for the second model. In this case we were fortunate enough to get a general solution of the FLRW equations. The obtained scale factor and the matter energy density were plotted and compared to those of the standard Λ CDM model. We also obtained the plots for the deceleration parameter for this model in a comparative

scenario with the Λ CDM model. The transition from a decelerating to an accelerating universe was clearly evident and the deviation of the trajectories from those of the Λ CDM model was also noted.

Finally we explored the energy conditions in the background of the theory. The basic energy conditions WEC, NEC, DEC and SEC were derived for the theory and also for the two specific toy-models. In the first model we had two free parameters n and η after we fixed H and ρ with their current values from the observations. Similarly in the second model we had three free parameters f_0 , m and n after the fixation from the observations. Basically four constraints were obtained for each model using the energy conditions which are sufficient to put bounds on the free parameters of both the models. Thus the models were well-constrained by the energy conditions. Finally from the study we conclude that the theory is perfectly suitable to describe the cosmological dynamics of both the early and the late universe without resorting to dark energy. There is scope for further development of the theory, which will be undertaken in future projects. The standout feature of EMSSTG is that the field equations contain terms which arise from the quantum gravity effects and thus are responsible for the avoidance of the singularity. So this theory is a singularity free cosmologically viable theory. Moreover the non-linear density terms in the equations dominate in the early universe and gradually fade away at later times. So the effect of modified gravity is predominantly felt in the early universe and it eases out to give the standard FLRW effects at late times.

Acknowledgments

The author acknowledges the Inter University Centre for Astronomy and Astrophysics (IUCAA), Pune, India for granting visiting associateship.

7 Appendix

7.1 Expressions for ς and ϑ in section 5.2

$$\begin{aligned} \varsigma = & (\xi\rho_0^2)^{-n} [-2f_0 ((6H_0^2)^m + 4(6H_0^2)^n (n-1)) (1 - 3(m+1)n + 2n^2) (\xi\rho_0^2)^n \chi \\ & + 6nH_0^2 (-6k\rho_0 - f_0 (3(6H_0^2)^m (3m-1) + 2(6H_0^2)^n m (n-1)) (\xi\rho_0^2)^n)] \end{aligned} \quad (7.1)$$

$$\vartheta = f_0 [-2mn(6H_0^2)^{1+n} + ((6H_0^2)^m + 2(6H_0^2)^n (n-1)) 2(2n-1)\chi] \quad (7.2)$$

where

$$\chi = \left(\frac{6^{-m}(\xi\rho_0^2)^{n-1}}{f_0\rho_0(\xi(2m-1) - 2\rho(3\rho-2)n)} \right)^{1/m}$$

References

- [1] A. Einstein :- *Annalen der Physik* **354**, 769 (1916).
- [2] D. F. Watson, E. A. Stanley, Davidson C.:- *Philosophical Transactions of the Royal Society of London: Series A* **220** 291 (1920)
- [3] L. D. Landau, E. M. Lifshitz, *The Classical Theory of Fields* (Pergamon Press, Oxford, 1970)
- [4] B. P. Abbott et al., LIGO Scientific Collaboration and Virgo Collaboration:- *Phys. Rev. Lett.* **116**, 061102 (2016).
- [5] The Event Horizon Telescope Collaboration :- *ApJL* **875** L1 (2019)
- [6] C.M. Will :- *Living Rev. Relat.* **17**, 4 (2014)

- [7] M. Ishak :- *Living Rev. Relat.* **22**, 1 (2019).
- [8] S. Perlmutter et. al. :- *Astrophys. J.* **517** 565 (1999).
- [9] A. G. Riess et al. :- *Astron. J.* **116** 1009 (1998).
- [10] P. Brax :- *Rep. Prog. Phys.* **81** 016902 (2018)
- [11] S. Nojiri, S. D. Odintsov, V. K. Oikonomou :- *Phys. Rep.* **692** 1 (2017).
- [12] S. Nojiri and S. D. Odintsov :- *Int. J. Geom. Methods Mod. Phys.* **04** 115 (2007).
- [13] S. Capozziello, R. D'Agostino, O. Luongo:- *Int. J. Mod. Phys. D* **28** 1930016 (2019).
- [14] H. A. Buchdahl : *Month. Not. R. Astron. Soc.* **150**, 1 (1970)
- [15] L. Amendola, D. Polarski, S. Tsujikawa :- *Phys. Rev. Lett.* **98** 131302 (2007).
- [16] T. P. Sotiriou :- *Classical Quantum Gravity* **23** 5117 (2006).
- [17] Y.-S. Song, W. Hu, I. Sawicki :- *Phys. Rev. D* **75** 044004 (2007).
- [18] S. Nojiri, S. D. Odintsov :- *Phys. Rev. D* **74** 086005 (2006).
- [19] P. Rudra :- *Nucl. Phys. B* **956** 115014 (2020)
- [20] P. Rudra :- *Commun. Theor. Phys.* **66** 149 (2016)
- [21] T. P. Sotiriou, V. Faraoni :- *Rev. Mod. Phys.* **82** 451 (2010).
- [22] A. De Felice, S. Tsujikawa :- *Living Rev. Relativity* **13** 3 (2010).
- [23] T. Harko, F. S. N. Lobo :- *Eur. Phys. J. C.* **70** 373 (2010).
- [24] R. Ribeiro, J. Pramos :- *Phys. Rev. D* **90** 124065 (2014).
- [25] R. P. L. Azevedo, J. Pramos :- *Phys. Rev. D* **94** 064036 (2016).
- [26] B. Pourhassan, P. Rudra :- *Phys. Rev. D* **101** 084057 (2020).
- [27] T. Harko, F. S. N. Lobo, S. Nojiri, S. D. Odintsov :- *Phys. Rev. D.* **84** 024020 (2011).
- [28] M. Sharif, M. Zubair :- *JCAP* **03** 028 (2012).
- [29] E. H. Baffou, M. J. S. Houndjo, M. E. Rodrigues, A. V. Kpadonou, J. Tossa :- *Phys. Rev. D* **92** 8, 084043 (2015).
- [30] P. Rudra :- *Eur. Phys. J. Plus* **130** 4, 66 (2015).
- [31] H. Shabani, M. Farhoudi :- *Phys. Rev. D* **88** 044048 (2013).
- [32] F. G. Alvarenga, A. de la Cruz-Dombriz, M. J. S. Houndjo, M. E. Rodrigues, D. Sez-Gmez :- *Phys. Rev. D* **87** 10, 103526 (2013).
- [33] P. Rudra, K. Giri :- *Nucl. Phys. B* **967** 115428 (2021)
- [34] P. Rudra :- arxiv: 2006.00228 [gr-qc]
- [35] N. Katirci, M. Kavuk, :- *Eur. Phys. J. Plus* **129**, 163(2014)
- [36] M. Roshan, F. Shojai :- *Phys. Rev. D.* **94** 044002 (2016).
- [37] A. Ashtekar, T. Pawłowski, P. Singh :- *Phys. Rev. D* **74**, 084003 (2006).
- [38] C. V. R. Board, J. D. Barrow :- *Phys. Rev. D* **96**, 123517 (2017).
- [39] O. Akarsu, N. Katirci, S. Kumar, R. C. Nunes, M. Sami :- *Phys. Rev. D* **98** 6 (2018)
- [40] S. Bahamonde, M. Marciu, P. Rudra :- *Phys. Rev. D* **100**, 083511 (2019).
- [41] C. Ranjit, P. Rudra, S. Kundu :- *Annals Phys.* **428** 168432 (2021).
- [42] P. Rudra, B. Pourhassan :- *Phys. Dark Univ.* **33** 100849 (2021).

- [43] O. Akarsu, N. Katirci, S. Kumar :- *Phys. Rev. D* **97** 2 (2018)
- [44] O. Akarsu, J. D. Barrow , S. Cikintoglu, K. Y. Eksi, N. Katirci :- *Phys. Rev. D* **97** 12 (2018)
- [45] P. H. R. S. Moraes , P. K. Sahoo :- *Phys. Rev. D* **97** 2 (2018)
- [46] N. Nari, M. Roshan :- *Phys. Rev. D* **98** 2 (2018)
- [47] A. Keskin :- *AIP Conf. Proc.* **2042** 1 (2018)
- [48] O. Akarsu, J. D. Barrow, C. V. R. Board, N. M. Uzun, J. A. Vazquez :- *Eur. Phys. J. C.* **79** 10, 846 (2019)
- [49] B. Riemann, Habilitationsschrift, 1854. Abhandlungen der Kniglichen Gesellschaft der Wissenschaften zu Gttingen **13**, 1 (1867)
- [50] H. Weyl :- *Sitzungsber. Preuss. Akad. Wiss.* **465**, 1 (1918).
- [51] E. Cartan, C. R. Acad :- *Sci.* **174**, 593 (1922).
- [52] E. Cartan :- *Ann. Ec. Norm.* **40**, 325 (1923).
- [53] E. Cartan :- *Ann. Ec. Norm.* **41**, 1 (1924).
- [54] E. Cartan :- *Ann. Ec. Norm.* **42**, 17 (1925).
- [55] F. W. Hehl, P. von derHeyde, G. D. Kerlick, J. M. Nester : *Rev. Mod. Phys.* **48**, 393 (1976).
- [56] D. Puetzfeld, R. Tresguerres :- *Class. Quant. Gravity* **18**, 677 (2001).
- [57] D. Putzfeld :- *Class. Quant. Gravity* **19**, 4463 (2002).
- [58] D. Puetzfeld :- *Class. Quant. Gravity* **19**, 3263 (2002).
- [59] T. Y. Moon, P. Oh, J.S. Sohn :- *JCAP* **11**, 005 (2010).
- [60] M. Novello, S.E. Perez Bergliaffa :- *Phys. Rep.* **463**, 127 (2008).
- [61] R. Weitzenbeck :- *Invariantentheorie* (Noordhoff, Groningen, 1923)
- [62] A. Einstein :- Preussische Akademie der Wissenschaften, Phys.-math. Klasse, Sitzungsberichte 1928, 217 (1928).
- [63] C. Moller :- *Mat. Fys. Skr. Dan. Vid. Selsk.* **1**, 10 (1961)
- [64] C. Pellegrini, J. Plebanski :- *Mat. Fys. Skr. Dan. Vid. Selsk.* **2**, 4 (1963)
- [65] K. Hayashi, T. Shirafuji :- *Phys. Rev. D* **19**, 3524 (1979)
- [66] Y-F. Cai, S. Capozziello, M. D. Laurentis, E. N. Saridakis :- *Rep. Prog. Phys.* **79** 106901 (2016)
- [67] M. Krssak, R. J. van den Hoogen, J. G. Pereira, C. G. Bohmer, A. A. Coley :- *Class. Quantum Grav.* **36** 183001 (2019)
- [68] R. Ferraro, F. Fiorini :- *Phys. Rev. D* **75**, 084031 (2007)
- [69] R. Ferraro, F. Fiorini :- *Phys. Rev. D* **78**, 124019 (2008)
- [70] G. R. Bengochea, R. Ferraro :- *Phys. Rev. D* **79**, 124019 (2009)
- [71] E. V. Linder :- *Phys. Rev. D* **81**, 127301 (2010)
- [72] T. Harko, F. S. N. Lobo, G. Otalora, E. N. Saridakis :- *Phys. Rev. D* **89**, 124036 (2014)
- [73] S. Bahamonde, C. G. Boehmer, M. Krssak :- *Phys. Lett. B* **775**, 37 (2017)
- [74] M. Jamil, D. Momeni, R. Myrzakulov, P. Rudra :- *J. Phys. Soc. Jap.* **81** 114004 (2012)
- [75] P. Rudra :- *Astrophys. Space Sci.* **357** 135 (2015)
- [76] C. Ranjit, P. Rudra :- *Int. J. Mod. Phys. D* **25** 1650008 (2016)
- [77] Z. Haghani, T. Harko, H. R. Sepangi, S. Shahidi :- *JCAP* **10**, 061 (2012)

- [78] Z. Haghani, T. Harko, H. R. Sepangi, S. Shahidi :- *Phys. Rev. D* **88**, 044024 (2013)
- [79] J. M. Nester, H.-J. Yo :- *Chin. J. Phys.* **37**, 113 (1999)
- [80] J. Beltran Jimenez, L. Heisenberg, T. Koivisto :- *Phys. Rev.* **98**, 044048 (2018)
- [81] J. Lu, X. Zhao, G. Chee :- *Eur. Phys. J. C* **79**, 530 (2019)
- [82] R. Lazkoz, F. S. N. Lobo, M. Ortiz-Bao, V. Salzano :- *Phys. Rev. D* **100** 104027 (2019)
- [83] M. Adak, M. Kalay, O. Sert :- *Int. J. Mod. Phys. D* **15**, 619 (2006).
- [84] M. Adak :- *Turk. J. Phys.* **30**, 379 (2006).
- [85] M. Adak, . Sert, M. Kalay, M. Sari :- *Int. J. Mod. Phys. A* **28**, 1350167 (2013).
- [86] I. Mol :- *Advances in Applied Clifford Algebras* **27**, 2607 (2017).
- [87] T. Harko, T. S. Koivisto, F. S. N. Lobo, G. J. Olmo, D. Rubiera-Garcia :- *Phys. Rev. D* **98** 084043 (2018)
- [88] Y. Xu, G. Li, T. Harko, S-D Liang :- *Eur. Phys. J. C.* **79**, 708 (2019)
- [89] J. M. Nester, H.-J. Yo :- *Chin. J. Phys.* **37**, 113 (1999).
- [90] T. Harko, T. S. Koivisto, F. S. N. Lobo, G. J. Olmo, D. Rubiera-Garcia :- *Phys. Rev. D* **98**, 084043 (2018)
- [91] C. Rovelli :- *Living Rev. Relativ.* **1**, 1 (1998)
- [92] R. Maartens, K. Koyama :- *Living Rev. Relativity* **13**, 5 (2010)
- [93] S. Capozziello, S. Nojiri, S. D. Odintsov :- *Phys. Lett. B* **781**, 99 (2018).
- [94] Planck Collaboration :- *Astron. Astrophys.* **641**, A6 (2020)
- [95] S. Capozziello, R. D'Agostino, O. Luongo :- *Int. J. Mod. Phys. D* **28**, 1930016 (2019).
- [96] https://wmap.gsfc.nasa.gov/universe/uni_matter.html