

Dynamic Cantor Derivative Logic

David Fernández-Duque[§] and Yoàv Montacute[†]

[§]Department of Mathematics, Ghent University
david.fernandezduque@ugent.be

[†]Computer Laboratory, University of Cambridge
yoav.montacute@cl.cam.ac.uk

September 14, 2022

Abstract

Topological semantics for modal logic based on the Cantor derivative operator gives rise to derivative logics, also referred to as d -logics. Unlike logics based on the topological closure operator, d -logics have not previously been studied in the framework of dynamical systems, which are pairs (X, f) consisting of a topological space X equipped with a continuous function $f: X \rightarrow X$.

We introduce the logics **wK4C**, **K4C** and **GLC** and show that they all have the finite Kripke model property and are sound and complete with respect to the d -semantics in this dynamical setting. In particular, we prove that **wK4C** is the d -logic of all dynamic topological systems, **K4C** is the d -logic of all T_D dynamic topological systems, and **GLC** is the d -logic of all dynamic topological systems based on a scattered space. We also prove a general result for the case where f is a homeomorphism, which in particular yields soundness and completeness for the corresponding systems **wK4H**, **K4H** and **GLH**.

The main contribution of this work is the foundation of a general proof method for finite model property and completeness of dynamic topological d -logics. Furthermore, our result for **GLC** constitutes the first step towards a proof of completeness for the trimodal topo-temporal language with respect to a finite axiomatisation – something known to be impossible over the class of all spaces.

1 Introduction

Dynamic (topological) systems are mathematical models of processes that may be iterated indefinitely. Formally, they are defined as pairs $\langle \mathfrak{X}, f \rangle$ consisting of a topological space $\mathfrak{X} = \langle X, \tau \rangle$ and a continuous function $f: X \rightarrow X$; the intuition is that points in the space \mathfrak{X} ‘move’ along their orbit, $x, f(x), f^2(x), \dots$ which usually simulates changes in time. *Dynamic topological logic (DTL)* combines modal logic and its topological semantics with linear temporal logic (see Pnueli [22]) in order to reason about dynamical systems in a decidable framework.

Due to their rather broad definition, dynamical systems are routinely used in many pure and applied sciences, including computer science. To cite some

recent examples, in data-driven dynamical systems, data-related problems may be solved through data-oriented research in dynamical systems as suggested by Brunton and Kutz [4]. Weinan [25] proposes a dynamic theoretic approach to machine learning where dynamical systems are used to model nonlinear functions employed in machine learning. Lin and Antsaklis’s [19] hybrid dynamical systems have been at the centre of research in control theory, artificial intelligence and computer-aided verification. Mortveit and Reidys’s [21] sequential dynamical systems generalise aspects of systems such as cellular automata, and also provide a framework through which we can study dynamical processes in graphs. Another example of dynamical systems and computer science can be found in the form of linear dynamical systems, i.e. systems with dynamics given by a linear transformation. Examples of such systems in computer science include Markov chains, linear recurrence sequences and linear differential equations. Moreover, there are known strong connections between dynamical systems and algorithms. This may be found for example in the work of Hanrot, Pujol and Stehlé [14], and in the work of Chu [5].

Such applications raise a need for effective formal reasoning about topological dynamics. Here, we may take a cue from modal logic and its topological semantics. The study of the latter dates back to McKinsey and Tarski [20], who proved that the modal logic **S4** is complete for a wide class of spaces, including the real line. Artemov, Davoren and Nerode [1] extended **S4** with a ‘next’ operator in the spirit of **LTL**, producing the logic **S4C**. They proved that this logic is sound and complete with respect to the class of all dynamic topological systems. The system **S4C** was enriched with the ‘henceforth’ tense by Kremer and Mints, who dubbed the new logic *dynamic topological logic* (**DTL**). Later, Konev et al. [15] showed that **DTL** is undecidable, and Fernández-Duque [9] showed that it is not finitely axiomatisable on the class of all dynamic topological systems.

The aforementioned work on dynamic topological logic interprets the modal operator \diamond as a closure operator. However, McKinsey and Tarski had already contemplated semantics that are instead based on the Cantor derivative [20]: the *Cantor derivative* of a set A , usually denoted by $d(A)$, is the set of points x such that x is in the closure of $A \setminus \{x\}$ (see Section 2). This interpretation is often called *d-semantics* and the resulting logics are called *d-logics*. These logics were first studied in detail by Esakia, who showed that the *d*-logic **wK4** is sound and complete with respect to the class of all topological spaces [6]. It is well-known that semantics based on the Cantor derivative are more expressive than semantics based on the topological closure. For example, consider the property of a space \mathfrak{X} being *dense-in-itself*, meaning that \mathfrak{X} has no isolated points (see Section 3.2). The property of being dense-in-itself cannot be expressed in terms of the closure operator, but it *can* be expressed in topological *d*-semantics by the formula $\diamond\top$.

Logics based on the Cantor derivative appear to be a natural choice for reasoning about dynamical systems. However, there are no established results of completeness for such logics in the setting of dynamical systems, i.e. when a topological space is equipped with a continuous function. Our goal is to prove the finite Kripke model property, completeness and decidability of logics with the Cantor derivative operator and the ‘next’ operator \circ over some prominent classes of dynamical systems: namely, those based on arbitrary spaces, on T_D spaces (spaces validating the 4 axiom $\Box p \rightarrow \Box\Box p$) and on *scattered spaces* (see Section 3.2 for definitions). The reason for considering scattered spaces is to

circumvent the lack of finite axiomatisability of **DTL** by restricting to a suitable subclass of all dynamical systems. In the study of dynamical systems and topological modal logic, one often works with *dense-in-themselves* spaces. This is a sensible consideration when modelling physical spaces, as Euclidean spaces are dense-in-themselves. However, as we will see in Section 3.2, some technical issues that arise when studying **DTL** over the class of all spaces disappear when restricting our attention to scattered spaces, which in contrast have many isolated points. Further, we consider dynamical systems where f is a *homeomorphism*, i.e. where f^{-1} is also a continuous function. Such dynamical systems are called *invertible*.

The basic dynamic d -logic we consider is **wK4C**, which consists of **wK4** and the temporal axioms for the continuous function f . In addition, we investigate two extensions of **wK4C**: **K4C** and **GLC**. As we will see, **K4C** is the d -logic of all T_D dynamical systems, and **GLC** is the d -logic of all dynamical systems based on a scattered space. Unlike the generic logic of the trimodal topo-temporal language \mathcal{L}_{\diamond}^* , we conjecture that a complete finite axiomatisation for **GLC**, extended with axioms for the ‘henceforth’ operator, will not require changes to the trimodal language. This logic is of special interest to us as it would allow for the first finite axiomatisation and completeness results for a logic based on the trimodal topo-temporal language.

Outline. This paper is structured as follows: in Section 2 we give the required definitions and notations necessary to understand the paper. In Section 3 we provide some background on prior work on the topic of dynamic topological logics. Moreover, we motivate our interest in **GLC**, the most unusual logic we work with.

In Section 4 we present the canonical model, and in Section 5 we construct a ‘finitary’ accessibility relation on it. Both are then used in Section 6 in order to develop a proof technique that, given the right modifications, would work for many d -logics above **wK4C**. In particular, we use it to prove the finite model property, soundness and completeness for the d -logics **wK4C**, **K4C** and **GLC**, with respect to the appropriate classes of Kripke models.

In Section 7 we prove topological d -completeness of **K4C**, **wK4C** and **GLC** with respect to the appropriate classes of dynamical systems. In Section 8 we present logics for systems with homeomorphisms and provide a general completeness result which, in particular, applies to the d -logics **wK4H**, **K4H** and **GLH**. Finally, in Section 9 we provide some concluding remarks.

2 Preliminaries

In this section we review some basic notions required for understanding this paper. We work with the general setting of *derivative spaces*, in order to unify the topological and Kripke semantics of our logics.

Definition 2.1 (topological space). A topological space is a pair $\mathfrak{X} = \langle X, \tau \rangle$, where X is a set and τ is a subset of $\wp(X)$ that satisfies the following conditions:

- $X, \emptyset \in \tau$;
- if $U, V \in \tau$, then $U \cap V \in \tau$;

- if $\mathcal{U} \subseteq \tau$, then $\bigcup \mathcal{U} \in \tau$.

The elements of τ are called open sets, and we say that τ forms a topology on X . A complement of an open set is called a *closed* set.

The main operation on topological spaces we are interested in is the *Cantor derivative*.

Definition 2.2 (Cantor derivative). Let $\mathfrak{X} = \langle X, \tau \rangle$ be a topological space. Given $S \subseteq X$, the *Cantor derivative* of S is the set $d(S)$ of all limit points of S , i.e. $x \in d(S) \iff \forall U \in \tau$ s.t. $x \in U$, $(U \cap S) \setminus \{x\} \neq \emptyset$. We may write $d(S)$ or dS indistinctly.

Given subsets $A, B \subseteq X$, it is not difficult to verify that $d = d_\tau$ satisfies the following properties:

1. $d(\emptyset) = \emptyset$;
2. $d(A \cup B) = d(A) \cup d(B)$;
3. $dd(A) \subseteq A \cup d(A)$.

In fact, these conditions lead to a more general notion of *derivative spaces*:¹

Definition 2.3 (derivative space). A *derivative space* is a pair $\langle X, \rho \rangle$, where X is a set and $\rho: \wp(X) \rightarrow \wp(X)$ is a map satisfying properties 1-3 (with ρ in place of d).

When working with more than one topological space, we will often denote the map ρ on the topological space $\langle X, \tau \rangle$ by ρ_X . The intended derivative spaces discussed in this paper are of the form $\langle X, d_\tau \rangle$. However, there are other examples of derivative spaces. The standard topological closure may be defined by $c(A) = A \cup d(A)$. Then, $\langle X, c \rangle$ is also a derivative space, which satisfies the additional property $A \subseteq c(A)$ (and, *a fortiori*, $cc(A) = c(A)$), which together form Kuratowski's axioms; we call such derivative spaces *closure spaces*. Similarly to the Cantor derivative, we will often denote the closure of the topological space $\langle X, \tau \rangle$ by c_τ .

Another example of derivative spaces comes from weakly transitive Kripke frames. For the sake of succinctness, we call these frames *derivative frames*. Below and throughout the text, we write $\exists x \sqsupset y \varphi$ instead of $\exists x(y \sqsubset x \wedge \varphi)$, and adopt a similar convention for the universal quantifier and other relational symbols.

Definition 2.4 (derivative frame). A *derivative frame* is a pair $\mathfrak{F} = \langle W, \sqsubset \rangle$ where W is a non-empty set and \sqsubset is a *weakly transitive* relation on W , meaning that $w \sqsubset v \sqsubset u$ implies that $w \sqsubseteq u$, where \sqsubseteq is the reflexive closure of \sqsubset .

We chose the notation \sqsubset because it is suggestive of a transitive relation, but remains ambiguous regarding reflexivity, as there may be irreflexive and reflexive points. Note that \sqsubset is weakly transitive iff \sqsubseteq is transitive. Given $A \subseteq W$, we define \downarrow_{\sqsubset} as a map $\downarrow_{\sqsubset}: \wp(W) \rightarrow \wp(W)$ such that

$$\downarrow_{\sqsubset}(A) = \{w \in W : \exists v \sqsupset w(v \in A)\}.$$

The following is then readily verified:

¹Derivative spaces are a special case of *derivative algebras* introduced by Esakia [8], where $\wp(X)$ is replaced by an arbitrary Boolean algebra.

Lemma 2.5. *If $\langle W, \sqsubset \rangle$ is a derivative frame, then $\langle W, \downarrow_{\sqsubset} \rangle$ is a derivative space.*

There is a connection between derivative frames and topological spaces. Given a derivative frame $\langle W, \sqsubset \rangle$, we define a topology τ_{\sqsubset} on W such that $U \in \tau_{\sqsubset}$ iff U is upwards closed under \sqsubset , in the sense that if $w \in U$ and $v \sqsupseteq w$ then $v \in U$. Topologies of this form are *Aleksandroff topologies*. The following is well known and easily verified.

Lemma 2.6. *Let $\langle W, \sqsubset \rangle$ be a derivative frame and $\tau = \tau_{\sqsubset}$. Then, $d_{\tau} = \downarrow_{\sqsubset}$ iff \sqsubset is irreflexive and $c_{\tau} = \downarrow_{\sqsubset}$ iff \sqsubset is reflexive.*

Dynamical systems consist of a topological space equipped with a continuous function. Recall that if $\langle X, \tau \rangle$ and $\langle Y, \nu \rangle$ are topological spaces and $f: X \rightarrow Y$, then f is *continuous* if whenever $U \in \nu$, it follows that $f^{-1}(U) \in \tau$. The function f is *open* if $f(V)$ is open whenever V is open, and f is a *homeomorphism* if f is continuous, open and bijective. It is well known (and not hard to check) that f is continuous iff $c_{\tau}f^{-1}(A) \subseteq f^{-1}c_{\nu}(A)$ for all $A \subseteq Y$. By unfolding the definition of the closure operator, this becomes $f^{-1}(A) \cup d_{\tau}f^{-1}(A) \subseteq f^{-1}(A) \cup f^{-1}d_{\nu}(A)$, or equivalently, $d_{\tau}f^{-1}(A) \subseteq f^{-1}(A) \cup f^{-1}d_{\nu}(A)$. We thus arrive at the following definition.

Definition 2.7. Let $\langle X, \rho_X \rangle$ and $\langle Y, \rho_Y \rangle$ be derivative spaces. We say that $f: X \rightarrow Y$ is *continuous* if for all $A \subseteq Y$, $\rho_X f^{-1}(A) \subseteq f^{-1}(A) \cup f^{-1}\rho_Y(A)$. We say that f is a *homeomorphism* if it is bijective and $\rho_X f^{-1}(A) = f^{-1}\rho_Y(A)$.

It is worth checking that these definitions coincide with their standard topological counterparts.

Lemma 2.8. *If $\langle X, \tau \rangle$ and $\langle Y, \nu \rangle$ are topological spaces with Cantor derivatives d_{τ} and d_{ν} , respectively, and $f: X \rightarrow Y$, then*

1. *if f is continuous as a function between topological spaces, it is continuous as a function between derivative spaces, and*
2. *if f is a homeomorphism as a function between topological spaces, it is a homeomorphism as a function between derivative spaces.*

We are particularly interested in the case where $X = Y$, which leads to the notion of *dynamic derivative system*.

Definition 2.9. A *dynamic derivative system* is a triple $\mathfrak{S} = \langle X, \rho, f \rangle$, where $\langle X, \rho \rangle$ is a derivative space and $f: X \rightarrow X$ is continuous. If f is a homeomorphism, we say that \mathfrak{S} is *invertible*.

If $\mathfrak{S} = \langle X, \rho, f \rangle$ is such that $\rho = d_{\tau}$ for some topology τ , we say that \mathfrak{S} is a *dynamic topological system* and identify it with the triple $\langle X, \tau, f \rangle$. If $\rho = \downarrow_{\sqsubset}$ for some weakly transitive relation \sqsubset , we say that \mathfrak{S} is a *dynamic Kripke frame* and identify it with the triple $\langle X, \sqsubset, f \rangle$.

It will be convenient to characterise dynamic Kripke frames in terms of the relation \sqsubset .

Definition 2.10 (monotonicity and weak monotonicity). Let $\langle W, \sqsubset \rangle$ be a derivative frame. A function $f: W \rightarrow W$ is *monotonic* if $w \sqsubset v$ implies $f(w) \sqsubset f(v)$, and *weakly monotonic* if $w \sqsubset v$ implies $f(w) \sqsubseteq f(v)$.

The function f is *persistent* if it is a bijection and for all $w, v \in W$, $w \sqsubset v$ if and only if $f(w) \sqsubset f(v)$. We say that a Kripke frame is *invertible* if it is equipped with a persistent function.

Lemma 2.11. *If $\langle W, \sqsubset \rangle$ is a derivative frame and $f: W \rightarrow W$, then*

1. *if f is weakly monotonic then it is continuous with respect to \downarrow_{\sqsubset} , and*
2. *if f is persistent then it is a homeomorphism with respect to \downarrow_{\sqsubset} .*

Our goal is to reason about various classes of dynamic derivative systems using the logical framework defined in the next section.

3 Dynamic Topological Logics

In this section we discuss dynamic topological logic in the general setting of dynamic derivative systems. Given a non-empty set PV of propositional variables, the language $\mathcal{L}_{\diamond}^{\circ}$ is defined recursively as follows:

$$\varphi ::= p \mid \varphi \wedge \psi \mid \neg\varphi \mid \diamond\varphi \mid \circ\varphi,$$

where $p \in PV$. It consists of the Boolean connectives \wedge and \neg , the temporal modality \circ , and the modality \diamond for the derivative operator with its dual $\square := \neg\diamond\neg$. The interior modality may be defined as $\boxdot\varphi := \varphi \wedge \square\varphi$.

Definition 3.1 (semantics). A *dynamic derivative model* (DDM) is a quadruple $\mathfrak{M} = \langle X, \rho, f, \nu \rangle$ where $\langle X, \rho, f \rangle$ is a dynamic derivative system and $\nu: PV \rightarrow \wp(X)$ is a valuation function assigning a subset of X to each propositional letter in PV . Given $\varphi \in \mathcal{L}_{\diamond}^{\circ}$, we define the truth set $\|\varphi\| \subseteq X$ of φ inductively as follows:

- $\|p\| = \nu(p)$;
- $\|\neg\varphi\| = X \setminus \|\varphi\|$;
- $\|\varphi \wedge \psi\| = \|\varphi\| \cap \|\psi\|$;
- $\|\diamond\varphi\| = \rho(\|\varphi\|)$;
- $\|\circ\varphi\| = f^{-1}(\|\varphi\|)$.

We write $\mathfrak{M}, x \models \varphi$ if $x \in \|\varphi\|$, and $\mathfrak{M} \models \varphi$ if $\|\varphi\| = X$. We may write $\|\cdot\|_{\mathfrak{M}}$ or $\|\cdot\|_{\nu}$ instead of $\|\cdot\|$ when working with more than one model or valuation.

We define other connectives (e.g. \vee, \rightarrow) as abbreviations in the usual way. The fragment of $\mathcal{L}_{\diamond}^{\circ}$ that includes only \diamond will be denoted by \mathcal{L}_{\diamond} . Since our definition of the semantics applies to any derivative space and a general operator ρ , we need not differentiate in our results between d -logics, logics based on closure semantics and logics based on relational semantics. Instead, we indicate the specific class of derivative spaces to which the result applies.

In order to keep with the familiar axioms of modal logic, it is convenient to discuss the semantics of \square . Accordingly, we define the dual of the derivative, called the *co-derivative*.

Definition 3.2 (co-derivative). Let $\langle X, \rho \rangle$ be a derivative space. For each $S \subseteq X$ we define $\hat{\rho}(S) := X \setminus \rho(X \setminus S)$ to be the *co-derivative* of S .

The co-derivative satisfies the following properties, where $A, B \subseteq X$:

1. $\hat{\rho}(X) = X$;
2. $A \cap \hat{\rho}(A) \subseteq \hat{\rho}(A)$;

$$3. \hat{\rho}(A \cap B) = \hat{\rho}(A) \cap \hat{\rho}(B).$$

It can readily be checked that for each dynamic derivative model $\langle X, \rho, f, \nu \rangle$ and all formulas φ , $\|\Box\varphi\| = \hat{\rho}(\|\varphi\|)$. The co-derivative can be used to define the standard *interior* of a set, given by $i(A) = A \cap \hat{\rho}(A)$ for each $A \subseteq X$. This implies that $U \subseteq \hat{\rho}(U)$ for each open set U , but not necessarily $\hat{\rho}(U) \subseteq U$. Next, we discuss the systems of axioms that are of interest to us. Let us list the axiom schemes and rules that we will consider in this paper:

$$\begin{array}{ll} \text{Taut} := \text{All propositional tautologies} & \text{Next}_\wedge := \bigcirc(\varphi \wedge \psi) \leftrightarrow \bigcirc\varphi \wedge \bigcirc\psi \\ \mathbf{K} := \Box(\varphi \rightarrow \psi) \rightarrow (\Box\varphi \rightarrow \Box\psi) & \mathbf{C} := \bigcirc\varphi \wedge \Box\bigcirc\varphi \rightarrow \Box\bigcirc\varphi \\ \mathbf{T} := \Box\varphi \rightarrow \varphi & \mathbf{H} := \Box\bigcirc\varphi \leftrightarrow \bigcirc\Box\varphi \\ \mathbf{w4} := \varphi \wedge \Box\varphi \rightarrow \Box\Box\varphi & \mathbf{MP} := \frac{\varphi \quad \varphi \rightarrow \psi}{\psi} \\ \mathbf{L} := \Box(\Box\varphi \rightarrow \varphi) \rightarrow \Box\varphi & \mathbf{Nec}_\Box := \frac{\varphi}{\Box\varphi} \\ \mathbf{4} := \Box\varphi \rightarrow \Box\Box\varphi & \mathbf{Nec}_\bigcirc := \frac{\varphi}{\bigcirc\varphi} \\ \mathbf{Next}_\neg := \neg\bigcirc\varphi \leftrightarrow \bigcirc\neg\varphi & \end{array}$$

The ‘base modal logic’ over \mathcal{L}_\diamond is given by

$$\mathbf{K} := \text{Taut} + \mathbf{K} + \mathbf{MP} + \mathbf{Nec}_\Box,$$

but we are mostly interested in proper extensions of \mathbf{K} . Let Λ and Λ' be logics over languages \mathcal{L} and \mathcal{L}' respectively. We say that Λ extends Λ' if $\mathcal{L}' \subseteq \mathcal{L}$ and all the axioms and rules of Λ' are derivable in Λ . A logic over \mathcal{L}_\diamond is *normal* if it extends \mathbf{K} . If Λ is a logic and φ is a formula, $\Lambda + \varphi$ is the least extension of Λ which contains every substitution instance of φ as an axiom.

We then define $\mathbf{wK4} := \mathbf{K} + \mathbf{w4}$, $\mathbf{K4} := \mathbf{K} + \mathbf{4}$, $\mathbf{S4} := \mathbf{K4} + \mathbf{T}$ and $\mathbf{GL} := \mathbf{K4} + \mathbf{L}$. These logics are well known and characterise certain classes of topological spaces and Kripke frames which we review below. In addition, for a logic Λ over \mathcal{L}_\diamond , $\Lambda\mathbf{F}$ is the logic over \mathcal{L}_\diamond^2 given by²

$$\Lambda\mathbf{F} := \Lambda + \mathbf{Next}_\neg + \mathbf{Next}_\wedge + \mathbf{Nec}_\bigcirc.$$

This simply adds axioms of linear temporal logic to Λ , which hold whenever \bigcirc is interpreted using a function. Finally, we define $\Lambda\mathbf{C} := \Lambda\mathbf{F} + \mathbf{C}$ and $\Lambda\mathbf{H} := \Lambda\mathbf{F} + \mathbf{H}$, which as we will see correspond to derivative spaces with a continuous function or a homeomorphism respectively. The following is well known and dates back to McKinsey and Tarski [20].

Theorem 3.3. *$\mathbf{S4}$ is the logic of all topological closure spaces, the logic of all transitive, reflexive derivative frames, and the logic of the real line with the standard closure.*

Analogously, Esakia demonstrated that $\mathbf{wK4}$ is the logic of topological derivative spaces [6].

²Logics of the form $\Lambda\mathbf{F}$ correspond to dynamical systems with a possibly discontinuous function. We will not discuss discontinuous systems in this paper; see [1] for more information.

Theorem 3.4. **wK4** is the logic of all topological derivative spaces, as well as the logic of all weakly-transitive derivative frames.

The logic **K4** includes the axiom $\Box p \rightarrow \Box\Box p$, which is not valid over the class of all topological spaces. The class of spaces satisfying this axiom is denoted by T_D , defined as the class of spaces in which every singleton is the result of an intersection between an open set and a closed set. Moreover, Esakia showed that this is the logic of transitive derivative frames [8].

Theorem 3.5. **K4** is the logic of all T_D topological derivative spaces, as well as the logic of all transitive derivative frames.

Many familiar topological spaces, including Euclidean spaces, satisfy the T_D property, making **K4** central in the study of topological modal logic. A somewhat more unusual class of spaces, which is nevertheless of particular interest to us, is the class of *scattered spaces*.

Definition 3.6 (scattered space). A topological space $\langle X, \tau \rangle$ is *scattered* if for every $S \subseteq X$, $S \subseteq d(S)$ implies $S = \emptyset$.

This is equivalent to the more common definition of a scattered space where a topological space is called scattered if every non-empty subset has an isolated point. Scattered spaces are closely related to converse well-founded relations. Below, recall that $\langle W, \sqsubset \rangle$ is *converse well-founded* if there is no infinite sequence $w_0 \sqsubset w_1 \sqsubset \dots$ of elements in W .

Lemma 3.7. If $\langle W, \sqsubset \rangle$ is an irreflexive frame, then $\langle W, \tau_{\sqsubset} \rangle$ is scattered iff \sqsubset is converse well-founded.

Theorem 3.8 (Simmons [23] and Esakia [7]). **GL** is the logic of all scattered topological derivative spaces, as well as the logic of all converse well-founded derivative frames and the logic of all finite, transitive, irreflexive derivative frames.

Aside from its topological interpretation, the logic **GL** is of particular interest as it is also the logic of provability in Peano arithmetic, as was shown by Boolos [3]. Meanwhile, logics with the C and H axioms correspond to classes of dynamical systems.

Lemma 3.9.

1. If Λ is sound for a class of derivative spaces Ω , then $\Lambda\mathbf{C}$ is sound for the class of dynamic derivative systems $\langle X, \rho, f \rangle$, where $\langle X, \rho \rangle \in \Omega$ and f is continuous.
2. If Λ is sound for a class of derivative spaces Ω , then $\Lambda\mathbf{H}$ is sound for the class of dynamic derivative systems $\langle X, \rho, f \rangle$, where $\langle X, \rho \rangle \in \Omega$ and f is a homeomorphism.

The above lemma is easy to verify from the definitions of continuous functions and homeomorphisms in the context of derivative spaces (Definition 2.7).

3.1 Prior Work

The study of dynamic topological logic originates with Artemov, Davoren and Nerode, who observed that it is possible to reason about dynamical systems within modal logic. They introduced the logic **S4C** and proved that it is decidable, as well as sound and complete for the class of all dynamic closure systems (i.e. dynamic derivative systems based on a closure space). Kremer and Mints [17] considered the logic **S4H** and showed that it is sound and complete for the class of dynamic closure systems where f is a homeomorphism.

The latter also suggested adding the ‘henceforth’ operator, $*$, from Pnueli’s linear temporal logic (**LTL**) [22], leading to the language we denote by $\mathcal{L}_{\diamond}^{\circ*}$. The resulting trimodal system was named *dynamic topological logic* (**DTL**). Kremer and Mints offered an axiomatisation for **DTL**, but Fernández-Duque proved that it is incomplete; in fact, **DTL** is not finitely axiomatisable [9]. Fernández-Duque also showed that **DTL** has a natural axiomatisation when extended with the *tangled closure* [12]. In contrast, Konev et al. established that **DTL** over the class of dynamical systems with a homeomorphism is non-axiomatisable [16].

3.2 The tangled closure on scattered spaces

Our interest in considering the class of scattered spaces within dynamic topological logic is motivated by results of Fernández-Duque [12]. He showed that the set of valid formulas of $\mathcal{L}_{\diamond}^{\circ*}$ over the class of all dynamic closure systems is not finitely axiomatisable. Nevertheless, he found a natural (yet infinite) axiomatisation by introducing the *tangled closure* and adding it to the language of **DTL** [10]. Here, we use the more general *tangled derivative*, as defined by Goldblatt and Hodkinson [13].

Definition 3.10 (tangled derivative). Let $\langle X, \rho \rangle$ be a derivative space and let $\mathcal{S} \subseteq \wp(X)$. Given $A \subseteq X$, we say that \mathcal{S} is tangled in A if for all $S \in \mathcal{S}$, $A \subseteq \rho(S \cap A)$. We define the *tangled derivative* of \mathcal{S} as

$$\mathcal{S}^* := \bigcup \{A \subseteq X : \mathcal{S} \text{ is tangled in } A\}.$$

The *tangled closure* is then the special case of the tangled derivative where ρ is a closure operator, i.e. $\rho = c$. Fernández-Duque’s axiomatisation is based on the extended language $\mathcal{L}_{\diamond}^{\circ*}$. This language is obtained by extending $\mathcal{L}_{\diamond}^{\circ*}$ with the following operation.

Definition 3.11 (tangled language). We define $\mathcal{L}_{\diamond}^{\circ*}$ by extending the recursive definition of $\mathcal{L}_{\diamond}^{\circ*}$ in such a way that if $\varphi_1, \dots, \varphi_n \in \mathcal{L}_{\diamond}^{\circ*}$, then $\diamond\{\varphi_1, \dots, \varphi_n\} \in \mathcal{L}_{\diamond}^{\circ*}$. The semantic clauses are then extended so that on every model \mathfrak{M} ,

$$\|\diamond\{\varphi_1, \dots, \varphi_n\}\| = \{\|\varphi_1\|, \dots, \|\varphi_n\|\}^*.$$

The logic **DGL** is an extension of **GLC** that includes the temporal operator $*$. Unlike the complete axiomatisation of **DTL** that requires the tangled operator, in the case of **DGL**, we should be able to avoid this and use the original spatial operator \diamond alone. This is due to the following:

Theorem 3.12. *Let $\mathfrak{X} = \langle X, \tau \rangle$ be a scattered space and $\{\varphi_1, \dots, \varphi_n\}$ a set of formulas. Then*

$$\diamond\{\varphi_1, \dots, \varphi_n\} \equiv \perp.$$

Proof. Let ν be a valuation on \mathfrak{X} , $\mathfrak{M} = \langle \mathfrak{X}, \nu \rangle$, and suppose that $\mathfrak{M}, x \models \diamond\{\varphi_1, \dots, \varphi_n\}$ for some $x \in X$. Then by definition there exists $S \subseteq X$ s.t. $x \in S$ and $S \subseteq d(S \cap \llbracket \varphi_i \rrbracket)$, for all $i \leq n$. It follows that $S \subseteq d(S)$. Since \mathfrak{X} is scattered, then $S = \emptyset$ in contradiction. It follows that $\mathfrak{M}, x \not\models \diamond\{\varphi_1, \dots, \varphi_n\}$ for all $x \in X$. \square

This leads to the conjecture that the axiomatic system of Kremer and Mints [17], combined with **GL**, will lead to a finite axiomatisation for **DGL**. While such a result requires techniques beyond the scope of the present work, the completeness proof we present here for **GLC** is an important first step. Before proving topological completeness for this and the other logics we have mentioned, we show that they are complete and have the finite model property for their respective classes of dynamic derivative frames.

4 The Canonical Model

The first step in our Kripke completeness proof will be a fairly standard canonical model construction. A *maximal Λ -consistent set* (Λ -MCS) w is a set of formulas that is Λ -consistent, i.e. $w \not\vdash_{\Lambda} \perp$, and every set of formulas that properly contains it is Λ -inconsistent.

Given a logic Λ over $\mathcal{L}_{\diamond}^{\circ}$, let $\mathfrak{M}_c^{\Lambda} = \langle W_c, \sqsubset_c, g_c, \nu_c \rangle$ be the canonical model for Λ , where

1. W_c is the set of all Λ -MCSs;
2. $w \sqsubset_c v$ iff for all formulas φ , if $\Box\varphi \in w$, then $\varphi \in v$;
3. $g_c(w) = \{\varphi : \bigcirc\varphi \in w\}$;
4. $\nu_c(p) = \{w : p \in w\}$.

It can easily be verified that **wK4C** defines the class of all weakly transitive Kripke models with a weakly monotonic map. Moreover, **K4C** defines the class of all transitive Kripke models with a weakly monotonic map. We call these models **wK4C** models and **K4C** models respectively.

Lemma 4.1. *If Λ extends **wK4C**, then the canonical model for Λ is a **wK4C** model. If Λ extends **K4C**, then the canonical model of Λ is a **K4C** model.*

Proof. Let $\mathfrak{M}_c^{\Lambda} = \langle W, \sqsubset, g, \nu \rangle$. Suppose that Λ extends **wK4C**. We prove that g is weakly monotonic. Suppose $w \sqsubset v$ and $g(w) \neq g(v)$. Since $g(w) \neq g(v)$, there exists φ such that $\varphi \in g(w)$ and $\varphi \notin g(v)$. We consider an arbitrary $\Box\psi \in g(w)$ then clearly $\Box(\psi \vee \varphi) \in g(w)$ and $(\psi \vee \varphi) \in g(w)$. In particular, $\bigcirc\Box(\psi \vee \varphi) \in w$ and $\bigcirc(\psi \vee \varphi) \in w$. Since $(\bigcirc\Box p \wedge \bigcirc p) \rightarrow \Box\bigcirc p \in \mathbf{wK4C}$, then $\Box\bigcirc(\varphi \vee \psi) \in w$. Since $w \sqsubset v$, then $\bigcirc(\varphi \vee \psi) \in v$ and $(\varphi \vee \psi) \in g(v)$. Since $\varphi \notin g(v)$, then $\psi \in g(v)$ and because $\Box\psi$ is arbitrary, then $g(w) \sqsubset g(v)$, as required. Hence g is weakly monotonic.

We prove that \sqsubset is weakly transitive. Suppose $w \sqsubset v \sqsubset u$ and $w \neq u$. From $w \neq u$ it follows that there exists φ such that $\varphi \in w$ and $\varphi \notin u$. We consider an arbitrary $\Box\psi \in w$, then clearly $\Box(\psi \vee \varphi) \in w$ and $(\psi \vee \varphi) \in w$. Since $(\varphi \wedge \Box\varphi) \rightarrow \Box\Box\varphi \in \mathbf{wK4C}$, then $\Box\Box(\psi \vee \varphi) \in w$. Then $w \sqsubset v \sqsubset u$ implies

$(\psi \vee \varphi) \in u$ and as $\varphi \notin u$ we have $\psi \in u$. Since $\Box\psi$ is arbitrary, then $w \sqsubset u$ holds, as required.

It follows that g is weakly monotonic and \sqsubset is weakly transitive and hence \mathfrak{M}_c^Λ is a **wK4C** model.

Suppose that Λ extends **K4C**, then weak monotonicity holds as before since **K4C** extends **wK4C**. Therefore, we only need to prove that \sqsubset is transitive. Suppose $w \sqsubset v \sqsubset u$. We consider an arbitrary $\Box\psi \in w$. Since $\Box p \rightarrow \Box\Box p \in \mathbf{wK4C}$, then $\Box\Box\psi \in w$. Then, $w \sqsubset v \sqsubset u$ implies $\psi \in u$ and since $\Box\psi$ is arbitrary, then $w \sqsubset u$, as required.

It follows that g is weakly monotonic and \sqsubset is transitive and thus \mathfrak{M}_c^Λ is a **K4C** model. \square

It is a well-known fact that the transitivity axiom $\Box p \rightarrow \Box\Box p$ is derivable in **GL** (see [24]). Therefore, **GLC** extends the system **K4C**. The proofs of the following two lemmas are standard and can be found for example in [2].

Lemma 4.2 (existence lemma). *Let Λ be a normal modal logic and let $\mathfrak{M}_c^\Lambda = \langle W_c, \sqsubset_c, g_c, \nu_c \rangle$. Then, for every $w \in W_c$ and every formula φ in Λ , if $\Diamond\varphi \in w$ then there exists a point $v \in W_c$ such that $w \sqsubset_c v$ and $\varphi \in v$.*

Lemma 4.3 (truth lemma). *Let Λ be a normal modal logic. For every $w \in W_c$ and every formula φ in Λ ,*

$$\mathfrak{M}_c^\Lambda, w \models \varphi \text{ iff } \varphi \in w.$$

Corollary 4.4. *The logic **wK4C** is sound and complete with respect to the class of all weakly monotonic dynamic derivative frames, and **K4C** is sound and complete with respect to the class of all weakly monotonic, transitive dynamic derivative frames.*

5 A finitary accessibility relation

One key ingredient in our finite model property proof will be the construction of a ‘finitary’ accessibility relation \sqsubset_Φ on the canonical model. This accessibility relation will have the property that each point has finitely many successors, yet the existence lemma will hold for formulas in a prescribed finite set Φ .

We define the \sqsubset_c -cluster $C(w)$ for each point $w \in W_c$ as

$$C(w) = \{w\} \cup \{v : w \sqsubset_c v \sqsubset_c w\}.$$

Definition 5.1 (φ -final set). A set w is said to be a φ -final set (or point) if w is an MCS, $\varphi \in w$, and whenever $w \sqsubset_c v$ and $\varphi \in v$, it follows that $v \in C(w)$.

Let us write \sqsubseteq_c for the reflexive closure of \sqsubset_c . It will be convenient to characterise \sqsubseteq_c in the canonical model syntactically. Recall that $\Box\varphi := \varphi \wedge \Box\varphi$.

Lemma 5.2. *If Λ extends **wK4C** and $w, v \in W_c$, then $w \sqsubseteq_c v$ if and only if whenever $\Box\varphi \in w$, it follows that $\Box\varphi \in v$.*

Proof. First assume that $w \sqsubseteq_c v$. Obviously if $w = v$ then $\Box\varphi \in w$ implies $\Box\varphi \in v$. If $\Box\varphi \in w$ then from $\Box\varphi \in w$ we obtain $\varphi \in v$ and by w4 we have $\Box\Box\varphi \in w$ and so $\Box\varphi \in v$.

Conversely, assume that $\Box\varphi \in w$ implies $\Box\varphi \in v$. If $w = v$, then $w \sqsubseteq_c v$. Otherwise, $w \neq v$, and we can find $\psi \in w \setminus v$. Suppose that $\Box\varphi \in w$. Then $\Box(\varphi \vee \psi) \in w$ and so $\Box(\varphi \vee \psi) \in v$. It follows that $\varphi \vee \psi \in v$. But $\psi \notin v$, so $\varphi \in v$; since φ was arbitrary, then $w \sqsubseteq_c v$, as required. \square

The following version of Zorn's lemma will be used to prove an important existence property.

Lemma 5.3 (Zorn's Lemma). *Let (A, \leq) be a preordered set where A is non-empty. A chain is a subset $C \subseteq A$ whose elements are totally ordered by \leq . Suppose that every chain C has an upper bound in A . Then, A has a \leq -maximal element.*

The following proves a sufficient condition for the existence of a final point accessible from a given point.

Lemma 5.4. *If $\Diamond\varphi \in w$, then there is φ -final point v such that $w \sqsubseteq_c v$.*

Proof. Suppose that \mathcal{C} is an \sqsubseteq_c -chain in $A := \sqsubseteq_c(w) \cap \|\varphi\|$ (where $\sqsubseteq_c(w) = \{v \in W_c : w \sqsubseteq_c v\}$). We show that there is an upper bound of \mathcal{C} that belongs to A .

Choose a formula θ as follows: If w is an irreflexive world (i.e. $w \not\sqsubseteq_c w$), then choose θ so that $\Box\theta \wedge \neg\theta \in w$. Otherwise, w is reflexive and we simply set $\theta = \top$. Let Γ be the set

$$\{\Box\psi : \exists x \in \mathcal{C}(\Box\psi \in x)\}.$$

Suppose $\Box\psi_1, \dots, \Box\psi_n \in \Gamma$. For each i there is $w_i \in \mathcal{C}$ such that $\Box\psi_i \in w_i$. Since \mathcal{C} is a chain, for some j we have that $w_j = \max_{i=1}^n w_i$. Then, $\varphi, \theta, \Box\psi_1, \dots, \Box\psi_n \in w_j$, hence $\{\varphi, \theta, \Box\psi_1, \dots, \Box\psi_n\}$ is consistent. Since $\{\Box\psi_1, \dots, \Box\psi_n\}$ is an arbitrary finite subset of Γ , we have that $\Gamma \cup \{\varphi, \theta\}$ is consistent.

By the Lindenbaum lemma we can extend $\Gamma \cup \{\varphi, \theta\}$ to an MCS, which we denote by w^* . Suppose $\Box\psi \in v \in \mathcal{C}$, then by construction $\Box\psi \in w^*$. It follows by Lemma 5.2 that $v \sqsubseteq_c w^*$. Hence w^* is an upper bound for \mathcal{C} . Note that the assumption that $\Diamond\varphi \in w$ implies that there is some $v \sqsupset_c w$ such that $\varphi \in v$, that is, $v \in A$. Hence $w \sqsubseteq_c v \sqsubseteq_c w^*$, which yields $w \sqsubseteq_c w^*$. To see that indeed $w \sqsubseteq_c w^*$, if w is reflexive there is nothing to prove, otherwise since $\theta \in w^*$ we have that $w \neq w^*$. Moreover $\varphi \in w^*$ by construction, so $w^* \in A$.

Thus, by Lemma 5.3 we conclude that A contains a maximal element, and this maximal element is clearly a φ -final point above w . \square

We are now ready to prove the main result of this section regarding the existence of the finitary relation \sqsubseteq_Φ .

Lemma 5.5. *Let Λ extend **wK4C** and Φ be a finite set of formulas closed under subformulas. There is an auxiliary relation \sqsubseteq_Φ on the canonical model of Λ such that:*

- (i) \sqsubseteq_Φ is a subset of \sqsubseteq_c ;
- (ii) For each $w \in W$, the set $\sqsubseteq_\Phi(w)$ is finite;
- (iii) If $\Diamond\varphi \in w \cap \Phi$, then there exists $v \in W$ with $w \sqsubseteq_\Phi v$ and $\varphi \in v$;

(iv) If $w \sqsubset_c v \sqsubset_c w$ then $\sqsubset_\Phi(w) \subseteq \sqsubset_\Phi(v)$;

(v) \sqsubset_Φ is weakly transitive. Moreover, if Λ extends **K4C** then \sqsubset_Φ is transitive, and if Λ extends **GLC** then \sqsubset_Φ is irreflexive.

Proof. Let C be any cluster of points in W and define

$$\sqsubset_c(C) = \bigcup \{\sqsubset_c(v) : v \in C\}.$$

We construct the weakly transitive relation \sqsubset_Φ as follows: Using Lemma 5.4 we use the axiom of choice to choose a function that for each formula φ and each cluster C such that $\diamond\varphi \in \bigcup C$, assigns a φ -final point $w(\varphi, C)$ such that $w(\varphi, C) \in \sqsubset_c(C)$. We choose a second point $w'(\varphi, C)$, possibly equal to $w(\varphi, C)$, such that

- if $\diamond\varphi \in w(\varphi, C)$, then $w'(\varphi, C)$ is any φ -final point such that $w(\varphi, C) \sqsubset_c w'(\varphi, C)$;
- otherwise, $w'(\varphi, C) = w(\varphi, C)$.

Set $u \sqsubset_\Phi^0 v$ iff $u \sqsubset_c v$ and there exists $\psi \in \Phi$ such that $\diamond\psi \in u$ and $v \in \{w(\psi, C(u)), w'(\psi, C(u))\}$. Let \sqsubset_Φ be the weakly transitive closure of \sqsubset_Φ^0 .

It is clear that (i), (iii) and (iv) follow directly from the construction: (i) follows from the fact that \sqsubset_Φ is the transitive closure of \sqsubset_Φ^0 ; (iii) follows from Lemma 5.1; (iv) follows from the fact that $v \in C(w)$ and by assuming that $v \sqsubset_\Phi u$ and unraveling the definition of \sqsubset_Φ .

We continue to verify conditions (ii) and (v). For condition (ii), first observe that $\sqsubset_\Phi^0(u)$ is finite by construction, as it contains at most two elements for each $\varphi \in \Phi$. Now, if $u \sqsubset_\Phi v$ then there is a sequence

$$u \sqsubset_\Phi^0 v_1 \sqsubset_\Phi^0 \dots \sqsubset_\Phi^0 v_n = v.$$

By taking a minimal such sequence, we may assume that it is injective. Consider the tree consisting of all such sequences (ordered by the initial segment relation). This is a finitely-branching tree, as $\sqsubset_\Phi^0(x)$ is always finite. Moreover, if $\sqsubset_\Phi(u)$ is infinite, then this tree is infinite. By König's lemma, there is an infinite sequence

$$u \sqsubset_\Phi^0 v_1 \sqsubset_\Phi^0 v_2 \sqsubset_\Phi^0 \dots$$

By definition of \sqsubset_Φ^0 , for each $i \in \omega$ there is $\varphi_i \in \Phi$ such that v_{i+1} is φ_i -final. Since $\sqsubset_\Phi^0 \subseteq \sqsubset_c$, we have that $v_i \sqsubset_c v_j$ whenever $i \leq j$. Since Φ is finite, there is some $\theta \in \Phi$ such that v_i is θ -final for infinitely many values of i . Let i_0 be the least such value. If $i > i_0$ is any other such value, $v_{i_0} \sqsubset_c v_i$ together with θ -finality of v_{i_0} yields $v_i \sqsubset_c v_{i_0}$. Thus $v_i \in C(v_{i_0})$ and $v_i \sqsubset_\Phi^0 v_{i+1}$, which by definition of \sqsubset_Φ^0 yields $v_{i+1} \in \sqsubset_\Phi^0(v_{i_0})$. But $\sqsubset_\Phi^0(v_{i_0})$ is finite, contradicting that the chain is infinite and injective.

Finally, we verify condition (v). The relation \sqsubset_Φ is weakly transitive by definition, and if Λ extends **K4C**, we have that $x \sqsubset_\Phi y \sqsubset_\Phi z$ implies $x \sqsubset_\Phi z$, which in the case that $x = z$ implies that $x \sqsubset_c x$ and also that x, y are in the same cluster. This together with $y \sqsubset_\Phi z$ implies that $x \sqsubset_\Phi z$, as \sqsubset_Φ is cluster-invariant.

Now let Λ extend **GLC**, and suppose that w is φ -final. We claim that $\diamond\varphi \notin w$, which immediately yields $w \neq \sqsubset_\Phi w$, as needed. By the contrapositive

of L , we have that $\diamond\varphi \rightarrow \diamond(\varphi \wedge \Box\neg\varphi) \in w$. If $\diamond\varphi \in w$, then $\diamond(\varphi \wedge \Box\neg\varphi) \in w$. It follows that there is $v \sqsupset_{\Phi} w$ with $\varphi \wedge \Box\neg\varphi \in v$. But $\Box\neg\varphi \in v$ implies that $v \not\sqsupset_{\Phi} w$, contradicting the φ -finality of w . So $\diamond\varphi \notin w$ and w is irreflexive, as required. \square

6 Stories and Φ -morphisms

In this section we show that the logics **wK4C**, **K4C** and **GLC** have the finite model property by constructing finite models and truth preserving maps from these models to the canonical model.

If \sqsubseteq is a weakly transitive relation on A , $\langle A, \sqsubseteq \rangle$ is tree-like if whenever $a \sqsubseteq c$ and $b \sqsubseteq c$, it follows that $a \sqsubseteq b$ or $b \sqsubseteq a$. We will use labelled tree-like structures called *moments* to record the ‘static’ information at a point; that is, the structure involving \sqsubseteq , but not f .

Definition 6.1 (moment). A Λ -*moment* is a structure $\mathbf{m} = \langle |\mathbf{m}|, \sqsubseteq_{\mathbf{m}}, \nu_{\mathbf{m}}, r_{\mathbf{m}} \rangle$, where $\langle |\mathbf{m}|, \sqsubseteq_{\mathbf{m}} \rangle$ is a finite tree-like Λ frame with a root $r_{\mathbf{m}}$, and $\nu_{\mathbf{m}}$ is a valuation on $|\mathbf{m}|$.

In order to also record ‘dynamic’ information, i.e. information involving the transition function, we will stack up several moments together to form a ‘story’. Below, \sqcup denotes a disjoint union.

Definition 6.2 (story). A *story* (with duration I) is a structure $\mathfrak{S} = \langle |\mathfrak{S}|, \sqsubseteq_{\mathfrak{S}}, f_{\mathfrak{S}}, \nu_{\mathfrak{S}}, r_{\mathfrak{S}} \rangle$ such that there are $I < \omega$, moments $\mathfrak{S}_i = \langle |\mathfrak{S}_i|, \sqsubseteq_i, \nu_i, r_i \rangle$ for each $i \leq I$, and functions $(f_i)_{i < I}$ such that:

1. $|\mathfrak{S}| = \sqcup_{i \leq I} |\mathfrak{S}_i|$;
2. $\sqsubseteq_{\mathfrak{S}} = \sqcup_{i \leq I} \sqsubseteq_i$;
3. $\nu_{\mathfrak{S}}(p) = \sqcup_{i \leq I} \nu_i(p)$ for each variable p ;
4. $r_{\mathfrak{S}} = r_0$;
5. $f_{\mathfrak{S}} = \text{Id}_I \cup \sqcup_{i < I} f_i$ with $f_i: |\mathfrak{S}_i| \rightarrow |\mathfrak{S}_{i+1}|$ being a weakly monotonic map such that $f_i(r_i) = r_{i+1}$ for all $i < I$ (we say that f_i is *root preserving*), and Id_I is the identity on $|\mathfrak{S}_I|$.




Figure 1: An example of a **GL**-story. The squiggly arrows represent the relation $\sqsubseteq_{\mathfrak{S}}$ while the straight arrows represent the function $f_{\mathfrak{S}}$. Each vertical slice represents a **GL**-moment. In the case of other types of stories, we may also have clusters besides singletons.

We often omit the subindices \mathbf{m} or \mathfrak{S} when this does not lead to confusion. We may also assign different notations to the components of a moment, so that if we write $\mathbf{m} = \langle W, \sqsubset, \nu, x \rangle$, it is understood that $W = |\mathbf{m}|$, $\sqsubset = \sqsubset_{\mathbf{m}}$, etc.

Recall that a *p-morphism* between Kripke models is a type of map that preserves validity. It can be defined in the context of dynamic derivative frames as follows:

Definition 6.3 (dynamic *p*-morphism). Let $\mathfrak{M} = \langle W_{\mathfrak{M}}, \sqsubset_{\mathfrak{M}}, g_{\mathfrak{M}} \rangle$ and $\mathfrak{N} = \langle W_{\mathfrak{N}}, \sqsubset_{\mathfrak{N}}, g_{\mathfrak{N}} \rangle$ be dynamic derivative frames. Let $\pi: W_{\mathfrak{M}} \rightarrow W_{\mathfrak{N}}$. We say that π is a *dynamic p-morphism* if $w \sqsubset_{\mathfrak{M}} v$ implies that $\pi(w) \sqsubset_{\mathfrak{N}} \pi(v)$, $\pi(w) \sqsubset_{\mathfrak{N}} u$ implies that there is $v \sqsupset_{\mathfrak{M}} w$ with $\pi(v) = u$, and $\pi \circ g_{\mathfrak{M}} = g_{\mathfrak{N}} \circ \pi$.

It is then standard that if $\pi: W_{\mathfrak{M}} \rightarrow W_{\mathfrak{N}}$ is a surjective, dynamic *p*-morphism, then any formula valid on \mathfrak{M} is also valid on \mathfrak{N} . However, our relation \sqsubset_{Φ} will allow us to weaken these conditions and still obtain maps that preserve the truth of (some) formulas.

Definition 6.4 (Φ -morphism). Fix a logic Λ and let $\mathfrak{M}_c^{\Lambda} = \langle W_c, \sqsubset_c, g_c, \nu_c \rangle$ and \mathfrak{S} be a story of duration I . A map $\pi: |\mathfrak{S}| \rightarrow W_c$ is called a *dynamic Φ -morphism* if for all $x \in |\mathfrak{S}|$ the following conditions are satisfied:

1. $x \in \nu_{\mathfrak{S}}(p) \iff p \in \pi(x)$;
2. If $x \in |\mathfrak{S}_i|$ for some $i < I$, then $g_c(\pi(x)) = \pi(f_{\mathfrak{S}}(x))$;
3. If $x \sqsubset_{\mathfrak{S}} y$ then $\pi(x) \sqsubset_c \pi(y)$;
4. If $\pi(x) \sqsubset_{\Phi} v$ for some $v \in W_c$, then there exists $y \in |\mathfrak{S}|$ such that $x \sqsubset_{\mathfrak{S}} y$ and $v = \pi(y)$.

If we drop condition 2, we say that π is a *Φ -morphism*.

We now show that a dynamic Φ -morphism π preserves the truth of formulas of suitable \circ -depth, where the latter is defined as usual in terms of nested occurrences of \circ in a formula φ .

Lemma 6.5 (truth preservation). *Let \mathfrak{S} be a story of duration I and $x \in |\mathfrak{S}_0|$. Let π be a dynamic Φ -morphism to the canonical model of some normal logic Λ over $\mathcal{L}_{\diamond}^{\circ}$. Suppose that $\varphi \in \Phi$ is a formula of \circ -depth at most I . Then $\varphi \in \pi(x)$ iff $x \in \|\varphi\|_{\mathfrak{S}}$.*

Proof. We prove the more general claim that if φ has \circ -depth at most $I - j$ and $x \in |\mathfrak{S}_j|$, then $\varphi \in \pi(x)$ iff $x \in \|\varphi\|_{\mathfrak{S}}$. We proceed by induction on the complexity of φ . For the base case, suppose $p \in \pi(x)$. Then by the definition of a dynamic Φ -morphism we have $x \in \nu_{\mathfrak{S}}(p)$ as required. The other direction follows similarly, and since the Boolean connectives follow by standard arguments, then we focus on modalities.

If $\Box\varphi \in \pi(x)$ and $y \sqsupset_{\mathfrak{S}} x$ then $\pi(y) \sqsupset_c \pi(x)$, so $\varphi \in \pi(y)$ and the induction hypothesis and the fact that $y \in |\mathfrak{S}_j|$ (as \mathfrak{S}_j is closed under $\sqsubset_{\mathfrak{S}}$) yields $y \in \|\varphi\|_{\mathfrak{S}}$. Since y was arbitrary, $x \in \|\Box\varphi\|_{\mathfrak{S}}$.

If $\Diamond\varphi \in \pi(x)$ then there exists $v \in W_c$ such that $\pi(x) \sqsubset_{\Phi} v$ and $\varphi \in v$. By the definition of a dynamic Φ -morphism there exists $v' \in |\mathfrak{S}|$ such that $x \sqsubset_{\mathfrak{S}} v'$ and $v = \pi(v')$. By the induction hypothesis $v' \in \|\varphi\|_{\mathfrak{S}}$ and thus $x \in \|\Diamond\varphi\|_{\mathfrak{S}}$.

If $\circ\varphi \in \pi(x)$ then $\varphi \in g_c(\pi(x))$. By the definition of a dynamic Φ -morphism we get $g_c(\pi(x)) = \pi(f_{\mathfrak{S}}(x))$. Note that $f_{\mathfrak{S}}(x) \in |\mathfrak{S}_{j+1}|$, but the \circ -depth of φ is bounded by $I - j - 1$. From the induction hypothesis we get $f_{\mathfrak{S}}(x) \in \|\varphi\|_{\mathfrak{S}}$. Hence by definition $x \in \|\circ\varphi\|_{\mathfrak{S}}$.

If $x \in \|\circ\varphi\|_{\mathfrak{S}}$, then $f_{\mathfrak{S}}(x) \in \|\varphi\|_{\mathfrak{S}}$. The bound on \circ -depth can be checked as above, so from the induction hypothesis it follows that $\varphi \in \pi(f_{\mathfrak{S}}(x))$ and by the definition of a dynamic Φ -morphism we get $\pi(f_{\mathfrak{S}}(x)) = g_c(\pi(x))$. Therefore $\circ\varphi \in \pi(x)$. \square

We next demonstrate that for every point w in the canonical model, there exists a suitable moment \mathfrak{m} and a Φ -morphism mapping \mathfrak{m} to w . In order to do this, we define a procedure for constructing new moments from smaller ones.

Definition 6.6 (moment construction). Let $\Lambda \in \{\mathbf{wK4}, \mathbf{K4}, \mathbf{GL}\}$ and $C' \cup \{x\} \subseteq C(x)$ for some x in the canonical model \mathfrak{M}_c^Λ . Let $\vec{\mathfrak{a}} = \langle \mathfrak{a}_m \rangle_{m < N}$ be a sequence of moments. We define a structure $\mathfrak{n} = \left(\frac{\vec{\mathfrak{a}}}{C'}\right)_x$ as follows:

1. $|\mathfrak{n}| = C' \sqcup \bigsqcup_{m < N} |\mathfrak{a}_m|$;
2. $y \sqsubset_{\mathfrak{n}} z$ if either
 - $y, z \in C'$, $\Lambda \neq \mathbf{GL}$ and $y \sqsubset_c z$,
 - $y \in C'$ and $z \in |\mathfrak{a}_m|$ for some m , or
 - $y, z \in |\mathfrak{a}^m|$ and $y \sqsubset_{\mathfrak{a}_m} z$ for some m ;
3. $\nu_{\mathfrak{n}}(p) = \{x \in C' : x \in \nu_c(p)\} \sqcup \bigsqcup_{m < N} \nu_{\mathfrak{a}_m}(p)$;
4. $r_{\mathfrak{n}} = x$.

The moment construction for some logic Λ can be used to produce Λ -moments.

Lemma 6.7. *Let $\Lambda \in \{\mathbf{wK4}, \mathbf{K4}, \mathbf{GL}\}$ and $C' \subseteq C(x)$ for some x in the canonical model \mathfrak{M}_c^Λ , where C' is a singleton if $\Lambda = \mathbf{GL}$ and finite otherwise. Let $\vec{\mathfrak{a}} = \langle \mathfrak{a}^m \rangle_{m < N}$ be a sequence of Λ -moments. Then, $\left(\frac{\vec{\mathfrak{a}}}{C'}\right)_x$ is a Λ -moment.*

Proof. If $\Lambda = \mathbf{wK4}$ then \sqsubset is easily seen to be weakly transitive since each \mathfrak{a}_i is weakly transitive and the root sees all other points. The transitivity of \sqsubset where $\Lambda \supseteq \mathbf{K4}$ follows by the same reasoning. The irreflexivity of \sqsubset where $\Lambda = \mathbf{GL}$ follows from the fact that C' is an irreflexive singleton and each \mathfrak{a}_i is irreflexive. \square

We use this moment construction to show the existence of a Φ -morphism between a moment and the canonical model. Below, we denote by $C_{\Phi}(w)$ the \sqsubset_{Φ} -cluster of w , i.e.

$$C_{\Phi}(w) = \{w\} \cup \{v : w \sqsubset_{\Phi} v \sqsubset_{\Phi} w\}.$$

Lemma 6.8. *Given $\Lambda \in \{\mathbf{wK4C}, \mathbf{K4C}, \mathbf{GLC}\}$, for all $w \in W_c$ there exists a Λ -moment \mathfrak{m} and a Φ -morphism $\pi : |\mathfrak{m}| \rightarrow W_c$ such that $\pi(r_{\mathfrak{m}}) = w$.*

Proof. We prove the stronger claim, that there is a moment \mathbf{m} and a map $\pi : |\mathbf{m}| \rightarrow W_c$ that is a p -morphism on the structure (W_c, \sqsubset_Φ) (we will say that π is a p -morphism with respect to \sqsubset_Φ). Let \sqsubset_Φ^1 be the strict \sqsubset_Φ successor, i.e. $w \sqsubset_\Phi^1 v$ iff $w \sqsubset_\Phi v$ and $\neg(v \sqsubset_\Phi w)$. Since \sqsubset_Φ^1 is converse well-founded, we can assume inductively that for each v such that $w \sqsubset_\Phi^1 v$, there is a moment \mathbf{m}_v and a p -morphism $\pi_v : |\mathbf{m}_v| \rightarrow W_c$ with respect to \sqsubset_Φ that maps the root of \mathbf{m}_v to v . Accordingly, we define a moment

$$\mathbf{m} = \left(\begin{array}{c} \{\mathbf{m}_v : v \sqsubset_\Phi^1 w\} \\ C_\Phi(w) \end{array} \right)_w ;$$

note that $C_\Phi(w)$ is finite and, if $\Lambda = \mathbf{GLC}$, it is a singleton, so \mathbf{m} is a Λ -moment by Lemma 6.7.

Next we define a map $\pi : |\mathbf{m}| \rightarrow W$ as

$$\pi(x) = \begin{cases} x & \text{if } x \in C_\Phi(w), \\ \pi_v(x) & \text{if } x \in |\mathbf{m}_v|. \end{cases}$$

We prove that π is a p -morphism for \sqsubset_Φ . First assume that $\pi(x) \sqsubset_\Phi v$; we must show that there is $y \sqsubset_{\mathbf{m}} x$ so that $\pi(y) = v$. Either $x \in C_\Phi(w)$ or $x \in |\mathbf{m}_u|$ for some u . In the first case, we consider two sub-cases. If $v \in C_\Phi(w)$ as well, then $\pi(v) = v$ and we may set $y := v$. If not, $\pi(x) \sqsubset_\Phi^1 v$, and by letting y be the root of \mathbf{m}_v , we see that $\pi(y) = v$. If instead $x \in |\mathbf{m}_u|$ for some u , then by assumption π_u is a p -morphism, immediately yielding the desired y (also satisfying $y \in |\mathbf{m}_u|$).

Now, suppose $x \sqsubset_{\mathbf{m}} y$. We check that $\pi(x) \sqsubset_\Phi \pi(y)$. There are two cases to consider: First suppose that $x \in |\mathbf{m}_u|$ for some u and $y \in |\mathbf{m}_v|$ for some v . Then by the definition of the moment construction operation, $u = v$. By the induction hypothesis, since π_u is a p -morphism, then $\pi(x) \sqsubset_\Phi \pi(y)$. Otherwise, suppose that $x \in C_\Phi(w)$. If $y \in C_\Phi(w)$ we immediately obtain $\pi(x) \sqsubset_\Phi \pi(y)$, since $C_\Phi(w)$ is precisely the \sqsubset_Φ -cluster of w (with the same accessibility relation). Otherwise, $y \in |\mathbf{m}_v|$ for some v , and we let y' be the root of \mathbf{m}_v , so that $\pi(y') = v$. Then, $\pi(x) \sqsubset_\Phi \pi(y')$. If $y = y'$ we are done. If $y \neq y'$, then since by the induction hypothesis π_v is a p -morphism with respect to \sqsubset_Φ , then $y' \sqsubset y$ implies $\pi(y') \sqsubset_\Phi \pi(y)$. By the (weak) transitivity of \sqsubset_Φ we have that either $\pi(x) \sqsubset_\Phi \pi(y)$ and we are done, or if $\Lambda = \mathbf{wK4C}$, possibly $\pi(x) = \pi(y)$. But then, $\pi(y') \sqsubset_\Phi \pi(x)$, contradicting that $\pi(x) \sqsubset_\Phi^1 \pi(y')$. \square

We next define the notions of *pre- Φ -morphism* and *quotient moment* that will be essential for the rest of the proof.

Definition 6.9 (Λ -bottom). Let \mathbf{m} be a moment and $\pi : |\mathbf{m}| \rightarrow W_c$. We say that $x \in |\mathbf{m}|$ is *at the Λ -bottom* for $\Lambda \in \{\mathbf{wK4}, \mathbf{K4}, \mathbf{GL}\}$ if

- $\Lambda \neq \mathbf{GL}$ and $\pi(x) \in C(\pi(r_{\mathbf{m}}))$, or
- $\Lambda = \mathbf{GL}$ and for all $y \sqsubset_{\mathbf{m}} x$, $\pi(y) = \pi(x)$.

We will refer to ‘ Λ -bottom’ simply as ‘bottom’ when this does not lead to confusion.

We say that $\pi : |\mathbf{m}| \rightarrow W_c$ is a *pre- Φ -morphism* if it fulfils conditions 1 and 4 of a Φ -morphism (Definition 6.4), and $x \sqsubset_{\mathbf{m}} y$ implies that either $\pi(x) \sqsubset_c \pi(y)$ or x, y are at the bottom.

Definition 6.10 (quotient moment). Let $\Lambda \in \{\mathbf{wK4C}, \mathbf{K4C}, \mathbf{GLC}\}$. Let \mathfrak{m} be a Λ -moment and $\pi : |\mathfrak{m}| \rightarrow W_c$ be a pre- Φ -morphism. We define $x \sim y$ if either $x = y$, or x, y are at the bottom and $\pi(x) = \pi(y)$. Given $y \in |\mathfrak{m}|$ we set $[y] = \{z : z \sim y\}$.

The *quotient moment* \mathfrak{m}/π of \mathfrak{m} and its respective map $[\pi] : |\mathfrak{m}/\pi| \rightarrow W_c$ are defined as follows, where $x, y \in |\mathfrak{m}|$:

- (a) $|\mathfrak{m}/\pi| = \{[x] : x \in |\mathfrak{m}|\}$;
- (b) $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$ iff one of the following conditions is satisfied:
 - x, y are at the bottom, $\pi(x) \sqsubset_c \pi(y)$, and $\Lambda \neq \mathbf{GLC}$;
 - x is at the bottom and y is not at the bottom;
 - x, y are not at the bottom and $x \sqsubset_{\mathfrak{m}} y$;
- (c) $\nu_{\mathfrak{m}/\pi}(p) = \{[x] : x \in \nu_{\mathfrak{m}}(p)\}$;
- (d) $r_{\mathfrak{m}/\pi} = [r_{\mathfrak{m}}]$;
- (e) $[\pi]([x]) = \pi(x)$.

The following proposition shows that the property of being at the bottom is downward-closed.

Proposition 6.11. *Let \mathfrak{m} be a moment and $\pi : |\mathfrak{m}| \rightarrow W_c$ be a pre- Φ -morphism. Let $x, y \in |\mathfrak{m}|$. If x is at the Λ -bottom and $y \sqsubset_{\mathfrak{m}} x$, then y is at the Λ -bottom.*

Proof. Suppose $\Lambda \neq \mathbf{GLC}$. If x is at the bottom, we have that $\pi(x) \in C(\pi(r_{\mathfrak{m}}))$. If moreover $y \sqsubset_{\mathfrak{m}} x$, then since π is a pre- Φ -morphism, we have that either $\pi(y) \sqsubset_c \pi(x)$ or y is at the bottom. In the latter case there is nothing to prove. Otherwise, $r_{\mathfrak{m}} \sqsubset_{\mathfrak{m}} y \sqsubset_{\mathfrak{m}} x$ and y is not at the bottom. Then $\pi(r_{\mathfrak{m}}) \sqsubset_c \pi(y) \sqsubset_c \pi(x)$ and hence $\pi(y) \in C(r_{\mathfrak{m}})$. This is a contradiction and so y is at the bottom.

Suppose $\Lambda = \mathbf{GLC}$. Then by the transitivity of $\sqsubset_{\mathfrak{m}}$ it follows that y is also at the bottom. \square

The quotient moment and its respective map hold some essential properties for the derivation of this section's main result. The idea is that by constructing a Λ -moment m with an associated pre- Φ -morphism π , we get that \mathfrak{m}/π is still a Λ -moment, but now $[\pi]$ is a proper Φ -morphism. The next few results make this intuition precise.

Lemma 6.12. *For $\Lambda \in \{\mathbf{wK4}, \mathbf{K4}, \mathbf{GL}\}$, if \mathfrak{m} is any Λ -moment and $\pi : |\mathfrak{m}| \rightarrow W_c$ is a pre- Φ -morphism, then \mathfrak{m}/π is a Λ -moment and $[\cdot] : |\mathfrak{m}| \rightarrow |\mathfrak{m}/\pi|$ is weakly monotonic and root-preserving.*

Proof. For the cases where $\Lambda = \mathbf{wK4}$ and $\Lambda = \mathbf{K4}$ it is easy to verify that $|\mathfrak{m}/\pi|$ is weakly transitive and transitive, respectively. For the case where $\Lambda = \mathbf{GL}$ suppose for contradiction that $[x] \sqsubset_{\mathfrak{m}/\pi} [x]$. Then it must be the case that x is not at the bottom and $x \sqsubset_{\mathfrak{m}} x$ in contradiction to \mathfrak{m} being a \mathbf{GL} -moment.

Next we check that $[\cdot]$ is weakly monotonic. Suppose that $x \sqsubset_{\mathfrak{m}} y$. By Proposition 6.11, if y is at the bottom, so is x . Thus in order to prove that $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$, there are three cases to consider. If x, y are at the bottom, either $\pi(x) = \pi(y)$, so that $[x] = [y]$, or else $\Lambda \neq \mathbf{GLC}$ and $\pi(x) \sqsubset_c \pi(y)$, hence

$[x] \sqsubset_{\mathfrak{m}/\pi} [y]$ by definition of $\sqsubset_{\mathfrak{m}/\pi}$. If x is at the bottom and y is not at the bottom, then by definition $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$. If neither is at the bottom, $x \sqsubset_{\mathfrak{m}} y$ yields $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$. In all cases we have that $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$. \square

Proposition 6.13. *If \mathfrak{m} is a moment and $\pi: |\mathfrak{m}| \rightarrow W_c$ is a pre- Φ -morphism, then the map $[\pi]$ is a well-defined Φ -morphism.*

Proof. To show that $[\pi]$ is well defined, suppose that for some $x, y \in |\mathfrak{m}|$ we have $x \neq y$ and $[x] = [y]$. By definition of \sim it follows that $\pi(x) = \pi(y)$, as required. It remains to show that $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$ implies $[\pi]([x]) \sqsubset_c [\pi]([y])$. Suppose $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$. If both x, y are at the bottom, then $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$ may only occur when $\pi(x) \sqsubset_c \pi(y)$. If neither x nor y are at the bottom then $[x] \sqsubset_{\mathfrak{m}/\pi} [y]$ may only occur when $x \sqsubset_{\mathfrak{m}} y$, which implies that $\pi(x) \sqsubset_c \pi(y)$ since π is a pre- Φ -morphism. The case where y is at the bottom but x is not is impossible by Lemma 6.11. So we are left with the case where x is at the bottom but y is not. But then $\pi(x) = \pi(r_{\mathfrak{m}}) \sqsubset_c \pi(y)$, where $\pi(r_{\mathfrak{m}}) \sqsubset_c \pi(y)$ is obtained from the definition of pre- Φ -morphism, using the fact that $r_{\mathfrak{m}} \sqsubset_{\mathfrak{m}} y$ and y is not at the bottom. We conclude that $[\pi]$ satisfies Condition 3 of a Φ -morphism.

For Condition 4, suppose that $[\pi]([x]) \sqsubset_{\Phi} v$. Since $[\pi]([x]) = \pi(x)$, there is $y \sqsupset_{\mathfrak{m}} x$ such that $\pi(y) = v$. Inspecting the definition of $\sqsubset_{\mathfrak{m}/\pi}$ yields that $[\pi]([x]) \sqsubset_{\mathfrak{m}/\pi} [\pi]([y])$, except possibly when $\Lambda = \mathbf{GLC}$ and x, y are at the bottom. However, in this case we would have $v = \pi(x)$, but \sqsubset_{Φ} is irreflexive, so this is impossible. We conclude that Condition 4 is also satisfied. \square

Thus from a pre- Φ -morphism we can produce a suitable Φ -morphism. It remains to provide tools for producing dynamic Φ -morphisms.

Proposition 6.14. *Fix $\Lambda \in \{\mathbf{wK4C}, \mathbf{K4C}, \mathbf{GLC}\}$ and let $\mathfrak{M}_c^{\Lambda} = \langle W, \sqsubset, g, \nu \rangle$. Let \mathfrak{m} and \mathfrak{n} be moments, $f: |\mathfrak{m}| \rightarrow |\mathfrak{n}|$ be root-preserving, weakly monotonic, and $\pi: |\mathfrak{m}| \rightarrow W$ and $\rho: |\mathfrak{n}| \rightarrow W$ be such that π is a Φ -morphism, ρ is a pre- Φ -morphism, and $g \circ \pi = \rho \circ f$. Let $[f]$ be given by $[f](x) = [f(x)]$. Then, $[f]: |\mathfrak{m}| \rightarrow |\mathfrak{n}/\rho|$ is weakly monotonic and root-preserving and satisfies $g \circ \pi = [\rho] \circ [f]$.*

Proof. Checking that $g \circ \pi = [\rho] \circ [f]$ is routine, and $[f]$ is weakly monotonic and root-preserving since it is a composition of maps with these properties in view of Lemma 6.12. \square

Using these properties, we can prove the existence of an appropriate story that maps to the canonical model. This is based on the following useful lemma:

Lemma 6.15. *Fix $\Lambda \in \{\mathbf{wK4C}, \mathbf{K4C}, \mathbf{GLC}\}$ and let $\mathfrak{M}_c^{\Lambda} = \langle W_c, \sqsubset_c, g_c, \nu_c \rangle$. Let \mathfrak{m} be a Λ -moment and suppose that there exists a Φ -morphism $\pi: |\mathfrak{m}| \rightarrow W_c$. Then, there exists a Λ -moment \mathfrak{n} , a weakly monotonic map $f: |\mathfrak{m}| \rightarrow |\mathfrak{n}|$, and a Φ -morphism $\rho: |\mathfrak{n}| \rightarrow W_c$ such that $g_c \circ \pi = \rho \circ f$.*

Proof. We proceed by induction on the height of \mathfrak{m} . Let C be the cluster of $r_{\mathfrak{m}}$ and let $\vec{\mathfrak{a}} = \langle \mathfrak{a}_n \rangle_{n < N}$ be the generated sub-models of the immediate strict successors of $r_{\mathfrak{m}}$; note that each \mathfrak{a}_n is itself a moment of smaller height. By the induction hypothesis, there exist moments $\langle \mathfrak{a}'_n \rangle_{n < N}$, root-preserving, weakly monotonic maps $f_n: |\mathfrak{a}_n| \rightarrow |\mathfrak{a}'_n|$, and Φ -morphism $\rho_n: |\mathfrak{a}'_n| \rightarrow W_c$ such that $g_c \circ \pi_n = \rho_n \circ f_n$. Moreover, for each $v \sqsupset_{\Phi} g_c(r_{\mathfrak{m}})$, by Lemma 6.8 there

are \mathbf{b}_v and a Φ -morphism $\rho_v: |\mathbf{b}_v| \rightarrow W_c$ mapping the root of \mathbf{b}_v to v . Let $D = g_c\pi(C) \cup C_\Phi(g_c(x))$, and let $\hat{\mathbf{n}} = \left(\begin{smallmatrix} \bar{\mathbf{a}} * \bar{\mathbf{b}} \\ D \end{smallmatrix}\right)_{g_c(w)}$. It is not difficult to verify that the maps ρ_v can be used to define a pre- Φ -morphism $\hat{\rho}$ from $\hat{\mathbf{n}}$ to W_c . Let

$$\hat{f}(w) = \begin{cases} g_c(w) & \text{if } w \in C, \\ f_n(w) & \text{if } w \in |\mathbf{a}_n|. \end{cases}$$

It is easy to see that $\hat{f}: |\mathbf{m}| \rightarrow |\hat{\mathbf{n}}|$ is weakly monotonic and satisfies $g_c \circ \pi = \hat{\rho} \circ \hat{f}$. Setting $\mathbf{n} = \hat{\mathbf{n}}/\hat{\rho}$, $f = [\hat{f}]$ and $\rho = [\hat{\rho}]$, Lemma 6.12 tells us that \mathbf{n} is a Λ -moment and Propositions 6.13 and 6.14 imply that f and ρ have the desired properties. \square

Proposition 6.16. *Fix $\Lambda \in \{\mathbf{wK4C}, \mathbf{K4C}, \mathbf{GLC}\}$. Given $I < \omega$ and $w \in W_c$, there is a story \mathfrak{S} of duration I and a dynamic Φ -morphism $\pi: |\mathfrak{S}| \rightarrow W_c$ with $w = \pi(r_{\mathfrak{S}})$.*

Proof. Proceed by induction on I . For $I = 0$, this is essentially Lemma 6.8. Otherwise, by the induction hypothesis, assume that a story $\hat{\mathfrak{S}}$ of depth I and dynamic p -morphism $\hat{\pi}$ exist. By Lemma 6.15, there is a moment \mathfrak{S}_{I+1} , map $f_I: |\mathfrak{S}_I| \rightarrow |\mathfrak{S}_{I+1}|$, and Φ -morphism $\pi_{I+1}: |\mathfrak{S}_{I+1}| \rightarrow W_c$ commuting with f_I . We define \mathfrak{S} by adding \mathfrak{S}_{I+1} to $\hat{\mathfrak{S}}$ in order to obtain the desired story. \square

It follows that any satisfiable formula is also satisfiable on a finite story, hence satisfiable on a finite model, yielding the main result of this section.

Theorem 6.17. *The logics $\mathbf{wK4C}$, $\mathbf{K4C}$ and \mathbf{GLC} are sound and complete for their respective class of finite dynamic Λ -frames.*

Proof. Soundness follows from Lemma 3.9 and the well known soundness results reviewed in Section 3. Let $\Lambda \in \{\mathbf{wK4C}, \mathbf{K4C}, \mathbf{GLC}\}$ and suppose $\Lambda \not\vdash \varphi$. Then in the canonical model $\mathfrak{M}_c^\Lambda = \langle W, \sqsubset, g, \nu \rangle$, there is $w \in W$ that refutes φ . By Proposition 6.16, there is a story \mathfrak{S} and a dynamic Φ -morphism $\pi: |\mathfrak{S}| \rightarrow W$ such that $w = \pi(r)$, where r is a root of \mathfrak{S} . It follows that $\mathfrak{S}, r \not\models \varphi$. Recall that \mathfrak{S} is a finite dynamic derivative frame. In the case of $\mathbf{K4C}$, \mathfrak{S} is also transitive, and in the case of \mathbf{GLC} , also transitive and irreflexive. Then $\mathbf{wK4C}$ is complete with respect to finite dynamic derivative frames, $\mathbf{K4C}$ with respect to finite, transitive, dynamic derivative frames, and \mathbf{GLC} is complete with respect to finite, irreflexive, transitive dynamic derivative frames. \square

7 Topological d -completeness

In this section we establish completeness results for classes of dynamic topological systems with continuous functions. We begin with the logic \mathbf{GLC} , simply because the topological d -completeness for \mathbf{GLC} is almost immediate, given that a \mathbf{GLC} model is already a dynamic derivative model based on a scattered space via the standard up-set topology.

Theorem 7.1. *\mathbf{GLC} is the d -logic of all dynamic topological systems based on a scattered space. Furthermore, it has the finite model property for this class.*

Proof. Soundness follows from Theorem 3.8 and Lemma 3.9. For completeness, suppose that $\not\vdash_{\mathbf{GLC}} \varphi$. Then by Theorem 6.17 there is a finite model $\mathfrak{M} = \langle W, \sqsubset, g, \nu \rangle$ based on a transitive, irreflexive dynamic derivative frame such that $\mathfrak{M} \not\models \varphi$. But by Lemmas 2.6 and 3.7, setting $\tau = \tau_{\sqsubset}$, we see that $d_{\tau} = d_{\sqsubset}$ and $\langle W, \tau, f \rangle$ is a scattered dynamic topological system, which is identical to \mathfrak{M} as a dynamic derivative system and thus also refutes φ . \square

In order to prove topological d -completeness for $\mathbf{wK4C}$, we first provide a definition and some generalisations of well-known results. We use similar constructions as in e.g. [8].

Definition 7.2. Let $\mathfrak{F} = \langle W, \sqsubset, g \rangle$ be a $\mathbf{wK4C}$ -frame and let W^i and W^r be the sets of irreflexive and reflexive points respectively. We define a new frame $\mathfrak{F}_{\oplus} = \langle W_{\oplus}, \sqsubset_{\oplus}, g_{\oplus} \rangle$, where

1. $W_{\oplus} = (W^i \times \{0\}) \cup (W^r \times \{0, 1\})$;
2. $(w, i) \sqsubset_{\oplus} (v, j)$ iff $w \sqsubset v$ and $(w, i) \neq (v, j)$;
3. $g_{\oplus}(w, i) = (g(w), 0)$.

The following is standard [8] and easily verified.

Proposition 7.3. *If $\mathfrak{F} = \langle W, \sqsubset, g \rangle$ is any dynamic derivative frame, then \mathfrak{F}_{\oplus} is an irreflexive dynamic derivative frame and $\pi: W_{\oplus} \rightarrow W$ given by $\pi(w, i) = w$ is a surjective, dynamic p -morphism.*

Proposition 7.3 allows us to obtain topological completeness from Theorem 6.17, as Lemma 2.6 tells us that irreflexive Kripke frames are essentially Cantor derivative spaces. We thus obtain the following:

Theorem 7.4. *$\mathbf{wK4C}$ is the d -logic of all dynamic topological systems. Furthermore, it has the finite model property for this class.*

Proof. Soundness follows from Lemma 3.9. For completeness, suppose $\mathbf{wK4C} \not\vdash \varphi$. Then by Theorem 6.17 there exists a finite $\mathbf{wK4C}$ -frame $\mathfrak{F} = \langle W, \sqsubset, g \rangle$ such that $\mathfrak{F} \not\models \varphi$. But then, by Proposition 7.3, $\mathfrak{F}_{\oplus} = \langle W_{\oplus}, \sqsubset_{\oplus}, g_{\oplus} \rangle$ is an irreflexive dynamic derivative system and $\pi: W_{\oplus} \rightarrow W$ is a surjective, dynamic p -morphism. But by Lemma 2.6 and by setting $\tau = \tau_{\sqsubset}$ we see that $d_{\sqsubset} = d_{\tau}$, so \mathfrak{F}_{\oplus} can be seen as a dynamic topological system with the same semantics, also falsifying φ . \square

Finally we turn our attention to $\mathbf{K4C}$. Unlike the other two logics, $\mathbf{K4C}$ (or even $\mathbf{K4}$) does not have the topological finite model property, despite having the Kripke finite model property. In the case of Aleksandroff spaces, the class T_D is easy to describe.

Lemma 7.5. *The Aleksandroff space of a $\mathbf{K4}$ -frame $\mathfrak{F} = \langle W, \sqsubset \rangle$ is T_D if and only if \sqsubset is antisymmetric, in the sense that $w \sqsubset v \sqsubset w$ implies $w = v$.*

If we moreover want the Kripke and the d -semantics to coincide on \mathfrak{F} , we need \sqsubset to be irreflexive. Thus we wish to ‘unwind’ \mathfrak{F} to get rid of all non-trivial clusters. The following construction achieves this.

Definition 7.6. Let $\mathfrak{F} = \langle W, \sqsubset, g \rangle$ be a dynamic **K4** frame. We define a new frame $\vec{\mathfrak{F}} = \langle \vec{W}, \vec{\sqsubset}, \vec{g} \rangle$, where

- \vec{W} is the set of all finite sequences (w_0, \dots, w_n) , where $w_i \sqsubset w_{i+1}$ for all $i < n$;
- $\mathbf{w} \vec{\sqsubset} \mathbf{v}$ iff \mathbf{w} is a strict initial segment of \mathbf{v} ;
- $\vec{g}(w_0, \dots, w_n)$ is the subsequence of $(g(w_0), \dots, g(w_n))$, obtained by deleting every entry that is equal to its immediate predecessor.

We then define a map $\pi: \vec{W} \rightarrow W$, where $\pi(w_0, \dots, w_n) = w_n$.

In the definition of \vec{g} , note that g is only weakly monotonic, so it may be that, for instance, $g(w_0) = g(w_1)$; in this case, we include only one copy of $g(w_0)$ to ensure that $\vec{g}(\mathbf{w}) \in \vec{W}$.

Proposition 7.7. *If $\mathfrak{F} = \langle W, \sqsubset, g \rangle$ is any dynamic **K4** frame, then $\vec{\mathfrak{F}}$ is an antisymmetric and irreflexive dynamic **K4** frame. Moreover, $\pi: \vec{W} \rightarrow W$ is a surjective, dynamic p -morphism.*

Proof. It is easy to see that $\vec{\mathfrak{F}}$ is antisymmetric and irreflexive. In order to show that $\vec{\mathfrak{F}}$ is a dynamic **K4** frame we need to show that it is transitive and \vec{g} is a weakly monotonic function. Suppose $\mathbf{w} \vec{\sqsubset} \mathbf{v} \vec{\sqsubset} \mathbf{u}$, then clearly since \mathbf{w} is an initial segment of \mathbf{v} , it is also an initial segment of \mathbf{u} and therefore $\mathbf{w} \vec{\sqsubset} \mathbf{u}$ and thus $\vec{\sqsubset}$ is transitive. The functionality of \vec{g} follows from the functionality of g , and weak monotonicity can be checked from the definitions using the weak monotonicity of g . Finally, we show that π is a surjective p -morphism. It is clearly surjective. Next, we note that if $\mathbf{w} \vec{\sqsubset} \mathbf{v}$ then transitivity plus $w_n = v_n$ yields $w_n \sqsubset v_i$ for all $i \in (n, m]$, and in particular $w_n \sqsubset v_m$, that is, $\pi(\mathbf{w}) \sqsubset \pi(\mathbf{v})$. Moreover, if $\pi(\mathbf{w}) \sqsubset u$, then $w_n \sqsubset u$ by definition of π , which means that $\mathbf{u} := (w_1, \dots, w_n, u) \in \vec{W}$ is such that $\mathbf{w} \vec{\sqsubset} \mathbf{u}$ and $\pi(\mathbf{u}) = u$. Furthermore, $\pi(\vec{g}(w_0, \dots, w_n)) = g(w_n) = g(\pi(w_0, \dots, w_n))$ and thus π is a p -morphism. \square

Similar constructions have already appeared in e.g. [15]. From this we obtain the following:

Theorem 7.8. **K4C** is the d -logic of all dynamic topological systems based on a T_D space, as well as the d -logic of all dynamic topological systems based on an Aleksandroff T_D space.

Proof. Soundness follows from soundness for **wK4C** and Lemma 7.5. For completeness, if **K4C** $\not\models \varphi$, by Theorem 6.17 there is a **K4C** model $\mathfrak{M} = \langle W, \sqsubset, g, V \rangle$ falsifying φ . Let $\mathfrak{F} = \langle W, \sqsubset, g \rangle$. But by Proposition 7.7, $\pi: \vec{W} \rightarrow W$ is a surjective p -morphism, which implies that φ is not valid on $\vec{\mathfrak{F}}$. By Proposition 7.7, $\vec{\sqsubset}$ is antisymmetric, hence by Lemma 7.5, the Aleksandroff topology of $\vec{\mathfrak{F}}$ is T_D ; moreover, $\vec{\sqsubset}$ is also irreflexive, so we see that φ is falsified on this T_D Aleksandroff space as well. \square

8 Invertible Systems

Recall that if $\langle X, \tau \rangle$ is a topological space and $f: X \rightarrow X$ is a function, then f is a *homeomorphism* if it is a bijection and both f and f^{-1} are continuous.

Unlike in the continuous case, for logics with homeomorphisms, every formula is equivalent to a formula where all occurrences of \circ are applied to atoms. More formally, we say that a formula is in \circ -normal form if it is of the form $\varphi(\circ^{k_0}p_0, \dots, \circ^{k_n}p_n)$, where $\varphi(p_0, \dots, p_n)$ does not contain \circ , the p_i 's are variables, and the k_i 's are natural numbers. It is readily observed that the axiom H allows us to 'push' all instances of \circ to the propositional level. Thus we obtain the following useful representation lemma:

Lemma 8.1. *For every logic Λ and every formula φ , there is a formula φ' in \circ -normal form such that $\Lambda\mathbf{H} \vdash \varphi \leftrightarrow \varphi'$.*

Proof. This is a standard proof by induction on the complexity of φ . In particular, note that the axioms $\Box\circ p \rightarrow \circ\Box p$ and $\circ\Box p \rightarrow \Box\circ p$ permit the interchangeability between \circ and \Box , thus allow us to push \circ towards the atomic part of the formula. \square

As we shall see shortly, $\Lambda\mathbf{H}$ inherits completeness and the finite model property almost immediately from Λ . If $\mathfrak{X} = \langle X, \rho_X \rangle$ and $\mathfrak{Y} = \langle Y, \rho_Y \rangle$ are derivative spaces, then the *topological sum* is defined as $\mathfrak{X} \oplus \mathfrak{Y} = \langle X \oplus Y, \rho_X \oplus \rho_Y \rangle$, where $X \oplus Y$ is the disjoint union of the two sets, and $\rho_X \oplus \rho_Y$ is given by $(\rho_X \oplus \rho_Y)A = \rho_X(A \cap X) \cup \rho_Y(A \cap Y)$. We say that a class Ω of derivative spaces is *closed under sums* if whenever $\mathfrak{A}, \mathfrak{B} \in \Omega$, it follows that $\mathfrak{A} \oplus \mathfrak{B} \in \Omega$.

Given a derivative space $\mathfrak{A} = \langle A, \rho \rangle$, we may write \mathfrak{A}^n instead of $\mathfrak{A} \oplus \mathfrak{A} \oplus \dots \oplus \mathfrak{A}$ (n times), and if \mathfrak{A} has domain A , we may identify the domain of \mathfrak{A}^n with $A^n = A \times \{0, \dots, n-1\}$. It should be clear that if $\mathfrak{A} \in \Omega$ and Ω is closed under sums, then $\mathfrak{A}^n \in \Omega$. Let $(m)_n$ denote the remainder of m modulo n . We then define a dynamical structure $\mathfrak{A}^{(n)} = (\mathfrak{A}^n, f)$, where $f: A^n \rightarrow A^n$ is given by $f(w, i) = (w, (i+1)_n)$.

Lemma 8.2. *Let Ω be a class of derivative spaces closed under sums. Then, if $\mathfrak{A} \in \Omega$, it follows that $\mathfrak{A}^{(n)}$ is an invertible dynamic derivative system based on an element of Ω .*

Proof. Suppose $\mathfrak{A} = \langle X, d \rangle \in \Omega$ is a derivative space. Let $\mathfrak{A}^{(n)} = (X^n, d', f)$. Clearly f is a bijection as it defines disjoint cycles. If $Y \subseteq X^n$, let $Y_i = \{x \in X : (x, i) \in Y\}$. Then, $d'f^{-1}Y = \bigcup_{i < n} (dY_i, (i-1)_n) = f^{-1} \bigcup_{i < n} (dY_i, i) = f^{-1}d'Y$. So, $\mathfrak{A}^{(n)}$ is a dynamic derivative frame. \square

For our proof of completeness, we need the notion of *extended valuation*.

Definition 8.3. Let $\mathfrak{A} = \langle A, \rho \rangle$ be a derivative space. Let PV° be the set of all expressions $\circ^i p$, where $i \in \mathbb{N}$ and p is a propositional variable. An *extended valuation* on \mathfrak{A} is a relation $\nu \subseteq \text{PV}^\circ \times A$.

If ν is an extended valuation on \mathfrak{A} , we define a valuation on $\mathfrak{A}^{(n)}$ so that for any variable p and $(w, i) \in A^n$, $(w, i) \in \nu^{(n)}(p)$ if and only if $w \in \nu(\circ^i p)$.

Lemma 8.4. *Let $\mathfrak{A} = \langle A, \rho \rangle$ be any derivative space and ν be any extended valuation on \mathfrak{A} . Let $w \in A$, $i < n$, and φ be any formula in \circ -normal form which has \circ -depth less than $n - i$. Then, $\langle \mathfrak{A}^{(n)}, \nu^{(n)} \rangle, (w, i) \models \varphi$ if and only if $\langle \mathfrak{A}, \nu \rangle, w \models \varphi$.*

Proof. We start with the right to left direction and we only show the case where $\varphi = \circ\psi$. Since φ is in \circ -normal form then it is of the form $\circ^j p$. Since $j < n - i$ and by definition $(w, j) \in \nu^{(n)} p$, then $\langle \mathfrak{A}^{(n)}, \nu^{(n)} \rangle, (w, i) \models \circ^j p$. The other direction follows similarly. The rest of the modalities and connectives follow by a standard induction on the build of φ . \square

It then readily follows that φ is consistent if and only if φ' is consistent, which implies that φ' is satisfiable on Ω . This is equivalent to φ being satisfiable on the class of invertible systems based on Ω . From this, we obtain the following general result.

Theorem 8.5. *Let Λ be complete for a class Ω of derivative spaces. Then, if Ω is closed under sums, it follows that $\Lambda\mathbf{H}$ is complete for the class of invertible systems based on Ω .*

Proof. Assume that Λ and Ω are as in the statement of the theorem. Let φ be any formula, and using Lemma 8.1, let φ' be a $\Lambda\mathbf{H}$ -equivalent formula in \circ -normal form. Let n be the \circ -depth of φ' . If φ is consistent, then φ' is consistent over Λ . By the assumption, φ' is satisfied on some $\mathfrak{A} = \langle A, \rho \rangle \in \Omega$. Suppose it is satisfied on a point $w_* \in A$. By Lemma 8.2, $\mathfrak{A}^{(n)}$ is an invertible system based on an element of Ω , which by Lemma 8.4 satisfies φ' on $(w_*, 0)$. But $\Lambda\mathbf{H}$ is sound for the class of invertible systems based on Ω , hence φ is also satisfied on $(w_*, 0)$. \square

Corollary 8.6.

1. **wK4H** is sound and complete for
 - (a) The class of all finite invertible dynamic **wK4** frames.
 - (b) The class of all finite invertible dynamic topological systems with Cantor derivative.
2. **K4H** is sound and complete for
 - (a) The class of all finite invertible dynamic **K4** frames.
 - (b) The class of all invertible, T_D dynamic topological systems with Cantor derivative.
3. **GLH** is sound and complete for
 - (a) The class of all finite invertible dynamic **GL** frames.
 - (b) The class of all finite invertible, scattered dynamic topological systems with Cantor derivative.

9 Conclusion

We have recast dynamic topological logic in the more general setting of derivative spaces and established the seminal results for the $\{\diamond, \circ\}$ -fragment for the variants of the standard logics with the Cantor derivative in place of the topological closure. Semantics on the Cantor derivative give rise to a richer family of modal logics than their counterparts based on closure. This is evident in the

distinction between e.g. the logics **wKC** and **K4C**, both of which collapse to **S4C** when replaced by their closure-based counterparts. This line of research goes hand in hand with recent trends that consider the Cantor derivative as the basis of topological semantics [11, 18].

There are many natural problems that remain open. The logic **S4C** is complete for the Euclidean plane. In the context of d -semantics, the logic of the Euclidean plane is a strict extension of **K4**, given that punctured neighbourhoods are *connected* in the sense that they cannot be split into two disjoint, non-empty open sets. Thus one should not expect **K4C** to be complete for the plane. This raises the question: what is the dynamic d -logic of Euclidean spaces in general, and of the plane in particular?

The d -semantics also poses new lines of inquiry with respect to the class of functions considered. We have discussed continuous functions and homeomorphisms. Artemov et al. [1] also considered arbitrary functions, and we expect that the techniques used by them could be modified without much issue for d -semantics. On the other hand, in the setting of closure-based logics, the logic of spaces with continuous, open maps that are not necessarily bijective coincides with the logic of spaces with a homeomorphism. This is no longer true in the d -semantics setting, as the validity of the H axiom requires injectivity. Along these lines, the d -logic of *immersions* (i.e. continuous, injective functions) would validate the original continuity axiom $\circ\Box p \rightarrow \Box\circ p$. Thus there are several classes of dynamical systems whose closure-based logics coincide, but are split by d -semantics.

Finally, there is the issue of extending our language to the trimodal language with the ‘henceforth’ operator. It is possible that the d -logic of all dynamic topological systems may be axiomatised using the tangled derivative, much as the tangled closure was used to provide an axiomatisation of the closure-based **DTL**. However, given that the tangled derivative is made trivial on scattered spaces, we conjecture that the trimodal d -logic based on this class has a natural, finite axiomatisation. The work presented here is an important first step towards proving this.

References

- [1] S. Artemov, J. Davoren, and A. Nerode. Modal logics and topological semantics for hybrid systems. *Technical Report MSI 97-05*, 1997.
- [2] Patrick Blackburn, Maarten de Rijke, and Yde Venema. *Modal Logic*, volume 53 of *Cambridge Tracts in Theoretical Computer Scie.* Cambridge University Press, Cambridge, 2001.
- [3] George Boolos. The logic of provability. *The American Mathematical Monthly*, 91(8):470–480, 1984. [arXiv:https://doi.org/10.1080/00029890.1984.11971467](https://doi.org/10.1080/00029890.1984.11971467), [doi:10.1080/00029890.1984.11971467](https://doi.org/10.1080/00029890.1984.11971467).
- [4] Steven L Brunton and J Nathan Kutz. *Data-driven science and engineering: Machine learning, dynamical systems, and control*. Cambridge University Press, 2019.

- [5] Moody T Chu. Linear algebra algorithms as dynamical systems. *Acta Numerica*, 17:1–86, 2008.
- [6] L. Esakia. Weak transitivity–restitution. In Nauka, editor, *Study in Logic*, volume 8, pages 244–254, 2001. In Russian.
- [7] Leo Esakia. Diagonal constructions, Löb’s formula and Cantor’s scattered spaces. *Studies in logic and semantics*, 132(3):128–143, 1981.
- [8] Leo Esakia. Intuitionistic logic and modality via topology. *Ann. Pure Appl. Log.*, 127(1-3):155–170, 2004. doi:10.1016/j.apal.2003.11.013.
- [9] D. Fernández-Duque. Non-finite axiomatizability of dynamic topological logic. *ACM Transactions on Computational Logic*, 15(1):4:1–4:18, 2014. URL: <http://doi.acm.org/10.1145/2489334>, doi:10.1145/2489334.
- [10] David Fernández-Duque. Tangled modal logic for spatial reasoning. In Toby Walsh, editor, *IJCAI 2011, Proceedings of the 22nd International Joint Conference on Artificial Intelligence, Barcelona, Catalonia, Spain, July 16-22, 2011*, pages 857–862. IJCAI/AAAI, 2011. doi:10.5591/978-1-57735-516-8/IJCAI11-149.
- [11] David Fernández-Duque and Petar Iliev. Succinctness in subsystems of the spatial μ -calculus. *Journal of Applied Logics*, 5(4):827–874, 2018.
- [12] David Fernández-Duque. A sound and complete axiomatization for dynamic topological logic. *The Journal of Symbolic Logic*, 77(3):947–969, 2012. doi:10.2178/js1/1344862169.
- [13] R. Goldblatt and I. Hodkinson. Spatial logic of tangled closure operators and modal mu-calculus. *Annals of Pure and Applied Logic*, 168(5):1032 – 1090, 2017.
- [14] Guillaume Hanrot, Xavier Pujol, and Damien Stehlé. Analyzing blockwise lattice algorithms using dynamical systems. In *Annual Cryptology Conference*, pages 447–464. Springer, 2011.
- [15] B. Konev, R. Kontchakov, F. Wolter, and M. Zakharyashev. Dynamic topological logics over spaces with continuous functions. In G. Governatori, I. Hodkinson, and Y. Venema, editors, *Advances in Modal Logic*, volume 6, pages 299–318, London, 2006. College Publications.
- [16] Boris Konev, Roman Kontchakov, Frank Wolter, and Michael Zakharyashev. On dynamic topological and metric logics. *Studia Logica*, 84(1):129–160, 2006. doi:10.1007/s11225-006-9005-x.
- [17] P. Kremer and G. Mints. Dynamic topological logic. *Annals of Pure and Applied Logic*, 131:133–158, 2005.
- [18] Andrey Kudinov. Topological modal logics with difference modality. In Guido Governatori, Ian M. Hodkinson, and Yde Venema, editors, *Advances in Modal Logic 6*, pages 319–332. College Publications, 2006. URL: <http://www.aiml.net/volumes/volume6/Kudinov.ps>.

- [19] Hai Lin and Panos J. Antsaklis. Hybrid dynamical systems: An introduction to control and verification. *Found. Trends Syst. Control*, 1(1):1–172, March 2014. doi:10.1561/2600000001.
- [20] J.C.C. McKinsey and A. Tarski. The algebra of topology. *Annals of Mathematics*, 2:141–191, 1944.
- [21] Henning S. Mortveit and Christian M. Reidys. *An Introduction to Sequential Dynamical Systems*. Springer-Verlag, Berlin, Heidelberg, 2007.
- [22] A. Pnueli. The temporal logic of programs. In *Proceedings 18th IEEE Symposium on the Foundations of CS*, pages 46–57, 1977.
- [23] H. Simmons. Topological aspects of suitable theories. *Proceedings of the Edinburgh Mathematical Society*, 19(4):383–391, 1975. doi:10.1017/S001309150001049X.
- [24] Craig Smorynski. *Self-reference and modal logic*. Springer, 1985.
- [25] E Weinan. A proposal on machine learning via dynamical systems. *Communications in Mathematics and Statistics*, 5(1):1–11, 2017.