

Modular curves and their pseudo-analytic cover

B.Zilber*and C.Daw†

January 12, 2022

We find a natural $L_{\omega_1, \omega}$ -axiomatisation Σ of a structure on the upper half-plane \mathbb{H} as the covering space of modular curves. The main theorem states that Σ has a unique model in every uncountable cardinal.

The proof relies heavily on the theory of complex multiplication and the work on Langland’s conjecture on the conjugation of Shimura varieties. We also use the earlier work on a related problem by C.Daw and A.Harris. The essential difference between the setting of this work and that of the current paper is that the former was in the language which named the CM-points of the modular curves while our results here are over \mathbb{Q} .

1 Introduction

1.1 Motivation and origins. The original aim of the paper is the extension of the project *category of non-elementary theories of analytic covers* from “abelian” cases, such as $\exp : \mathbb{C} \rightarrow \mathbb{G}_m$ or $\exp_\Lambda : \mathbb{C} \rightarrow E_\Lambda$ (elliptic curve), to hyperbolic curves and possibly wider, see e.g. the survey part in [1] for some history and references.

The first obstruction for this project is that an adequate formalism (that is the language) in which such an analytic cover can be properly presented is not easy to determine. Some attempts in this direction were by M.Gavrilovich [2], as well as later attempts by A.Harris [3]. The one which we found satisfactory and applied here is based on the formalism close to the one applied in the recent [4].

*Supported by EPSRC Program grant “Symmetries and Correspondences”

†Supported by EPSRC New Investigator award EP/S029613/1

More specifically, we formalise e.g. the case $\exp : \mathbb{C} \rightarrow \mathbb{G}_m$ as a structure with two sorts \mathbb{U} and \mathbb{F} , where \mathbb{U} is the complex numbers with the \mathbb{Q} -module structure and the distinguished subgroup $2\pi i\mathbb{Z}$, and \mathbb{F} is the complex numbers as the field. Then $\exp : \mathbb{U} \rightarrow \mathbb{G}_m(\mathbb{F})$ is the map between the two sorts. But in fact along with \exp we automatically get in abelian cases the family

$$\exp_n : \mathbb{U} \rightarrow \mathbb{G}_m(\mathbb{F}), \quad \exp_n : x \mapsto \exp\left(\frac{x}{n}\right)$$

which of course agrees with the system of finite covers

$$\text{pr}_{k,n} : \mathbb{G}_m(\mathbb{F}) \rightarrow \mathbb{G}_m(\mathbb{F}); \quad y \mapsto y^{\frac{k}{n}}, \quad \text{when } n|k.$$

This we describe axiomatically by simple $L_{\omega_1, \omega}$ sentences. In terms of model-theoretic classification such a structure is a *fusion between a locally modular structure \mathbb{U} and an algebraically closed field \mathbb{F}* . The case $\exp_\Lambda : \mathbb{C} \rightarrow E_\Lambda$ similarly represents a fusion between a locally modular structure \mathbb{U}_Λ , and an algebraically closed field \mathbb{F} , where \mathbb{U}_Λ is a \mathbb{Q} -module with distinguished \mathbb{Z} -module Λ with an alternating form on it (accounting for the Weil pairing).

1.2 It is not clear a priori why such $L_{\omega_1, \omega}$ -theories should be categorical in uncountable cardinals and the fact that they are must be of some significance.

The geometric value of the project is perhaps in the fact that the formulation of the categorical theory of the universal cover of a variety \mathbb{X} (essentially the description of \mathbb{U}) is essentially a formulation of a complete formal **invariant** of \mathbb{X} . Indeed, the most interesting outcome of the earlier works was establishing an equivalence between categoricity of the cover of E_Λ and the conjunction of the two arithmetic facts:

- (i) the complete classification of the Galois action on the torsion of E_Λ (the open image theorem by J.-P.Serre) and
- (ii) the Kummer theory for E_Λ (M.Bashmakov and K.Ribet)

For abelian varieties the success of the program depends on the extension of analogues of (i) and (ii) to abelian varieties, and (ii) is known due to K.Ribet and M.Larsen. However, an analogue of Serre's theorem for abelian varieties is an open problem and therefore the best categoricity result here is under the assumption that the language names points of *the kernel* of the exponential map. This automatically removes the problem of determining the Galois action on the torsion points at the cost of weakening the formal invariant of \mathbb{X}/k , the theory of the cover of \mathbb{X}/k , to the formal invariant of

\mathbb{X}/K where K is the field obtained from k by adjoining all torsion points of \mathbb{X} .

The case of hyperbolic curves \mathbb{X} was first considered by A.Harris [3] and C.Daw and A.Harris [13] in the context of modular curves $\mathbb{Y}(\Gamma)$ in the formalism (the choice of the language) which has names for each element of $\mathrm{GL}_2^+(\mathbb{Q})$ acting on the upper half plane \mathbb{H} by Möbius transformations. The proof of categoricity in this setting required essentially Serre's open image theorem for products of non-CM elliptic curves. Interestingly, in the analysis of $(\mathbb{H}, j_N, \mathbb{Y}(N))$ Serre's theorem plays rather the role of (ii), while (i) is not needed since naming elements of $\mathrm{GL}_2(\mathbb{Q})$ fixes the special points (CM-points). This has a cost: one works out the formal invariant of $\mathbb{Y}(\Gamma)$ over the extension of the natural field of definition by special points.

1.3 Our setting. Our current interest is the case of the universal cover of hyperbolic curves, e.g. the complex curve $\mathbb{X} = \mathbf{P}^1 \setminus \{0, 1, \infty\}$. However, before approaching this case we set ourselves a simpler task of the *cover of the modular curve $\mathbb{Y}(N)$ universal in the class of modular curves*, which means that our structure incorporates the analytic covering maps

$$j_n : \mathbb{H} \rightarrow \mathbb{Y}(n), \text{ for all } n \text{ such that } N|n$$

agreeing with the algebraic finite covers

$$\mathrm{pr}_{n,m} : \mathbb{Y}(n) \rightarrow \mathbb{Y}(m), \text{ for all } n, m \text{ such that } N|m, m|n.$$

In fact it is enough to classify the case $N = 1$, $\mathbb{Y}(1) = \mathbb{Y} = \mathbf{A}^1$ the affine line (the fact that for some n the covers $j_n : \mathbb{H} \rightarrow \mathbb{Y}(n)$ are ramified does not matter in our setting).

Note that $\mathbb{Y}(2) = \mathbf{P}^1 \setminus \{0, 1, \infty\}$ when we consider $\mathbb{Y}(2)$ as an algebraic curve (without the level structure).

The important difference with the case of the proper universal cover is that, instead of the profinite completion $\hat{\Gamma}(N)$ of the respective fundamental group, in the modular setting one has the group $\tilde{\Gamma}(N)$, the completion in the topology based on congruence subgroups, which for $N = 1$ gives us

$$\tilde{\Gamma}(1) = \tilde{\Gamma} = \mathrm{SL}(2, \hat{\mathbb{Z}}),$$

where

$$\hat{\mathbb{Z}} = \lim \mathbb{Z}/n\mathbb{Z},$$

the projective limit of residue rings.

1.4 The key problem, similar to earlier cases, is in classifying the saturated version of \mathbb{H} in the structure $(\mathbb{H}, j_n, \mathbb{Y}(n))_{n \in \mathbb{N}}$ which is essentially reducible to understanding the structure on the projective limit

$$\tilde{\mathbb{H}} := \varprojlim \Gamma(n) \backslash \mathbb{H} \cong \varprojlim \mathbb{Y}(n).$$

This includes giving a detailed enough description of the action of the automorphism group $\text{Aut } \mathbb{C}$ on $\tilde{\mathbb{H}}$ and, in particular, its action on $\tilde{\mathbb{H}}(\text{CM})$, the special points of the structure (equivalently, the action of $\text{Gal}(\mathbb{Q}(\text{CM}) : \mathbb{Q})$ on $\tilde{\mathbb{H}}(\text{CM})$), the analogue of (i) of 1.2.

It turns out that this problem is closely connected to the problem/conjecture on conjugation of Shimura varieties posed by R.Langlands in [5] and solved in various degrees of generality by G.Shimura, P.Deligne, M.Borovoi, J.Milne and others; see [8] in a most general form, and [9] for the case of Shimura curves. In the interpretation of Borovoi [7] Langlands' conjecture amounts to characterising $\tilde{\mathbb{H}}$ as a structure with \mathbb{H} , its substructure, and describing the way in which \mathbb{H} is conjugated inside $\tilde{\mathbb{H}}$ by $\text{Aut } \mathbb{C}$ (in the case of general Shimura varieties). We believe that the categorical axiomatisation of $(\mathbb{H}, j_n, \mathbb{Y}(n))_{n \in \mathbb{N}}$ established in the paper implies Langlands' conjecture (for modular curves) but leave the proof of this statement for future work.

The theory resulting from Langlands' Conjecture proves uniqueness of so called *canonical models of Shimura varieties*. This involves an advanced theory of complex multiplication and Artin's reciprocity map. The results allow one to identify the action of $\text{Aut } \mathbb{C}$ on a single CM-point (that is the Galois orbit of the point), see [8], 12.8. We need a stronger result: our paper [10] goes further and identifies the action of $\text{Gal}(\mathbb{Q}(\text{CM}) : \mathbb{Q})$ on $\tilde{\mathbb{H}}(\text{CM})$, at the same time describing all the relations between CM-points. One of the interesting model-theoretic forms of the main result of [10] is that *the field $\mathbb{Q}(\text{CM})$ as a structure is $L_{\omega_1, \omega}$ -bi-interpretable with a certain structure formulated purely in terms of the ring \mathbb{A}_f of finite adeles over \mathbb{Q} .*

Theorem 4.1 in section 4 below formulates the main result of [10] in the form suitable for the current purposes. In particular, for a certain locally modular structure of trivial type on the set $\tilde{\mathbb{H}}$, which we term $\tilde{\mathbb{H}}^{\text{Pure}}$, the theorem states:

$$\text{Aut } \tilde{\mathbb{H}}^{\text{Pure}}(\text{CM}) \cong \text{Gal}(\mathbb{Q}(\text{CM}) : \mathbb{Q}).$$

The structure can be formulated in terms of the action of a large subgroup \tilde{G} of $\text{GL}_2(\mathbb{A}_f)$ on $\tilde{\mathbb{H}}$ and has a "standard" version \mathbb{H}^{Pure} which is based on

the actual upper half-plane \mathbb{H} and is given in terms of the action of $\mathrm{GL}_2^+(\mathbb{Q})$ and complex multiplication.

1.5 With Theorem 4.1 in hand we can apply the model-theoretic techniques on categoricity developed in earlier works (see e.g. the survey in [1]) and in particular the proof in [13] which we follow quite closely (and so use the Serre open image theorem).

The axioms Σ of the resulting $L_{\omega_1, \omega}$ -theory consist, as in abelian examples, of three parts:

- A The axioms describing $\mathbb{U}^{\mathrm{Pure}}$ (the abstract version of $\mathbb{H}^{\mathrm{Pure}}$), a locally modular structure of trivial type with the action of the group G isomorphic to $\mathrm{GL}_2^+(\mathbb{Q})$, and its subgroups corresponding to $\Gamma(n)$.
- B The axioms describing an algebraically closed field F of characteristic 0 and curves $\mathbb{Y}(n) \subset \mathbf{P}^3(F)$.
- C The axioms describing $j_n : \mathbb{U}^{\mathrm{Pure}} \rightarrow \mathbb{Y}(n)$ obtained via the translation of relations on $\mathbb{U}^{\mathrm{Pure}}$ to special relations on the $\mathbb{Y}(n)$.

Main Theorem. *The above system of axioms is satisfied by the standard complex structure \mathbb{H} and has, up to isomorphism, a unique model in any uncountable cardinal.*

1.6 The first author wishes to express his gratitude to J.Derakhshan for his interest in the work and his help with some of the mathematical issues in the paper.

The second author would like to thank the EPSRC for its support via a New Investigator Award (EP/S029613/1). He would also like to thank the University of Oxford for having him as a Visiting Research Fellow.

2 Groups acting on \mathbb{H} .

2.1 Groups G and Γ and their generators.

Generators of $\Gamma := \mathrm{SL}_2(\mathbb{Z})$ are represented by matrices

$$\mathbf{s} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{t} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}:$$

$$\mathbf{s}^2 = -\mathbf{I}, \quad (\mathbf{st})^3 = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}^3 = -\mathbf{I}.$$

Generators of $G := \mathrm{GL}_2^+(\mathbb{Q})$:

$$\mathbf{s}, \mathbf{t} \text{ and } \mathbf{d}_q := \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix}, q \in \mathbb{Q}_+$$

satisfying:

$$\begin{aligned} \mathbf{d}_q \mathbf{d}_r &= \mathbf{d}_{qr}, \\ \mathbf{s} \mathbf{d}_q &= q \mathbf{d}_q^{-1} \mathbf{s}, \\ \mathbf{d}_n \mathbf{t} &= \mathbf{t}^n \mathbf{d}_n, \text{ for } n \in \mathbb{N}. \end{aligned}$$

We will use

$$\begin{aligned} \mathbf{d}'_q &:= \mathbf{s} \mathbf{d}_q \mathbf{s}^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & q \end{pmatrix}, q \in \mathbb{Q}_+ \\ \mathbf{t}_- &= \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \end{aligned}$$

which satisfies

$$\mathbf{st} \mathbf{s}^{-1} = \mathbf{t}_-; \quad \mathbf{d}_n \mathbf{t}_-^n = \mathbf{t}_- \mathbf{d}_n, \text{ for } n \in \mathbb{N},$$

and subgroups

$$\Delta(\mathbb{Q}_+) = \{\mathbf{d}_q : q \in \mathbb{Q}_+\} \text{ and } \Delta'(\mathbb{Q}_+) = \{\mathbf{d}'_q : q \in \mathbb{Q}_+\}. \quad (1)$$

Similar notation also make sense for the multiplicative group of a commutative ring R ,

$$\Delta(R^\times) = \{\mathbf{d}_q : q \in R^\times\} \text{ and } \Delta'(R^\times) = \{\mathbf{d}'_q : q \in R^\times\}.$$

2.2 Remark. Note that all automorphisms of $\mathrm{GL}_2^+(\mathbb{Q})$ are inner and the only non-identity automorphism which fixes Δ' element-wise and preserves the subgroup Γ is the involution

$$g \mapsto \check{g} := \mathbf{d}_{-1} \cdot g \cdot \mathbf{d}_{-1}.$$

2.3 Special points on \mathbb{H} . Let

$$E = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G : (d-a)^2 < -4bc \right\} = \{e \in G : \mathrm{tr}^2 e < 4 \det e\},$$

the set of *elliptic* transformations.

These are exactly the elements for which there is a unique fixed point $\tau_e \in \mathbb{H}$ which can be found by solving the equation

$$cx^2 + (d - a)x - b = 0, \quad \tau_e := x, \quad \Im x > 0. \quad (2)$$

Note that elements of the centre

$$Z = \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \in G \right\}$$

act as identity on \mathbb{H} .

2.4 Remark. (i) E is invariant under automorphisms of G since $g \cdot e \cdot g^{-1}$ fixes $g\tau_e$ and thus belongs to E .

(ii) The subgroup $\text{St}_\tau \subset G$ fixing point $\tau = \tau_e$ is definable by the condition that $\text{St}_\tau = C(e)$, the centraliser of e in the group. This is a general fact for Mobius transformations over any field of characteristic 0.

3 Projective limit

3.1 We consider the structure $\tilde{\mathbb{H}}$, the projective limit of structures

$$\tilde{\mathbb{H}} = \varprojlim \Gamma(N) \backslash \mathbb{H}, \quad (3)$$

along the projective system of

$$\Gamma(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma, \quad a \equiv d \equiv 1 \pmod{N}, \quad c \equiv b \equiv 0 \pmod{N} \right\},$$

normal subgroups of $\Gamma = \text{SL}(2, \mathbb{Z})$.

The structures $\Gamma(N) \backslash \mathbb{H}$ are identified with the classical complex modular curves $\mathbb{Y}(N)$ (without the full level N structure). These are non-singular algebraic curves and therefore can be realised as quasi-projective algebraic curves in $\mathbf{P}^3(\mathbb{C})$. Moreover, the realisation can be obtained over \mathbb{Q} , see e.g. section 4 of [11] or the paper [12], section 2. Another argument for this fact is given in [10], 3.2(8).

The classical modular functions

$$j_N : \mathbb{H} \rightarrow \mathbb{Y}(N); \quad \tau \mapsto \Gamma(N) \cdot \tau$$

are holomorphic and the curves $\mathbb{Y}(N)$ are finite coverings of $\mathbb{Y}(N/d)$, for $d|N$, via the projection maps

$$\mathrm{pr}_{N,N/d} : j_N(\tau) \mapsto j_{N/d}(\tau), \quad \mathbb{Y}(N) \rightarrow \mathbb{Y}(N/d). \quad (4)$$

3.2 The definition (3) implies the existence of maps

$$j_N : \tilde{\mathbb{H}} \rightarrow \Gamma(N) \backslash \mathbb{H} = \mathbb{Y}(N)$$

which we are going to use below.

By definition, any $\tau \in \tilde{\mathbb{H}}$ is uniquely determined by the sequence

$$j_n(\tau) \in \mathbb{Y}(n), \quad n \in \mathbb{N}$$

and the sequence has the property

$$\mathrm{pr}_{n,n/d}(j_n(\tau)) = j_{n/d}(\tau), \quad \text{for each } d|n.$$

3.3 Lemma. *Any $\tau \in \mathbb{H}$ is uniquely determined by the sequence*

$$j_n(\tau) \in \mathbb{Y}(n) : \quad n \in \mathbb{N}.$$

This gives the canonical embedding

$$\mathbb{H} \hookrightarrow \tilde{\mathbb{H}}.$$

Proof. The first statement follows from the fact that $\bigcap_N \Gamma(N) = \{1\}$. The second statement is the consequence of the fact that the sequence satisfies

$$\mathrm{pr}_{n,n/d}(j_n(\tau)) = j_{n/d}(\tau), \quad \text{for each } d|n.$$

□

3.4 Remark. The system of covers (4) is not étale. However, by removing finitely many points on $\mathbb{Y}(1)$ and all the points on $\mathbb{Y}(n)$ over these one gets smooth curves $\mathbb{Y}^-(n)$ and a projective system of covers

$$\mathrm{pr}_{n,n/d} : \mathbb{Y}^-(n) \rightarrow \mathbb{Y}^-(n/d)$$

which is étale.

Since the construction of $\tilde{\Gamma}$ depends on generic fibres we have the same $\tilde{\Gamma}$ for the construction corresponding to the system of étale covers.

3.5 The projective limit as the structure $\tilde{\mathbb{H}}^{\text{Full}}$.

The analysis and study of the projective limit of modular curves $\mathbb{Y}(n)$ with the n -level structure defined over $\mathbb{Q}(\zeta_n)$ is readily reducible to the study of *canonical models of Shimura varieties*, see [8], and more specifically canonical models of modular curves, [9]. The case of curves $\mathbb{Y}(n)$ over \mathbb{Q} as above in the setting appropriate for our purposes is studied in [10]. The structure $\tilde{\mathbb{H}}^{\text{Full}}$ is described therein as the quotient $\Delta(\hat{\mathbb{Z}}^\times) \backslash \mathbf{S}^{\text{Full}}$ of a more fundamental structure \mathbf{S}^{Full} associated with the Shimura datum $(\mathbb{H} \cup -\mathbb{H}, \text{GL}_2)$.

The main conclusions, see [10], 3.22, are as follows:

(a) There is a group \tilde{G} acting on $\tilde{\mathbb{H}}$.

$$\tilde{G} \cong \Delta(\mathbb{Q}_+) \cdot \text{SL}_2(\mathbb{A}_f) \subset \text{GL}_2(\mathbb{A}_f),$$

where \mathbb{A}_f is the ring of finite adeles.

One can speak about an action of $\mathbf{g} \in \Delta(\mathbb{Q}_+) \cdot \text{SL}_2(\mathbb{A}_f)$ on $\tilde{\mathbb{H}}$ once an isomorphism $\varphi : \Delta(\mathbb{Q}_+) \cdot \text{SL}_2(\mathbb{A}_f) \rightarrow \tilde{G}$ is provided. Call φ a **naming isomorphism**. The naming isomorphisms form the family

$$\{\varphi_\mu : \Delta(\mathbb{Q}_+) \cdot \text{SL}_2(\mathbb{A}_f) \rightarrow \tilde{G} \mid \mu \in \hat{\mathbb{Z}}^\times\}; \quad \text{set } \mathbf{g}^\mu := \varphi_\mu(\mathbf{g}),$$

which satisfies

$$\mathbf{g}^{\mu\lambda} = (\mathbf{d}_\lambda^{-1} \cdot \mathbf{g} \cdot \mathbf{d}_\lambda)^\mu$$

where the conjugation by \mathbf{d}_λ is in the ambient group $\text{GL}_2(\mathbb{A}_f)$.

(b) Any $\mathbf{g} \in \Delta(\mathbb{Q}_+) \cdot \text{SL}_2(\mathbb{A}_f)$ determines the 0-definable subset of \tilde{G} :

$$\mathbf{g}^\Delta = \{\mathbf{g}^\mu : \mu \in \hat{\mathbb{Z}}^\times\}.$$

Each \mathbf{g}^μ gives rise to the sequence of algebraic curves defined over $\bar{\mathbb{Q}}$,

$$C_{\mathbf{g},N}^\mu \subset \mathbb{Y}(N) \times \mathbb{Y}(N); \quad C_{\mathbf{g},N}^\mu := \{\langle j_N(u), j_N(\mathbf{g}^\mu \cdot u) \rangle : u \in \tilde{\mathbb{H}}\}$$

(here $C_{\mathbf{g},N}^\mu$ corresponds to $C_{\mathbf{g},\approx K}^\mu$ of [10]) and the *limit curve*

$$C_{\mathbf{g}}^\mu \subset \tilde{\mathbb{H}} \times \tilde{\mathbb{H}}; \quad C_{\mathbf{g}}^\mu := \{(u, \mathbf{g}^\mu u) : u \in \tilde{\mathbb{H}}\}.$$

(c) For a fixed $\mathbf{g} \in \Delta(\mathbb{Q}_+) \cdot \text{SL}_2(\mathbb{A}_f)$ we obtain the finite family of curves on $\mathbb{Y}(N) \times \mathbb{Y}(N)$

$$\{C_{\mathbf{g},N}^\mu, \mu \in \hat{\mathbb{Z}}^\times\}$$

where

$$\mu - \lambda \in N\hat{\mathbb{Z}} \implies C_{\mathbf{g},N}^\mu = C_{\mathbf{g},N}^\lambda.$$

These curves are irreducible components of the algebraic curve $C_{\mathbf{g},N}$ defined over \mathbb{Q} :

$$C_{\mathbf{g},N} = \bigcup_{\mu \in \hat{\mathbb{Z}}^\times} C_{\mathbf{g},N}^\mu \subset \mathbb{Y}(N) \times \mathbb{Y}(N).$$

In the limit one obtains the the infinite-component limit curve on $\tilde{\mathbb{H}} \times \tilde{\mathbb{H}}$:

$$C_{\mathbf{g}} = \bigcup_{\mu \in \hat{\mathbb{Z}}^\times} C_{\mathbf{g}}^\mu.$$

$C_{\mathbf{g}}$ is defined over \mathbb{Q} too.

(d) The irreducible components $C_{\mathbf{g},N}^\mu$ of $C_{\mathbf{g},N}$ are Galois conjugated over \mathbb{Q} .

Clearly $C_{\mathbf{g}} = C_{\mathbf{g}'}$ for $\mathbf{g}' = \mathbf{g}^\mu$. The image of $C_{\mathbf{g}}$ under $j_N \times j_N$ is $C_{\mathbf{g},N}$.

(e) For any $\mathbf{g} \in \Delta(\mathbb{Q}_+) \cdot \mathrm{SL}_2(\mathbb{A}_f)$, the definable 4-ary relation on $\tilde{\mathbb{H}}$

$$\mathrm{Comp}_{\mathbf{g}}(s_1, s_2, t_1, t_2) := \exists h \in \tilde{\mathbb{G}} \ s_2 = h \cdot s_1 \ \& \ t_2 = h \cdot t_1 \ \& \ C_{\mathbf{g}}(s_1, s_2)$$

determines the condition that $\langle s_1, s_2 \rangle$ and $\langle t_1, t_2 \rangle$ belong to the same irreducible component of a $C_{\mathbf{g}}$.

For each N the relation $\mathrm{Comp}_{\mathbf{g},N}$ on $\mathbb{Y}(N)$ is the image of $\mathrm{Comp}_{\mathbf{g}}$ under j_N , the relation which determines the decomposition of the algebraic curve $C_{\mathbf{g},N}$ into irreducible components. This relation is invariant under $\mathrm{Gal}_{\mathbb{Q}}$ and so definable over \mathbb{Q} .

We also have on $\tilde{\mathbb{H}}$ the definable equivalence

$$j_N(s_1) = j_N(s_2),$$

which can be equivalently given by:

$$\exists \gamma \in \tilde{\Gamma}(n) \ s_2 = \gamma \cdot s_1.$$

(f) The points s in $\tilde{\mathbb{H}}$ which are fixed by a $g \in \tilde{\mathbb{G}} \setminus \varphi_\mu(Z)$ (Z is the centre of $\mathrm{GL}_2^+(\mathbb{Q})$) will be called special, or CM-points of $\tilde{\mathbb{H}}$. $j_N(s)$ is a CM-point in $\mathbb{Y}(N)$ and, in particular, is algebraic.

The paper [10] describes the binary relations on $\tilde{\mathbb{H}}$ written as

$$\langle t_1, t_2 \rangle \in \mathrm{tp}(s_1, s_2)$$

defined for each pair s_1, s_2 of CM-points. The relation is valid if and only if there is an automorphism σ of the projective system (4) such that $\sigma : \langle s_1, s_2 \rangle \mapsto \langle t_1, t_2 \rangle$. The relations are invariant under automorphisms of the projective system (i.e. defined over \mathbb{Q}) for each choice of s_1, s_2 .

The image of the relation under j_N is the relation on $\mathbb{Y}(N)$

$$\langle y_1, y_2 \rangle \in \text{tp}_N(x_1, x_2)$$

which holds if and only if $\langle y_1, y_2 \rangle$ is Galois conjugated to $\langle x_1, x_2 \rangle$ over \mathbb{Q} .

3.6 Definition. $\tilde{\mathbb{H}}^{\text{Pure}}$ is the structure with the universe $\tilde{\mathbb{H}}$ and relations $C_{\mathbf{g}}(s_1, s_2)$, $\text{Comp}_{\mathbf{g}}(s_1, s_2, t_1, t_2)$ ($\mathbf{g} \in \Delta(\mathbb{Q}_+) \cdot \text{SL}_2(\mathbb{A}_f)$), $j_N(s_1) = j_N(s_2)$ and $\langle t_1, t_2 \rangle \in \text{tp}(s_1, s_2)$.

$\tilde{\mathbb{H}}^{\text{Full}}$ is the multisorted structure with sorts $\tilde{\mathbb{H}}$ and $\mathbb{Y}(N)$, $N \in \mathbb{N}$, and relations:

- on $\tilde{\mathbb{H}}$ the relations of $\tilde{\mathbb{H}}^{\text{Pure}}$;
- on the $\mathbb{Y}(N)$ the Zariski closed relations defined over \mathbb{Q} ;
- the maps $j_N : \tilde{\mathbb{H}} \rightarrow \mathbb{Y}(N)$ and $\text{pr}_{N, N/d} : \mathbb{Y}(N) \rightarrow \mathbb{Y}(N/d)$.

$\tilde{\mathbb{H}}^{\text{Pure}}(\text{CM})$ and $\tilde{\mathbb{H}}^{\text{Full}}(\text{CM})$ are substructures of the structures with universes restricted to their special points.

\mathbb{H}^{Full} and \mathbb{H}^{Pure} are “standard” substructures of $\tilde{\mathbb{H}}^{\text{Full}}$ and $\tilde{\mathbb{H}}^{\text{Pure}}$, respectively, with the sort \mathbb{H} in place of $\tilde{\mathbb{H}}$.

In particular, relations $C_{\mathbf{g}}$ and $\text{Comp}_{\mathbf{g}}$ are empty in \mathbb{H}^{Full} unless $\mathbf{g} \in \text{GL}_2^+(\mathbb{Q})$. Also, by [10], 3.24.3, the relation $\langle t_1, t_2 \rangle \in \text{tp}(s_1, s_2)$ on \mathbb{H} can be expressed in terms of actions by elements of $\text{GL}_2^+(\mathbb{Q})$: if s_1, s_2 are fixed points of respective $g_1, g_2 \in \text{GL}_2^+(\mathbb{Q})$ then either $t_1 = s_1$, $t_2 = s_2$, or t_1, t_2 are fixed points of the conjugates $\check{g}_1 := \mathbf{d}_{-1}g_1\mathbf{d}_{-1}$ and $\check{g}_2 := \mathbf{d}_{-1}g_2\mathbf{d}_{-1}$ respectively.

Remarks. (i) Since components of the $C_{\mathbf{g}}(s_1, s_2)$ are graphs of actions of elements of the group \tilde{G} , we often look at $\tilde{\mathbb{H}}$ as a \tilde{G} -set.

(ii) A corollary of definitions is that the relations $C_{\mathbf{g}}$, $\text{Comp}_{\mathbf{g}}$, and $\langle t_1, t_2 \rangle \in \text{tp}(s_1, s_2)$ are projective limits of the relations $C_{\mathbf{g}, N}$, $\text{Comp}_{\mathbf{g}, N}$, and $\langle t_1, t_2 \rangle \in \text{tp}_N(s_1, s_2)$ on the $\mathbb{Y}(N)$. That is the relations on the sort $\tilde{\mathbb{H}}$ are positive-type-definable in terms of pull-backs of the respective relations on the $\mathbb{Y}(N)$ along with the pull-back of equality.

We thus can, up to $L_{\omega_1, \omega}$ -bi-interpretability, identify $\tilde{\mathbb{H}}^{\text{Pure}}$ as the structure given by the pull-backs of $C_{\mathbf{g}, N}$, $\text{Comp}_{\mathbf{g}, N}$, $\langle t_1, t_2 \rangle \in \text{tp}_N(s_1, s_2)$ and equality.

Call it **the pull-backs version of $\tilde{\mathbb{H}}^{\text{Pure}}$** . In the same sense we speak about the pull-backs version of \mathbb{H}^{Pure} , a substructure of the pull-backs version of $\tilde{\mathbb{H}}^{\text{Pure}}$.

3.7 Lemma. *The pull-backs versions of $\tilde{\mathbb{H}}^{\text{Pure}}$ and of \mathbb{H}^{Pure} are locally modular of trivial type.*

The structures satisfy

$$\mathbb{H}^{\text{Pure}} < \tilde{\mathbb{H}}^{\text{Pure}}.$$

Proof. The structure \mathbb{H}^{Pure} satisfies the assumptions of Theorems 4.11 and 4.14 of [1]. It follows that its theory has quantifier elimination and is of trivial type. $\tilde{\mathbb{H}}^{\text{Pure}}$ is then its elementary extension in the obvious way. \square

3.8 Remarks. It is easy to see that the centre Z of $\text{GL}_2^+(\mathbb{Q})$ acts on $\tilde{\mathbb{H}}$ trivially.

Note that in the definitions above the curves $\mathbb{Y}(N)$ are curves over \mathbb{C} (defined over \mathbb{Q}) and so the points of $\tilde{\mathbb{H}}$ are limits of \mathbb{C} -points.

However the definitions and results are valid in the context of curves $\mathbb{Y}(N)$ over F , an abstract algebraically closed field of characteristic 0. In this case $\mathbb{Y}_F(N)$ are curves over F .

3.9 Define $\tilde{\mathbb{U}}_F^{\text{Pure}}$ and $\tilde{\mathbb{U}}_F^{\text{Full}}$ to be the respective structures obtained as the projective limit of the $\mathbb{Y}_F(N)$, for an arbitrary algebraically closed field of characteristic 0.

3.10 Remark. Note that

$$\tilde{\mathbb{H}}^{\text{Pure}}(\text{CM}) = \tilde{\mathbb{U}}_F^{\text{Pure}}(\text{CM})$$

since both sides are the structures obtained by taking the projective limit of the respective substructures $\mathbb{Y}(n)^{\text{Pure}}(\text{CM})$ on the curves. The same is true for the full structures:

$$\tilde{\mathbb{H}}^{\text{Full}}(\text{CM}) = \tilde{\mathbb{U}}_F^{\text{Full}}(\text{CM}).$$

3.11 Define

$$\tilde{\mathbb{E}} = \{e \in \tilde{\mathbb{G}} \setminus Z \exists u \in \tilde{\mathbb{U}} e \cdot u = u\}.$$

Set

$$\tilde{G}_* = (\tilde{G}, \tilde{\Gamma}, \tilde{E}, \{\mathbf{d}_q, \mathbf{d}'_q : q \in \mathbb{Q}_+\})$$

the structure on the group \tilde{G} with distinguished subgroup $\tilde{\Gamma}$, distinguished subset \tilde{E} and distinguished elements \mathbf{d}_q and \mathbf{d}'_q .

Analogously,

$$G_* = (G, \Gamma, E, \{\mathbf{d}_q, \mathbf{d}'_q : q \in \mathbb{Q}_+\}).$$

Clearly,

$$G_* \subset \tilde{G}_*$$

as structures.

3.12 Remark. There is an embedding

$$\tilde{G} \subset \mathrm{GL}_2(\mathbb{A}_f)$$

which is an identity on the diagonal elements $\mathbf{d}_q, \mathbf{d}'_q : q \in \mathbb{Q}_+$. It is easy to see that such an embedding is determined uniquely, up to the conjugation by $\mathbf{d}_\mu, \mu \in \hat{\mathbb{Z}}^\times$.

In particular, we may assume that elements g of \tilde{G} are also elements of $\mathrm{GL}_2(\mathbb{A}_f)$ and thus the conjugation by an element $\mathbf{d}_\lambda \in \Delta$,

$$g \mapsto \mathbf{d}_\lambda \cdot g \cdot \mathbf{d}_\lambda^{-1}$$

is well defined.

3.13 Lemma. *Consider the natural embedding $G_* \subset \tilde{G}_*$ and let*

$$\psi : G_* \rightarrow G'_* \subset \tilde{G}_*$$

be a partial isomorphism of \tilde{G}_ .*

ψ can be extended to an automorphism $\tilde{\psi} : \tilde{G}_ \rightarrow \tilde{G}_*$. Moreover, there is $\lambda \in \hat{\mathbb{Z}}^\times$ such that*

$$\tilde{\psi} : g \mapsto \mathbf{d}_\lambda \cdot g \cdot \mathbf{d}_\lambda^{-1}, \text{ for all } g \in \tilde{G}.$$

Proof. Recall that the subgroup $\Gamma \subset G$ is generated by elements \mathbf{s} and \mathbf{t} , and G is generated over Γ by elements $\mathbf{d}_q, q \in \mathbb{Q}_+$.

Let $\mathbf{s}' = \psi(\mathbf{s}) \in \Gamma' \subset G'$. Then

$$\mathbf{s}' \in \tilde{G}, \quad \mathbf{s}' \cdot \mathbf{d}_{-1} = \mathbf{d}'_{-1} \mathbf{s}' \text{ and } \mathbf{s}'^2 = -\mathbf{I} = \mathbf{d}_{-1} \cdot \mathbf{d}'_{-1}.$$

It is easy to see that the three equations imply that, for some $\lambda \in \widehat{\mathbb{Z}}^\times$,

$$\mathbf{s}' = \begin{pmatrix} 0 & \lambda \\ -\lambda^{-1} & 0 \end{pmatrix} = \mathbf{d}_\lambda \cdot \mathbf{s} \cdot \mathbf{d}_\lambda^{-1}.$$

Let $\mathbf{t}' = \psi(\mathbf{t}) \in G'$. Then for each $n \in \mathbb{N}_{>0}$,

$$\mathbf{t}' \in \tilde{\Gamma}, \quad \mathbf{d}_n \cdot \mathbf{t}' \cdot \mathbf{d}_n^{-1} = \mathbf{t}'^n \quad \text{and} \quad (\mathbf{s}'\mathbf{t}')^3 = \mathbf{I}.$$

It follows

$$\mathbf{t}' = \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} = \mathbf{d}_\lambda \cdot \mathbf{t} \cdot \mathbf{d}_\lambda^{-1}.$$

Thus, G' is the group generated by \mathbf{s}' , \mathbf{t}' and \mathbf{d}_q , $q \in \mathbb{Q}_+$, and

$$\psi : g \mapsto \mathbf{d}_\lambda \cdot g \cdot \mathbf{d}_\lambda^{-1} \quad \text{for all } g \in G.$$

Take

$$\tilde{\psi} : g \mapsto \mathbf{d}_\lambda \cdot g \cdot \mathbf{d}_\lambda^{-1} \quad \text{for all } g \in \tilde{G}.$$

It is clear that $\tilde{\psi}$ preserves $\tilde{\Gamma}$, \mathbf{d}_q and \mathbf{d}'_q , $q \in \mathbb{Q}_+$.

Finally note that $g \in \tilde{E}$ if and only if $\tilde{\psi}(g) \in \tilde{E}$. This follows from the description of fixed points in 3.22D(b) of [10].

□

4 The action of $\text{Aut } F$ on $\tilde{U}_F^{\text{Full}}$.

The following is essentially a corollary of the main results of [10].

4.1 Theorem.

(i) $\text{Gal}(\mathbb{Q}(\text{CM}) : \mathbb{Q})$ acts naturally on $\tilde{U}_F^{\text{Pure}}(\text{CM})$, and

$$\text{Gal}(\mathbb{Q}(\text{CM}) : \mathbb{Q}) = \text{Aut } \tilde{U}_F^{\text{Pure}}(\text{CM}).$$

(ii) Any automorphism of $\tilde{U}_F^{\text{Pure}}(\text{CM})$ extends to an automorphism of $\tilde{U}_F^{\text{Full}}$.

(iii) For $s_1, \dots, s_n, t_1, \dots, t_n \in \tilde{U}_F^{\text{Pure}}(\text{CM})$:

$$\exists \sigma \in \text{Aut } \tilde{U}_F^{\text{Pure}}(\text{CM}) \quad \bigwedge_{i=1}^n \sigma : s_i \mapsto t_i \quad \Leftrightarrow \quad \models \bigwedge_{i=1}^n \langle t_1, t_i \rangle \in \text{tp}(s_1, s_i).$$

Proof. We rely heavily on [10].

We use the fact 3.10:

$$\tilde{\mathbb{H}}^{\text{Pure}}(\text{CM}) = \tilde{\mathbb{U}}_{\text{F}}^{\text{Pure}}(\text{CM}).$$

Given that by definition $\tilde{\mathbb{H}}^{\text{Pure}} = \mathbf{S}_{\approx}^{\text{pure}}$, 3.21 gives us (i) of the theorem. In particular, any automorphism σ of $\tilde{\mathbb{U}}_{\text{F}}^{\text{Pure}}(\text{CM})$ is induced by an automorphism $\rho \in \text{Aut F}$ acting on the CM-points, that is $\sigma = \rho|_{\text{CM}}$. But

$$\text{Aut } \tilde{\mathbb{U}}_{\text{F}}^{\text{Full}} = \text{Aut F}$$

by construction. In other words σ lifts to an automorphism ρ of $\tilde{\mathbb{U}}_{\text{F}}^{\text{Full}}$. This proves (ii).

Finally, Theorem 3.23 implies (iii). \square

4.2 Lemma. *Any $\psi \in \text{Aut } \tilde{\mathbb{G}}_*$ can be extended to $\tilde{\psi} \in \text{Aut } \tilde{\mathbb{U}}_{\text{F}}^{\text{Pure}}(\text{CM})$.*

Proof As noted above $\tilde{\mathbb{U}}_{\text{F}}^{\text{Pure}}(\text{CM}) = \tilde{\mathbb{H}}^{\text{Pure}}(\text{CM})$ so we may argue in the setting of $\tilde{\mathbb{H}}$.

By 3.13 ψ has the form $\mathbf{g}^{\mu} \mapsto \mathbf{g}^{\mu \cdot \lambda} = \mathbf{d}_{\lambda} \cdot \mathbf{g}^{\mu} \cdot \mathbf{d}_{\lambda}^{-1}$, for some $\lambda \in \hat{\mathbb{Z}}^{\times}$. And by 3.5(d) ψ is induced by a $\sigma \in \text{Gal}_{\mathbb{Q}}$. In its turn σ acts on the CM-points of $\tilde{\mathbb{U}}_{\text{F}}$ and induces a $\tilde{\psi} \in \text{Aut } \tilde{\mathbb{U}}_{\text{F}}^{\text{Pure}}(\text{CM})$, as required. \square

5 Axiomatisation of $\mathbb{H}_{\text{F}}^{\text{Full}}$ and $\mathbb{U}_{\text{F}}^{\text{Full}}$.

5.1 The language and axioms Σ .

The language $\mathcal{L}(j_{n \in \mathbb{N}})$ is 3-sorted, with sorts \mathbb{U}, \mathbb{G} and \mathbb{F} . The structure on \mathbb{F} is that of a field given in a standard ring language, the structure on \mathbb{G} and \mathbb{U} is that of a group acting on \mathbb{U} with distinguished subsets $\mathbb{E}, \Gamma, \{\mathbf{d}_q, q \in \mathbb{Q}^{\times}\}, \{\mathbf{d}'_q, q \in \mathbb{Q}^{\times}\}$. Note that $\{\mathbf{d}_q, q \in \mathbb{Q}^{\times}\} = \Delta(\mathbb{Q})$ with all its elements named and the same for $\{\mathbf{d}'_q, q \in \mathbb{Q}^{\times}\} = \Delta'(\mathbb{Q})$. Note that

$$\text{GL}_2^+(\mathbb{Q}) = \Delta(\mathbb{Q}_{>0}) \cdot \text{SL}_2(\mathbb{Z}) = \Delta'(\mathbb{Q}_{>0}) \cdot \text{SL}_2(\mathbb{Z})$$

and so group \mathbb{G} is isomorphic to $\text{GL}_2^+(\mathbb{Q})$ with distinguished elements of Δ, Δ' and subgroup $\Gamma \cong \text{SL}_2(\mathbb{Z})$ will have the same structure.

Note that $\text{GL}_2^+(\mathbb{Q})$ is invariant under the involutive transformation

$$g \mapsto \check{g} := \mathbf{d}_{-1} \cdot g \cdot \mathbf{d}_{-1}$$

where $\mathbf{d}_{-1} \in \mathrm{GL}_2(\mathbb{Q}) \subset \mathrm{GL}_2(\mathbb{A}_f)$.

The maps j_n have \mathbb{U} as their domain and have quasi-projective curves $\mathbb{Y}(n)$ as their range.

We consider the following system of axioms Σ which is split into five groups of axioms :

Group axioms:

$$(\mathrm{G}, \Gamma, \mathrm{E}, \{\mathbf{d}_q : q \in \mathbb{Q}_{>0}\}) \cong (\mathrm{GL}_2^+(\mathbb{Q}), \mathrm{SL}_2(\mathbb{Z}), \mathrm{E}(\mathbb{Q}), \{\mathbf{d}_q : q \in \mathbb{Q}_{>0}\}). \quad (5)$$

Note that $\Gamma(n)$ is definable from the data, namely the subgroup $\Gamma_0(n)$ of matrices of the form $\mathbf{g} = \mathbf{t}^{nm}$ are definable by condition

$$\exists \gamma \in \Gamma \quad \mathbf{g} = \mathbf{d}_n \cdot \gamma \cdot \mathbf{d}_n^{-1},$$

and $\Gamma(n)$ can be defined as the normal closure of $\Gamma_0(n)$.

Let $Z = \mathrm{Centre}(\mathrm{G})$.

Action axiom: G acts on \mathbb{U} ;

$$\begin{aligned} \forall g \in \mathrm{G} \setminus (\mathrm{E} \cup Z) \quad \forall u \in \mathbb{U} \quad g \cdot u \neq u, \\ \forall g \in Z \quad \forall u \in \mathbb{U} \quad g \cdot u = u, \\ \forall e \in \mathrm{E} \quad \exists! u_e \in \mathbb{U} \quad e \cdot u_e = u_e. \end{aligned} \quad (6)$$

Fibre formula:

$$\forall u, v \in \mathbb{U} \quad j_n(u) = j_n(v) \leftrightarrow \exists \gamma \in \Gamma(n) \quad v = \gamma \cdot u \quad (7)$$

ACF₀ axioms and sorts $\mathbb{Y}(n)$:

$$\mathrm{F} \models \mathrm{ACF}_0 \quad (8)$$

and

$$\mathbb{Y}(n) \subset \mathbf{P}^3(\mathrm{F}); \quad \mathrm{pr}_{n,m} : \mathbb{Y}(n) \rightarrow \mathbb{Y}(m), \quad \text{for } m|n$$

are given by specific equations over \mathbb{Q} .

Axioms - Functional equations:

$$j_n : \mathbb{U} \twoheadrightarrow \mathbb{Y}(n); \quad \mathrm{pr}_{n,m} \circ j_n = j_m \quad \text{for each } m|n; \quad (9)$$

$$\begin{aligned} \text{For each } \mathbf{g} \in \mathrm{GL}_2^+(\mathbb{Q}), \quad n \in \mathbb{N}, \quad \text{and } \mu \in \hat{\mathbb{Z}}^\times, \quad \exists g \in \mathbf{g}^\Delta \quad \forall u, v \in \mathbb{U}: \\ \langle j_n(u), j_n(v) \rangle \in C_{\mathbf{g},n}^\mu \leftrightarrow \exists v', u' \in \mathbb{U} \quad v' = g \cdot u' \quad \& \quad j_n(u') = j_n(u) \quad \& \quad j_n(v') = j_n(v) \end{aligned} \quad (10)$$

For each n and any $g_1 \in \mathbf{g}_1^\Delta, \dots, g_m \in \mathbf{g}_m^\Delta$:

$$\begin{aligned} & \{ \langle j_n(u), j_n(g_1 u), \dots, j_n(g_m u) \rangle : u \in \mathbb{U} \} = \\ & = \text{an irreducible component of} \\ & \{ \langle y, y_1, \dots, y_m \rangle \in \mathbb{Y}(n)^{m+1} : \bigwedge_{i=1}^m \langle y, y_i \rangle \in C_{\mathbf{g}_i, n} \} \end{aligned} \quad (11)$$

5.2 Commentary. Axioms (5), (6) and (7) describe the structure $\mathbb{U}_F^{\text{Pure}}$.

Axiom (8) describes the algebro-geometric part of $\mathbb{U}_F^{\text{Full}}$, the structure on the F-curves $\mathbb{Y}(n)$.

Axioms (9)-(11) describes the connection between $\mathbb{U}_F^{\text{Pure}}$ and the $\mathbb{Y}(n)$ in terms of the j_n . More precisely, (10) states, in accordance with 3.5, that an algebraic curve of the form $C_{\mathbf{g}, n}^\mu$ is the image of the graph of g under j_n for some $g \in \mathbf{g}^\Delta$. Combining this with (11) for $m = 1$ and using the fact that curves $C_{\mathbf{g}, n}^\mu$ are irreducible the axioms imply that for all $g \in \mathbf{g}^\Delta$ the image of the graph of g under j_n is of the form $C_{\mathbf{g}, n}^\mu$ for some $\mu \in \hat{\mathbb{Z}}^\times$.

Note that for a given $\mathbf{g}_1, \dots, \mathbf{g}_m$ the algebraic subvariety $\{ \langle y, y_1, \dots, y_m \rangle : \bigwedge_{i=1}^m \langle y, y_i \rangle \in C_{\mathbf{g}_i, n} \}$ of $\mathbb{Y}(n)^{m+1}$ of (11) is well-defined (although its explicit representation in terms of equations could be hard to obtain) and thus there is a well-defined finite list of its irreducible components.

5.3 Lemma. $\mathbb{H}_\mathbb{C}^{\text{Full}}$ is a model of axioms Σ .

Proof. The axioms just describe formally the structure $\mathbb{H}_\mathbb{C}^{\text{Full}}$ following 3.5. \square

5.4 Lemma. For any model $\mathbb{U}_F^{\text{Full}}$ of Σ there is an embedding

$$i : \mathbb{U}_F^{\text{Full}} \hookrightarrow \tilde{\mathbb{U}}_F^{\text{Full}} \quad (12)$$

which is the identity on F .

Proof. $\tilde{\mathbb{U}}$ in $\tilde{\mathbb{U}}_F^{\text{Full}}$ is given as the universal cover of $\mathbb{Y}(F)$, with respect to the class of modular curves. By axiom (9) \mathbb{U} covers every principal modular curve $\mathbb{Y}(n)$ by $j_n : \mathbb{U} \rightarrow \mathbb{Y}(n)$. Hence there is a unique map $i : \mathbb{U} \rightarrow \tilde{\mathbb{U}}$ such that

$$j_n(i(u)) = j_n(u). \quad (13)$$

Now we construct an embedding of the groups $G \rightarrow \tilde{G}$ which agrees with the action and takes $\Gamma(n)$ to $\tilde{\Gamma}(n)$. In order to do this recall that (10)-(11) imply that there is a bijective correspondence between elements of $g \in G$ and

the sequences $C_{\mathbf{g},n}^\mu$, $n \in \mathbb{N}$, where μ depends on g (see 5.2 above). On the other hand the sequence $C_{\mathbf{g},n}^\mu$, $n \in \mathbb{N}$, determines a limit curve $C_{\mathbf{g}}^\mu \subset \tilde{\mathbb{H}}^2$ (see 3.5(b)) and thus an element, call it $i(g)$, of \tilde{G} together with its action on $\tilde{\mathbb{H}}$.

Note that (13) together with the correspondnce of group actions imply that

$$j_n(\gamma \cdot u) = j_n(u) \Leftrightarrow j_n(i(\gamma) \cdot i(u)) = j_n(i(u))$$

and hence by (6) $\gamma \in \Gamma(n) \Leftrightarrow i(\gamma) \in \tilde{\Gamma}(n)$.

Finally the structure induced on the $\mathbb{Y}(n)_F$ from F is the same by virtue of (13) and axioms (8) and (9).

□

Given a model $\mathbb{U}_F^{\text{Full}}$ of Σ and $g \in E$ we denote u_g the unique fixed point of g in \mathbb{U} .

Given $g_1, \dots, g_k \in G$, we denote $\text{Type}(g_1, \dots, g_k)$, the type of the tuple in G , to be the formula in terms of the structure G_* such that: if $G'_* \cong G_*$ and $g'_1, \dots, g'_k \in G'$ then

$\langle g'_1, \dots, g'_k \rangle$ satisfies $\text{Type}(g_1, \dots, g_k)$ if and only if there exists an isomorphism $\psi : G_* \rightarrow G'_*$, $\langle g_1, \dots, g_k \rangle \mapsto \langle g'_1, \dots, g'_k \rangle$.

Since elements g_i of $G = \text{GL}_2^+(\mathbb{Q})$ can be written as the group words $g_i = w_i(\mathbf{s}, \mathbf{t}, \mathbf{d}_q)$, it is immediate from 3.13 that

$$\text{Type}(g_1, \dots, g_k) \equiv \exists s, t \in \Gamma \ s^2 = -\mathbf{I} \ \& \ (st)^3 = \mathbf{I} \ \& \ g_i = w_i(s, t, \mathbf{d}_q).$$

We will write \mathbb{U}_F without superscript when speaking on models of (5)-(11).

5.5 Proposition. *Let \mathbb{U}_F and \mathbb{U}'_F be two models of Σ .*

Let $\psi : G_ \rightarrow G'_*$ be an isomorphism of respective groups with extra structure and $g_1, \dots, g_k \in G$, $g'_1, \dots, g'_k \in G'$ such that $\psi : g_i \mapsto g'_i$, $i = 1, \dots, k$.*

Then, the k -tuples $\langle j_N(u_{g_1}), \dots, j_N(u_{g_k}) \rangle$ and $\langle j_N(u_{g'_1}), \dots, j_N(u_{g'_k}) \rangle$ are Galois conjugated.

Moreover, the algebraic type of $\langle j_N(u_{g_1}), \dots, j_N(u_{g_k}) \rangle$ over \mathbb{Q} is determined by $\text{Type}(g_1, \dots, g_k)$, the type of $\langle g_1, \dots, g_k \rangle$ in G_ .*

Proof. By 5.4 we have two embeddings

$$\mathbb{U}_F \subset \tilde{\mathbb{U}}_F^{\text{Full}} \text{ and } \mathbb{U}'_F \subset \tilde{\mathbb{U}}_F^{\text{Full}}$$

into the projective limit of curves $\mathbb{Y}(n)$ over F .

By 3.13 ψ can be extended to $\tilde{\psi} \in \text{Aut } \tilde{G}_*$. By 4.2 and 4.1(i) this extends to an automorphism of $\tilde{U}_F^{\text{Full}}(\text{CM})$, that is an automorphism $\psi \in \text{Gal}(\mathbb{Q}(\text{CM}) : \mathbb{Q})$ acting naturally on the CM-points of the curves $\mathbb{Y}(n)$ and respectively on $\tilde{U}_F^{\text{Full}}(\text{CM})$. Thus the special points $j_N(u_{g_i})$ are Galois conjugated to the special points $j_N(u_{g'_i})$.

Moreover, since the algebraic type of $j_N(u_{g_1}), \dots, j_N(u_{g_k})$ is determined in the substructure $\tilde{U}_F^{\text{Full}}(\text{CM})$ of $\tilde{U}_F^{\text{Full}}$ and the substructure coincides with $\tilde{H}(\text{CM})$, the type is independent on the choice of the model U_F . \square

5.6 Corollary. *Let $\bar{g} = \langle g_1, \dots, g_m \rangle \in G^m$. Then, in a model U_F of Σ , the set*

$$Z_{\bar{g},N} = \{ \langle j_N(v), j_N(g_1v), \dots, j_N(g_nv) \rangle : v \in \mathbb{U} \}$$

is an irreducible algebraic curve defined over the extension of \mathbb{Q} by some special points $j_N(u_{e_1}), \dots, j_N(u_{e_k})$.

If $\psi \in \text{Aut } \tilde{U}_F^{\text{Full}}$, $\psi : g_i \mapsto g'_i$, $e_l \mapsto e'_l$ $i = 1, \dots, m$, $l = 1, \dots, k$ then $Z_{\bar{g},N}$ is Galois conjugated to $Z_{\bar{g}',N}$.

Moreover, the algebraic type of $Z_{\bar{g},N}$ over \mathbb{Q} is determined by the type of $\langle g_1, \dots, g_k \rangle$ in G_ only in all models of Σ .*

Proof. $Z_{\bar{g},N}$ is an irreducible algebraic curve by axiom (11). Curves of this form were studied in [3] and [13]; they are special subvarieties containing infinitely many special points and thus are defined over an extension of \mathbb{Q} by special points.

Since ψ induces a Galois automorphism on CM-points, the statement of the corollary follows. \square

Given a model U_F of Σ and a naming isomorphism

$$\varphi : (\text{GL}_2^+(\mathbb{Q}), \text{SL}_2(\mathbb{Z}), \text{E}(\mathbb{Q}), \{\mathbf{d}_q : q \in \mathbb{Q}_{>0}\}) \rightarrow (\text{G}, \Gamma, \text{E}, \{\mathbf{d}_q : q \in \mathbb{Q}_{>0}\})$$

define $U_{F,\varphi}$ to be the expansion of the structure U_F by names $\mathbf{g} \in \text{GL}_2^+(\mathbb{Q})$ for each element $\varphi(\mathbf{g}) \in \text{G}$. We call the later structure an inessential G -expansions of the former. The language of $U_{F,\varphi}$ is the extension of the language $\mathcal{L}(j_{n \in \mathbb{N}})$ by the names $\mathbf{g} \in \text{GL}_2^+(\mathbb{Q})$ and the names for $j_N(u_{\mathbf{g}})$, for $\mathbf{g} \in E$. The latter are names of certain elements of $\mathbb{Y}(N)$, tuples in $\mathbb{Q} \subseteq F$, so essentially name points of the field which are special. We refer to this language as $\mathcal{L}(j_{n \in \mathbb{N}}, \mathbf{g} \in \text{GL}_2^+(\mathbb{Q}))$. Below we are interested in reducts $U_{F,\varphi}^{(N)}$ of $U_{F,\varphi}$ which ignore all the j_n and $\mathbb{Y}(n)$ except when $n = N$. In this case we use the sublanguage $\mathcal{L}(j_N, \mathbf{g} \in \text{GL}_2^+(\mathbb{Q}))$ of $\mathcal{L}(j_{n \in \mathbb{N}}, \mathbf{g} \in \text{GL}_2^+(\mathbb{Q}))$.

5.7 Lemma. *Two models \mathbb{U}_F and \mathbb{U}'_F of Σ are isomorphic if and only if, for some φ, φ' , $\mathbb{U}_{F,\varphi}$ is isomorphic to $\mathbb{U}'_{F,\varphi'}$.*

Proof. An isomorphism $\alpha : \mathbb{U}_F \rightarrow \mathbb{U}'_F$ induces by the group axiom (5) the isomorphism of groups $\varphi : \mathrm{GL}_2^+(\mathbb{Q}) \rightarrow G$ and $\varphi' : \mathrm{GL}_2^+(\mathbb{Q}) \rightarrow G$ such that $\varphi(g) \cdot u = \varphi'(g) \cdot \alpha(u)$ and the G_* structures respected. This gives us the isomorphism of named structures $\mathbb{U}_{F,\varphi}$ and $\mathbb{U}'_{F,\varphi'}$ which coincides with α on \mathbb{U} and F .

Conversely, an isomorphism $\alpha : \mathbb{U}_F \rightarrow \mathbb{U}'_F$ becomes the isomorphism $\mathbb{U}_F \rightarrow \mathbb{U}'_F$ if we ignore the naming of elements of G and G' . \square

6 The structure over the modular curve $\mathbb{Y}(N)$.

6.1 The structure \mathbb{U}_F and the A.Harris axioms.

[3] and [13] study substructures $\mathbb{U}_F^{(N)}$ of \mathbb{U}_F which are restrictions of the latter to sorts \mathbb{U} , F and $\mathbb{Y}(N)$, for a given N . Since elements g of G are named, the special points of \mathbb{U} and $\mathbb{Y}(N)$ get names u_g and $j_N(u_g)$. Now the field F has to be considered a field with constant symbols for CM-points (and hence the automorphisms of the structure fix the subfield generated by the CM-points).

Below we present axioms $\Sigma(N)$ for $\mathbb{U}_F^{(N)}$ in language $\mathcal{L}(j_N, \mathbf{g} \in \mathrm{GL}_2^+(\mathbb{Q}))$ in a form which essentially copies axioms of [3] for $\mathbb{U}_F^{(1)}$. One of the differences with [3] and [13] is that we use $\mathrm{GL}_2^+(\mathbb{Q})$ where they used $\mathrm{GL}_2^+(\mathbb{Q})/Z$.

Note that, for a finite tuple $\bar{\mathbf{g}} = \langle \mathbf{g}_1, \dots, \mathbf{g}_n \rangle \in \mathrm{GL}_2^+(\mathbb{Q})^n$, the algebraic curve $Z_{\bar{\mathbf{g}}}$ is uniquely defined by 5.6 in the standard model $\mathbb{H}^{\mathrm{Full}}$. We use these below in axiom (18).

Axioms $\Sigma(N)$:

Action axioms: For each $\mathbf{g} \in \mathrm{GL}_2^+(\mathbb{Q}) \setminus (E \cup Z)$,

$$\forall u \in \mathbb{U} \quad \mathbf{g} \cdot u \neq u \tag{14}$$

For each $\mathbf{g} \in Z$,

$$\forall u \in \mathbb{U} \quad \mathbf{g} \cdot u = u$$

For each $\mathbf{e} \in E$,

$$\exists! u_e \in \mathbb{U} \quad \mathbf{e} \cdot u_e = u_e; \tag{15}$$

The field axioms:

$$F \models \text{ACF}_0 \tag{16}$$

and

$$\mathbb{Y}(N) \subset \mathbf{P}^3(F)$$

is given by specific equations over \mathbb{Q} .

Functional equations:

For any $\mathbf{e}_1, \dots, \mathbf{e}_k \in E$, the special points $j(u_{e_1}), \dots, j(u_{e_n})$ satisfy

$$p_{\bar{\mathbf{e}}}(j_N(u_{e_1}), \dots, j_N(u_{e_k})) = 0, \tag{17}$$

for a polynomial $p_{\bar{\mathbf{e}}}(X_1, \dots, X_k)$ over \mathbb{Q} if this holds in the standard model.

For every finite tuple $\bar{\mathbf{g}} = \langle \mathbf{g}_1, \dots, \mathbf{g}_m \rangle \in \text{GL}_2^+(\mathbb{Q})^m$, and the algebraic curve $Z_{\bar{\mathbf{g}}}$ defined over $\mathbb{Q}(j(u_{e_1}), \dots, j(u_{e_k}))$

$$\begin{aligned} &\text{the map } u \mapsto \langle j_N(u), j_N(\mathbf{g}_1 u), \dots, j_N(\mathbf{g}_m u) \rangle \\ &\text{is a surjection onto the algebraic curve } Z_{\bar{\mathbf{g}}, N}. \end{aligned} \tag{18}$$

The fibre formula:

$$\forall u_1, u_2 \in \mathbb{U} \quad j_N(u_1) = j_N(u_2) \leftrightarrow \bigvee_{\gamma \in \Gamma(N)} u_2 = \gamma \cdot u_1. \tag{19}$$

6.2 Theorem ([3], [13]) *The set of axioms $\Sigma(N)$ has a unique model in every uncountable cardinal.*

S. Eterovich in [15] extended the approach further to Shimura varieties \mathcal{A}_2 and \mathcal{A}_3 , moduli spaces of abelian varieties of dimensions 2 and 3 respectively.

6.3 Lemma. *Suppose \mathbb{U}_F is a model of Σ . Then $\mathbf{U}_F^{(N)}$ is a model of $\Sigma(N)$.*

Proof. (14), (15) and (19) are immediate.

(17) follows from 5.5 and (18) from 5.6 since the tuples $\langle j_N(e_1), \dots, j_N(e_k) \rangle$ and the curve $Z_{\bar{\mathbf{g}}, N}$ are determined by $\mathbf{e}_1, \dots, \mathbf{e}_n$ and $\mathbf{g}_1, \dots, \mathbf{g}_m$ in the standard model. \square

6.4 6.2 together with 6.3 fails to imply categoricity of Σ .

Nevertheless, the arguments from section 4 and 5 of [13] are sufficient to prove categoricity of Σ . We proceed with the arguments below.

We fix notation for the subfield

$$L_0 = \mathbb{Q}(\text{CM}) \subset F.$$

6.5 Lemma. *Let \bar{a} be an m -element subset of points of $\mathbb{Y}(1) = \mathbf{A}^1(\mathbb{F}) = \mathbb{F}$ and $L = L_0(\bar{a})$. There exists a homomorphism*

$$\mathfrak{a} : \text{Aut}(\mathbb{F}/L) \rightarrow \tilde{\Gamma}^m$$

such that the number of $\text{Aut}(\mathbb{F}/L)$ -orbits in $j^{-1}(\bar{a})$ is equal to the index of $\mathfrak{a}(\text{Aut}(\mathbb{F}/L))$ in $\tilde{\Gamma}^m$.

Proof. Consider the canonical action of $\text{Aut} \mathbb{F}$ on $\tilde{\mathbb{U}}$ and $\tilde{\Gamma}$.

Note that $\text{Aut}(\mathbb{F}/L)$ fixes $\tilde{\Gamma}$ point-wise since it fixes all the CM-points.

Let $\bar{u} \in \tilde{\mathbb{U}}^m$, $\bar{u} \in j^{-1}(\bar{a})$, $\sigma \in \text{Aut}(\mathbb{F}/L)$. Since $\bar{a} \subset L$, $\sigma(\bar{u})$ belongs to the same fibre $j^{-1}(\bar{a})$ and hence

$$\sigma(\bar{u}) = \gamma_{\sigma, \bar{u}} \cdot \bar{u}, \text{ for some } \gamma_{\sigma, \bar{u}} \in \tilde{\Gamma}^m.$$

Claim. $\gamma_{\sigma, \bar{u}}$ does not depend on the choice of \bar{u} in $j^{-1}(\bar{a})$. The map

$$\sigma \mapsto \gamma_{\sigma}; \quad \text{Aut}(\mathbb{F}/L) \rightarrow \tilde{\Gamma}^m$$

is a group homomorphism.

This follows by group-theoretic calculations.

By the same calculation, for a $\bar{v} \in j^{-1}(\bar{a})$,

$$\exists \sigma \in \text{Aut}(\mathbb{F}/L) \sigma \bar{u} = \bar{v} \Leftrightarrow \exists \gamma \in \mathfrak{a}(\text{Aut}(\mathbb{F}/L)) \gamma \bar{u} = \bar{v}.$$

Hence $\text{Aut}(\mathbb{F}/L)$ -orbits are in a bijective correspondence with $\mathfrak{a}(\text{Aut}(\mathbb{F}/L))$ -cosets in $\tilde{\Gamma}^m$.

□

6.6 Proposition. *Σ is κ -categorical for all uncountable κ if and only if the following two conditions hold:*

- (i) *given \bar{a} such that its co-ordinates $a_i \notin L_0$ and in distinct Hecke orbits, the image of $\text{Aut}(\mathbb{F}/L)$ in $\tilde{\Gamma}^m$ is of finite index;*
- (ii) *given an algebraically closed subfield $K \subset \mathbb{F}$ and \bar{a} such that its co-ordinates $a_i \notin K$ and in distinct Hecke orbits, the image of $\text{Aut}(\mathbb{F}/K(\bar{a}))$ in $\tilde{\Gamma}^m$ is of finite index.*

Remark. The necessary condition (ii) is equivalent to a weaker condition which assumes that $\text{tr.deg} K(\bar{a})/K = 1$.

(i) and (ii) are effectively the conditions 4.7 and 4.8 of [13].

Proof of the Proposition. The same as the combined proofs of Theorems 4.1 and 4.9 in [13]. □

6.7 Proposition. *The conditions (i) and (ii) of 6.6 are satisfied.*

Proof. See 5.1 and 5.2 of [13]. \square

Remark. The key reference in the proof is the open image theorem of Serre, [14].

As a corollary we get the main theorem.

6.8 Theorem. Σ is κ -categorical for all uncountable κ .

References

- [1] B.Zilber, *Model theory of special subvarieties and Schanuel-type conjectures* Annals of Pure and Applied Logic, Volume 167, Issue 10, October 2016, pp. 1000-1028
- [2] M.Gavrilovich, **Model theory of the universal covering spaces of complex algebraic varieties**, PhD Thesis, Oxford, 2006
- [3] A.Harris, *Categoricity and covering spaces*, PhD Thesis, Oxford, 2013
- [4] R.Abdolazhadi and B.Zilber, *Definability, interpretation and the étale fundamental groups*, arXiv:1906.05052v2
- [5] R.Langlands, *Automorphic representations, Shimura varieties, and motives, Ein Märchen*, In **Proc. Sympos. Pure Math.**, Vol. 33, Part 2, Amer. Math. Soc, Providence, R. I., 1979, pp. 205-246
- [6] P.Deligne, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*, **Automorphic forms, representations and L-functions** (Part 2), Proc. Sympos. Pure Math., XXXIII, AMS, Providence, R.I., 1979, pp. 247–289
- [7] M.Borovoi, *Conjugation of Shimura varieties*. In:”**Proc. Internat. Congr. Math.**, Berkeley, 1986”, AMS, 1987, pp. 783 – 790.

- [8] J.Milne, *Introduction to Shimura varieties*, In **Harmonic Analysis, the trace formula and Shimura varieties**, Clay Math. Proc., 2005, pp.265–378
- [9] J.Milne, **Canonical models of Shimura curves**, Notes, 2003 author’s web-site
- [10] B.Zilber and C.Daw, *Canonical models of modular curves and Galois action on CM-points*
- [11] N.D. Elkies, *The Klein quartic in number theory*, pp. 51–102 in **The Eightfold Way: The Beauty of Klein’s Quartic Curve**, S.Levy, ed.; Cambridge Univ. Press, 1999
- [12] F.Bars, A.Kontogeorgis, and X.Xarles, *Bielliptic and hyperelliptic modular curves $X(N)$ and the group $\text{Aut } X(N)$* , Acta Arithmetica 161(3), 2012
- [13] C.Daw and A.Harris, *Categoricity of modular and Shimura curves*, Journal of the Institute of Mathematics of Jussieu, v. 16, 5, 2017, pp.1075 –1101
- [14] J.-P. Serre, *Propriétés galoisiennes des points d’ordre fini des courbes elliptiques*, Invent. Math. 15 (1972), 259–331
- [15] S.Eterovich, *Categoricity of Shimura varieties*, DPhil Thesis, Oxford, 2019; axiv:1803.06700
- [16] B.Zilber, *Covers of the multiplicative group of an algebraically closed field of characteristic zero*, J. London Math. Soc. (2), 74(1):41–58, 2006