

Possibility of Primordial black holes Collision with Earth and the Consequences of this Collision

Sohrab Rahvar*

¹*Physics Department, Sharif University, P.O.Box 11365-9161, Azadi Avenue, Tehran, Iran*

26 July 2021

ABSTRACT

In the context of the existence of Primordial black holes (PBHs), they may compose a fraction of the dark matter of the Universe. Assuming that PBHs fill the dark content of the Milky Way Galaxy in the Galactic halo and dark disk, we calculate the probability of collision of the PBHs with Earth. This collision has different consequences as heating the interior of the earth through dynamical friction and accretion processes. In this work, we calculate the rate of collisions and a fraction of black holes that can be trapped inside the earth. Finally, we compare the danger of PBH collision with the asteroid impacts on Earth.

Key words: black hole physics, Earth

1 INTRODUCTION

The existence of black holes as the singularities of space-time is one of the predictions of general relativity (Penrose 1965). These objects have been observed indirectly for decades by different methods (Samimi et al 1979; Shipman et al 1975). Recent observations of gravitational waves by the LIGO and Virgo detectors revealed the existence of massive binary black holes where merging of black holes generate the burst of gravitational waves (Abbott et al. 2016; Abbott et al 2019). The mass of binary black holes derived from the analysis of gravitational wave data indicates that the mass should be around tens of solar masses (Abbott et al 2017, 2020). It has been known that the mass of black holes from the astrophysical process is around a few solar masses and this mass is much small compared to the mass of binary black holes derived from the LIGO observations. One of the possibilities for the production of these large mass black holes could be through the process of Primordial Black (PBH) formation in the early universe (Sasaki et al. 2016). The interpretation of LIGO data based on this hypothesis is under investigation (Khalouei et al. 2021).

There are several scenarios to describe the formation of PBHs as the collapse of over dense regions in the early universe where in this case we expect a width mass function for PBHs (Kühnel & Freese 2017; Niemeyer & Jedamzik 1998; Carr & Kühnel 2020). Parts of window for the mass function of the PBHs have been excluded by various observations such as microlensing observations of compact objects along the Large and Small Magellanic Clouds and Galactic Bulge (Green 2016). However, there are still two open regions for

this window for the small mass (i.e. $M < 10^{-7} M_{\odot}$) and large masses (i.e. $M > 10^5 M_{\odot}$). While the black holes are evaporating by the Hawking radiation, however large mass black holes from the early times can still survive from the radiation. Assuming the existence of PBHs from the early Universe, in this work, we examine the scenario of collision of the PBHs with the earth and investigate the consequences of this collision.

In Section (2), we calculate the rate of PBHs collision with the Earth as a function of their masses. In Section (3), we investigate the dissipation process from the collision of black holes with earth. Also we calculate the probability of trapping black holes at the interior of the earth. In Section (4) we study the energy release during the collision of black holes with earth and compare it with the impact of asteroids. The conclusion is given in (5).

2 RATE OF PBH COLLISION WITH EARTH

Let us assume a fraction of dark matter is made of PBHs with a mass function of $f(M)$ for PBHs. For simplicity in the calculation, we take a Dirac-Delta mass function $\delta(m - m_b)$ for the black holes, however, we will consider a mass function in this calculation when we are calculating the overall rate of collisions with earth. For the mass density of dark matter, $\rho_{DM}(x)$, the number density of black holes is $n_{bh} = f\rho_{DM}(x)/m_{bh}$ where f is the fraction of halo is made of black holes. For PBHs moving inside the dark halo with the dispersion velocity associated with the objects inside the halo, PBHs may collide with the stars, planets, and the earth. Since the dark component of our galaxy is in

* E-mail: rahvar@sharif.edu

the halo and dark disk, the collisions can happen from these two different structures.

We take a black hole as a projectile approaching the earth. The maximum amount of angular momentum of a black hole at the closest approach to earth is $\ell = R_e m v_{max}$ where v_{max} is the maximum velocity that the black hole can achieve in a tangent collision with earth and R_e is the radius of the earth. On the other hand, the angular momentum of the black hole with respect to earth at infinity is $\ell = b m v$ where b is the impact parameter and v is the relative velocity of the black hole with respect to the earth at infinity. Using the conservation of angular momentum and conservation of energy, the impact parameter obtain as

$$b = R_e \left(1 + \frac{2GM_e}{R_e v^2}\right)^{1/2}, \quad (1)$$

where M_e is the mass of Earth. The cross-section associated to this collision is $\sigma = \pi R_e^2 (1 + v_{es}^2/v^2)$ where v_{es} is the escape velocity from earth. From the cross-section, the number density of black holes in space, and relative velocity of black holes with respect to the earth, we can calculate the impact rate of black holes with the earth. Let us take a normalized probability distribution function for the velocity of black holes as $p(v)$, the rate of collisions of the black holes with the earth is

$$\frac{dN}{dt} = \int J(v) \sigma(v) p(v) dv^3, \quad (2)$$

where $J(v) = n_{bh} v$ is the current of black holes and

$$p(v) = \frac{1}{(2\pi\sigma^2)^{3/2}} \exp(-v^2/2\sigma^2) \quad (3)$$

is the normalized distribution function of objects in the Galactic halo (Binney & Tremaine 2008). The result of integration is

$$\frac{dN}{dt} = \sqrt{2\pi} n_{bh} R_e^2 \frac{(v_{es}^2 + 2\sigma^2)}{\sigma}. \quad (4)$$

The local dark matter of Galaxy is composed from the galactic halo structure and local dark disk. The standard halo model provides the density of halo dark matter to be $\rho_h = 8 \times 10^{-3} M_\odot pc^{-3}$ with the dispersion velocity of objects inside halo, $\sigma = 220 \text{ km s}^{-1}$ (Binney & Tremaine 2008). On the other hand the investigation on the dynamics of stars perpendicular to the galactic disk with the Gaia data provides the local density of Dark Disk (DD), $\rho_{DD} = 0.016 M_\odot pc^{-3}$ (Buch et al. 2019). The dispersion velocity of thin disk model which is compatible with the other observations are as $\sigma_r = 27.4 \text{ km s}^{-1}$, $\sigma_\theta = 20.8 \text{ km s}^{-1}$ and $\sigma_z = 16.3 \text{ km s}^{-1}$ where overall dispersion $\sigma^2 = \sigma_r^2 + \sigma_\theta^2 + \sigma_z^2$ is $\sigma = 38.1 \text{ km s}^{-1}$ (Moniez et al. 2017). We adapt this dispersion velocity for the black holes in the dark disk. Substituting the numerical values in equation (4), the rate of collision of the black holes with earth from the galactic halo and dark disk is

$$\left(\frac{dN}{dt}\right)_{halo} = 7.2 \times 10^{-9} f \left(\frac{M_{bh}}{10^{23} \text{ gr}}\right)^{-1} \text{Gyr}^{-1}, \quad (5)$$

$$\left(\frac{dN}{dt}\right)_{DD} = 2.6 \times 10^{-9} f \left(\frac{M_{bh}}{10^{23} \text{ gr}}\right)^{-1} \text{Gyr}^{-1}, \quad (6)$$

where adding the contribution of the halo and dark disk, the overall rate of collision is

$$\frac{dN}{dt} \simeq 10^{-8} f \left(\frac{M_{bh}}{10^{23} \text{ gr}}\right)^{-1} \text{Gyr}^{-1}. \quad (7)$$

In order to have at least one collision of PBHs with earth per one billion year, we have to put the constrain of $M_{bh} < 10^{15} f \text{ gr}$. This mass range for PBHs still is not constrained by the other observations.

The black holes lose their masses through the Hawking radiation as (Hawking 1974)

$$\frac{dM}{dt} = -\frac{\hbar c^4}{1530\pi G^2} \frac{1}{M^2} \quad (8)$$

where the solution of this equation results in $M(t) = (M_i^3 - Ct)^{1/3}$ where $C = 10^{32} \text{ gr/yr}$ and minimum initial mass of black holes that survive at present time is $M_{bh}^{(initial)} > 10^{14} \text{ gr}$. Let us take a lognormal mass function of PBHs as (Carr et al. 2017; Kannike et al. 2017; Dolgov and Silk 1993)

$$\psi(M_i) = \frac{1}{\sqrt{2\pi}\sigma M_i} \exp\left[-\frac{\log^2(M_i/M_c)}{2\sigma^2}\right], \quad (9)$$

where M_c represents the peak of $\psi(M)M$ function. Black holes with the mass in the order of 10^{14} gr will lose half of their masses during the life-time of the Universe. So for the black holes within the range of $10^{14} \text{ gr} < M < 10^{15} \text{ gr}$ we would expect black holes survived from the Hawking evaporation and have one collision per 10^9 year.

We can imagine various physical processes for the collision of the low-mass PBHs with earth. The effect of this collision on earth may have geological traces as well as effects on living species on earth. During the passage of PBH from the earth (i) black holes can decelerate due to dynamical friction, also (ii) accrete matter from the interior of the earth on it. The deceleration due to dynamical friction is a consequence of gravitational interaction with the material of earth as well as the momentum transfer due to the accretion of matter on it. On the other hand, we expect the radiation due to the accretion where the overall effects could generate heat at the interior of the earth.

We define a deceleration time scale (t_d) during the passage of a PBH through the earth. For the case that it is shorter than the crossing time scale t_c of the PBH through the earth (i.e. $t_d < t_c$) the PBH will lose its energy, spiral, and sink into the center of the earth and finally could be trapped by the earth. The accretion of matter on the black hole will heat the interior of the earth and makes the black hole grow and finally, all the earth will be swallowed by the black hole. We will discuss the threshold mass of black holes that can be trapped inside the earth and calculate the rate of these events compare with the lifetime of the Universe. For the case that PBH crosses the earth and leave from the other side to space, black holes just transfer part of its energy to the earth.

We define the crossing time scale of black holes through the earth as $t_c = R_e/\bar{v}$ where the average velocity in terms of dispersion velocity from the Maxwell distribution is $\bar{v} = \sqrt{3}\sigma$. For the dispersion velocity of halo $\sigma(halo) = 220 \text{ km s}^{-1}$ and dispersion velocity of dark disk $\sigma(DD) = 38 \text{ km s}^{-1}$, the corresponding crossing time scales are $t_c(halo) = 17 \text{ s}$ and $t_c(DD) = 97 \text{ s}$, respectively. We can calculate the distribution function for the normalized transit time by defining $\hat{t} = (R_e/v)/t_c = \sqrt{3}\sigma/v$. We use the Maxwell-Boltzmann distribution for the velocity of black holes, then the distribution function for \hat{t} obtain as

$$p(\hat{t})d\hat{t} = \sqrt{\frac{6}{\pi}} \frac{1}{\hat{t}^2} \exp\left(-\frac{3}{2\hat{t}^2}\right)d\hat{t} \quad (10)$$




Figure 1. Normalized distribution function for the normalized crossing-time of black holes (i.e. $\hat{t} = t/t_c$) that cross the earth during collision.

Figure (1) shows this distribution function in terms of normalized transit time-scale. In the next section, we will compare the transit time-scale with the time-scales of dissipation processes to evaluate the impact of a black hole on the Earth.

3 DISSIPATION PROCESSES AND HEATING THE INTERIOR OF THE EARTH

The dissipation process while PBH crossing the earth happens due to two different mechanisms of dynamical friction (Chandrasekhar 1943) and accretion of the matter by the black holes. The deceleration due to the dynamical friction is a result of the interaction of matter in the medium by a black hole and is given by

$$\frac{d\mathbf{v}_{bh}}{dt} = -\frac{4\pi \ln(\Lambda) G^2 \rho_e M_{bh}}{v_{bh}^3} \left[\operatorname{erf}(X) - \frac{2X}{\sqrt{\pi}} e^{-X^2} \right] \mathbf{v}_{bh}, \quad (11)$$

where $X = v_{bh}/(\sqrt{2}\sigma_e)$ and σ_e is the dispersion velocity of particles inside the earth, ρ_e is the density of earth and $\ln \Lambda$ is the column logarithm and is given by

$$\ln \Lambda = \ln \frac{b_{max} \sigma_e^2}{G(M_{bh} + m)}, \quad (12)$$

where b_{max} is the maximum impact parameter that we use the radius of earth (i.e. $b_{max} = R_e$) and m is the mass particles inside the earth. Equation (12) simplifies to $\ln \Lambda = \ln(R_e \sigma_e^2 / (GM_{bh}))$ where substituting the numerical values results in

$$\ln \Lambda = 10.3 + \ln\left(\frac{M_{bh}}{10^{23} \text{gr}}\right). \quad (13)$$

The time-scale for deceleration of black holes while passing through the earth as a result of dynamical friction is defined by $\tau_{DF} = v_{bh}/\dot{v}_{bh}$. By taking into account that $v_{bh} \gg \sigma_e$, the time scale simplifies to

$$t_{DF} = \frac{v_{bh}^3}{4\pi G^2 M_{bh} \rho_e \ln \Lambda}, \quad (14)$$

where for the black holes from the halo and dark disk, the numerical values of this time scale is

$$t_{DF} = C \left(\frac{M_{bh}}{10^{23} \text{gr}} \right)^{-1} \left(1 + \frac{1}{10} \ln \frac{M_{bh}}{10^{23} \text{gr}} \right)^{-1} s, \quad (15)$$

where the parameter C depending on the black holes from the halo or dark disk is, $C_{halo} = 3.8 \times 10^{10}$ and $C_{DD} = 1.95 \times 10^8$, respectively.

Here we compare the dynamical friction time-scale (t_{DF}) with the crossing-time scale of the black holes through the earth (t_c). Let us define the parameter of $Y = t_{DF}/t_c$ where substituting the parameters results in

$$Y = \frac{v_{bh}^4}{A^4}. \quad (16)$$

where $A^4 = 4\pi R_e G^2 M_{bh} \rho_b \ln \Lambda$. We have two phases for Y where for $Y > 1$ dissipation is not effective and black holes can cross the earth without sinking inside the earth and for $Y < 1$ dissipation is effective. For the velocity of black holes in the halo (dN/dv_{bh}) follows a Maxwell-Boltzmann distribution,

$$\frac{dN}{dv_{bh}} dv_{bh} = \frac{4\pi v_{bh}^2}{(2\pi\sigma^2)^{3/2}} \exp\left(-\frac{v_{bh}^2}{2\sigma^2}\right) dv_{bh}, \quad (17)$$

we change the variable from v_{bh} to Y , the result is

$$\frac{dN}{dY} dY = \frac{1}{\sqrt{2\pi}} \kappa^3 Y^{-1/4} \exp\left(-\frac{\kappa^2 Y^{1/2}}{2}\right) dY, \quad (18)$$

where

$$\kappa = A/\sigma = 6.25 \times 10^{-3} \left(\frac{\sigma}{220 \text{km/s}} \right)^{-1} \left(\frac{M}{10^{23} \text{gr}} \right)^{1/4}.$$

Integrating over Y from zero to one, results in the probability of black holes sink in the earth. The numerical value of this probability is

$$P(Y < 1) = 1.3 \times 10^{-7} \left(\frac{\sigma}{220 \text{km/s}} \right)^{-3} \left(\frac{M}{10^{23} \text{gr}} \right)^{3/4}.$$

If we multiply the probability function to the total number of collisions of black holes originated from the disk and halo (equations (5) and (6)), we obtain the rate of black hole

collisions with earth that can be sinked in the earth

$$\left(\frac{dN}{dt}\right)_h(Y < 1) = 9.4 \times 10^{-16} f\left(\frac{M_{bh}}{10^{23} \text{gr}}\right)^{-1/4} \text{Gyr}^{-1}, \quad (19)$$

$$\left(\frac{dN}{dt}\right)_d(Y < 1) = 1.3 \times 10^{-13} f\left(\frac{M_{bh}}{10^{23} \text{gr}}\right)^{-1/4} \text{Gyr}^{-1}. \quad (20)$$

Since the rate of black holes that could be trapped in the earth, originated from the halo is three orders of magnitude smaller than that of the dark disk, we can ignore the contribution from the halo. For black holes with the mass of $M \sim 10^{15}$ gr, we expect the rate of black holes that can be trapped in the earth is $(dN/dt)(Y < 1) \simeq 10^{-11} \text{Gyr}^{-1}$. Taking into account the possible smallest black holes with the Planck mass $M_{bh} = 10^5$ gr, where charged black holes can survive from the Hawking evaporation (Balbinot 1986), the rate of smallest black holes that can be trapped inside the Earth during the lifetime of the Earth is almost zero. Concluding this section, we ensure that all the black holes colliding with earth will cross its interior and leave space.

4 THE BLACK HOLES INTERACTION AND ENERGY RELEASE

In the previous section, we have shown that the probability that a black hole collision with the earth and can be trapped at the interior of the Earth is almost zero. However, those black holes that cross the earth can heat the interior of the earth with various mechanisms.

In what follows we investigate the amount of energy release of black holes in addition to the amount of accretion during the passage of the black holes through the earth. The lose of kinetic energy during this passage which transfers to heat is $Q \simeq M_{bh} \dot{v}_{bh} R_e$. Using equation (11) for the deceleration of black holes due to dynamical friction and using the numerical values, the average heat production is

$$\begin{aligned} \bar{Q}_{Halo} &= 3.7 \times 10^{21} \left(\frac{v_{bh}}{220 \text{km s}^{-1}}\right)^{-2} \left(\frac{M_{bh}}{10^{23} \text{gr}}\right)^2 J, \\ \bar{Q}_{disk} &= 1.2 \times 10^{24} \left(\frac{v_{bh}}{38 \text{km s}^{-1}}\right)^{-2} \left(\frac{M_{bh}}{10^{23} \text{gr}}\right)^2 J, \end{aligned} \quad (21)$$

where depending on the velocity of black holes from the halo or dark disk the energy loss would be different. For the mass range of $M_{bh} < 10^{15}$ gr, the number of interactions of primordial black holes with the earth is larger than one event per Gyr where for this range of masses the heat production due to the dynamical friction is $Q_{halo} < 10^5 J$ and $Q_{disk} < 10^8 J$.

The other source of heat production is the accretion of matter into the black holes while passing through the Earth. The maximum amount of energy released from the accreting matter around the black hole is given by the Eddington limit (Eddington 1916), where equating the radiation pressure with the gravitation pressure results in

$$L_{Edd} = \frac{4\pi G M_{bh} m_p}{\sigma_{Th}}, \quad (22)$$

where σ_{Th} is Thompson cross section for photons and electrons. Using the numerical values, the Eddington limit is

$$L_{Edd} = 6.5 \times 10^{27} \left(\frac{M_{bh}}{10^{23} \text{gr}}\right) \text{erg/s}, \quad (23)$$

For a black hole with the mass in the order of $M_{bh} = 10^{15}$ gr,

using the crossing time-scale through the earth (i.e. $t_c = R_e/\bar{v}$), the corresponding energy from the radiation, $Q = L_{Edd} t_c$ is

$$Q_{Halo}^{(rad)} = 2.2 \times 10^{22} \left(\frac{M_{bh}}{10^{23} \text{gr}}\right) J, \quad (24)$$

$$Q_{disk}^{(rad)} = 1.3 \times 10^{23} \left(\frac{M_{bh}}{10^{23} \text{gr}}\right) J, \quad (25)$$

where for the mass range of $M_{bh} < 10^{15}$ gr, the heat production due to accretion in the black holes sourced from halo and disk would be $Q_{halo} < 10^{14} J$ and $Q_{disk} < 10^{15} J$. Comparing the energy release from the accretion with the dissipation of energy due to dynamical friction in equation (21), we can neglect the heat production due to the dynamical friction.

In order to have an idea about the amount of energy that is released from the black hole impact, we compare it with the energy released from the asteroid impacts the earth. For an asteroid impact with the crater size of D on earth, the energy release is about $E \simeq 10^{16} (D/\text{km})^3 J$ (Dence et al 1977). On the other hand, the rate of the impacts is $\Gamma_{asteroid} = 6.64 \times 10^4 (D/\text{km})^{2.55} \text{Gyr}^{-1}$ (Hergarten & Kenkmann 2015) where for an asteroid with the crater size of km, the energy release is almost equal to the impact of a black hole with the mass of $M_{bh} \simeq 10^{15}$ gr. However, we note that the rate of an asteroid impact is four orders of magnitude larger than that of PBH impact. The impact of the asteroid and a black hole with the earth also has two different physical features. While an asteroid during the entrance to the atmosphere burn and has an explosive impact on the surface of the earth, we expect that a black hole has different features in the impact.

For a black hole with the size of Schwarzschild radius, R_s and the relative velocity of v_{bh} with respect to the earth, the cross-section that materials accrete into the horizon of a black hole, similar to equation (1) is

$$\sigma_{acc} = \pi R_s^2 \left(1 + \frac{2GM_{bh}}{R_e v_{bh}^2}\right), \quad (26)$$

where using the numerical values results in

$$\sigma_{acc} = 1.6 \times 10^{-7} \left(\frac{M}{10^{23} \text{gr}}\right)^2 \text{m}^2. \quad (27)$$

This accretion of matter results in radiation which is given by equations (24) and (25).

Since the black holes are moving through to the earth, the radiating energy around the Schwarzschild radius transfer away from the trajectory of black holes across a cylinder. This radiation can increase the temperature of the earth in the crust around the cylinder with the center of the trajectory of black holes by the amount of $Q = mC\Delta T$ where C is the heat capacity of rocks. Assuming $T = 10^3 \text{C}$ for the melting of rocks. The total mass of the crust that will be melted during this heating would be $m = Q/(C\Delta T)$, where using the heat-capacity of rock $C = 2 \times 10^3 J/\text{kgK}$ and $\Delta T = 10^3$ as the temperature for melting rock, the amount of rocks that can be melted due to heat production by the accretion from equation (24) and (25) is

$$m_{Halo}^{(melt)} = 1.1 \times 10^{16} \left(\frac{M_{bh}}{10^{23} \text{gr}}\right) \text{kg}, \quad (28)$$

$$m_{disk}^{(melt)} = 5.9 \times 10^{16} \left(\frac{M_{bh}}{10^{23} \text{gr}}\right) \text{kg}. \quad (29)$$

From equation (7), for the mass range of $M_{bh} < 10^{15}$ gr, the probability of black holes collision is larger than one collision per Giga year where the corresponding melting mass is $m^{(melt)} \simeq 5 \times 10^8$ kg.

We assume a typical trajectory of the black holes equal to the radius of earth and density of interior matter in the order of $\rho = 3\text{gr/cm}^3$, a cylinder that can be melted due to the passage of black holes has a radius of $R \sim 10$ cm. The geological investigations of the deep drilling program on earth crust show a rapid temperature gradient in the crust (Emmerman & Lauterjung 1997). We expect that such a long melting hole due to the passage of black holes will rapidly dissipate the energy and earth crust will be recovered where just in the upper parts of crust we may distinguish an unusual defect of the melted rocks along a straight line.

5 CONCLUSION

In this work, we investigated the scenario of collision of primordial black holes with earth. We calculated the rate of impact assuming that hundred percent of dark matter is made of these objects. Assuming a Dirac-Delta mass function, for the black holes that survived from the Hawking radiation, black holes with the mass $M < 10^{15}$ gr have the chance of more than one impact per billion years with Earth. Since the velocity of black holes in the halo and the dark disk is high, the dissipation process inside the earth is not effective to decelerate the black holes and black holes do not sink inside the earth where in this case black holes could heat the interior of the earth and finally swallow the whole mass of earth. The number of this event for $M \simeq 10^{15}$ gr is about 10^{-11}Gyr^{-1} . This calculation assures that probability of primordial black holes being trapped inside the earth is almost zero.

We also calculated the energy released by a black hole collision with earth and compare it with the impact of asteroids on the earth. We have shown that the accretion process by the black holes is the dominant process for energy release. The amount of energy from this collision is comparable with a kilometer size asteroid where it happens four orders of magnitude more frequently than a black hole collision. We conclude that the likelihood of the dangerous impact of primordial black holes with the earth is very low.

ACKNOWLEDGMENTS

I would like to thank Shant Baghran for his useful comments. This research was supported by Sharif University of Technology's Office of Vice President for Research under Grant No. G950214.

REFERENCES

Abbott, B.P. et al., 2016, Phys. Rev. Lett. 116 (6): 061102.
 Abbott, R. et al. 2017, Astrophys. J. Lett. 851, L35
 Abbott B. P. et al. 2019, Physical Review X. 9 (3), 031040
 Abbott, R. et al., 2020, Phys. Rev. Lett. 125, 101102
 Balbinot, R., 1986, Class. Quant. Grav. 3, L1
 Binney J. and Tremaine, S., 2008, Galactic Dynamics: Second Edition, Princeton University Press

Buch, J., Leung J. S. C & Fan, J. 2019, JCAP, 026, 2019
 Carr, B., Raidal, M., Tenkanen, T., Vaskonen, V. and Veermäe, H., 2017, Phys. Rev. D 96, 023514
 Carr, B. ; Kühnel, F., 2020, Annual Review of Nuclear and Particle Science, 70, 355
 Chandrasekhar, S., 1943, Astrophysical Journal **97**, 255
 Dence, M. C., Grieve, R. A. F. & Robertson, P. B. Terrestrial impact structures: principal characteristics and energy consideration. In: Roddy, D.J. Pepin, R.O., and Merrill, R.B. (eds). Impact and explosion cratering. Pergamon Press, Oxford, UK, pp 247-275 (1977).
 Dolgov A. and Silk, J., 1993, Phys. Rev. D 47, 4244
 Eddington, A. S. 1916, MNRAS, 77: 16
 Emmerman R. & Lauterjung, J., 1997, Journal of Geophysical Research: Solid Earth. 102 (B8), 18179
 Green, A. M., 2016, Phys. Rev. D 94, 063530
 Hawking, S. W., 1974, Nature. 248 (5443): 30
 Hergarten S. & Kenkmann, T., 2015, Earth and Planetary Science Letters, 425, 187
 Kannike, K., Marzola, L., Raidal M. , and Veermäe, H., 2017 , JCAP 09, 020
 Khalouei, E., Ghodsi, H., Rahvar, S. & Abedi, J., 2021, Phys. Rev. D 103, 084001
 Kühnel F. and Freese, K., 2017, Phys. Rev. D 95, 083508
 Moniez, M. et al. A&A **604**, A124 (2017)
 Niemeyer J. C. and Jedamzik, K., 1998, Phys. Rev. Lett. 80, 5481
 Penrose, R., 1965, Physical Review Letters. 14 (3), 57.
 Samimi, J. et al, 1979, Nature 278, 434.
 Sasaki, M., Suyama, T., Tanaki, T., 2016, Physical Review Letters. 117, 061101
 Shipman, H. L., Yu, Z., Du, Y.W., 1975, Astrophysical Letters, 16 (1), 9