

# Helical magnetogenesis with reheating phase from higher curvature coupling and baryogenesis

Kazuharu Bamba,<sup>1</sup> Sergei D. Odintsov,<sup>2,3</sup> Tanmoy Paul<sup>4,5</sup> Debaprasad Maity<sup>6</sup>

<sup>1)</sup> *Division of Human Support System, Faculty of Symbiotic Systems Science, Fukushima University, Fukushima 960-1296, Japan*

<sup>2)</sup> *Institute of Space Sciences (IEEC-CSIC) C. Can Magrans s/n 08193 Bellaterra (Barcelona), Spain*

<sup>3)</sup> *ICREA, Passeig Luis Companys 23, 08010 Barcelona, Spain*

<sup>4)</sup> *Department of Physics, Chandernagore College, Hooghly - 712 136, India.*

<sup>5)</sup> *International Laboratory for Theoretical Cosmology, TUSUR, 634050 Tomsk, Russia.*

<sup>6)</sup> *Department of Physics, Indian Institute of Technology Guwahati, Assam 781039, India*

We investigate the generation of helical magnetic fields and address the baryon asymmetry of the universe from an inflationary magnetogenesis scenario, in which the conformal and parity symmetries of the electromagnetic field are broken through its coupling the Ricci scalar and the Gauss-Bonnet invariant via the dual field tensor, so that the generated magnetic field can have a helical nature. It is demonstrated that if a reheating phase with non-zero e-fold numbers following inflation is taken into account, the energy density of the magnetic fields as well as their helicity spectrum evolve differently compared to those for the instantaneous reheating case. As a result, it is shown that in presence of the reheating phase, both the field strength of the generated magnetic fields and the resultant baryon asymmetry of the universe can be compatible with the observations. Furthermore, we find a viable constraint on the reheating equation of state parameter  $\omega_{\text{eff}}$  from the data of the cosmic microwave background radiation as  $0.284 \lesssim \omega_{\text{eff}} \lesssim 0.2975$ , which is consistent with a negligible Schwinger backreaction.

## I. INTRODUCTION

Large scale magnetic field and baryon asymmetry of the universe are some of the important unresolved issues in the standard model of cosmology and particle physics. Despite of large attempts being made to understand those problems, it is still an active area of research. Although both the issues are seemingly disconnected, endeavour towards unified theoretical description would be extremely interesting. In order to proceed, one needs to go beyond standard model of cosmology and particle physics. Various astrophysical observations indicate that the universe is magnetized over a wide range of length scales. The magnetic fields have been detected in galaxies, galaxy clusters and even in intergalactic voids [1–6]. On the other hand, it is observationally known that the universe carries a net excess of baryons over antibaryons [7–17]. Such baryon asymmetry of the universe can be realized from the Sakharov’s three conditions [18]: (1) baryon number non-conservation, (2) Charge(C) and Charge parity(CP) violation, and (3) departure from thermal equilibrium.

Our current understanding on the origin of large scale magnetic fields are broadly classified into: (1) the astrophysical origin associated with dynamo mechanism [19–21] and (2) the primordial origin where the magnetic field is generated from the primordial quantum fluctuations during the inflationary epoch [22–55], or during the bouncing phase of the universe [56–61]. Among all the models proposed so far, the inflationary magnetogenesis earned significant attention due to the fact that the inflation concomitantly solves the well known flatness and horizon problems and predicts almost scale invariant curvature perturbation which is in agreement with the Planck data [62–67] to a great precision. Due to this background curvature fluctuation the standard Maxwell’s action may produce magnetic field out of quantum vacuum, but its magnitude will naturally be very feeble and lie far below the observational constraints. The reason behind this feeble production of magnetic field is the conformal invariant nature of the standard electromagnetic theory and conformal flat nature of the background spacetime. Therefore, simplest way to enhance the strength of the magnetic field is to introduce the conformal non-invariant coupling in the electromagnetic sector. Several inflationary magnetogenesis models have been proposed so far, where the conformal non-invariant and non-minimal electromagnetic coupling has been introduced through the scalar field such as inflaton, axion and also through higher curvature terms [22–36, 38–55, 68–73]. Depending upon the basic physical requirements, those couplings can be classified into: parity invariant and parity violating magnetogenesis scenario. Interestingly, parity violating and non-conformally coupled electromagnetic field can generate helical magnetic field which has been shown to play crucial role in generating baryon asymmetry in the universe. However the construction of inflationary magnetogenesis is riddled with some severe theoretical problems, such as, the backreaction problem and the strong coupling problem respectively. The backreaction problem arises when the electromagnetic field has considerable backreaction on the background inflationary spacetime, in particular, when the electromagnetic energy density becomes larger or comparable to the background energy density. And, the strong coupling problem occurs when the effective electric charge becomes

strong during inflation and makes the perturbative calculation unreliable. Therefore in order to get viable inflationary magnetogenesis models, the backreaction and the strong coupling problem should be resolved in the model (one may see [22, 28, 51, 73, 74]). Apart from the inflationary set-up, the bouncing magnetogenesis models, where the magnetic field generation occurs from a non-singular bounce scenario, has also been proposed in the literature [56–61]. In most of the bouncing magnetogenesis models, the electromagnetic field couples with the bouncer (who drives the bounce), which in turn spoils the conformal symmetry of the electromagnetic action. Similar to inflation, the bounce cosmology is also able to generate an observationally compatible primordial scalar and tensor power spectrum. Here we would like to mention that most of the bounce models are plagued with some severe problems, like, the BKL instability associated with the growth of background anisotropy during the contracting stage of the universe, the instability of scalar and tensor perturbations around the bounce [75–80]. However such problems can be resolved, to some extent, in some modified gravity theories including the higher curvature or extra dimensional braneworld model [77, 81–86]. However, in this paper our focus will be on the inflationary magnetogenesis.

Coming back to the relation between magnetogenesis and baryogenesis; it is well known that the presence of magnetic field leads to: (a) the dynamics of plasma to be out-of-thermal-equilibrium as in thermal equilibrium the photon distribution must be thermal which is incompatible with existence of long range magnetic field, (b) violation of time reversal (T) or in other words CP symmetry as well as C symmetry, and (c) being a vector quantity, the presence of magnetic field breaks the rotational invariance or SO(3) symmetry [87]. Those are precisely the Sakharov’s conditions for the successful generation of baryogenesis. Therefore, once we assume the genesis of magnetic field in the early universe that can act as a potential source for the production of baryon asymmetry in the universe. Motivated by this, Giovannini and Shaposhnikov [9, 10] first investigated the generation of baryon asymmetry through the abelian anomaly in the form of  $U(1)$  Chern-Simons term which assumes non-zero vacuum expectation value due to helical magnetic field. This in turn produces net baryon asymmetry though the non-conservation of baryon number current. Related proposals are investigated in [88–91]. Hence, it is important to have non-zero helicity of the magnetic field which is varying with the cosmological evolution. Axion- electrodynamics has been widely studied in this regard. Non-trivial dynamics of axion field produces helical magnetic field which finally leads to the baryon asymmetry of the universe [48]. Recently, the baryogenesis has been addressed from the production of helical magnetic fields, where the electromagnetic field gets coupled with the background Riemann tensor by the dual field tensor [92]. Such coupling between the electromagnetic field and the Riemann tensor gives rise to helical nature of magnetic field, which in turn leads to a net baryonic number density as of the order  $n_B/s_0 \sim 10^{-10}$ , that is consistent with the cosmic microwave background (CMB) observations (where  $n_B$  is the net baryonic number density and  $s_0$  is the entropy density of the present universe). From a different perspective, the article [93] discussed the baryogenesis from magnetic fields, however from MHD amplification.

In the present work, we propose a helical magnetogenesis scenario from inflationary set-up and address the baryon asymmetry of the universe (BAU), from higher curvature coupling of the electromagnetic (EM) field. In particular, the EM field couples with the background higher curvature term(s), namely with the Ricci scalar and the Gauss-Bonnet invariant, via the dual field tensor. Such non-minimal coupling in the EM action breaks the conformal and the parity symmetry, however preserves the  $U(1)$  invariance, of the EM field, which in turn leads to the gauge field production from the primordial Bunch-Davies vacuum. Moreover the positive and negative helicity modes of the EM field get different amplitudes along their cosmological evolution, and thus give rise to helical nature of the magnetic field. The Chern-Simons number stored in the helical magnetic fields is directly connected to the net baryonic number density of the universe. In regard to the background spacetime, it is considered to be the scalar-Einstein-Gauss-Bonnet gravity theory which is known to provide viable inflationary scenario for suitable choices of scalar field potential and Gauss-Bonnet coupling function [94–104]. Owing to the fact that the non-minimal coupling of the EM field occurs via the dual field tensor, the kinetic term of the EM field remains canonical even in presence of such non-minimal coupling, and thus the strong coupling problem is naturally resolved in the present magnetogenesis model. In such scenario, we explore the cosmological evolution of the EM field and consequently determine the net baryon density of the universe. Most of the earlier magnetogenesis scenarios did not explore the effect of reheating phase until recently. Reheating can be important in this regard has recently been realized in [43], and further explored in [36–38]. During the expansion, the universe enters into a reheating phase after inflation and depending on the reheating dynamics, we consider two different cases: (1) the instantaneous reheating case where the universe suddenly jumps to the radiation dominated epoch after the end of inflation, and thus the reheating era has zero e-fold number, and (2) the case where the reheating era possesses a non-zero e-fold number, in particular, we consider the Kamionkowski reheating mechanism that parametrize the reheating dynamics by a non-zero e-fold number, a constant equation of state parameter and a given reheating temperature [105] (see also [106–115]). These two cases make qualitative differences in the evolution of the helicity power spectrum. Specifically, the presence of the reheating phase with non-zero e-fold number leads to an enhancement in the magnetic strength and consequently in the helicity spectrum at present universe, compared to that of in the instantaneous reheating case. Due to such enhancement, the theoretical predictions of net baryonic density seems to be consistent with the observational constraints, and thus the model provides a natural explanation

of BAU. Here we would like to mention that the baryogenesis from helical magnetic fields with curvature couplings have been proposed earlier, however in quite different contexts [92]. It may be noted that in our present analysis, we include the higher curvature Gauss-Bonnet coupling in the set-up and also discuss the possible effects of the reheating phase in the production of helical magnetic field and consequently in the baryogenesis, which makes the present model essentially different from earlier ones.

The paper is organized as follows: in Sec. II, we describe the model; in Sec. III, we give the general expressions for the EM power spectra and the helicity spectrum in the present; Sec. IV is reserved for the solution of the vector potential, the electromagnetic energy density and the helicity spectrum during inflation. The corresponding calculations during the reheating phase are carried out in Sec. V and Sec. VI, which correspond to the cases of instantaneous reheating and a Kamionkowski like reheating model, respectively. The Schwinger backreaction issue and the necessary constraints are discussed in Sec. VII. The paper ends with some conclusions.

## II. SCALAR GAUSS-BONNET GRAVITY WITH NON-MINIMALLY COUPLED ELECTROMAGNETIC FIELD

The action is,

$$S = S_{grav} + S_{em}^{(can)} + S_{CB} \quad , \quad (1)$$

where  $S_{grav}$ , namely the gravitational action, determines the background evolution and a non-minimally coupled electromagnetic (EM) field propagates over the background spacetime, where  $S_{em}^{(can)}$  and  $S_{CB}$  represent the canonical kinetic term and the non-minimal coupling function of the EM field respectively. The EM field couples with the higher curvature of the background spacetime, which actually provides the non-minimal coupling term in the action. The  $S_{grav}$  has the following form,

$$S_{grav} = \int d^4x \sqrt{-g} \left[ \frac{R}{2\kappa^2} - \frac{1}{2} \partial_\mu \Phi \partial_\nu \Phi - V(\Phi) - \xi(\Phi) \mathcal{G} \right] \quad , \quad (2)$$

where  $\kappa^2 = 8\pi G$  ( $G$  is the Newton's constant). Note that the theory (2) represents string-inspired Gauss-Bonnet gravity, where  $R$  is the Ricci scalar,  $\mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta}$  symbolizes the Gauss-Bonnet (GB) invariant and  $\xi(\phi)$  is the Gauss-Bonnet coupling with  $\phi$  being the scalar field under consideration. Moreover  $V(\phi)$  denotes the potential term of the scalar field. The coupling between the Gauss-Bonnet curvature and the scalar field, namely  $\xi(\phi)$ , ensures the non-topological character of the Gauss-Bonnet term in the above action and thus has non-trivial contributions in the background evolution. The canonical kinetic term of the EM field is given by,

$$S_{em}^{(can)} = \int d^4x \sqrt{-g} \left[ -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right] \quad , \quad (3)$$

where  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$  is the EM field tensor and  $A_\mu$  is the corresponding field. In regard to the  $S_{CB}$ , as mentioned earlier, the EM field non-minimally couples with the spacetime curvature of the background spacetime, in particular,

$$S_{CB} = \int d^4x \sqrt{-g} f(R, \mathcal{G}) \left[ \lambda F_{\mu\nu} \tilde{F}^{\mu\nu} \right] \quad , \quad (4)$$

where  $\tilde{F}^{\mu\nu} = \epsilon^{\mu\nu\alpha\beta} F_{\alpha\beta}$  with  $\epsilon^{\mu\nu\alpha\beta}$  is the four dimensional Levi-Civita tensor and defined by  $\epsilon^{\mu\nu\alpha\beta} = \frac{1}{\sqrt{-g}} [\mu\nu\alpha\beta]$ , the  $[\mu\nu\alpha\beta]$  denotes the completely antisymmetric permutation having  $[0123] = 1$ . The EM coupling function  $f(R, \mathcal{G})$  is considered to depend on the background Ricci scalar and Gauss-Bonnet curvature respectively, particularly the form of  $f(R, \mathcal{G})$  is given by,

$$f(R, \mathcal{G}) = \kappa^{2q} (R^q + \mathcal{G}^{q/2}) \quad , \quad (5)$$

with  $q$  being a parameter of the model which is considered to be less than unity. This is a viable consideration; otherwise for  $q > 1$ , the conformal breaking function  $f(R, \mathcal{G})$  becomes higher dimensional operator (higher than six) and hence gets more suppressed by the Planck mass. Moreover, later we will show that such consideration (i.e.  $q < 1$ ) is indeed compatible with the CMB observations of present magnetic strength. It may be observed that the EM field gets the non-minimal coupling via the dual field tensor, in particular by  $F_{\mu\nu} \tilde{F}^{\mu\nu}$  controlled by the dimensionless parameter  $\lambda$ , which acts as a parity violating agent in the EM field action and ensures the helical nature of the magnetic field, as we will see later. Here it deserves mentioning that the  $S_{CB}$  spoils the conformal

invariance of the EM field, although preserves the U(1) invariance of the gauge field. The broken conformal symmetry is the key ingredient in the magnetogenesis scenario, otherwise the EM field energy density redshifts as  $1/a^4$  with the cosmological expansion of the universe and results to a very feeble magnetic strength at present epoch – not compatible with the CMB observation at all. During the early universe when the spacetime curvature is high, the conformal breaking coupling leads to a non-trivial contribution to the EM field equations, however at late times (in particular at the end of inflation) the  $S_{CB}$  will not contribute and consequently the EM field behaves as standard Maxwellian theory. Due to the presence of  $F_{\mu\nu}\tilde{F}^{\mu\nu}$  ( $=\epsilon^{\mu\nu\alpha\beta}F_{\mu\nu}F_{\alpha\beta}$ ) in Eq.(4), the conformal breaking coupling does not contain the term like  $(A'_i)^2$  (where the prime denotes the derivative with respect to the conformal time) and thus the kinetic term of the EM field, i.e  $(A'_i)^2$ , comes only through the  $S_{em}^{(can)}$  given in Eq.(3). This in turn indicates that the EM kinetic term remains canonical in the present magnetogenesis model where the EM field gets coupled with the spacetime curvature through the corresponding dual field tensor. As a consequence, the model is free from the strong coupling problem. Moreover we will show later that the EM field energy density has negligible backreaction on the background inflationary FRW spacetime, which ensures the resolution of the backreaction issue in the present context.

In regard to the background spacetime, the scalar-Einstein-GB gravity theory, i.e the  $S_{grav}$ , is known to provide viable inflationary scenario for suitable choices of scalar field potential, with the observable quantities like the spectral index, tensor-to-scalar ratio being compatible with the Planck 2018 constraints [7]. In particular, the authors of [99] showed that for a quadratic power law form of the scalar potential, the scalar-Einstein-Gauss-Bonnet theory leads to a de-Sitter expansion of the universe, while for an exponential scalar potential, the universe expands with a decelerating stage; thereby smoothly interpolating these two eras yields a successful inflation with a natural exit mechanism in the context of scalar-Einstein-GB theory. Moreover the presence of the scalar potential ensures the stability of the scalar and tensor perturbations in GB theory. Here we would like to mention that, unlike to scalar-tensor or F(R) theory, the speed of the tensor perturbation ( $c_T^2$ ) in the Gauss-Bonnet gravity theory is different than unity and the deviation of  $c_T^2$  from the value 1 is entirely controlled by the GB coupling function  $\xi(\phi)$ . The condition  $c_T^2 \neq 1$  indicates that the gravitational wave speed is different compared to the speed of light in vacuum (which is unity in natural units) and thus is not compatible with the observations of the event GW170817. However there exists certain class of GB coupling functions for which the  $c_T^2$  becomes unity in the scalar-Einstein-GB theory and hence the GB model gets compatible with the GW170817. The early universe phenomenology in the scalar-Einstein-GB theory compatible with GW170817 are discussed in [116–119].

In such magnetogenesis set-up, we aim to determine the magnetic strength at the present epoch and consequently the baryon asymmetry of the universe from the helical nature of the magnetic field. Varying the action (1) with respect to  $A_\mu$ , we get the field equation for the EM field as,

$$\partial_\alpha [\sqrt{-g} \{g^{\mu\alpha} g^{\nu\beta} F_{\mu\nu} - 8\lambda f(R, \mathcal{G}) \epsilon^{\mu\nu\alpha\beta} \partial_\mu A_\nu\}] = 0 \quad , \quad (6)$$

where it may be noticed that the conformal breaking coupling modifies the EM field equation compared to the standard Maxwell's equation. A spatially flat FRW metric ansatz will fulfill our purpose in the present context, i.e

$$ds^2 = a^2(\eta) [-d\eta^2 + d\vec{x}^2] \quad , \quad (7)$$

where  $a(\eta)$  is the scale factor of the universe with  $\eta$  being the conformal time and related to the cosmic time ( $t$ ) by  $d\eta = \frac{dt}{a(\eta)}$ . For this metric ansatz, the temporal and spatial components of the EM field Eq.(6) become,

$$\partial^i [\partial_i A_0 - \partial_0 A_i] = 0 \quad (8)$$

and

$$\delta^{ij} \partial_0 [h^2(R, G) F_{0j}] - \delta^{ij} \partial^l [h^2(R, G) F_{lj}] = 0 \quad (9)$$

respectively. Due to the antisymmetric nature of  $\epsilon^{\mu\nu\alpha\beta}$ , the temporal component Eq.(8) seems to free from the effect of the coupling function  $f(R, \mathcal{G})$ . Here we consider the Coulomb gauge condition, in particular  $\partial_i A^i = 0$ , due to which, Eq.(8) leads to the solution as  $A_0 = 0$ . Therefore in effect of  $\partial_i A^i = 0$  and  $A_0 = 0$ , the spatial component Eq.(9) of the EM field takes the following form,

$$A_i''(\eta, \vec{x}) - \partial_t \partial^l A_i - 8\lambda f'(R, \mathcal{G}) \epsilon_{ijk} \partial_j A_k = 0 \quad , \quad (10)$$

where  $\epsilon_{ijk} = [0ijk]$ . As mentioned earlier, the scalar-Einstein-GB theory (i.e the  $S_{grav}$  we considered in Eq.(2)) is able to drive successful inflationary scenario during the early universe for suitable choices of  $V(\phi)$  and  $\xi(\phi)$  respectively.

Keeping this in mind, we consider a quasi-de-Sitter inflationary background spacetime in the present context, where the scale factor has the following form,

$$a(\eta) = \left( \frac{-\eta}{\eta_0} \right)^{\beta+1} \quad \text{with} \quad \beta = -2 - \epsilon = -3 + \frac{\mathcal{H}'}{\mathcal{H}^2} . \quad (11)$$

Here, and also in the subsequent calculation, a prime denotes  $\frac{d}{d\eta}$ ,  $\mathcal{H}$  is the conformal Hubble parameter and defined by  $\mathcal{H} = a'/a$ . Moreover  $\epsilon$  is known as the slow roll parameter having the expression  $\epsilon = -\frac{\mathcal{H}'}{\mathcal{H}^2} + 1$ . Eq.(11) evidents that for  $\epsilon = 0$ , the scale factor becomes  $a(\eta) \propto (-\eta)^{-1}$  which results to the de-Sitter evolution of the universe. The scale factor (11) immediately leads to the conformal Hubble parameter, Ricci scalar and the Gauss-Bonnet invariant as,

$$\mathcal{H} = \frac{\beta + 1}{\eta} \quad (12)$$

and

$$\begin{aligned} R &= \frac{6}{a^2} (\mathcal{H}' + \mathcal{H}^2) = \frac{6\beta(\beta + 1)}{\eta_0^2} \left( \frac{-\eta}{\eta_0} \right)^{2\epsilon} \\ \mathcal{G} &= \frac{24}{a^4} \mathcal{H}^2 \mathcal{H}' = -\frac{24(\beta + 1)^3}{\eta_0^4} \left( \frac{-\eta}{\eta_0} \right)^{4\epsilon} , \end{aligned} \quad (13)$$

respectively. The cosmic Hubble parameter (defined by  $H = \dot{a}/a$ , where an overdot represents  $\frac{d}{dt}$ ) is related to the conformal Hubble parameter as  $H = \frac{1}{a} \mathcal{H}$  and thus we can express  $H$  as,

$$H = \frac{1}{\eta_0} \exp \left( -\frac{\epsilon N}{1 + \epsilon} \right), \quad (14)$$

in terms of the e-folding number (symbolized by  $N$ ). The e-folding number up-to the time  $\eta$  is defined as  $N = \int^\eta aH d\eta$ , where  $N = 0$  refers the instance of the beginning of inflation which, in the present context, we consider to happen when the CMB scale mode crosses the Hubble horizon. Therefore Eq.(14) indicates that at the beginning of inflation, the cosmic Hubble parameter acquires  $H = 1/\eta_0$ , we will denote the initial cosmic Hubble parameter by  $H_0$  in the later calculations, i.e  $H_0 = \eta_0^{-1}$ . Plugging the above expressions of  $R$  and  $\mathcal{G}$  into Eq.(5) and a little bit of simplification yield the form of  $f(R, \mathcal{G})$  as,

$$f(R, \mathcal{G}) = \kappa^{2q} \left\{ \frac{[6\beta(\beta + 1)]^q + [-24(\beta + 1)^3]^{q/2}}{\eta_0^{2q}} \right\} \left( \frac{-\eta}{\eta_0} \right)^{2\epsilon q} . \quad (15)$$

It may be observed from the above expression that the condition  $\epsilon = 0$  leads to the function  $f(R, \mathcal{G})$  as a constant, for which the EM field Eq.(10) becomes similar to the standard Maxwell's equation. This is however expected because for constant  $f(R, \mathcal{G})$ , the conformal breaking term  $\mathcal{L}_{CB} = \sqrt{-g} f(R, \mathcal{G}) \epsilon^{\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta}$  present in the action becomes a total surface term and thus gives no contributions in the field equations. As a result, the EM action preserves the conformal symmetry and thus the EM energy density decays by  $1/a^4$  with the cosmological expansion of the universe, which in turn results to a very feeble magnetic amplitude at the current epoch. Thus the present model where the EM field couples with the background spacetime curvature via the term like  $\sim \sqrt{-g} f(R, \mathcal{G}) \epsilon^{\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta}$ , requires  $\epsilon \neq 0$  in order to break the conformal invariance of the EM field. Hence in order to generate sufficient strength of magnetic field and consequently the baryon asymmetry of the present universe in such higher curvature helical magnetogenesis set-up, we consider the background inflationary scenario as a quasi-de-Sitter one, in which case  $\epsilon \neq 0$  and  $\epsilon < 1$ , in the subsequent calculations.

### III. ELECTRIC, MAGNETIC AND HELICITY POWER SPECTRA

In this section we aim to calculate the energy-momentum tensor of the EM field and subsequently we will determine the individual energy density of electric and magnetic field. In this regard, it may be mentioned that the electric and magnetic field are frame dependent, and for this purpose, here we consider the *comoving observer*, in which case the four velocity is given by  $u^\mu = (a^{-1}(\eta), 0, 0, 0)$  in the  $(\eta, \vec{x})$  coordinate system. The calculation of electric, magnetic

energy density at time  $\eta$  requires the energy-momentum tensor and the state (at  $\eta$ ) of the EM field respectively. We will consider that the EM field starts from the Bunch-Davies vacuum state at distant past (i.e the “no-particle” state), which is indeed compatible with the equation of motion, as we will show in the later section. Moreover, since we work in the Heisenberg picture, the state of the field remains fixed over time; however due to the interaction between the EM field and the background FRW spacetime, the particle production occurs and thus the EM field does not remain in *vacuum* at later time. From action (1), we determine the energy-momentum tensor of the EM field as,

$$\begin{aligned} T_{\alpha\beta} &= -\frac{2}{\sqrt{-g}} \frac{\delta}{\delta g^{\alpha\beta}} \left[ \sqrt{-g} \left\{ -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \lambda f(R, \mathcal{G}) \epsilon^{\mu\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta} \right\} \right] \\ &= -\frac{1}{4} \{ g_{\alpha\beta} F_{\mu\nu} F^{\mu\nu} - 4g^{\mu\nu} F_{\mu\alpha} F_{\nu\beta} \} + \frac{2\lambda}{\sqrt{-g}} [\mu\nu\rho\sigma] F_{\mu\nu} F_{\rho\sigma} \left( \frac{\delta f(R, \mathcal{G})}{\delta g^{\alpha\beta}} \right) , \end{aligned} \quad (16)$$

where we use  $\frac{\delta\sqrt{-g}}{\delta g^{\alpha\beta}} = -\frac{1}{2}\sqrt{-g}g_{\alpha\beta}$  and  $[\nu\rho\sigma]$  represents the antisymmetric permutation. The energy density of the EM field can be obtained from the (0,0) component of the above  $T_{\alpha\beta}$ , in particular the energy density in FRW spacetime is given by  $T_0^0 = -\frac{1}{a^2}T_{00}$ . Eq.(16) leads to,

$$T_0^0 = -\frac{1}{2a^4} (A'_i)^2 - \frac{1}{4a^4} F_{ij} F_{ij} - \frac{8\lambda}{a^6} \epsilon_{ijk} A'_i F_{jk} \left( \frac{\delta f(R, \mathcal{G})}{\delta g^{00}} \right) . \quad (17)$$

The quantity  $\frac{\delta f}{\delta g^{00}}$  is calculated as,

$$\frac{\delta f(R, \mathcal{G})}{\delta g^{00}} = \frac{\partial f}{\partial R} \frac{\delta R}{\delta g^{00}} + \frac{\partial f}{\partial \mathcal{G}} \frac{\delta \mathcal{G}}{\delta g^{00}} = -3q\kappa^{2q}\mathcal{H}' \left[ R^{q-1} + \frac{2\mathcal{H}^2}{a^2} \mathcal{G}^{\frac{q}{2}-1} \right] , \quad (18)$$

where in the second equality, we use the explicit form of  $f(R, \mathcal{G})$ , i.e  $f(R, \mathcal{G}) = \kappa^{2q} (R^q + \mathcal{G}^{q/2})$ . Moreover using the expressions of  $\mathcal{H}$ ,  $R$  and  $\mathcal{G}$  (from Eqs.(12) and (13)), we simplify Eq.(18) and finally lands to,

$$\frac{1}{a^2} \left( \frac{\delta f(R, \mathcal{G})}{\delta g^{00}} \right) = -\frac{q}{4} \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta+1)]^q + [-24(\beta+1)^3]^{q/2} \right\} \left( \frac{-\eta}{\eta_0} \right)^{2\epsilon q} .$$

Therefore the energy density of EM field from Eq.(17) turns out to be,

$$T_0^0 = -\frac{1}{2a^4} (A'_i)^2 - \frac{1}{4a^4} F_{ij} F_{ij} + \frac{2\lambda q}{a^4} \epsilon_{ijk} A'_i F_{jk} \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta+1)]^q + [-24(\beta+1)^3]^{q/2} \right\} \left( \frac{-\eta}{\eta_0} \right)^{2\epsilon q} . \quad (19)$$

We are now in a position to separate the energy density for electric and magnetic fields and for this purpose, one needs to recall that we consider the comoving observer as the required frame in the present context. As a consequence, the temporal derivative of  $A_i$  contributes to the electric energy density, while the magnetic energy density comes through the spatial derivative of  $A_i$ . Therefore the self energy density for electric ( $\rho(\vec{E})$ ) and magnetic ( $\rho(\vec{B})$ ) fields come through the terms containing  $(A'_i)^2$  and  $F_{ij} F_{ij}$  respectively of Eq.(19), while the third term of the above  $T_0^0$ , that contains  $A'_i F_{jk}$ , represents the interaction energy density between the electric and magnetic field ( $\rho_{\text{int}}(\vec{E}, \vec{B})$ ). In particular, we can write

$$\begin{aligned} \rho(\vec{E}) &= -\frac{1}{2a^4} (A'_i)^2 , & \rho(\vec{B}) &= -\frac{1}{4a^4} F_{ij} F_{ij} \\ \rho_{\text{int}}(\vec{E}, \vec{B}) &= \frac{2\lambda q}{a^4} \epsilon_{ijk} A'_i F_{jk} \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta+1)]^q + [-24(\beta+1)^3]^{q/2} \right\} \left( \frac{-\eta}{\eta_0} \right)^{2\epsilon q} . \end{aligned} \quad (20)$$

It may be observed that  $\rho_{\text{int}}(\vec{E}, \vec{B})$  gets proportional to the non-minimal coupling parameter  $\lambda$ , which indicates that the interaction energy density between electric and magnetic fields is generated entirely due to the presence of the parity violating term  $\mathcal{L}_{CB} = \sqrt{-g}\lambda f(R, \mathcal{G}) \epsilon^{\nu\alpha\beta} F_{\nu\alpha} F_{\beta}$  in the action. This is unlike to the magnetogenesis model discussed in [36] where the action preserves the parity of the EM field, and thus the interaction energy density between electric and magnetic fields do not seem to arise in such models. Coming back to our present model, the possible effects of  $\mathcal{L}_{CB}$  are also incorporated within  $\rho(\vec{E})$  and  $\rho(\vec{B})$ , however in a implicit way through the solution of  $A_i(\eta, \vec{x})$ . As mentioned earlier, the parity violating term  $\mathcal{L}_{CB}$  in the action leads to the helical nature of the magnetic field, in which case the helicity can be quantified by the quantity  $\rho_h = -A_i B^i$ , where  $B^i$  specifies the magnetic field. At this stage, we proceed to quantize the EM field, i.e  $A_i$  is promoted to a hermitian operator  $\hat{A}_i$ , and moreover the field is

considered to start from the Bunch-Davies vacuum state (symbolized by  $|0\rangle$ ) at distant past. Due to the Heisenberg picture, the state of the field remains at  $|0\rangle$  with time. As a result, the expectation value of electric energy density in the state  $|0\rangle$  is given by,

$$\langle \rho(\vec{E}) \rangle = -\frac{1}{2a^4} \langle 0|(A'_i)^2|0\rangle . \quad (21)$$

Similarly the expectation values of magnetic and the interaction energy density read as,

$$\langle \rho(\vec{B}) \rangle = -\frac{1}{4a^4} \langle 0|F_{ij}F_{ij}|0\rangle , \quad (22)$$

and

$$\langle \rho_{\text{int}}(\vec{E}, \vec{B}) \rangle = \frac{2\lambda q}{a^4} \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta+1)]^q + [-24(\beta+1)^3]^{q/2} \right\} \left( \frac{-\eta}{\eta_0} \right)^{2\epsilon q} \langle 0|\epsilon_{ijk}A'_iF_{jk}|0\rangle , \quad (23)$$

respectively. Moreover the helicity energy density has the expectation as  $\langle \rho_h \rangle = -\langle 0|A_iB^i|0\rangle$ . Expanding  $\hat{A}_i(\eta, \vec{x})$  in terms of Fourier basis as,

$$\hat{A}_i(\eta, \vec{x}) = \int \frac{d\vec{k}}{(2\pi)^3} \sum_{r=1,2} \epsilon_{ri} \left[ \hat{b}_r(\vec{k})A_r(k, \eta)e^{i\vec{k}\cdot\vec{x}} + \hat{b}_r^+(\vec{k})A_r^*(k, \eta)e^{-i\vec{k}\cdot\vec{x}} \right] , \quad (24)$$

where  $\vec{k}$  is the Fourier mode momentum or equivalently the EM wave vector,  $r$  is the polarization index,  $\vec{\epsilon}_+$  and  $\vec{\epsilon}_-$  are two polarization vectors, and  $A_r(k, \eta)$  is the EM mode function corresponds to the mode  $\vec{k}$ . Due to the helical nature of the magnetic field, it will be useful if we work with the helicity basis set in the present context, in which case the polarization vectors are given by:  $\vec{\epsilon}_+ = \frac{1}{\sqrt{2}}(1, i, 0)$  and  $\vec{\epsilon}_- = \frac{1}{\sqrt{2}}(1, -i, 0)$  respectively. The Coulomb gauge condition, i.e.  $\partial_i A^i = 0$  immediately leads to  $\vec{k}\cdot\vec{\epsilon}_+ = \vec{k}\cdot\vec{\epsilon}_- = 0$ , which reveals that the EM propagation vector is perpendicular to the plane spanned by  $(\vec{\epsilon}_+, \vec{\epsilon}_-)$  vectors. The polarization vectors also satisfy the relation  $\epsilon_{ijk}k_j\epsilon_{\pm k} = \mp ik\epsilon_{\pm i}$  with  $k = |\vec{k}|$ , which will be useful later in solving the EM mode function. Furthermore  $\hat{b}_r(\vec{k})$  and  $\hat{b}_r^+(\vec{k})$  are the creation and annihilation operators of  $\eta \rightarrow -\infty$  respectively, in particular they are defined over the vacuum state of distant past by the condition  $\hat{b}_r(\vec{k})|0\rangle = 0 \forall \vec{k}$ . The quantization rule is imposed by the commutation relation of creation and annihilation operators as,

$$[\hat{b}_p(\vec{k}), \hat{b}_r^+(\vec{k}')] = \delta_{pr}\delta(\vec{k} - \vec{k}') . \quad (25)$$

Using the Fourier mode decomposition of the EM field along with the commutation relation of Eq.(25), the expectation values of various energy densities come as,

$$\begin{aligned} \langle \rho(\vec{E}) \rangle &= \sum_{r=1,2} \int \frac{dk}{2\pi^2} \frac{k^2}{a^4} |A'_r(k, \eta)|^2 \\ \langle \rho(\vec{B}) \rangle &= \sum_{r=1,2} \int \frac{dk}{2\pi^2} \frac{k^4}{a^4} |A_r(k, \eta)|^2 \end{aligned} \quad (26)$$

and

$$\langle \rho_{\text{int}}(\vec{E}, \vec{B}) \rangle = -4\lambda q \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta+1)]^q + [-24(\beta+1)^3]^{q/2} \right\} \left( \frac{-\eta}{\eta_0} \right)^{2\epsilon q} \sum_{r=1,2} \int \frac{dk}{2\pi^2} \frac{k^3}{a^4} |A_r(k, \eta)A'_r(k, \eta)| \quad (27)$$

respectively. Similarly, the helicity energy density in the Fourier basis is given by,

$$\langle \rho_h \rangle = \int \frac{dk}{2\pi^2} \frac{k^3}{a^3} \left\{ |A_+(k, \eta)|^2 - |A_-(k, \eta)|^2 \right\} . \quad (28)$$

Due to the presence of the parity violating term in the action, the two modes of the EM field, i.e.  $A_+(k, \eta)$  and  $A_-(k, \eta)$ , evolve differently and leads to a non-zero helicity in the present context. This is in contrast to [36] where the EM field couples with the background spacetime curvature via  $\sim \sqrt{-g}f(R, \mathcal{G})F_{\mu\nu}F^{\mu\nu}$ , in which case the  $A_+(k, \eta)$  and  $A_-(k, \eta)$  evolve in a similar fashion and thus the helicity vanishes trivially. Coming back to our present model,

Eq.(42) immediately leads to the power spectra of electric and magnetic fields (defined by the energy density within unit logarithmic interval of  $k$ ) as,

$$\mathcal{P}(\vec{E}) = \frac{\partial \langle \rho(\vec{E}) \rangle}{\partial \ln k} = \sum_{r=1,2} \frac{k}{2\pi^2} \frac{k^2}{a^4} |A'_r(k, \eta)|^2 \quad , \quad \mathcal{P}(\vec{B}) = \frac{\partial \langle \rho(\vec{B}) \rangle}{\partial \ln k} = \sum_{r=1,2} \frac{k}{2\pi^2} \frac{k^4}{a^4} |A_r(k, \eta)|^2 \quad . \quad (29)$$

Furthermore, the helicity power spectrum is determined from Eq.(28), and given by,

$$\mathcal{P}_h = \frac{\partial \langle \rho_h \rangle}{\partial \ln k} = \frac{k}{2\pi^2} \frac{k^3}{a^3} \left\{ |A_+(k, \eta)|^2 - |A_-(k, \eta)|^2 \right\} \quad . \quad (30)$$

Here it may be observed that the scale dependence of  $\mathcal{P}(\vec{E})$ ,  $\mathcal{P}(\vec{B})$  and  $\mathcal{P}_h$  are different. In particular, the electric power spectrum gets the scale dependency from the  $k^3$  factor as well as from the time derivative of the EM mode function, while the magnetic power spectrum has the scale dependency due to the  $k^5$  factor and also due to the mode function itself. On the other hand, the scale dependency of the helicity spectrum comes through the  $k^3$  factor and from the difference between the two EM mode functions. Therefore it is very unlikely, that the electric, magnetic and helicity power spectra have similar dependency on  $k$ . In order to extract the scale dependency of such power spectra, we need to solve the EM mode functions from Eq.(10), which is the subject of the next section. The solution of the mode functions are also necessary to determine the magnetic strength and consequently the baryon asymmetry of the present universe.

#### IV. SOLUTION OF THE MODE FUNCTION AND SCALE DEPENDENCE OF THE POWER SPECTRA DURING INFLATION

In this section we aim to determine the solution of EM mode functions and the corresponding power spectra during the inflationary era when the scale factor behaves as of Eq.(11). For this purpose, we need Eq.(10) which, due to the Fourier decomposition of  $A_i(\eta, \vec{x})$ , takes the following form:

$$A''_{\pm}(k, \eta) + (k^2 \mp 8\lambda k f'(R, \mathcal{G})) A_{\pm}(k, \eta) = 0 \quad , \quad (31)$$

where  $f'(R, \mathcal{G}) = \frac{df}{d\eta}$ . Eq.(31) clearly evidents that the two mode functions, i.e  $A_+(k, \eta)$  and  $A_-(k, \eta)$ , obey different equation of motion and hence have different cosmological evolution. This is however expected solely due to the reason that the EM field couples with the spacetime curvature via its dual tensor, which indeed violates the parity of the EM field. The form of  $f(R, \mathcal{G})$  of Eq.(5) immediately leads to the  $f'(R, \mathcal{G})$  as follows,

$$f'(R, \mathcal{G}) = (2\epsilon q) \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta + 1)]^q + [-24(\beta + 1)^3]^{q/2} \right\} \left( \frac{1}{\eta} \right) \quad .$$

Plugging back this expression of  $f'(R, \mathcal{G})$  into Eq.(31), we get

$$A''_{\pm}(k, \eta) + \left( k^2 \mp \frac{Bk}{\eta} \right) A_{\pm}(k, \eta) = 0 \quad , \quad (32)$$

where  $B$  has the following form,

$$B = (16\epsilon q \lambda) \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta + 1)]^q + [-24(\beta + 1)^3]^{q/2} \right\} \quad . \quad (33)$$

Here it may be mentioned that the quantity  $B$  is proportional with  $\epsilon$ ,  $q$  and thus for  $\epsilon = 0$  or  $q = 0$ , both the EM mode functions seem to satisfy the standard Maxwell's equation in vacuum. This is expected, because for  $\epsilon = 0$  or  $q = 0$ , the function  $f(R, \mathcal{G})$  becomes constant (with time), due to which the conformal breaking term  $\mathcal{L}_{CB} = \sqrt{-g} \lambda f(R, \mathcal{G}) \epsilon^{\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta}$  present in the action becomes a total surface term (as mentioned earlier) and thus gives no contributions to the field equations. Thereby the non-zero values of  $\epsilon$  (the slow roll parameter) and  $q$  modify the EM field equations non-trivially by Eq.(32), compared to the standard Maxwell's equation. Eq.(32) is solved for  $A_{\pm}(k, \eta)$  and the solutions are following:

$$A_+(k, \eta) = e^{ik\eta} (-k\eta) \left\{ C_1 U \left( 1 + \frac{iB}{2}, 2, -2ik\eta \right) + C_2 1F1 \left( 1 + \frac{iB}{2}, 2, -2ik\eta \right) \right\} \quad ,$$



$$A_-(k, \eta) = e^{ik\eta} (-k\eta) \left\{ C_3 U \left( 1 - \frac{iB}{2}, 2, -2ik\eta \right) + C_4 {}_1F_1 \left( 1 - \frac{iB}{2}, 2, -2ik\eta \right) \right\}, \quad (34)$$

where  $C_i$  ( $i = 1, 2, 3, 4$ ) are integration constants,  $U(a, b, z)$  is the confluent hypergeometric function and  ${}_1F_1(a, b, z)$  symbolizes the Kummer confluent hypergeometric function. The integration constants can be determined from the initial Bunch-Davies vacuum state of the EM mode functions, in particular,  $\lim_{|k\eta| \gg 1} A_\pm(k, \eta) = \frac{1}{\sqrt{2k}} e^{-ik\eta}$ . Such initial condition is compatible with Eq.(32) and the demonstration goes as follows: Eq.(32), in the deep sub-Hubble regime when the modes are within the Hubble radius, can be written as  $A_\pm''(k, \eta) + k^2 A_\pm(k, \eta) = 0$  which is indeed satisfied by the Bunch-Davies solution. Therefore in order to determine  $C_i$ , one needs to calculate the limiting forms of  $A_\pm(k, \eta)$  at  $|k\eta| \gg 1$ . For this purpose we use the following asymptotic expressions of the hypergeometric functions as,

$$\begin{aligned} \lim_{|k\eta| \gg 1} U \left( 1 \pm \frac{iB}{2}, 2, -2ik\eta \right) &= (-2ik\eta)^{-1 \mp iB/2}, \\ \lim_{|k\eta| \gg 1} {}_1F_1 \left( 1 \pm \frac{iB}{2}, 2, -2ik\eta \right) &= \frac{1}{\Gamma(1 \mp iB/2)} (2ik\eta)^{-1 \mp iB/2} + \frac{e^{-2ik\eta}}{\Gamma(1 \pm iB/2)} (-2ik\eta)^{-1 \pm iB/2}, \end{aligned}$$

due to which, the EM mode functions from Eq.(32) take the following forms in the deep sub-Hubble regime,

$$\begin{aligned} \lim_{|k\eta| \gg 1} A_+(k, \eta) &= e^{ik\eta} (-2i)^{-1-iB/2} \left\{ C_1 - \frac{C_2(-1)^{iB/2}}{\Gamma(1-iB/2)} \right\} + e^{-ik\eta} (2i)^{-1+iB/2} \frac{C_2}{\Gamma(1+iB/2)}, \\ \lim_{|k\eta| \gg 1} A_-(k, \eta) &= e^{ik\eta} (-2i)^{-1+iB/2} \left\{ C_3 - \frac{C_4(-1)^{-iB/2}}{\Gamma(1+iB/2)} \right\} + e^{-ik\eta} (2i)^{-1-iB/2} \frac{C_4}{\Gamma(1-iB/2)}, \end{aligned} \quad (35)$$

where  $\Gamma(z)$  is the gamma function. Owing to the Bunch-Davies condition, the coefficient of the positive frequency part of  $A_\pm(k, \eta)$  matches with  $\frac{1}{\sqrt{2k}}$  and that of the negative frequency part must vanish. As a result, Eq.(35) yields four algebraic equations of  $C_i$ , from which one easily determines the integration constants as,

$$\begin{aligned} C_1 &= \frac{2^{1-iB/2} i}{\sqrt{2k}} \left( \frac{\Gamma(1+iB/2)}{\Gamma(1-iB/2)} \right) e^{-\pi B/4}, & C_2 &= \frac{2^{1-iB/2} i}{\sqrt{2k}} \Gamma(1+iB/2) e^{\pi B/4}, \\ C_3 &= \frac{2^{1+iB/2} i}{\sqrt{2k}} \left( \frac{\Gamma(1-iB/2)}{\Gamma(1+iB/2)} \right) e^{\pi B/4}, & C_4 &= \frac{2^{1+iB/2} i}{\sqrt{2k}} \Gamma(1-iB/2) e^{-\pi B/4}, \end{aligned} \quad (36)$$

respectively, with recall,  $B$  is given in Eq.(33). Thus as a whole,  $A_\pm(k, \eta)$  follow the solutions of Eq.(34) where  $C_i$  are obtained in Eq.(36). Consequently Eq.(34) leads to the  $A_\pm(k, \eta)$  in the superhorizon limit, when the modes are outside of the Hubble radius satisfying  $|k\eta| \ll 1$ , as,

$$\lim_{|k\eta| \ll 1} A_+(k, \eta) = \frac{C_1}{2i\Gamma(1+iB/2)}, \quad \lim_{|k\eta| \ll 1} A_-(k, \eta) = \frac{C_3}{2i\Gamma(1-iB/2)}. \quad (37)$$

Moreover  $A'_\pm(k, \eta)$  (which are necessary for electric power spectrum) in the superhorizon regime are obtained as,

$$\begin{aligned} \lim_{|k\eta| \ll 1} \frac{dA_+}{d(-k\eta)} &= C_2 + \frac{C_1}{\Gamma(iB/2)} \{ \gamma + \ln(-2k\eta) + iB/2 \} \\ \lim_{|k\eta| \ll 1} \frac{dA_-}{d(-k\eta)} &= C_4 + \frac{C_3}{\Gamma(-iB/2)} \{ \gamma + \ln(-2k\eta) - iB/2 \}, \end{aligned} \quad (38)$$

where  $\gamma$  is the Euler's constant with numerical value  $\simeq 0.577216$ . Plugging back the above expressions into Eq.(29) immediately yields the electric and magnetic power spectra in the superhorizon regime as,

$$\frac{\partial \rho(\vec{E})}{\partial \ln k} = \frac{2H_0^4 \cosh(\pi B/2)}{\pi^2 |\Gamma(iB/2)|^2} (-k\eta)^4 [\ln(-k\eta)]^2 \quad (39)$$

and

$$\frac{\partial \rho(\vec{B})}{\partial \ln k} = \frac{H_0^4 \cosh(\pi B/2)}{2\pi^2 |\Gamma(1+iB/2)|^2} (-k\eta)^4 \quad (40)$$

respectively. Similarly, for the interaction energy density between electric and magnetic fields, we get

$$\begin{aligned} \frac{\partial \rho_{\text{int}}}{\partial \ln k} &= - \left( \frac{4\lambda q}{\pi^2} \right) \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta+1)]^q + [-24(\beta+1)^3]^{q/2} \right\} \left( \frac{-\eta}{\eta_0} \right)^{2\epsilon q} \\ &\times \left( \frac{H_0^4 \cosh(\pi B/2)}{|\Gamma(iB/2)\Gamma(1+iB/2)|} \right) (-k\eta)^4 [\ln(-k\eta)] . \end{aligned} \quad (41)$$

Eqs.(39) and (40) clearly demonstrate that the electric and magnetic power spectra are not scale invariant during the superhorizon inflationary era, however they have different scale dependence, in particular, the electric power spectrum scales as  $k^4 [\ln(-k\eta)]^2$  while the magnetic spectrum goes as  $k^4$ . The logarithmic scale behaviour of the electric spectrum is a distinct feature of the present magnetogenesis model compared to various previous magnetogenesis models where both the electric and magnetic spectra scale by the power law behaviour [36, 37]. Furthermore both the  $\frac{\partial \rho(\vec{E})}{\partial \ln k}$  and  $\frac{\partial \rho(\vec{B})}{\partial \ln k}$  tends to zero as  $|k\eta| \rightarrow 0$ , which is also true for the interaction power spectrum as evident from Eq.(41). Such characteristics of the power spectra hints to the resolution of the backreaction problem in the present magnetogenesis model.

However in order to ensure that the backreaction of the EM field on the background spacetime stays small during the inflation, we first consider the electric field energy density at  $\eta = \eta_c$ , which is given by,

$$\rho(\vec{E}, \eta_c) = \int_{k_i}^{k_c} \frac{\partial \rho(\vec{E})}{\partial \ln k} d \ln k = \frac{H_0^4 \cosh(\pi B/2)}{16\pi^2 |\Gamma(iB/2)|^2} \left\{ 1 - e^{-4N_c} (1 + 4N_c + 8N_c^2) \right\} , \quad (42)$$

where  $k_i$  and  $k_c$  are the mode momentum which cross the horizon at the beginning of inflation and at  $\eta = \eta_c$  respectively. As mentioned earlier, the beginning of inflation is considered to be the instance when the CMB scale mode crosses the horizon, and thus  $k_i = k_{\text{CMB}} \approx 10^{-40} \text{ GeV}$ . Moreover we have  $|\eta_c| = k_c^{-1}$  where  $\eta_c$  is any intermediate time instance during the inflation and thus  $k_c$  is greater than the CMB scale momentum. The quantity  $N_c$  represents the inflationary e-folding number up-to  $\eta = \eta_c$ , i.e  $N_c = \ln \left( \frac{a_c}{a_i} \right)$  with  $a_c = a(\eta_c)$  and  $a_i = a(\eta_i)$ . Similarly using Eq.(40), we derive the energy density stored in the magnetic field at  $\eta = \eta_c$  as,

$$\rho(\vec{B}, \eta_c) = \int_{k_i}^{k_c} \frac{\partial \rho(\vec{B})}{\partial \ln k} d \ln k = \frac{H_0^4 \cosh(\pi B/2)}{8\pi^2 |\Gamma(1+iB/2)|^2} \left\{ 1 - e^{-4N_c} \right\} , \quad (43)$$

where, as well as in Eq.(42), we use  $\eta_0^{-1} = H_0$ . The energy density due to the interaction between the electric and magnetic fields at  $\eta = \eta_c$  comes from Eq.(41), and is determined as,

$$\begin{aligned} \rho_{\text{int}}(\eta_c) &= \int_{k_i}^{k_c} \frac{\partial \rho_{\text{int}}}{\partial \ln k} d \ln k = \left( \frac{\lambda q}{4\pi^2} \right) \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta+1)]^q + [-24(\beta+1)^3]^{q/2} \right\} \left( \frac{-\eta_c}{\eta_0} \right)^{2\epsilon q} \\ &\times \left( \frac{H_0^4 \cosh(\pi B/2)}{|\Gamma(iB/2)\Gamma(1+iB/2)|} \right) \left\{ 1 - e^{-4N_c} (1 + 4N_c) \right\} . \end{aligned} \quad (44)$$

Therefore the total electromagnetic energy density at  $\eta = \eta_c$  becomes

$$\rho_{em}(\eta_c) = \rho(\vec{E}, \eta_c) + \rho(\vec{B}, \eta_c) + \rho_{\text{int}}(\eta_c) , \quad (45)$$

where  $\rho(\vec{E}, \eta_c)$ ,  $\rho(\vec{B}, \eta_c)$  and  $\rho_{\text{int}}(\eta_c)$  are obtained in Eqs.(42), (43) and (44) respectively. In order to resolve the backreaction issue, we need to show that the total EM energy density becomes lower than the background energy density which is given by  $\rho_{\text{bg}} = 3H^2 M_{\text{Pl}}^2$ , where  $M_{\text{Pl}}$  is the reduced Planck mass. Since the total e-folding of the inflation (symbolized by  $N_f$ ) is nearly  $N_f = 60$ , we safely consider  $N_c \leq 60$ , and due to the fact that  $\epsilon < 1$  and  $q < 1$ , Eq.(45) reveals that the total EM energy density during the inflation is of the order  $\sim H_0^4$ , i.e,

$$\rho_{em}(\eta_c) \sim H_0^4 . \quad (46)$$

The energy scale at the beginning of inflation is generally considered to be  $H_0 = 10^{-5} M_{\text{Pl}}$  and moreover the Hubble parameter at the end of inflation can be determined from  $H = H_0 \exp \left( -\frac{\epsilon N_f}{1+\epsilon} \right)$  as established in Eq.(14). Therefore Eq.(46) leads to the inequality that  $\rho_{em} \ll \rho_{\text{bg}} = 3H^2 M_{\text{Pl}}^2$ , in particular, the ratio of  $\rho_{em}$  and  $\rho_{\text{bg}}$  is found to be:  $\frac{\rho_{em}}{\rho_{\text{bg}}} \approx 10^{-4}$  for  $\epsilon = 0.1$ ,  $N_f = 60$  and  $\frac{\rho_{em}}{\rho_{\text{bg}}} \approx 10^{-8}$  for  $\epsilon = 0.01$ ,  $N_f = 60$  etc. This confirms that the EM field

has negligible backreaction on the background inflationary spacetime, leading to the resolution of the backreaction problem in the present context.

In regard to the helicity of the magnetic field, Eq.(30) leads to the helicity power spectrum during the inflation as,

$$\frac{\partial \langle \rho_h \rangle}{\partial \ln k} = -\frac{1}{2\pi^2} \left( \frac{k^3}{a^3} \right) \left( \frac{\sinh(\pi B/2)}{|\Gamma(1+iB/2)|^2} \right), \quad (47)$$

where we use the the superhorizon solutions of  $A_{\pm}(k, \eta)$ , given in Eq.(37). Several points need to be mentioned from Eq.(47) : (1) the comoving helicity remains conserved with the expansion of the universe during the inflationary super-Hubble regime. (2) Since the quantity  $B$  is proportional to the conformal breaking parameter  $\lambda$  (see Eq.(33)), the helicity spectrum vanishes for  $\lambda = 0$ . This is due to the reason that for  $\lambda = 0$ , both the EM mode functions have identical evolution, and as a result, the helicity of the magnetic field becomes zero. (3) Owing to the fact that  $B > 0$  (see Eq.(33)), the helicity power spectrum during the inflation seems to be negative, which indicates that the conformal breaking term present in the EM action enhances the amplitude of  $A_{-}(k, \eta)$  compared to that of  $A_{+}(k, \eta)$  in the inflationary era.

## V. PRESENT MAGNETIC STRENGTH AND MAGNETIC HELICITY FOR THE INSTANTANEOUS REHEATING CASE

In this section, we aim to calculate the magnetic strength and consequently the magnetic helicity at current universe. For this purpose, we need to know about the conductivity of the universe during and after the inflation. During the inflation, the universe remains a poor electrical conductor where the conductivity is low, however in the post-inflationary era, the universe becomes a good conductor. The large conductivity during the post-inflation lies in the consideration that the universe experiences an *instantaneous reheating* phase just after the inflation, in which case, the e-fold number of the reheating epoch is zero and thus the universe makes a sudden jump from the inflation to a radiation dominated epoch where the Hubble parameter evolves as  $H \propto t^{-1}$  with  $t$  being the cosmic time (here we would like to mention that at a later section, we will relax the assumption of ‘‘instantaneous reheating’’ and will consider a reheating epoch having non-zero e-fold number after inflation). The high conductivity in the post-inflation phase leads to a electric current of the form  $J^i = \sigma E^i$  (where  $\sigma$  is the electrical conductivity and  $E^i$  is the electric field) which in turn acts as a source term in the EM field equation. Due to the presence of such source term and owing to the large  $\sigma$ , the solution of the EM field after inflation behaves as  $A_{\pm}(k, \eta) = C_{\pm} + D_{\pm} e^{-\sigma t}$  (with  $C_{\pm}$  and  $D_{\pm}$  being the integration constants), where the term  $e^{-\sigma t}$  becomes soon negligible and thus  $A_{\pm}(k, \eta)$  becomes constant with time. As a result, the electric field becomes vanishingly small and the magnetic field remains the dominant piece in the electromagnetic energy density. Moreover we mentioned earlier that at the late time, the conformal breaking coupling  $f(R, \mathcal{G})$  do not contribute and thus the EM action restores the conformal symmetry of the EM field. In particular, the function  $f(R, \mathcal{G})$  is considered to be zero from the end of inflation, which is also connected from the continuity point of view, as Eq.(5) leads to  $f(R, \mathcal{G})$  tends to zero as  $|k\eta| \rightarrow 0$ , i.e at the end of inflation. From the conformal symmetry of the EM field, we may argue that the EM energy density decays as  $\frac{1}{a^4}$  with the expansion of the universe, or equivalently the magnetic energy density goes by  $1/a^4$ , as the electric field practically vanishes in the post-inflationary epoch. As a result, the magnetic energy density at the end of inflation is related to that of at the present epoch by the following relation,

$$\left. \frac{\partial \rho(\vec{B})}{\partial \ln k} \right|_0 = \left( \frac{a_f}{a_0} \right)^4 \left. \frac{\partial \rho(\vec{B})}{\partial \ln k} \right|_{\eta_f}, \quad (48)$$

where the suffix ‘f’ and ‘0’ denote the end instance of inflation and the current time of universe respectively. Moreover, due to the reason that the EM mode functions become constant with time, the helicity power spectrum evolves as  $1/a^3$  in the post-inflationary era, and thus  $\mathcal{P}_h$  at the end of inflation is connected to that of at the present epoch by following:

$$\left. \frac{\partial \rho_h}{\partial \ln k} \right|_0 = \left( \frac{a_f}{a_0} \right)^3 \left. \frac{\partial \rho_h}{\partial \ln k} \right|_{\eta_f}. \quad (49)$$

Eqs.(40) and (48) lead to the magnetic field’s current amplitude as,

$$B_0 = \frac{1}{\sqrt{\pi}} \left\{ \frac{\cosh(\pi B/2)}{|\Gamma(1+iB/2)|^2} \right\}^{\frac{1}{2}} \left( \frac{a_f}{a_0} \right)^2 H_0^2 (-k\eta_f)^2, \quad (50)$$

and moreover, Eqs.(30) and (49) yield the helicity power spectrum at present epoch as,

$$\left. \frac{\partial \rho_h}{\partial \ln k} \right|_0 = -\frac{1}{2\pi^2} \left( \frac{a_f}{a_0} \right)^3 \left( \frac{\sinh(\pi B/2)}{|\Gamma(1+iB/2)|^2} \right) H_0^3 (-k\eta_f)^3, \quad (51)$$

with, recall that  $\mathcal{B} = (16\epsilon q\lambda) \left( \frac{\kappa}{\eta_0} \right)^{2q} \left\{ [6\beta(\beta+1)]^q + [-24(\beta+1)^3]^{q/2} \right\}$  and  $k$  denotes the mode-momentum on which we are interested to estimate the magnetic strength. In order to estimate  $B_0$  from Eq.(50), we need to know  $\frac{a_f}{a_0}$  and  $|k\eta_f|$ . In regard to  $\frac{a_f}{a_0}$ , we use the entropy conservation in the post-inflation era, i.e  $gT^3 a^3 = \text{constant}$ , where  $g$  symbolizes the effective relativistic degrees of freedom and  $T$  is the corresponding temperature – this finally results to  $\frac{a_0}{a_f} = 10^{30} (H_f/10^{-5} M_{\text{Pl}})^{1/2}$ , where  $H_f$  is the Hubble parameter at the end of inflation and can be determined from Eq.(14) as  $H_f = H_0 \exp\left(-\frac{\epsilon N_f}{1+\epsilon}\right)$ , with  $N_f$  being the total e-fold of the inflationary phase. The set:  $H_0 = 10^{-5} M_{\text{Pl}}$ ,  $\epsilon = 0.1$  and  $N_f = 60$  immediately leads to  $H_f \approx 5 \times 10^{-8} M_{\text{Pl}}$ . Moreover, for the purpose of determining  $|k\eta_f|$ , the relation  $|\eta_f| = 1/k_f$  stands to be useful where  $k_f$  is the momentum of the mode which crosses the Hubble horizon at the end instance of inflation, and  $k$  lies around the CMB scale where we want to determine  $B_0$ . As a consequence,  $|k\eta_f|$  is given by  $|k\eta_f| = \exp(-N_f)$ . Using such expressions of  $\frac{a_f}{a_0}$  and  $|k\eta_f|$ , along with  $q = 0.5$  and  $\lambda \sim 1$ , we calculate the present amplitude of the magnetic field from Eq.(50) as,

$$B_0 \Big|_{\text{CMB}} \approx 10^{-62} \text{G}, \quad (52)$$

where the conversion  $1\text{G} = 1.95 \times 10^{-20} \text{GeV}^2$  is used. Eq.(52) provides the theoretical prediction of magnetic field's current amplitude in the context of our magnetogenesis model. In order to confront the model with observations, we use the CMB results, according to which, the present magnetic field around the CMB scale lies within  $10^{-10} \text{G} \lesssim B_0 \lesssim 10^{-22} \text{G}$ . However Eq.(52) demonstrates that the theoretically predicted  $B_0$  lies far below than the observational constraints. Thus we may argue that the magnetogenesis model, where the EM field gets non-minimally coupled by the term  $\mathcal{L}_{CB} = \sqrt{-g} \lambda f(R, \mathcal{G}) \epsilon^{\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta}$ , seems to be unable to generate sufficient magnetic strength at present epoch of the universe. This argument is not only confined to the value  $q = 0.5$ , however holds true for the entire possible range of  $0 < q < 1$ .

Here we would like to mention that the above arguments are valid in the case of *instantaneous reheating* where, as mentioned earlier, the reheating epoch has zero e-fold and thus the universe gets a large conductivity after the end of inflation. However it would be more physical if the universe undergoes through a reheating phase having a non-zero e-fold number. Motivated by this, in the next section, we will relax the assumption of ‘‘instantaneous reheating’’ and will consider a prolonged reheating stage in-between the end of inflation and the beginning of radiation dominated epoch.

## VI. PRESENT MAGNETIC STRENGTH AND MAGNETIC HELICITY FOR A KAMIONKOWSKI LIKE REHEATING MODEL WITH NON-ZERO E-FOLDING NUMBER

Instead of instantaneous reheating (that we have considered in Sec.[V]), it would be more physical if the universe undergoes through a reheating phase which has a non-zero e-fold number. In the case of instantaneous reheating, the electrical conductivity immediately turns on after the end of inflation and thus the electric field practically goes to zero. However if the reheating phase is considered to have a non-zero e-fold number, then there is no reason to consider a large conductivity immediately after the inflation. Indeed, the conductivity remains non-zero and small, due to which, the existence of strong electric field induces the magnetic field during the stage between the end of inflation and the end of reheating [43]. As a result, the magnetic field's strength and consequently the helicity spectrum at present universe may become larger compared to the instantaneous reheating case. In such situation, it is important to calculate the magnetic field's strength and the helicity spectrum at current time by considering the reheating phase with a non-zero e-fold number. This is our aim in the present section.

In regard to the reheating dynamics, we consider the conventional reheating mechanism proposed by Kamionkowski et al. [105], where the main idea is to parametrize the reheating dynamics by – a non-zero e-fold number, a constant equation of state (EoS) parameter and a given reheating temperature. Clearly, the Hubble parameter during the reheating era is connected to that of at the end of inflation by the constant EoS parameter (symbolized by  $\omega_{\text{eff}}$ ), in particular,  $H \propto a^{-\frac{3}{2}(1+\omega_{\text{eff}})}$  in the reheating stage. Moreover the inflaton energy density is supposed to instantaneously convert to radiation energy density after the end of reheating, which indicates the beginning of radiation dominated

epoch. The characterized quantities of the reheating phase, i.e the e-fold number ( $N_{\text{re}}$ ) and the temperature ( $T_{\text{re}}$ ) can be expressed in terms of  $\omega_{\text{eff}}$  and of some inflationary parameters by following [105, 106],

$$N_{\text{re}} = \frac{4}{(1 - 3\omega_{\text{eff}})} \left[ -\frac{1}{4} \ln \left( \frac{45}{\pi^2 g_{\text{re}}} \right) - \frac{1}{3} \ln \left( \frac{11g_{s,\text{re}}}{43} \right) - \ln \left( \frac{k}{a_0 T_0} \right) - \ln \left( \frac{(3H_f^2 M_{\text{Pl}}^2)^{1/4}}{H_0} \right) - N_f \right] , \quad (53)$$

$$T_{\text{re}} = H_0 \left( \frac{43}{11g_{s,\text{re}}} \right)^{\frac{1}{3}} \left( \frac{a_0 T_0}{k} \right) \exp [ - (N_f + N_{\text{re}}) ] , \quad (54)$$

where  $T_0 = 2.75\text{K}$  is the present temperature of the universe,  $\frac{k}{a_0}$  is the pivot scale generally taken as  $\approx 0.02\text{Mpc}^{-1}$  in case of large scale modes and  $a_0$  is the present cosmological scale factor. Here, for simplicity, we have taken both the values of the degrees of freedom (dof) as  $g_{s,\text{re}} = g_{\text{re}} \approx 100$ , where  $g_{s,\text{re}}$  is the dof for entropy at reheating and  $g_{\text{re}}$  is the effective number of relativistic species upon thermalization. With this reheating model in hand, we now solve the electromagnetic mode function and consequently determine the power spectra during the reheating epoch, in the next subsection.

### A. Solution of the mode function and power spectra during the reheating epoch

We mentioned earlier that the function  $f(R, \mathcal{G})$  is considered to be zero after the end of inflation, which is also connected from the continuity point of view, as Eq.(5) clearly indicates that  $f(R, \mathcal{G})$  tends to zero as  $|k\eta| \rightarrow 0$ . As a consequence, the production of the gauge field from the quantum vacuum stops, which is accounted by the fact that the Bogoliubov coefficients in the post-inflation become explicitly time independent and get equal to that of at the end of inflation. Therefore, in the reheating epoch, the EM field follows the standard Maxwell's equation, in particular,

$$A_{\pm}^{(re)''}(k, \eta) + k^2 A_{\pm}^{(re)}(k, \eta) = 0 , \quad (55)$$

where  $A_{\pm}^{(re)}(k, \eta)$  is the EM mode function during the reheating phase. Eq.(55) points that, during the reheating era, the electrical conductivity of the universe is considered to be negligible, in particular to be zero. However due to the Schwinger effect, the assumption of zero conductivity needs proper justification, which we will investigate in Sec.[VII]. Solving Eq.(55), one gets,

$$A^{(re)}(k, \eta) = \frac{1}{\sqrt{2k}} \left[ c_{\pm} e^{-ik(\eta-\eta_f)} + d_{\pm} e^{ik(\eta-\eta_f)} \right] , \quad (56)$$

where  $c_{\pm}$  and  $d_{\pm}$  are integration constants, that can be determined from the continuity conditions of the EM field at the junction between the end of inflation and the beginning of reheating, i.e,

$$A_{\pm}^{(re)}(k, \eta_f) = A_{\pm}(k, \eta_f) \quad \text{and} \quad A_{\pm}^{(re)'}(k, \eta_f) = A'_{\pm}(k, \eta_f), \quad (57)$$

with  $A_{\pm}(k, \eta_f)$  is the EM mode functions at the end of inflation and have the solution of Eq.(34) for  $\eta = \eta_f$ . Such continuity relations immediately lead to the integration constants in terms of  $A_{\pm}(k, \eta_f)$  and  $A'_{\pm}(k, \eta_f)$  as,

$$\begin{aligned} c_{\pm} &= \sqrt{\frac{k}{2}} A_{\pm}(k, \eta_f) + \frac{i}{\sqrt{2k}} A'_{\pm}(k, \eta_f) \\ d_{\pm} &= \sqrt{\frac{k}{2}} A_{\pm}(k, \eta_f) - \frac{i}{\sqrt{2k}} A'_{\pm}(k, \eta_f) . \end{aligned} \quad (58)$$

Plugging the explicit solutions of  $A_{\pm}(k, \eta)$  into Eq.(58) and after a little bit of simplification, we get the following forms of  $c_{\pm}$  and  $d_{\pm}$  as,

$$\begin{aligned} c_+ &= -\frac{i}{2} \sqrt{\frac{k}{2}} e^{ik\eta_f} \left\{ 2C_2 {}_1F1 \left( 1 + \frac{iB}{2}, 2, -2ik\eta_f \right) - C_2 k\eta_f (2i - B) {}_1F1 \left( 2 + \frac{iB}{2}, 3, -2ik\eta_f \right) \right. \\ &\quad \left. + 2C_1 \left[ U \left( \frac{iB}{2}, 0, -2ik\eta_f \right) - U \left( \frac{iB}{2}, 1, -2ik\eta_f \right) \right] \right\} , \end{aligned}$$

$$\begin{aligned}
c_- &= -\frac{i}{2}\sqrt{\frac{k}{2}}e^{ik\eta_f}\left\{2C_4 {}_1F1\left(1-\frac{iB}{2}, 2, -2ik\eta_f\right) - C_4 k\eta_f(2i+B) {}_1F1\left(2-\frac{iB}{2}, 3, -2ik\eta_f\right)\right. \\
&\quad \left.+ 2C_3\left[U\left(-\frac{iB}{2}, 0, -2ik\eta_f\right) - U\left(-\frac{iB}{2}, 1, -2ik\eta_f\right)\right]\right\}, \\
d_+ &= \frac{i}{2}\sqrt{\frac{k}{2}}e^{ik\eta_f}\left\{2C_2 {}_1F1\left(1+\frac{iB}{2}, 2, -2ik\eta_f\right) + C_2 k\eta_f(2i+B) {}_1F1\left(1+\frac{iB}{2}, 3, -2ik\eta_f\right)\right. \\
&\quad \left.- 2C_1 U\left(\frac{iB}{2}, 1, -2ik\eta_f\right)\right\}, \\
d_- &= \frac{i}{2}\sqrt{\frac{k}{2}}e^{ik\eta_f}\left\{2C_4 {}_1F1\left(1-\frac{iB}{2}, 2, -2ik\eta_f\right) + C_4 k\eta_f(2i-B) {}_1F1\left(1-\frac{iB}{2}, 3, -2ik\eta_f\right)\right. \\
&\quad \left.- 2C_3 U\left(-\frac{iB}{2}, 1, -2ik\eta_f\right)\right\} \tag{59}
\end{aligned}$$

respectively, with  $C_i$  ( $i = 1, 2, 3, 4$ ) are obtained in Eq.(36). One of the important quantities in magnetogenesis scenario are the Bogoliubov coefficients which actually account the production of gauge particles from the quantum vacuum. In particular, due to the interaction of the EM field with the background FRW spacetime, the Hamiltonian of the EM field becomes explicit time dependent, which in turn makes the electromagnetic vacuum state explicit time dependent. Thus if the EM field starts from the vacuum state at distant past (namely the Bunch-Davies state which we consider in the present context), then it will not remain in vacuum at a later time. This corresponds to the production of gauge particles from the Bunch-Davies state and the Bogoliubov coefficients provide the number of produced particles at a certain time. More precisely, the Bogoliubov coefficients at time  $\eta$  relate the mode functions correspond to the vacuum state of distant past with the set of mode functions that lead to the instantaneous vacuum of the field at the  $\eta$ . Based on this, the Bogoliubov coefficients during the reheating epoch, in the present context, are determined as,

$$\begin{aligned}
\alpha_{\pm}(k, \eta) &= \sqrt{\frac{k}{2}} A_{\pm}^{(re)}(k, \eta) + \frac{i}{\sqrt{2k}} A'_{\pm}{}^{(re)}(k, \eta), \\
\beta_{\pm}(k, \eta) &= \sqrt{\frac{k}{2}} A_{\pm}^{(re)}(k, \eta) - \frac{i}{\sqrt{2k}} A'_{\pm}{}^{(re)}(k, \eta) \quad . \tag{60}
\end{aligned}$$

Eqs.(56) and (60) immediately result  $\alpha_{\pm}$  and  $\beta_{\pm}$  as,

$$\alpha_{\pm}(k, \eta) = c_{\pm} e^{-ik(\eta-\eta_f)} \quad , \quad \beta_{\pm}(k, \eta) = d_{\pm} e^{ik(\eta-\eta_f)} \tag{61}$$

which relate the Bogoliubov coefficients with the integration constants that appear in the solution of  $A_{\pm}^{(re)}(k, \eta)$ . The absolute value of  $\beta_{\pm}$  (i.e  $|\beta_{\pm}|$ ) during the reheating era seems to be explicit time independent and equal to that of at the end of inflation, i.e  $|\beta_{\pm}(k, \eta)| = |\beta_{\pm}(k, \eta_f)| = d_{\pm}$  (where  $\eta$  is any time during reheating), from Eq.(61). This is a direct consequence of the fact that the coupling function  $f(R, \mathcal{G})$  is considered to be zero in the reheating stage, due to which the EM action restores the conformal symmetry and thus the EM field equations in the background FRW spacetime become similar to that of in the Minkowski spacetime. As a result, the gauge production ceases to occur after the inflation, which is actually reflected through the time independency of  $|\beta_{\pm}|$ . Moreover Eq.(61) leads to  $c_{\pm} = \alpha_{\pm}(k, \eta_f)$  and  $d_{\pm} = \beta_{\pm}(k, \eta_f)$ , and thus from Eq.(56),

$$A_{\pm}^{(re)}(k, \eta) = \frac{1}{\sqrt{2k}} \left[ \alpha_{\pm}(k, \eta_f) e^{-ik(\eta-\eta_f)} + \beta_{\pm}(k, \eta_f) e^{ik(\eta-\eta_f)} \right] \quad . \tag{62}$$

Plugging the above solution of  $A_{\pm}^{(re)}(k, \eta)$  into Eq.(29) along with the relation  $|\alpha_{\pm}|^2 - |\beta_{\pm}|^2 = 1$  yield the magnetic and electric power spectra during the reheating phase as,

$$\begin{aligned}
\frac{\partial \rho(\vec{B})}{\partial \ln k} &= \frac{1}{2\pi^2} \sum_{r=+,-} \frac{k^5}{a^4} |A_r^{(re)}(k, \eta)|^2 \\
&= \frac{1}{2\pi^2} \sum_{r=+,-} \left( \frac{k^4}{a^4} \right) \left[ |\alpha_r(k, \eta_f)|^2 + |\beta_r(k, \eta_f)|^2 + 2|\alpha_r(k, \eta_f) \beta_r(k, \eta_f)| \cos \{ \theta_1^{(r)} - \theta_2^{(r)} - 2k(\eta - \eta_f) \} \right] \tag{63}
\end{aligned}$$

and

$$\begin{aligned} \frac{\partial \rho(\vec{E})}{\partial \ln k} &= \frac{1}{2\pi^2} \sum_{r=1,2} \frac{k^2}{a^4} |A_r^{(re)}(k, \eta)|^2 \\ &= \frac{1}{2\pi^2} \sum_{r=+,-} \left( \frac{k^4}{a^4} \right) \left[ |\alpha_r(k, \eta_f)|^2 + |\beta_r(k, \eta_f)|^2 - 2|\alpha_r(k, \eta_f) \beta_r(k, \eta_f)| \cos \{ \theta_1^{(r)} - \theta_2^{(r)} - 2k(\eta - \eta_f) \} \right] \end{aligned} \quad (64)$$

respectively, where  $\theta_1^{(r)} = \text{Arg} [\alpha_r(k, \eta_f)]$  and  $\theta_2^{(r)} = \text{Arg} [\beta_r(k, \eta_f)]$ . Consequently, the total electromagnetic power spectrum turns out to be,

$$\frac{\partial \rho_{em}}{\partial \ln k} = \frac{\partial \rho(\vec{B})}{\partial \ln k} + \frac{\partial \rho(\vec{E})}{\partial \ln k} = \frac{1}{\pi^2} \sum_{r=+,-} \left( \frac{k^4}{a^4} \right) \left[ |\alpha_r(k, \eta_f)|^2 + |\beta_r(k, \eta_f)|^2 \right]. \quad (65)$$

It may be observed from the above expressions that the total EM power spectrum redshifts by  $\frac{1}{a^4}$  with the expansion of the universe, however the individual magnetic and electric spectra do not evolve as  $\frac{1}{a^4}$  due to the presence of the time dependent factor  $k(\eta - \eta_f)$  within the cosine argument of Eqs.(63) and (64) respectively. Thus the comoving EM energy density seems to be conserved in the reheating era, which, once again, is a consequence of the fact that the EM action restores the conformal symmetry in the post-inflationary epoch. Furthermore Eqs.(62) and (30) lead to the helicity power spectrum in-between the end of inflation and the end of reheating as,

$$\begin{aligned} P^{(h)}(k, \eta) &= \left( \frac{k^3}{\pi^2 a^3} \right) \left\{ |\beta_+|^2 - |\beta_-|^2 + \sqrt{1 + |\beta_+|^2} |\beta_+| \cos \{ \theta_1^{(+)} - \theta_2^{(+)} - 2k(\eta - \eta_f) \} \right. \\ &\quad \left. - \sqrt{1 + |\beta_-|^2} |\beta_-| \cos \{ \theta_1^{(-)} - \theta_2^{(-)} - 2k(\eta - \eta_f) \} \right\}. \end{aligned} \quad (66)$$

Here we would like to mention that both the EM mode functions follow the same equation of motion (see Eq.(55)) in the reheating era, however during inflation, the conformal coupling  $f(R, \mathcal{G})$  differs the amplitude of  $A_+(k, \eta)$  compared to that of  $A_-(k, \eta)$ , and as a result,  $A_+(k, \eta_f) \neq A_-(k, \eta_f)$  and  $A'_+(k, \eta_f) \neq A'_-(k, \eta_f)$ , i.e positive and negative helicity modes obey different initial conditions at the beginning of reheating (recall,  $\eta_f$  represents the end instance of inflation or equivalently the start of reheating). Thus as a whole,  $A_{\pm}(k, \eta)$  have same equations of motion but different initial conditions during the reheating stage, in effect of which,  $A_+(k, \eta)$  and  $A_-(k, \eta)$  get different and makes a non-zero helicity during the same, as given in Eq.(66).

Following, we determine the explicit forms of Bogoliubov coefficients and the factor  $k(\eta - \eta_f)$  present in the expressions of various kinds of power spectra.

- **Determination of  $\alpha_{\pm}(k, \eta_f)$  and  $\beta_{\pm}(k, \eta_f)$ :** In view of Eqs.(58) and (61), we can express  $\alpha_r(k, \eta_f)$  and  $\beta_r(k, \eta_f)$  as,

$$\begin{aligned} \alpha_r(k, \eta_f) &= \sqrt{\frac{k}{2}} \left[ A_r(k, \eta_f) + i \left. \frac{dA_r}{d(k\eta)} \right|_{\eta_f} \right], \\ \beta_r(k, \eta_f) &= \sqrt{\frac{k}{2}} \left[ A_r(k, \eta_f) - i \left. \frac{dA_r}{d(k\eta)} \right|_{\eta_f} \right], \end{aligned} \quad (67)$$

with recall, the index  $r$  specifies the two polarization modes of the EM field. The  $A_r(k, \eta)$  and  $\frac{dA_r}{d(k\eta)}$  in the super-Hubble scale are given in Eqs.(37) and (38) respectively. Due to the presence of  $\ln(-k\eta)$  in the expression of  $\frac{dA_r}{d(k\eta)}$  and owing to the fact that inflation ends at  $\eta \rightarrow 0^-$ ,  $\frac{dA_r}{d(k\eta)}$  become larger than  $A_r(k, \eta_f)$  and thus the Bogoliubov coefficients from Eq.(67) can be written as,

$$\begin{aligned} \alpha_+(k, \eta_f) &\simeq i\sqrt{\frac{k}{2}} \left( \left. \frac{dA_+}{d(k\eta)} \right|_{\eta_f} \right) = i\sqrt{\frac{k}{2}} \left[ C_2 + \frac{C_1}{\Gamma(iB/2)} \{ \gamma + \ln(-2k\eta_f) + iB/2 \} \right], \\ \alpha_-(k, \eta_f) &\simeq i\sqrt{\frac{k}{2}} \left( \left. \frac{dA_-}{d(k\eta)} \right|_{\eta_f} \right) = i\sqrt{\frac{k}{2}} \left[ C_4 + \frac{C_3}{\Gamma(-iB/2)} \{ \gamma + \ln(-2k\eta_f) - iB/2 \} \right], \\ \beta_+(k, \eta_f) &\simeq -i\sqrt{\frac{k}{2}} \left( \left. \frac{dA_+}{d(k\eta)} \right|_{\eta_f} \right) = -i\sqrt{\frac{k}{2}} \left[ C_2 + \frac{C_1}{\Gamma(iB/2)} \{ \gamma + \ln(-2k\eta_f) + iB/2 \} \right], \end{aligned}$$

$$\beta_-(k, \eta_f) \simeq -i\sqrt{\frac{k}{2}} \left( \frac{dA_-}{d(k\eta)} \Big|_{\eta_f} \right) = -i\sqrt{\frac{k}{2}} \left[ C_4 + \frac{C_3}{\Gamma(-iB/2)} \{ \gamma + \ln(-2k\eta_f) - iB/2 \} \right], \quad (68)$$

where the constants  $C_i$  are determined in Eq.(36). Thereby the relative phase factors between  $\alpha_r(\eta_f)$  and  $\beta_r(\eta_f)$  seem to be nearly equal to  $\pi$ , in particular  $\text{Arg}[\alpha_r(k, \eta_f)\beta_r^*(k, \eta_f)] \simeq \pi$  (with  $*$  denotes the complex conjugate); and moreover the absolute value of the Bogoliubov coefficients at  $\eta = \eta_f$  (namely  $|\alpha_r(\eta_f)|^2$  and  $|\beta_r(\eta_f)|^2$ ) get much larger than unity. Later, by numerical calculation, we will show that such cases indeed hold true in the present context.

- **Determination of “ $\eta - \eta_f$ ” during reheating:** The factor  $k(\eta - \eta_f)$  in Eqs.(63) and (66) leads to the non-conventional dynamics of magnetic and helicity energy density. The constant equation of state during reheating dynamics results to this special term as,

$$k(\eta - \eta_f) = \frac{2k}{(3\omega_{\text{eff}} + 1)} \left[ \frac{1}{aH} - \frac{1}{a_f H_f} \right], \quad (69)$$

where we use  $\eta - \eta_f = \int_{a_f}^a \frac{da}{a^2 H}$  and the quantities with suffix ‘f’ represent the respective quantities at the end of inflation.

With the above expressions of  $\alpha_r(k, \eta_f)$ ,  $\beta_r(k, \eta_f)$  and  $k(\eta - \eta_f)$ : the magnetic, electric and helicity power spectra during the reheating era (from Eqs.(63), (64) and (66)) turn out to be,

$$\frac{\partial \rho(\vec{B})}{\partial \ln k} = \frac{1}{\pi^2} \left( \frac{k^4}{a^4} \right) \sum_{r=+,-} |\beta_r(k, \eta_f)|^2 \left\{ \frac{4k}{3\omega_{\text{eff}} + 1} \left( \frac{1}{aH} - \frac{1}{a_f H_f} \right) \right\}^2, \quad (70)$$

$$\frac{\partial \rho(\vec{E})}{\partial \ln k} = \frac{1}{\pi^2} \left( \frac{k^4}{a^4} \right) \sum_{r=+,-} |\beta_r(k, \eta_f)|^2, \quad (71)$$

and

$$\frac{\partial \rho_h}{\partial \ln k} = \frac{1}{\pi^2} \left( \frac{k^3}{a^3} \right) \left( |\beta_+(k, \eta_f)|^2 - |\beta_-(k, \eta_f)|^2 \right) \left\{ \frac{4k}{3\omega_{\text{eff}} + 1} \left( \frac{1}{aH} - \frac{1}{a_f H_f} \right) \right\}^2, \quad (72)$$

respectively. Therefore the magnetic power spectrum during the reheating seems to be controlled by two terms: (1) the conventional one that redshifts by  $1/a^4$  with the expansion of the universe and (2) the term which is proportional to  $(a^3 H)^{-2}$  emerged from  $(\frac{k}{aH})^2$  present in the expression of Eq.(70). Since the universe, in the reheating phase, undergoes through a deceleration, we can safely consider  $\omega_{\text{eff}} > -\frac{1}{3}$  and thus the term proportional to  $(a^3 H)^{-2}$  becomes the dominant contribution in the magnetic power spectrum. In regard to the helicity power spectrum ( $\mathcal{P}_h$ ) during reheating dynamics, Eq.(72) reveals that, similar to the magnetic spectrum,  $\mathcal{P}_h$  is also controlled by two terms: (1) the term that evolves by  $1/a^3$  and (2) the other one which is proportional to  $1/(a^5 H^2)$ . However due to the constant equation of state, the Hubble parameter during the reheating behaves (in terms of scale factor) as  $H \propto a^{-\frac{3}{2}(1+\omega_{\text{eff}})}$  and thus the term  $1/(a^5 H^2)$  is further proportional to  $\propto a^{-2+3\omega_{\text{eff}}}$ . Therefore, due to  $\omega_{\text{eff}} > -1/3$ , the term  $1/(a^5 H^2) \propto a^{-2+3\omega_{\text{eff}}}$  proves to be the dominant piece (compared to the other one that goes by  $1/a^3$ ) in the helicity power spectrum of the reheating stage. Such special terms in the magnetic as well as in the helicity power spectra provide the main difference in the magnetogenesis scenario between the instantaneous reheating case and the case where the reheating phase has a non-zero e-fold number. For the magnetic power spectrum ( $\mathcal{P}(\vec{B})$ ): in the instantaneous reheating case, it evolves by  $\frac{1}{a^4}$  from the very end of inflation to the present epoch; however in the case of a non-zero e-fold reheating phase,  $\mathcal{P}(\vec{B})$  is controlled by  $1/(a^6 H^2)$  in-between the end of inflation to the end of reheating and then goes as  $1/a^4$  from the end of reheating to the present epoch. On other hand, for the helicity power spectrum: in the instantaneous reheating case, the comoving helicity remains conserved (i.e  $\mathcal{P}_h \propto 1/a^3$ ) after inflation and until today; however the evolution of  $\mathcal{P}_h$  becomes different when the reheating phase is considered to have a non-zero e-fold number, in particular,  $\mathcal{P}_h \propto 1/(a^5 H^2)$  from the end of inflation to the end of reheating and then  $\mathcal{P}_h \propto 1/a^3$  until the present epoch. Thereby the presence of a reheating phase results to an enhancement in the magnetic field’s current amplitude as well as in the present helicity strength, compared to that of in the instantaneous reheating case. Finally we would like to mention that unlike to the magnetic or helicity power spectra, the electric one decays by  $1/a^4$  during the reheating era, as evident from Eq.(71).



## B. Current magnetic strength and baryon asymmetry of the universe: Constraints on $\omega_{\text{eff}}$

The presence of a reheating phase with non-zero e-fold number results to the existence of an electric field in the post-inflationary evolution of the universe till the time when the universe becomes purely conductive and this generally occurs at the end of reheating. Such electric field in turn induces the magnetic field according to the Faraday's law of induction [43], due to which, the magnetic field energy density in the reheating epoch redshifts slower than that of in the case of instantaneous reheating. Therefore it is important to investigate whether the presence of a reheating phase having non-zero e-fold yields a sufficient amount of magnetic strength and consequently the helicity at the present time. We will investigate this in the present subsection. Owing to the *constant* equation of state in the Kamionkowski reheating dynamics (symbolized by  $\omega_{\text{eff}}$ ), the Hubble parameter during the reheating phase evolves as  $H \propto a^{-\frac{3}{2}(1+\omega_{\text{eff}})}$ , and thus the Hubble parameter at the end of reheating is connected to that of at the end of inflation as,

$$H_{re} = H_f \left( \frac{a_{re}}{a_f} \right)^{-\frac{3}{2}(1+\omega_{eff})}, \quad (73)$$

where the suffix 're' denotes the end instance of reheating. Therefore  $\ln\left(\frac{a_{re}}{a_f}\right) = N_{re}$  represents the e-fold number of the reheating era, the  $N_{re}$  in terms of the reheating parameters is given in Eq.(53). The above expression of  $H_{re}$  immediately leads to the magnetic and helicity power spectra (from Eqs.(70) and (72)) at the end of reheating as,

$$\left. \frac{\partial \rho(\vec{B})}{\partial \ln k} \right|_{re} = \frac{1}{\pi^2} \left( \frac{k^4}{a_{re}^4} \right) \sum_{r=+,-} |\beta_r(k, \eta_f)|^2 \left\{ \frac{4}{3\omega_{\text{eff}} + 1} \left( \frac{k}{a_f H_f} \right) \left[ \left( \frac{H_f}{H_{re}} \right)^{\frac{3\omega_{\text{eff}}+1}{3\omega_{\text{eff}}+3}} - 1 \right] \right\}^2. \quad (74)$$

and

$$\left. \frac{\partial \rho_h}{\partial \ln k} \right|_{re} = \frac{1}{\pi^2} \left( \frac{k^3}{a_{re}^3} \right) (|\beta_+(k, \eta_f)|^2 - |\beta_-(k, \eta_f)|^2) \left\{ \frac{4}{3\omega_{\text{eff}} + 1} \left( \frac{k}{a_f H_f} \right) \left[ \left( \frac{H_f}{H_{re}} \right)^{\frac{3\omega_{\text{eff}}+1}{3\omega_{\text{eff}}+3}} - 1 \right] \right\}^2. \quad (75)$$

respectively. After the end of reheating, the universe becomes a good conductor, which in turn makes the electric field zero. Moreover due to the conformal invariance, the comoving EM energy density remains conserved as the universe expands, or equivalently the magnetic energy density evolves by  $1/a^4$ , as the electric field is vanishingly small after the end of reheating. Consequently the helicity power spectrum decays by  $1/a^3$  (i.e the comoving helicity is conserved) during the post-reheating era. Hence, the present magnetic and the present helicity power spectra is connected to that of at the end of reheating by following:

$$\left. \frac{\partial \rho(\vec{B})}{\partial \ln k} \right|_0 = \left( \frac{a_{re}}{a_0} \right)^4 \left. \frac{\partial \rho(\vec{B})}{\partial \ln k} \right|_{re} \quad \text{and} \quad \left. \frac{\partial \rho_h}{\partial \ln k} \right|_0 = \left( \frac{a_{re}}{a_0} \right)^3 \left. \frac{\partial \rho_h}{\partial \ln k} \right|_{re}, \quad (76)$$

where the suffix '0' denotes the present time of the universe. Using the above expression along with Eqs.(74) and (75), we get the magnetic field's current amplitude and the helicity power spectrum at present epoch as,

$$B_0 = \frac{\sqrt{2}}{\pi} \left( \frac{k}{a_0} \right)^2 \sqrt{|\beta_+(k, \eta_f)|^2 + |\beta_-(k, \eta_f)|^2} \left\{ \frac{4}{3\omega_{\text{eff}} + 1} \left( \frac{k}{a_f H_f} \right) \left[ \left( \frac{H_f}{H_{re}} \right)^{\frac{3\omega_{\text{eff}}+1}{3\omega_{\text{eff}}+3}} - 1 \right] \right\}, \quad (77)$$

and

$$\left. \frac{\partial \rho_h}{\partial \ln k} \right|_0 = \frac{1}{\pi^2} \left( \frac{k^3}{a_0^3} \right) (|\beta_+(k, \eta_f)|^2 - |\beta_-(k, \eta_f)|^2) \left\{ \frac{4}{3\omega_{\text{eff}} + 1} \left( \frac{k}{a_f H_f} \right) \left[ \left( \frac{H_f}{H_{re}} \right)^{\frac{3\omega_{\text{eff}}+1}{3\omega_{\text{eff}}+3}} - 1 \right] \right\}^2, \quad (78)$$

where in Eq.(77), we use  $\left. \frac{\partial \rho(\vec{B})}{\partial \ln k} \right|_0 = \frac{1}{2} B_0^2$ , with  $B_0$  being the present amplitude of the magnetic field. Here we need to recall that  $\left. \frac{\partial \rho_h}{\partial \ln k} \right|_0$  yields the net baryon density in present universe ( $n_B$ ). Therefore it may be observed that both the  $B_0$  and  $n_B$  depends on the reheating dynamics (through  $H_{re}$  and  $\omega_{\text{eff}}$ ) as well as on the background inflationary dynamics (through  $H_f$  and the absolute value of the Bogoliubov coefficients i.e  $|\beta_{\pm}(k, \eta_f)|^2$ ). As a result,

the magnetic field's current strength and the net baryon density of the present universe get a one-to-one correspondence with the reheating equation of state parameter. This in turn helps to probe the reheating dynamics, in particular, by confronting the theoretical predictions with the observational constraints given by  $10^{-22}\text{G} \lesssim B_0 \lesssim 10^{-10}\text{G}$  and  $\frac{n_B}{s_0} = (6.09 \pm 0.06) \times 10^{-10}$  respectively ( $s_0$  represents the entropy density of the present universe). Plugging the expressions of  $\beta_{\pm}(k, \eta_f)$  from Eq.(68) to Eq.(78) and after a bit of simplification, the helicity power spectrum turns out to be,

$$\begin{aligned} \left. \frac{\partial \rho_h}{\partial \ln k} \right|_0 = & - \frac{2}{\pi^2} \left( \frac{k^3}{a_0^3} \right) \sinh \left( \frac{\pi B}{2} \right) \left\{ \frac{4}{3\omega_{\text{eff}} + 1} \left( \frac{k}{a_f H_f} \right) \left[ \left( \frac{H_f}{H_{re}} \right)^{\frac{3\omega_{\text{eff}} + 1}{3\omega_{\text{eff}} + 3}} - 1 \right] \right\}^2 \\ & \times \left\{ [\gamma + \ln(-2k\eta_f)]^2 + \left( \frac{B}{2} \right)^2 - \left| \Gamma \left( \frac{iB}{2} \right) \Gamma \left( 1 + \frac{iB}{2} \right) \right|^2 \right\} . \end{aligned} \quad (79)$$

To derive the above expression, we use the forms of  $C_i$  ( $i = 1, 2, 3, 4$ ) given in Eq.(36). It may be observed from Eq.(79) that for  $B = 0$ , the helicity strength at present universe becomes zero; this is however expected because the quantity  $B$  is proportional to the parity violating parameter (see Eq.(33)) and thus for  $B = 0$  the EM action gets the parity symmetry of the EM field, due to which the helicity vanishes identically.

In regard to the generation of BAU from helical magnetic fields, the net baryonic number density ( $n_B$ ) is defined by [10, 120],

$$n_B = \frac{e^2}{4\pi^2} a(\eta) n_{\text{CS}} , \quad (80)$$

where  $n_{\text{CS}}$  is the net Chern-Simons number density stored in the helical magnetic fields and given by  $n_{\text{CS}} = \langle 0 | K^0 | 0 \rangle$ , with  $K^0 = \frac{1}{a^4(\eta)} \epsilon^{ijk} \partial_j A_k$ . Thereby, in Fourier space,  $n_B$  can be expressed as,

$$n_B = \frac{e^2}{4\pi^2} \int_{k_0}^{k_f} d \ln k \left( \frac{\partial \rho_h}{\partial \ln k} \right) , \quad (81)$$

where  $k_f$  and  $k_0$  are the momentum of the modes which re-enter the horizon at the end of inflation and at present epoch respectively, and moreover,  $k_f = 1/|\eta_f|$  (with  $\eta_f$  being the end instance of inflation) and  $k_0$  is around the CMB scale momentum. Therefore in deriving  $n_B$ , we consider such modes which are within the horizon at present time, in particular, ranging from  $k_f$  to  $k_0$ . As a result, we arrive at the following expression of  $n_B$ ,

$$\begin{aligned} n_B = & \frac{e^2}{10^3 \pi^4} \left( \frac{1}{a_0} \right)^3 \sinh \left( \frac{\pi B}{2} \right) \left\{ \frac{4}{3\omega_{\text{eff}} + 1} \left( \frac{1}{a_f H_f} \right) \left[ \left( \frac{H_f}{H_{re}} \right)^{\frac{3\omega_{\text{eff}} + 1}{3\omega_{\text{eff}} + 3}} - 1 \right] \right\}^2 \times \left[ k^5 \left\{ 8 + 25B^2 + 20\gamma(-2 + 5\gamma) \right. \right. \\ & \left. \left. - 100\pi^2 \text{csch}^2 \left( \frac{B}{10\pi} \right) + 20 \ln 2(-2 + 10\gamma + 5 \ln 2) + 20 \ln(-k\eta_f)(-2 + 10\gamma + 5 \ln(-4k\eta_f)) \right\} \right] \Big|_{k_f}^{k_0} , \end{aligned} \quad (82)$$

with, recall that  $\gamma$  is the Euler's constant  $\simeq 0.577216$ . Moreover  $B$ , in terms of the model parameters  $\lambda$  and  $q$ , is given in Eq.(33). The model parameters are considered in such a way that the  $k$  independent terms within the curly bracket of the above expression vanishes, in particular,

$$8 + 25B^2 + 20\gamma(-2 + 5\gamma) - 100\pi^2 \text{csch}^2 \left( \frac{B}{10\pi} \right) + 20 \ln 2(-2 + 10\gamma + 5 \ln 2) = 0 . \quad (83)$$

Such consideration has a phenomenological motivation from the perspective of the observational compatibility of the model; we will show that the condition in Eq.(83) indeed leads to the compatibility between the theoretical predictions of the model and the observational results, which in turn argues the phenomenological viability of the above condition. Furthermore, Eq.(83), seems to suggest that the net baryonic number density gets its main contribution from large scale modes of the magnetic field. Eq.(83) has the solution  $B \simeq 13.74$ , due to which, Eq.(33) results to the following constraint on the model parameters as,

$$(16\epsilon q \lambda) \left( \frac{H_0}{M_{\text{Pl}}} \right)^{2q} \left\{ [6\beta(\beta + 1)]^q + [-24(\beta + 1)^3]^{q/2} \right\} = 13.74 , \quad (84)$$

where we need to recall that  $\lambda$  is the conformal breaking parameter,  $q$  appears in the form of  $f(R, \mathcal{G})$ ,  $\epsilon$  is the slow roll parameter and  $\eta_0^{-1} = H_0$ . In order to get the allowed ranges of  $\lambda$  and  $q$ , we give the plots of  $\lambda$  vs  $q$  from the constraint Eq.(84), in Fig.[1] where  $\epsilon$  acquires five different values from  $\epsilon = 0.01$  (the uppermost plot) to  $\epsilon = 0.1$  (the lowermost plot) and  $H_0 = 10^{-5} M_{\text{Pl}}$ . Therefore for any values of  $0.01 < \epsilon < 0.1$ , the respective plot of  $\lambda$  vs  $q$  lies in-between the uppermost and the lowermost plots. The figure clearly demonstrates that the model parameters in the present context (i.e  $\lambda$ ,  $q$ ) are allowed to have a wide range of values from the constraint relation (84). Now, we investigate whether such ranges of the parameters lead to the theoretical consistency of  $B_0$  and  $n_B$  with the observational results.




FIG. 1:  $\lambda$  vs  $q$  for various values of  $\epsilon$ , from Eq.(84). From above to below:  $\epsilon = 0.01, 0.02, 0.03, 0.05, 0.1$  respectively and  $H_0 = 10^{-5} M_{\text{Pl}}$ . Moreover for any values of  $0.01 < \epsilon < 0.1$ , the respective plot of  $\lambda$  vs  $q$  lies in-between the uppermost and the lowermost plots. This clearly justifies that the model parameters are allowed to have a wide range of values from the constraint relation (84).

However, before doing so, let us rewrite the net baryon density using Eqs.(82) and (83),

$$n_B = \frac{e^2}{10^3 \pi^4} \left( \frac{k_0}{a_0} \right)^3 \sinh \left( \frac{\pi B}{2} \right) \left\{ \frac{4}{3\omega_{\text{eff}} + 1} \left( \frac{k_0}{a_f H_f} \right) \left[ \left( \frac{H_f}{H_{\text{re}}} \right)^{\frac{3\omega_{\text{eff}} + 1}{3\omega_{\text{eff}} + 3}} - 1 \right] \right\}^2 \times \left[ 20 \ln(-k_0 \eta_f) \left( -2 + 10\gamma + 5 \ln(-4k_0 \eta_f) \right) \right], \quad (85)$$

where we use  $k_f \eta_f = -1$ , and consequently  $\ln(-k_f \eta_f) = 0$ . It will be more useful if we express the net baryon density in terms of the magnetic power spectrum by using Eq.(77), and is given by,

$$n_B = \frac{e^2}{10^3 \pi^2} \left( \frac{a_0}{k_0} \right) \sinh \left( \frac{\pi B}{2} \right) \left( \frac{\partial \rho(\vec{B})}{\partial \ln k} \right)_0 \times \left[ \frac{20 \ln(-k_0 \eta_f) \left( -2 + 10\gamma + 5 \ln(-4k_0 \eta_f) \right)}{|\beta_+(k_0, \eta_f)|^2 + |\beta_-(k_0, \eta_f)|^2} \right]. \quad (86)$$

Thus as a whole, Eqs.(77) and (86) are the key equations to determine the magnetic field's current strength and the net baryon density of the universe. Having obtained such expressions, we now confront the model with the CMB observations which put a constraint on the present magnetic strength as  $10^{-22} \text{G} \lesssim B_0 \lesssim 10^{-10} \text{G}$ . In regard to this, we need  $h_f/H_{\text{re}}$  which in turn depends on  $N_{\text{re}}$  due to the Eq.(73). The reheating e-fold number ( $N_{\text{re}}$ ) is given in Eq.(53) pointing the fact that the  $N_{\text{re}}$  depends on  $H_0$  and  $H_f$ , i.e on the cosmic Hubble parameter at the beginning and at the end of inflation respectively. Here we consider  $H_0 = 10^{-5} M_{\text{Pl}}$ , and due to Eq.(14), i.e  $H = H_0 \exp\left(-\frac{\epsilon N}{1+\epsilon}\right)$ , the quantity  $H_f$  is estimated as  $H_f = 5 \times 10^{-8} M_{\text{Pl}}$  for  $\epsilon = 0.1$  and  $N_f = 58$ . Consequently, the model parameters ( $\lambda$  and  $q$ ) acquire the values according to the ranges demonstrated in Fig.[1], in particular for  $\epsilon = 0.1$ ,  $\lambda$  and  $q$  lie on the lowermost plot of Fig.[1]. With such considerations, we plot the current magnetic strength ( $B_0$ ) vs the reheating equation of state parameter ( $\omega_{\text{eff}}$ ) by using Eq.(77), in Fig.[2]. Furthermore, such numerical values of these quantities lead to the absolute value of the Bogoliubov coefficients at  $\eta = \eta_f$  as much larger than unity, namely  $|\beta_{\pm}(\eta_f)|^2 \gg 1$ .

The Fig.[2] clearly demonstrates that  $B_0$  lies within the observational constraints if the reheating equation of state parameter situates within  $0.284 \lesssim \omega_{\text{eff}} \lesssim 0.301$ . In the upper x-axis of the figure, we put the reheating temperature

( $T_{\text{re}}$ ) for different values of  $\omega_{\text{eff}}$  by using Eq.(54), which, due to the BBN constraint is limited by  $T_{\text{re}} \lesssim 10^{-2}\text{GeV}$ . Such constraint on  $T_{\text{re}}$  in turn sets a lower range of  $\omega_{\text{eff}}$  given by  $0.284 \lesssim \omega_{\text{eff}} \lesssim 0.2975$ , and thus the remaining range of  $\omega_{\text{eff}}$  becomes disfavoured which is shown by the shaded region in the figure. Thus as a whole, in order to satisfy both the CMB observations on  $B_0$  and the BBN constraint on  $T_{\text{re}}$ , the possible regime of  $\omega_{\text{eff}}$  seems to be:  $0.284 \lesssim \omega_{\text{eff}} \lesssim 0.2975$ .




FIG. 2:  $B_0$  vs  $\omega_{\text{eff}}$  for  $\frac{k_0}{a_0} = 0.02\text{Mpc}^{-1}$ . The reheating temperature ( $T_{\text{re}}$ ) for different values of  $\omega_{\text{eff}}$  is shown in the upper x-axis.

In regard to estimate the net baryon density, we will use Eq.(86) where  $n_{\text{B}}$  has been expressed in terms of the present magnetic power spectrum. However it is more convenient to get the ratio of net baryon density to the entropy density of the present universe, i.e  $n_{\text{B}}/s_0$ , which is dimensionless and has observational constraints as  $n_{\text{B}}/s_0 = (6.09 \pm 0.06) \times 10^{-10}$  from the CMB data. The current entropy density ( $s_0$ ) is given by,

$$s_0 = 2.97 \times 10^3 \left( \frac{T_0}{2.75 [K]} \right)^3 \text{ cm}^{-3} , \quad (87)$$

with  $T_0 = 2.75\text{K}$  is the present temperature of the cosmic microwave background (CMB) radiation. Fig.[2] depicts that the model predicts sufficient magnetic strength for a suitable regime of the reheating EoS parameter. Using this information along with the parameter values that we considered earlier (i.e  $H_0 = 10^{-5}M_{\text{Pl}}$ ,  $\epsilon = 0.1$ ,  $N_{\text{f}} = 58$  and  $H_f = 5 \times 10^{-8}M_{\text{Pl}}$ : recall, they are connected by Eq.(14)), we estimate  $n_{\text{B}}/s_0$  for different values of  $B_0$ , with  $\frac{k_0}{a_0} = 0.02\text{Mpc}^{-1}$ . These are given in Table[I] where the corresponding values of  $\omega_{\text{eff}}$  are incorporated. From Table[I], we see that the magnetic fields of current strength  $\sim 10^{-18}\text{G}$  leads to the resultant value of  $n_{\text{B}}/s_0$  as of the order  $\sim 10^{-10}$ , which is indeed consistent with the observational constraints. Therefore, in the present context where the EM field couples with the higher curvature term(s) via its dual field tensor, the net baryon density induced by the helical magnetic fields seems to be compatible with the CMB observations.

Current magnetic field: $B_0$ [Gauss]	Reheating EoS parameter: $\omega_{\text{eff}}$	$n_{\text{B}}/s_0$ (Dimensionless)
$3.084 \times 10^{-18}$	$\approx 0.2915$	$6.09 \times 10^{-10}$
$3.074 \times 10^{-18}$	$\approx 0.2915$	$6.05 \times 10^{-10}$
$3.092 \times 10^{-18}$	$\approx 0.2916$	$6.12 \times 10^{-10}$
$3.097 \times 10^{-18}$	$\approx 0.2916$	$6.14 \times 10^{-10}$

TABLE I: The ratio of net baryon density to the present entropy density (i.e  $n_{\text{B}}/s_0$ ) for different magnetic strength  $B_0$  with  $\frac{k_0}{a_0} = 0.02\text{Mpc}^{-1}$ . The corresponding values of  $\omega_{\text{eff}}$  are also shown in the table.

Thus as a whole, the presence of the reheating phase with non-zero e-fold number enhances the strength of magnetic field and the helicity as well, compared to that of in the instantaneous reheating scenario; by which, the magnetic field's current amplitude and consequently the net baryonic number density get consistent with the observational results when the universe undergoes through a reheating era after the end of inflation.

## VII. SCHWINGER BACKREACTION AND NECESSARY CONSTRAINTS

In Sec.[VI], the universe during the reheating era is considered to be a bad conductor, in particular we take the electrical conductivity to be zero, and thus the EM field satisfies the Maxwell's equations in vacuum. However the presence of strong electric field induces the Schwinger production during the reheating stage, which in turn backreacts the magnetogenesis scenario [43, 121, 122]. Therefore due to the Schwinger backreaction, the assumption of zero conductivity needs a proper justification, which we intend to perform in the present section. For this purpose, we start with a non-zero value of the electrical conductivity and investigate the condition for negligible Schwinger backreaction.

The Ampere-Maxwell equation of the EM field with a non-zero conductivity (symbolized by  $\sigma$ ) reads as,

$$\epsilon_{ijl}\partial_j B_l = E'_i + \left( a\sigma + \frac{a'}{a} \right) E_i \quad (88)$$

where  $\epsilon_{ijl}$  is the 3-Levi Civita index. Therefore the condition by which the Schwinger backreaction can be neglected is,

$$|a\sigma| < \frac{a'}{a} = aH \quad \text{or,} \quad |\sigma| < H \quad . \quad (89)$$

In progressing further, one needs a certain form of  $\sigma$  in the background of reheating dynamics where the Hubble parameter behaves as  $H \propto a^{-\frac{3}{2}(1+\omega_{\text{eff}})}$ . A complete analysis of  $\sigma$  in a generic FRW spacetime is beyond the scope of this paper. However the electrical conductivity in FRW background can be obtained from that of in the Minkowski background by replacing the elapsed time of the electric field in the Minkowski spacetime to  $H^{-1}$  in FRW spacetime, as explained in [43]. As a consequence, the conductivity in FRW spacetime becomes,

$$\sigma = \frac{1}{12\pi^3} \frac{|e^3 E|}{H} \exp \left[ -\frac{\pi m^2}{|eE|} \right] \quad (90)$$

where  $m$  and  $e$  are the mass and the electrical charge of the charged particles respectively produced by the Schwinger effect. It may be mentioned that the form of conductivity in Eq.(90) is also valid in a de-Sitter background spacetime under a strong electric field approximation [121]. Solving Eq.(90) for the electric field,

$$\frac{|E|}{H^2} = \frac{12\pi^3}{e^3} \left( \frac{\sigma}{H} \right) \exp \left\{ W \left( \frac{e^2}{12\pi^2} \frac{m^2 H}{\sigma} \right) \right\} \quad (91)$$

where  $W(x)$  is the Lambert W-function and the solution of  $We^W = x$ . For low massive charged particles i.e for  $m^2 \ll H^2$ , the W-function can be approximated as  $W(0 < x \ll 1) \approx x$  and thus the above equation can be expressed as,

$$\frac{|E|}{H^2} = \frac{12\pi^3}{e^3} \left( \frac{\sigma}{H} \right) \quad . \quad (92)$$

Thereby the above expression translates the condition of Eq.(89) to an upper bound of the electric field during the reheating phase as,

$$\frac{|E|}{12\pi^3 H^2} < \frac{1}{e^3} \approx 1 \quad . \quad (93)$$

Now, Eq.(71) evidents that the electric field during the reheating depends on  $|\beta_{\pm}(\eta_f)|^2$  which are obtained in Eq.(68). Moreover the Hubble parameter in the reheating dynamics evolves as,

$$H^2 = H_f^2 \left( \frac{a}{a_f} \right)^{-3(1+\omega_{\text{eff}})} \quad , \quad (94)$$

where recall,  $H_f$  is the Hubble parameter at the end of inflation and  $\omega_{\text{eff}}$  is the effective EoS parameter during the reheating phase. Due to such expressions of  $|E|$  and  $H^2$ , the left hand side of Eq.(93) turns out to be,

$$\left( \frac{\sqrt{2}}{12\pi^3} \right) \sqrt{|\beta_+(\eta_f)|^2 + |\beta_-(\eta_f)|^2} \left( \frac{a}{a_f} \right)^{1+3\omega_{\text{eff}}} e^{-2N_f} < 1 \quad , \quad (95)$$

with  $N_f$  being the e-fold number of the inflationary era. The left hand side of the above inequality seems to increase with the scale factor; thus in order to investigate the inequality during the entire reheating epoch, we put  $a = a_{re}$  in Eq.(95), i.e evaluating at the end of reheating, and we get,

$$\mathcal{M} = \left( \frac{\sqrt{2}}{12\pi^3} \right) \sqrt{|\beta_+(\eta_f)|^2 + |\beta_-(\eta_f)|^2} \left( \frac{a_{re}}{a_f} \right)^{1+3\omega_{eff}} e^{-2N_f} , \quad (96)$$

which is the maximum value that can be achieved by the left hand side of Eq.(95) during the reheating phase. The factor  $\frac{a_{re}}{a_f}$  is actually related to the reheating e-fold number by  $\frac{a_{re}}{a_f} = e^{N_{re}}$ , and thus the quantity  $\mathcal{M}$  can be expressed as,

$$\mathcal{M} = \left( \frac{\sqrt{2}}{12\pi^3} \right) \left( \sqrt{|\beta_+(\eta_f)|^2 + |\beta_-(\eta_f)|^2} \right) \exp \{ -2N_f + (1 + 3\omega_{eff}) N_{re} \} . \quad (97)$$

The reheating e-fold number, in terms of  $\omega_{eff}$  and  $N_f$ , is given in Eq.(53), by plugging which into Eq.(97) yields the final form of  $\mathcal{M}$ :

$$\mathcal{M} = \left( \frac{\sqrt{2}}{12\pi^3} \right) \left( \sqrt{|\beta_+(\eta_f)|^2 + |\beta_-(\eta_f)|^2} \right) \exp \left\{ - \left( \frac{6 + 6\omega_{eff}}{1 - 3\omega_{eff}} \right) N_f - \frac{4(1 + 3\omega_{eff})}{(1 - 3\omega_{eff})} \left[ \frac{1}{4} \ln \left( \frac{45}{\pi^2 g_{re}} \right) + \frac{1}{3} \ln \left( \frac{11g_{s,re}}{43} \right) + \ln \left( \frac{k}{a_0 T_0} \right) + \ln \left( \frac{(3H_f^2 M_{Pl}^2)^{1/4}}{H_0} \right) \right] \right\} \quad (98)$$

Eq.(98) clearly indicates that the quantity  $\mathcal{M}$  depends on the inflationary parameters like  $N_f$ ,  $|\beta_{\pm}(\eta_f)|^2$  as well as on the reheating parameters like  $\omega_{eff}$ . Here we need to recall the set of parametric values that we considered earlier, in particular:  $H_0 = 10^{-5} M_{Pl}$ ,  $\epsilon = 0.1$ ,  $N_f = 58$  respectively, for which  $H_f = 5 \times 10^{-8} M_{Pl}$  due to the relation  $H = H_0 \exp \left( -\frac{\epsilon N}{1+\epsilon} \right)$ , and moreover, the conformal breaking parameters  $\lambda$  and  $q$  are shown in Fig.[1]. With such set of values of the parameters, along with the expression of  $|\beta_{\pm}(\eta_f)|^2$  (see Eq.(68)), we give a plot of  $\mathcal{M}$  vs  $\omega_{eff}$  in Fig.[3].




FIG. 3:  $\mathcal{M}$  vs  $\omega_{eff}$ , with  $g_{re} = g_{s,re} = 100$  and  $T_0 = 2.3 \times 10^{-13} \text{GeV}$ .

Fig.[3] clearly demonstrates that within  $0.284 \lesssim \omega_{eff} \lesssim 0.2975$ , which makes the magnetic field's present strength ( $B_0$ ) and consequently the net baryonic density ( $n_B$ ) compatible with the CMB observations (see Fig.[2]), the quantity  $\mathcal{M}$  lies well below than unity. This in turn confirms the inequality of Eq.(93) during the entire reheating epoch. Therefore we may argue that in the regime  $0.284 \lesssim \omega_{eff} \lesssim 0.2975$ , the Schwinger backreaction or equivalently the electrical conductivity in the reheating era can be neglected and thus the EM field evolution obeys the standard Maxwell's equations in vacuum. Thus the viable regime of the reheating EoS parameter is given by

$0.284 \lesssim \omega_{\text{eff}} \lesssim 0.2975$ , which makes the model compatible with the CMB observations of  $B_0$  and  $n_B$ , and also leads to a negligible Schwinger backreaction.

Before concluding, we would like to mention that the production of the gauge field in the present magnetogenesis scenario may source the curvature perturbation in the super Hubble scales [123–127]. The calculation of such induced curvature power spectrum and its consistency with the Planck constraints will be considered elsewhere.

### VIII. CONCLUSION

We provide a viable helical magnetogenesis scenario from inflation and explain the baryon asymmetry of the universe. The electromagnetic (EM) field couples with the background higher curvature term(s), in particular with the Ricci scalar and the Gauss-Bonnet (GB) invariant, via the dual field tensor. Such non-minimal coupling of the EM field breaks the conformal and the parity symmetry, however preserves the  $U(1)$  invariance, of the electromagnetic action. As a result, the positive and negative helicity modes get different amplitudes along their cosmological evolution, and thus the magnetic field gets helical in nature, which is further connected to the net baryonic number density of the universe. The background spacetime is governed by the well studied scalar-Einstein-Gauss-Bonnet gravity theory which is known to lead to a viable inflationary scenario for suitable choices of scalar field potential and the GB coupling function. The model has some good features: first – during the early universe when the curvature is large, the conformal breaking coupling remains significant and results to a non-trivial contribution to the EM field equations, however during the late times, the curvature becomes low, and as a consequence, the conformal breaking term does not actually contribute, particularly from the end of inflation. Second – due to the fact that the non-minimal coupling of the EM field occurs via the dual field tensor, the kinetic term of the EM field remains canonical even in presence of such non-minimal coupling. This leads to the resolution of strong coupling problem in the present model. As a third strong point of our model – the EM field is found to have a negligible backreaction on the background spacetime, thus the background inflationary evolution does not harm by the presence of the EM field and consequently the backreaction issue is resolved in the model.

In such scenario, we explore the cosmological evolution of electric and magnetic fields, starting from the inflationary era to the present epoch. Consequently we address the helicity power spectrum, that arises due to the difference in the amplitudes between the positive and negative helicity modes of the EM field. During the evolution, the universe enters to a reheating era after the end of inflation and depending on the reheating dynamics, we consider two different cases: (1) the instantaneous reheating case where the universe instantaneously jumps to the radiation dominated epoch as the inflation ends, i.e the e-fold number of the instantaneous reheating era is zero. (2) As a second case, the reheating phase is considered to have a non-zero e-fold number, in particular, we consider the conventional reheating mechanism proposed by Kamionkowski et al. [105], where the reheating phase is parametrized by a non-zero e-fold number ( $N_{\text{re}}$ ), a constant equation of state parameter ( $\omega_{\text{eff}}$ ) and a given reheating temperature ( $T_{\text{re}}$ ).

In the case of instantaneous reheating, as the universe suddenly jumps from inflation to radiation era, the conductivity of the universe becomes huge in the post-inflationary stage and thus the electric field dies out quickly. Moreover, the EM action restores the conformal symmetry after inflation, the EM energy density evolves as  $a^{-4}$  with the expansion of the universe, or equivalently the magnetic energy density redshifts as  $a^{-4}$  as the electric field practically vanishes. Consequently the helicity power spectrum goes by  $a^{-3}$ , i.e the comoving helicity seems to be conserved with cosmic time in the post-inflation phase. As a result, the theoretical predictions for the magnetic field's current strength ( $B_0$ ) as well as the net baryonic number density of the universe ( $n_B$ ) are found to lie far below than the CMB observational constraints which are given by:  $10^{-22}\text{G} \lesssim B_0 \lesssim 10^{-10}\text{G}$  and  $\frac{n_B}{s_0} = (6.09 \pm 0.06) \times 10^{-10}$  respectively (with  $s_0$  being the entropy density of the present universe).

However the evolution of electric and magnetic fields become completely different when the reheating phase is considered to have a non-zero e-fold number. As mentioned, we consider a Kamionkowski like reheating mechanism where, due to the constant EoS parameter symbolized by  $\omega_{\text{eff}}$ , the Hubble parameter during the reheating era is given by  $H \propto a^{-\frac{3}{2}(1+\omega_{\text{eff}})}$ . Moreover the electrical conductivity of the universe during the reheating is taken to be zero, due to which, the existence of electric field in turn induces the magnetic field. Then, after the end of reheating, the universe enters to a radiation era and the conductivity gets a huge value, which in turn makes the electric field practically zero. Specifically, in regard to the magnetic power spectrum – it evolves by  $1/(a^6 H^2)$  during the reheating stage, and after that, as  $1/a^4$  till the present epoch. On other hand, the helicity power spectrum goes as  $1/(a^5 H^2)$  in the reheating era, and then, as  $1/a^3$  from the end of reheating to the present time. Based on these arguments, we argue that the presence of a reheating phase, with non-vanishing e-fold number, leads to an enhancement in the current magnetic strength as well as in the helicity power spectrum of the present universe, compared to that of in the instantaneous reheating case where the magnetic energy density and the helicity spectrum decay by  $a^{-4}$  and  $a^{-3}$  respectively from the very end of inflation.

As a result, in the case where the reheating era have a non-zero number of e-folds, the current magnetic strength ( $B_0$ ) and the net baryon density ( $n_B$ ) are found to be compatible with the observational constraints, for a suitable regime of reheating parameters. In particular, we get  $0.284 \lesssim \omega_{\text{eff}} \lesssim 0.2975$ , in order to make compatible the theoretical predictions of the present model with the CMB observations of  $B_0$  and  $n_B$  respectively. Such range of the reheating EoS parameter also seems to be consistent with the BBN constraint of the reheating temperature. Finally we would like to mention that due to the assumption of zero conductivity during the reheating era, we investigate the Schwinger backreaction issue in the present context. It turns out that the range  $0.284 \lesssim \omega_{\text{eff}} \lesssim 0.2975$ , for which the model becomes compatible with the observational data of  $B_0$  and  $n_B$ , is indeed accordant with a negligible Schwinger backreaction to the EM field evolution.

### Acknowledgments

This work was supported by MINECO (Spain), project PID2019-104397GB-I00 (SDO). The work of KB was supported in part by the JSPS KAKENHI Grant Number JP21K03547.

- 
- [1] D. Grasso and H. R. Rubinstein, Phys. Rept. **348** (2001), 163-266 [arXiv:astro-ph/0009061 [astro-ph]].
  - [2] R. Beck, Space Sci. Rev. **99** (2001), 243-260 [arXiv:astro-ph/0012402 [astro-ph]].
  - [3] L. M. Widrow, Rev. Mod. Phys. **74** (2002), 775-823 [arXiv:astro-ph/0207240 [astro-ph]].
  - [4] A. Kandus, K. E. Kunze and C. G. Tsagas, Phys. Rept. **505** (2011), 1-58 [arXiv:1007.3891 [astro-ph.CO]].
  - [5] R. Durrer and A. Neronov, Astron. Astrophys. Rev. **21** (2013), 62 [arXiv:1303.7121 [astro-ph.CO]].
  - [6] K. Subramanian, Rept. Prog. Phys. **79** (2016) no.7, 076901 [arXiv:1504.02311 [astro-ph.CO]].
  - [7] Y. Akrami *et al.* [Planck], Astron. Astrophys. **641** (2020), A10 doi:10.1051/0004-6361/201833887 [arXiv:1807.06211 [astro-ph.CO]].
  - [8] A. Riotto and M. Trodden, Ann. Rev. Nucl. Part. Sci. **49** (1999), 35-75 doi:10.1146/annurev.nucl.49.1.35 [arXiv:hep-ph/9901362 [hep-ph]].
  - [9] M. Giovannini and M. E. Shaposhnikov, Phys. Rev. Lett. **80** (1998), 22-25 doi:10.1103/PhysRevLett.80.22 [arXiv:hep-ph/9708303 [hep-ph]].
  - [10] M. Giovannini and M. E. Shaposhnikov, Phys. Rev. D **57** (1998), 2186-2206 doi:10.1103/PhysRevD.57.2186 [arXiv:hep-ph/9710234 [hep-ph]].
  - [11] M. Dine and A. Kusenko, Rev. Mod. Phys. **76** (2003), 1 doi:10.1103/RevModPhys.76.1 [arXiv:hep-ph/0303065 [hep-ph]].
  - [12] J. M. Cline, [arXiv:hep-ph/0609145 [hep-ph]].
  - [13] B. Garbrecht, Prog. Part. Nucl. Phys. **110** (2020), 103727 doi:10.1016/j.ppnp.2019.103727 [arXiv:1812.02651 [hep-ph]].
  - [14] S. D. Odintsov and V. K. Oikonomou, Phys. Lett. B **760** (2016), 259-262 doi:10.1016/j.physletb.2016.06.074 [arXiv:1607.00545 [gr-qc]].
  - [15] S. D. Odintsov and V. K. Oikonomou, EPL **116** (2016) no.4, 49001 doi:10.1209/0295-5075/116/49001 [arXiv:1610.02533 [gr-qc]].
  - [16] K. Bamba, C. Q. Geng and S. H. Ho, Phys. Lett. B **664** (2008), 154-156 doi:10.1016/j.physletb.2008.05.027 [arXiv:0712.1523 [hep-ph]].
  - [17] K. Bamba, Phys. Rev. D **74** (2006), 123504 doi:10.1103/PhysRevD.74.123504 [arXiv:hep-ph/0611152 [hep-ph]].
  - [18] A. D. Sakharov, Pisma Zh. Eksp. Teor. Fiz. **5** (1967), 32-35 doi:10.1070/PU1991v034n05ABEH002497
  - [19] R. M. Kulshrud and E. G. Zweibel, Rept. Prog. Phys. **71** (2008), 0046091 [arXiv:0707.2783 [astro-ph]].
  - [20] A. Brandenburg and K. Subramanian, Phys. Rept. **417** (2005), 1-209 [arXiv:astro-ph/0405052 [astro-ph]].
  - [21] K. Subramanian, Astron. Nachr. **331** (2010), 110-120 [arXiv:0911.4771 [astro-ph.CO]].
  - [22] R. Sharma, S. Jagannathan, T. R. Seshadri and K. Subramanian, Phys. Rev. D **96** (2017) no.8, 083511 [arXiv:1708.08119 [astro-ph.CO]].
  - [23] R. Sharma, K. Subramanian and T. R. Seshadri, Phys. Rev. D **97** (2018) no.8, 083503 [arXiv:1802.04847 [astro-ph.CO]].
  - [24] R. K. Jain and M. S. Sloth, Phys. Rev. D **86** (2012), 123528 [arXiv:1207.4187 [astro-ph.CO]].
  - [25] R. Durrer, L. Hollenstein and R. K. Jain, JCAP **03** (2011), 037 [arXiv:1005.5322 [astro-ph.CO]].
  - [26] S. Kanno, J. Soda and M. a. Watanabe, JCAP **12** (2009), 009 [arXiv:0908.3509 [astro-ph.CO]].
  - [27] L. Campanelli, Int. J. Mod. Phys. D **18** (2009), 1395-1411 [arXiv:0805.0575 [astro-ph]].
  - [28] V. Demozzi, V. Mukhanov and H. Rubinstein, JCAP **08** (2009), 025 [arXiv:0907.1030 [astro-ph.CO]].
  - [29] V. Demozzi and C. Ringeval, JCAP **05**, 009 (2012) [arXiv:1202.3022 [astro-ph.CO]].
  - [30] K. Bamba and S. D. Odintsov, JCAP **0804** (2008) 024 [arXiv:0801.0954 [astro-ph]].
  - [31] K. Bamba, S. Nojiri and S. D. Odintsov, Phys. Rev. D **77**, 123532 (2008) [arXiv:0803.3384 [hep-th]].
  - [32] K. Bamba, C. Q. Geng and L. W. Luo, JCAP **1210**, 058 (2012) [arXiv:1208.0665 [astro-ph.CO]].
  - [33] K. Bamba and M. Sasaki, JCAP **02** (2007), 030 [arXiv:astro-ph/0611701 [astro-ph]].
  - [34] K. Bamba and J. Yokoyama, Phys. Rev. D **69** (2004), 043507 [arXiv:astro-ph/0310824 [astro-ph]].
  - [35] K. Bamba and J. Yokoyama, Phys. Rev. D **70** (2004), 083508 [arXiv:hep-ph/0409237 [hep-ph]].



- [36] K. Bamba, E. Elizalde, S. D. Odintsov and T. Paul, JCAP **04** (2021), 009 doi:10.1088/1475-7516/2021/04/009 [arXiv:2012.12742 [gr-qc]].
- [37] D. Maity, S. Pal and T. Paul, JCAP **05** (2021), 045 doi:10.1088/1475-7516/2021/05/045 [arXiv:2103.02411 [hep-th]].
- [38] M. R. Haque, D. Maity and S. Pal, [arXiv:2012.10859 [hep-th]].
- [39] M. Giovannini, Phys. Lett. B **819** (2021), 136444 doi:10.1016/j.physletb.2021.136444 [arXiv:2104.03891 [hep-th]].
- [40] M. Giovannini, Class. Quant. Grav. **35** (2018) no.8, 084003 doi:10.1088/1361-6382/aab17d [arXiv:1712.07598 [astro-ph.CO]].
- [41] M. Giovannini, Int. J. Mod. Phys. D **13** (2004), 391-502 doi:10.1142/S0218271804004530 [arXiv:astro-ph/0312614 [astro-ph]].
- [42] M. Gasperini, M. Giovannini and G. Veneziano, Phys. Rev. Lett. **75**, 3796-3799 (1995) doi:10.1103/PhysRevLett.75.3796 [arXiv:hep-th/9504083 [hep-th]].
- [43] T. Kobayashi and M. S. Sloth, Phys. Rev. D **100** (2019) no.2, 023524 [arXiv:1903.02561 [astro-ph.CO]].
- [44] B. Ratra, Astrophys. J. Lett. **391** (1992), L1-L4
- [45] P. A. R. Ade *et al.* [Planck], Astron. Astrophys. **594** (2016), A19 [arXiv:1502.01594 [astro-ph.CO]].
- [46] D. Chowdhury, L. Sriramkumar and M. Kamionkowski, JCAP **10** (2018), 031 [arXiv:1807.07477 [astro-ph.CO]].
- [47] T. Vachaspati, Phys. Lett. B **265** (1991), 258-261
- [48] M. S. Turner and L. M. Widrow, Phys. Rev. D **37** (1988), 2743
- [49] K. Takahashi, K. Ichiki, H. Ohno and H. Hanayama, Phys. Rev. Lett. **95** (2005), 121301 [arXiv:astro-ph/0502283 [astro-ph]].
- [50] I. Agullo and J. Navarro-Salas, [arXiv:1309.3435 [gr-qc]].
- [51] R. J. Z. Ferreira, R. K. Jain and M. S. Sloth, JCAP **10** (2013), 004 [arXiv:1305.7151 [astro-ph.CO]].
- [52] K. Atmjeet, T. R. Seshadri and K. Subramanian, Phys. Rev. D **91** (2015), 103006 [arXiv:1409.6840 [astro-ph.CO]].
- [53] R. Sharma, [arXiv:2102.09358 [astro-ph.CO]].
- [54] P. Adshedy, J. T. Giblin, T. R. Scully and E. I. Sfakianakis, JCAP **12** (2015), 034 [arXiv:1502.06506 [astro-ph.CO]].
- [55] P. Adshedy, J. T. Giblin, T. R. Scully and E. I. Sfakianakis, JCAP **10** (2016), 039 [arXiv:1606.08474 [astro-ph.CO]].
- [56] E. Frion, N. Pinto-Neto, S. D. P. Vitenti and S. E. Perez Bergliaffa, Phys. Rev. D **101** (2020) no.10, 103503 [arXiv:2004.07269 [gr-qc]].
- [57] D. Chowdhury, L. Sriramkumar and R. K. Jain, Phys. Rev. D **94** (2016) no.8, 083512 [arXiv:1604.02143 [gr-qc]].
- [58] D. Chowdhury, L. Sriramkumar and M. Kamionkowski, JCAP **01** (2019), 048 [arXiv:1807.05530 [astro-ph.CO]].
- [59] P. Qian, Y. F. Cai, D. A. Easson and Z. K. Guo, Phys. Rev. D **94** (2016) no.8, 083524 [arXiv:1607.06578 [gr-qc]].
- [60] R. Koley and S. Samtani, JCAP **04** (2017), 030 [arXiv:1612.08556 [gr-qc]].
- [61] F. A. Membrilla, Nucl. Phys. B **885** (2014), 196-224 [arXiv:1312.2162 [astro-ph.CO]].
- [62] A.H. Guth; Phys.Rev. D23 347-356 (1981).
- [63] A. D. Linde, Contemp. Concepts Phys. **5** (1990) 1 [hep-th/0503203].
- [64] D. Langlois, hep-th/0405053.
- [65] A. Riotto, ICTP Lect. Notes Ser. **14** (2003) 317 [hep-ph/0210162].
- [66] D. Baumann, [arXiv:0907.5424 [hep-th]].
- [67] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, Phys. Rept. **692** (2017), 1-104 doi:10.1016/j.physrep.2017.06.001 [arXiv:1705.11098 [gr-qc]].
- [68] C. Caprini and L. Sorbo, JCAP **10** (2014), 056 [arXiv:1407.2809 [astro-ph.CO]].
- [69] T. Kobayashi, JCAP **05** (2014), 040 [arXiv:1403.5168 [astro-ph.CO]].
- [70] K. Atmjeet, I. Pahwa, T. R. Seshadri and K. Subramanian, Phys. Rev. D **89** (2014) no.6, 063002 [arXiv:1312.5815 [astro-ph.CO]].
- [71] T. Fujita, R. Namba, Y. Tada, N. Takeda and H. Tashiro, JCAP **05** (2015), 054 [arXiv:1503.05802 [astro-ph.CO]].
- [72] L. Campanelli, Eur. Phys. J. C **75** (2015) no.6, 278 [arXiv:1503.07415 [gr-qc]].
- [73] G. Tasinato, JCAP **03** (2015), 040 [arXiv:1411.2803 [hep-th]].
- [74] D. Nandi, [arXiv:2103.03159 [astro-ph.CO]].
- [75] R. H. Brandenberger, arXiv:1206.4196 [astro-ph.CO].
- [76] R. Brandenberger and P. Peter, arXiv:1603.05834 [hep-th].
- [77] D. Battefeld and P. Peter, Phys. Rept. **571** (2015) 1 [arXiv:1406.2790 [astro-ph.CO]].
- [78] M. Novello and S. E. P. Bergliaffa, "Bouncing Cosmologies," Phys. Rept. **463** (2008) 127 [arXiv:0802.1634 [astro-ph]].
- [79] Y. F. Cai, Sci. China Phys. Mech. Astron. **57** (2014) 1414 [arXiv:1405.1369 [hep-th]].
- [80] S. Nojiri, S. D. Odintsov, V. K. Oikonomou and T. Paul, Phys. Rev. D **100** (2019) no.8, 084056 [arXiv:1910.03546 [gr-qc]].
- [81] Y. F. Cai, T. t. Qiu, R. Brandenberger and X. m. Zhang, Phys. Rev. D **80** (2009), 023511 [arXiv:0810.4677 [hep-th]].
- [82] E. Elizalde, S. D. Odintsov and T. Paul, Eur. Phys. J. C **80** (2020) no.1, 10 [arXiv:1912.05138 [gr-qc]].
- [83] E. Elizalde, S. D. Odintsov, V. K. Oikonomou and T. Paul, Nucl. Phys. B **954** (2020), 114984 [arXiv:2003.04264 [gr-qc]].
- [84] K. Bamba, A. N. Makarenko, A. N. Myagky and S. D. Odintsov, Phys. Lett. B **732** (2014), 349-355 [arXiv:1403.3242 [hep-th]].
- [85] S. D. Odintsov, V. K. Oikonomou and T. Paul, Class. Quant. Grav. **37** (2020) no.23, 235005 [arXiv:2009.09947 [gr-qc]].
- [86] I. Banerjee, T. Paul and S. SenGupta, [arXiv:2011.11886 [gr-qc]].
- [87] S. Davidson, Phys. Lett. B **380** (1996), 253-256 doi:10.1016/0370-2693(96)00501-1 [arXiv:astro-ph/9605086 [astro-ph]].
- [88] M. Joyce and M. E. Shaposhnikov, Phys. Rev. Lett. **79** (1997), 1193-1196 doi:10.1103/PhysRevLett.79.1193 [arXiv:astro-ph/9703005 [astro-ph]].
- [89] C. Thompson, Phys. Lett. B **422** (1998), 61-68 doi:10.1016/S0370-2693(98)00053-7

- [90] T. Vachaspati and G. B. Field, *Phys. Rev. Lett.* **73** (1994), 373-376 doi:10.1103/PhysRevLett.73.373 [arXiv:hep-ph/9401220 [hep-ph]].
- [91] V. A. Rubakov and M. E. Shaposhnikov, *Usp. Fiz. Nauk* **166** (1996), 493-537 doi:10.1070/PU1996v039n05ABEH000145 [arXiv:hep-ph/9603208 [hep-ph]].
- [92] A. Kushwaha and S. Shankaranarayanan, [arXiv:2103.05339 [hep-ph]].
- [93] M. Giovannini, *Eur. Phys. J. C* **81** (2021), 81 doi:10.1140/epjc/s10052-021-09282-7 [arXiv:2103.04137 [astro-ph.CO]].
- [94] S. Nojiri, S. D. Odintsov and M. Sasaki, *Phys. Rev. D* **71** (2005), 123509 doi:10.1103/PhysRevD.71.123509 [arXiv:hep-th/0504052 [hep-th]].
- [95] B. Li, J. D. Barrow and D. F. Mota, *Phys. Rev. D* **76** (2007), 044027 [arXiv:0705.3795 [gr-qc]].
- [96] B. M. Carter and I. P. Neupane, *JCAP* **06** (2006), 004 [arXiv:hep-th/0512262 [hep-th]].
- [97] S. Nojiri, S. Odintsov, V. Oikonomou, N. Chatzarakis and T. Paul, *Eur. Phys. J. C* **79** (2019) no.7, 565 [arXiv:1907.00403 [gr-qc]].
- [98] A. de la Cruz-Dombriz and D. Saez-Gomez, *Class. Quant. Grav.* **29** (2012), 245014 [arXiv:1112.4481 [gr-qc]].
- [99] S. Chakraborty, T. Paul and S. SenGupta, *Phys. Rev. D* **98** (2018) no.8, 083539 [arXiv:1804.03004 [gr-qc]].
- [100] P. Kanti, R. Gannouji and N. Dadhich, *Phys. Rev. D* **92** (2015) no.4, 041302 [arXiv:1503.01579 [hep-th]].
- [101] P. Kanti, R. Gannouji and N. Dadhich, *Phys. Rev. D* **92** (2015) no.8, 083524 [arXiv:1506.04667 [hep-th]].
- [102] S. D. Odintsov and V. K. Oikonomou, *Phys. Rev. D* **98** (2018) no.4, 044039 [arXiv:1808.05045 [gr-qc]].
- [103] E. N. Saridakis, *Phys. Rev. D* **97** (2018) no.6, 064035 [arXiv:1707.09331 [gr-qc]].
- [104] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov and S. Zerbini, *Phys. Rev. D* **73** (2006), 084007 [arXiv:hep-th/0601008 [hep-th]].
- [105] L. Dai, M. Kamionkowski and J. Wang, *Phys. Rev. Lett.* **113** (2014), 041302 [arXiv:1404.6704 [astro-ph.CO]].
- [106] J. L. Cook, E. Dimastrogiovanni, D. A. Easson and L. M. Krauss, *JCAP* **04** (2015), 047 [arXiv:1502.04673 [astro-ph.CO]].
- [107] A. Albrecht, P. J. Steinhardt, M. S. Turner and F. Wilczek, *Phys. Rev. Lett.* **48** (1982), 1437
- [108] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos and K. A. Olive, *JCAP* **07** (2015), 050 [arXiv:1505.06986 [hep-ph]].
- [109] Y. Ueno and K. Yamamoto, *Phys. Rev. D* **93** (2016) no.8, 083524 [arXiv:1602.07427 [astro-ph.CO]].
- [110] M. Eshaghi, M. Zarei, N. Riazi and A. Kiasatpour, *Phys. Rev. D* **93** (2016) no.12, 123517 [arXiv:1602.07914 [astro-ph.CO]].
- [111] D. Maity and P. Saha, *JCAP* **07** (2019), 018 [arXiv:1811.11173 [astro-ph.CO]].
- [112] M. R. Haque, D. Maity and P. Saha, *Phys. Rev. D* **102** (2020) no.8, 083534 [arXiv:2009.02794 [hep-th]].
- [113] A. Di Marco, P. Cabella and N. Vittorio, *Phys. Rev. D* **95** (2017) no.10, 103502 [arXiv:1705.04622 [astro-ph.CO]].
- [114] M. Drewes, J. U. Kang and U. R. Mun, *JHEP* **11** (2017), 072 [arXiv:1708.01197 [astro-ph.CO]].
- [115] A. Di Marco, G. Pradisi and P. Cabella, *Phys. Rev. D* **98** (2018) no.12, 123511 [arXiv:1807.05916 [astro-ph.CO]].
- [116] S. D. Odintsov and V. K. Oikonomou, *Phys. Lett. B* **797** (2019), 134874 [arXiv:1908.07555 [gr-qc]].
- [117] S. D. Odintsov, V. K. Oikonomou and F. P. Fronimos, *Nucl. Phys. B* **958** (2020), 115135 [arXiv:2003.13724 [gr-qc]].
- [118] S. D. Odintsov, V. K. Oikonomou and F. P. Fronimos, *Annals Phys.* **424** (2021), 168359 doi:10.1016/j.aop.2020.168359 [arXiv:2011.08680 [gr-qc]].
- [119] S. D. Odintsov, V. K. Oikonomou and F. P. Fronimos, *Annals Phys.* **420** (2020), 168250 doi:10.1016/j.aop.2020.168250 [arXiv:2007.02309 [gr-qc]].
- [120] N. D. Barrie and A. Kobakhidze, *JHEP* **09** (2014), 163 doi:10.1007/JHEP09(2014)163 [arXiv:1401.1256 [hep-ph]].
- [121] T. Kobayashi and N. Afshordi, *JHEP* **10** (2014), 166 doi:10.1007/JHEP10(2014)166 [arXiv:1408.4141 [hep-th]].
- [122] C. Stahl, *Nucl. Phys. B* **939** (2019), 95-104 doi:10.1016/j.nuclphysb.2018.12.017 [arXiv:1806.06692 [hep-th]].
- [123] T. Fujita and S. Yokoyama, *JCAP* **09** (2013), 009 doi:10.1088/1475-7516/2013/09/009 [arXiv:1306.2992 [astro-ph.CO]].
- [124] T. Fujita and R. Namba, *Phys. Rev. D* **94** (2016) no.4, 043523 doi:10.1103/PhysRevD.94.043523 [arXiv:1602.05673 [astro-ph.CO]].
- [125] K. Bamba, *Phys. Rev. D* **91** (2015), 043509 doi:10.1103/PhysRevD.91.043509 [arXiv:1411.4335 [astro-ph.CO]].
- [126] N. Barnaby, R. Namba and M. Peloso, *Phys. Rev. D* **85** (2012), 123523 doi:10.1103/PhysRevD.85.123523 [arXiv:1202.1469 [astro-ph.CO]].
- [127] R. J. Z. Ferreira, R. K. Jain and M. S. Sloth, *JCAP* **06** (2014), 053 doi:10.1088/1475-7516/2014/06/053 [arXiv:1403.5516 [astro-ph.CO]].