

Compact elastic objects in general relativity

Artur Alho,¹ José Natário,¹ Paolo Pani,² and Guilherme Raposo^{3,4,2}

¹*Center for Mathematical Analysis, Geometry and Dynamical Systems,*

Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

²*Dipartimento di Fisica, Sapienza Università di Roma & INFN Roma1, Piazzale Aldo Moro 5, 00185, Roma, Italy*

³*Centre for Research and Development in Mathematics and Applications (CIDMA),
Campus de Santiago, 3810-183 Aveiro, Portugal*

⁴*CENTRA, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal*

We introduce a rigorous and general framework to study systematically self-gravitating elastic materials within general relativity, and apply it to investigate the existence and viability, including radial stability, of spherically symmetric elastic stars. We present the mass-radius ($M - R$) diagram for various families of models, showing that elasticity contributes to increase the maximum mass and the compactness up to a $\mathcal{O}(10\%)$ factor, thus supporting compact stars with mass well above two solar masses. Some of these elastic stars can reach compactness as high as $GM/(c^2R) \approx 0.35$ while remaining stable under radial perturbations and satisfying all energy conditions and subluminal wave propagation, thus being physically viable models of stars with a light ring. We provide numerical evidence that radial instability occurs for central densities larger than that corresponding to the maximum mass, as in the perfect-fluid case. Elasticity may be a key ingredient to build consistent models of exotic ultracompact objects and black-hole mimickers, and can also be relevant for a more accurate modelling of the interior of neutron stars.

Introduction. Astrophysical compact objects are typically idealized as self-gravitating (often perfect) fluids, wherein (isotropic) pressure prevents gravitational collapse. However, while degenerate fermions behave as a weakly-interacting gas at relatively small densities, nuclear interactions and QCD effects become crucial inside relativistic stars. Thus, it is reasonable to expect that the perfect-fluid idealization will eventually break down, at least to some extent, and that solid phases of matter may be relevant for astrophysical compact objects. This is indeed the situation in the crust of a neutron star [1, 2], whose fundamental constituents are largely unknown, especially in the core [3].

A natural generalization of fluid models is to consider elastic materials [4–8]. In the theory of relativistic elasticity the configuration of a body is described by a map Ψ from the spacetime (\mathcal{M}, g) to the 3-dimensional Riemannian material space (\mathcal{B}, γ) , and it is often not clear how the configuration map relates to the matter fields, which makes it hard to close the system of equations in a natural way. In addition to these formal approaches, elasticity has also been studied perturbatively to model the crust of a neutron star [1, 2]. In this letter, we introduce a systematic, physics-oriented approach to the problem of self-gravitating elastic materials, which is based on a General-Relativity (GR) extension of the stored energy functions for well-known Newtonian models [9, 10]. Our approach allows to build (partially) elastic compact objects in a simple – yet general – way and to assess their viability in the strong-gravity regime.

Beside offering a more accurate description of the stellar interior [11–13], elasticity might play a crucial role in constructing consistent models of exotic compact objects and black-hole mimickers within GR and extensions thereof [14, 15]. Under certain hypotheses [14, 16, 17], Buchdhal’s theorem [18] states that self-

gravitating, perfect-fluid GR solutions satisfy the following bound on the compactness: $M/R \leq 4/9$, where M is the mass and R is the radius of the star (henceforth we use $G = c = 1$ units). Buchdhal’s theorem assumes that matter is described by a perfect fluid, and can be extended to mildly anisotropic fluids for which the radial pressure is larger than the tangential one [16, 17]. Indeed, compact objects made of strongly anisotropic fluids (e.g., gravastars [19] and anisotropic stars [20]) can have higher compactness and a continuous BH limit, $M/R \rightarrow 1/2$ [21–24]. However, the viability of such ultracompact models is questionable, since they either violate some of the energy conditions [25], or feature superluminal speed of sound or ad-hoc thin-shells within the fluid (see [14] for a discussion). On the other hand, viable models like boson stars (which also feature anisotropies in the matter fields) are not significantly more compact than an ordinary perfect-fluid neutron star in the static case [26].

Since elastic materials feature shears and anisotropies, it is natural to ask whether completely viable models of *ultracompact* [14] elastic stars can be built. In this letter we will show that this is the case. Viable elastic stars can have $M/R > 1/3$, thus featuring the same Schwarzschild photon sphere at radial coordinate $r = 3M$, a crucial property to mimic the phenomenology of black holes [14, 15, 27, 28]. We will also show that elastic stars can exceed the Buchdhal’s bound on the compactness, but only in their unstable and superluminal branch, at least for the class of materials under consideration.

Setup. We focus on spherical symmetry and study both static solutions and their dynamical radial perturbations. More details and models will be given in a companion paper [29]. Static spherically symmetric self-gravitating relativistic bodies are described in Schwarzschild coordinates

by the Tolman-Oppenheimer-Volkoff (TOV) equations

$$\frac{dp_{\text{rad}}}{dr} = \frac{2}{r}(p_{\text{tan}} - p_{\text{rad}}) - (p_{\text{rad}} + \rho) \frac{d\alpha}{dr}, \quad (1)$$

$$\frac{d\alpha}{dr} = \frac{e^{2\beta}}{r} \left(\frac{m}{r} + 4\pi r^2 p_{\text{rad}} \right), \quad (2)$$

where $e^{-2\beta(r)} = 1 - \frac{2m(r)}{r}$, $p_{\text{rad}}(r)$ and $p_{\text{tan}}(r)$ are the radial and tangential pressures, respectively, $\rho(r)$ is the energy density, $\alpha(r)$ the relativistic gravitational potential, and $m(r) = 4\pi \int_0^r \rho(u)u^2 du$ is the Misner-Sharp mass. The line element then reads $ds^2 = -e^{2\alpha(r)} dt^2 + e^{2\beta(r)} dr^2 + r^2 d\Omega^2$, where $d\Omega^2$ is the metric of the unit 2-sphere, and the stress-energy tensor that sources Einstein's equations is given by $T_{\nu}^{\mu} = \text{diag}(\rho, p_{\text{rad}}, p_{\text{tan}}, p_{\text{tan}})$. The TOV system is closed by postulating a constitutive equation relating the pressures and the density.

Relativistic elasticity is based on a variational principle wherein the Lagrangian density is covariant under spacetime diffeomorphisms, and consists of the sum of the rest-frame energy density for the undeformed material and a (deformation) potential energy density, so that it coincides with the total energy density ρ measured by an observer at rest with respect to the material [30, 31]. For homogeneous and isotropic elastic materials and under spherical symmetry, the Lagrangian is given by [29]

$$\hat{\rho}(\delta, \eta) = \delta(\rho_0 + \hat{w}(\delta, \eta)), \quad (3)$$

where the potential energy density (also called stored energy function, by analogy with the Newtonian case) is $w(r) = \hat{w}(\delta(r), \eta(r))$, so that $\rho(r) = \hat{\rho}(\delta(r), \eta(r))$. Here

$$\delta(r) = \frac{n(r)}{n_0}, \quad \eta(r) = \frac{3}{r^3} \int_0^r e^{\beta(u)} \delta(u) u^2 du, \quad (4)$$

where $n(r)$ is the (conserved) particle number density in the physical (deformed) state, and $n_0 > 0$ and $\rho_0 > 0$ are the particle number density and energy density in the reference material frame, respectively. The reference state is an idealized state with $n = n_0$ and $\beta = 0$ (corresponding to a flat material metric), that is, $(\delta, \eta) = (1, 1)$.

For a given choice of \hat{w} , the constitutive functions are

$$\hat{p}_{\text{rad}}(\delta, \eta) = \delta \partial_{\delta} \hat{\rho} - \hat{\rho}, \quad (5a)$$

$$\hat{q}(\delta, \eta) \equiv \hat{p}_{\text{tan}} - \hat{p}_{\text{rad}} = \frac{3}{2} \eta \partial_{\eta} \hat{\rho}. \quad (5b)$$

Note that the perfect fluid case is included for any Lagrangian such that $\partial_{\eta} \hat{\rho} = 0$ (equivalently $\partial_{\eta} \hat{w} = 0$).

The function $\hat{w}(\delta, \eta)$ satisfies the natural reference state condition (state of zero energy)

$$\hat{w}(1, 1) = 0, \quad (6)$$

while compatibility with linear elasticity requires

$$\partial_{\delta}^2 \hat{w}(1, 1) = \lambda + 2\mu, \quad \partial_{\eta}^2 \hat{w}(1, 1) = -\partial_{\eta\delta}^2 \hat{w}(1, 1) = \frac{4}{3}\mu, \quad (7)$$

where λ, μ are the Lamé parameters. The radial and tangential pressures satisfy the reference state condition

$$\hat{p}_{\text{rad}}(1, 1) = \hat{p}_{\text{tan}}(1, 1) = p_0. \quad (8)$$

The reference state is said to be stress-free (pre-stressed) if $p_0 = 0$ ($p_0 > 0$).

In terms of the variables $(\delta(r), \eta(r), m(r))$, the TOV equations become a closed first-order system,

$$\partial_{\delta} \hat{p}_{\text{rad}} \frac{d\delta}{dr} = \frac{3}{r} \partial_{\eta} \hat{p}_{\text{rad}} (\eta - e^{\beta} \delta) + \frac{2}{r} \hat{q} - (\hat{p}_{\text{rad}} + \hat{\rho}) \frac{e^{2\beta}}{r} \left(\frac{m}{r} + 4\pi r^2 \hat{p}_{\text{rad}} \right), \quad (9a)$$

$$\frac{d\eta}{dr} = -\frac{3}{r} (\eta - e^{\beta} \delta), \quad \frac{dm}{dr} = 4\pi r^2 \hat{\rho}, \quad (9b)$$

supplemented by (3) and the equations of state (5). In the Newtonian limit we recover the results of [9, 10].

For regular solutions we have $\lim_{r \rightarrow 0^+} \delta(r) = \delta_c$ and $\lim_{r \rightarrow 0^+} e^{\beta(r)} = 1$, implying $\lim_{r \rightarrow 0^+} \eta(r) = \delta_c$. Regularity at the origin imposes $p_{\text{rad}}(0) = p_{\text{tan}}(0)$, which in turn implies

$$\hat{q}(\delta, \delta) = 0 \quad \text{for all } \delta > 0. \quad (10)$$

Material models. The above formalism is general for any given function \hat{w} satisfying Eqs. (6), (7), (8), and (10). Here we focus on elastic constitutive functions that are continuous deformations of relativistic fluid models, in particular relativistic polytropes (other examples are given in [29]). The simplest such material model is defined by

$$\hat{\rho}(\delta, \eta) = (1 - \kappa n) \rho_0 \delta + \kappa n \rho_0 \delta^{1+\frac{1}{n}} + \varepsilon \rho_0 (\delta - \eta)^2, \quad (11)$$

which contains three dimensionless parameters κ, n and ε . For $\varepsilon = 0$ we have a perfect fluid with a polytropic equation of state with polytropic index n and reference state pressure $p_0 = \kappa \rho_0$.

The Newtonian limit of the previous quadratic model leads to equations of motion which are invariant under homologous transformations [32] only for $n = 1$. It is possible to generalize this stored energy function to one that leads to Newtonian equations of motion which are invariant under homologous transformations for general polytropic index n [29]:

$$\hat{w}(\delta, \eta) = w_0 + \eta^{\frac{1}{n}} \left[w_1 + w_2 \left(\frac{\delta}{\eta} \right)^{-1} + w_3 \left(\frac{\delta}{\eta} \right)^{\frac{1}{s}} \right], \quad (12)$$

where $w_0 \equiv -n\kappa\rho_0$, $w_1 \equiv \frac{(n-s)(1+n)}{n} \rho_0 \kappa - 2s\rho_0\varepsilon$, $w_2 \equiv \frac{(s-n)}{(1+s)n} \rho_0 \kappa + \frac{2s}{1+s} \rho_0 \varepsilon$, $w_3 \equiv \frac{(1+n)s^2}{(1+s)n} \rho_0 \kappa + \frac{2s^2}{1+s} \rho_0 \varepsilon$. Here s can be interpreted as the shear index; when $s = n$ and $\varepsilon = 0$ we recover the usual relativistic polytropes. This function $\hat{w}(\delta, \eta)$ is similar to the power-law stored energy function (2.18) in [33], which belongs to a more general class of polytropic elastic materials introduced




FIG. 1. Mass-radius diagram for the quadratic elastic model (11) with $n = 1/2$ and $K = 6 \times 10^4 M_\odot^4$ (left) and for the two-parameter elastic model (12) with $n = 1$ and $K = 100 M_\odot^2$ (right). Solid (dashed) curves correspond to configurations with subluminal (superluminal) wave propagation. For some regions of the parameter space (cf. green curve of left panel) there are configurations below the maximum mass featuring a photon sphere ($R < 3M$) and subluminal wave propagation.

in [34]. However, in contrast to these models, \hat{w} defined in Eq. (12) leads to a pre-stressed material model from the outset, with pressure $p_0 = \kappa\rho_0$, and has therefore a proper relativistic generalisation. When $s = n = 1$ it becomes the simple quadratic model (11). In both models, the first Lamé parameter is $\lambda = \kappa\rho_0(1 + \frac{1}{n}) - \varepsilon\rho_0$, and the shear modulus is $\mu = \frac{3}{2}\varepsilon\rho_0$, which implies $\varepsilon \geq 0$.

When the model is stress-free, $p_0 = 0$, the reference state $(\delta, \eta) = (1, 1)$ is uniquely defined. However, pre-stressed models (such as the ones we are considering) do not have a preferred reference state. In this case a different reference state, compressed or expanded with respect to the original reference state, provides an equivalent description of the material, moving from the parameters $(\rho_0, \kappa, \varepsilon)$ to new parameters $(\hat{\rho}_0, \hat{\kappa}, \hat{\varepsilon})$. The choice of reference state is thus akin to a gauge choice (see [29] for more details).

It can be shown that in the fluid case $\hat{p}_{\text{rad}} = \hat{p}_{\text{tan}} = \kappa\rho_0\delta^{1+\frac{1}{n}} = K\hat{\sigma}^{1+\frac{1}{n}}$, where $\hat{\sigma}$ is the baryon density and $K = \kappa(1 - \kappa n)^{-\frac{n+1}{n}}\rho_0^{-\frac{1}{n}}$. The latter quantity is in fact invariant under renormalization of the reference state. Moreover, changing K only affects the equilibrium configuration with an overall scale, and thus does not affect the value of dimensionless quantities such as the compactness. Another invariant quantity under renormalization is

$$E = \frac{\varepsilon}{\kappa} \left(\frac{\kappa}{1 - \kappa n} \right)^{1-n} \quad \text{or} \quad E = \frac{\varepsilon}{\kappa}, \quad (13)$$

in the case of model (11) or model (12), respectively. In the latter case, E can be written explicitly in terms of the Poisson ratio ν as $E = \left(\frac{n+1}{n}\right) \left(\frac{1-2\nu}{1+\nu}\right)$. When written in terms of the parameters (K, E, n, s) , the TOV equations become independent of ρ_0 , as expected.

Equilibrium configurations. The equations (9) for the stellar structure should be solved by requiring regularity

of the functions at the center of the star. For a given choice of \hat{w} , the solutions form a one-parameter family in terms of δ_c , or, equivalently, of the central density $\rho(0) = \hat{\rho}(\delta_c, \delta_c) = \rho_c$. The radius R of the star is defined by the condition $p_{\text{rad}}(R) = 0$, whereas $\rho(R)$ and $p_{\text{tan}}(R)$ do not need to vanish. Due to Birkhoff's theorem, the metric at $r > R$ (where $\rho = p_{\text{rad}} = p_{\text{tan}} = 0$) is the standard Schwarzschild metric with $m(R) = M$ and $\alpha(r) = -\beta(r)$. (See [5, 7, 29, 35, 36] for the problem of C^1 regularity of relativistic spherical stars.)

Within GR, physically viable matter fields should satisfy the following energy conditions [25]:

$$\begin{aligned} \text{SEC} : \quad & \rho + p_{\text{rad}} + 2p_{\text{tan}} \geq 0; \quad \rho + p_{\text{rad}} \geq 0; \quad \rho + p_{\text{tan}} \geq 0; \\ \text{WEC} : \quad & \rho \geq 0; \quad \rho + p_{\text{rad}} \geq 0; \quad \rho + p_{\text{tan}} \geq 0; \\ \text{NEC} : \quad & \rho + p_{\text{rad}} \geq 0; \quad \rho + p_{\text{tan}} \geq 0; \\ \text{DEC} : \quad & \rho \geq |p_{\text{rad}}|; \quad \rho \geq |p_{\text{tan}}|; \end{aligned}$$

for the strong, weak, null, and dominant energy condition, respectively. Some further restrictions come from requiring causal wave propagation within the material. The square speed of elastic longitudinal and transverse waves in the *radial* direction are given by [6]

$$c_L^2(\delta, \eta) = \frac{\delta\partial_\delta \hat{p}_{\text{rad}}}{\hat{\rho} + \hat{p}_{\text{rad}}}, \quad (15)$$

$$c_T^2(\delta, \eta) = \frac{\hat{p}_{\text{tan}} - \hat{p}_{\text{rad}}}{(\hat{\rho} + \hat{p}_{\text{tan}})(1 - \delta^2/\eta^2)}, \quad (16)$$

whereas in the *tangential* direction

$$\tilde{c}_L^2(\delta, \eta) = \frac{\delta\partial_\delta \hat{p}_{\text{tan}} + 3\eta\partial_\eta \hat{p}_{\text{tan}}}{\hat{\rho} + \hat{p}_{\text{tan}}}, \quad (17)$$

$$\tilde{c}_T^2(\delta, \eta) = \frac{\hat{p}_{\text{rad}} - \hat{p}_{\text{tan}}}{(\hat{\rho} + \hat{p}_{\text{rad}})(1 - \eta^2/\delta^2)}. \quad (18)$$

Due to spherical symmetry, there is a single transverse mode in both radial and tangential directions, which van-

ishes in the perfect-fluid case, since $\hat{p}_{\text{tan}} = \hat{p}_{\text{rad}}$. Overall, causality requires $0 \leq c_{L,T}^2 \leq 1$, and $0 \leq \tilde{c}_{L,T}^2 \leq 1$.

Figure 1 shows the $M - R$ diagram for some representative examples of the aforementioned elastic models. Larger values of E and of $s - n$ increase the maximum mass and maximum compactness of star. In particular, a one-parameter deformation of the $n = 1/2$ polytrope (left panel) supports configurations with a light ring ($R < 3M$) and exceeding the Buchdahl's limit ($R < 9M/4$). However, a relevant question is whether the equilibrium configurations satisfy the various constraints imposed by the subluminality of wave propagation in the radial and tangential directions. Solid (dashed) parts of each curve in Fig. 1 correspond to subluminality (superluminality). Depending on the model, the transition can occur before or after the maximum mass, which corresponds to the heaviest stable configuration under radial perturbations, as discussed below. In general, as the central density increases, some wave speeds (predominantly \tilde{c}_L , i.e. the speed of longitudinal waves along the tangential direction) become superluminal within the star, as shown in Fig. 2 for a configuration that is almost marginally causal. These viability requirements set an upper bound on the maximum mass and compactness of viable equilibrium solutions. In particular, in all models under consideration, the Buchdahl limit is exceeded only for radially unstable or superluminal configurations. On the other hand, we found ultracompact configurations ($R < 3M$) for both models (11) and (12), although the second case requires a fine tuning of the parameters and the compactness never exceeds $M/R = 1/3$ by more than 1% [29].

For the simple quadratic model (11) with $n = 1/2$, we find stable and causal configurations with compactness as large as $M/R \approx 0.35$, which is reached for $E = \mathcal{O}(10^{-1})$. Interestingly, in this case the shear modulus is approximately

$$\mu \approx 7 \times 10^{26} \left(\frac{\rho_0}{10^{11} \text{ g/cm}^3} \right)^2 \sqrt{\frac{K}{10^5 M_\odot^4}} \left(\frac{E}{0.1} \right) \text{ erg/cm}^3, \quad (19)$$

which is in the typical range of values for lattice models describing the neutron-star crust [1].

Radial stability. Linear radial perturbations of relativistic elastic balls have been treated in [37] using a Lagrangian approach. In [29] we consider a more traditional Eulerian approach, also including perturbations of a (possibly non-flat) material metric. The perturbation variables are $\zeta(t, r) \equiv r^2 e^{-\alpha \xi}$ and $\chi(t, r) = -e^{\beta+2\alpha} (p_{\text{rad}})_L$, where ξ is the usual radial displacement in the perturbed configuration and $(p_{\text{rad}})_L$ is the Lagrangian perturbation of the radial pressure. Making the ansatz $\zeta(t, r) = e^{\pm i\omega t} \zeta(r)$, $\chi(t, r) = e^{\pm i\omega t} \chi(r)$ leads to an eigenvalue problem for a system of first-order ordinary differential equations

$$\delta \partial_\delta \hat{p}_{\text{rad}} \frac{d\zeta}{dr} = e^{-(3\alpha+\beta)} r^2 \chi - \frac{3}{r} \eta \partial_\eta \hat{p}_{\text{rad}} \zeta, \quad (20a)$$

$$\delta \partial_\delta \hat{p}_{\text{rad}} \frac{d\chi}{dr} = \frac{3}{r} \eta \partial_\eta \hat{p}_{\text{rad}} \chi - [Q_1 + Q_2 \omega^2] \zeta, \quad (20b)$$




FIG. 2. Sound speeds (upper panel) and density and pressure profiles (bottom panel) for the quadratic elastic model with $n = 1/2$, $K = 6 \times 10^4 M_\odot^4$. We compare the perfect-fluid case ($E = 0$, dashed lines) with an elastic configuration with $E = 10^{-1}$ (solid lines). The latter configuration features a light ring ($M/R \approx 0.35$) and the wave speeds are always subluminal.

where

$$Q_1 = \frac{e^{3\alpha+\beta}}{r^2} \left[\frac{4}{r} (\delta \partial_\delta \hat{q} - \hat{q})^2 + \delta \partial_\delta \hat{p}_{\text{rad}} \left(\frac{2}{r^2} \hat{q} - \frac{6}{r^2} \eta \partial_\eta \hat{q} - \frac{4}{r} \frac{dp_{\text{rad}}}{dr} + \left(\frac{6}{r} \hat{q} - \frac{dp_{\text{rad}}}{dr} \right) \frac{d\alpha}{dr} - 8\pi (\hat{\rho} + \hat{p}_{\text{rad}}) e^{2\beta} \hat{p}_{\text{rad}} \right) \right], \quad (21a)$$

$$Q_2 = \frac{e^{\alpha+3\beta}}{r^2} \delta \partial_\delta \hat{p}_{\text{rad}} (\hat{\rho} + \hat{p}_{\text{rad}}), \quad (21b)$$

subject to the standard boundary conditions [38] $\lim_{r \rightarrow 0^+} \zeta = 0$ and $\lim_{r \rightarrow R} \chi = 0$.

As a representative example, in Fig. 3 we show the eigenvalues ω^2 obtained numerically for model (11) as a function of the mass for $n = 1/2$ and a range of values of E . Within numerical accuracy the zero crossing corresponds to the point of the maximum mass. We found the same result in all models under investigation [29]. Based on this numerical evidence, we conclude that elastic stars beyond the maximum-mass configuration are radially unstable, as in the perfect-fluid case [38].

Discussion. Anisotropies are ubiquitous in physical systems and seem also a key ingredient to built solutions for ultracompact self-gravitating objects. However, introducing anisotropies in GR is often based on ad-hoc models which might also suffer from pathologies, e.g. violation of the energy conditions or superluminal wave propagation. We presented a practical, yet rigorous and general, framework to build relativistic compact objects containing elastic materials. All configurations that we have con-




FIG. 3. Squared frequency eigenvalues for the radial stability analysis of elastic stars in the quadratic model (11) with $n = 1/2$ and different values of E as a function of the mass or of the compactness (see inset). Solid (dashed) lines correspond to stable (unstable) configurations with $\omega^2 > 0$ ($\omega^2 < 0$). In all cases the zero crossing corresponds to the maximum mass within numerical accuracy.

structed have $p_{\text{tan}} \geq p_{\text{rad}}$ within the star. Interestingly, this condition violates the mild-anisotropy assumption of the Buchdahl’s theorem [16, 17], and in fact elasticity allows for compact objects which can exceed the Buchdahl’s limit, $M/R > 4/9$. However, in all models we have explored this limit is reached only in the branch that is unstable against radial perturbations. Indeed, we advocate the importance of checking the (nontrivial) viability of the matter fields for anisotropic configurations, in particular the causality conditions (of longitudinal and transverse waves along the radial and tangential directions) and the radial stability of the solution. While these requirements limit the maximum mass and compactness of elastic stars, we showed that it is nevertheless possible to obtain viable ultracompact configurations featuring a light ring. Elastic stars are one of the few GR models of ultracompact objects that satisfy all the above viability requirements and come from a first-principles Lagrangian approach. The only other known example in the static case are quark stars [17, 39–41] that however require strange mat-

ter. Due to the photon sphere, nonradial perturbations of ultracompact elastic stars can feature gravitational-wave echoes [42–44], as in the quark star case [17, 39–41]. Horizonless compact objects with an unstable light ring feature a second, stable, inner photon sphere, which might be prone to nonlinear instabilities [27, 45, 46]. Investigating this problem in detail was so far hampered by the lack of viable and first-principle solutions, but it can be done within our framework. Indeed, by solving the null geodesic equations we have confirmed that ultracompact elastic stars do feature a stable photon ring.

Elasticity tends to support more massive and compact configurations. For the models at hand the maximum mass can increase up to $\approx 25\%$ relative to the perfect-fluid case while the material remains viable. Interestingly, this suggests that static neutron star models, that can reach $M \approx 2M_\odot$ in the perfect-fluid case, could potentially be as massive as $M \approx 2.5M_\odot$ when elasticity is included, without violating any physical requirement. Such heavy neutron stars would be compatible with the exotic secondary object of the gravitational-wave event GW190814 [47].

Further natural applications of our framework include considering: (i) other models of elastic materials, including non-flat reference material metrics; (ii) consistent multilayer solutions, e.g. made of a perfect-fluid interior and an outer elastic crust [1, 2, 48]; (iii) models that deform generic barotropic fluids beyond the polytropic equation of state. We will report on these applications elsewhere [29]. Finally, we focused here on spherical symmetry but our approach can be extended to less symmetric configurations. In particular, future work will also focus on rotating and deformed [48] elastic solutions.

Acknowledgments. A.A. and J.N. were partially supported by FCT/Portugal through project UIDP/MAT/04459/2020. P.P. acknowledges financial support provided under the European Union’s H2020 ERC, Starting Grant agreement no. DarkGRA–757480, and under the MIUR PRIN and FARE programmes (GW-NEXT, CUP: B84I20000100001), and support from the Amaldi Research Center funded by the MIUR program ‘Dipartimento di Eccellenza’ (CUP: B81I18001170001).

-
- [1] N. Chamel and P. Haensel, ‘‘Physics of Neutron Star Crusts,’’ *Living Rev. Rel.* **11** (2008) 10, [arXiv:0812.3955](https://arxiv.org/abs/0812.3955) [astro-ph].
- [2] L. Suleiman, M. Fortin, J. L. Zdunik, and P. Haensel, ‘‘Influence of the crust on the neutron star macrophysical quantities and universal relations,’’ [arXiv:2106.12845](https://arxiv.org/abs/2106.12845) [astro-ph.HE].
- [3] J. M. Lattimer and M. Prakash, ‘‘The physics of neutron stars,’’ *Science* **304** (2004) 536–542, [arXiv:astro-ph/0405262](https://arxiv.org/abs/astro-ph/0405262) [astro-ph].
- [4] B. Carter and H. Quintana, ‘‘Foundations of general

relativistic high-pressure elasticity theory,’’ *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* **331** no. 1584, (1972) 57–83.

<http://www.jstor.org/stable/78402>.

- [5] J. Park, ‘‘Spherically Symmetric Static Solutions of the Einstein Equations with Elastic Matter Source,’’ *General Relativity and Gravitation* **32** (Feb., 2000) 235–252, [arXiv:gr-qc/9810010](https://arxiv.org/abs/gr-qc/9810010) [gr-qc].
- [6] M. Karlovini and L. Samuelsson, ‘‘Elastic stars in general relativity. 1. Foundations and equilibrium models,’’ *Class. Quant. Grav.* **20** (2003) 3613–3648,

- [arXiv:gr-qc/0211026](https://arxiv.org/abs/gr-qc/0211026).
- [7] J. Frauendiener and A. Kabobel, “The Static spherically symmetric body in relativistic elasticity,” *Class. Quant. Grav.* **24** (2007) 4817–4838, [arXiv:0707.0110](https://arxiv.org/abs/0707.0110) [gr-qc].
- [8] H. Andréasson and S. Calogero, “Spherically symmetric steady states of John elastic bodies in general relativity,” *Class. Quant. Grav.* **31** (2014) 165008, [arXiv:1403.2565](https://arxiv.org/abs/1403.2565) [gr-qc].
- [9] A. Alho and S. Calogero, “Multi-body spherically symmetric steady states of Newtonian self-gravitating elastic matter,” *Commun. Math. Phys.* **371** no. 3, (2019) 975–1004, [arXiv:1807.03062](https://arxiv.org/abs/1807.03062) [math-ph].
- [10] A. Alho and S. Calogero, “Static self-gravitating Newtonian elastic balls,” *Arch. Ration. Mech. Anal.* **238** (2020) 639–669, [arXiv:1907.09970](https://arxiv.org/abs/1907.09970) [math-ph].
- [11] K. Rajagopal and R. Sharma, “Crystallography of three-flavor quark matter,” *Physical Review D* **74** no. 9, (Nov, 2006) . <http://dx.doi.org/10.1103/PhysRevD.74.094019>.
- [12] K. Rajagopal and R. Sharma, “The crystallography of strange quark matter,” *Journal of Physics G: Nuclear and Particle Physics* **32** no. 12, (Nov, 2006) S483–S490. <http://dx.doi.org/10.1088/0954-3899/32/12/S61>.
- [13] M. G. Alford, A. Schmitt, K. Rajagopal, and T. Schäfer, “Color superconductivity in dense quark matter,” *Reviews of Modern Physics* **80** no. 4, (Nov, 2008) 1455–1515. <http://dx.doi.org/10.1103/RevModPhys.80.1455>.
- [14] V. Cardoso and P. Pani, “Testing the nature of dark compact objects: a status report,” *Living Rev. Rel.* **22** no. 1, (2019) 4, [arXiv:1904.05363](https://arxiv.org/abs/1904.05363) [gr-qc].
- [15] R. Carballo-Rubio, F. Di Filippo, S. Liberati, and M. Visser, “Phenomenological aspects of black holes beyond general relativity,” *Phys. Rev. D* **98** no. 12, (2018) 124009, [arXiv:1809.08238](https://arxiv.org/abs/1809.08238) [gr-qc].
- [16] P. Karageorgis and J. G. Stalker, “Sharp bounds on $2m/r$ for static spherical objects,” *Class. Quant. Grav.* **25** (2008) 195021, [arXiv:0707.3632](https://arxiv.org/abs/0707.3632) [gr-qc].
- [17] A. Urbano and H. Veermäe, “On gravitational echoes from ultracompact exotic stars,” *JCAP* **04** (2019) 011, [arXiv:1810.07137](https://arxiv.org/abs/1810.07137) [gr-qc].
- [18] H. A. Buchdahl, “General Relativistic Fluid Spheres,” *Phys. Rev.* **116** (1959) 1027.
- [19] P. O. Mazur and E. Mottola, “Gravitational vacuum condensate stars,” *Proc. Nat. Acad. Sci.* **101** (2004) 9545–9550, [arXiv:gr-qc/0407075](https://arxiv.org/abs/gr-qc/0407075) [gr-qc].
- [20] G. Raposo, P. Pani, M. Bezares, C. Palenzuela, and V. Cardoso, “Anisotropic stars as ultracompact objects in General Relativity,” *Phys. Rev. D* **99** no. 10, (2019) 104072, [arXiv:1811.07917](https://arxiv.org/abs/1811.07917) [gr-qc].
- [21] P. Pani, “I-Love-Q relations for gravastars and the approach to the black-hole limit,” *Phys. Rev.* **D92** no. 12, (2015) 124030, [arXiv:1506.06050](https://arxiv.org/abs/1506.06050) [gr-qc]. [Erratum: Phys. Rev. D95, no. 4, 049902(2017)].
- [22] N. Uchikata and S. Yoshida, “Slowly rotating thin shell gravastars,” *Class. Quant. Grav.* **33** no. 2, (2016) 025005, [arXiv:1506.06485](https://arxiv.org/abs/1506.06485) [gr-qc].
- [23] N. Uchikata, S. Yoshida, and P. Pani, “Tidal deformability and I-Love-Q relations for gravastars with polytropic thin shells,” *Phys. Rev.* **D94** no. 6, (2016) 064015, [arXiv:1607.03593](https://arxiv.org/abs/1607.03593) [gr-qc].
- [24] P. Beltracchi, P. Gondolo, and E. Mottola, “Slowly rotating gravastars,” [arXiv:2107.00762](https://arxiv.org/abs/2107.00762) [gr-qc].
- [25] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2011.
- [26] S. L. Liebling and C. Palenzuela, “Dynamical Bosen Stars,” *Living Rev. Rel.* **15** (2012) 6, [arXiv:1202.5809](https://arxiv.org/abs/1202.5809) [gr-qc].
- [27] V. Cardoso, L. C. B. Crispino, C. F. B. Macedo, H. Okawa, and P. Pani, “Light rings as observational evidence for event horizons: long-lived modes, ergoregions and nonlinear instabilities of ultracompact objects,” *Phys. Rev. D* **90** no. 4, (2014) 044069, [arXiv:1406.5510](https://arxiv.org/abs/1406.5510) [gr-qc].
- [28] P. V. P. Cunha, C. A. R. Herdeiro, and E. Radu, “Fundamental photon orbits: black hole shadows and spacetime instabilities,” *Phys. Rev. D* **96** no. 2, (2017) 024039, [arXiv:1705.05461](https://arxiv.org/abs/1705.05461) [gr-qc].
- [29] A. Alho, J. Natário, P. Pani, and G. Raposo, “Spherically symmetric elastic bodies in general relativity: a detailed analysis (in preparation),”.
- [30] R. Beig and B. G. Schmidt, “Relativistic elasticity,” *Class. Quant. Grav.* **20** (2003) 889–904, [arXiv:gr-qc/0211054](https://arxiv.org/abs/gr-qc/0211054).
- [31] J. D. Brown, “Elasticity Theory in General Relativity,” *Class. Quant. Grav.* **38** no. 8, (2021) 085017, [arXiv:2004.03641](https://arxiv.org/abs/2004.03641) [gr-qc].
- [32] S. Chandrasekhar, *An Introduction to the Study of Stellar Structure*. London: Cambridge University Press, 1939.
- [33] A. Alho, S. Calogero, and A. Liljenberg, “Self-gravitating static balls of power-law elastic matter,” *arXiv e-prints* (Apr., 2021) [arXiv:2104.11131](https://arxiv.org/abs/2104.11131), [arXiv:2104.11131](https://arxiv.org/abs/2104.11131) [math-ph].
- [34] S. Calogero, “On self-gravitating polytropic elastic balls,” *arXiv e-prints* (Apr., 2021) [arXiv:2104.11126](https://arxiv.org/abs/2104.11126), [arXiv:2104.11126](https://arxiv.org/abs/2104.11126) [math.AP].
- [35] A. D. Rendall and B. G. Schmidt, “Existence and properties of spherically symmetric static fluid bodies with a given equation of state,” *Class. Quant. Grav.* **8** (1991) 985–1000.
- [36] T. W. Baumgarte and A. D. Rendall, “Regularity of spherically symmetric static solutions of the Einstein equations,” *Classical and Quantum Gravity* **10** no. 2, (Feb., 1993) 327–332.
- [37] M. Karlović, L. Samuelsson, and M. Zarroug, “Elastic stars in general relativity. 2. Radial perturbations,” *Class. Quant. Grav.* **21** (2004) 1559–1581, [arXiv:gr-qc/0309056](https://arxiv.org/abs/gr-qc/0309056).
- [38] K. D. Kokkotas and J. Ruoff, “Radial oscillations of relativistic stars,” *Astron. Astrophys.* **366** (2001) 565, [arXiv:gr-qc/0011093](https://arxiv.org/abs/gr-qc/0011093).
- [39] M. Mannarelli and F. Tonelli, “Gravitational wave echoes from strange stars,” *Phys. Rev. D* **97** no. 12, (2018) 123010, [arXiv:1805.02278](https://arxiv.org/abs/1805.02278) [gr-qc].
- [40] J. Bora and U. D. Goswami, “Radial oscillations and gravitational wave echoes of strange stars for various equations of state,” *Mon. Not. Roy. Astron. Soc.* **502** no. 2, (2021) 1557–1568, [arXiv:2007.06553](https://arxiv.org/abs/2007.06553) [gr-qc].
- [41] C. Zhang, “Gravitational wave echoes from interacting quark stars,” [arXiv:2107.09654](https://arxiv.org/abs/2107.09654) [hep-ph].
- [42] V. Cardoso, E. Franzin, and P. Pani, “Is the gravitational-wave ringdown a probe of the event horizon?,” *Phys. Rev. Lett.* **116** no. 17, (2016) 171101, [arXiv:1602.07309](https://arxiv.org/abs/1602.07309) [gr-qc]. [Erratum: Phys. Rev. Lett. 117, no. 8, 089902(2016)].
- [43] V. Cardoso and P. Pani, “Tests for the existence of black

- holes through gravitational wave echoes,” *Nature Astron.* **1** no. 9, (2017) 586–591, [arXiv:1709.01525 \[gr-qc\]](#).
- [44] P. Pani and V. Ferrari, “On gravitational-wave echoes from neutron-star binary coalescences,” *Class. Quant. Grav.* **35** no. 15, (2018) 15LT01, [arXiv:1804.01444 \[gr-qc\]](#).
- [45] P. V. P. Cunha, E. Berti, and C. A. R. Herdeiro, “Light-Ring Stability for Ultracompact Objects,” *Phys. Rev. Lett.* **119** no. 25, (2017) 251102, [arXiv:1708.04211 \[gr-qc\]](#).
- [46] R. Ghosh and S. Sarkar, “Light rings of stationary spacetimes,” [arXiv:2107.07370 \[gr-qc\]](#).
- [47] R. Abbott *et al.*, “GW190814: Gravitational Waves from the Coalescence of a 23 Solar Mass Black Hole with a 2.6 Solar Mass Compact Object,” *Astrophys. J.* **896** no. 2, (2020) L44, [arXiv:2006.12611 \[astro-ph.HE\]](#).
- [48] G. Raposo and P. Pani, “Axisymmetric deformations of neutron stars and gravitational-wave astronomy,” *Phys. Rev. D* **102** no. 4, (2020) 044045, [arXiv:2002.02555 \[gr-qc\]](#).