

DIAGRAMMATICS FOR F_4

RAJ GANDHI, ALISTAIR SAVAGE, AND KIRILL ZAINOULLINE

ABSTRACT. We define a diagrammatic monoidal category, together with a full and essentially surjective monoidal functor from this category to the category of modules over the exceptional Lie algebra of type F_4 . In this way, we obtain a set of diagrammatic tools for studying type F_4 representation theory that are analogous to those of the oriented and unoriented Brauer categories in classical type.

1. INTRODUCTION

If $V = \mathbb{C}^d$ denotes the natural module for the complex general linear Lie algebra \mathfrak{gl}_d , Schur–Weyl duality states that the natural actions of \mathfrak{gl}_d and the symmetric group \mathfrak{S}_k on $V^{\otimes k}$ commute and generate each other’s centralizers. This classic result can be extended to the definition of a full monoidal functor from the diagrammatic *oriented Brauer category* \mathcal{OB}_d of [BCNR17] to the category of \mathfrak{gl}_d -modules. This functor sends the two generating objects \uparrow and \downarrow of \mathcal{OB}_d to V and its dual V^* . Since $\mathrm{Hom}_{\mathcal{OB}_d}(\uparrow^{\otimes k}) \cong \mathbb{C}\mathfrak{S}_k$, we have an induced surjective algebra homomorphism

$$\mathbb{C}\mathfrak{S}_k \twoheadrightarrow \mathrm{End}_{\mathfrak{gl}_d}(V^{\otimes k}),$$

recovering one of the statements of Schur–Weyl duality. After passing to the additive Karoubi envelope $\mathrm{Kar}(\mathcal{OB}_d)$ of \mathcal{OB}_d , the above functor is also essentially surjective. It follows that $\mathfrak{gl}_d\text{-mod}$ is a quotient of $\mathrm{Kar}(\mathcal{OB}_d)$ by a tensor ideal. This observation allows one to use powerful and intuitive diagrammatic techniques in the study of the representation theory of the general linear Lie algebra. It also leads to the definition of Deligne’s interpolating categories.

Analogues of the above theory also exist in types BCD . Here the oriented Brauer category is replaced by the *Brauer category* of [LZ15], whose endomorphism algebras are *Brauer algebras*. However, analogous techniques in *exceptional* type are not as well developed. For type G_2 , the diagrammatic category has been described by Kuperberg [Kup94, Kup96]. (In fact, Kuperberg treats the quantum case; see below.) Invariant tensors for classical and exceptional semisimple Lie algebras have been computed diagrammatically by Cvitanović [Cvi08], but the approach there is rather different, inspired by the language of Feynman diagrams in quantum field theory. This approach has been further investigated for exceptional Lie algebras in [MT07, Wen03, Wes03].

The goal of the current paper is to develop a diagrammatic category for type F_4 analogous to the oriented and unoriented Brauer categories in classical type. Hints of the defining relations appear in the aforementioned papers. In particular, several of the equations deduced in the current paper can be found in [Cvi08, Ch. 19] and [Thu04] in a different language. However, a complete treatment from the monoidal category point of view seems to be new.

Given a field \mathbb{k} of characteristic zero, we define a strict monoidal \mathbb{k} -linear category $\mathcal{F} = \mathcal{F}_{\alpha, \delta}$, depending on two parameters $\alpha, \delta \in \mathbb{k}$. (In fact, up to isomorphism, the category is independent of α ; see Remark 2.2.) We consider the strict \mathbb{k} -linear monoidal category generated by a single object $\mathbb{1}$ and four morphisms

$$\begin{array}{c} \diagup \diagdown : \mathbb{1}^{\otimes 2} \rightarrow \mathbb{1}, \quad \times : \mathbb{1}^{\otimes 2} \rightarrow \mathbb{1}^{\otimes 2}, \quad \cup : \mathbb{1} \rightarrow \mathbb{1}^{\otimes 2}, \quad \cap : \mathbb{1}^{\otimes 2} \rightarrow \mathbb{1}, \end{array}$$

2020 *Mathematics Subject Classification.* 18M05, 18M30, 17B10, 17B25, 20G41.

Key words and phrases. Monoidal category, string diagram, exceptional Lie algebra, F_4 , Albert algebra.

where $\mathbb{1}$ is the unit object. These morphisms are subject to certain relations, which we split into two families. We denote by $\mathcal{T} = \mathcal{T}_{\alpha, \delta}$ the category obtained by imposing the first family of relations; see Definition 2.1. These relations imply, in particular, that the category is symmetric monoidal and strict pivotal, the trivalent vertex is symmetric, and the generating object $\mathbb{1}$ is symmetrically self-dual of categorical dimension δ . To obtain the category \mathcal{F} , we then impose three additional relations; see Definition 2.5. The first of these is the relation

$$(1.1) \quad \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} + \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array} = \frac{2\alpha}{\delta + 2} \left(\begin{array}{c} | \\ | \end{array} + \begin{array}{c} \cup \\ \cap \end{array} + \begin{array}{c} \times \\ \times \end{array} \right),$$

while the other two express the square and pentagon in terms of acyclic diagrams.

When $\mathbb{k} = \mathbb{C}$, we define (Theorems 5.1 and 5.3) a monoidal functor

$$\Phi: \mathcal{F}_{7/3, 26} \rightarrow \mathfrak{g}\text{-mod},$$

where \mathfrak{g} is the complex simple Lie algebra of type F_4 . The generating object $\mathbb{1}$ is sent to the natural \mathfrak{g} -module V , which is 26-dimensional. The compact Lie group G corresponding to \mathfrak{g} is the group of algebra automorphisms of the Albert algebra which, over the complex numbers, is the unique exceptional Jordan algebra. The natural module V can be identified with the traceless part of the Albert algebra. Multiplication in the Albert algebra gives rise to a \mathfrak{g} -module homomorphism $V^{\otimes 2} \rightarrow V$, which is the image under Φ of the trivalent vertex $\begin{array}{c} \diagup \\ \diagdown \diagup \end{array}$. The morphism $\begin{array}{c} \times \\ \times \end{array}$ corresponds to the symmetric braiding in $\mathfrak{g}\text{-mod}$, and \cap is sent to the natural invariant bilinear form on V coming from the trace on the Albert algebra. The morphism \cup is also determined by this bilinear form. In this way, the category \mathcal{F} can also be viewed as a diagrammatic calculus for the Albert algebra. The relation (1.1) corresponds to the Cayley–Hamilton theorem for V ; see Remark 5.4.

The functor Φ is defined only when $\delta = 26$, since that is the dimension of the natural \mathfrak{g} -module V . However, the diagrammatic category $\mathcal{F}_{\alpha, \delta}$ is defined for any $\delta \neq -2$. (When $\delta = -2$, the preliminary category \mathcal{T} collapses to the Temperley–Lieb category; see Proposition 2.7.) The importance of the case $\delta = 26$ can be seen purely diagrammatically. It corresponds to the fact that V is not a summand of the tensor square of the first fundamental representation of \mathfrak{g} . See Remark 6.7.

Since the category $\mathfrak{g}\text{-mod}$ is idempotent complete, the functor Φ induces a functor

$$\text{Kar}(\Phi): \text{Kar}(\mathcal{F}_{7/3, 26}) \rightarrow \mathfrak{g}\text{-mod}.$$

Then $\text{Kar}(\Phi)$ is full since Φ is. In addition, we show (Proposition 5.5) that $\text{Kar}(\Phi)$ is essentially surjective. Thus $\mathfrak{g}\text{-mod}$ is equivalent to a quotient of the diagrammatic category \mathcal{F} by a tensor ideal. In fact, we conjecture that $\text{Kar}(\Phi)$ is also faithful, and hence an equivalence of categories. We also give (Conjecture 5.8) conjectural bases for the morphism spaces in \mathcal{F} .

One immediate consequence of the fullness of Φ (Corollary 5.6) is that we have surjective algebra homomorphisms

$$\text{End}_{\mathcal{F}}(\mathbb{1}^{\otimes k}) \twoheadrightarrow \text{End}_{\mathfrak{g}}(V^{\otimes k}), \quad k \in \mathbb{N}.$$

In other words, the endomorphism algebras in \mathcal{F} play the role in type F_4 that the oriented and unoriented Brauer algebras play in the classical types.

In classical types, quantum versions of the relevant diagrammatic categories exist. In type A , the quantum analogue of the oriented Brauer category is the framed HOMFLYPT skein category. In types BCD , the analogue of the unoriented Brauer category is the Kauffman skein category. As their names suggest, both categories are closely related to important knot invariants. In type G_2 , the connection to the corresponding Reshetikhin–Turaev invariant is discussed in [Kup94, Kup96]. There should also exist a quantum analogue of the category \mathcal{F} , related to the Reshetikhin–Turaev invariant in type F_4 . These quantum diagrammatics should also be related to a quantum version of the Albert algebra.

The category \mathcal{F} is also a first step towards a category of webs for type F_4 . The main goal in the theory of webs is to give a presentation, in terms of generators and relations, for the full monoidal subcategory of the category of modules for a quantized enveloping algebra, generated by the fundamental modules. Such presentations are typically in terms of diagrammatic categories known as *web categories*. Web categories were first developed for rank two simple complex Lie algebras by Kuperberg [Kup96]. Then, in more general type A , they were described by Cautis–Kamnitzer–Morrison [CKM14]. More recently, the type C case has been treated in [BERT21]; see also [Wes08]. The degenerate (that is, $q = 1$) web category for type F_4 should be the full monoidal subcategory of $\text{Kar}(\mathcal{F})$ generated by objects corresponding to the four fundamental modules. We explicitly identify three of these objects in Section 6.

Acknowledgements. The research of R.G. was supported by an Ontario Graduate Scholarship and a Canada Graduate Scholarship from the Natural Sciences and Engineering Research Council of Canada (NSERC). The research of A.S. and K.Z. was supported by NSERC Discovery Grants RGPIN-2017-03854 and RGPIN-2015-04469, respectively. We thank Erhard Neher for helpful conversations and Bruce Westbury for useful remarks on an earlier version of this paper.

2. THE DIAGRAMMATIC CATEGORY

We fix a field \mathbb{k} of characteristic zero. All categories are \mathbb{k} -linear and all algebras and tensor products are over \mathbb{k} unless otherwise specified. We let $\mathbb{1}$ denote the unit object of a monoidal category. For objects X and Y in a category \mathcal{C} , we denote by $\mathcal{C}(X, Y)$ the vector space of morphisms from X to Y .

Definition 2.1. Fix $\alpha, \delta \in \mathbb{k}$. Let $\mathcal{T} = \mathcal{T}_{\alpha, \delta}$ be the strict monoidal category generated by the object $\mathbb{1}$ and generating morphisms

$$(2.1) \quad \frown : \mathbb{1} \otimes \mathbb{1} \rightarrow \mathbb{1}, \quad \bowtie : \mathbb{1} \otimes \mathbb{1} \rightarrow \mathbb{1} \otimes \mathbb{1}, \quad \cup : \mathbb{1} \rightarrow \mathbb{1} \otimes \mathbb{1}, \quad \cap : \mathbb{1} \otimes \mathbb{1} \rightarrow \mathbb{1},$$

subject to the following relations:

$$(2.2) \quad \cup \cap = \text{id}, \quad \Upsilon := \cup \frown = \frown \cup, \quad \cap \bowtie = \bowtie \cap,$$

$$(2.3) \quad \bowtie = \text{id}, \quad \bowtie \bowtie = \bowtie \bowtie, \quad \cup \bowtie = \bowtie \cup, \quad \bowtie \cup = \cup \bowtie,$$

$$(2.4) \quad \cap \cup = \cap, \quad \cup \frown = \frown, \quad \frown \cup = \alpha \text{id}, \quad \circ = \delta \mathbb{1}_{\mathbb{1}}, \quad \circ = 0.$$

Remark 2.2. As long as α is invertible and has a square root in \mathbb{k} , we can rescale \frown by $\alpha^{-1/2}$ to see that $\mathcal{T}_{\alpha, \delta}$ is isomorphic to $\mathcal{T}_{1, \delta}$. However, it will be useful in the forthcoming applications to include the parameter α in our definition. In particular, we will be most interested in the case where $\alpha = \frac{7}{3}$ and $\delta = 26$.

Remark 2.3. The relations in Definition 2.1 all have conceptual category-theoretic meanings:

- The relations (2.3) and the last equality in (2.2) correspond to the fact that the crossing endows \mathcal{F} with the structure of a symmetric monoidal category.
- The first two equalities in (2.2), together with the first equality in (2.4) assert that the generating object $\mathbb{1}$ is symmetrically self-dual. Then the fourth equality in (2.2) implies that \mathcal{F} is strict pivotal. (See the discussion after Proposition 2.4.)
- The second equality in (2.4) can be viewed as stating that \frown corresponds to a commutative binary operation on $\mathbb{1}$.
- The fourth relation in (2.4) states that $\mathbb{1}$ has categorical dimension δ . (See Section 6 for further discussion of categorical dimension.)

- If we want $\mathbb{1}$ to be a simple object (more precisely, $\mathcal{F}(\mathbb{1}, \mathbb{1}) = \mathbb{k}\mathbb{1}$), not isomorphic to $\mathbb{1}$, then the fifth equality corresponds to the fact that there are no nonzero morphisms $\mathbb{1} \rightarrow \mathbb{1}$, while the left-hand side of the third equality in (2.4) must be a scalar multiple (which we denote by α) of the identity $\mathbb{1}$.

We define

$$(2.5) \quad \times := \begin{array}{c} \diagup \\ | \\ \diagdown \end{array}.$$

Proposition 2.4. *The following relations hold in \mathcal{T} :*

$$(2.6) \quad \curvearrowright = \curvearrowleft = \curvearrowright, \quad \curvearrowright := \curvearrowright = \curvearrowright, \quad \curvearrowleft := \cup = \cup,$$

$$(2.7) \quad \cup = \cup, \quad \cup \cap = \times = \times, \quad \cup \cap = \times = \times.$$

Proof. The first and second equalities in (2.6) follow immediately from the first four equalities in (2.2). Then, using the first and second equalities in (2.6), we have

$$\curvearrowright = \cap = \curvearrowright,$$

proving the fourth equality in (2.6). The proof of the sixth equality in (2.6) is analogous.

To prove the first equality in (2.7), we use (2.2) to compute

$$\cup = \cup \cup = \cup \cup = \cup.$$

The second and third equalities in (2.7) now follow from sliding the crossing over the cup or cap, and then using the first two equalities in (2.2).

It remains to prove the fourth equality in (2.7). Using the second and third relations in (2.2) and the first two relations in (2.6) to rotate trivalent vertices, we have

$$\cup \cap = \begin{array}{c} \diagup \\ | \\ \diagdown \end{array}.$$

Now, composing the fourth equality in (2.3) on the bottom with \times and on the top with \times , and then using the first equality in (2.3), we have

$$\begin{array}{c} \times \\ \diagdown \\ \diagup \end{array} = \begin{array}{c} \diagup \\ | \\ \diagdown \end{array}.$$

Using this and the second equality in (2.4), we have

$$\begin{array}{c} \diagup \\ | \\ \diagdown \end{array} = \begin{array}{c} \diagup \\ | \\ \diagdown \end{array} = \times,$$

completing the verification of the fourth equality in (2.7). \square

It follows from (2.2), (2.6) and (2.7) that the cups and caps equip \mathcal{F} with the structure of a *strict pivotal* category. Intuitively, this means that morphisms are invariant under ambient isotopy fixing the boundary. Thus, for example, it makes sense to allow horizontal strands in diagrams:

$$(2.8) \quad \begin{array}{c} \diagup \\ | \\ \diagdown \end{array} := \begin{array}{c} \diagup \\ | \\ \diagdown \end{array} = \begin{array}{c} \diagup \\ | \\ \diagdown \end{array}.$$

In addition, since the object $\mathbb{1}$ is self-dual, the cups and caps yield natural isomorphisms

$$(2.9) \quad \mathcal{T}(\mathbb{1}^{\otimes m}, \mathbb{1}^{\otimes n}) \cong \mathcal{T}(\mathbb{1}^{\otimes(m+n)}, \mathbb{1}), \quad n, m \in \mathbb{N}.$$

Definition 2.5. Fix $\alpha, \delta \in \mathbb{k}$, with $\delta \neq -2$. Let $\mathcal{F} = \mathcal{F}_{\alpha, \delta}$ be the strict monoidal category obtained from $\mathcal{T}_{\alpha, \delta}$ by imposing the following three additional relations:

$$(2.10) \quad \begin{array}{c} \diagup \diagdown + \diagdown \diagup + \times \\ \diagup \diagdown + \diagdown \diagup + \times \end{array} = \frac{2\alpha}{\delta+2} \left(\begin{array}{c} | \\ | + \cup + \times \end{array} \right),$$

$$(2.11) \quad \begin{array}{c} \square \\ \square \end{array} = \frac{\alpha^2(\delta+14)}{2(\delta+2)^2} \left(\begin{array}{c} | \\ | + \cup \end{array} \right) + \frac{\alpha(\delta-6)}{2(\delta+2)} \left(\begin{array}{c} \diagup \diagdown + \diagdown \diagup \end{array} \right) + \frac{3\alpha^2(2-\delta)}{2(\delta+2)^2} \times,$$

$$(2.12) \quad \begin{array}{c} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \\ \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \end{array} = \frac{\alpha(10-\delta)}{4(\delta+2)} \left(\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} + \begin{array}{c} \diagdown \diagup \\ \diagdown \diagup \end{array} + \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array} + \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} \right) \\ - \frac{\alpha^2(\delta+30)}{8(\delta+2)^2} \left(\begin{array}{c} \diagdown \\ \diagdown + | \diagup + \diagup + \cup + \diagdown \end{array} \right) \\ + \frac{3\alpha^2(\delta-2)}{8(\delta+2)^2} \left(\begin{array}{c} \diagup \\ \diagup + \diagdown + \diagdown + \diagdown + \diagdown \end{array} \right).$$

Remark 2.6. We will see in Remark 5.4 that (2.10) corresponds to the Cayley–Hamilton theorem for traceless 3×3 octonionic matrices (see (4.9) and (4.10)).

Before proceeding further, let us motivate the assumption in Definition 2.5 that $\delta \neq -2$. In fact, we could multiply both sides of (2.10) to (2.12) by an appropriate power of $\delta+2$ to clear this factor from the denominators. For instance, (2.10) then becomes

$$(2.13) \quad (\delta+2) \left(\begin{array}{c} \diagup \diagdown + \diagdown \diagup + \times \\ \diagup \diagdown + \diagdown \diagup + \times \end{array} \right) = 2\alpha \left(\begin{array}{c} | \\ | + \cup + \times \end{array} \right).$$

Then it makes sense to allow $\delta = -2$. However, in this case, the category collapses considerably, as we now show.

Recall that, for $\delta \in \mathbb{k}$, the *Temperley–Lieb category* \mathcal{TL}_δ is the strict monoidal category generated by the object $|$ and morphisms \cup, \cap , subject to the first two equalities in (2.2) and the fourth equality in (2.4).

Proposition 2.7. *The category obtained from $\mathcal{T}_{\alpha, -2}$ by imposing the relation (2.13), with $\delta = -2$, is isomorphic to \mathcal{TL}_{-2} .*

Proof. When $\delta = -2$, (2.13) gives $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = - \begin{array}{c} | \\ | - \cup \end{array}$. Composing on the top with $\begin{array}{c} \diagdown \\ \diagdown \end{array}$ and using the second and fifth equalities in (2.4) then gives $\begin{array}{c} \diagdown \\ \diagdown \end{array} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = 0$. Is it then a straightforward exercise to see that all the relations in Definition 2.1 are either trivial or follow from the first two equalities in (2.2) and the fourth equality in (2.4). \square

3. DIMENSION RESTRICTIONS

In this section we show that, with some mild restrictions on δ , any quotient of the category \mathcal{T} with certain conditions on the dimensions of the morphism spaces $|^{\otimes 2} \rightarrow |^{\otimes 2}$ and $|^{\otimes 2} \rightarrow |^{\otimes 3}$ satisfies the additional relations (2.10) to (2.12), and hence is also a quotient of \mathcal{F} . Later, in Section 5, we will use this result show that the functor $\mathcal{T} \rightarrow \mathfrak{g}\text{-mod}$ defined in Theorem 5.1 factors through \mathcal{F} (Theorem 5.3).

Recall that an *ideal* in a \mathbb{k} -linear category \mathcal{C} is a collection \mathcal{I} of vector subspaces $\mathcal{I}(X, Y)$ of $\mathcal{C}(X, Y)$ for all $X, Y \in \text{Ob}(\mathcal{C})$ such that

$$\mathcal{C}(Y, Z) \circ \mathcal{I}(X, Y) \subseteq \mathcal{I}(X, Z) \quad \text{and} \quad \mathcal{I}(X, Y) \circ \mathcal{C}(Z, X) \subseteq \mathcal{I}(Z, Y)$$

for all $X, Y, Z \in \text{Ob}(\mathcal{C})$. If, in addition, \mathcal{C} is a monoidal category, then we say \mathcal{I} is a *tensor ideal* if it is an ideal and

$$1_Z \otimes \mathcal{I}(X, Y) \subseteq \mathcal{I}(Z \otimes X, Z \otimes Y) \quad \text{and} \quad \mathcal{I}(X, Y) \otimes 1_Z \subseteq \mathcal{I}(X \otimes Z, Y \otimes Z)$$

for all $X, Y, Z \in \text{Ob}(\mathcal{C})$. If \mathcal{I} is a tensor ideal, it follows that $f \otimes g$ and $g \otimes f$ belong to \mathcal{I} for arbitrary morphisms f in \mathcal{I} and g in \mathcal{C} . If \mathcal{I} is a tensor ideal of \mathcal{C} , then the *quotient category* \mathcal{C}/\mathcal{I} is the category with

$$\text{Ob}(\mathcal{C}/\mathcal{I}) = \text{Ob}(\mathcal{C}), \quad (\mathcal{C}/\mathcal{I})(X, Y) = \mathcal{C}(X, Y)/\mathcal{I}(X, Y).$$

The composition and tensor product in \mathcal{C}/\mathcal{I} are induced by those in \mathcal{C} .

Theorem 3.1. *Assume $\delta \notin \{-2, 2, 6, 10\}$. If \mathcal{I} is a tensor ideal of \mathcal{T} such that*

$$(3.1) \quad \dim((\mathcal{T}/\mathcal{I})(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2})) = 5 \quad \text{and} \quad \dim((\mathcal{T}/\mathcal{I})(\mathbb{1}^{\otimes 3}, \mathbb{1}^{\otimes 2})) \leq 15$$

then relations (2.10) to (2.12) hold in \mathcal{T}/\mathcal{I} .

The proof of Theorem 3.1 will occupy the remainder of this section. We break the proof into a series of smaller results.

We assume for the remainder of this section that $\delta \neq -2$.

Consider the linear operators

$$\begin{aligned} \text{Rot}: \mathcal{T}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2}) &\rightarrow \mathcal{T}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2}), & \text{Rot} \left(\begin{array}{|c|} \hline \boxed{f} \\ \hline \end{array} \right) &= \begin{array}{|c|} \hline \boxed{f} \\ \hline \cup \\ \hline \end{array}, \\ \text{Switch}: \mathcal{T}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2}) &\rightarrow \mathcal{T}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2}), & \text{Switch} \left(\begin{array}{|c|} \hline \boxed{f} \\ \hline \end{array} \right) &= \begin{array}{|c|} \hline \boxed{f} \\ \hline \times \\ \hline \end{array}. \end{aligned}$$

In other words,

$$\text{Rot}(f) = (\cap \otimes \mathbb{1}_1^{\otimes 2}) \circ (\mathbb{1}_1 \otimes f \otimes \mathbb{1}_1) \circ (\mathbb{1}_1^{\otimes 2} \otimes \cup), \quad \text{Switch}(f) = f \circ \times.$$

Any tensor ideal of \mathcal{T} or \mathcal{F} is invariant under Rot. We have

$$(3.2) \quad \begin{aligned} \text{Rot} \left(\begin{array}{|c|} \hline | \\ \hline \end{array} \right) &= \cup, & \text{Rot} \left(\begin{array}{|c|} \hline \cup \\ \hline \end{array} \right) &= |, & \text{Rot}(\times) &= \times, \\ \text{Rot}(\succ) &= \times, & \text{Rot}(\times) &= \succ, & \text{Rot}(\times) &= \times. \end{aligned}$$

and

$$(3.3) \quad \begin{aligned} \text{Switch} \left(\begin{array}{|c|} \hline | \\ \hline \end{array} \right) &= \times, & \text{Switch}(\times) &= |, & \text{Switch} \left(\begin{array}{|c|} \hline \cup \\ \hline \end{array} \right) &= \cup, \\ \text{Switch}(\succ) &= \times, & \text{Switch}(\times) &= \succ, & \text{Switch} \left(\begin{array}{|c|} \hline \times \\ \hline \end{array} \right) &= \times. \end{aligned}$$

Proposition 3.2. *If \mathcal{I} is a tensor ideal of \mathcal{T} such that*

$$(3.4) \quad \dim((\mathcal{T}/\mathcal{I})(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2})) = 5,$$

then (2.10) holds in \mathcal{T}/\mathcal{I} , and the morphisms

$$(3.5) \quad |, \cup, \times, \succ, \times$$

give a basis for $(\mathcal{T}/\mathcal{I})(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2})$.

Proof. Suppose \mathcal{I} is a tensor ideal of \mathcal{T} satisfying (3.4), and let $\mathcal{C} = \mathcal{T}/\mathcal{I}$. Then, in \mathcal{C} , there must be a linear dependence relation involving the six morphisms

$$(3.6) \quad |, \cup, \times, \succ, \times, \times.$$

Let us for a moment view the diagrams in (3.6) as graphs (with \times being a 4-valent vertex) embedded in the plane. The operators Rot and Switch act on this set as in (3.2) and (3.3). It follows that Rot and Switch generate an action of the symmetric group \mathfrak{S}_3 on the 6-dimensional space U spanned by (3.6) (viewed only as embedded planar graphs). The space U decomposes as a direct sum

$$U = U_1 \oplus U_2 \oplus U_3 \oplus U_4,$$

where

$$U_1 = \text{span}_{\mathbb{k}} \left\{ \left| \begin{array}{c} | \\ \cup \\ | \end{array} \right. + \times \right\} \quad \text{and} \quad U_2 = \text{span}_{\mathbb{k}} \left\{ \begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} + \times \right\}$$

are copies of the trivial \mathfrak{S}_3 -module,

$$U_3 = \text{span}_{\mathbb{k}} \left\{ \left| \begin{array}{c} | \\ \cup \\ | \end{array} \right. - 2 \times, \left| \begin{array}{c} | \\ | \\ \cup \end{array} \right. \right\} \quad \text{and} \quad U_4 = \text{span}_{\mathbb{k}} \left\{ \begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} - 2 \times, \begin{array}{c} \diagup \\ \diagdown \end{array} - \begin{array}{c} \diagdown \\ \diagup \end{array} \right\}$$

are copies of the unique simple \mathfrak{S}_3 -module of dimension 2 (that is, the Specht module corresponding to the partition $(2, 1)$), and there is an isomorphism $U_3 \xrightarrow{\cong} U_4$ sending the given basis of U_3 to the given basis of U_4 . If some linear combination u of the elements (3.6) is zero in \mathcal{T}/\mathcal{I} , then all elements of the \mathfrak{S}_3 -submodule generated by u are also zero in \mathcal{T}/\mathcal{I} .

First consider the case where $\mathcal{I}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2}) \cap (U_3 \oplus U_4) \neq \{0\}$. By the above discussion, \mathcal{I} then contains at least the span of the vectors

$$\lambda \left(\begin{array}{c} \diagup \\ \diagdown \end{array} - \begin{array}{c} \diagdown \\ \diagup \end{array} \right) + \mu \left(\left| \begin{array}{c} | \\ | \\ \cup \end{array} \right. \right) \quad \text{and} \quad \lambda \left(\begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} - 2 \times \right) + \mu \left(\left| \begin{array}{c} | \\ \cup \\ | \end{array} \right. - 2 \times \right)$$

for some $\lambda, \mu \in \mathbb{k}$, not both zero. Now, if $\lambda = 0$, then we have $\left| \begin{array}{c} | \\ | \\ \cup \end{array} \right. = \cup$. Composing on the top with $\begin{array}{c} \diagup \\ \diagdown \end{array}$ and using the fourth relation in (2.4), we then get $\begin{array}{c} \diagup \\ \diagdown \end{array} = 0$. It is then clear that (3.4) cannot be satisfied.

Thus, we may suppose $\lambda \neq 0$, in which case relations of the form

$$(3.7) \quad \begin{array}{c} \diagup \\ \diagdown \end{array} - \begin{array}{c} \diagdown \\ \diagup \end{array} = \mu \left(\left| \begin{array}{c} | \\ | \\ \cup \end{array} \right. \right) \quad \text{and} \quad \begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} - 2 \times = \mu \left(\left| \begin{array}{c} | \\ \cup \\ | \end{array} \right. - 2 \times \right)$$

hold in \mathcal{T}/\mathcal{I} . From the first equation in (3.7), we have

$$\begin{array}{c} \diagup \\ \diagdown \end{array} = \begin{array}{c} \diagdown \\ \diagup \end{array} + \mu \left(\left| \begin{array}{c} | \\ | \\ \cup \end{array} \right. \right),$$

which allows us to reduce the length of any cycle of length at least three, or break open the cycle. (The part of a cycle that would be replaced is indicated by dotted lines.) Thus, $(\mathcal{T}/\mathcal{I})(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2})$ is spanned by acyclic diagrams. The second relation in (3.7) then allows us to eliminate \times . This implies that $\left| \begin{array}{c} | \\ \cup \\ | \end{array} \right.$, \cup , $\begin{array}{c} \diagup \\ \diagdown \end{array}$, and $\begin{array}{c} \diagdown \\ \diagup \end{array}$ span $(\mathcal{T}/\mathcal{I})(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2})$, since these are the only planar acyclic trivalent graphs connecting four endpoints. This contradicts our hypothesis (3.4).

We now know that $\mathcal{I}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2}) \subseteq U_1 \oplus U_2$. So \mathcal{I} contains at least the span of the vectors

$$\lambda \left(\left| \begin{array}{c} | \\ \cup \\ | \end{array} \right. + \times \right) + \mu \left(\begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} + \times \right)$$

for some $\lambda, \mu \in \mathbb{k}$, not both zero. If $\mu = 0$, then we have

$$\left| \begin{array}{c} | \\ \cup \\ | \end{array} \right. + \times = 0 \quad \text{in } \mathcal{T}/\mathcal{I}.$$

Composing on the bottom with \cup yields $(\delta + 2)\cup = 0$. If $\cup = 0$, then $1_1 = 0$ by the first relation in (2.2), and so \mathcal{T}/\mathcal{I} is the trivial category. Hence $\delta = -2$ and \mathcal{T}/\mathcal{I} is a quotient of the Temperley–Lieb \mathcal{TL}_{-2} category by Proposition 2.7. Since $\dim(\mathcal{TL}_{-2}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2})) = 2$, this contradicts our hypothesis (3.4).

We may thus assume $\mu \neq 0$. Hence a relation of the form

$$\begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} + \times = \lambda \left(\left| \begin{array}{c} | \\ \cup \\ | \end{array} \right. + \times \right)$$

holds in \mathcal{T}/\mathcal{I} . Now, composing on the bottom with \cup and using (2.4), we have

$$2\alpha\cup = \lambda(\delta + 2)\cup,$$

which implies that $\lambda = \frac{2\alpha}{\delta+2}$. (As explained above, we cannot have $\cup = 0$.) Therefore, (2.10) holds in \mathcal{T}/\mathcal{I} .

It remains to prove that the morphisms (3.5) give a basis for $(\mathcal{T}/\mathcal{I})(\mathbb{I}^{\otimes 2}, \mathbb{I}^{\otimes 2})$. In light of our assumption (3.4), it suffices to show that the morphisms (3.5) are linearly independent. In fact, this already follows from the above discussion. As we saw above, no nonzero element of $U_3 \oplus U_4$ can be zero in \mathcal{T}/\mathcal{I} , and the space $U_1 \oplus U_2$ has dimension one in \mathcal{T}/\mathcal{I} . Therefore, in \mathcal{T}/\mathcal{I} , the span of the morphisms (3.5) is 5-dimensional, as required. \square

Remark 3.3. Composing the relations in (3.7) on the bottom with \cup shows that $(\delta-1)\lambda = \alpha$. (We assume here that $\cup \neq 0$, since, as we saw in the proof of Proposition 3.2, $\cup = 0$ would imply that \mathcal{T}/\mathcal{I} is trivial.) Thus, there is a category \mathcal{D} obtained from \mathcal{T} by imposing the additional relations

$$(3.8) \quad \begin{array}{c} \diagup \\ \diagdown \end{array} - \begin{array}{c} \diagdown \\ \diagup \end{array} = \frac{\alpha}{\delta-1} \left(\begin{array}{c} | \\ | \end{array} - \begin{array}{c} \cup \\ \cup \end{array} \right) \quad \text{and} \quad \begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} - 2 \begin{array}{c} \times \\ \times \end{array} = \frac{\alpha}{\delta-1} \left(\begin{array}{c} | \\ | \end{array} + \begin{array}{c} \cup \\ \cup \end{array} - 2 \begin{array}{c} \times \\ \times \end{array} \right).$$

(We have assumed here that $\delta \neq 1$, since $\delta = 1$ leads to a rather uninteresting category where $\mathbb{I}^{\otimes 2} \cong \mathbb{1}$.)

Other than the Temperley–Lieb category \mathcal{TL}_{-2} and the category with $\mathbb{I}^{\otimes 2} \cong \mathbb{1}$, the above argument shows that there are precisely *two* quotients of \mathcal{T} whose morphisms spaces $\mathbb{I}^{\otimes 2} \rightarrow \mathbb{I}^{\otimes 2}$ have dimension less than or equal to 5. Whereas the goal of the current paper is to examine the category \mathcal{F} and relate it to the representation theory of the Lie algebra of type F_4 , the authors do not know of the significance of the category \mathcal{D} in representation theory. We feel this category merits further investigation.

Lemma 3.4. *If \mathcal{I} is a tensor ideal of \mathcal{T} such that (2.10) holds in \mathcal{T}/\mathcal{I} , then the relation*

$$(3.9) \quad \begin{array}{c} \diagup \\ \diagdown \end{array} = \frac{\alpha(2-\delta)}{2(\delta+2)} \begin{array}{c} \diagdown \\ \diagup \end{array}$$

holds in \mathcal{T}/\mathcal{I} . In particular, (3.9) holds in \mathcal{F} .

Proof. Relation (3.9) follows by composing (2.10) on the top with $\begin{array}{c} \diagdown \\ \diagup \end{array}$, then using (2.4) and (2.7). \square

The first two relations in (2.3) imply that we have an algebra homomorphism

$$(3.10) \quad \mathbb{k}\mathfrak{S}_n \rightarrow \mathcal{T}(\mathbb{I}^{\otimes n}, \mathbb{I}^{\otimes n}),$$

sending the simple transposition $s_i \in \mathfrak{S}_n$ to the crossing of the i -th and $(i+1)$ -st strands. We will denote the image of the complete symmetrizers and antisymmetrizers under this homomorphism by white and black rectangles, respectively:

$$(3.11) \quad \begin{array}{c} \cdots \\ \text{white rectangle} \\ \cdots \end{array} = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \begin{array}{c} \cdots \\ \sigma \\ \cdots \end{array}, \quad \begin{array}{c} \cdots \\ \text{black rectangle} \\ \cdots \end{array} = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\ell(\sigma)} \begin{array}{c} \cdots \\ \sigma \\ \cdots \end{array},$$

where the diagrams contain n strings, \mathfrak{S}_n is the symmetric group on n letters, and $\ell(\sigma)$ is the length of the permutation σ . It then follows from (2.3) and (2.4) that

$$(3.12) \quad \begin{array}{c} \cup \\ \text{white rectangle} \\ \cup \end{array} = \cap, \quad \begin{array}{c} \cup \\ \text{black rectangle} \\ \cup \end{array} = 0, \quad \begin{array}{c} \times \\ \text{white rectangle} \\ \times \end{array} = \begin{array}{c} \diagup \\ \diagdown \end{array}, \quad \begin{array}{c} \times \\ \text{black rectangle} \\ \times \end{array} = - \begin{array}{c} \diagdown \\ \diagup \end{array}, \quad \begin{array}{c} \diagdown \\ \text{white rectangle} \\ \diagup \end{array} = \begin{array}{c} \diagdown \\ \diagup \end{array}, \quad \begin{array}{c} \diagdown \\ \text{black rectangle} \\ \diagup \end{array} = 0.$$

It also follows from (2.7) that

$$(3.13) \quad \begin{array}{c} \diagup \\ \text{white rectangle} \\ \diagdown \end{array} = \begin{array}{c} \diagdown \\ \text{white rectangle} \\ \diagup \end{array}, \quad \begin{array}{c} \diagup \\ \text{black rectangle} \\ \diagdown \end{array} = \begin{array}{c} \diagdown \\ \text{black rectangle} \\ \diagup \end{array}.$$

Lemma 3.5. *If \mathcal{I} is a tensor ideal of \mathcal{T} satisfying (3.4), then the relation (2.11) holds in \mathcal{T}/\mathcal{I} .*

Proof. By Proposition 3.2 and Lemma 3.4, the morphisms (3.5) give a basis for $(\mathcal{T}/\mathcal{I})(\mathbb{I}^{\otimes 2}, \mathbb{I}^{\otimes 2})$ and (3.9) holds. Since $\begin{array}{c} \diagdown \\ \diagup \end{array}$ is invariant under Rot, we must have a relation in \mathcal{T}/\mathcal{I} of the form

$$(3.14) \quad \begin{array}{c} \diagdown \\ \diagup \end{array} = \beta_1 \left(\begin{array}{c} | \\ | \end{array} + \begin{array}{c} \cup \\ \cup \end{array} \right) + \beta_2 \left(\begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} \right) + \beta_3 \begin{array}{c} \times \\ \times \end{array}.$$

Attaching a symmetrizer to the bottom of the diagrams in (2.10) and using (3.12) gives

$$(3.15) \quad 2 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|} \hline \diagup \\ \hline \diagdown \\ \hline \end{array} = \frac{2\alpha}{\delta+2} \left(\begin{array}{|c|} \hline \cup \\ \hline \end{array} + 2 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \right).$$

Attaching a $\begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array}$ to the top of the diagrams in (3.15) and using (2.4) and (3.9) gives

$$(3.16) \quad \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \\ \hline \end{array} = \frac{\alpha}{\delta+2} \left(\alpha \begin{array}{|c|} \hline \cup \\ \hline \end{array} + 2 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \right) + \frac{\alpha(\delta-2)}{4(\delta+2)} \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \stackrel{(3.15)}{=} \frac{4\alpha^2}{(\delta+2)^2} \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + \frac{(\delta+4)\alpha^2}{(\delta+2)^2} \begin{array}{|c|} \hline \cup \\ \hline \end{array} + \frac{(\delta-6)\alpha}{4(\delta+2)} \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array}.$$

On the other hand, attaching a symmetrizer to the bottom of (3.14) gives

$$(3.17) \quad \begin{array}{|c|} \hline \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \\ \hline \end{array} = \beta_1 \left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|} \hline \cup \\ \hline \end{array} \right) + \beta_2 \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right) + \beta_3 \begin{array}{|c|} \hline \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \\ \hline \end{array} \\ \stackrel{(3.15)}{=} \left(\beta_1 + \beta_3 + \frac{2\alpha\beta_2}{\delta+2} \right) \begin{array}{|c|} \hline \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \\ \hline \end{array} + \left(\beta_1 + \frac{\alpha\beta_2}{\delta+2} \right) \begin{array}{|c|} \hline \cup \\ \hline \end{array} + \frac{\beta_2}{2} \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array}.$$

Since the morphisms (3.5) are linearly independent, comparing (3.16) and (3.17) gives

$$\frac{4\alpha^2}{(\delta+2)^2} = \beta_1 + \beta_3 + \frac{2\alpha}{\delta+2}\beta_2, \quad \frac{(\delta+4)\alpha^2}{(\delta+2)^2} = \beta_1 + \frac{\alpha}{\delta+2}\beta_2, \quad \frac{(\delta-6)\alpha}{4(\delta+2)} = \frac{1}{2}\beta_2.$$

Solving this linear system for β_1 , β_2 , and β_3 gives the coefficients of relation (2.11). \square

Lemma 3.6. *If \mathcal{I} is a tensor ideal of \mathcal{T} satisfying (3.1), and $\delta \notin \{2, 6, 10\}$, then the relation (2.12) holds in \mathcal{T}/\mathcal{I} .*

Proof. Since the pentagon is rotation invariant, the assumption (3.1) implies that we must have a relation of the form

$$(3.18) \quad \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \\ \hline \end{array} \\ \hline \end{array} = \gamma_1 \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right) + \gamma_2 \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \cup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \cup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \cup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right) \\ + \gamma_3 \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right).$$

(More precisely, given any relation involving the diagrams appearing in (3.18), we can sum over its images under Rot^n , $0 \leq n \leq 5$, to get a relation of the form (3.18).) Let $\theta = \frac{\alpha(2-\delta)}{2(\delta+2)}$, so that $\begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} = \theta \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array}$. Composing with $\begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array}$ on the rightmost two strings at the top of (3.18) gives

$$\begin{array}{|c|} \hline \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \\ \hline \end{array} = \gamma_1 \left((\theta + \alpha) \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right) + \gamma_2 \left(\alpha \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \cup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right) \\ + \gamma_3 \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + 2 \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} + \alpha \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right).$$

Thus, using (2.10) to eliminate $\begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array}$, we have

$$(\theta\gamma_0 - \gamma_1) \begin{array}{|c|} \hline \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \\ \hline \end{array} = \left(\alpha\gamma_2 + \frac{4\alpha}{\delta+2}\gamma_3 \right) \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \cup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right) + ((\theta + \alpha)\gamma_1 + \gamma_2 - \gamma_3) \left(\begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array} \right) + \frac{\alpha(\delta+6)}{\delta+2}\gamma_3 \begin{array}{|c|} \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline \diagdown \\ \hline \diagup \\ \hline \end{array} \\ \hline \end{array}.$$

Comparing to (2.11) and using the fact that the morphisms (3.5) are linearly independent, this gives

$$\begin{aligned} \frac{\alpha(\delta+14)}{2(\delta+2)^2}(\theta\gamma_0 - \gamma_1) &= \gamma_2 + \frac{4}{\delta+2}\gamma_3, \\ \frac{\alpha(\delta-6)}{2(\delta+2)}(\theta\gamma_0 - \gamma_1) &= (\theta + \alpha)\gamma_1 + \gamma_2 - \gamma_3, \\ \frac{3\alpha(2-\delta)}{2(\delta+2)}(\theta\gamma_0 - \gamma_1) &= (\delta+6)\gamma_3. \end{aligned}$$

Assuming $\delta \neq 2$, this is equivalent to the linear system

$$(3.19) \quad \begin{aligned} \gamma_2 + \frac{\delta + 30}{3(\delta - 2)}\gamma_3 &= 0, & \frac{\alpha(\delta + 6)}{2(\delta + 2)}\gamma_1 + \gamma_2 + \frac{\delta^2 - 3\delta - 30}{3(\delta - 2)}\gamma_3 &= 0, \\ \frac{3\alpha(2 - \delta)}{2(\delta + 2)}\gamma_1 + (\delta + 6)\gamma_3 &= \frac{3\alpha^2(2 - \delta)^2}{4(\delta + 2)^2}\gamma_0. \end{aligned}$$

Subtracting the first equation above from the second gives

$$\frac{\alpha(\delta + 6)}{2(\delta + 2)} + \frac{(\delta + 6)(\delta - 10)}{3(\delta - 2)}\gamma_3 = 0 \iff \gamma_3 = \frac{3\alpha(2 - \delta)}{2(\delta + 2)(\delta - 10)}\gamma_1.$$

as long as $\delta \neq 6, 10$. Combining with (3.19) then gives

$$\begin{aligned} \frac{3\alpha(2 - \delta)}{2(\delta + 2)}\gamma_1 + \frac{3\alpha(\delta + 6)(2 - \delta)}{2(\delta + 2)(\delta - 10)}\gamma_1 &= \frac{3\alpha^2(2 - \delta)^2}{4(\delta + 2)^2}\gamma_0 \\ \implies -\frac{3\alpha(\delta - 2)^2}{(\delta + 2)(\delta - 10)}\gamma_1 &= \frac{3\alpha^2(2 - \delta)^2}{4(\delta + 2)^2}\gamma_0 \implies \gamma_1 = \frac{\alpha(10 - \delta)}{4(\delta + 2)}\gamma_0. \end{aligned}$$

Thus

$$\gamma_3 = \frac{3\alpha(2 - \delta)}{2(\delta + 2)(\delta - 10)}\gamma_1 = \frac{3\alpha^2(\delta - 2)}{8(\delta + 2)^2}\gamma_0 \quad \text{and} \quad \gamma_2 = \frac{\delta + 30}{3(2 - \delta)}\gamma_3 = -\frac{\alpha^2(\delta + 30)}{8(\delta + 2)^2}\gamma_0.$$

We see that the relation (3.18) becomes trivial if $\gamma_0 = 0$, so we may assume $\gamma_0 \neq 0$. Then, dividing (3.18) by γ_0 gives the relation (2.12). \square

Remark 3.7. When $\delta \in \{2, 6, 10\}$, there exist other solutions to the linear system appearing in the proof of Lemma 3.6. We are not sure of the role of these other categories in representation theory. Compare to Remark 3.3.

Proof of Theorem 3.1. The theorem now follows immediately from Proposition 3.2 and Lemmas 3.5 and 3.6. \square

Remark 3.8. When combined with Remark 2.3, Theorem 3.1 implies that every pivotal symmetric monoidal category \mathcal{C} generated by a symmetric self-dual object $\mathbb{1}$ and a rotationally invariant symmetric morphism $\mathbb{1} \rightarrow \mathbb{1}^{\otimes 3}$, and with $\dim \mathcal{C}(\mathbb{1}, \mathbb{1}^{\otimes n})$ equal to 1, 0, 1, 1, 5, 15 for $n = 0, 1, 2, 3, 4, 5$, respectively, is a quotient of \mathcal{F} for some value of δ . Similar categories, with different conditions on the dimensions, were classified in [MPS17]. The corresponding statement for the quantum G_2 link invariant is given in [Kup94, Th. 2.1].

4. THE ALBERT ALGEBRA AND THE LIE GROUP OF TYPE F_4

In this section, we will develop some properties of the Albert algebra and the Lie group and Lie algebra of type F_4 that will be used in the sequel. For further details, we refer the reader to [Ada96, Ch. 16]. Let

$$A = \left\{ \begin{pmatrix} \lambda_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \lambda_2 & x_1 \\ x_2 & \bar{x}_1 & \lambda_3 \end{pmatrix} : \lambda_i \in \mathbb{R}, x_i \in \mathbb{O} \right\}$$

denote the set of 3×3 self-adjoint matrices over the octonions \mathbb{O} , equipped with the bilinear operation

$$a \circ b := \frac{1}{2}(ab + ba), \quad a, b \in A,$$

where the juxtaposition ab denotes usual matrix multiplication. Thus A is one of the three real Albert algebras. Note that this algebra is commutative and unital, but *not* associative. We have $\dim_{\mathbb{R}}(A) = 27$. Eventually, we will be interested in the complexification $\mathbb{C} \otimes_{\mathbb{R}} A$, which is the

unique simple exceptional complex Jordan algebra, up to isomorphism. However, since many of our preliminary arguments are valid over \mathbb{R} , we state them in that setting.

Let $\text{tr}: A \rightarrow \mathbb{R}$ denote the trace map, so that

$$(4.1) \quad \text{tr} \begin{pmatrix} \lambda_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \lambda_2 & x_1 \\ x_2 & \bar{x}_1 & \lambda_3 \end{pmatrix} = \lambda_1 + \lambda_2 + \lambda_3.$$

For $a \in A$, let

$$L_a: A \rightarrow A, \quad b \mapsto a \circ b,$$

denote the \mathbb{R} -linear map given by left multiplication by a .

Lemma 4.1. *For $a \in A$, we have*

$$(4.2) \quad \text{tr}(a) = \frac{1}{9} \text{Tr}(L_a),$$

where $\text{Tr}(L_a)$ denotes the usual trace of the linear operator L_a on the 27-dimensional real vector space A .

Proof. Let E_{ij} denote the matrix with a 1 in the (i, j) -entry and all other entries equal to zero. Since both sides of (4.2) are \mathbb{R} -linear in a , it suffices to consider the cases where $a = xE_{ij} + \bar{x}E_{ji}$ for $x \in \mathbb{O}$, $1 \leq i, j \leq 3$. Consider the basis of A given by the elements

$$yE_{kl} + \bar{y}E_{lk}, \quad k \leq l,$$

where y runs over a basis of \mathbb{O} if $k \neq l$, and $y = \frac{1}{2}$ if $k = l$. Now

$$L_a(yE_{kl} + \bar{y}E_{lk}) = \frac{1}{2} (\delta_{jk}xyE_{il} + \delta_{jl}x\bar{y}E_{ik} + \delta_{ik}\bar{x}yE_{jl} + \delta_{il}\bar{x}\bar{y}E_{jk}).$$

We see that the $yE_{ij} + \bar{y}E_{ji}$ component of this is zero unless $i = j = k$ or $i = j = l$. Thus $\text{tr}(L_a)$ is zero unless a is a diagonal matrix. On the other hand, if $a = \lambda E_{ii}$, $\lambda \in \mathbb{R}$, then L_a acts as multiplication by $\frac{\lambda}{2}(\delta_{ik} + \delta_{il})$ on the subspace $xE_{kl} + \bar{x}E_{lk}$, $x \in \mathbb{O}$, $k \neq l$, and multiplication by λ on the subspace $\mathbb{R}E_{ii}$. Thus $\text{tr}(L_a) = (1 + \frac{1}{2}(16))\lambda = 9\lambda$. \square

Let G denote the group of algebra automorphisms of A . Thus G is the compact connected real Lie group of type F_4 .

Lemma 4.2. *We have $\text{tr}(ga) = \text{tr}(a)$ for all $a \in A$ and $g \in G$.*

Proof. For $a \in A$ and $g \in G$, we have

$$(gL_a g^{-1})(b) = g(a \circ (g^{-1}b)) = (ga) \circ b = L_{ga}(b).$$

Thus $\text{tr}(ga) = \frac{1}{9} \text{Tr}(L_{ga}) = \frac{1}{9} \text{Tr}(gL_a g^{-1}) = \frac{1}{9} \text{Tr}(L_a) = \text{tr}(a)$. \square

Corollary 4.3. *The symmetric bilinear form*

$$(4.3) \quad B: A \otimes A \rightarrow \mathbb{R}, \quad B(a \otimes b) := \text{tr}(a \circ b),$$

is nondegenerate and G -invariant.

We will sometimes write $B(a, b)$ for $B(a \otimes b)$.

Proof. Direct computation shows that, if a is the matrix appearing in (4.1), then

$$B(a, a) = \sum_{i=1}^3 (\lambda_i^2 + \|x_i\|^2),$$

which is nonzero when $a \neq 0$. Hence B is nondegenerate. Since

$$B(ga, gb) = \text{tr}((ga) \circ (gb)) = \text{tr}(g(a \circ b)) = \text{tr}(a \circ b) = B(a \otimes b),$$

we also see that B is G -invariant. □

It follows from Corollary 4.3 that we have a decomposition of G -modules

$$(4.4) \quad A = \mathbb{R}1_A \oplus V_{\mathbb{R}}, \quad V_{\mathbb{R}} := \ker(\text{tr}).$$

Then $V_{\mathbb{R}}$ is the 26-dimensional irreducible G -module ([Ada96, Cor. 16.2]). Let

$$(4.5) \quad \pi: A \rightarrow V_{\mathbb{R}}, \quad \pi(a) = a - \frac{1}{3} \text{tr}(a)1_A$$

be the projection along the decomposition (4.4).

For $x \in \mathbb{O}$, let $\text{Re}(x)$ denote its real part. It is straightforward to verify that

$$(4.6) \quad \text{Re}(xy) = \text{Re}(yx), \quad \text{Re}((xy)z) = \text{Re}(x(yz)), \quad x, y, z \in \mathbb{O}.$$

For $X \in \text{Mat}_{3 \times 3}(\mathbb{O})$, let $\text{tr}_{\mathbb{R}} = \text{Re}(\text{tr}(X))$.

Lemma 4.4. *For $X, Y, Z \in \text{Mat}_{3 \times 3}(\mathbb{O})$, we have*

$$(4.7) \quad \text{tr}_{\mathbb{R}}(XY) = \text{tr}_{\mathbb{R}}(YX) \quad \text{and} \quad \text{tr}_{\mathbb{R}}((XY)Z) = \text{tr}_{\mathbb{R}}(X(YZ)).$$

Proof. Since both sides of both equalities to be proved are \mathbb{R} -linear in X , Y , and Z , it suffices to consider the case where $X = xE_{ij}$, $Y = yE_{kl}$, and $Z = zE_{mn}$, for $x, y, z \in \mathbb{O}$. For the first equality, we have

$$\text{tr}_{\mathbb{R}}(XY) = \delta_{jk}\delta_{il} \text{Re}(xy) \stackrel{(4.6)}{=} \delta_{jk}\delta_{il} \text{Re}(yx) = \text{tr}_{\mathbb{R}}(YX).$$

For the second equality, we have

$$\text{tr}_{\mathbb{R}}((XY)Z) = \delta_{jk}\delta_{lm}\delta_{in} \text{Re}((xy)z) \stackrel{(4.6)}{=} \delta_{jk}\delta_{lm}\delta_{in} \text{Re}(x(yz)) = \text{tr}_{\mathbb{R}}(X(YZ)). \quad \square$$

Lemma 4.5. *We have*

$$(4.8) \quad \text{tr}((a \circ b) \circ c) = \text{tr}(a \circ (b \circ c)), \quad a, b, c \in A.$$

Proof. For $a, b, c \in A$, we have

$$\begin{aligned} 4 \text{tr}((a \circ b) \circ c) &= 4 \text{tr}_{\mathbb{R}}((a \circ b) \circ c) = \text{tr}_{\mathbb{R}}((ab)c + (ba)c + c(ab) + c(ba)) \\ &\stackrel{(4.7)}{=} \text{tr}_{\mathbb{R}}(a(bc) + a(cb) + (bc)a + (cb)a) = 4 \text{tr}_{\mathbb{R}}(a \circ (b \circ c)) = 4 \text{tr}(a \circ (b \circ c)). \quad \square \end{aligned}$$

Lemma 4.6. *For $a \in V_{\mathbb{R}}$, we have*

$$(4.9) \quad \pi(\pi(a \circ a) \circ a) = \frac{1}{6} \text{tr}(a \circ a)a.$$

Proof. Since $\text{tr}(a) = 0$, [Ada96, Th. 16.6(iii)] (which is essentially the Cayley–Hamilton theorem for A) implies that

$$0 = (a \circ a) \circ a - \frac{1}{2} \text{tr}(a \circ a)a - \frac{1}{3} \text{tr}((a \circ a) \circ a) \stackrel{(4.5)}{=} \pi((a \circ a) \circ a) - \frac{1}{2} \text{tr}(a \circ a)a.$$

Now, since $\pi(a) = a$, we have

$$\pi((a \circ a) \circ a) \stackrel{(4.5)}{=} \pi(\pi(a \circ a) \circ a) + \frac{1}{3} \text{tr}(a \circ a)a.$$

The identity (4.9) now follows. □

Corollary 4.7. *For $a, b, c \in V_{\mathbb{R}}$, we have*

$$(4.10) \quad \pi(\pi(a \circ b) \circ c) + \pi(\pi(b \circ c) \circ a) + \pi(\pi(a \circ c) \circ b) = \frac{1}{6} (\text{tr}(b \circ c)a + \text{tr}(a \circ b)c + \text{tr}(a \circ c)b).$$

Proof. This is the polarization of the identity (4.9). That is, we replace a in (4.9) by $\lambda_a a + \lambda_b b + \lambda_c c$, expand, and take the $\lambda_a \lambda_b \lambda_c$ terms. □

Fix a basis \mathbf{B}_V of $V_{\mathbb{R}}$. By Corollary 4.3, there exists a dual basis $\mathbf{B}_V^{\vee} = \{b^{\vee} : b \in \mathbf{B}_V\}$ defined by

$$\mathrm{tr}(a^{\vee} \circ b) = \delta_{a,b}, \quad a, b \in \mathbf{B}_V.$$

We can extend this to a basis $\mathbf{B}_A := \mathbf{B}_V \sqcup \{1\}$ of A , with dual basis $\mathbf{B}_V^{\vee} \sqcup \{\frac{1}{3}\}$, i.e. $1^{\vee} = \frac{1}{3}$. Note that the elements $\sum_{b \in \mathbf{B}_V} b \otimes b^{\vee} \in V_{\mathbb{R}} \otimes V_{\mathbb{R}}$ and $\sum_{b \in \mathbf{B}_A} b \otimes b^{\vee} \in A \otimes A$ are both independent of the choice of bases.

Lemma 4.8. *For $a \in A$, we have*


$$(4.11) \quad \sum_{b \in \mathbf{B}_A} a \circ b \otimes b^{\vee} = \sum_{b \in \mathbf{B}_A} b \otimes b^{\vee} \circ a.$$

Proof. We have

$$\sum_{b \in \mathbf{B}_A} a \circ b \otimes b^{\vee} = \sum_{b, c \in \mathbf{B}_A} \mathrm{tr}(c^{\vee} \circ (a \circ b)) c \otimes b^{\vee} \stackrel{(4.8)}{=} \sum_{b, c \in \mathbf{B}_A} c \otimes \mathrm{tr}((c^{\vee} \circ a) \circ b) b^{\vee} = \sum_{c \in \mathbf{B}_A} c \otimes c^{\vee} \circ a. \quad \square$$

Let $\mathfrak{g} = \mathbb{C} \otimes_{\mathbb{R}} \mathfrak{g}_{\mathbb{R}}$ be the complexification of the Lie algebra $\mathfrak{g}_{\mathbb{R}}$ of G , and let $V = \mathbb{C} \otimes_{\mathbb{R}} V_{\mathbb{R}}$ be the corresponding natural \mathfrak{g} -module. Let $\mathfrak{g}\text{-mod}$ denote the category of finite-dimensional \mathfrak{g} -modules. We continue to denote by tr and B the complexification of the maps (4.1) and (4.3). Then \mathbf{B}_V is also a \mathbb{C} -basis of V with dual basis \mathbf{B}_V^{\vee} . We will continue to use the bar notation $\bar{}$ to denote the conjugation of the octonions, extended to their complexification by \mathbb{C} -linearity.

We conclude this section by recalling some basic facts about the representation theory of \mathfrak{g} . Consider the following labeling of the nodes of the Dynkin diagram of type F_4 :



Let $\omega_1, \omega_2, \omega_3, \omega_4$ denote the corresponding fundamental weights. For a dominant integral weight λ , let V_{λ} denote the simple \mathfrak{g} -module of highest weight λ . In particular $V = V_{\omega_4}$, while V_{ω_1} is the adjoint representation. We have tensor product decompositions

$$(4.12) \quad V^{\otimes 2} = V_0 \oplus V_{\omega_1} \oplus V_{\omega_3} \oplus V_{\omega_4} \oplus V_{2\omega_4},$$

$$(4.13) \quad V^{\otimes 3} = V_0 \oplus V_{\omega_1}^{\oplus 2} \oplus V_{\omega_2} \oplus V_{\omega_3}^{\oplus 4} \oplus V_{\omega_4}^{\oplus 5} \oplus V_{\omega_1 + \omega_4}^{\oplus 3} \oplus V_{\omega_3 + \omega_4}^{\oplus 2} \oplus V_{2\omega_4}^{\oplus 3} \oplus V_{3\omega_4}.$$

This follows, for example from the table given in [MPR90, Ch. 11, Table 7]. By Schur's lemma, we thus have

$$(4.14) \quad \dim \mathrm{Hom}_{\mathfrak{g}}(V^{\otimes 2}, V^{\otimes 2}) = 5 \quad \text{and} \quad \dim \mathrm{Hom}_{\mathfrak{g}}(V^{\otimes 3}, V^{\otimes 2}) = 15$$

The importance of these dimensions is the assumption (3.1) in Theorem 3.1.

5. THE FUNCTOR

In this section we describe a natural functor from the category \mathcal{F} to the category $\mathfrak{g}\text{-mod}$ of finite-dimensional modules over the complex Lie algebra \mathfrak{g} of type F_4 . We do this by first defining a functor from \mathcal{T} , and then showing that it factors through \mathcal{F} . Throughout this section we work over the field $\mathbb{k} = \mathbb{C}$.

Theorem 5.1. *There is a unique monoidal functor*

$$\Phi: \mathcal{T}_{7/3,26} \rightarrow \mathfrak{g}\text{-mod}$$

given on objects by $l \mapsto V$ and on morphisms by

$$(5.1) \quad \Phi(\frown): V \otimes V \rightarrow V, \quad a \otimes b \mapsto \pi(a \circ b),$$

$$(5.2) \quad \Phi(\otimes): V \otimes V \rightarrow V \otimes V, \quad a \otimes b \mapsto b \otimes a,$$

$$(5.3) \quad \Phi(\cup): \mathbb{C} \rightarrow V \otimes V, \quad 1 \mapsto \sum_{b \in \mathbf{B}_V} b \otimes b^\vee,$$

$$(5.4) \quad \Phi(\cap): V \otimes V \rightarrow \mathbb{C}, \quad a \otimes b \mapsto B(a \otimes b) = \text{tr}(a \circ b).$$

Furthermore

$$(5.5) \quad \Phi(\Upsilon): V \rightarrow V \otimes V, \quad a \mapsto \sum_{b \in \mathbf{B}_V} b \otimes \pi(b^\vee \circ a) = \sum_{b \in \mathbf{B}_V} \pi(a \circ b) \otimes b^\vee.$$

Proof. We must verify that Φ respects the defining relations in Definition 2.1. The verification of the relations (2.3) and the first two equalities in (2.2) is straightforward.

For the third equality in (2.2), we first use (4.11) to see that, for $a \in V$,

$$\sum_{b \in \mathbf{B}_V} a \circ b \otimes b^\vee + a \otimes \frac{1}{3} = \sum_{b \in \mathbf{B}_V} b \otimes b^\vee + \frac{1}{3} \otimes a.$$

Applying $\pi \otimes \pi$ yields

$$\sum_{b \in \mathbf{B}_V} \pi(a \circ b) \otimes b^\vee = \sum_{b \in \mathbf{B}_V} b \otimes \pi(b^\vee \circ a).$$

Thus, for $a \in V$, we have

$$\Phi\left(\bigcup\right)(a) = \sum_{b \in \mathbf{B}_V} b \otimes \pi(b^\vee \circ a) = \sum_{b \in \mathbf{B}_V} \pi(a \circ b) \otimes b^\vee = \Phi\left(\bigcap\right)(a).$$

This shows that Φ preserves the third equality in (2.2) and that it satisfies (5.5). For the fourth equality in (2.2), we compute

$$\Phi\left(\bigcap\right)(a \otimes b \otimes c) = B(a \otimes c)b = \Phi\left(\bigcap\right)(a \otimes b \otimes c).$$

The fact that Φ respects the first two relations in (2.4) follows immediately from the fact that $a \circ b = b \circ a$ for $a, b \in V$. Now consider the third relation in (2.4). Since V is an irreducible \mathfrak{g} -module, there exists $\alpha \in \mathbb{C}$ such that

$$\Phi\left(\bigcirc\right)(a) = \alpha a \quad \text{for all } a \in V,$$

and we must show that $\alpha = \frac{7}{3}$. It suffices to show this for some specific choice of a , so we choose $a = E_{11} - E_{22}$. Now, choose the basis

$$\mathbf{B}_V = \{b_1 = \frac{1}{\sqrt{2}}E_{11} - \frac{1}{\sqrt{2}}E_{22}, b_2 = \frac{1}{\sqrt{6}}E_{11} + \frac{1}{\sqrt{6}}E_{22} - \frac{2}{\sqrt{6}}E_{33}\} \sqcup \mathbf{B}'_V, \quad \text{where}$$

$$\mathbf{B}'_V = \{\frac{1}{2}(xE_{ij} + \bar{x}E_{ji}) : 1 \leq i < j \leq 3, x \in \mathbf{B}_\mathbb{O}\},$$

and where $\mathbf{B}_\mathbb{O}$ is the usual basis of \mathbb{O} (in particular, $\bar{x}x = x\bar{x} = 1$ for $x \in \mathbf{B}_\mathbb{O}$). Then \mathbf{B}_V is an orthonormal basis for V , that is, $b^\vee = b$ for all $b \in \mathbf{B}_V$. Furthermore, for $b = \frac{1}{2}(xE_{ij} + \bar{x}E_{ji})$, with $x \in \mathbf{B}_\mathbb{O}$ and $1 \leq i < j \leq 3$, we have

$$(a \circ b) \circ b^\vee = \frac{1}{2}(\delta_{i1} - \delta_{i2} - \delta_{j2})b \circ b = \frac{1}{4}(\delta_{i1} - \delta_{i2} - \delta_{j2})(E_{ii} + E_{jj}).$$

Therefore,

$$\begin{aligned} \sum_{b \in \mathbf{B}_V} (a \circ b) \circ b^\vee &= (a \circ b_1) \circ b_1 + (a \circ b_2) \circ b_2 + \sum_{b \in \mathbf{B}'_V} (a \circ b) \circ b \\ &= \frac{1}{2}(E_{11} - E_{22}) + \frac{1}{6}(E_{11} - E_{22}) + 2 \sum_{i < j} (\delta_{i1} - \delta_{i2} - \delta_{j2})(E_{ii} + E_{jj}) = \frac{8}{3}a. \end{aligned}$$

Thus we have

$$\begin{aligned} \Phi \left(\begin{array}{c} \diagup \\ \circ \\ \diagdown \end{array} \right) (a) &= \sum_{b \in \mathbf{B}_V} \pi(\pi(a \circ b) \circ b^\vee) \stackrel{(4.5)}{=} \sum_{b \in \mathbf{B}_V} \pi((a \circ b) \circ b^\vee) - \frac{1}{3} \sum_{b \in \mathbf{B}_V} B(a \otimes b) \pi(b^\vee) \\ &= \sum_{b \in \mathbf{B}_V} \pi((a \circ b) \circ b^\vee) - \frac{1}{3} \pi(a) = \frac{8}{3} a - \frac{1}{3} a = \frac{7}{3} a, \end{aligned}$$

as desired.

That Φ respects the fourth equality in (2.4) follows immediately from the fact that $\dim_{\mathbb{C}} V = 26$. Since $\Phi \left(\begin{array}{c} \diagup \\ \circ \\ \diagdown \end{array} \right)$ is a homomorphism of \mathfrak{g} -modules from the trivial module to V , it must be zero by Schur's lemma, proving that Φ respects the fifth equality in (2.4). \square

For a linear category \mathcal{C} , let $\text{Kar}(\mathcal{C})$ denote its additive Karoubi envelope. Thus, objects of $\text{Kar}(\mathcal{C})$ are pairs (X, e) , where X is an object in the additive envelope $\text{Add}(\mathcal{C})$ of \mathcal{C} , and $e \in \mathcal{C}(X, X)$ is an idempotent endomorphism. Morphisms in $\text{Kar}(\mathcal{C})$ are given by

$$\text{Kar}(\mathcal{C})((X, e), (X', e')) = e' \text{Add}(\mathcal{C})(X, X') e.$$

Composition is as in \mathcal{C} .

Proposition 5.2. *The functor Φ is full.*

Proof. We follow the method of the proof of [Kup96, Th. 5.1]. Since the category $\mathfrak{g}\text{-mod}$ is idempotent complete, we have an induced functor

$$(5.6) \quad \text{Kar}(\mathcal{T}_{7/3,26}) \rightarrow \mathfrak{g}\text{-mod}.$$

Let \mathcal{C} be the image of this functor. Then \mathcal{C} is a rigid symmetric monoidal category. We claim that $\text{End}_{\mathcal{C}}(V^{\otimes n})$ is a semisimple algebra for all $n \geq 0$. Indeed, consider the conjugate-linear contravariant monoidal endofunctor Ξ of \mathcal{F} determined on objects by $\mathbb{1} \mapsto \mathbb{1}$ and on morphisms by

$$\begin{array}{c} \diagup \\ \diagdown \end{array} \mapsto \begin{array}{c} \diagdown \\ \diagup \end{array}, \quad \begin{array}{c} \diagdown \\ \diagup \end{array} \mapsto \begin{array}{c} \diagup \\ \diagdown \end{array}, \quad \cup \mapsto \cap, \quad \cap \mapsto \cup.$$

Intuitively, Ξ is given by reflecting diagrams in the vertical axis and taking the complex conjugate of all coefficients. Then Φ intertwines Ξ with the hermitian adjoint. It follows that $\text{End}_{\mathcal{C}}(V^{\otimes n})$ is closed under hermitian adjoint, and hence is semisimple. Thus \mathcal{C} satisfies the hypotheses of the Tannaka–Krein duality theorem [Kir76, p. 177], and must be the category of finite-dimensional representations of some compact group H . Since all morphisms of \mathcal{C} are homomorphisms of G -modules, we have $G \subseteq H$. On the other hand, G is precisely the group of automorphisms of V preserving $\Phi \left(\begin{array}{c} \diagup \\ \diagdown \end{array} \right)$ and $\Phi \left(\begin{array}{c} \diagdown \\ \diagup \end{array} \right)$. Thus $G = H$ and so (5.6) is full. Viewing \mathcal{T} as a full subcategory of $\text{Kar}(\mathcal{T})$ in the usual way, we conclude that Φ is full. \square

Theorem 5.3. *The functor Φ from Theorem 5.1 factors through $\mathcal{F}_{7/3,26}$.*

Proof. Let \mathcal{I} be the kernel of the functor Φ . Then \mathcal{I} is a tensor ideal of \mathcal{T} , and we must show that the relations (2.10) to (2.12) are satisfied in \mathcal{T}/\mathcal{I} . By Proposition 5.2 and (4.14), it follows that condition (3.1) is satisfied. Then the theorem follows from Theorem 3.1. \square

Remark 5.4. It is also possible to give a more direct proof that the image of (2.10) under Φ holds in $\mathfrak{g}\text{-mod}$. Bending the top right endpoint to the bottom of the diagrams by tensoring on the right with $\mathbb{1}$ and attaching a cup to the two rightmost endpoints at the top of the diagram (an operation which is invertible by the first relation in (2.2)), we see that (2.10) is equivalent to

$$\begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} + \begin{array}{c} \diagup \\ \diagdown \\ \diagup \\ \diagdown \end{array} = \frac{1}{6} \left(\begin{array}{c} \diagup \\ \cap \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \cap \\ \diagup \end{array} + \begin{array}{c} \diagup \\ \cap \\ \diagup \\ \diagdown \end{array} \right).$$

The fact that Φ respects this relation follows from (4.10) and the fact that the operation \circ on A is commutative. Hence, we see that (2.10) essentially corresponds to the Cayley–Hamilton theorem for the Albert algebra.

Since the category $\mathfrak{g}\text{-mod}$ is idempotent complete, we have an induced functor

$$(5.7) \quad \text{Kar}(\Phi): \text{Kar}(\mathcal{F}_{7/3,26}) \rightarrow \mathfrak{g}\text{-mod}.$$

Proposition 5.5. *The functor $\text{Kar}(\Phi)$ of (5.7) is full and essentially surjective.*

Proof. Fullness follows from Proposition 5.2, so it remains to show that $\text{Kar}(\Phi)$ is essentially surjective. If $\lambda = \sum_{i=1}^4 \lambda_i \omega_i$, with $\lambda_i \in \mathbb{Z}_{\geq 0}$, then V_λ is the submodule of $\bigotimes_{i=1}^4 V_{\omega_i}^{\otimes \lambda_i}$ generated by the one-dimensional λ weight space. Since the category $\mathfrak{g}\text{-mod}$ is semisimple, this implies that V_λ is a direct summand of $\bigotimes_{i=1}^4 V_{\omega_i}^{\otimes \lambda_i}$. Therefore, it suffices to show that the image of $\text{Kar}(\Phi)$ contains the fundamental representations V_{ω_i} for $i \in \{1, 2, 3, 4\}$. We see from (4.12) that V_{ω_1} and V_{ω_3} are contained in $V_{\omega_4}^{\otimes 2}$. It also follows from [MPR90, Ch. 11, Table 7] that V_{ω_2} is contained in $V_{\omega_3} \otimes V_{\omega_4}$. \square

Corollary 5.6. *We have a surjective algebra homomorphism*

$$\mathcal{F}_{7/3,26}(\mathbb{1}^{\otimes k}, \mathbb{1}^{\otimes k}) \twoheadrightarrow \text{End}_{\mathfrak{g}}(V^{\otimes k}), \quad k \in \mathbb{N}.$$

Corollary 5.6 implies that the endomorphism algebras of \mathcal{F} play the role in type F_4 that the group algebra of the symmetric group (or the oriented Brauer algebras if one includes the dual of the natural module) plays in type A and that the Brauer algebras play in types BCD .

We conclude this section with some conjectures.

Conjecture 5.7. *The functor $\text{Kar}(\Phi)$ is faithful, and hence is an equivalence of categories.*

A string diagram built from the generating morphisms (2.1) via tensor product and composition can be viewed as a graph. Here we view \frown as a trivalent vertex and \times as two edges (that is, we do *not* view the crossing as a vertex). We say such a graph is *component-planar* if its connected components are planar graphs. For example, (3.5) is a complete list of the 5 acyclic component-planar graphs $\mathbb{1}^{\otimes 2} \rightarrow \mathbb{1}^{\otimes 2}$, and there are precisely 15 acyclic component-planar graphs $\mathbb{1}^{\otimes 2} \rightarrow \mathbb{1}^{\otimes 3}$:

$$(5.8) \quad \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}, \begin{array}{c} \text{Y} \\ \text{Y} \\ \text{Y} \end{array}.$$

These are the morphisms appearing on the right-hand side of (2.12).

Conjecture 5.8. *For $m, n \in \mathbb{N}$, a basis for $\mathcal{F}(\mathbb{1}^{\otimes m}, \mathbb{1}^{\otimes n})$ is given by the component-planar graphs whose cycles are all of length at least six.*

Recall from (4.14) that

$$\dim \text{Hom}_{\mathfrak{g}}(V^{\otimes 2}, V^{\otimes 2}) = 5 \quad \text{and} \quad \dim \text{Hom}_{\mathfrak{g}}(V^{\otimes 3}, V^{\otimes 2}) = 15.$$

One can also count that there are 70 component-planar graphs $\mathbb{1}^{\otimes 3} \rightarrow \mathbb{1}^{\otimes 3}$ whose cycles are all of length at least six, and that $\dim \text{Hom}_{\mathfrak{g}}(V^{\otimes 3}, V^{\otimes 3}) = 70$ (using (4.13) and Schur's lemma). Thus, it would follow from Proposition 5.2, (2.9), and Conjecture 5.8 that the functor Φ induces isomorphisms

$$\mathcal{F}(\mathbb{1}^{\otimes m}, \mathbb{1}^{\otimes n}) \xrightarrow{\cong} \text{Hom}_{\mathfrak{g}}(V^{\otimes m}, V^{\otimes n}) \quad \text{for } m + n \leq 6.$$

This is related to Conjecture 5.7.

6. FUNDAMENTAL MODULES

Our goal in this final section is to describe the objects in $\text{Kar}(\mathcal{F})$ sent, under the functor $\text{Kar}(\Phi)$ of (5.7), to the four fundamental \mathfrak{g} -modules. Some of our intermediate results will be valid for more general α and δ . However, throughout this section we suppose that

$$\alpha \neq 0 \quad \text{and} \quad \delta \notin \{-10, -2, 0\}.$$

We continue to work over the field $\mathbb{k} = \mathbb{C}$.

Recall the definition of the antisymmetrizers in (3.11). Let

$$e_0 = \frac{1}{\delta} \cup, \quad e_1 = \frac{8}{\delta + 10} \left(\begin{array}{|c|} \hline \blacksquare \\ \hline \end{array} + \frac{\delta + 2}{4\alpha} \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \right),$$

$$e_3 = \frac{\delta + 2}{\delta + 10} \left(\begin{array}{|c|} \hline \blacksquare \\ \hline \end{array} - \frac{2}{\alpha} \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \right), \quad e_4 = \frac{1}{\alpha} \times, \quad \tilde{e} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} - \frac{1}{\delta} \cup - \frac{1}{\alpha} \times.$$

Lemma 6.1. *We have $ef, fe \in \mathbb{k}e$ for all*

$$f \in \left\{ \begin{array}{|c|} \hline | \\ \hline \end{array}, \cup, \times, \times, \times \right\}, \quad e \in \{e_0, e_1, e_3, e_4, \tilde{e}\}.$$

Proof. Since the given choices of e and f are invariant under rotation by 180° , it suffices to show that $ef \in \mathbb{k}e$. This is a straightforward computation. For example,

$$\begin{array}{|c|} \hline | \\ \hline \end{array} \circ e_1 = e_1, \quad \cup \circ e_1 \stackrel{(3.12)}{=} \stackrel{(3.13)}{=} 0, \quad \times \circ e_1 \stackrel{(3.12)}{=} \stackrel{(3.13)}{=} -e_1, \quad \times \circ e_1 \stackrel{(3.12)}{=} \stackrel{(3.13)}{=} 0, \quad \times \circ e_1 \stackrel{(2.11)}{=} \stackrel{(3.12)}{=} \frac{\alpha}{2} e_1$$

and

$$\begin{array}{|c|} \hline | \\ \hline \end{array} \circ \tilde{e} = \tilde{e}, \quad \cup \circ \tilde{e} \stackrel{(3.12)}{=} \stackrel{(2.4)}{=} 0, \quad \times \circ \tilde{e} \stackrel{(3.12)}{=} \stackrel{(2.4)}{=} e, \quad \times \circ \tilde{e} \stackrel{(3.12)}{=} \stackrel{(2.4)}{=} 0,$$

$$\times \circ \tilde{e} \stackrel{(2.4)}{=} \stackrel{(3.9)}{=} \frac{1}{2} (\times + \times) - \frac{\alpha}{\delta} \cup + \frac{\delta - 2}{2(\delta + 2)} \times \stackrel{(2.10)}{=} \frac{2\alpha}{\delta + 2} \tilde{e}.$$

The computations for $e \in \{e_0, e_3, e_4\}$ are similar. □

Lemma 6.2. *We have a decomposition*

$$(6.1) \quad 1_{\mathbb{1} \otimes \mathbb{1}} = e_0 + e_1 + e_3 + e_4 + \tilde{e}$$

of $1_{\mathbb{1} \otimes \mathbb{1}}$ as a sum of pairwise orthogonal idempotents.

Proof. A straightforward computation verifies (6.1). Next we show that e_0, e_1, e_3, e_4 are idempotent. We have $e_0^2 = e_0$ and $e_4^2 = e_4$ by the fourth and third relations in (2.4), respectively. Next,

$$e_1^2 = \frac{64}{(\delta + 10)^2} \left(\begin{array}{|c|} \hline \blacksquare \\ \hline \end{array} + \frac{\delta + 2}{4\alpha} \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} + \frac{(\delta + 2)^2}{16\alpha^2} \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \right)$$

$$\stackrel{(2.11)}{=} \stackrel{(3.12)}{=} \frac{64}{(\delta + 10)^2} \left(\frac{\delta + 10}{8} \begin{array}{|c|} \hline | \\ \hline \end{array} + \frac{(\delta + 10)(\delta + 2)}{32\alpha} \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \right) = e_1.$$

Since

$$e_3 = \begin{array}{|c|} \hline \blacksquare \\ \hline \end{array} - e_1,$$

it then follows that $e_3^2 = e_3$.

Now, by Lemma 6.1, we have $ef \in \mathbb{k}e \cap \mathbb{k}f$ for $e, f \in \{e_0, e_1, e_3, e_4, \tilde{e}\}$. Since no two of the idempotents e_0, e_1, e_3, e_4, e are scalar multiples of each other, it follows that they are orthogonal. □

Remark 6.3. If we knew that the morphisms (3.5) spanned $\mathcal{F}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2})$, then it would follow from Lemma 6.1 that

$$\dim_{\mathbb{k}} (e\mathcal{F}(\mathbb{1}^{\otimes 2}, \mathbb{1}^{\otimes 2})e) = 1$$

and hence that e_0, e_1, e_3, e_4 are primitive. For example, this would be the case if Conjecture 5.8 holds.

Recall from Section 5 our conventions for labeling of weights of \mathfrak{g} and that $V = V_{\omega_4}$.

Theorem 6.4. *Suppose $\alpha = \frac{7}{3}$ and $\delta = 26$. Then*

$$\text{Kar}(\Phi)(\mathbb{I}^{\otimes 2}, e_1) = V_{\omega_1}, \quad \text{Kar}(\Phi)(\mathbb{I}^{\otimes 2}, e_3) = V_{\omega_3}, \quad \text{Kar}(\Phi)(\mathbb{1}, \mathbb{1}_1) = V_{\omega_4}, \quad \text{Kar}(\Phi)(\mathbb{I}^{\otimes 2}, \tilde{e}) = V_{2\omega_4}.$$

Proof. In \mathfrak{g} -mod, we have the tensor product decomposition (4.12) and dimensions

$$\dim V_0 = 1, \quad \dim V_{\omega_1} = 52, \quad \dim V_{\omega_3} = 273, \quad \dim V_{\omega_4} = 26, \quad \dim V_{2\omega_4} = 324.$$

See, for example, [MPR90, Ch. 6, Table 2]. By Lemma 6.2, we also have the decomposition

$$V^{\otimes 2} \cong \text{Kar}(\Phi)(\mathbb{I}^{\otimes 2}, e_0) \oplus \text{Kar}(\Phi)(\mathbb{I}^{\otimes 2}, e_1) \oplus \text{Kar}(\Phi)(\mathbb{I}^{\otimes 2}, e_3) \oplus \text{Kar}(\Phi)(\mathbb{I}^{\otimes 2}, e_4) \oplus \text{Kar}(\Phi)(\mathbb{I}^{\otimes 2}, \tilde{e}).$$

Thus, if we can show that the images of the given elements of $\text{Kar}(\mathcal{F})$ have the expected dimensions, the theorem follows. Since the dimension of a \mathfrak{g} -module is the trace of its identity endomorphism, the dimension of $\text{Kar}(\Phi)(\mathbb{I}^{\otimes n}, e)$ is the image under Φ of

$$\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \boxed{e} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \in \mathcal{F}(\mathbb{1}, \mathbb{1}),$$

where there are n strands. (Here, and in what follows, we identify $\lambda \mathbb{1}_{\mathbb{1}}$ with $\lambda \in \mathbb{k}$.)

Note that

$$\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \frac{1}{2} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} - \frac{1}{2} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \stackrel{(2.4)}{=} \frac{\delta(\delta-1)}{2},$$

$$\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \frac{1}{2} \begin{array}{c} \text{---} \\ \text{---} \end{array} - \frac{1}{2} \begin{array}{c} \text{---} \\ \text{---} \end{array} \stackrel{(2.4)}{=} 0 - \frac{1}{2} \begin{array}{c} \text{---} \\ \text{---} \end{array} \stackrel{(2.4)}{=} -\frac{\alpha\delta}{2}.$$

Therefore, we have

$$\dim(\mathbb{I}^{\otimes 2}, e_1) = \frac{8}{\delta+10} \left(\frac{\delta(\delta-1)}{2} - \frac{\delta(\delta+2)}{8} \right) = \frac{3\delta(\delta-2)}{\delta+10} = 52,$$

$$\dim(\mathbb{I}^{\otimes 2}, e_3) = \frac{\delta+2}{\delta+10} \left(\frac{\delta(\delta-1)}{2} + \delta \right) = \frac{\delta(\delta+1)(\delta+2)}{2(\delta+10)} = 273.$$

It is also clear that $\dim(\mathbb{1}, \mathbb{1}_1) = \delta = 26$ and that $\dim(\mathbb{I}^{\otimes 2}, e_0) = 1$. It then follows that $\dim(\mathbb{I}^{\otimes 2}, \tilde{e}) = 26^2 - 1 - 52 - 273 - 26 = 324$. \square

Note that Theorem 6.4 does not describe an object in $\text{Kar}(\mathcal{F})$ that is mapped to the second fundamental representation V_{ω_2} . In \mathfrak{g} -mod, the lowest tensor power of V in which V_{ω_2} appears is $V^{\otimes 3}$. Thus, we expect that there exists an idempotent $e_2 \in \mathcal{F}(\mathbb{I}^{\otimes 3}, \mathbb{I}^{\otimes 3})$ such that $\text{Kar}(\Phi)(\mathbb{I}^{\otimes 3}, e_2) = V_{\omega_2}$. (Such an idempotent is guaranteed to exist if Conjecture 5.7 holds.) In fact, we can say a bit more. A computation similar to the ones in the proof of Theorem 6.4 shows that

$$(6.2) \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \frac{\delta(\delta-1)(\delta-2)}{6} = 2600 = 1274 + 273 + 1053 \quad \text{when } \delta = 26.$$

The tensor product decomposition of $V^{\otimes 3}$ is given in (4.13). Since

$$\begin{aligned} \dim V_0 &= 1, \quad \dim V_{\omega_1} = 52, \quad \dim V_{\omega_2} = 1274, \quad \dim V_{\omega_3} = 273, \quad \dim V_{\omega_4} = 26, \\ \dim V_{\omega_1+\omega_4} &= 1053, \quad \dim V_{\omega_3+\omega_4} = 4096, \quad \dim V_{2\omega_4} = 324, \quad \text{and} \quad \dim V_{3\omega_4} = 2652, \end{aligned}$$

(see, for example, [MPR90, Ch. 6, Table 2]) the only submodule of $V^{\otimes 3}$ of dimension 2600 is $V_{\omega_2} \oplus V_{\omega_3} \oplus V_{\omega_1+\omega_4}$. Thus, we expect that the antisymmetrizer $\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$ decomposes as a sum of three

orthogonal minimal idempotents, one of which is e_2 . (This is guaranteed to happen if Conjecture 5.7 holds.) Unfortunately, the computations required to find this idempotent explicitly are unwieldy.

Lemma 6.5. *We have*

$$\begin{array}{c} \diagup \\ \boxed{e_1} \quad \boxed{e_1} \\ \diagdown \end{array} = \frac{3\alpha(2-\delta)(\delta-26)}{4(\delta+10)^2} 1_1.$$

Proof. We compute

$$\begin{aligned} \begin{array}{c} \text{---} \\ \diagup \\ \boxed{e_1} \quad \boxed{e_1} \\ \diagdown \end{array} &= \frac{1}{4} \left(\begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} + \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} \right) \stackrel{(2.4)}{=} \frac{\alpha(\delta-2)}{4} 1_1, \\ \begin{array}{c} \text{---} \\ \diagup \\ \boxed{e_1} \quad \text{---} \\ \diagdown \end{array} &= \frac{1}{4} \left(\begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} + \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} \right) \stackrel{(2.4)}{=} \frac{1}{4} \left(0 - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} + \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} \right) \stackrel{(2.4)}{=} \frac{\alpha^2(\delta-2)}{8(\delta+2)} 1_1, \\ \begin{array}{c} \text{---} \\ \diagup \\ \boxed{e_1} \quad \text{---} \\ \diagdown \end{array} &= \frac{1}{4} \left(\begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} + \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} \right) \stackrel{(2.4)}{=} \frac{1}{4} \left(\alpha \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} - \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} + \begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} \right) \\ &\stackrel{(2.4)}{=} \frac{\alpha^3(2-\delta)(3\delta+2)}{16(\delta+2)^2} 1_1. \end{aligned}$$

Thus

$$\begin{aligned} \begin{array}{c} \diagup \\ \boxed{e_1} \quad \boxed{e_1} \\ \diagdown \end{array} &= \frac{64}{(\delta+10)^2} \left(\begin{array}{c} \diagup \\ \boxed{e_1} \\ \diagdown \end{array} + \frac{\delta+2}{4\alpha} \begin{array}{c} \text{---} \\ \diagup \\ \boxed{e_1} \quad \text{---} \\ \diagdown \end{array} + \frac{\delta+2}{4\alpha} \begin{array}{c} \text{---} \\ \diagup \\ \boxed{e_1} \quad \text{---} \\ \diagdown \end{array} + \frac{(\delta+2)^2}{16\alpha^2} \begin{array}{c} \text{---} \\ \diagup \\ \boxed{e_1} \quad \text{---} \\ \diagdown \end{array} \right) \\ &= \frac{3\alpha(2-\delta)(\delta-26)}{4(\delta+10)^2} 1_1. \end{aligned} \quad \square$$

Corollary 6.6. *If $\alpha \neq 0$ and $\delta \notin \{-2, -10, 2, 26\}$, then*

$$\frac{4(\delta+10)^2}{3\alpha(2-\delta)(\delta-26)} \begin{array}{c} \boxed{e_1} \quad \boxed{e_1} \\ \diagup \quad \diagdown \\ \boxed{e_1} \quad \boxed{e_1} \end{array}$$

is a nonzero idempotent endomorphism. In particular, 1 is a direct summand of $(1^{\otimes 2}, e_1)^{\otimes 2}$ in $\text{Kar}(\mathcal{F})$.

Remark 6.7. Corollary 6.6 and Theorem 6.4 explain the importance of the choice $\delta = 26$. The \mathfrak{g} -module V_{ω_1} is the adjoint representation, and $V_{\omega_1}^{\otimes 2}$ does not contain a copy of the natural representation $V = V_{\omega_4}$. In fact, a lengthy computation shows that the morphism

$$\begin{array}{c} \diagup \\ \boxed{e_1} \quad \boxed{e_1} \\ \diagdown \end{array}$$

can be written as a linear combination of the morphisms (5.8) with the two leftmost top strands brought to the bottom using caps. The coefficients in this linear combination all vanish *if and only if* $\delta = 26$.

Remark 6.8. Suppose we have an idempotent $e_2 \in \mathcal{F}(\mathbb{I}^{\otimes 3}, \mathbb{I}^{\otimes 3})$ such that $\text{Kar}(\Phi)(\mathbb{I}^{\otimes 3}, e_2) = V_{\omega_2}$. Let $\mathcal{W} = \mathcal{W}(\alpha, \delta)$ be the full monoidal subcategory of $\text{Kar}(\mathcal{F}_{\alpha, \delta})$ generated by

$$(6.3) \quad (\mathbb{I}^{\otimes 2}, e_1), \quad (\mathbb{I}^{\otimes 3}, e_2), \quad (\mathbb{I}^{\otimes 2}, e_3), \quad \mathbb{1}.$$

Then morphisms in \mathcal{W} can be depicted as string diagrams with strands labeled by elements of $\{1, 2, 3, 4\}$, with a strand labeled i corresponding to the identity morphism of the i -th object in the list (6.3). The category \mathcal{W} should be a degenerate (that is, $q = 1$) web category of type F_4 .

REFERENCES

- [Ada96] J. F. Adams. *Lectures on exceptional Lie groups*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1996. With a foreword by J. Peter May, Edited by Zafer Mahmud and Mamoru Mimura.
- [BCNR17] J. Brundan, J. Comes, D. Nash, and A. Reynolds. A basis theorem for the affine oriented Brauer category and its cyclotomic quotients. *Quantum Topol.*, 8(1):75–112, 2017. [arXiv:1404.6574](#), [doi:10.4171/QT/87](#).
- [BERT21] E. Bodish, B. Elias, D.E.V. Rose, and L. Tatham. Type C webs. 2021. [arXiv:2103.14997](#).
- [CKM14] S. Cautis, J. Kamnitzer, and S. Morrison. Webs and quantum skew Howe duality. *Math. Ann.*, 360(1-2):351–390, 2014. [arXiv:1210.6437](#), [doi:10.1007/s00208-013-0984-4](#).
- [Cvi08] P. Cvitanović. *Group theory*. Princeton University Press, Princeton, NJ, 2008. Birdtracks, Lie’s, and exceptional groups. [doi:10.1515/9781400837670](#).
- [Kir76] A. A. Kirillov. *Elements of the theory of representations*. Grundlehren der Mathematischen Wissenschaften, Band 220. Springer-Verlag, Berlin-New York, 1976. Translated from the Russian by Edwin Hewitt.
- [Kup94] G. Kuperberg. The quantum G_2 link invariant. *Internat. J. Math.*, 5(1):61–85, 1994. [arXiv:math/9201302](#), [doi:10.1142/S0129167X94000048](#).
- [Kup96] G. Kuperberg. Spiders for rank 2 Lie algebras. *Comm. Math. Phys.*, 180(1):109–151, 1996. URL: <http://projecteuclid.org/euclid.cmp/1104287237>, [arXiv:q-alg/9712003](#).
- [LZ15] G. I. Lehrer and R. B. Zhang. The Brauer category and invariant theory. *J. Eur. Math. Soc. (JEMS)*, 17(9):2311–2351, 2015. [arXiv:1207.5889](#), [doi:10.4171/JEMS/558](#).
- [MPR90] W. G. McKay, J. Patera, and D. W. Rand. *Tables of representations of simple Lie algebras. Vol. I*. Université de Montréal, Centre de Recherches Mathématiques, Montreal, QC, 1990. Exceptional simple Lie algebras, With a foreword by R. V. Moody.
- [MPS17] S. Morrison, E. Peters, and N. Snyder. Categories generated by a trivalent vertex. *Selecta Math. (N.S.)*, 23(2):817–868, 2017. [arXiv:1501.06869](#), [doi:10.1007/s00029-016-0240-3](#).
- [MT07] N. J. MacKay and A. Taylor. Rational R -matrices, centralizer algebras and tensor identities for e_6 and e_7 exceptional families of Lie algebras. *J. Math. Phys.*, 48(10):103507, 14, 2007. [arXiv:math/0608248](#), [doi:10.1063/1.2779960](#).
- [Thu04] D. P. Thurston. The F_4 and E_6 families have a finite number of points. 2004. URL: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.118.497&rep=rep1&type=pdf>.
- [Wen03] H. Wenzl. On tensor categories of Lie type E_N , $N \neq 9$. *Adv. Math.*, 177(1):66–104, 2003. [doi:10.1016/S0001-8708\(02\)00048-8](#).
- [Wes03] B. Westbury. Invariant tensors and diagrams. In *Proceedings of the Tenth Oporto Meeting on Geometry, Topology and Physics (2001)*, volume 18, pages 49–82, 2003. [doi:10.1142/S0217751X03017968](#).
- [Wes08] B. Westbury. Invariant tensors for the spin representation of $\mathfrak{so}(7)$. *Math. Proc. Cambridge Philos. Soc.*, 144(1):217–240, 2008. [arXiv:math/0601209](#), [doi:10.1017/S0305004107000722](#).

(R.G.) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, OTTAWA, ON K1N 6N5,
CANADA

Email address: rgand037@uottawa.ca

(A.S.) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, OTTAWA, ON K1N 6N5,
CANADA

URL: alistairsavage.ca, *ORCID:* orcid.org/0000-0002-2859-0239

Email address: alistair.savage@uottawa.ca

(K.Z.) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, OTTAWA, ON K1N 6N5,
CANADA

URL: mysite.science.uottawa.ca/kzaynull/

Email address: kzaynull@uottawa.ca