

# THE CATEGORY OF SILVA SPACES IS NOT INTEGRAL

MARIANNE LAWSON<sup>1</sup> AND SVEN-AKE WEGNER<sup>2</sup>

**ABSTRACT.** We establish that the category of Silva spaces, aka LS-spaces, formed by countable inductive limits of Banach spaces with compact linking maps as objects and linear and continuous maps as morphisms, is not an integral category. The result carries over to the category of PLS-spaces, i.e., countable projective limits of LS-spaces—which contains prominent spaces of analysis such as the space of distributions and the space of real analytic functions. As a consequence, we obtain that both categories neither have enough projective nor enough injective objects. All results hold true when ‘compact’ is replaced by ‘weakly compact’ or ‘nuclear’. This leads to the categories of PLS-, PLS<sub>w</sub>- and PLN-spaces, which are examples of ‘inflation exact categories with admissible cokernels’ as recently introduced by Henrard, van Roosmalen, Kvamme and the second-named author.

## 1. INTRODUCTION

Integral categories were introduced by Rump [14] in 2001 and have since caught attention on the one hand in representation theory, see, e.g., Brüstle, Hassoun, Tattar [2], or Nakaoka [13], and on the other hand in functional analysis, see Hassoun, Shah, Wegner [9]. By definition, a pre-abelian category  $\mathcal{A}$  is *right integral*, if monomorphisms pushout to monomorphism, meaning whenever  $f: X \rightarrow Y$  is monic and  $g: X \rightarrow Z$  is arbitrary, then in the diagram below

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ g \downarrow & \text{PO} & \downarrow j \\ Z & \xrightarrow{h} & P \end{array}$$

the morphism  $h$  is again monic. *Left integral* categories are defined dually.

The aim of this paper is to establish that the categories LS, PLS, LS<sub>w</sub>, PLS<sub>w</sub>, LN and PLN, whose definitions will be given in due course, are all neither right nor left integral, and to outline consequences of this. The latter categories have proved to be very important in functional analysis as they arise naturally when Ext<sup>1</sup>-techniques are used to solve linear partial differential equations, see, in particular, Wengenroth [17] and Dierolf, Sieg [6]. We will also put the above categories in the context of *inflation exact categories* that were introduced recently by Henrard, van Roosmalen, Kvamme, Wegner [10].

To keep this article short, we refer to [9] for a detailed discussion of how integral categories fit into the hierarchy of semi- and quasi-abelian categories, and for a long list of examples and non-examples from functional analysis.

## 2. CORE RESULT

A Hausdorff locally convex space  $X$  is said to be an *LS-space* (named after e Silva [8]), if there exists a spectrum of Banach spaces  $X_1 \rightarrow X_2 \rightarrow X_3 \rightarrow \dots$  linked by linear, injective and compact maps, such that  $X = \text{ind}_{n \in \mathbb{N}} X_n$  is the locally convex inductive limit of the spectrum. We denote by LS the category which has the LS-spaces as objects and linear and continuous maps as morphisms. In LS the kernel of a morphism  $f: X \rightarrow Y$  is the space  $f^{-1}(0) \subseteq X$  endowed with the subspace topology and the cokernel is the quotient  $Y/\overline{f(X)}$  endowed with the locally convex quotient topology. It is classic [17, p. 96] that LS is quasi-abelian, that a morphism is a kernel if and only if it is injective and has a closed range, and that it is a cokernel if and only if it is surjective. Our main result concerning LS is the following.

**Theorem 1.** The category of LS-spaces is neither left nor right integral.

*Proof.* Since the category of LS-spaces is quasi-abelian, it is right integral if and only if it is left integral by [14, Cor. on p. 173] (see also [9, Prop. 2.6]). We provide a counterexample to show that LS is not

2020 *Mathematics Subject Classification*: Primary 18E05, 46A13; Secondary 46M10, 46A45, 18G80.

*Key words and phrases*: LS-space, PLS-space, integral category, injective and projective objects.

<sup>1</sup> Teesside University, School of Computing, Engineering & Digital Technologies, TS1 3BX Middlesbrough, UK.

<sup>2</sup> Corresponding author: Universität Hamburg, Fachbereich Mathematik, Bundesstraße 55, 20146 Hamburg, Germany, phone: +49 (0) 40 42838 76 57, e-mail: sven.wegner@uni-hamburg.de.

right integral. For a decreasing sequence  $V = (v_n)_{n \in \mathbb{N}}$  of strictly positive functions  $v_n: \mathbb{N} \rightarrow \mathbb{R}$  we denote by

$$k^1(V) := \operatorname{ind}_{n \in \mathbb{N}} \ell^1(v_n)$$

the Köthe co-echelon space of order one. The steps in the inductive limit are the Banach spaces

$$\ell^1(v_n) := \{x \in \mathbb{K}^{\mathbb{N}} \mid \|x\|_{v_n} := \sum_{j=1}^{\infty} v_n(j)|x(j)| < \infty\}$$

and the linking maps are the inclusions. Due to a classic result by Köthe, see Bierstedt [1, Proposition 14], the space  $k^1(V)$  is an LS-space if and only if the sequence  $V$  satisfies condition

$$(S) \quad \forall n \exists m > n: \lim_{j \rightarrow \infty} \frac{v_m(j)}{v_n(j)} = 0.$$

We now define two concrete sequences  $V = (v_n)_{n \in \mathbb{N}}$  and  $W = (w_n)_{n \in \mathbb{N}}$  by putting  $v_n(j) := j^{-n}$  and  $w_n := e^{-jn}$ . Then both sequences enjoy (S) and since  $e^{-jn} \leq j^{-n}$  holds for all  $n, j \in \mathbb{N}$  we get

$$\forall x \in \ell^1(v_n): \|x\|_{w_n} = \sum_{j=1}^{\infty} e^{-jn} |x(j)| \leq \sum_{j=1}^{\infty} j^{-n} |x(j)| = \|x\|_{v_n} < \infty.$$

This means that  $\ell^1(v_n) \subseteq \ell^1(w_n) \subseteq k^1(W)$  holds for every  $n \in \mathbb{N}$  with continuous inclusion. But then  $k^1(V) \subseteq k^1(W)$  must also hold with continuous inclusion by the universal property of the inductive limit, see, e.g., Meise, Vogt [12, Prop. 24.7]. Let us denote the inclusion map by  $i: k^1(V) \rightarrow k^1(W)$ .

Next, we define the map

$$\theta: k^1(V) \rightarrow \mathbb{K}, \quad \theta(x) := \sum_{j=1}^{\infty} e^{-j} \cdot x(j)$$

which is well-defined and continuous: Indeed, for  $n \in \mathbb{N}$  we choose  $C > 0$  such that  $e^{-j} \leq Cj^{-n}$  holds for all  $j \in \mathbb{N}$ . Then we get

$$\forall x \in \ell^1(v_n): |\theta(x)| \leq \sum_{j=1}^{\infty} e^{-j} |x(j)| \leq C \sum_{j=1}^{\infty} j^{-n} |x(j)| = C \|x\|_{v_n} < \infty$$

which means that  $\theta: \ell^1(v_n) \rightarrow \mathbb{K}$  is well-defined and continuous. Now we again use [12, Prop. 24.7].

We continue by forming the pushout of  $i$  along  $\theta$  which leads, in view of what we noted about cokernels in LS right before Theorem 1, to the diagram

$$\begin{array}{ccc} k^1(V) & \xrightarrow{i} & k^1(W) \\ \theta \downarrow & \text{PO} & \downarrow \psi \\ \mathbb{K} & \xrightarrow{\varphi} & \frac{k^1(W) \oplus \mathbb{K}}{\operatorname{ran}[i - \theta]^T} \end{array} \quad (1)$$

where  $\varphi$  and  $\psi$  are the natural maps. We claim now that  $\varphi$  is not injective and thus not a monomorphism in LS. This can be seen by considering the sequence  $(x^{(k)})_{k \in \mathbb{N}} \subseteq k^1(V)$  defined by  $x^{(k)}(j) = (e^1/k, e^2/k, \dots, e^k/k, 0, \dots)$ . Applying  $\theta$  to each member of the sequence leads to

$$\theta(x^{(k)}) = \sum_{j=1}^{\infty} e^{-j} \cdot x^{(k)}(j) = \sum_{j=1}^k e^{-j} \cdot e^j/k = \sum_{j=1}^k 1/k = 1.$$

Applying  $i$  means considering  $(x^{(k)})_{k \in \mathbb{N}} \subseteq k^1(W)$ , and since the sequence has only finitely many non-zero entries, it is contained in every step space. We consider it in  $\ell^1(w_2)$ , where we obtain

$$\|x^{(k)}\|_{w_2} = \sum_{j=1}^{\infty} e^{-2j} |x^{(k)}(j)| = \sum_{j=1}^k e^{-2j} \cdot e^j/k \leq 1/k \cdot \sum_{j=1}^{\infty} e^{-j} \xrightarrow{k \rightarrow \infty} 0$$

from whence it follows that  $(x^{(k)})_{k \in \mathbb{N}}$  converges to zero in  $k^1(W)$ . Combining the above implies

$$\begin{bmatrix} 0 \\ -1 \end{bmatrix} \in \overline{\left\{ \begin{bmatrix} i \\ -\theta \end{bmatrix}(x) \in k^1(W) \oplus \mathbb{K} \mid x \in k^1(V) \right\}}$$

which means that  $\varphi(-1) = \begin{bmatrix} 0 \\ -1 \end{bmatrix} + \overline{\text{ran}[i \ -\theta]^T} = 0$  and establishes the claim.  $\square$

**Remark 2.** The method used in the proof above is a refinement of that one applied in [9, Thm. 3.2] and was developed by the first-named author in her thesis [11].

### 3. CONSEQUENCES AND FURTHER RESULTS

A Hausdorff locally convex space  $X$  is a *PLS-space* if  $X = \text{proj}_{n \in \mathbb{N}} X_n$  is the projective limit of a strongly reduced spectrum  $\cdots \rightarrow X_3 \rightarrow X_2 \rightarrow X_1$  of LS-spaces. We denote by **PLS** the category with PLS-spaces as objects and linear and continuous maps as morphisms. For a morphism  $f: X \rightarrow Y$  the kernel is computed by the same formula as in **LS** but the cokernel is the completion of  $Y/f(X)$ , see Sieg [15, Proof of Prop. 3.1.3]. Since **LS**  $\subseteq$  **PLS** reflects cokernels we can nevertheless recycle our counterexample from Theorem 1 as well as the dual counterexample (which must exist as **LS** is quasi-abelian) and obtain:

**Corollary 3.** The category of PLS-spaces is neither left nor right integral.  $\square$

Applying Buan, Marsh [3, Prop. 3.9] (see also [9, Prop. 5.2]) leads to:

**Corollary 4.** The categories of LS-spaces and PLS-spaces neither have enough projective objects nor do they have enough injective objects.  $\square$

In contrast to **LS**, the category **PLS** is not even left semi-abelian [15, Prop. 3.1.6]. It thus follows from [14, p. 169, Cor. 1] (see also [9, Prop. 2.5]) that **PLS** does not have enough quasi-projectives (replace ‘epimorphism’ with ‘cokernel’ everywhere in the definition). It is open if **PLS** has enough quasi-injectives, or, taking its natural exact structure  $\mathbb{E} := \mathbb{E}_{\max} = \mathbb{E}_{\text{top}}$ , see [15, Prop. 3.2.1], into account, enough  $\mathbb{E}$ -injectives in the notation of [2, 4] (‘inflation’ instead of ‘monomorphism’).

Replacing ‘compact’ with ‘weakly compact’ or ‘nuclear’ in the definition of LS-spaces leads to the categories of  $\text{LS}_w$ -spaces, and  $\text{LN}$ -spaces, respectively. Taking strongly reduced countable projective limits of spaces in the two classes gives rise to  $\text{PLS}_w$ -spaces and  $\text{PLN}$ -spaces. We have the following full inclusions

$$\begin{array}{ccccc} \text{PLN} & \subseteq & \text{PLS} & \subseteq & \text{PLS}_w \\ \cup & & \cup & & \cup \\ \text{LN} & \subseteq & \text{LS} & \subseteq & \text{LS}_w \end{array} \quad (2)$$

of pre-abelian categories. Indeed, in  $\text{LN}$  and  $\text{PLN}$  kernels and cokernels are computed by the same formulas as in **LS** and **PLS**, respectively (for kernels this follows from Domański, Vogt [7, Prop. 1.2], for cokernels in  $\text{LN}$  use [7, Prop. 1.3], and for cokernels in  $\text{PLN}$  proceed as in [15, Proof of Prop. 3.1.3]). Since the Köthe matrices  $V$  and  $W$ , which we employed in Theorem 1, satisfy condition

$$(N) \quad \forall n \exists m > n: \sum_{j=1}^{\infty} \frac{v_m(j)}{v_n(j)} < \infty,$$

it follows from a classic result by Grothendieck, see Bierstedt [1, Prop. 15], that  $k^1(V)$  and  $k^1(W)$  are indeed  $\text{LN}$ -spaces. The diagram (1) is thus a pushout in  $\text{LN}$  and in  $\text{PLN}$ . Since  $\text{LN} \subseteq \text{LS}$  reflects kernels and cokernels,  $\text{LN}$  is quasi-abelian and thus there exists a dual counterexample in  $\text{LN}$ . Since  $\text{LN} \subseteq \text{PLN}$  reflects kernels and cokernels the example applies in  $\text{PLN}$ , too. We thus get:

**Corollary 5.** The categories of  $\text{LN}$ -spaces and  $\text{PLN}$ -spaces are neither left nor right integral. Furthermore, they neither have enough projective objects nor enough injective objects.  $\square$

In the categories  $\text{LS}_w$  and  $\text{PLS}_w$  cokernels compute again as in **LS** and **PLS**, respectively. Kernels however do not: Given a morphism  $f: \text{ind}_{n \in \mathbb{N}} X_n \rightarrow \text{ind}_{n \in \mathbb{N}} Y_n$  of  $\text{LS}_w$ -spaces the kernel is

$$\text{ind}_{n \in \mathbb{N}} (f^{-1}(0) \cap X_n) \rightarrow \text{ind}_{n \in \mathbb{N}} X_n \quad (3)$$

where the topology on the left is in general strictly finer than the subspace topology [6, Rmk. 3.1.1 and Cor. 3.1.3(i)]. For morphisms between  $\text{PLS}_w$ -spaces the same has to be done for every step in the projective limit [6, Prop. 3.1.5]. Nevertheless, in both categories it is true that a morphism  $f$  is monic if and only if it is injective, and  $\text{LS} \subseteq \text{LS}_w$  and  $\text{PLS} \subseteq \text{PLS}_w$  reflect kernels (as, for  $\text{LS}$ -spaces, the topology defined in (3) coincides with the subspace topology). Moreover,  $\text{LS}_w$  is quasi-abelian [6, p. 1307 and Cor. 3.1.3(i)]. By recycling our example from Theorem 1 a third time we arrive at:

**Corollary 6.** The categories of  $\text{LS}_w$ -spaces and  $\text{PLS}_w$ -spaces are neither left nor right integral. Neither of them have enough projective objects nor enough injective objects.  $\square$

We mentioned already that the categories in the bottom row of (2) are quasi-abelian. For the top row this is not the case and indeed a particular aim of [6, 15] (see also [5, 16]) was to establish natural exact structures on  $\text{PLS}$ ,  $\text{PLS}_w$  and  $\text{PLN}$ . We conclude our article by relating the latter approach to the notion of *inflation-exact categories with admissible cokernels* as defined recently in [10]. This new notion allows for the use of homological algebra with respect to the class  $\mathbb{C}_{\text{all}}$  of all kernel-cokernel pairs in cases where the latter is not an exact structure in the classical Quillen sense.

**Theorem 7.** Endowed with the class  $\mathbb{C}_{\text{all}}$  of all kernel-cokernel pairs as conflations, the categories  $\text{PLS}$ ,  $\text{PLS}_w$  and  $\text{PLN}$  are inflation-exact categories with admissible cokernels. The class  $\mathbb{C}_{\text{all}}$  is not an exact structure.

*Proof.* All three categories are right quasi-abelian but not left quasi-abelian in the notation of [9, Dfn. 2.3]. For  $\text{PLS}_w$  this follows from [6, Prop. 3.1.9]. For  $\text{PLS}$  and  $\text{PLN}$  the ‘right’ part can be proved in exactly the same way. The ‘not left’ part for  $\text{PLS}$  follows from the fact that  $\text{PLS}$  is not left semi-abelian [15, Prop. 3.1.6]. As the example uses a quotient of the space  $\mathcal{D}'(\Omega)$ , it applies to  $\text{PLN}$ , too. Using the dual of [10, Rmk. 4.10(1)] establishes the claim.  $\square$

We refer to [10] for various implications of Theorem 7, including the means for defining the derived category. The following illustrates the trade-off that comes with switching from a (natural) exact structure  $\mathbb{E}$  to the conflation structure  $\mathbb{C}_{\text{all}}$  on the categories covered by Theorem 7:

**Corollary 8.** Let  $\mathcal{A} \in \{\text{PLS}, \text{PLS}_w, \text{PLN}\}$ .

- (i) There is a natural triangle equivalence  $\mathbf{D}^b(\mathcal{A}, \mathbb{C}_{\text{all}}) \rightarrow \mathbf{D}^b(\mathcal{RH}(\mathcal{A}))$  with  $\mathcal{RH}(\mathcal{A})$  being the heart of a natural t-structure on the derived category.
- (ii) The natural triangle functor  $\mathbf{D}^b(\mathcal{A}, \mathbb{E}) \rightarrow \mathbf{D}^b(\mathcal{A}, \mathbb{C}_{\text{all}})$  is neither an equivalence for the maximal exact structure  $\mathbb{E} = \mathbb{E}_{\text{max}}$  nor for any other exact structure.

*Proof.* The statements can be proved by copying and dualising [10, Thm. 9.3] and [10, Thm. 9.6].  $\square$

## REFERENCES

1. K. D. Bierstedt, *An introduction to locally convex inductive limits*, Geometrical methods of the theory of Fréchet spaces, 1988, pp. 35–133.
2. T. Brüstle, S. Hassoun, and A. Tattar, *Intersections, sums, and the Jordan-Hölder property for exact categories*, J. Pure Appl. Algebra **225** (2021), no. 11, 106724.
3. A. B. Buan and R. J. Marsh, *From triangulated categories to module categories via localization II: calculus of fractions*, J. Lond. Math. Soc. (2) **86** (2012), no. 1, 152–170.
4. T. Bühler, *Exact categories*, Expo. Math. **28** (2010), no. 1, 1–69.
5. S. Crivei, *Maximal exact structures on additive categories revisited*, Math. Nachr. **285** (2012), no. 4, 440–446.
6. B. Dierolf and D. Sieg, *A homological approach to the splitting theory of  $\text{PLS}_w$  spaces*, J. Math. Anal. Appl. **433** (2016), no. 2, 1305–1328.
7. P. Domański and D. Vogt, *A splitting theory for the space of distributions*, Stud. Math. **140** (2000), no. 1, 57–77.
8. J. S. e Silva, *Su certi classi di spazi localmente convessi importanti per le applicazioni*, Rend. Mat. e delle sue Appl. **14** (1955), 388–414.
9. S. Hassoun, A. Shah, and S. Wegner, *Examples and non-examples of integral categories and the admissible intersection property*, Cahiers Topologie Géom. Différentielle Catég. **62** (2021), no. 3, 329–354.
10. R. Henrard, A-C. van Roosmalen, S. Kvamme, and S.-A. Wegner, *The left heart and exact hull of an additive regular category*, (2021), arXiv:2105.11483.
11. M. Lawson, *Integral categories in functional analysis*, BSc Thesis, Teesside University, Middlesbrough, UK, 2021.
12. R. Meise and D. Vogt, *Introduction to Functional Analysis*, Oxford Graduate Texts in Mathematics, vol. 2, The Clarendon Press, Oxford University Press, New York, 1997.
13. H. Nakaoka, *General heart construction for twin torsion pairs on triangulated categories*, J. Algebra **374** (2013), 195–215.
14. W. Rump, *Almost abelian categories*, Cahiers Topologie Géom. Différentielle Catég. **42** (2001), no. 3, 163–225.
15. D. Sieg, *A homological approach to the splitting theory of  $\text{PLS}$ -spaces*, PhD Thesis, Universität Trier, Germany, 2010.
16. D. Sieg and S.-A. Wegner, *Maximal exact structures on additive categories*, Math. Nachr. **284** (2011), no. 16, 2093–2100.
17. J. Wengenroth, *Derived Functors in Functional Analysis*, Springer-Verlag, Berlin, 2003.