

WASSERSTEIN CONDITIONAL INDEPENDENCE TESTING

ANDREW WARREN

ABSTRACT. We introduce a test for the conditional independence of random variables X and Y given a random variable Z , specifically by sampling from the joint distribution (X, Y, Z) , binning the support of the distribution of Z , and conducting multiple p -Wasserstein two-sample tests. Under a p -Wasserstein Lipschitz assumption on the conditional distributions $\mathcal{L}_{X|Z}$, $\mathcal{L}_{Y|Z}$, and $\mathcal{L}_{(X,Y)|Z}$, we show that it is possible to control the Type I and Type II error of this test, and give examples of explicit finite-sample error bounds in the case where the distribution of Z has compact support.

Consider random variables X, Y, Z . That is, $(\Omega, \mathcal{A}, \mathbb{P})$ is our sample space, and $X : \Omega \rightarrow \mathbb{R}^{d_X}$, $Y : \Omega \rightarrow \mathbb{R}^{d_Y}$, and $Z : \Omega \rightarrow \mathbb{R}^{d_Z}$ are all Borel functions, with distributions $\mathcal{L}_X := \mathbb{P} \circ X^{-1}$ (and similar for Y and Z) as well as joint distribution $\mathcal{L}_{(X,Y,Z)} := \mathbb{P} \circ (X, Y, Z)^{-1}$. We are interested in testing the conditional dependence relations between X , Y , and Z . For a general overview of some of the many ways in which conditional (in)dependence arises in statistics, we refer the reader to the classic article by Dawid [Daw79]. More recently, conditional independence testing has found a central role in the areas of causal inference and causal discovery; for a summary of this line of work, see the recent text by Peters et al. [PJS17], as well as the foundational works by Spirtes et al. [SGS+01] and Pearl [Pea09].

Numerous practical methods for testing conditional independence have been proposed — for instance, Heinze-Deml et al. [HDPM18] describe six classes of conditional independence test which have been implemented in the R package `CondIndTests`.

Nonetheless, general theoretical guarantees for conditional independence testing — consistency results for general joint distributions $\mathcal{L}_{(X,Y,Z)}$, sample complexity, power estimates, and so on — have been hard to come by. Recently, Shah and Peters [SP20] have demonstrated that this is not an accident: indeed, they show that for general joint distribution $\mathcal{L}_{(X,Y,Z)}$, any valid test of conditional independence does not have power against any alternative. (This result has also been reproved, using techniques from optimal transport, in [NBW20].) Consequently, it is of interest to find *mild* assumptions on $\mathcal{L}_{(X,Y,Z)}$ (or, similarly, on the various conditional distributions between the variables X , Y , and Z) under which a general test for conditional independence does, indeed, exist.

Here, our strategy is to present regularity conditions on the conditional distributions $\mathcal{L}_{X|Z=z}$, $\mathcal{L}_{Y|Z=z}$, and $\mathcal{L}_{(X,Y)|Z=z}$, so that if we bin the conditioning variable Z using a sufficiently fine, finite partition V_1, \dots, V_J of $\text{supp}(\mathcal{L}_Z)$, then for all $j = 1, \dots, J$ and all $z \in V_j$,

$$\mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z} \approx \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}, \text{ and } \mathcal{L}_{(X,Y)|Z=z} \approx \mathcal{L}_{(X,Y)|Z \in V_j}$$

where \approx denotes closeness with respect to some suitable metric on the space of probability measures. If this holds, then we can replace the (intractable in general) problem of directly comparing the measures $\mathcal{L}_{(X,Y)|Z=z}$ and $\mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}$ for every z in the range of Z , with the (costly, but in principle tractable) problem of comparing $\mathcal{L}_{(X,Y)|Z \in V_j}$ and $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$ for every bin V_j — in other words, we seek regularity conditions that allow for conditional independence to be tested

indirectly using multiple two-sample independence tests. Of course, one must also ensure that the two-sample tests comparing $\mathcal{L}_{(X,Y)|Z \in V_j}$ and $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$ are themselves feasible (in the sense, for instance, of having explicit finite-sample control on the Type I & II error).

Our work uses the formalism of *optimal transport*, in particular the *Wasserstein distance* W_p , in an essential way. As a metric structure on the space of probability measures, the Wasserstein distance is only slightly stronger than weak convergence, so the stipulation that (for instance) “for small V_j , $W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z \in V_j}) \approx 0$ ” is comparatively easy to satisfy. At the same time, it is also possible to use the Wasserstein distance to conduct two-sample tests of quite general probability measures.

The structure of the paper is as follows. In Section 1, we give an overview of the necessary background from optimal transport, and discuss how to use the Wasserstein distance to conduct two-sample independence tests. In Section 2 we review the general infeasibility of conditional independence testing, but show that if the conditional distributions $\mathcal{L}_{X|Z=z}$, $\mathcal{L}_{Y|Z=z}$, and $\mathcal{L}_{(X,Y)|Z=z}$ are *Lipschitz* functions of z when the space of probability measures is equipped with the W_p metric, then binning the conditioning variable Z incurs a small, explicit error in the W_p metric. We then explain how this allows for a “ W_p multiple two-sample” conditional independence test, with explicit finite-sample control of the Type I and Type II error, under a wide variety of mild auxiliary regularity assumptions (which are needed so that the two-sample tests are themselves feasible). Additionally, in Section 3 we give a “plug-in” estimator for the W_p -Lipschitz constants of $\mathcal{L}_{X|Z=z}$, $\mathcal{L}_{Y|Z=z}$, and $\mathcal{L}_{(X,Y)|Z=z}$.

Note. While the work presented in this article was in progress, the author learned of the recent article [NBW20], which also considers conditional independence testing under smoothness assumptions on the conditional distributions $\mathcal{L}_{(X,Y)|Z}$, $\mathcal{L}_{X|Z}$, and $\mathcal{L}_{Y|Z}$. The two works are similar in spirit, but complementary. On the one hand, as discussed below in Section 2, our background assumptions are weaker than those used in [NBW20]; on the other hand, [NBW20] provides delicate minimax estimates which are currently unavailable in our setting.

1. TWO-SAMPLE INDEPENDENCE TESTING WITH THE WASSERSTEIN DISTANCE

We first consider testing whether $X \perp\!\!\!\perp Y$. Note that this amounts to a two-sample test between the distributions $\mathcal{L}_{(X,Y)}$ and $\mathcal{L}_X \otimes \mathcal{L}_Y$. Furthermore, note that given the ability to sample from $\mathcal{L}_{(X,Y)}$, we can always artificially sample from $\mathcal{L}_X \otimes \mathcal{L}_Y$ as well — for instance, one can sample from $\mathcal{L}_{(X,Y)}$ $2n$ times, and for the first n samples discard the Y values, and for the second n samples discard the X values, and then $(x_1, y_{n+1}), (x_2, y_{n+2}), \dots, (x_n, y_{2n})$ will be drawn from $\mathcal{L}_X \otimes \mathcal{L}_Y$.

Specifically, we consider Wasserstein-based two sample tests (cf. [RGTC17], which considers some stronger results in the univariate case). We do not attempt to give a general survey on optimal transport/Wasserstein distances here (a more than sufficient background reference is [San15]), but we give a short motivational discussion. Recall that for $p \in [1, \infty)$, the p -Wasserstein distance on the space $\mathcal{P}_p(U)$ of Borel probability measures with finite p -th moments on the domain $U \subseteq \mathbb{R}^d$ is given by

$$W_p(\mu, \nu) := \inf_{\gamma \in \Pi(\mu, \nu)} \left(\int_{U^2} \|x - y\|^p d\gamma(x, y) \right)^{1/p}$$

where $\Pi(\mu, \nu)$ is the space of all *couplings* of μ and ν , namely all probability measures on U^2 with marginals μ and ν . For $p \in [1, \infty)$, the p -Wasserstein distance metrizes convergence in distribution (a.k.a. narrow convergence, a.k.a. weak convergence of probability measures, etc.). In the jargon of optimal transport, a measure $\gamma \in \Pi(\mu, \nu)$ is said to be a *transport plan* between μ and ν , and

moreover γ is said to be an *optimal plan* if the infimum in the definition of W_p is attained at γ , that is, $W_p(\mu, \nu) = \left(\int_{U^2} \|x - y\|^p d\gamma(x, y) \right)^{1/p}$. It is a basic feature of the theory that, given arbitrary measures $\mu, \nu \in \mathcal{P}_p(U)$ on a “reasonable” domain U (specifically, we require that U be a Polish space), there exists an optimal plan γ between μ and ν , but γ need not be unique.

If, moreover, we restrict our attention to *absolutely continuous measures* (say with respect to the Lebesgue measure on U), then the p -Wasserstein distances enjoy another interpretation, in terms of optimal transport maps rather than optimal couplings. If we write $T_{\#}\mu$ for the pushforward of μ by the map T (that is, the measure defined by $T_{\#}\mu(A) := \mu(T^{-1}(A))$), then it turns out that when μ is absolutely continuous,

$$W_p(\mu, \nu) = \inf_{T: T_{\#}\mu = \nu} \left(\int_U \|x - T(x)\|^p d\mu(x) \right)^{1/p}.$$

There is an extensive theory studying these *transport maps* — under what circumstances the infimum in the previous expression is attained, whether the map T instantiating the infimum is unique, what regularity properties it possesses, and so on. This theoretical apparatus will not play a role in our arguments. However, there is one further feature of the p -Wasserstein distances that will be very helpful for us, which is the following. Suppose that x_1, x_2, \dots are i.i.d. samples from μ . Let $\mu_n := \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$ denote the n th *empirical measure* for μ . (Note that μ_n is random, but μ is fixed.) One variant of the law of large numbers (namely, the Glivenko-Cantelli theorem) states that μ_n converges to μ in distribution with probability 1; accordingly, $W_p(\mu_n, \mu) \rightarrow 0$ almost surely, when $p \in [1, \infty)$ (and, it turns out, when $p = \infty$ as well). In fact, $W_p(\mu_n, \mu)$ possesses known sample complexity bounds: for instance, if the domain U on which μ resides is compact, d -dimensional and Hausdorff regular, and (for instance) μ is absolutely continuous w.r.t. the volume measure on U , then for any $p \in [1, d/2)$ the W_p sample complexity of μ is no worse than $O(n^{-1/d})$ [WB19, Theorem 1 and Proposition 8] (see also discussion in the introduction of [AG19] or [NWR19] for a brief overview of similar results). Significant improvements on this bound are possible in special cases, for example if μ is actually supported on a lower-dimensional surface, or possesses smooth density, or satisfies one of an assortment of other regularity conditions (see e.g. [BGV07, WB19]). Note also that if $\mu \neq \nu$, then the convergence of $W_p(\mu_n, \nu_n)$ to $W_p(\mu, \nu)$ was recently shown to occur at a faster rate than $O(n^{-1/d})$ in general [CRL⁺20, MNW21].

Now let’s return our focus to (two-sample) independence testing for X and Y .

Let’s be slightly more explicit about what this test looks like. The null hypothesis H_0 will be that X and Y are indeed independent, so $\mathcal{L}_{(X,Y)} = \mathcal{L}_X \otimes \mathcal{L}_Y$. We first approximate both $\mathcal{L}_{(X,Y)}$ and $\mathcal{L}_X \otimes \mathcal{L}_Y$ with empirical measures based on samples, for instance

$$\frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)} \quad \text{and} \quad \frac{1}{n} \sum_{i=1}^n \delta_{(x_{n+i}, y_{2n+i})}$$

respectively. Under the null hypothesis, we use the fact that $\mathcal{L}_{(X,Y)} = \mathcal{L}_X \otimes \mathcal{L}_Y$ to deduce that

$$\begin{aligned} W_p \left(\frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)}, \frac{1}{n} \sum_{i=1}^n \delta_{(x_{n+i}, y_{2n+i})} \right) \\ \leq W_p \left(\frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)}, \mathcal{L}_{(X,Y)} \right) + W_p \left(\mathcal{L}_X \otimes \mathcal{L}_Y, \frac{1}{n} \sum_{i=1}^n \delta_{(x_{n+i}, y_{2n+i})} \right) \end{aligned}$$

and observe that

$$W_p \left(\frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)}, \mathcal{L}_{(X, Y)} \right) \xrightarrow{n \rightarrow \infty} 0, \quad W_p \left(\mathcal{L}_X \otimes \mathcal{L}_Y, \frac{1}{n} \sum_{i=1}^n \delta_{(x_{n+i}, y_{2n+i})} \right) \xrightarrow{n \rightarrow \infty} 0.$$

This motivates the following definition:

Definition 1. (*p*-Wasserstein Two Sample Independence Test) The null hypothesis H_0 is that $\mathcal{L}_{(X, Y)} = \mathcal{L}_X \otimes \mathcal{L}_Y$. We then set some “level” $\varepsilon_0 > 0$, and say that we reject the null hypothesis if

$$W_p \left(\frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)}, \frac{1}{n} \sum_{i=1}^n \delta_{(x_{n+i}, y_{2n+i})} \right) \geq \varepsilon_0,$$

and we fail to reject otherwise.

In other words, the test above prescribes using the *plug-in estimator* $W_p(\widehat{\mathcal{L}}_{(X, Y)}, \widehat{\mathcal{L}}_X \otimes \widehat{\mathcal{L}}_Y)$ for $W_p(\mathcal{L}_{(X, Y)}, \mathcal{L}_X \otimes \mathcal{L}_Y)$.

Remark. Other finite-sample estimators for the W_p distance between two measures have been considered in the literature, and may even be more desirable in some situations (see discussion, for instance, in [CRL⁺20]). We restrict our attention to the plug-in estimator only in the interest of simplicity.

To emphasize, the false rejection rate under the null is governed by the previously stated W_p sample complexity bounds for empirical measures. On the other hand, the false acceptance rate under the alternative (that is, the probability of the two empirical measures being within ε_0 of each other, even though $\mathcal{L}_{(X, Y)} \neq \mathcal{L}_X \otimes \mathcal{L}_Y$) depends on which distribution $\mathcal{L}_{(X, Y)}$ actually is; more specifically, it depends on $W_p(\mathcal{L}_{(X, Y)}, \mathcal{L}_X \otimes \mathcal{L}_Y)$.

Before proceeding to testing procedures for conditional independence in Section 2 below, we give a more explicit analysis of the “Wasserstein two-sample” independence test described above. Namely: how must ε_0 be chosen in order to ensure a given p -value and power function for the test? Morally, this is “just” an application of the sample complexity estimates for W_p already mentioned; note that $\mathcal{L}_{(X, Y)}$ and $\mathcal{L}_X \otimes \mathcal{L}_Y$ reside in $\mathbb{R}^{d_X + d_Y}$, so the “off-the-shelf” W_p sample complexity for such a two-sample test is $O(n^{-1/(d_X + d_Y)})$ (for $p \in [1, \frac{1}{2}(d_X + d_Y)]$). However, the “off-the-shelf” sample complexity is an *asymptotic* result, and is therefore unsuitable e.g. for the construction of an explicit confidence interval. At the same time, while there is notable recent progress on *central limit theorems* for the empirical Wasserstein distance, for instance for the asymptotic distribution of the Wasserstein distance $W_p(\mu_n, \nu)$ where μ_n is an empirical measure for μ [dBL19, dBGSL21], the result therein are also insufficient to construct explicit confidence intervals without more information regarding the quantity $\mathbb{E}W_p(\mu_n, \nu)$.

We therefore proceed by an alternative analysis, which has essentially two ingredients:

- (1) Given a measure μ and an empirical measure μ_n for μ , we desire an explicit upper bound on the *expected error* introduced by passing to the empirical measure μ_n , that is, the quantity $\mathbb{E}W_p(\mu, \mu_n)$. In what follows, we call this the *expectation bound*.
- (2) Additionally, we would like a *concentration inequality* indicating how the random quantity $W_p(\mu, \mu_n)$ is distributed around its expectation $\mathbb{E}W_p(\mu, \mu_n)$.

If we have these two ingredients, it is straightforward to produce an upper bound on the probability that, under the null, the test indicated in Definition 1 will nonetheless be rejected.

For an example of a suitable concentration inequality, let us mention the following:

Proposition 2 ([WB19, Proposition 20]). *Let (X, d) be a Polish metric space, and let $\mu \in \mathcal{P}_p(X)$ be a measure supported on a set of diameter at most D . Let μ_n denote an empirical measure for μ . Then,*

$$\mathbb{P} \left[W_p^p(\mu, \mu_n) \geq \mathbb{E} W_p^p(\mu, \mu_n) + t \right] \leq \exp \left(-\frac{2nt^2}{D^{2p}} \right).$$

Remark. Weed and Bach only state their concentration inequality in the case where $D = 1$. However the modification of their argument for general diameter is quite routine; for completeness, we rerun the argument with general diameter in Appendix B.

It is also possible to extend the argument for Proposition 2 to the case of a measure μ with unbounded support on a Polish metric space (X, d) , at least provided μ also lies in $\mathcal{P}_q(X)$ for some $q > p$.

Theorem 3 (W_p empirical measure concentration inequality with higher moment bound). *Let (X, d) be a Polish metric space with (arbitrary) distinguished point $e \in X$. Let $\mu \in \mathcal{P}(X)$ with $(\int_X d(e, x)^q d\mu(x))^{1/q} \leq M_q$ for some $q > p \geq 1$. Let $\mu_n = \sum_{i=1}^n \delta_{x_i}$ be a (random) empirical measure for μ . Furthermore, let $\eta \in (0, 1)$, and let*

$$\bar{\mu} \upharpoonright B(e, M_q \eta^{-1/q}) := \frac{1}{\mu(B(e, M_q \eta^{-1/q}))} \mu \upharpoonright B(e, M_q \eta^{-1/q})$$

denote the normalized restriction of μ to $B(e, M_q \eta^{-1/q})$, and let $\mu_n \mid (X_1, \dots, X_n) \in B(e, M_q \eta^{-1/q})^n$ denote the conditional empirical measure where $x_i \in B(0, M_q \eta^{-1/q})^n$ for all $i = 1, \dots, n$. Then, for any $t > 0$, it holds that

$$\begin{aligned} \mathbb{P} \left[W_p^p(\mu_n, \mu) \geq \mathbb{E} \left[\left(W_p \left(\mu_n \mid (X_1, \dots, X_n) \in B(0, M_q \eta^{-1/q})^n, \bar{\mu} \upharpoonright B(0, M_q \eta^{-1/q}) \right) + c_{p,q} M_q \eta^{1/p-1/q} \right)^p \right] + t \right] \\ \leq 1 - (1 - \eta)^n + \exp \left(-\frac{2n(t - (1 - (1 - \eta)^n)((M_q \eta^{-1/q})^p)_+^2)}{(M_q \eta^{-1/q})^{2p}} \right) \end{aligned}$$

where $c_{p,q} := 2^{1/q} \frac{2^{1/p-1/q}}{2^{1/p-1/q}-1} + \frac{2^{1/p}}{2^{1/p}-1}$.

This concentration inequality is of independent interest; we give a proof in Appendix B.

It turns out that of the two ingredients — the expectation bound, and the concentration inequality — it is the former that is the more challenging one, and indeed has been the subject of ongoing investigation by numerous authors, such as [BLG14, DSS13, FG15, Lei20, WB19]. In particular, it is known that estimating the error introduced by passing to empirical measures for distributions $\mathcal{L}_{(X,Y)}$ and $\mathcal{L}_X \otimes \mathcal{L}_Y$ depends sensitively on the dimension $d_X + d_Y$; we first address the case where $d_X + d_Y \geq 3$.

Theorem 4 ([DSS13, Theorem 1]). *Let μ be a measure on \mathbb{R}^d with $d \geq 3$, and let μ_n be an empirical measure for μ . Let $p \in [1, d/2)$ and $q > dp/(d-p)$. Then there exists a constant $\kappa_{p,q,d}$ depending only on p, q , and d , such that*

$$\mathbb{E} [W_p^p(\mu_n, \mu)]^{1/p} \leq \kappa_{p,q,d} \left[\int_{\mathbb{R}^d} \|x\|^q d\mu(x) \right]^{1/q} n^{-1/d}$$

where $\|\cdot\|$ is a norm on \mathbb{R}^d and $\kappa_{p,q,d}$ is given explicitly in [DSS13].

Observe that by combining the estimates

$$\mathbb{P}[|W_p^p(\mu_n, \mu) - \mathbb{E}[W_p^p(\mu_n, \mu)]| \geq t] \leq \exp(-2nt^2/D^{2p})$$

and

$$\mathbb{E}[W_p^p(\mu_n, \mu)]^{1/p} \leq \kappa_{p,q,d} \left[\int_{\mathbb{R}^d} \|x\|^q d\mu(x) \right]^{1/q} n^{-1/d}$$

it follows directly that in the case where μ has support with diameter at most D (and therefore $\int_{\mathbb{R}^d} \|x\|^q d\mu(x) < \infty$),

$$\begin{aligned} \mathbb{P}[W_p^p(\mu_n, \mu) \geq t + \mathbb{E}[W_p^p(\mu_n, \mu)]] &\leq \mathbb{P}\left[W_p^p(\mu_n, \mu) \geq t + \kappa_{p,q,d}^p \left[\int_{\mathbb{R}^d} \|x\|^q d\mu(x) \right]^{p/q} n^{-p/d}\right] \\ &\leq \exp(-2nt^2/D^{2p}). \end{aligned}$$

For instance (although this is far from the most efficient upper bound in $n!$), if n is large enough that $\kappa_{p,q,d}^p \left[\int_{\mathbb{R}^d} \|x\|^q d\mu(x) \right]^{p/q} n^{-p/d} \leq \frac{1}{2} \left(\frac{\varepsilon}{2}\right)^p$, then putting $t = \frac{1}{2} \left(\frac{\varepsilon}{2}\right)^p$ gives

$$\mathbb{P}\left[W_p^p(\mu_n, \mu) \geq \left(\frac{\varepsilon}{2}\right)^p\right] \leq \exp\left(-\frac{2n\varepsilon^{2p}}{4^{p+1}D^{2p}}\right)$$

or equivalently

$$\mathbb{P}\left[W_p(\mu_n, \mu) \geq \frac{\varepsilon}{2}\right] \leq \exp\left(-\frac{2n\varepsilon^{2p}}{4^{p+1}D^{2p}}\right).$$

Lastly, note that if μ_n and μ'_n are both empirical measures drawn from the same distribution μ , and $W_p(\mu_n, \mu'_n) \geq \varepsilon$, then it must be the case that either $W_p(\mu_n, \mu) \geq \frac{\varepsilon}{2}$ or $W_p(\mu'_n, \mu) \geq \frac{\varepsilon}{2}$. Hence,

$$\mathbb{P}[W_p(\mu_n, \mu'_n) \geq \varepsilon] \leq 2 \exp\left(-\frac{2n\varepsilon^{2p}}{4^{p+1}D^{2p}}\right).$$

Likewise, if μ has unbounded support but still satisfies a higher q th moment bound, as per Theorem 4, one can combine Theorems 3 and 4 in a similar manner: see Corollary 14 below.

Theorems 3 and 4 are sufficient when $\mu \in \mathcal{P}_p(\mathbb{R}^d)$ with $d \geq 3$. However, in our situation, it is also important to address the case where $d = 2$ (that is: both the variables X and Y are univariate). This case must be handled by special analysis, which we conduct, in the case of a simple compact domain, in Example 15, below, in Appendix B.

It is also possible to explicitly bound the type II error of the test given in Definition 1, as a function of $W_p(\mathcal{L}_{(X,Y)}, \mathcal{L}_X \otimes \mathcal{L}_Y)$. Indeed, suppose that $W_p(\mathcal{L}_{(X,Y)}, \mathcal{L}_X \otimes \mathcal{L}_Y) \geq \varepsilon_0 + \delta_0$. Then, if a type II error occurs, namely $W_p\left(\frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)}, \frac{1}{n} \sum_{i=1}^n \delta_{(x_{n+i}, y_{2n+i})}\right) < \varepsilon_0$, it must be the case that at least one of the following holds:

$$W_p\left(\mathcal{L}_{(X,Y)}, \frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)}\right) > \frac{\delta_0}{2} \text{ or } W_p\left(\mathcal{L}_X \otimes \mathcal{L}_Y, \frac{1}{n} \sum_{i=1}^n \delta_{(x_{n+i}, y_{2n+i})}\right) > \frac{\delta_0}{2}.$$

But we can upper bound the probability of each of these events using exactly the same analysis — expectation bound combined with concentration inequality — that we have just described.

Additionally, let us remark on the task of actually computing the estimator $W_p(\hat{\mu}_n, \hat{\nu}_n)$ for $W_p(\mu, \nu)$. Computing the optimal coupling for $\hat{\mu}_n$ and $\hat{\nu}_n$ takes the form of a classical (discrete) *optimal matching* problem, which has been extensively studied and is amenable to the standard methods of linear programming. State-of-the-art computational methods for this problem are surveyed, for instance, in [PC19]; for example, fast methods based on *Sinkhorn's algorithm*, discussed

therein, achieve a computational cost of $O(n^2 \log n)$. Therefore, when conducting a Wasserstein two-sample test, one must work with empirical measures supported on a *moderate* number n of points, so that the test achieves a small enough p -value, while at the same time the estimator itself is feasible to compute.

In what follows, we (to some extent) work *modulo* the existence of a Wasserstein two-sample (independence) test with a given p -value and power function. However, we hope that from the previous discussion, the reader is convinced that such a test is feasible, albeit subject to ongoing technical improvements, *viz.* in terms of expectation bounds, concentration inequalities, and computational complexity.

2. CONDITIONAL INDEPENDENCE TESTING VIA BINNING

Now we consider conditional independence testing. That is, we ask whether $X \perp\!\!\!\perp Y \mid Z$. At the level of distributions, this amounts to comparing $\mathcal{L}_{(X,Y)\mid Z=z}$ with $\mathcal{L}_{X\mid Z=z} \otimes \mathcal{L}_{Y\mid Z=z}$ for *every* z in the range of Z . (Note that these distributions should be understood as disintegrations of the joint distributions $\mathcal{L}_{(X,Y,Z)}$ and $\mathcal{L}_{(X,Z)}$ and $\mathcal{L}_{(Y,Z)}$ w.r.t Z , e.g. in the sense of *regular conditional distributions*.) That is, conditional independence testing may be understood as (at least implicitly) requiring a *continuum* number of two-sample tests.

It should now be clear to the reader that without further assumptions on the random variables X, Y , and Z (and their joint distribution), finite sample conditional independence testing is futile. Indeed, it may be that for all of the (finitely many) values z of Z that we actually get to sample, it is the case that $\mathcal{L}_{(X,Y)\mid Z=z} = \mathcal{L}_{X\mid Z=z} \otimes \mathcal{L}_{Y\mid Z=z}$; but for some z' out of our sample, $\mathcal{L}_{(X,Y)\mid Z=z'}$ and $\mathcal{L}_{X\mid Z=z'} \otimes \mathcal{L}_{Y\mid Z=z'}$ are very far apart as distributions. Compare with the analogous concern for $\mathcal{L}_{(X,Y)}$ versus $\mathcal{L}_X \otimes \mathcal{L}_Y$; here, we have a guarantee from the (convergence of measures version of the) law of large numbers that empirical measures converge to the measures they are drawn from almost surely, with a deviation rate/concentration bounds given by Sanov's theorem or similar. We have no such luck here, simply from the fact that $\mathcal{L}_{(X,Y)\mid Z=z}$ and $\mathcal{L}_{(X,Y)\mid Z=z'}$ are different distributions with no *a priori* relationship. (The proof in [SP20] which shows that conditional independence testing is not feasible in general, proceeds by way of a different, but in some sense related, argument: there, they construct two joint distributions of random variables (X, Y, Z) and $(\tilde{X}, \tilde{Y}, \tilde{Z})$ where each of $\{X, \tilde{X}\}$, $\{Y, \tilde{Y}\}$, and $\{Z, \tilde{Z}\}$ are close with high probability, hence costly to distinguish with finite samples, yet (X, Y, Z) and $(\tilde{X}, \tilde{Y}, \tilde{Z})$ have different conditional dependence relations.)

So in order to construct a conditional independence test which, say, comes with any kind of finite sample guarantee, some auxiliary assumptions are necessary. In particular, if we knew that all the marginal distributions varied in some continuous fashion with z , we would be able to use the fact that $\mathcal{L}_{(X,Y)\mid Z=z} \approx \mathcal{L}_{(X,Y)\mid Z=z'}$ whenever $z \approx z'$ (and likewise for the families of distributions $\mathcal{L}_{X\mid Z=z}$ and $\mathcal{L}_{Y\mid Z=z}$), to deduce some overall estimate of conditional dependence by binning the variable Z , and then performing a (possibly very large) number of separate two-sample independence tests.

In what follows, we consider the special case where \mathcal{L}_Z is compactly supported inside \mathbb{R}^{d_Z} ; however, we do not assume compact support for the conditional distributions $\mathcal{L}_{(X,Y)\mid Z=z}$.

We also make the following type of assumption: consider the function

$$\begin{aligned} \mathbb{R}^{d_Z} &\rightarrow \mathcal{P}(\mathbb{R}^{d_X+d_Y}) \\ z &\mapsto \mathcal{L}_{(X,Y)\mid Z=z}. \end{aligned}$$

We ask that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ is Lipschitz continuous, provided we equip $\mathcal{P}(\mathbb{R}^{d_X+d_Y})$ with the W_p metric; in other words,

$z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ is L_{XY} -Lipschitz \iff

$$(\forall z, z' \in \mathbb{R}^{d_Z} \cap \text{supp}(\mathcal{L}_Z)) W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z'}) \leq L_{XY}|z - z'|.$$

Observe that requiring that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ be L -Lipschitz not only imposes that $z \approx z' \implies \mathcal{L}_{(X,Y)|Z=z} \approx_{W_p} \mathcal{L}_{(X,Y)|Z=z'}$, it does so in a quantitative fashion: if we know that $|z - z'| < \varepsilon/L_{XY}$, then we can say that $W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z'}) < \varepsilon$.

In addition, we will also require that the functions

$$z \mapsto \mathcal{L}_{X|Z=z} \text{ and } z \mapsto \mathcal{L}_{Y|Z=z}$$

be Lipschitz continuous, in other words,

$$\begin{aligned} (\forall z, z' \in \mathbb{R}^{d_Z} \cap \text{supp}(\mathcal{L}_Z)) \quad W_p(\mathcal{L}_{X|Z=z}, \mathcal{L}_{X|Z=z'}) &\leq L_X|z - z'|; \\ (\forall z, z' \in \mathbb{R}^{d_Z} \cap \text{supp}(\mathcal{L}_Z)) \quad W_p(\mathcal{L}_{Y|Z=z}, \mathcal{L}_{Y|Z=z'}) &\leq L_Y|z - z'|. \end{aligned}$$

In what follows, it turns out that we will need to set some small parameters (specifically the diameter of bins for the range of Z) to be small compared to the inverse of *all three* optimal Lipschitz constants simultaneously.

Remark. There are, of course, *many* inequivalent metrics of interest on the space of probability measures. Why choose the W_p metric (for some $p \in [1, \infty)$) in particular? Indeed, the recent article [NBW20], also concerned with conditional independence testing, makes extensive use of the total variation (TV) metric rather than a metric coming from an optimal transport problem. To some extent, the choice of metric on the space of probability measures is simply a modeling decision, but let us emphasize a topological point in favor of the W_p metrics. On compact sets, convergence in W_p is equivalent to convergence in distribution, a.k.a. convergence in the probabilist's weak topology; on the other hand, the TV norm metrizes *strong* convergence, equivalently convergence in L^1 . Since the strong topology is (as the same suggests) stronger than the weak topology, it follows that *fewer* functions from \mathbb{R}^d into the space of probability measures are strongly continuous than weakly continuous. In our particular situation, this means that on a compact domain, the requirement that conditional distributions of the form $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ be W_p -continuous is less stringent than requiring that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ be TV-continuous. This difference in strength of assumption is especially evident in the case where the conditional distributions $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ (and $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$) are *mutually singular* for different values of z , as is the case, for example, when the conditional distributions are supported on disjoint hypersurfaces parametrized by z . For emphasis, we give a couple of concrete instances of this type in the following example.

Example 5. (examples of conditional distribution which is W_p -Lipschitz but not TV-continuous)
(1) Consider the slightly trivial case where $Y = X$ and $Z = X$. In this case, $\mathcal{L}_{(X,Y)|Z=z} = \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z} = \delta_z \otimes \delta_z$. In particular, if (z_n) is a sequence which converges to z in \mathbb{R}^{d_Z} (but $z_n \neq z$ for all $n \in \mathbb{N}$), then $(\delta_{z_n} \otimes \delta_{z_n})$ converges to $\delta_z \otimes \delta_z$ in the weak topology as well as in W_p (for every $p \in [1, \infty)$). However, $TV(\delta_{z_n} \otimes \delta_{z_n}, \delta_z \otimes \delta_z) \not\rightarrow 0$, since the total variation distance between *any* two mutually singular measures is identically 1.

(2) Suppose that X and Y are both random variables in \mathbb{R}^2 (equipped with polar coordinates) where the radial variable for X and Y is deterministically coupled. In other words, there is some random variable $Z : \Omega \rightarrow \mathbb{R}_+$ where $X = (Z, \Theta_1(Z, \cdot))$ and $Y = (Z, \Theta_2(Z, \cdot))$ where $\Theta_1(r, \cdot)$ and $\Theta_2(r, \cdot)$ are both ‘‘random angles’’, that is, random variables from Ω into $[0, 2\pi)$, which may depend

on the radial variable r . In this case, X and Y may both have full support in \mathbb{R}^2 ; but the conditional distributions $\mathcal{L}_{X|Z=z}$ and $\mathcal{L}_{Y|Z=z}$ are always supported within the sphere of radius z , and so $\mathcal{L}_{X|Z=z}$ and $\mathcal{L}_{Y|Z=z}$ (and also $\mathcal{L}_{(X,Y)|Z=z}$) are mutually singular for all distinct z and z' . It follows that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ and $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$ cannot be TV -continuous, but may still be W_p -continuous if $\Theta_1(r, \cdot)$ and $\Theta_2(r, \cdot)$ vary continuously with r (we demonstrate this immediately below, in Example 6, in the particular case where $\Theta_1(r, \cdot)$ and $\Theta_2(r, \cdot)$ are Lipschitz continuous in r).

It is certainly worth asking how W_p -Lipschitz type assumptions relate to the candidate functional relations between X , Y , and Z in the causal inference setting. For instance, if there is no direct functional relationship between X and Y but $X = f(Z) + E_1$ and $Y = g(Z) + E_2$ (where E_1 and E_2 are Gaussian noise variables), how do the W_p -Lipschitz assumptions on the conditional distributions we have just detailed relate to the regularity properties of f and g ?

Example 6 (Additive noise model; Z causes X and Y but $X \perp\!\!\!\perp Y \mid Z$). Suppose we are given random variables X, Y , and Z , and we know that $X = f(Z) + E_1$ and $Y = g(Z) + E_2$, where E_1 and E_2 are independent and identically distributed noise variables (for instance, E_1 and E_2 may be (truncated) Gaussians), and f and g are (deterministic) Lipschitz continuous functions. We claim that in this circumstance, all of $z \mapsto \mathcal{L}_{X|Z=z}$, $z \mapsto \mathcal{L}_{Y|Z=z}$, and $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ are Lipschitz continuous with respect to W_p . Note that in this case, if we condition on $Z = z_0$, then X has the distribution $f(z_0) + E_1$ and Y has the distribution $g(z_0) + E_2$. Thus, if both f and g are L -Lipschitz, the distributions of $X|Z = z_0$ and $X|Z = z_1$ are identical, *modulo* a shift by a constant at most ε , provided that $|z_0 - z_1| < \varepsilon/L$, and similarly for $Y|Z = z_0$ and $Y|Z = z_1$ respectively. Note that since $\mathcal{L}_{X|Z=z_0} = f(z_0) + \mathcal{L}_{E_1}$, it follows that the shift map

$$x \mapsto x + f(z_1) - f(z_0)$$

pushes $\mathcal{L}_{X|Z=z_0}$ onto $\mathcal{L}_{X|Z=z_1}$. Consequently, this shift is an admissible transport map in the optimal transport problem, and it follows that

$$W_p(\mathcal{L}_{X|Z=z_0}, \mathcal{L}_{X|Z=z_1}) \leq \left(\int_{\Omega} |f(z_1) - f(z_0)|^p d\mathcal{L}_{X|Z=z_0} \right)^{1/p}.$$

But since $|f(z_1) - f(z_0)| < \varepsilon$, it follows that $W_p(\mathcal{L}_{X|Z=z_0}, \mathcal{L}_{X|Z=z_1}) < \varepsilon$. By an identical argument, it also holds that $W_p(\mathcal{L}_{Y|Z=z_0}, \mathcal{L}_{Y|Z=z_1}) < \varepsilon$. And, of course, in this case, X and Y are, in fact conditionally independent given Z , so since $\mathcal{L}_{(X,Y)|z} = \mathcal{L}_{X|z} \otimes \mathcal{L}_{Y|z}$ we see that an analogous estimate applies to the joint distribution; that is, that since the product shift map

$$(x, y) \mapsto (x + f(z_1) - f(z_0), y + g(z_1) - g(z_0))$$

pushes $\mathcal{L}_{X|z_0} \otimes \mathcal{L}_{Y|z_0}$ onto $\mathcal{L}_{X|z_1} \otimes \mathcal{L}_{Y|z_1}$, we have the transport map upper bound

$$\begin{aligned} W_p^p(\mathcal{L}_{(X,Y)|z_0}, \mathcal{L}_{(X,Y)|z_1}) &\leq \int_{\Omega^2} |(f(z_1), g(z_1)) - (f(z_0), g(z_0))|^p d\mathcal{L}_{X|Z=z_0} \otimes d\mathcal{L}_{Y|Z=z_0} \\ &\leq 2^{p-1} \left(\int_{\Omega^2} (|f(z_1) - f(z_0)|^p + |g(z_1) - g(z_0)|^p) d\mathcal{L}_{X|Z=z_0} \otimes d\mathcal{L}_{Y|Z=z_0} \right) \\ &= 2^{p-1} \int_{\Omega} |f(z_1) - f(z_0)|^p d\mathcal{L}_{X|Z=z_0} + 2^{p-1} \int_{\Omega} |g(z_1) - g(z_0)|^p d\mathcal{L}_{Y|Z=z_0} \\ &\leq (2\varepsilon)^p. \end{aligned}$$

Remark. On the other hand, if we consider a non-additive noise model, then it is less clear whether there is clean a relationship between the Lipschitz property of $z \mapsto \mathcal{L}_{(X,Y)|z}$ and the functional form of the causal relations between X, Y , and Z .

As promised, the Lipschitz assumption allows us to perform a conditional independence test on (X, Y, Z) using binning. Let us now summarize how this works, before proceeding with a series of technical arguments. The idea is to divide the range of Z into cells V_j of diameter at most ε/L where L is some function of L_X , L_Y , and L_{XY} (here one could use, say, cubes or Voronoi cells), and compute empirical estimates of the conditional distributions $\mathcal{L}_{(X,Y)|z \in V_j}$ and $\mathcal{L}_{X|z \in V_j} \otimes \mathcal{L}_{Y|z \in V_j}$. Then, for each cell V_j , we compute the Wasserstein distance between those two empirical distributions; if for each V_j the distance between the two empirical distributions is small, then we can conclude that with high probability, X and Y are conditionally independent given “ Z discretized onto the cells V_j ”. A fancier way to say this is that if \mathcal{V} is the σ -algebra in the sample space generated by the partition $Z^{-1}\{V_j\}$, we are testing whether $X \perp\!\!\!\perp Y \mid \mathbb{E}[Z \mid \mathcal{V}]$. Finally, our Lipschitz assumption will allow us to deduce a relationship between our test for $X \perp\!\!\!\perp Y \mid \mathbb{E}[Z \mid \mathcal{V}]$, and whether $X \perp\!\!\!\perp Y \mid Z$.

Crucially, we must explicitly control the error introduced by replacing Z with $\mathbb{E}[Z \mid \mathcal{V}]$. At the heuristic level, the situation is rather straightforward. Indeed, suppose that our joint and conditional distributions have densities; consider (x_1, y_1, z_1) and (x_2, y_2, z_2) , both samples from the joint density $p(x, y, z)$. Suppose that z_1 and z_2 belong to the same cell, and thus $z_2 \in B_{\varepsilon/L}(z_1)$. We want to pretend that up to some small error, (x_2, y_2) is a sample from $p(x, y|z_1)$ rather than $p(x, y|z_2)$, as is actually the case. If we had an “optimal transport oracle”, we could ask for the optimal transport map T from $p(x, y|z_2)$ to $p(x, y|z_1)$, and then just compute $T((x_2, y_2))$; then this *would* be a sample from $p(x, y|z_1)$. Instead, we do nothing to (x_2, y_2) , but since $p(x, y|z_2)$ and $p(x, y|z_1)$ are close in the Wasserstein sense, we know that $\|T - id\|$ is small in the corresponding norm; this will allow us to say something like

$$W_p \left(\frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)|z_i \in V_j}, p(x, y|z_1) \right) \leq \varepsilon.$$

Despite this pleasing heuristic motivation, phrased in terms of optimal maps and samples, it turns out to be more direct, from a technical standpoint, to work with optimal plans and population measures, which is how we will proceed in the remainder of the section. Our analysis depends on the following lemma (stated with under rather general assumptions, since the proof is no more difficult):

Lemma 7. *Let $z \mapsto \mu_z$ be a Borel measurable function from a Polish space S , to $(\mathcal{P}_p(U), W_p)$, where (U, d) is a Polish metric space, and let $V \subseteq S$ be a Borel set inside S . Let λ be a probability measure on V , and let $\nu \in W_p(U)$. Then,*

$$W_p^p \left(\int_V \mu_z d\lambda(z), \nu \right) \leq \int_V W_p^p(\mu_z, \nu) d\lambda(z).$$

In fact, the proof of such a result is more or less indicated by the contents of [Vil08, Ch. 4 & 5]; however, the argument therein omits a number of technical points, so we nonetheless give a rather explicit proof in Appendix A.

We can now easily prove the following:

Proposition 8. *Suppose that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ be L_{XY} -Lipschitz with respect to W_p . Suppose that $\{V_j\}_{j=1}^J$ is a measurable partition of the support of \mathcal{L}_Z , and that $\text{diam}(V_j) \leq \varepsilon/L_{XY}$ for every bin*

V_j in the partition. Then, for all $z_0 \in V_j$, it holds that

$$W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{(X,Y)|Z=z_0}) \leq \varepsilon.$$

Proof. In the setting of Lemma 7, let $V = V_j$, $U = \mathbb{R}^{d_X} \times \mathbb{R}^{d_Y}$, and

$$\mu_z = \mathcal{L}_{(X,Y)|Z=z} \text{ and } \nu = \mathcal{L}_{(X,Y)|Z=z_0}.$$

Furthermore, let $\lambda = \frac{1}{\mathcal{L}_Z(V_j)} \mathcal{L}_Z \upharpoonright V_j$. Under the assumptions of the proposition,

$$W_p^p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z_0}) \leq \varepsilon^p \text{ for all } z, z_0 \in V_j.$$

Since the function $z \mapsto \mu_z$ is Lipschitz continuous from $\text{supp}(\mathcal{L}_Z)$ (which is closed inside \mathbb{R}^{d_Z} , hence a Polish space) to $(\mathcal{P}_p(\mathbb{R}^{d_X+d_Y}), W_p)$, it is automatically Borel measurable. Lemma 7 then indicates that

$$\begin{aligned} W_p^p \left(\int_{V_j} \mathcal{L}_{(X,Y)|Z=z} d\mathcal{L}_Z(z), \mathcal{L}_{(X,Y)|Z=z_0} \right) &\leq \int_{V_j} W_p^p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z_0}) d\mathcal{L}_Z(z) \\ &\leq \int_{V_j} \varepsilon^p d\mathcal{L}_Z(z) \\ &= \varepsilon^p. \end{aligned}$$

Since $\int_{V_j} \mathcal{L}_{(X,Y)|Z=z} d\mathcal{L}_Z(z) = \mathcal{L}_{(X,Y)|Z \in V_j}$, the proposition is proved. \square

We also need an analogous proposition, but for the measures $\mathcal{L}_{X|Z=z_0} \otimes \mathcal{L}_{Y|Z=z_0}$ and $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$.

Proposition 9. *Suppose that $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$ are L_X - and L_Y -Lipschitz (resp.) with respect to W_p . Suppose that $\{V_j\}_{j=1}^J$ is a measurable partition of the support of \mathcal{L}_Z , and that for all bins V_j ,*

$$\text{diam}(V_j) \leq \frac{\varepsilon}{(2^{p-1} (L_X^p + L_Y^p))^{1/p}}.$$

Then

$$W_p(\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}, \mathcal{L}_{X|Z=z_0} \otimes \mathcal{L}_{Y|Z=z_0}) \leq \varepsilon.$$

Proof. The argument is similar to that for the previous proposition, but with a couple of additional complications.

Let γ_X be an optimal transport plan between $\mathcal{L}_{X|Z \in V_j}$ and $\mathcal{L}_{X|Z=z_0}$, and similarly for γ_Y . Then, $\gamma_X \otimes \gamma_Y$ is a transport plan between $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$ and $\mathcal{L}_{X|Z=z_0} \otimes \mathcal{L}_{Y|Z=z_0}$; consequently,

$$\begin{aligned} W_p^p(\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}, \mathcal{L}_{X|Z=z_0} \otimes \mathcal{L}_{Y|Z=z_0}) &\leq \int_{(\mathbb{R}^{d_X+d_Y})^2} \|(x, y) - (x_0, y_0)\|^p d(\gamma_X \otimes \gamma_Y)((x, y), (x_0, y_0)) \\ &= \int_{(\mathbb{R}^{d_X})^2} \int_{(\mathbb{R}^{d_Y})^2} \|(x, y) - (x_0, y_0)\|^p d\gamma_X(x, x_0) d\gamma_Y(y, y_0) \\ &= \int_{(\mathbb{R}^{d_X})^2} \int_{(\mathbb{R}^{d_Y})^2} \left(\sqrt{(x-x_0)^2 + (y-y_0)^2} \right)^p d\gamma_X(x, x_0) d\gamma_Y(y, y_0). \end{aligned}$$

Now, since $(x-x_0)^2 + (y-y_0)^2 \leq (x-x_0)^2 + 2|x-x_0||y-y_0| + (y-y_0)^2$, it follows that

$$\sqrt{(x-x_0)^2 + (y-y_0)^2} \leq \sqrt{(|x-x_0| + |y-y_0|)^2}$$

and so

$$\int_{(\mathbb{R}^{d_X})^2} \int_{(\mathbb{R}^{d_Y})^2} \left(\sqrt{(x-x_0)^2 + (y-y_0)^2} \right)^p d\gamma_X d\gamma_Y \leq \int_{(\mathbb{R}^{d_X})^2} \int_{(\mathbb{R}^{d_Y})^2} (|x-x_0| + |y-y_0|)^p d\gamma_X d\gamma_Y.$$

In turn, $(|x-x_0| + |y-y_0|)^p \leq 2^{p-1} (|x-x_0|^p + |y-y_0|^p)$, so

$$\begin{aligned} \int_{(\mathbb{R}^{d_X})^2} \int_{(\mathbb{R}^{d_Y})^2} (|x-x_0| + |y-y_0|)^p d\gamma_X d\gamma_Y &\leq \int_{(\mathbb{R}^{d_X})^2} \int_{(\mathbb{R}^{d_Y})^2} 2^{p-1} (|x-x_0|^p + |y-y_0|^p) d\gamma_X d\gamma_Y \\ &= 2^{p-1} \left(\int_{(\mathbb{R}^{d_X})^2} |x-x_0|^p d\gamma_X + \int_{(\mathbb{R}^{d_Y})^2} |y-y_0|^p d\gamma_Y \right) \\ &= 2^{p-1} (W_p^p(\mathcal{L}_{X|Z \in V_j}, \mathcal{L}_{X|Z=z_0}) + W_p^p(\mathcal{L}_{Y|Z \in V_j}, \mathcal{L}_{Y|Z=z_0})). \end{aligned}$$

Since both $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$ are Lipschitz continuous (with respective Lipschitz constants L_X and L_Y), we know that for $z, z_0 \in V_j$,

$$W_p(\mathcal{L}_{X|Z=z}, \mathcal{L}_{X|Z=z_0}) \leq L_X |z - z_0| \leq \frac{L_X}{(2^{p-1} (L_X^p + L_Y^p))^{1/p}} \varepsilon$$

and likewise

$$W_p(\mathcal{L}_{Y|Z=z}, \mathcal{L}_{Y|Z=z_0}) \leq L_Y |z - z_0| \leq \frac{L_Y}{(2^{p-1} (L_X^p + L_Y^p))^{1/p}} \varepsilon.$$

Therefore, we can invoke Lemma 7 with $\mu_z = \mathcal{L}_{X|Z=z}$, $\nu = \mathcal{L}_{X|Z=z_0}$, and conclude, as in the proof of the previous proposition, that

$$\begin{aligned} W_p^p(\mathcal{L}_{X|Z \in V_j}, \mathcal{L}_{X|Z=z_0}) &\leq \int_{V_j} W_p^p(\mathcal{L}_{X|Z=z}, \mathcal{L}_{X|Z=z_0}) d\mathcal{L}_Z(z) \\ &\leq \int_{V_j} \left(\frac{L_X}{(2^{p-1} (L_X^p + L_Y^p))^{1/p}} \varepsilon \right)^p d\mathcal{L}_Z(z) \\ &= \left(\frac{L_X}{(2^{p-1} (L_X^p + L_Y^p))^{1/p}} \varepsilon \right)^p \end{aligned}$$

and, by identical reasoning,

$$W_p^p(\mathcal{L}_{Y|Z \in V_j}, \mathcal{L}_{Y|Z=z_0}) \leq \left(\frac{L_Y}{(2^{p-1} (L_X^p + L_Y^p))^{1/p}} \varepsilon \right)^p.$$

Therefore,

$$\begin{aligned} W_p^p(\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}, \mathcal{L}_{X|Z=z_0} \otimes \mathcal{L}_{Y|Z=z_0}) &\leq 2^{p-1} \left(\frac{L_X}{(2^{p-1} (L_X^p + L_Y^p))^{1/p}} \varepsilon \right)^p \\ &\quad + 2^{p-1} \left(\frac{L_Y}{(2^{p-1} (L_X^p + L_Y^p))^{1/p}} \varepsilon \right)^p \\ &= \varepsilon^p \end{aligned}$$

as desired. \square

Remark. Using essentially the same computation as in the proof above, one can show that if $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$ are L_X - and L_Y -Lipschitz (resp.) with respect to W_p , then $z \mapsto \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}$ is Lipschitz with constant $(2^{p-1}(L_X^p + L_Y^p))^{1/p}$. Of course, under the null hypothesis, $\mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z} = \mathcal{L}_{(X,Y)|Z=z}$ for all z , so in particular it must be the case that $(2^{p-1}(L_X^p + L_Y^p))^{1/p} = L_{XY}$. Under the alternative, however, there is no *a priori* relationship between L_X , L_Y , and L_{XY} .

Corollary 10. *Suppose that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ be L_{XY} -Lipschitz with respect to W_p , and that $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$ are L_X - and L_Y -Lipschitz (resp.) with respect to W_p . Suppose that $\{V_j\}_{j=1}^J$ is a measurable partition of the support of \mathcal{L}_Z , and that for every bin V_j ,*

$$\text{diam}(V_j) \leq \frac{\varepsilon}{4 \max \left\{ L_{XY}, (2^{p-1}(L_X^p + L_Y^p))^{1/p} \right\}}.$$

Then,

$$\left| W_p \left(\mathcal{L}_{(X,Y)|Z=z_0}, \mathcal{L}_{X|Z=z_0} \otimes \mathcal{L}_{Y|Z=z_0} \right) - W_p \left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j} \right) \right| \leq \frac{\varepsilon}{2}.$$

Proof. This follows directly from Propositions 8 and 9, together with the triangle inequality. \square

The significance of this corollary is the following. If, for every bin V_j , we perform a W_p two-sample test on the measures $\mathcal{L}_{(X,Y)|Z \in V_j}$ and $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$, then (under our running assumptions) this amounts to a W_p two-sample test on the measures $\mathcal{L}_{(X,Y)|Z=z_0}$ and $\mathcal{L}_{X|Z=z_0} \otimes \mathcal{L}_{Y|Z=z_0}$, for every $z_0 \in V_j$. Consequently, under our assumptions, “ W_p conditional independence testing” is reducible to a large, but finite, number of W_p two-sample tests.

To put this more formally, we now state a “metatheorem”, indicating how to aggregate Wasserstein two-sample tests on the bins V_j , with given Type I and Type II error, into a conditional independence test.

Theorem 11. *Let $X : \Omega \rightarrow \mathbb{R}^{d_X}$, $Y : \Omega \rightarrow \mathbb{R}^{d_Y}$, and $Z : \Omega \rightarrow \mathbb{R}^{d_Z}$ be random variables. Suppose that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ is L_{XY} -Lipschitz with respect to W_p , and that $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$ are L_X - and L_Y -Lipschitz (resp.) with respect to W_p . Suppose the bins $\{V_j\}_{j=1}^J$ form a measurable partition of the support of \mathcal{L}_Z , and that for every bin V_j ,*

$$\text{diam}(V_j) \leq \frac{\varepsilon}{4 \max \left\{ L_{XY}, (2^{p-1}(L_X^p + L_Y^p))^{1/p} \right\}}.$$

Let $W_p \left(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j} \right)$ denote the plug-in estimator for $W_p \left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j} \right)$, and let n_j denote the number of data points drawn from each of $\mathcal{L}_{(X,Y)|Z \in V_j}$ and $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$.

Then:

- (1) *(Control of Type I error) Suppose that for a given $j \in \{1, \dots, J\}$, n_j is sufficiently large that for $p_j : \mathbb{N} \rightarrow [0, 1]$,*

$$\mathbb{P} \left(W_p \left(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j} \right) \geq W_p \left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j} \right) + \frac{\varepsilon}{2} \right) \leq p_j(n_j)$$

Then for all $z \in V_j$,

$$\mathbb{P} \left(W_p \left(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j} \right) \geq W_p \left(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z} \right) + \varepsilon \right) \leq p_j(n_j)$$

and: under the null hypothesis that for all z in the support of \mathcal{L}_Z , $W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}) = 0$, it holds that

$$\mathbb{P}\left(\exists j \in \{1, \dots, J\}, W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) \geq \varepsilon\right) \leq \sum_{j=1}^J p_j(n_j).$$

(2) (Control of Type II error) Suppose that for a given $j \in \{1, \dots, J\}$, n_j is sufficiently large that for $\varepsilon, \delta_j > 0$ and $\alpha_j : \mathbb{R}_+ \rightarrow [0, 1]$,

$$\begin{aligned} \mathbb{P}\left(W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) < W_p\left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}\right) - \delta_j + \frac{\varepsilon}{2}\right) \\ \leq \alpha_j\left(\delta_j - \frac{\varepsilon}{2}, n_j\right). \end{aligned}$$

Then for all $z \in V_j$,

$$\begin{aligned} \mathbb{P}\left(W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) < W_p\left(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}\right) - \delta_j + \varepsilon\right) \\ \leq \alpha_j\left(\delta_j - \frac{\varepsilon}{2}, n_j\right) \end{aligned}$$

and: under the alternative hypothesis that for every $j \in \{1, \dots, J\}$, there exists a $z \in V_j$ such that $W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}) \geq \delta_j + \varepsilon$, it holds that

$$\mathbb{P}\left(\forall j \in \{1, \dots, J\}, W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) < \varepsilon\right) \leq \prod_{j=1}^J \alpha_j\left(\delta_j - \frac{\varepsilon}{2}, n_j\right).$$

Proof. (1) This is immediate from Corollary 10, together with a Bonferroni correction.

(2) This is also an immediate consequence of Corollary 10, together with the fact that the bins V_j are disjoint, and hence the events

$$W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) < W_p\left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}\right) - \delta_j + \frac{\varepsilon}{2},$$

for $j \in \{1, \dots, J\}$, are independent. Explicitly, independence implies that

$$\begin{aligned} & \mathbb{P}\left(\forall j \in \{1, \dots, J\}, W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) < W_p\left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}\right) - \delta_j + \frac{\varepsilon}{2}\right) \\ &= \prod_{j=1}^J \mathbb{P}\left(W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) < W_p\left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}\right) - \delta_j + \frac{\varepsilon}{2}\right) \\ &\leq \prod_{j=1}^J \alpha_j\left(\delta_j - \frac{\varepsilon}{2}, n_j\right) \end{aligned}$$

and, under the stipulated alternative, namely $W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}) \geq \delta_j + \varepsilon$ for some $z \in V_j$, we have by Corollary 10 that $W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}) \geq \delta_j + \frac{\varepsilon}{2}$, and hence

$$\begin{aligned} & \mathbb{P}\left(\forall j \in \{1, \dots, J\}, W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) < \varepsilon\right) \\ &\leq \mathbb{P}\left(\forall j \in \{1, \dots, J\}, W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}\right) < W_p\left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}\right) - \delta_j + \frac{\varepsilon}{2}\right). \end{aligned}$$

Hence

$$\mathbb{P}\left(\forall j \in \{1, \dots, J\}, W_p\left(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j}\right) < \varepsilon\right) \leq \prod_{j=1}^J \alpha_j\left(\delta_j - \frac{\varepsilon}{2}, n_j\right)$$

as desired. \square

Remark. Note that J , the number of bins partitioning $\text{supp}(\mathcal{L}_Z)$, is on the order of the $\varepsilon/(4 \max\{L_{XY}, (2^{p-1}(L_X^p + L_Y^p))^{1/p}\})$ -covering number of $\text{supp}(\mathcal{L}_Z)$, and so if $\text{supp}(\mathcal{L}_Z)$ is geometrically “simple” (i.e. has small isoperimetric ratio compared to a d_Z -ball), then J is on the order of

$$\left(8 \cdot \text{diam}(\text{supp}(\mathcal{L}_Z)) \max\left\{L_{XY}, (2^{p-1}(L_X^p + L_Y^p))^{1/p}\right\} \sqrt{d}/\varepsilon\right)^d.$$

Example 12. Let us see how the preceding theorem caches out in a more specific instance, where we have made more specific regularity assumptions on the random variables X , Y , and Z , and the distributions $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ and $z \mapsto \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}$. Whatever the case, if we are to employ an expectation bound or concentration inequality (or analogous substitutes) that require some additional regularity assumptions, we must check that those regularity properties are preserved when we pass from $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ and $z \mapsto \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}$ to the binned distributions $j \mapsto \mathcal{L}_{(X,Y)|Z \in V_j}$ and $j \mapsto \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$ (since it is more natural to impose regularity assumptions on the (unbinned) conditional distributions $\mathcal{L}_{(X,Y)|Z=z}$, $\mathcal{L}_{X|Z=z}$, and $\mathcal{L}_{Y|Z=z}$).

Here, consider the case where X and Y are both compactly supported inside a ball of radius D inside $\mathbb{R}^{d_X} \times \mathbb{R}^{d_Y}$, and assume that $d_X + d_Y \geq 3$. Therefore we have access to the expectation bound of Dereich et al. stated in Theorem 4, and the concentration inequality of Weed & Bach stated in Proposition 2; let us check that the requisite assumptions for these two bounds are stable under binning. (For discussion of the case where $d_X + d_Y = 2$, we refer the reader to Example 15 Appendix B.)

Indeed, recall the expectation bound from Theorem 4, namely, that if $p \in [1, d/2)$ and $q > dp/(d-p)$, then there exists a constant $\kappa_{p,q,d}$ depending only on p, q , and d , such that

$$\mathbb{E}[W_p^p(\mu, \mu_n)]^{1/p} \leq \kappa_{p,q,d} \left[\int_{\mathbb{R}^d} \|x\|^q d\mu(x) \right]^{1/q} n^{-1/d}.$$

So suppose that there is some uniform M_{XY} such that

$$\left[\int_{\mathbb{R}^{d_X+d_Y}} \|(x, y)\|^q d\mu(x, y) \right]^{1/q} \leq M_{XY}$$

whenever $\mu = \mathcal{L}_{(X,Y)|Z=z}$ and $z \in \text{supp}(\mathcal{L}_Z)$. We claim that the same moment bound then holds for $\mathcal{L}_{(X,Y)|z \in V_j}$ also, at least provided that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ is continuous w.r.t. W_q (in addition to being L_{XY} -Lipschitz w.r.t. W_p ; since convergence in W_q is equivalent to weak convergence plus convergence of q th moments, it suffices, for instance, to assume yet another uniform moment bound, for any $q' > q$: see [Bil95, Corollary following Theorem 25.12]). Indeed,

$$\int_{\mathbb{R}^{d_X+d_Y}} \|(x, y)\|^q d\mu(x, y) = W_q^q(\mu, \delta_0)$$

and so, by Lemma 7,

$$\begin{aligned} \int_{\mathbb{R}^{d_X+d_Y}} \|(x, y)\|^q d\mathcal{L}_{(X,Y)|Z \in V_j}(x, y) &= W_q^q(\mathcal{L}_{(X,Y)|z \in V_j}, \delta_0) \\ &\leq \int_{V_j} W_q^q(\mathcal{L}_{(X,Y)|Z=z}, \delta_0) d\mathcal{L}_Z(z) \\ &\leq \int_{V_j} M_{XY}^q d\mathcal{L}_Z(z) \\ &= M_{XY}^q. \end{aligned}$$

Likewise, it is obvious that if, for all $z \in \text{supp}\mathcal{L}_Z$, $\mathcal{L}_{(X,Y)|Z=z}$ is supported inside some ball $B(0, D)$, then the same is true for each $\mathcal{L}_{(X,Y)|Z \in V_j}$. And identical reasoning applies to both $\mathcal{L}_{X|Z=z}$ and $\mathcal{L}_{Y|Z=z}$. Finally, observe that

$$\int_{\mathbb{R}^{d_X+d_Y}} \|(x, y)\|^q d\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}(x, y) \leq 2^{q-1} \left(\int_{\mathbb{R}^{d_X}} \|x\|^q d\mathcal{L}_{X|Z \in V_j}(x) + \int_{\mathbb{R}^{d_Y}} \|y\|^q d\mathcal{L}_{Y|Z \in V_j}(y) \right)$$

so, if $\mathcal{L}_{X|Z \in V_j}$ has q th moment bound M_X and $\mathcal{L}_{Y|Z \in V_j}$ has q th moment bound M_Y , it follows that $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$ has q th moment bound $(2^{q-1}(M_X^q + M_Y^q))^{1/q}$.

Thus, if we assume that $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ and $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$ are:

- (1) L_{XY} -, L_X -, and L_Y -Lipschitz with respect to W_p for some $p \in [1, \infty)$,
- (2) continuous with respect to W_q for some $q > (d_X + d_Y)p / (d_X + d_Y - p)$, and
- (3) have q th moments uniformly bounded by M_{XY} and M_X and M_Y respectively, and
- (4) have bounded support¹ with radius D ,

Then we can use both the expectation bound from Dereich et al., as well as the concentration inequality from Weed & Bach, in the case of the empirical measures of $\mathcal{L}_{(X,Y)|Z \in V_j}$ and $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$.

Indeed, we saw in the previous section that under the null hypothesis, if μ_n and μ'_n are both samples from the same measure μ , and n is sufficiently large that $(\kappa_{p,q,d} [\int_{\mathbb{R}^d} \|x\|^q d\mu(x)]^{1/q} n^{-1/d})^p \leq \frac{1}{2} (\frac{\varepsilon}{4})^p$, then combining the expectation bound from Theorem 4 and the concentration inequality from Proposition 2 results in the estimate

$$\begin{aligned} \mathbb{P} \left[W_p(\mu_n, \mu'_n) \geq \frac{\varepsilon}{2} \right] &\leq \mathbb{P} \left[W_p(\mu_n, \mu) \geq \frac{\varepsilon}{4} \right] + \mathbb{P} \left[W_p(\mu'_n, \mu) \geq \frac{\varepsilon}{4} \right] \\ &\leq 2 \exp \left(-\frac{2n\varepsilon^{2p}}{4^{2p+1}D^{2p}} \right). \end{aligned}$$

So, by the preceding theorem, it holds under the null that if for all $j \in \{1, \dots, J\}$, n_j is sufficiently large that

$$\left(\kappa_{p,q,d} \left[\int_{\mathbb{R}^{d_X+d_Y}} \|(x, y)\|^q d\mathcal{L}_{(X,Y)|Z \in V_j}(x, y) \right]^{1/q} n_j^{-1/d} \right)^p \leq \left(\kappa_{p,q,d} M_{XY} n_j^{-1/d} \right)^p \leq \frac{1}{2} \left(\frac{\varepsilon}{4} \right)^p,$$

¹Note: this implies automatically that (3) holds, but we run the analysis with M_X , M_Y , M_{XY} given more generally, in case one has access to tighter moment bounds than those implied automatically from the diameter of the support.

it holds that

$$\mathbb{P} \left[(\exists j \in \{1, \dots, J\}) W_p(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j}) \geq \varepsilon \right] \leq \sum_{j=1}^J 2 \exp \left(-\frac{2n_j \varepsilon^{2p}}{4^{2p+1} D^{2p}} \right)$$

which is precisely a bound on the Type I error.

Let us now turn to the Type II error. As discussed in the previous section, we reason as follows. Our alternative hypothesis is that for each $j \in \{1, \dots, J\}$, there exists a $z \in V_j$ such that $W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}) \geq \delta_j + \varepsilon$; hence $W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}) \geq \delta_j + \frac{\varepsilon}{2}$. If

$$W_p(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j}) < W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}) - \delta_j + \frac{\varepsilon}{2}$$

then it must be the case that

$$W_p(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \mathcal{L}_{(X,Y)|Z \in V_j}) \geq \frac{\delta_j}{2} - \frac{\varepsilon}{4} \text{ or } W_p(\widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}) \geq \frac{\delta_j}{2} - \frac{\varepsilon}{4}.$$

In the first case, using $t = \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p$ in Proposition 2, we see that

$$\mathbb{P} \left[W_p(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \mathcal{L}_{(X,Y)|Z \in V_j}) \geq \frac{\delta_j}{2} - \frac{\varepsilon}{4} \right] \leq \exp \left(-\frac{2n_j (\delta_j - \frac{\varepsilon}{2})^{2p}}{4^{2p+1} D^{2p}} \right)$$

provided that n_j is sufficiently large that

$$\left(\kappa_{p,q,d} \left[\int_{\mathbb{R}^d \times \mathbb{R}^d} \|(x,y)\|^q d\mathcal{L}_{(X,Y)|Z \in V_j}(x,y) \right]^{1/q} n^{-1/d} \right)^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p$$

or, in terms of the moment bound M_{XY} , that

$$\left(\kappa_{p,q,d} M_{XY} n^{-1/d} \right)^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p.$$

In the second case, similarly,

$$\mathbb{P} \left[W_p(\widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}) \geq \frac{\delta_j}{2} - \frac{\varepsilon}{4} \right] \leq \exp \left(-\frac{2n_j (\delta_j - \frac{\varepsilon}{2})^{2p}}{4^{2p+1} D^{2p}} \right)$$

provided that n_j is sufficiently large that

$$\left(\kappa_{p,q,d} \left[\int_{\mathbb{R}^d \times \mathbb{R}^d} \|(x,y)\|^q d\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}(x,y) \right]^{1/q} n^{-1/d} \right)^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p$$

or, in terms of our moment bound, that

$$\left(\kappa_{p,q,d} (2^{q-1} (M_X^q + M_Y^q))^{1/q} n^{-1/d} \right)^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p.$$

Therefore,

$$\begin{aligned} \mathbb{P} \left[W_p(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j}) \leq W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}) - \delta_j + \frac{\varepsilon}{2} \right] \\ \leq 2 \exp \left(-\frac{2n_j (\delta_j - \frac{\varepsilon}{2})^{2p}}{4^{2p+1} D^{2p}} \right) \end{aligned}$$

provided that n_j is sufficiently large that

$$\left(\max \left\{ M_{XY}, (2^{q-1}(M_X^q + M_Y^q))^{1/q} \right\} \kappa_{p,q,d} n_j^{-1/d} \right)^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p.$$

Combining these estimates across bins, we conclude that

$$\mathbb{P} \left(\forall j \in \{1, \dots, J\}, W_p \left(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j} \right) < \varepsilon \right) \leq \prod_{j=1}^J 2 \exp \left(- \frac{2n_j (\delta_j - \frac{\varepsilon}{2})^{2p}}{4^{2p+1} D^{2p}} \right)$$

provided that for all $j \in \{1, \dots, J\}$, $\left(\max \left\{ M_{XY}, (2^{q-1}(M_X^q + M_Y^q))^{1/q} \right\} \kappa_{p,q,d} n_j^{-1/d} \right)^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p$.

Remark. Using the concentration inequality from Theorem 3, it is also possible to extend the analysis from the previous example to the case where \mathcal{L}_X and \mathcal{L}_Y have unbounded support (but still satisfy a higher moment bound).

3. PLUG-IN ESTIMATION OF THE LIPSCHITZ CONSTANT

We have seen that the role played by the Lipschitz constants L_X , L_Y , and L_{XY} is that they specify a bin size which is sufficient to introduce only a small discretization error. In specific applications, there may well be domain knowledge that allows us to upper bound the L 's in an *a priori* fashion; however, testing whether a function (with continuous domain) is Lipschitz, or testing for L given the fact that the function is Lipschitz with *some* constant, is not possible in general, at least if one requires any sort of finite-sample guarantees. (To see why this is the case, it suffices to consider some standard pathological function from real analysis, such as the Weierstrass function.)

Remark. In a situation similar to the one from Example 6 above, some *a priori*/expert knowledge about L may come from the functional form of a causal model under investigation, e.g. if one considers an additive noise model as we did there. Additionally, it may be possible to deduce an upper bound on L analytically, for instance if the conditional distributions are known to satisfy a suitable *functional inequality* (e.g. of Poincaré or log-Sobolev type) — for such a result, we refer the reader to [DM20, Theorem 2.1].

However, it is still of interest to provide a consistent *estimator* of L , should no guess based on domain knowledge be available.

To that end, we note that it *is* possible to compute the Lipschitz constant (w.r.t. the W_p metric) of the *discrete* function

$$V_j \mapsto \widehat{\mathcal{L}}_{(X,Y)|Z \in V_j} := \frac{1}{|\{i : z_i \in V_j\}|} \sum_{i: z_i \in V_j} \delta_{(x_i, y_i)}$$

which, given a bin, returns the empirical conditional distribution for that bin. Briefly, this is because the space of bins V_j can be viewed as a discrete space, and in such a setting one can simply compute the Lipschitz constant by brute force. (Note, however, that there is work on *efficient* estimation of Lipschitz constants for discrete functions, for instance [AJMR16].)

Therefore, we propose using the “plug-in estimator” \widehat{L}_{XY} for the Lipschitz constant L_{XY} , namely

$$\widehat{L}_{XY} := \text{Lip} \left(V_j \mapsto \widehat{\mathcal{L}}_{(X,Y)|Z \in V_j} \right)$$

where $\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}$ resides in a Wasserstein space $W_p(\mathbb{R}^{d_X + d_Y})$, and the distance between two bins V_j and $V_{j'}$ is given by the Euclidean distance between their centroids. In exactly the same fashion,

we also define the plug-in estimators \widehat{L}_X and \widehat{L}_Y for L_X and L_Y , respectively, by

$$\begin{aligned}\widehat{L}_X &:= \text{Lip}\left(V_j \mapsto \widehat{\mathcal{L}_{X|Z \in V_j}}\right); \\ \widehat{L}_Y &:= \text{Lip}\left(V_j \mapsto \widehat{\mathcal{L}_{Y|Z \in V_j}}\right).\end{aligned}$$

We now claim the following.

Proposition 13. \widehat{L}_{XY} is a consistent estimator, in the sense that if $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ is L_{XY} -Lipschitz, then almost surely, $\widehat{L}_{XY} \rightarrow L_{XY}$ as both the number of data points goes to infinity, and the diameter of the bins goes to zero. The same holds true for \widehat{L}_X and \widehat{L}_Y .

Proof. The proof is identical for all three estimators; we therefore proceed only for the estimator \widehat{L}_{XY} .

Previously, in Proposition 8, we have seen that if $\text{diam}(V_j) \leq \varepsilon/L_{XY}$, then

$$W_p\left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{(X,Y)|Z=z_0}\right) < \varepsilon.$$

At the same time, for n sufficiently large we have that $\mathcal{L}_{(X,Y)|Z \in V_j} \approx_{W_p} \widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}$, with probability going to 1 as n goes to ∞ .

Therefore, we proceed as follows. Pick z_0 and z'_0 , and without loss of generality, take $\varepsilon < \frac{L_{XY}}{2}|z_0 - z'_0|$. Under this circumstance, if $z_0 \in V_j$ and $z'_0 \in V_{j'}$, then $V_j \neq V_{j'}$ from the diameter assumption on V_j and $V_{j'}$.

We know that

$$W_p\left(\mathcal{L}_{(X,Y)|Z=z_0}, \mathcal{L}_{(X,Y)|Z=z'_0}\right) \leq L_{XY}|z_0 - z'_0|.$$

At the same time, by the triangle inequality and the binning consistency estimate, we have (still with $z_0 \in V_j$ and $z'_0 \in V_{j'}$) that

$$W_p\left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{(X,Y)|Z \in V_{j'}}\right) \leq W_p\left(\mathcal{L}_{(X,Y)|Z=z_0}, \mathcal{L}_{(X,Y)|Z=z'_0}\right) + 2\varepsilon.$$

Hence in particular,

$$W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{(X,Y)|Z \in V_{j'}}}\right) \leq L_{XY}|z_0 - z'_0| + 2\varepsilon.$$

Using the triangle inequality again, and the fact that $\mathcal{L}_{(X,Y)|Z \in V_j} \approx_{W_p} \widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}$ and $\mathcal{L}_{(X,Y)|Z \in V_{j'}} \approx_{W_p} \widehat{\mathcal{L}_{(X,Y)|Z \in V_{j'}}}$ for n very large (with probability approaching 1 as $n \rightarrow \infty$), we have that (also with probability approaching 1 as $n \rightarrow \infty$)

$$W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{(X,Y)|Z \in V_{j'}}}\right) \lesssim L_{XY}|z_0 - z'_0| + 2\varepsilon.$$

Now, this analysis holds for any z_0 and z'_0 ; in particular we may take z_0 and z'_0 to be the centroids of V_j and $V_{j'}$ (although note this may change ε). In this case, the distance between V_j and $V_{j'}$ that we have specified is none other than $|z_0 - z'_0|$. Taking ε to be much smaller than the sample complexity error, we conclude that (with probability approaching 1 as $n \rightarrow \infty$)

$$W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{(X,Y)|Z \in V_{j'}}}\right) \lesssim L_{XY}|z_0 - z'_0|$$

where the symbol \lesssim conceals only an error term arising from the sample complexity of $\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}$. Therefore, for enough data points and small enough bins, we have asymptotically that (with probability approaching 1 as $n \rightarrow \infty$)

$$\sup_{j,j'} W_p\left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{(X,Y)|Z \in V_{j'}}}\right) \lesssim L_{XY} \cdot \text{dist}(V_j, V_{j'})$$

which means that $\limsup_{n,\varepsilon} \widehat{L}_{XY} \leq L_{XY}$, almost surely.

It remains, therefore, to show that $L_{XY} \leq \widehat{L}_{XY}$. The argument is very similar. Take V_j and $V_{j'}$ to be arbitrary. We know that

$$W_p(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{(X,Y)|Z \in V_{j'}}) \leq \widehat{L}_{XY} \cdot \text{dist}(V_j, V_{j'})$$

simply from the definition of \hat{L} . For a very large number of data points n , we have that $\mathcal{L}_{(X,Y)|Z \in V_j} \approx_{W_p} \widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}$ and $\mathcal{L}_{(X,Y)|Z \in V_{j'}} \approx_{W_p} \widehat{\mathcal{L}}_{(X,Y)|Z \in V_{j'}}$ (with probability approaching 1 as $n \rightarrow \infty$), and so

$$W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{(X,Y)|Z \in V_{j'}}) \lesssim \widehat{L}_{XY} \cdot \text{dist}(V_j, V_{j'}).$$

Here also, the symbol \lesssim conceals only an error term arising from the sample complexity of $\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}$.

Let z_0 and z'_0 be the centroids of V_j and $V_{j'}$. Then (with probability approaching 1 as $n \rightarrow \infty$)

$$W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{(X,Y)|Z \in V_{j'}}) \lesssim \widehat{L}_{XY} |z_0 - z'_0|.$$

From the triangle inequality and the binning consistency estimate, we have that (with probability approaching 1 as $n \rightarrow \infty$)

$$W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z'}) \lesssim \widehat{L}_{XY} |z - z'| + 2\varepsilon \quad \forall z \in V_j, z' \in V_{j'}.$$

Now, note that

$$|z_0 - z|, |z'_0 - z'| < \varepsilon / L_{XY}$$

from the diameter estimate on the bins. Therefore, (with probability approaching 1 as $n \rightarrow \infty$)

$$W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z'}) \lesssim \widehat{L}_{XY} |z - z'| + \left(2 + 2\frac{\widehat{L}_{XY}}{L_{XY}}\right) \varepsilon \quad \forall z \in V_j, z' \in V_{j'}.$$

Quantifying over all bins, we conclude that for all z and z' , (with probability approaching 1 as $n \rightarrow \infty$)

$$W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z'}) \lesssim \widehat{L}_{XY} |z - z'| + \left(2 + 2\frac{\widehat{L}_{XY}}{L_{XY}}\right) \varepsilon.$$

Sending ε to be small compared to the sample complexity error (i.e., take the bins V_j and $V_{j'}$ to be very small, but such that they still contain z and z' respectively), and using the fact that $\limsup_{n,\varepsilon} \widehat{L}_{XY} \leq L_{XY}$ a.s. from above, we deduce that (with probability approaching 1 as $n \rightarrow \infty$)

$$W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z'}) \lesssim \widehat{L}_{XY} |z - z'|.$$

Therefore, (with probability approaching 1 as $n \rightarrow \infty$)

$$\sup_{z,z'} W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{(X,Y)|Z=z'}) \lesssim \widehat{L}_{XY} |z - z'|$$

and hence from the definition of L_{XY} , we have $L_{XY} \leq \liminf_{n,\varepsilon} \widehat{L}_{XY}$ almost surely, as desired. \square

4. DISCUSSION

It should be emphasized that the class of joint distributions (X, Y, Z) for which a conditional independence test, constructed in the manner of Theorem 11, is feasible, is quite general. As we have already discussed, the W_p -Lipschitz continuity we require is a weaker condition than the TV -Lipschitz continuity assumption from [NBW20]; nor do we assume that any of the distributions involved have density with respect to the Lebesgue measure; nor do we assume that the joint distribution is produced by any sort of parametric model. Nonetheless, we mention several possible extensions of the work in this article, stated in approximate order of increasing difficulty.

- (1) *Relaxing the Lipschitz continuity assumption.* One might entertain other quantitative smoothness conditions on the maps $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$, $z \mapsto \mathcal{L}_{X|Z=z}$, and $z \mapsto \mathcal{L}_{Y|Z=z}$, besides Lipschitz continuity, such as Hölder continuity. Indeed, if one takes the Hölder exponent and constant as given, the proofs in Section 2 still go through with only minor adjustments to the epsilon management. Likewise, if one takes the Hölder exponent as given, it is straightforward to adapt the arguments in Section 3 so as to produce a consistent estimator for the Hölder constant.

In principle, more general quantitative smoothness assumptions (such as an explicitly given modulus of continuity) are also feasible; what our analysis ultimately demands is *some* sort of quantitative information about the continuity of the maps $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$, $z \mapsto \mathcal{L}_{X|Z=z}$, and $z \mapsto \mathcal{L}_{Y|Z=z}$, so that we can appropriately fix the diameter of the bins.

- (2) *Allowing the support of \mathcal{L}_Z to be unbounded.* The restriction of the results in Section 2 to the case where \mathcal{L}_Z has bounded support inside \mathbb{R}^{dz} excludes many natural situations (take for instance Z to be a Gaussian random variable!). We propose two possible ways to relax this assumption and work with non-compactly supported \mathcal{L}_Z , one of which amounts to “moving the goalposts” and the other of which amounts to “adding a much stronger assumption somewhere else”.

For the first, observe that since \mathcal{L}_Z is automatically tight, for every $\varepsilon > 0$ there exists a compact set K_ε such that $\mathcal{L}_Z(\mathbb{R}^{dz} \setminus K_\varepsilon) < \varepsilon$. Therefore, while it is not possible to form a measurable partition of $\text{supp}(\mathcal{L}_Z)$ with a finite number of bins and where the bins all have uniformly diameter (as is required by Theorem 11), what *is* possible is to form such a partition on K_ε . Then, Theorem 11 indicates how to test the conditional independence of X and Y given Z , *conditional on* $Z \in K_\varepsilon$. This does *not* allow us to test whether $\mathcal{L}_{(X,Y)|Z=z}$ is equal to $\mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}$ for *every* nonzero z , but rather for all z lying in a set K_ε where $\mathbb{P}(Z \in K_\varepsilon) \geq 1 - \varepsilon$. Call this “ $(1 - \varepsilon)$ -conditional independence testing”; this discussion shows that, *if* we are able to come up with an explicit such K_ε for Z , and X , Y , and Z satisfy the other assumptions of Theorem 11, then $(1 - \varepsilon)$ -conditional independence testing is feasible.

As for the second: suppose that, instead of $z \mapsto \mathcal{L}_{(X,Y)|Z=z}$ and $z \mapsto \mathcal{L}_{X|Z=z}$ and $z \mapsto \mathcal{L}_{Y|Z=z}$ merely being W_p -Lipschitz maps, we require that these maps moreover have a local Lipschitz constant which quickly goes to zero as $|z| \rightarrow \infty$. In this situation, it becomes permissible for the bins partitioning \mathcal{L}_Z to have larger and larger diameter as $|z| \rightarrow \infty$; for bins $z \in V_j$ where all z are “very far” from 0, it is even permissible for the bin to be unbounded towards infinity, and still have small discretization error. We are not aware of an application area in conditional independence testing where such an “asymptotically very W_p smooth” assumption would be natural, but from from an *analytical* standpoint

such an assumption is sufficient for W_p conditional independence testing to be feasible with unbounded \mathcal{L}_Z .

- (3) *Data-dependent bins.* In Section 2, we assumed that the support of \mathcal{L}_Z was first partitioned into bins, and then samples are drawn from the joint distribution (X, Y, Z) . In this setting, Theorem 11 then indicates how many data points must lie in each bin in order to have global control of the Type I & II error for the conditional independence test. Or (what is much the same thing), the bins are produced *independently* from the data.

In a target application where data is plentiful and the bottleneck is the computational cost of the empirical Wasserstein distance (see discussion in Section 1), this is not too concerning. However, in general it might be desirable to first collect “as many samples as one can”, and *then* partition the space where \mathcal{L}_Z resides in such a way that each bin has “enough” samples, and then seek various finite sample guarantees. In this situation, the set of bins (equivalently, the measurable partition induced by the bins) becomes a random variable which depends on the samples. This *significantly* complicates much of the analysis — see, for instance, the work Canonne et al. [CDKS18], which offers a conditional independence test for random variables which take values in a *finite set*, and which *does* offer a binning scheme where the bins are allocated in a data-dependent fashion. In our (continuous) setting, we leave such analysis to future work.

- (4) *Random variables taking values in spaces other than $\mathbb{R}^{d_X} \times \mathbb{R}^{d_Y} \times \mathbb{R}^{d_Z}$.* We have stipulated that the random variables X , Y , and Z are Euclidean-valued for concreteness. However, the general theory of Wasserstein spaces/optimal transport can be carried out in the rather general setting of complete separable metric spaces [AG13, Vil03], and some of the statistical results in, for instance, [BLG14, WB19] are stated for more general metric spaces. Most tantalizing is the recent result of Lei [Lei20], which offers an expected error bound, similar to the one of Dereich et al. from [DSS13], in an infinite-dimensional setting. (However, [Lei20] does not provide explicit constants, which our statistical applications would certainly require.)

Unfortunately, in (say) an infinite-dimensional Banach space one runs into the difficulty that (for say $\varepsilon < \frac{1}{2}$) the ε -covering number of the unit ball is infinite, so even if \mathcal{L}_Z has bounded support, binning the support of \mathcal{L}_Z seems challenging! (On the other hand, if we assume that \mathcal{L}_Z has *compact* support, then this essentially means that \mathcal{L}_Z resides in a finite-dimensional subspace, and the putative infinite-dimensional setting trivializes.) One possible strategy is to exploit the *tightness* of (Radon) probability measures — that is, even in an infinite dimensional space, if \mathcal{L}_Z is Radon we know that for every $\varepsilon > 0$, there exists a compact K_ε with $\mathcal{L}_Z(K_\varepsilon) > 1 - \varepsilon$. If, moreover, modeling assumptions tell us that Z must e.g. be concentrated in some fashion that actually *tells us* a compact set which serves as a K_ε , it is conceivable that, with a suitable infinite-dimensional concentration inequality, and a more explicit error estimate resembling the one from [Lei20], that one could perform “ $(1 - \varepsilon)$ -conditional independence testing” as per (2). However we expect that the requisite analysis to construct such a test would be challenging.

ACKNOWLEDGMENTS

The author thanks Tudor Manole, Dejan Slepčev, Larry Wasserman, and Kun Zhang for helpful discussions.

REFERENCES

- [AG13] Luigi Ambrosio and Nicola Gigli, *A user's guide to optimal transport*, Modelling and optimisation of flows on networks, Springer, 2013, pp. 1–155.
- [AG19] Luigi Ambrosio and Federico Glaudo, *Finer estimates on the 2-dimensional matching problem*, Journal de l'École polytechnique — Mathématiques **6** (2019), 737–765 (en).
- [AJMR16] Pranjali Awasthi, Madhav Jha, Marco Molinaro, and Sofya Raskhodnikova, *Testing lipschitz functions on hypergrid domains*, Algorithmica **74** (2016), no. 3, 1055–1081.
- [BGV07] François Bolley, Arnaud Guillin, and Cédric Villani, *Quantitative concentration inequalities for empirical measures on non-compact spaces*, Probability Theory and Related Fields **137** (2007), no. 3-4, 541–593.
- [Bil95] P. Billingsley, *Probability and measure*, Wiley Series in Probability and Statistics, Wiley, 1995.
- [BLG14] Emmanuel Boissard and Thibaut Le Gouic, *On the mean speed of convergence of empirical and occupation measures in wasserstein distance*, Annales de l'IHP Probabilités et statistiques **50** (2014), no. 2, 539–563.
- [BS78] Dimitri P Bertsekas and Steven E Shreve, *Stochastic optimal control: The discrete-time case*, Academic Press, 1978.
- [CDKS18] Clément L. Canonne, Ilias Diakonikolas, Daniel M. Kane, and Alistair Stewart, *Testing conditional independence of discrete distributions*, 2018 Information Theory and Applications Workshop, ITA 2018, San Diego, CA, USA, February 11-16, 2018, IEEE, 2018, pp. 1–57.
- [Çin11] Erhan Çinlar, *Probability and stochastics*, vol. 261, Springer Science & Business Media, 2011.
- [Com15] Richard Combes, *An extension of mcdiarmid's inequality*, arXiv preprint arXiv:1511.05240 (2015).
- [CRL+20] Lenaïc Chizat, Pierre Roussillon, Flavien Léger, François-Xavier Vialard, and Gabriel Peyré, *Faster wasserstein distance estimation with the sinkhorn divergence*, Neural Information Processing Systems, 2020.
- [Daw79] A Philip Dawid, *Conditional independence in statistical theory*, Journal of the Royal Statistical Society: Series B (Methodological) **41** (1979), no. 1, 1–31.
- [dBGSL21] Eustasio del Barrio, Alberto González-Sanz, and Jean-Michel Loubes, *Central limit theorems for general transportation costs*, arXiv preprint arXiv:2102.06379 (2021).
- [dBL19] Eustasio del Barrio and Jean-Michel Loubes, *Central limit theorems for empirical transportation cost in general dimension*, The Annals of Probability **47** (2019), no. 2, 926–951.
- [DM20] Emanuele Dolera and Edoardo Mainini, *Lipschitz continuity of probability kernels in the optimal transport framework*, arXiv preprint arXiv:2010.08380 (2020).
- [DSS13] Steffen Dereich, Michael Scheutzow, and Reik Schottstedt, *Constructive quantization: Approximation by empirical measures*, Annales de l'IHP Probabilités et statistiques **49** (2013), no. 4, 1183–1203.
- [FG15] Nicolas Fournier and Arnaud Guillin, *On the rate of convergence in wasserstein distance of the empirical measure*, Probability Theory and Related Fields **162** (2015), no. 3, 707–738.
- [HDPM18] Christina Heinze-Deml, Jonas Peters, and Nicolai Meinshausen, *Invariant causal prediction for nonlinear models*, Journal of Causal Inference **6** (2018), no. 2.
- [Kec12] Alexander Kechris, *Classical descriptive set theory*, vol. 156, Springer Science & Business Media, 2012.
- [Lei20] Jing Lei, *Convergence and concentration of empirical measures under wasserstein distance in unbounded functional spaces*, Bernoulli **26** (2020), no. 1, 767–798.
- [MNW21] Tudor Manole and Jonathan Niles-Weed, *Sharp convergence rates for empirical optimal transport with smooth costs*, arXiv preprint arXiv:2106.13181 (2021).
- [NBW20] Matey Neykov, Sivaraman Balakrishnan, and Larry Wasserman, *Minimax optimal conditional independence testing*, arXiv preprint arXiv:2001.03039 (2020).
- [NWR19] Jonathan Niles-Weed and Philippe Rigollet, *Estimation of wasserstein distances in the spiked transport model*, arXiv preprint arXiv:1909.07513 (2019).
- [PC19] Gabriel Peyré and Marco Cuturi, *Computational optimal transport: With applications to data science*, Foundations and Trends in Machine Learning **11** (2019), no. 5-6, 355–607.
- [Pea09] Judea Pearl, *Causality*, Cambridge university press, 2009.
- [PJS17] Jonas Peters, Dominik Janzing, and Bernhard Schölkopf, *Elements of causal inference: foundations and learning algorithms*, The MIT Press, 2017.
- [RGTC17] Aaditya Ramdas, Nicolás García Trillos, and Marco Cuturi, *On Wasserstein two-sample testing and related families of nonparametric tests*, Entropy **19** (2017), no. 2, 47.
- [San15] Filippo Santambrogio, *Optimal transport for applied mathematicians*, Springer, 2015.

- [SGS⁺01] Peter Spirtes, Clark Glymour, Richard Scheines, et al., *Causation, prediction, and search*, The MIT Press, 2001.
- [SP20] Rajen D Shah and Jonas Peters, *The hardness of conditional independence testing and the generalised covariance measure*, *Annals of Statistics* **48** (2020), no. 3, 1514–1538.
- [Vil03] Cédric Villani, *Topics in optimal transportation*, no. 58, American Mathematical Soc., 2003.
- [Vil08] ———, *Optimal transport: old and new*, vol. 338, Springer Science & Business Media, 2008.
- [WB19] Jonathan Weed and Francis Bach, *Sharp asymptotic and finite-sample rates of convergence of empirical measures in wasserstein distance*, *Bernoulli* **25** (2019), no. 4A, 2620–2648.

APPENDIX A. PROOF OF LEMMA 7

Before proceeding, we require the following technical measure-theoretic result, which we will use to handle the potential non-uniqueness of optimal transport plans in our setting.

Theorem (Arsenin-Kunugui uniformization theorem [Kec12, Theorem 35.46 (ii)]). *Let X and Y be Polish spaces. Let $R \subseteq X \times Y$ be a Borel set. Suppose that for every $x \in X$, $\{y \in Y : (x, y) \in R\}$ is a countable union of compact sets. Then the projection of R onto X is a Borel subset of X , and there exists a uniformization of R , that is, a Borel function $f : \text{proj}_X R \rightarrow Y$ such that $\{x, f(x)\} \in R$ for all $x \in \text{proj}_X R$.*

Proof of Lemma 7. Let $V \subseteq Z$ where Z is some Polish space.

Given any $z \in Z$, define $\text{Opt}(\mu_z, \nu)$ to be the set of optimal transport plans between the probability measures μ_z and ν . Since $\mathcal{P}(U)$ is a Polish space (when equipped with the narrow topology), it is known that for fixed z , $\text{Opt}(\mu_z, \nu)$ is compact inside the Polish space $\mathcal{P}(U \times U)$ [Vil08, Corr. 5.21]. Also this set is always non-empty [San15, Theorem 1.7]. Moreover, it follows from [Vil08, Theorem 5.20] that the set

$$\text{Opt}(\cdot, \nu) := \{\gamma \in \mathcal{P}(U \times U) : \exists \mu \in \mathcal{P}(U), \pi \in \text{Opt}(\mu, \nu)\}$$

is closed inside $\mathcal{P}(U \times U)$, and is therefore itself a Polish space.

Now, let consider the projection map

$$\text{proj}_1 : \text{Opt}(\cdot, \nu) \rightarrow \mathcal{P}(U)$$

which, given a transport plan $\gamma \in \text{Opt}(\mu, \nu)$, returns the first marginal of γ , namely μ . It is not hard to see that proj_1 is narrowly continuous: indeed, consider some bounded continuous function φ on U^2 that only depends on the first coordinate; in this case,

$$\int_{U^2} \varphi \gamma = \int_U \varphi \text{proj}_1 \gamma$$

so in particular, for any narrowly convergent sequence γ_n of transport plans (with limit γ), we have

$$\int \varphi \gamma_n = \int \varphi \text{proj}_1 \gamma_n; \quad \int \varphi \gamma = \int \varphi \text{proj}_1 \gamma$$

which implies that $\int \varphi \text{proj}_1 \gamma_n \rightarrow \int \varphi \text{proj}_1 \gamma$ for any bounded continuous φ on U^2 , as desired.

Since proj_1 is narrowly continuous, it follows that the transpose of the graph of proj_1 , namely the set

$$\text{Gr}^T \text{proj}_1 := \{(\mu, \gamma) \in \mathcal{P}(U) \times \text{Opt}(\cdot, \nu) : \mu = \text{proj}_1 \gamma\},$$

is closed (when both $\text{Opt}(\cdot, \nu)$ and $\mathcal{P}(U)$ are equipped with the narrow topology), and so, in particular, is Borel. Since both $\text{Opt}(\cdot, \nu)$ and $\mathcal{P}(U)$ are Polish, it follows from the Arsenin-Kunugui uniformization theorem that there exists a Borel function OptSelect from $\text{proj}_1 \text{Opt}(\cdot, \nu)$ to $\text{Opt}(\cdot, \nu)$,

whose graph is contained within $\text{Gr}^T \text{proj}_1$. But since $\text{proj}_1 \text{Opt}(\cdot, \nu) = \mathcal{P}(U)$ (since $\text{Opt}(\mu, \nu)$ is non-empty for every $\mu, \nu \in \mathcal{P}(U)$) it follows that the domain of OptSelect is all of $\mathcal{P}(U)$. In other words, $\text{OptSelect}(\mu)$ is a *Borel measurable selection of a transport plan in $\text{Opt}(\mu, \nu)$* .

Now, we have assumed that the function $z \mapsto \mu_z$ is Borel, so by composing $z \mapsto \mu_z$ with OptSelect , we get a Borel measurable mapping

$$z \mapsto \gamma_z : \text{proj}_1 \gamma_z = \mu_z.$$

We also make the following measure-theoretic observation. The weak topology on probability measures has the property that for every open set O , the evaluation map $\mu \mapsto \mu(O)$ is continuous; in particular, this implies (by the monotone class theorem; see also [BS78, Prop. 7.25]) that for every Borel set B , the evaluation map $\mu \mapsto \mu(B)$ is Borel. Consequently, for our measurable selection $z \mapsto \gamma_z$, it holds simultaneously that (1) for every $z \in Z$, γ_z is a probability measure (on the space $U \times U$), and that (2) for fixed Borel set B , the map $z \mapsto \gamma_z(B)$ is Borel (since it is the composition of Borel maps). In other words, our measurable selection γ_z is a (Borel measurable) *stochastic kernel*.

We claim that for any Borel measurable $V \subseteq Z$, the measure

$$\int_V \gamma_z d\lambda(z)$$

is a transport plan with first marginal $\int_V \mu_z d\lambda(z)$. Indeed, take any measurable set $B \subseteq U$; then, since γ_z is a stochastic kernel, we can use the following elementary property of stochastic kernels (see e.g. [Cm11, Thm. I.6.3]):

$$\begin{aligned} \int_{U \times U} 1_{A \times U} d \left(\int_V \gamma_z d\lambda(z) \right) &= \int_V \left(\int_{U \times U} 1_{A \times U} d\gamma_z \right) d\lambda(z) \\ &= \int_V \left(\int_U 1_A d\mu_z \right) d\lambda(z) \\ &= \int_U 1_A d \left(\int_V \mu_z d\lambda(z) \right) \end{aligned}$$

and so $(\int_V \gamma_z d\lambda(z))(A \times U) = (\int_V \mu_z d\lambda(z))(A)$. Moreover, by the same reasoning, we see that the second marginal is ν .

Now, compute as follows. Since $\int_V \gamma_z d\lambda(z)$ is a transport plan between $\int_V \mu_z d\lambda(z)$ and ν , it holds that

$$W_p^p \left(\int_V \mu_z d\lambda(z), \nu \right) \leq \int_{U^2} d(u_1, u_2)^p d \left(\int_V \gamma_z d\lambda(z) \right) (u_1, u_2).$$

Using [Cm11, Thm. I.6.3] again, we deduce that

$$\int_{U^2} d(u_1, u_2)^p d \left(\int_V \gamma_z d\lambda(z) \right) (u_1, u_2) = \int_V \left(\int_{U^2} d(u_1, u_2)^p d\gamma_z(u_1, u_2) \right) d\lambda(z).$$

But note that

$$\int_{U^2} d(u_1, u_2)^p d\gamma_z(u_1, u_2) = W_p^p(\mu_z, \nu).$$

Therefore,

$$\int_V \left(\int_{U^2} d(u_1, u_2)^p d\gamma_z(u_1, u_2) \right) d\lambda(z) = \int_V W_p^p(\mu_z, \nu) d\lambda(z)$$

and so

$$W_p^p \left(\int_V \mu_z d\lambda(z), \nu \right) \leq \int_V W_p^p(\mu_z, \nu) d\lambda(z)$$

as desired. This completes the proof. \square

APPENDIX B. ADDITIONAL RESULTS

B.1. Concentration of $W_p^p(\mu_n, \mu)$. We first prove Proposition 2 in the case where the support of μ has general diameter (as opposed to a diameter of 1, as in [WB19]). We emphasize that this is extremely close to what is already done therein; consider this proof more of a sanity check.

Proof of Proposition 2. Let $c(x, y) = d(x, y)^p$. Kantorovich duality tells us that

$$W_p^p(\mu_n, \mu) = \max_{f \in C_b} \left[\int f d\mu_n - \int f^c d\mu \right]$$

and

$$f(x) = \inf_y [f^c(y) + d(x, y)^p]$$

and

$$f^c(y) = \sup_x [f(x) - d(x, y)^p].$$

Chose $f \in C_b$ so that it attains the maximum for $\int f d\mu_n - \int f^c d\mu$. f is defined only up to a constant, so without loss of generality take $\sup_x f(x) = D^p$. Consequently,

$$f^c(y) \geq \sup_x [D^p - d(x, y)^p] \geq 0$$

and therefore

$$f(x) \geq \inf_y [0 + d(x, y)^p] \geq 0.$$

Thus, we can take the $f \in C_b$ which attains the maximum for $\int f d\mu_n - \int f^c d\mu$ to have the property that $0 \leq f(x) \leq D^p$ for all x in the domain.

Now, let X_1, \dots, X_n be i.i.d. draws from μ , so that $\mu_n = \frac{1}{n} \sum_{i=1}^n X_i$. Define the random function

$$\begin{aligned} w(X_1, \dots, X_n) &:= W_p^p(\mu_n, \mu) \\ &= \sup_{f \in C_b} \left[\int f d\mu_n - \int f^c d\mu \right] \\ &= \sup_{\substack{f \in C_b \\ 0 \leq f \leq D^p}} \left[\frac{1}{n} \sum_{i=1}^n f(X_i) - \int f^c d\mu \right]. \end{aligned}$$

Let $j \in \{1, \dots, n\}$. Observe that given any x_1, \dots, x_n and x'_j ,

$$\begin{aligned} w(x_1, \dots, x_j, \dots, x_n) - w(x_1, \dots, x'_j, \dots, x_n) &\leq \sup_{\substack{f \in C_b \\ 0 \leq f \leq D^p}} \left[\frac{1}{n} \sum_{i=1}^n f(x_i) - \int f^c d\mu \right] \\ &\quad - \sup_{\substack{f' \in C_b \\ 0 \leq f' \leq D^p}} \left[\frac{1}{n} \sum_{\substack{i=1 \\ i \neq j}}^n f'(x_i) + \frac{1}{n} f'(x'_j) - \int (f')^c d\mu \right]. \end{aligned}$$

Let

$$\tilde{f} \in \operatorname{argmax}_{\substack{f \in C_b \\ 0 \leq f \leq D^p}} \left[\frac{1}{n} \sum_{i=1}^n f(x_i) - \int f^c d\mu \right].$$

Note that

$$\left[\frac{1}{n} \sum_{\substack{i=1 \\ i \neq j}}^n \tilde{f}(x_i) + \frac{1}{n} \tilde{f}(x_j) - \int (\tilde{f})^c d\mu \right] \leq \sup_{\substack{f' \in C_b \\ 0 \leq f' \leq D^p}} \left[\frac{1}{n} \sum_{\substack{i=1 \\ i \neq j}}^n f'(x_i) + \frac{1}{n} f'(x_j) - \int (f')^c d\mu \right]$$

and thus

$$\begin{aligned} & \sup_{\substack{f \in C_b \\ 0 \leq f \leq D^p}} \left[\frac{1}{n} \sum_{i=1}^n f(x_i) - \int f^c d\mu \right] - \sup_{\substack{f' \in C_b \\ 0 \leq f' \leq D^p}} \left[\frac{1}{n} \sum_{\substack{i=1 \\ i \neq j}}^n f'(x_i) + \frac{1}{n} f'(x_j) - \int (f')^c d\mu \right] \\ & \leq \left[\frac{1}{n} \sum_{i=1}^n \tilde{f}(x_i) - \int \tilde{f}^c d\mu \right] - \left[\frac{1}{n} \sum_{\substack{i=1 \\ i \neq j}}^n \tilde{f}(x_i) + \frac{1}{n} \tilde{f}(x_j) - \int (\tilde{f})^c d\mu \right] \\ & \leq \frac{1}{n} [\tilde{f}(x_j) - \tilde{f}(x'_j)] \\ & \leq \frac{D^p}{n}. \end{aligned}$$

It therefore follows from McDiarmid's inequality that

$$\mathbb{P} [W_p^p(\mu_n, \mu) \geq \mathbb{E}W_p^p(\mu_n, \mu) + t] \leq \exp \left(-\frac{2nt^2}{D^{2p}} \right)$$

as desired. \square

Let us now adapt the previous proof to the case where μ is not necessarily bounded, using Combes's variant on McDiarmid's inequality, proved in [Com15]. Mostly we are interested in a concentration inequality under the same assumptions as in Theorem 4, namely that $d \geq 3$, $p \in [1, d/2)$, and μ has a q th moment bound for $q > dp/(d-p)$.

Let's recall the statement of Combes's variant on McDiarmid's inequality.

Theorem (Combes [Com15]). *Let $(\mathcal{X}_1, \dots, \mathcal{X}_n)$ be sets and define $\mathcal{X} := \prod_{i=1}^n \mathcal{X}_i$. Let $X := (X_1, \dots, X_n)$ be independent random variables with $X_i \in \mathcal{X}_i$. Let $w : \mathcal{X} \rightarrow \mathbb{R}$, so that $w(X)$ is a random variable.*

Let $\mathcal{Y} \subset \mathcal{X}$. Define $\mathbf{p} := 1 - \mathbb{P}[X \in \mathcal{Y}]$ and $m = \mathbb{E}[w(X) \mid X \in \mathcal{Y}]$. Let $\vec{c} := (c_1, \dots, c_n) \in \mathbb{R}_+^n$ and define $\bar{c} = \sum_{i=1}^n c_i$. Suppose that: for all $j \in \{1, \dots, n\}$, for all $(y, y') \in \mathcal{Y}^2$ with $y_i = y'_i$ for all $i \neq j$, it holds that $|w(y) - w(y')| \leq c_j$. Then, for all $t > 0$, it holds that

$$\mathbb{P}[w(X) \geq m + t] \leq \mathbf{p} + \exp \left(-\frac{2(t - \mathbf{p}\bar{c})_+^2}{\sum_{i=1}^n c_i^2} \right).$$

We will apply Combes's inequality in the case where: (X_1, \dots, X_n) are i.i.d. samples from some probability measure μ (and we denote $\mu_n = \frac{1}{n} \sum_{i=1}^n X_i$), $\mathcal{Y} = B(e, R)^n$ for some "large" radius R and arbitrary "origin" $e \in X$, and $w(X) = W_p^p(\mu_n, \mu)$. In order to apply Combes's inequality, we

must compute c_i , m , and \mathfrak{p} . We do so in the situation where μ has bounded q th moment, where $q > p$.

Proof of Theorem 3. First, recall that every (Radon) probability measure μ on a Polish space is automatically tight - that is, for every $\eta \in (0, 1)$ there exists a compact K_η such that $\mu(K_\eta) > 1 - \eta$. If, moreover, μ has bounded q th moment, we can say more. Indeed, suppose that $\mu(X \setminus B(e, R)) \geq \eta$. Then,

$$\left(\int_X d(e, x)^q d\mu(x) \right)^{1/q} \geq R\eta^{1/q}.$$

It follows that if we know that $(\int_X d(e, x)^q d\mu(x))^{1/q} \leq M_q$, then $R\eta^{1/q} \leq M_q$, in other words, the maximal radius at which $\mu(X \setminus B(e, R)) \geq \eta$ is $M_q\eta^{-1/q}$. Likewise, if we have n i.i.d. draws from μ , we know that for each draw, the probability of landing outside of $B(e, M_q\eta^{-1/q})$ is at most η . It follows that the probability of all the draws landing inside $B(e, M_q\eta^{-1/q})$ — which is the quantity $1 - \mathfrak{p}$ — is at least $(1 - \eta)^n$. Hence $\mathfrak{p} \leq 1 - (1 - \eta)^n$.

To compute m , we will need to upper bound $W_p^p(\mu, \frac{1}{\mu(B(e, M_q\eta^{-1/q}))} \mu \upharpoonright B(e, M_q\eta^{-1/q}))$ in the case where $p < q$. So, compute as follows. We have seen that $\mu(B(e, M_q(\frac{\eta}{2^i})^{-1/q})) \geq 1 - \frac{\eta}{2^i}$. It follows that

$$\mu \left(B(e, M_q \left(\frac{\eta}{2^i} \right)^{-1/q} \setminus B(e, M_q \left(\frac{\eta}{2^{i-1}} \right)^{-1/q}) \right) \leq \mu \left(\mathbb{R}^d \setminus B(e, M_q \left(\frac{\eta}{2^{i-1}} \right)^{-1/q}) \right) < \frac{\eta}{2^{i-1}}.$$

At the same time, regardless of the choice of transport plan between $\frac{1}{\mu(B(e, M_q\eta^{-1/q}))} \mu \upharpoonright B(e, M_q\eta^{-1/q})$ and μ , when transporting the mass in μ which resides in $B(e, M_q(\frac{\eta}{2^i})^{-1/q}) \setminus B(e, M_q(\frac{\eta}{2^{i-1}})^{-1/q})$ onto $\frac{1}{\mu(B(e, M_q\eta^{-1/q}))} \mu \upharpoonright B(e, M_q\eta^{-1/q})$, the distance traveled by this mass is at most $M_q(\frac{\eta}{2^i})^{-1/q} + M_q\eta^{-1/q}$ (for each particle of mass), and the total mass transported from $B(e, M_q(\frac{\eta}{2^i})^{-1/q}) \setminus B(e, M_q(\frac{\eta}{2^{i-1}})^{-1/q})$ is less than $\frac{\eta}{2^{i-1}}$. Consequently, the W_p transportation cost to transport this portion of the mass in μ is less than $\left(M_q(\frac{\eta}{2^i})^{-1/q} + M_q\eta^{-1/q} \right) \left(\frac{\eta}{2^{i-1}} \right)^{1/p}$.

At the same time, since the mass in $\frac{1}{\mu(B(e, M_q\eta^{-1/q}))} \mu \upharpoonright B(e, M_q\eta^{-1/q})$ is everywhere greater than the mass in $\mu \upharpoonright B(e, M_q\eta^{-1/q})$, it follows that we can choose a transport plan from μ to $\frac{1}{\mu(B(e, M_q\eta^{-1/q}))} \mu \upharpoonright B(e, M_q\eta^{-1/q})$ will leave all of μ 's mass within $B(e, M_q\eta^{-1/q})$ unmoved. Therefore, we can upper bound

$$\begin{aligned} W_p(\mu, \frac{1}{\mu(B(e, M_q\eta^{-1/q}))} \mu \upharpoonright B(e, M_q\eta^{-1/q})) &< \sum_{i=1}^{\infty} \left(M_q \left(\frac{\eta}{2^i} \right)^{-1/q} + M_q\eta^{-1/q} \right) \left(\frac{\eta}{2^{i-1}} \right)^{1/p} \\ &= M_q\eta^{1/p-1/q} \sum_{i=1}^{\infty} \left(\frac{1}{(2^i)^{-1/q}(2^{i-1})^{1/p}} + \frac{1}{(2^{i-1})^{1/p}} \right) \\ &= M_q\eta^{1/p-1/q} \sum_{i=1}^{\infty} \left(\frac{2^{1/q}}{(2^{i-1})^{-1/q}(2^{i-1})^{1/p}} + \frac{1}{(2^{i-1})^{1/p}} \right) \\ &= M_q\eta^{1/p-1/q} \left(2^{1/q} \frac{2^{1/p-1/q}}{2^{1/p-1/q} - 1} + \frac{2^{1/p}}{2^{1/p} - 1} \right). \end{aligned}$$

Letting $c_{p,q} := 2^{1/q} \frac{2^{1/p-1/q}}{2^{1/p-1/q}-1} + \frac{2^{1/p}}{2^{1/p}-1}$, we see that

$$W_p\left(\mu, \frac{1}{\mu(B(e, M_q \eta^{-1/q}))} \mu \upharpoonright B(e, M_q \eta^{-1/q})\right) < c_{p,q} M_q \eta^{1/p-1/q}.$$

Note that $c_{p,q} \rightarrow \infty$ as $q \searrow p$.

We are now in a position to compute c_i and m . First, note that the random measure μ_n , but conditioned on the event $(X_1, \dots, X_n) = X \in B(e, M_q \eta^{-1/q})^n$, is an empirical measure for $\bar{\mu} \upharpoonright B(e, M_q \eta^{-1/q})$. Thus, reasoning as in the proof of Proposition 2, we see that

$$W_p^p\left(\mu_n \mid (X_1, \dots, X_n) \in B(e, M_q \eta^{-1/q})^n, \bar{\mu} \upharpoonright B(e, M_q \eta^{-1/q})\right) = \sup_{\substack{f \in \mathcal{C}_b \\ 0 \leq f \leq (M_q \eta^{-1/q})^p}} \left[\frac{1}{n} \sum_{i=1}^n f(X_i) - \int f^c d\mu \right]$$

and likewise, conditioned on the event $X \in B(e, M_q \eta^{-1/q})^n$, the proof of Proposition 2 shows that for all $j \in \{1, \dots, n\}$,

$$w(x_1, \dots, x_j, \dots, x_n) - w(x_1, \dots, x'_j, \dots, x_n) \leq \frac{(M_q \eta^{-1/q})^p}{n} := c_j.$$

Hence $\bar{c} := \sum_{i=1}^m c_i = (M_q \eta^{-1/q})^p$, and $\sum_{i=1}^n c_i^2 = \frac{(M_q \eta^{-1/q})^{2p}}{n}$. At the same time, by the triangle inequality,

$$\begin{aligned} m &:= \mathbb{E} W_p^p\left(\mu_n \mid (X_1, \dots, X_n) \in B(0, M_q \eta^{-1/q})^n, \mu\right) \leq \\ \mathbb{E} &\left[\left(W_p\left(\mu_n \mid (X_1, \dots, X_n) \in B(e, M_q \eta^{-1/q})^n, \bar{\mu} \upharpoonright B(e, M_q \eta^{-1/q})\right) + W_p\left(\mu, \bar{\mu} \upharpoonright B(e, M_q \eta^{-1/q})\right) \right)^p \right] \\ &\leq \mathbb{E} \left[\left(W_p\left(\mu_n \mid (X_1, \dots, X_n) \in B(e, M_q \eta^{-1/q})^n, \bar{\mu} \upharpoonright B(e, M_q \eta^{-1/q})\right) + c_{p,q} M_q \eta^{1/p-1/q} \right)^p \right]. \end{aligned}$$

Therefore, Combes's inequality tells us that: for all $\eta \in (0, 1)$ and $t > 0$,

$$\begin{aligned} \mathbb{P} \left[W_p^p(\mu_n, \mu) \geq \mathbb{E} \left[\left(W_p\left(\mu_n \mid (X_1, \dots, X_n) \in B(e, M_q \eta^{-1/q})^n, \bar{\mu} \upharpoonright B(e, M_q \eta^{-1/q})\right) + c_{p,q} M_q \eta^{1/p-1/q} \right)^p \right] + t \right] \\ \leq \mathbb{P} \left[W_p^p(\mu_n, \mu) \geq m + t \right] \leq 1 - (1 - \eta)^n + \exp \left(- \frac{2n (t - (1 - (1 - \eta)^n) ((M_q \eta^{-1/q})^p))_+^2}{(M_q \eta^{-1/q})^{2p}} \right). \end{aligned}$$

□

It is possible to combine this concentration inequality with the expectation bound from Theorem 4 in a straightforward way:

Corollary 14. *Under the same assumptions as the preceding theorem, suppose, in addition, that $X = \mathbb{R}^d$, $p \in [1, d/2)$, and $q > dp/(d-p)$. Then, for all $\eta \in (0, 1)$ and $t > 0$,*

$$\begin{aligned} \mathbb{P} \left[W_p^p(\mu_n, \mu) \geq \left(\kappa_{p,q,d} M_q (1 + \eta)^{1/q} n^{-1/d} + c_{p,q} M_q \eta^{1/p-1/q} \right)^p + t \right] \\ \leq 1 - (1 - \eta)^n + \exp \left(- \frac{2n (t - (1 - (1 - \eta)^n) ((M_q \eta^{-1/q})^p))_+^2}{(M_q \eta^{-1/q})^{2p}} \right). \end{aligned}$$

Proof. If μ has moment bound M_q , then it follows that $\bar{\mu} \upharpoonright B(0, M_q \eta^{-1/q})$ has upper moment bound $M_q(1 + \eta)^{1/q}$ (because in the worst case, μ is concentrated near $\partial B(0, M_q \eta^{-1/q})$.) So, we know, using Theorem 4, that under the requisite additional assumptions of Theorem 4, namely that $p \in [1, d/2)$ and $q > dp/(d - p)$, it holds that

$$\mathbb{E} W_p \left(\mu_n \mid (X_1, \dots, X_n) \in B(0, M_q \eta^{-1/q})^n, \bar{\mu} \upharpoonright B(0, M_q \eta^{-1/q}) \right) \leq \kappa_{p,q,d} M_q (1 + \eta)^{1/q} n^{-1/d}$$

and so it follows, in this circumstance, that that

$$\begin{aligned} \mathbb{P} \left[W_p^p(\mu_n, \mu) \geq \left(\kappa_{p,q,d} M_q (1 + \eta)^{1/q} n^{-1/d} + c_{p,q} M_q \eta^{1/p-1/q} \right)^p + t \right] \\ \leq 1 - (1 - \eta)^n + \exp \left(- \frac{2n (t - (1 - (1 - \eta)^n) ((M_q \eta^{-1/q})^p)_+^2)}{(M_q \eta^{-1/q})^{2p}} \right). \end{aligned}$$

□

B.2. Analysis of the case where $d_X + d_Y = 2$.

Example 15. (expectation bound and concentration for μ supported on a simple 2d compact domain) For expectation bounds in the case where $d = 2$, we have the *asymptotic* result due to Ambrosio & Glaudo [AG19] that if μ_n and μ'_n are two empirical measures for μ , it holds that

$$\lim_{n \rightarrow \infty} \frac{n}{\log(n)} \cdot \mathbb{E} [W_2^2(\mu_n, \mu'_n)] = \frac{1}{2\pi}.$$

As for *explicit, non-asymptotic* results of this kind, the only one we are aware of is [WB19, Proposition 12]. Here, we will use this proposition to acquire an expectation bound that is asymptotically loose (indeed, the exponential order is worse than the one from [AG19]), but is usable as a finite sample bound.

We recall that the ϵ -covering number of a set $S \subseteq \mathbb{R}^d$ is the minimum number of balls of radius ϵ which cover S ; we notate this number by $\mathcal{N}_\epsilon(S)$. For a measure μ , we also define the (ϵ, τ) covering number by

$$\mathcal{N}_\epsilon(\mu, \tau) := \inf \{ \mathcal{N}_\epsilon(S) : \mu(S) \geq 1 - \tau \}.$$

Then, [WB19, Proposition 12] says that

$$\mathbb{E} [W_p^p(\mu, \hat{\mu}_n)] \leq (9^p + 3) \sqrt{\frac{m_n}{n}}; \quad m_n := \inf_{\epsilon > 0} \max \{ \mathcal{N}_\epsilon(\mu, \epsilon^p), n \epsilon^{2p} \}.$$

This bound has the advantage of giving an estimate when μ is not absolutely continuous. However, let's specify it to the case where μ is supported on the unit square in \mathbb{R}^2 . In this case, it is known that

$$\mathcal{N}_\epsilon([0, 1]^2) \leq \frac{2\sqrt{3}}{9\epsilon^2} + \frac{8}{\pi\epsilon} + 16.$$

Therefore, since $\mathcal{N}_\epsilon(\mu, \epsilon^p) \leq \mathcal{N}_\epsilon([0, 1]^2)$, we can upper bound m_n as follows:

$$m_n \leq \min_{\epsilon} \max \left\{ \frac{2\sqrt{3}}{9\epsilon^2} + \frac{8}{\pi\epsilon} + 16, n \epsilon^{2p} \right\}.$$

The minimum (in terms of ϵ) on the right hand side is attained at the point where the two expressions in braces are equal, which is the same as

$$\epsilon^{2p+2} - \frac{1}{n} \left[\frac{2\sqrt{3}}{9} + \frac{8}{\pi}\epsilon + 16\epsilon^2 \right] = 0.$$

Let $\bar{\epsilon}_n$ denote the solution to this equation for each n ; Taking $\bar{\epsilon}_n$ and plugging it into $n\epsilon^{2p}$ then gives us a bound for m_n . In general we must solve for $\bar{\epsilon}_n$ numerically for each n , but for instance, with $n = 10^6$ and $p = 1$ we have $\bar{\epsilon}_n \approx 0.026$; and moreover, $\bar{\epsilon}_n \rightarrow 0$ as $n \rightarrow \infty$. In any case, we have the upper bound

$$\mathbb{E}[W_p^p(\mu, \hat{\mu}_n)] \leq (9^p + 3)\bar{\epsilon}_n^p.$$

Lastly, combining this with the concentration inequality from Proposition 2, we deduce that if n is large enough that $(9^p + 3)\bar{\epsilon}_n^p \leq \frac{1}{2}\eta^p$, then (note in this case $D = \sqrt{2}$)

$$\mathbb{P} \left[W_p^p(\mu_n, \mu) \geq \frac{1}{2}\eta^p + t \right] \leq \exp \left(-\frac{nt^2}{2^{p-1}} \right)$$

and so, putting $t = \frac{1}{2}\eta^p$, we see that

$$\mathbb{P} [W_p(\mu_n, \mu) \geq \eta] \leq \exp \left(-\frac{n\eta^{2p}}{4 \cdot 2^{p-1}} \right).$$

Finally, if μ_n and μ'_n are both independent samples from μ , it follows that if n is large enough that $(9^p + 3)\bar{\epsilon}_n^p \leq \frac{1}{2}\eta^p$, we have the upper bound

$$\mathbb{P} [W_p(\mu_n, \mu'_n) \geq \eta] \leq 2 \exp \left(-\frac{n\eta^{2p}}{2^{p+1}} \right).$$

Similar analysis can be conducted for measures $\mu \in \mathcal{P}(\mathbb{R}^2)$ which are compactly supported on any compact domain K which is specified explicitly enough that an upper bound on the ε covering number $\mathcal{N}_\varepsilon(K)$ can be computed.

Let us discuss how this analysis can be combined with Theorem 11. Indeed, suppose that for all $z \in \text{supp}(\mathcal{L}_Z)$, $\mathcal{L}_{(X,Y)|Z=z}$ and $\mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}$ are supported on the unit square. It follows that the same is true for the binned conditional distributions $\mathcal{L}_{(X,Y)|Z \in V_j}$ and $\mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}$. Therefore, reasoning as in Example 12, if $\text{supp}(\mathcal{L}_Z)$ is binned using a partition $\{V_j\}_{j=1}^J$ with diameter bound

$$\text{diam}(V_j) \leq \frac{\varepsilon}{4 \max \left\{ L_{XY}, (2^{p-1} (L_X^p + L_Y^p))^{1/p} \right\}},$$

and for all $j \in \{1, \dots, J\}$, we have n_j large enough that if n is large enough that $(9^p + 3)\bar{\epsilon}_n^p \leq \frac{1}{2}(\frac{\eta}{2})^p$, then it holds, under the null, that

$$\mathbb{P} \left[(\exists j \in \{1, \dots, J\}) W_p(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j}) \geq \varepsilon \right] \leq 2 \sum_{i=1}^J \exp \left(-\frac{n_j \varepsilon^{2p}}{2^{3p+1}} \right).$$

In other words, we have finite sample control of the type I error.

As for the type II error, consider the alternative where, for each $j \in \{1, \dots, J\}$, there exists a $z \in V_j$ such that $W_p(\mathcal{L}_{(X,Y)|Z=z}, \mathcal{L}_{X|Z=z} \otimes \mathcal{L}_{Y|Z=z}) \geq \delta_j + \varepsilon$. Hence $W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}) \geq \delta_j + \frac{\varepsilon}{2}$ (using the diameter bound on V_j). If

$$W_p(\widehat{\mathcal{L}}_{(X,Y)|Z \in V_j}, \widehat{\mathcal{L}}_{X|Z \in V_j} \otimes \widehat{\mathcal{L}}_{Y|Z \in V_j}) < W_p(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j}) - \delta_j + \frac{\varepsilon}{2}$$

then it must be the case that

$$W_p \left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \mathcal{L}_{(X,Y)|Z \in V_j} \right) \geq \frac{\delta_j}{2} - \frac{\varepsilon}{4} \text{ or } W_p \left(\widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j} \right) \geq \frac{\delta_j}{2} - \frac{\varepsilon}{4}.$$

In the first case, putting $t = \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p$ in Proposition 2, and recalling that $D = \sqrt{2}$, we see that

$$\begin{aligned} \mathbb{P} \left[W_p \left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \mathcal{L}_{(X,Y)|Z \in V_j} \right) \geq \frac{\delta_j}{2} - \frac{\varepsilon}{4} \right] &\leq \exp \left(-\frac{2n_j \left(\delta_j - \frac{\varepsilon}{2} \right)^{2p}}{4^{2p+1} D^{2p}} \right) \\ &= \exp \left(-\frac{n_j \left(\delta_j - \frac{\varepsilon}{2} \right)^{2p}}{2^{5p+1}} \right) \end{aligned}$$

provided that n_j is sufficiently large that

$$(9^p + 3)\bar{\varepsilon}_{n_j}^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p.$$

In the second case, similarly,

$$\mathbb{P} \left[W_p \left(\widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j} \right) \geq \frac{\delta_j}{2} - \frac{\varepsilon}{4} \right] \leq \exp \left(-\frac{n_j \left(\delta_j - \frac{\varepsilon}{2} \right)^{2p}}{2^{5p+1}} \right)$$

also provided that n_j is sufficiently large that $(9^p + 3)\bar{\varepsilon}_{n_j}^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p$. Therefore,

$$\begin{aligned} \mathbb{P} \left[W_p \left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}} \right) \leq W_p \left(\mathcal{L}_{(X,Y)|Z \in V_j}, \mathcal{L}_{X|Z \in V_j} \otimes \mathcal{L}_{Y|Z \in V_j} \right) - \delta_j + \frac{\varepsilon}{2} \right] \\ \leq 2 \exp \left(-\frac{n_j \left(\delta_j - \frac{\varepsilon}{2} \right)^{2p}}{2^{5p+1}} \right) \end{aligned}$$

provided that n_j is sufficiently large that $(9^p + 3)\bar{\varepsilon}_{n_j}^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p$. Combining these estimates across bins, we conclude that

$$\mathbb{P} \left(\forall j \in \{1, \dots, J\}, W_p \left(\widehat{\mathcal{L}_{(X,Y)|Z \in V_j}}, \widehat{\mathcal{L}_{X|Z \in V_j}} \otimes \widehat{\mathcal{L}_{Y|Z \in V_j}} \right) < \varepsilon \right) \leq \prod_{j=1}^J 2 \exp \left(-\frac{n_j \left(\delta_j - \frac{\varepsilon}{2} \right)^{2p}}{2^{5p+1}} \right)$$

provided that for all $j \in \{1, \dots, J\}$, $(9^p + 3)\bar{\varepsilon}_{n_j}^p \leq \frac{1}{2} \left(\frac{\delta_j - \frac{\varepsilon}{2}}{4} \right)^p$.