

# CATCHING A FAST ROBBER ON A GRID

WILLIAM B. KINNERSLEY AND NIKOLAS TOWNSEND

ABSTRACT. We consider a variant of Cops and Robbers in which the robber may traverse as many edges as he likes in each turn, with the constraint that he cannot pass through any vertex occupied by a cop. We study this model on several classes of grid-like graphs. In particular, we determine the cop numbers for two-dimensional Cartesian grids and tori up to an additive constant, and we give asymptotic bounds for the cop numbers of higher-dimensional grids and hypercubes.

## 1. INTRODUCTION

*Cops and Robbers* is a well-studied model of pursuit and evasion in which a team of one or more cops attempts to capture a single robber on a graph  $G$ . The cops and robber occupy vertices of  $G$ ; multiple cops may occupy the same vertex, and both players know the positions of the cops and robber. The cops begin the game by choosing their initial positions, after which the robber chooses his initial position in response. Thenceforth the game proceeds in *rounds*, each consisting of a *cop turn* followed by a *robber turn*. On the cops' turn, each cop may either remain on her current vertex or move to a neighboring vertex; likewise, on the robber's turn, he may remain on his current vertex or move to a neighboring vertex. If any cop ever occupies the same vertex as the robber, we say that the cop *captures* the robber, and the cops win. Conversely, the robber wins if he can perpetually evade capture.

Given a graph  $G$ , it is natural to ask how many cops are needed to guarantee that the cops win the game on  $G$ . The *cop number* of  $G$ , denoted  $c(G)$ , is the minimum number of cops needed to guarantee capture of a robber on  $G$  no matter how the robber plays. Quilliot [16], as well as Nowakowski and Winkler [14], independently introduced the game of Cops and Robbers and characterized graphs with cop number 1. Aigner and Fromme [1] first introduced the notion of the cop number; since then, the cop number has been extensively studied. For more background on the game, we direct the reader to [5].

Cops and Robbers has spawned many variants, each of which models pursuit and evasion in a slightly different context. One natural variant permits the robber to move faster than the cops: a robber with *speed*  $s$  (for some positive integer  $s$ ) may, on each robber turn, traverse any cop-free path of length at most  $s$ ; in other words, the robber may take up to  $s$  steps on his turn, provided that he never passes through a vertex occupied by a cop. (Note that the case  $s = 1$  corresponds to the original model of Cops and Robbers.) A robber with *speed*  $\infty$  may follow any cop-free path he likes, regardless of its length. As in the usual model of Cops and Robbers, it is natural to ask how many cops are needed to capture a robber with speed  $s$ ; this quantity is the *speed- $s$  cop number* of  $G$ , denoted  $c_s(G)$ . When  $s = \infty$ , we refer to  $c_\infty(G)$  as the *infinite-speed cop number*.

---

2020 *Mathematics Subject Classification*. Primary 05C57.

*Key words and phrases*. pursuit-evasion games, cops and robbers, Cartesian grids, hypercubes.

The speed- $s$  cop number was formally introduced by Fomin, Golovach, Kratochvíl, Nisse, and Suchan in [7]. They investigated the computational complexity of computing  $c_s$  when  $s \geq 2$ ; additionally, they proved that  $c_2(P_n \square P_n) = \Omega(\sqrt{\log n})$ , thereby showing that  $c_2$  is unbounded on the class of planar graphs (in contrast to the usual cop number, which is bounded above by 3 [1]). Frieze, Krivilevich, and Loh [8] constructed  $n$ -vertex graphs with speed- $s$  cop number in  $\Omega(n^{(s-3)/(s-2)})$ ; this result was subsequently improved by Alon and Mehrabian [2], who constructed graphs with speed- $s$  cop number in  $\Omega(n^{s/(s+1)})$ . The infinite-speed cop number in particular has received a great deal of attention.  $c_\infty$  has been studied on many classes of graphs, including planar graphs [3], chordal graphs [13], interval graphs [13], expander graphs [12], random graphs [3, 12], and Cartesian products of paths and cycles [12]. Those graphs for which  $c_\infty(G) = 1$  were characterized in [12], and a construction of graphs for which  $c_\infty(G) = \Theta(V(G))$  was given in [8]. Results on the computational complexity of determining  $c_\infty(G)$  appear in [6] and [7].

In this paper, we analyze the infinite-speed game on several families of Cartesian products of paths and cycles, extending upon earlier work of Mehrabian in [12]. Mehrabian showed that when  $G$  is the  $k$ -fold Cartesian product of  $n$ -vertex paths – that is, the  $k$ -dimensional Cartesian grid – we have  $\frac{1}{4k^2}n^{k-1} \leq c_\infty(G) \leq n^{k-1}$ ; when  $G$  is the  $k$ -fold Cartesian product of  $n$ -vertex cycles, he showed  $\frac{1}{2k^2}n^{k-1} \leq c_\infty(G) \leq 2n^{k-1}$ . Finally, when  $G$  is the  $k$ -dimensional hypercube – that is, the  $k$ -fold product of  $P_2$  – he proved that for some constants  $c_1$  and  $c_2$ , we have  $\frac{c_1}{k^{3/2}}2^k \leq c_\infty(G) \leq \frac{c_2}{k}2^k$ . We aim to tighten these bounds.

In Section 2, we consider two-dimensional Cartesian grids. We show in Theorem 2.4 that  $c_\infty(P_n \square P_n)$  is  $n - 1$  when  $n$  is odd, and either  $n - 1$  or  $n$  when  $n$  is even. Similar arguments show that the infinite-speed cop number for the two-dimensional Cartesian torus,  $C_n \square C_n$ , is between  $2n - 24$  and  $2n$  (provided  $n \geq 18$ ); see Theorem 2.7. Section 3 deals with higher-dimensional grids, on which the game is substantially more complex. Our main result in this section is that for sufficiently large  $n$ , we have  $0.7172n^2 \leq c_\infty(P_n \square P_n \square P_n) \leq (0.75 + o(1))n^2$ . (Refer to Theorem 3.2 for the upper bound and Theorem 3.7 for the lower bound.) Our upper bound generalizes to grids of higher dimension: see Theorem 3.8. Finally, in Section 4, we consider hypercubes. We use a potential function argument to improve upon the lower bound from [12]: our Theorem 4.3 states that when  $G$  is the  $k$ -dimensional hypercube,  $c_\infty(G) \geq \frac{2^{k-3}}{k \ln k}$ .

## 2. TWO-DIMENSIONAL GRIDS

We begin the paper with a discussion of the infinite-speed robber game played on two-dimensional Cartesian grids, i.e. graphs of the form  $P_m \square P_n$ . Maamoun and Meyniel [10] showed that in the speed-1 game, two cops suffice to capture a robber on any two-dimensional grid. One might expect that as the speed of the robber increases, so too does the number of cops needed to capture him. This is indeed the case: Fomin et al. [7] showed that  $c_2(P_n \square P_n) = \Omega(\sqrt{\log n})$ . In this section, we determine the infinite-speed cop number of  $P_n \square P_n$  up to an additive constant; in particular, we show that  $c_\infty(P_n \square P_n) \in \{n - 1, n\}$ .

We view the vertex set of  $P_n \square P_n$  as the set  $\{(x, y) : 1 \leq x, y \leq n\}$ . The  $i$ th column of  $P_n \square P_n$  is the path consisting of all vertices of the form  $(i, y)$  for  $1 \leq y \leq n$ , and the  $j$ th row is the path consisting of vertices  $(x, j)$  for  $1 \leq x \leq n$ . The left half of  $P_n \square P_n$  consists of

the first  $\lceil n/2 \rceil$  columns, while the *right half* consists of the last  $\lfloor n/2 \rfloor$ . We say that vertex  $(x, y)$  is *above* (resp. below, left of, right of)  $(x', y')$  if  $y < y'$  (resp.  $y > y'$ ,  $x < x'$ ,  $x > x'$ ).

Mehrabian [12] observed that  $c_\infty(P_n \square P_n) \leq n$ , since  $n$  cops can begin by occupying every vertex in the first row, then move to occupy every vertex in the second row, then every vertex in the third row, and so on; the robber is forced to remain below the cops and must eventually be captured.

**Theorem 2.1** ([12], Theorem 5.1(b)). *For all positive integers  $n$ , we have  $c_\infty(P_n \square P_n) \leq n$ .*

When  $n$  is odd, this upper bound can be slightly improved.

**Theorem 2.2.** *If  $n$  is odd and at least 3, then  $c_\infty(P_n \square P_n) \leq n - 1$ .*

*Proof.* We give a strategy for  $n - 1$  cops to capture a robber on  $G$ . At the beginning of the game, for each  $k$  in  $\{1, \dots, (n - 1)/2\}$ , two cops begin on vertex  $(2k, 2k)$ . Suppose the robber begins on vertex  $(x, y)$ . We consider three cases.

- *Case 1:  $x < y$ .* For each  $k$  in  $\{1, \dots, (n - 1)/2\}$ , the two cops on vertex  $(2k, 2k)$  move to vertices  $(2k - 1, 2k)$  and  $(2k, 2k + 1)$ . Cops now occupy vertices  $(1, 2), (2, 3), \dots, (n - 1, n)$ . Let  $S$  denote the set of cop positions, and note that  $S$  is a cut-set of  $G$ : the component of  $G - S$  above the cops contains all vertices  $(i, j)$  with  $i \geq j$ , while the component below contains all vertices  $(i, j)$  with  $i < j - 1$ . Since  $x < y$ , either some cop has moved onto the robber's vertex (and the cops have won), or the robber is in the component below the cops. On the robber's turn, he cannot pass through any vertex occupied by a cop, so he must remain in the component below the cops. On the following cop turn, if the robber has not yet been captured, then the cops in columns  $1, 2, \dots, n - 2$  each move down one step. On the cops' next turn – supposing again that the robber has not yet been captured – the cops in columns  $1, 2, \dots, n - 3$  each move down one step. The cops continue in this manner until the robber is captured. (Refer to Figure 1.)
- *Case 2:  $x > y$ .* For this case, the cops can employ a strategy symmetric to that used in Case 1.
- *Case 3:  $x = y$ .* If  $x$  and  $y$  are even, then the robber occupies the same vertex as some cop and the game is already over, so suppose instead that  $x = y = 2k - 1$  for some  $k$  in  $\{1, \dots, (n + 1)/2\}$ . If  $k = 1$ , then the two cops on vertex  $(2, 2)$  move to vertices  $(1, 2)$  and  $(2, 1)$ . The robber is forced to remain on vertex  $(1, 1)$ , so the cops can capture him on their ensuing turn. Likewise, if  $k = (n + 1)/2$ , then the cops on  $(n - 1, n - 1)$  move to  $(n - 1, n)$  and  $(n, n - 1)$  and win on their next turn. Finally, suppose  $2 \leq k \leq (n - 1)/2$ . The two cops on  $(2k - 2, 2k - 2)$  move to  $(2k - 1, 2k - 2)$  and  $(2k - 2, 2k - 1)$ , while the two cops on  $(2k, 2k)$  move to  $(2k, 2k - 1)$  and  $(2k - 1, 2k)$ . The cops now occupy all four neighbors of the the robber's vertex, so once again he must remain on that vertex and the cops can capture him on their next turn. (Refer to Figure 1.)

□

Having established an upper bound on  $c_\infty(P_n \square P_n)$ , we seek to prove a matching lower bound. As it turns out, this is substantially more difficult: proving a lower bound on  $c_\infty$




FIGURE 1. Cop strategy in Theorem 2.2 (for  $n = 5$ ). From left: cops' initial position; cop response to robber below main diagonal; cop response to robber on main diagonal.

requires giving a strategy for the robber to evade a large number of cops, and the freedom of movement enjoyed by an infinite-speed robber enables fairly intricate strategies.

In general, the robber will want to seek out portions of the graph that contain no cops. Given a graph  $G$ , we call a subgraph  $H$  of  $G$  *empty* at some point during the game if, at that time, no cops occupy vertices of  $H$ . We say that the robber *can reach* or *has access to* a vertex  $v$  if there is a cop-free path joining the robber's current position to  $v$ ; the robber can reach a subgraph  $H$  of  $G$  if he can reach any vertex of  $H$ .

**Theorem 2.3.** *For all positive integers  $n$ , we have  $c_\infty(P_n \square P_n) \geq n - 1$ .*

*Proof.* The claim is clear by inspection when  $n \leq 3$ , so suppose  $n \geq 4$ . Let  $G = P_n \square P_n$ ; we give a strategy for the robber to evade  $n - 2$  cops on  $G$ . Let  $v_0$  denote the robber's initial position, and for  $i \geq 1$ , let  $v_i$  denote the robber's position after his  $i$ th move. The robber will choose each  $v_i$  so that the following properties are satisfied:

- (1) If the robber can reach an empty row of  $G$  on his  $i$ th turn, then he can reach  $v_i$ .
- (2) No cop is adjacent to  $v_i$ .
- (3) After the cops'  $(i + 1)$ st turn, the robber can reach an empty row of  $G$  from  $v_i$ .

If the robber can choose the  $v_i$  in this way, then he can perpetually evade the cops: property (2) implies that the robber cannot be captured on the cops'  $(i + 1)$ st turn, and properties (1) and (3) together imply that the robber can always reach  $v_{i+1}$  from  $v_i$ , i.e. that he can legally move to  $v_{i+1}$  with his  $(i + 1)$ st turn.

Fix  $i \geq 0$ , and consider the state of the game after the cops'  $i$ th turn. We explain how the robber chooses  $v_i$ . We consider two cases.

**Case 1:** For some  $k$  in  $\{2, 3, \dots, n - 2\}$ , either the first  $k$  or the last  $k$  rows of  $G$  contain no more than  $k - 2$  cops between them. Without loss of generality, suppose the first  $k$  rows contain at most  $k - 2$  cops. Within these rows, either the left half or the right half must contain at most  $\lfloor \frac{k-2}{2} \rfloor$  cops. Suppose this is true of the right half; a similar argument suffices if it is true for the left half. We refer to the first  $k$  rows of the right half of  $G$  as the robber's *sector* of  $G$  and denote this subgraph by  $S$ .

If the first two rows of  $S$  contain no cops, then the robber chooses  $v_i$  to be any vertex in the first row of  $S$  that doesn't belong to the first column of  $S$ . This choice of  $v_i$  clearly satisfies property (2). It satisfies property (1) as well: if the robber can reach an empty row of  $G$ , then he can use this empty row to reach an empty column (which must exist since there are  $n$  columns but only  $n - 2$  cops), and he can use this empty column to reach the

first row of  $G$  and subsequently  $v_i$ . Finally, to see that it satisfies property (3), note that since the first two rows of  $G$  are empty before the cops'  $(i + 1)$ st turn, the first row remains empty after the cops'  $(i + 1)$ st turn.

Suppose instead that the first two rows of  $S$  contain at least one cop.  $S$  contains  $k$  rows and at most  $\lfloor \frac{k-2}{2} \rfloor$  cops. Note that  $S$  must have at least 4 rows: it contains no more than  $\lfloor \frac{k-2}{2} \rfloor$  cops, but the first two rows contain at least one cop, hence  $\lfloor \frac{k-2}{2} \rfloor \geq 1$  and so  $k \geq 4$ . We claim that there must be some four consecutive rows within  $S$  that contain at most one cop between them. To prove this, let  $k = 4d + r$  for some positive integer  $d$  and some  $r \in \{0, \dots, 3\}$ ; we consider two cases.

- Suppose  $r \in \{0, 1\}$ . In this case,  $\lfloor \frac{k-2}{2} \rfloor = 2d - 1$ . Partition the first  $4d$  rows of  $S$  into  $d$  groups of four rows apiece; since there are at most  $2d - 1$  cops, at least one of these groups of four rows must have fewer than two cops.
- Suppose  $r \in \{2, 3\}$ . By assumption, the first two rows of  $S$  contain at least one cop. Hence, the next  $k - 2$  rows contain at least  $\lfloor \frac{k-2}{2} \rfloor - 1$  cops, which simplifies to  $2d - 1$ . As in the previous case, we may partition the next  $4d$  rows of  $S$  into groups of four, one of which must have fewer than two cops.

The robber now chooses  $v_i$  to be any vertex within the second and third of these four rows that is not currently adjacent to a cop and does not belong to the first column of  $S$ . (To see that this is possible, first note that since  $S$  has at least four rows, we must have  $n \geq 6$ , hence  $S$  has at least three columns. Even after discarding the first column of  $S$ , at least two columns remain; the robber has at least four potential destinations to choose from, and a single cop cannot be adjacent to all of them.)

The robber's choice of  $v_i$  clearly satisfies property (2). To see that it satisfies property (1), suppose the robber can reach an empty row of  $G$ . Since  $G$  has  $n$  columns and at most  $n - 2$  cops, it must have an empty column, which the robber can reach using the empty row. There must also be an empty row among the first  $k$  rows of  $G$ ; the robber can reach such a row using the empty column of  $G$ . Next, the robber can use this empty row to reach an empty column of  $S$  (which must exist since  $S$  contains at least  $\lfloor n/2 \rfloor$  columns and at most  $\lfloor \frac{k-2}{2} \rfloor$  cops). Finally, since  $v_i$  is either in or adjacent to an empty row of  $S$ , the robber can use the empty columns of  $S$  to reach this row and subsequently  $v_i$ .

It remains to be shown that the robber's choice of  $v_i$  satisfies property (3). Let  $S'$  denote  $S$  with the last row and first column removed. Consider the state of the game after the cops'  $(i + 1)$ st turn. Although some cops may have entered the last row or first column of  $S$  on their turn, no cops outside  $S$  can have entered  $S'$ . Consequently,  $S'$  has at most  $\lfloor \frac{k-2}{2} \rfloor$  cops. Before the cops' move,  $v_i$  belonged to either the second or third of four consecutive rows in  $S'$  with at most one cop between them; hence, after the cops' move, the second and third rows still have at most one cop between them. Consequently, either the row of  $S'$  that contains  $v_i$  is empty, or the row directly above or below  $v_i$  is empty. Either way, the robber can clearly reach an empty row of  $S'$ . Note that  $S'$  contains at least  $\lfloor n/2 \rfloor - 1$  columns and, as noted above, has at most  $\lfloor \frac{k-2}{2} \rfloor$  cops. Since

$$\left\lfloor \frac{k-2}{2} \right\rfloor = \left\lfloor \frac{k}{2} \right\rfloor - 1 \leq \left\lfloor \frac{n-2}{2} \right\rfloor - 1 = \left\lfloor \frac{n}{2} \right\rfloor - 2 < \left\lfloor \frac{n}{2} \right\rfloor - 1,$$

$S'$  has at least one empty column. Since the robber can reach an empty row of  $S'$ , he can thus also reach this empty column of  $S'$ . Now note that the first  $k$  rows of  $G$  contained at most  $k - 2$  cops before the cops' move; some cops may have entered the  $k$ th row of  $G$ , but the first  $k - 1$  rows must still have at most  $k - 2$  cops. Consequently, one of the first  $k - 1$  rows of  $G$  is empty. Since the robber can reach an empty column of  $S'$ , he can use it to reach this empty row of  $G$ .

Thus, the robber's choice of  $v_i$  satisfies properties (1), (2), and (3), as claimed.

**Case 2:** There is no  $k$  in  $\{2, 3, \dots, n - 2\}$  such that either the first  $k$  or the last  $k$  rows of  $G$  contain at most  $k - 2$  cops. In this case, the first two rows of  $G$  cannot both be empty, since then we could take  $k = 2$  and use the first two rows of  $G$ . The first two rows of  $G$  cannot contain two or more cops between them, since then we could take  $k = n - 2$  and use the last  $n - 2$  rows of  $G$ . Thus, the first two rows of  $G$  contain exactly one cop between them. Similarly, the last two rows of  $G$  must also contain exactly one cop between them. Every other row of  $G$  must contain exactly one cop: if row  $j$  were to contain more than one cop, for some  $j$  in  $\{3, \dots, n - 2\}$ , then either the first  $j - 1$  rows would contain at most  $j - 3$  cops, or the last  $n - j$  rows would contain at most  $n - j - 2$  cops. By symmetry, we may also assume that there is no  $k$  in  $\{2, 3, \dots, n - 2\}$  such that either the first  $k$  or the last  $k$  columns of  $G$  contain at most  $k - 2$  cops; otherwise, the robber could choose  $v_i$  as in Case 1, except with the roles of rows and columns switched. (Note that properties (1)-(3) are symmetric with respect to rows and columns, because the robber can reach an empty row if and only if he can reach an empty column.) Hence, we may also conclude that the first two columns contain exactly one cop, the last two columns contain exactly one cop, and every other column contains exactly one cop.

The robber chooses  $v_i$  to be any vertex that belongs to whichever of the first two rows is empty, is not adjacent to a cop, and is not in the first or last column of  $G$ . It is clear that this choice of  $v_i$  satisfies property (2). It also satisfies property (1): if the robber can reach an empty row of  $G$ , then he can reach an empty column, use this column to reach whichever of the first two rows is empty, and proceed from there to  $v_i$ .

Finally, consider property (3). After the cops' move, if either the first or second row is empty, then the robber can clearly reach an empty row and property (3) is satisfied. Otherwise, it must be that the cop in row 3 moved up into row 2. Say this cop occupies column  $k$ . We claim that the robber can either reach all columns to the left of column  $k$  or all columns to the right. Since neither of the first two rows is empty, each must contain exactly one cop. If the robber is strictly to the left of column  $k$ , then he can move into the second row (if he isn't already there) and use it to reach all columns to the left of column  $k$ . Similarly, if he is to the right of column  $k$ , then he can reach all columns to the right. Finally, if the robber is in column  $k$ , then he must be in row 1; if the cop in row 1 is to the left of the robber then the robber can reach all columns to the right of column  $k$ , and otherwise he can reach all columns to the left.

Without loss of generality, suppose the robber can reach all columns to the left of column  $k$ . Before the cops' move, one of the first two columns was empty; in particular, there was an empty column to the left of column  $k$ . We claim that there is still an empty column to the left of column  $k$ ; it would then follow that the robber can reach an empty column, from which he could reach an empty row, so property (3) would be satisfied.

If column 1 is empty then we are done, so suppose otherwise. Choose the least  $\ell$  greater than 1 such that the cop in column  $\ell$  did not move to the left; note that  $\ell \leq k$ , since the cop in column  $k$  moved up, not left. In order for neither of the first two columns to be empty, it must be that the cop in column 3 moved left into column 2. Now, in order for column 3 not to be empty, the cop in column 4 must have moved left into column 3, and so on. It follows that column  $\ell - 1$  must be empty, since the cop in column  $\ell - 1$  moved left, but the cop in column  $\ell$  did not. Since  $\ell - 1 < k$ , we have shown that there is an empty column to the left of column  $k$ , as claimed. It follows that the robber's choice of  $v_i$  satisfies properties (1), (2), and (3).  $\square$

The following theorem summarizes our bounds on  $c_\infty(P_n \square P_n)$ .

**Theorem 2.4.** *If  $n \leq 2$ , then  $c_\infty(P_n \square P_n) = n$ ; if  $n$  is odd and at least 3, then  $c_\infty(P_n \square P_n) = n - 1$ ; if  $n$  is even and at least 4, then  $c_\infty(P_n \square P_n) \in \{n - 1, n\}$ .*

*Proof.* Let  $G = P_n \square P_n$ . If  $n \leq 2$ , then by inspection  $c_\infty(G) = n$ . If  $n$  is odd and at least 3, then Theorems 2.2 and 2.3 together give  $n - 1 \leq c_\infty(G) \leq n - 1$ ; if  $n$  is even and at least 4, then Theorem 2.1 and 2.3 together give  $n - 1 \leq c_\infty(G) \leq n$ .  $\square$

Although we have thus far focused exclusively on  $n \times n$  grids, the game behaves quite similarly on general  $m \times n$  grids, as we next show.

**Theorem 2.5.** *If  $m, m', n$ , and  $n'$  are positive integers with  $m \leq m'$  and  $n \leq n'$ , then  $c_\infty(P_m \square P_n) \leq c_\infty(P_{m'} \square P_{n'})$ .*

*Proof.* Let  $G = P_m \square P_n$ , let  $G' = P_{m'} \square P_{n'}$ , and let  $C = c_\infty(P_m \square P_n)$ ; we give a strategy for the robber to evade  $C - 1$  cops on  $G'$ . The robber imagines that he is playing the game on  $G$  and uses a strategy for that game to guide his play on  $G'$ . At each point in the game, each cop occupies a vertex of  $G'$ , but the robber will imagine that she occupies a vertex of  $G$ : in particular, a cop at vertex  $(x', y')$  in  $G'$  will be imagined to occupy vertex  $(x, y)$  in  $G$ , where  $x = \min\{x', m\}$  and  $y = \min\{y', n\}$ . The robber's strategy is now straightforward: he simply follows a winning strategy for the game on  $G$ . Whenever his strategy calls for him to move to a vertex  $(x, y)$  in  $G$ , he moves to  $(x, y)$  in  $G'$ .

Note that the robber's moves are all legal in the game on  $G'$ : if there is a cop-free path in  $G$  from the robber's current position to some vertex  $v$ , then the corresponding path in  $G'$  is also cop-free. The cops' imagined moves in  $G$  are also legal, since traversing a single edge in  $G'$  corresponds either to traversing a single edge in  $G$  or to sitting still. Finally, if a cop ever captures the robber on  $G'$  then she captures the robber in the imagined game on  $G$  as well: since she occupies the same vertex  $(x, y)$  as the robber in  $G'$ , we must have  $x \leq m$  and  $y \leq n$ , so she occupies  $(x, y)$  in  $G$  as well and thus captures the robber in  $G$ .

Since the robber's strategy enables him to perpetually evade  $C - 1$  cops on  $G$ , it also allows him to perpetually evade  $C - 1$  cops on  $G'$ .  $\square$

The cop strategy used to show that  $c_\infty(P_n \square P_n) \leq n$  extends naturally to  $P_m \square P_n$  and shows that  $c_\infty(P_m \square P_n) \leq \min\{m, n\}$ . Additionally, Theorems 2.4 and 2.5 together show that  $c_\infty(P_m \square P_n) \geq \min\{m, n\} - 1$ .

**Corollary 2.6.** *For all positive integers  $m$  and  $n$ ,*

$$\min\{m, n\} - 1 \leq c_\infty(P_m \square P_n) \leq \min\{m, n\}.$$

We conclude the section with a brief look at the discrete torus  $C_n \square C_n$ . We can establish lower and upper bounds on  $c_\infty(C_n \square C_n)$  with arguments very similar to those used for  $P_n \square P_n$ . As with  $P_n \square P_n$ , we view the vertex set of  $C_n \square C_n$  as the set  $\{(x, y) : 1 \leq x, y \leq n\}$ . The  $i$ th column is the cycle consisting of all vertices of the form  $(i, y)$  for  $1 \leq y \leq n$ , and the  $j$ th row is the cycle consisting of  $(x, j)$  for  $1 \leq x \leq n$ .

**Theorem 2.7.** *For all  $n \geq 18$ , we have  $2n - 24 \leq c_\infty(C_n \square C_n) \leq 2n$ .*

*Proof.* Let  $n \geq 18$  and let  $G = C_n \square C_n$ .

For the upper bound, cops begin by occupying every vertex on the first and  $n$ th rows of  $G$ . On each subsequent cop turn, those cops who began on the first row move down, and those cops who began on the  $n$ th row move up. The robber cannot pass through the two rows of cops and hence will eventually be captured.

For the lower bound, we give a strategy for the robber to evade  $2n - 25$  cops on  $G$ . Let  $n = 6d + r$ , for some positive integer  $d$  and  $r \in \{0, \dots, 5\}$ .

We partition the rows of  $G$  into six *bands*, each consisting of either  $d$  or  $d + 1$  consecutive rows. We call a row or column of  $G$  *nearly empty* if there exists at most one cop in that row or column. Let  $v_0$  denote the robber's initial position, and for  $i \geq 1$ , let  $v_i$  denote the robber's position after his  $i$ th move. The robber will choose each  $v_i$  such that the following properties hold:

- (1) If the robber can reach a nearly empty row of  $G$ , then he can reach  $v_i$ .
- (2) No cop is adjacent to  $v_i$ .
- (3) After the cops'  $(i + 1)$ st move, the robber can reach a nearly empty row of  $G$  from  $v_i$ .

As in the proof of Theorem 2.3, property (2) ensures that the robber cannot lose on the cops'  $(i + 1)$ st turn, while properties (1) and (3) together ensure that, on his  $(i + 1)$ st turn, he can reach  $v_{i+1}$  from  $v_i$ . Thus if the robber can choose the  $v_i$  to have these three properties, then he can evade the cops indefinitely.

Fix  $i \geq 0$  and consider the state of the game after the cops'  $i$ th turn. We explain how the robber chooses  $v_i$ . Since there are at most  $2n - 25$  cops, there must exist a band of  $d$  rows with at most  $2d - 5$  cops or a band of  $d + 1$  rows with at most  $2(d + 1) - 5$  cops. In either case, this band has at least three nearly empty rows. Since

$$2(d + 1) - 5 = 2d - 3 < \frac{6d + r}{3} = \frac{n}{3},$$

the band must have three consecutive empty columns. The robber chooses  $v_i$  to be the vertex at the intersection of the second of these three columns and the second nearly empty row in the band. Clearly,  $v_i$  satisfies property (2); we will next show that it also satisfies properties (1) and (3).

To show that  $v_i$  satisfies property (1), suppose the robber can reach a nearly empty row of  $G$ . The robber would like to move to  $v_i$ , which is located on a nearly empty row in some band of  $G$ . Since there are fewer than  $2n - 4$  cops playing on  $G$ , there must be at least three nearly empty columns in  $G$ . At most one of these three columns may contain a cop in the robber's current row, and at most one of these columns may contain a cop in the row containing  $v_i$ . Thus, using the nearly empty row, the robber can access at least one nearly empty column with no cop in either the robber's current row nor in the row containing  $v_i$ ; using that column, he can reach the nearly empty row that contains  $v_i$  and, consequently,  $v_i$  itself, so (1) is satisfied.

Finally, to show that  $v_i$  satisfies property (3), suppose the cops have made their  $(i + 1)$ st move. Before the cops' move,  $v_i$  was in the middle of three consecutive empty columns in some band. Thus, after the cops' move,  $v_i$  is still in a column of the band that is empty except perhaps for the first and last rows in the band (since some cops may have moved into these rows on the cops' move). Let  $k$  denote the number of rows in the band. There were at most  $2k - 5$  cops in the band before the cops' move, so disregarding the first and last rows of the band, there are still at most  $2k - 5$  cops spread across  $k - 2$  rows. At least one of these rows must be nearly empty; the robber can reach this row using his current column in the band, so property (3) is satisfied.  $\square$

### 3. HIGHER-DIMENSIONAL GRIDS

We now turn our attention to three-dimensional Cartesian grids. We begin with the following observation; we omit the proof, since it is extremely similar to that of Theorem 2.5.

**Theorem 3.1.** *For positive integers  $m, m', n, n', p,$  and  $p'$ , if  $m \leq m'$ ,  $n \leq n'$ , and  $p \leq p'$ , then  $c_\infty(P_m \square P_n \square P_p) \leq c_\infty(P_{m'} \square P_{n'} \square P_{p'})$ .*

Henceforth, we focus on the  $n \times n \times n$  grid  $P_n \square P_n \square P_n$ . Let  $G = P_n \square P_n \square P_n$ . We view the vertex set of  $G$  as the set  $\{(x, y, z) : 0 \leq x, y, z \leq n - 1\}$ . A *row* of  $G$  is a path  $R_{y,z}$  consisting of those vertices whose second and third coordinates are  $y$  and  $z$ , respectively. Similarly, a *column* of  $G$  is a path  $C_{x,z}$  consisting of all vertices whose first and third coordinates are  $x$  and  $z$ , and a *shaft* of  $G$  is a path  $S_{x,y}$  consisting of all vertices whose first and second coordinates are  $x$  and  $y$ , respectively. The  $k$ th *plane* of  $G$  is the subgraph induced by those vertices whose third coordinate is  $k$ .

Our first main result in this section is an upper bound on  $c_\infty(P_n \square P_n \square P_n)$ . In what follows, the following intuition will be helpful for the reader. Often, an effective cop strategy works (very roughly) as follows. First, the cops occupy a cut-set of the graph, thereby splitting the graph into two or more components. They then identify the component in which the robber is located and gradually close in, making the robber's component smaller and smaller, until he is eventually captured. To execute such a strategy with as few cops as possible, the cops will often want to find a "small" cut-set  $S$  in the graph  $G$  that minimizes the size of the largest component of  $G - S$ . In particular, the cops may wish to find a small cut-set  $S$  that splits  $G - S$  into two components, each containing roughly half the vertices of  $G$ .

**Theorem 3.2.** *For all positive integers  $n$ , we have  $c_\infty(P_n \square P_n \square P_n) \leq (0.75 + o(1))n^2$ .*

*Proof.* Since we seek an asymptotic bound, by Theorem 3.1 it suffices to consider the case where  $n$  is odd. Let  $G = P_n \square P_n \square P_n$ ; we give a strategy for  $(0.75 + o(1))n^2$  cops to catch a robber on  $G$ . Loosely, the cops' strategy works as follows. The cops will be divided into two groups: a group of  $(3n^2 + 1)/4$  *blocking cops* and a group of  $(n + 1)/2$  *reserve cops*. The blocking cops will occupy a cut-set  $S$  of  $G$ , which we refer to as the cops' *blockade*. The robber must choose an initial position in some component of  $G - S$ , which we refer to as the *robber's component*. The blockade prevents the robber from leaving his component. Over the course of several rounds, the cops will gradually move the blockade so as to repeatedly decrease the size of the robber's component, leading to the robber's eventual capture.

For  $m \in \{0, 1, \dots, 3(n-1)\}$ , let  $S_m$  be the set of all vertices in  $G$  whose coordinates sum to  $m$ . Let  $m^* = 3(n-1)/2$ ; the cops' initial blockade will be the set  $S_{m^*}$ . Initially, one blocking cop begins on each vertex of  $S_{m^*}$ . To count the number of cops needed to occupy all of  $S_{m^*}$ , we must count the number of nonnegative-integer triples  $(x, y, z)$  such that  $x + y + z = m^*$ . This is equivalent to the number of  $m^*$ -combinations with repetition from 3 elements, which is  $\binom{m^*+2}{2}$ . However, not every such triple corresponds to a vertex in  $G$ , since  $x, y$ , or  $z$  might exceed  $n-1$ . The number of these triples in which  $x \geq n$  is equivalent to the number of  $(m^*-n)$ -combinations with repetition from 3 elements, which is  $\binom{m^*-n+2}{2}$ ; by symmetry, this also counts the number of triples in which  $y \geq n$  and those in which  $z \geq n$ . Consequently, we have

$$\begin{aligned}
|S| &= \binom{m^*+2}{2} - 3\binom{m^*-n+2}{2} \\
&= \frac{(m^*+2)(m^*+1)}{2} - 3\frac{(m^*-n+2)(m^*-n+1)}{2} \\
&= \frac{(3n+1)(3n-1)}{8} - 3\frac{(n+1)(n-1)}{8} \\
&= \frac{9n^2-1}{8} - \frac{3n^2-3}{8} \\
&= \frac{3n^2+1}{4},
\end{aligned}$$

so there are enough blocking cops to occupy all vertices of  $S_{m^*}$ . The reserve cops choose their initial positions arbitrarily.

$G - S_{m^*}$  contains two components, namely the subgraph of  $G$  induced by those vertices whose coordinates sum to less than  $m^*$  and the subgraph consisting of those vertices whose coordinates sum to more than  $m^*$ ; suppose without loss of generality that the robber begins in the former component.

Henceforth, the game will progress in *phases*, each of which will consist of a finite number of rounds. At the end of each phase, the blocking cops will move to decrease the size of the robber's component, while the reserve cops will help reinforce the blockade. In the first phase, one reserve cop moves to each of the  $(n+1)/2$  vertices of  $S_{m^*-1}$  whose third coordinate is  $n-1$ . This may take several rounds; however, note that while the reserve cops are moving into position, the cops' blockade prevents the robber from leaving his component. Once the reserve cops are in position, each reserve cop and each blocking cop on the 0th plane remain in place, while all other blocking cops move down in the graph (i.e. they move so as to decrease their third coordinate).

Note that cops now occupy all vertices of  $S_{m^*-1}$ ; we reclassify those cops in  $S_{m^*-1}$  as "blocking cops" and the remaining cops as "reserve cops". In the next phase, the reserve cops move to cover all vertices of  $S_{m^*-2}$  whose third coordinate is  $n-1$ , after which point the cops can shift the blockade downward to  $S_{m^*-2}$ . The cops iterate this strategy, repeatedly shifting the blockade downward. The robber's component gets smaller with each phase, so eventually he must be caught.

□

Theorem 3.2 established an upper bound on  $c_\infty(P_n \square P_n \square P_n)$ ; we now seek a lower bound. As with  $P_n \square P_n$ , the lower bound is considerably more complicated than the upper bound, so we start with a few preliminary lemmas.

The cops' strategy in Theorem 3.2 revolved around splitting  $P_n \square P_n \square P_n$  into two equally-sized pieces as efficiently as possible, then slowly shrinking the robber's component. To evade capture, it is crucial that the robber avoid this sort of situation. As such, in the proof of Theorem 3.7, our lower bound on  $c_\infty(P_n \square P_n \square P_n)$ , it will be useful to know how efficiently a fixed number cops can split the graph into two pieces; even if the cops lack the numbers needed to split the graph into equally-sized pieces, we would like to know how close they can come.

Thus motivated, we introduce the following notation.

**Definition 3.3.** Fix positive integers  $k$  and  $C$  and integers  $n_1, n_2, \dots, n_k$  all at least 2, and let  $G = P_{n_1} \square P_{n_2} \square \dots \square P_{n_k}$ . Define  $L(C; n_1, \dots, n_k)$  to be the minimum, over all subsets  $S$  of  $V(G)$  with  $|S| \leq C$ , of the maximum size of a component in  $G - S$ .

It is clear from the definition that  $L(C; n_1, \dots, n_k)$  is nonincreasing in  $C$ . Suppose  $C$  is sufficiently small that  $L(C; n_1, \dots, n_k) > |V(G)|/2$ , and suppose we play on  $G$  with  $C$  cops. In this situation, for every possible set  $S$  of cop positions, some component of  $G - S$  contains at least half the vertices of  $G$ ; we refer to this component as the *large component* of  $G$ . (Note that the large component has size at least  $L(C; n_1, \dots, n_k)$ .)

Furthermore, let  $H$  denote the union of all components of  $G - S$  other than the large component. We sometimes refer to  $H$  as the *small component* of  $G$ . (This is an abuse of notation because  $H$  need not be connected; however, it will turn out that when the cops play "efficiently", the small component is in fact connected.) In the proof of Theorem 3.7, we will need to be able to bound the size of the small and large components of various subgraphs of  $P_n \square P_n \square P_n$ , while playing against various numbers of cops. To facilitate this, we introduce the following notation.

**Definition 3.4.** For a positive integer  $m$ , define

- $C_m(n_1, \dots, n_k) = |\{(x_1, \dots, x_k) : x_1 + \dots + x_k = m \text{ and } 0 \leq x_i < n_i \text{ for all } i\}|$
- $S_m(n_1, \dots, n_k) = |\{(x_1, \dots, x_k) : x_1 + \dots + x_k < m \text{ and } 0 \leq x_i < n_i \text{ for all } i\}|$
- $L_m(n_1, \dots, n_k) = |\{(x_1, \dots, x_k) : x_1 + \dots + x_k > m \text{ and } 0 \leq x_i < n_i \text{ for all } i\}|$

To motivate the choice of notation, let us remark that (as we will see shortly), when playing on  $P_{n_1} \square P_{n_2} \square \dots \square P_{n_k}$  with  $C_m(n_1, \dots, n_k)$  cops, the size of the small component will be at most  $S_m(n_1, \dots, n_k)$  and the size of the large component will be at least  $L_m(n_1, \dots, n_k)$ .

Bollobás and Leader in [4] completely characterized the sets  $S$  of fixed size that maximize the size of the small component (and hence minimize the size of the large component). We will not need the full power of their result; the following special case will suffice.

**Lemma 3.5.** *If  $C \leq C_m(n_1, \dots, n_k)$ , then  $L(C; n_1, \dots, n_k) \geq L_m(n_1, \dots, n_k)$ .*

*Proof.* By Theorem 8 in [4], when  $C = C_m(n_1, \dots, n_k)$ , the maximum size of the small component is  $S_m(n_1, \dots, n_k)$ , realized when  $S = \{(x_1, \dots, x_k) \in V(G) : x_1 + \dots + x_k = m\}$ . The large component consists of all vertices in  $G$  that do not belong to  $S$  and are not occupied

by cops; consequently,

$$L(C; n_1, \dots, n_k) = \prod_{i=1}^k n_i - S_m(n_1, \dots, n_k) - C_m(n_1, \dots, n_k) = L_m(n_1, \dots, n_k).$$

When  $C < C_m(n_1, \dots, n_k)$ , the claim holds because  $L(C; n_1, \dots, n_k)$  is nonincreasing in  $C$ .  $\square$

Intuitively, Lemma 3.5 states that the most “efficient” way for a limited number of cops to divide  $G$  into two components is to occupy the set of vertices whose coordinates sum to  $m$ , for some  $m$ .

In the lemma below, we determine some specific values of  $C_m$  and  $S_m$  that will be useful later.

**Lemma 3.6.** *Fix a positive integer  $n$  and let  $m = c(n - 1)$  for some positive real number  $c$ .*

(a) *If  $c \leq 1$ , then:*

- $C_m(n, n) = (c + o(1))n$ , and
- $S_m(n, n) = \left(\frac{1}{2}c^2 + o(1)\right)n^2$ .

(b) *If  $n$  is even and  $0.5 \leq c \leq 1$ , then:*

- $C_m(n/2 - 1, n/2 - 1, n) = \left(-\frac{1}{2}c^2 + c - \frac{1}{4} + o(1)\right)n^2$ , and
- $S_m(n/2 - 1, n/2 - 1, n) = \left(-\frac{1}{6}c^3 + \frac{1}{2}c^2 - \frac{1}{4}c + \frac{1}{24} + o(1)\right)n^3$ .

(c) *If  $n$  is even and  $0.5 \leq c \leq 1$ , then:*

- $C_m(n/2 - 1, n, n) = \left(\frac{1}{2}c - \frac{1}{8} + o(1)\right)n^2$ , and
- $S_m(n/2 - 1, n, n) = \left(\frac{1}{4}c^2 - \frac{1}{8}c + \frac{1}{48} + o(1)\right)n^3$ .

(d) *If  $c \leq 1$ , then:*

- $C_m(n, n, n) = \left(\frac{1}{2}c^2 + o(1)\right)n^2$ , and
- $S_m(n, n, n) = \left(\frac{1}{6}c^3 + o(1)\right)n^3$ .

*If instead  $1 < c \leq 1.5$ , then:*

- $C_m(n, n, n) = \left(-c^2 + 3c - \frac{3}{2} + o(1)\right)n^2$ , and
- $S_m(n, n, n) = \left(-\frac{1}{3}c^3 + \frac{3}{2}c^2 - \frac{3}{2}c + \frac{1}{2} + o(1)\right)n^3$ .

*Proof.*

(a)  $C_m(n, n)$  represents the number of ordered pairs  $(x, y)$  with  $0 \leq x, y \leq n - 1$  such that  $x + y = m$ ; consequently,

$$C_m(n, n) = m + 1 = (c + o(1))n.$$

Similarly,  $S_m(n, n)$  represents the number of ordered pairs  $(x, y)$  with  $0 \leq x, y \leq n-1$  such that  $x + y \leq m$ . This is equivalent to the number of  $m$ -combinations with repetition from 3 elements, which is  $\binom{m+2}{2}$ . Consequently,

$$S_m(n, n) = \binom{m+2}{2} = \left(\frac{1}{2}c^2 + o(1)\right)n^2.$$

- (b)  $C_m(n/2 - 1, n/2 - 1, n)$  represents the number of ordered triples  $(x, y, z)$  with  $0 \leq x, y \leq n/2 - 2$  and  $0 \leq z \leq n - 1$  such that  $x + y + z = m$ . The number of nonnegative integer ordered triples  $(x, y, z)$  satisfying  $x + y + z = m$  is equivalent to the number of  $m$ -combinations with repetition from 3 elements, which is  $\binom{m+2}{2}$ . All such triples satisfy  $x, y, z \geq 0$ , as well as  $z \leq n - 1$  (since  $m \leq n - 1$ ); however, they do not all satisfy  $x, y \leq n/2 - 2$ . The number of these triples that *do not* satisfy  $x \leq n/2 - 2$  is equivalent to the number of  $(m - n/2 + 1)$ -combinations with repetition from 3 elements, which is  $\binom{m-n/2+3}{2}$ ; by symmetry, this is also the number of triples not satisfying  $y \leq n/2 - 2$ . Thus, we have

$$C_m(n/2 - 1, n/2 - 1, n) = \binom{m+2}{2} - 2\binom{m-n/2+3}{2} = \left(-\frac{1}{2}c^2 + c - \frac{1}{4} + o(1)\right)n^2.$$

We can compute  $S_m(n/2 - 1, n/2 - 1, n)$  similarly. This time, we seek the number of solutions to  $x + y + z \leq m$  satisfying  $0 \leq x, y \leq n/2 - 2$  and  $0 \leq z \leq n - 1$ . The number of nonnegative integer solutions to  $x + y + z \leq m$  is equivalent to the number of  $m$ -combinations from 4 elements with repetition, which is  $\binom{m+3}{3}$ ; the number not satisfying  $x \leq n/2 - 2$  (and, by symmetry, the number not satisfying  $y \leq n/2 - 2$ ) is  $\binom{m-n/2+4}{3}$ . Thus,

$$S_m(n/2 - 1, n/2 - 1, n) = \binom{m+3}{3} - 2\binom{m-n/2+4}{3} = \left(-\frac{1}{6}c^3 + \frac{1}{2}c^2 - \frac{1}{4}c + \frac{1}{24} + o(1)\right)n^3.$$

- (c) We can apply an argument similar to that in part (b); the only difference is that when computing both  $C_m$  and  $S_m$ , we seek triples  $(x, y, z)$  satisfying  $0 \leq x \leq n/2 - 2$  and  $0 \leq y, z \leq n - 1$ , rather than  $0 \leq x, y \leq n/2 - 2$  and  $0 \leq z \leq n - 1$ . As in part (b), we thus have

$$C_m(n/2 - 1, n, n) = \binom{m+2}{2} - \binom{m-n/2+3}{2} = \left(\frac{1}{2}c - \frac{1}{8} + o(1)\right)n^2$$

and

$$S_m(n/2 - 1, n, n) = \binom{m+3}{3} - \binom{m-n/2+4}{3} = \left(\frac{1}{4}c^2 - \frac{1}{8}c + \frac{1}{48} + o(1)\right)n^3.$$

- (d) We can proceed as in parts (b) and (c), except that now we must satisfy the constraints  $0 \leq x, y, z \leq n - 1$ . As before, the number of solutions to  $x + y + z = m$  is simply  $\binom{m+2}{2}$  and the number of solutions to  $x + y + z \leq m$  is  $\binom{m+3}{3}$ . If  $c \leq 1$ , then the constraints do not exclude any solutions, so  $C_m(n, n, n) = \binom{m+2}{2}$  and  $S_m(n, n, n) = \binom{m+3}{3}$ ; if instead  $1 < c \leq 1.5$ , then we must exclude solutions in which  $x, y, \text{ or } z$  exceeds  $n - 1$ , hence  $C_m(n, n, n) = \binom{m+2}{2} - 3\binom{m-n+2}{2}$  and  $S_m(n, n, n) = \binom{m+3}{3} - 3\binom{m-n+3}{3}$ .

□

We are nearly ready to establish a lower bound on  $c_\infty(P_n \square P_n \square P_n)$ , but first we present some definitions that will be used in the proof. Let  $G = P_n \square P_n \square P_n$ . For a subgraph  $H$  of  $G$ , let  $\partial(H)$  denote the set of vertices in  $H$  that have neighbors outside of  $H$ , and let  $\text{int}(H)$  denote the subgraph induced by  $V(H) - \partial(H)$ .

For a family  $\mathcal{H} = \{H_1, H_2, \dots, H_k\}$  of subgraphs of  $G$ , we call  $H_j$  the *sparsest* member of  $\mathcal{H}$  if  $H_j$  contains the fewest cops among members of  $\mathcal{H}$ . (A family may contain more than one sparsest subgraph.)

Let  $T$  denote the set of vertices of  $G$  of the form  $(x, y, z)$  with  $z \geq \lceil (n-1)/2 \rceil$ . The *top half* of  $G$  is the subgraph of  $G$  induced by  $T$ , while the *bottom half* is the subgraph induced by  $\overline{T}$ . Similarly, let  $F$  be the set of vertices  $(x, y, z)$  with  $y \geq \lceil (n-1)/2 \rceil$  and let  $R$  be the set of vertices  $(x, y, z)$  with  $x \geq \lceil (n-1)/2 \rceil$ ; the *front half* of  $G$  is the subgraph induced by  $F$ , the *back half* is the subgraph induced by  $\overline{F}$ , the *right half* is the subgraph induced by  $R$ , and the *left half* is the subgraph induced by  $\overline{R}$ . The *top-right quadrant* is the intersection of the top half and the right half; the *top-left quadrant*, *front-right quadrant*, etc. are defined analogously. Finally, the *top-right-front octant* is the intersection of the top, right, and front halves of  $G$ ; other octants are defined similarly.

We are finally ready to prove our lower bound on  $c_\infty(P_n \square P_n \square P_n)$ .

**Theorem 3.7.** *For sufficiently large  $n$ , we have  $c_\infty(P_n \square P_n \square P_n) > 0.7172n^2$ .*

*Proof.* We suppose for convenience that  $n$  is divisible by 10; the claim for  $n$  not divisible by 10 can be handled with a similar argument in conjunction with Theorem 3.1. Let  $G = P_n \square P_n \square P_n$ . We will give a strategy for the robber to evade  $\lfloor 0.7172n^2 \rfloor$  cops on  $G$ .

We begin with a high-level overview of the robber's strategy. Let  $v_0$  denote the robber's initial position, and for  $i \geq 1$ , let  $v_i$  denote the robber's position after his  $i$ th move. The robber will choose each  $v_i$  so as to satisfy the following properties:

- (1)  $v_i$  is in the large component of  $G$  before the cops'  $(i+1)$ st move.
- (2) No cop is adjacent to  $v_i$ .
- (3) No matter how the cops move in their  $(i+1)$ st turn,  $v_i$  will remain in the large component.

The robber will play as follows. To begin the game, he will look at the cops' initial positions and choose his own initial position  $v_0$  satisfying properties (1)-(3). Since  $v_0$  satisfies property (2), the cops cannot capture him on their first turn. After the cops move, the robber will choose  $v_1$ ; he aims to move to  $v_1$ , so we must show that he can actually reach  $v_1$  from  $v_0$ . Since  $v_0$  satisfied property (3) before the cops' turn, it must still belong to the large component of  $G$ . Since  $v_1$  satisfies property (1), it also belongs to the large component of  $G$ . Thus, since both  $v_0$  and  $v_1$  belong to the large component, there must be a cop-free path from  $v_0$  to  $v_1$ , hence the robber can reach  $v_1$ . Once again, since  $v_1$  satisfies property (2), the cops cannot capture the robber on their ensuing turn. The robber waits until the cops move, chooses  $v_2$ , moves there, and so forth. He can repeat this process, and thus evade capture, indefinitely.

Fix  $i \geq 0$ ; we explain how to choose  $v_i$ . Consider the state of the game immediately after the cops'  $i$ th turn (or, if  $i = 0$ , immediately after their initial placement). First, the robber identifies the sparsest half of  $G$ , which we denote  $H_i$ . Next, he identifies the sparsest quadrant of  $G$  contained within  $H_i$ ; denote this quadrant by  $Q_i$ . Finally, he identifies the sparsest octant of  $G$  contained within  $Q_i$ ; denote this octant by  $O_i$ .

Since  $O_i$  contains at most one-eighth of the total number of cops, it contains at most  $0.08965n^2$  cops. The  $n/2$  planes in  $O_i$  can be partitioned into  $n/10$  groups of five consecutive planes. One of these groups must contain at most  $0.8965n$  cops; let  $X_i$  denote the subgraph induced by some such group of planes.

The robber seeks a subgrid of  $O_i$  containing very few cops. Since

$$0.8965 \cdot n \cdot \frac{5}{\frac{n}{2}} = 8.965 < 9,$$

before the cops' turn,  $X_i$  must contain an  $n/2 \times 5 \times 5$  subgrid (that is, a subgrid isomorphic to  $P_{n/2} \square P_5 \square P_5$ ) containing fewer than nine cops. (Refer to Figure 2.) Let  $\hat{X}_i$  denote this subgrid. Provided that  $n$  is sufficiently large, within  $\hat{X}_i$  there must be a  $3 \times 5 \times 5$  subgrid  $\widetilde{X}_i$  that contains no cops. The subgraph  $\text{int}(\widetilde{X}_i)$  is (or contains) a  $3 \times 3 \times 1$  grid; the robber chooses  $v_i$  to be the vertex in the center of this grid.




FIGURE 2. Important subgraphs of  $O_i$ . From left:  $X_i$ ,  $\hat{X}_i$ , and  $\widetilde{X}_i$ .

Property (2) holds since all vertices adjacent to  $v_i$  belong to  $\widetilde{X}_i$ , which has no cops. We next verify properties (1) and (3). Note that property (3) actually implies property (1): if we can show that  $v_i$  must be in the large component of  $G$  after the cops'  $(i+1)$ st move no matter how they move, then  $v_i$  must have been in the large component of  $G$  before the cops'  $(i+1)$ st move, since all of the cops could have chosen to stay put on their  $(i+1)$ st move. We will now argue that (3), and therefore also (1), holds for  $v_i$ .

The remainder of the proof contains some rather technical details. To provide some intuition for the reader, we begin with a rough outline of what follows. First, we argue that  $v_i$  must belong to the large component of  $\text{int}(O_i)$ ; we do so by showing that a robber at  $v_i$  has access to more vertices of  $\text{int}(O_i)$  than the small component could possibly contain. We next argue that since the large component of  $\text{int}(O_i)$  is larger than the small component of  $\text{int}(Q_i)$ , the robber has access to more vertices than are in the small component of  $\text{int}(Q_i)$  and thus  $v_i$  must be in the large component of  $\text{int}(Q_i)$ . We use similar arguments to prove that  $v_i$  must belong to the large component of  $\text{int}(H_i)$  and, finally, to the large component of  $G$ .

We will now argue that  $v_i$  must belong to the large component of  $\text{int}(O_i)$ , or, equivalently, that a robber at  $v_i$  must be in the large component of  $\text{int}(O_i)$ . There were no cops in  $\widetilde{X}_i$  before the cops' move, so there can be no cops in  $\text{int}(\widetilde{X}_i)$  after the cops' move. Hence, the

robber can reach all vertices of  $\text{int}(\widetilde{X}_i)$  and can thus reach at least nine columns of  $\text{int}(\widehat{X}_i)$ . Likewise, there were at most eight cops in  $\widehat{X}_i$  before the cops' move, so there are at most eight cops in  $\text{int}(\widehat{X}_i)$  after the cops' move; thus one of the nine columns of  $\text{int}(\widehat{X}_i)$  that the robber can reach must be empty. Moreover, the robber can reach all but at most eight vertices in some column of the second plane of  $\text{int}(\widehat{X}_i)$ : either the empty column already belongs to the second plane, or it is directly above or below some column in the second plane, enabling the robber to reach any cop-free vertex in that column.

We aim to show that the robber is in the large component of  $\text{int}(O_i)$ . Toward this end, we first claim that he can reach an empty row or column within some plane  $P$  of  $\text{int}(O_i)$  that contains fewer than  $0.3n$  cops. There were at most  $0.8965n$  cops in  $X_i$  before the cops' turn, so there are at most  $0.8965n$  cops in  $\text{int}(X_i)$  after the cops' turn. Since there are at least three planes in  $\text{int}(X_i)$ , one of the first three planes must have fewer than  $0.3n$  cops; let  $P$  denote some such plane. We have already shown that the robber can reach an empty column in  $\text{int}(X_i)$ ; let  $P'$  denote the plane containing this column. If  $P' = P$ , then we are done. If instead  $P$  and  $P'$  are adjacent planes, then the robber can use the empty column in  $P'$  to reach an empty row in  $P$  and proceed from there to an empty column in  $P$ . Finally, suppose  $P$  and  $P'$  are not adjacent. It must be that  $P$  is either the first or third plane of  $\text{int}(\widehat{X}_i)$ . We argued above that the robber can reach all but at most eight vertices of some column in the second plane of  $\text{int}(\widehat{X}_i)$ ; using this column, he can reach all but at most eight rows of  $P$ . Since  $P$  has  $0.5n - 1$  columns and fewer than  $0.3n$  cops, it must have at least  $0.2n$  empty rows. If  $n$  is sufficiently large, then  $P$  has more than eight empty rows, so the robber can reach at least one, after which he may use this empty row to reach an empty column of  $P$ . In any case, the robber can reach an empty column of  $P$ , as claimed.

To prove that the robber must be in the large component of  $\text{int}(O_i)$ , we will show that the number of vertices of  $\text{int}(O_i)$  to which he has access exceeds the number of vertices in the small component of  $\text{int}(O_i)$ . As noted above,  $P$  must have at least  $0.2n$  empty rows and  $0.2n$  empty columns. The empty rows collectively contain at least  $0.1n^2 - 0.2n$  vertices, the empty columns also contain at least  $0.1n^2 - 0.2n$ , and their intersection contains at most  $0.04n^2$ . Consequently, the empty rows and columns collectively contain at least  $0.16n^2 - 0.4n$  vertices, and the robber can access all of these. Since  $\text{int}(O_i)$  contains at most  $0.08965n^2$  cops, no more than  $0.08965n^2$  shafts can be nonempty, so the robber has access (through  $P$ ) to at least  $0.07035n^2 - 0.4n$  empty shafts in  $\text{int}(O_i)$ . Each of these shafts contains  $n/2 - 1$  vertices, so together they contain at least  $(0.07035n^2 - 0.4n) \cdot (0.5n - 1)$  vertices.

Note that  $\text{int}(O_i)$  is a  $(n/2 - 1) \times (n/2 - 1) \times (n/2 - 1)$  grid. Hence, applying Lemma 3.6(d) with  $m = n/2 - 2$ , we have

$$C_m(n/2 - 1, n/2 - 1, n/2 - 1) = \left(\frac{1}{2} \cdot (1)^2 + o(1)\right) \cdot \left(\frac{n}{2} - 1\right)^2 = \left(\frac{1}{8} + o(1)\right) n^2$$

and

$$S_m(n/2 - 1, n/2 - 1, n/2 - 1) = \left(\frac{1}{6} \cdot (1)^3 + o(1)\right) \cdot \left(\frac{n}{2} - 1\right)^3 = \left(\frac{1}{48} + o(1)\right) n^3,$$

which is less than  $0.021n^3$  provided that  $n$  is sufficiently large. The number of cops in  $\text{int}(O_i)$  is at most  $0.08965n^2$ , which is smaller than  $C_m(n/2 - 1, n/2 - 1, n/2 - 1)$ . Thus, by Lemma 3.5, the small component of  $\text{int}(O_i)$  has fewer than  $0.021n^3$  vertices. The robber has access to more than  $0.021n^3$  vertices, so he cannot be in the small component of  $\text{int}(O_i)$ ; he must

therefore be in the large component of  $\text{int}(O_i)$ . For future reference, note also that the large component of  $\text{int}(O_i)$  must contain more than  $0.104n^3$  vertices (provided  $n$  is sufficiently large), since  $\text{int}(O_i)$  itself contains  $(n/2 - 1)^3$  vertices, of which fewer than  $0.021n^3$  belong to the small component and at most  $0.08965n^2$  are occupied by cops. Consequently, the robber can access more than  $0.104n^3$  vertices in  $\text{int}(O_i)$ .

We next argue that the robber must be in the large component of  $\text{int}(Q_i)$ . The number of cops in  $\text{int}(Q_i)$  is at most  $0.1793n^2$ , which is less than  $0.2n^2$ . Note that  $\text{int}(Q_i)$  is an  $n \times (n/2 - 1) \times (n/2 - 1)$  grid. Applying Lemma 3.6(b) using  $m = \frac{3}{4}(n - 1)$ , we have

$$C_m(n/2 - 1, n/2 - 1, n) = \left( -\frac{1}{2} \cdot \left(\frac{3}{4}\right)^2 + \frac{3}{4} - \frac{1}{4} + o(1) \right) n^2 = \left( \frac{7}{32} + o(1) \right) n^2,$$

which is greater than  $0.2n^2$  (provided  $n$  is sufficiently large). In addition,

$$S_m(n/2 - 1, n/2 - 1, n) = \left( -\frac{1}{6} \cdot \left(\frac{3}{4}\right)^3 + \frac{1}{2} \cdot \left(\frac{3}{4}\right)^2 - \frac{1}{4} \cdot \frac{3}{4} + \frac{1}{24} + o(1) \right) n^3 = \left( \frac{25}{384} + o(1) \right) n^3,$$

which is less than  $0.1n^3$  (once again, provided  $n$  is sufficiently large). As noted earlier, the robber can reach more than  $0.1n^3$  vertices in  $\text{int}(O_i)$ , and hence more than  $0.1n^3$  vertices in  $\text{int}(Q_i)$ ; thus, by Lemma 3.5, he must be in the large component of  $\text{int}(Q_i)$ . Note that  $\text{int}(Q_i)$  contains  $(\frac{1}{4} + o(1))n^3$  vertices, of which fewer than  $0.1n^3$  belong to the small component and only a quadratic number are occupied by cops; hence the large component of  $\text{int}(Q_i)$  contains more than  $0.15n^3$  vertices (provided  $n$  is sufficiently large).

A similar argument shows that the robber must be in the large component of  $\text{int}(H_i)$ . The number of cops in  $\text{int}(H_i)$  is at most  $0.3586n^2$ . By Lemma 3.6(c), using  $m = 0.96721(n - 1)$ , we have

$$C_m(n/2 - 1, n, n) = \left( \frac{1}{2} \cdot 0.96721 - \frac{1}{8} + o(1) \right) n^2 > 0.3586n^2$$

and

$$S_m(n/2 - 1, n, n) = \left( \frac{1}{4} \cdot 0.96721^2 - \frac{1}{8} \cdot 0.96721 + \frac{1}{48} + o(1) \right) n^3 < 0.133807n^3$$

(provided, as usual, that  $n$  is sufficiently large). The robber can reach more than  $0.133807n^3$  vertices of  $\text{int}(Q_i)$ , so he must be in the large component of  $\text{int}(H_i)$ . Of the  $(\frac{1}{2} + o(1))n^3$  vertices in  $\text{int}(H_i)$ , under  $0.133807n^3$  belong to the small component and at most a quadratic number are occupied by cops, so the large component contains at least  $0.366193n^3$  vertices (assuming once again that  $n$  is sufficiently large).

Finally, we show that the robber must be in the large component of  $G$ , which will complete the proof. By Lemma 3.6(d) with  $m = 1.3189(n - 1)$ , we have

$$C_m(n, n, n) = \left( -1.3189^2 + 3 \cdot 1.3189 - \frac{3}{2} + o(1) \right) n^2 > 0.7172n^2$$

and

$$S_m(n, n, n) = \left( -\frac{1}{3} \cdot 1.3189^3 + \frac{3}{2} \cdot 1.3189^2 - \frac{3}{2} \cdot 1.3189 + \frac{1}{2} + o(1) \right) n^3 < 0.36616n^3$$

(as always, provided that  $n$  is sufficiently large). The robber can reach at least  $0.366193n^3$  vertices in  $\text{int}(H_i)$ , and thus in  $G$ ; this is greater than the number of vertices in the small component of  $G$ . Hence the robber (and, by extension,  $v_i$ ) must be in the large component in  $G$ . We have thus shown that  $v_i$  satisfies properties (1) and (3), which completes the proof.  $\square$

**3.1. Four or more dimensions.** We conclude the section with a brief look at higher-dimensional grids. The ideas used to prove Theorem 3.2 extend naturally to higher dimensions, as we next demonstrate.

**Theorem 3.8.** *If  $G$  is the  $d$ -fold Cartesian product of copies of  $P_n$ , then*

$$c_\infty(G) \leq (1 + o(1)) \sum_{k=0}^{\lfloor d/2 \rfloor} (-1)^k \binom{d}{k} \binom{\lfloor d(n+1)/2 \rfloor - kn - 1}{d-1}.$$

*Proof.* As in the proof of Theorem 3.2, we suppose for the sake of simplicity that  $n$  is odd; a similar argument suffices when  $n$  is even. For nonnegative integers  $a$  and  $b$ , let  $S_{a,b}$  denote the set of vertices in the  $a$ -fold Cartesian product of copies of  $P_n$  whose coordinates sum to  $b$ . We claim that

$$|S_{a,b}| = \sum_{k=0}^{\lfloor a/2 \rfloor} (-1)^k \binom{a}{k} \binom{b - kn + a - 1}{a-1}.$$

$|S_{a,b}|$  counts the number of solutions to  $x_1 + x_2 + \cdots + x_a = b$  with each  $x_i$  in  $\{0, \dots, n-1\}$ . The number of nonnegative-integer solutions to  $x_1 + x_2 + \cdots + x_a = b$  is  $\binom{b+a-1}{a-1}$ ; however, we must exclude solutions for which some  $x_i$  exceeds  $n-1$ . For  $k \in \{1, \dots, \lfloor a/2 \rfloor\}$ , the number of solutions in which  $k$  different  $x_i$  exceed  $n-1$  is equivalent to the number of nonnegative-integer solutions to  $x_1 + x_2 + \cdots + x_a = b - kn$ , which is  $\binom{b-kn+a-1}{a-1}$ . Consequently, by the inclusion-exclusion principle,

$$|S_{a,b}| = \sum_{k=0}^{\lfloor a/2 \rfloor} (-1)^k \binom{a}{k} \binom{b - kn + a - 1}{a-1},$$

as claimed.

We are now ready to present the cops' strategy. Let  $m = \lfloor d(n-1)/2 \rfloor$ . We begin with  $|S_{d,m}| + |S_{d-1,m-n}|$  cops, of which  $|S_{d,m}|$  are designated *blocking cops* and the remainder are designated *reserve cops*. Initially, one blocking cop occupies each vertex of  $S_{d,m}$  and the reserve cops position themselves arbitrarily. Let  $v$  denote the robber's initial position. Let  $H$  denote the component of  $G - S_{d,m}$  consisting of all vertices whose coordinates sum to less than  $m$ , and suppose that the robber begins in  $H$  (a similar argument works if he begins in the other component of  $G - S_{d,m}$ ). Over the course of several turns, reserve cops move to occupy every vertex in  $H$  whose coordinates sum up to  $m-1$  and whose first coordinate is  $n-1$ . (Note that there are precisely  $|S_{d-1,m-n}|$  such vertices, so there are enough reserve cops to occupy all of them.) The robber must remain in  $H$  during this time, since the blocking cops separate  $H$  from  $G - H$ .

Once the reserve cops are in position, all blocking cops whose first coordinate is at least 1 move so as to decrease their first coordinate. The cops now occupy all vertices of  $S_{d,m-1}$ . All cops in  $S_{d,m-1}$  are now designated blocking cops, while the rest are designated reserve cops. The cops repeat this process: the reserve cops move to occupy vertices with first coordinate

$n - 1$  whose coordinates sum to  $m - 2$ , the blocking cops decrease their first coordinates, and so on. The size of the robber's component gradually decreases, until he is eventually captured.  $\square$

We remark that the techniques used to establish a lower bound on  $c_\infty(P_n \square P_n \square P_n)$  should also extend to the  $d$ -dimensional grid, but a general analysis seems difficult; rather, the argument would need to be specialized to each particular value of  $d$ .

#### 4. HYPERCUBES

We now turn our attention to the infinite-speed cop number of hypercubes. The  $n$ -dimensional *hypercube*  $Q_n$  is the graph whose vertex set consists of all  $n$ -bit binary strings, with two vertices adjacent if and only if the corresponding strings differ in exactly one bit. (Equivalently, it is the  $n$ -fold Cartesian product of copies of  $K_2$ .) Mehrabian [12] established the following bounds on  $c_\infty(Q_n)$ :

**Theorem 4.1** ([12], Theorem 9.1(e)). *For all  $n \geq 1$ , there exist positive constants  $\eta_1$  and  $\eta_2$  such that*

$$\eta_1 \cdot \frac{2^n}{n^{3/2}} \leq c_\infty(Q_n) \leq \eta_2 \cdot \frac{2^n}{n}.$$

The upper bound in Theorem 4.1 is straightforward: the infinite-speed cop number of a graph is bounded above by the domination number – since the cops can start on a dominating set and capture the robber with their first move – and it is well-known that  $\gamma(Q_n) \leq \frac{2^{n+1}}{n+1}$ . The lower bound, however, is more intricate and relies on a connection between the infinite-speed cop number of a graph and its treewidth. We obtain a stronger lower bound using a different approach.

Before proceeding, we state a technical lemma.

**Lemma 4.2.** *Let  $n$  be even, and suppose we play the infinite-speed robber game on  $Q_n$  with  $C$  cops. For sufficiently large  $n$ , if  $C \leq \frac{2^{n-3}}{n \ln n}$ , then the small component has fewer than  $\frac{1}{2} \binom{n}{n/2}$  vertices.*

*Proof.* It is a well-known consequence of Stirling's approximation that  $\binom{n}{n/2} \sim \frac{2^n}{\sqrt{n\pi/2}}$ ; we will make use of this fact below.

Let  $k = \frac{n}{2} - \frac{1}{2}\sqrt{n(\ln n + \ln \ln n)}$ . We claim that  $C \leq \binom{n}{k}$ . Stirling's approximation yields (see [17], p. 66, equation (5.41)):

$$\begin{aligned}
\binom{n}{k} &\sim \binom{n}{n/2} \exp\left(-(\sqrt{n(\ln n + \ln \ln n)})^2/(2n)\right) \\
&= \binom{n}{n/2} \exp\left(-\frac{1}{2}(\ln n + \ln \ln n)\right) \\
&= \binom{n}{n/2} \cdot \frac{1}{\sqrt{n \ln n}} \\
&\sim \frac{2^n}{\sqrt{n\pi/2}} \cdot \frac{1}{\sqrt{n \ln n}} \\
&= \frac{1}{\sqrt{\pi/2}} \cdot \frac{2^n}{n\sqrt{\ln n}}.
\end{aligned}$$

Thus  $C \leq \binom{n}{k}$  provided that  $n$  is sufficiently large. It now follows by Harper's theorem on vertex-isoperimetry in  $Q_n$  [9] that the size of the small component cannot exceed  $\sum_{i=0}^k \binom{n}{i}$ . We bound this sum using the Chernoff bound, in a form originally stated by Okamoto ([15], Theorem 1). Let  $X$  be a binomial random variable with parameters  $n$  and  $p = 1/2$ . Now

$$\begin{aligned}
\sum_{i=0}^k \binom{n}{i} &= 2^n \cdot P(X \leq k) \\
&= 2^n \cdot P\left(X \leq \left(\frac{1}{2} - \frac{1}{2}\sqrt{\frac{\ln n + \ln \ln n}{n}}\right)n\right) \\
&\leq 2^n \exp\left(-2\left(\frac{1}{2}\sqrt{\frac{\ln n + \ln \ln n}{n}}\right)^2 n\right) \\
&= 2^n \exp\left(-\frac{1}{2}(\ln n + \ln \ln n)\right) \\
&= \frac{2^n}{\sqrt{n \ln n}}.
\end{aligned}$$

In particular, the size of the small component is less than  $\frac{1}{2}\binom{n}{n/2}$  for sufficiently large  $n$ .  $\square$

We are now ready to establish a lower bound on  $c_\infty(Q_n)$ .

**Theorem 4.3.** *For sufficiently large  $n$ , we have  $c_\infty(Q_n) \geq \frac{2^{n-3}}{n \ln n}$ .*

*Proof.* We give a strategy for the robber to avoid  $\frac{2^{n-3}}{n \ln n} - 1$  or fewer cops. Intuitively, on each robber turn, the robber wants to move to a vertex with few enough cops nearby that the cops cannot cut off the robber's escape routes. To formalize the robber's strategy, we use a potential function. Suppose the robber currently occupies vertex  $v$ . For a cop  $c$  at vertex  $u$ , define that cop's *potential*  $P(c)$  as follows. If in fact  $u = v$ , meaning that the cop and robber occupy the same vertex, then  $P(c) = 1$ ; otherwise,  $P(c) = \binom{n}{\text{dist}(u,v)-1}^{-1}$ . Moreover, we define the *total potential*,  $\Phi$ , by  $\Phi = \sum_c P(c)$ , where the sum is taken over all cops  $c$ .

Note that any cop on or adjacent to the robber's vertex contributes 1 to the total potential. Hence, if  $\Phi < 1$  at the end of a robber turn, then there are no cops in the closed neighborhood of the robber's vertex, so he cannot be captured on the next cop turn. In general, the smaller the value of  $\Phi$ , the fewer cops there are "near" the robber; as such, the robber will aim to keep  $\Phi$  as small as possible. Before outlining the robber's strategy, we prove the following statement: if  $\Phi < 1/2$  at the end of a robber turn, then the robber's vertex belongs to the large component of  $Q_n$ , and it will continue to belong to the large component even after the next cop turn.

Suppose  $\Phi < 1/2$  at the end of a robber turn and suppose (by symmetry) that the robber currently occupies the vertex whose coordinates are all zero. We claim that this vertex is currently in the large component and remains in the large component after the cops take their turn. By Lemma 4.2, the size of the small component is strictly less than  $\frac{1}{2} \binom{n}{n/2}$ . Hence it suffices to show that, after the cops' move, the robber can reach at least this many vertices.

Let  $S$  denote the set of all vertices at distance  $n/2$  from the robber and let  $\mathcal{P}$  denote the collection of all shortest paths from the robber's vertex to a vertex in  $S$ . As the robber traverses such a path, each step corresponds to changing some coordinate of his position from 0 to 1. Consequently, each path in  $\mathcal{P}$  corresponds to changing  $n/2$  coordinates from 0 to 1, in some order; there are  $\binom{n}{n/2}$  ways to decide which coordinates to change and  $(n/2)!$  possible orders in which to change them, so

$$|\mathcal{P}| = \binom{n}{n/2} \cdot (n/2)! = \frac{n!}{(n/2)!(n/2)!} \cdot (n/2)! = \frac{n!}{(n/2)!}.$$

Not all of these paths can actually be traversed by the robber, because they may contain one or more vertices occupied by cops. Say that a path in  $\mathcal{P}$  is *blocked* by a cop if that cop occupies some vertex on the path. We claim that if  $\Phi < 1/2$  at the end of the robber's turn, then fewer than half of the paths will be blocked after the cops' ensuing turn. To see this, first note that a cop at distance  $k$  from the robber's vertex (where  $k \leq n/2$ ) blocks exactly  $k! \cdot \binom{n-k}{n/2-k} \cdot (n/2 - k)!$  paths in  $\mathcal{P}$ : there are  $k!$  shortest paths from the robber's vertex to the cop's, and there are  $\binom{n-k}{n/2-k} \cdot (n/2 - k)!$  shortest paths from the cop's vertex to a vertex in  $S$ . This quantity simplifies:

$$k! \cdot \binom{n-k}{n/2-k} \cdot (n/2 - k)! = k! \cdot \frac{(n-k)!}{(n/2-k)!(n/2)!} \cdot (n/2 - k)! = \frac{k!(n-k)!}{(n/2)!}.$$

The total number of paths blocked by the cops is at most the sum of this quantity over all cops. For a cop  $c$ , let  $d(c)$  denote that cop's distance from the robber's vertex. We now have

$$\begin{aligned}
\text{Total paths blocked} &\leq \sum_c \frac{(d(c))!(n-d(c))!}{(n/2)!} \\
&= \sum_c \frac{n!}{n!} \cdot \frac{(d(c))!(n-d(c))!}{(n/2)!} \\
&= \frac{n!}{(n/2)!} \sum_c \frac{1}{\binom{n}{d(c)}} \\
&\leq \frac{n!}{(n/2)!} \sum_c P(c) \\
&= \frac{n!}{(n/2)!} \cdot \Phi \\
&< \frac{n!}{(n/2)!} \cdot \frac{1}{2},
\end{aligned}$$

where the inequality on the fourth line follows because a cop that is currently at distance  $d(c)$  was at distance  $d(c) + 1$  or less prior to the cops' turn and thus contributed at least  $\binom{n}{d(c)+1-1}^{-1}$  to  $\Phi$ .

Consequently, at least half of the shortest paths from the robber's vertex to  $S$  are unblocked, so the robber can reach at least half of the vertices in  $S$ . Since  $|S| = \binom{n}{n/2}$ , the robber cannot be in the small component (which, as mentioned above, has size less than  $\frac{1}{2} \binom{n}{n/2}$ ).

We next claim that on each robber turn, if the robber is within the large component of  $Q_n$ , then he can move so as to make the total potential less than  $1/2$ . Let  $\Phi_v$  denote the value of the total potential when the robber moves to  $v$ ; we aim to show that  $\Phi_v < 1/2$  for some vertex  $v$  in the large component. We begin by computing  $\sum_{v \in V(G)} \Phi_v$ . For each cop  $c$  and for all  $k \in \{0, 1, \dots, n\}$ , there are  $\binom{n}{k}$  vertices at distance  $k$  from  $c$ ;  $c$  exerts a potential of  $\binom{n}{k-1}^{-1}$  on each such vertex provided  $k \geq 1$  and a potential of 1 otherwise. Consequently, between all of the vertices in the graph, the cop exerts an aggregate potential of

$$\begin{aligned}
1 + \sum_{k=1}^n \frac{\binom{n}{k}}{\binom{n}{k-1}} &= 1 + \sum_{k=1}^n \frac{n-k+1}{k} \\
&= 1 + \sum_{k=1}^n \left( \frac{n+1}{k} - 1 \right) \\
&= 1 - n + (n+1) \sum_{k=1}^n \frac{1}{k} \\
&\leq 1 - n + (n+1)(\ln n + 1) \\
&\leq 2n \ln n \quad (\text{provided } n \geq 3).
\end{aligned}$$

Since there are  $\frac{2^{n-3}}{n \ln n}$  cops, the aggregate potential exerted by all cops is at most

$$\frac{2^{n-3}}{n \ln n} \cdot 2n \ln n = 2^{n-2}.$$

Consequently, there are at most  $2^{n-1}$  vertices  $u$  such that  $\Phi_u \geq 1/2$ , so there are at least  $2^{n-1}$  vertices  $v$  such that  $\Phi_v < 1/2$ . Since the large component contains more than  $2^{n-1}$  vertices, we have  $\Phi_{v^*} < 1/2$  for at least one vertex  $v^*$  in the large component. Since the robber is currently in the large component, there is a cop-free path from his current vertex to  $v^*$ ; consequently, the robber may move to  $v^*$ .

Finally, we explain the robber's actual strategy. Initially, the robber begins at any vertex  $v$  such that  $\Phi_v < 1/2$ . Note that the cops cannot capture the robber on their ensuing turn, since any cop within distance 1 of the robber would contribute 1 to the potential, contradicting  $\Phi_v < 1/2$ . Moreover, since the potential is less than  $1/2$ , the robber will be in the large component after the cops' move. On the robber's next turn, he moves to a vertex  $v^*$  such that  $\Phi_{v^*} < 1/2$ . He then repeats this process indefinitely, repeatedly moving so as to keep the potential below  $1/2$ .  $\square$

## REFERENCES

- [1] M. Aigner and M. Fromme, A game of cops and robbers, *Discrete Appl. Math.* 8 (1984), 1–12.
- [2] N. Alon and A. Mehrabian, On a generalization of Meyniel's conjecture on the Cops and Robbers game, *Electron. J. Combin.* 18 (2011), no. 1, Paper 19, 7 pp.
- [3] N. Alon, A. Mehrabian, Chasing a fast robber on planar graphs and random graphs, *J. Graph Theory* 78 (2015), no. 2, 81–96.
- [4] B. Bollobás and I. Leader, Compressions and isoperimetric inequalities, *J. Combin Theory Ser. A* 56 (1991), 47–62.
- [5] A. Bonato and R.J. Nowakowski, *The Game of Cops and Robbers on Graphs*, American Mathematical Society, Providence, Rhode Island, 2011.
- [6] D. Dereniowski, T. Gavenčiak, and J. Kratochvíl, Cops, a fast robber and defensive domination on interval graphs, *Theor. Comp. Sci.* 794 (2019), 47–58.
- [7] F. Fomin, P. Golovach, J. Kratochvíl, and K. Suchan, Pursuing a fast robber on a graph, *Theor. Comput. Sci.* 411 (2010), 1167–1181.
- [8] A. Frieze, M. Krivilevich, and P. Loh, Variations on cops and robbers, *J. Graph Theory* 69 (2012), no. 4, 383–402.
- [9] L. Harper, Optimal numberings and isoperimetric problems on graphs, *J. Comb. Theory* 1 (1966), no. 3, 385–393.
- [10] M. Maamoun and H. Meyniel, On a game of policemen and robber, *Discrete Appl. Math.* 17 (1987), 307–309.
- [11] A. Mehrabian, Lower bounds for the cop number when the robber is fast, *Combin. Probab. Comput.* 20 (2011), no. 4, 617–621.
- [12] A. Mehrabian, Cops and robber game with a fast robber on expander graphs and random graphs, *Ann. Comb.* 16 (2012), no. 4, 829–846.
- [13] A. Mehrabian, The fast robber on interval and chordal graphs, *Discrete Appl. Math.* 180 (2015), 188–193.
- [14] R.J. Nowakowski and P. Winkler, Vertex-to-vertex pursuit in a graph, *Discrete Math.* 43 (1983), 235–239.
- [15] M. Okamoto, Some inequalities relating to the partial sum of binomial probabilities, *Ann Inst Stat Math* 10 (1959), 29–35.
- [16] A. Quilliot, Jeux et pointes fixes sur les graphes, Thèse de 3ème cycle, Université de Paris VI, 1978, 131–145.
- [17] J. Spencer, L. Florescu, *Asymptopia*, Student mathematical library 71, AMS, Providence, RI, 2014.

- [18] P. D. Seymour and R. Thomas, Graph searching and a min-max theorem for treewidth, *J. Combin. Theory Ser. B* 58 (1993), no. 1, 22–33.
- [19] D.B. West, *Introduction to Graph Theory, 2nd edition*, Prentice Hall, 2001.

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICAL SCIENCES, UNIVERSITY OF RHODE ISLAND, UNIVERSITY OF RHODE ISLAND, KINGSTON, RI,USA, 02881  
*Email address:* billk@uri.edu

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICAL SCIENCES, UNIVERSITY OF RHODE ISLAND, UNIVERSITY OF RHODE ISLAND, KINGSTON, RI,USA, 02881  
*Email address:* townsendn@uri.edu