

Time-Slicing Path-integral in Curved Space

Mingnan Ding^{1*} and Xiangjun Xing^{1,2,3†}

¹ *Wilczek Quantum Center, School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai, 200240 China*

² *T.D. Lee Institute, Shanghai Jiao Tong University, Shanghai, 200240 China*

³ *Shanghai Research Center for Quantum Sciences, Shanghai 201315 China*

(Dated: January 13, 2022)

Path integrals constitute powerful representations for both quantum and stochastic dynamics. Yet despite many decades of intensive studies, there is no consensus on how to formulate them for dynamics in curved space, or how to make them covariant with respect to nonlinear transform of variables. In this work, we construct rigorous and covariant formulations of time-slicing path integrals for quantum and classical stochastic dynamics in curved space. We first establish a rigorous criterion for correct *time-slice actions* of path integrals (Lemma 1). This implies the existence of infinitely many equivalent representations for time-slicing path integral. We then show that, for any dynamics with second order generator, all time-slice actions are asymptotically equivalent to a Gaussian (Lemma 2). Using these results, we further construct a continuous family of equivalent actions parameterized by an interpolation parameter $\alpha \in [0, 1]$ (Lemma 3). The action generically contains a *spurious drift* term linear in $\Delta \mathbf{x}$, whose concrete form depends on α . Finally we also establish the covariance of our path-integral formalism, by demonstrating how the action transforms under nonlinear transform of variables. The $\alpha = 0$ representation of time-slice action is particularly convenient because it is Gaussian and invariant, as long as $\Delta \mathbf{x}$ transforms according to *Ito's formula*.

I. INTRODUCTION

Path integral as a representation of quantum mechanics was first envisaged by Dirac in his renowned book [1], and was developed systematically by Feynman [2, 3] in 1948. Since then, through the hands of many outstanding physicists and mathematicians, it has been transformed into the arguably most powerful tool for theoretical physics [4]. The applications of path integral methods range from quantum mechanics [5] to quantum field theory [6], quantum open systems [7, 8], and quantum gravity [9], from Brownian motion to general classical stochastic processes [10, 11], as well as polymer physics [12], and even financial study [13, 14]. The formalism of path integral not only help shaping our intuition about quantum and classical fluctuations, but also played a key role in the synthesis of quantum field theory with statistical field theory in the last century.

Consider a 1d quantum particle with Hamiltonian $\hat{H} = \hat{p}^2/2m + V(x)$. The Green's function is defined by

$$\begin{aligned} i\hbar \frac{\partial}{\partial \tau} G(x, \tau | x_0, 0) &= \hat{H} G(x, \tau | x_0, 0), \\ G(x, 0 | x_0, 0) &= \delta(x - x_0). \end{aligned} \quad (1)$$

The path-integral representation of $G(x, \tau | x_0, 0)$ is

$$G(x, \tau | x_0, 0) = \int_{(x_0, 0)}^{(x, \tau)} Dx(\tau) e^{\frac{i}{\hbar} S_{\text{cl}}[x(\tau)]}, \quad (2)$$

where the integral means summation over all paths with initial condition $x(0) = x_0$ and final condition $x(\tau) = x$,

and $S_{\text{cl}}[x(\tau)]$ is the classical action:

$$S_{\text{cl}}[x(\tau)] = \int_0^\tau ds L(x, \dot{x}) = \int_0^\tau ds \left[\frac{m^2}{2} \dot{x}^2 - V(x) \right]. \quad (3)$$

One must specify how to sum over paths in order to assign a precise meaning to the path integral. Feynman approximated the original path $x(\tau)$ by a sequence of straight-line segments that pass through $\{x_k = x(k\Delta\tau), k = 0, 1, \dots, N\}$, with $\Delta\tau = \tau/N$. This leads to approximation of the action Eq. (3) as a discrete sum:

$$S_{\text{cl}}[x(\tau)] \approx \sum_k \Delta\tau \left[\frac{m}{2} \left(\frac{x_{k+1} - x_k}{\Delta\tau} \right)^2 - V(\bar{x}_k) \right], \quad (4)$$

which turns into a Riemann integration in the limit $\Delta\tau \rightarrow 0$. Here \bar{x}_k is an arbitrary point between x_k and x_{k+1} , whose choices have vanishing influence in the limit $\Delta\tau \rightarrow 0$, as long as both $V(x)$ and the path $x(\tau)$ are Riemann integrable. The path-integral (2) is then transformed into integration over $N-1$ coordinates $\{x_k, k = 1, \dots, N-1\}$:

$$C \int \prod_{k=1}^{N-1} dx_k \exp \frac{i\Delta\tau}{\hbar} \sum_k \left[\frac{m}{2} \left(\frac{x_{k+1} - x_k}{\Delta\tau} \right)^2 - V(\bar{x}_k) \right], \quad (5)$$

where the normalization constant C can be fixed by a reference problem, e.g., that of a free particle. Feynman carried out explicit calculations of Eq. (5) for free particle and harmonic oscillator, and obtained results consistent with direct solutions to Schrödinger equations.

The prescription of calculating path integral using Eqs. (4) and (5) is usually called *time-slicing*, and is one of many possible *regularization schemes* which make the path integral Eq. (2) finite and calculable. In field theories, other regularization schemes such as momentum

*Electronic address: dmnphy@sjtu.edu.cn

†Electronic address: xxing@sjtu.edu.cn

cut-off or dimensional regularization are more popular. We shall not discuss them in this work.

Because of the imaginary nature of the exponent in Eq. (5), the integral is not convergent in the usual sense. To improve the convergence, people often analytically continue the time into the imaginary axis, $\tau = -it$, and transform Eqs. (1) into:

$$-\hbar \frac{\partial}{\partial t} G(x, t|x_0, 0) = \hat{H}G(x, t|x_0, 0), \quad (6a)$$

$$G(x, 0|x_0, 0) = \delta(x - x_0). \quad (6b)$$

The integral Eq. (5) then becomes

$$C \int \prod_{k=1}^{N-1} dx_k \exp - \sum_k A(x_{k+1}, x_k; \Delta t), \quad (7a)$$

where $A(x_{k+1}, x_k; \Delta t)$, which shall be called the *time-slice action*, is given by

$$A(x_{k+1}, x_k; \Delta t) = \frac{\Delta t}{\hbar} \left[\frac{m}{2} \left(\frac{x_{k+1} - x_k}{\Delta t} \right)^2 + V(\bar{x}_k) \right] \quad (7b)$$

For $V(x) = 0$, Eqs. (6) reduce to the classical diffusion equation, and Eqs. (7) defines the celebrated *Wiener measure* [15] in the path space of classical Brownian motion. A mathematically rigorous theory of path integral Eqs. (7) as a representation of Eqs. (6) was established by Kac [16]. It is well known that in classical Brownian motion $\Delta x \sim \Delta t^{1/2}$, and hence typical paths are continuous but non-differentiable everywhere [17]. Using this property, one easily sees that the choice of \bar{x}_k in Eq. (4) influences the action by an amount scaling as $\Delta t^{1/2}$, which vanishes in the continuum limit. Time-slicing path integral formulation of classical Langevin dynamics was developed by Onsager and Machlup [19, 20].

There are however many dynamic processes, either quantum or classical, happening in curved spaces. There are also processes happening in Euclidean space but are better formulated using curvilinear coordinates, as well as stochastic processes with multiplicative noises, which resemble very much dynamics in curved space. For these problems, it has been known for long time that the above time-slicing prescription of path integral breaks down. Consider for example a quantum particle moving in a curved manifold. The covariant classical action and covariant quantum Hamiltonian are respectively

$$S_{\text{cl}}[\mathbf{x}(t)] = \int dt \left[\frac{m}{2} g_{ij}(\mathbf{x}) \dot{x}^i \dot{x}^j - V(\mathbf{x}) \right], \quad (8)$$

$$\hat{H}(\mathbf{x}) = -\frac{\hbar^2}{2m\sqrt{g}} \frac{\partial}{\partial x^i} \sqrt{g} g^{ij} \frac{\partial}{\partial x^j} + V(\mathbf{x}), \quad (9)$$

where g^{ij} is the metric inverse of g_{ij} , and $g = (\det g_{ij}) = (\det g^{ij})^{-1}$ [64]. The Green's function then is defined by:

$$-\hbar \frac{\partial}{\partial t} G(\mathbf{x}, t|\mathbf{x}_0, 0) = \hat{H}(\mathbf{x}) G(\mathbf{x}, t|\mathbf{x}_0, 0), \quad (10a)$$

$$G(\mathbf{x}, 0|\mathbf{x}_0, 0) = \delta(\mathbf{x}, \mathbf{x}_0), \quad (10b)$$

where $\delta(\mathbf{x}, \mathbf{x}_0)$ is a covariant Dirac delta function, to be defined below in Eq. (17).

In many previous works, people tried to establish the following path integral representation for Eq. (10):

$$G(\mathbf{x}, t|\mathbf{x}_0, 0) \sim C \int \prod_{k=1}^{N-1} d\mu(x_k) \quad (11)$$

$$\exp - \frac{\Delta t}{\hbar} \sum_k \left[\frac{m}{2} g_{ij}(\bar{\mathbf{x}}_k) \frac{\Delta x_k^i \Delta x_k^j}{\Delta t^2} - V(\bar{\mathbf{x}}_k) - \delta V(\bar{\mathbf{x}}_k) \right],$$

where $\Delta \mathbf{x}_k \equiv \mathbf{x}_{k+1} - \mathbf{x}_k$, and $\delta V(\bar{\mathbf{x}}_k)$ is usually called the extra term [21–24]. Note, however, unlike the case of Euclidean space, the choice of $\bar{\mathbf{x}}_k$ in $g_{ij}(\bar{\mathbf{x}}_k)$ does matter. Typical variation of $\bar{\mathbf{x}}_k$ scales as $\Delta \mathbf{x}$ and leads to correction of the classical action estimated as follows

$$\Delta t \sum_k \frac{\Delta \mathbf{x}^3}{\Delta t^2} \sim \sum_k \sqrt{\Delta t} \sim \Delta t^{-1/2}, \quad (12)$$

which diverges in the continuum limit. Such a correction cannot be compensated by $\sum_k \delta V(\bar{\mathbf{x}}_k) \Delta t$, which scales as $O(1)$. Common choices for $\bar{\mathbf{x}}_k$ are $\bar{\mathbf{x}}_k = (1 - \alpha)\mathbf{x}_k + \alpha\mathbf{x}_{k+1}$, where $\alpha \in [0, 1]$ shall be called the *interpolation parameter* in this work. Many theories with $\alpha = 0$ (pre-point), $\alpha = 1/2$ (mid-point), and $\alpha = 1$ (post-point) have been worked out, yet the results often do not agree with each other.

In fact, the most serious mistake in Eq. (11) is that the action misses a part linear in $\Delta \mathbf{x}_k$, summation of which also scales as $\Delta t^{-1/2}$. This term leads to non-vanishing average of $\Delta \mathbf{x}_k$, and shall be called *quantum spurious drift*, for reasons that will become clear below. We will also see that, while the precise form of quantum spurious drift depends on the particular choice of α , for a generic multi-dimensional model, it can not be cancelled by a judicious choice of coordinate systems, or by tuning of α .

Similar difficulties also arise in path integral representation of classical Markov dynamics in curved space, which can be described either by the covariant Langevin equation [25, 26]

$$dx^i + \left(L^{ij} \partial_j U - \frac{1}{\sqrt{g}} \partial_i \sqrt{g} L^{ij} \right) dt = b^{i\mu} dW_\mu(t), \quad (13)$$

or by the covariant Fokker-Planck equation [25]:

$$\partial_t p = \hat{L}_{\text{FP}} p = \frac{1}{\sqrt{g}} \partial_i \sqrt{g} L^{ij} (\partial_j + (\partial_j U)) p, \quad (14)$$

where $p = p(\mathbf{x}, t)$ is the invariant pdf of slow variables, and $b^{i\mu} b^{j\mu} = L^{ij} + L^{ji}$. The term $-\frac{1}{\sqrt{g}} \partial_i \sqrt{g} L^{ij}$ is called the *spurious drift*, and has been the target of intensive study for long time [18, 26, 45, 53, 58].

Note that Eq. (14) has the same form as the Schrödinger equation (10) in imaginary time, with the Fokker-Planck operator

$$\hat{L}_{\text{FP}} = \frac{1}{\sqrt{g}} \partial_i \sqrt{g} L^{ij} (\partial_j + (\partial_j U)), \quad (15)$$

playing the role of negative Hamiltonian. Note however \bar{L}_{FP} is generically non-Hermitian. If the space is curved or the coordinates are curvilinear (both imply that g_{ij} depends on \mathbf{x}), or if the noise-amplitudes $b^{i\mu}$ are state-dependent (multiplicative noises), then the same difficulty discussed above also show up, and the proper path integral representation is still under debate.

Earlier studies of path integral in curved space and in curvilinear coordinates were carried out by Dewitt [27] and by Edwards and Gulyaev [28]. Edwards and Gulyaev argued that the action of path integral is not invariant under usual rules of calculus when making nonlinear variable transformation, yet did not supply sufficient details. Since then, many authors tried to construct path integral representations for quantum and/or classical problems in curved space [29–42], or to study Langevin dynamics with multiplicative noises [43–49], or to understand how path integral should transform under nonlinear transformation of variables [50–52]. In spite of the large number of papers published on the subjects, however, there is still no sign of convergence on opinions. One school of researchers tried to solve the problems by introducing an extra term, as shown in Eq. (11) [21–24]. Many authors still hold the opinion that actions of path integrals are invariant, or at least can be made invariant via possible revision of time-slicing scheme. See, for example, Refs. [50–52] for the most recent effort on one dimensional case. Comparison of previous works are difficult because of the diversity of methods and conventions used, and also because of the lack of mathematical details. More seriously, many authors used Ito’s rule $dW(t)dW(t) = dt$ freely in the derivation of action, which is very dangerous. (See Sec.V D for an illustration of mistakes resulting from this application of Ito’s rule.)

The approach we shall take in this work is markedly different from all previous works. Using the method of asymptotic analysis, we shall first establish three lemmas with reasonable degree of mathematical rigor. These lemmas constitute the foundations of all our results. With Lemma 1, we establish the criterion for the correctness of time-slicing path integrals. This lemma also explains why there are so many seemingly different but equivalent representations of time-slicing path integrals. With Lemma 2, we prove that there is a Gaussian representation of time-slicing path integrals. With Lemma 3, we explicitly construct a one-parameter family of equivalent representations for time-slicing path integrals, parameterized by $\alpha \in [0, 1]$. The time-slicing path integral, whether it is for the quantum or for the classical stochastic problem, can then be obtained by straightforward application of these three lemmas. The action given in Eq. (26) is the most general form, which agrees with previous works [53–55] under the specific choices of interpolation parameters.

For classical Markov processes, we shall further study the connection between time-slicing path integral representation and Langevin dynamics representation. As is well-known, there are infinite number of equivalent representations for a given Langevin dynamics, each param-




FIG. 1: Time-slicing: approximation of continuous path $x(t)$ by piecewise linear path. An intermediate point \mathbf{x}_α needs to be chosen between \mathbf{x}_0 and \mathbf{x}_1 to evaluate the potential $V(\mathbf{x})$.

eterized by another parameter $\bar{\alpha}$, which specifies where the noise amplitudes are evaluated. The case of Ito representation ($\bar{\alpha} = 0$) is particularly simple, because it implies a linear relation between rate of slow variables and noises, conditioned on the slow variables at earlier time. Exploiting equivalence between different representations of Langevin dynamics, we shall derive a two-parameter family of equivalent representations for time-slicing path integral.

The last major issue we shall address is the covariance property. Our path integral formalism is constructed without making any assumption on the coordinate system, and the action is formulated in terms of tensor objects whose transformations under nonlinear transform of coordinates are well understood [65]. (As a comparison, the main results of Ref. [56] are established only for normal coordinates.) Hence our theory is covariant in the sense that all results are applicable to genetic coordinate systems, and transformation between different coordinate systems are well-defined. Nonetheless, there is an (often confusing) issue of transformation rule of the action of path integral. In contrast with the classical action (8), we shall find that the time-slice action of path integral is generically not invariant under usual rules of calculus. The invariance of action is spoiled both by the integration measure and, more importantly, by the non-differentiable nature of typical paths that dominate path integrals. More importantly, we shall explicitly show that the $\alpha = 0$ time-slice action can be made invariant if we replace the usual chain rule by Ito’s formula, which is more natural in view of the non-differentiable nature of typical paths.

The remaining of this work is organized as follows. In Section II, we establish three Lemmas, which constitute the base of all later results. In Section III, we apply three lemmas to obtain a continuous family of equivalent representations for time-slicing path integral. We also revisit the problem of Edwards and Gulyaev and discuss the geometric origin of spurious drift. In Section IV, we study the connection between classical nonlinear Langevin dynamics and time-slicing path integral, and obtain a more

general family of equivalent path integral representations for classical Markov processes. In Section V, we discuss the covariance property of time-slicing path integral. Finally, in Section VI, we draw the concluding remark and outline future research directions. In Appendix A, we present the details of proof for Lemma 3.

II. THREE LEMMAS ABOUT TSGFS

Consider a general Riemannian manifold where the quantum dynamics takes place. Let $\mathbf{x} = \{x^1, x^2, \dots, x^d\}$ be the coordinate system, $g_{ij}(\mathbf{x})$ the metric tensor, $g(\mathbf{x}) = \det(g_{ij}(\mathbf{x}))$ its determinant. We further define the volume measure $d\mu(\mathbf{x})$ and the invariant volume measure $dv(\mathbf{x})$ as

$$d\mu(\mathbf{x}) \equiv dx^1 dx^2 \dots dx^d, \quad (16a)$$

$$dv(\mathbf{x}) \equiv \sqrt{g(\mathbf{x})} d\mu(\mathbf{x}), \quad (16b)$$

as well as the covariant Dirac delta function as

$$\delta(\mathbf{x}, \mathbf{x}_0) \equiv \frac{1}{\sqrt{g(\mathbf{x}_0)}} \prod_{i=1}^d \delta(x^i - x_0^i) = \delta(\mathbf{x}_0, \mathbf{x}), \quad (17)$$

where $\delta(x^i - x_0^i)$ is the usual 1d Dirac delta function. For an arbitrary function $f(\mathbf{x})$ on the manifold, we have

$$\int f(\mathbf{x}) \delta(\mathbf{x}, \mathbf{x}_0) dv(\mathbf{x}) = f(\mathbf{x}_0). \quad (18)$$

We consider dynamics generated by a general second order partial differential operator

$$\hat{L}(\mathbf{x}) = \partial_i \partial_j D^{ij}(\mathbf{x}) - \partial_i F^i(\mathbf{x}) + \Phi(\mathbf{x}), \quad (19)$$

where repeated indices are summed over, $D^{ij}(\mathbf{x}) = D^{ji}(\mathbf{x})$ are symmetric and non-singular at all \mathbf{x} , and $\partial_i = \partial/\partial x^i$ act on everything to the right. We do not show time-dependence of $D^{ij}(\mathbf{x})$, $F^i(\mathbf{x})$, $\Phi(\mathbf{x})$, even though our theory is valid if these functions depend on time explicitly. We are interested in the Green's function:

$$G(\mathbf{x}, t|\mathbf{x}_0; 0) = e^{\hat{L}(\mathbf{x})t} \delta(\mathbf{x}, \mathbf{x}_0), \quad (20)$$

which is also known as propagator, transition amplitude (quantum mechanics), and transition probability (classical stochastic processes) etc.

We cut the time interval $[0, t]$ into N slices with duration $\Delta t = t/N$, and introduce the time-slice Green's function (TSGF):

$$G(\mathbf{x}|\mathbf{x}_0; \Delta t) = e^{\hat{L}(\mathbf{x})\Delta t} \delta(\mathbf{x}, \mathbf{x}_0), \quad (21)$$

with $\Delta t = t/N$. Eq. (20) can be rewritten into

$$G(\mathbf{x}, t|\mathbf{x}_0; 0) = \int dv_{N-1} \dots dv_1 G(\mathbf{x}_N|\mathbf{x}_{N-1}; \Delta t) \dots G(\mathbf{x}_2|\mathbf{x}_1; \Delta t) G(\mathbf{x}_1|\mathbf{x}_0; \Delta t), \quad (22)$$

where we have defined $\mathbf{x}_N = \mathbf{x}$, and $dv_k = dv(\mathbf{x}_k)$. The discretized paths $\mathbf{x}_N, \mathbf{x}_{N-1}, \dots, \mathbf{x}_1, \mathbf{x}_0$ formed the cylinder set [17], which can be used to construct rigorously the sigma algebra of the path space. To construct path integral representation of the dynamics, we make Δt infinitesimal, and express TSGF $G(\mathbf{x}|\mathbf{x}_0; \Delta t)$ in an exponential form. In the limit $\Delta t \rightarrow 0$, Eq. (22) becomes the time-slicing path integral.

For classical stochastic process, $G(\mathbf{x}|\mathbf{x}_0; \Delta t)dv(\mathbf{x})$ can be understood as a probability distribution function for variables \mathbf{x} , with \mathbf{x}_0 and Δt playing the roles of parameters. Integral of $G(\mathbf{x}|\mathbf{x}_0; \Delta t)dv(\mathbf{x})$ yields unity, as guaranteed by probability conservation. For quantum mechanics, $G(\mathbf{x}|\mathbf{x}_0; \Delta t)$ is generally not normalized.

In this section, we will establish three general properties of this TSGF, which will be the base of all later results. What we need is to make the RHS of Eq. (22) a faithful representation of the LHS in the limit $\Delta t \rightarrow 0$. We first note that the TSGF $G(\mathbf{x}|\mathbf{x}_0; \Delta t)$ is fully characterized by its moments, which can be expanded in terms of Δt . For this purpose, we do not need to calculate the TSGF $G(\mathbf{x}|\mathbf{x}_0, \Delta t)$ rigorously, but only need to calculate it such that all moments of $\mathbf{x} - \mathbf{x}_0$ are correct up to the order of Δt . Hence we obtain the following Lemma:

Lemma 1 *In order for the time-slicing path integral, i.e., the LHS of Eq. (22), to generate the dynamics defined by Eq. (20), we only need to calculate the TSGFs $G(\mathbf{x}_k|\mathbf{x}_{k-1}, \Delta t)$ such that all moments of $\mathbf{x}_k - \mathbf{x}_{k-1}$ are correct up to Δt . Higher order terms do not influence the dynamics in the continuum limit.*

According to Lemma 1, if two representations of TSGF yield equal moments up to order Δt , they generate the same continuum dynamics in the limit $\Delta t \rightarrow 0$, and hence are equivalent for the purpose of path integral representation. The validity of this Lemma is self-evident. But its importance can hardly be overrated. Not only it explains why there are many equivalent path integral representations for the same dynamics, it also provides a rigorous and operational criterion for proving equivalence (or inequivalence) between different representations.

Let us calculate the moments of $G(\mathbf{x}|\mathbf{x}_0; \Delta t)$ explicitly. Let $M = i_1 + \dots + i_m$, then a M -th order moment can be written as

$$\begin{aligned} & \langle \Delta x^{i_1} \dots \Delta x^{i_m} \rangle \\ &= \int dv(\mathbf{x}) \Delta x^{i_1} \dots \Delta x^{i_m} G(\mathbf{x}|\mathbf{x}_0; \Delta t) \\ &= \int dv(\mathbf{x}) \Delta x^{i_1} \dots \Delta x^{i_m} e^{\hat{L}(\mathbf{x})\Delta t} \delta(\mathbf{x}, \mathbf{x}_0) \\ &= \int dv(\mathbf{x}) \Delta x^{i_1} \dots \Delta x^{i_m} \left[1 + \Delta t \hat{L}(\mathbf{x}) + \dots \right] \delta(\mathbf{x}, \mathbf{x}_0). \end{aligned} \quad (23)$$

For all $M > 0$, the term zero-th order in Δt in the above expansion vanishes because of the delta function. The term first order in Δt can be calculated via integration by

parts. Because $\hat{L}(\mathbf{x})$ is a second order differential operator, we see that for all $M \geq 3$, the term first order in Δt also vanishes, again because of the delta function. Hence we conclude that all M -th order moments of $G(\mathbf{x}|\mathbf{x}_0; \Delta t)$ with $M \geq 3$ are at least of order of Δt^2 . But according to Lemma 1, these moments do not influence the continuum dynamics, and we might as well set them to zero, or anything as long as they are at least of order Δt^2 . But a distribution with all $M \geq 3$ order moments vanishing is Gaussian. Hence we arrive at the following Lemma:

Lemma 2 *There is representation of $G(\mathbf{x}|\mathbf{x}_0; \Delta t)$ such that $G(\mathbf{x}|\mathbf{x}_0; \Delta t)dv(\mathbf{x})$ is Gaussian.*

As is well known, a Gaussian representation can be easily constructed using the zeroth, first, and second order moments of $G(\mathbf{x}|\mathbf{x}_0; \Delta t)$.

Next, we state a Lemma which establishes a continuous family of representations for TSGF that are equivalent to each other in the sense of Lemma 1. The proof of this Lemma is technically very complicated, and is presented in Appendix A.

Lemma 3 *Let $D^{ij}(\mathbf{x}) = D^{ji}(\mathbf{x})$ be a $d \times d$ symmetric positive definite matrix, $F^i(\mathbf{x})$ a vector, and $\Phi(\mathbf{x})$ a scalar, all functions of \mathbf{x} . Let $\Delta \mathbf{x} = \mathbf{x} - \mathbf{x}_0$, and $\mathbf{x}_\alpha = \mathbf{x}_0 + \alpha(\mathbf{x} - \mathbf{x}_0)$ with $\alpha \in [0, 1]$. The following one-parameter family of distributions are equivalent to each other, in the sense that all their moments of $\Delta \mathbf{x}$ are equal up to order $O(\Delta t)$:*

$$\frac{d\mu(\mathbf{x})}{\sqrt{(4\pi\Delta t)^d \det D^{ij}(\mathbf{x}_\alpha)}} \exp \left\{ - \left(\Delta x^i - F^i(\mathbf{x}_\alpha)\Delta t + 2\alpha\partial_k D^{ik}(\mathbf{x}_\alpha)\Delta t \right) \frac{D_{ij}^{-1}(\mathbf{x}_\alpha)}{4\Delta t} \left(\Delta x^j - F^j(\mathbf{x}_\alpha)\Delta t + 2\alpha\partial_l D^{jl}(\mathbf{x}_\alpha)\Delta t \right) - \alpha\partial_i F^i(\mathbf{x}_\alpha)\Delta t + \alpha^2\partial_i\partial_j D^{ij}(\mathbf{x}_\alpha)\Delta t - \Phi(\mathbf{x}_\alpha)\Delta t \right\}. \quad (24)$$

Note that all functions inside the bracket are evaluated at \mathbf{x}_α . The $\alpha = 0$ version of Eq. (24) is Gaussian in \mathbf{x} :

$$\frac{d\mu(\mathbf{x})}{\sqrt{(4\pi\Delta t)^d \det D^{ij}(\mathbf{x}_0)}} \exp \left\{ - \left(\Delta x^i - F^i(\mathbf{x}_0)\Delta t \right) \frac{D_{ij}^{-1}(\mathbf{x}_0)}{4\Delta t} \left(\Delta x^j - F^j(\mathbf{x}_0)\Delta t \right) - \Phi(\mathbf{x}_0)\Delta t \right\}. \quad (25)$$

There is an extension of Lemma 3. We can obtain a more general two-parameter family of equivalent representations, where D^{ij} is evaluated at $\mathbf{x}_{\alpha_1} = \mathbf{x} + \alpha_1\Delta \mathbf{x}$, and whereas F^i is evaluated at $\mathbf{x}_{\alpha_2} = \mathbf{x} + \alpha_2\Delta \mathbf{x}$:

$$\frac{d\mu(\mathbf{x})}{\sqrt{(4\pi\Delta t)^d \det D^{ij}(\mathbf{x}_{\alpha_1})}} \exp \left\{ - \left(\Delta x^i - F^i(\mathbf{x}_{\alpha_2})\Delta t + 2\alpha_1\partial_k D^{ik}(\mathbf{x}_{\alpha_1})\Delta t \right) \frac{D_{ij}^{-1}(\mathbf{x}_{\alpha_1})}{4\Delta t} \left(\Delta x^j - F^j(\mathbf{x}_{\alpha_2})\Delta t + 2\alpha_1\partial_l D^{jl}(\mathbf{x}_{\alpha_1})\Delta t \right) - \alpha_2\partial_i F^i(\mathbf{x}_{\alpha_2})\Delta t + \alpha_1^2\partial_i\partial_j D^{ij}(\mathbf{x}_{\alpha_1})\Delta t - \Phi\Delta t \right\}. \quad (26)$$

The place of evaluation for Φ has no influence in the limit $\Delta t \rightarrow 0$. Proof of this is exactly the same as Lemma 3. This result provides more flexibility in implementation of time-slicing.

III. THE α -REPRESENTATION

We will now construct a continuous family of equivalent representations for action of time-slicing path integral, both for quantum mechanics and for classical stochastic processes. These representations are characterized by an interpolation parameter $\alpha \in [0, 1]$. We first use Lemma 2 to derive a Gaussian representation, then use Lemma 3 to construct arbitrary α -representation.

A. Quantum mechanics in curved space

Using the coordinate representation and imaginary time, the TSGF is given by

$$G(\mathbf{x}|\mathbf{x}_0; \Delta t) = e^{-\Delta t \hat{H}(\mathbf{x})/\hbar} \delta(\mathbf{x}, \mathbf{x}_0), \quad (27)$$

where \hat{H} is given in Eq. (9). The moments of $G(\mathbf{x}|\mathbf{x}_0; \Delta t)$ are calculated up to the order of Δt , by expanding the exponential and using integration by parts:

$$\langle 1 \rangle = \int d\mu(\mathbf{x}) \sqrt{g(\mathbf{x})} e^{-\Delta t \hat{H}(\mathbf{x})/\hbar} \delta(\mathbf{x}, \mathbf{x}_0) = 1 - \frac{\Delta t}{\hbar} V(\mathbf{x}_0) + O(\Delta t^2), \quad (28a)$$

$$\langle \Delta x^k \rangle = \int d\mu(\mathbf{x}) \sqrt{g(\mathbf{x})} \Delta x^k e^{-\Delta t \hat{H}(\mathbf{x})/\hbar} \delta(\mathbf{x}, \mathbf{x}_0) = \frac{\hbar \Delta t}{2} \left(\frac{1}{\sqrt{g}} \partial_j (\sqrt{g} g^{jk}) \right)_0 + O(\Delta t^2), \quad (28b)$$

$$\langle \Delta x^k \Delta x^l \rangle = \int d\mu(\mathbf{x}) \sqrt{g(\mathbf{x})} \Delta x^k \Delta x^l e^{-\Delta t \hat{H}(\mathbf{x})/\hbar} \delta(\mathbf{x}, \mathbf{x}_0) = \hbar \Delta t g^{ij}(\mathbf{x}_0) + O(\Delta t^2), \quad (28c)$$

where $(\dots)_0$ means that all functions inside the bracket are evaluated at \mathbf{x}_0 .

Using these moments, we can construct a Gaussian expression for the TSGF:

$$dv(\mathbf{x}) G(\mathbf{x}|\mathbf{x}_0; \Delta t) = \frac{\sqrt{g(\mathbf{x}_0)} d\mu(\mathbf{x})}{(2\pi\hbar\Delta t)^{d/2}} e^{-A^0(\mathbf{x}, \mathbf{x}_0; \Delta t)}, \quad (29a)$$

where the *time-slice action* $A^0(\mathbf{x}, \mathbf{x}_0; \Delta t)$ is given by

$$A^0(\mathbf{x}, \mathbf{x}_0; \Delta t) = \left[\Delta x^i - \frac{\hbar \Delta t}{2} \left(\frac{1}{\sqrt{g}} \partial_k \sqrt{g} g^{ki} \right)_0 \right] \frac{g_{ij}(\mathbf{x}_0)}{2\hbar \Delta t} \left[\Delta x^j - \frac{\hbar \Delta t}{2} \left(\frac{1}{\sqrt{g}} \partial_l \sqrt{g} g^{jl} \right)_0 \right] + \frac{\Delta t}{\hbar} V(\mathbf{x}_0), \quad (29b)$$

$$= \left[\Delta x^i + \frac{\hbar \Delta t}{2} g^{kl}(\mathbf{x}_0) \Gamma_{kl}^i(\mathbf{x}_0) \right] \frac{g_{ij}(\mathbf{x}_0)}{2\hbar \Delta t} \left[\Delta x^j + \frac{\hbar \Delta t}{2} g^{mn}(\mathbf{x}_0) \Gamma_{mn}^j(\mathbf{x}_0) \right] + \frac{\Delta t}{\hbar} V(\mathbf{x}_0), \quad (29c)$$

where all functions are evaluated at \mathbf{x}_0 , and Γ_{mn}^j is the Cristopher symbol, constructed from the metric tensor:

$$\Gamma_{ij}^k \equiv \frac{1}{2} g^{kl} (\partial_i g_{jl} + \partial_j g_{il} - \partial_l g_{ij}), \quad (30a)$$

and we have used the contracting relations:

$$g^{kl} \Gamma_{kl}^i = -\frac{1}{\sqrt{g}} \partial_k (\sqrt{g} g^{ik}). \quad (30b)$$

The distribution Eq. (29a) is Gaussian, because the action $A^0(\mathbf{x}, \mathbf{x}_0; \Delta t)$ is quadratic in \mathbf{x} , and the prefactor

$\sqrt{g(\mathbf{x}_0)} d\mu(\mathbf{x})$ is independent of \mathbf{x} . One interesting feature about Eqs. (29b) is that the average of $\Delta \mathbf{x}$ is non-vanishing if the metric tensor is not constant. This is what we call the *quantum spurious drift*, which has been missed by many previous studies on path integral representation of quantum mechanics in curved space.

Now let us compare Eqs. (29b) with (25), and make the identification $F^i = (\hbar/2\sqrt{g}) \partial_k (\sqrt{g} g^{ki})$, $D^{ij} = \hbar g^{ij}/2$, and $\Phi = V/\hbar$. Applying Lemma 3, we find the following one-parameter family of representations for TSGF, all of which equivalent to Eqs. (29):

$$dv(\mathbf{x}) G(\mathbf{x}|\mathbf{x}_0; \Delta t) = \frac{\sqrt{g(\mathbf{x}_\alpha)} d\mu(\mathbf{x})}{(2\pi\hbar\Delta t)^{d/2}} e^{-A^\alpha(\mathbf{x}, \mathbf{x}_0; \Delta t)}, \quad (31a)$$

$$A^\alpha(\mathbf{x}, \mathbf{x}_0; \Delta t) = \left[\Delta x^i - \hbar \Delta t \left(\frac{1}{2\sqrt{g}} \partial_k \sqrt{g} g^{ik} - \alpha \partial_k g^{ik} \right)_\alpha \right] \frac{g_{ij}(\mathbf{x}_\alpha)}{2\hbar \Delta t} \left[\Delta x^j - \hbar \Delta t \left(\frac{1}{2\sqrt{g}} \partial_l \sqrt{g} g^{jl} - \alpha \partial_l g^{jl} \right)_\alpha \right] + \frac{\hbar \alpha}{2} \Delta t \left(\partial_i \sqrt{1/g} \partial_k \sqrt{g} g^{ki} \right)_\alpha - \frac{\hbar \alpha^2}{2} \Delta t (\partial_i \partial_j g^{ij})_\alpha + \frac{\Delta t}{\hbar} V(\mathbf{x}_\alpha). \quad (31b)$$

where $(\dots)_\alpha$ means that all functions are evaluated at \mathbf{x}_α . For the special case $\alpha = 1/2$, the quantum spurious drift reduces to

$$\hbar \left(\frac{1}{2\sqrt{g}} \partial_k \sqrt{g} g^{ik} - \alpha \partial_k g^{ik} \right) \rightarrow \frac{\hbar}{2} g^{ik} \partial_k \log \sqrt{g} = \frac{\hbar}{2} g^{ik} \Gamma_{kj}^j. \quad (32)$$

It is seen from Eq. (31b) that whilst the detailed form of quantum spurious drift depends on the choice of α , it is always non-vanishing for a generic multi-dimensional model. The lesson we learn here is that a typical quantum trajectory in curved space or in curvilinear coordinates behaves as a biased random walk.

B. Classical Markov processes

We can now apply the same procedure to the classical case, whose TSGF is

$$G(\mathbf{x}|\mathbf{x}_0; \Delta t) = e^{-\Delta t \hat{L}_{\text{FP}}(\mathbf{x})} \delta(\mathbf{x}, \mathbf{x}_0), \quad (33)$$

where the Fokker-Planck operator $\hat{L}_{\text{FP}}(\mathbf{x})$ is given in

Eq. (15). The zeroth order moments can be calculated straightforwardly:

$$\langle 1 \rangle = \int d\mu(\mathbf{x}) \sqrt{g(\mathbf{x})} e^{\Delta t \hat{L}(\mathbf{x})} \delta(\mathbf{x}, \mathbf{x}_0) = 1 + O(\Delta t^2) \quad (34a)$$

The meaning of this result is conservation of probability by Fokker-Planck equation. The first and second order moments can also be calculated:

$$\langle \Delta x^k \rangle = \Delta t \left(\frac{1}{\sqrt{g}} \partial_j (\sqrt{g} L^{kj}) - L^{kj} \partial_j U \right)_0 + O(\Delta t^2), \quad (34b)$$

$$\langle \Delta x^k \Delta x^l \rangle = 2\Delta t B^{kl}(\mathbf{x}_0) + O(\Delta t^2), \quad (34c)$$

where $(\dots)_0$ means that all functions inside the bracket are evaluated at \mathbf{x}_0 .

Using these moments, we obtain a Gaussian representation for the TSGF:

$$dv(\mathbf{x}) G(\mathbf{x}|\mathbf{x}_0; \Delta t) = \frac{d\mu(\mathbf{x}) e^{-A^0(\mathbf{x}, \mathbf{x}_0; \Delta t)}}{\sqrt{(4\pi\Delta t)^d \det B^{ij}(\mathbf{x}_0)}}, \quad (35)$$

where the time-slice action $A^0(\mathbf{x}, \mathbf{x}_0; \Delta t)$ is:

$$A^0(\mathbf{x}, \mathbf{x}_0; \Delta t) = \left[\Delta x^i - \Delta t \left(\frac{1}{\sqrt{g}} \partial_k (\sqrt{g} L^{ik}) - L^{ik} \partial_k U \right) \right] \frac{B_{ij}^{-1}(\mathbf{x}_0)}{4\Delta t} \left[\Delta x^j - \Delta t \left(\frac{1}{\sqrt{g}} \partial_l (\sqrt{g} L^{jl}) - L^{jl} \partial_l U \right) \right], \quad (36)$$

where $(B^{-1})_{ij}$ is the inverse matrix of B^{ij} , and all functions are evaluated at the initial point \mathbf{x}_0 .

We can compare Eqs. (36) with (25), and invoke Lemma 3, so that we obtain a one-parameter family of representations that are equivalent to Eqs. (36):

$$dv(\mathbf{x}) G(\mathbf{x}|\mathbf{x}_0; \Delta t) = \frac{d\mu(\mathbf{x}) e^{-A^\alpha(\mathbf{x}, \mathbf{x}_0; \Delta t)}}{\sqrt{(4\pi\Delta t)^d \det B^{ij}(\mathbf{x}_\alpha)}}, \quad (37a)$$

$$A^\alpha(\mathbf{x}, \mathbf{x}_0; \Delta t) = \left[\Delta x^i - \Delta t (F^i - 2\alpha \partial_k B^{ik})_\alpha \right] \frac{B_{ij}^{-1}(\mathbf{x}_\alpha)}{4\Delta t} \left[\Delta x^j - \Delta t (F^j - 2\alpha \partial_l B^{jl})_\alpha \right] + \alpha (\partial_i F^i)_\alpha \Delta t - \alpha^2 (\partial_i \partial_j B^{ij})_\alpha \Delta t, \quad (37b)$$

where all functions are evaluated at $\mathbf{x}_\alpha = \mathbf{x} + \alpha \Delta \mathbf{x}$, and F^i is

$$F^i = \Delta t \sqrt{1/g} (\partial_k \sqrt{g} L^{ik}) - \Delta t L^{ik} \partial_k U. \quad (38)$$

More generally, we can apply the extension of Lemma 3, and obtain a two-parameter family of equivalent representations for the TSGF:

$$dv(\mathbf{x}) G(\mathbf{x}|\mathbf{x}_0; \Delta t) = \frac{d\mu(\mathbf{x}) e^{-A^{\alpha_1, \alpha_2}(\mathbf{x}, \mathbf{x}_0; \Delta t)}}{\sqrt{(4\pi\Delta t)^d \det B^{ij}(\mathbf{x}_{\alpha_1})}}, \quad (39)$$

$$A^{\alpha_1, \alpha_2}(\mathbf{x}, \mathbf{x}_0; \Delta t) = \alpha_2 \partial_i F^i(\mathbf{x}_{\alpha_2}) \Delta t - \alpha_1^2 \partial_i \partial_j B^{ij}(\mathbf{x}_{\alpha_1}) \Delta t + \left(\Delta x^i - F^i(\mathbf{x}_{\alpha_2}) \Delta t + 2\alpha_1 \partial_l B^{il}(\mathbf{x}_{\alpha_1}) \Delta t \right) \frac{B_{ij}^{-1}(\mathbf{x}_{\alpha_1})}{4\Delta t} \left(\Delta x^j - F^j(\mathbf{x}_{\alpha_2}) \Delta t + 2\alpha_1 \partial_m B^{jm}(\mathbf{x}_{\alpha_1}) \Delta t \right)$$

where B^{ij} is evaluated at $\mathbf{x}_{\alpha_1} = \mathbf{x} + \alpha_1 \Delta \mathbf{x}$, and F^i at $\mathbf{x}_{\alpha_2} = \mathbf{x} + \alpha_2 \Delta \mathbf{x}$.

C. Two Examples

1. The problem of Edwards-Gulyaev

We consider the problem studied by Edwards and Gulyaev [28]. Consider a free particle moves in a flat plane. The quantum Hamiltonian is

$$\hat{H} = -\frac{1}{2}(\partial_x^2 + \partial_y^2) = -\frac{1}{2} \left(\frac{1}{r} \partial_r r \partial_r + \frac{1}{r^2} \partial_\phi^2 \right). \quad (40)$$

We will illustrate the above results using two simple examples.

where we have set $\hbar = m = 1$. The classical action is

$$S_{\text{cl}} = \frac{1}{2} \int dt (\dot{x}^2 + \dot{y}^2) = \frac{1}{2} \int dt (\dot{r}^2 + r^2 \dot{\phi}^2). \quad (41)$$

The TSGF in the Cartesian coordinates is

$$d\mu(x, y) G(x, y | x_0, y_0; \Delta t) = \frac{d\mu(x, y) e^{-(\Delta x^2 + \Delta y^2)/(2\Delta t)}}{2\pi\Delta t}, \quad (42)$$

where $\Delta x = x - x_0$, $\Delta y = y - y_0$, and $d\mu(x, y) = dx dy$.

Edwards and Gulyaev transform Eq. (42) to the polar coordinates

$$r = \sqrt{x^2 + y^2}, \phi = \arctan(y/x) \quad (43)$$

while keeping Δt finite, and obtain

$$\begin{aligned} & dv(r, \phi) G(r, \phi | r_0, \phi_0; \Delta t) \\ &= \frac{dv(r, \phi) e^{-[r^2 + r_0^2 - 2rr_0 \cos(\phi - \phi_0)]/(2\Delta t)}}{2\pi\Delta t}, \end{aligned} \quad (44)$$

where $dv(r, \phi) = r dr d\phi = dx dy = d\mu(x, y)$.

Defining $\Delta r = r - r_0$, $\Delta \phi = \phi - \phi_0$, and realizing that $\Delta r, \Delta \phi \sim \sqrt{\Delta t}$, we may expand the negative exponent of Eq. (44) up to order of Δt and obtain:

$$\frac{1}{2\Delta t} \left(\Delta r^2 + r_0^2 \Delta \phi^2 + r_0 \Delta r \Delta \phi^2 - \frac{1}{12} r_0^2 \Delta \phi^4 \right) + O(\Delta t^{3/2}) \quad (45)$$

If we only keep the first two terms in the bracket in Eq. (45), we would obtain the classical action (41) of an infinitesimal step:

$$\frac{1}{2\Delta t} (\Delta r^2 + r_0^2 \Delta \phi^2). \quad (46)$$

However, because $\Delta r, \Delta \phi \sim \sqrt{\Delta t}$, the third and fourth terms in Eq. (45) scale respectively as $\Delta t^{1/2}$ and Δt (taking into account the factor $1/\Delta t$ outside the bracket), and hence can not be ignored according to Lemma 1. Historically, Edwards and Gulyaev noticed the importance of the fourth term, but missed the third term which is even more important.

This problem can be easily solved using our method. We can treat it as either a quantum case or a classical case. As a quantum case, we note that the polar coordinate version of Eq. (40) can be written as Eq. (9) with $\hbar = m = 1$ and $V = 0$, and

$$g_{ij} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}, \quad g^{ij} = \begin{pmatrix} 1 & 0 \\ 0 & r^{-2} \end{pmatrix}, \quad g(r) = r^2. \quad (47)$$

Substituting these back into Eqs. (29a) and (29b), we find the TSGF (with $d\mu(r, \phi) = r dr d\phi$):

$$\begin{aligned} & dv(r, \phi) G(r, \phi | r_0, \phi_0; \Delta t) \\ &= \frac{r_0 d\mu(r, \phi) e^{-\frac{(\Delta r - \frac{\Delta t}{2r_0})^2 + r_0^2 \Delta \phi^2}{2\Delta t}}}{2\pi\Delta t}. \end{aligned} \quad (48)$$




FIG. 2: Because of the curved nature of the coordinate lines, the probability of $dr > 0$ is larger than that of $dr < 0$.

Note that Eq. (48) looks very different from Eq. (44). In particular, Eq. (48) is Gaussian in Δr and $\Delta \phi$, because the volume measure $r_0 d\mu(r, \phi)$ does not depend on r, ϕ . By contrast, Eq. (44) is clearly not Gaussian. However Lemmas 1 and 2 guarantee that these two distributions are equivalent in the sense that they share the same moments up to order Δt , which are given by

$$\begin{aligned} \langle 1 \rangle &= 1, \\ \langle \Delta r \rangle &= \frac{\Delta t}{2r_0}, \quad \langle \Delta \phi \rangle = 0, \\ \sigma_{\Delta r}^2 &= \Delta t, \quad \sigma_{\Delta \phi}^2 = \frac{\Delta t}{r_0^2}. \end{aligned} \quad (49)$$

The most salient feature of Eqs. (49) is that the average of Δr is non-vanishing. It however is anti-proportional to r_0 and hence diminishes with increasing r_0 . A non-vanishing average of Δr may sound counter-intuitive, but is in an inevitable consequence of polar coordinates. To see this, consider a particle starting from (r_0, ϕ_0) , and diffuse isotropically. It is clear that the probability of r increasing is larger than that of r decreasing, due to the curved nature of the coordinate lines, as illustrated in Fig. (2).

The Hamiltonian Eq. (40) can also be understood as negative of the Fokker-Planck operator Eq. (15) with $U = 0$ and $L^{ij} = B^{ij} = \frac{1}{2} g^{ij}$. We can then calculate the TSGF using Eqs. (36), and obtain the same result Eq. (48).

Finally, we could also rewrite Eq. (48) by Lemma 3 as

$$\begin{aligned} & dv(r, \phi) G(r, \phi | r_0, \phi_0; \Delta t) \\ &= \frac{r_{1/2} d\mu(r, \phi)}{2\pi\Delta t} \exp \left[-\frac{(\Delta r - \frac{\Delta t}{2r_{1/2}})^2 + r_{1/2}^2 \Delta \phi^2}{2\Delta t} + \frac{dt}{4r_{1/2}^2} \right] \end{aligned} \quad (50)$$

where $r_{1/2} = (r + r_0)/2$. Compared to Eq. (48), there is only an additional term proportional to dt in the action which is used to ensure normalization. The spurious drift has the same form since $\partial_i g^{ij} = 0$ for all i, j .

2. Free particle on a unit sphere

Consider a quantum particle moving on a 2-sphere with unit radius. The Hamiltonian is

$$\hat{H} = -\frac{1}{2\sin\theta}\partial_\theta\sin\theta\partial_\theta - \frac{1}{2\sin^2\theta}\partial_\phi^2, \quad (51)$$

which is in the form of Eq. (9) with $\mathbf{x} = (\theta, \phi)$, $g(\mathbf{x}) = \sin^2\theta$, and

$$g_{ij} = \begin{pmatrix} 1 & 0 \\ 0 & \sin^2\theta \end{pmatrix}, \quad g^{ij} = \begin{pmatrix} 1 & 0 \\ 0 & \sin^{-2}\theta \end{pmatrix}. \quad (52)$$

The TSGF can then be obtained using Eqs. (29):

$$\begin{aligned} & G(\theta, \phi|\theta_0, \phi_0; \Delta t)dv(\theta, \phi) \\ &= \frac{\sin\theta_0 d\mu(\theta, \phi)}{2\pi\Delta t} e^{-[(\Delta\theta - \cot(\theta_0)\Delta t/2)^2 + \sin^2\theta_0\Delta\phi^2]/2\Delta t}, \end{aligned} \quad (53)$$

where $d\mu(\theta, \phi) = d\theta d\phi$, and $dv(\theta, \phi) = \sin\theta d\theta d\phi$. For small θ , these results reduce to Eq. (48) with $\theta \rightarrow r$, as it should be.

IV. PATH-INTEGRAL FROM LANGEVIN EQUATION

It is well known that classical Markov processes can be represented by either a Fokker-Planck equation, or a Langevin equation, or a path-integral representation. The transformation between different versions of Langevin theories and the corresponding Fokker-Planck theories is well-understood, and is discussed in many standard textbooks, see for example Refs. [57, 58]. In the preceding section, we have established the correspondence between the Fokker-Planck theory and the α -representation of time-slicing path integral, i.e. Eq. (36). In this section, we will show that these representations can be obtained from the Ito-Langevin dynamics in a remarkably simple way, due to the linear relation between differential of slow variables $d\mathbf{x}$ and Wiener noises dW_μ . This simplicity is not shared by the Stratonovich-Langevin theory, or other types of Langevin theory, the relation between $d\mathbf{x}$ and noises is nonlinear. Nonetheless, we will also establish the connection between the path integral action and $\bar{\alpha}$ -Langevin dynamics, and find an equivalent family of path integral actions parameterized by two parameters $\alpha, \bar{\alpha}$. As we will see, the resulting action is much more complicated. Furthermore, for genetic multi-dimensional model, there is no way to cancel the spurious drift terms by tuning of α and $\bar{\alpha}$.

A. Path-integral from Ito-Langevin equation

We start with the Ito-Langevin equation

$$dx^i = F^i(\mathbf{x}, t)dt + b^{i\mu}(\mathbf{x})dW_\mu(t) \quad (54)$$

where $d\mathbf{x} \equiv \mathbf{x}(t+dt) - \mathbf{x}(t)$ is the infinitesimal evolution of \mathbf{x} during time step dt , and $dW_\mu(t)$, $\mu = 1, 2, 3 \dots m$ are m -dimensional Wiener noises, which are Gaussian and white, acting on \mathbf{x} during the time interval $(t, t+dt)$:

$$dW_\mu(t)dW_\nu(t) = \delta_{\mu\nu}dt. \quad (55)$$

This result is widely known as *Ito's rule* in the field of stochastic dynamics [57]. The product between the noise amplitudes $b^{i\mu}(\mathbf{x})$ and the Wiener noises $dW_\mu(t)$ is defined in Ito's sense [57], which means that $F^i(\mathbf{x}, t)$ and $b^{i\mu}(\mathbf{x})$ in Eq. (54) are evaluated at $\mathbf{x}(t)$. It then follows that Eq. (54) defines a linear relation between $d\mathbf{x}(t)$ and $dW_\mu(t)$. Hence $d\mathbf{x}(t)$ is also Gaussian, whose average and variance can be directly obtained from Eq. (54):

$$\langle dx^i \rangle = F^i(\mathbf{x}, t)dt, \quad (56a)$$

$$\left\langle (dx^i - F^i dt)(dx^j - F^j dt) \right\rangle = 2B^{ij}(\mathbf{x})dt. \quad (56b)$$

The Ito-Langevin equation (54) is mathematically equivalent to the Fokker-Planck equation:

$$\partial_t p = -\frac{1}{\sqrt{g}}\partial_i(F^i\sqrt{g}p) + \frac{1}{\sqrt{g}}\partial_i\partial_j(B^{ij}\sqrt{g}p), \quad (57)$$

where $p(\mathbf{x}, t)\sqrt{g(\mathbf{x})}d^d\mathbf{x}$ is the invariant differential probability of slow variables defined in the Introduction, and the symmetric matrix $B^{ij}(\mathbf{x})$, assumed non-singular, is related to the noise amplitudes $b^{i\mu}$ in Eq. (54) via

$$B^{ij}(\mathbf{x}) \equiv \frac{1}{2}b^{i\mu}(\mathbf{x})b^{j\nu}(\mathbf{x}) = B^{ji}(\mathbf{x}). \quad (58)$$

In two preceding publications [25, 26], we and collaborator formulated a covariant Ito-Langevin theory, where F^i in Ito-Langevin equation (54) is parameterized as

$$F^i = \frac{1}{\sqrt{g}}(\partial_t\sqrt{g}L^i) - L^j\partial_j U, \quad (59)$$

where U is related to the steady state pdf via $U = -\log p^{\text{SS}}$, and L^{ij} can be decomposed a symmetric part B^{ij} and an antisymmetric part Q^{ij} , the latter being related to the steady state current via: $J_{\text{SS}}^i = \partial_j(Q^{ij}e^{-U})$. The Ito-Langevin equation (54) and Fokker-Planck equation (57) then can be rewritten into the covariant forms, Eqs. (13) and (14). The precise meaning of covariance is discussed in Refs. [26], and will be discussed in more detailed in Sec. V.

Since $d\mathbf{x}$ as determined from Ito-Langevin dynamics is Gaussian, we can directly write down its probability distribution using its first and second order moments, given in Eqs. (56). Let $\mathbf{x}_1 \equiv \mathbf{x} + d\mathbf{x}$, and $d\mu(\mathbf{x}_1) = d^d\mathbf{x}_1$ the infinitesimal volume element at \mathbf{x}_1 and $dv(\mathbf{x}_1) = \sqrt{g(\mathbf{x}_1)}d\mu(\mathbf{x}_1)$, we have

$$dv(\mathbf{x}_1) G(\mathbf{x}_1|\mathbf{x}; dt) = \frac{d\mu(\mathbf{x}_1) \exp \left\{ - (dx^i - F^i(\mathbf{x}, t) dt) \frac{B_{ij}^{-1}(\mathbf{x})}{4dt} (dx^j - F^j(\mathbf{x}, t) dt) \right\}}{\sqrt{(4\pi)^d \det B^{ij}(\mathbf{x}) dt}}. \quad (60)$$

Note that Eq. (60) refers to transition from \mathbf{x} to $\mathbf{x}_1 = \mathbf{x} + d\mathbf{x}$ during the time interval dt , whereas in Eq. (36), the transition is from \mathbf{x}_0 to $\mathbf{x}_0 + \Delta\mathbf{x} = \mathbf{x}$ during the time interval Δt . With the correspondence of notations $(dt, d\mathbf{x}, \mathbf{x}, \mathbf{x}_1) \leftrightarrow (\Delta t, \Delta\mathbf{x}, \mathbf{x}_0, \mathbf{x})$, and F^i given by Eq. (59), Eq. (60) indeed reduces to Eqs. (35) and (36).

B. Path-integral for Stratonovich-Langevin and $\bar{\alpha}$ -Langevin

Many physicists prefer Stratonovich version of Langevin equation:

$$dx^i = F_S^i(\mathbf{x}, t) dt + b^{i\mu}(\mathbf{x}, t) \circ dW_\mu(t), \quad (61a)$$

where the product $b^{i\mu}(\mathbf{x}, t) \circ dW_\mu(t)$ is defined in Stratonovich's sense:

$$b^{i\mu}(\mathbf{x}, t) \circ dW_\mu(t) \equiv b^{i\mu}(\mathbf{x} + d\mathbf{x}/2, t) dW_\mu(t). \quad (61b)$$

Because of the hidden and generally nonlinear dependence of Eq. (61b) on $d\mathbf{x}$, the relation between $d\mathbf{x}$ and the Wiener noises $dW_\mu(t)$, as defined by Eq. (61a) is no longer linear. This leads to substantial complexity in the calculation of pdf for $d\mathbf{x}$. Using Ito's formula, one can easily prove [57] that the Stratonovich-Langevin equation (61a) is equivalent to the Ito-Langevin equation (54) with the following correspondence:

$$F^i(\mathbf{x}, t) = F_S^i(\mathbf{x}, t) + \frac{1}{2} b^{j\mu}(\mathbf{x}, t) \partial_j b^{i\mu}(\mathbf{x}, t). \quad (62)$$

More generally there is a continuous family of representations, which we shall call $\bar{\alpha}$ -Langevin equation here, with arbitrary $\bar{\alpha} \in [0, 1]$:

$$dx^i = F_{\bar{\alpha}}^i(\mathbf{x}, t) dt + b^{i\mu}(\mathbf{x}, t) \circ_{\bar{\alpha}} dW_\mu(t), \quad (63a)$$

where the product $b^{i\mu}(\mathbf{x}, t) \circ_{\bar{\alpha}} dW_\mu(t)$ is defined as

$$b^{i\mu}(\mathbf{x}, t) \circ_{\bar{\alpha}} dW_\mu(t) \equiv b^{i\mu}(\mathbf{x} + \bar{\alpha} d\mathbf{x}, t) dW_\mu(t), \quad (63b)$$

which means that the noise amplitudes $b^{i\mu}$ are evaluated at an intermediate point $\mathbf{x}_{\bar{\alpha}} \equiv \mathbf{x} + \bar{\alpha} d\mathbf{x}$. It is well known [57] that the Stratonovich-Langevin equation (63a) is equivalent to Ito-Langevin equation (54) with the following correspondence:

$$F^i(\mathbf{x}, t) = F_{\bar{\alpha}}^i(\mathbf{x}, t) + \bar{\alpha} b^{j\mu}(\mathbf{x}, t) \partial_j b^{i\mu}(\mathbf{x}, t). \quad (63c)$$

It is important to note that the parameter $\bar{\alpha}$ introduced in Langevin equation (63b) is independent of the parameter α we introduce earlier in path integral action in Sec. III. Whilst in many previous works people tried to identify these two parameters, there is no *a priori* reason for them to be the same.

We can use Eq. (62) to replace $F^i(\mathbf{x}, t)$ in Eq. (60) in terms of $F_S^i(\mathbf{x}, t)$, and obtain an equivalent expression for the TSGF:

$$dv(\mathbf{x}_1) G(\mathbf{x}_1|\mathbf{x}; dt) = \frac{d\mu(\mathbf{x}_1) \exp \left\{ - (dx^i - F_S^i dt - \frac{1}{2} b^{k\mu} \partial_k b^{i\mu} dt) \frac{B_{ij}^{-1}}{4dt} (dx^j - F_S^j dt - \frac{1}{2} b^{l\nu} \partial_l b^{j\nu} dt) \right\}}{\sqrt{(4\pi)^d \det B^{ij}}}, \quad (64)$$

where the functions F_S, B, B^{-1} and b are evaluated at \mathbf{x} . Further applying Lemma 3, we obtain an equivalent representation for $G(\mathbf{x}'|\mathbf{x}; dt)$ where all functions are evaluated at the Stratonovich point $\mathbf{x}_{1/2} = \mathbf{x} + d\mathbf{x}/2$:

$$dv(\mathbf{x}_1) G(\mathbf{x}_1|\mathbf{x}; dt) = \frac{d\mu(\mathbf{x}_1)}{\sqrt{(4\pi)^d \det B^{ij}(\mathbf{x}_{1/2})}} \exp \left\{ \begin{aligned} & - \left(dx^i - F_S^i(\mathbf{x}_{1/2}) dt + \frac{1}{2} b^{i\mu}(\mathbf{x}_{1/2}) \partial_k b^{k\mu}(\mathbf{x}_{1/2}) dt \right) \frac{B_{ij}^{-1}(\mathbf{x}_{1/2})}{4dt} \left(dx^j - F_S^j(\mathbf{x}_{1/2}) dt + \frac{1}{2} b^{i\mu}(\mathbf{x}_{1/2}) \partial_k b^{k\mu}(\mathbf{x}_{1/2}) dt \right) \\ & - \frac{dt}{2} \partial_i F_S^i + \frac{dt}{8} \left((\partial_i b^{i\mu})(\partial_j b^{j\mu}) - (\partial_i b^{j\mu})(\partial_j b^{i\mu}) \right). \end{aligned} \right. \quad (65)$$

Recall that it does not matter whether we evaluate the last two terms in Eq. (65) (both linear in dt) at \mathbf{x} or at $\mathbf{x}_{1/2}$.

More generally, we may use the $\bar{\alpha}$ -version of Langevin equation (63) and evaluate all functions at $\mathbf{x}_\alpha = \mathbf{x} + \alpha d\mathbf{x}$. The resulting TSGF is even more complicated:

$$dv(\mathbf{x}_1) G(\mathbf{x}_1|\mathbf{x}; dt) = \frac{d\mu(\mathbf{x}_1)}{(4\pi)^{d/2} \sqrt{\det B^{ij}(\mathbf{x}_\alpha)}} \exp \left\{ \begin{aligned} & - \left(dx^i - F_\alpha^i dt - \bar{\alpha} b^{k\mu} \partial_k b^{i\mu} dt + 2\alpha \partial_k B^{ik} dt \right)_\alpha \frac{B_{ij}^{-1}(\mathbf{x}_\alpha)}{4 dt} \left(dx^j - F_\alpha^j dt - \bar{\alpha} b^{\nu l} \partial_l b^{j\nu} dt + 2\alpha \partial_l B^{jl} dt \right)_\alpha \\ & - \alpha \partial_i F_\alpha^i dt - \alpha \bar{\alpha} \partial_i (b^{k\mu} \partial_k b^{i\mu}) dt + \alpha^2 \partial_i \partial_j B^{ij} dt \end{aligned} \right\}. \quad (66)$$

The spurious drift is now given by $(\bar{\alpha} b^{k\mu} \partial_k b^{i\mu} - 2\alpha \partial_k B^{ik}) dt$, which depends both on α and on $\bar{\alpha}$. It can be seen that for a general multi-dimensional model, there is no way to cancel the spurious drift by tuning of parameters $\alpha, \bar{\alpha}$. For $\alpha = \bar{\alpha} = 1/2$, Eq. (66) reduces to Eq. (65). For $\alpha = \bar{\alpha} = 0$, Eq. (66) reduces to Eq. (60), which is the simplest of all representations.

V. COVARIANCE

The principle of covariance dictates that all basic equations and quantities must be represented in tensor forms, whose validity is independent of choice of coordinate systems. The concept of covariance has served as a cornerstone for the major parts of modern theoretical physics, including general relativity and gauge theories of elementary interactions. The fundamental assumption underlying the principle of covariance is that laws of physics are objective, whereas choices of coordinate system are subjective. Change of coordinate system does not influence the validity of physical laws, but only leads to equivalent representations of the same laws.

For physical theories defined on curved manifolds, various tensors can be classified as covariant and contra-variant according to specific transformation laws they obey under nonlinear transformation of coordinates. It is well accepted that all physics equations should be written explicitly in a covariant or contra-variant format, which guarantees their validity in arbitrary coordinate system, given that they are valid in one particular coordinate system. Discovery of the covariant form of physical equations however may not be a trivial task.

The issue of covariance of path integral has caused substantial confusion in the past. Even though a large number of works were published on the topic of path integral in curved space [29, 52, 59–63], very few of these works addressed explicitly how path integral transform under nonlinear change of coordinates. Edwards and Gulyaev [28] argued that the usual chain rule of calculus is not applicable in coordinate transformation of path integral, but did not supply detail. Deininghaus

and Graham [56] developed a covariant theory for path integral which is applicable only for normal coordinate systems, which is of little use. More recently, Cugliandolo *et. al.* [50–52] tried to make path integral covariant under usual calculus rules by adjusting the interpolation parameter α . It is not clear whether this method works for multi-dimensional systems.

We first note that our α representations of TSGF, Eqs. (31b), (37c), and (39), are expressed exclusively in terms of tensor objects. For the quantum case, they are the covariant metric tensor g_{ij} , and the potential $V(\mathbf{x})$ which is a scalar. For the classical case, these objects are the second rank contra-variant tensor L^{ij} , the covariant metric tensor g_{ij} , and the scalar $U(\mathbf{x})$. When one makes a nonlinear transformation of coordinates, these tensor objects transform in a well-known fashion given below

$$p(\mathbf{x}) \rightarrow p'(\mathbf{x}') = p(\mathbf{x}). \quad (67a)$$

$$V(\mathbf{x}) \rightarrow V'(\mathbf{x}') = V(\mathbf{x}), \quad (67b)$$

$$U(\mathbf{x}) \rightarrow U'(\mathbf{x}') = U(\mathbf{x}), \quad (67c)$$

$$g_{ij}(\mathbf{x}) \rightarrow g'_{ab}(\mathbf{x}') = \frac{\partial x^i}{\partial x'^a} g_{ij}(\mathbf{x}) \frac{\partial x^j}{\partial x'^b}, \quad (67d)$$

$$b^{i\mu}(\mathbf{x}) \rightarrow b'^{a\mu}(\mathbf{x}') = \frac{\partial x'^a}{\partial x^i} b^{i\mu}(\mathbf{x}), \quad (67e)$$

$$L^{ij}(\mathbf{x}) \rightarrow L'^{ab}(\mathbf{x}') = \frac{\partial x'^a}{\partial x^i} L^{ij}(\mathbf{x}) \frac{\partial x'^b}{\partial x^j}. \quad (67f)$$

We can then use the new tensors to assemble the new α representation, which has the same form as the old representation, but with all tensors and coordinates replaced by their transformed versions. For example, the transformed quantum TSGF is

$$dv(\mathbf{x}') G'(\mathbf{x}'|\mathbf{x}'_0; \Delta t) = \frac{\sqrt{g(\mathbf{x}'_0)} d\mu(\mathbf{x}')}{(2\pi\hbar\Delta t)^{d/2}} e^{-A'^0(\mathbf{x}', \mathbf{x}'_0; \Delta t)}, \quad (68a)$$

$$\begin{aligned}
A'^0(\mathbf{x}, \mathbf{x}_0; \Delta t) &= \left[\Delta x'^a - \frac{\hbar \Delta t}{2} \left(\frac{1}{\sqrt{g'}} \partial'_c \sqrt{g'} g'^{ca} \right)_0 \right] \frac{g'_{ab}(\mathbf{x}'_0)}{2\hbar \Delta t} \left[\Delta x'^b - \frac{\hbar \Delta t}{2} \left(\frac{1}{\sqrt{g'}} \partial'_d \sqrt{g'} g'^{db} \right)_0 \right] + \frac{\Delta t}{\hbar} V'(\mathbf{x}'_0), \\
&= \left[\Delta x'^a + \frac{\hbar \Delta t}{2} g'^{cd}(\mathbf{x}'_0) \Gamma'_{cd}{}^a(\mathbf{x}'_0) \right] \frac{g'_{ab}(\mathbf{x}'_0)}{2\hbar \Delta t} \left[\Delta x'^b + \frac{\hbar \Delta t}{2} g'^{ef}(\mathbf{x}'_0) \Gamma'_{ef}{}^j(\mathbf{x}'_0) \right] + \frac{\Delta t}{\hbar} V'(\mathbf{x}'_0), \quad (68b)
\end{aligned}$$

which is the counterpart of Eqs. (29). Hence our results of TSGF valid for arbitrary choices of coordinate systems.

On the other hand, one may wonder whether the action of path integral Eq. (29) itself transforms as a scalar, i.e., i.e., whether we have

$$A'^0(\mathbf{x}', \mathbf{x}'_0; \Delta t) \stackrel{?}{=} A^0(\mathbf{x}, \mathbf{x}_0; \Delta t). \quad (69)$$

The expectation of equality seems very natural, since the classical action (8) is known to be a scalar under nonlinear change of coordinates. On the quantum side, since both \hat{H} and $\delta(\mathbf{x}, \mathbf{x}_0)$ are scalars, the TSGF as defined by Eq. (27) for finite transition time Δt , must also be scalars. More precisely, we expect that under nonlinear transform of variables, the Green's function transforms should be invariant with usual rules of calculus:

$$G'(\mathbf{x}'; t|\mathbf{x}'_0, t_0) = G(\mathbf{x}; t|\mathbf{x}_0, t_0), \quad (70)$$

where the LHS is the Green's function formulated in the new coordinates, whereas the RHS is that formulated in the old coordinates. This argument seems to suggest that the path integral actions, which we formulated in Eqs. (31b), (37c), and (39), should also behave as scalars under NTV.

There is however an important difference between the volume measures in the LHS and RHS of Eq. (31a). Whereas $\sqrt{g(\mathbf{x})}d^d x$ in the LHS is the invariant volume measure at point \mathbf{x} , in the RHS we have $\sqrt{g(\mathbf{x}_\alpha)}d^d x$, which is not a scalar. This is because $d^d x$ is an infinitesimal volume element at point \mathbf{x} , whereas $\sqrt{g(\mathbf{x}_\alpha)}$ is evaluated at a different point $\mathbf{x}_\alpha = \mathbf{x}_0 + \alpha(\mathbf{x} - \mathbf{x}_0)$. Their matrices of transformation differ from each other. Analogous statement can also be made for the classical results, Eqs. (37c). Hence we conclude that for $\alpha \neq 1$, the actions of path integral do not transform as a scalar, if the usual rules of calculus are used. This discussion seems to suggest that the $\alpha = 1$ representation of time-slicing path integral is superior because its action transforms as a scalar. However, this representation of TSGF is not Gaussian, because all functions are evaluated at \mathbf{x} . In fact only the $\alpha = 0$ representation is Gaussian, from which we can directly read off average and variance of $\Delta \mathbf{x}$. This makes it most convenient for numerical implementation and also for analytic discussion.

A. $\alpha = 0$ Quantum Path Integral

Let us discuss the transformation of $\alpha = 0$ representation more carefully. Eqs. (68) and (29) are equivalent to each other, in the sense that the moments of \mathbf{x}' and \mathbf{x}

they yield are consistent with each other up to the order of Δt . More precisely, suppose we want to calculate the average of dx'^a , we can do it in two different ways. Firstly we can treat dx'^a as a function of $\Delta \mathbf{x}$ and \mathbf{x}_0 , which is derivable from the coordinate transformation $x'^a(\mathbf{x})$ by Taylor expansion:

$$\Delta x'^a = \frac{\partial x'^a}{\partial x^i} \Delta x^i + \frac{1}{2} \frac{\partial^2 x'^a}{\partial x^i \partial x^j} \Delta x^i \Delta x^j + \dots \quad (71)$$

If \mathbf{x} and \mathbf{x}' are both smooth functions of t , $\Delta \mathbf{x}'$ and $\Delta \mathbf{x}$ would be both linear in Δt . Then we can stop at the first order in the RHS of Eq. (71), and conclude that $\Delta \mathbf{x}$ transforms as a contra-variant vector. However typical paths of path integrals are continuous but non-differentiable, and $\Delta \mathbf{x}$ scales as $\Delta t^{1/2}$. Hence in order to achieve precision up to the order Δt , we have to keep the second order term in Eq. (71), which is linear in Δt . Taking average of Eq. (71) we find

$$\begin{aligned}
\langle \Delta x'^a \rangle &= \frac{\hbar \Delta t}{2} \left(\frac{\partial x'^a}{\partial x^i} \frac{1}{\sqrt{g}} \partial_k \sqrt{g} g^{ki} + \frac{\partial^2 x'^a}{\partial x^i \partial x^j} g^{ij} \right)_0 \\
&= \frac{\partial x'^a}{\partial x^i} \langle \Delta x^i \rangle + \frac{\hbar \Delta t}{2} \left(\frac{\partial^2 x'^a}{\partial x^i \partial x^j} g^{ij} \right)_0. \quad (72)
\end{aligned}$$

Alternatively, we can also directly read off the average of $\Delta x'^a$ from Eq. (68), the transformed version of TSGF:

$$\langle \Delta x'^a \rangle = \frac{\hbar \Delta t}{2} \left(\frac{1}{\sqrt{g'}} \partial'_d \sqrt{g'} g'^{db} \right)_0. \quad (73)$$

The RHS of Eqs. (73) and (72) must be identical. In fact, using of Eq. (67d) in Eq. (72), and using the identity (proved in the appendix of Ref. [26]):

$$\frac{\partial}{\partial x'^a} \left(\frac{\partial x'^a}{\partial x^i} J \right) = 0, \quad (74)$$

where $J = \det \left(\frac{\partial x'^a}{\partial x^i} \right)$ is the Jacobian, we find

$$\begin{aligned}
&\frac{\hbar \Delta t}{2} \left(\frac{1}{\sqrt{g'}} \partial'_d \sqrt{g'} g'^{db} \right) \\
&= \frac{\hbar \Delta t}{2} \left(\frac{\partial x'^a}{\partial x^i} \frac{1}{\sqrt{g}} \partial_k \sqrt{g} g^{ki} + \frac{\partial^2 x'^a}{\partial x^i \partial x^j} g^{ij} \right). \quad (75)
\end{aligned}$$

This is the transformation rule of the quantum spurious drift. Hence we conclude that spurious drift transforms does not as a vector, but much similar to the affine connections in Riemannian geometry.

But there is an even more interesting observation. Comparing Eqs. (68b) with (29c), and noticing that the

metric tensor g_{ij} is covariant, we see that if we further impose the following transformation law:

$$\Delta x'^a = \frac{\partial x'^a}{\partial x^i} \Delta x^i + \frac{\hbar}{2} g^{ij} \frac{\partial^2 x'^a}{\partial x^i \partial x^j} \Delta t, \quad (76)$$

the linear combination

$$\Delta x^i - \frac{\hbar \Delta t}{2} \left(\frac{1}{\sqrt{g}} \partial_k \sqrt{g} g^{ki} \right) = \Delta x^i + \frac{\hbar \Delta t}{2} g^{kl}(\mathbf{x}_0) \Gamma_{kl}^i(\mathbf{x}_0)$$

transforms as a contra-variant vector and the action Eq. (29c) transforms as a scalar, i.e., it remains invariant under nonlinear change of coordinates. We call Eq. (76) the *quantum Ito's formula* because it involves the Planck constant \hbar , and is closely related to Ito's formula in classical stochastic processes, as we will show momentarily.

It can be seen that $\alpha \neq 0$ representations of path-integral action do not transform as scalars under Ito's formula. In principle, there may be other rules which make these actions invariant, but we do not know the existence of any such rule. We note however, Ito's formula is unique in the sense that it encodes the transformations of both average and variance of $\Delta \mathbf{x}$ under nonlinear change of coordinates.

B. $\alpha = 0$ Classical Stochastic Processes

In two previous works, we and collaborator have established the covariance of both Ito-Langevin theory and Fokker-Planck theory [25, 26]. More specifically, we have shown that under the transformation rules Eqs. (67), the Fokker-Planck equation (14) is transformed into:

$$\partial_t p' = \frac{1}{\sqrt{g'}} \partial'_a \sqrt{g'} L'^{ab} ((\partial'_b U') + \partial'_b) p' = L'_{\text{FP}} p', \quad (77)$$

which has the same form as Eq. (14). It can be seen from here that the Fokker-Planck operator transforms also as a scalar:

$$L'_{\text{FP}}(\mathbf{x}') = L_{\text{FP}}(\mathbf{x}). \quad (78)$$

Furthermore, under the transformation rules Eqs. (67) together with *Ito's formula* [57]:

$$dx'^a = \frac{\partial x'^a}{\partial x^i} dx^i + B^{ij} \frac{\partial^2 x'^a}{\partial x^i \partial x^j} dt, \quad (79)$$

the Ito-Langevin equation (13) is transformed into:

$$dx'^a + L'^{ab} \partial'_b U' dt - \frac{1}{\sqrt{g'}} \partial'_b \sqrt{g'} L'^{ab} dt = b'^{a\alpha} dW^\alpha, \quad (80)$$

which again has the same form as Eq. (13). Ito's formula (79) was deprive by using Eq. (13) and Ito's rule (55), and keeping terms up to $O(dt)$ [57].

Note that in accordance with the standard notations of Langevin dynamics, we use notations $d\mathbf{x}$, $d\mathbf{x}'$ and dt , instead of $\Delta \mathbf{x}$, $\Delta \mathbf{x}'$ and Δt as in Sec. V A. Note also that

Ito's formula Eq. (79) is identical to the quantum Ito's formula Eq. (76) under the correspondence $B^{ij} \leftrightarrow \hbar g^{ij}/2$. The classical analogue of Eq. (72) is

$$\langle dx'^a \rangle = \frac{\partial x'^a}{\partial x^i} \langle dx^i \rangle + \frac{\partial^2 x'^a}{\partial x^i \partial x^j} B^{ij} dt, \quad (81)$$

which can also be obtained by averaging Eq. (79).

But according to Eqs. (56), (59), the average of dx^i is

$$\langle dx^i \rangle = -L^{ij} \partial_j U dt + \frac{1}{\sqrt{g}} \partial_j \sqrt{g} L^{ij} dt, \quad (82a)$$

whereas in the transformed theory, we have

$$\langle dx'^a \rangle = -L'^{ab} \partial'_b U' dt + \frac{1}{\sqrt{g'}} \partial'_b \sqrt{g'} L'^{ab} dt. \quad (82b)$$

Substituting the preceding two results back into Eq. (81), we obtain:

$$\begin{aligned} & -L'^{ab} \partial'_b U' dt + \frac{1}{\sqrt{g'}} \partial'_b \sqrt{g'} L'^{ab} dt \\ &= \frac{\partial x'^a}{\partial x^i} \left(-L^{ij} \partial_j U dt + \frac{1}{\sqrt{g}} \partial_j \sqrt{g} L^{ij} dt \right) + \frac{\partial^2 x'^a}{\partial x^i \partial x^j} B^{ij} dt. \end{aligned} \quad (83)$$

Combining Eqs. (79) with (83), we see that even though neither term in the LHS of Eq. (13) transforms as a vector, the linear combination does transform homogeneously and as a contra-variant vector under NTV. Ito-Langevin equation (13) is frequently written in an alternative form:

$$dx^i = \left(-L^{ij} \partial_j U + \frac{1}{\sqrt{g}} \partial_i \sqrt{g} L^{ij} \right) dt + b^{i\mu} dW_\mu(t), \quad (84)$$

where the systematic force appear in RHS. This form is however less convenient, because neither side of it transforms as a vector.

We hence conclude that the time-slice action $A^0(\mathbf{x}, \mathbf{x}_0, \Delta t)$ for classical Markov process, given in Eq. (36), also transform as a scalar under nonlinear change of coordinates, if $\Delta \mathbf{x}$ transform according to Ito's formula (79).

We must emphasize however Ito's formula (79), and the closely related Ito's rule (55) must be used with extreme caution. In many previous works, Ito's rule (55) was used freely in the derivation of path-integral action. This may result in errors. In Sec. V D, we explicitly illustrate errors due to application of Ito's rule, using the problem of Edwards-Gulyaev as an example.

C. Edwards-Gulyaev revisited

Let us revisit the example of 2d free particle in polar coordinates using Langevin theory. The Langevin equations of a free particle in Cartesian coordinates are

$$dx = dW_1(t); \quad dy = dW_2(t). \quad (85)$$

This can also be written in the covariant form with $U = 0$, $g_{ij} = L^{ij} = \delta_{ij}$. The transition probability is just

$$d\mu(x_1, y_1)G(x_1, y_1|x, y; dt) = \frac{d\mu(x_1, y_1)}{2\pi} \exp -\frac{dx^2 + dy^2}{2dt}, \quad (86)$$

where, as in Eq. (16), $d\mu(x_1, y_1) = dx_1 dy_1$. Now transform to the polar coordinate $r = \sqrt{x^2 + y^2}$, $\phi = \arctan(y/x)$. Ito's rule

$$dW_1(t)^2 = dW_2(t)^2 = dt, \quad dW_1(t)dW_2(t) = 0, \quad (87)$$

must be invoked for transformations from dx, dy to $dr, d\phi$:

$$\begin{aligned} dr &= \cos \phi dx + \sin \phi dy + \frac{1}{2} \frac{\partial^2 r}{\partial x^2} dx^2 + \frac{1}{2} \frac{\partial^2 r}{\partial y^2} dy^2 \\ &= \cos \phi dx + \sin \phi dy + \frac{dt}{2r}; \end{aligned} \quad (88a)$$

$$\begin{aligned} d\phi &= -\frac{\sin \phi}{r} dx + \frac{\cos \phi}{r} dy + \frac{\partial^2 \phi}{\partial x^2} dx^2 + \frac{\partial^2 \phi}{\partial y^2} dy^2 \\ &= -\frac{\sin \phi}{r} dx + \frac{\cos \phi}{r} dy. \end{aligned} \quad (88b)$$

Using these to express dx, dy in terms of $dr, d\phi$ in Eqs. (85), we obtain the Ito-Langevin equations in polar coordinates:

$$dr = \frac{1}{2r} dt + b^{1m} dW_m; \quad (89a)$$

$$d\phi = b^{2m} dW_m. \quad (89b)$$

where the matrix b^{im} is given by

$$(b^{im}) = \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi/r & \cos \phi/r \end{pmatrix}. \quad (90)$$

So we can write the transition probability as

$$\begin{aligned} dv(r_1, \phi_1)G(r_1, \phi_1|r, \phi, dt) \\ = \frac{rd\mu(r_1, \phi_1)}{2\pi dt} \exp -\left(\frac{(dr - dt/(2r))^2}{2dt} + \frac{r^2 d\phi^2}{2dt} \right), \end{aligned} \quad (91)$$

where $dv(r_1, \phi_1) = r_1 dr_1 d\phi_1$ and $d\mu = dr_1 d\phi_1$. This result is identical to Eqs. (48), under the correspondence of notations: $(dr_1, d\phi_1, dt, r, \phi) \leftrightarrow (\Delta r, \Delta \phi, \Delta t, r_0, \phi_0)$.

The Ito-Langevin equations (89) can be transformed into Stratonovich-Langevin form:

$$dr = \frac{1}{2r} dt + b^{1m} \circ dW_m; \quad (92)$$

$$d\phi = b^{2m} \circ dW_m. \quad (93)$$

which is formally identical to the Ito form, except that the Ito products are replaced by the Stratonovich products. The path-integral representation written with $\alpha = 1/2$ can also be obtained using Eq. (65):

$$\begin{aligned} dv(r_1, \phi_1)G(r_1, \phi_1|r, \phi; dt) \\ = \frac{r_{1/2} d\mu(r_1, \phi_1)}{2\pi dt} \exp \left[-\frac{(dr - \frac{dt}{2r_{1/2}})^2}{2dt} + \frac{dt}{4r_{1/2}^2} \right] \end{aligned} \quad (94)$$

This result is identical to Eq. (94) since it describes the same dynamics with all functions evaluated at the same point $r_{1/2} = r + dr/2$.

D. Improper use of Ito's rule

Here we use the problem of Edwards-Gulyaev to illustrate the mistake caused by improper use of Ito's rule in the derivation of time-slice action. Using Eq. (45) the action can be written as

$$\frac{1}{2dt} \left(dr^2 + r^2 d\phi^2 + r dr d\phi^2 - \frac{1}{12} r^2 d\phi^4 \right) \quad (95)$$

Now if we use Ito's rule to replace the cross term $r dr d\phi^2$ by $dr dt/r^2$ and rewrite the action as

$$\frac{1}{2dt} \left(dr^2 + r^2 d\phi^2 + dr dt/r - \frac{1}{12} r^2 d\phi^4 \right) \quad (96)$$

This new action is not correctly normalized. The error is a function of r . This is shown by direct calculation

$$\begin{aligned} &\int \frac{dv(r_1, \phi_1)}{2\pi dt} e^{-\frac{1}{2dt}(dr^2 + r^2 d\phi^2 + dr dt/r - \frac{1}{12} r^2 d\phi^4)} \\ &= \int \frac{(r + dr)d\mu(r_1, \phi_1)}{2\pi dt} e^{-\frac{1}{2dt}(dr^2 + r^2 d\phi^2)} \\ &\quad \left(1 - dr/(2r) + \frac{dr^2}{8r^2} + \frac{r^3 d\phi^4}{24dt} \right) + O(dt^{3/2}) \\ &= 1 - \frac{dt}{4r^2} + O(dt^{3/2}). \end{aligned} \quad (97)$$

This example shows that the application of Ito's rule to the derivation of action may lead to mistake.

VI. CONCLUSION AND OUTLOOK

Let us summarize the main results we have achieved in this work: (1) With Lemma 1 we have established a rigorous criterion for equivalence between different time-slicing path integrals. (2) With Lemma 2 we have demonstrated the existence of a Gaussian representation for the time-slice Green's function. (3) With Lemma 3 we constructed a continuous family of equivalent time-slicing path-integral actions, parameterized by an interpolation parameter $\alpha \in [0, 1]$. (4) We have established the connection between Langevin dynamics and time-slicing path integral, and have clarified the origin and transformation rules of spurious drift. (5) We have shown why in general path integral actions do not transform as scalars under usual rules of tensor calculus. We have also explicitly demonstrated that the $\alpha = 0$ representation of path integral action transforms as a scalar if $\Delta \mathbf{x}$ transform according to Ito's formula, while all other ingredients transform according to the usual rules of tensor calculus. These results resolve several major confusions

regarding time-slicing path integral in curved space, and provide a rigorous, practical, and manifestly covariant calculation scheme.

The transformation rule of spurious drift, Eq. (75) for quantum case and Eq. (83) for classical case, resembles affine connections in differential geometry. Together with the quantum Ito formula, they hint at some intriguing geometric effects of both quantum mechanics and stochastic processes in curved space, which we shall explore in the future. We also plan to study the covariance properties of path integral formalism for Non-linear Sigma model and Non-Abelian gauge field theories in curved spaces. More

importantly, we shall apply the path-integral methods developed here to study concrete quantum and classical stochastic processes in curved space.

VII. ACKNOWLEDGEMENT

The authors acknowledge support from NSFC via grant 11674217 as well as Shanghai Municipal Science and Technology Major Project (Grant No.2019SHZDZX01). X.X. is also thankful to additional support from a Shanghai Talent Program.

-
- [1] Dirac, Paul Adrien Maurice. The principles of quantum mechanics. No. 27. Oxford university press, 1981.
 - [2] Feynman, R. P. Space-Time Approach to Non-Relativistic Quantum Mechanics. *Reviews of Modern Physics* 20.2 (1948): 367-387.
 - [3] Feynman, Richard P., Albert R. Hibbs, and Daniel F. Styer. *Quantum mechanics and path integrals*. Courier Corporation, 2010.
 - [4] Kleinert, Hagen. *Quantum equivalence principle. Functional Integration*. Springer, Boston, MA, 1997. 67-92.
 - [5] Zinn-Justin, Jean. *Path integrals in quantum mechanics*. Oxford University Press, 2010.
 - [6] Zee, Anthony. *Quantum field theory in a nutshell*. Vol. 7. Princeton university press, 2010.
 - [7] Feynman, Richard Phillips, and F. L. Vernon Jr. "The theory of a general quantum system interacting with a linear dissipative system." *Annals of physics* 281.1-2 (2000): 547-607.
 - [8] Weiss, Ulrich. *Quantum dissipative systems*. Vol. 13. World scientific, 2012.
 - [9] Hawking, Stephen W. *The path-integral approach to quantum gravity. General relativity*. 1979.
 - [10] Wio, Horacio S. *Path integrals for stochastic processes: An introduction*. World Scientific, 2013.
 - [11] Chernyak, Vladimir Y., Michael Chertkov, and Christopher Jarzynski. Path-integral analysis of fluctuation theorems for general Langevin processes. *Journal of Statistical Mechanics: Theory and Experiment* 2006.08 (2006): P08001.
 - [12] De Gennes, Pierre-Gilles, and Pierre-Gilles Gennes. *Scaling concepts in polymer physics*. Cornell university press, 1979.
 - [13] Kleinert, Hagen. *Path integrals in quantum mechanics, statistics, polymer physics, and financial markets*. World scientific, 2009.
 - [14] Linetsky, Vadim. The path integral approach to financial modeling and options pricing. *Computational Economics* 11.1 (1997): 129-163.
 - [15] Wiener, Norbert. *Differential-Space*. *Journal of Mathematics and Physics* 2.1-4 (1923): 131-174.
 - [16] Kac, Mark. On some connections between probability theory and differential and integral equations. *Proceedings of the second Berkeley symposium on mathematical statistics and probability*. University of California Press, 1951.
 - [17] Glimm, James, and Arthur Jaffe. *Quantum physics: a functional integral point of view*. Springer Science & Business Media, 2012.
 - [18] Hänggi, P. Connection between deterministic and stochastic descriptions of non-linear systems. *Physica Acta* 53, 491-496 (1980).
 - [19] Onsager, Lars, and Stefan Machlup. Fluctuations and irreversible processes. *Physical Review* 91.6 (1953): 1505.
 - [20] Machlup, Stefan, and Lars Onsager. Fluctuations and irreversible process. II. Systems with kinetic energy. *Physical Review* 91.6 (1953): 1512.
 - [21] Bastianelli, Fiorenzo, and Peter Van Nieuwenhuizen. *Path integrals and anomalies in curved space*. Cambridge University Press, 2006.
 - [22] Salomonson, Per. When does a non-linear point transformation generate an extra $O(\hbar^2)$ potential in the effective Lagrangian?. *Nuclear Physics B* 121.3 (1977): 433-444.
 - [23] Gervais, J-L., and A. Jevicki. Point canonical transformations in the path integral. *Nuclear Physics B* 110.1 (1976): 93-112.
 - [24] Apfeldorf, Karyn M., and Carlos Ordonez. Coordinate redefinition invariance and 'extra' terms. *Nuclear Physics B* 479.1-2 (1996): 515-526.
 - [25] Mingnan Ding and Xiangjun Xing. Covariant Non-equilibrium Thermodynamics for Small Systems. arXiv:2105.14534
 - [26] Mingnan Ding, Zhanchun Tu, and Xiangjun Xing. Covariant formulation of nonlinear Langevin theory with multiplicative Gaussian white noises. *Physical Review Research* 2.3 (2020): 033381.
 - [27] DeWitt, Bryce S. *Dynamical theory in curved spaces. I. A review of the classical and quantum action principles. Reviews of modern physics* 29.3 (1957): 377.
 - [28] Edwards, Samuel Frederick, and Y. V. Gulyaev. Path integrals in polar co-ordinates. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* 279.1377 (1964): 229-235.
 - [29] D. W. McLaughlin and L. S. Schulman. *Path Integrals in Curved Spaces*. *Journal of Mathematical Physics* 12, 2520 (1971).
 - [30] C. Grosche and F. Steiner. Path integrals on curved manifolds. *Zeitschrift für Physik C Particles and Fields* 36, 699 (1987).
 - [31] P. Hänggi. Path integral solutions for non-Markovian processes. *Zeitschrift für Physik B Condensed Matter* 75, 275 (1989).
 - [32] H. Dekker. On the functional integral for generalized

- Wiener processes and nonequilibrium phenomena. *Physica A: Statistical Mechanics and its Applications* 85, 598 (1976).
- [33] R. F. Fox. Functional-calculus approach to stochastic differential equations. *Phys. Rev. A* 33, 467 (1986).
- [34] D. Durr and A. Bach. The Onsager-Machlup function as Lagrangian for the most probable path of a diffusion process. *Commun.Math. Phys.* 60, 153 (1978).
- [35] J. Alfaro and P. H. Damgaard. Field transformations, collective coordinates and BRST invariance. *Annals of Physics* 202, 398 (1990).
- [36] A. Inoue and Y. Maeda. On integral transformations associated with a certain Lagrangian-as a prototype of quantization. *J. Math. Soc. Japan* 37, 219 (1985).
- [37] Grosche, Christian. Path integration and separation of variables in spaces of constant curvature in two and three dimensions. *Fortschritte der Physik/Progress of Physics* 42.6 (1994): 509-584.
- [38] Lecheheb, A., M. Merad, and T. Boudjedaa. Path integral treatment for a Coulomb system constrained on D-dimensional sphere and hyperboloid. *Annals of Physics* 322.5 (2007): 1233-1246.
- [39] Tisza, Laszlo, and Irwin Manning. Fluctuations and irreversible thermodynamics. *Physical Review* 105.6 (1957): 1695.
- [40] Bhagwat, K. V., Dinkar C. Khandekar, and Shilpa V. Lawande. Path integral methods and their applications. World Scientific, 1993.
- [41] Dekker, H. Functional integration and the Onsager-Machlup Lagrangian for continuous Markov processes in Riemannian geometries. *Physical Review A* 19.5 (1979): 2102.
- [42] Janssen, Hans-Karl. On a Lagrangean for classical field dynamics and renormalization group calculations of dynamical critical properties. *Zeitschrift für Physik B Condensed Matter* 23.4 (1976): 377-380.
- [43] Adib, Artur B. Stochastic actions for diffusive dynamics: Reweighting, sampling, and minimization. *The Journal of Physical Chemistry B* 112.19 (2008): 5910-5916.
- [44] Langouche, Flor, Dirk Roekaerts, and Enrique Tirapegui. Functional integration and semiclassical expansions. Vol. 10. Springer Science & Business Media, 2013.
- [45] Itami, Masato, and Shin-ichi Sasa. Universal form of stochastic evolution for slow variables in equilibrium systems. *Journal of Statistical Physics* 167.1 (2017): 46-63.
- [46] Arnold, Peter. Symmetric path integrals for stochastic equations with multiplicative noise. *Physical Review E* 61.6 (2000): 6099.
- [47] Calisto, H., and E. Tirapegui. Comment on ‘Symmetric path integrals for stochastic equations with multiplicative noise’. *Physical Review E* 65.3 (2002): 038101.
- [48] Langouche, F., D. Roekaerts, and E. Tirapegui. On the path integral solution of the master equation. *Physics Letters A* 68.5-6 (1978): 418-420.
- [49] Tang, Ying, Ruoshi Yuan, and Ping Ao. Summing over trajectories of stochastic dynamics with multiplicative noise. *The Journal of chemical physics* 141.4 (2014): 044125.
- [50] Cugliandolo, Leticia F., and Vivien Lecomte. Rules of calculus in the path integral representation of white noise Langevin equations: the Onsager-Machlup approach. *Journal of Physics A: Mathematical and Theoretical* 50.34 (2017): 345001.
- [51] Cugliandolo, Leticia F., Vivien Lecomte, and Frédéric Van Wijland. Building a path-integral calculus. (2018).
- [52] Cugliandolo, Leticia F., Vivien Lecomte, and Frédéric Van Wijland. Building a path-integral calculus: a covariant discretization approach. *Journal of Physics A: Mathematical and Theoretical* 52.50 (2019): 50LT01.
- [53] Lau, Andy WC, and Tom C. Lubensky. State-dependent diffusion: Thermodynamic consistency and its path integral formulation. *Physical Review E* 76.1 (2007): 011123.
- [54] Haken, H. Generalized Onsager-Machlup function and classes of path integral solutions of the Fokker-Planck equation and the master equation. *Zeitschrift für Physik B Condensed Matter* 24.3 (1976): 321-326.
- [55] Wissel, C. Manifolds of equivalent path integral solutions of the Fokker-Planck equation. *Zeitschrift für Physik B Condensed Matter* 35.2 (1979): 185-191.
- [56] Deininghaus, U., and R. Graham. Nonlinear point transformations and covariant interpretation of path integrals. *Zeitschrift für Physik B Condensed Matter* 34.2 (1979): 211-219.
- [57] Gardiner, Crispin W. Handbook of stochastic methods. Vol. 3. Berlin: springer, 1985.
- [58] Van Kampen, Nicolaas Godfried. Stochastic Processes in Physics and Chemistry, 3rd ed., Elsevier, 2007.
- [59] Weiss, U. Operator ordering schemes and covariant path integrals of quantum and stochastic processes in curved space. *Zeitschrift für Physik B Condensed Matter* 30.4 (1978): 429-436.
- [60] Graham, Robert. Path integral formulation of general diffusion processes. *Zeitschrift für Physik B Condensed Matter* 26.3 (1977): 281-290.
- [61] Graham, Robert. Covariant formulation of nonequilibrium statistical thermodynamics. *Zeitschrift für Physik B Condensed Matter* 26.4 (1977): 397-405.
- [62] Kerler, W. Definition of path integrals and rules for nonlinear transformations. *Nuclear Physics B* 139.3 (1978): 312-326.
- [63] Hirshfeld, Allen C. Canonical and covariant path integrals. *Physics Letters A* 67.1 (1978): 5-8.
- [64] It has been known [13] that there are many quantum Hamiltonians (9) corresponding to the same classical action (8). These quantum Hamiltonians differ from each other by a term linear in Ricci scalar curvature R , which can be absorbed into the potential energy $V(\mathbf{x})$.
- [65] These transformations are usually called *point transformations* in Hamiltonian mechanics. We do not need to study the more general type of canonical transformations which mix coordinates and momenta, because we do not study phase space integral.

Appendix A: Proof of Lemma 3

We want to prove that the following α -dependent time-slice Green's function (TSGF)

$$\begin{aligned} \mathcal{G}_\alpha(\mathbf{x}; \mathbf{x}_0, \Delta t) d\mu(\mathbf{x}) &= \frac{d\mu(\mathbf{x})}{\sqrt{(4\pi\Delta t)^N D(\mathbf{x}_\alpha)}} \exp \left\{ \right. \\ &\quad - \left(\Delta x^i - F^i(\mathbf{x}_\alpha)\Delta t + 2\alpha\partial_i D^{il}(\mathbf{x}_\alpha)\Delta t \right) \frac{D_{ij}^{-1}(\mathbf{x}_\alpha)}{4\Delta t} \left(\Delta x^j - F^j(\mathbf{x}_\alpha)\Delta t + 2\alpha\partial_k D^{jk}(\mathbf{x}_\alpha)\Delta t \right) \\ &\quad \left. - \alpha\partial_i F^i(\mathbf{x}_\alpha)\Delta t + \alpha^2\partial_i\partial_j D^{ij}(\mathbf{x}_\alpha)\Delta t - \Phi(\mathbf{x}_\alpha)\Delta t \right\}, \end{aligned} \quad (\text{A1})$$

with $d\mu(\mathbf{x}) = d^d x$ and $\mathbf{x}_\alpha = \mathbf{x} + \alpha\Delta\mathbf{x}_0$ is asymptotically independent of α . We only need to prove that Eq. (A1) is equivalent to the $\alpha = 0$ version:

$$\mathcal{G}(\mathbf{x}; \mathbf{x}_0, \Delta t) d\mu(\mathbf{x}) = \frac{d\mu(\mathbf{x})}{\sqrt{(4\pi\Delta t)^N D(\mathbf{x}_0)}} \exp \left\{ - \left(\Delta x^i - F^i(\mathbf{x}_0, t)\Delta t \right) \frac{D_{ij}^{-1}(\mathbf{x}_0)}{4\Delta t} \left(\Delta x^j - F^j(\mathbf{x}_0, t)\Delta t \right) - \Phi(\mathbf{x}_0)\Delta t \right\}, \quad (\text{A2})$$

Here and below we use $D(\mathbf{x})$ to denote the determinant of the symmetric matrix $D^{ij}(\mathbf{x})$. Note here that \mathbf{x} are the stochastic variables and \mathbf{x}_0 are fixed parameters. $\Delta\mathbf{x} = \mathbf{x} - \mathbf{x}_0$ have averages $F^i(\mathbf{x}_0)dt$ and variances $2D^{ij}(\mathbf{x}_0)\Delta t$. All higher order moments are zero. Note that Eq. (A2) is not normalized in usual sense if $\Phi(\mathbf{x}) \neq 0$.

Such an equivalent class of infinitesimal time transition probability/amplitude was derived using formal methods by Wissel in 1979 [55]. This result however has not received much attention, due to the lack of mathematical rigor in Wissel's derivation. Our proof of Eq. (A1) is based on asymptotic analyses, and is mathematically rigorous.

Because of Lemma 1, we only need to show that Eqs. (A2) and (A1) have equal moments up to the order Δt . Mathematically we want to prove:

$$\int (\mathcal{G}_\alpha - \mathcal{G}) d\mathbf{x} = o(\Delta t), \quad (\text{A3a})$$

$$\int (\mathcal{G}_\alpha - \mathcal{G}) \Delta x^i d\mathbf{x} = o(\Delta t), \quad (\text{A3b})$$

$$\int (\mathcal{G}_\alpha - \mathcal{G}) \Delta x^i \Delta x^j d\mathbf{x} = o(\Delta t), \quad (\text{A3c})$$

$$\dots\dots\dots = o(\Delta t). \quad (\text{A3d})$$

Since $\Delta\mathbf{x} \sim \sqrt{\Delta t}$, we see that the difference between $\Phi(\mathbf{x}_0)\Delta t$ and $\Phi(\mathbf{x}_\alpha)\Delta t$ is of order $\Delta t^{3/2}$ and hence makes no contribution to the dynamics in the continuum limit. It then follows that we only need to prove the above conditions for the case $\Phi = 0$. Below we will set $\Phi = 0$.

It is then convenient to introduce a set of scaled variables ξ^i via

$$\Delta x^i = \xi^i \sqrt{\Delta t} + F^i(\mathbf{x}_0)\Delta t, \quad (\text{A4})$$

which makes it clear that both the average and variance of $\Delta\mathbf{x}$ are of order Δt . The joint pdf Eq. (A2) then becomes

$$p(\boldsymbol{\xi}; \mathbf{x}) d^d \xi \equiv \frac{d^d \xi}{\sqrt{(4\pi)^d D(\mathbf{x}_0)}} e^{-\xi^i D_{ij}^{-1}(\mathbf{x}_0) \xi^j / 4}, \quad (\text{A5})$$

with shows that $\boldsymbol{\xi}$ have vanishing averages and variances $2D^{ij}(\mathbf{x})$. We can similarly rewrite Eq. (A1) in terms of $\boldsymbol{\xi}$ and Δt through Eq. (A4). We further expand the exponent in terms of Δt , up the first order. Higher order terms do not contribute to the continuum dynamics. The resulting expression is

$$p_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x}_0) d^d \xi = \frac{d^d \xi}{\sqrt{(4\pi)^d \det D_{ij}(\mathbf{x}_0)}} e^{-\xi^i D_{ij}^{-1}(\mathbf{x}_0) \xi^j / 4 - \delta\mathcal{A}_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x}_0)}, \quad (\text{A6a})$$

$$\delta\mathcal{A}_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x}_0) = \mathcal{A}^{1/2}(\boldsymbol{\xi}; \mathbf{x}_0) \sqrt{\Delta t} + \mathcal{A}^1(\boldsymbol{\xi}; \mathbf{x}_0) \Delta t + \dots, \quad (\text{A6b})$$

where $\delta\mathcal{A}_\alpha$ gives all non-Gaussian corrections to the distribution:

$$\mathcal{A}^{1/2} = -\frac{\alpha}{4} (\mathbf{D}^{-1} (\partial_k \mathbf{D}) \mathbf{D}^{-1})_{ij} \xi^i \xi^j \xi^k + \frac{\alpha}{2} \text{Tr}(\mathbf{D}^{-1} \partial_i \mathbf{D}) \xi^i + \alpha D_{ij}^{-1} (\partial_l D^{jl}) \xi^i, \quad (\text{A7a})$$

$$\mathcal{A}^1 = \mathcal{A}^{1,1} + \mathcal{A}^{1,2} + \mathcal{A}^{1,3} + \mathcal{A}^{1,4} + \partial_i F^i - \alpha^2 \partial_i \partial_j D^{ij}, \quad (\text{A7b})$$

$$\mathcal{A}^{1,1} = -\frac{\alpha}{4} (\mathbf{D}^{-1} (\partial_k \mathbf{D}) \mathbf{D}^{-1})_{ij} F^k \xi^i \xi^j + \frac{\alpha}{2} \text{Tr}(\mathbf{D}^{-1} (\partial_i \mathbf{D})) F^i, \quad (\text{A7c})$$

$$\mathcal{A}^{1,2} = -\frac{\alpha}{2} (\partial_k F^i) D_{ij}^{-1} \xi^k \xi^j, \quad (\text{A7d})$$

$$\mathcal{A}^{1,3} = -\frac{\alpha^2}{8} (\mathbf{D}^{-1} (\partial_i \partial_j \mathbf{D}) \mathbf{D}^{-1})_{lm} \xi^i \xi^j \xi^l \xi^m + \alpha^2 D_{ij}^{-1} (\partial_k \partial_l D^{ik}) \xi^j \xi^l + \frac{\alpha^2}{4} \text{Tr}(\mathbf{D}^{-1} \partial_i \partial_j \mathbf{D}) \xi^i \xi^j, \quad (\text{A7e})$$

$$\begin{aligned} \mathcal{A}^{1,4} &= \frac{\alpha^2}{4} (\mathbf{D}^{-1} (\partial_i \mathbf{D}) \mathbf{D}^{-1} (\partial_j \mathbf{D}) \mathbf{D}^{-1})_{mn} \xi^i \xi^j \xi^m \xi^n - \frac{\alpha^2}{4} \text{Tr}(\mathbf{D}^{-1} (\partial_i \mathbf{D}) \mathbf{D}^{-1} (\partial_j \mathbf{D})) \xi^i \xi^j \\ &\quad - \alpha^2 (\partial_l D^{il}) (\mathbf{D}^{-1} (\partial_k \mathbf{D}) \mathbf{D}^{-1})_{ij} \xi^k \xi^j + \alpha^2 (\partial_l D^{il}) D_{ij}^{-1} (\partial_m D^{jm}), \end{aligned} \quad (\text{A7f})$$

where it is understood that all functions in Eqs. (A7) are evaluated at \mathbf{x}_0 .

Note that in the limit $\Delta t \rightarrow 0$, every term in the RHS of Eq. (A6b) vanishes, and Eq. (A6a) reduces to Eq. (A5), which is Gaussian. It then follows that all $n > 2$ order moments κ_n of $\boldsymbol{\xi}$ calculated from Eq. (A6a) vanish in the limit $\Delta t \rightarrow 0$. Using Eq. (A4), we see that all n -th order moments of $\Delta \mathbf{x}$ (for $n > 2$) calculated from Eq. (A1) are at least of order $\Delta t^{3/2}$, independent of α . It then follows that we only need to prove the first three conditions of Eqs. (A3).

In terms of ξ , the first condition, Eq. (A3a) becomes

$$\int p_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x}) d^d \xi - \int p(\boldsymbol{\xi}; \mathbf{x}) d^d \xi = o(\Delta t). \quad (\text{A8})$$

When calculating the first integral above, we can expand $p_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x})$ in terms of Δt using Eqs. (A6). To the right order, then Eq. (A8) becomes

$$\int \left[\frac{1}{2} (\mathcal{A}^{1/2})^2 - \mathcal{A}^1 \right] p(\boldsymbol{\xi}; \mathbf{x}) d^d \xi = o(1). \quad (\text{A9})$$

The proof of this condition is very complicated, and is presented in Sec. A 1.

Using Eq. (A4), the second condition, Eq. (A3b), can be rewritten as

$$\int (p_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x}) - p(\boldsymbol{\xi}; \mathbf{x})) (\xi^i \sqrt{\Delta t} + F^i(\mathbf{x}) \Delta t) d^d \xi = o(\Delta t). \quad (\text{A10})$$

But $F^i(\mathbf{x}) \Delta t$ is independent of $\boldsymbol{\xi}$, and hence drops out due to Eq. (A8). On the other hand, average of ξ^i with respect to $p(\boldsymbol{\xi}; \mathbf{x})$ vanishes identically. Hence Eq. (A3b) reduces to

$$\int p_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x}) \xi^i d^d \xi = o(\sqrt{\Delta t}). \quad (\text{A11})$$

Again using Eqs. (A6b) to expand $p_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x})$ in terms of Δt , we further rewrite Eq. (A11) into

$$\int \mathcal{A}^{1/2}(\boldsymbol{\xi}; \mathbf{x}) \xi^i p(\boldsymbol{\xi}; \mathbf{x}) d^d \xi = o(1). \quad (\text{A12})$$

Let us define for arbitrary function $f(\boldsymbol{\xi})$:

$$\langle f(\boldsymbol{\xi}) \rangle \equiv \int p(\boldsymbol{\xi}; \mathbf{x}) f(\boldsymbol{\xi}) d^d \xi \quad (\text{A13})$$

Because $p(\boldsymbol{\xi}; \mathbf{x}) d^d \xi$ is Gaussian, Eq. (A12) then can be calculated using Wick's theorem:

$$\begin{aligned} &\int \mathcal{A}^{1/2}(\boldsymbol{\xi}; \mathbf{x}) \xi^i p(\boldsymbol{\xi}; \mathbf{x}) d^d \xi = \langle \mathcal{A}^{1/2}(\boldsymbol{\xi}; \mathbf{x}) \xi^i \rangle \\ &= -\frac{\alpha}{4} (\mathbf{D}^{-1} (\partial_k \mathbf{D}) \mathbf{D}^{-1})_{ij} \langle \xi^i \xi^j \xi^k \xi^l \rangle + \frac{\alpha}{2} \text{Tr}(\mathbf{D}^{-1} \partial_l \mathbf{D}) \langle \xi^i \xi^l \rangle + \alpha D_{kj}^{-1} (\partial_l D^{jl}) \langle \xi^i \xi^k \rangle + o(1) \\ &= -\alpha (\mathbf{D}^{-1} (\partial_k \mathbf{D}) \mathbf{D}^{-1})_{lj} (D^{ij} D^{kl} + D^{ik} D^{jl} + D^{il} D^{jk}) + \alpha D^{il} D_{jk}^{-1} \partial_l D^{jk} + 2\alpha D_{kj}^{-1} (\partial_l D^{jl}) D^{ik} + o(1) \\ &= -\alpha (\partial_k D^{ki} + D^{ik} D_{lm}^{-1} \partial_k D^{ml} + \partial_k D^{ik}) + \alpha D^{il} D_{jk}^{-1} \partial_l D^{jk} + 2\alpha \partial_l D^{il} + o(1) \\ &= o(1). \end{aligned} \quad (\text{A15})$$

Finally, in terms of $\boldsymbol{\xi}$, the third condition, Eq. (A3c), becomes

$$\int p_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x}) \xi^i \xi^j d^d \xi - \int p(\boldsymbol{\xi}; \mathbf{x}) \xi^i \xi^j d^d \xi = o(1). \quad (\text{A16})$$

But this is trivially valid, since we already know that as $\Delta t \rightarrow 0$, $p_\alpha(\boldsymbol{\xi}; \Delta t, \mathbf{x})$ converges to $p(\boldsymbol{\xi}; \mathbf{x})$. Hence we have proved that Eqs. (A1) and (A2) are equivalent to each other.

1. Proof of Eq. (A9)

We will prove here the condition Eq. (A9), which can be rewritten into the following form:

$$\left\langle \mathcal{A}^1 - \frac{1}{2} \left(\mathcal{A}^{1/2} \right)^2 \right\rangle = o(1), \quad (\text{A17})$$

which means that the LHS vanishes in the limit $\Delta t \rightarrow 0$.

We have split \mathcal{A}^1 in Eq. (A7b) into six parts. Let us calculate the average of each part:

$$\begin{aligned} \langle \mathcal{A}^{1,1} \rangle &= -\frac{1}{4} (\alpha \mathbf{D}^{-1} (\partial_k \mathbf{D}) \mathbf{D}^{-1})_{ij} F_1^k \langle \xi^i \xi^j \rangle + \frac{1}{2} \text{Tr}(\alpha \mathbf{D}^{-1} (\partial_i \mathbf{D})) F^i \\ &= -\frac{\alpha}{2} D^{ij} D_{im}^{-1} (\partial_k D^{ml}) D_{lj}^{-1} F^k + \frac{\alpha}{2} D_{jk}^{-1} (\partial_i D^{jk}) F^i = 0, \end{aligned} \quad (\text{A18})$$

$$\langle \mathcal{A}^{1,2} \rangle = -\frac{\alpha}{2} (\partial_k F^i) D_{ij}^{-1} \langle \xi^k \xi^j \rangle = -\frac{\alpha}{2} 2D^{kj} D_{ij}^{-1} \partial_k F^i = -\partial_i F^i, \quad (\text{A19})$$

$$\begin{aligned} \langle \mathcal{A}^{1,3} \rangle &= -\frac{\alpha^2}{8} D_{lm}^{-1} (\partial_i \partial_j D^{mn}) \langle \xi^i \xi^j \xi^l \xi^n \rangle + \alpha^2 D_{ij}^{-1} (\partial_k \partial_l D^{ik}) \langle \xi^j \xi^l \rangle + \frac{\alpha^2}{4} \text{Tr}(\mathbf{D}^{-1} (\partial_i \partial_j \mathbf{D}) \mathbf{D}^{-1}) \langle \xi^i \xi^j \rangle \\ &= -\frac{\alpha^2}{2} D_{lm}^{-1} (\partial_i \partial_j D^{mn}) (D^{ij} D^{ln} + D^{il} D^{jn} + D^{in} D^{jl}) + 2\alpha^2 D_{ij}^{-1} (\partial_k \partial_l D^{ik}) D^{jl} + \frac{\alpha^2}{2} D^{ij} D_{kl}^{-1} \partial_i \partial_j D^{lk} \\ &= \alpha^2 \partial_i \partial_j D^{ij}. \end{aligned} \quad (\text{A20})$$

Hence we have

$$\langle \mathcal{A}^{1,1} + \mathcal{A}^{1,2} + \mathcal{A}^{1,3} + \partial_i F^i - \alpha^2 \partial_i \partial_j D^{ij} \rangle = 0. \quad (\text{A21})$$

Hence to prove Eq. (A17), we only need to show

$$\langle \mathcal{A}^{1,4} \rangle - \frac{1}{2} \left\langle \left(\mathcal{A}^{1/2} \right)^2 \right\rangle = 0. \quad (\text{A22})$$

Using Eq. (A7f) and Wick's theorem, we have

$$\begin{aligned} \langle \mathcal{A}^{1,4} \rangle &= \frac{\alpha^2}{4} (\mathbf{D}^{-1} (\partial_i \mathbf{D}) \mathbf{D}^{-1} (\partial_j \mathbf{D}) \mathbf{D}^{-1})_{mn} \langle \xi^i \xi^j \xi^m \xi^n \rangle - \frac{\alpha^2}{4} \text{Tr}(\mathbf{D}^{-1} (\partial_i \mathbf{D}) \mathbf{D}^{-1} (\partial_j \mathbf{D})) \langle \xi^i \xi^j \rangle \\ &\quad - \alpha^2 (\partial_l D^{il}) (\mathbf{D}^{-1} (\partial_k \mathbf{D}) \mathbf{D}^{-1})_{ij} \langle \xi^k \xi^j \rangle + \alpha^2 (\partial_l D^{il}) D_{ij}^{-1} (\partial_m D^{jm}) \\ &= \alpha^2 \left[(\partial_m D^{jk}) D_{kl}^{-1} (\partial_n D^{lr}) D_{rj}^{-1} D^{mn} + (\partial_m D^{mk}) D_{kl}^{-1} (\partial_n D^{ln}) + (\partial_m D^{sk}) D_{kl}^{-1} (\partial_s D^{lm}) \right. \\ &\quad \left. - \frac{1}{2} D_{ik}^{-1} (\partial_s D^{kl}) D_{im}^{-1} (\partial_t D^{mi}) D^{st} - 2(\partial_s D^{st}) D_{ij}^{-1} (\partial_l D^{jl}) + (\partial_l D^{il}) D_{ij}^{-1} (\partial_k D^{jk}) \right] \\ &= \alpha^2 \left[\frac{1}{2} (\partial_m D^{jk}) D_{kl}^{-1} (\partial_n D^{lr}) D_{rj}^{-1} D^{mn} + (\partial_m D^{sk}) D_{kl}^{-1} (\partial_s D^{lm}) \right]. \end{aligned} \quad (\text{A23})$$

Using Eq. (A7a) and expanding, and further using Wick's theorem, we find

$$\frac{1}{2} \left\langle \left(\mathcal{A}^{1/2} \right)^2 \right\rangle = \alpha^2 [\langle \mathcal{A}^{2,1} \rangle + \langle \mathcal{A}^{2,2} \rangle + \langle \mathcal{A}^{2,3} \rangle + \langle \mathcal{A}^{2,4} \rangle], \quad (\text{A24})$$

$$\langle \mathcal{A}^{2,1} \rangle = \frac{1}{4} D^{ij} D_{mn}^{-1} (\partial_i D^{mn}) D_{st}^{-1} (\partial_j D^{st}) + (\partial_l D^{jl}) D_{jm}^{-1} (\partial_n D^{mn}) + D_{kl}^{-1} (\partial_m D^{kl}) (\partial_n D^{mn}),$$

$$\langle \mathcal{A}^{2,2} \rangle = \frac{1}{4} D_{il}^{-1} (\partial_k D^{lm}) D_{mj}^{-1} D_{sp}^{-1} (\partial_r D^{pq}) D_{qt}^{-1} (D^{ij} D^{kr} D^{st} + \text{other 14 pairing terms}),$$

$$\langle \mathcal{A}^{2,3} \rangle = -\frac{1}{2} D_{lj}^{-1} (\partial_i D^{jl}) D_{pm}^{-1} (\partial_k D^{nm}) D_{nq}^{-1} (D^{ik} D^{pq} + D^{ip} D^{kq} + D^{iq} D^{pk}),$$

$$\langle \mathcal{A}^{2,4} \rangle = -D_{ij}^{-1} (\partial_l D^{lj}) D_{pm}^{-1} (\partial_k D^{nm}) D_{nq}^{-1} (D^{ik} D^{pq} + D^{ip} D^{kq} + D^{iq} D^{pk}). \quad (\text{A25})$$

Adding up all these terms (which demands great patience), we find

$$\frac{1}{2} \left\langle \left(\mathcal{A}^{1/2} \right)^2 \right\rangle = \frac{\alpha^2}{2} (\partial_m D^{jk}) D_{kl}^{-1} (\partial_n D^{lr}) D_{rj}^{-1} D^{mn} + \alpha^2 (\partial_m D^{sk}) D_{kl}^{-1} (\partial_s D^{lm}). \quad (\text{A26})$$

Combining this with Eq. (A23), we obtain Eq. (A9) as expected. This completes our proof of equivalence between Eqs. (A1) and (A2).
