

Coupled and uncoupled sign-changing spikes of singularly perturbed elliptic systems

Mónica Clapp* and Mayra Soares†

July 11, 2022

Abstract

We study the existence and asymptotic behavior of solutions having positive and sign-changing components to the singularly perturbed system of elliptic equations

$$\begin{cases} -\varepsilon^2 \Delta u_i + u_i = \mu_i |u_i|^{p-2} u_i + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i, \\ u_i \in H_0^1(\Omega), \quad u_i \neq 0, \quad i = 1, \dots, \ell, \end{cases}$$

in a bounded domain Ω in \mathbb{R}^N , with $N \geq 4$, $\varepsilon > 0$, $\mu_i > 0$, $\lambda_{ij} = \lambda_{ji} < 0$, $\alpha_{ij}, \beta_{ij} > 1$, $\alpha_{ij} = \beta_{ji}$, $\alpha_{ij} + \beta_{ij} = p \in (2, 2^*)$, and $2^* := \frac{2N}{N-2}$.

If Ω is the unit ball we obtain solutions with a prescribed combination of positive and nonradial sign-changing components exhibiting two different types of asymptotic behavior as $\varepsilon \rightarrow 0$: solutions whose limit profile is a rescaling of a solution with positive and nonradial sign-changing components of the limit system

$$\begin{cases} -\Delta u_i + u_i = \mu_i |u_i|^{p-2} u_i + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i, \\ u_i \in H^1(\mathbb{R}^N), \quad u_i \neq 0, \quad i = 1, \dots, \ell, \end{cases}$$

and solutions whose limit profile is a solution of the uncoupled system, i.e., after rescaling and translation, the limit profile of the i -th component is a positive or a nonradial sign-changing solution to the equation

$$-\Delta u + u = \mu_i |u|^{p-2} u, \quad u \in H^1(\mathbb{R}^N), \quad u \neq 0.$$

KEYWORDS: Nonlinear elliptic system, singularly perturbed, weakly coupled, competitive, positive and sign-changing spikes.

MSC2020: 35J57 (35B06, 35B25, 35B40, 47J30).

1 Introduction

We consider the following system of singularly perturbed elliptic equations

$$(\mathcal{S}_{\varepsilon, \Omega}) \quad \begin{cases} -\varepsilon^2 \Delta u_i + u_i = \mu_i |u_i|^{p-2} u_i + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i, \\ u_i \in H_0^1(\Omega), \quad u_i \neq 0, \quad i = 1, \dots, \ell, \end{cases}$$

where $\varepsilon > 0$ is a small parameter, Ω is a bounded smooth domain in \mathbb{R}^N which contains the origin, $N \geq 2$, $\mu_i > 0$, $\lambda_{ij} = \lambda_{ji} < 0$, $\alpha_{ij}, \beta_{ij} > 1$, $\alpha_{ij} = \beta_{ji}$, $\alpha_{ij} + \beta_{ij} = p \in (2, 2^*)$, and 2^* is the critical Sobolev exponent (i.e., $2^* := \frac{2N}{N-2}$ if $N \geq 3$ and $2^* := \infty$ if $N = 2$).

*M. Clapp was supported by CONACYT (Mexico) through the grant for the research project A1-S-10457.

†M. Soares was supported by UNAM-DGAPA (Mexico) through a postdoctoral fellowship.

This system arises as a model for various physical phenomena, in particular in the study of standing waves for a mixture of Bose–Einstein condensates of ℓ different hyperfine states which overlap in space, see for example [10]. Here we consider the case in which the interaction between particles in the same state is attractive ($\mu_i > 0$) and the interaction between particles in any two different states is repulsive ($\lambda_{ij} < 0$).

In their seminal paper [12] Lin and Wei described the behavior of positive least energy solutions for the system $(\mathcal{S}_{\varepsilon, \Omega})$ with cubic nonlinearity ($\alpha_{ij} = \beta_{ij} = 2$) as $\varepsilon \rightarrow 0$. They showed that each component is a spike, i.e., a rescaling of the positive ground state solution to the problem

$$(\mathcal{P}_i) \quad \begin{cases} -\Delta u + u = \mu_i |u|^{p-2} u, \\ u \in H^1(\mathbb{R}^N), \quad u \neq 0, \end{cases}$$

and that, as $\varepsilon \rightarrow 0$, the centers of the spikes approach a sphere-packing position in Ω , i.e., a configuration of points maximizing the distances among them and to the boundary of Ω . Multiple positive solutions and a numerical description of them is given in [25].

Our main objective is to study the existence and profile of solutions to $(\mathcal{S}_{\varepsilon, \Omega})$ some of whose components can be positive while others change sign. It is reasonable to expect that there will be solutions with sign-changing spikes, i.e., solutions whose sign-changing components look like rescalings of a sign-changing solution to the problem (\mathcal{P}_i) . The existence of nonradial sign-changing solutions to (\mathcal{P}_i) was first established by Bartsch and Willem in [2] for dimensions $N = 4$ and $N \geq 6$, and by Lorca and Ubilla for $N = 5$ in [16], taking advantage of some symmetry properties of \mathbb{R}^N . Other solutions of this type were found in [7].

On the other hand, rescaling the components by $\tilde{u}_i(x) := u_i(\varepsilon x)$ the system $(\mathcal{S}_{\varepsilon, \Omega})$ becomes

$$\begin{cases} -\Delta u_i + u_i = \mu_i |u_i|^{p-2} u_i + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i, \\ u_i \in H_0^1(\Omega_\varepsilon), \quad u_i \neq 0, \quad i = 1, \dots, \ell, \end{cases}$$

in $\Omega_\varepsilon := \{x \in \mathbb{R}^N : \varepsilon x \in \Omega\}$. As $\varepsilon \rightarrow 0$ these domains cover the whole space \mathbb{R}^N . So it is natural to ask whether the system $(\mathcal{S}_{\varepsilon, \Omega})$ has a solution that, after rescaling, approaches a solution to the system

$$(\mathcal{S}_{\infty, \ell}) \quad \begin{cases} -\Delta u_i + u_i = \mu_i |u_i|^{p-2} u_i + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i, \\ u_i \in H^1(\mathbb{R}^N), \quad u_i \neq 0, \quad i = 1, \dots, \ell. \end{cases}$$

As shown by Lin and Wei in [13, Theorem 1], this system does not have a ground state solution. On the other hand, Sirakov showed in [23] that it does have a positive least energy radial solution (i.e., every component u_i is positive and radial); see also [4, 26] and the references therein.

One might expect to obtain solutions with positive and sign-changing components for the system $(\mathcal{S}_{\varepsilon, \Omega})$ whose limit profile is a solution of the same type for the system $(\mathcal{S}_{\infty, \ell})$. For this last system with cubic nonlinearity (hence, $N \leq 3$) Sato and Wang [22] established the existence of least energy solutions of this type whose components are radial; see also [5, 14]. The following result provides solutions with nonradial sign-changing components. It is proved in Section 3.

Theorem 1.1. *Let $N = 4$ or $N \geq 6$. Then, for any given $0 \leq m \leq \ell$, the system $(\mathcal{S}_{\infty, \ell})$ has a solution $\mathbf{w} = (w_1, \dots, w_\ell)$ whose first m components w_1, \dots, w_m are positive and whose last $\ell - m$ components w_{m+1}, \dots, w_ℓ are nonradial and change sign. Furthermore, \mathbf{w} satisfies*

$$(1.1) \quad \begin{cases} w_i(z_1, z_2, x) = w_i(e^{i\vartheta} z_1, e^{i\vartheta} z_2, gx) & \text{for all } \vartheta \in [0, 2\pi), g \in O(N-4), i = 1, \dots, \ell, \\ w_i(z_1, z_2, x) = w_i(z_2, z_1, x) & \text{if } i = 1, \dots, m, \\ w_i(z_1, z_2, x) = -w_i(z_2, z_1, x) & \text{if } i = m+1, \dots, \ell, \end{cases}$$

for all $(z_1, z_2, x) \in \mathbb{C} \times \mathbb{C} \times \mathbb{R}^{N-4} \equiv \mathbb{R}^N$, and it has least energy among all nontrivial solutions with these symmetry properties.

The symmetries of the sign-changing components w_{m+1}, \dots, w_ℓ are basically the same as those considered in [2]. So one may wonder whether one can obtain a similar result for $N = 5$ using the symmetries introduced in [16] or [7]. As we shall see, this is not possible. A key role is played by the space of fixed points of the group of symmetries involved, see Theorem 3.4.

For the singularly perturbed system $(\mathcal{S}_{\varepsilon, \Omega})$ in a domain Ω having suitable symmetries we obtain solutions with a prescribed combination of positive and sign-changing components exhibiting two different types of asymptotic behavior as $\varepsilon \rightarrow 0$. We get solutions whose limit profile is a rescaling of a solution to the limit system $(\mathcal{S}_{\infty, \ell})$ with positive and sign-changing components, and solutions whose limit profile is a solution to the uncoupled system, i.e., the limit profile of the i -th component is a rescaling of a positive or a sign-changing solution to the problem (\mathcal{P}_i) . To illustrate our results, let us focus on the case where Ω is the open unit ball $B_1(0)$ in \mathbb{R}^N centered at the origin. For $\varepsilon > 0$ and $u \in H^1(\mathbb{R}^N)$ let

$$\|u\|_\varepsilon^2 := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} [\varepsilon^2 |\nabla u|^2 + u^2] \quad \text{and} \quad \|u\| := \|u\|_1.$$

The following two theorems are special cases of Theorem 4.1, which is stated and proved in Section 4.

Theorem 1.2. *Let $N = 4$ or $N \geq 6$, and $\Omega = B_1(0)$. Then, for any given $0 \leq m \leq \ell$ and any sequence (ε_k) of positive numbers converging to zero, there exists a solution $\widehat{\mathbf{u}}_k = (\widehat{u}_{1k}, \dots, \widehat{u}_{\ell k})$ to the system $(\mathcal{S}_{\varepsilon_k, \Omega})$ whose first m components are positive and whose last $\ell - m$ components are nonradial and change sign, with the following limit profile:*

There exists a fully nontrivial solution $\mathbf{w} = (w_1, \dots, w_\ell)$ to the system $(\mathcal{S}_{\infty, \ell})$ such that, after passing to a subsequence,

$$\lim_{k \rightarrow \infty} \|\widehat{u}_{ik} - w_i(\varepsilon_k^{-1} \cdot)\|_{\varepsilon_k} = 0 \quad \text{for all } i = 1, \dots, \ell.$$

The first m components of \mathbf{w} are positive, its last $\ell - m$ components are nonradial and change sign, and \mathbf{w} satisfies (1.1). Therefore,

$$\lim_{k \rightarrow \infty} \sum_{i=1}^{\ell} \|\widehat{u}_{ik}\|_{\varepsilon_k}^2 = \sum_{i=1}^{\ell} \|w_i\|^2 =: \widehat{\mathbf{c}}_m.$$

Theorem 1.3. *Let $N \geq 5$ and $\Omega = B_1(0)$. Then, for any given $0 \leq m \leq \ell$ and any sequence (ε_k) of positive numbers converging to zero, there exists a solution $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k})$ to the system $(\mathcal{S}_{\varepsilon_k, \Omega})$ whose first m components are positive and whose last $\ell - m$ components are nonradial and change sign, with the following limit profile:*

For each $i = 1, \dots, \ell$, there exist a sequence (ξ_{ik}) in $B_1(0)$ and a nontrivial solution v_i to the problem (\mathcal{P}_i) such that, after passing to a subsequence,

$$\lim_{k \rightarrow \infty} \varepsilon_k^{-1} \text{dist}(\xi_{ik}, \partial B_1(0)) = \infty, \quad \lim_{k \rightarrow \infty} \varepsilon_k^{-1} |\xi_{ik} - \xi_{jk}| = \infty \text{ if } i \neq j, \quad \lim_{k \rightarrow \infty} \|u_{ik} - v_i(\varepsilon_k^{-1}(\cdot - \xi_{ik}))\|_{\varepsilon_k} = 0.$$

The functions v_1, \dots, v_m are positive and radial, while the functions v_{m+1}, \dots, v_ℓ are sign-changing, nonradial and satisfy

$$(1.2) \quad \begin{cases} v_i(z_1, z_2, x) = v_i(e^{i\vartheta} z_1, e^{i\vartheta} z_2, gx) & \text{for all } \vartheta \in [0, 2\pi), g \in O(N-4), \\ v_i(z_1, z_2, x) = -v_i(z_2, z_1, x), \end{cases}$$

for all $(z_1, z_2, x) \in \mathbb{C} \times \mathbb{C} \times \mathbb{R}^{N-4} \equiv \mathbb{R}^N$, $i = m+1, \dots, \ell$. Furthermore,

$$\lim_{k \rightarrow \infty} \sum_{i=1}^{\ell} \|u_{ik}\|_{\varepsilon_k}^2 = \sum_{i=1}^{\ell} \|v_i\|^2 =: \mathbf{c}_m,$$

satisfies $\mathbf{c}_m < \widehat{\mathbf{c}}_m$, with $\widehat{\mathbf{c}}_m$ as in Theorem 1.2, if $N \geq 6$.

These last two results exhibit some interesting facts. The solutions given by Theorem 1.3 are least energy solutions to the system $(\mathcal{S}_{\varepsilon,\Omega})$ having some specific symmetries described in Example 2.1(ii) below. Their components behave as expected in the repulsive case $\lambda_{ij} < 0$, i.e., they concentrate at points that are far from each other and from the boundary of the ball. On the other hand, the solutions given by Theorem 1.2 behave as shown by Lin and Wei [12, Theorem 1.1(2)] for the attractive case $\lambda_{ij} > 0$, i.e., all components concentrate at the origin. The solutions given by Theorem 1.2 enjoy the same symmetries as those given by Theorem 1.3 (and more), but they have higher energy. This shows that the sign of the interaction coefficient λ_{ij} is not determinant in the segregation behavior of higher energy solutions.

Another interesting feature of the solutions given by Theorem 1.2 is that the limit profiles of their components are coupled, i.e., they solve the system $(\mathcal{S}_{\infty,\ell})$, up to rescaling. A similar behavior can be seen for least energy positive radial solutions to $(\mathcal{S}_{\varepsilon,\Omega})$ in the unit ball: their limit profile is a positive least energy radial solution to the system $(\mathcal{S}_{\infty,\ell})$, see Theorem 4.3. As far as we know, these are the first examples in the literature that show this kind of asymptotic behavior.

To prove our results we follow the approach introduced in [7]. Using concentration compactness techniques, we carry out a careful analysis of the behavior of minimizing sequences for the system $(\mathcal{S}_{\infty,\ell})$ that have a specific type of symmetries, described in Section 2. The symmetries can be chosen to produce a change of sign by construction. As $\varepsilon \rightarrow 0$ the components of the system concentrate at fixed points of the group action. If the space of fixed points is trivial, all components will necessarily concentrate at the origin, but they will move far away from each other when the fixed-point space has positive dimension. A detailed description is given in Theorem 3.3.

We note that for $N = 3$ there are no symmetries with the properties required to produce nonradial sign-changing solutions. For the single equation (\mathcal{P}_i) the existence of this kind of solutions was shown by Musso, Pacard and Wei in [18] using the Lyapunov-Schmidt reduction procedure.

We wish to point out that solutions with sign-changing components for a system related to the existence of optimal partitions for the eigenvalue problem were obtained by Tavares and Terracini in [24]. For the system $(\mathcal{S}_{1,\Omega})$ with fixed $\varepsilon = 1$ and cubic nonlinearity (hence, $N \leq 3$) in a bounded domain Ω , existence and multiplicity of sign-changing and semi-nodal solutions (some components change sign and others are positive) have been established by several authors, see, e.g., [5, 21, 22].

This paper is organized as follows: in Section 2 we introduce the symmetric variational setting for systems with positive and sign-changing components and establish the existence of minimizers for the system $(\mathcal{S}_{\varepsilon,\Omega})$. In Section 3 we give a detailed description of the symmetric minimizing sequences for the system $(\mathcal{S}_{\infty,\ell})$ and prove Theorem 1.1. In Section 4 we state and prove a general result for the system $(\mathcal{S}_{\varepsilon,\Omega})$ and derive Theorems 1.2 and 1.3 from it.

2 Minimizers with positive and sign-changing components

Let G be a closed subgroup of the group $O(N)$ of linear isometries of \mathbb{R}^N and denote by $Gx := \{gx : g \in G\}$ the G -orbit of $x \in \mathbb{R}^N$. Let $\phi : G \rightarrow \mathbb{Z}_2 := \{-1, 1\}$ be a continuous homomorphism of groups with the following property:

(A₁) If ϕ is surjective, then there exists $x_0 \in \mathbb{R}^N$ such that $Kx_0 \neq Gx_0$ where $K := \ker \phi$.

Let Θ be an open subset of \mathbb{R}^N which is G -invariant, i.e., $Gx \subset \Theta$ for every $x \in \Theta$. Then, a function $u : \Theta \rightarrow \mathbb{R}$ is called G -invariant if it is constant on Gx for every $x \in \Theta$ and will be called ϕ -equivariant if

$$u(gx) = \phi(g)u(x) \quad \text{for all } g \in G, x \in \Theta.$$

Define

$$H_0^1(\Theta)^\phi := \{u \in H_0^1(\Theta) : u \text{ is } \phi\text{-equivariant}\}.$$

Assumption (A₁) guarantees that $H_0^1(\Theta)^\phi$ has infinite dimension, see [3]. If $\phi \equiv 1$ is the trivial homomorphism, then $H_0^1(\Theta)^\phi$ is the space of G -invariant functions in $H_0^1(\Theta)$. On the other hand, if ϕ is surjective, then every nontrivial function $u \in H_0^1(\Theta)^\phi$ is nonradial and changes sign. The following examples are of relevance to our main results.

Examples 2.1. Consider the following examples of groups of isometries and homomorphisms satisfying (A_1) :

(i) For any group G the trivial homomorphism $\phi \equiv 1$ is not surjective, so (A_1) is trivially satisfied.

(ii) Let Γ be the group generated by $\{e^{i\vartheta} : \vartheta \in [0, 2\pi)\} \cup \{\tau\}$ acting on \mathbb{R}^N by

$$e^{i\vartheta}(z_1, z_2, x) = (e^{i\vartheta}z_1, e^{i\vartheta}z_2, x) \quad \text{and} \quad \tau(z_1, z_2, x) = (z_2, z_1, x) \quad \forall (z_1, z_2, x) \in \mathbb{C} \times \mathbb{C} \times \mathbb{R}^{N-4} \cong \mathbb{R}^N,$$

and let $\phi : \Gamma \rightarrow \mathbb{Z}_2$ be the homomorphism given by $\phi(e^{i\vartheta}) := 1$ and $\phi(\tau) := -1$. The kernel of ϕ is the group $K := \{e^{i\vartheta} : \vartheta \in [0, 2\pi)\}$ and the point $x_0 = (1, 0, 0) \in \mathbb{C} \times \mathbb{C} \times \mathbb{R}^{N-4}$ is such that

$$Kx_0 = \{(e^{i\vartheta}, 0, 0) : \vartheta \in [0, 2\pi)\} \quad \text{and} \quad \Gamma x_0 = \{(e^{i\vartheta}, 0, 0) : \vartheta \in [0, 2\pi)\} \cup \{(0, e^{i\vartheta}, 0) : \vartheta \in [0, 2\pi)\},$$

so (A_1) is satisfied.

(iii) Let $G := \Gamma \times O(N-4)$ with Γ as in (ii) and $g \in O(N-4)$ acting as

$$g(z_1, z_2, x) = (z_1, z_2, gx) \quad \forall (z_1, z_2, x) \in \mathbb{C} \times \mathbb{C} \times \mathbb{R}^{N-4} \cong \mathbb{R}^N.$$

Let $\phi : G \rightarrow \mathbb{Z}_2$ be the homomorphism given by $\phi(e^{i\vartheta}) := 1$, $\phi(\tau) := -1$ and $\phi(g) := 1$ for $g \in O(N-4)$. Then $K := \ker \phi = \{e^{i\vartheta} : \vartheta \in [0, 2\pi)\} \times O(N-4)$ and, if $x_0 = (1, 0, 0) \in \mathbb{C} \times \mathbb{C} \times \mathbb{R}^{N-4}$,

$$Kx_0 = \{(e^{i\vartheta}, 0, 0) : \vartheta \in [0, 2\pi)\} \quad \text{and} \quad Gx_0 = \{(e^{i\vartheta}, 0, 0) : \vartheta \in [0, 2\pi)\} \cup \{(0, e^{i\vartheta}, 0) : \vartheta \in [0, 2\pi)\},$$

so (A_1) is satisfied.

For the remaining of this section we fix a closed subgroup G of $O(N)$ and, for each $i = 1, \dots, \ell$, a continuous homomorphism $\phi_i : G \rightarrow \mathbb{Z}_2$ satisfying (A_1) .

If Θ is a G -invariant open subset of \mathbb{R}^N , we consider the system

$$(\mathcal{J}_{\varepsilon, \Theta}^\phi) \quad \begin{cases} -\varepsilon^2 \Delta u_i + u_i = \mu_i |u_i|^{p-2} u_i + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i, \\ u_i \in H_0^1(\Theta)^{\phi_i}, \quad u_i \neq 0, \quad i = 1, \dots, \ell, \end{cases}$$

with $\varepsilon > 0$, $\mu_i > 0$, $\lambda_{ij} = \lambda_{ji} < 0$, $\alpha_{ij}, \beta_{ij} > 1$, $\alpha_{ij} = \beta_{ji}$ and $\alpha_{ij} + \beta_{ij} = p \in (2, 2^*)$.

As usual, we consider $H_0^1(\Theta)$ as a subspace of $H^1(\mathbb{R}^N)$, identifying a function $u \in H_0^1(\Theta)$ with its trivial extension to \mathbb{R}^N . Set

$$\mathcal{H}^\ell(\Theta) := H_0^1(\Theta)^{\phi_1} \times \dots \times H_0^1(\Theta)^{\phi_\ell},$$

and denote an element in $\mathcal{H}^\ell(\Theta)$ by $\mathbf{u} = (u_1, \dots, u_\ell)$. For each $\varepsilon > 0$ we define

$$\|\mathbf{u}\|_{\ell, \varepsilon} := \left(\sum_{i=1}^{\ell} \|u_i\|_{\varepsilon}^2 \right)^{1/2}, \quad \text{where} \quad \|u_i\|_{\varepsilon}^2 := \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \left[\varepsilon^2 |\nabla u_i|^2 + u_i^2 \right].$$

If $\varepsilon = 1$ we write $\|\mathbf{u}\|_{\ell}$ instead of $\|\mathbf{u}\|_{\ell, 1}$. Note that $\|\cdot\|_{\varepsilon}$ is a norm in $H_0^1(\Theta)$ equivalent to the standard one $\|\cdot\| := \|\cdot\|_1$. Therefore, $\|\cdot\|_{\ell, \varepsilon}$ is a norm in $\mathcal{H}^\ell(\Theta)$ for every $\varepsilon > 0$ and all of these norms are equivalent.

Consider the functional $\mathcal{J}_{\varepsilon}^{\ell} : \mathcal{H}^{\ell}(\Theta) \rightarrow \mathbb{R}$ given by

$$\mathcal{J}_{\varepsilon}^{\ell}(\mathbf{u}) := \frac{1}{2} \sum_{i=1}^{\ell} \|u_i\|_{\varepsilon}^2 - \frac{1}{p} \sum_{i=1}^{\ell} \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \mu_i |u_i|^p - \frac{1}{2} \sum_{\substack{i,j=1 \\ j \neq i}}^{\ell} \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \lambda_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}},$$

which is of class \mathcal{C}^1 . Since $\lambda_{ij} = \lambda_{ji}$, $\beta_{ij} = \alpha_{ji}$ and $\alpha_{ij} + \beta_{ij} = p$, its partial derivatives are given by

$$\begin{aligned}
\partial_i \mathcal{J}_\varepsilon^\ell(\mathbf{u})v &= \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \left(\varepsilon^2 \nabla u_i \cdot \nabla v + u_i v \right) - \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \mu_i |u_i|^{p-2} u_i v \\
&\quad - \frac{1}{2} \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i v - \frac{1}{2} \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \lambda_{ij} \alpha_{ji} |u_i|^{\alpha_{ji}-2} u_i v |u_j|^{\beta_{ji}} \\
(2.1) \quad &= \frac{1}{\varepsilon^N} \left[\int_{\mathbb{R}^N} (\varepsilon^2 \nabla u_i \cdot \nabla v + u_i v) - \int_{\mathbb{R}^N} \mu_i |u_i|^{p-2} u_i v - \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i v \right],
\end{aligned}$$

for $v \in H_0^1(\Theta)^{\phi_i}$ and $i = 1, \dots, \ell$. So, by the principle of symmetric criticality, the solutions to system $(\mathcal{J}_{\varepsilon, \Theta}^\phi)$ are the critical points of $\mathcal{J}_\varepsilon^\ell$ whose components u_i are nontrivial. They belong to the Nehari-type set

$$\mathcal{N}_\varepsilon^\ell(\Theta) := \left\{ \mathbf{u} \in \mathcal{H}^\ell(\Theta) : u_i \neq 0, \partial_i \mathcal{J}_\varepsilon^\ell(\mathbf{u})u_i = 0, \forall i = 1, \dots, \ell \right\}.$$

Note that

$$(2.2) \quad \partial_i \mathcal{J}_\varepsilon^\ell(\mathbf{u})u_i = \|u_i\|_\varepsilon^2 - \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \mu_i |u_i|^p - \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}}.$$

Define

$$c_\varepsilon^\ell(\Theta) := \inf_{\mathbf{u} \in \mathcal{N}_\varepsilon^\ell(\Theta)} \mathcal{J}_\varepsilon^\ell(\mathbf{u}).$$

From (2.1) one sees that

$$(2.3) \quad \mathcal{J}_\varepsilon^\ell(\mathbf{u}) = \frac{p-2}{2p} \sum_{i=1}^{\ell} \|u_i\|_\varepsilon^2 = \frac{p-2}{2p} \|\mathbf{u}\|_{\ell, \varepsilon}^2 \quad \text{if } \mathbf{u} = (u_1, u_2, \dots, u_\ell) \in \mathcal{N}_\varepsilon^\ell(\Theta).$$

Remark 2.2. For $\mathbf{u} = (u_1, \dots, u_\ell) \in \mathcal{H}^\ell(\Theta)$ and $\varepsilon > 0$, define $\tilde{u}_i(z) := u_i(\varepsilon z)$. It is straightforward to verify that $\tilde{u}_i \in H^1(\mathbb{R}^N)^{\phi_i}$ and

$$\|u_i\|_\varepsilon^2 = \|\tilde{u}_i\|^2, \quad \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} |u_i|^p = \int_{\mathbb{R}^N} |\tilde{u}_i|^p, \quad \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}} = \int_{\mathbb{R}^N} |\tilde{u}_j|^{\alpha_{ij}} |\tilde{u}_i|^{\beta_{ij}}.$$

For $\mathbf{u} = (u_1, \dots, u_\ell) \in \mathcal{H}^\ell(\Theta)$ and $\mathbf{t} = (t_1, \dots, t_\ell) \in (0, \infty)^\ell$, we write $\mathbf{t}\mathbf{u} := (t_1 u_1, \dots, t_\ell u_\ell)$. Our next results provide some useful information about the Nehari-type set $\mathcal{N}_\varepsilon^\ell(\Theta)$.

Lemma 2.3. *The following statements hold true:*

(i) $\mathcal{N}_\varepsilon^\ell(\Theta) \neq \emptyset$ and there exists a constant $c_0 > 0$, independent of ε , such that

$$\|u_i\|_\varepsilon^2 > c_0 \quad \text{for every } \mathbf{u} = (u_1, \dots, u_\ell) \in \mathcal{N}_\varepsilon^\ell(\Theta), \quad i = 1, \dots, \ell.$$

Therefore $\mathcal{N}_\varepsilon^\ell(\Theta)$ is a closed subset of $\mathcal{H}^\ell(\Theta)$ and $0 < c_\varepsilon^\ell(\Theta) < \infty$.

(ii) For each $\mathbf{u} \in \mathcal{H}^\ell(\Theta)$ such that

$$(2.4) \quad \int_{\mathbb{R}^N} \mu_i |u_i|^p + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}} > 0 \quad \forall i = 1, \dots, \ell,$$

there exists a unique $\mathbf{t}_\mathbf{u} \in (0, \infty)^\ell$, such that $\mathbf{t}_\mathbf{u}\mathbf{u} \in \mathcal{N}_\varepsilon^\ell(\Theta)$. Furthermore,

$$\mathcal{J}_\varepsilon^\ell(\mathbf{t}_\mathbf{u}\mathbf{u}) = \max_{\mathbf{t} \in (0, \infty)^\ell} \mathcal{J}_\varepsilon^\ell(\mathbf{t}\mathbf{u}).$$

Proof. (i) Since ϕ_i satisfies (A_1) , for each $i = 1, \dots, \ell$ one can choose $u_i \in C_c^\infty(\Theta) \cap H_0^1(\Theta)^{\phi_i}$, $u_i \neq 0$, such that $\text{supp}(u_i) \cap \text{supp}(u_j) = \emptyset$ for all $j \neq i$, and then take $t_i > 0$ such that $v_i := t_i u_i$ satisfies $\|v_i\|_\varepsilon^2 = \varepsilon^{-N} \int_{\mathbb{R}^N} \mu_i |v_i|^p$. Since $\text{supp}(v_i) \cap \text{supp}(v_j) = \emptyset$ for all $j \neq i$, setting $\mathbf{v} := (v_1, \dots, v_\ell)$ we get that $\partial_i \mathcal{J}_\varepsilon^\ell(\mathbf{v}) v_i = 0$ and $v_i \neq 0$ for all $i = 1, \dots, \ell$. Therefore $\mathbf{v} \in \mathcal{N}_\varepsilon^\ell(\Theta)$.

Let $\mathbf{u} = (u_1, \dots, u_\ell) \in \mathcal{N}_\varepsilon^\ell(\Theta)$ and set $\tilde{u}_i(z) := u_i(\varepsilon z)$. Then, since $\lambda_{ij} < 0$ for every pair $i \neq j$, from Remark 2.2 and Sobolev's embedding theorem we derive

$$\|\tilde{u}_i\|^2 \leq \int_{\mathbb{R}^N} \mu_i |\tilde{u}_i|^p \leq C \mu_i \|\tilde{u}_i\|^p,$$

with C a positive constant, depending only on N and p . This yields the estimate in statement (i) and completes its proof.

(ii) Fix $\mathbf{u} = (u_1, \dots, u_\ell) \in \mathcal{H}^\ell(\Theta)$ and define $J_{\mathbf{u}}^\ell : (0, \infty)^\ell \rightarrow \mathbb{R}$ by

$$J_{\mathbf{u}}^\ell(\mathbf{t}) := \mathcal{J}_\varepsilon^\ell(\mathbf{t}\mathbf{u}) = \sum_{i=1}^{\ell} a_{u,i} t_i^2 - \sum_{i=1}^{\ell} b_{u,i} t_i^p + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} d_{u,ij} t_j^{\alpha_{ij}} t_i^{\beta_{ij}},$$

where $\mathbf{t} = (t_1, \dots, t_\ell) \in (0, \infty)^\ell$,

$$a_{u,i} := \frac{1}{2} \|u_i\|_\varepsilon^2, \quad b_{u,i} := \frac{1}{p\varepsilon^N} \int_{\mathbb{R}^N} \mu_i |u_i|^p, \quad d_{u,ij} := \frac{1}{2\varepsilon^N} \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \int_{\mathbb{R}^N} |\lambda_{ij}| \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}}.$$

If $u_i \neq 0$ for all $i = 1, \dots, \ell$, then, as

$$t_i \partial_i J_{\mathbf{u}}^\ell(\mathbf{t}) = \partial_i \mathcal{J}_\varepsilon^\ell(\mathbf{t}\mathbf{u})[t_i u_i], \quad i = 1, \dots, \ell,$$

we have that $\mathbf{t}\mathbf{u} \in \mathcal{N}_\varepsilon^\ell(\Theta)$ iff \mathbf{t} is a critical point of $J_{\mathbf{u}}^\ell$. Note that (2.4) implies that $u_i \neq 0$ for all $i = 1, \dots, \ell$. Hence, (ii) follows from [8, Lemmas 2.1 and 2.2]. \square

Theorem 2.4. *If there exists $\mathbf{u} \in \mathcal{N}_\varepsilon^\ell(\Theta)$ such that $\mathcal{J}_\varepsilon^\ell(\mathbf{u}) = c_\varepsilon^\ell(\Theta)$, then $\mathbf{u} = (u_1, \dots, u_\ell)$ is a solution to system $(\mathcal{S}_{\varepsilon, \Theta}^\phi)$.*

Proof. The proof is practically the same as that of [8, Theorem 3.4(a)] with the obvious modifications. \square

Definition 2.5. *A solution to the system $(\mathcal{S}_{\varepsilon, \Theta}^\phi)$ such that $\mathcal{J}_\varepsilon^\ell(\mathbf{u}) = c_\varepsilon^\ell(\Theta)$ will be called a least energy solution to $(\mathcal{S}_{\varepsilon, \Theta}^\phi)$.*

Remark 2.6. If $\phi_i \equiv 1$ for every i in some subset I of $\{1, \dots, \ell\}$ and \mathbf{u} is a least energy solution to $(\mathcal{S}_{\varepsilon, \Theta}^\phi)$, then, setting $v_i := |u_i|$ if $i \in I$ and $v_i := u_i$ if $i \notin I$, we have that $v_i \in H_0^1(\Theta)^{\phi_i}$, $\mathbf{v} = (v_1, \dots, v_\ell) \in \mathcal{N}_\varepsilon^\ell(\Theta)$ and $\mathcal{J}_\varepsilon^\ell(\mathbf{v}) = \mathcal{J}_\varepsilon^\ell(\mathbf{u})$. Hence, \mathbf{v} is a least energy solution to $(\mathcal{S}_{\varepsilon, \Theta}^\phi)$ whose i -th component is positive for every $i \in I$.

Theorem 2.7 (Existence of minimizers). *If Θ is a bounded G -invariant domain in \mathbb{R}^N , then, for each $\varepsilon > 0$, the system $(\mathcal{S}_{\varepsilon, \Theta}^\phi)$ has a least energy solution.*

Proof. Fix $\varepsilon > 0$ and let $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k}) \in \mathcal{N}_\varepsilon^\ell(\Theta)$ satisfy $\mathcal{J}_\varepsilon^\ell(\mathbf{u}_k) \rightarrow c_\varepsilon^\ell(\Theta)$. From (2.3) we see that (u_{ik}) is bounded in $H_0^1(\Theta)$ for every $i = 1, \dots, \ell$. So, after passing to a subsequence, $u_{ik} \rightharpoonup u_i$ weakly in $H_0^1(\Theta)^{\phi_i}$ and $u_{ik} \rightarrow u_i$ strongly in $L^p(\Theta)$. From (2.2) and Lemma 2.3 we get that

$$0 < c_0 \leq \|u_{ik}\|_\varepsilon^2 = \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \mu_i |u_{ik}|^p + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \frac{1}{\varepsilon^N} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_{jk}|^{\alpha_{ij}} |u_{ik}|^{\beta_{ij}}$$

and, passing to the limit, we obtain

$$0 < c_0 \leq \int_{\mathbb{R}^N} \mu_i |u_i|^p + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}} \quad \text{for all } i = 1, \dots, \ell.$$

Set $\mathbf{u} := (u_1, \dots, u_\ell)$. By Lemma 2.3 there exists $\mathbf{t} = (t_1, \dots, t_\ell) \in (0, \infty)^\ell$ such that $\mathbf{t}\mathbf{u} \in \mathcal{N}_\varepsilon^\ell(\Theta)$ and

$$c_\varepsilon^\ell(\Theta) \leq \mathcal{J}_\varepsilon^\ell(\mathbf{t}\mathbf{u}) \leq \liminf_{k \rightarrow \infty} \mathcal{J}_\varepsilon^\ell(\mathbf{t}\mathbf{u}_k) \leq \lim_{k \rightarrow \infty} \mathcal{J}_\varepsilon^\ell(\mathbf{u}_k) = c_\varepsilon^\ell(\Theta),$$

because $\mathbf{u}_k \in \mathcal{N}_\varepsilon^\ell(\Theta)$ for all $k \in \mathbb{N}$. Thus, $\lim_{k \rightarrow \infty} \mathcal{J}_\varepsilon^\ell(\mathbf{t}\mathbf{u}_k) = \mathcal{J}_\varepsilon^\ell(\mathbf{t}\mathbf{u})$. As $u_{ik} \rightarrow u_i$ strongly in $L^p(\Theta)$, this implies that $\lim_{k \rightarrow \infty} \|\mathbf{t}\mathbf{u}_k\|_{\ell, \varepsilon}^2 = \|\mathbf{t}\mathbf{u}\|_{\ell, \varepsilon}^2$. Since $\mathbf{u}_k \rightharpoonup \mathbf{u}$ weakly in $\mathcal{H}^\ell(\Theta)$, we derive that $\mathbf{u}_k \rightarrow \mathbf{u}$ strongly in $\mathcal{H}^\ell(\Theta)$. As a consequence, $\mathbf{u} \in \mathcal{N}_\varepsilon^\ell(\Theta)$ and $\mathcal{J}_\varepsilon^\ell(\mathbf{u}) = c_\varepsilon^\ell(\Theta)$. \square

3 The limit system

We fix a closed subgroup G of $O(N)$ and, for each $i = 1, \dots, \ell$, a continuous homomorphism $\phi_i : G \rightarrow \mathbb{Z}_2$. We assume throughout this section that the ϕ_i satisfy (A_1) and that G has the following property:

(A_2) For every $x \in \mathbb{R}^N$, the G -orbit of x is either infinite, or $Gx = \{x\}$.

Our aim is to study the behavior of minimizing sequences for the functional related to the system

$$(\mathcal{S}_{\infty, \ell}^\phi) \quad \begin{cases} -\Delta u_i + u_i = \mu_i |u_i|^{p-2} u_i + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \lambda_{ij} \beta_{ij} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}-2} u_i, \\ u_i \in H^1(\mathbb{R}^N)^{\phi_i}, \quad u_i \neq 0, \quad 1 < i \leq \ell. \end{cases}$$

We write $\|\cdot\|$ for the standard norm in $H^1(\mathbb{R}^N)$. Set $\mathcal{H}^\ell := H^1(\mathbb{R}^N)^{\phi_1} \times \dots \times H^1(\mathbb{R}^N)^{\phi_\ell}$, $\mathcal{J}_\infty^\ell := \mathcal{J}_1^\ell : \mathcal{H}^\ell \rightarrow \mathbb{R}$ and $\mathcal{N}_\infty^\ell := \mathcal{N}_1^\ell(\mathbb{R}^N)$ as in the previous section, and let

$$c_\infty^\ell := \inf_{\mathbf{u} \in \mathcal{N}_\infty^\ell} \mathcal{J}_\infty^\ell(\mathbf{u}).$$

For each $i = 1, \dots, \ell$, consider the problem

$$(\mathcal{P}_i^{\phi_i}) \quad \begin{cases} -\Delta u + u = \mu_i |u|^{p-2} u, \\ u \in H^1(\mathbb{R}^N)^{\phi_i}, \quad u \neq 0. \end{cases}$$

Its solutions are the critical points of the functional $J_i : H^1(\mathbb{R}^N)^{\phi_i} \rightarrow \mathbb{R}$ given by

$$J_i(u) := \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla u|^2 + u^2) - \frac{1}{p} \int_{\mathbb{R}^N} |u|^p,$$

belonging to the Nehari manifold $\mathcal{N}_i := \{u \in H^1(\mathbb{R}^N)^{\phi_i} : u \neq 0, J_i'(u)u = 0\}$. We set

$$c_i := \inf_{u \in \mathcal{N}_i} J_i(u).$$

A solution \mathbf{u} to $(\mathcal{S}_{\infty, \ell}^\phi)$ satisfying $\mathcal{J}_\infty^\ell(\mathbf{u}) = c_\infty^\ell$ will be called a *least energy solution to $(\mathcal{S}_{\infty, \ell}^\phi)$* . Similarly, a solution u to $(\mathcal{P}_i^{\phi_i})$ satisfying $J_i(u) = c_i$ will be called a *least energy solution to $(\mathcal{P}_i^{\phi_i})$* .

The G -fixed-point space

$$\text{Fix}(G) := \{x \in \mathbb{R}^N : gx = x \text{ for all } g \in G\}$$

plays an important role on the existence or nonexistence of a least energy solution to the system $(\mathcal{S}_{\infty, \ell}^\phi)$. We start with the following result.

Proposition 3.1. *The following statements hold true:*

$$(i) \quad c_\infty^\ell \geq \sum_{i=1}^{\ell} c_i.$$

$$(ii) \quad \text{If } \text{Fix}(G) \text{ has positive dimension, then } c_\infty^\ell = \sum_{i=1}^{\ell} c_i.$$

$$(iii) \quad \text{If } \ell \geq 2 \text{ and } c_\infty^\ell = \sum_{i=1}^{\ell} c_i, \text{ then } c_\infty^\ell \text{ is not attained.}$$

Proof. (i) : Let $\mathbf{u} = (u_1, \dots, u_\ell) \in \mathcal{N}_\infty^\ell$. As $\lambda_{ij} < 0$, from (2.2) and Lemma 2.3 (i) we get that

$$0 < \|u_i\|^2 \leq \int_{\mathbb{R}^N} \mu_i |u_i|^p \quad \text{for all } i = 1, \dots, \ell.$$

Hence, there exist $t_1, \dots, t_\ell \in (0, \infty)$ such that $t_i u_i \in \mathcal{N}_i$. Setting $\mathbf{t} := (t_1, \dots, t_\ell)$ and applying Lemma 2.3 (ii) we obtain

$$\sum_{i=1}^{\ell} c_i \leq \sum_{i=1}^{\ell} J_i(t_i u_i) \leq \mathcal{J}_\infty^\ell(\mathbf{t}\mathbf{u}) \leq \mathcal{J}_\infty^\ell(\mathbf{u}),$$

as claimed.

(ii) : For each $i = 1, \dots, \ell$, let $v_i \in \mathcal{N}_i$. As $\text{Fix}(G)$ has positive dimension we may choose $\xi_{ik} \in \text{Fix}(G)$ such that $|\xi_{ik} - \xi_{jk}| \rightarrow \infty$ as $k \rightarrow \infty$ if $i \neq j$. Define $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k})$ by $u_{ik}(x) := v_i(x - \xi_{ik})$. Note that, as $\xi_{ik} \in \text{Fix}(G)$, we have that $u_{ik} \in H^1(\mathbb{R}^N)^{\phi_i}$. Furthermore, $u_{ik} \in \mathcal{N}_i$, $J_i(u_{ik}) = J_i(v_i)$ and

$$\int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_{jk}|^{\alpha_{ij}} |u_{ik}|^{\beta_{ij}} = \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |v_j(\cdot + \xi_{ik} - \xi_{jk})|^{\alpha_{ij}} |v_i|^{\beta_{ij}} = o(1).$$

Therefore,

$$0 < \frac{2p}{p-2} c_i \leq \|u_{ik}\|^2 = \int_{\mathbb{R}^N} \mu_i |u_{ik}|^p + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_{jk}|^{\alpha_{ij}} |u_{ik}|^{\beta_{ij}} + o(1).$$

Applying Lemma 2.3 (ii) we see that, for k large enough, there exists $\mathbf{t}_k := (t_{1k}, \dots, t_{\ell k}) \in (0, \infty)^\ell$ such that $\mathbf{t}_k \mathbf{u}_k \in \mathcal{N}_\infty^\ell$, and $t_{ik} \rightarrow 1$ for all $i = 1, \dots, \ell$. Hence,

$$c_\infty^\ell \leq \mathcal{J}_\infty^\ell(\mathbf{t}_k \mathbf{u}_k) = \sum_{i=1}^{\ell} J_i(t_{ik} u_{ik}) + o(1) = \sum_{i=1}^{\ell} J_i(v_i) + o(1).$$

This shows that $\sum_{i=1}^{\ell} c_i \geq c_\infty^\ell$ and from (i) we obtain $\sum_{i=1}^{\ell} c_i = c_\infty^\ell$.

(iii) : Let $c_\infty^\ell = \sum_{i=1}^{\ell} c_i$. To show that this value is not attained, we argue by contradiction. Assume that $\mathbf{u} = (u_1, \dots, u_\ell) \in \mathcal{N}_\infty^\ell$ and $\mathcal{J}_\infty^\ell(\mathbf{u}) = c_\infty^\ell$. There are two possibilities. If for every i there exists $j \neq i$ such that $\int_{\mathbb{R}^N} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}} \neq 0$, then $\|u_i\|^2 < \int_{\mathbb{R}^N} \mu_i |u_i|^p$ for all $i = 1, \dots, \ell$ and, hence, there exists $t_i \in (0, 1)$ such that $t_i u_i \in \mathcal{N}_i$ and

$$c_i \leq J_i(t_i u_i) = \frac{p-2}{2p} \|t_i u_i\|^2 < \frac{p-2}{2p} \|u_i\|^2.$$

It follows that

$$c_\infty^\ell = \mathcal{J}_\infty^\ell(\mathbf{u}) = \frac{p-2}{2p} \sum_{i=1}^{\ell} \|u_i\|^2 > \sum_{i=1}^{\ell} c_i,$$

contradicting our assumption. On the other hand, if there exists i such that $\int_{\mathbb{R}^N} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}} = 0$ for all $j \neq i$, then

$$\|u_i\|^2 = \int_{\mathbb{R}^N} \mu_i |u_i|^p \quad \text{and} \quad J_i(u_i) = c_i.$$

Hence, u_i is a nontrivial solution to the problem

$$-\Delta w + w = \mu_i |w|^{p-2} w, \quad w \in H^1(\mathbb{R}^N).$$

But $\int_{\mathbb{R}^N} |u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}} = 0$ also implies that $|u_j|^{\alpha_{ij}} |u_i|^{\beta_{ij}} = 0$ a.e. in \mathbb{R}^N . As $u_j \neq 0$ for all j , we have that $u_i = 0$ in some subset of positive measure of \mathbb{R}^N . This contradicts the unique continuation principle. \square

The following lemma will play an important role in the proof of Theorem 3.3.

Lemma 3.2. *For any given sequence (y_k) in \mathbb{R}^N , there exist a sequence (ζ_k) in \mathbb{R}^N and a constant $C > 0$ such that, up to a subsequence,*

$$\text{dist}(Gy_k, \zeta_k) \leq C \quad \text{for all } k \in \mathbb{N},$$

and

- either $\zeta_k \in \text{Fix}(G)$ for all $k \in \mathbb{N}$,
- or there exist $g_n \in G$, $n \in \mathbb{N}$, such that $\lim_{k \rightarrow \infty} |g_{n_1} \zeta_k - g_{n_2} \zeta_k| = \infty$ for any pair $n_1, n_2 \in \mathbb{N}$ with $n_1 \neq n_2$.

Proof. This is the statement of [6, Lemma 3.2] for a group G satisfying (A_2) . \square

We are ready to prove the main result of this section.

Theorem 3.3. *Let $\ell \geq 2$ and $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k}) \in \mathcal{N}_\infty^\ell$ be such that $\mathcal{J}_\infty^\ell(\mathbf{u}_k) \rightarrow c_\infty^\ell$.*

- (I) *If $\text{Fix}(G) = \{0\}$, then there exists a least energy solution $\mathbf{w} = (w_1, \dots, w_\ell)$ to the system $(\mathcal{P}_{\infty, \ell}^\phi)$ such that, after passing to a subsequence,*

$$\lim_{k \rightarrow \infty} \|u_{ik} - w_i\| = 0 \quad \text{for all } i = 1, \dots, \ell,$$

and $c_\infty^\ell > \sum_{i=1}^{\ell} c_i$. Moreover, if $u_{ik} \geq 0$ for all $k \in \mathbb{N}$, then $w_i \geq 0$.

- (II) *If $\text{Fix}(G)$ has positive dimension, then, for each $i = 1, \dots, \ell$, there exist a sequence (ξ_{ik}) in $\text{Fix}(G)$ and a least energy solution v_i to the problem $(\mathcal{P}_i^{\phi_i})$ such that, after passing to a subsequence,*

$$\lim_{k \rightarrow \infty} |\xi_{ik} - \xi_{jk}| = \infty \text{ if } i \neq j, \quad \lim_{k \rightarrow \infty} \|u_{ik} - v_i(\cdot - \xi_{ik})\| = 0 \text{ for } i, j = 1, \dots, \ell,$$

and $c_\infty^\ell = \sum_{i=1}^{\ell} c_i$. Moreover, if $u_{ik} \geq 0$ for all $k \in \mathbb{N}$, then $v_i \geq 0$.

Proof. We split the proof into three steps.

STEP 1. *The choice of the concentration points.*

As $\mathbf{u}_k \in \mathcal{N}_\infty^\ell$ and $\lambda_{ij} < 0$, from Lemma 2.3 we get a constant $c_0 > 0$ such that

$$(3.1) \quad \int_{\mathbb{R}^N} \mu_i |u_{ik}|^p > c_0 \quad \text{for all } k \in \mathbb{N} \text{ and } i = 1, \dots, \ell.$$

Invoking [8, Theorem 3.3] and Ekeland's variational principle [29, Theorem 8.5] we may assume without loss of generality that $(\mathcal{J}_\infty^\ell)'(\mathbf{u}_k) \rightarrow 0$ in the dual space $(\mathcal{H}^\ell)'$. From (3.1) and Lions' lemma [29, Lemma 1.21] we see that there exists $\theta > 0$ such that

$$\limsup_{k \rightarrow \infty} \sup_{y \in \mathbb{R}^N} \int_{B_1(y)} \mu_i |u_{ik}|^p > 2\theta > 0 \quad \text{for all } i = 1, \dots, \ell.$$

So, passing to a subsequence, we may choose $y_{ik} \in \mathbb{R}^N$ satisfying

$$(3.2) \quad \int_{B_1(y_{ik})} \mu_i |u_{ik}|^p > \theta \quad \text{for all } k \in \mathbb{N} \text{ and } i = 1, \dots, \ell.$$

Applying Lemma 3.2 to (y_{ik}) we obtain a sequence (ζ_{ik}) and a constant C such that $\text{dist}(Gy_{ik}, \zeta_{ik}) \leq C$ for all k and i . Since $|u_{ik}|$ is G -invariant, taking $g_{ik} \in G$ such that $|g_{ik}y_{ik} - \zeta_{ik}| = \text{dist}(Gy_{ik}, \zeta_{ik})$ we get

$$(3.3) \quad \int_{B_{C+1}(\zeta_{ik})} \mu_i |u_{ik}|^p \geq \int_{B_1(g_{ik}y_{ik})} \mu_i |u_{ik}|^p \geq \int_{B_1(y_{ik})} \mu_i |u_{ik}|^p > \theta > 0.$$

We claim that $\zeta_{ik} \in \text{Fix}(G)$. Otherwise, by the alternative in Lemma 3.2, for every $M \in \mathbb{N}$ we can choose $g_{i1}, \dots, g_{iM} \in G$ such that

$$|g_{in}\zeta_{ik} - g_{in'}\zeta_{ik}| \geq 2(C+1) \quad \text{for all pairs } n \neq n', \quad n, n' = 1, \dots, M,$$

and for sufficiently large $k \in \mathbb{N}$. As a consequence, $B_{C+1}(g_{in}\zeta_{ik}) \cap B_{C+1}(g_{in'}\zeta_{ik}) = \emptyset$ and, so,

$$\int_{\mathbb{R}^N} \mu_i |u_{ik}|^p \geq \sum_{n=1}^M \int_{B_{C+1}(g_{in}\zeta_{ik})} \mu_i |u_{ik}|^p = M \int_{B_{C+1}(\zeta_{ik})} \mu_i |u_{ik}|^p > M\theta,$$

which is a contradiction, because M is arbitrary and, as (\mathbf{u}_k) is bounded in \mathcal{H}^ℓ , the sequence (u_{ik}) is bounded in $L^p(\mathbb{R}^N)$. This shows that $\zeta_{ik} \in \text{Fix}(G)$.

After passing to a subsequence and reordering the components, there are integers

$$0 =: \ell_0 < \ell_1 < \dots < \ell_n := \ell$$

such that, setting $I_r := (\ell_r, \ell_{r+1}]$ for $r = 0, \dots, n-1$, we have

$$\begin{aligned} (|\zeta_{ik} - \zeta_{jk}|) \text{ is bounded} & \quad \text{if } i, j \in I_r \text{ for some } r, \\ |\zeta_{ik} - \zeta_{jk}| \rightarrow \infty & \quad \text{if } i \in I_r \text{ and } j \in I_{r'} \text{ with } r \neq r'. \end{aligned}$$

Now we replace ζ_{ik} by $\xi_{ik} := \zeta_{\ell_{r+1}k}$ for every $i \in I_r$, and we fix $R > 0$ large enough so that $B_{C+1}(\zeta_{ik}) \subset B_R(\xi_{ik})$ for all $k \in \mathbb{N}$ and $i = 1, \dots, \ell$. Then these new points satisfy

$$\begin{aligned} \xi_{ik} & \in \text{Fix}(G), \\ \xi_{ik} = \xi_{jk} & \text{ for all } k & \quad \text{if } i, j \in I_r \text{ for some } r, \\ |\xi_{ik} - \xi_{jk}| & \rightarrow \infty & \quad \text{if } i \in I_r \text{ and } j \in I_{r'} \text{ with } r \neq r', \end{aligned}$$

and from (3.3) we obtain

$$(3.4) \quad \int_{B_R(\xi_{ik})} \mu_i |u_{ik}|^p > \theta > 0 \quad \text{for all } k \in \mathbb{N} \text{ and } i = 1, \dots, \ell.$$

Note that, if $\text{Fix}(G) = \{0\}$, then $\xi_{ik} = 0$ for all k and i , and there is only one interval $I_0 := (0, \ell]$.

STEP 2. *A solution to a subsystem.*

Define

$$w_{ik}(x) := u_{ik}(x + \xi_{ik}).$$

Observe that $w_{ik} \in H^1(\mathbb{R}^N)^{\phi_i}$, because $\xi_{ik} \in \text{Fix}(G)$. Since the sequence (w_{ik}) is bounded in $H^1(\mathbb{R}^N)$, passing to a subsequence, we have that $w_{ik} \rightharpoonup w_i$ weakly in $H^1(\mathbb{R}^N)$, $w_{ik} \rightarrow w_i$ in $L^p_{\text{loc}}(\mathbb{R}^N)$ and $w_{ik} \rightarrow w_i$ a.e. in \mathbb{R}^N . Then, $w_i \geq 0$ if $u_{ik} \geq 0$ for all $k \in \mathbb{N}$. As, by (3.4),

$$\int_{B_R(0)} \mu_i |w_{ik}|^p = \int_{B_R(\xi_{ik})} \mu_i |u_{ik}|^p > \theta > 0 \quad \text{for all } k \in \mathbb{N} \text{ and } i = 1, \dots, \ell,$$

we deduce that $w_i \neq 0$ for all $i = 1, \dots, \ell$. Furthermore,

$$(3.5) \quad \frac{p-2}{2p} \sum_{i=1}^{\ell} \|w_i\|^2 \leq \lim_{k \rightarrow \infty} \frac{p-2}{2p} \sum_{i=1}^{\ell} \|w_{ik}\|^2 = \lim_{k \rightarrow \infty} \frac{p-2}{2p} \sum_{i=1}^{\ell} \|u_{ik}\|^2 = \lim_{k \rightarrow \infty} \mathcal{J}_{\infty}^{\ell}(\mathbf{u}_k) = c_{\infty}^{\ell}.$$

Fix $r \in \{0, \dots, n-1\}$, take ξ_{ik} with $i \in I_r$ and define $\mathbf{u}_k^r = (u_{1k}^r, \dots, u_{\ell k}^r)$ by

$$u_{jk}^r(x) := u_{jk}(x + \xi_{ik}) = \begin{cases} w_{jk}(x) & \text{if } j \in I_r, \\ w_{jk}(x + \xi_{ik} - \xi_{jk}) & \text{if } j \notin I_r. \end{cases}$$

After passing to a subsequence, we have that $u_{jk}^r \rightharpoonup u_j^r$ weakly in $H^1(\mathbb{R}^N)$. Note that $u_j^r = w_j \neq 0$ if $j \in I_r$. Set $\mathbf{u}^r := (u_1^r, \dots, u_{\ell}^r)$. Let $\varphi \in C_c^{\infty}(\mathbb{R}^N)$ and define $\widehat{\varphi}_k(x) := \varphi(x - \xi_{ik})$. Then we have that $\partial_j \mathcal{J}_{\infty}^{\ell}(\mathbf{u}_k) \widehat{\varphi}_k = \partial_j \mathcal{J}_{\infty}^{\ell}(\mathbf{u}_k^r) \varphi$ and, as $(\mathcal{J}_{\infty}^{\ell})'(\mathbf{u}_k) \rightarrow 0$, we deduce that

$$0 = \lim_{k \rightarrow \infty} \partial_j \mathcal{J}_{\infty}^{\ell}(\mathbf{u}_k) \widehat{\varphi}_k = \lim_{k \rightarrow \infty} \partial_j \mathcal{J}_{\infty}^{\ell}(\mathbf{u}_k^r) \varphi = \partial_j \mathcal{J}_{\infty}^{\ell}(\mathbf{u}^r) \varphi \quad \text{for all } j = 1, \dots, \ell.$$

This shows that \mathbf{u}^r is a critical point of $\mathcal{J}_{\infty}^{\ell} : \mathcal{H}^{\ell} \rightarrow \mathbb{R}$. Hence, its nontrivial components give a fully nontrivial solution to the subsystem

$$(3.6) \quad \begin{cases} -\Delta v_m + v_m = \mu_m |v_m|^{p-2} v_m + \sum_{\substack{j \in \widehat{I}_r \\ j \neq m}} \lambda_{mj} \beta_{mj} |v_j|^{\alpha_{mj}} |v_m|^{\beta_{mj}-2} v_m, \\ v_m \in H^1(\mathbb{R}^N)^{\phi_m}, \quad v_m \neq 0, \quad m \in \widehat{I}_r, \end{cases}$$

where $\widehat{I}_r := \{m : 1 \leq m \leq \ell, u_m^r \neq 0\}$. Note that $\widehat{I}_r \supset I_r$.

STEP 3. *The conclusion.*

Now we distinguish two cases.

(I) : Let $\text{Fix}(G) = \{0\}$. Then $\xi_{ik} = 0$ and $w_{ik} = u_{ik}$ for all k , $\widehat{I}_0 = \{1, \dots, \ell\}$ and $\mathbf{w} := (w_1, \dots, w_{\ell})$ is a fully nontrivial solution to the system (3.6) with $r = 0$. From (3.5) we derive

$$c_{\infty}^{\ell} \leq \mathcal{J}_{\infty}^{\ell}(\mathbf{w}) = \frac{p-2}{2p} \sum_{i=1}^{\ell} \|w_i\|^2 \leq \lim_{k \rightarrow \infty} \frac{p-2}{2p} \sum_{i=1}^{\ell} \|u_{ik}\|^2 = c_{\infty}^{\ell}.$$

As $u_{ik} \rightharpoonup w_i$ weakly in $H^1(\mathbb{R}^N)$, this implies that $u_{ik} \rightarrow w_i$ strongly in $H^1(\mathbb{R}^N)$ and that $\mathcal{J}_{\infty}^{\ell}(\mathbf{w}) = c_{\infty}^{\ell}$.

Then, by Proposition 3.1(iii), we have that $c_{\infty}^{\ell} > \sum_{i=1}^{\ell} c_i$, as claimed.

(II) : Let $\text{Fix}(G)$ have positive dimension. Since for each $r \in \{0, \dots, n-1\}$, the nontrivial components of \mathbf{u}^r satisfy (3.6), multiplying the i -th equation by w_i we get

$$(3.7) \quad \|w_i\|^2 = \int_{\mathbb{R}^N} \mu_i |w_i|^p + \sum_{\substack{j \in \widehat{I}_r \\ j \neq i}} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_j^r|^{\alpha_{ij}} |w_i|^{\beta_{ij}} \leq \int_{\mathbb{R}^N} \mu_i |w_i|^p.$$

Hence, there exists $t_i \in (0, 1]$ such that $t_i w_i \in \mathcal{N}_i$ and, so,

$$c_i \leq \frac{p-2}{2p} \|t_i w_i\|^2 \leq \frac{p-2}{2p} \|w_i\|^2 \quad \text{for every } i = 1, \dots, \ell.$$

From (3.5) and Proposition 3.1 we derive

$$(3.8) \quad \sum_{i=1}^{\ell} c_i \leq \frac{p-2}{2p} \sum_{i=1}^{\ell} \|w_i\|^2 \leq \lim_{k \rightarrow \infty} \frac{p-2}{2p} \sum_{i=1}^{\ell} \|w_{ik}\|^2 = c_{\infty}^{\ell} = \sum_{i=1}^{\ell} c_i.$$

It follows that $\frac{p-2}{2p} \|w_i\|^2 = c_i$ and $t_i = 1$, so w_i is a nontrivial least energy solution to the problem $(\mathcal{P}_i^{\phi_i})$ for each $i = 1, \dots, \ell$. Furthermore, (3.7) yields

$$\int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |u_j^r|^{\alpha_{ij}} |w_i|^{\beta_{ij}} = 0 \quad \text{for every } j \neq i.$$

Therefore, $u_j^r = 0$ for every $j \neq i$ and, so, $\widehat{I}_r = I_r = \{i\}$. This implies that $|\xi_{ik} - \xi_{jk}| \rightarrow \infty$ for all $j \neq i$. Finally, as $w_{ik} \rightharpoonup w_i$ weakly in $H^1(\mathbb{R}^N)$, it follows from (3.8) that $w_{ik} \rightarrow w_i$ strongly in $H^1(\mathbb{R}^N)$. This implies that $\lim_{k \rightarrow \infty} \|u_{ik} - w_i(\cdot - \xi_{ik})\| = 0$, and the proof is complete. \square

From Theorem 3.3 and Proposition 3.1(ii)-(iii) we immediately obtain the following result.

Theorem 3.4. *Let $\ell \geq 2$. The system $(\mathcal{S}_{\infty, \ell}^{\phi})$ has a least energy solution if and only if $\text{Fix}(G) = \{0\}$.*

Theorem 1.1 follows from the previous one.

Proof of Theorem 1.1. The group $G := \Gamma \times O(N-4)$ introduced in Example 2.1(iii) satisfies (A_2) if either $N = 4$ or $N \geq 6$, and $\text{Fix}(G) = \{0\}$. For $i = 1, \dots, m$ we set $\phi_i \equiv 1$, and for $i = m+1, \dots, \ell$ we take ϕ_i as in Example 2.1(iii). Applying Theorem 3.4 we obtain a least energy solution $\mathbf{w} = (w_1, \dots, w_{\ell})$ to the system $(\mathcal{S}_{\infty, \ell})$ satisfying (1.1). By Remark 2.6, we may take w_1, \dots, w_m to be positive. The symmetries ensure that the last $\ell - m$ components are nonradial and change sign. \square

To further emphasize the role of the fixed points, we prove the following result.

Theorem 3.5. *If $\text{Fix}(G) = \{0\}$, then the system $(\mathcal{S}_{\infty, \ell}^{\phi})$ has infinitely many solutions.*

The key ingredient is the following fact.

Lemma 3.6. *If G satisfies (A_2) and $\text{Fix}(G) = \{0\}$, then the embedding $H^1(\mathbb{R}^N)^{\phi} \hookrightarrow L^p(\mathbb{R}^N)$ is compact for any homomorphism $\phi : G \rightarrow \mathbb{Z}_2$ and $p \in (2, 2^*)$.*

Proof. Let $K := \ker \phi$. Since the K -orbit of every $x \in \mathbb{R}^N \setminus \{0\}$ is infinite, the embedding $H^1(\mathbb{R}^N)^K \hookrightarrow L^p(\mathbb{R}^N)$ is compact, where $H^1(\mathbb{R}^N)^K$ denotes the space of K -invariant functions in $H^1(\mathbb{R}^N)$; see, e.g., [8, Lemma 4.3]. Since $H^1(\mathbb{R}^N)^{\phi} \subset H^1(\mathbb{R}^N)^K$, our claim follows. \square

Proof of Theorem 3.5. We note that the results in [8, Section 3] hold true if we take $\mathcal{H} := \mathcal{H}^\ell$. A standard argument using Lemma 3.6 shows that $\mathcal{J}_\infty^\ell : \mathcal{H}^\ell \rightarrow \mathbb{R}$ satisfies the Palais-Smale condition at every level $c \in \mathbb{R}$. As in [8, Section 3] we set

$$\mathcal{U} := \{\mathbf{u} \in \mathcal{H}^\ell : \|u_i\| = 1 \text{ for } i = 1, \dots, \ell, \text{ and there exists } \mathbf{t} \in (0, \infty)^\ell \text{ such that } \mathbf{t}\mathbf{u} \in \mathcal{N}_\infty^\ell\},$$

and we define $\Psi : \mathcal{U} \rightarrow \mathbb{R}$ by $\Psi(u) := \mathcal{J}_\infty^\ell(\mathbf{t}\mathbf{u})$ for $\mathbf{t} \in (0, \infty)^\ell$ such that $\mathbf{t}\mathbf{u} \in \mathcal{N}_\infty^\ell$. Such \mathbf{t} is unique.

It follows from [8, Theorem 3.3] that $\Psi \in C^1(\mathcal{U}, \mathbb{R})$ and it satisfies the Palais-Smale condition at every level $c \in \mathbb{R}$. Exactly the same argument used to prove [8, Lemma 4.5] shows that $\text{genus}(\mathcal{U}) = \infty$ (one simply replaces $H_0^1(\Omega)^G$ with $H^1(\mathbb{R}^N)^{\phi_i}$). Now [8, Theorem 3.4] asserts that $(\mathcal{S}_{\infty, \ell}^\phi)$ has an unbounded sequence of solutions. \square

We observe that, while the last $\ell - m$ components of the solutions given by the Theorem 3.5 change sign by construction, this theorem does not give any information about the sign of the first m components.

4 The asymptotic behavior of minimizers

As before, we fix a closed subgroup G of $O(N)$ satisfying (A_2) and, for each $i = 1, \dots, \ell$, a continuous homomorphism $\phi_i : G \rightarrow \mathbb{Z}_2$ satisfying (A_1) . Furthermore, we assume throughout that Ω is a G -invariant bounded domain in \mathbb{R}^N and that $0 \in \Omega$.

Our aim is to prove the following result.

Theorem 4.1. *For any given sequence (ε_k) of positive numbers converging to zero, there exists a least energy solution $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k})$ to the system $(\mathcal{S}_{\varepsilon_k, \Omega}^\phi)$ with the following limit profile:*

(I) *If $\text{Fix}(G) = \{0\}$, then there exists a least energy solution $\mathbf{w} = (w_1, \dots, w_\ell)$ to the system $(\mathcal{S}_{\infty, \ell}^\phi)$ such that, after passing to a subsequence,*

$$\lim_{k \rightarrow \infty} \|u_{ik} - w_i(\varepsilon_k^{-1} \cdot)\|_{\varepsilon_k} = 0 \quad \text{for all } i = 1, \dots, \ell,$$

and $\lim_{\varepsilon \rightarrow 0} c_\varepsilon^\ell(\Omega) = c_\infty^\ell$. Moreover, $w_i \geq 0$ if $u_{ik} \geq 0$ for all $k \in \mathbb{N}$.

(II) *If $\text{Fix}(G)$ has positive dimension, then, for each $i = 1, \dots, \ell$, there exist a sequence (ξ_{ik}) in $\Omega \cap \text{Fix}(G)$ and a least energy solution v_i to the problem $(\mathcal{P}_i^{\phi_i})$ such that, after passing to a subsequence,*

$$\lim_{k \rightarrow \infty} \varepsilon_k^{-1} \text{dist}(\zeta_{ik}, \partial\Omega) = \infty, \quad \lim_{k \rightarrow \infty} \varepsilon_k^{-1} |\zeta_{ik} - \zeta_{jk}| = \infty \text{ if } i \neq j, \quad \lim_{k \rightarrow \infty} \|u_{ik} - v_i(\varepsilon_k^{-1}(\cdot - \zeta_{ik}))\|_{\varepsilon_k} = 0,$$

for $i, j = 1, \dots, \ell$, and $\lim_{\varepsilon \rightarrow 0} c_\varepsilon^\ell(\Omega) = \sum_{i=1}^{\ell} c_i$. Moreover, $v_i \geq 0$ if $u_{ik} \geq 0$ for all $k \in \mathbb{N}$.

We start with the proof of the following statement.

Lemma 4.2. $\lim_{\varepsilon \rightarrow 0} c_\varepsilon^\ell(\Omega) = c_\infty^\ell$.

Proof. Let $\mathbf{u} = (u_1, \dots, u_\ell) \in \mathcal{N}_\varepsilon^\ell(\Omega)$ and define $\tilde{u}_i(z) := u_i(\varepsilon z)$. By Remark 2.2, $\tilde{\mathbf{u}} = (\tilde{u}_1, \dots, \tilde{u}_\ell) \in \mathcal{N}_\infty^\ell$ and $\mathcal{J}_\varepsilon^\ell(\mathbf{u}) = \mathcal{J}_\infty^\ell(\tilde{\mathbf{u}}) \geq c_\infty^\ell$. Therefore,

$$(4.1) \quad c_\infty^\ell \leq \liminf_{\varepsilon \rightarrow 0} c_\varepsilon^\ell.$$

To prove the opposite inequality, let $\mathbf{w} = (w_1, \dots, w_\ell) \in \mathcal{N}_\infty^\ell$. Fix $r > 0$ such that $B_r(0) \subset \Omega$ and a radial cut-off function $\chi \in C_c^\infty(\mathbb{R}^N)$ satisfying $\chi(x) = 1$ for $|x| \leq \frac{r}{2}$ and $\chi(x) = 0$ for $|x| > r$. For $\varepsilon > 0$ define $\mathbf{w}_\varepsilon = (w_{1\varepsilon}, \dots, w_{\ell\varepsilon})$ by

$$w_{i\varepsilon}(x) := w_i\left(\frac{x}{\varepsilon}\right)\chi(x) \quad i = 1, \dots, \ell.$$

Since $w_i \in H^1(\mathbb{R}^N)^{\phi_i}$ and χ is radial, we have that $w_{i\varepsilon} \in H^1(\mathbb{R}^N)^{\phi_i}$. As $\mathbf{w} \in \mathcal{N}_\infty^\ell$, from (2.2) and Lemma 2.3(i) we get

$$0 < c_0 \leq \|w_i\|^2 = \mu_i |w_i|_p^p + \sum_{\substack{j=1 \\ j \neq i}}^{\ell} \int_{\mathbb{R}^N} \lambda_{ij} \beta_{ij} |w_j|^{\alpha_{ij}} |w_i|^{\beta_{ij}} \quad \text{for all } i = 1, \dots, \ell.$$

Note that $w_{i\varepsilon} \rightarrow w_i$ in $H^1(\mathbb{R}^N)$ as $\varepsilon \rightarrow 0$. Therefore, for sufficiently small $\varepsilon > 0$, Lemma 2.3(ii) yields $\mathbf{t}_\varepsilon = (t_{1\varepsilon}, \dots, t_{\ell\varepsilon}) \in (0, \infty)^\ell$ such that $\mathbf{t}_\varepsilon \mathbf{w}_\varepsilon \in \mathcal{N}_\varepsilon^\ell(\Omega)$, and $t_{i\varepsilon} \rightarrow 1$ as $\varepsilon \rightarrow 0$. Thus,

$$c_\varepsilon^\ell(\Omega) \leq \mathcal{J}_\varepsilon^\ell(\mathbf{t}_\varepsilon \mathbf{w}_\varepsilon) = \frac{p-2}{2p} \|\mathbf{t}_\varepsilon \mathbf{w}_\varepsilon\|_{\ell, \varepsilon}^2 \rightarrow \frac{p-2}{2p} \|\mathbf{w}\|_\ell^2 = \mathcal{J}_\infty^\ell(\mathbf{w}) \quad \text{as } \varepsilon \rightarrow 0,$$

and, as a consequence,

$$(4.2) \quad \limsup_{\varepsilon \rightarrow 0} c_\varepsilon^\ell(\Omega) \leq c_\infty^\ell.$$

Inequalities (4.1) and (4.2) yield the desired identity. \square

Proof of Theorem 4.1. Applying Theorem 2.7 we obtain a least energy solution $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k})$ to the system $(\mathcal{S}_{\varepsilon_k, \Omega}^\phi)$ for every $k \in \mathbb{N}$. Define $\tilde{\mathbf{u}}_k = (\tilde{u}_{1k}, \dots, \tilde{u}_{\ell k})$ by taking $\tilde{u}_{ik}(z) := u_{ik}(\varepsilon_k z)$. From Remark 2.2 and Lemma 4.2 we see that $\tilde{\mathbf{u}}_k \in \mathcal{N}_\infty^\ell$ and $\mathcal{J}_\infty^\ell(\tilde{\mathbf{u}}_k) \rightarrow c_\infty^\ell$. Now we distinguish two cases:

(I) If $\text{Fix}(G) = \{0\}$, applying Theorem 3.3 to $\tilde{\mathbf{u}}_k$ we obtain a least energy solution $\mathbf{w} = (w_1, \dots, w_\ell)$ to the system $(\mathcal{S}_{\infty, \ell}^\phi)$ such that $\|u_{ik} - w_i(\varepsilon_k^{-1} \cdot)\|_{\varepsilon_k} = \|\tilde{u}_{ik} - w_i\| \rightarrow 0$. The identity $\lim_{\varepsilon \rightarrow 0} c_\varepsilon^\ell(\Omega) = c_\infty^\ell$ was proved in Lemma 4.2.

(II) If $\text{Fix}(G)$ has positive dimension, Theorem 3.3 applied to $\tilde{\mathbf{u}}_k$ yields a sequence (ξ_{ik}) in $\text{Fix}(G)$ and a least energy solution v_i to the problem $(\mathcal{P}_i^{\phi_i})$ for each $i = 1, \dots, \ell$, such that, setting $\zeta_{ik} := \varepsilon_k \xi_{ik}$, we get

$$\lim_{k \rightarrow \infty} \varepsilon_k^{-1} |\zeta_{ik} - \zeta_{jk}| = \infty \quad \text{if } i \neq j, \quad \text{and} \quad \|u_{ik} - v_i(\varepsilon_k^{-1}(\cdot - \zeta_{ik}))\|_{\varepsilon_k} = \|\tilde{u}_{ik} - v_i(\cdot - \xi_{ik})\| \rightarrow 0,$$

for $i, j = 1, \dots, \ell$. We are left with showing that $\zeta_{ik} \in \Omega$ and

$$\lim_{k \rightarrow \infty} \varepsilon_k^{-1} \text{dist}(\zeta_{ik}, \partial\Omega) = \infty \quad \text{for all } i = 1, \dots, \ell.$$

To this end, we recall that the points ξ_{ik} satisfy (3.4), i.e., there exist $\theta, R > 0$ such that

$$(4.3) \quad \int_{B_R(\xi_{ik})} \mu_i |\tilde{u}_{ik}|^p > \theta > 0 \quad \text{for all } k \in \mathbb{N} \text{ and } i = 1, \dots, \ell.$$

Note that $\text{supp}(\tilde{u}_{ik}) \subset \Omega_{\varepsilon_k}$, where $\Omega_{\varepsilon_k} := \{z \in \mathbb{R}^N : \varepsilon_k z \in \Omega\}$. So (4.3) implies that $\text{dist}(\xi_{ik}, \Omega_{\varepsilon_k}) < R$ for every $k \in \mathbb{N}$ and every i . A standard argument shows that $\text{dist}(\xi_{ik}, \partial\Omega_{\varepsilon_k}) \rightarrow \infty$, otherwise v_i would be zero in some open subset of \mathbb{R}^N , contradicting the unique continuation principle. In particular, we have that $\xi_{ik} \in \Omega_{\varepsilon_k}$ for all $k \in \mathbb{N}$. Rescaling yields $\zeta_{ik} \in \Omega$ and $\lim_{k \rightarrow \infty} \varepsilon_k^{-1} \text{dist}(\zeta_{ik}, \partial\Omega) = \infty$ for all $i = 1, \dots, \ell$, as claimed. This completes the proof. \square

Next, we derive Theorems 1.2 and 1.3 from Theorem 4.1.

Proof of Theorem 1.2. The group $G := \Gamma \times O(N-4)$ in Example 2.1(iii) satisfies (A_2) if $N = 4$ or $N \geq 6$, and $\text{Fix}(G) = \{0\}$. For $i = 1, \dots, m$ we take $\phi_i \equiv 1$, and for $i = m+1, \dots, \ell$ we take $\phi_i := \phi$ as in Example 2.1(iii). By Theorem 4.1 there exist a least energy solution $\hat{\mathbf{u}}_k = (\hat{u}_{1k}, \dots, \hat{u}_{\ell k})$ to the system $(\mathcal{S}_{\varepsilon_k, B_1(0)}^\phi)$ and a least energy solution $\mathbf{w} = (w_1, \dots, w_\ell)$ to the system $(\mathcal{S}_{\infty, \ell}^\phi)$ satisfying (1.1) such that, after passing to a subsequence,

$$\lim_{k \rightarrow \infty} \|\hat{u}_{ik} - w_i(\varepsilon_k^{-1} \cdot)\|_{\varepsilon_k} = 0 \quad \text{for all } i = 1, \dots, \ell,$$

and $\lim_{k \rightarrow \infty} \sum_{i=1}^{\ell} \|\widehat{u}_{ik}\|_{\varepsilon_k}^2 = \lim_{\varepsilon \rightarrow 0} c_\varepsilon^\ell(B_1(0)) = c_\infty^\ell = \sum_{i=1}^{\ell} \|w_i\|^2$. By Remark 2.6, we may take $\widehat{u}_{1k}, \dots, \widehat{u}_{mk}$ to be positive. Then, w_1, \dots, w_m are also positive. The symmetries ensure that the last $\ell - m$ components of $\widehat{\mathbf{u}}_k$ and \mathbf{w} are nonradial and change sign. \square

Proof of Theorem 1.3. The group $G := \Gamma$ in Example 2.1(ii) satisfies (A_2) , and $\text{Fix}(G) = \{0\} \times \mathbb{R}^{N-4}$ has positive dimension if $N \geq 5$. For $i = 1, \dots, m$ we take $\phi_i \equiv 1$, and for $i = m+1, \dots, \ell$ we take $\phi_i := \phi$ as in Example 2.1(ii). By Theorem 4.1, there exist a least energy solution $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k})$ to the system $(\mathcal{S}_{\varepsilon_k, B_1(0)}^\phi)$ and, for each $i = 1, \dots, \ell$, a sequence (ζ_{ik}) in $B_1(0) \cap \text{Fix}(G)$ and a least energy solution w_i to the problem $(\mathcal{P}_i^{\phi_i})$ such that, after passing to a subsequence,

$$\lim_{k \rightarrow \infty} \varepsilon_k^{-1} \text{dist}(\zeta_{ik}, \partial B_1(0)) = \infty, \quad \lim_{k \rightarrow \infty} \varepsilon_k^{-1} |\zeta_{ik} - \zeta_{jk}| = \infty \text{ if } i \neq j, \quad \lim_{k \rightarrow \infty} \|u_{ik} - w_i(\varepsilon_k^{-1}(\cdot - \zeta_{ik}))\|_{\varepsilon_k} = 0,$$

$$\text{for } i, j = 1, \dots, \ell, \text{ and } \lim_{k \rightarrow \infty} \sum_{i=1}^{\ell} \|u_{ik}\|_{\varepsilon_k}^2 = \lim_{\varepsilon \rightarrow 0} c_\varepsilon^\ell(B_1(0)) = \sum_{i=1}^{\ell} c_i = \sum_{i=1}^{\ell} \|w_i\|^2.$$

By Remark 2.6, we may take u_{1k}, \dots, u_{mk} to be positive. Then, w_1, \dots, w_m are also positive. It is well known that problem (\mathcal{P}_i) has a unique positive radial solution v_i and that any other positive solution is a translation of it [11]. Hence, there exists $\vartheta_i \in \mathbb{R}^N$ such that $v_i(y) = w_i(y + \vartheta_i)$ for every $y \in \mathbb{R}^N$ and $i = 1, \dots, m$.

Now take $i \in \{m+1, \dots, \ell\}$. Then, $w_i \in H^1(\mathbb{R}^N)^\phi$ with ϕ as in Example 2.1(ii), i.e., w_i satisfies

$$(4.4) \quad w_i(z_1, z_2, x) = w_i(e^{i\vartheta} z_1, e^{i\vartheta} z_2, x) \text{ for all } \vartheta \in [0, 2\pi), \quad w_i(z_1, z_2, x) = -w_i(z_2, z_1, x),$$

for all $(z_1, z_2, x) \in \mathbb{C} \times \mathbb{C} \times \mathbb{R}^{N-4}$. We claim that there exists $\vartheta_i \in \text{Fix}(G)$ such that $v_i(y) := w_i(y + \vartheta_i)$ satisfies (1.2). To prove this claim, observe that, if $\pi_j : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is the reflection on the hyperplane $\{(z, x_1, \dots, x_{N-4}) \in \mathbb{C}^2 \times \mathbb{R}^{N-4} : x_j = a\}$, $a \in \mathbb{R}$, then the function

$$\bar{w}(z, x) := \begin{cases} w_i(z, x) & \text{if } x_j > a, \\ (w_i \circ \pi_j)(z, x) & \text{if } x_j < a, \end{cases}$$

satisfies (4.4). With this remark and following Lopes' argument [15] (see also [29, Appendix C]) one shows that, for each $j = 1, \dots, N-4$, there exists $\theta_{ij} \in \mathbb{R}$ such that w_i is invariant with respect to the reflection on the hyperplane $x_j = \theta_{ij}$. Setting $\theta_i = (\theta_{i1}, \dots, \theta_{i, N-4})$ we infer that the function $v_i(z, x) := w_i(z, x + \theta_i)$ satisfies $v_i(z, x) = v_i(z, gx)$ for all $g \in O(N-4)$ and $(z, x) \in \mathbb{C}^2 \times \mathbb{R}^{N-4}$. Hence, setting $\vartheta_i = (0, \theta_i) \in \mathbb{C}^2 \times \mathbb{R}^{N-4}$, we have that $v_i(y) = w_i(y + \vartheta_i)$ satisfies (1.2), as claimed.

Let $\xi_{ik} := \zeta_{ik} + \varepsilon_k \vartheta_i$, where ϑ_i are the points defined in the previous paragraphs, so that $v_i(y) := w_i(y + \vartheta_i)$ is radial if $i = 1, \dots, m$ and it satisfies (1.2) if $i = m+1, \dots, \ell$. Note that, as $\varepsilon_k \rightarrow 0$, we have that $\xi_{ik} \in B_1(0)$ for k large enough. From the corresponding statements for ζ_{ik} and w_i one immediately derives

$$\lim_{k \rightarrow \infty} \varepsilon_k^{-1} \text{dist}(\xi_{ik}, \partial B_1(0)) = \infty, \quad \lim_{k \rightarrow \infty} \varepsilon_k^{-1} |\xi_{ik} - \xi_{jk}| = \infty \text{ if } i \neq j, \quad \lim_{k \rightarrow \infty} \|u_{ik} - v_i(\varepsilon_k^{-1}(\cdot - \xi_{ik}))\|_{\varepsilon_k} = 0,$$

$$\text{for } i, j = 1, \dots, \ell, \text{ and } \lim_{k \rightarrow \infty} \sum_{i=1}^{\ell} \|u_{ik}\|_{\varepsilon_k}^2 = \lim_{\varepsilon \rightarrow 0} c_\varepsilon^\ell(B_1(0)) = c_\infty^\ell = \sum_{i=1}^{\ell} c_i = \sum_{i=1}^{\ell} \|v_i\|^2.$$

The inequality $c_m < \widehat{c}_m$ follows from Proposition 3.1. \square

Finally, regarding positive solutions in the the unit ball $B_1(0)$ in \mathbb{R}^N , we derive the following result.

Theorem 4.3. *Let $N \geq 2$, $\Omega = B_1(0)$ and (ε_k) be a sequence of positive numbers converging to zero. Then there exist two solutions $\widehat{\mathbf{u}}_k = (\widehat{u}_{1k}, \dots, \widehat{u}_{\ell k})$ and $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k})$ to the system $(\mathcal{S}_{\varepsilon_k, \Omega})$ whose components \widehat{u}_{ik} and u_{ik} are positive and satisfy:*

- \widehat{u}_{ik} is radial and, after passing to a subsequence,

$$\lim_{k \rightarrow \infty} \|\widehat{u}_{ik} - \widehat{\omega}_i(\varepsilon_k^{-1} \cdot)\|_{\varepsilon_k} = 0 \quad \text{for all } i = 1, \dots, \ell,$$

where $\widehat{\omega}_k = (\widehat{\omega}_{1k}, \dots, \widehat{\omega}_{\ell k})$ is a least energy positive radial solution to the system $(\mathcal{S}_{\infty, \ell})$ in \mathbb{R}^N .

- For each $i = 1, \dots, \ell$ there is a sequence (ξ_{ik}) in $B_1(0)$ such that, after passing to a subsequence,

$$\lim_{k \rightarrow \infty} \varepsilon_k^{-1} \text{dist}(\zeta_{ik}, \partial\Omega) = \infty, \quad \lim_{k \rightarrow \infty} \varepsilon_k^{-1} |\zeta_{ik} - \zeta_{jk}| = \infty \text{ if } i \neq j, \quad \lim_{k \rightarrow \infty} \|u_{ik} - v_i(\varepsilon_k^{-1}(\cdot - \zeta_{ik}))\|_{\varepsilon_k} = 0,$$

for $i, j = 1, \dots, \ell$, where v_i is the positive radial solution to the problem (\mathcal{P}_i) .

Proof. The solutions $\widehat{\mathbf{u}}_k$ are obtained applying Theorem 4.1 with $G = O(N)$ and $\phi_i \equiv 1$ for all $i = 1, \dots, \ell$. The solutions $\mathbf{u}_k = (u_{1k}, \dots, u_{\ell k})$ are obtained applying Theorem 4.1 to the trivial group $G = \{1\}$ and $\phi_i \equiv 1$ for all $i = 1, \dots, \ell$. \square

The solutions \mathbf{u}_k were obtained and fully described in [12]. In addition, as mentioned in the introduction, the existence of a least energy radial solution to the system $(\mathcal{S}_{\infty, \ell})$ was shown by Sirakov in [23]. Theorem 4.3 gives an alternative proof of this fact. Finally, we point out that the existence of multiple positive solutions, both in \mathbb{R}^N and in bounded domains, has been established, for instance, in [1, 9, 19, 27, 28]. Positive solutions of singularly perturbed systems with potentials have been obtained, e.g., in [17, 20].

References

- [1] Bartsch, Thomas; Dancer, Norman; Wang, Zhi-Qiang: A Liouville theorem, a-priori bounds, and bifurcating branches of positive solutions for a nonlinear elliptic system. *Calc. Var. Partial Differential Equations* 37 (2010), no. 3-4, 345–361.
- [2] Bartsch, Thomas; Willem, Michel: Infinitely many nonradial solutions of a Euclidean scalar field equation. *J. Funct. Anal.* 117 (1993), no. 2, 447–460.
- [3] Bracho, Javier; Clapp, Mónica; Marzantowicz, Waclaw: Symmetry breaking solutions of nonlinear elliptic systems. *Topol. Methods Nonlinear Anal.* 26 (2005), no. 1, 189–201.
- [4] Byeon, Jaeyoung; Lee, Youngae; Moon, Sang-Hyuck: Partly clustering solutions of nonlinear Schrödinger systems with mixed interactions. *J. Funct. Anal.* 280 (2021), no. 12, Paper No. 108987, 54 pp.
- [5] Chen, Zhijie; Lin, Chang-Shou; Zou, Wenming: Infinitely many sign-changing and semi-nodal solutions for a nonlinear Schrödinger system. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 15 (2016), 859–897.
- [6] Cingolani, Silvia; Clapp, Mónica; Secchi, Simone: Multiple solutions to a magnetic nonlinear Choquard equation. *Z. Angew. Math. Phys.* 63 (2012), no. 2, 233–248.
- [7] Clapp, Mónica; Srikanth, P. N.: Entire nodal solutions of a semilinear elliptic equation and their effect on concentration phenomena. *J. Math. Anal. Appl.* 437 (2016), no. 1, 485–497.
- [8] Clapp, Mónica; Szulkin, Andrzej: A simple variational approach to weakly coupled competitive elliptic systems. *NoDEA Nonlinear Differential Equations Appl.* 26 (2019), no. 4, Paper No. 26, 21 pp.
- [9] Dancer, E. N.; Wei, Juncheng; Weth, Tobias: A priori bounds versus multiple existence of positive solutions for a nonlinear Schrödinger system. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 27 (2010), no. 3, 953–969.
- [10] Esry, B. D.; Greene, Chris H.; Burke, Jr., James P.; Bohn, John L: Hartree-Fock theory for double condensates, *Phys. Rev. Lett.* 78 (1997), 3594–3597.

- [11] Kwong, Man Kam: Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbb{R}^n . *Arch. Rational Mech. Anal.* 105 (1989), no. 3, 243–266.
- [12] Lin, Tai-Chia; Wei, Juncheng: Spikes in two coupled nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 22 (2005), no. 4, 403–439.
- [13] Lin, Tai-Chia; Wei, Juncheng: Ground state of N coupled nonlinear Schrödinger equations in \mathbb{R}^n , $n \leq 3$. *Comm. Math. Phys.* 255 (2005), no. 3, 629–653.
- [14] Liu, Jiaquan; Liu, Xiangqing; Wang, Zhi-qiang: Multiple mixed states of nodal solutions for nonlinear Schrödinger systems. *Calc. Var. Partial Differential Equations* 52 (2015), no. 3-4, 565–586.
- [15] Lopes, Orlando: Radial symmetry of minimizers for some translation and rotation invariant functionals. *J. Differential Equations* 124 (1996), no. 2, 378–388.
- [16] Lorca, Sebastián; Ubilla, Pedro: Symmetric and nonsymmetric solutions for an elliptic equation on \mathbb{R}^N . *Nonlinear Anal.* 58 (2004), no. 7-8, 961–968.
- [17] Montefusco, Eugenio; Pellacci, Benedetta; Squassina, Marco: Semiclassical states for weakly coupled nonlinear Schrödinger systems. *J. Eur. Math. Soc. (JEMS)* 10 (2008), no. 1, 47–71.
- [18] Musso, Monica; Pacard, Frank; Wei, Juncheng: Finite-energy sign-changing solutions with dihedral symmetry for the stationary nonlinear Schrödinger equation. *J. Eur. Math. Soc. (JEMS)* 14 (2012), no. 6, 1923–1953.
- [19] Noris, Benedetta; Ramos, Miguel: Existence and bounds of positive solutions for a nonlinear Schrödinger system. *Proc. Amer. Math. Soc.* 138 (2010), no. 5, 1681–1692.
- [20] Pomponio, Alessio: Coupled nonlinear Schrödinger systems with potentials. *J. Differential Equations* 227 (2006), no. 1, 258–281.
- [21] Sato, Yohei; Wang, Zhi-Qiang: On the multiple existence of semi-positive solutions for a nonlinear Schrödinger system. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 30 (2013), no. 1, 1–22
- [22] Sato, Yohei; Wang, Zhi-Qiang: On the least energy sign-changing solutions for a nonlinear elliptic system. *Discrete Contin. Dyn. Syst.* 35 (2015), no. 5, 2151–2164.
- [23] Sirakov, Boyan: Least energy solitary waves for a system of nonlinear Schrödinger equations in \mathbb{R}^n . *Comm. Math. Phys.* 271 (2007), no. 1, 199–221.
- [24] Tavares, Hugo; Terracini, Susanna: Sign-changing solutions of competition-diffusion elliptic systems and optimal partition problems. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 29 (2012), no. 2, 279–300.
- [25] Wang, Weichung; Wu, Tsung-Fang; Liu, Chien-Hsiang: On the multiple spike solutions for singularly perturbed elliptic systems, *Discrete Continuous Dynamical Systems Series B* 18 (2013), no 1, 237-258.
- [26] Wei, Juncheng; Wu, Yuanze: Ground states of nonlinear Schrödinger systems with mixed couplings. *J. Math. Pures Appl.* (9) 141 (2020), 50–88.
- [27] Wei, Juncheng; Weth, Tobias: Nonradial symmetric bound states for a system of coupled Schrödinger equations. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* 18 (2007), no. 3, 279–293.
- [28] Wei, Juncheng; Weth, Tobias: Radial solutions and phase separation in a system of two coupled Schrödinger equations. *Arch. Ration. Mech. Anal.* 190 (2008), no. 1, 83–106.
- [29] Willem, Michel: *Minimax theorems. Progress in Nonlinear Differential Equations and their Applications* 24. Birkhäuser Boston, Inc., Boston, MA (1996).

Mónica Clapp and Mayra Soares
Instituto de Matemáticas
Universidad Nacional Autónoma de México
Circuito Exterior, Ciudad Universitaria
04510 Coyoacán, Ciudad de México, México
monica.clapp@im.unam.mx
ssc_mayra@im.unam.mx