

HÖLDER REGULARITY FOR NON-VARIATIONAL POROUS MEDIA TYPE EQUATIONS

HÉCTOR A. CHANG-LARA AND MAKSON S. SANTOS

ABSTRACT. We present a Krylov-Safonov regularity approach for the Hölder regularity of viscosity solutions to non-variational porous media type equations.

1. INTRODUCTION

Consider the evolution problem modeled by a continuous and non-negative function $u \in \mathcal{C}(B_1 \times (-1, 0])$ and driven by an equation of the form

$$(1.1) \quad \partial_t u = \sum_{i,j=1}^n u a_{ij} \partial_{ij} u + b_{ij} \partial_i u \partial_j u \quad \text{in } B_1 \times (-1, 0]$$

where $a_{ij} = a_{ij}(x), b_{ij} = b_{ij}(x) \in \mathbb{R}^{n \times n}$ are bounded measurable coefficients that satisfy the uniform ellipticity hypothesis

$$\sum_{i,j=1}^n a_{ij} \xi_i \xi_j, \sum_{i,j=1}^n b_{ij} \xi_i \xi_j \in [\lambda, \Lambda] \subseteq (0, \infty) \quad \forall \xi \in \partial B_1$$

(denoted by $(a_{ij}), (b_{ij}) \in [\lambda, \Lambda] \subseteq (0, \infty)$ from now on). The degeneracy of the diffusion as u goes to zero is certainly the most interesting feature of these equations from the regularity point of view. This is responsible of the finite speed of propagation for the support of the solution, and gives one of the fundamental examples in free boundary problems.

In this article we pursue the regularity theory for these type of equations. To illustrate the variety of challenges encountered, consider the solution $(e \cdot x + t)_+$ for some $e \in \mathbb{R}^n$ such that $\sum_{i,j=1}^n b_{ij} e_i e_j = 1$. This traveling front exhibits two interesting behaviors: The impossibility of a Harnack inequality, and the fact that we should not expect solutions to be better than Lipschitz. Here is our main theorem.

Theorem 1.1. *Let $(a_{ij}), (b_{ij}) \in [\lambda, \Lambda] \subseteq (0, \infty)$ and $u \in \mathcal{C}(B_1 \times (-1, 0])$ be a non-negative solution of (1.1). Then*

$$\|u\|_{\mathcal{C}^\alpha(B_{1/2} \times (-1/2, 0])} \leq C \|u\|_{\mathcal{C}^0(B_1 \times (-1, 0])}.$$

where $\alpha \in (0, 1)$ and $C > 0$ are constants depending only on λ, Λ , and the dimension n .

The classical example to keep in mind is the porous media equation (PME), historically developed from material sciences and fluid dynamics. Here u is a density being transported by a potential flow and modeled by the continuity equation $\partial_t u - \operatorname{div}(u Dp) = 0$. The pressure p gets related to the density u by a constitutive relation which takes the form $p = \phi(u)$, in particular if $p = u$ we get the Boussinesq equation $\partial_t u = u \Delta u + |Du|^2$.

Besides the classical motivation in fluid dynamics, the PME appears in other very interesting settings such as biological models [18, 26] and differential geometry [31, 13]. For a detailed exposition of the motivating problems we recommend the first chapters in the book [29] by Vázquez.

Due to its diverse applications and fascinating nonlinear structure, the PME and its generalizations have attracted the attention of many authors over the years. Most of the developments account for distributional weak solutions, including unique solvability, regularity, finite speed of propagation, and asymptotic behavior. The regularity of solutions in several dimensions was established in 1979 by Caffarelli and Friedman [6, 7] based on a priori estimates due to Aronson and Bénéilan [1] around the same time. The theory was then quickly extended to the two phase Stefan problem by Caffarelli and Evans [5] and very general singular equations in divergence form by Ziemer [32], DiBenedetto [14, 15], and Sacks [27]; all of them around 1982 and based on the de Giorgi-Nash-Moser approach. Besides Chapter 7 in the book of Vázquez [29], we also recommend the books by DiBenedetto, Urbano, and Vespi [16], and Urbano [28] for a detailed and pedagogical analysis on the regularity theory of degenerate equations.

Meanwhile the degeneracy under the variational structure is well understood, the non-variational counterpart still offers some open questions. In this case the natural setting for the existence, uniqueness, and stability theorems are the viscosity solutions developed also in the eighties and nineties. For the classical PME, this treatment was started by Caffarelli and Vázquez in 1996 [10] and extended by Brändle and Vázquez in 2005 [2].

This work continues this line of research by analyzing the regularity of solutions to PME type equations in non-divergence form. In this sense our methods belong to the Krylov-Safonov regularity theory, originally presented in a probabilistic setting in [23, 24]. The main distinction between the variational and non-variational approach is that the energy estimates are replaced by the Aleksandrov-Bakelman-Pucci (ABP) principle. Standard references are the book by Gilbarg and Trudinger [17] or the one by Cabré and Caffarelli [4].

Let us emphasize that because of the degeneracy, we cannot expect Harnack's inequality to hold in this scenario. However, as u approaches zero, the evolution is driven by an eikonal equation which controls the speed of propagation of the level sets. We like to think about it in the following terms, either the equation falls in a uniformly elliptic regime or the eikonal mechanism takes care of the regularity. This is also, in broad sense, the point of view taken for the regularity theory of the Stefan equation in [5].

To make this idea precise, we take into account the scale invariance of the equation and pursue a diminish of oscillation lemma. For a solution u taking values in $[0, 1]$ we consider two alternative scenarios in measure: Either $\{u \leq 1/2\}$ covers a positive fraction of the domain or $\{u > 1/2\}$ takes most of it instead. The first case leads to an improvement of the oscillation from above and it is the easiest to handle; in this instance the equation is uniformly elliptic. For the second case we show that u becomes positive in a smaller cylinder (and from then on falls in a uniformly elliptic regime) by proving an ABP-type measure estimate adapting Mooney's clever approach in [25] to degenerate parabolic equations.

In the preliminary Section 2 we review the notion of viscosity solution, give a precise statement of our theorem, and finally state the lemmas that build our result. Section 3 takes care of the improvement from below, meanwhile in Section 4 we do the improvement from

above. In the concluding Section 5 we recapitulate the whole strategy and comment on some further extensions.

Acknowledgment: Both authors were supported by CONACyT-MEXICO Grant A1-S-48577.

2. PRELIMINARIES

2.1. Geometric configurations. The *open ball* in \mathbb{R}^n of radius r centered at x is denoted by $B_r(x) := \{y \in \mathbb{R}^n : |y - x| < r\}$ and by default $B_r := B_r(0)$. The *open cube* of length l and centered at $x = (x_1, \dots, x_n)$ is

$$Q_l(x) := (x_1 - l/2, x_1 + l/2) \times \dots \times (x_n - l/2, x_n + l/2), \quad Q_l := Q_l(0)$$

2.2. Operators. Given $[\lambda, \Lambda] \subseteq (0, \infty)$ we define the Pucci's extremal operators acting on $(M_{ij}) \in \mathbb{R}^{n \times n}$ as

$$\mathcal{M}^+ M := \sup_{(a_{ij}) \in [\lambda, \Lambda]} \sum_{i,j=1}^n a_{ij} M_{ij} = \sum_{e \in \text{eig}(M_{ij})} \Lambda e_+ - \lambda e_-, \quad \mathcal{M}^- M := -\mathcal{M}^+(-M)$$

where $\text{eig}(M_{ij})$ is the set of eigenvalues and $e_{\pm} = \max(\pm e, 0)$ are the positive and negative parts.

This family of operators allows us to give a precise meaning to the equation (1.1) which is from now on understood in terms of the following inequalities

$$u \mathcal{M}^-(D^2 u) + \lambda |Du|^2 \leq \partial_t u \leq u \mathcal{M}^+(D^2 u) + \Lambda |Du|^2.$$

Remark 2.1. *These inequalities are an invariant family under horizontal translations and the following scaling transformation for any exponent $\alpha \in \mathbb{R}$. If u satisfies any of the inequalities above in a domain $\Omega \times (t_i, t_f]$ then $v(x, t) = r^{-\alpha} u(rx, r^\beta t)$ also satisfies the corresponding inequality in $r^{-1} \Omega \times (r^{-\beta} t_i, r^{-\beta} t_f]$ provided that $\beta = 2 - \alpha$.*

2.3. Viscosity solutions. In this section $F = F(M, p, z)$ will be a continuous function representing the fully-nonlinear operator driving the equation $\partial_t u = F(D^2 u, Du, u)$ over $\Omega \times (t_i, t_f]$. We should mainly keep in mind $F = z \mathcal{M}^\pm M + b|p|^2$. Our goal is to define a weak notion of solution for this dynamic.

A *classical sub-solution* of

$$\partial_t u \leq F(D^2 u, Du, u) \text{ in } \Omega \times (t_i, t_f]$$

is a non-negative function $u \in \mathcal{C}(\Omega \times (t_i, t_f])$ that is smooth over $\text{spt } u \cap \Omega \times (t_i, t_f]$, with $\partial \text{spt } u \cap \Omega \times (t_i, t_f]$ also (locally) smooth, and such that it satisfies in the classical point-wise sense

$$\partial_t u \leq F(D^2 u, Du, u) \text{ in } \text{spt } u \cap \Omega \times (t_i, t_f].$$

We say that the sub-solution is *strict* if the inequality is also strict at every point. (*Strict classical super-solutions* are defined in a similar way but with respect to the opposite inequalities.

The notion of viscosity solutions is given in terms of the *comparison principle* that we should certainly expect for operators of the form $F = z \mathcal{M}^\pm M + b|p|^2$ with $z \geq 0$. To be

precise, a (classical) sub-solution u that starts smaller than a (classical) super-solution v at some time, and never cross it on the boundary of the domain, must remain smaller in the interior at future times. Otherwise we get a contradiction at the first contact point using the strong comparison principle or Hopf's Lemma, if the first contact point happens to be on the intersection of the free boundaries. Let us make this idea rigorous.

Consider a (parabolic) neighborhood $\mathcal{N} = B_r(x_0) \times (t_0 - r, t_0]$ of (x_0, t_0) . For a pair of non-negative functions $u, v \in \mathcal{C}(\mathcal{N})$ such that $u \leq v$ and $(x_0, t_0) \in \text{spt } u \cap \Omega \times (t_i, t_f]$ (and hence also $\in \text{spt } v \cap \Omega \times (t_i, t_f]$), we say that v *touches* u from above at (x_0, t_0) if $v(x_0, t_0) = u(x_0, t_0)$. We define the contact from below in an analogous way.

Let $D \subseteq \mathbb{R}^n \times \mathbb{R}$ be open (in the parabolic sense), i.e. for every $(x_0, t_0) \in D$ there exists some $\mathcal{N} = B_r(x_0) \times (t_0 - r, t_0] \subseteq D$. We say that $u \in \mathcal{C}(D)$ is a *viscosity sub-solution* of

$$\partial_t u \leq F(D^2 u, Du, u) \text{ in } D$$

if for every $\mathcal{N} = B_r(x_0) \times (t_0 - r, t_0] \subseteq D$, u cannot be touched from above by a strict classical super-solution of the same problem in \mathcal{N} . The notion of *viscosity super-solution* is given in similar way by ruling out the contact from below with strict classical sub-solutions. Finally, *viscosity solutions* are those which are simultaneously viscosity sub and super-solutions with respect to the same operator.

Most of the time our domains are just cylinders, however in the proof of Lemma 3.3 we also consider solutions in arbitrary open sets.

The existence, uniqueness and stability of viscosity solutions to free boundary problems of the form $\partial_t u = F(D^2 u, Du, u)$ is a delicate issue that we will not pursue in this article. Whenever we invoke the comparison principle at least one of the solutions will be classical, so the conclusions just follow from the definitions. A careful analysis of the well-posedness by the viscosity approach has been studied for instance in [10, 2] for equations of the form

$$\partial_t u = a(u)\Delta u + |Du|^2$$

with even weaker assumption on the diffusion a than we have (Brändle and Vázquez allow a to be sub-linear as $u \rightarrow 0^+$). Here we would like to highlight as well a closely related line of research for several free boundary problems such as Hele-Shaw and Stefan [21], Richards equation [22], or the PME with drift [20].

2.4. Main result and overview of the strategy. Here is the precise version of the theorem stated at the introduction.

Theorem 2.2. *Given $[\lambda, \Lambda] \subseteq (0, \infty)$ there exist $\alpha \in (0, 1)$ and $C > 0$ such that if $u \in \mathcal{C}(B_1 \times (-1, 0])$ is a non-negative function that satisfies the following inequalities in the viscosity sense*

$$\begin{cases} \partial_t u \geq u\mathcal{M}^-(D^2 u) + \lambda|Du|^2 & \text{in } B_1 \times (-1, 0] \\ \partial_t u \leq u\mathcal{M}^+(D^2 u) + \Lambda|Du|^2 & \text{in } B_1 \times (-1, 0] \end{cases}$$

then $u \in \mathcal{C}^\alpha(B_{1/2} \times (-1/2, 0])$ with an estimate of the form

$$\|u\|_{\mathcal{C}^\alpha(B_{1/2} \times (-1/2, 0])} \leq C\|u\|_{\mathcal{C}^0(B_1 \times (-1, 0])}.$$

By the translation and scale invariance of the problem (Remark 2.1) it suffices to establish the decay of oscillation reflected in the next two lemmas. We consider two possible

alternatives for u in measure in order to discriminate in which direction does the oscillation improve. First we state the most delicate case, whether u is a super-solution taking values in $[0, 1]$, that is larger than $1/2$ most of the time. In this case the oscillation improves from below and then the equation becomes uniformly parabolic. The following lemma is proven in the next section.

Lemma 2.3 (Improvement from below). *Given $[\lambda, \Lambda] \subseteq (0, \infty)$ there exist $\eta, \theta \in (0, 1)$ such that if $u \in \mathcal{C}(B_1 \times (-1, 0])$ is a non-negative function, taking values in $[0, 1]$, that satisfies the following super-solution inequality in the viscosity sense*

$$\partial_t u \geq u\mathcal{M}^-(D^2u) + \lambda|Du|^2 \text{ in } B_1 \times (-1, 0]$$

then

$$|\{u > 1/2\} \cap B_1 \times (-1, 0]| \geq (1 - \eta)|B_1 \times (-1, 0]| \quad \Rightarrow \quad \inf_{B_{1/2} \times (-1/2, 0]} u \geq \theta.$$

Once fixed the fraction η in terms of the previous lemma we can consider the alternative scenario, namely u being less than $1/2$ an η -fraction of the measure. The following lemma is proved in Section 4.

Lemma 2.4 (Improvement from above). *Given $[\lambda, \Lambda] \subseteq (0, \infty)$ and $\eta > 0$, there exists $\theta \in (0, 1)$ such that if $u \in \mathcal{C}(B_1 \times (-1, 0])$ is a non-negative function, taking values in $[0, 1]$, that satisfies the following sub-solution inequality in the viscosity sense*

$$\partial_t u \leq u\mathcal{M}^+(D^2u) + \Lambda|Du|^2 \text{ in } B_1 \times (-1, 0]$$

then

$$|\{u \leq 1/2\} \cap B_1 \times (-1, 0]| \geq \eta|B_1 \times (-1, 0]| \quad \Rightarrow \quad \sup_{B_{1/2} \times (-1/2, 0]} u \leq (1 - \theta).$$

3. IMPROVEMENT FROM BELOW

In this section we prove the contrapositive of Lemma 2.3 in a convenient geometric configuration. Let us anticipate that the choice $R = 5\sqrt{n}$ and $T = 2n$ will be explained later on.

Lemma 3.1 (Measure estimate). *Given $[\lambda, \Lambda] \subseteq (0, \infty)$ there exist $\eta \in (0, 1)$ and $M > 0$ such that if $u \in \mathcal{C}(B_R \times (-T, 1])$ is a non-negative viscosity super-solution*

$$\partial_t u \geq u\mathcal{M}^-(D^2u) + \lambda|Du|^2 \text{ in } B_R \times (-T, 1].$$

then

$$u(0, 1) \leq 1 \quad \Rightarrow \quad |\{u \leq M\} \cap B_R \times [-T, 0]| \geq \eta|B_R \times [-T, 0]|$$

Lemma 2.3 follows from this result by standard scaling and covering arguments. The idea in the uniformly elliptic or parabolic regime is known as the ABP Lemma. It consists on showing that there is a set of positive measure where u can be touched by a family of paraboloids [3].

3.1. Paraboloids and contact sets. A concave *paraboloid* centered at the *vertex* $(y, s) \in \mathbb{R}^n \times \mathbb{R}$, of opening $\alpha > 0$ in space, and $\beta > 0$ in time, is the quadratic function

$$P_{y,s}^{\alpha,\beta}(x, t) := -\frac{\alpha}{2}|x - y|^2 + \beta(t - s).$$

In order to simplify the notation we may use $P_{y,s}^\alpha := P_{y,s}^{\alpha,\alpha}$, $P^{\alpha,\beta} := P_{0,0}^{\alpha,\beta}$, and $P^\alpha := P^{\alpha,\alpha}$.

Remark 3.2. *The paraboloids are related by the Lipschitz (in space and time) scaling transformation*

$$P_{y,s}^{r\alpha,\beta}(x, t) = r^{-1}P_{ry,rs}^{\alpha,\beta}(rx, rt)$$

Keep in mind that by Remark 2.1, this Lipschitz transformation also leaves invariant the equations under consideration.

We define the *contact set* of $u \in \mathcal{C}(\Omega \times (t_i, t_f])$ with respect to a set of vertices $B \subseteq \mathbb{R}^n \times \mathbb{R}$ as

$$A^{\alpha,\beta}(B) := \{(x, t) \in \Omega \times (t_i, t_f] : \exists (y, s) \in B \text{ such that } P_{y,s}^{\alpha,\beta} \text{ touches } u \text{ from below at } (x, t)\}$$

We may shorten the notation to $A^{\alpha,\beta}(\{(x, t)\}) := A^{\alpha,\beta}(x, t)$ and $A^\alpha(B) := A^{\alpha,\alpha}(B)$.

3.2. ABP-type lemma. In the following lemma the parameters $\alpha = 100 \max(1, \lambda)$ and $m = \min(1, 1/(n\Lambda))$ are fixed. The reason for this choice will be made clear later on. For the moment notice the following facts used in the proof below:

$$A^\alpha(0, 0) \subseteq \text{spt}(P^\alpha)_+ \subseteq B_{\sqrt{2}} \times (0, 1] \subseteq B_{R-1} \times (0, 1]$$

and under the hypothesis $u(0, 1) \leq 1$ from Lemma 3.1 we get that $\alpha \geq 1$ implies $A^\alpha(0, 0) \neq \emptyset$. Hence, the the main point of the hypothesis in the next lemma is whether at least one of the points in $A^\alpha(0, 0)$ falls in the uniformly elliptic regime.

Lemma 3.3 (ABP). *Given $[\lambda, \Lambda] \subseteq (0, \infty)$ there exist $M, \eta > 0$ such that if $u \in \mathcal{C}(B_R \times (0, 1])$ is a non-negative viscosity super-solution*

$$(3.2) \quad \partial_t u \geq u\mathcal{M}^-(D^2u) \text{ in } B_R \times (0, 1].$$

then

$$A^\alpha(0, 0) \cap \{u > m\} \neq \emptyset \quad \Rightarrow \quad |\{u \leq M\} \cap B_R \times (0, 1]| \geq \eta$$

Proof. As a first step we will construct a set B of vertices in terms of the contact point $(x_0, t_0) \in A^\alpha(0, 0) \cap \{u > m\} \subseteq B_{R-1} \times (0, 1]$ such that

$$A^{2\alpha}(B) \subseteq B_R \times (0, 1] \quad \text{and} \quad A^{2\alpha}(x, t) \cap \{u > m/2\} \neq \emptyset \quad \forall (x, t) \in B.$$

Notice that $A^{2\alpha}(B) \subseteq \{u < M\}$ for some $M = C\alpha$ sufficiently large. The second step in the proof will be to establish a lower bound on $|A^{2\alpha}(B)|$.

Let $r = m/(8\alpha)$ such that

$$\begin{aligned} B_{3r}(x_0) \times (t_0 - r^2, t_0] &\subseteq \{P^\alpha > m/2\} \cap \mathbb{R}^n \times (t_0 - r^2, t_0] \\ &\subseteq \{u > m/2\} \cap \mathbb{R}^n \times (t_0 - r^2, t_0] \subseteq B_R \times (0, 1]. \end{aligned}$$

Consider the family of paraboloids

$$P_{x/2, t/2}^{2\alpha} = P^\alpha + P_{x,t}^\alpha \quad (x, t) \in B_r(x_0) \times (t_0 - r^2, t_0 - r^2/2]$$




FIGURE 1. Set of vertices.

which determines our set of vertices (see Figure 1)

$$B := B_{r/2}(x_0/2) \times (t_0/2 - r^2/2, t_0/2 - r^2/4).$$

Notice that

$$\{P_{x,t}^\alpha > 0\} \cap \mathbb{R}^n \times (t, t_0] \subseteq B_{2r}(x) \times (t, t_0] \subseteq B_{3r}(x_0) \times (t, t_0] \subseteq \{u > m/2\} \cap \mathbb{R}^n \times (t, t_0]$$

Hence, $P_{x/2, t/2}^{2\alpha} \leq P^\alpha$ over $(\mathbb{R}^n \setminus B_{3r}(x_0)) \times (t, t_0]$. This means that if $P_{x/2, t/2}^{2\alpha}$ has contact with u in the time interval $(t, t_0]$, then it will necessarily occur over the set $\{u > m/2\}$. Also $P_{x,t}^\alpha(x_0, t_0) = -\frac{\alpha}{2}|x_0 - x|^2 + \alpha(t_0 - t) \geq 0$, therefore $P_{x/2, t/2}^{2\alpha}$ will definitively have contact with u over the time interval $(t, t_0]$. This verifies the desired properties on B and finishes the first step.

Recall that by B and α being bounded and $A^{2\alpha}(B) \subseteq \{u > m/2\}$ we get that $A^{2\alpha}(B) \subseteq \{u \in (m/2, M)\}$, for some M sufficiently large. Over this regime we get to replace the factor u in front of Pucci operator by a constant giving us, for some larger ellipticity interval $[\bar{\lambda}, \bar{\Lambda}] \subseteq (0, \infty)$ and $\bar{\mathcal{M}}^- = \mathcal{M}^-(\bar{\lambda}, \bar{\Lambda})$

$$\partial_t u \geq \bar{\mathcal{M}}^- u \text{ in } \{u \in (m/2, M)\} \cap B_R \times (0, 1].$$

From now on we will assume, without loss of generality, that u is semi-concave in space and time, otherwise regularize using the inf-convolution (see [30, Lemma 4.2] which refers back to [12, Section 8]). Under this regularity we can define (up to a set of measure zero) the map from the contact points $(x, t) \in A^{2\alpha}(B)$ to the unique vertex $(y, s) \in B$ such that $P_{y,s}^{2\alpha}$ touches u from below at (x, t)

$$(x, t) \mapsto (y, s) = \left(x + \frac{1}{2\alpha} Du(x, t), t - \frac{1}{2\alpha} u(x, t) - \frac{1}{4\alpha^2} |Du(x, t)|^2\right).$$

The facts proven on the previous step about contact set guarantee that the map is well defined and surjective. The determinant of the Jacobian can be computed as

$$\det \begin{pmatrix} I + \frac{1}{2\alpha} D^2 u & \frac{1}{2\alpha} D \partial_t u \\ -\frac{1}{2\alpha} Du^T - \frac{1}{2\alpha^2} Du^T D^2 u & 1 - \frac{1}{2\alpha} \partial_t u - \frac{1}{2\alpha^2} Du \cdot D \partial_t u \end{pmatrix} = \det \begin{pmatrix} I + \frac{1}{2\alpha} D^2 u & \frac{1}{2\alpha} D \partial_t u \\ 0 & 1 - \frac{1}{2\alpha} \partial_t u \end{pmatrix}.$$

By invoking the area formula for Lipschitz maps we get that

$$|B| = \iint_{A^{2\alpha}(B)} (1 - \frac{1}{2\alpha} \partial_t u) \det(I + \frac{1}{2\alpha} D^2 u).$$

At the contact set we immediately have that $D^2 u \geq -2\alpha I$, and $\partial_t u \leq 2\alpha$. Recall that over $\{u \in (m/2, M)\}$, u satisfies a uniformly elliptic equation and then we can use the equation to bound the derivatives from the other side

$$2\alpha \geq \partial_t u \geq \bar{\lambda} |(D^2 u)^+| - \bar{\Lambda} |(D^2 u)^-| \geq -C\alpha.$$

By applying this estimate to the area formula we get the desired inequality. \square

The previous proof can also be adapted to the following version needed in the iterative procedure for the next section. Keep in mind that the equations are invariant by Lipschitz rescalings, see Remark 2.1 for $\alpha = 1$ and Remark 3.2 for the paraboloids. The choice $\tau = n$ will be explained later on.

Corollary 3.4. *Given $[\lambda, \Lambda] \subseteq (0, \infty)$ there exist $M, \eta > 0$ such that for any $k \in \mathbb{N}$ it holds that if $u \in \mathcal{C}(B_{2^{-k}R} \times (0, 2^{-k}(2\tau + 2)))$ is a non-negative viscosity super-solution*

$$\partial_t u \geq u \mathcal{M}^-(D^2 u) \text{ in } B_{2^{-k}R} \times (0, 2^{-k+1}(2\tau + 2)).$$

then

$$A^{2^k \alpha, \alpha}(0, 0) \cap \{u > 2^{-k}m\} \neq \emptyset \quad \Rightarrow \quad \frac{|\{u \leq M\} \cap B_{2^{-k}R} \times (0, 2^{-k}(2\tau + 2))|}{|B_{2^{-k}R} \times (0, 2^{-k}(2\tau + 2))|} \geq \eta$$

As in the previous lemma we need to have some distance between the contact point and the boundary of the domain. This is the case if

$$\text{spt}(P^\alpha)_+ \cap \mathbb{R}^n \times (0, 2\tau + 2] \subseteq B_{R-1} \times (0, 2\tau + 2]$$

for which it suffices to ask $R - 1 \geq 2\sqrt{\tau + 1}$.

3.3. Measure estimate at a fixed time. The previous ABP-type lemma only applies if we know that the contact point falls on the uniformly elliptic regime. In this section we lift such requirement by implementing a selection algorithm iterated over dyadic cubes as in [25]. The result of this procedure is (in the worst case) a measure estimate in space and at time $t = -2n$.

3.3.1. Algorithm and parameters. Let us settle some notation first. The unit cube $Q_1 = (-1/2, 1/2)^n \subseteq \mathbb{R}^n$ decomposes into 2^n congruent and disjoint sub-cubes $Q_{2^{-1}}(x_j^1)$ to which we refer as the first generation. To be precise,

$$x_j^1 := \frac{\sqrt{n}}{4}(j_1, \dots, j_n) \text{ where } j := (j_1, j_2, \dots, j_n) \in \{\pm 1\}^n \text{ is a multi-index.}$$




FIGURE 2. Geometric configuration for the selection algorithm.

The k^{th} generation consists of 2^{nk} congruent and disjoint sub-cubes $\{Q_{2^{-k}}(x_j^k)\}$ generated from a similar decomposition for each cube in the previous generation. We could give a precise rule to define x_j^k as before, however we will not need these details in our arguments, so we prefer not to distract the main flow of ideas with these rather technical aspect.

We say that a cube $Q_{2^{-k}}(x_j^k)$ is a descendant of $Q_{2^{-(k-1)}}(x_{j'}^{k-1})$ if the first cube is one of the 2^n sub-cubes obtained from the decomposition of the second. Reciprocally, we say that $Q_{2^{-(k-1)}}(x_{j'}^{k-1})$ is the progenitor of $Q_{2^{-k}}(x_j^k)$. Each cube $Q_{2^{-k}}(x_j^k)$ is associated with a time interval

$$(t_i^k := t_i^{k-1} - 2^{-(k-1)}\tau, t_f^k := t_i^k + 2^{-k}(2\tau + 2)], \quad t_i^0 = 0$$

to be determined once we declare τ , and with the only exception $t_f^0 = 1$, see Figure 2.

The algorithm goes as follows: Select the cube $Q_{2^{-k}}(x_j^k)$ if none of its predecessors have been previously chosen and the paraboloid $P_{x_j^k, t_i^k}^{2^k\alpha, \alpha}$ touches the super-solution over $\{u > 2^{-k}m\}$, such that Corollary 3.4 can be applied. In order to get the inductive hypothesis whenever the cube fails to be chosen, we need to fix the constants α, m, τ, R conveniently.

Let us carefully review the geometric configuration over $Q_{2^{-k}}(x_j^k) \times (t_i^k, t_f^k]$. First we take a system of coordinates such that we identify the cylinder $Q_{2^{-k}}(x_j^k) \times (t_i^k, t_f^k]$ with $Q_1 \times (0, 2\tau + 2]$.

This is the reason why Corollary 3.4 is given over the time interval $(0, 2^{-k}(2\tau + 2)]$ instead of $(0, 2^{-k}]$.

1. Inductive hypothesis: Under the contact assumption

$$A^\alpha(0, 0) \cap \mathbb{R}^n \times (0, 2\tau + 2] \neq \emptyset,$$

the alternative hypothesis to the ABP-type Corollary 3.4, which determines whenever we should proceed to the next generation of sub-cubes, is

$$(3.3) \quad \emptyset \neq A^\alpha(0, 0) \cap \mathbb{R}^n \times (0, 2\tau + 2] \subseteq \{u \leq m\}.$$

To continue the iteration we need to have that for every multi-index $j \in \{\pm 1\}^n$

$$(3.4) \quad A^{2\alpha, \alpha}(x_j^1, -\tau) \cap \mathbb{R}^n \times (-\tau, 1] \neq \emptyset.$$

Our goal at this point is to justify our choices for the constants so that we can guarantee this statement.

2. Localization via the eikonal equation (fixing α and m): So far we have not used the eikonal mechanism of the equation

$$\partial_t u \geq u\mathcal{M}^-(D^2u) + \lambda|\mathbf{D}u|^2.$$

Under the hypothesis (3.3) we will now take advantage of the finite speed of propagation to localize the contact set. Indeed, if we take $\alpha = 100 \max(1, 1/\lambda)$ and $m = \min(1, 1/(n\Lambda))$ then by evaluating the super-solution equation on the paraboloid P^α for a point $(x_0, t_0) \in A^\alpha(0, 0) \subseteq \{u \leq m\}$

$$\alpha \geq -m\Lambda n\alpha + \lambda\alpha^2|x_0|^2 \quad \Rightarrow \quad |x_0| \leq 1/4.$$

Also after time $t > 1$, $\alpha \geq 2$ implies that $P^\alpha(\cdot, t) > 1$ in $B_{1/4}$ so that $t_0 \in (0, 1]$. In other words, by appropriately choosing α and m we now see that (3.3) implies the following localization of the contact set

$$(3.5) \quad \emptyset \neq A^\alpha(0, 0) \cap \mathbb{R}^n \times (0, 2\tau + 2] \subseteq \{u \leq m\} \cap B_{1/4} \times (0, 1]$$

3. Fixing τ and R : We use (3.5) to finally get (3.4) for any multi-index.

The contact would follow if $P_{x_j^1, -\tau}^{2\alpha, \alpha} > m$ in $B_{1/4} \times \{0\}$, for which it suffices to fix $\tau = n$. On the other hand, we have already discussed that this contact happens strictly inside the domain if we take $R = 5\sqrt{n} \geq 2\sqrt{\tau + 1} + 1$.

4. Output of the algorithm: As a result we get a family of selected dyadic cubes in each scale which we can identify by its centers $\mathcal{C}_k \subseteq \{x_i^k\}$.

This selection leads to at least one of the following two alternatives: Either

$$(3.6) \quad \left| \bigcup_{\substack{k \geq 0, \\ x_i^k \in \mathcal{C}_k}} Q_{2^{-k}}(x_i^k) \right| > 1/2,$$

or by the continuity of u

$$|\{u(\cdot, -2n) = 0\} \cap Q_1| \geq 1/2.$$

The second case already gives us a subset of $Q_1 \times \{-2n\}$ with positive measure in space where the super-solution is bounded (actually vanishes).




FIGURE 3. An elementary geometric fact.

3.3.2. *Fixed time measure estimate.* In the scenario given by (3.6), what we can conclude is that there is a subset of B_R with positive measure (in space), such that for each of its points the following maximal density (measured over time intervals) is universally bounded away from zero

$$\Theta_M(x) := \sup_{t \in [-2n, 1]} \frac{|\{u(x, \cdot) \leq M\} \cap [-2n, t]|}{|[-2n, t]|}.$$

Notice that as $t \rightarrow -2n$, a lower bound on Θ_M only recovers information about u being bounded at time $t = -2n$.

Corollary 3.5 (Fixed time measure estimate). *Given $[\lambda, \Lambda] \subseteq (0, \infty)$ there exist $M, \eta, c > 0$ such that if $u \in C(B_{5\sqrt{n}} \times [-2n, 1])$ is a non-negative viscosity super-solution*

$$\partial_t u \geq u\mathcal{M}^-(D^2u) + \lambda|Du|^2 \text{ in } B_{5\sqrt{n}} \times (-2n, 1].$$

then $A^\alpha(0, 0) \neq \emptyset$ implies at least one of the following alternatives

$$|\{u(\cdot, -2n) = 0\} \cap Q_1| \geq 1/2 \quad \text{or} \quad |\{\Theta_M \geq \eta\} \cap B_{5\sqrt{n}}| \geq c$$

Proof. Assume that the first alternative does not hold such that (3.6) has to be true instead.

We pass from the cubes to the balls where we have the density estimates

$$\left| \bigcup_{\substack{k \geq 0, \\ x_i^k \in \mathcal{C}_k}} B_{2^{-k}R}(x_j^k) \right| \geq \left| \bigcup_{\substack{k \geq 0, \\ x_i^k \in \mathcal{C}_k}} Q_{2^{-k}}(x_j^k) \right| > 1/2.$$

By replacing η by an even smaller fraction we can rephrase the conclusion of Corollary 3.4 for each $x_i^k \in \mathcal{C}_k$ as

$$|\{u \leq M\} \cap B_{2^{-k}R}(x_j^k) \times (-2n, t_f^k]| \geq 2\eta|B_{2^{-k}R}(x_j^k) \times (-2n, t_f^k]|$$

such that,

$$|\{\theta_k > \eta\} \cap B_{2^{-k}R}(x_j^k)| \geq \eta|B_{2^{-k}R}(x_j^k)|, \quad \theta_k(x) := \frac{|\{u(x, \cdot) \leq M\} \cap (-2n, t_f^k]|}{|(-2n, t_f^k]|}.$$

See Figure 3.

Even though the balls in this union are not disjoint, we can use Vitali's Covering Lemma to extract a disjoint sub-covering, still with measure bounded away from zero by some constant depending only on the dimension. With this we now conclude the estimate. \square

3.4. Integration in time. As a final step to get the measure estimate Lemma 3.1, we need to apply Corollary 3.5 in a whole interval of times and integrate from there a set of positive measure in *space and time* where the super-solution is bounded.

Corollary 3.6. *Given $[\lambda, \Lambda] \subseteq (0, \infty)$ there exists $M, c > 0$ such that if $u \in C(B_{5\sqrt{n}} \times [-2n, 1])$ is a non-negative viscosity super-solution*

$$\partial_t u \geq u\mathcal{M}^-(D^2u) + \lambda|Du|^2 \text{ in } B_{5\sqrt{n}} \times (-2n, 1].$$

then $A^\alpha(0, t) \neq \emptyset$ for each $t \in [0, 1/2]$ implies that

$$|\{u \leq M\} \cap B_{5\sqrt{n}} \times [-2n, 1]| \geq c$$

Proof. By applying Corollary 3.5 to $u(\cdot, \cdot - t)$ we get that one of the following alternatives hold for each $t \in [0, 1/2]$

$$|\{u(\cdot, -2n + t) = 0\} \cap Q_1| \geq 1/2 \quad \text{or} \quad |\{\Theta_M(\cdot, -2n + t) \geq \eta\} \cap B_{5\sqrt{n}}| \geq c$$

where

$$\Theta_M(x, t) := \sup_{s \in [t, 1]} \frac{|\{u(x, \cdot) \leq M\} \cap [t, s]|}{|[t, s]|}.$$

If the set of $t \in [0, 1/2]$ where the first alternative holds has length at least $1/4$ we conclude by saying that u actually vanishes in a set of measure at least $1/8$ inside $B_{5\sqrt{n}} \times [-2n, 1]$. So let us consider instead that the set of $t \in [0, 1/2]$ where the second alternative holds has length at least $1/4$. In other words, for some c perhaps even smaller

$$|\{\Theta_M \geq \eta\} \cap B_{5\sqrt{n}} \times [-2n, -2n + 1/2]| \geq 2c|B_{5\sqrt{n}} \times [-2n, -2n + 1/2]|.$$

such that

$$|\{\theta > c\} \cap B_{5\sqrt{n}}| \geq c|B_{5\sqrt{n}}|, \quad \theta(x) := \frac{|\{\Theta_M(x, \cdot) \geq \eta\} \cap [-2n, -2n + 1/2]|}{|[-2n, -2n + 1/2]|}.$$

Lastly, let us fix $x \in \{\theta > c\} \cap B_{5\sqrt{n}}$ and invoke Vitali's Covering Lemma to show that

$$|\{u(x, \cdot) \leq M\} \cap [-2n, 1]| \geq c\eta/20.$$

Indeed, there is a covering of $\{\Theta_M(x, \cdot) \geq \eta\} \cap [-2n, -2n + 1/2]$ (a one dimensional set with length $> c/2$) with intervals that intersect $\{u(x, \cdot) \leq M\}$ in a positive fraction. Namely by the definition of Θ_M , for every $t \in \{\Theta_M(x, \cdot) \geq \eta\} \cap [-2n, -2n + 1/2]$ there is some $s = s(t)$ such that

$$\frac{|\{u(x, \cdot) \leq M\} \cap [t, s]|}{|[t, s]|} \geq \eta/2.$$

By Vitali's we can assume that the covering is disjoint and still has its measure bounded from below by $c/10$.

Finally we integrate this information for every $x \in \{\theta > c\} \cap B_{5\sqrt{n}}$ to get the desired estimate in space and time. \square

4. IMPROVEMENT FROM ABOVE

As a last step towards the proof of Theorem 2.2 we need to establish the improvement of the upper bound provided a weak control in measure for a sub-solution.

Proof of Lemma 2.4. It can be directly checked from the definition by testing with smooth super-solutions, that the truncation $w := \max(1/2, u)$ is a viscosity sub-solution to

$$\partial_t w \leq \bar{\mathcal{M}}^+(D^2 w) + \Lambda |Dw|^2 \quad \text{in } B_1 \times (-1, 0] \quad (\bar{\mathcal{M}}^+ = \mathcal{M}^+(\lambda/2, \Lambda)).$$

Then we consider $v := e^{Aw}$ such that, again from the definition of sub-solution by applying the same transformation to the test functions,

$$\frac{\partial_t v}{Av} \leq \bar{\mathcal{M}}^+ \left(\frac{D^2 v}{Av} - \frac{Dv \otimes Dv}{Av^2} \right) + \frac{\Lambda |Dv|^2}{A^2 v^2} \quad \text{in } B_1 \times (-1, 0].$$

By taking A sufficiently large in terms of the ellipticity constants, we get to cancel the gradient terms in order to recover the heat equation

$$\partial_t v \leq \bar{\mathcal{M}}^+(D^2 v) \quad \text{in } B_1 \times (-1, 0].$$

The result now follow by applying the weak Harnack's inequality on $(e^A - v)$. □

5. CONCLUSION

At this point the proof of our main result, namely Theorem 2.2, is completed. Let us recapitulate:

1. In order to establish an interior Hölder estimates it suffices to establish the modulus of continuity at a generic interior point and then use the translation and scale invariance of the equation (Remark 2.1).

2. For the diminish of oscillation we consider a solution u over the domain $B_1 \times (-1, 0]$ and taking values in $[0, 1]$. For the fraction $\eta \in (0, 1)$ determined in Lemma 2.3 we get the following alternative scenarios in measure:

Lemma 2.3 states that if $\{u > 1/2\}$ covers at least a fraction $(1 - \eta)$ of the domain (most of it), then we show that the (zero) lower bound can be raised in a smaller cylinder. If this ever happens the degeneracy disappears and the equation can be made uniformly elliptic after a change of variables as in the proof of Lemma 2.4 in Section 4.

Lemma 2.4 states that if instead $\{u \leq 1/2\}$ covers at least a fraction η of the domain (some of it), then we show that the upper bound can be lowered in a smaller cylinder. It may be the case that the solution vanishes at the center of the cylinder and then the oscillation always diminishes from above.

3. The most delicate analysis happens when we want to establish the improvement form below. Lemma 3.1 is the contrapositive version of Lemma 2.3 in a convenient geometric configuration for the ensuing computations. Instead of having the infimum of u over some set towards the future, we have a punctual control, however this implies the statement for the infimum after using a standard covering argument.

4. We use a dyadic selection algorithm based on the ABP principle. Each dyadic cube is associated with a concave paraboloid to test the super-solution from below. If it happens to

touch over the uniformly elliptic regime, then we collect a set of positive measure in space and time where the solution is bounded. Otherwise the contact has to happen over the set where u is small and the eikonal mechanism prevents the horizontal spread of these points. Corollary 3.5 exhibits the output of this procedure which, in the worst case, states that u is bounded at some fixed time in a set of positive measure (but only in space).

5. To finish the proof of Lemma 3.1 we need to integrate the sets obtained by Corollary 3.5 over a range of times. This idea is made precise in Corollary 3.6 by using Vitali's Covering Lemma one more time.

6. Once the fraction $\eta \in (0, 1)$ has been fixed by Lemma 2.3, the alternative case given by Lemma 2.4 is proven (it actually holds for any fraction). Given that we are dealing with a sub-solution in this case, truncation from below at a positive level set preserves the equation and actually free us from the degeneracy. A classical change of variables allow us to transform the equation into the heat equation with bounded measurable coefficients for which the weak Harnack inequality give us the desired improvement.

5.1. Further directions. Keeping in mind the interest established in [2] on equations of the form $\partial_t u = a(u)\Delta u + |Du|^2$ with a sublinear as $u \rightarrow 0^+$, we considered extending our result to $\partial_t u = u^\alpha \Delta u + |Du|^\beta$. The scaling of this equation seems quite restrictive. For instance, if our goal is to show a diminish of oscillation of order γ in space we find the constrain $(\alpha + 1)\gamma - 2 = \beta\gamma - \beta$ on γ , unless $\alpha + 1 = \beta = 2$ which is our original case. Otherwise, we need to find an strategy to show directly a γ -Hölder modulus of continuity (in space) with $\gamma = (2 - \beta)/(\alpha + 1 - \beta)$, which is not necessarily a small exponent. For $\alpha > 1$ (the case we can relate with [2]) we get that $\gamma \rightarrow 0^+$ if $\beta \rightarrow 2^-$, perhaps there is an opportunity to show a result in such range by a perturbative approach.

The two phase PME is also a well understood problem from the variational point of view since the eighties [5, 32, 14, 15, 27]. However, as far as we know, the viscosity solution approach for this particular type of two-phase free boundary problem has not been developed. Expanding the variational equation $\partial_t u = \operatorname{div}(|u|Du)$, we see that the non-variational form should say instead the following: Each phase $u_\pm = \max(\pm u, 0)$ satisfies the un-signed PME outside of the free boundary points connecting the two phases

$$\partial_t u = u\Delta u + |Du|^2 \text{ in } (\operatorname{spt} u_\pm \setminus \operatorname{spt} u_\mp) \cap \Omega \times (t_i, t_f]$$

and at the two-phase free boundary points we get

$$\frac{\partial_t u}{|Du_+|} = -\frac{\partial_t u}{|Du_-|} = |Du_+| - |Du_-| \text{ in } \operatorname{spt} u_\pm \cap \operatorname{spt} u_\mp \cap \Omega \times (t_i, t_f]$$

Besides the challenges that the well-posedness may present, we believe that our treatment for the regularity theory requires some substantial modifications. For instance, the ABP Lemma 3.3 uses as test functions paraboloids that start growing from the zero level set. We could consider paraboloids that start from smaller level sets, however it is not clear how to use the eikonal equation to prevent the horizontal spread of the contact set whenever the contact happens at $\{u \in (-m, 0]\}$.

An interesting problem arises from models with non-local interactions. Let us recall that the continuity equation $\partial_t u - \operatorname{div}(uDp) = 0$ models the evolution of the density u driven by the pressure p . The case $p = -(-\Delta)^{-\sigma}u$ with $\sigma \in (0, 1)$ gives the fractional PME proposed

by Caffarelli and Vázquez in [11]. Meanwhile the regularity of the solution was established by the de Giorgi method in [9], the regularity of the free boundary remains largely open. The difficulty resides on the presence of the non-local drift term $Dp \cdot Du$ which prevents the comparison principle. Preliminary computations show that our technique could be extended to the equation

$$\partial_t u = -u(-\Delta)^{1-\sigma} u + |Du|^2$$

where the dangerous presence of the non-local drift is replaced by the classical and local one, with the trade off of destroying the variational structure of the problem.

The regularity of the free boundary for evolution problems has a well developed non-variational approach, the book by Caffarelli and Salsa [8] is our recommended source. For the PME a very elegant improvement of flatness strategy was recently implemented by Kienzler, Koch and Vázquez in [19]. The regularity of a broader family of PME type equations may now open the possibility of extending this higher regularity for the free boundary to the corresponding non-variational class.

REFERENCES

- [1] Donald G. Aronson and Philippe Bénilan. Régularité des solutions de l'équation des milieux poreux dans \mathbf{R}^N . *C. R. Acad. Sci. Paris Sér. A-B*, 288(2):A103–A105, 1979.
- [2] Cristina Brändle and Juan Luis Vázquez. Viscosity solutions for quasilinear degenerate parabolic equations of porous medium type. *Indiana Univ. Math. J.*, 54(3):817–860, 2005.
- [3] Xavier Cabré. Nondivergent elliptic equations on manifolds with nonnegative curvature. *Comm. Pure Appl. Math.*, 50(7):623–665, 1997.
- [4] Luis Caffarelli and Xavier Cabré. *Fully nonlinear elliptic equations*, volume 43 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1995.
- [5] Luis Caffarelli and L. Craig Evans. Continuity of the temperature in the two-phase Stefan problem. *Arch. Rational Mech. Anal.*, 81(3):199–220, 1983.
- [6] Luis Caffarelli and Avner Friedman. Continuity of the density of a gas flow in a porous medium. *Trans. Amer. Math. Soc.*, 252:99–113, 1979.
- [7] Luis Caffarelli and Avner Friedman. Regularity of the free boundary of a gas flow in an n -dimensional porous medium. *Indiana Univ. Math. J.*, 29(3):361–391, 1980.
- [8] Luis Caffarelli and Sandro Salsa. *A geometric approach to free boundary problems*, volume 68 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [9] Luis Caffarelli, Fernando Soria, and Juan Luis Vázquez. Regularity of solutions of the fractional porous medium flow. *J. Eur. Math. Soc. (JEMS)*, 15(5):1701–1746, 2013.
- [10] Luis Caffarelli and Juan Luis Vázquez. Viscosity solutions for the porous medium equation. In *Differential equations: La Pietra 1996 (Florence)*, volume 65 of *Proc. Sympos. Pure Math.*, pages 13–26. Amer. Math. Soc., Providence, RI, 1999.
- [11] Luis Caffarelli and Juan Luis Vázquez. Nonlinear porous medium flow with fractional potential pressure. *Arch. Ration. Mech. Anal.*, 202(2):537–565, 2011.
- [12] Michael G. Crandall, Hitoshi Ishii, and Pierre-Louis Lions. User's guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [13] P. Daskalopoulos and R. Hamilton. Regularity of the free boundary for the porous medium equation. *J. Amer. Math. Soc.*, 11(4):899–965, 1998.
- [14] Emmanuele DiBenedetto. Continuity of weak solutions to certain singular parabolic equations. *Ann. Mat. Pura Appl. (4)*, 130:131–176, 1982.
- [15] Emmanuele DiBenedetto. Continuity of weak solutions to a general porous medium equation. *Indiana Univ. Math. J.*, 32(1):83–118, 1983.
- [16] Emmanuele DiBenedetto, José Miguel Urbano, and Vincenzo Vespri. Current issues on singular and degenerate evolution equations. In *Evolutionary equations. Vol. I*, Handb. Differ. Equ., pages 169–286. North-Holland, Amsterdam, 2004.

- [17] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [18] Morton E. Gurtin and Richard C. MacCamy. On the diffusion of biological populations. *Math. Biosci.*, 33(1-2):35–49, 1977.
- [19] Clemens Kienzler, Herbert Koch, and Juan Luis Vázquez. Flatness implies smoothness for solutions of the porous medium equation. *Calculus of Variations and Partial Differential Equations*, 57(1):18, 2018.
- [20] Inwon C. Kim and Helen K. Lei. Degenerate diffusion with a drift potential: a viscosity solutions approach. *Discrete Contin. Dyn. Syst.*, 27(2):767–786, 2010.
- [21] Inwon C. Kim and Norbert Požár. Viscosity solutions for the two-phase Stefan problem. *Comm. Partial Differential Equations*, 36(1):42–66, 2011.
- [22] Inwon C. Kim and Norbert Požár. Nonlinear elliptic-parabolic problems. *Arch. Ration. Mech. Anal.*, 210(3):975–1020, 2013.
- [23] Nikolai V. Krylov and Mikhail V. Safonov. An estimate for the probability of a diffusion process hitting a set of positive measure. *Dokl. Akad. Nauk SSSR*, 245(1):18–20, 1979.
- [24] Nikolai V. Krylov and Mikhail V. Safonov. A property of the solutions of parabolic equations with measurable coefficients. *Izv. Akad. Nauk SSSR Ser. Mat.*, 44(1):161–175, 239, 1980.
- [25] Connor Mooney. Harnack inequality for degenerate and singular elliptic equations with unbounded drift. *J. Differential Equations*, 258(5):1577–1591, 2015.
- [26] Víctor Padrón. Effect of aggregation on population recovery modeled by a forward-backward pseudoparabolic equation. *Trans. Amer. Math. Soc.*, 356(7):2739–2756, 2004.
- [27] Paul E. Sacks. Continuity of solutions of a singular parabolic equation. *Nonlinear Anal.*, 7(4):387–409, 1983.
- [28] José Miguel Urbano. *The method of intrinsic scaling*, volume 1930 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2008.
- [29] Juan Luis Vázquez. *The porous medium equation*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2007.
- [30] Yu Wang. Small perturbation solutions for parabolic equations. *Indiana Univ. Math. J.*, 62(2):671–697, 2013.
- [31] Lang-Fang Wu. A new result for the porous medium equation derived from the Ricci flow. *Bull. Amer. Math. Soc. (N.S.)*, 28(1):90–94, 1993.
- [32] William P. Ziemer. Interior and boundary continuity of weak solutions of degenerate parabolic equations. *Trans. Amer. Math. Soc.*, 271(2):733–748, 1982.

DEPARTMENT OF MATHEMATICS, CIMAT, GUANAJUATO, MEXICO

Email address: hector.chang@cimat.mx

DEPARTMENT OF MATHEMATICS, CIMAT, GUANAJUATO, MEXICO

Email address: makson.santos@cimat.mx