

Arcsine and Darling–Kac laws for piecewise linear random interval maps

Genji HATA⁽¹⁾ and Kouji YANO⁽²⁾⁽³⁾

Abstract

We give examples of piecewise linear random interval maps satisfying arcsine and Darling–Kac laws, which are analogous to Thaler’s arcsine and Aaronson’s Darling–Kac laws for the Boole transform. They are constructed by random switch of two piecewise linear maps with attracting or repelling fixed points, which behave as if they were indifferent fixed points of a deterministic map.

Keywords and phrases: arcsine law; Darling–Kac law; random dynamical system; infinite ergodic theory; Markov partition

AMS 2010 subject classifications: 37A50 (37H12 60F05)

1 Introduction

Let \mathcal{A} and \mathcal{N} denote random variables with arcsine and standard normal distributions, respectively:

$$P(\mathcal{A} \in du) = \frac{du}{\pi\sqrt{u(1-u)}} \quad (0 < u < 1), \quad P(\mathcal{N} \in du) = e^{-u^2/2} \frac{du}{\sqrt{2\pi}} \quad (u \in \mathbb{R}). \quad (1.1)$$

It is well-known that a simple symmetric random walk $\{W_n\}_{n=0}^\infty$ on \mathbb{Z} satisfies *Lévy’s arcsine law* and *Darling–Kac law*, respectively:

$$\frac{1}{N} \sum_{n=0}^{N-1} 1_{\{W_n > 0\}} \xrightarrow[N \rightarrow \infty]{d} \mathcal{A} \quad \text{and} \quad \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} 1_{\{W_n \in E\}} \xrightarrow[N \rightarrow \infty]{d} \frac{\#E}{\sqrt{\pi}} |\mathcal{N}| \quad (1.2)$$

for all bounded set $E \subset \mathbb{Z}$, where $\#E$ denotes the number of elements of E and \xrightarrow{d} denotes the convergence in distribution. Note that the uniform measure $\mu = \sum_{k \in \mathbb{Z}} \delta_k$ on \mathbb{Z} is a unique σ -finite invariant measure for the random walk $\{W_n\}$:

$$\mathbb{E}\mu(\cdot + W_1) = \mu(\cdot). \quad (1.3)$$

Here and hereafter by uniqueness of an invariant measure we mean uniqueness up to constant multiplicity.

⁽¹⁾SAPIX

⁽²⁾Graduate School of Science, Kyoto University.

⁽³⁾The research of Kouji Yano was supported by JSPS KAKENHI grant no.’s JP19H01791 and JP19K21834 and by JSPS Open Partnership Joint Research Projects grant no. JPJSBP120209921. This research was supported by RIMS and by ISM.

Thaler [35] and Aaronson [1] obtained analogous results for the *Boole transform* $S : \mathbb{R} \rightarrow \mathbb{R}$ defined as $S(x) = x - 1/x$. We know that the Lebesgue measure $\lambda(dx) = dx$ on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$, where $\mathcal{B}(\cdot)$ denotes the family of all Borel sets, is S -invariant:

$$\lambda \circ S^{-1} = \lambda. \quad (1.4)$$

We also know that a λ -a.c. σ -finite S -invariant measure is unique. Thaler's arcsine and Aaronson's Darling–Kac laws can be stated as follows, respectively: For any random initial point Θ in \mathbb{R} with λ -a.c. density, it holds that

$$\frac{1}{N} \sum_{n=0}^{N-1} 1_{\{S^n(\Theta) > 0\}} \xrightarrow[N \rightarrow \infty]{d} \mathcal{A} \quad \text{and} \quad \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} 1_{\{S^n(\Theta) \in E\}} \xrightarrow[N \rightarrow \infty]{d} \frac{\lambda(E)}{\sqrt{\pi}} |\mathcal{N}| \quad (1.5)$$

for all Borel set E with $\lambda(E) < \infty$.

For our purpose we change variables by a diffeomorphism $\Phi : (0, 1) \rightarrow \mathbb{R}$ which is defined as $\Phi(x) = \frac{2x-1}{x(1-x)}$. The Boole transform is changed into $T = \Phi^{-1} \circ S \circ \Phi : [0, 1] \rightarrow [0, 1]$ given as

$$T(x) = \frac{x(1-x)}{1-x-x^2} \quad (0 < x < 1/2), \quad T(x) = 1 - T(1-x) \quad (1/2 < x < 1). \quad (1.6)$$

For T the points 0 and 1 are *indifferent fixed points*:

$$[T(0+) = 0, T'(0+) = 1] \quad \text{and} \quad [T(1-) = 1, T'(1-) = 1]. \quad (1.7)$$

The unique λ -a.c. σ -finite T -invariant measure is given as

$$\mu(dx) = \Phi'(x)dx = \left(\frac{1}{x^2} + \frac{1}{(1-x)^2} \right) dx \quad \text{on } [0, 1]. \quad (1.8)$$

We note that μ has infinite mass near 0 and 1 and has finite mass away from 0 and 1. This fact shows that the orbit of T stays most of time near the indifferent fixed points 0 and 1. Thaler's arcsine and Aaronson's Darling–Kac laws can be restated as follows: For any random initial point Θ in $[0, 1]$ with λ -a.c. density, it holds that

$$\frac{1}{N} \sum_{n=0}^{N-1} 1_{\{T^n(\Theta) > 1/2\}} \xrightarrow[N \rightarrow \infty]{d} \mathcal{A} \quad \text{and} \quad \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} 1_{\{T^n(\Theta) \in E\}} \xrightarrow[N \rightarrow \infty]{d} \frac{\mu(E)}{\sqrt{\pi}} |\mathcal{N}| \quad (1.9)$$

for all Borel set E with $\mu(E) < \infty$.

Random maps

For two deterministic interval maps $\tau_1, \tau_2 : [0, 1] \rightarrow [0, 1]$ and a constant $0 < p < 1$, we consider the random map

$$T = \begin{cases} \tau_1 & \text{(with probability } p), \\ \tau_2 & \text{(with probability } 1 - p). \end{cases} \quad (1.10)$$

A measure μ on $[0, 1]$ is called T -invariant if μ is not a zero measure and

$$(\mathbb{E}\mu \circ T^{-1} =) p\mu \circ \tau_1^{-1} + (1 - p)\mu \circ \tau_2^{-1} = \mu. \quad (1.11)$$

Let $\{T_n\}_{n=1}^\infty$ be an i.i.d. sequence of random maps with $T_1 \stackrel{d}{=} T$ and we define

$$T^{(n)} = T_n \circ T_{n-1} \circ \cdots \circ T_1. \quad (1.12)$$

The resulting random map $T^{(n)}$ can be regarded as the n -fold composition of the random map T . Our aim is to obtain arcsine and Darling–Kac laws for random maps which are analogous to those for the Boole transform.

Main results: Hata map

Based on Hata [17], we adopt the two deterministic maps

$$\tau_1(x) = \begin{cases} x/2 & (0 < x < 1/2) \\ 2x - 1 & (1/2 < x < 1) \end{cases}, \quad \tau_2(x) = \begin{cases} 2x & (0 < x < 1/2) \\ (x + 1)/2 & (1/2 < x < 1) \end{cases}. \quad (1.13)$$

Note that 0 for τ_1 and 1 for τ_2 are attracting fixed points:

$$\tau_1^n(x) \xrightarrow[n \rightarrow \infty]{} 0, \quad \tau_2^n(x) \xrightarrow[n \rightarrow \infty]{} 1 \quad \text{for } 0 < x < 1 \text{ except for } 1/2, \quad (1.14)$$




Figure 1: $\tau_1(x)$




Figure 2: $\tau_2(x)$

in the sense that

$$[T(0+) = 0, \mathbb{E} \log |T'(0+)| = 0] \quad \text{and} \quad [T(1-) = 1, \mathbb{E} \log |T'(1-)| = 0]. \quad (1.15)$$

We expect that the attracting and repelling effects are balanced in this case.

We define the λ -partition α of $[0, 1]$ for the Hata map as $\alpha = \{I_k^-, I_k^+\}_{k=0}^\infty$ with

$$I_k^- = \left(\frac{1}{2^{k+2}}, \frac{1}{2^{k+1}} \right), \quad I_k^+ = \left(1 - \frac{1}{2^{k+1}}, 1 - \frac{1}{2^{k+2}} \right). \quad (1.16)$$

Here we say that α is a λ -partition of $[0, 1]$ if α is a disjoint family of Borel subsets of $[0, 1]$ such that $\sum_{s \in \alpha} s = [0, 1] \bmod \lambda$ and $0 < \lambda(s) < \infty$ for all $s \in \alpha$. Note that we sometimes write \sum instead of \cup for disjoint union.

Theorem 1.1. *Regarding the Hata map with $p = 1/2$, there exists a unique λ -a.c. σ -finite T -invariant measure. It is given as*

$$\mu(dx) = \sum_{s \in \alpha} \frac{\rho(s)}{\lambda(s)} 1_s(x) dx \quad (1.17)$$

with

$$\rho(I_k^-) = \rho(I_k^+) = (2^{k+1} - 1) \cdot 2^{-k-2}, \quad k = 0, 1, 2, \dots \quad (1.18)$$

Consequently, μ has infinite mass near 0 and 1 and has finite mass away from 0 and 1.

Theorem 1.1 will be slightly generalized and proved in Theorem 5.1.

This result shows that the indifferent-like fixed points 0 and 1 for the random map T behave as if they were indifferent fixed points of a deterministic map. We now state our arcsine and Darling–Kac laws.

Theorem 1.2. *Regarding the Hata map with $p = 1/2$, for any random initial point Θ in $[0, 1]$ with λ -a.c. density which is independent of the random maps $\{T_n\}$, it holds that*

$$\frac{1}{N} \sum_{n=0}^{N-1} 1_{\{T^{(n)}(\Theta) > 1/2\}} \xrightarrow[N \rightarrow \infty]{d} \mathcal{A}, \quad \text{and} \quad \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} 1_{\{T^{(n)}(\Theta) \in E\}} \xrightarrow[N \rightarrow \infty]{d} \frac{\mu(E)}{\sqrt{\pi}} |\mathcal{N}| \quad (1.19)$$

for all Borel set E with $\mu(E) < \infty$, where by \xrightarrow{d} we mean the convergence in distribution on the extended probability space.

The proof of Theorem 1.2 will be divided into the following steps:

- (i) We show irreducible recurrence of the Markov chain on the λ -partition.
- (ii) We modify the λ -partition to be a Markov partition for the skew-product.

- (iii) We verify conjugacy between the random dynamical system and the Markov chain.
- (iv) We study the wandering rate asymptotics and check that the assumptions of Thaler–Zweimüller’s theorems [36] are satisfied.

Lévy’s arcsine and the Darling–Kac laws have been extended for dynamical systems by Thaler [35] and by Aaronson [1], respectively. Their results were generalized in a common framework by Thaler–Zweimüller [36] and Zweimüller [38]. Aaronson’s Darling–Kac law was generalized to convergence on a functional space by Aaronson [2] and Owada–Samorodnitsky [27]. In the case of several indifferent fixed points, a joint-distributional generalization of Thaler’s arcsine law was obtained by Sera–Yano [33]. Recently Sera [32] obtained a functional and joint-distributional generalization of Thaler’s arcsine and Aaronson’s Darling–Kac laws.

Pelikan and mBGI maps

Among others, we would like to focus on the two random maps with infinite invariant measures obtained by Pelikan [28] and by Boyarsky–Góra–Islam [10]. For other results on random maps with infinite T -invariant measures, see Bahsoun–Bose–Duan [8, 9], Bahsoun–Bose [7], Gharaei–Homburg [16], Abbasi–Gharaei–Homburg [4], Inoue [20], Toyokawa [37] and Homburg–Kalle–Ruziboev–Verbitskiy–Zeegers [18].

First, we take up

$$\tau_1(x) = 2x \bmod 1, \quad \tau_2(x) = x/2. \tag{1.20}$$

and we call the corresponding random map T the *Pelikan map*. Note that both 0 and 1 for τ_1 are repelling fixed points and 0 for τ_2 is an attracting fixed point. Although Pelikan [28] excluded the case $p = 1/2$, we choose $p = 1/2$ so that 0 is an indifferent-like fixed point in the sense that

$$[T(0+) = 0, \mathbb{E} \log |T'(0+)| = 0]. \tag{1.21}$$

The λ -partition for the Pelikan map is defined as $\alpha = \{I_k\}_{k=0}^\infty$ with

$$I_k = \left(\frac{1}{2^{k+1}}, \frac{1}{2^k} \right). \tag{1.22}$$

Theorem 1.3. *Regarding the Pelikan map with $p = 1/2$, there exists a unique λ -a.c. σ -finite T -invariant measure. It is given as $\mu(dx) = \sum_{s \in \alpha} \frac{\rho(s)}{\lambda(s)} 1_s(x) dx$ with*

$$\rho(I_k) = 2^{-2} \cdot (2^{k+1} - 1) \cdot 2^{-k-1}, \quad k = 0, 1, 2, \dots \tag{1.23}$$

Consequently, μ has infinite mass near 0 and has finite mass away from 0.

Theorem 1.3 will be slightly generalized and proved in Theorem 5.3. Note that Pelikan [28, Section 4] implicitly found the invariant measure, while uniqueness was not mentioned.

Although the arcsine law does not make sense, we obtain the Darling–Kac law as follows.

Theorem 1.4. *Regarding the Pelikan map with $p = 1/2$, for any random initial point Θ in $[0, 1]$ with λ -a.c. density which is independent of the random maps $\{T_n\}$, it holds that*

$$\frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} 1_{\{T^{(n)}(\Theta) \in E\}} \xrightarrow[N \rightarrow \infty]{d} \frac{\mu(E)}{\sqrt{\pi}} |\mathcal{N}| \quad (1.24)$$

for all Borel set E with $\mu(E) < \infty$.

The proof of Theorem 1.4 will be divided into the same steps as those of Theorem 1.2.

Second, we take up

$$\tau_1(x) = \begin{cases} 2x & (0 < x < 1/4) \\ 1 - 2x & (1/4 < x < 1/2) \\ 2 - 2x & (1/2 < x < 3/2) \\ 2x - 1 & (3/2 < x < 1) \end{cases}, \quad \tau_2(x) = \begin{cases} 1 - x/4 & (0 < x < 1/2) \\ (1 - x)/4 & (1/2 < x < 1) \end{cases}. \quad (1.25)$$

and we call the corresponding random map T the *modified Boyarsky–Góra–Islam map* or *mBGI map* in short. Note that both 0 and 1 are repelling fixed points for τ_1 and the two-point set $\{0, 1\}$ is an attractor for τ_2 . We choose $p = 2/3$ so that the two-point set $\{0, 1\}$ is an *indifferent-like attractor* in the sense that

$$T\{0, 1\} = \{0, 1\}, \quad \mathbb{E} \log |T'(0+)| = 0, \quad \mathbb{E} \log |T'(1-)| = 0. \quad (1.26)$$

We introduce the λ -partition α of $[0, 1]$ for the mBGI map as $\alpha = \{I_k^-, I_k^+\}_{k=0}^{\infty}$ with

$$I_k^- = \left(\frac{1}{2^{k+2}}, \frac{1}{2^{k+1}} \right), \quad I_k^+ = \left(1 - \frac{1}{2^{k+1}}, 1 - \frac{1}{2^{k+2}} \right). \quad (1.27)$$

Theorem 1.5. *Regarding the mBGI map with $p = 2/3$, there exists a unique λ -a.c. σ -finite T -invariant measure. It is given as $\mu(dx) = \sum_{s \in \alpha} \frac{\rho(s)}{\lambda(s)} 1_s(x) dx$ with*

$$\rho(I_k^-) = \rho(I_k^+) = \frac{3\sqrt{2}}{8} \left(\frac{8}{3} \cdot 2^k + \frac{1}{3} \cdot (-1)^k - 1 \right) \cdot 2^{-k-2}, \quad k = 0, 1, 2, \dots \quad (1.28)$$

Consequently, μ has infinite mass near 0 and 1 and has finite mass away from 0 and 1.

Theorem 1.5 will be slightly generalized and proved in Theorem 5.2.

Although we will see in Remark 6.5 that the arcsine law does not make sense, we obtain the Darling–Kac law as follows.

Theorem 1.6. *Regarding the mBGI map with $p = 2/3$, for any random initial point Θ in $[0, 1]$ with λ -a.c. density which is independent of the random maps $\{T_n\}$, it holds that*

$$\frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} 1_{\{T^{(n)}(\Theta) \in E\}} \xrightarrow[N \rightarrow \infty]{d} \frac{\mu(E)}{\sqrt{\pi}} |\mathcal{N}| \quad (1.29)$$

for all Borel set E with $\mu(E) < \infty$.

The proof of Theorem 1.6 will be divided into the same steps as those of Theorem 1.2.

Intermittency

The *intermittency* has been introduced in statistical physics by Pomeau–Manneville [30] and Manneville [23], which describes irregular transitions between the *laminar phase* and the *turbulent burst*. As mathematical models of intermittency, dynamical systems with indifferent fixed points have been studied by many researchers such as Takahashi [34], Collet–Ferrero [13], Collet–Galves–Schmitt [15], Collet–Galves [14], Mori [24], Campanino–Isola [11, 12], Liverani–Saussol–Vaienti [22], Pollicott–Yuri [29], Hu [19] and Munday–Knight [25]. The Boole and Hata maps may be regarded also as mathematical models of intermittency thanks to indifferent-like fixed point(s). For other studies of statistical physics about random maps, see Ashwin–Aston–Nicol [6], Akimoto–Aizawa [5] and Sato–Klages [31].

Organization

This paper is organized as follows. In Section 2 we review the general theory assuring that a dynamical system is conjugate to a graph shift and to a Markov chain. In Section 3 we discuss irreducible recurrence of the Markov chain on the λ -partition. In Section 4 we discuss skew-product representation and verify the Markov partition condition. In Section 5 we prove the conjugacy and show unique existence of a.c. invariant measure. In Section 6 we investigate the wandering rate asymptotics to complete the proofs of our limit theorems.

Acknowledgements

The authors would like to express our gratitude to Shafiqul Islam, Jon Aaronson and Tomoki Inoue for useful advice. The first author thanks Yushi Nakano, Fumihiko Nakamura, Hisayoshi Toyokawa and Toru Sera for discussions which helped improve the earlier versions of this paper. He also thanks Yuzuru Sato and Takuma Akimoto for drawing his attention to the papers [6], [5] and [31].

2 Conjugacy induced by a Markov partition

2.1 Conjugacy to a graph shift

Let us review the general theory which ensures that a dynamical system with a Markov partition is conjugate to a graph shift. We complement Chapter 4 of [3] with the proofs which were omitted there.

Let (X, λ, T) be a non-singular dynamical system, i.e. $(X, \mathcal{B}(X), \lambda)$ is a standard measure space and $[\lambda \circ T^{-1}(B) = 0 \text{ whenever } \lambda(B) = 0]$. Note that, for any measure μ on $\mathcal{B}(X)$ which is equivalent to λ , the system (X, μ, T) is again a non-singular dynamical system.

We call α a λ -*partition* if α is a finite or countable disjoint family of Borel sets of X such that

$$[\sum_{s \in \alpha} s = X \text{ mod } \lambda] \text{ and } [0 < \lambda(s) < \infty \text{ for all } s \in \alpha]. \quad (2.1)$$

A λ -partition α is called a *Markov partition* if the following conditions are satisfied:

- (M1) The σ -field $\mathcal{G} := \sigma(T^{-n}s : s \in \alpha, n \geq 0)$ coincides with $\mathcal{B}(X) \text{ mod } \lambda$;
- (M2) The map $T : s \mapsto Ts$ is invertible λ -a.e. for all $s \in \alpha$;
- (M3) The forward image Ts belongs to $\sigma(\alpha) \text{ mod } \lambda$ for all $s \in \alpha$.

We denote the space of sequences of α by

$$Y := \{\mathbf{s} = (s_0, s_1, s_2, \dots) : s_0, s_1, s_2, \dots \in \alpha\}. \quad (2.2)$$

Although each element of the λ -partition α is a subset of X , we regard it as a point when we consider the space Y . Let θ denote the shift operator of Y , that is,

$$\theta : Y \ni (s_0, s_1, s_2, \dots) \mapsto (s_1, s_2, \dots) \in Y. \quad (2.3)$$

A *cylinder* of Y is given of the form

$$[s_0, s_1, \dots, s_n] = \{(s_0, s_1, \dots, s_n, t_{n+1}, t_{n+2}, \dots) \in Y : t_{n+1}, t_{n+2}, \dots \in \alpha\} \quad (2.4)$$

and we equip Y with the product topology, so that the Borel field $\mathcal{B}(Y)$ is generated by the family of cylinders.

We introduce the *graph shift* by

$$Y_\lambda := \{\mathbf{s} \in Y : \lambda(s_0 \cap T^{-1}s_1) > 0, \lambda(s_1 \cap T^{-1}s_2) > 0, \dots\}. \quad (2.5)$$

We note that $\mathcal{B}(Y_\lambda)$ is generated by the family of cylinders which are contained in Y_λ . We define a surjection $\phi : X \rightarrow Y_\lambda$ by

$$\phi(x) = \mathbf{s} \text{ with } x \in s_0, T(x) \in s_1, T^2(x) \in s_2, \dots \quad (2.6)$$

This map ϕ is Borel measurable since

$$\phi^{-1}[s_0, s_1, \dots, s_n] = s_0 \cap T^{-1}s_1 \cap \dots \cap T^{-n}s_n \in \mathcal{B}(X). \quad (2.7)$$

The following theorem can be found in [3, Proposition 4.2.3], although the proof was omitted there.

Theorem 2.1. *Let μ be a measure on X which is equivalent to λ . Then, with $\nu = \mu \circ \phi^{-1}$, the Borel surjection $\phi : (X, \mu, T) \rightarrow (Y_\lambda, \nu, \theta)$ is a conjugacy, i.e. the following hold:*

(i) $\phi \circ T = \theta \circ \phi$;

(ii) ϕ is essentially a Borel isomorphism.

Proof. Let us prove $\phi^{-1}\mathcal{B}(Y_\lambda) = \mathcal{B}(X) \bmod \mu$. It is obvious that $\phi^{-1}\mathcal{B}(Y_\lambda) \subset \mathcal{B}(X)$. For the converse inclusion, we take arbitrary $s_0, s_1, \dots, s_n \in \alpha$ such that $[s_0, s_1, \dots, s_n] \subset Y_\lambda$, and we have

$$s_0 \cap T^{-1}s_1 \cap \dots \cap T^{-n}s_n = \phi^{-1}[s_0, s_1, \dots, s_n] \in \phi^{-1}\mathcal{B}(Y_\lambda). \quad (2.8)$$

If $[s_0, s_1, \dots, s_n] \subset Y_\lambda$ fails, we have $\lambda(s_{k-1} \cap T^{-1}s_k) = 0$ for some k , so that

$$\mu(s_0 \cap T^{-1}s_1 \cap \dots \cap T^{-n}s_n) \leq \mu(T^{-(k-1)}(s_{k-1} \cap T^{-1}s_k)) = 0, \quad (2.9)$$

because $\lambda(T^{-(k-1)}(s_{k-1} \cap T^{-1}s_k)) = 0$ by the non-singularity of T and by the equivalence between μ and λ . By Condition (M1), we conclude that $\mathcal{B}(X) \subset \phi^{-1}\mathcal{B}(Y_\lambda) \bmod \mu$.

Since ϕ is a Borel surjection, we may apply von Neumann's selection theorem to ensure that there exist $Y'_\lambda \in \mathcal{B}(Y_\lambda)$ with $Y'_\lambda = Y_\lambda \bmod \nu$ and a Borel section $\psi : Y'_\lambda \rightarrow X$ such that $\phi \circ \psi(y) = y$ for all $y \in Y'_\lambda$. We write $X' = \psi(Y'_\lambda)$ and hence $\psi : Y'_\lambda \rightarrow X'$ and $\phi : X' \rightarrow Y'_\lambda$ are Borel isomorphisms.

We now verify that $X' = X \bmod \mu$. Let $s \in \alpha$. Since $\phi^{-1}\mathcal{B}(Y_\lambda) = \mathcal{B}(X) \bmod \mu$, there exist B and $B' \in \mathcal{B}(Y_\lambda)$ such that $\phi^{-1}(B) \subset X' \cap s \subset \phi^{-1}(B')$ and $\mu(\phi^{-1}(B') \cap \phi^{-1}(B)^c) = 0$. We thus have $\nu(B' \cap B^c) = \mu \circ \phi^{-1}(B' \cap B^c) = 0$. We now have

$$\mu(X' \cap s) = \nu(B') \geq \nu(\phi(X' \cap s)) = \nu(Y'_\lambda \cap \phi(s)) = \mu \circ \phi^{-1}(\phi(s)) \geq \mu(s), \quad (2.10)$$

which shows $X' \cap s = s \bmod \mu$. We therefore conclude that $X' = X \bmod \mu$. \square

2.2 Markov chain

Let us study the condition that a graph shift is a Markov chain. We introduce the transition probability on Y as

$$p(s, t) = \frac{\lambda(s \cap T^{-1}t)}{\lambda(s)}, \quad s, t \in \alpha. \quad (2.11)$$

Note that the graph shift Y_λ can be rewritten as

$$Y_\lambda := \{\mathbf{s} \in Y : p(s_0, s_1) > 0, p(s_1, s_2) > 0, \dots\}. \quad (2.12)$$

We apply Kolmogorov's extension theorem to see that, for any $s \in \alpha$, there exists a unique probability measure ν_s on Y such that

$$\nu_s([s_0, s_1, \dots, s_n]) = \delta_s(s_0)p(s_0, s_1)p(s_1, s_2) \cdots p(s_{n-1}, s_n) \quad (2.13)$$

for all cylinders, where δ_s denotes the Dirac mass at s . For a measure μ on X such that α is a μ -partition, we define a measure ν_μ on Y by

$$\nu_\mu(B) = \sum_{s \in \alpha} \mu(s) \nu_s(B), \quad B \in \mathcal{B}(Y). \quad (2.14)$$

Note that ν_μ is supported on the graph shift Y_λ .

By Theorem 2.1, we need to check the condition $[\nu_\mu = \mu \circ \phi^{-1}]$ in order to obtain the conjugacy between (X, μ, T) and $(Y_\lambda, \nu_\mu, \theta)$. For a σ -finite measure μ on X , we write $\widehat{T}_\mu : L^1(\mu) \rightarrow L^1(\mu)$ for the Perron–Frobenius operator of T with respect to μ :

$$\int_X \widehat{T}_\mu f \cdot g \, d\mu = \int_X f \cdot g \circ T \, d\mu, \quad f \in L^1(\mu), \, g \in L^\infty(\mu). \quad (2.15)$$

Proposition 2.2. *It holds that the condition*

$$\nu_\lambda = \lambda \circ \phi^{-1} \quad (2.16)$$

is satisfied if and only if

$$\widehat{T}_\lambda 1_s = \sum_{t \in \alpha} c_\lambda(s, t) 1_t \text{ with } c_\lambda(s, t) = \frac{\lambda(s)p(s, t)}{\lambda(t)}, \quad s \in \alpha. \quad (2.17)$$

Proof. (i) Suppose $\nu_\lambda = \lambda \circ \phi^{-1}$. For $s_1, \dots, s_n \in \alpha$, we set

$$g = 1_{s_1 \cap T^{-1}s_2 \cap \dots \cap T^{-(n-1)}s_n}. \quad (2.18)$$

We then have

$$\int_X \widehat{T}_\lambda 1_s \cdot g \, d\lambda = \int_X 1_{s \cap T^{-1}s_1 \cap T^{-2}s_2 \cap \dots \cap T^{-n}s_n} \, d\lambda = \nu_\lambda([s, s_1, \dots, s_n]) \quad (2.19)$$

$$= \frac{\lambda(s)p(s, s_1)}{\lambda(s_1)} \cdot \lambda(s_1)p(s_1, s_2) \cdots p(s_{n-1}, s_n) = \int_X \left(\sum_{t \in \alpha} \frac{\lambda(s)p(s, t)}{\lambda(t)} 1_t \right) \cdot g \, d\lambda, \quad (2.20)$$

which shows (2.17), since $\mathcal{G} = \mathcal{B}(X) \bmod \lambda$.

(ii) Suppose (2.17) is satisfied. By (2.7), the condition $\nu_\lambda = \lambda \circ \phi^{-1}$ is equivalent to

$$\nu_\lambda([s_0, s_1, \dots, s_n]) = \lambda(s_0 \cap T^{-1}s_1 \cap \dots \cap T^{-n}s_n), \quad s_0, \dots, s_n \in \alpha \quad (2.21)$$

for all $n \geq 0$, since $\mathcal{G} = \mathcal{B}(X) \bmod \lambda$. We shall prove (2.21) by induction in $n \geq 0$. The case $n = 0$ is obvious. Suppose (2.21) is true in the case of $n \geq 0$. In the case of $n + 1$, we have

$$\lambda(s_0 \cap T^{-1}s_1 \cap \cdots \cap T^{-n}s_n \cap T^{-(n+1)}s_{n+1}) \quad (2.22)$$

$$= \int_X 1_{s_0} \cdot 1_{s_1 \cap T^{-1}s_2 \cap \cdots \cap T^{-(n-1)}s_n \cap T^{-n}s_{n+1}} \circ T \, d\lambda \quad (2.23)$$

$$= \int_X \widehat{T}_\lambda 1_{s_0} \cdot 1_{s_1 \cap T^{-1}s_2 \cap \cdots \cap T^{-(n-1)}s_n \cap T^{-n}s_{n+1}} \, d\lambda \quad (2.24)$$

$$= \frac{\lambda(s_0)p(s_0, s_1)}{\lambda(s_1)} \lambda(s_1 \cap T^{-1}s_2 \cap \cdots \cap T^{-(n-1)}s_n \cap T^{-n}s_{n+1}). \quad (2.25)$$

By the induction assumption, we have

$$(2.25) = \frac{\lambda(s_0)p(s_0, s_1)}{\lambda(s_1)} \nu_\lambda([s_1, s_2, \dots, s_n, s_{n+1}]) \quad (2.26)$$

$$= \lambda(s_0)p(s_0, s_1)p(s_1, s_2) \cdots p(s_n, s_{n+1}) \quad (2.27)$$

$$= \nu_\lambda([s_0, s_1, \dots, s_n, s_{n+1}]), \quad (2.28)$$

which shows (2.21) in the case of $n + 1$. \square

Proposition 2.3. *Suppose $\nu_\lambda = \lambda \circ \phi^{-1}$ is satisfied. Let μ be a measure on X which is λ -a.c. Then it holds that $\nu_\mu = \mu \circ \phi^{-1}$ is satisfied if and only if μ has a constant density on each $s \in \alpha$ with respect to λ . In this case, we have*

$$\widehat{T}_\mu 1_s = \sum_{t \in \alpha} c_\mu(s, t) 1_t \text{ with } c_\mu(s, t) = \frac{\mu(s)p(s, t)}{\mu(t)}, \quad s \in \alpha, \quad (2.29)$$

where we understand that $c_\mu(s, t) = 0$ if $\mu(s) = 0$. Moreover, μ is T -invariant if and only if ν_μ is θ -invariant.

Proof. (i) Suppose $\nu_\mu = \mu \circ \phi^{-1}$ is satisfied. Then, for $s_0, \dots, s_n \in \alpha$,

$$\mu(s_0 \cap T^{-1}s_1 \cap \cdots \cap T^{-n}s_n) = \nu_\mu([s_0, s_1, \dots, s_n]) \quad (2.30)$$

$$= \mu(s_0) \nu_{s_0}([s_0, s_1, \dots, s_n]) = \frac{\mu(s_0)}{\lambda(s_0)} \nu_\lambda([s_0, s_1, \dots, s_n]). \quad (2.31)$$

Since $\mathcal{G} = \mathcal{B}(X) \bmod \lambda$ and since μ is λ -a.c., we see that

$$\mu(dx) = \sum_{s \in \alpha} \frac{\mu(s)}{\lambda(s)} 1_s(x) \lambda(dx). \quad (2.32)$$

(ii) Suppose μ has a constant density on each $s \in \alpha$ with respect to λ . Then, for $s_0, \dots, s_n \in \alpha$,

$$\nu_\mu([s_0, s_1, \dots, s_n]) = \frac{\mu(s_0)}{\lambda(s_0)} \nu_\lambda([s_0, s_1, \dots, s_n]) \quad (2.33)$$

$$= \frac{\mu(s_0)}{\lambda(s_0)} \lambda(s_0 \cap T^{-1}s_1 \cap \cdots \cap T^{-n}s_n) = \mu(s_0 \cap T^{-1}s_1 \cap \cdots \cap T^{-n}s_n), \quad (2.34)$$

which yields $\nu_\mu = \mu \circ \phi^{-1}$, since $\mathcal{B}(Y)$ is generated by cylinders.

Let $s \in \alpha$ and $g \in L^\infty(\mu)$. For $t \in \alpha$ with $\mu(t) = 0$, it is obvious that $\widehat{T}_\mu 1_s = 0$ on t . For $t \in \alpha$ with $\mu(t) > 0$, we have $g1_t \in L^\infty(\lambda)$, so that

$$\int_X \left(\widehat{T}_\mu 1_s \right) 1_t \cdot g \, d\mu = \int_X 1_s \cdot (g1_t) \circ T \, d\mu = \frac{\mu(s)}{\lambda(s)} \int_X 1_s \cdot (g1_t) \circ T \, d\lambda \quad (2.35)$$

$$= \frac{\mu(s)}{\lambda(s)} \int_X \widehat{T}_\lambda 1_s \cdot (g1_t) \, d\lambda = \frac{\mu(s)p(s,t)}{\lambda(t)} \int_X g1_t \, d\lambda = \int_X \frac{\mu(s)p(s,t)}{\mu(t)} 1_t \cdot g \, d\mu. \quad (2.36)$$

We thus obtain (2.29).

(iii) The last assertion is obvious by $\phi \circ T = \theta \circ \phi$. □

2.3 Conservative ergodicity

It is well-known (see [3, Proposition 1.2.2]) that the dynamical system (X, λ, T) is *conservative ergodic* if and only if

$$\sum_{n=1}^{\infty} 1_B \circ T^n = \infty \quad \lambda\text{-a.e. for all } B \in \mathcal{B}(X) \text{ with } \lambda(B) > 0. \quad (2.37)$$

This property can be discussed in terms of the Markov chain.

We say that a stochastic matrix $P = (p(s, t))_{s, t \in \alpha}$ is *irreducible* if for any $s, t \in \alpha$ there exists $n \geq 1$ such that

$$p^{(n)}(s, t) := \sum_{s_1, \dots, s_{n-1} \in \alpha} p(s, s_1)p(s_1, s_2) \cdots p(s_{n-2}, s_{n-1})p(s_{n-1}, t) > 0 \quad (2.38)$$

(we understand that $p^{(1)}(s, t) = p(s, t)$). We say that an irreducible stochastic matrix P is *recurrent* (resp. *transient*) if

$$\sum_{n=1}^{\infty} p^{(n)}(s, s) = \infty \quad (\text{resp. } < \infty) \quad \text{for some } s \in \alpha, \quad (2.39)$$

and in this case (2.39) holds for all $s \in \alpha$. If we denote the first hitting time of $t \in \alpha$ by

$$\varphi_t(\mathbf{s}) = \inf\{n \geq 1 : s_n = t\} \quad (\mathbf{s} = (s_0, s_1, s_2, \dots) \in Y), \quad (2.40)$$

then it is well-known (see, e.g. [26, Theorems 1.5.3 and 1.5.7]) that the condition (2.39) can be replaced by

$$\nu_s(\varphi_s < \infty) = 1 \quad (\text{resp. } < 1) \quad \text{for some } s \in \alpha. \quad (2.41)$$

It is also well-known (see e.g. [26, Theorems 1.7.5 and 1.7.6]) that, if the stochastic matrix P is irreducible recurrent, then

$$\nu_s(\varphi_t < \infty) = 1 \quad \text{for all } s, t \in \alpha, \quad (2.42)$$

and there exists a unique σ -finite P -invariant measure on α :

$$\rho(t) = \sum_{s \in \alpha} \rho(s)p(s, t), \quad t \in \alpha. \quad (2.43)$$

Proposition 2.4. *Suppose $\nu_\lambda = \lambda \circ \phi^{-1}$ is satisfied. Suppose, in addition, the stochastic matrix P is irreducible recurrent. Let ρ be a unique σ -finite P -invariant measure on α . Define the measure μ on X by*

$$\mu(dx) = \sum_{s \in \alpha} \frac{\rho(s)}{\lambda(s)} 1_s(x) dx. \quad (2.44)$$

Then it holds that the dynamical system (X, μ, T) is conservative ergodic, and that the measure μ is a unique λ -a.c. σ -finite T -invariant measure.

Proof. By Proposition 2.3, we see that $\mu \circ \phi^{-1} = \nu_\mu$. By [3, Theorem 4.5.3], we see that the dynamical system $(Y_\lambda, \nu_\mu, \theta)$ is conservative ergodic. Hence, so is (X, μ, T) , by the conjugacy.

The T -invariance of μ follows from the θ -invariance of ν_μ , by the conjugacy. The uniqueness follows from [3, Theorem 1.5.6]. \square

3 Markov chain on the λ -partition

We want to investigate irreducible recurrence or transience of the Markov chains on the λ -partitions. We follow the notation of Section 2:

$$Y = \{(s_0, s_1, s_2, \dots) : s_0, s_1, s_2, \dots \in \alpha\}. \quad (3.1)$$

We define the transition probability on α as

$$p(s, t) = \frac{\mathbb{E}\lambda(s \cap T^{-1}t)}{\lambda(s)} = \frac{p\lambda(s \cap \tau_1^{-1}t) + (1-p)\lambda(s \cap \tau_2^{-1}t)}{\lambda(s)}, \quad s, t \in \alpha. \quad (3.2)$$

For a measure μ on X , we write ν_μ for the measures on Y satisfying

$$\nu_\mu([s_0, s_1, \dots, s_n]) = \mu(s_0)p(s_0, s_1)p(s_1, s_2) \cdots p(s_{n-1}, s_n). \quad (3.3)$$

Let us study the three random maps.

Proposition 3.1. *Regarding the Hata map with $p = 1/2$, the stochastic matrix $P = (p(s, t))_{s, t \in \alpha}$ given in (3.2) is irreducible recurrent and has a unique P -invariant measure ρ given as*

$$\rho(I_k^-) = \rho(I_k^+) = (2^{k+1} - 1) \cdot 2^{-k-2}, \quad k = 0, 1, 2, \dots, \quad (1.18)$$

where I_k^- 's and I_k^+ 's have been introduced in (1.16).

Proof. We sometimes omit “mod λ ” in the identities among subsets of X . Note that

$$\tau_1 I_k^- = I_{k+1}^- \quad (k \geq 0), \quad \tau_2 I_k^- = \begin{cases} I_{k-1}^- & (k \geq 1) \\ \sum_{j=0}^{\infty} I_j^+ & (k = 0) \end{cases}, \quad (3.4)$$

$$\tau_1 I_k^+ = \begin{cases} I_{k-1}^+ & (k \geq 1) \\ \sum_{j=0}^{\infty} I_j^- & (k = 0) \end{cases}, \quad \tau_2 I_k^+ = I_{k+1}^+ \quad (k \geq 0). \quad (3.5)$$

After an easy computation we have

$$p(s, t) = \begin{cases} \frac{1}{2} & \text{if } (s, t) = (I_k^-, I_{k+1}^-) \text{ or } (I_k^+, I_{k+1}^+) \text{ for } k \geq 0 \\ \frac{1}{2} & \text{if } (s, t) = (I_k^-, I_{k-1}^-) \text{ or } (I_k^+, I_{k-1}^+) \text{ for } k \geq 1 \\ \frac{1}{2^{j+2}} & \text{if } (s, t) = (I_0^-, I_j^+) \text{ or } (I_0^+, I_j^-) \text{ for } j \geq 0 \end{cases} \quad (3.6)$$

(Note that $\frac{1}{2} + \sum_{j=0}^{\infty} \frac{1}{2^{j+2}} = 1$). It is obvious that P is irreducible. So we need only to show $\nu_{I_0^+}(\varphi_{I_0^+} < \infty) = 1$ (see Section 2.3).

Note that the Markov chain with the transition probability P starting from I_k^- for $k \geq 1$ and stopped at the first hitting of I_0^- is identical in law to the simple symmetric random walk on \mathbb{Z} starting from k and stopped at the first hitting of 0. Since the simple symmetric random walk on \mathbb{Z} is irreducible recurrent, we have

$$\nu_{I_k^-}(\varphi_{I_0^-} < \infty) = 1, \quad k \geq 1. \quad (3.7)$$

By the Markov property, we have

$$\nu_{I_0^+}(\varphi_{I_0^-} < \infty) = p(I_0^+, I_0^-) + \sum_{k \geq 1} p(I_0^+, I_k^-) \nu_{I_k^-}(\varphi_{I_0^-} < \infty) = \sum_{k \geq 0} p(I_0^+, I_k^-) = 1. \quad (3.8)$$

We also have $\nu_{I_0^-}(\varphi_{I_0^+} < \infty) = 1$ by symmetry. By the strong Markov property, we obtain

$$\nu_{I_0^+}(\varphi_{I_0^+} < \infty) \geq \nu_{I_0^+}(\varphi_{I_0^-} < \infty) \nu_{I_0^-}(\varphi_{I_0^+} < \infty) = 1, \quad (3.9)$$

which shows $\nu_{I_0^+}(\varphi_{I_0^+} < \infty) = 1$, and hence P is recurrent.

Let us prove that the measure ρ given in (1.18) is P -invariant. By symmetry, the P -invariance is equivalent to the recurrence relation:

$$\rho(I_k^-) = \frac{1}{2} \rho(I_{k+1}^+) + \frac{1}{2} \rho(I_{k-1}^+) + \frac{1}{2^{k+2}} \rho(I_0^-), \quad k = 0, 1, 2, \dots, \quad (3.10)$$

if we understand that $\rho(I_{-1}^-) = 0$. We can easily verify that the measure ρ given in (1.18) satisfies this recurrence relation. \square

Proposition 3.2. *Regarding the Pelikan map with $p = 1/2$, the stochastic matrix $P = (p(s, t))_{s, t \in \alpha}$ given in (3.2) is irreducible recurrent and has a unique P -invariant measure ρ given as*

$$\rho(I_k) = 2^{-2} \cdot (2^{k+1} - 1) \cdot 2^{-k-1}, \quad k = 0, 1, 2, \dots, \quad (1.23)$$

where I_k 's have been introduced in (1.22).

Proof. Note that

$$\tau_1 I_k = \begin{cases} I_{k-1} & (k \geq 1) \\ \sum_{j=0}^{\infty} I_j & (k = 0) \end{cases}, \quad \tau_2 I_k = I_{k+1} \quad (k \geq 0). \quad (3.11)$$

After an easy computation we have

$$p(s, t) = \begin{cases} \frac{1}{2} & \text{if } (s, t) = (I_k, I_{k-1}) \text{ or } (I_k, I_{k+1}) \text{ for } k \geq 1 \\ \frac{1}{2^{j+2}} & \text{if } (s, t) = (I_0, I_j) \text{ for } j = 0, 2, 3, \dots \\ \frac{1}{2} + \frac{1}{2^3} & \text{if } (s, t) = (I_0, I_1) \end{cases} \quad (3.12)$$

It is obvious that P is irreducible. So we need only to show $\nu_{I_0}(\varphi_{I_0} < \infty) = 1$ (see Section 2.3).

Since the Markov chain with the transition probability P starting from I_k for $k \geq 1$ and stopped at the first hitting of I_0 is identical in law to the simple symmetric random walk on \mathbb{Z} starting from k and stopped at the first hitting of 0, we have

$$\nu_{I_k}(\varphi_{I_0} < \infty) = 1, \quad k \geq 1. \quad (3.13)$$

By the Markov property, we have

$$\nu_{I_0}(\varphi_{I_0} < \infty) = p(I_0, I_0) + \sum_{k \geq 1} p(I_0, I_k) \nu_{I_k}(\varphi_{I_0} < \infty) = \sum_{k \geq 0} p(I_0, I_k) = 1, \quad (3.14)$$

which shows that P is recurrent.

The P -invariance of ρ is equivalent to the recurrence relation:

$$\rho(I_k) = \frac{1}{2} \rho(I_{k+1}) + \frac{1}{2} \rho(I_{k-1}) + \frac{1}{2^{k+2}} \rho(I_0), \quad k = 0, 1, 2, \dots, \quad (3.15)$$

if we understand that $\rho(I_{-1}) = 0$. We can easily verify that the measure ρ given in (1.23) satisfies this recurrence relation. \square

Proposition 3.3. *Regarding the mBGI map with $p = 2/3$, the stochastic matrix $P = (p(s, t))_{s, t \in \alpha}$ given in (3.2) is irreducible recurrent and has a unique P -invariant measure ρ given as*

$$\rho(I_k^-) = \rho(I_k^+) = \frac{3\sqrt{2}}{8} \left(\frac{8}{3} \cdot 2^k + \frac{1}{3} \cdot (-1)^k - 1 \right) \cdot 2^{-k-2}, \quad k = 0, 1, 2, \dots, \quad (1.28)$$

where I_k^- 's and I_k^+ 's have been introduced in (1.27).

Proof. Note that

$$\tau_1 I_k^- = \begin{cases} I_{k-1}^- & (k \geq 1) \\ \sum_{j=0}^{\infty} I_j^- & (k = 0) \end{cases}, \quad \tau_1 I_k^+ = \begin{cases} I_{k-1}^+ & (k \geq 1) \\ \sum_{j=0}^{\infty} I_j^+ & (k = 0) \end{cases} \quad (3.16)$$

and

$$\tau_2 I_k^- = I_{k+2}^+ \quad (k \geq 0), \quad \tau_2 I_k^+ = I_{k+2}^- \quad (k \geq 0). \quad (3.17)$$

It is easy to see that

$$p(s, t) = \begin{cases} \frac{1}{3} & \text{if } (s, t) = (I_k^-, I_{k+2}^+) \text{ or } (I_k^+, I_{k+2}^-) \text{ for } k \geq 0 \\ \frac{2}{3} & \text{if } (s, t) = (I_k^-, I_{k-1}^-) \text{ or } (I_k^+, I_{k-1}^+) \text{ for } k \geq 1 \\ \frac{2}{3} \cdot \frac{1}{2^{j+1}} & \text{if } (s, t) = (I_0^-, I_j^-) \text{ or } (I_0^+, I_j^+) \text{ for } j \geq 0 \end{cases} \quad (3.18)$$

and that P is irreducible. So we need only to show $\nu_{I_0^-}(\varphi_{I_0^-} < \infty) = 1$ (see Section 2.3).

We define the transition probability $Q = (q(k, j))_{k, j=0}^\infty$ on $\{0, 1, 2, \dots\}$ as

$$q(k, j) = \begin{cases} p(I_k^\pm, I_j^-) + p(I_k^\pm, I_j^+) & \text{if } k \geq 1 \\ 1 & \text{if } k = j = 0 \end{cases} = \begin{cases} 1/3 & \text{if } j = k + 2 \\ 2/3 & \text{if } j = k - 1 \\ 1 & \text{if } k = j = 0 \end{cases}. \quad (3.19)$$

Let $(\{W'_n\}, \{\mathbb{P}_k\})$ denote the random walk on \mathbb{Z} such that $\mathbb{P}_k(W'_n = k) = 1$ and

$$\mathbb{P}_k(W'_n - W'_{n-1} = 2) = 1/3, \quad \mathbb{P}_k(W'_n - W'_{n-1} = -1) = 2/3. \quad (3.20)$$

Then the transition probability Q is nothing else but that of the random walk $\{W'_n\}$ stopped at its first hitting time of 0, which will be denoted by $\varphi_0^{W'}$. Since $\mathbb{E}_k[W'_n - W'_{n-1}] = 0$, we see by [21, Proposition 9.14] that $\limsup_{n \rightarrow \infty} W'_n = \limsup_{n \rightarrow \infty} (-W'_n) = \infty$ a.s., which shows that $\{W'_n\}$ is irreducible recurrent. Hence we have

$$\nu_{I_k^\pm}(\varphi_{I_0^-} < \infty \text{ or } \varphi_{I_0^+} < \infty) = \mathbb{P}_k(\varphi_0^{W'} < \infty) = 1, \quad k \geq 1. \quad (3.21)$$

This implies $\nu_{I_0^\pm}(\varphi_{I_0^-} < \infty \text{ or } \varphi_{I_0^+} < \infty) = 1$, and consequently $a^\pm + b^\pm = 1$ with

$$a^\pm := \nu_{I_0^\pm}(\varphi_{I_0^-} < \infty, \varphi_{I_0^-} < \varphi_{I_0^+}), \quad b^\pm := \nu_{I_0^\pm}(\varphi_{I_0^+} < \infty, \varphi_{I_0^+} < \varphi_{I_0^-}). \quad (3.22)$$

By the strong Markov property, we have

$$\nu_{I_0^+}(\varphi_{I_0^-} < \infty) = a^+ + b^+ \nu_{I_0^+}(\varphi_{I_0^-} < \infty). \quad (3.23)$$

Since $1 - b^+ = a^+ \geq p(I_0^+, I_2^-)p(I_2^-, I_1^-)p(I_1^-, I_0^-) > 0$, we obtain $\nu_{I_0^+}(\varphi_{I_0^-} < \infty) = 1$. We now obtain

$$\nu_{I_0^-}(\varphi_{I_0^-} < \infty) = a^- + b^- \nu_{I_0^+}(\varphi_{I_0^-} < \infty) = a^- + b^- = 1, \quad (3.24)$$

which shows P is recurrent.

The P -invariance of ρ is equivalent to the recurrence relations:

$$\rho(I_k^-) = \frac{1}{3}\rho(I_{k-2}^+) + \frac{2}{3}\rho(I_{k+1}^-) + \frac{2}{3} \cdot \frac{1}{2^{k+1}}\rho(I_0^-), \quad k = 0, 1, 2, \dots, \quad (3.25)$$

$$\rho(I_k^+) = \frac{1}{3}\rho(I_{k-2}^-) + \frac{2}{3}\rho(I_{k+1}^+) + \frac{2}{3} \cdot \frac{1}{2^{k+1}}\rho(I_0^+), \quad k = 0, 1, 2, \dots, \quad (3.26)$$

if we understand that $\rho(I_{-1}^\pm) = \rho(I_{-2}^\pm) = 0$. We can easily verify that the measure ρ given in (1.28) satisfies this recurrence relation. \square

4 Skew-product representation and Markov partition

Let us recall that a random interval map

$$T = \begin{cases} \tau_1 & \text{(with probability } p), \\ \tau_2 & \text{(with probability } 1 - p) \end{cases} \quad (4.1)$$

for two deterministic maps $\tau_1, \tau_2 : [0, 1] \rightarrow [0, 1]$ can be represented by skew-product. In what follows, we write $X = [0, 1]$ and write λ for the Lebesgue measure on X . Let us write $(\Omega, \mathcal{B}(\Omega), \mathbb{P})$ for the coin flip, i.e.,

$$\Omega = \{\omega = (\omega_1, \omega_2, \dots) : \omega_1, \omega_2, \dots \in \{\tau_1, \tau_2\}\} \quad (4.2)$$

equipped with the product topology, $\mathcal{B}(\Omega)$ for the Borel field of Ω , and

$$\mathbb{P}(\omega \in \Omega : \omega_1 = \chi_1, \dots, \omega_n = \chi_n) = p^{N_n^1(\chi)}(1-p)^{N_n^2(\chi)} \quad (4.3)$$

for all $\chi_1, \dots, \chi_n \in \{\tau_1, \tau_2\}$ and all n , where $N_n^1(\chi) = \#\{k = 1, \dots, n : \chi_k = \tau_1\}$ and $N_n^2(\chi) = \#\{k = 1, \dots, n : \chi_k = \tau_2\}$. For $B \in \mathcal{B}(X)$, we denote $\tilde{B} = B^\sim = \Omega \times B$, and we equip $\tilde{X} = \Omega \times X$ with the product topology of Ω and X . We write $\mathcal{B}(\tilde{X})$ for the Borel field of \tilde{X} and write $\tilde{\lambda} = \mathbb{P} \otimes \lambda$ for the product measure of \mathbb{P} and λ . We define the deterministic transform $\tilde{T} : \tilde{X} \rightarrow \tilde{X}$ as

$$\tilde{T}(\omega, x) = (\theta\omega, \omega_1 x), \quad \omega = (\omega_1, \omega_2, \dots) \in \Omega, \quad x \in X. \quad (4.4)$$

Here $\theta : \Omega \rightarrow \Omega$ stands for the shift operator: $\theta(\omega_1, \omega_2, \dots) = (\omega_2, \omega_3, \dots)$. The random maps $\{T_n\}$ and $\{T^{(n)}\}$ in Section 1 are obtained as

$$T_n(\omega)(x) = \omega_n(x), \quad T^{(n)}(\omega)(x) = \omega_n \omega_{n-1} \cdots \omega_1(x), \quad (4.5)$$

so that $\tilde{T}^n(\omega, x) = (\theta_n \omega, T^{(n)}(\omega)(x))$. For a measure μ on X , it is easy to see that $\tilde{\mu} := \mathbb{P} \otimes \mu$ is \tilde{T} -invariant if and only if μ is T -invariant in the sense of (1.11).

Theorem 4.1. *Regarding the Hata map with $p = 1/2$, we introduce the $\tilde{\lambda}$ -partition of \tilde{X} defined by*

$$\tilde{\alpha} = \{\tilde{s} : s \in \alpha\} = \left\{ \tilde{I}_k^-, \tilde{I}_k^+ \right\}_{k=0}^\infty, \quad (4.6)$$

where I_k^- 's and I_k^+ 's have been introduced in (1.16). Then it is a Markov partition for the dynamical system $(\tilde{X}, \tilde{\lambda}, \tilde{T})$, i.e. the following assertions hold:

- (i) The σ -field $\mathcal{G} := \sigma\left(\tilde{T}^{-n}(\tilde{s}) : \tilde{s} \in \tilde{\alpha}, n \geq 0\right)$ coincides with $\mathcal{B}(\tilde{X}) \bmod \tilde{\lambda}$;
- (ii) The map $\tilde{T} : \tilde{s} \mapsto \tilde{T}\tilde{s}$ is invertible $\tilde{\lambda}$ -a.e. for all $\tilde{s} \in \tilde{\alpha}$;
- (iii) The forward image $\tilde{T}\tilde{s}$ belongs to $\sigma(\tilde{\alpha}) \bmod \tilde{\lambda}$ for all $\tilde{s} \in \tilde{\alpha}$.

Proof. We sometimes omit “mod $\tilde{\lambda}$ ” in the identities among subsets of \tilde{X} . We prove Claims (ii) and (iii) at once. For $(\omega, x) \in \tilde{I}_k^- \in \tilde{\alpha}$ with $k \geq 1$, we have

$$\tilde{T}(\omega, x) = (\theta\omega, \omega_1(x)) \in \tilde{I}_{k+1}^- \text{ or } \tilde{I}_{k-1}^- \text{ according as } \omega_1 = \tau_1 \text{ or } \tau_2. \quad (4.7)$$

We thus see that $\tilde{T} : \tilde{I}_k^- \rightarrow \tilde{T}(\tilde{I}_k^-) = \tilde{I}_{k+1}^- + \tilde{I}_{k-1}^- \in \sigma(\tilde{\alpha})$ is invertible as

$$\tilde{T}^{-1}(\omega, x) = \begin{cases} ((\tau_1, \omega), 2x) & ((\omega, x) \in \tilde{I}_{k+1}^-), \\ ((\tau_2, \omega), x/2) & ((\omega, x) \in \tilde{I}_{k-1}^-). \end{cases} \quad (4.8)$$

For $(\omega, x) \in \tilde{I}_0^- \in \tilde{\alpha}$, we have

$$\tilde{T}(\omega, x) = (\theta\omega, \omega_1(x)) \in \tilde{I}_1^- \text{ or } (1/2, 1)^\sim \text{ according as } \omega_1 = \tau_1 \text{ or } \tau_2. \quad (4.9)$$

Since $\tilde{T}(\tilde{I}_0^-) = \tilde{I}_1^- + (1/2, 1)^\sim = \tilde{I}_1^- + \sum_{k=0}^\infty \tilde{I}_k^+ \in \sigma(\tilde{\alpha})$, we thus see that $\tilde{T} : \tilde{I}_0^- \rightarrow \tilde{T}(\tilde{I}_0^-)$ is invertible $\tilde{\lambda}$ -a.e. as

$$\tilde{T}^{-1}(\omega, x) = \begin{cases} ((\tau_1, \omega), 2x) & ((\omega, x) \in \tilde{I}_1^-), \\ ((\tau_2, \omega), x/2) & ((\omega, x) \in \tilde{I}_k^+, k \geq 0). \end{cases} \quad (4.10)$$

By symmetry we obtain a similar result for $\tilde{T} : \tilde{I}_k^+ \rightarrow \tilde{T}(\tilde{I}_k^+)$.

Let us now prove Claim (i). For $\omega = (\omega_1, \omega_2, \dots)$ and for $x \in X$, we write $\xi_n(\omega, x) = \omega_n$. It suffices to show

$$\{\xi_n = \tau_1\}, \{\xi_n = \tau_2\} \in \mathcal{G}, \quad n = 1, 2, \dots, \quad (4.11)$$

$$\left(\frac{i}{2^m}, \frac{i+1}{2^m}\right)^\sim \in \mathcal{G}, \quad i = 0, 1, \dots, 2^m - 1, \quad m = 0, 1, 2, \dots \quad (4.12)$$

To obtain $\{\xi_1 = \tau_1\} \in \mathcal{G}$, we have

$$\{\xi_1 = \tau_1\} = \sum_{k \geq 0} \{\xi_1 = \tau_1\} \cap \tilde{I}_k^- + \sum_{k \geq 0} \{\xi_1 = \tau_1\} \cap \tilde{I}_k^+ \quad (4.13)$$

$$= \sum_{k \geq 0} \left(\tilde{T}^{-1}\tilde{I}_{k+1}^-\right) \cap \tilde{I}_k^- + \sum_{k \geq 1} \left(\tilde{T}^{-1}\tilde{I}_{k-1}^+\right) \cap \tilde{I}_k^+ + \left(\sum_{j \geq 0} \tilde{T}^{-1}\tilde{I}_j^-\right) \cap \tilde{I}_0^+ \in \mathcal{G}, \quad (4.14)$$

and hence we obtain

$$\{\xi_n = \tau_1\} = \tilde{T}^{-(n-1)}\{\xi_1 = \tau_1\} \in \mathcal{G}. \quad (4.15)$$

Since $\{\xi_n = \tau_2\} = \{\xi_n = \tau_1\}^c \in \mathcal{G}$, we have proved (4.11).

We proceed to show (4.12) by induction. The case $m = 0$ is obvious; $(0, 1)^\sim = \tilde{X} \in \mathcal{G}$. Suppose (4.12) holds for m and let us prove that (4.12) holds also for $m + 1$. Set

$$\eta_m = \begin{cases} 1 & (\xi_n = \tau_1) \\ -1 & (\xi_n = \tau_2) \end{cases} \quad (n \geq 1) \quad (4.16)$$

and

$$W_n = \eta_1 + \cdots + \eta_n, \quad n \geq 1. \quad (4.17)$$

Note that $(\{W_n\}, \mathbb{P})$ is the simple symmetric random walk on \mathbb{Z} starting from 0. For $k \in \mathbb{Z}$, we denote the first hitting time of k for the random walk by

$$\varphi_k^W = \inf\{n \geq 1 : W_n = k\}. \quad (4.18)$$

Since $(\{W_n\}, \mathbb{P})$ is irreducible recurrent, we have $\mathbb{P}(\varphi_k^W < \infty) = 1$ for all $k \in \mathbb{Z}$.

For any $i = 0, \dots, 2^m - 1$ so that $\frac{i+1}{2^{m+1}} \leq \frac{1}{2}$, we want to show $(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}})^\sim \in \mathcal{G}$. We divide it into

$$\left(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}}\right)^\sim = \sum_{n=1}^{\infty} \left(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}}\right)^\sim \cap \{\varphi_{-1}^W = n\}, \quad (4.19)$$

where φ_{-1}^W can be regarded as the first hitting time of m for the random walk starting from $m+1$. On the event $\{\varphi_{-1}^W = n\}$, we have $\xi_{j-1} \circ \cdots \circ \xi_1(x) \in (0, \frac{i+1}{2^{m+1}})$ for all $j \leq n$, so that $\xi_j \circ \xi_{j-1} \circ \cdots \circ \xi_1(x) = 2^{-W_j}x$ for all $j \leq n$. Hence

$$\left(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}}\right)^\sim \cap \{\varphi_{-1}^W = n\} = \left(\tilde{T}^{-n} \left(\frac{i}{2^m}, \frac{i+1}{2^m}\right)^\sim\right) \cap \{\varphi_{-1}^W = n\}. \quad (4.20)$$

Since $\tilde{T}^{-n}(\frac{i}{2^m}, \frac{i+1}{2^m})^\sim \in \mathcal{G}$ by the assumption of induction and since

$$\{\varphi_{-1}^W = n\} \in \sigma(\xi_1, \dots, \xi_n) \in \mathcal{G}, \quad (4.21)$$

we obtain $(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}})^\sim \in \mathcal{G}$ for $i = 0, 1, \dots, 2^m - 1$. By symmetry, we also obtain $(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}})^\sim \in \mathcal{G}$ for $i = 2^m, 2^m + 1, \dots, 2^{m+1} - 1$. We have now obtained (4.12).

The proof is therefore complete. \square

Let us study the Pelikan map with $p = 1/2$. Unfortunately, the $\tilde{\lambda}$ -partition $\tilde{\alpha} = \{\tilde{s} : s \in \alpha\}$ is not a Markov partition for $(\tilde{X}, \tilde{\mu}, \tilde{T})$, because the map $\tilde{T} : \tilde{I}_0 \rightarrow \tilde{T}\tilde{I}_0$ is not invertible. To overcome this difficulty, we write

$$\tilde{s}^{\tau_1} = \{\xi_1 = \tau_1\} \cap \tilde{s}, \quad \tilde{s}^{\tau_2} = \{\xi_1 = \tau_2\} \cap \tilde{s}, \quad s \in \alpha. \quad (4.22)$$

Theorem 4.2. *Regarding the Pelikan map with $p = 1/2$, we introduce the $\tilde{\lambda}$ -partition of \tilde{X} defined by*

$$\tilde{\alpha}^* = \left\{ \tilde{I}_k^{\tau_1}, \tilde{I}_k^{\tau_2} \right\}_{k=0}^{\infty}, \quad (4.23)$$

where I_k 's have been introduced in (1.22). Then it is a Markov partition for the dynamical system $(\tilde{X}, \tilde{\lambda}, \tilde{T})$.

Proof. Claims (ii) and (iii) of Theorem 4.1 are obvious, since $\tilde{T}(\omega, x) = (\theta\omega, \chi(x))$ for $(\omega, x) \in s^x$ with $s \in \alpha$ and $\chi \in \{\tau_1, \tau_2\}$. For Claim (i), it suffices to show

$$\left(\frac{i}{2^m}, \frac{i+1}{2^m}\right)^\sim \in \mathcal{G}, \quad i = 0, 1, \dots, 2^m - 1, \quad m = 0, 1, 2, \dots \quad (4.24)$$

We show (4.24) by induction. The case $m = 0$ is obvious. Suppose (4.24) holds for m . Recall that φ_k^W is the first hitting time of k for the simple symmetric random walk $(\{W_n\}, \mathbb{P})$, which has been introduced in (4.18). In the same way as Theorem 4.1 we see that, for any $i = 0, \dots, 2^m - 1$ so that $\frac{i+1}{2^{m+1}} \leq \frac{1}{2}$,

$$\left(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}}\right)^\sim = \sum_{n=1}^{\infty} \left(\tilde{T}^{-n} \left(\frac{i}{2^m}, \frac{i+1}{2^m} \right)^\sim \right) \cap \{\varphi_{-1}^W = n\} \in \mathcal{G}, \quad (4.25)$$

and for $i = 2^m, 2^m + 1, \dots, 2^{m+1} - 1$ so that $\frac{i}{2^{m+1}} \geq \frac{1}{2}$,

$$\left(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}}\right)^\sim = \sum_{n=1}^{\infty} \left(\tilde{T}^{-n} \left(\frac{i-2^m}{2^m}, \frac{i+1-2^m}{2^m} \right)^\sim \right) \cap \{\varphi_{-1}^W = n\} \in \mathcal{G}. \quad (4.26)$$

We have now obtained (4.24). The proof is therefore complete. \square

Theorem 4.3. *Regarding the mBGI map with $p = 2/3$, we introduce the $\tilde{\lambda}$ -partition of \tilde{X} defined by*

$$\tilde{\alpha} = \{\tilde{s} : s \in \alpha\} = \left\{ \tilde{I}_k^-, \tilde{I}_k^+ \right\}_{k=0}^{\infty}, \quad (4.27)$$

where I_k^- 's and I_k^+ 's have been introduced in (1.27). Then it is a Markov partition for the dynamical system $(\tilde{X}, \tilde{\lambda}, \tilde{T})$.

Proof. Claims (ii) and (iii) of Theorem 4.1 are obvious. As it is obvious that $\{\xi_n = \tau_1\}$ and $\{\xi_n = \tau_2\}$ belong to \mathcal{G} , for Claim (i) it suffices to show by induction that

$$\mathcal{I}_m := \left\{ \left(\frac{i}{2^m}, \frac{i+1}{2^m} \right)^\sim \right\}_{\substack{i=0,1,\dots,2^m-1 \\ m=0,1,2,\dots}} \subset \mathcal{G}. \quad (4.28)$$

The cases $m = 0$ and 1 are obvious. Suppose (4.28) holds for $m \geq 1$ and let us prove that (4.28) holds also for $m + 1$. Set

$$\eta'_n = \begin{cases} -1 & (\xi_n = \tau_1) \\ 2 & (\xi_n = \tau_2) \end{cases} \quad (n \geq 1) \quad (4.29)$$

and

$$W'_n = \eta'_1 + \dots + \eta'_n, \quad n \geq 1. \quad (4.30)$$

For $k \in \mathbb{Z}$, we denote the first hitting time of k for the random walk by

$$\varphi_k^{W'} = \inf\{n \geq 1 : W'_n = k\}. \quad (4.31)$$

As we have mentioned it in the proof of Proposition 3.3, we see that $(\{W'_n\}, \mathbb{P})$ is an irreducible recurrent random walk on \mathbb{Z} , so that we have $\mathbb{P}(\varphi_k^{W'} < \infty) = 1$ for all $k \in \mathbb{Z}$. For any $i = 0, \dots, 2^{m+1} - 1$, we have we divide

$$\left(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}}\right)^\sim = \sum_{n=1}^{\infty} \left(\frac{i}{2^{m+1}}, \frac{i+1}{2^{m+1}}\right)^\sim \cap \{\varphi_{-1}^{W'} = n\} \quad (4.32)$$

$$= \sum_{n=1}^{\infty} \left(\tilde{T}^{-n} \left(\frac{i}{2^m}, \frac{i+1}{2^m}\right)^\sim\right) \cap \{\varphi_{-1}^{W'} = n\} \in \mathcal{G}, \quad (4.33)$$

by the assumption of induction and by the fact that $\{\varphi_{-1}^{W'} = n\} \in \sigma(\xi_1, \dots, \xi_n) \in \mathcal{G}$. We have now obtained (4.28).

The proof is therefore complete. \square

5 Invariant measure

We want to show that our dynamical system $(\tilde{X}, \tilde{\mu}, \tilde{T})$ is conjugate to an irreducible recurrent Markov chain.

For the Hata map with $p = 1/2$, we have seen in Theorem 4.1 that $\tilde{\alpha}$ is a Markov partition, and so we see by Theorem 2.1 that our dynamical system $(\tilde{X}, \tilde{\lambda}, \tilde{T})$ is conjugate to $(\tilde{Y}, \tilde{\lambda} \circ \tilde{\phi}^{-1}, \theta)$ on the graph shift

$$\tilde{Y} := \{\tilde{\mathbf{s}} = (\tilde{s}_0, \tilde{s}_1, \tilde{s}_2, \dots) : \tilde{s}_0, \tilde{s}_1, \tilde{s}_2, \dots \in \tilde{\alpha}\} \quad (5.1)$$

with the Borel bijection $\tilde{\phi} : \tilde{X} \rightarrow \tilde{Y}$ defined as

$$\tilde{\phi}(\omega, x) = \tilde{\mathbf{s}} \text{ with } \tilde{T}^n(\omega, x) \in \tilde{s}_n \text{ for all } n. \quad (5.2)$$

Note that the graph shift \tilde{Y} can be identified with Y given in (3.1). The transition probability on $\tilde{\alpha}$ defined as

$$\tilde{p}(\tilde{s}, \tilde{t}) = \frac{\tilde{\lambda}(\tilde{s} \cap \tilde{T}^{-1}\tilde{t})}{\tilde{\lambda}(\tilde{s})}, \quad s, t \in \alpha \quad (5.3)$$

is nothing else but the transition probability p on α defined in (3.2). For a measure μ on X , we write $\tilde{\nu}_\mu$ for the measure on \tilde{Y} satisfying

$$\tilde{\nu}_\mu([\tilde{s}_0, \tilde{s}_1, \dots, \tilde{s}_n]) = \tilde{\mu}(\tilde{s}_0) \tilde{p}(\tilde{s}_0, \tilde{s}_1) \tilde{p}(\tilde{s}_1, \tilde{s}_2) \cdots \tilde{p}(\tilde{s}_{n-1}, \tilde{s}_n), \quad (5.4)$$

which equals to $\nu_\mu([s_0, s_1, \dots, s_n])$ defined in (3.3).

The following theorem slightly generalizes Theorem 1.1, because it asserts conservative ergodicity.

Theorem 5.1. *Regarding the Hata map with $p = 1/2$, it holds that $\tilde{\nu}_\lambda = \tilde{\lambda} \circ \tilde{\phi}^{-1}$. Consequently, the dynamical system $(\tilde{X}, \tilde{\lambda}, \tilde{T})$ is conservative ergodic, and for the measure given in (1.17) and (1.18), the measure $\tilde{\mu}$ is a unique $\tilde{\lambda}$ -a.c. σ -finite \tilde{T} -invariant measure.*

Proof. We write \tilde{T}_λ^\wedge for the Perron–Frobenius operator of \tilde{T} with respect to $\tilde{\lambda}$. For $A \in \mathcal{B}(\Omega)$, $B \in \mathcal{B}(X)$ and $s \in \alpha$, we have

$$\int_{\tilde{X}} \tilde{T}_\lambda^\wedge 1_{\tilde{s}} \cdot 1_{A \times B} d\tilde{\lambda} = \int_{\tilde{X}} 1_{\tilde{s}} \cdot 1_{A \times B} \circ \tilde{T} d\tilde{\lambda} \quad (5.5)$$

$$= \int_{\Omega} \int_X 1_s(x) \cdot 1_A(\theta\omega) 1_B(\omega_1(x)) \mathbb{P}(d\omega) dx \quad (5.6)$$

$$= \mathbb{P}(A) \cdot \frac{1}{2} \sum_{i=1,2} \int_X 1_s(\tau_i^{-1}(x)) \cdot 1_B(x) (\tau_i^{-1})' dx \quad (5.7)$$

$$= \mathbb{P}(A) \cdot \frac{1}{2} \sum_{i=1,2} \int_X \sum_{t \in \alpha} 1_{\tau_i s \cap t}(x) \cdot 1_B(x) (\tau_i^{-1})' dx \quad (5.8)$$

$$= \int_{\tilde{X}} \sum_{t \in \alpha} \left(\frac{1}{2} \sum_{i=1,2} (\tau_i^{-1})' 1_{(\tau_i s \cap t)^\sim} \right) \cdot 1_{A \times B} d\tilde{\lambda}. \quad (5.9)$$

Since $(\tau_i s \cap t)^\sim = \tilde{t}$ or \emptyset (see (iii) of Theorem 4.1) and since $(\tau_i^{-1})'$ is constant on t , we obtain the representation

$$\tilde{T}_\lambda^\wedge 1_{\tilde{s}} = \sum_{t \in \alpha} c(s, t) 1_{\tilde{t}}, \quad s \in \alpha \quad (5.10)$$

for some function $c(s, t)$ on $\alpha \times \alpha$. This shows $\tilde{\nu}_\lambda = \tilde{\lambda} \circ \tilde{\phi}^{-1}$ by Proposition 2.2.

The remainder of the proof is therefore obvious by Propositions 2.4 and 3.1. \square

The following theorem slightly generalizes Theorem 1.5.

Theorem 5.2. *Regarding the mBGI map with $p = 2/3$, it holds that $\tilde{\nu}_\lambda = \tilde{\lambda} \circ \tilde{\phi}^{-1}$. Consequently, the dynamical system $(\tilde{X}, \tilde{\lambda}, \tilde{T})$ is conservative ergodic, and for the measure given in (1.28) the measure $\tilde{\mu}$ is a unique $\tilde{\lambda}$ -a.c. σ -finite \tilde{T} -invariant measure.*

The proof is almost the same as that of Theorem 5.1, and so we omit it.

For the Pelikan map with $p = 1/2$, we have seen in Theorem 4.2 that $\tilde{\alpha}^*$ is a Markov partition, and so our dynamical system $(\tilde{X}, \tilde{\lambda}, \tilde{T})$ is conjugate to $(\tilde{Y}^*, \tilde{\lambda} \circ (\tilde{\phi}^*)^{-1}, \theta)$ on the graph shift

$$\tilde{Y}^* := \{ \tilde{\mathbf{s}}^* = (\tilde{s}_0^{\chi_0}, \tilde{s}_1^{\chi_1}, \tilde{s}_2^{\chi_2}, \dots) : \tilde{s}_0^{\chi_0}, \tilde{s}_1^{\chi_1}, \tilde{s}_2^{\chi_2}, \dots \in \tilde{\alpha}^* \} \quad (5.11)$$

with the Borel bijection $\tilde{\phi}^* : \tilde{X} \rightarrow \tilde{Y}^*$ defined as

$$\tilde{\phi}^*(\omega, x) = \tilde{\mathbf{s}}^* \text{ with } \tilde{T}^n(\omega, x) \in \tilde{s}_n^{\chi_n} \text{ for all } n. \quad (5.12)$$

The transition probability on $\tilde{\alpha}^*$ defined as

$$\tilde{p}^*(\tilde{s}^{\chi_1}, \tilde{t}^{\chi_2}) = \frac{\tilde{\lambda}(\tilde{s}^{\chi_1} \cap \tilde{T}^{-1}\tilde{t}^{\chi_2})}{\tilde{\lambda}(\tilde{s}^{\chi_1})} = \frac{1}{2} \frac{\lambda(s \cap \chi_1^{-1}t)}{\lambda(s)}, \quad s, t \in \alpha, \quad \chi_1, \chi_2 \in \{\tau_1, \tau_2\}. \quad (5.13)$$

Note that $\tilde{p}^*(\tilde{s}^\chi, \tilde{t}^{\tau_1}) = \tilde{p}^*(\tilde{s}^\chi, \tilde{t}^{\tau_2})$ and

$$\tilde{p}^*(\tilde{s}^{\tau_1}, \tilde{t}^{\tau_1}) + \tilde{p}^*(\tilde{s}^{\tau_2}, \tilde{t}^{\tau_1}) = p(s, t), \quad s, t \in \alpha, \quad (5.14)$$

where $p(s, t)$ is given in (3.2). For a measure μ on X , we write $\tilde{\nu}_\mu^*$ for the measure on \tilde{Y}^* satisfying

$$\tilde{\nu}_\mu^*([\tilde{s}_0^{\chi_0}, \tilde{s}_1^{\chi_1}, \dots, \tilde{s}_n^{\chi_n}]) = \tilde{\mu}(\tilde{s}_0^{\chi_0})\tilde{p}^*(\tilde{s}_0^{\chi_0}, \tilde{s}_1^{\chi_1})\tilde{p}^*(\tilde{s}_1^{\chi_1}, \tilde{s}_2^{\chi_2}) \cdots \tilde{p}^*(\tilde{s}_{n-1}^{\chi_{n-1}}, \tilde{s}_n^{\chi_n}). \quad (5.15)$$

The following theorem slightly generalizes Theorem 1.3.

Theorem 5.3. *Regarding the Pelikan map with $p = 1/2$, it holds that $\tilde{\nu}_\lambda^* = \tilde{\lambda} \circ (\tilde{\phi}^*)^{-1}$. Consequently, the dynamical system $(\tilde{X}, \tilde{\lambda}, \tilde{T})$ is conservative ergodic, and for the measure given in (1.23), the measure $\tilde{\mu}$ is a unique $\tilde{\lambda}$ -a.c. σ -finite \tilde{T} -invariant measure.*

Proof. Let us first prove that the irreducible stochastic matrix \tilde{P}^* corresponding to the transition probability \tilde{p}^* is recurrent. For any $s_0 \in \alpha$ and $\chi_0 \in \{\tau_1, \tau_2\}$,

$$\sum_{n=1}^{\infty} (\tilde{p}^*)^{(n+1)}(\tilde{s}_0^{\chi_0}, \tilde{s}_0^{\chi_0}) \quad (5.16)$$

$$= \sum_{n=1}^{\infty} \sum_{\substack{s_1, \dots, s_n \in \alpha \\ \chi_1, \dots, \chi_n = \tau_1, \tau_2}} \tilde{p}^*(\tilde{s}_0^{\chi_0}, \tilde{s}_1^{\chi_1}) \cdots \tilde{p}^*(\tilde{s}_{n-1}^{\chi_{n-1}}, \tilde{s}_n^{\chi_n}) \tilde{p}^*(\tilde{s}_n^{\chi_n}, \tilde{s}_0^{\chi_0}) \quad (5.17)$$

$$= \sum_{n=1}^{\infty} \sum_{\substack{s_1, \dots, s_n \in \alpha \\ \chi_1, \dots, \chi_n = \tau_1, \tau_2}} \tilde{p}^*(\tilde{s}_0^{\chi_0}, \tilde{s}_1^{\tau_1}) \cdots \tilde{p}^*(\tilde{s}_{n-1}^{\chi_{n-1}}, \tilde{s}_n^{\tau_1}) \tilde{p}^*(\tilde{s}_n^{\chi_n}, \tilde{s}_0^{\tau_1}) \quad (5.18)$$

$$= \sum_{n=1}^{\infty} \sum_{s_1, \dots, s_n \in \alpha} \tilde{p}^*(\tilde{s}_0^{\chi_0}, \tilde{s}_1^{\tau_1}) p(s_1, s_2) \cdots p(s_{n-1}, s_n) p(s_n, s_0) \quad (5.19)$$

$$= \sum_{s_1 \in \alpha} \tilde{p}^*(\tilde{s}_0^{\chi_0}, \tilde{s}_1^{\tau_1}) \sum_{n=1}^{\infty} \sum_{s_2, \dots, s_n \in \alpha} p(s_1, s_2) \cdots p(s_{n-1}, s_n) p(s_n, s_0), \quad (5.20)$$

which is divergent since $P = (p(s, t))_{s, t \in \alpha}$ is irreducible recurrent by Proposition 3.2. Thus we see that \tilde{P}^* is recurrent.

Hence there exists a unique \tilde{P}^* -invariant measure $\tilde{\rho}^*$ on $\tilde{\alpha}^*$. We then have

$$\tilde{\rho}^*(\tilde{t}^\chi) = \sum_{s \in \alpha} \sum_{i=1,2} \tilde{\rho}^*(\tilde{s}^{\tau_i}) \tilde{p}^*(\tilde{s}^{\tau_i}, \tilde{t}^\chi) = \sum_{s \in \alpha} \sum_{i=1,2} \tilde{\rho}^*(\tilde{s}^{\tau_i}) \tilde{p}^*(\tilde{s}^{\tau_i}, \tilde{t}^{\tau_1}), \quad (5.21)$$

and so we have $\tilde{\rho}^*(\tilde{t}^{\tau_1}) = \tilde{\rho}^*(\tilde{t}^{\tau_2})$. We then have

$$\tilde{\rho}^*(\tilde{t}^{\tau_1}) = \sum_{s \in \alpha} \sum_{i=1,2} \tilde{\rho}^*(\tilde{s}^{\tau_1}) \tilde{p}^*(\tilde{s}^{\tau_i}, \tilde{t}^{\tau_1}) = \sum_{s \in \alpha} \tilde{\rho}^*(\tilde{s}^{\tau_1}) p(s, t), \quad (5.22)$$

and hence by Proposition 3.2 we may redefine $\tilde{\rho}^*(\tilde{t}^{\tau_1}) = \tilde{\rho}^*(\tilde{t}^{\tau_2}) = \rho(t)/2$ for all $t \in \alpha$, noting that the uniqueness is up to constant multiplicity.

We write \tilde{T}_λ^\wedge for the Perron–Frobenius operator of \tilde{T} with respect to $\tilde{\lambda}$. For $A \in \mathcal{B}(\Omega)$, $B \in \mathcal{B}(X)$ and $s \in \alpha$ and $\chi \in \{\tau_1, \tau_2\}$, we obtain, by a similar argument to Theorem 5.1,

$$\int_{\tilde{X}} = \int_{\tilde{X}} \sum_{t \in \alpha} \left(\frac{1}{2} (\chi^{-1})' 1_{(\chi s \cap t)^\sim} \right) \cdot 1_{A \times B} d\tilde{\lambda}. \quad (5.23)$$

Since $(\chi s \cap t)^\sim = \tilde{t}$ or \emptyset and since $(\chi^{-1})'$ is constant on t , we obtain the representation

$$\tilde{T}_\lambda^\wedge 1_{\tilde{s}^\chi} = \sum_{t \in \alpha} c(\tilde{s}^\chi, t) 1_{\tilde{t}}, \quad \tilde{s}^\chi \in \tilde{\alpha}^* \quad (5.24)$$

for some function $c(\tilde{s}^\chi, t)$ on $\tilde{\alpha}^* \times \alpha$. This shows $\tilde{\nu}_\lambda^* = \tilde{\lambda} \circ (\tilde{\phi}^*)^{-1}$ by Proposition 2.2.

By Proposition 2.4, we see that the measure

$$\tilde{\mu}(d\omega dx) = \sum_{\substack{s \in \alpha \\ \chi \in \{\tau_1, \tau_2\}}} \frac{\tilde{\rho}^*(\tilde{s}^\chi)}{\tilde{\lambda}(\tilde{s}^\chi)} 1_{\tilde{s}^\chi}(\omega, x) \mathbb{P}(d\omega) dx = \sum_{s \in \alpha} \frac{\rho(s)}{\lambda(s)} 1_s(x) \mathbb{P}(d\omega) dx \quad (5.25)$$

is a unique $\tilde{\lambda}$ -a.c. σ -finite \tilde{T} -invariant measure $\tilde{\mu}$ and the dynamical system $(\tilde{X}, \tilde{\mu}, \tilde{T})$ is conservative ergodic. The remainder of the proof is therefore obvious. \square

6 Limit theorems

6.1 Hata map

We now proceed to prove Theorem 1.2 for the Hata map with $p = 1/2$.

Proof of Theorem 1.2. We reduce to Thaler–Zweimüller’s result [36]. Let us check that the assumptions of Theorems 3.1 and 3.2 of [36] are all satisfied. We set

$$J := \left(\frac{1}{4}, \frac{3}{4} \right) = I_0^- + I_0^+, \quad R^- := \left(0, \frac{1}{4} \right) = \sum_{k=1}^{\infty} I_k^-, \quad R^+ := \left(\frac{3}{4}, 1 \right) = \sum_{k=1}^{\infty} I_k^+ \quad (6.1)$$

so that $J + R^- + R^+ = X$, where the identities hold mod λ . By the fact that

$$\tau_1^{-1} R^- = \left(0, \frac{1}{8} \right) \subset R^-, \quad \tau_2^{-1} R^- = \left(0, \frac{5}{8} \right) \subset J + R^- \quad \text{mod } \lambda \quad (6.2)$$

and by symmetry, we see that \tilde{J} dynamically separates \tilde{R}^- and \tilde{R}^+ , in the sense that

$$(\omega, x) \in \tilde{R}^\pm \text{ and } \tilde{T}^2(\omega, x) \in \tilde{R}^\mp \text{ imply } \tilde{T}(\omega, x) \in \tilde{J}. \quad (6.3)$$

We may call J the *junction* and R^+ and R^- the *rays*. We denote the first return time by

$$\varphi_{\tilde{J}}(\omega, x) := \inf \left\{ n \geq 1 : \tilde{T}^n(\omega, x) \in \tilde{J} \right\} \quad (6.4)$$

and set

$$\tilde{J}_0 := \tilde{J}, \quad \tilde{J}_n := \tilde{J}^c \cap \{\varphi_{\tilde{J}} = n\} \quad (n = 1, 2, \dots). \quad (6.5)$$

(Note that \tilde{J}_n cannot be of the form $\tilde{J}_n = \Omega \times J_n$.) For $N = 1, 2, \dots$, we denote the wandering rate of \tilde{J} by

$$w_N(\tilde{J}) := \sum_{n=0}^{N-1} \tilde{\mu}(\tilde{J} \cap \{\varphi_{\tilde{J}} > n\}) = \int_{\tilde{J}} \left(\sum_{n=0}^{N-1} \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n} \right) d\tilde{\mu} \quad (6.6)$$

and the wandering rate of \tilde{J} through \tilde{R}^\pm by

$$w_N(\tilde{J}, \tilde{R}^\pm) := \sum_{n=0}^{N-1} \tilde{\mu}(\tilde{J} \cap \tilde{T}^{-1} \tilde{R}^\pm \cap \{\varphi_{\tilde{J}} > n\}) = \tilde{\mu}(\tilde{J} \cap \tilde{T}^{-1} \tilde{R}^\pm) + \sum_{n=1}^{N-1} \tilde{\mu}(\tilde{J}_n \cap \tilde{R}^\pm). \quad (6.7)$$

Recall that φ_k^W is the first hitting time of k for the simple symmetric random walk $(\{W_n\}, \mathbb{P})$, which has been introduced in (4.18). We prove the following lemma about the Perron–Frobenius operator.

Lemma 6.1. *For $n \geq 1$, it holds that*

$$\tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n \cap \tilde{R}^\pm}(\omega, x) = c_n 1_{I_0^\pm}(x), \quad \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n}(\omega, x) = c_n 1_J(x), \quad \tilde{\lambda}\text{-a.e.}, \quad (6.8)$$

where

$$c_n := \sum_{k=1}^{\infty} (2^{k+1} - 1) 2^{-k} \mathbb{P}(\varphi_{-k}^W = n). \quad (6.9)$$

Proof. By symmetry and by $\tilde{J}_n \subset \tilde{R}^+ + \tilde{R}^-$, it suffices to show

$$\int_{\tilde{X}} g_1(\omega) g_2(x) \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n \cap \tilde{R}^-}(\omega, x) \mathbb{P}(d\omega) \mu(dx) = c_n \mathbb{E}[g_1] \int_{I_0^-} g_2(y) \mu(dy) \quad (6.10)$$

for all $g_1 \in L^\infty(\mathbb{P})$ and $g_2 \in L^\infty(\mu)$. For $k \geq 1$, we have

$$\tilde{J}_n \cap \tilde{I}_k^- = \tilde{I}_k^- \cap \tilde{T}^{-1} \tilde{R}^- \cap \dots \cap \tilde{T}^{-(n-1)} \tilde{R}^- \cap \tilde{T}^{-n} \tilde{J} = \tilde{I}_k^- \cap \{\varphi_{-k}^W = n\}, \quad (6.11)$$

where φ_{-k}^W can be regarded as the first hitting time of 0 for the random walk starting from k . The left hand side of (6.10) equals to

$$\int_{\tilde{X}} g_1(\theta^n \omega) g_2(\omega_n \circ \omega_{n-1} \circ \cdots \circ \omega_1(x)) 1_{\tilde{J}_n}(\omega, x) 1_{R^-}(x) \mathbb{P}(d\omega) \mu(dx) \quad (6.12)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_{\tilde{X}} g_2(\omega_n \circ \omega_{n-1} \circ \cdots \circ \omega_1(x)) 1_{\{\varphi_{-k}^W = n\}}(\omega) 1_{I_k^-}(x) \mathbb{P}(d\omega) \mu(dx) \quad (6.13)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_X g_2(2^k x) 1_{I_k^-}(x) \mu(dx) \mathbb{P}(\varphi_{-k}^W = n) \quad (6.14)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_X g_2(y) 1_{I_0^-}(y) \frac{\rho(I_k^-)}{\lambda(I_k^-)} \cdot \frac{\lambda(I_0^-)}{\rho(I_0^-)} \cdot 2^{-k} \mu(dy) \mathbb{P}(\varphi_{-k}^W = n), \quad (6.15)$$

which equals to the right hand side of (6.10). Here, the equality (6.14) can be obtained by the same argument as that of the proof of (4.12) of Theorem 4.1. \square

The proof of Theorem 1.2 is completed by proving the following lemma.

Lemma 6.2. *The following assertions hold as $N \rightarrow \infty$:*

$$(i) \quad w_N(\tilde{J}) \sim \sqrt{\frac{2}{\pi}} N^{1/2} \quad \text{and} \quad w_N(\tilde{J}, \tilde{R}^\pm) \sim \frac{1}{2} \sqrt{\frac{2}{\pi}} N^{1/2};$$

$$(ii) \quad \frac{1}{w_N(\tilde{J})} \sum_{n=0}^{N-1} \tilde{T}^{\wedge n} 1_{\tilde{J}_n}(\omega, x) \rightarrow 2 \cdot 1_J(x) \quad \text{uniformly on } \tilde{J};$$

$$(iii) \quad \frac{1}{w_N(\tilde{J}, \tilde{R}^\pm)} \sum_{n=0}^{N-1} \tilde{T}^{\wedge n} 1_{\tilde{J}_n \cap \tilde{R}^\pm}(\omega, x) \rightarrow 4 \cdot 1_{I_0^\pm}(x) \quad \text{uniformly on } \tilde{J}.$$

Here by $a_N \sim b_N$ we mean $a_N/b_N \rightarrow 1$.

Proof. Let $0 < z < 1$. By the strong Markov property and stationarity, we have

$$\mathbb{E} z^{\varphi_{-1}^W} = \frac{z}{2} \mathbb{E} z^{\varphi_{-2}^W} + \frac{z}{2} = \frac{z}{2} (\mathbb{E} z^{\varphi_{-1}^W})^2 + \frac{z}{2}, \quad (6.16)$$

and so we obtain, for any $k \geq 1$,

$$\mathbb{E} z^{\varphi_{-k}^W} = \left(\mathbb{E} z^{\varphi_{-1}^W} \right)^k = \left(\frac{1 - \sqrt{1 - z^2}}{z} \right)^k. \quad (6.17)$$

Let us study the asymptotic behavior. As $z \uparrow 1$, we have

$$w := 1 - \mathbb{E} z^{\varphi_{-1}^W} = \frac{\sqrt{1-z}(\sqrt{1+z} - \sqrt{1-z})}{z} \sim \sqrt{2} \sqrt{1-z}. \quad (6.18)$$

The generating function of $\{c_n\}$ can be computed as

$$\sum_{n=1}^{\infty} c_n z^n = \sum_{k=1}^{\infty} (2^{k+1} - 1) 2^{-k} \sum_{n=1}^{\infty} z^n \mathbb{P}(\varphi_{-k}^W = n) \quad (6.19)$$

$$= \sum_{k=1}^{\infty} (2^{k+1} - 1) 2^{-k} (1-w)^k \quad (6.20)$$

$$= \frac{2(1-w)}{w} - \frac{1-w}{1+w} \sim \sqrt{\frac{2}{1-z}} \quad \text{as } z \uparrow 1. \quad (6.21)$$

By the Tauberian theorem (see [36, Proposition 4.2]), we obtain

$$\sum_{n=1}^{N-1} c_n \sim \frac{\sqrt{2}}{\Gamma(3/2)} N^{1/2} = 2\sqrt{\frac{2}{\pi}} N^{1/2} \quad \text{as } N \rightarrow \infty. \quad (6.22)$$

(i) By Lemma 6.1, we have

$$w_N(\tilde{J}, \tilde{R}^{\pm}) = \tilde{\mu}(\tilde{J} \cap \tilde{T}^{-1} \tilde{R}^{\pm}) + \sum_{n=1}^{N-1} \tilde{\mu}(\tilde{J}_n \cap \tilde{R}^{\pm}) \quad (6.23)$$

$$= \tilde{\mu}(\tilde{J} \cap \tilde{T}^{-1} \tilde{R}^{\pm}) + \sum_{n=1}^{N-1} \int_{\tilde{X}} \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n \cap \tilde{R}^{\pm}}(\omega, x) d\tilde{\mu} \quad (6.24)$$

$$= \frac{1}{2} \mu\left(\left(\frac{1}{4}, \frac{5}{8}\right)\right) + \sum_{n=1}^{N-1} c_n \cdot \mu(I_0^{\pm}) \quad (6.25)$$

$$= \frac{3}{16} + \frac{1}{4} \sum_{n=0}^{N-1} c_n \sim \frac{1}{2} \sqrt{\frac{2}{\pi}} N^{1/2} \quad \text{as } N \rightarrow \infty. \quad (6.26)$$

We also obtain

$$w_N(\tilde{J}) = w_N(\tilde{J}, \tilde{R}^+) + w_N(\tilde{J}, \tilde{R}^-) + \tilde{\mu}(\tilde{J} \cap \tilde{T}^{-1} \tilde{J}) \sim \sqrt{\frac{2}{\pi}} N^{1/2} \quad \text{as } N \rightarrow \infty. \quad (6.27)$$

(iii) Again by Lemma 6.1, we obtain

$$\frac{1}{w_N(\tilde{J}, \tilde{R}^{\pm})} \sum_{n=1}^{N-1} \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n \cap \tilde{R}^{\pm}}(\omega, x) = \frac{\sum_{n=1}^{N-1} c_n}{w_N(\tilde{J}, \tilde{R}^{\pm})} \cdot 1_{I_0^{\pm}}(x) \rightarrow 4 \cdot 1_{I_0^{\pm}}(x) \quad (6.28)$$

as $N \rightarrow \infty$, where the convergence occurs uniformly in (ω, x) .

(ii) This claim is obvious by (iii) and (i). \square

6.2 Pelikan map

Let us now prove Theorem 1.4 for the Pelikan map with $p = 1/2$.

Proof of Theorem 1.4. Let us check that the assumptions of Theorems 3.1 and 3.2 of [36] are all satisfied for

$$J := \left(\frac{1}{2}, 1\right) = I_0. \quad (6.29)$$

Let \tilde{J}_n 's and $w_N(\tilde{J})$ be those of (6.5) and (6.6), respectively.

The proof is completed by proving the following lemma.

Lemma 6.3. *For $n \geq 1$, it holds that*

$$\tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n}(\omega, x) = 2c_n 1_J(x), \quad \tilde{\lambda}\text{-a.e.}, \quad (6.30)$$

where c_n is given in (6.9). It also holds that

$$w_N(\tilde{J}) \sim \sqrt{\frac{2}{\pi}} N^{1/2} \quad \text{as } N \rightarrow \infty. \quad (6.31)$$

Proof. For any $g_1 \in L^\infty(\mathbb{P})$ and $g_2 \in L^\infty(\mu)$, we have

$$\int_{\tilde{X}} g_1(\omega) g_2(x) \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n}(\omega, x) \mathbb{P}(d\omega) \mu(dx) \quad (6.32)$$

$$= \int_{\tilde{X}} g_1(\theta^n \omega) g_2(\omega_n \circ \omega_{n-1} \circ \cdots \circ \omega_1(x)) 1_{\tilde{J}_n}(\omega, x) \mathbb{P}(d\omega) \mu(dx) \quad (6.33)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_{\tilde{X}} g_2(\omega_n \circ \omega_{n-1} \circ \cdots \circ \omega_1(x)) 1_{\{\varphi_{-k}^W = n\}}(\omega) 1_{I_k}(x) \mathbb{P}(d\omega) \mu(dx) \quad (6.34)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_X g_2(2^k x) 1_{I_k}(x) \mu(dx) \mathbb{P}(\varphi_{-k}^W = n) \quad (6.35)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_X g_2(y) 1_{I_0}(y) \frac{\rho(I_k)}{\lambda(I_k)} \cdot \frac{\lambda(I_0)}{\rho(I_0)} \cdot 2^{-k} \mu(dy) \mathbb{P}(\varphi_{-k}^W = n), \quad (6.36)$$

which yields (6.30). We thus obtain

$$w_N(\tilde{J}) = \int_{\tilde{J}} \sum_{n=0}^{N-1} \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n} d\tilde{\mu} = \sum_{n=0}^{N-1} 2c_n \mu(J) = \frac{1}{2} \sum_{n=0}^{N-1} c_n, \quad (6.37)$$

which yields (6.31). □

6.3 mBGI map

Let us now prove Theorem 1.6 for the mBGI map with $p = 2/3$.

Proof of Theorem 1.6. We take

$$J := \left(\frac{1}{4}, \frac{3}{4}\right) = I_0^- + I_0^+. \quad (6.38)$$

Using the following lemma, we can complete the proof of Theorem 1.6 in the same way as that of the proof of Theorem 1.4.

Lemma 6.4. *For $n \geq 1$, it holds that*

$$\tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n}(\omega, x) = c'_n 1_J(x), \quad \tilde{\lambda}\text{-a.e.}, \quad (6.39)$$

where

$$c'_n := \sum_{k=1}^{\infty} \frac{3\sqrt{2}}{16} \left(\frac{8}{3} + \frac{1}{3} \cdot \left(-\frac{1}{2}\right)^k - \left(\frac{1}{2}\right)^k \right) \mathbb{P}(\varphi_{-k}^{W'} = n). \quad (6.40)$$

It also holds that

$$w_N(\tilde{J}) \sim \sqrt{\frac{2}{\pi}} N^{1/2} \quad \text{as } N \rightarrow \infty. \quad (6.41)$$

Proof. Recall that $\varphi_k^{W'}$ is the first hitting time of k for the simple symmetric random walk $(\{W'_n\}, \mathbb{P})$, which has been introduced in (4.31). For $k \geq 1$, we have

$$\tilde{J}_n \cap (\tilde{I}_k^- + \tilde{I}_k^+) = (\tilde{I}_k^- + \tilde{I}_k^+) \cap \tilde{T}^{-1} \tilde{J}^c \cap \dots \cap \tilde{T}^{-(n-1)} \tilde{J}^c \cap \tilde{T}^{-n} \tilde{J} \quad (6.42)$$

$$= (\tilde{I}_k^- + \tilde{I}_k^+) \cap \{\varphi_{-k}^{W'} = n\}. \quad (6.43)$$

For any $g_1 \in L^\infty(\mathbb{P})$ and $g_2 \in L^\infty(\mu)$, we have

$$\int_{\tilde{X}} g_1(\omega) g_2(x) \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n}(\omega, x) \mathbb{P}(d\omega) \mu(dx) \quad (6.44)$$

$$= \int_{\tilde{X}} g_1(\theta^n \omega) g_2(\omega_n \circ \omega_{n-1} \circ \dots \circ \omega_1(x)) 1_{\tilde{J}_n}(\omega, x) \mathbb{P}(d\omega) \mu(dx) \quad (6.45)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_{\tilde{X}} g_2(\omega_n \circ \omega_{n-1} \circ \dots \circ \omega_1(x)) 1_{\{\varphi_{-k}^{W'} = n\}}(\omega) 1_{I_k^- + I_k^+}(x) \mathbb{P}(d\omega) \mu(dx) \quad (6.46)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_{\Omega} \mathbb{P}(d\omega) 1_{\{\varphi_{-k}^{W'} = n\}}(\omega) \int_X g_2(\omega_n \circ \omega_{n-1} \circ \dots \circ \omega_1(x)) 1_{I_k^- + I_k^+}(x) \mu(dx). \quad (6.47)$$

Let us write

$$r(k) := \frac{\rho(I_k^\pm)}{\lambda(I_k^\pm)} = \frac{8}{3} \cdot 2^k + \frac{1}{3} \cdot (-1)^k - 1, \quad k = 0, 1, 2, \dots, \quad (6.48)$$

where $\rho(I_k^\pm)$ has been given in (1.28). Noting that, for any $g \in L^\infty(\mathbb{P})$, we have

$$\int_X g(\tau_1(x)) 1_{I_k^- + I_k^+}(x) \mu(dx) = \int_X g(y) 1_{I_{k-1}^- + I_{k-1}^+}(y) \frac{r(k)}{r(k-1)} \cdot \frac{1}{2} \mu(dy) \quad (k \geq 1), \quad (6.49)$$

$$\int_X g(\tau_2(x)) 1_{I_{k+2}^- + I_{k+2}^+}(x) \mu(dx) = \int_X g(y) 1_{I_{k+2}^- + I_{k+2}^+}(y) \frac{r(k)}{r(k+2)} \cdot 4 \mu(dy) \quad (k \geq 0). \quad (6.50)$$

We thus obtain

$$(6.47) = \mathbb{E}[g_1] \sum_{k=1}^{\infty} \int_{\Omega} \mathbb{P}(d\omega) 1_{\{\varphi_{-k}^{W'}=n\}}(\omega) \int_X g_2(y) 1_{I_0^-+I_0^+}(y) \frac{r(k)}{r(0)} \cdot 2^{-k} \mu(dy) \quad (6.51)$$

$$= \mathbb{E}[g_1] \sum_{k=1}^{\infty} \frac{r(k)}{r(0)} \cdot 2^{-k} \mathbb{P}(\varphi_{-k}^{W'} = n) \int_X g_2(y) 1_J(y) \mu(dy), \quad (6.52)$$

which yields (6.39).

For $0 < z < 1$, we have

$$\mathbb{E}z^{\varphi_{-1}^{W'}} = \frac{z}{3} \mathbb{E}z^{\varphi_{-3}^{W'}} + \frac{2z}{3} = \frac{z}{3} \left(\mathbb{E}z^{\varphi_{-1}^{W'}} \right)^3 + \frac{2z}{3}. \quad (6.53)$$

Let $w := 1 - \mathbb{E}z^{\varphi_{-1}^{W'}}$. We then have

$$\frac{1-w}{z} = \frac{1}{3}(1-w)^3 + \frac{2}{3} = 1-w+w^2-\frac{1}{3}w^3 \quad (6.54)$$

and

$$w^2 - \frac{1}{3}w^3 = \frac{1-w}{z} - (1-w) = \frac{1-w}{z} \cdot (1-z). \quad (6.55)$$

Since $w \downarrow 0$ as $z \uparrow 1$, we obtain

$$w = \sqrt{w^2} = \sqrt{\frac{1}{1-w/3} \cdot \frac{1-w}{z} \cdot (1-z)} \sim \sqrt{1-z} \quad \text{as } z \uparrow 1. \quad (6.56)$$

The generating function of $\{c'_n\}$ can be computed as

$$\sum_{n=1}^{\infty} c'_n z^n = \sum_{k=1}^{\infty} \frac{3\sqrt{2}}{16} \left(\frac{8}{3} + \frac{1}{3} \cdot \left(-\frac{1}{2}\right)^k - \left(\frac{1}{2}\right)^k \right) \sum_{n=1}^{\infty} z^n \mathbb{P}(\varphi_{-k}^{W'} = n) \quad (6.57)$$

$$= \sum_{k=1}^{\infty} \frac{3\sqrt{2}}{16} \left(\frac{8}{3} + \frac{1}{3} \cdot \left(-\frac{1}{2}\right)^k - \left(\frac{1}{2}\right)^k \right) (1-w)^k \quad (6.58)$$

$$= \frac{3\sqrt{2}}{16} \left(\frac{8}{3} \cdot \frac{1-w}{w} + \frac{1}{3} \cdot \frac{1-w}{3-w} - \frac{1-w}{1+w} \right) \sim \frac{1}{2} \sqrt{\frac{2}{1-z}} \quad \text{as } z \uparrow 1. \quad (6.59)$$

By the Tauberian theorem (see [36, Proposition 4.2]), we obtain

$$\sum_{n=1}^{N-1} c'_n \sim \frac{\sqrt{2}}{2} \cdot \frac{1}{\Gamma(3/2)} N^{1/2} = \sqrt{\frac{2}{\pi}} N^{1/2} \quad \text{as } N \rightarrow \infty. \quad (6.60)$$

We thus obtain

$$w_N(\tilde{J}) = \int_{\tilde{J}} \sum_{n=0}^{N-1} \tilde{T}_{\tilde{\mu}}^{\wedge n} 1_{\tilde{J}_n} d\tilde{\mu} = \sum_{n=0}^{N-1} c'_n \mu(J) = \sum_{n=0}^{N-1} c'_n, \quad (6.61)$$

which yields (6.41). □

Remark 6.5. One may expect that we could prove the arcsine law by taking

$$R^- = \left(0, \frac{1}{4}\right) = \sum_{k=1}^{\infty} I_k^-, \quad R^+ = \left(\frac{3}{4}, 1\right) = \sum_{k=1}^{\infty} I_k^+. \quad (6.62)$$

Unfortunately, we fail to do it, because $\tau_2^{-1}R^+ \supset I_k^-$ for $k \geq 2$, so that \tilde{J} does not dynamically separate \tilde{R}^- and \tilde{R}^+ .

References

- [1] J. Aaronson. The asymptotic distributional behaviour of transformations preserving infinite measures. *J. Analyse Math.*, 39:203–234, 1981.
- [2] J. Aaronson. Random f -expansions. *Ann. Probab.*, 14(3):1037–1057, 1986.
- [3] J. Aaronson. *An introduction to infinite ergodic theory*, volume 50 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [4] N. Abbasi, M. Gharaei, and A. J. Homburg. Iterated function systems of logistic maps: synchronization and intermittency. *Nonlinearity*, 31(8):3880–3913, 2018.
- [5] T. Akimoto and Y. Aizawa. Subexponential instability in one-dimensional maps implies infinite invariant measure. *Chaos*, 20(3):033110, 7, 2010.
- [6] P. Ashwin, P. J. Aston, and M. Nicol. On the unfolding of a blowout bifurcation. *Phys. D*, 111(1-4):81–95, 1998.
- [7] W. Bahsoun and C. Bose. Mixing rates and limit theorems for random intermittent maps. *Nonlinearity*, 29(4):1417–1433, 2016.
- [8] W. Bahsoun, C. Bose, and Y. Duan. Decay of correlation for random intermittent maps. *Nonlinearity*, 27(7):1543–1554, 2014.
- [9] W. Bahsoun, C. Bose, and Y. Duan. Rigorous pointwise approximations for invariant densities of non-uniformly expanding maps. *Ergodic Theory Dynam. Systems*, 35(4):1028–1044, 2015.
- [10] A. Boyarsky, P. Góra, and M. S. Islam. Randomly chosen chaotic maps can give rise to nearly ordered behavior. *Phys. D*, 210(3-4):284–294, 2005.
- [11] M. Campanino and S. Isola. Statistical properties of long return times in type I intermittency. *Forum Math.*, 7(3):331–348, 1995.
- [12] M. Campanino and S. Isola. Infinite invariant measures for non-uniformly expanding transformations of $[0, 1]$: weak law of large numbers with anomalous scaling. *Forum Math.*, 8(1):71–92, 1996.

- [13] P. Collet and P. Ferrero. Some limit ratio theorem related to a real endomorphism in case of a neutral fixed point. *Ann. Inst. H. Poincaré Phys. Théor.*, 52(3):283–301, 1990.
- [14] P. Collet and A. Galves. Statistics of close visits to the indifferent fixed point of an interval map. *J. Statist. Phys.*, 72(3-4):459–478, 1993.
- [15] P. Collet, A. Galves, and B. Schmitt. Unpredictability of the occurrence time of a long laminar period in a model of temporal intermittency. *Ann. Inst. H. Poincaré Phys. Théor.*, 57(3):319–331, 1992.
- [16] M. Gharaei and A. J. Homburg. Random interval diffeomorphisms. *Discrete Contin. Dyn. Syst. Ser. S*, 10(2):241–272, 2017.
- [17] G. Hata. Arcsine law for a piecewise linear random map. *Master Thesis, Kyoto University*, 2019.
- [18] A. J. Homburg, C. Kalle, M. Ruziboev, E. Verbitskiy, and B. Zeegers. Critical intermittency in random interval maps. 2021. Preprint, arXiv:2107.02556.
- [19] H. Hu. Decay of correlations for piecewise smooth maps with indifferent fixed points. *Ergodic Theory Dynam. Systems*, 24(2):495–524, 2004.
- [20] T. Inoue. First return maps of random maps and invariant measures. *Nonlinearity*, 33(1):249–275, 2020.
- [21] O. Kallenberg. *Foundations of modern probability*. Probability and its Applications (New York). Springer-Verlag, New York, second edition, 2002.
- [22] C. Liverani, B. Saussol, and S. Vaienti. A probabilistic approach to intermittency. *Ergodic Theory Dynam. Systems*, 19(3):671–685, 1999.
- [23] P. Manneville. Intermittency, self-similarity and $1/f$ spectrum in dissipative dynamical systems. *J. Physique*, 41(11):1235–1243, 1980.
- [24] M. Mori. On the intermittency of a piecewise linear map (Takahashi model). *Tokyo J. Math.*, 16(2):411–428, 1993.
- [25] S. Munday and G. Knight. Escape rate scaling in infinite measure preserving systems. *J. Phys. A*, 49(8):085101, 12, 2016.
- [26] J. R. Norris. *Markov chains*, volume 2 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 1998. Reprint of 1997 original.
- [27] T. Owada and G. Samorodnitsky. Functional central limit theorem for heavy tailed stationary infinitely divisible processes generated by conservative flows. *Ann. Probab.*, 43(1):240–285, 2015.
- [28] S. Pelikan. Invariant densities for random maps of the interval. *Trans. Amer. Math. Soc.*, 281(2):813–825, 1984.

- [29] M. Pollicott and M. Yuri. Statistical properties of maps with indifferent periodic points. *Comm. Math. Phys.*, 217(3):503–520, 2001.
- [30] Y. Pomeau and P. Manneville. Intermittent transition to turbulence in dissipative dynamical systems. *Comm. Math. Phys.*, 74(2):189–197, 1980.
- [31] Y. Sato and R. Klages. Anomalous diffusion in random dynamical systems. *Phys. Rev. Lett.*, 122(17):174101, 6, 2019.
- [32] T. Sera. Functional limit theorem for occupation time processes of intermittent maps. *Nonlinearity*, 33(3):1183–1217, 2020.
- [33] T. Sera and K. Yano. Multiray generalization of the arcsine laws for occupation times of infinite ergodic transformations. *Trans. Amer. Math. Soc.*, 372(5):3191–3209, 2019.
- [34] Y. Takahashi. Power spectrum and Fredholm determinant related to intermittent chaos. In *Stochastic processes in physics and engineering (Bielefeld, 1986)*, volume 42 of *Math. Appl.*, pages 357–379. Reidel, Dordrecht, 1988.
- [35] M. Thaler. A limit theorem for sojourns near indifferent fixed points of one-dimensional maps. *Ergodic Theory Dynam. Systems*, 22(4):1289–1312, 2002.
- [36] M. Thaler and R. Zweimüller. Distributional limit theorems in infinite ergodic theory. *Probab. Theory Related Fields*, 135(1):15–52, 2006.
- [37] H. Toyokawa. On the existence of a σ -finite acim for a random iteration of intermittent Markov maps with uniformly contractive part. *Stoch. Dyn.*, 21(3):Paper No. 2140003, 14, 2021.
- [38] R. Zweimüller. Infinite measure preserving transformations with compact first regeneration. *J. Anal. Math.*, 103:93–131, 2007.