

SEPARATING TOPOLOGICAL RECURRENCE FROM MEASURABLE RECURRENCE: EXPOSITION AND EXTENSION OF KRIZ'S EXAMPLE

JOHN T. GRIESMER

ABSTRACT. We prove that for every infinite set $E \subset \mathbb{Z}$, there is a set $S \subset E - E$ which is a set of topological recurrence and not a set of measurable recurrence. This extends a result of Igor Kriz, proving that there is a set of topological recurrence which is not a set of measurable recurrence. Our construction follows Kriz's closely, and this paper can be considered an exposition of the original argument.

1. STRUCTURE OF DIFFERENCE SETS

We write \mathbb{Z} for the group of integers. If $A \subset \mathbb{Z}$, we write $A - A$ for the *difference set*, defined to be $\{a_1 - a_2 : a_1, a_2 \in A\}$. If $A, B \subset \mathbb{Z}$, we write $A + B$ for the *sumset*, $\{a + b : a \in A, b \in B\}$. The *upper asymptotic density* (or just *upper density*) of A is $\bar{d}(A) := \limsup_{n \rightarrow \infty} \frac{|A \cap \{1, \dots, n\}|}{n}$. If the limit exists, we write $d(A)$ in place of $\bar{d}(A)$.

1.1. Kriz's theorem. We say that $S \subset \mathbb{Z}$ is a *set of density recurrence* (or “is *density recurrent*”) if for all $A \subset \mathbb{Z}$, with $\bar{d}(A) > 0$, there exists $a, b \in A$ such that $b - a \in S$. In other words, $(A - A) \cap S \neq \emptyset$ whenever $\bar{d}(A) > 0$. The set $\{n^2 : n \in \mathbb{N}\}$ of perfect squares is density recurrent, as proved by Furstenberg ([9], via ergodic theory) and Sárközy ([18], using the circle method). This result, along with earlier work by Bogoliouboff [4] and Følner [5], motivated further investigation into the structure of difference sets.

Another result influencing this investigation is van der Waerden's theorem on arithmetic progressions [20]: if \mathbb{N} is partitioned into finitely many sets A_1, \dots, A_r , then at least one of the sets A_j contains arithmetic progressions of every finite length. The “density version” of van der Waerden's theorem is Szemerédi's theorem [19], which says that if $\bar{d}(A) > 0$, then A contains arithmetic progressions of every finite length. These results suggested a possible deep connection between partition results and density results. To formalize this suggestion, we say set $S \subset \mathbb{Z}$ is a *set of chromatic recurrence* (or “is *chromatically recurrent*”) if for every partition $\{A_1, \dots, A_r\}$ of \mathbb{Z} into finitely many sets, there is a cell

Date: February 18, 2022.

A_j of the partition such that $(A_j - A_j) \cap S \neq \emptyset$. Upper density is finitely subadditive, so if \mathbb{Z} is partitioned into finitely many sets, then at least one of the cells has positive upper density, and its difference set therefore intersects every density recurrent set. Thus every density recurrent set is also chromatically recurrent. Bergelson asked in [2], and Furstenberg asked in [8], whether the converse holds: is every chromatically recurrent set also density recurrent? Igor Kriz gave a negative answer in the following theorem from [13]. The main purpose of this article is to give an expository proof.

Theorem 1.1. *There is a set of integers which is chromatically recurrent and not density recurrent.*

Readers familiar with [13] will see that the methods there can be combined with Lemma 6.2 below to prove the following generalization of Theorem 1.1. Our second purpose in this article is to give an explicit proof of this generalization.

Theorem 1.2. *If E is an infinite set of integers, then there is a subset $S \subset E - E$ such that S is chromatically recurrent and not density recurrent.*

Theorem 1.1 may be stated as “there is a set of topological recurrence which is not a set of measurable recurrence.” Likewise, Theorem 1.2 says that if $E \subset \mathbb{Z}$ is infinite, then there is subset of $E - E$ which is a set of topological recurrence and not a set of measurable recurrence; see [3] for a general discussion of the equivalences between various recurrence properties.

There are several proofs of Theorem 1.1 ([6],[16],[17],[21]), all of which follow the same broad outline as [13] and overcome the essential difficulties in a similar way. Hopefully our exposition will help readers solve some related open problems, or find a fundamentally different approach to Theorems 1.1 and 1.2.

1.2. Outline of the article. In §2 we prove a version of Theorem 1.1 for groups of the form $(\mathbb{Z}/2\mathbb{Z})^d$. This introduces one of the key ideas in a clearer setting.

In §3 we state and prove Lemma 3.6, one of two main lemmas required for the proof of Theorem 1.1. We then state the other main lemmas, Lemmas 3.7 and 3.8, and prove Theorems 1.1 and 1.2. The latter two lemmas are proved in §4, with arguments similar to those in §2, together with some known results on uniform distribution, presented in §7.

The proofs of Theorems 1.1 and 1.2 are mostly elementary. The only highly nontrivial result required, for both proofs, is a lower bound for chromatic numbers of Kneser graphs, discussed in §2.1. Theorem 1.2 uses a well known result of Weyl on uniform distribution

in \mathbb{T}^d ; we provide a proof in §7. We also use, without proof, the elementary binomial estimate $\lim_{n \rightarrow \infty} \binom{n}{\lfloor n/2 \rfloor} 2^{-n} = 0$.

1.3. Acknowledgements. This article summarizes lectures given by the author in an informal seminar with Nishant Chandgotia, Aryaman Jal, Abhishek Khetan, and Andy Parrish. I thank them for their interest and encouragement.

2. A MODEL SETTING

While Theorems 1.1 and 1.2 are about sets of integers, one of our main arguments is easier to develop in finite vector spaces over the field with two elements. Forrest takes a similar approach in [6], [7].

Let Γ be a finite abelian group and $S \subset \Gamma$. We say that S is

- *δ -density recurrent* if $(A - A) \cap S \neq \emptyset$ for all $A \subset \Gamma$ having $|A| > \delta|\Gamma|$.
- *δ -nonrecurrent* if there exists $A \subset \Gamma$ having $|A| > \delta|\Gamma|$ such that $(A - A) \cap S = \emptyset$.
- *k -chromatically recurrent* if for every partition of Γ into k sets A_1, A_2, \dots, A_k , the intersection $(A_j - A_j) \cap S$ is nonempty for some $j \leq k$.

If $d \in \mathbb{N}$, we write \mathbb{F}_2^d for the group $(\mathbb{Z}/2\mathbb{Z})^d$, the product of d copies of $\mathbb{Z}/2\mathbb{Z}$. Elements of \mathbb{F}_2^d will be written as $\mathbf{x} = (x_1, \dots, x_d)$, where $x_j \in \{0, 1\}$. The remainder of this section is dedicated to proving the following analogue of Theorem 1.1.

Theorem 2.1. *Let $\delta < \frac{1}{2}$ and $k \in \mathbb{N}$. For all sufficiently large d , there exists $S \subset \mathbb{F}_2^d$ such that S is k -chromatically recurrent and δ -nonrecurrent.*

The next section summarizes background for the proof of Theorem 2.1.

2.1. Cayley graphs and Kneser graphs. We adopt the usual terminology from graph theory: a *graph* \mathcal{G} is a set V , whose elements are called *vertices*, together with a set E of unordered pairs of elements of V , called *edges*. A *vertex coloring* of \mathcal{G} with k colors (briefly, a k -coloring) is a function $f : V \rightarrow \{1, \dots, k\}$. We say that f is *proper* if $f(v_1) \neq f(v_2)$ for every edge $\{v_1, v_2\} \in E$. The *chromatic number* of \mathcal{G} is the smallest k such that there is a proper k -coloring of G .

Let $r \leq n \in \mathbb{N}$. The *Kneser graph* $KG(n, r)$ is the graph whose vertices are the r -element subsets of $\{1, \dots, n\}$, with two vertices $A, B \subset \{1, \dots, n\}$ joined by an edge if and only if $A \cap B = \emptyset$. Note that $KG(n, r)$ has no edges if $2r > n$. M. Kneser proved that the chromatic number of $KG(n, r)$ is no greater than $n - 2r + 2$, and conjectured that this is the correct value when $2r \leq n$. Lovász proved Kneser's conjecture in [14],

and Bárány [1] provided a more elementary proof. An even shorter proof was provided by Greene in [10]. Matousek's book [15] gives a detailed exposition of these proofs and subsequent developments of the techniques they introduced.

Theorem 2.2 ([14]). *If $2r \leq n \in \mathbb{N}$, then the chromatic number of $KG(n, r)$ is $n - 2r + 2$.*

Examination of our proofs shows that we need only the following corollary of Theorem 2.2.

Corollary 2.3. *If $n_k \rightarrow \infty$, $r_k \rightarrow \infty$, and $r_k^2/n_k \rightarrow 0$, then the chromatic number of $KG(2n_k + r_k, n_k)$ tends to ∞ .*

Surprisingly, there are no known proofs of nontrivial lower bounds for the chromatic number of $KG(n, r)$, besides those that prove Theorem 2.2. A substantially different proof of Corollary 2.3 could lead to progress on problems in recurrence.

Given an abelian group Γ and a subset $S \subset \Gamma$, the *Cayley graph based on S* , denoted $\text{Cay}(S)$, is the graph whose vertex set is Γ , with two vertices x, y joined by an edge if $x - y \in S$ or $y - x \in S$. It follows immediately from the definitions that S is k -chromatically recurrent if and only if the chromatic number of $\text{Cay}(S)$ is strictly greater than k . To prove S is k -chromatically recurrent, it therefore suffices to prove that the chromatic number of $\text{Cay}(S)$ is at least $k + 1$.

2.2. Hamming balls in \mathbb{F}_2^d . For $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{F}_2^d$, we define

$$w(\mathbf{x}) := |\{j \leq d : x_j \neq 0\}|.$$

Given $k \leq d$ and $\mathbf{y} \in \mathbb{F}_2^d$, the *Hamming ball of radius k around \mathbf{y}* is

$$H_k(\mathbf{y}) := \{\mathbf{x} \in \mathbb{F}_2^d : w(\mathbf{y} - \mathbf{x}) \leq k\}.$$

So $H_k(\mathbf{y})$ is the set of $\mathbf{x} = (x_1, \dots, x_d)$ such that $x_j \neq y_j$ for at most k coordinates j .

Let $\mathbf{0} = (0, \dots, 0)$ and $\mathbf{1} = (1, \dots, 1) \in \mathbb{F}_2^d$. The following identities are easy to verify from the definitions:

$$(2.1) \quad H_k(\mathbf{0}) - H_k(\mathbf{0}) = H_{2k}(\mathbf{0}),$$

$$(2.2) \quad H_k(\mathbf{0}) \cap H_r(\mathbf{1}) = \emptyset \text{ iff } k + r < d.$$

Lemma 2.4. *Let $k \in \mathbb{N}$ and $\delta < \frac{1}{2}$. If $d \in \mathbb{N}$ is sufficiently large, then $H_k(\mathbf{1}) \subset \mathbb{F}_2^d$ is δ -nonrecurrent.*

We will use the following well known estimates on binomial coefficients: setting $M_d := \max_{0 \leq j \leq d} \binom{d}{j}$, we have $\lim_{d \rightarrow \infty} M_d/2^d = 0$. The identities $\binom{d}{j} = \binom{d}{d-j}$ and $\sum_{j=0}^d \binom{d}{j} = 2^d$ then imply that for fixed $k \in \mathbb{N}$ and $\delta < \frac{1}{2}$, we have the following for all large enough d :

$$(2.3) \quad \sum_{j=0}^{\lfloor d/2 \rfloor - k} \binom{d}{j} > \delta 2^d.$$

Proof of Lemma 2.4. Fix $k \in \mathbb{N}$ and $\delta < \frac{1}{2}$. Choose d to be sufficiently large that (2.3) holds. Let $A = H_{\lfloor d/2 \rfloor - k}(\mathbf{0})$, so that A is the set of elements $(x_1, \dots, x_d) \in \mathbb{F}_2^d$ having at most $\lfloor d/2 \rfloor - k$ entries equal to 1. Now $|A|$ is given by the sum in (2.3), so $|A| > \delta |\mathbb{F}_2^d|$. Equation (2.1) implies $A - A = H_{2(\lfloor d/2 \rfloor - k)}(\mathbf{0})$, which is disjoint from $H_k(\mathbf{1})$ by (2.2). So we have shown that $H_k(\mathbf{1})$ is δ -nonrecurrent. \square

Lemma 2.5. *Let $k, d \in \mathbb{N}$ with $2k \leq d$. The Cayley graph $\text{Cay}(H_{2k+1}(\mathbf{1}))$ in \mathbb{F}_2^d contains a subgraph isomorphic to $KG(d, \lfloor d/2 \rfloor - k)$. Consequently, $H_{2k+1}(\mathbf{1})$ is $2k$ -chromatically recurrent.*

Proof. Let \mathcal{G} denote the Cayley graph $\text{Cay}(H_{2k+1}(\mathbf{1}))$ and let $r = \lfloor d/2 \rfloor - k$. To each $C \subset \{1, \dots, d\}$ having cardinality r , we will assign an element $\mathbf{x}_C \in \mathbb{F}_2^d$, and we will show that if such C, C' are disjoint (meaning they are joined by an edge in $KG(d, r)$), then $\mathbf{x}_C - \mathbf{x}_{C'} \in H_{2k+1}(\mathbf{1})$ (meaning \mathbf{x}_C and $\mathbf{x}_{C'}$ are joined by an edge in \mathcal{G}). The set of \mathbf{x}_C so chosen will thereby determine a subgraph of $\text{Cay}(H_{2k+1}(\mathbf{1}))$ isomorphic to $KG(d, r)$.

For each $C \subset \{1, \dots, d\}$ having $|C| = r$, we let $\mathbf{x}_C := 1_C$. This is the characteristic function of C , viewed as an element of \mathbb{F}_2^d , so \mathbf{x}_C has exactly r entries equal to 1. Now if $C \cap C' = \emptyset$, then $\mathbf{x}_C - \mathbf{x}_{C'}$ has exactly $2r$ entries equal to 1. Since $2r \geq d - (2k + 1)$, this means $\mathbf{x}_C - \mathbf{x}_{C'} \in H_{2k+1}(\mathbf{1})$, so that \mathbf{x}_C and $\mathbf{x}_{C'}$ are joined by an edge of \mathcal{G} .

According to Theorem 2.2, the chromatic number of $KG(d, r)$ is at least $d - 2r + 2$. With $r = \lfloor d/2 \rfloor - k$, we get $d - 2r + 2 \geq 2k + 1$, so \mathcal{G} has chromatic number at least $2k + 1$, which implies that H_{2k+1} is $2k$ -chromatically recurrent. \square

Proof of Theorem 2.1. Fix $\delta \in (0, \frac{1}{2})$ and $k \in \mathbb{N}$. Combining Lemmas 2.4 and 2.5, we get that for all sufficiently large d , the Hamming ball $H_{2k+1}(\mathbf{1}) \subset \mathbb{F}_2^d$ is δ -nonrecurrent and k -chromatically recurrent. \square

3. PROOF OF THEOREMS 1.1 AND 1.2

3.1. Assembling finite pieces. We now return to \mathbb{Z} and prove Theorems 1.1 and 1.2. These theorems involve sets which are not density recurrent, but do have some other

recurrence property. Lemma 3.6 provides a general procedure for building such sets from finite pieces. To make this idea precise, we need the following definitions.

Let $S \subset \mathbb{Z}$. We say that S is

- δ -density recurrent if $(A - A) \cap S \neq \emptyset$ for all $A \subset \mathbb{Z}$ having $\bar{d}(A) > \delta$.
- δ -nonrecurrent if there exists $A \subset \mathbb{Z}$ having $\bar{d}(A) > \delta$ such that $(A - A) \cap S = \emptyset$. In other words, S is not δ -density recurrent.
- k -chromatically recurrent if for every partition of \mathbb{Z} into k sets A_1, A_2, \dots, A_k , the intersection $(A_j - A_j) \cap S$ is nonempty for some $j \leq k$.

Equivalently, S is k -chromatically recurrent if for every function $f : \mathbb{Z} \rightarrow \{1, \dots, k\}$, there exist $a, b \in \mathbb{Z}$ such that $f(a) = f(b)$ and $b - a \in S$.

Remark 3.1. Note that $(A - A) \cap S = \emptyset$ if and only if $A \cap (A + S) = \emptyset$. This equivalence will be used from time to time without comment. Likewise we will use the following trivial observations, true for all $A, B \subset \mathbb{Z}$ and $t \in \mathbb{Z}$.

- $(A - t) - (A - t) = A - A$.
- $(A \cap B) - t = (A - t) \cap (B - t)$.

If $m \in \mathbb{N}$, we write $[m]$ for the interval $\{0, 1, \dots, m - 1\}$ in \mathbb{Z} .

Definition 3.2. If $B \subset \mathbb{Z}$ is finite, $m \in \mathbb{N}$, and $B \subset [m]$, we say that (B, m) witnesses the δ -nonrecurrence of S if $|B| > \delta m$ and $B \cap (B + S) = \emptyset$, $B + S \subset [m]$, and $B + S + S \subset [m]$.

The condition $B + S + S \subset [m]$ may seem unmotivated, but it will be used in the proof of Lemma 3.6.

Lemma 3.3. *If $A \subset \mathbb{Z}$ and $m \in \mathbb{N}$, then there is a $t \in \mathbb{Z}$ such that $|A \cap ([m] + t)| \geq \bar{d}(A)m$.*

Proof. Let $m \in \mathbb{N}$ and let

$$\delta := \sup_{t \in \mathbb{Z}} \frac{|A \cap ([m] + t)|}{m},$$

so that $|A \cap ([m] + t)| \leq \delta m$ for every $t \in \mathbb{Z}$. Write $[N]$ as a union of $\lfloor N/m \rfloor$ disjoint intervals $I_1, \dots, I_{\lfloor N/m \rfloor}$ of length m , together with another (possibly empty) interval $I_{\lfloor N/m \rfloor + 1}$ of length at most m . Then $|A \cap I_j| \leq \delta m$ for every j , so

$$|A \cap [N]| \leq \sum_{j=1}^{\lfloor N/m \rfloor + 1} |A \cap I_j| \leq \lfloor N/m \rfloor \delta m + m \leq \delta N + m.$$

Then $\limsup_{N \rightarrow \infty} \frac{|A \cap [N]|}{N} \leq \delta$, meaning $\bar{d}(A) \leq \delta$. Since there are only finitely many possible values of $\frac{|A \cap ([m] + t)|}{m}$, the supremum is attained. \square

Lemma 3.4. *If $S \subset \mathbb{Z}$ is finite and δ -nonrecurrent, then for all sufficiently large m , there is a set $B \subset [m]$ such that (B, m) witnesses the δ -nonrecurrence of S .*

Proof. Assuming $\delta > 0$ and that S is finite and δ -nonrecurrent, there is set $A \subset \mathbb{Z}$ such that $\bar{d}(A) > \delta$ and $(A - A) \cap S = \emptyset$. Fix such an A . Let $k = \max\{|n| : n \in S\}$, and choose m_0 sufficiently large that $\bar{d}(A)m - 2k > \delta m$ whenever $m > m_0$.

Let $m > m_0$. We will find an interval of length m and a set $B \subset [m]$ such that (B, m) witnesses the δ -nonrecurrence of S . By Lemma 3.3, choose $t \in \mathbb{Z}$ so that $|A \cap ([m] + t)| \geq \bar{d}(A)m$. Then $|(A-t) \cap [m]| \geq \bar{d}(A)m$. Let $B = (A-t) \cap [m-2k]$, so that $|B| \geq \bar{d}(A)m - 2k$. Then $|B| > \delta m$, by our choice of m_0 . Since B is contained in a translate of A , we have $B - B \subset A - A$, meaning $B - B$ is disjoint from S . \square

The next lemma will be used in conjunction with Lemma 3.6 below. If $S \subset \mathbb{Z}$ and $m \in \mathbb{N}$, we write mS for $\{mn : n \in S\}$.

Lemma 3.5. *Let $k, m \in \mathbb{N}$. If $S \subset \mathbb{Z}$ is k -chromatically recurrent then so is mS .*

Proof. Suppose $m \in \mathbb{N}$, S is k -chromatically recurrent, and let $f : \mathbb{Z} \rightarrow \{1, \dots, k\}$. Form a new coloring $\tilde{f} : \mathbb{Z} \rightarrow \{1, \dots, k\}$ by $\tilde{f}(n) = f(mn)$. Since S is k -chromatically recurrent, there exists $a, b \in \mathbb{Z}$ such that $b - a \in S$ and $\tilde{f}(a) = \tilde{f}(b)$. This means $f(ma) = f(mb)$, and $mb - ma \in mS$. Since f was an arbitrary function into $\{1, \dots, k\}$, this shows that mS is k -chromatically recurrent. \square

Here is the first key lemma for our constructions, a variant of Lemma 3.2 in [13]. We continue to use $[m]$ to denote $\{0, \dots, m-1\}$.

Lemma 3.6. *Let $\delta, \eta \in (0, \frac{1}{2})$, and let $E, F \subset \mathbb{N}$ be finite sets which are δ -nonrecurrent and η -nonrecurrent, respectively. Then for all sufficiently large m , $E \cup mF$ is $2\delta\eta$ -nonrecurrent.*

Furthermore, if (A, m) witnesses the δ -nonrecurrence of E , then for all sufficiently large $l \in \mathbb{N}$, there exists $C \subset [lm]$ with $A \subset C$ such that (C, lm) witnesses the $2\delta\eta$ -nonrecurrence of $E \cup mF$.

In practice we will apply Lemma 3.6 with η close to $\frac{1}{2}$, so that the quantity $2\delta\eta$ in the conclusion will be close to δ in the hypothesis.

Proof. Let E, F, A , and m be as in the hypothesis of the lemma. Since (A, m) witnesses the δ -nonrecurrence of E , we have $A \subset [m]$ and $|A| > \delta m$. Write $|A|$ as $\delta'm$, so that

$\delta' > \delta$. Let $k = \max(E \cup mF)$, and choose l_0 so that for all $l > l_0$, we have

$$(3.1) \quad 2\delta'\eta lm - 2k > 2\delta\eta lm.$$

Let $l \in \mathbb{N}$ be greater than l_0 and large enough (by Lemma 3.4) that there is a set $B \subset [l]$ such that (B, l) witnesses the η -nonrecurrence of F ; fix such a B . We will form C as a union of translates of A . Each such translate $A + t$ will lie in one of the mutually disjoint intervals

$$I_0 = [0, m - 1], I_1 = [m, 2m - 1], \dots, I_{l-1} = [(l-1)m, lm - 1],$$

and will be arranged so that if $A + t \subset I_j$, then $A + t + E \subset I_j$. First fix arbitrary elements $e \in E$ and $f \in F$. Let

$$C_1 := \bigcup_{b \in B} A + mb, \quad C_2 := \bigcup_{b \in B} A + e + m(b + f), \quad C := (C_1 \cup C_2) \cap [lm - 2k].$$

We will prove that

$$(3.2) \quad |C| > 2\delta\eta lm,$$

$$(3.3) \quad C + (E \cup mF) + (E \cup mF) \subset [lm],$$

$$(3.4) \quad C \cap (C + (E \cup mF)) = \emptyset.$$

To compute $|C|$, note that each of the constituent sets $A + mb$ forming C_1 are mutually disjoint, since $A + mb \subset I_b$, and the intervals I_j are mutually disjoint. Likewise, the sets $A + e + m(b + f)$ forming C_2 are mutually disjoint: we have $A + E \subset I_0$, which implies $A + e + m(b + f) \subset I_0 + m(b + f) = I_{b+f}$. Thus $|C_1| = |C_2| = |A||B|$. Furthermore, $I_b \cap I_{b'+f} = \emptyset$ whenever $b, b' \in B$, since $b \neq b' + f$ for such b, b' (i.e. we assumed $B \cap (B + F) = \emptyset$). We therefore have $(A + mb) \cap (A + e + m(b' + f)) = \emptyset$ for all $b, b' \in B$. Thus C_1 and C_2 are disjoint, and we get that

$$|C| \geq 2|A||B| - 2k > 2\delta'\eta lm - 2k > 2\delta\eta lm, \quad \text{by (3.1)}$$

which proves (3.2). The containment (3.3) follows from the containment $C \subset [lm - 2k]$ and our choice of k .

We now prove (3.4), by showing $C \cap (C + E) = \emptyset$ and $C \cap (C + mF) = \emptyset$. To prove that $C \cap (C + E) = \emptyset$, we first show that

$$(3.5) \quad C \cap (C + E) = \bigcup_{j=0}^{l-1} (C \cap I_j) \cap ((C \cap I_j) + E).$$

To see this, note that for each j , the intersection $C \cap I_j$ is one of $A + mj$, $A + e + mj$, or \emptyset . In each case, we have $(C \cap I_j) + E \subset I_j$, by our assumption that $A + E \subset [m]$ and

$A + E + E \subset [m]$. Since $A \cap (A + E) = \emptyset$, we also have $(C \cap I_j) \cap ((C \cap I_j) + E) = \emptyset$ in each case, so (3.5) implies $C \cap (C + E) = \emptyset$.

To prove $C \cap (C + mF) = \emptyset$, we will prove $C_i \cap (C_j + mF) = \emptyset$ for each $i, j \in \{1, 2\}$.

To prove that $C_1 \cap (C_1 + mF) = \emptyset$, first note that C_1 is contained in the union $\bigcup_{b \in B} I_b$, while $C_1 + mF$ is contained in $\bigcup_{j \in B + F} I_j$. The disjointness of B from $B + F$ and the mutual disjointness of the intervals I_j then implies $C_1 \cap (C_1 + mF) = \emptyset$. To see that $C_2 \cap (C_2 + mF) = \emptyset$, it now suffices to observe that C_2 is a translate of C_1 .

Now we show that $C_1 \cap (C_2 + mF) = \emptyset$, by showing that for each $b, b' \in B$,

$$(3.6) \quad (A + mb) \cap (A + e + m(b' + f) + mF) = \emptyset.$$

To see this, note that for each $t \in F$, $A + bm \subset I_b$ and $A + m(b' + f) + mt \subset I_{b'+f+t}$, and these intervals do not overlap unless they are equal, meaning $b = b' + f + t$. In this case the intersection in (3.6) simplifies to $(A + mb) \cap (A + e + mb)$, which is a translate of $A \cap (A + e)$, and therefore empty, as $e \in E$. A similar argument shows that $C_2 \cap (C_1 + mF) = \emptyset$.

Since C is a union of translates of A , we may replace C with one of its translates $C - t$ ($t \geq 0$), to get $A \subset C$ and maintain the inclusions in (3.3). Together with (3.2) and (3.4), this shows that (C, lm) witnesses the $2\delta\eta$ -nonrecurrence of $E \cup mF$. \square

3.2. Proof of Theorem 1.1. Here is our second key ingredient, which we prove in §4.

Lemma 3.7. *For all $k \in \mathbb{N}$ and all $\delta \in (0, \frac{1}{2})$, there is a finite set $S \subset \mathbb{Z}$ such that S is k -chromatically recurrent and δ -nonrecurrent.*

We now prove Theorem 1.1 by combining this lemma with Lemma 3.6. The proof will use no additional information about the sets provided by Lemma 3.7.

Proof of Theorem 1.1. Let $\delta \in (0, \frac{1}{2})$. We will find a chromatically recurrent set S and a set $C \subset \mathbb{Z}$ with $\bar{d}(C) \geq \delta$ such that $(C - C) \cap S = \emptyset$, meaning S is not density recurrent. To build S and C , we will use Lemmas 3.6 and 3.7 to find increasing sequences of sets $S_1 \subset S_2 \subset \dots$, $C_1 \subset C_2 \subset \dots$, and intervals $[m_1], [m_2], \dots$, with $m_k \rightarrow \infty$, so that the following conditions hold for all $k \in \mathbb{N}$:

- (i) S_k is k -chromatically recurrent,
- (ii) $C_k \subset [m_k]$, $C_k + S_k \subset [m_k]$, and $|C_k| > \delta m_k$,
- (iii) $(C_k - C_k) \cap S_k = \emptyset$.

Having constructed these, we let $C := \bigcup_{k=1}^{\infty} C_k$ and $S := \bigcup_{k=1}^{\infty} S_k$. Then $(C - C) \cap S = \emptyset$; otherwise for some $k \in \mathbb{N}$ we would have $c - c' = s$ for some $c, c' \in C_k$ and $s \in S_k$. Item

(ii) implies $\bar{d}(C) \geq \delta$, as $\frac{|C_k \cap [m_k]|}{m_k} > \delta$ for every k . Item (i) implies S is chromatically recurrent, being k -chromatically recurrent for every $k \in \mathbb{N}$.

To find S_k and C_k , we first choose $S_1 = \{1\}$, $m_1 = 2$, and $C_1 = \{0\}$, so that (i)-(iii) are trivially satisfied with $k = 1$.

To perform the inductive step, we assume the sets S_k and C_k have been constructed to satisfy (i)-(iii), and write $|C_k|$ as $\delta_k m_k$. Since C_k satisfies (ii), we have $\delta_k > \delta$, and we choose $\eta < \frac{1}{2}$ so that $2\delta_k \eta > \delta$. Lemma 3.7 provides a finite set S' which is $(k+1)$ -chromatically recurrent and η -nonrecurrent. Apply Lemma 3.6 to find $l \geq 2$ and $C_{k+1} \subset [lm_k]$ such that (C_{k+1}, lm_k) witnesses the $2\delta_k \eta$ -nonrecurrence of $S_k \cup m_k S'$. Finally, Lemma 3.5 implies $m_k S$ is $(k+1)$ -chromatically recurrent. Setting $m_{k+1} = lm_k$ and $S_{k+1} = S_k \cup m_k S$, we get that (i)-(iii) are satisfied with $k+1$ in place of k . \square

3.3. Proof of Theorem 1.2. The following modification of Lemma 3.7 is proved in §4.

Lemma 3.8. *Let $k, m \in \mathbb{N}$ and $\delta < \frac{1}{2}$. If $E \subset \mathbb{Z}$ is infinite, then there is a finite δ -nonrecurrent set $S \subset \mathbb{N}$ such that $mS \subset E - E$ and mS is k -chromatically recurrent.*

Theorem 1.2 may be proved by following the proof of Theorem 1.1 verbatim, with two modifications:

- Let $S_1 = \{t\}$ where $t \in E - E$ satisfies $t > (\frac{1}{2} - \delta)^{-1}$, let $C_1 = [t - 1]$, and let $m_1 = 2t$. Then $|C_1|/m_1 = (t - 1)/(2t) = \frac{1}{2} - \frac{1}{2t} > \delta$, and it is easy to check that (i)-(iii) are satisfied with $k = 1$.
- Cite Lemma 3.8 in place of Lemma 3.7. \square

4. CHROMATIC RECURRENCE AND δ -NONRECURRENCE IN \mathbb{T}^d

The remainder of this article contains the proofs of Lemmas 3.7 and 3.8. Very roughly, the proofs proceed by copying the Hamming balls $H_k(\mathbf{1})$ from \mathbb{F}_2^d into \mathbb{Z} , passing through \mathbb{T}^d as an intermediate step. To be more precise, we fix some notation.

Let \mathbb{R} denote the real numbers with the usual topology and let \mathbb{T} denote \mathbb{R}/\mathbb{Z} with the quotient topology. For $x \in \mathbb{T}$, let \tilde{x} be the unique element of $[0, 1)$ such that $\tilde{x} + \mathbb{Z} = x$, and write $\|x\|$ for $\min_{n \in \mathbb{Z}} |\tilde{x} - n|$.

When defining subsets of \mathbb{T} , we identify subintervals of $[0, 1) \subset \mathbb{R}$ with their images in \mathbb{T} under the quotient map.

Let $G_d := \{0, 1/2\}^d \subset \mathbb{T}^d$, so that G_d is a subgroup isomorphic to \mathbb{F}_2^d . With the natural identification, $\mathbf{1} \in \mathbb{F}_2^d$ is identified with $\frac{1}{2} := (1/2, 1/2, \dots, 1/2) \in \mathbb{T}^d$. We will use the

notation $H_k(\mathbf{t})$ for Hamming balls around elements of G_d , so $H_k(\frac{1}{2})$ is, by definition, the set of $(x_1, \dots, x_d) \in G_d$ where at most k entries x_i are not equal to $1/2$.

Let V_ε denote the open box $\{\mathbf{x} \in \mathbb{T}^d : \max \|x_j\| < \varepsilon\}$. For $\boldsymbol{\alpha} \in \mathbb{T}^d$, define

$$(4.1) \quad \tilde{H}(\boldsymbol{\alpha}; k, \varepsilon) := \{n \in \mathbb{Z} : n\boldsymbol{\alpha} \in H_k(\frac{1}{2}) + V_\varepsilon\}.$$

We call $\tilde{H}(\boldsymbol{\alpha}; k, \varepsilon)$ the ε -copy of $H_k(\mathbf{1})$ determined by $\boldsymbol{\alpha}$.

The next lemma records the key properties of the $\tilde{H}(\boldsymbol{\alpha}; k, \varepsilon)$. When $\boldsymbol{\alpha} \in \mathbb{T}^d$ and $E \subset \mathbb{Z}$, we write $E\boldsymbol{\alpha}$ for $\{n\boldsymbol{\alpha} : n \in E\}$.

- Lemma 4.1.** (i) *Let $\delta < \frac{1}{2}$ and $k \in \mathbb{N}$. For all sufficiently large $d \in \mathbb{N}$ there is an $\varepsilon > 0$ such that $\tilde{H}(\boldsymbol{\alpha}; k, \varepsilon)$ is δ -nonrecurrent whenever $\boldsymbol{\alpha} \in \mathbb{T}^d$.*
- (ii) *If $\mathbb{Z}\boldsymbol{\alpha}$ is dense in \mathbb{T}^d and $\varepsilon > 0$, then there is a finite subset of $\tilde{H}(\boldsymbol{\alpha}; 2k + 1, \varepsilon)$ which is $2k$ -chromatically recurrent.*
- (iii) *If $E \subset \mathbb{Z}$ is such that $E\boldsymbol{\alpha}$ is dense in \mathbb{T}^d and $\varepsilon > 0$, then there is a finite subset of $\tilde{H}(\boldsymbol{\alpha}; 2k + 1, \varepsilon) \cap (E - E)$ which is $2k$ -chromatically recurrent.*

Parts (ii) and (iii) will be proved in §5. The remainder of this section is dedicated to the proof of Part (i).

Given $\varepsilon > 0$, let $I_\varepsilon = [\varepsilon, \frac{1}{2} - \varepsilon] \subset \mathbb{T}$, and let $I_\varepsilon^d \subset \mathbb{T}^d$ be its d -fold cartesian power. Observe that

$$(4.2) \quad \{\mathbf{h} + I_\varepsilon^d : \mathbf{h} \in G_d\} \text{ is mutually disjoint.}$$

Write μ for Haar probability measure on \mathbb{T}^d . Given $A \subset G_d$, let $A_\varepsilon^\square := A + I_\varepsilon^d$, so that (4.2) implies

$$(4.3) \quad \mu(A_\varepsilon^\square) = |A|(\frac{1}{2} - 2\varepsilon)^d.$$

Lemma 4.2. *If $\mathbf{t} \in G_d$ and $A \subset G_d$, then $A_\varepsilon^\square \cap (A_\varepsilon^\square + \mathbf{t}) = (A \cap (A + \mathbf{t}))_\varepsilon^\square$.*

Proof. We prove only that $A_\varepsilon^\square \cap (A_\varepsilon^\square + \mathbf{t}) \subseteq (A \cap (A + \mathbf{t}))_\varepsilon^\square$, as the reverse containment is easy to check.

Assuming $\mathbf{x} \in A_\varepsilon^\square$, there is an $\mathbf{a} \in A$ such that $\mathbf{x} \in \mathbf{a} + I_\varepsilon^d$, while assuming $\mathbf{x} \in A_\varepsilon^\square + \mathbf{t}$ we find $\mathbf{a}' \in A$ such that $\mathbf{x} \in \mathbf{a}' + \mathbf{t} + I_\varepsilon^d$. Thus $\mathbf{x} \in (\mathbf{a} + I_\varepsilon^d) \cap (\mathbf{a}' + \mathbf{t} + I_\varepsilon^d)$, and the mutual disjointness of the $\mathbf{h} + I_\varepsilon^d$ mentioned in (4.2) then implies $\mathbf{a} = \mathbf{a}' + \mathbf{t}$. It follows that $\mathbf{x} \in (A \cap (A + \mathbf{t}))_\varepsilon^\square$, as desired. \square

The important consequence of Lemma 4.2 is that if $A, R \subset G_d$ with $A \cap (A + R) = \emptyset$, then $A_\varepsilon^\square \cap (A_\varepsilon^\square + R) = \emptyset$.

A *box* in \mathbb{T}^d is a product $J_1 \times J_2 \times \cdots \times J_d$ of intervals $J_i \subset \mathbb{T}$. Note that the sets A_ε^\square defined above are finite unions of boxes.

The following standard fact about uniform distribution is proved in §7.

Lemma 4.3. *Let $\alpha \in \mathbb{T}^d$ be such that $\mathbb{Z}\alpha$ is dense in \mathbb{T}^d . If $A \subset \mathbb{T}^d$ is a finite union of boxes and $C := \{n \in \mathbb{Z} : n\alpha \in A\}$ then $d(C) = \mu(A)$, where μ is Haar probability measure on \mathbb{T}^d .*

Lemma 4.4. *Let $B, U \subset \mathbb{T}^d$ and assume $\mathbb{Z}\alpha$ is dense in \mathbb{T}^d . If B is a finite union of boxes with $\mu(B) > \delta$ and $B \cap (B + U) = \emptyset$, then $S := \{n : n\alpha \in U\}$ is δ -nonrecurrent.*

Proof. Let $C := \{n : n\alpha \in B\}$. By Lemma 4.3, $d(C) = \mu(B) > \delta$. To see that $C \cap (C + S) = \emptyset$, note that if $n \in C \cap (C + S)$, we have $n\alpha \in B$ and $n\alpha \in B + U$, meaning $n\alpha \in B \cap (B + U)$, which violates our hypothesis. \square

Lemma 4.5. *If $R \subset G_d$ is δ -nonrecurrent and $\alpha \in \mathbb{T}^d$, then for sufficiently small ε , $S := \{n \in \mathbb{Z} : n\alpha \in R + V_\varepsilon\}$ is δ -nonrecurrent.*

Proof. Let $A \subset G_d$ be such that $A \cap (A + R) = \emptyset$ and $|A| > \delta 2^d$. Choose $\varepsilon > 0$ small enough that $|A|(\frac{1}{2} - 2\varepsilon)^d > \delta$. Let $B := A_\varepsilon^\square$, so that $\mu(B) = |A|(\frac{1}{2} - 2\varepsilon)^d > \delta$. Now $B \cap (B + R) = \emptyset$, by Lemma 4.2. Since B is compact, we also have $B \cap (B + R + V_\varepsilon) = \emptyset$, provided ε is sufficiently small. Lemma 4.4 then implies S is δ -nonrecurrent. \square

Proof of Lemma 4.1 Part (i). Let $\delta < \frac{1}{2}$ and $k \in \mathbb{N}$. By Lemma 2.4, choose $d \in \mathbb{N}$ so that $H_k(\mathbf{1})$ is a δ -nonrecurrent subset of \mathbb{F}_2^d . Then $H_k(\frac{1}{2})$ is a δ -nonrecurrent subset of G_d , and Lemma 4.5 provides an $\varepsilon > 0$ so that $\{n \in \mathbb{Z} : n\alpha \in H_k(\frac{1}{2}) + V_\varepsilon\}$ is δ -nonrecurrent. This δ -nonrecurrent set is, by definition, $\tilde{H}(\alpha; k, \varepsilon)$. \square

5. LIFTING CHROMATIC RECURRENCE FROM TOPOLOGICAL GROUPS

Here we show how a homomorphism from \mathbb{Z} into a topological abelian group K can be used to copy Cayley graphs from K into \mathbb{Z} ; we maintain the conventions of §2.1.

For this section we fix a discrete abelian group G , a topological abelian group K (not necessarily compact, not necessarily metrizable), and a homomorphism $\rho : G \rightarrow K$ with $\rho(G)$ dense in K . We also fix $E \subset G$ with $\rho(E)$ dense in K . The only case we will use in this article is $G = \mathbb{Z}$, $K = \mathbb{T}^d$, and $\rho(n) = n\alpha$ for some $\alpha \in \mathbb{T}^d$, so readers can specialize to this setting at will.

Lemma 5.1. *Assume G , ρ , E , and K are as specified above. If $U \subset K$ is open, every finite subgraph \mathcal{G} of $\text{Cay}(U)$ has an isomorphic copy in $\text{Cay}(\rho^{-1}(U) \cap (E - E))$.*

Proof. To prove the lemma it suffices to prove that if V is a finite subset of K , then there exist $\{g_v : v \in V\} \subset E$ such that for each $v, v' \in V$, we have

$$(5.1) \quad v - v' \in U \implies g_v - g_{v'} \in \rho^{-1}(U).$$

So let V be a finite subset of K . Let $S := (V - V) \cap U$, and let W be a neighborhood of 0 in K so that $S + W \subset U$. Choose a neighborhood W' of 0 so that $W' - W' \subset W$. For each $v \in V$, choose $g_v \in E$ so that $\rho(g_v) \in v + W'$; this is possible since $\rho(E)$ is dense in K . We now prove (5.1) holds with these g_v . Assuming $v - v' \in U$, we have

$$\rho(g_v) - \rho(g_{v'}) \in v + W' - (v' + W') = (v - v') + (W' - W') \subset v - v' + W \subset U,$$

so $g_v - g_{v'} \in \rho^{-1}(U)$. This proves (5.1) \square

Since chromatic number is invariant under isomorphism of graphs, the following corollary is immediate.

Corollary 5.2. *If $U \subset K$ and $\text{Cay}(U)$ has a finite subgraph with chromatic number $\geq k$, then $\text{Cay}(\rho^{-1}(U) \cap (E - E))$ has chromatic number $\geq k$.*

Proof of Lemma 4.1 Parts (ii) and (iii). Part (ii) follows from Part (iii), so we only prove Part (iii).

Let $2k \leq d \in \mathbb{N}$, let $E \subset \mathbb{Z}$, and let $\alpha \in \mathbb{T}^d$ with $E\alpha$ dense in \mathbb{T}^d . We use Corollary 5.2, with the open set $U = H_{2k+1}(\frac{1}{2}) + V_\varepsilon$ and $\rho(n) := n\alpha$. Since $H_{2k+1}(\frac{1}{2}) \subset U$, $\text{Cay}(U)$ contains the finite subgraph $\text{Cay}(H_{2k+1}(\frac{1}{2}))$ (with vertex set G_d). The latter graph is isomorphic to $\text{Cay}(H_{2k+1}(\mathbf{1}))$, and by Lemma 2.5, it has chromatic number at least $2k + 1$. Now Corollary 5.2 implies $\text{Cay}(\rho^{-1}(U) \cap (E - E))$ has chromatic number at least $2k + 1$. Examining the definitions, we see that $\rho^{-1}(U) = \tilde{H}(\alpha; k, \varepsilon)$, and we get that $\tilde{H}(\alpha; k, \varepsilon) \cap (E - E)$ is $2k$ -chromatically recurrent. \square

6. PROOF OF LEMMAS 3.7 AND 3.8

Lemma 3.7 is an immediate consequence of Parts (i) and (ii) of Lemma 4.1. For Lemma 3.8, we need an elementary result on recurrence properties of dilates. Given $S \subset \mathbb{Z}$ write S/m for the set $\{n \in \mathbb{Z} : mn \in S\}$ ($= \{n/m : n \in S\} \cap \mathbb{Z}$).

Lemma 6.1. *If $m \in \mathbb{N}$ and $S \subset \mathbb{Z}$ is δ -nonrecurrent, then S/m is also δ -nonrecurrent.*

Proof. Let $S \subset \mathbb{Z}$ be δ -nonrecurrent, and choose $A \subset \mathbb{Z}$ such that $(A - A) \cap S = \emptyset$ and $\bar{d}(A) > \delta$. The subadditivity of \bar{d} implies that for some $t \in \{0, \dots, m - 1\}$, the set $A_t := A \cap (t + m\mathbb{Z})$ satisfies $\bar{d}(A_t) > \delta/m$. The elements of $A_t - t$ are all divisible by m ,

so $A' := (A_t - t)/m$ satisfies $\bar{d}(A') > \delta$. Furthermore, $m(A' - A') \subset A - A$, so $m(A' - A')$ is disjoint from S , meaning $A' - A'$ is disjoint from S/m . \square

We recall the statement of Lemma 3.8: if $k, m \in \mathbb{N}$, $\delta < \frac{1}{2}$, and $E \subset \mathbb{Z}$ is an infinite set, then there is a finite δ -nonrecurrent set $S \subset \mathbb{N}$ such that mS is k -chromatically recurrent and $mS \subset E - E$. Our proof will use the following result from the theory of uniform distribution, which we prove in §7.

Lemma 6.2. *Let $E \subset \mathbb{Z}$ be an infinite set and $d \in \mathbb{N}$. For μ -almost every $\alpha \in \mathbb{T}^d$, the set $E\alpha := \{n\alpha : n \in E\}$ is dense in \mathbb{T}^d .*

Proof of Lemma 3.8. Fix $\delta < \frac{1}{2}$. Let $E \subset \mathbb{Z}$ be infinite and $m \in \mathbb{N}$. Let $E' \subset E$ be an infinite subset with $m|(b - a)$ for all $a, b \in E'$ (i.e. the elements of E' are mutually congruent mod m).

By Part (i) of Lemma 4.1, choose $d \in \mathbb{N}$ large enough and $\varepsilon > 0$ small enough that $\tilde{H} := \tilde{H}(\alpha; 2k + 1, \varepsilon)$ is δ -nonrecurrent. We then apply Part (iii) of Lemma 4.1 to find a finite subset $S_0 \subset \tilde{H} \cap (E' - E')$ which is $2k$ -chromatically recurrent.

Since the elements of E' are mutually congruent mod m , we have $S_0 \subset m\mathbb{Z}$, and the previous paragraph shows that S_0 is $2k$ -chromatically recurrent and δ -nonrecurrent. Let $S := S_0/m$. By Lemma 6.1 S is also δ -nonrecurrent. Now S is the desired set: we have $mS = S_0 \subset E' - E' \subset E - E$, S is δ -nonrecurrent, and mS is $2k$ -chromatically recurrent. \square

7. DENSENESS AND UNIFORM DISTRIBUTION IN \mathbb{T}^d

The material in this section is due to Kronecker and to Weyl; the standard reference is [12]. We present it here to make the proof of Lemma 6.2 self-contained. For this section we fix $d \in \mathbb{N}$ and write μ for Haar probability measure on \mathbb{T}^d .

A sequence $\alpha_1, \alpha_2, \dots$ of points in \mathbb{T}^d is called *uniformly distributed* if

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(\alpha_n) = \int f d\mu$$

for every continuous $f : \mathbb{T}^d \rightarrow \mathbb{C}$. A straightforward approximation argument shows that a sequence $(\alpha_n)_{n \in \mathbb{N}}$ is uniformly distributed if and only if for every box $R := \prod_{j=1}^d [a_j, b_j]$ in \mathbb{T}^d , we have

$$\lim_{N \rightarrow \infty} \frac{|\{n \in \{1, \dots, N\} : \alpha_n \in R\}|}{N} = \mu(R).$$

Consequently, the terms of a uniformly distributed sequence are dense in \mathbb{T}^d .

The above criterion immediately yields the following lemma.

Lemma 7.1. *If $A \subset \mathbb{T}^d$ is a finite union of boxes and $(\alpha_n)_{n \in \mathbb{N}}$ is uniformly distributed then $C := \{n : \alpha_n \in A\}$ has $d(C) = \mu(A)$.*

We write \mathcal{S}^1 for the group of complex numbers of modulus 1; the group operation on \mathcal{S}^1 is multiplication and the topology is the one inherited from \mathbb{C} . A *character* of \mathbb{T}^d is a continuous homomorphism $\chi : \mathbb{T}^d \rightarrow \mathcal{S}^1$; the set of characters of \mathbb{T}^d is denoted by $\widehat{\mathbb{T}}^d$. A well known fact from basic harmonic analysis is that every character of \mathbb{T}^d has the form $\chi(x_1, \dots, x_d) := \exp(2\pi i(n_1 x_1 + \dots + n_d x_d))$, where n_1, \dots, n_d are integers. The *trivial character* satisfies $\chi(\mathbf{x}) = 1$ for all $\mathbf{x} \in \mathbb{T}^d$. It is easy to check that if χ is a nontrivial character and $m \in \mathbb{Z}$ then $\mathbf{x} \mapsto \chi(m\mathbf{x})$ is another nontrivial character. Furthermore, the characters of \mathbb{T}^d are mutually orthogonal in $L^2(\mu)$.

The *Weyl criterion* [12, p.62] for uniform distribution says that a sequence of points $(\alpha_n)_{n \in \mathbb{N}}$ in \mathbb{T}^d is uniformly distributed if and only if

$$(7.1) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \chi(\alpha_n) = 0 \quad \text{for every nontrivial character } \chi \in \widehat{\mathbb{T}}^d.$$

We now use Weyl criterion to prove Lemma 4.3, which says that if $(n\alpha)_{n \in \mathbb{N}}$ is dense in \mathbb{T}^d , $A \subset \mathbb{T}^d$ is a finite union of boxes, and $C := \{n \in \mathbb{Z} : n\alpha \in A\}$, then $\bar{d}(C) = \mu(A)$.

Proof of Lemma 4.3. By Lemma 7.1, it suffices to prove that if $(n\alpha)_{n \in \mathbb{N}}$ is dense in \mathbb{T}^d , then (7.1) is satisfied by $\alpha_n = n\alpha$. To prove this, we need only show that $\chi(\alpha) \neq 1$, for then we may apply the identity $\chi(n\alpha) = \chi(\alpha)^n$ and simplify partial sums $S_N := \sum_{n=1}^N \chi(n\alpha)$ as $\chi(\alpha) \frac{1 - \chi(\alpha)^N}{1 - \chi(\alpha)}$. Since $|\chi(\alpha)| = 1$, the latter quantity is bounded in modulus by a number independent of N , whence $\lim_{N \rightarrow \infty} \frac{1}{N} S_N = 0$. To see that $\chi(\alpha) \neq 1$, note that the denseness of $(n\alpha)_{n \in \mathbb{N}}$ in \mathbb{T}^d implies that $\{\chi(n\alpha) : n \in \mathbb{N}\}$ is dense in the image of χ (since χ is continuous), and in particular $\chi(n\alpha) \neq 1$ for some n . This implies $\chi(\alpha)^n \neq 1$, so $\chi(\alpha) \neq 1$. \square

The next lemma forms most of the proof of Lemma 6.2.

Lemma 7.2. *If μ is a probability measure and $(\phi_j)_{j \in \mathbb{N}}$ are mutually orthogonal elements of $L^2(\mu)$ pointwise bounded in modulus by 1, then $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N \phi_j(x) = 0$ for μ -a.e. x .*

Proof. Let $S_N(x) := \frac{1}{N} \sum_{j=1}^N \phi_j(x)$. It suffices to prove that $\sum_{N=1}^{\infty} |S_N(x)|^2$ converges μ -almost everywhere, since the convergence of the sum implies the terms $|S_N(x)|$ converge

to 0. Expanding $\int |S_N(x)|^2 d\mu(x)$ and using the fact that $\int \phi_i \bar{\phi}_j d\mu = 0$ for $i \neq j$, we get

$$\int |S_N(x)|^2 d\mu(x) = \frac{1}{N^2} \sum_{j,k=1}^N \int \phi_j \bar{\phi}_k d\mu = \frac{1}{N^2} \sum_{j=1}^N \|\phi_j\|^2 \leq \frac{1}{N}.$$

Thus $\sum_{N=1}^{\infty} \int |S_{N^2}(x)|^2 d\mu \leq \sum_{N=1}^{\infty} \frac{1}{N^2} < \infty$. The monotone convergence theorem then implies that the sum $\sum_{N=1}^{\infty} |S_{N^2}(x)|^2$ is finite for μ -almost every x , and we conclude that

$$(7.2) \quad \lim_{N \rightarrow \infty} S_{N^2}(x) = 0$$

for μ -almost every x . We will prove that $\lim_{n \rightarrow \infty} S_n(x) = 0$ for every x satisfying (7.2). To see this, fix such an x , an $\varepsilon > 0$, and choose N large enough that $|S_{N^2}(x)| < \varepsilon/2$ and $|1 - \frac{N^2}{(N+1)^2}| < \varepsilon/2$. Then for every n between N^2 and $(N+1)^2$, we have

$$\begin{aligned} |S_n(x)| &= \left| \frac{1}{n} \sum_{j=1}^{N^2} \phi_j(x) + \frac{1}{n} \sum_{j=N^2+1}^n \phi_j(x) \right| \leq \frac{N^2}{n} \left| \frac{1}{N^2} \sum_{j=1}^{N^2} \phi_j(x) \right| + \frac{1}{n} |n - N^2| \\ &\leq |S_{N^2}(x)| + \left| 1 - \frac{N^2}{(N+1)^2} \right| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

Since ε was arbitrary, this shows that $S_n(x) \rightarrow 0$ as $n \rightarrow \infty$. \square

Theorem 7.3. *Let $(s_n)_{n \in \mathbb{N}}$ be a sequence of distinct integers and $d \in \mathbb{N}$. For μ -almost every α in \mathbb{T}^d , the sequence $(s_n \alpha)_{n \in \mathbb{N}}$ is uniformly distributed in \mathbb{T}^d .*

Proof. By the Weyl criterion it suffices to prove that for μ -almost every α , we have

$$(7.3) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \chi(s_n \alpha) = 0$$

for every nontrivial character χ of \mathbb{T}^d . Since there are only countably many characters of \mathbb{T}^d , it suffices to prove that for all nontrivial characters χ , equation (7.3) holds for μ -almost every α . To prove this, note that when χ is nontrivial, the functions ϕ_n defined by $\phi_n(\mathbf{x}) := \chi(s_n \mathbf{x})$ are mutually distinct characters, and $(\phi_n)_{n \in \mathbb{N}}$ is therefore a sequence of mutually orthogonal functions in $L^2(\mu)$. The limit (7.3) now follows from Lemma 7.2. \square

We now prove Lemma 6.2, which says that for every infinite set $E \subset \mathbb{Z}$ all $d \in \mathbb{N}$ and μ -almost every $\alpha \in \mathbb{T}^d$, the set $E\alpha := \{n\alpha : n \in E\}$ is dense in \mathbb{T}^d .

Proof of Lemma 6.2. Enumerating E as $(e_n)_{n \in \mathbb{N}}$, we see that the lemma is a corollary of Theorem 7.3, since the terms of a uniformly distributed sequence must be dense in \mathbb{T}^d . \square

8. QUESTIONS

A proof or disproof of the following conjecture would be very interesting in relation to the results of [11].

Conjecture 8.1. *If $S \subset \mathbb{Z}$ is a set of density recurrence, then there is a set $S' \subset S$ such that S' is a set of chromatic recurrence and not a set of density recurrence.*

Perhaps the proof of Theorem 1.2 can be extended to prove Conjecture 8.1. If so, the key lemma to generalize is Part (iii) of Lemma 4.1. Specifically, the required generalization would replace $E - E$ with an arbitrary set of chromatic recurrence, resulting in the following conjecture. We use the notation defined in (4.1).

Conjecture 8.2. *Assume $S \subset \mathbb{Z}$ is a set of chromatic recurrence, $k \in \mathbb{N}$, and $C > 0$. Then there exist infinitely many $d \in \mathbb{N}$ and $\alpha \in \mathbb{T}^d$, such that for all $\varepsilon > 0$, the intersection*

$$S \cap \tilde{H}(\alpha; C\sqrt{d}, \varepsilon)$$

is a set of k -chromatic recurrence.

See the remarks following Theorem 2.2 above to motivate the appearance of $C\sqrt{d}$ here.

REFERENCES

1. I. Bárány, *A short proof of Kneser's conjecture*, J. Combin. Theory Ser. A **25** (1978), no. 3, 325–326. MR 514626
2. Vitaly Bergelson, *Ergodic Ramsey theory*, Logic and combinatorics (Arcata, Calif., 1985), Contemp. Math., vol. 65, Amer. Math. Soc., Providence, RI, 1987, pp. 63–87. MR 891243
3. Vitaly Bergelson and Randall McCutcheon, *Recurrence for semigroup actions and a non-commutative Schur theorem*, Topological dynamics and applications (Minneapolis, MN, 1995), Contemp. Math., vol. 215, Amer. Math. Soc., Providence, RI, 1998, pp. 205–222.
4. N. Bogoliouboff, *Sur quelques propriétés arithmétiques des presque-périodes*, Ann. Chaire Phys. Math. Kiev **4** (1939), 185–205. MR 20164
5. Erling Følner, *Note on a generalization of a theorem of Bogoliouboff*, Math. Scand. **2** (1954), 224–226. MR 69188
6. Alan H. Forrest, *Recurrence in dynamical systems: A combinatorial approach*, ProQuest LLC, Ann Arbor, MI, 1990, Thesis (Ph.D.)—The Ohio State University.
7. ———, *The construction of a set of recurrence which is not a set of strong recurrence*, Israel J. Math. **76** (1991), no. 1-2, 215–228. MR 1177341 (93j:28023)
8. H. Furstenberg, *Recurrence in ergodic theory and combinatorial number theory*, Princeton University Press, Princeton, N.J., 1981, M. B. Porter Lectures. MR 603625 (82j:28010)

9. Harry Furstenberg, *Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions*, J. Analyse Math. **31** (1977), 204–256. MR 498471
10. Joshua E. Greene, *A new short proof of Kneser’s conjecture*, Amer. Math. Monthly **109** (2002), no. 10, 918–920. MR 1941810
11. John T. Griesmer, *Special cases and equivalent forms of Katznelson’s problem on recurrence*, 2021, Draft.
12. L. Kuipers and H. Niederreiter, *Uniform distribution of sequences*, Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, 1974, Pure and Applied Mathematics. MR 0419394
13. Igor Kříž, *Large independent sets in shift-invariant graphs: solution of Bergelson’s problem*, Graphs Combin. **3** (1987), no. 2, 145–158.
14. L. Lovász, *Kneser’s conjecture, chromatic number, and homotopy*, J. Combin. Theory Ser. A **25** (1978), no. 3, 319–324. MR 514625
15. Jiří Matoušek, *Using the Borsuk-Ulam theorem*, Universitext, Springer-Verlag, Berlin, 2003, Lectures on topological methods in combinatorics and geometry, Written in cooperation with Anders Björner and Günter M. Ziegler. MR 1988723
16. Randall McCutcheon, *Three results in recurrence*, Ergodic theory and its connections with harmonic analysis (Alexandria, 1993), London Math. Soc. Lecture Note Ser., vol. 205, Cambridge Univ. Press, Cambridge, 1995, pp. 349–358. MR 1325710 (96e:28015)
17. ———, *Elemental methods in ergodic Ramsey theory*, Lecture Notes in Mathematics, vol. 1722, Springer-Verlag, Berlin, 1999. MR 1738544
18. A. Sárközy, *On difference sets of sequences of integers. I*, Acta Math. Acad. Sci. Hungar. **31** (1978), no. 1-2, 125–149. MR 466059
19. E. Szemerédi, *On sets of integers containing no k elements in arithmetic progression*, Acta Arith. **27** (1975), 199–245. MR 369312
20. B.L. van der Waerden, *Beweis einer boudetschen vermutung*, Nieuw Arch. Wisk. **15** (1927), 212–216.
21. Benjamin Weiss, *Single orbit dynamics*, CBMS Regional Conference Series in Mathematics, vol. 95, American Mathematical Society, Providence, RI, 2000. MR 1727510

Email address: `jtgriesmer@gmail.com`

DEPARTMENT OF APPLIED MATHEMATICS AND STATISTICS, COLORADO SCHOOL OF MINES, GOLDEN,
COLORADO