

SPECTRAL PROPERTIES OF SUBSTITUTIONS ON COMPACT ALPHABETS

NEIL MAÑIBO, DAN RUST, AND JAMES J. WALTON

ABSTRACT. We consider substitutions on compact alphabets and provide sufficient conditions for the diffraction to be pure point, absolutely continuous and singular continuous. This allows one to construct examples for which the Koopman operator on the associated function space has specific spectral components. For abelian bijective substitutions, we provide a dichotomy result regarding the spectral type of the diffraction. We also provide the first example of a substitution which has countably infinite Lebesgue spectral components and countably infinite singular continuous components. Lastly, we give a non-constant length substitution on a countably infinite alphabet which gives rise to substitutive Delone sets of infinite type. This extends the spectral theory of substitutions on finite alphabets and Delone sets of finite type with inflation symmetry.

1. INTRODUCTION

Substitutions over infinite alphabets have already been considered in a variety of contexts [12, 14, 28, 29], but very few accounts on their spectral theory are available. One difficulty is that classical methods such as Perron–Frobenius theory are no longer immediately available in the infinite alphabet setting, and so new techniques must be developed [27]. Even then, it is possible that the maximal spectral type of the underlying system is generated by an infinite family of functions, which is usually the case.

In this work, we consider continuous substitutions on a compact alphabet and develop the spectral theory for certain families of representative examples, which are infinite alphabet generalisations of the Thue–Morse, period doubling and Rudin–Shapiro substitutions. We demonstrate how similar arguments to those implemented in the finite alphabet setting allow one to completely determine their spectral type and, at the same time, show that new phenomena can occur.

It is well known that there is a correspondence between the diffraction spectrum and dynamical spectrum. Indeed, this has been rigorously established for systems with pure point spectrum in [6, 23], for systems with mixed spectrum in the finite local complexity (FLC) setting in [7] and for general dynamical systems in [25]. We employ a diffraction-based approach and study the diffraction of weighted point sets which arise from elements of the subshift. This entails choosing an appropriate set of weight functions, which we get for free when the alphabet has an extra group structure. In these cases, the characters of the associated group provide complete access to the admissible spectral types.

We establish sufficient conditions for when the corresponding diffraction measure is pure point and purely singular continuous for bijective substitutions on compact abelian alphabets, which are seen as compact alphabet generalisations of the Thue–Morse substitution; see Theorem 3.18 for equivalent conditions for singular continuity. These spectral results allow one to distinguish bijective abelian substitutions which admit periodic factors. These results generalise the ones in [5, 9, 15]. In the finite alphabet case, it is conjectured that bijective substitutions always have singular spectrum. This has been confirmed for substitutions on a binary alphabet [5] and for abelian substitutions [9] but remains open for the non-abelian case; see [9, 28].

We also consider the compact alphabet analogues of the period doubling substitution and show that they have pure point spectrum (both in the dynamical and diffraction sense), which is done in Theorem 3.23, extending a famous result due to Dekking [11]. The subshifts they generate can be seen as generalisations of Toeplitz shifts over compact alphabets.

2010 *Mathematics Subject Classification.* 37B10, 37A30, 52C23, 42A16.

Key words and phrases. substitutions, compact alphabets, diffraction, dynamical spectrum, Delone sets.

To complete the picture, we provide an analogue of the Rudin–Shapiro substitution on a compact alphabet. The alphabet is $\mathcal{A} = S^1 \times \{0, 1\}$, and the corresponding dynamical system is an infinite-index extension of the odometer $(\mathbb{Z}_2, +1)$ by S^1 ; compare [1, 16, 19]. Its spectral theory is completely determined by the defining spin matrix W . From W , one can explicitly define the skew product which induces the \mathbb{Z} -action on $\mathbb{Z}_2 \times S^1$. Using the results in [16], we prove in Theorem 3.29 that this example has countably infinite Lebesgue components and countably infinite singular continuous components in its dynamical spectrum. To the authors’ knowledge, this is the first substitutive example with such a spectral decomposition; obviously one cannot have this for finite alphabets where the maximal spectral type is generated by only finitely many correlation measures [9, 28].

All of the previously mentioned families are constant-length, and so one can say that, for all of them, infinite local complexity arises in a purely combinatorial fashion. We end with an example of a non-constant length substitution with uniquely ergodic subshift with alphabet $\mathbb{N}_0 \cup \{\infty\}$. It admits natural tile lengths which are bounded from above and are bounded away from zero [27]. This allows one to construct a tiling of \mathbb{R} with infinitely many prototile lengths. Here, infinite local complexity also manifests geometrically apart from the combinatorial complexity induced by the alphabet. From this tiling, one can derive a Delone set of infinite type which still has an inflation symmetry in the sense of Lagarias [22].

2. SUBSTITUTIONS ON COMPACT ALPHABETS

In this section, we recall notions and results regarding substitutions on compact alphabets. For a more detailed treatment and proofs of the dynamical results, we refer the reader to [27]. Let \mathcal{A} be a compact Hausdorff space which we call an *alphabet* and whose elements we call *letters*. Let $\mathcal{A}^+ = \bigsqcup_{n \geq 1} \mathcal{A}^n$ denote the set of all finite (non-empty) *words* over the alphabet \mathcal{A} , where \mathcal{A}^n has the product topology and \mathcal{A}^+ has the topology inherited from a disjoint union. Let $\mathcal{A}^* = \mathcal{A}^+ \sqcup \{\varepsilon\}$, where ε is the *empty word*. *Concatenation* is a binary operation $\mathcal{A}^* \times \mathcal{A}^* \rightarrow \mathcal{A}^*$ given by $(\varepsilon, u) \mapsto u$, $(u, \varepsilon) \mapsto u$ and $(u_1 \cdots u_n, v_1 \cdots v_m) \mapsto u_1 \cdots u_n v_1 \cdots v_m$, where $u = u_1 \cdots u_n \in \mathcal{A}^n$ and $v = v_1 \cdots v_m \in \mathcal{A}^m$. Without loss of generality, we write uv as short hand for the concatenation of u and v . Note that concatenation is a continuous operation. Let $\mathcal{A}^{\mathbb{Z}}$ denote the set of bi-infinite sequences over \mathcal{A} with the product topology. We use a vertical line $|$ to denote the position between the -1 st and 0 th element of a bi-infinite sequence, and so we write $w = \cdots w_{-2}w_{-1}|w_0w_1 \cdots$. The space $\mathcal{A}^{\mathbb{Z}}$ is compact by Tychonoff’s theorem. The left shift $\sigma: \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$ is defined by $\sigma(w)_n = w_{n+1}$. A subshift $X \subseteq \mathcal{A}^{\mathbb{Z}}$ is a subspace of the full shift which is closed and σ -invariant.

Definition 2.1. Let $\varrho: \mathcal{A} \rightarrow \mathcal{A}^+$ be a continuous function. We call such a function a *substitution* on \mathcal{A} . We say ϱ is a substitution of *constant length* n if $\varrho(\mathcal{A}) \subseteq \mathcal{A}^n$.

If \mathcal{A} is connected, then by continuity, ϱ is necessarily of constant length. Hence, constant-length substitutions are fairly ‘typical’ in the compact alphabet setting, where connected alphabets such as manifolds can be considered.

Definition 2.2. Let $\varrho: \mathcal{A} \rightarrow \mathcal{A}^+$ be a substitution. We say a word $u \in \mathcal{A}^n$ is *admitted* by ϱ if there exist $a \in \mathcal{A}$, $k \geq 1$ such that $\varrho^k(a)$ contains u as a subword. We write

$$\mathcal{L}^n(\varrho) := \overline{\{u \in \mathcal{A}^n \mid u \text{ is admitted by } \varrho\}}$$

and define $\mathcal{L}(\varrho) := \bigsqcup \mathcal{L}^n(\varrho)$. We call $\mathcal{L}(\varrho)$ the *language* of ϱ and call words in $\mathcal{L}(\varrho)$ *legal*. We note here that we have to take the closure to form \mathcal{L}^n , which is automatically satisfied for finite alphabets. We define a closed, shift-invariant subspace of $\mathcal{A}^{\mathbb{Z}}$ by

$$X_\varrho := \{w \in \mathcal{A}^{\mathbb{Z}} \mid \forall i \leq j, w_{[i,j]} \in \mathcal{L}(\varrho)\}$$

and call X_ϱ the *subshift* associated with ϱ [27].

The following definition is the natural generalisation of primitivity to infinite alphabets, adapted from the work of Durand, Ormes and Petite [12]; see also Frank and Sadun [17] and Queffélec [28].

Definition 2.3. Let $\varrho: \mathcal{A} \rightarrow \mathcal{A}^+$ be a substitution. We say ϱ is *primitive* if, for every open set $U \subset \mathcal{A}$, there exists a $p = p(U) \geq 0$ such that for all $a \in \mathcal{A}$, some letter of $\varrho^p(a)$ is in U .

Let $E = C(\mathcal{A})$ be the Banach space of real-valued continuous functions over \mathcal{A} equipped with the sup-norm. The analogue of the transpose of the substitution matrix in the finite alphabet setting is the operator $M: E \rightarrow E$ given by $Mf(a) = \sum_{b \in \varrho(a)} f(b)$. Note that M is a bounded and positive operator on E , but is no longer necessarily a compact operator.

Theorem 2.4 ([27]). *For ϱ a primitive substitution on a compact Hausdorff alphabet \mathcal{A} ,*

- (1) (X_ϱ, σ) is minimal.
- (2) ϱ admits a unique (up to scaling) natural length function $\ell \in E$ which satisfies $\ell > 0$ and $M\ell(a) = \lambda\ell(a)$, where λ is the spectral radius of M . □

The second part of Theorem 2.4 is the compact alphabet analogue for the existence of a left PF eigenvector. The existence of a unique positive right PF eigenvector to λ is more subtle, and requires one to look at the dual operator $M^*: E^* \rightarrow E^*$, where E^* is the Banach dual of E . Here, M^* is defined via $M^*\psi(f) = \psi(Mf)$ for $\psi \in E^*$. The existence of σ -invariant measures on X_ϱ depends on the existence of M^* -invariant functionals in E^* [17, 27, 28]. For certain primitive substitutions, one has the following sufficient criteria for the subshift X_ϱ to be uniquely ergodic.

Theorem 2.5 ([27]). *Let ϱ be a substitution on a compact metrisable alphabet \mathcal{A} . Suppose that \mathcal{A} contains at least one isolated point, ϱ is primitive, recognisable and, for each isolated point $a \in \mathcal{A}$, $\varrho(a)$ contains only isolated points. Then, (X_ϱ, σ) is uniquely ergodic.* □

Theorem 2.6 ([28, Cor. 12.2]). *Let \mathcal{A} be a compact metrisable alphabet and ϱ a constant-length substitution over \mathcal{A} . Suppose ϱ is primitive and that the columns $\{\varrho_i\}_{0 \leq i \leq L-1}$ of ϱ generate an equicontinuous semigroup. Then, (X_ϱ, σ) is uniquely ergodic.* □

3. SPECTRAL THEORY

3.1. Diffraction and dynamical spectrum. For our spectral analysis, we restrict to substitutions of constant length, and so the corresponding point sets are supported on \mathbb{Z} . Let Λ_w be a coloured point set in \mathbb{Z} , which is the geometric realisation of $w \in X_\varrho$. Here, the label set is the alphabet \mathcal{A} and we can define a weighted Dirac comb ω via $\omega = \sum u(w_z)\delta_z$, where $u: \mathcal{A} \rightarrow \mathbb{C}$ is a bounded function. It follows from a general result regarding weighted lattice combs that the approximating autocorrelations for ω admit at least one limit point $\gamma_u = \sum_{m \in \mathbb{Z}} \eta(m)\delta_m$ [3, 8]. When the subshift is uniquely ergodic, one has that γ_u is the unique autocorrelation for all elements in the subshift X_ϱ , which is true for all cases we treat here. The diffraction measure is its Fourier transform $\widehat{\gamma}_u$. This is a positive measure which splits into pure point, absolutely continuous and singular continuous components $\widehat{\gamma}_{\text{pp}}$, $\widehat{\gamma}_{\text{ac}}$ and $\widehat{\gamma}_{\text{sc}}$, not all of which vanish. We refer the reader to [4] for an exposition of diffraction theory.

For a fixed weight function u , the autocorrelation coefficient $\eta_u(m)$ for $m \in \mathbb{Z}$ is given by

$$\eta_u(m) = \lim_{N \rightarrow \infty} \frac{1}{2N+1} \sum_{j=-N}^N u(w_j) \overline{u(w_{j+m})}.$$

It follows from a general result of Strungaru [30, Thm. 4.1] for weighted Dirac combs with Meyer set support Λ that the autocorrelation γ admits the generalised Eberlein decomposition

$$(1) \quad \gamma = \gamma_s + \gamma_{0a} + \gamma_{0s},$$

where γ_s, γ_{0a} , and γ_{0s} are pure point measures supported on $\Lambda - \Lambda$ that satisfy

$$\widehat{\gamma}_s = \widehat{\gamma}_{\text{pp}}, \quad \widehat{\gamma}_{0a} = \widehat{\gamma}_{\text{ac}}, \quad \widehat{\gamma}_{0s} = \widehat{\gamma}_{\text{sc}}.$$

Here, γ_s is the strongly almost periodic part and $\gamma_{0a} + \gamma_{0s}$ is the 0-weakly almost periodic part. If it can be shown that γ is a strongly almost periodic measure, it follows that $\widehat{\gamma}$ is a pure point measure. An equivalent criterion for γ to be strongly almost periodic is given by the following general result.

Theorem 3.1 ([8, Thm. 5]). *Let ω be a translation-bounded measure on a σ -compact locally compact abelian group G whose autocorrelation γ_ω exists and is of the form $\gamma_\omega = \sum_{m \in \Delta} \eta(m) \delta_m$, where Δ is Meyer. Then the autocorrelation is strongly almost periodic if and only if the set P_ε of ε -almost periods of $\eta(m)$ is relatively dense in Δ .* \square

Remark 3.2. Recently, Lenz, Spindeler and Strungaru proved an equivalent condition for pure point diffraction at the level of the measure ω , which requires ω to be *mean almost periodic* [26]. Here, we stick with the criterion in Theorem 3.1 because the autocorrelation coefficients are always complex-valued, meaning we can leverage the metric in \mathbb{C} , whereas mean almost periodicity requires comparing w_m and w_{j+m} , which might need some extra work if the alphabet is not metrisable. \diamond

Diffraction is intimately connected to the dynamical spectrum of X_ϱ . By dynamical spectrum, we mean the spectrum of the Koopman operator $U: L^2(X_\varrho, \mu) \rightarrow L^2(X_\varrho, \mu)$ associated with the shift action, with $U(f) = f \circ \sigma$. For $f \in L^2(X_\varrho, \mu)$, its *spectral measure* is the unique positive measure ρ_f on \mathbb{T} which satisfies $\langle U^n f | f \rangle = \int_{\mathbb{T}} z^n d\rho_f(z)$ for $n \in \mathbb{Z}$. A measure $\rho_{\max} \in \mathcal{M}(\mathbb{T})$ is called the *spectral measure of maximal type* of U if $\rho_f \ll \rho_{\max}$ for all f , where \ll denotes the absolute continuity relation for measures. By a slight abuse of notation, when $H \subseteq L^2(X, \mu)$ is a translation-invariant subspace, we write ρ_{\max}^H for the maximal spectral type of H , where $\rho_f \ll \rho_{\max}^H$ for all $f \in H$. Equivalently, this is the maximal spectral type of the subrepresentation $U|_H$.

3.2. Bijective Substitutions. We introduce the following definition as a natural extension of the notion from the finite alphabet case:

Definition 3.3. Given a constant-length substitution $\varrho: \mathcal{A} \rightarrow \mathcal{A}^+$, we say that ϱ is *bijective* if each column $\varrho_j(a) := (\varrho(a))_j$, for $0 \leq j \leq L-1$, is a bijection, and thus by continuity of ϱ a homeomorphism of \mathcal{A} . Let $\text{hom}(\mathcal{A})$ denote the set of homeomorphisms of \mathcal{A} , equipped with the compact-open topology. We let \mathcal{C} denote the subgroup of $\text{hom}(\mathcal{A})$ generated by the columns. The closure of \mathcal{C} in $\text{hom}(\mathcal{A})$ is denoted $\mathcal{G} := \overline{\mathcal{C}}$. We call ϱ *compact bijective* if \mathcal{G} is compact. A compact bijective substitution is called *abelian* if \mathcal{G} is abelian.

Remark 3.4. In the finite alphabet case, a bijective substitution is abelian if and only if the group generated by the columns is abelian. \diamond

Remark 3.5. In the general case, ϱ is abelian if and only if the columns commute and generate a relatively compact subgroup in $\text{hom}(\mathcal{A})$. Indeed, the algebraic closure of a commuting subset is abelian, as is the topological closure of an abelian subgroup. \diamond

Lemma 3.6. *Suppose that ϱ is primitive and compactly bijective. Then \mathcal{G} acts transitively on \mathcal{A} .*

Proof. Let $a, b \in \mathcal{A}$. We wish to find some $g \in \mathcal{G}$ with $g(a) = b$. Let $U \subset \mathcal{A}$ be an open set containing b . By primitivity, there is some $N \in \mathbb{N}$ so that $\varrho^N(a)$ contains a letter of U . Equivalently, we have that $(\varrho_{i_0} \circ \cdots \circ \varrho_{i_{N-1}})(a) \in U$ where each ϱ_{i_i} is a column of ϱ . It follows that we may construct a map $f_U \in \mathcal{C}$ so that $f_U(a) \in U$.

The open sets U containing b form a directed set, and so we have a net (f_U) . By compactness, we have that $f_U \rightarrow f$ for some $f \in \mathcal{G}$. Since $(f_U) \rightarrow f$ with respect to the compact-open topology it certainly converges pointwise, so in particular $(f_U(a))_U \rightarrow f(a)$. Then clearly $f(a) = b$, since $f_U(a) \in U$, where U is an arbitrary open set containing b . \square

Proposition 3.7. *Suppose that ϱ is primitive and compactly bijective. Then $\mathcal{A} \cong \mathcal{G}/H$, where $H \leq \mathcal{G}$ is a compact subgroup. With respect to this identification, we have*

$$\varrho[g] = (\varrho_0, \varrho_1, \dots, \varrho_{L-1})[g] = [\varrho_0 \cdot g][\varrho_1 \cdot g] \cdots [\varrho_{L-1} \cdot g],$$

where $[g]$ denotes the coset of $g \in \mathcal{G}$ in \mathcal{G}/H .

Proof. The evaluation map $C(X, Y) \times X \rightarrow Y$ is always continuous for X locally compact and Hausdorff and $C(X, Y)$ equipped with the compact-open topology. Hence the action $\mathcal{G} \times \mathcal{A} \rightarrow \mathcal{A}$ of \mathcal{G} on \mathcal{A} , given by $g \cdot a := g(a)$, is continuous. Since \mathcal{G} acts transitively on \mathcal{A} by Lemma 3.6, we have an identification between the \mathcal{G} -actions on \mathcal{A} and on \mathcal{G}/H by (left) multiplication, where

$$H := \text{stab}(p) := \{g \in \mathcal{G} \mid g(p) = p\}$$

and $p \in \mathcal{A}$ is an arbitrary basepoint. Stabiliser subgroups of topological group actions are always closed, so H is compact.

These actions are intertwined via the bijection

$$\phi: \mathcal{G}/H \rightarrow \mathcal{A}, \quad \phi[g] := g(p)$$

with inverse $a \mapsto [g]$, where g is such that $g(p) = a$ (uniquely defined up to coset representative). The map $\mathcal{G} \rightarrow \mathcal{A}$ defined by $g \mapsto g(p)$ is continuous, since, as above, evaluation is continuous. Hence, this descends to a continuous map on the quotient \mathcal{G}/H , so ϕ is continuous. By assumption, \mathcal{A} is Hausdorff and, since \mathcal{G} is compact, so is \mathcal{G}/H , hence ϕ is a homeomorphism.

By assumption, we have that $\varrho = (\varrho_0, \varrho_1, \dots, \varrho_{L-1})$, where each ϱ_i is a column and thus is an element of \mathcal{G} . Finally, writing ϱ as a substitution on \mathcal{G}/H by the identification ϕ , we have $\varrho[g]_i = \varrho(g(p))_i = \varrho_i(g(p)) = [\varrho_i \cdot g]$ and the result follows. \square

Corollary 3.8. *Let ϱ be a primitive, abelian, compactly bijective substitution. Then, we may equip \mathcal{A} with the structure of a compact group so that each column of the substitution is a group rotation. In particular, there exist $\beta_i \in \mathcal{A}$ so that*

$$\varrho(a) = (\beta_0 \cdot a)(\beta_1 \cdot a) \cdots (\beta_{L-1} \cdot a).$$

Proof. In applying Proposition 3.7 we have $H = \text{stab}(p)$ where p is an arbitrary base point. Then $H = \{\text{id}\}$ since ϱ is abelian. Indeed, let $g \in \text{stab}(p)$ and $x \in \mathcal{A}$ be arbitrary. By Lemma 3.6, the action of \mathcal{G} on \mathcal{A} is transitive, so there is some $h \in \mathcal{G}$ with $h(p) = x$. Then $g(x) = (gh)(p) = (hg)(p) = h(p) = x$ and thus $g = \text{Id}$. It follows from Proposition 3.7 that we may identify $\mathcal{A} \cong \mathcal{G}$ and that each column is a group rotation. \square

It follows that a constant-length substitution is compactly bijective and abelian if and only if we may equip \mathcal{A} with the structure of an abelian topological group so that each column of ϱ is a group translation. We will assume this structure for the remainder of this section. We let $S^1 \subset \mathbb{C}$ denote the unit circle, also considered as an abelian group. A unitary multiplicative *group character* χ is a continuous group homomorphism $\chi: \mathcal{A} \rightarrow S^1$. Here, we will use the characters χ as our weight functions.

Definition 3.9. Let ϱ be a substitution on a compact alphabet \mathcal{A} and let X_ϱ be the associated subshift. We call $x \in X_\varrho$ a *pseudo-fixed point* of ϱ if $\varrho(x) = \sigma^s(x)$, where $0 \leq s \leq L-1$.

For finite alphabets, the existence of a bi-infinite fixed point is guaranteed i.e. $x \in X_\varrho$ such that $\varrho^p(x) = x$ for some power $p \in \mathbb{N}$, which is convenient for diffraction. This fails in general for infinite alphabets. Here, we show that it suffices to consider pseudo-fixed points, as the associated recurrence relations are similar to those for two-sided fixed points. Pseudo-fixed points can be seen as true fixed points of the corresponding affine substitution; see [4, Sec. 4.8.1].

Proposition 3.10. *Let ϱ be a primitive constant-length substitution on a compact Hausdorff abelian group, such that all the columns are group translations. Then there exists another such substitution ϱ' such that $X_{\varrho'} = X_\varrho$ and ϱ' admits a pseudo-fixed point in X_ϱ .*

Proof. We write $\varrho := \varrho_0 \varrho_1 \cdots \varrho_{L-1}$, where $\varrho_i: \mathcal{A} \rightarrow \mathcal{A}$ are group translations. For $m \in \mathbb{N}$ we can define $\varrho' := \varrho^m \beta_s^{-1}$, where $\varrho_s^m(\theta) = \theta \beta_s$ for some $0 \leq s \leq L^m-1$, so that $\varrho'_s = \text{id}$. We see that ϱ and ϱ' define the

same subshift because they generate the same language. More explicitly, one has $(\varrho')^n(\theta\beta_j^n) = \varrho'^{mn}(\theta)$ for $\theta \in \mathcal{A}$ and $n \in \mathbb{N}$. For large enough m , we can take $1 \leq j \leq L^m - 2$ so that, by construction, ϱ' has a fixed internal letter at the j th position, i.e., $\varrho'(\theta)_j = \theta$. Using this, we construct a sequence of points which converges to a pseudo-fixed point.

Let $x^{(0)} \in X$ be such that $x_{[-j, L-1-j]}^{(0)} = \varrho(\theta)$ for some $\theta \in \mathcal{A}$. It then follows that $x_0^{(0)} = \theta$. If we substitute, we get that $(\sigma^{-s} \circ \varrho')(x_0)_{[-j, L-1-j]} = \varrho'(\theta)$. One can then consider the sequence $\{x^{(n)}\}_{n \geq 0}$, with $x^{(n)} = (\sigma^{-s} \circ \varrho')^n(x^{(0)})$. Due to the growing nested subwords around 0, this sequence converges to a point $x \in X$, which by construction satisfies $(\sigma^{-s} \circ \varrho')(x) = x$ and so x is a pseudo-fixed point for ϱ' . \square

Hence for us, without loss of generality, we may assume that ϱ admits a pseudo-fixed point.

Example 3.11. Let $\mathcal{A} = S^1$ and consider the substitution $\varrho: [\theta] \mapsto [\theta\alpha] [\theta\beta] [\theta\gamma]$, with $\alpha, \beta, \gamma \in \mathcal{A}$. We can instead consider

$$\varrho := \varrho': [\theta] \mapsto [\theta\alpha\beta^{-1}] [\theta] [\theta\gamma\beta^{-1}],$$

which has a fixed internal letter at position $s = 1$. Fix $\theta \in \mathcal{A}$ and let $x^{(0)}$ be the bi-infinite word given by $x^{(0)} = \dots [\theta\alpha\beta^{-1}] | [\theta] [\theta\beta^{-1}\gamma] \dots$. Here, the vertical line $|$ signifies the location of the origin. Substituting and shifting by $-s = -1$ yields

$$x^{(1)} = (\sigma^{-1} \circ \varrho)(x^{(0)}) = \dots [\theta\alpha\gamma^{-2}\beta] [\theta\alpha\beta^{-1}] | [\theta] [\theta\beta^{-1}\gamma] [\theta\alpha\beta^{-2}\gamma] \dots$$

By iterating the process, one fixes the letters in more positions, eventually exhausting all $m \in \mathbb{Z}$, which yields the limit x with $\varrho(x) = \sigma(x)$. \diamond

Let w be a pseudo-fixed point of ϱ and $\chi \in \widehat{\mathcal{A}}$ be a unitary multiplicative character. We consider the weighted Dirac comb $\omega_\chi = \sum_{m \in \mathbb{Z}} \chi(w_m) \delta_m$. The following result provides a sufficient condition for absence of absolutely continuous diffraction; compare [10, Thm. 6] for one-sided sequences.

Proposition 3.12. *Let ϱ be a substitution of constant length L on a compact Hausdorff abelian group \mathcal{A} and let w be a pseudo-fixed point of ϱ . Suppose ϱ_j is a group translation for $0 \leq j \leq L-1$. Then, for any unitary multiplicative character $\chi \in \widehat{\mathcal{A}}$, the weighted Dirac comb $\omega = \sum_{m \in \mathbb{Z}} \chi(w_m) \delta_m$ has no absolutely continuous diffraction.*

Proof. Since w is a pseudo-fixed point of ϱ , one has $\varrho(w) = \sigma^s(w)$ for some $0 \leq s \leq L-1$, meaning the entries of w satisfy the recurrence $w_{Lm+k-s} = w_m \beta_k$, for all $0 \leq k \leq L-1$ and $m \in \mathbb{Z}$. Fix $0 \leq k \leq L-1$, and let $\{\beta_j\}_{0 \leq j \leq L-1}$ be the translations defining ϱ . For any $m \in \mathbb{Z}$, one has

$$\begin{aligned} \eta(Lm+k) &= \lim_{N \rightarrow \infty} \frac{1}{2N+1} \sum_{r=-s}^{L-1-s} \sum_{j=-\lfloor \frac{N}{L} \rfloor - \varphi_r^-}^{\lfloor \frac{N}{L} \rfloor + \varphi_r^+} \chi(w_{Lj+r}) \overline{\chi(w_{L(j+m)+r+k})} \\ &= \lim_{N \rightarrow \infty} \frac{1}{2N+1} \sum_{r=-s}^{L-1-s} \sum_{j=-\lfloor \frac{N}{L} \rfloor - \varphi_r^-}^{\lfloor \frac{N}{L} \rfloor + \varphi_r^+} \chi(w_j w_{j+m+f(r,k)}^{-1}) \chi(\beta_{(r+s) \bmod L}) \overline{\chi(\beta_{(r+k+s) \bmod L})} \\ (2) \quad &= \frac{1}{L} \sum_{r=-s}^{L-1-s} \eta(m+f(r,k)) \chi(\beta_{(r+s) \bmod L}) \overline{\chi(\beta_{(r+k+s) \bmod L})}, \end{aligned}$$

where

$$(3) \quad f(r,k) := \begin{cases} 0, & r+k \leq L-1-s, \\ \lfloor \frac{r+k+s}{L} \rfloor, & \text{otherwise,} \end{cases} \quad \text{and} \quad \varphi_r^\pm := \begin{cases} 1, & N - L \lfloor \frac{N}{L} \rfloor \mp r \geq L, \\ 0, & 0 \leq N - L \lfloor \frac{N}{L} \rfloor \mp r \leq L-1, \\ -1, & \text{otherwise.} \end{cases}$$

Note that these linear recurrences are homogeneous and that, for $0 \leq m \leq L-1$, the values of $\eta(m)$ depend solely on the values $\eta(0)$ and $\eta(1)$. In particular, an explicit computation yields

$$(4) \quad \eta(1) = \frac{\eta(0) \sum_{r=0}^{L-2} \chi(\beta_r) \overline{\chi(\beta_{(r+1) \bmod L}}}{L(1 - \frac{1}{L} \chi(\beta_{L-1}) \overline{\chi(\beta_0)})}$$

$$(5) \quad \eta(L-1) = \frac{1}{L} \eta(0) \chi(\beta_0) \overline{\chi(\beta_{L-1})} + \frac{1}{L} \sum_{r=1}^{L-1} \eta(1) \chi(\beta_r) \overline{\chi(\beta_{(r+L-1) \bmod L})}$$

$$(6) \quad \eta(L^\ell m) = \eta(m), \quad \text{for any } m \in \mathbb{Z} \text{ and } \ell \in \mathbb{N}.$$

The key takeaway here is that the relations (4)-(6) no longer depend on s . Hence, our analysis proceeds as if we had a true fixed point. Due to the homogeneity of the relations, and the mutual measure-theoretic orthogonality of $(\widehat{\gamma})_{\text{pp}}$, $(\widehat{\gamma})_{\text{ac}}$ and $(\widehat{\gamma})_{\text{sc}}$, one finds that the individual components γ_s , γ_{0a} and γ_{0s} of the autocorrelation satisfy these relations independently; compare [5, Prop. 2].

We then consider $\gamma_{0a} = \sum_{m \in \mathbb{Z}} \eta_{\text{ac}}(m) \delta_m$ which gives rise to the ac component. By Eq. (6) and the Riemann–Lebesgue Lemma, one has $\eta_{\text{ac}}(m) = 0$, for all $m \geq 1$; compare [4, Prop. 8.6] and [15, Prop. 3.1]. To prove the claim, it remains to show that $\eta_{\text{ac}}(0) = 0$. Since $\eta_{\text{ac}}(1) = 0$, if $\eta_{\text{ac}}(0) \neq 0$, Eq. (5) yields $\eta_{\text{ac}}(L-1) = \frac{1}{L} \eta_{\text{ac}}(0) \chi(\beta_0) \overline{\chi(\beta_{L-1})} \neq 0$, which contradicts the previous observation. Hence, $\gamma_{0a} = 0$ and consequently $\widehat{\gamma}_{\text{ac}} = 0$. \square

Remark 3.13. The set of recurrences in Eq. (2) imply that the sequence $\{\eta(m)\}_{m \geq 0}$ associated to such a substitution is L -regular over \mathbb{C} in the sense of Allouche and Shallit; compare [2, Ex. 14]. \diamond

We now give a sufficient criterion for the absence of pure point diffraction components. In our setting, the set P_ε of ε -almost periods for η is given by

$$(7) \quad P_\varepsilon = \{m \in \mathbb{Z} \mid |\eta(0) - \eta(m)|^{1/2} < \varepsilon\}.$$

Theorem 3.14. *Let ϱ be as in Proposition 3.12 such that $|\eta(1)| < 1$. Then, the diffraction associated to ω does not have a pure point component.*

Proof. Suppose $\widehat{\gamma}_{\text{pp}} \neq 0$. Then, by Eq. (1), $\gamma_s = \sum_{m \in \mathbb{Z}} \eta_{\text{pp}}(m) \delta_m \neq 0$. Because γ_s is a strongly almost periodic measure, Theorem 3.1 implies that the set P_ε of ε -almost periods of $\eta_{\text{pp}}(m)$ given in Eq. (7) must be relatively dense, for all $\varepsilon > 0$. We show that $|\eta(1)| < 1$ implies that $|\eta_{\text{pp}}(m)|$ is uniformly bounded away from $|\eta_{\text{pp}}(0)|$, which directly contradicts the relative denseness of P_ε , for all $\varepsilon \leq \varepsilon'$, for some ε' .

The autocorrelation coefficients of each of the three components of γ must satisfy Eq. (2) independently since the corresponding component measures are mutually singular. With this, the assumption on $\eta(1)$ and Eq. (4) imply $|\eta_{\text{pp}}(1)| < |\eta_{\text{pp}}(0)|$. Suppose $|\eta_{\text{pp}}(0)| - |\eta_{\text{pp}}(1)| > c$, for some $c > 0$. Restricting the recurrences in Eq. (2) to γ_s one gets for $1 \leq k \leq L-1$,

$$|\eta_{\text{pp}}(k)| \leq \frac{1}{L} \sum_{r=0}^{L-1} \left| \eta_{\text{pp}}(f(r, k)) \right| = \frac{L-k}{L} |\eta_{\text{pp}}(0)| + \frac{k}{L} |\eta_{\text{pp}}(1)| < |\eta_{\text{pp}}(0)| - \frac{k}{L} c$$

which means

$$(8) \quad |\eta_{\text{pp}}(0)| - |\eta_{\text{pp}}(k)| > \frac{1}{L} c.$$

Now let $L \leq k = Lm + j \leq L^2 - 1$, with $1 \leq m \leq L-2$ and $0 \leq j \leq L-1$. Using the same arguments as in the previous step, one obtains

$$\begin{aligned} |\eta_{\text{pp}}(k)| &\leq \frac{1}{L} \sum_{r=0}^{L-1} \left| \eta_{\text{pp}}(m + f(r, j)) \right| = \frac{L-j}{L} |\eta_{\text{pp}}(m)| + \frac{j}{L} |\eta_{\text{pp}}(m+1)| \\ &< \frac{L-j}{L} \left(|\eta_{\text{pp}}(0)| - \frac{m}{L} c \right) + \frac{j}{L} \left(|\eta_{\text{pp}}(1)| - \frac{m+1}{L} c \right) \\ &< |\eta_{\text{pp}}(0)| - \frac{Lm+j}{L^2} c. \end{aligned}$$

Since $\frac{1}{L} \leq \frac{Lm+i}{L^2} \leq \frac{L^2-1}{L^2}$, we recover Eq. (8) for all $L \leq k \leq L^2 - 1$. It is straight-forward to show via an inductive argument that Eq. (8) holds for all $k \in \mathbb{Z}$. This means

$$|\eta_{\text{pp}}(0) - \eta_{\text{pp}}(k)| \geq \left| |\eta_{\text{pp}}(0)| - |\eta_{\text{pp}}(k)| \right| > \frac{1}{L}c,$$

which implies $P_\varepsilon = \emptyset$, for all $\varepsilon < \sqrt{\frac{c}{L}}$, contradicting the relative denseness property required for pure point diffraction. Hence, $\gamma_s = 0$ and $\widehat{\gamma}_{\text{pp}} = 0$. \square

The following corollary is immediate from Proposition 3.12 and Theorem 3.14.

Corollary 3.15. *Let ϱ be as in Proposition 3.12, w a pseudo-fixed point of ϱ , and ω the associated weighted Dirac comb via the weight function $\chi \in \widehat{\mathcal{A}}$. If $|\eta(1)| < 1$, the diffraction $\widehat{\gamma}$ of ω is purely singular continuous.* \square

Lemma 3.16. *Let ϱ be as in Proposition 3.12. Fix $\chi \in \widehat{\mathcal{A}}$ and suppose further $\chi(\beta_0) \neq \chi(\beta_{L-1})$. Then, one has $|\eta(1)| < 1$.*

Proof. This follows directly from Eq. (4) since, $|\eta(1)| \leq \frac{|\eta(0)|(L-1)}{|L - \chi(\beta_{L-1})\overline{\chi(\beta_0)}|} < 1$, where the last inequality holds since the denominator satisfies $|L - \chi(\beta_{L-1})\overline{\chi(\beta_0)}| > L - 1$, which follows from the assumption $\chi(\beta_{L-1})\overline{\chi(\beta_0)} = \chi(\beta_{L-1}\beta_0^{-1}) \neq 1$. \square

Remark 3.17. Note that when $\mathcal{A} = S^1$ the condition in Lemma 3.16 is satisfied for all $\chi \in \widehat{\mathcal{A}}$ whenever $\beta_0\beta_{L-1}^{-1}$ is an irrational rotation because $\ker(\chi)$ is finite for all $\chi \in \widehat{S^1}$. \diamond

Theorem 3.18. *Let ϱ be a primitive substitution of length L on a compact Hausdorff abelian group \mathcal{A} such that all of its columns are group translations. Let w be a pseudo-fixed point of ϱ and consider $\chi \in \widehat{\mathcal{A}}$. Then, the following are equivalent:*

- (1) $\widehat{\gamma}_\chi$ is not purely singular continuous;
- (2) $|\eta(1)| = 1$;
- (3) The Dirac comb $\omega_\chi = \sum_{m \in \mathbb{Z}} \chi(w_m)\delta_m$ can be generated by a substitution ϱ_{cyc} on a finite cyclic group \mathcal{A} of order n , where ϱ is of the form

$$(9) \quad \varrho_{\text{cyc}}: [\theta] \mapsto [\theta\beta_0] [\theta\beta_0\beta_1] \cdots [\theta\beta_0\beta_1^{n-1}] [\theta\beta_0],$$

where $\beta_0, \beta_1 \in \mathcal{A}$ and $\text{ord}(\beta_1) = n$;

- (4) ω_χ is periodic;
- (5) $\widehat{\gamma}_\chi$ is pure point.

Before we go to the proof of Theorem 3.18, we will need the following Lemma; see [21, Cor. 4.3].

Lemma 3.19. *The substitution ϱ_{cyc} of Eq. (9) defines a periodic subshift.*

Proof. Every 2-letter subword of a substituted letter is of the form $[\theta][\beta_1\theta]$ for some $\theta \in \mathcal{A}$. Moreover, given a pair of this form,

$$\varrho_{\text{cyc}}([\theta] \bullet [\beta_1\theta]) = [\theta\beta_0] \cdots [\theta\beta_0] \bullet [\beta_1\theta\beta_0] \cdots [\beta_1\theta\beta_0],$$

where we emphasise the location of the boundary of supertiles. Note that the pair meeting over the supertile boundary is of the same form. Every 2-letter word generated by ϱ_{cyc} is either as a subword of some $\varrho_{\text{cyc}}(\theta)$, or as the pair at the supertile boundary of $\varrho_{\text{cyc}}(\theta\theta')$, where $\theta\theta'$ is also generated by ϱ_{cyc} . It follows that all legal 2-letter words are of the form $[\theta][\beta_1\theta]$. Then periodicity follows from $\text{ord}(\beta_1) = \#\mathcal{A} = n$. \square

Proof of Theorem 3.18.

- (i) (3) \implies (4) follows from Lemma 3.19.
- (ii) (4) \implies (5) is immediate from the periodicity of ω_χ .
- (iii) (5) \implies (1) is clear.

(iii) **(1)** \implies **(2)** follows from Theorem 3.14.

It then remains to show that **(2)** implies **(3)**. Suppose $|\eta(1)| = 1$. We write $\varrho := \varrho_0 \varrho_1 \cdots \varrho_{L-1}$, with $\varrho_i(\theta) := \theta \beta_i$ for $\beta_i \in \mathcal{A}$. From the proof of Proposition 3.10, we can assume that $\beta_s = \text{id}$ for some $1 \leq s \leq L-2$, since multiplying all β_i with β_s^{-1} yields a substitution which generates the same subshift. From Lemma 3.16, we can assume that $\chi(\beta_0) = \chi(\beta_{L-1})$, since otherwise $|\eta(1)| < 1$. From Eq. (4), one must then have

$$\left| \sum_{r=0}^{L-2} \chi(\beta_r) \overline{\chi(\beta_{(r+1) \bmod L})} \right| = L-1,$$

which only holds when $\chi(\beta_j \beta_{j+1}^{-1})$ is constant for all $0 \leq j \leq L-2$. Since $\beta_s = \text{id}$, we obtain $\chi(\beta_{s+1}^{-1}) = \chi(\beta_{s+1} \beta_{s+2}^{-1})$. Continuing inductively yields $\chi(\beta_r) = \chi(\beta_{s+1})^{r-s}$ for $s+1 \leq r \leq L-1$. We also have $\chi(\beta_r) = \chi(\beta_{s+1})^{L-(s+1-r)}$ for $0 \leq r \leq s$. It follows that $\chi(\beta_{s+1})^{L-1} = \text{id}$, so $\chi(\beta_{s+1}) \in S^1$ must be a root of unity of order at most $L-1$. Now, let $g_0 = \chi(\beta_0)$, $g = \chi(\beta_{s+1})$ and $n = \text{ord}(\chi(\beta_{s+1}))$ and consider $\mathcal{A}' = \langle g \rangle \simeq C_n$. One can check that the substitution $\varrho' : C_n \rightarrow C_n^+$ given by

$$\varrho' : [a] \mapsto [ag_0] [ag_0g] [ag_0g^2] \cdots [ag_0g^{n-1}] [ag_0]$$

is primitive and yields the exact same recurrence relations for the autocorrelation $\eta(m)$ as ϱ , which completes the proof. \square

As a remarkable consequence, for a pseudo-fixed point w of an abelian bijective substitution ϱ on a compact alphabet, if $\chi(w_j)$ admits infinitely many complex values, then $|\eta(1)| < 1$ and the diffraction $\widehat{\gamma}_\chi$ is purely singular continuous.

Example 3.20. Let $\mathcal{A} = S^1$ and define the substitution $\varrho_1 : \mathcal{A} \mapsto \mathcal{A}^+$ via

$$\varrho_1 : [\theta] \mapsto [\theta] \ [\theta] \ [\theta\alpha] \ [\theta]$$

where $\alpha \in \mathcal{A}$. Consider the corresponding bi-infinite fixed point w associated to the legal seed $1|1$. It suffices to look at the one-sided fixed point, which satisfies the recurrences

$$(10) \quad \begin{array}{ll} w_{4m} = w_m & w_{4m+2} = \alpha w_m \\ w_{4m+1} = w_m & w_{4m+3} = w_m \end{array}$$

where w_m is the letter at position $m \in \mathbb{N}$. Here, we write $\alpha = e^{2\pi i \varphi}$ for some $\varphi \in [0, 1)$.

Let $\chi_n \in \widehat{S^1}$ be given by $\chi_n(\theta) = \theta^n$ for $\theta \in S^1$. One can show by direct computation that $\eta_n(1) = \eta_n(4^r) = \frac{1}{3} + \frac{2}{3} \cos(2\pi n \varphi)$ for any $r \in \mathbb{N}$. This implies that $|\eta(1)| = 1$ only when $\varphi \in \mathbb{Q}$ with $n\varphi \in \mathbb{Z}$. It follows from Theorem 3.18 that, for all $\alpha \in (0, 1)$ not satisfying this condition, the diffraction of ω_{χ_n} is purely singular continuous for any $x \in X_\varrho$. When α is an n th root of unity, $\omega_{\chi_n} = \delta_{\mathbb{Z}}$, which obviously has pure point diffraction. \diamond

Example 3.21 (Substitution with a non-trivial pure point factor). Here, we demonstrate how the characters and their corresponding diffraction can reveal interesting factors. Let $\mathcal{A} = C_2 \times S^1$ with $C_2 = \{e, g\}$ and consider ϱ which is given by $\varrho : [\theta] \mapsto [\theta\beta_0][\theta\beta_1][\theta\beta_2]$ with $\beta_0 = \beta_2 = (e, 1)$ and $\beta_1 = (g, \alpha)$, where α is an irrational rotation. Let $\chi_1 \otimes \chi_2 \in \widehat{\mathcal{A}} \simeq C_2 \times \mathbb{Z}$. When χ_1 is non-trivial and χ_2 is trivial, this character induces a factor map to the substitution on C_2 given by $\varrho' : [a] \mapsto [a][ag][a]$, which by Theorem 3.18 is periodic. When χ_2 is non-trivial, $\chi_1 \otimes \chi_2(w)$ always admits infinitely many complex values due to the irrationality of α and hence has singular continuous diffraction by Theorem 3.18. This means the maximal equicontinuous factor of ϱ must be $(\mathbb{Z}_3 \times C_2, T_1 \times T_2)$, where \mathbb{Z}_3 is the 3-adic integers and T_1 is the $+1$ -map and C_2 is endowed with the group multiplication $T_2 : h \rightarrow gh$ by g ; compare [15] and Section 3.4. \diamond

3.3. Substitutions with coincidences.

Definition 3.22. Let ϱ be a constant-length substitution on a compact alphabet \mathcal{A} . We say that ϱ admits a *coincidence* if for some $0 \leq j \leq L-1$, $\varrho(\theta)_j = \beta_0$, for all $\theta \in \mathcal{A}$, where β_0 is a constant.

As in Proposition 3.10, it is routine to show that when one has at least one coincidence and the other columns are group translations, ϱ also admits a pseudo-fixed point.

Theorem 3.23. Let ϱ be a substitution of length L on a compact Hausdorff abelian group \mathcal{A} such that $\varrho(\theta)_j$ is either a group translation or is constant, for $0 \leq j \leq L-1$, with at least one constant column. Then, for any $\chi \in \widehat{\mathcal{A}}$, the weighted Dirac comb $\omega_\chi = \sum_{m \in \mathbb{Z}} \chi(w_m) \delta_m$ arising from a pseudo-fixed point w has pure point diffraction.

Proof. Without loss of generality, choose s to be the minimal position with a coincidence and let $w \in X_\varrho$ be such that $\varrho(w) = \sigma^s(w)$. Suppose that there are p coincident columns in ϱ , which are at positions $\{c_1 = s, c_2, \dots, c_p\}$.

Consider $\eta(Lm)$, which can be written as

$$\eta(Lm) = \lim_{N \rightarrow \infty} \frac{1}{2N+1} \sum_{r=-s}^{L-1-s} \sum_{j=-\lfloor \frac{N}{L} \rfloor - \varphi_r}^{\lfloor \frac{N}{L} \rfloor + \varphi_r} \chi \left(w_{Lj+r} w_{L(j+m)+r}^{-1} \right),$$

where φ_r is the same as in Section 3.2. The constant columns will yield p recurrence relations for w_j , which read $w_{Lm+c_i-s} = \beta_{c_i}$. For the remaining positions $0 \leq k \leq L-1$, which are group translations, one has $w_{Lm+k-s} = w_m \beta_k$ as in the bijective case. Taking these recurrences together yields

$$\begin{aligned} \eta(Lm) &= \lim_{N \rightarrow \infty} \frac{1}{2N+1} \left[p \sum_{j=-\lfloor \frac{N}{L} \rfloor - \varphi_r}^{\lfloor \frac{N}{L} \rfloor + \varphi_r} \chi(e) + (L-p) \sum_{j=-\lfloor \frac{N}{L} \rfloor - \varphi_r}^{\lfloor \frac{N}{L} \rfloor + \varphi_r} \chi(w_j w_{j+m}^{-1}) \right] \\ (11) \quad &= \frac{p}{L} + \frac{L-p}{L} \eta(m). \end{aligned}$$

The unitarity of χ implies that $\eta(0) = 1$. To use Theorem 3.1, we then need to show that

$$P_\varepsilon = \{m \in \mathbb{Z} \mid |1 - \eta(m)|^{1/2} < \varepsilon\}$$

is relatively dense, for all $\varepsilon > 0$. To this end, let $\eta(m) = a + ib \in \mathbb{C}$. Applying Eq. 11 iteratively yields

$$\eta(L^\ell m) = \left(\frac{L-p}{L} a_{\ell-1} + \frac{p}{L} \right) + i \left(\frac{L-p}{L} \right)^\ell b,$$

where $a_\ell = \frac{L-p}{L} a_{\ell-1} + \frac{p}{L}$ with $a_0 = a$. Since $|\eta(m)| \leq 1$ for all m , one has $a, b \in [-1, 1]$, implying that, for any fixed $m \in \mathbb{Z}$, $\Re(\eta(L^\ell m)) \rightarrow 1$ and $\Im(\eta(L^\ell m)) \rightarrow 0$ as $\ell \rightarrow \infty$, with uniform convergence in $[-1, 1]$. Given $\varepsilon > 0$, and $z = L^\ell m$ for some $\ell \in \mathbb{N}$ and $m \in \mathbb{Z}$, one has

$$|1 - \eta(z)|^{1/2} = \left| \left(1 - \frac{L-p}{L} a_{\ell-1} + \frac{p}{L} \right) - i \left(\frac{L-p}{L} \right)^\ell b \right|^{1/2} \leq |\varepsilon_1^{(\ell)} + \varepsilon_2^{(\ell)}|^{1/2} \leq \sqrt{2} \max \left\{ \sqrt{\varepsilon_1^{(\ell)}}, \sqrt{\varepsilon_2^{(\ell)}} \right\}.$$

One can then choose $\ell_0 \in \mathbb{N}$ such that for all $\ell > \ell_0$, $\max \left\{ \sqrt{\varepsilon_1^{(\ell)}}, \sqrt{\varepsilon_2^{(\ell)}} \right\} < \frac{\varepsilon}{\sqrt{2}}$. Since this choice of ℓ_0 can be made to cover all possible values of $a, b \in [-1, 1]$, one has $|1 - \eta(z)|^{1/2} < \varepsilon$ for the elements of the set $\mathcal{Z}^{(\ell_0)} := \{z \in \mathbb{Z} \mid z = L^\ell m, m \in \mathbb{Z}, \ell \geq \ell_0\}$. Clearly, $\mathcal{Z}^{(\ell_0)}$ is relatively dense in \mathbb{Z} , and hence $\widehat{\gamma}_\chi$ is pure point by Theorem 3.1. \square

The following result follows from [6, Thm. 7], which establishes the equivalence of a measure-preserving dynamical system (X, S, μ) having pure point dynamical spectrum and the diffraction being pure point (which in fact holds without any ergodicity assumptions on μ).

Corollary 3.24. *Let ϱ be as in Theorem 3.23 such that one of the columns is a group translation of infinite order. Then ϱ is primitive and (X_ϱ, σ) is strictly ergodic with pure point dynamical spectrum. \square*

Here, minimality follows from primitivity and unique ergodicity follows again from Theorem 2.6 since the columns of ϱ generate an equicontinuous semigroup.

Example 3.25. Consider the substitution $\varrho_2: \mathcal{A} \rightarrow \mathcal{A}^+$

$$\varrho_2: [\theta] \mapsto [\theta] [1] [\theta\alpha] [\theta],$$

where $\mathcal{A} = S^1$, and the corresponding bi-infinite fixed point w associated to the legal seed $1|1$. For this example, the recurrence relations are given by

$$(12) \quad \begin{aligned} w_{4m} &= w_m & w_{4m+2} &= \alpha w_m \\ w_{4m+1} &= 1 & w_{4m+3} &= w_m. \end{aligned}$$

As before, we assign the weight $\chi_n(\theta) = \theta^n$ to each point of type θ . Theorem 3.23 tells us that the corresponding diffraction for ϱ_2 is pure point. \diamond

Remark 3.26. Substitutions covered in Theorem 3.23 generate sequences exhibiting a generalised Toeplitz structure; see [13, Sec. 4.3] on generalised Toeplitz sequences over compact alphabets. \diamond

3.4. A substitution with countably infinite Lebesgue and countably infinite singular continuous spectral components. Here, we provide a generalisation of the Rudin–Shapiro substitution on infinite alphabets. Let $\mathcal{A} = G \times \mathcal{D}$, with $\mathcal{D} = \{0, 1\}$ and $G = S^1$. Let $\alpha \in S^1$ be an irrational rotation and consider $\varrho: \mathcal{A} \rightarrow \mathcal{A}^+$ given by

$$(13) \quad \varrho: \begin{cases} (\theta, 0) \mapsto (\theta, 0) (-\theta, 1) \\ (\theta, 1) \mapsto (\theta\alpha, 0) (\theta\alpha, 1). \end{cases}$$

Such a substitution is called a *spin substitution* by Frank and Mañibo [16]. For $\mathbf{a} = (\theta, \mathbf{d}) \in \mathcal{A}$, we call $\pi_G(\mathbf{a}) = \theta$ its *spin* and $\pi_{\mathcal{D}}(\mathbf{a}) = \mathbf{d}$ its *digit*. The distribution of the spins in the level-1 substituted words are determined by the spin matrix

$$W = \begin{pmatrix} 1 & -1 \\ \alpha & \alpha \end{pmatrix},$$

which satisfies $W_{ij} = \pi_G(\varrho(1, i)_j)$ for $i, j \in \mathcal{D}$. Primitivity follows from the irrationality of α . Note that this substitution is ‘recognisable’ because the image of any letter with digit 1 has the same spin for its first and second letter. One can show that the associated subshift is measure-theoretically isomorphic to an S^1 -extension of the dyadic odometer, i.e., $(X_\varrho, \mathbb{Z}, \mu) \simeq (\mathbb{Z}_2 \times S^1, \mathbb{Z}, \nu \times \mu_{\mathbb{H}})$, where ν and $\mu_{\mathbb{H}}$ are the Haar measures on \mathbb{Z}_2 and S^1 , respectively. The skew product which induces the \mathbb{Z} -action can be derived directly from W ; see [16]. It follows from the classical theory of group extensions that one has the induced splitting $L^2(X_\varrho, \mu) = \bigoplus_{\chi \in \widehat{S^1}} H_\chi$ with $H_\chi = L^2(\mathbb{Z}_2, \nu) \otimes \chi$. The subspaces H_χ are translation-invariant. The spectral type of functions in H_χ is determined by $\chi(W) := (\chi(W_{ij}))_{i,j}$ via the following result.

Theorem 3.27 ([16]). *Let ϱ be a primitive recognisable spin substitution on $\mathcal{A} = G \times \mathcal{D}$ with spin map W , where G is a compact abelian group.*

- (1) *If $\frac{1}{|\mathcal{D}|} \chi(W)$ is a unitary matrix, H_χ has Lebesgue spectral type of multiplicity $|\mathcal{D}|$.*
- (2) *If $\chi(W)$ is a rank-1 matrix, χ induces a factor map onto a bijective abelian substitution ϱ' . Moreover, the maximal spectral type of H_χ is either pure point or purely singular continuous, and is absolutely continuous to the maximal spectral type of ϱ' . \square*

Proposition 3.28. *Let ϱ be the spin substitution in Eq. (13). Let $\chi_n \in \widehat{S^1} \simeq \mathbb{Z}$, with $\chi_n(z) = z^n$. Consider the weighted Dirac comb $\omega_{\chi_n} = \sum_{m \in \mathbb{Z}} \chi_n(\pi_G(w_m)) \delta_m$, where $w \in X_\varrho$. Let $\widehat{\gamma}^{(n)}$ be the diffraction measure of ω_{χ_n} . Then one has the following.*

- (1) *If $n = 0$, the diffraction $\widehat{\gamma}^{(n)} = \delta_{\mathbb{Z}}$ and hence pure point.*

- (2) If $n \in 2\mathbb{Z} + 1$, the diffraction $\widehat{\gamma}^{(n)}$ is Lebesgue measure.
- (3) If $n \in 2\mathbb{Z} \setminus \{0\}$, the diffraction $\widehat{\gamma}^{(n)}$ is purely singular continuous and is a generalised Riesz product.

Proof. The first claim is straightforward. Before we proceed, we note that $\widehat{\gamma}^{(n)} = \rho_f * \delta_{\mathbb{Z}}$, where ρ_f is the spectral measure of the function $f: x \mapsto \chi_n(\pi_G(x_0))$, which is always in H_{χ_n} . To prove the second claim, note that $\frac{1}{2}\chi_n(W)$ is a unitary matrix whenever n is a nonzero odd integer. From Theorem 3.27, H_{χ} contains only functions whose spectral measures are absolutely continuous with respect to Lebesgue measure. For the third claim, the matrix $\chi(W)$ is rank-1 whenever n is even. For $n \neq 0$, the character χ_n induces a factor map from X_{ϱ} onto $(S^1)^{\mathbb{Z}}$ which identifies the letters in $\{(\zeta\alpha^n, 0)\}$ with $(1, 1)$, where ζ is an n th root of unity. The image of X_{ϱ} under χ_n can be realised as the subshift of the bijective substitution on S^1 given by $[\theta] \mapsto [\theta][\theta\alpha^n]$, which is primitive (since α is irrational) and recognisable. By Theorem 3.18, the diffraction measure $\widehat{\gamma}^{(n)}$ is purely singular continuous. The corresponding generalised Riesz product is given by $\rho = \prod_{m \geq 0} \frac{1}{2}|1 + \alpha^n e^{2^{m+1}\pi i t}|^2$, seen as a weak-* limit of absolutely continuous measures on \mathbb{T} , which arises from the relations $w_{2m} = w_m$ and $w_{2m+1} = \alpha^n w_m$; see [28, Prop. 4.13]. \square

Theorem 3.29. *Let ϱ be the spin substitution in Eq. (13) and consider $L^2(X_{\varrho}, \mu) = \bigoplus_{\chi_n \in \widehat{S^1}} H_{\chi_n}$. Let $\rho_{\max}^{(n)}$ denote the maximal spectral type of H_{χ_n} .*

- (1) If $n = 0$, $\rho_{\max}^{(n)}$ is pure point.
- (2) If $n \in 2\mathbb{Z} + 1$, $\rho_{\max}^{(n)}$ is Lebesgue measure.
- (3) If $n \in 2\mathbb{Z} \setminus \{0\}$, $\rho_{\max}^{(n)}$ is purely singular continuous.

Proof. It is a well known result for abelian group extensions that the restriction of U_T on any H_{χ_n} is spectrally pure, i.e., $\rho_{\max}^{(n)}$ is either pure point, absolutely continuous or singular continuous [19]. It then suffices to find the spectral type of a single function $f_n \in H_{\chi_n}$ to determine that of $\rho_{\max}^{(n)}$. From the proof of the previous proposition, we know that $f_n(x) = \chi_n(\pi_G(x_0))$ is in H_{χ_n} . Since the spectral type of σ_{f_n} is the same as the spectral type of the diffraction measure $\widehat{\gamma}^{(n)}$, the claim follows from Proposition 3.28. \square

Remark 3.30. As a consequence, one can form functions in $L^2(X, \mu)$ whose spectral measures have arbitrary (finite) number of absolutely continuous and singular continuous components. As an example, for the hyperlocal function $f(x) = (\chi_0 + \chi_1 + \chi_3 + \chi_4)(\pi_G(x_0))$, the spectral measure ρ_f decomposes into $\rho_f = \rho_0 + \rho_1 + \rho_3 + \rho_4$, where $\rho_i \perp \rho_j$ for $i \neq j$ and where ρ_0 is pure point, ρ_1 and ρ_3 are absolutely continuous, and ρ_4 is singular continuous. \diamond

3.5. Non-constant length example. Let $\mathcal{A} = \overline{\mathbb{N}_0} = \mathbb{N}_0 \cup \{\infty\}$ denote the one-point compactification of the natural numbers and consider the substitution

$$\varrho: \begin{cases} [0] \mapsto [0][1] \\ [n] \mapsto [0][n+1][n-1], & 0 < n < \infty \\ [\infty] \mapsto [0][\infty][\infty]. \end{cases}$$

This is a non-constant length substitution. One can easily check that it is primitive, which implies X_{ϱ} is minimal by Theorem 2.4. Moreover, since isolated points only get mapped to isolated points under ϱ , by Theorem 2.5, the associated subshift X_{ϱ} is uniquely ergodic. Primitivity also guarantees the existence of a strictly positive length function. In this case, the inflation factor is $\lambda = \frac{5}{2}$ and the length function $\ell: \mathcal{A} \rightarrow \mathbb{R}$ is given by $\ell(n) = 2 - 1/2^n$ and $\ell(\infty) = 2$. The letter frequencies are also well defined and are given by $\nu_n = \frac{1}{2^{n+1}}$ and $\nu_{\infty} = 0$. The existence of natural tile lengths allows one to construct a point set Λ from the bi-infinite fixed point $w = \varrho^{\infty}(\infty | 0)$. The length function is bounded from above and below, with $1 = \ell(0) < \ell(n) < \ell(n+1) < \ell(\infty) = 2$ for $n \in \mathcal{A}$. This implies that the point set Λ is both uniformly discrete and relatively dense, and hence is a Delone set. Since there are infinitely many distinct tile lengths, Λ is not Meyer. We call Λ a *Delone set of infinite type* with inflation symmetry, i.e., $\lambda\Lambda \subseteq \Lambda$. Lagarias proved in [22] that, if Λ is of finite type and has inflation symmetry, then λ must

be an algebraic integer. For the example above, $\lambda = \frac{5}{2}$ is algebraic but is not an integer. This leads to the following question:

Question 3.31. Let ϱ be a primitive substitution on a compact alphabet with inflation factor λ . Is it possible for λ to be a transcendental number?

For primitive substitutions on finite alphabets, it is well known that the diffraction measure of a weighted Dirac comb supported on a Delone set Λ arising from ϱ has non-trivial pure point component if and only if λ is Pisot [18]. Moreover, if $\widehat{\gamma}$ is pure point, Λ must be a Meyer set [24]. Note however that a general Delone set need not be Meyer for it to have a non-trivial pure point component; see [17, 20] for the “scrambled Fibonacci” example, which arises from a fusion rule with finite local complexity.

Most proofs of existence of non-trivial pure point component (Bragg peaks in the diffraction setting, eigenvalues in the dynamical setting) for point sets with some form of hierarchical structure rely on the Diophantine properties of the return vectors which generate the translation module. When the Delone set is of finite type, i.e., there are only finitely many distinct tile lengths, this \mathbb{Z} -module is finitely generated. For the example above, the translation module is infinitely generated and it is not clear whether the criteria for the existence of eigenvalues extend to this setting. This will be tackled in future work.

ACKNOWLEDGEMENTS

The authors would like to thank Michael Baake, Michael Coons, Natalie Priebe Frank, Franz Gähler, Christoph Richard and Nicolae Strungaru for helpful discussions. NM is funded by the German Research Foundation (DFG, Deutsche Forschungsgemeinschaft), via SFB 1283/2 2021–317210226. DR would like to acknowledge the support of the DFG via SFB1283/1. JW acknowledges the support of the EPSRC grant EP/R013691/1.

REFERENCES

- [1] I. Abou and P. Liardet, *Flots chaînés*, Proceedings of the Sixth Congress of Romanian Mathematicians. Vol. 1, Ed. Acad. Române, Bucharest, 2009, pp. 401–432.
- [2] J.-P. Allouche and J. Shallit, *The ring of k -regular sequences. II*, Theoret. Comput. Sci. **307** (2003), no. 1, 3–29.
- [3] M. Baake, *Diffraction of weighted lattice subsets*, Canad. Math. Bull. **45** (2002), no. 4, 483–498.
- [4] M. Baake and U. Grimm, *Aperiodic Order. Volume 1: A Mathematical Invitation*, Cambridge Univ. Press, 2013.
- [5] ———, *Squirals and beyond: substitution tilings with singular continuous spectrum*, Ergodic Theory Dynam. Systems **34** (2014), no. 4, 1077–1102.
- [6] M. Baake and D. Lenz, *Dynamical systems on translation bounded measures: pure point dynamical and diffraction spectra*, Ergodic Theory Dynam. Systems **24** (2004), no. 6, 1867–1893.
- [7] M. Baake, D. Lenz, and A. van Enter, *Dynamical versus diffraction spectrum for structures with finite local complexity*, Ergodic Theory Dynam. Systems **35** (2015), no. 7, 2017–2043.
- [8] M. Baake and R. V. Moody, *Weighted Dirac combs with pure point diffraction*, J. Reine Angew. Math. **573** (2004), 61–94.
- [9] A. Bartlett, *Spectral theory of \mathbb{Z}^d substitutions*, Ergodic Theory Dynam. Systems **38** (2018), no. 4, 1289–1341.
- [10] J. Coquet, T. Kamae, and M. Mendès France, *Sur la mesure spectrale de certaines suites arithmétiques*, Bull. Soc. Math. France **105** (1977), no. 4, 369–384.
- [11] F. M. Dekking, *The spectrum of dynamical systems arising from substitutions of constant length*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **41** (1977/78), 221–239.
- [12] F. Durand, N. Ormes, and S. Petite, *Self-induced systems*, J. Anal. Math. **135** (2018), no. 2, 725–756.
- [13] B. Eichinger and P. Gohlke, *Spectral properties of Schrödinger operators associated with almost minimal substitution systems*, Ann. Henri Poincaré **22** (2021), no. 5, 1377–1427.
- [14] S. Ferenczi, *Substitution dynamical systems on infinite alphabets*, Ann. Inst. Fourier (Grenoble) **56** (2006), no. 7, 2315–2343.
- [15] N. P. Frank, *Multidimensional constant-length substitution sequences*, Topology Appl. **152** (2005), no. 1-2, 44–69.
- [16] N. P. Frank and N. Mañibo, *Spectral theory of spin substitutions*, in preparation.
- [17] N. P. Frank and L. Sadun, *Fusion tilings with infinite local complexity*, Topology Proc. **43** (2014), 235–276.
- [18] F. Gähler and R. Klitzing, *The diffraction pattern of self-similar tilings*, The mathematics of long-range aperiodic order (R.V. Moody, ed.), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 489, Kluwer Acad. Publ., Dordrecht, 1997, pp. 141–174.

- [19] H. Helson, *Cocycles on the circle*, J. Operator Theory **16** (1986), no. 1, 189–199.
- [20] J. Kellendonk and L. Sadun, *Meyer sets, topological eigenvalues, and Cantor fiber bundles*, J. Lond. Math. Soc. **89** (2014), no. 1, 114–130.
- [21] J. Kellendonk and R. Yassawi, *The Ellis semigroup of bijective substitutions*, Groups Geom. Dyn., in press, <https://arxiv.org/pdf/1908.0569> (2020).
- [22] J. C. Lagarias, *Geometric models for quasicrystals I. Delone sets of finite type*, Discrete Comput. Geom. **21** (1999), no. 2, 161–191.
- [23] J.-Y. Lee, R. V. Moody, and B. Solomyak, *Pure point dynamical and diffraction spectra*, Ann. Henri Poincaré **3** (2002), no. 5, 1003–1018.
- [24] J.-Y. Lee and B. Solomyak, *Pure point diffractive substitution Delone sets have the Meyer property*, Discrete Comput. Geom. **39** (2008), no. 1-3, 319–338.
- [25] D. Lenz, *Spectral theory of dynamical systems as diffraction theory of sampling functions*, Monatsh. Math. **192** (2020), no. 3, 625–649.
- [26] D. Lenz, T. Spindeler, and N. Strungaru, *Pure point diffraction and mean, Besicovitch and Weyl almost periodicity*, preprint, <https://arxiv.org/abs/2006.10821> (2020).
- [27] N. Mañibo, D. Rust, and J. J. Walton, *Substitutions on compact alphabets*, in preparation.
- [28] M. Queffélec, *Substitution Dynamical Systems—Spectral Analysis*, second ed., LNM, vol. 1294, Springer, Berlin, 2010.
- [29] E. Rowland and R. Yassawi, *Profinite automata*, Adv. in Appl. Math. **85** (2017), 60–83.
- [30] N. Strungaru, *On the Fourier analysis of measures with Meyer set support*, J. Funct. Anal. **278** (2020), 108404, (30 pp.).

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD,
 POSTFACH 100131, 33501 BIELEFELD, GERMANY
Email address: cmanibo@math.uni-bielefeld.de

SCHOOL OF MATHEMATICS AND STATISTICS, THE OPEN UNIVERSITY,
 WALTON HALL, MILTON KEYNES, MK7 6AA, UK
Email address: dan.rust@open.ac.uk

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF NOTTINGHAM,
 KING'S MEADOW CAMPUS, LENTON LANE, NOTTINGHAM, NG7 2NR, UK
Email address: James.Walton@nottingham.ac.uk