

SPECIAL CASES AND EQUIVALENT FORMS OF KATZNELSON'S PROBLEM ON RECURRENCE

JOHN T. GRIESMER

ABSTRACT. We make the following three observations regarding a question popularized by Katznelson: is every subset of \mathbb{Z} which is a set of Bohr recurrence is also set of topological recurrence?

- (i) If G is a countable abelian group and $E \subset G$ is an I_0 set, then every subset of $E - E$ which is a set of Bohr recurrence is also a set of topological recurrence. In particular every subset of $\{2^n - 2^m : n, m \in \mathbb{N}\}$ which is a set of Bohr recurrence is a set of topological recurrence.
- (ii) Let \mathbb{Z}^ω be the direct sum of countably many copies of \mathbb{Z} with standard basis E . If every subset of $(E - E) - (E - E)$ which is a set of Bohr recurrence is also a set of topological recurrence, then every subset of every countable abelian group which is a set of Bohr recurrence is also a set of topological recurrence.
- (iii) Fix a prime p and let \mathbb{F}_p^ω be the direct sum of countably many copies of $\mathbb{Z}/p\mathbb{Z}$ with basis $(\mathbf{e}_i)_{i \in \mathbb{N}}$. If for every p -uniform hypergraph with vertex set \mathbb{N} and edge set \mathcal{F} having infinite chromatic number, the Cayley graph on \mathbb{F}_p^ω determined by $\{\sum_{i \in F} \mathbf{e}_i : F \in \mathcal{F}\}$ has infinite chromatic number, then every subset of \mathbb{F}_p^ω which is a set of Bohr recurrence is a set of topological recurrence.

1. RECURRENCE PROPERTIES

We consider a problem raised in Section 4 of [21], stated as Conjecture 1.2 below; see [13] for history and overview of this problem. Our main results are Proposition 2.2, a very special case of Conjecture 1.2, and Proposition 3.3, which reduces Conjecture 1.2 to a special case.

1.1. Definitions and main problem. As usual \mathbb{R} denotes the group of real numbers, \mathbb{Z} denotes the group of integers, and \mathbb{T} denotes \mathbb{R}/\mathbb{Z} with the quotient topology. The usual translation invariant metric on \mathbb{T}^d is defined as follows: for an element $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{T}^d$, we write $\|\mathbf{x}\|$ for $\max_{j \leq d} \min_{n \in \mathbb{Z}} |x_j - n|$ (adopting the usual abuses of notation). This makes $(\mathbf{x}, \mathbf{y}) \mapsto \|\mathbf{x} - \mathbf{y}\|$ a translation invariant metric on \mathbb{T}^d .

We write \mathcal{S}^1 for the circle group $\{z \in \mathbb{C} : |z| = 1\}$ with the operation of multiplication, which is isomorphic as a topological group to \mathbb{T} , under the identification

$t \in \mathbb{T} \leftrightarrow \exp(2\pi it) \in \mathcal{S}^1$. A *character* of G is a homomorphism $\chi : G \rightarrow \mathcal{S}^1$, and a *trigonometric polynomial* $p : G \rightarrow \mathbb{C}$ is a linear combination of characters.

If $A, B \subset G$, we write $A+B$ for $\{a+b : a \in A, b \in B\}$ and $A-A$ for $\{a-a' : a, a' \in A\}$. The *upper Banach density* of $A \subset G$ is

$$d^*(A) := \sup\{\lambda(1_A) : \lambda \text{ is an invariant mean on } l^\infty(G)\}.$$

See [3] or [1] for general exposition. We do not prove results concerning upper Banach density in this article, but we will refer to such results from the literature.

Definition 1.1. Let G be a countably infinite abelian group. If $S \subset G$, we say that S is a *set of*

- *density recurrence* if for every $A \subset G$ with $d^*(A) > 0$, we have $(A - A) \cap S \neq \emptyset$.
- *chromatic recurrence* if for every $r \in \mathbb{N}$ and every cover of G by r sets A_1, \dots, A_r , there is a $j \leq r$ such that $(A_j - A_j) \cap S \neq \emptyset$.

Equivalently, S is a set of chromatic recurrence if for every $r \in \mathbb{N}$ and every $f : G \rightarrow \{1, \dots, r\}$, there exists $a, b \in G$ such that $f(a) = f(b)$ and $b - a \in S$.

- *Bohr recurrence* if for every $d \in \mathbb{N}$, every homomorphism $\rho : G \rightarrow \mathbb{T}^d$, and every $\varepsilon > 0$, there exists $s \in S$ such that $\|\rho(s)\| < \varepsilon$.

Equivalently, S is a set of Bohr recurrence if for all $d \in \mathbb{N}$, every set χ_1, \dots, χ_d of d characters of G , and all $\varepsilon > 0$, there exists $s \in S$ such that $|\chi_j(s) - 1| < \varepsilon$ for each j .

The equivalence asserted in the third definition is due to the isomorphism of \mathbb{T} with \mathcal{S}^1 and the fact that a collection of d characters corresponds to a homomorphism $(\chi_1, \dots, \chi_d) : G \rightarrow (\mathcal{S}^1)^d$.

It is easy to verify that every set of density recurrence is a set of chromatic recurrence, and that every set of chromatic recurrence is a set of Bohr recurrence. Kriz [23] constructed a subset of \mathbb{Z} which is a set of chromatic recurrence and not a set of density recurrence. The question of whether every set of Bohr recurrence in \mathbb{Z} is a set of chromatic recurrence was suggested by Veech [27] (and implicitly by Følner [7]), and popularized by Katznelson in [21]; he did not formulate a conjecture as to which answer is correct. While [21] considers only the group \mathbb{Z} , the analogous question for arbitrary abelian groups is implicit in the surrounding literature. There is no countably infinite abelian group G where the answer is known. We state the question as a conjecture, to make the ensuing discussion more natural.

Conjecture 1.2. *Let G be a countably infinite abelian group. If $S \subset G$ is a set of Bohr recurrence, then S is a set of chromatic recurrence.*

1.2. Measurable recurrence and topological recurrence. In abelian groups, the sets of chromatic recurrence are precisely the sets of topological recurrence. Likewise, sets of density recurrence are sets of measurable recurrence and vice versa. Since this article never addresses dynamical systems per se, we use the terminology which is more closely aligned with our definitions. See [3] for a general discussion of the equivalences between various recurrence properties.

1.3. Outline. Our first result is Proposition 2.2, which says that if E belongs to a certain class of subsets of G , then every subset $S \subset E - E$ which is a set of Bohr recurrence is also a set of chromatic recurrence. A special case of this proposition says that if $S \subset \{2^n - 2^m : m, n \in \mathbb{N}\}$ is a set of Bohr recurrence, then S is a set of chromatic recurrence. This is in contrast to Kriz's construction [23], which proves that if $E \subset \mathbb{Z}$ is infinite, then there is a set $S \subset E - E$ which is a set of chromatic recurrence and not a set of density recurrence. Kriz [23] only stated that there is a subset of \mathbb{Z} which is a set of chromatic recurrence and not of density recurrence, but minor modifications of the proof yield the more general result. The article [18] explicitly proves the generalization.

In §3 we state Conjecture 3.2, the special case of Conjecture 1.2 where G is the direct sum of countably many copies of \mathbb{Z} and S is contained in $(\mathcal{E}_1 - \mathcal{E}_1) - (\mathcal{E}_1 - \mathcal{E}_1)$, where \mathcal{E}_1 is the standard basis of G . Proposition 3.3 shows that this special case implies the full conjecture. Conjecture 3.2 is superficially similar to Proposition 2.2, but we show in §5 that it is not susceptible to the same proof.

In §4 we consider the special case of Conjecture 1.2 where the ambient group is a vector space over a finite field of odd characteristic. Proposition 4.4 and Lemma 4.9 show that in this setting, Conjecture 1.2 can be reduced to the study of an easy-to-describe class of Bohr recurrent sets, first considered in [10] and [11] as a means to distinguish Bohr topologies on various groups.

Two open problems suggested by Propositions 2.2 and 3.3 are stated in §6

1.4. Previous work. Conjecture 1.2 has been studied in [12], [6], [19], [13], and [20]. The exposition in [13] thoroughly summarizes the history of the problem.

1.5. Acknowledgement. We thank Anh Le for suggesting the proof in Remark 2.6.

2. SUBSETS OF DIFFERENCE SETS

If G is an abelian group, we say $\psi : G \rightarrow \mathbb{C}$ is *uniformly almost periodic* if ψ is a uniform limit of trigonometric polynomials.

Definition 2.1. If G is an abelian group and $E \subset G$, we say that E is an I_0 set if for every bounded $f : E \rightarrow \mathbb{C}$, there is a uniformly almost periodic $\psi : G \rightarrow \mathbb{C}$ such that $\psi|_E = f$.

See [14], [22], or [24] for an overview of I_0 sets. An early result [26] on such sets says that if $A = \{a_1 < a_2 < \dots\} \subset \mathbb{Z}$ is lacunary (meaning $\liminf_{n \rightarrow \infty} a_{n+1}/a_n > 1$), then A is an I_0 set. Every infinite abelian group contains an I_0 set of the same cardinality as the group (see [22]).

Proposition 2.2. *Let G be a countable abelian group and let $E \subset G$ be an I_0 set. If $S \subset E - E$ and S is a set of Bohr recurrence, then S is a set of chromatic recurrence.*

We prove Proposition 2.2 after the following lemma, which provides an equivalent condition for E to be a set of Bohr recurrence.

Lemma 2.3. *Let G be an abelian group. The following are equivalent.*

- (1) S is a set of Bohr recurrence.
- (2) For every almost periodic $\psi : G \rightarrow \mathbb{C}$ and all $\varepsilon > 0$, there exists $s \in S$ such that $|\psi(g + s) - \psi(g)| < \varepsilon$ for all $g \in G$.

Proof. It suffices to prove that (1) implies the special case of (2) where ψ is a trigonometric polynomial p , since every almost periodic function may be uniformly approximated by these. To prove this, assume S is a set of Bohr recurrence, that $p = \sum_{j=1}^d c_j \chi_j$ is a trigonometric polynomial on G , and $\varepsilon > 0$. Let $M = \max_{j \leq d} |c_j|$ and let $\varepsilon' = \varepsilon/(Md)$. Since S is a set of Bohr recurrence, we may choose $s \in S$ such that $|\chi_j(s) - 1| < \varepsilon'$ for all $j \leq d$. Simplifying $|p(g + s) - p(g)|$ we get

$$\begin{aligned} \left| \sum_{j=1}^d c_j \chi_j(g + s) - c_j \chi_j(g) \right| &\leq \sum_{j=1}^d |c_j| |\chi_j(g + s) - \chi_j(g)| \\ &= \sum_{j=1}^d |c_j| |\chi_j(s) - 1| \\ &< dM\varepsilon' \\ &= \varepsilon. \end{aligned}$$

We will not need the fact that (2) implies (1), so we leave its proof as an exercise. \square

Proof of Proposition 2.2. Fixing G , E , and S in the hypothesis, it suffices to prove that for all $r \in \mathbb{N}$ and every function $f : G \rightarrow \{1, \dots, r\}$, there exists $a, b \in G$ such that $f(b) = f(a)$ and $b - a \in S$. So fix $r \in \mathbb{N}$ and $f : G \rightarrow \{1, \dots, r\}$. Since E is an I_0 set, we may choose a uniformly almost periodic $\psi : G \rightarrow \mathbb{C}$ such that $f(x) = \psi(x)$ for all $x \in E$. By Lemma 2.3, there is a $y \in S$ such that

$$(2.1) \quad |\psi(x + y) - \psi(x)| < 1 \quad \text{for all } x \in G.$$

Since $S \subset E - E$, we can write y as $b - a$, where $a, b \in E$. Setting $x = a$ in (2.1), we get $|\psi(b) - \psi(a)| < 1$. Then

$$|f(b) - f(a)| = |\psi(b) - \psi(a)| < 1.$$

Since f is integer valued, this implies $f(b) = f(a)$. As mentioned above, we have $b - a \in E$, so this proves that E is a set of chromatic recurrence. \square

To see that Proposition 2.2 is not vacuous, we mention the following classical result, which is essentially the Poincaré recurrence theorem; see [9], [2], or [8] for exposition.

Theorem 2.4. *Let G be a countable abelian group. If E is an infinite subset of G , then $\{a - b : a \neq b \in E\}$ is a set of density recurrence (and therefore a set of chromatic recurrence and a set of Bohr recurrence).*

Remark 2.5. If one could find an I_0 set E such that $G = E - E$, then Conjecture 1.2 would be a corollary of Proposition 2.2. But when G has infinite cardinality, the difference set of an I_0 set is never all of G ; see Lemma 2.11.

Remark 2.6. The proof of Proposition 2.2 is easily modified to prove the analogous statement with the following weaker hypothesis on E :

(*) Every bounded $f : E \rightarrow \mathbb{C}$ can be uniformly approximated by a trigonometric polynomial restricted to E .

The hypothesis (*) implies that E is an I_0 set, so we would obtain no greater generality by using it in Proposition 2.2. To see this implication, note that (*) implies that for any two disjoint subsets $E', E'' \subset E$ and all $\varepsilon > 0$, there is a trigonometric polynomial p with $|p(x) - 1| < \varepsilon$ for $x \in E'$ and $|p(x) - 0| < \varepsilon$ for $x \in E''$. This means that E' and E'' have disjoint closures in the Bohr compactification of G , and Proposition 3.4.1 of [14] then implies E is an I_0 set.

We thank Ahn Le for pointing out the argument above.

Remark 2.7. It is tempting to interpret Proposition 2.2 as saying “When E is an I_0 set, it is easy to prove that a subset of $E - E$ is a set of chromatic recurrence: just prove that it is Bohr recurrent.” Upon reflection, it seems more correct to say that “it is difficult to prove that a subset of $E - E$ is a set of Bohr recurrence: to do so one must prove that it is a set of chromatic recurrence.”

One may wonder if the conclusion of Proposition 2.2 can be improved to “... then S is a set of density recurrence.” But it cannot, as the following proposition shows.

Proposition 2.8 ([18], Theorem 1.2). *Let $E \subset \mathbb{Z}$ be infinite. There is a set $S \subset E - E$ such that S is a set of chromatic recurrence and not a set of density recurrence.*

In the hypothetical scenario where Conjecture 1.2 is false, Propositions 2.2 and 2.8 demonstrate that the relationship between Bohr recurrence and chromatic recurrence differs from the relationship between chromatic recurrence and measurable recurrence. Regardless of the status of Conjecture 1.2, the following problem seems interesting.

Problem 2.9. Extend Proposition 2.2 to a broader class of sets (broader than difference sets of finite unions of I_0 sets).

The parenthetical remark is included because we believe that extending the result to finite unions of I_0 sets will be relatively straightforward, using facts from [14], [22], or [24].

Remark 2.10. Proposition 2.2 is similar to Theorem 7.15 of [22], which solves a special case of an open problem about Sidon sets in \mathbb{Z} , under the additional assumption that the set under consideration is contained in the difference set $E - E$ of a lacunary set E .

The following lemma shows that difference sets of I_0 sets are small, and in particular never equal to the entire ambient group. If $S \subset G$ and $g \in G$, we say that g is a *Bohr limit point of S* if $(S \cap U) \setminus \{g\} \neq \emptyset$ for every Bohr neighborhood U of g . In other words, the closure $\overline{S \setminus \{g\}}$ in the Bohr topology contains g .

Lemma 2.11. *If G is an abelian group and $E \subset G$ is an I_0 set, then there is a Bohr neighborhood U of 0 in G such that 0 is the only Bohr limit point of $(E - E) \cap U$.*

Proof. Let $E \subset G$ be an I_0 set. By Theorems 5.3.1 and 5.3.9 of [14], E is a union of finitely many sets E_1, \dots, E_r such that for each j , the only Bohr limit point of $E_j - E_j$ is 0. This property is inherited by subsets, so we may assume that the E_j are mutually disjoint. Let

\tilde{E}_j be the closure of E_j in the Bohr compactification of G . By Proposition 3.4.1 of [14], the \tilde{E}_j are mutually disjoint. So for each $i, j, i \neq j$, there is a Bohr neighborhood $U_{i,j}$ of 0 such that $(E_i - E_j) \cap U_{i,j} = \emptyset$. Letting U be the intersection of these $U_{i,j}$, we get that $U \cap \bigcup_{i \neq j \leq r} E_i - E_j = \emptyset$. Now $U \cap (E - E) \subset \bigcup_{j=1}^r E_j - E_j$, and the only Bohr limit point of each $E_j - E_j$ is 0. So the only Bohr limit point of the union is 0, as well. Thus the only Bohr limit point of $U \cap (E - E)$ is 0. \square

3. ITERATED DIFFERENCES

In this section we state Conjecture 3.2 and prove that it implies Conjecture 1.2. This proof involves a classical result about Bohr neighborhoods, so we define those here.

We say that $B \subset G$ is a *Bohr neighborhood of 0* if there exists $d \in \mathbb{N}$, $\varepsilon > 0$, a homomorphism $\rho : G \rightarrow \mathbb{T}^d$, and a neighborhood U of 0 in \mathbb{T}^d such that B contains $\rho^{-1}(U)$. Note that $S \subset G$ is a set of Bohr recurrence if and only if $S \cap B \neq \emptyset$ for every Bohr neighborhood B of 0 in G .

The following theorem is a primary impetus for investigations of difference sets, and a key ingredient in the proof of Proposition 3.3 below. Given a subset A of an abelian group G , we write $\Delta_2(A)$ for the set $\{(a - b) - (c - d) : a, b, c, d \in A \text{ are mutually distinct}\}$.

Theorem 3.1 (Bogoliouboff [5]). *Let G be a countable abelian group. If G is written as a finite union of sets A_1, \dots, A_r , then for some $j \leq r$ the set $\Delta_2(A_j) \cup \{0\}$ contains a Bohr neighborhood of 0.*

Readers familiar with the literature will note that we have omitted bounds on the rank and radius of the Bohr neighborhood in Theorem 3.1, which may be of interest in quantitative approaches to Conjecture 1.2.

Often Theorem 3.1 is stated with $(A_j - A_j) - (A_j - A_j)$ in place of $\Delta_2(A_j)$, but imposing the additional restriction that the terms in the difference be distinct does not significantly alter the structure. Theorem 3.1, as stated here, is a special case of the main result of [15], which generalizes Theorem 6.1 of [4].

We now consider the group \mathbb{Z}^ω , the direct sum of countably many copies of \mathbb{Z} . We adopt the usual presentation: elements of \mathbb{Z}^ω are sequences (n_1, n_2, n_3, \dots) of elements of \mathbb{Z} where $n_j = 0$ for all but finitely many j . For each $j \in \mathbb{N}$, let \mathbf{e}_j be the element of \mathbb{Z}^ω where $(\mathbf{e}_j)_j = 1$ and $(\mathbf{e}_j)_k = 0$ if $j \neq k$. So $\mathbf{e}_1 = (1, 0, 0, \dots)$, $\mathbf{e}_2 = (0, 1, 0, 0, \dots)$, etc. We write \mathcal{E}_1 for $\{\mathbf{e}_j : j \in \mathbb{N}\}$. The set $\Delta_2(\mathcal{E}_1)$, contains elements such as $(0, 1, 0, -1, 1, -1, 0, \dots)$. The following conjecture, which we are unable to prove, is analogous to Proposition 2.2.

Conjecture 3.2. *If $S \subset \Delta_2(\mathcal{E}_1)$ and S is a set of Bohr recurrence, then S is a set of chromatic recurrence.*

Proposition 3.3. *Conjecture 3.2 implies Conjecture 1.2.*

Remark 3.4. One may hope for an easy proof of Conjecture 3.2 along the lines of our proof of Proposition 2.2. We point out an obstruction to this hope in §5.

The proof of Proposition 3.3 consists of the following two lemmas.

Lemma 3.5. *Let H and G be countable abelian groups, let $Q \subset H$, and let $\rho : H \rightarrow G$ be a homomorphism such that $\rho(Q) = G$. If $S \subset G$ is a set of Bohr recurrence, then $\rho^{-1}(S) \cap \Delta_2(Q)$ is a set of Bohr recurrence in H .*

Proof. Assuming H, G, Q , and ρ are as in the hypothesis of the lemma, we must prove that for all $d \in \mathbb{N}$, all homomorphisms $\psi : H \rightarrow \mathbb{T}^d$, and all $\varepsilon > 0$, there is an $h \in \rho^{-1}(S) \cap \Delta_2(Q)$ such that $\|\psi(h)\| < \varepsilon$. In other words, we must find distinct elements $q_1, q_2, q_3, q_4 \in Q$ such that

$$(3.1) \quad \rho(q_1 - q_2 - q_3 + q_4) \in S \quad \text{and} \quad \|\psi(q_1 - q_2 - q_3 + q_4)\| < \varepsilon.$$

So fix such d, ψ , and ε . To find q_i satisfying (3.1), write \mathbb{T}^d as a finite union of sets B_1, \dots, B_r having diameter at most $\varepsilon/4$. Then the preimages $\psi^{-1}(B_1), \dots, \psi^{-1}(B_r)$ cover H . In particular, the sets $C_j := \psi^{-1}(B_j) \cap Q$, $j \leq r$, cover Q . The hypothesis that $\rho(Q) = G$ now implies that the images $A_1 := \rho(C_1), \dots, A_r := \rho(C_r)$ cover G . Theorem 3.1 then implies that for some j , the set $\Delta_2(A_j)$ contains a Bohr neighborhood of 0 in G . Since we assumed S is a set of Bohr recurrence, we then have $S \cap \Delta_2(A_j) \neq \emptyset$. In particular, there exist mutually distinct elements $a_1, a_2, a_3, a_4 \in A_j$ and $s \in S$ such that $s = a_1 - a_2 - a_3 + a_4$. The definition of A_j allows us to write the latter equation as

$$(3.2) \quad s = \rho(q_1 - q_2 - q_3 + q_4), \quad \text{where each } q_i \in C_j.$$

Now the definition of C_j means that each q_i lies in Q and each $\psi(q_i)$ lies in B_j , which has diameter less than $\varepsilon/4$. This implies

$$(3.3) \quad \|\psi((q_1 - q_2) - (q_3 - q_4))\| \leq \|\psi(q_1) - \psi(q_2)\| + \|\psi(q_3) - \psi(q_4)\| < \varepsilon.$$

Now (3.2) and (3.3) together show that the q_i satisfy (3.1). \square

Lemma 3.6. *Let H and G be abelian groups and let $R \subset H$ be a set of chromatic recurrence. If $\rho : H \rightarrow G$ is a homomorphism, then $\rho(R)$ is a set of chromatic recurrence.*

Proof. Let $k \in \mathbb{N}$ and let $f : G \rightarrow \{1, \dots, k\}$ be a function. Then $\tilde{f} := f \circ \rho$ is a function from H to $\{1, \dots, k\}$. Since R is a set of chromatic recurrence, there exists $h_1, h_2 \in H$ such that $\tilde{f}(h_1) = \tilde{f}(h_2)$ and $h_2 - h_1 \in R$. By the definition of ρ , we have $f(\rho(h_1)) = f(\rho(h_2))$ and $\rho(h_2 - h_1) \in \rho(R)$, meaning $\rho(h_2) - \rho(h_1) \in \rho(R)$. Setting $a = \rho(h_1)$ and $b = \rho(h_2)$, we have found $a, b \in G$ such that $f(a) = f(b)$ and $b - a \in \rho(R)$. \square

Observation 3.7. If G is an abelian group and $f : \mathcal{E}_1 \rightarrow G$ is a function, then there is a homomorphism $\rho : \mathbb{Z}^\omega \rightarrow G$ such that $\rho|_{\mathcal{E}_1} = f$. Such a ρ can be defined uniquely by $\rho(\sum c_j \mathbf{e}_j) := \sum c_j f(\mathbf{e}_j)$.

Proof of Proposition 3.3. Let G be a countable abelian group, and let $S \subset G$ be a set of Bohr recurrence. By Observation 3.7 choose a homomorphism $\rho : \mathbb{Z}^\omega \rightarrow G$ such that $\rho(\mathcal{E}_1) = G$. Assuming Conjecture 3.2 holds, we will prove that $\rho^{-1}(S)$ is a set of chromatic recurrence. To see this, first apply Lemma 3.5 with $Q = \mathcal{E}_1$ to get that $\rho^{-1}(S) \cap \Delta_2(\mathcal{E}_1)$ is a set of Bohr recurrence. Now Conjecture 3.2 implies $\rho^{-1}(S) \cap \Delta_2(\mathcal{E}_1)$ is a set of chromatic recurrence, so $\rho^{-1}(S)$ is a set of chromatic recurrence, as well. By Lemma 3.6 we get that S is a set of chromatic recurrence. \square

Remark 3.8. Some special cases of Conjecture 3.2 are easy to prove. For example, $\Delta_2(\mathcal{E}_1)$ contains an infinite difference set $E - E$, where $E := \{\mathbf{e}_{2n} - \mathbf{e}_{2n+1} : n \in \mathbb{N}\}$. This E is an independent set¹, and therefore an I_0 set, in \mathbb{Z}^ω . Proposition 2.2 then proves that every subset of $E - E$ which is a set of Bohr recurrence is also a set of chromatic recurrence. We have not proved any case of Conjecture 3.2 beyond such trivial instances.

4. SPECIAL CASES IN \mathbb{F}_p^ω

Variations on Proposition 3.3 can be obtained from variations on Theorem 3.1; the resulting variations are especially appealing in vector spaces over finite fields.

Fix an odd prime p , let \mathbb{F}_p be the additive group $\mathbb{Z}/p\mathbb{Z}$, and let \mathbb{F}_p^ω denote the direct sum of countably many copies of \mathbb{F}_p . We adopt the usual presentation of \mathbb{F}_p^ω : elements are written as (x_1, x_2, x_3, \dots) , where each $x_i \in \mathbb{F}_p$, and $\mathbf{e}_1 = (1, 0, 0, \dots)$, $\mathbf{e}_2 = (0, 1, 0, \dots)$, \dots forms the standard basis.

Observation 4.1. Since every homomorphism $\rho : \mathbb{F}_p^\omega \rightarrow \mathbb{T}^d$ takes values in the finite subgroup of \mathbb{T}^d consisting of elements of order p , the Bohr neighborhoods of 0 in \mathbb{F}_p^ω are exactly the sets containing a finite index subgroup of \mathbb{F}_p^ω .

¹Meaning the only integer solutions n_i to $n_1 \mathbf{e}_1 + n_2 \mathbf{e}_2 + \dots + n_r \mathbf{e}_r = 0$ with mutually distinct $e_i \in E$ satisfy $n_1 \mathbf{e}_1 = n_2 \mathbf{e}_2 = \dots = n_r \mathbf{e}_r = 0$.

We write $\widehat{\mathbb{F}}_p^\omega$ for the set of characters of \mathbb{F}_p^ω , so that $\widehat{\mathbb{F}}_p^\omega$ is a group under pointwise multiplication. It is easy to verify that $\widehat{\mathbb{F}}_p^\omega$ is isomorphic to $\widehat{\mathbb{F}}_p^\mathbb{N}$, the product of countably many copies of \mathbb{F}_p . To see this, note that every element $\xi \in \widehat{\mathbb{F}}_p^\mathbb{N}$ induces a character $\chi(x) = e(\sum_{n \in \mathbb{N}} x_n \xi_n)$, where multiplication $x_n \xi_n$ is done in \mathbb{F}_p , and $e : \mathbb{F}_p \rightarrow \mathcal{S}^1$ is the homomorphism with $e(1) = e^{2\pi i/p}$. Conversely, every character of \mathbb{F}_p^ω has this form: given χ , define $\xi \in \widehat{\mathbb{F}}_p^\mathbb{N}$ to satisfy $e(\xi_n) = \chi(\mathbf{e}_n)$. One can verify that $\chi(x) = e(\sum_{n \in \mathbb{N}} x_n \xi_n)$ for every $x \in \mathbb{F}_p^\omega$.

For a finite subset $F \subset \mathbb{N}$, let $\mathbf{e}_F = \sum_{n \in F} \mathbf{e}_n$. Let $\mathcal{E}_d^{(p)} := \{\mathbf{e}_F : F \subset \mathbb{N}, |F| = d\}$. The following conjecture is a variant of Conjecture 3.2.

Conjecture 4.2. *Let p be prime and let $d \in \mathbb{N}$ be divisible by p . Every subset of $\mathcal{E}_d^{(p)}$ which is a set of Bohr recurrence is also set of chromatic recurrence.*

Here is the special case of Conjecture 1.2 for \mathbb{F}_p^ω .

Conjecture 4.3. *Every subset of \mathbb{F}_p^ω which is a set of Bohr recurrence is a set of chromatic recurrence.*

Clearly Conjecture 4.3 implies Conjecture 4.2. The purpose of this section is to prove the converse (assuming $d > 2$), and to prove Lemma 4.9, an appealing characterization of the Bohr recurrent subsets of $\mathcal{E}_p^{(p)}$.

Proposition 4.4. *Let p be prime and let $d > 2$ be divisible by p . The special case of Conjecture 4.2 for this d and p implies the special case of Conjecture 4.3 for the same p .*

Remark 4.5. Conjecture 4.2 is not vacuous: the set $\mathcal{E}_p^{(p)}$ itself is a set of a set of Bohr recurrence, which can be verified directly, or viewed as a consequence of Lemma 4.8, or of Lemma 4.9.

Naturally one wonders whether $\mathcal{E}_p^{(p)}$ itself is a set of chromatic recurrence. A straightforward extension of the Poincaré recurrence theorem shows that $\mathcal{E}_p^{(p)}$ is a set of density recurrence, and therefore also a set of chromatic recurrence; see [16] for discussion.

Remark 4.6. We are currently unable to prove that the special case of Conjecture 4.2 with $d = p = 2$ implies Conjecture 4.3. This special case of Conjecture 4.2 is true, as it is also a special case of Proposition 2.2.

We will use the following special case of the main result of [15].

Lemma 4.7. *Let p be prime and let $d > 2$ be divisible by p . If $A \subset \mathbb{F}_p^\omega$ has $d^*(A) > 0$, then the d -fold sumset*

$$\sum_{i=1}^d A := \{a_1 + a_2 + \cdots + a_d : a_i \in A \text{ are mutually distinct}\} \cup \{0\}$$

contains a Bohr neighborhood of 0. Consequently, if \mathbb{F}_p^ω is covered by finitely many sets A_1, \dots, A_r , then for some j the set $\sum_{i=1}^d A_j$ contains a Bohr neighborhood of 0.

The proof of Proposition 4.4 is similar to the proof of Proposition 3.3. It rests on the following variant of Lemma 3.5, whose proof uses Lemma 4.7 in place of Theorem 3.1.

Lemma 4.8. *Let p be prime, let $d > 2$ be divisible by p , and let $\rho : \mathbb{F}_p^\omega \rightarrow \mathbb{F}_p^\omega$ be a homomorphism such that $\rho(\mathcal{E}_1^{(p)}) = \mathbb{F}_p^\omega$. If $S \subset \mathbb{F}_p^\omega$ is a set of Bohr recurrence, then $\rho^{-1}(S) \cap \mathcal{E}_d^{(p)}$ is a set of Bohr recurrence, as well.*

Proof. Fix p, d, ρ , and S as in the hypothesis. We must prove that $S' := \rho^{-1}(S) \cap \mathcal{E}_d^{(p)}$ is a set of Bohr recurrence, which in this case simply means that S' has nonempty intersection with every finite index subgroup of \mathbb{F}_p^ω . Fix such a subgroup H , and enumerate the cosets of H as H_1, \dots, H_r . Let $C_j := H_j \cap \mathcal{E}_1^{(p)}$, so that $\mathcal{E}_1 = C_1 \cup \cdots \cup C_r$. Let $A_j := \rho(C_j)$, $j = 1, \dots, r$, so the hypothesis that $\rho(\mathcal{E}_1) = \mathbb{F}_p^\omega$ implies that the A_j cover \mathbb{F}_p^ω . Lemma 4.7 now implies that for some $j \leq r$, $\sum_{i=1}^d A_j$ contains a Bohr neighborhood of 0. Since S is a set of Bohr recurrence, we have that $S \cap \sum_{i=1}^d A_j$ is nonempty. In particular, we can find an $s \in S$ and mutually distinct $a_1, \dots, a_d \in A_j$ such that

$$s = a_1 + \cdots + a_d.$$

By the definition of A_j , this means there are mutually distinct $c_1, \dots, c_d \in C_j$ such that

$$(4.1) \quad s = \rho(c_1) + \cdots + \rho(c_d)$$

By the definition of C_j , these c_i all lie in the same coset of H , meaning $c_i + H = c_1 + H$ for each i . Since p divides d , we get that $c_1 + \cdots + c_d$ lies in H : to see this write $c_1 + \cdots + c_d + H = dc_1 + H = H$.

Now each $c_i \in \mathcal{E}_1$, so $c := c_1 + \cdots + c_d \in \mathcal{E}_d^{(p)}$. Equation (4.1) shows that $c \in \rho^{-1}(S)$, so we have proved that $\rho^{-1}(S) \cap \mathcal{E}_d^{(p)} \cap H \neq \emptyset$, as desired. \square

Proof of Proposition 4.4. Fix a prime p and a $d > 2$ divisible by p . Assuming Conjecture 4.2 is true for this p and d , we will prove that Conjecture 4.3 is true for this p , as well. Fix a set of Bohr recurrence $S \subset \mathbb{F}_p^\omega$. Let $\rho : \mathbb{F}_p^\omega \rightarrow \mathbb{F}_p^\omega$ be a homomorphism such that

$\rho(\mathcal{E}_1) = \mathbb{F}_p^\omega$ By Lemma 4.8, we have that $S' := \rho^{-1}(S) \cap \mathcal{E}_d^{(p)}$ is a set of Bohr recurrence. Our assumption that Conjecture 4.2 is true then implies that S' is a set of chromatic recurrence. Then Lemma 3.6 implies $\rho(S')$ also a set of chromatic recurrence, and the containment $\rho(S') \subset S$ implies S is, as well. \square

Now we characterize the Bohr recurrent subsets of $\mathcal{E}_p^{(p)}$. We call a collection \mathcal{F} of subsets of \mathbb{N} *partition regular*² if for every partition of \mathbb{N} into finitely many cells, at least one of the cells contains an element of \mathcal{F} . In other words, the hypergraph with vertex set \mathbb{N} and edge set \mathcal{F} has infinite chromatic number.

Lemma 4.9. *Let p be prime (not necessarily odd), let \mathcal{F} be a collection of subsets of \mathbb{N} each having cardinality p , and let $\mathcal{E}(\mathcal{F}) := \{\mathbf{e}_F : F \in \mathcal{F}\} \subset \mathbb{F}_p^\omega$. Then $\mathcal{E}(\mathcal{F})$ is a set of Bohr recurrence if and only if \mathcal{F} is partition regular.*

The implication “ $\mathcal{E}(\mathcal{F})$ is a set of Bohr recurrence if \mathcal{F} is partition regular” is a special case of Lemma 3.3 of [11] (originally proved in [10]). The converse is a special case of Lemma 3.5 of [11].

Proof. First suppose that \mathcal{F} is a partition regular collection of subsets of \mathbb{N} each having cardinality p . We will prove that $\mathcal{E}(\mathcal{F})$ is a set of Bohr recurrence by showing that for every finite set of characters $\chi_1, \dots, \chi_r \in \widehat{\mathbb{F}_p^\omega}$, there is an $\mathbf{e}_F \in \mathcal{E}(\mathcal{F})$ such that $\chi_j(\mathbf{e}_F) = 1$ for each j . So fix such a collection of characters χ_1, \dots, χ_r .

For $j \leq r$ and $m \leq p$, let $A_{j,m} := \{n \in \mathbb{N} : \chi_j(\mathbf{e}_n) = \exp(2\pi im/p)\}$. Let $\mathcal{P} = \{P_1, \dots, P_k\}$ be the partition of \mathbb{N} induced by the $A_{j,m}$ (i.e. the minimal nonempty elements of the algebra of subsets of \mathbb{N} generated by the $A_{j,m}$). Then each χ_j is constant on the elements of \mathcal{P} . By the partition regularity of \mathcal{F} , we may choose $F \in \mathcal{F}$ so that $F \subset P$ for some element P of \mathcal{P} . To see that $\chi_j(\mathbf{e}_F) = 1$ for each j , note that $\chi_j(x)$ is constant on $\{\mathbf{e}_n : n \in F\}$; call this constant s . Then

$$\chi_j(\mathbf{e}_F) = \chi_j\left(\sum_{n \in F} \mathbf{e}_n\right) = \prod_{n \in F} \chi_j(\mathbf{e}_n) = \prod_{n \in F} s = s^p = 1,$$

since χ_j takes values in the p^{th} roots of unity.

Now suppose that \mathcal{F} is not partition regular, meaning there is a partition of \mathbb{N} into cells C_1, \dots, C_r such that for all $F \in \mathcal{F}$, F is not contained in any of the C_i . We will find a finite collection of characters χ_1, \dots, χ_r such that for each $F \in \mathcal{F}$, there is a j with

²This is an abuse of terminology: we should say that “containing an element of \mathcal{F} is a partition regular property of subsets of \mathbb{N} .”

$|\chi_j(\mathbf{e}_F) - 1| \geq \varepsilon := |\exp(2\pi i/p) - 1|$. This shows that $\mathcal{E}(\mathcal{F})$ has empty intersection with the Bohr neighborhood of 0 with radius $\varepsilon/2$ determined by the χ_j .

Consider the characters $\chi_j \in \widehat{\mathbb{F}_p^\omega}$ defined by $\chi_j(x) := \prod_{n \in C_j} \exp(2\pi i x_n/p)$. This is well defined, since $x_n = 0$ for all but finitely many n . Then for $\mathbf{e}_F \in \mathcal{E}(\mathcal{F})$, we have $\chi_j(\mathbf{e}_F) = \exp(2\pi i |F \cap C_j|/p)$. Since F is not contained in any of the C_j , there is at least one j such that $0 < |F \cap C_j| < p$, so $\chi_j(\mathbf{e}_F) \neq 1$. Since the only possible values of a character on \mathbb{F}_p^ω are the p^{th} roots of unity, we get that $|\chi_j(\mathbf{e}_F) - 1| \geq |\exp(2\pi i/p) - 1|$ for this j . \square

Remark 4.10. The special case of Lemma 4.9 where $p = 2$ is the special case of Proposition 2.2 where $G = \mathbb{F}_2^\omega$ and $S = \mathcal{E}_1$.

As with Conjecture 3.2, we are unable to prove any nontrivial instance of Conjecture 4.2. One special case of this conjecture in \mathbb{F}_3^ω is given by $S_{3AP} := \{\mathbf{e}_n + \mathbf{e}_{n+d} + \mathbf{e}_{n+2d} : n, d \in \mathbb{N}\}$. The collection $\{n, n+d, n+2d : n, d \in \mathbb{N}\}$ is partition regular, by van der Waerden's theorem on arithmetic progressions, so Lemma 4.9 says that S_{3AP} is Bohr recurrent. The problem of proving (or refuting) that S_{3AP} is a set of chromatic recurrence seems interesting, but it is unclear whether this special case will be any easier than the full generality of Conjecture 4.2.

4.1. Cayley graphs. If G is a group and S is a subset of G , the *Cayley graph* associated to S is the graph $\text{Cay}(S)$ whose vertex set is G , where two vertices g, g' are joined by an edge if $g - g' \in S$ or $g' - g \in S$. It is easy to see that S is a set of chromatic recurrence if and only if the chromatic number of $\text{Cay}(S)$ is infinite - see [21] for further exposition.

Combining Proposition 4.4 and Lemma 4.9, we see that the following conjecture, for a fixed p , is equivalent to Conjecture 4.3, for that p .

Conjecture 4.11. *Let p be an odd prime. For every p -uniform hypergraph \mathcal{F} on \mathbb{N} having infinite chromatic number, the Cayley graph on \mathbb{F}_p^ω associated to $\mathcal{E}(\mathcal{F})$ also has infinite chromatic number.*

5. WHY CONJECTURES 3.2 AND 4.2 ARE NOT AS EASY AS PROPOSITION 2.2

The proof of Proposition 2.2 showed that when E is an I_0 set and $S \subset E - E$ is a set of Bohr recurrence, the graph with vertex set E and edge set $\{\{a, b\} : a, b \in E, b - a \in \pm S\}$ has infinite chromatic number. One may attempt to prove Conjecture 3.2 analogously by fixing a set $S \subset \mathcal{E}_4$ and considering the graph \mathcal{G} with vertices $\{\mathbf{e}_i - \mathbf{e}_j : i \neq j \in \mathbb{N}\} \subset \mathbb{Z}^\omega$,

joining two vertices by an edge if they differ by an element of S . This attempt would succeed if we could prove that \mathcal{G} has infinite chromatic number but it may have finite chromatic number, as we demonstrate below. Since \mathcal{G} is not all of $\text{Cay}(S)$, this example does not disprove Conjecture 3.2.

For the sake of visualization, we work in \mathbb{F}_2^ω rather than \mathbb{Z}^ω . We will find a set $S \subset \mathcal{E}_4 \subset \mathbb{F}_2^\omega$ which is Bohr recurrent, and where the subgraph of $\text{Cay}(E)$ induced by the elements of \mathcal{E}_2 has chromatic number 2.

Instead of the usual presentation of \mathbb{F}_2^ω , we consider $\text{Fin}(\mathbb{Z} \times \mathbb{Z})$, the collection of finite subsets of $\mathbb{Z} \times \mathbb{Z}$, with the group operation of symmetric difference. This is a countably infinite group, all of whose elements have order 2, and is therefore isomorphic to \mathbb{F}_2^ω . Let

$$S_\square := \{(n, m), (n + d, m), (n, m + d), (n + d, m + d)\} : n, m \in \mathbb{Z}, d \in \mathbb{N}\} \subset \text{Fin}(\mathbb{Z} \times \mathbb{Z}),$$

so that S_\square is the collection of 4 element subsets of the integer plane forming the vertices of a square with sides parallel to the coordinate axes. Gallai's multidimensional generalization of van der Waerden's theorem (see Chapter 42 of [25] for exposition) implies S_\square is partition regular. Lemma 4.9 then implies S_\square is a Bohr recurrent subset of $\text{Fin}(\mathbb{Z} \times \mathbb{Z})$.

Let \mathcal{G} be the subgraph of $\text{Cay}(S_\square)$ induced by Fin_2 , the 2-element subsets of $\mathbb{Z} \times \mathbb{Z}$. In other words, \mathcal{G} is the graph whose vertices are the 2-element subsets of $\mathbb{Z} \times \mathbb{Z}$, where two vertices $\{a, b\}$, $\{c, d\}$ are joined by an edge if their symmetric difference forms the vertices of a square with sides parallel to the coordinate axes. We will prove that \mathcal{G} has chromatic number 2, by showing that its connected components are each isomorphic to a either a singleton, a pair of vertices connected by an edge, or an infinite path. To see this, fix a two element set $v = \{(n, m), (r, s)\} \subset \mathbb{Z} \times \mathbb{Z}$, and note that:

- (i) If v is not two of the four vertices of some square parallel to the coordinate axes, then v is not connected to any other element of Fin_2 by an edge of \mathcal{G} .
- (ii) If v forms two diagonally opposed vertices of a square, such as $\{(n, m), (n + d, m + d)\}$, then v is connected in \mathcal{G} to exactly one element v' of Fin_2 , consisting of the other two vertices of the square, and this v' is connected only to v .
- (iii) If v forms two vertices on the same side of a square, such as $v = \{(n, m), (n, s)\}$, then v lies on the infinite path P in \mathcal{G} whose vertices are the pairs $\{(n + jd, m), (n + jd, s)\} : j \in \mathbb{Z}\}$, where $d = |y - w|$. Note that each element in P is connected to exactly two other elements of P . It is easy to check that P contains no cycle, so it is indeed an infinite path.

Thus the connected components of \mathcal{G} have chromatic number 2, and \mathcal{G} itself has chromatic number 2.

Remark 5.1. To address Conjecture 3.2 directly, one could consider the group $G = (\mathbb{Z} \times \mathbb{Z})^\omega$ with basis $\{\mathbf{e}_{i,j} : i, j \in \mathbb{Z}\}$, and the sets

$$S := \{\mathbf{e}_{n,m} + \mathbf{e}_{n+d,m} - \mathbf{e}_{n,m+d} - \mathbf{e}_{n+d,m+d} : n, m \in \mathbb{Z}, d \in \mathbb{N}\},$$

$$G_2 := \{k\mathbf{e}_{n,m} + \ell\mathbf{e}_{p,q} : n, m, p, q, k, \ell \in \mathbb{Z}\},$$

and prove that S is a set of Bohr recurrence in G while the subgraph of $\text{Cay}(S)$ induced by G_2 has chromatic number 2.

6. SUGGESTIONS

For an abelian group G , the Bohr topology of G is the smallest topology containing every translate of every Bohr neighborhood of 0. A set $S \subset G$ is dense in the Bohr topology if for all $g \in G$, the translate $S - g$ is a set of Bohr recurrence.

Propositions 2.2, 2.8, and Theorem 1.3 of [17] suggest the following problems.

Problem 6.1. Prove or disprove: there is a set $S \subset \mathbb{Z}$ which is dense in the Bohr topology of \mathbb{Z} such that if $S' \subset S$ is also dense in the Bohr topology, then for all $m \in \mathbb{Z}$, $S' - m$ is a set of chromatic recurrence.

Problem 6.2. Prove or disprove: if $S \subset \mathbb{Z}$ is a set of density recurrence, then there is an $S' \subset S$ which is a set of chromatic recurrence and not a set of density recurrence.

The variations where \mathbb{Z} is replaced by another countable abelian group are equally interesting.

REFERENCES

1. Mathias Beiglböck, Vitaly Bergelson, and Alexander Fish, *Sumset phenomenon in countable amenable groups*, Adv. Math. **223** (2010), no. 2, 416–432. MR 2565535 (2011a:11011)
2. Vitaly Bergelson, *The multifarious Poincaré recurrence theorem*, Descriptive set theory and dynamical systems (Marseille-Luminy, 1996), London Math. Soc. Lecture Note Ser., vol. 277, Cambridge Univ. Press, Cambridge, 2000, pp. 31–57. MR 1774423
3. Vitaly Bergelson and Randall McCutcheon, *Recurrence for semigroup actions and a non-commutative Schur theorem*, Topological dynamics and applications (Minneapolis, MN, 1995), Contemp. Math., vol. 215, Amer. Math. Soc., Providence, RI, 1998, pp. 205–222. MR 1603193
4. Vitaly Bergelson and Imre Z. Ruzsa, *Sumsets in difference sets*, Israel J. Math. **174** (2009), 1–18. MR 2581205 (2011b:11014)

5. N. Bogoliouboff, *Sur quelques propriétés arithmétiques des presque-périodes*, Ann. Chaire Phys. Math. Kiev **4** (1939), 185–205. MR 20164
6. Michael Boshernitzan and Eli Glasner, *On two recurrence problems*, Fund. Math. **206** (2009), 113–130. MR 2576263
7. Erling Følner, *Note on a generalization of a theorem of Bogoliouboff*, Math. Scand. **2** (1954), 224–226. MR 69188
8. Nikos Frantzikinakis and Randall McCutcheon, *Ergodic theory: Recurrence*, 2007.
9. H. Furstenberg, *Recurrence in ergodic theory and combinatorial number theory*, Princeton University Press, Princeton, N.J., 1981, M. B. Porter Lectures. MR 603625 (82j:28010)
10. Berit Nilsen Givens, *Hypergraphs and chromatic numbers, with applications to the Bohr topology*, ProQuest LLC, Ann Arbor, MI, 2003, Thesis (Ph.D.)—The University of Wisconsin - Madison. MR 2704616
11. Berit Nilsen Givens and Kenneth Kunen, *Chromatic numbers and Bohr topologies*, Topology Appl. **131** (2003), no. 2, 189–202. MR 1981873
12. Eli Glasner, *On minimal actions of Polish groups*, vol. 85, 1998, 8th Prague Topological Symposium on General Topology and Its Relations to Modern Analysis and Algebra (1996), pp. 119–125. MR 1617456
13. Daniel Glasscock, Andreas Koutsogiannis, and Florian K. Richter, *On katznelson’s question for skew product systems*, 2021.
14. Colin C. Graham and Kathryn E. Hare, *Interpolation and Sidon sets for compact groups*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, Springer, New York, 2013. MR 3025283
15. John T. Griesmer, *Bohr neighborhoods in generalized difference sets*, arXiv e-prints (2021), arXiv:2108.01302.
16. ———, *A few sets of recurrence and open problems*, 2021, Draft.
17. ———, *Separating bohr denseness from measurable recurrence*, Discrete Analysis (2021), arxiv.org/abs/2002.06994, to appear.
18. ———, *Separating topological recurrence from measurable recurrence: exposition and extension of Kriz’s example*, arXiv e-prints (2021), arXiv:2108.01642.
19. Sophie Grivaux and Maria Roginskaya, *Some new examples of recurrence and non-recurrence sets for products of rotations on the unit circle*, Czechoslovak Math. J. **63(138)** (2013), no. 3, 603–627. MR 3125645
20. Bernard Host, Bryna Kra, and Alejandro Maass, *Variations on topological recurrence*, Monatsh. Math. **179** (2016), no. 1, 57–89. MR 3439271
21. Y. Katznelson, *Chromatic numbers of Cayley graphs on \mathbb{Z} and recurrence*, Combinatorica **21** (2001), no. 2, 211–219, Paul Erdős and his mathematics (Budapest, 1999). MR 1832446
22. Kenneth Kunen and Walter Rudin, *Lacunarity and the Bohr topology*, Math. Proc. Cambridge Philos. Soc. **126** (1999), no. 1, 117–137. MR 1681658
23. Igor Kříž, *Large independent sets in shift-invariant graphs: solution of Bergelson’s problem*, Graphs Combin. **3** (1987), no. 2, 145–158. MR 932131

24. Anh N. Le, *Interpolation sets and nilsequences*, Colloq. Math. **Online First** (2020), 1–19, doi:10.4064/cm7937-9-2019.
25. Alexander Soifer, *The mathematical coloring book*, Springer, New York, 2009, Mathematics of coloring and the colorful life of its creators, With forewords by Branko Grünbaum, Peter D. Johnson, Jr. and Cecil Rousseau. MR 2458293
26. E. Strzelecki, *On a problem of interpolation by periodic and almost periodic functions*, Colloq. Math. **11** (1963), 91–99. MR 160080
27. William A. Veech, *The equicontinuous structure relation for minimal Abelian transformation groups*, Amer. J. Math. **90** (1968), 723–732. MR 232377

Email address: jtgriesmer@gmail.com