

NEF AND ABUNDANT DIVISORS, SEMIAMPLENESS AND CANONICAL BUNDLE FORMULA

PRIYANKUR CHAUDHURI

ABSTRACT. In this paper, we use canonical bundle formulas to prove various generalizations of an old theorem of Kawamata on the semiampleness of nef and abundant log canonical divisors. In particular, we show that for klt pairs (X, B) with $K_X + B$ effective, $L \in \text{Pic } X$ nef, nefness and abundance of $K_X + B + L$ implies semiampleness. This essentially generalizes Kawamata's theorem to the setting of generalized abundance.

1. INTRODUCTION

This paper will be concerned with the following:

Question 1. *Let (X, B) be a klt pair and $L \in \text{Pic } X$ nef such that $K_X + B + L$ is nef and abundant, then is $K_X + B + L$ semiample?*

This is clearly related to the following conjecture of Lazić and Peternell [12].

Conjecture 2. (*Generalized Abundance*) *Let (X, B) be a klt pair with $K_X + B$ pseudoeffective and L a nef Cartier divisor on X such that $K_X + B + L$ is also nef. Then there exists a semiample \mathbb{Q} -divisor M on X such that $K_X + B + L \equiv M$.*

Note that we are not assuming pseudoeffectivity of $(K_X + B)$ in our question. There are two reasons why one might expect this to be true. Firstly, in [3], the author has shown that generalized abundance does continue to hold for many uniruled varieties. Secondly, if $L \in \text{Pic } X$ is a large multiple of a strictly nef divisor, then under the above assumptions, one can show that $K_X + B + L$ is ample. So it is natural to ask for a version in which the pseudoeffectivity of $K_X + B$ is not required. The surface case of this question is very easily seen to be true. We now state our main results which settle certain special cases.

Theorem 3. *Let $(X, B + \mathbf{M})$ be a generalized klt pair, H a nef \mathbb{Q} -Cartier divisor on X such that $H - (K_X + B + \mathbf{M}_X)$ is nef and abundant. Assume further that \mathbf{M} is b -semiample, $\kappa(aH - (K_X + B + \mathbf{M}_X)) \geq 0$*

and $\nu(aH - (K_X + B + \mathbf{M}_X)) = \nu(H - (K_X + B + \mathbf{M}_X))$ for some $a \in \mathbb{Q}_{>1}$. Then H is semiample.

The case $\mathbf{M} = 0$ of the above theorem is due to Kawamata [9] and Fujino [7]. Our approach of using canonical bundle formula is similar to that of Fujino's.

Theorem 4. *Let (X, B) be a klt pair with $\kappa(K_X + B) \geq 0$, H and L nef \mathbb{Q} -Cartier divisors on X such that $H - (K_X + B + L)$ is nef and abundant. Assume further that $\kappa(aH - (K_X + B + L)) \geq 0$ and $\nu(aH - (K_X + B + L)) = \nu(H - (K_X + B + L))$ for some $a \in \mathbb{Q}_{>1}$. Then H is semiample.*

This will be obtained as a consequence of a version of a generalized canonical bundle formula (Theorem 14) to be proved in the third section. It essentially generalizes Kawamata's theorem to the setting of generalized abundance.

2. PRELIMINARIES

We will always work with \mathbb{Q} -divisors.

Definition 2.1 (Abundant divisors). Let X be a normal projective variety and D be a nef \mathbb{Q} -divisor with Iitaka dimension $\kappa(X, D)$. The *numerical dimension* of D is defined as $\nu(X, D) = \max\{e \mid D^e \neq 0\}$. In general, $\kappa(X, D) \leq \nu(X, D)$. D is *abundant* if $\kappa(X, D) = \nu(X, D)$. Examples include big nef divisors, semiample divisors and their pullbacks.

Definition 2.2 (b-divisors (see [4])). Let X be a normal variety. A *b-divisor* \mathbf{D} on X is a collection of Weil divisors \mathbf{D}_Y on higher birational models $Y \rightarrow X$ of X that are compatible under pushforward. \mathbf{D}_Y is called the *trace* of \mathbf{D} on Y . The *Cartier closure* of a \mathbb{Q} -Cartier \mathbb{Q} -divisor D on X is the b- \mathbb{Q} divisor $\overline{\mathbf{D}}$ with trace $\overline{\mathbf{D}}_Y := \rho^*(D)$ on a birational model $\rho : Y \rightarrow X$. A b- \mathbb{Q} -divisor \mathbf{D} on a normal variety X is *b- \mathbb{Q} -Cartier* if it is the Cartier closure of a \mathbb{Q} -Cartier \mathbb{Q} -divisor on some birational model of X ; in this situation, we say that \mathbf{D} *descends* to Y . \mathbf{D} is called *b-nef* (resp. *b-semiample*) if it descends to a nef (resp. semiample) \mathbb{Q} -divisor on some birational model over X . Let (X, B) be a sub-pair consisting of a normal variety X and a \mathbb{Q} -divisor B on X such that $K_X + B$ is \mathbb{Q} -Cartier. The *discrepancy b-divisor* $\mathbf{A} = \mathbf{A}(X, B)$ is defined by $K_Y = f^*(K_X + B) + \mathbf{A}_Y$ for all models $Y \xrightarrow{f} X$. (X, B) is called *sub-klt* (resp. *sub-lc*) if all coefficients of $\mathbf{A}_Y(X, B)$ are > -1 (resp. ≥ -1) for some (and hence, by [10, Lemma 2.30], for all) birational models $f : Y \rightarrow X$. Thus in particular, if X is

smooth, all coefficients of B are less than 1 and B has normal crossing support, then (X, B) is sub-klt. In case $B \geq 0$, we drop the prefix sub.

Definition 2.3 (Generalized pairs and their singularities(see [2])). A *generalized sub-pair* $(X, B + \mathbf{M})$ consists of a normal projective variety X , a \mathbb{Q} -divisor B and a \mathbb{Q} -b-divisor \mathbf{M} on X such that:

- (1) $K_X + B + \mathbf{M}_X$ is \mathbb{Q} -Cartier.
- (2) \mathbf{M} is b-nef.

When $B \geq 0$, we drop the prefix sub.

Let (X, B) be a generalized sub-pair and $Y \xrightarrow{\mu} X$ a higher birational model of X . Let B_Y be defined by $K_Y + B_Y + \mathbf{M}_Y = \mu^*(K_X + B + \mathbf{M}_X)$. We say that $(X, B + \mathbf{M})$ is *sub-generalized klt* (resp. *sub-generalized lc*) if every coefficient of B_Y is less than 1 (resp. less than or equal to 1). We will call them sub-gklt and sub-glc in short. The *generalized discrepancy b-divisor* $\mathbf{A}(X, B + \mathbf{M})$ is then defined by $\mathbf{A}_Y(X, B + \mathbf{M}) = -B_Y$. Note that \mathbf{M}_Y does not contribute to the singularities of $(Y, B_Y + \mathbf{M})$ once it descends to a nef divisor on Y , i.e. $\mathbf{A}(Y, B_Y + \mathbf{M}_Y) = \mathbf{A}(Y, B_Y)$ and thus $(Y, B_Y + \mathbf{M}_Y)$ is gklt (resp. glc) iff (Y, B_Y) is klt (resp. lc).

Definition 2.4 (Generalized klt-trivial fibrations). Let (X, B) be a pair consisting of a normal variety X and a \mathbb{Q} -divisor B on X . A morphism $f : (X, B) \rightarrow Y$, Y normal is *generalized klt-trivial* if

- (1) f is surjective with connected fibers,
- (2) (X, B) is sub-klt over the generic point of Y ,
- (3) there exists a \mathbb{Q} -Cartier \mathbb{Q} -divisor D on Y and a nef \mathbb{Q} -divisor L on X such that $K_X + B + L \sim_{\mathbb{Q}} f^*D$
- (4) there exists a log-resolution $\pi' : (X', B_{X'}) \rightarrow (X, B)$ such that if $f' : (X', B_{X'}) \rightarrow Y$ is the induced morphism, then $\text{rank } f'_* \mathcal{O}_{X'}(\lceil -B_{X'} \rceil) = 1$.

The following fact seems to be well-known to the experts. I will include a proof for the convenience of readers and for want of a reference.

Lemma 5. *Let $f : (X, B) \rightarrow Y$ be a generalized klt-trivial fibration, $\sigma : Y' \rightarrow Y$ a generically finite morphism from a normal variety Y' . Let X' denote the desingularization of the main component of $X \times_Y Y'$ and let $f' : (X', B_{X'}) \rightarrow Y'$ be the induced morphism. Then f' is generalized klt-trivial.*

Proof. Only item (4) in the above definition needs to be verified. Since $(X', B_{X'})$ is sub-klt, the rank is surely ≥ 1 . We only need to show it

is ≤ 1 and may assume that X itself is smooth. Let $\pi : (X', B_{X'}) \rightarrow (X, B)$ be the induced morphism.

- (1) Suppose first that σ is birational. Then there exists an effective, integral, π -exceptional divisor E' on X' such that $K_{X'} = \pi^*(K_X) + E'$. The condition $K_{X'} + B_{X'} = \pi^*(K_X + B)$ then gives $B_{X'} + E' = \pi^*B_X$. If we can show that $\sigma \circ f' : X' \rightarrow Y'$ satisfies $\text{rank}(\sigma \circ f')_* \mathcal{O}_{X'}(\lceil -B_{X'} \rceil) = 1$, then $\text{rank} f'_* \mathcal{O}_{X'}(\lceil -B_{X'} \rceil) = 1$ also since $\sigma \circ f'$ and f' have the same general fiber. Thus we may assume $\sigma = id$ and thus $f \circ \pi = f'$. Now,

$$\begin{aligned} f'_* \mathcal{O}_{X'}(\lceil -B_{X'} \rceil) &= f'_* \mathcal{O}_{X'}(\lceil -\pi^*B_X + E' \rceil) \\ &= f'_* \mathcal{O}_{X'}(\lceil -\pi^*B_X \rceil + E') \text{ (since } E' \text{ is integral)} \\ &\subset f'_* \mathcal{O}_{X'}(\pi^* \lceil -B_X \rceil + E') \\ &= f_* \pi_* \mathcal{O}_{X'}(\pi^*(\lceil -B_X \rceil) + E') = f_* \mathcal{O}_X(\lceil -B_X \rceil) \otimes \pi_* \mathcal{O}_{X'}(E') \\ &= f_* \mathcal{O}_X(\lceil -B_X \rceil) \text{ (by Hartog's theorem since } E' \text{ is} \\ &\quad \pi\text{-exceptional)}. \end{aligned}$$

Thus $\text{rank} f'_* \mathcal{O}_{X'}(\lceil -B_{X'} \rceil) \leq 1$.

- (2) Now suppose that σ is finite. In this case also, there exists an integral divisor $R' \geq 0$ such that $K_{X'} = \pi^*(K_X) + R'$ ($\text{Supp}(R')$ is the ramification locus of π). This gives $B_{X'} + R' = \pi^*B_X$. To show that $\text{rank} f'_* \mathcal{O}_{X'}(\lceil -B_{X'} \rceil) = 1$, it suffices to show that R' is not f' -horizontal. Now [8, Chapter 3, Proposition 10.1 (b)] implies that $f^{-1}(Sm(\sigma)) \subset Sm(\pi)$ (Sm denotes the smooth locus of the respective morphism) or in other words, $f(\text{Branch}(\pi)) = f(\pi(R')) \subset \text{Branch}(\sigma)$. Thus R' can not be f' -horizontal. □

3. NEF AND ABUNDANT DIVISORS

We start off by proving two easy results. They provide positive evidence for Question 1. Recall that an *almost strictly nef* (ASN) divisor means a birational pullback of a strictly nef Cartier divisor. See [3], [16], [13] for some recent developments around ASN divisors.

Lemma 6. *Let X be a normal projective variety, $M \in \text{Pic } X$ be almost strictly nef (ASN) and abundant. Then M is big.*

Proof. By [9, Proposition 2.1], there exist morphisms $X \xleftarrow{\mu} X' \xrightarrow{f} Y$, where μ is birational, f a contraction and X' and Y are smooth and projective such that

$$(7) \quad M' := \mu^*M = f^*(M_Y)$$

for some $M_Y \in \text{Pic } Y$ nef and big. If $\dim Y = \dim X$, then f is generically finite and M is already big. So assume that $\dim Y < \dim X$. By definition of ASN, there exists $X \xrightarrow{\pi} X_0$ birational and $M_0 \in \text{Pic } X_0$ strictly nef such that $\pi^*(M_0) = M$. Let $E \subset X$ be the exceptional locus of π . Then for all curves $C \not\subset E$, $(M \cdot C) = (\pi^*(M_0) \cdot C) > 0$. If $F \subset X'$ is a general fiber of f , then $F \not\subset E$. Then there is a curve $C \subset F$ such that $C \not\subset E$ and thus $(M \cdot C) > 0$. This is not possible by (7). Therefore, $\dim Y = \dim X$ and M is big. \square

Corollary 8. *Let (X, B) be a klt pair, $L \in \text{Pic } X$ strictly nef such that $K_X + B + t_0L$ is abundant for some $t_0 > 2 \cdot \dim X$. Then $K_X + B + tL$ is ample for all $t > 2 \cdot \dim X$.*

Proof. $K_X + B + t_0L$ is strictly nef by the cone theorem and abundant, hence big by above lemma. Then $2(K_X + B + t_0L) - (K_X + B)$ is nef and big and thus $K_X + B + t_0L$ is semiample by the basepoint free theorem. Now it is easy to see that a strictly nef and semiample divisor is ample, thus $K_X + B + t_0L$ is ample. The conclusion now follows from [14, Lemma 1.3] (or rather its log version which has the same proof). \square

The following generalization of the basepoint-free theorem will be indispensable for us. It can be proved by a minor modification of the usual arguments ([10, page 78]).

Theorem 9. ([7, Theorem 2.1]) *Let (X, B) be a sub-klt pair, H a nef \mathbb{Q} -Cartier divisor on X . Assume that*

- (1) $rH - (K_X + B)$ is nef and big for some $r \in \mathbb{N}$.
- (2) $H^0(\mathcal{O}_X([\mathbf{A}(X, B)] + j\bar{\mathbf{H}})) \subset H^0(\mathcal{O}_X(jH))$ for all $j \in \mathbb{N}$.

Then H is semiample.

Proposition 10. *Let $(X, B + \mathbf{M})$ be a sub-generalized klt pair, H nef \mathbb{Q} -Cartier divisor on X such that $H - (K_X + B + \mathbf{M}_X)$ is nef and abundant. Suppose also that $\kappa(aH - (K_X + B + \mathbf{M}_X)) \geq 0$ and $\nu(aH - (K_X + B + \mathbf{M}_X)) = \nu(H - (K_X + B + \mathbf{M}_X))$ for some $a \in \mathbb{Q}_{>1}$. Letting*

$$(X, B + \mathbf{M}_X) \xleftarrow{\mu} (Y, B_Y + \mathbf{M}_Y) \xrightarrow{f} Z$$

be the standard diagram induced by the Iitaka fibration of $H - (K_X + B + \mathbf{M}_X)$ (see [9, Proposition 2.1]), suppose that the moduli b -divisor \mathbf{M}_Z induced on Z by the generalized sub-pair $(Y, B_Y + \mathbf{M}_Y)$ is b -nef. Then H is semiample.

Proof. We argue along the lines of [7].

Recall first that [9, Proposition 2.1, 2.3] says that there exist smooth projective varieties Y and Z and morphisms

$$(X, B + \mathbf{M}_X) \xleftarrow{\mu} (Y, B_Y + \mathbf{M}_Y) \xrightarrow{f} Z$$

where μ is birational, f is a surjective morphism with connected fibers birational to the Iitaka fibration of $H - (K_X + B + \mathbf{M}_X)$, $\mu^*(K_X + B + \mathbf{M}_X) = K_Y + B_Y + \mathbf{M}_Y$, where B_Y is defined by $\mu_*(\mathbf{M}_Y) = \mathbf{M}_X$. We may assume that \mathbf{M}_Y descends to a nef Cartier divisor on Y (possibly after replacing Y by a higher model) and there exists $D_Z \in \text{Pic}(Z)$ nef and big such that, setting $H_Y = \mu^*(H)$, we have

$$\mu^*(H - (K_X + B + \mathbf{M}_X)) = H_Y - (K_Y + B_Y + \mathbf{M}_Y) \sim_{\mathbb{Q}} f^*(D_Z).$$

Also, there exists $H_Z \in \text{Pic}(Z)$ nef such that $H_Y \sim_{\mathbb{Q}} f^*(H_Z)$. Note that (Y, B_Y) is sub-klt and

$$\mathbf{A}_Y(Y, B_Y + \mathbf{M}_Y) = -B_Y$$

As noted above, there exists a nef and big divisor D_Z and a nef divisor H_Z on Z such that $K_Y + B_Y + \mathbf{M}_Y = f^*(H_Z - D_Z)$. Now by assumption, we can write $H_Z - D_Z \sim_{\mathbb{Q}} K_Z + B_Z + \mathbf{M}_Z$ such that

- (1) \mathbf{M}_Z is \mathbb{Q} -Cartier and b-nef.
- (2) (Z, B_Z) is sub-klt. B_Z is defined as follows. Let $D \subset Z$ be any prime divisor. Let

$$a_D := \sup\{t \mid (Y, B_Y + tf^*(D)) \text{ is sub-lc over the generic point of } D\}$$

and define $B_Z = \sum(1 - a_D)D$. We note that replacing Y and Z by higher models, we may assume that $(Y, B_Y) \xrightarrow{f} (Z, B_Z)$ satisfies standard SNC assumptions (see [6, Definition 4.4]). Then the definition of a_D requires that all coefficients of $B_Y + tf^*(D)$ are ≤ 1 . Now, (Y, B_Y) being sub-klt implies that all coefficients of B_Y are < 1 and thus that $a_D > 0$ for all D . (Z, B_Z) is thus sub-klt.

Now let $(Z', B_{Z'}) \rightarrow (Z, B_Z)$ be a higher model such that $\mathbf{M}_{Z'}$ is nef. Then $(Z', B_{Z'})$ is of course also sub-klt. Let Y' be the normalization of the main component of $Y \times_Z Z' \rightarrow Z'$. Replacing Y and Z by Y' and Z' , we may assume that \mathbf{M}_Z is nef on Z . We need:

Lemma 11. (See [4, Lemma 9.2.2]) *Let $f : (Y, B_Y) \rightarrow Z$ be a fibration such that (Y, B_Y) is sub-klt over the generic point of Z . Let \mathbf{B} be the induced discriminant \mathbb{Q} -b-divisor of Z . Then $\mathcal{O}_Z([- \mathbf{B}]) \subset f_* \mathcal{O}_Y([\mathbf{A}(Y, B_Y)])$.*

As in the proof of [4, Proposition 9.2.3], this shows that $\mathcal{O}_Z([- \mathbf{B} + \mathbf{D}]) \subset f_* \mathcal{O}_Y([\mathbf{A}(Y, B_Y) + f^* \mathbf{D}])$ holds for every \mathbb{Q} -b-Cartier \mathbb{Q} -b-divisor \mathbf{D}

on Y : we can replace Z by a higher model Z' on which \mathbf{D} is \mathbb{Q} -Cartier and replace Y by the normalization of the main component of $Y \times_Z Z'$ as before. Then $(Y, B_Y - f^*D)$ is sub-klt over the generic point of Z with discriminant \mathbb{Q} -b-divisor $\mathbf{B} - \mathbf{D}$ and

$$\mathbf{A}(Y, B_Y - f^*D) = \mathbf{A}(Y, B_Y) + f^*(D).$$

Now the above lemma applied to $f : (Y, B_Y - f^*D) \rightarrow Z$ gives

$$\mathcal{O}_Z(\lceil -\mathbf{B} + \mathbf{D} \rceil) \subset f_*\mathcal{O}_Y(\lceil \mathbf{A}(Y, B_Y) + f^*\mathbf{D} \rceil).$$

as claimed. Thus

$$\mathcal{O}_Z(\lceil \mathbf{A}(Z, B_Z) \rceil + j\overline{\mathbf{H}}_Z) \subset f_*\mathcal{O}_Y(\lceil \mathbf{A}(Y, B_Y) \rceil + j\overline{\mathbf{H}}_Y)$$

for all $j \in \mathbb{Z}$. Taking H^0 gives

$$\begin{aligned} H^0(Z, \mathcal{O}_Z(\lceil \mathbf{A}(Z, B_Z) \rceil + j\overline{\mathbf{H}}_Z)) &\subset H^0(Y, \mathcal{O}_Y(\lceil \mathbf{A}(Y, B_Y) \rceil + j\overline{\mathbf{H}}_Y)) \\ &= H^0(X, \mu_* (\mathcal{O}_Y(\lceil \mathbf{A}(Y, B_Y) \rceil + j\overline{\mathbf{H}}_Y))) = H^0(X, \lceil A(X, B) \rceil + jH) \\ &= H^0(X, \mathcal{O}_X(jH)) = H^0(Y, \mathcal{O}_Y(jH_Y)) \\ &= H^0(Y, \mathcal{O}_Y(jf^*(H_Z))) = H^0(Z, \mathcal{O}_Z(jH_Z)). \end{aligned}$$

Thus H_Z is semiample by Theorem 9. \square

Corollary 12. *Let $(X, B + \mathbf{M})$ be a generalized klt pair, H a nef \mathbb{Q} -Cartier divisor on X such that $H - (K_X + B + \mathbf{M}_X)$ is nef and abundant. Assume further that \mathbf{M} is b -semiample, $\kappa(aH - (K_X + B + \mathbf{M}_X)) \geq 0$ and $\nu(aH - (K_X + B + \mathbf{M}_X)) = \nu(H - (K_X + B + \mathbf{M}_X))$ for some $a \in \mathbb{Q}_{>1}$. Then H is semiample.*

Proof. As in the proof of the previous Proposition, there exist morphisms

$$(X, B + \mathbf{M}_X) \xleftarrow{\mu} (Y, B_Y + \mathbf{M}_Y) \xrightarrow{f} Z$$

where \mathbf{M}_Y is semiample, Y and Z are both smooth, there exists $D_Z \in \text{Pic } Z$ and nef and big $H_Z \in \text{Pic } Z$ such that $H_Y \sim_{\mathbb{Q}} f^*(H_Z)$,

$$\mu^*(H - (K_X + B + \mathbf{M}_X)) = H_Y - (K_Y + B_Y + \mathbf{M}_Y) \sim_{\mathbb{Q}} f^*(D_Z)$$

and

$$\mu^*(K_X + B_X + \mathbf{M}_X) = K_Y + B_Y + \mathbf{M}_Y \sim_{\mathbb{Q}} f^*(H_Z - D_Z)$$

such that $\text{Supp}(B_Y)$ is SNC. Note: $\lceil -B_Y \rceil \geq 0$. Since \mathbf{M}_Y is semi-ample, for m sufficiently large and divisible, there exists $H = D_1 + \cdots + D_k \in |m\mathbf{M}_Y|$ such D_i is smooth for all i , $D_i \cap D_j = \emptyset$ for all $i \neq j$ and $\text{Supp } H \cup \text{Supp } B_Y$ is SNC. By taking $m \gg 0$, $(1/m)H \in |\mathbf{M}_Y|_{\mathbb{Q}}$ can be made arbitrarily small, thus $(Y, B_Y + (1/m)H)$ is sub-klt. Now consider the induced fibration $(Y, B_Y + (1/m)H) \xrightarrow{f} Z$. Then

$$f_*\mathcal{O}_Y(\lceil -(B_Y + (1/m)H) \rceil) = f_*\mathcal{O}_Y(\lceil -B_Y \rceil)$$

since H can be chosen to have no components in common with B_Y . Clearly $\text{rank } f_*\mathcal{O}_Y(\lceil -B_Y \rceil) \geq 1$ since (Y, B_Y) is sub-klt.

Now take $A \in \text{Pic } Z$ sufficiently ample such that $f_*\mathcal{O}_Y(\lceil -B_Y \rceil) \otimes A$ is globally generated. Since D_Z is big,

$$f_*\mathcal{O}_Y(\lceil -B_Y \rceil) \otimes A \subset f_*\mathcal{O}_Y(\lceil -B_Y \rceil) \otimes \mathcal{O}_Z(mD_Z)$$

if $m \gg 0$. Now note that since $B_Y = \mu_*^{-1}(B) + F$ where F is μ -exceptional and

$$0 \leq \lceil -B_Y \rceil = \lceil \mu_*^{-1}(-B) \rceil + \lceil -F \rceil,$$

we can conclude that $\lceil -B_Y \rceil$ is an effective μ -exceptional divisor. Now

$$\begin{aligned} H^0(Z, f_*\mathcal{O}_Y(\lceil -B_Y \rceil) \otimes A) &\subset H^0(Z, f_*\mathcal{O}_Y(\lceil -B_Y \rceil) \otimes \mathcal{O}_Z(mD_Z)) = \\ &H^0(Y, \mathcal{O}_Y(\lceil -B_Y \rceil) \otimes (f^*(mD_Z))). \end{aligned}$$

We have

$$H^0(Y, \mathcal{O}_Y(\lceil -B_Y \rceil) \otimes \mathcal{O}_Y(f^*(mD_Z))) = H^0(\mathcal{O}_Y(E) \otimes f^*(mD_Z))$$

where $E \geq 0$ is μ -exceptional. This equals

$$\begin{aligned} H^0(Y, \mathcal{O}_Y(E) \otimes \mu^*(m(H - (K + X + B + \mathbf{M}_X)))) &= \\ H^0(X, m(H - (K_X + B + \mathbf{M}_X))) &= H^0(Z, mD_Z). \end{aligned}$$

This shows that $\text{rank } f_*\mathcal{O}_Y(\lceil -B_Y \rceil) = 1$ i.e. $(Y, B_Y + 1/mH) \xrightarrow{f} Z$ is a klt-trivial fibration. If B_Z and \mathbf{M}_Z denote the induced discriminant and the moduli divisors on Z , then by [1, Theorem 2.7], $H_Z - D_Z = K_Z + B_Z + \mathbf{M}_Z$ where \mathbf{M}_Z is b-nef. Thus we are done by the $\mathbf{M} = 0$ case of the Proposition 9 (which is essentially worked out in [7]). \square

Remark 13. The above corollary can also be proved by combining [6, Theorem 4.12] with Proposition 10.

The following is a generalized canonical bundle formula which is used for proving theorem 4. It may also be of independent interest.

Theorem 14. *Let (Y, B) be a sub klt pair, $L \in \text{Pic } X$ nef and $f : Y \rightarrow Z$ be a contraction such that $K_Y + B + L \sim_{\mathbb{Q}, f} 0$ and rank $f_* \mathcal{O}_Y(\lceil \mathbf{A}(Y, B) \rceil) = 1$. Suppose $K_Y + B \geq_{\mathbb{Q}} 0$. Then the moduli b-divisor \mathbf{M}_Z induced on Z is b-nef.*

Proof. We write

$$(15) \quad K_Y + B_Y + L_Y \sim_{\mathbb{Q}} f^*(D_Z) = f^*(K_Z + B_Z + \mathbf{M}_Z).$$

as before. Let $F \subset Y$ be a general fiber of f . Then $(K_Y + B_Y)|_F \geq 0$. Suppose $(K_Y + B_Y)|_F >_{\mathbb{Q}} 0$. Then there exists a curve $C \subset F$ such that $((K_Y + B_Y)|_F \cdot C) > 0$ and thus $((K_Y + B_Y + L_Y)|_F \cdot C) > 0$. This contradicts (15). Thus $(K_Y + B_Y)|_F \sim_{\mathbb{Q}} 0$, $L|_F \sim_{\mathbb{Q}} 0$, where F is a general fiber of f . Let $d \in \mathbb{N}$ such that $\mathcal{O}_F(dL_Y|_F) \cong \mathcal{O}_F$. Now by [12, Lemma 3.1], after replacing Y and Z with higher models, we can assume that L_Y is numerically trivial on every fiber of f . By performing a flattening base change, we can replace Y and Z again by higher models and assume that f is flat. Note that $\mathcal{O}_F(dL_Y|_F) \cong \mathcal{O}_F$ is still true for a general fiber F . Then by the semicontinuity theorem,

$$z \rightarrow h^0(\mathcal{O}(dL_Y)|_{Y_z})$$

is an upper semicontinuous function on Z . Thus

$$\{z \in Z | h^0(\mathcal{O}(dL_Y|_{Y_z})) \geq 1\}$$

is closed in Z . Since $\mathcal{O}_Y(dL_Y)$ is trivial on a general fiber and numerically trivial on all fibers, this shows that $\mathcal{O}_Y(dL_Y|_{Y_z}) \cong \mathcal{O}_{Y_z}$ for all $z \in Z$. Now there exists a finite morphism $\theta : Z' \rightarrow Z$, from a smooth projective variety Z' such that letting Y' denote the desingularization of the main component of $Y \times_Z Z'$, the induced morphism $f' : Y' \rightarrow Z'$ is semistable in codimension 1. Since $\mathcal{O}_{Y'}(dL_{Y'})$ is trivial on all fibers of f' and f' is flat with reduced connected fibers over a large open set, there exists $P_{Z'} \in \text{Pic}(Z')$ nef such that $f'^*(P_{Z'}) = \mathcal{O}_{Y'}(dL_{Y'})$ (same arguments as [8, Chapter 3, Exercise 12.4]). Let $\alpha : Z'' \rightarrow Z'$ be a Bloch-Gieseker cover with Z'' smooth such that $\alpha^*(P_{Z'}) = L_{Z''}^{\otimes d}$ for some $L_{Z''} \in \text{Pic}(Z'')$ (see [11, Theorem 4.1.10]). Then $L_{Y''} = f''^*(L_{Z''})$ where $f'' : Y'' \rightarrow Z''$ is obtained by base changing f .

We now argue along the lines of [1, Lemma 5.2(5)] to show that \mathbf{M}_Z is b-nef. Let $b \in \mathbb{N}$ smallest such that $b(K_{Y''} + B_{Y''} + L_{Y''}) \sim_{f''} 0$. There exists $\phi \in K(Y'')$ such that $K_{Y''} + B_{Y''} + L_{Y''} + 1/b(\phi) = f''^*(K_{Z''} + B_{Z''} + \mathbf{M}_{Z''})$. Let V'' be the desingularization of the normalization of Y'' in $K(Y'')(\phi^{1/b})$. Note that V'' is irreducible by the minimality of b . Let

$$\pi'' : (V'', B_{V''}) \xrightarrow{\nu''} (Y'', B_{Y''}) \xrightarrow{f''} Z''$$

be the induced set-up.

Let $\Sigma_{f''} \subset Z''$ be the maximal closed subvariety such that f'' is smooth over $Z'' \setminus \Sigma_{f''}$. Replacing Z'' by a log-resolution, we can assume that $\Sigma_{f''}$ has SNC support. Also by [1, Proposition 5.4], there exists $\tau : \overline{Z'} \rightarrow Z''$ finite such that $\overline{Z'}$ admits an SNC divisor supporting $\tau^{-1}(\Sigma_{f''})$ and the locus where τ is not etale such that the induced morphism $(\overline{V'}, B_{\overline{V'}}) \xrightarrow{\overline{\pi'}} \overline{Z'}$ is semistable in codimension 1. Then $\overline{\pi'}_*(\omega_{\overline{V'}/\overline{Z'}})$ is nef ([1, Theorem 4.4]). If $\mathbf{M}_{\overline{Z'}}$ is the induced moduli b-divisor on $\overline{Z'}$, then $\tau^*\mathbf{M}_{Z''} = \mathbf{M}_{\overline{Z'}}$ ([1, Proposition 5.5]) and $(\theta \circ \alpha)^*(\mathbf{M}_{Z''}) = \mathbf{M}_Z$ ([1, Lemma 5.1]). So it suffices to show that $\mathbf{M}_{\overline{Z'}}$ is nef. The category of generalized klt-trivial fibrations is closed under generically finite base changes (Lemma 5). Thus $\overline{f'} : (\overline{Y'}, B_{\overline{Y'}}) \rightarrow \overline{Z'}$ is again generalized klt-trivial.

To simplify notation, we thus replace $\overline{V'}, \overline{Y'}, \overline{Z'}, \overline{\nu'}, \overline{\pi'}, \overline{f'}$ by V, Y, Z, ν, π, f respectively.

$$\begin{aligned} & \text{Then } \pi_*(\omega_{V/Z}) \\ &= f_*(\bigoplus_{i=0}^{b-1} \mathcal{O}_Y(K_Y + \lceil i/b(\phi) \rceil) \otimes f^*\omega_Z^{-1}) \otimes f_*(\bigoplus_{i=0}^{b-1} \mathcal{O}_Y(\lfloor -i/b(\phi) \rfloor)) \\ & \text{([15, Lemma 2.3]). We have} \end{aligned}$$

$$f_*(\bigoplus_{i=0}^{b-1} \mathcal{O}_Y(\lfloor -i/b(\phi) \rfloor)) = \mathcal{O}_Z$$

and

$$i/b(\phi) = i(-K_{Y/Z} - B_Y - L_Y + f^*(B_Z + \mathbf{M}_Z)).$$

from which it follows that

$$\begin{aligned} & \pi_*(\omega_{V/Z}) \\ &= \bigoplus_{i=0}^{b-1} f_*\mathcal{O}_Y(\lceil (1-i)K_{Y/Z} - iB_Y - iL_Y + if^*(B_Z + \mathbf{M}_Z) \rceil) \end{aligned}$$

where the i -th summand on the right hand side is the eigensheaf corresponding to the eigenvalue ζ^i , where ζ is a primitive b -th root of unity, the action being that of $\mathbb{Z}/(b)$ on $K(V)$ and ψ is a rational function on V such that $\psi^b = \phi$ (see [5, page 23-24] for more details). By [1, Theorem 4.4], $\pi_*(\omega_{V/Z})$ is a nef vector bundle and hence so are the direct summands. Then

$$(16) \quad K_V + B_V + (\psi) = \pi^*(K_Z + B_Z + \mathbf{M}_Z - L_Z).$$

Let

$$\mathcal{L} = f_*\mathcal{O}_Y([-B_Y - L_Y + f^*(B_Z + \mathbf{M}_Z)])$$

be the ζ -eigensheaf. Then $\mathcal{L} \in \text{Pic } Z$ (it has rank 1 since $\text{rank } f_*\mathcal{O}_Y([-B_Y]) = 1$ and L_Y is trivial on all fibers).

Claim 17. There exists $U \subset Z$ large open such that $(-B_V + \pi^*B_Z)|_{\pi^{-1}(U)} \geq 0$.

Proof. By (16), we see that $(K_{V/Z})^h + B_V^h + (\psi)^h = 0$ (h denotes the horizontal part) and hence that B_V^h is a \mathbb{Z} -divisor. (V, B_V) being klt, $[-B_V^h] = -B_V^h \geq 0$. Let $B_V^v = \sum_j d_j P_j$ (v denotes the vertical part). We can ignore those P_j such that $\text{codim}(\pi(P_j)) \geq 2$ and consider only those that map to some prime component of B_Z . Recall that $B_Z = \sum_l \delta_l Q_l$ where

$$1 - \delta_l = \sup\{t|(Y, B_Y + t\pi^*(Q_l)) \text{ is sub-lc over the generic point of } Q_l\}.$$

which translates to $\sum_j d_j P_j + \pi^*(\sum_l (1 - \delta_l) Q_l) \leq 1$. Fix l and consider those P_j which map to Q_l . Since π is semistable in codimension 1, this leads to $\max_{P_j \rightarrow Q_l} (d_j) + 1 - \delta_l \leq 1$ with equality holding for some j . Thus $-d_j + \delta_l \geq 0$ for all j such that $\pi(P_j) = Q_l$. This proves the claim. \square

Also note the above proof shows $(-B_V + \pi^*B_Z)$ supports no fibers over codimension 1 points of Z . Let $H \subset V$ be a general fiber of π . Then $(\psi|_H) + K_H = -B_H \geq 0$ (Note everything is integral here). Thus ψ is a rational section of $\pi_*(\omega_{V/Z})$. Since $\psi \rightarrow \zeta \cdot \psi$ we conclude that ψ is a rational section of \mathcal{L} . Let $\mathcal{K}(Z)$ be the (constant) sheaf of rational functions on Z . Considering the injection $\mathcal{L} \hookrightarrow \mathcal{K}(Z)$ that sends a rational section e of \mathcal{L} to e/ψ shows that $\mathcal{L} \subset \mathcal{K}(Z) \cdot \psi$.

The above claim and (16) show that there exists $U \subset Z$ large open such that $\mathcal{O}_Z(\mathbf{M}_Z - L_Z) \cdot \psi|_U \subset \pi_*(\omega_{V/Z})|_U$ and thus $\mathcal{O}_Z(\mathbf{M}_Z - L_Z) \cdot \psi|_U \subset \mathcal{L}|_U$. Conversely, let $a \cdot \psi$ be a section of \mathcal{L} , $a \in K(Z)$. Then it is a section of $\pi_*(\omega_{V/Z})$, so $(\pi^*(a) \cdot \psi) + K_{V/Z} \geq 0$. Since

$$(18) \quad (\pi^*(a) \cdot \psi) + K_{V/Z} = \pi^*((a) + \mathbf{M}_Z - L_Z) + (-B_V + \pi^*B_Z)$$

we have $\pi^*((a) + \mathbf{M}_Z - L_Z) + (-B_V + \pi^*B_Z) \geq 0$. Since $-B_V + \pi^*B_Z$ contains no fibers over codimension 1 points of Z , this implies that $(a) + \mathbf{M}_Z - L_Z \geq 0$ and hence that $\mathcal{L} \subset \mathcal{O}_Z(\mathbf{M}_Z - L_Z) \cdot \psi$. We therefore conclude that

$$\mathcal{L}|_U = \mathcal{O}_Z(\mathbf{M}_Z - L_Z) \cdot \psi|_U.$$

Since U is large and \mathcal{L} is a line bundle, $\mathcal{L} = \mathcal{O}_Z(\mathbf{M}_Z - L_Z) \cdot \psi$. Since \mathcal{L} and L_Z are nef, we conclude that \mathbf{M}_Z is nef. \square

Corollary 19. *Let (X, B) be a klt pair with $\kappa(K_X + B) \geq 0$, H and L nef \mathbb{Q} -Cartier divisors on X such that $H - (K_X + B + L)$ is nef and abundant. Assume further that $\kappa(aH - (K_X + B + L)) \geq 0$ and $\nu(aH - (K_X + B + L)) = \nu(H - (K_X + B + L))$ for some $a \in \mathbb{Q}_{>1}$. Then H is semiample.*

Proof. Let $(X, B + L) \xleftarrow{\mu} (Y, B_Y + L_Y) \xrightarrow{f} Z$ be the standard diagram as before. Then as in the proof of Corollary 12, we can show that $\text{rank } f_*\mathcal{O}_Y([-B_Y]) = 1$. Now the conclusion follows from Proposition 10 and Theorem 14. \square

Remark 20. Thus in particular, if (X, B) is a klt pair with $K_X + B$ effective and if $L \in \text{Pic } X$ is nef such that $K_X + B + L$ nef and abundant, then $K_X + B + L$ is semiample. The non-vanishing conjecture ([12, page 1]) predicts that assuming $K_X + B$ pseudoeffective would suffice for this to hold.

4. ACKNOWLEDGEMENTS

I thank my advisor Patrick Brosnan for several enlightening discussions on the contents of this paper and Christopher Hacon, an email conversation with whom led to the start of this work. I would also like to express my gratitude to Caucher Birkar and Jason Starr for their valuable comments. The author was partially supported by the Hauptman fellowship during the preparation of this work.

REFERENCES

- [1] F. Ambro. *Shokurov's Boundary property*. *J. Differential Geometry*, 67 (2004), 229-255.
- [2] C. Birkar. *Generalised pairs in birational geometry*. arxiv.org/abs/2008.01008
- [3] P. Chaudhuri. *Strictly nef divisors and some remarks on a conjecture of Ser-rano*. <https://arxiv.org/pdf/2008.05009v4.pdf>
- [4] A. Corti (editor). *Flips for 3-folds and 4-folds*. *Oxford Lecture Series in Mathematics and its Applications*, 35. Oxford University Press, Oxford, 2007. x+189 pp
- [5] H. Esnault, E. Viehweg. *Lectures on Vanishing Theorems*. *DMV Seminar, 20*. Birkhäuser Verlag, Basel, 1992. vi+164 pp.
- [6] S. Filipazzi. *On a generalized canonical bundle formula and generalized adjunction*. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) 21 (2020), 1187-1221.
- [7] O. Fujino. *On Kawamata's theorem*. *EMS Ser. Congr. Rep.* Eur. Math. Soc., Zürich, 2011.

- [8] R. Hartshorne. *Algebraic Geometry. Graduate Texts in Mathematics*, No. 52. Springer-Verlag, New York-Heidelberg, 1977. xvi+496 pp
- [9] Y. Kawamata. *Pluricanonical systems on minimal algebraic varieties. Invent. Math.* 79 (1985), no. 3, 567–588.
- [10] J. Kollár, S. Mori. *Birational geometry of algebraic varieties*. With the collaboration of C. H. Clemens and A. Corti. Translated from the 1998 Japanese original. Cambridge Tracts in Mathematics, 134. Cambridge University Press, Cambridge, 1998. viii+254 pp. ISBN: 0-521-63277-3 14E3
- [11] R. Lazarsfeld. *Positivity in Algebraic Geometry 1 : Line bundles and Linear Series* Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 49. Springer-Verlag, Berlin, 2004. xviii+385 pp.
- [12] V. Lazić, Th. Peternell. *On Generalised Abundance, I.* arXiv:1808.00438v2 [math.AG]
- [13] H. Liu. *On strictly nef divisors on K -trivial fourfolds.* <https://arxiv.org/pdf/2105.07259.pdf>
- [14] F. Serrano. *Strictly nef divisors and Fano threefolds. J. Reine Angew. Math.* 1995, 187–206.
- [15] E. Viehweg. *Weak Positivity and the Additivity of the Kodaira Dimension for Certain Fibre Spaces. Advanced studies in Pure Mathematics* 1 (1983), 329-353.
- [16] G. Zhong. *Strictly nef divisors on singular threefolds.* <https://arxiv.org/pdf/2105.07681.pdf>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK,
MD 20742

Email address: pchaudhu@umd.edu