

SOFIC MEAN DIMENSION OF TYPICAL ACTIONS AND A COMPARISON THEOREM

LEI JIN AND YIXIAO QIAO

ABSTRACT. We refine two results in the paper entitled “Sofic mean dimension” by Hanfeng Li, improving two inequalities with two equalities, respectively, for sofic mean dimension of typical actions. On the one hand, we study sofic mean dimension of full shifts, for which, Li provided an upper bound which however is not optimal. We prove a more delicate estimate from above, which is optimal for sofic mean dimension of full shifts over arbitrary alphabets (i.e. compact metrizable spaces). Our refinement, together with the techniques (in relation to an estimate from below) in the paper entitled “Mean dimension of full shifts” by Masaki Tsukamoto, eventually allows us to get the exact value of sofic mean dimension of full shifts over any finite dimensional compact metrizable spaces. On the other hand, we investigate finite group actions. In contrast to the case that the acting group is infinite (and amenable), Li showed that if a finite group acts continuously on a finite dimensional compact metrizable space, then sofic mean dimension may be different from (strictly less than) the classical (i.e. amenable) mean dimension (an explicitly known value in this case). We strengthen this result by proving a sharp lower bound, which, combining with the upper bound, gives the exact value of sofic mean dimension for all the actions of finite groups on finite dimensional compact metrizable spaces. Furthermore, this equality leads to a satisfactory comparison theorem for those actions, deciding when sofic mean dimension would coincide with classical mean dimension. Moreover, our two results, in particular, verify for a typical class of sofic group actions that sofic mean dimension does not depend on sofic approximation sequences.

1. MAIN RESULTS

1.1. **Overview.** The purpose of this paper is to obtain the exact value of *sofic mean dimension* of a typical class of *sofic group actions*. We will state our main theorems very quickly (i.e. with a minimum introduction) in this section (Section 1). The definition of sofic mean dimension as well as all the other necessary terminologies are located in Section 2. We shall prove our main results in Section 3 and Section 4. A standard explanation in relation to two different approaches to the definition of sofic mean dimension, for the reader’s convenience, can be found in the appendix.

2010 *Mathematics Subject Classification.* 37B99; 54F45.

Key words and phrases. Sofic mean dimension; Full shift; Finite group action.

1.2. Convention. Throughout this paper the symbol \mathbb{N} is planned to denote the set of *positive* integers. All acting groups are always assumed to be *countable* and *discrete*. If we let a sofic group G act continuously on a compact metrizable space X and if we let Σ be a sofic approximation sequence for G , then we denote its sofic mean dimension by $\text{mdim}_\Sigma(X, G)$.

1.3. Full shifts. Possibly the most canonical and fundamental actions are the *full shifts*. For a compact metrizable space K and a group G we denote by (K^G, σ_G) the shift action of G on the product space K^G .

For \mathbb{Z} -actions the exact value of the mean dimension of full shifts was obtained satisfactorily by Tsukamoto [Tsu19] (provided that the alphabet K is *finite dimensional*). We notice that it is straightforward to generalize Tsukamoto's result to actions of amenable groups (e.g. this can be done with almost no additional effort with the help of [JQ]). But if we proceed to the case of sofic group actions (which are a much wider class of group actions), then the situation becomes nontrivial immediately.

Actually, Li [Li13, Theorem 7.1] showed (for any sofic approximation sequence Σ for G) that

$$\text{mdim}_\Sigma(K^G, \sigma_G) \leq \dim(K).$$

However, this upper bound for $\text{mdim}_\Sigma(K^G, \sigma_G)$ is not optimal in general. A main ingredient of this paper is to refine Li's upper bound with a more delicate estimate from above:

$$\text{mdim}_\Sigma(K^G, \sigma_G) \leq \inf_{n \in \mathbb{N}} \frac{\dim(K^n)}{n}.$$

Our approach is substantially different from Li's¹. We shall explain about our difficulty shortly in Remark 1.2. As we will see in a moment, our refinement gives an optimal upper bound for $\text{mdim}_\Sigma(K^G, \sigma_G)$ which applies to all alphabets K , which eventually allows us to get the exact value of sofic mean dimension of full shifts over any finite dimensional compact metrizable spaces. This is our first main result.

Theorem 1.1 (Main theorem 1). *Let K be a finite dimensional compact metrizable space. Let G be a sofic group and Σ a sofic approximation sequence for G . The following assertion is true:*

$$\text{mdim}_\Sigma(K^G, \sigma_G) = \begin{cases} \dim(K), & \text{if } K \text{ satisfies } \dim(K \times K) = 2 \dim(K) \\ \dim(K) - 1, & \text{otherwise} \end{cases}.$$

¹Roughly speaking, Li's method can be regarded as (in some sense, a kind of) *local* analysis, i.e. using the notion of *sofic metric mean dimension* which dominates sofic mean dimension from above. On the contrary, we shall adopt a *global* treatment of a sharp upper bound for $\text{mdim}_\Sigma(K^G, \sigma_G)$.

Remark 1.2. For mean dimension of \mathbb{Z} -actions, the above two types of inequalities for full shifts can be deduced directly from each other. More precisely, let us denote by $\text{mdim}(K^{\mathbb{Z}}, \sigma_{\mathbb{Z}})$ the (classical) mean dimension of the \mathbb{Z} -shift on $K^{\mathbb{Z}}$. The inequality $\text{dim}(K^n)/n \leq \text{dim}(K)$ (for $n \in \mathbb{N}$) is generally true. Conversely, if we assume that $\text{mdim}(K^{\mathbb{Z}}, \sigma_{\mathbb{Z}}) \leq \text{dim}(K)$ for any compact metrizable space K , then replacing the alphabet K by K^n for an arbitrary $n \in \mathbb{N}$ and noting that $((K^n)^{\mathbb{Z}}, \sigma_{\mathbb{Z}})$ is (dynamically) isomorphic to $(K^{\mathbb{Z}}, \sigma_{\mathbb{Z}}^n)$ we see that $\text{mdim}(K^{\mathbb{Z}}, \sigma_{\mathbb{Z}}) \leq \text{dim}(K^n)/n$ for all $n \in \mathbb{N}$. A point here is that the acting group \mathbb{Z} contains a *subgroup* $n\mathbb{Z}$ of index $n \in \mathbb{N}$ which is possible to be arbitrarily large, and furthermore, any subgroup action of \mathbb{Z} also forms a \mathbb{Z} -action.² Unfortunately, the same trick does not apply to the context of sofic group actions essentially, because in general we cannot expect an arbitrary group action to have this nice property and also because we do not have such an equality for sofic mean dimension of product actions with respect to the *same* sofic approximation sequence. Thus, Li's handling of $\text{mdim}_{\Sigma}(K^G, \sigma_G)$ is not adequate for the purpose of obtaining its exact value. This turns out to be one of the main obstacles that we have to overcome. To deal with an upper bound in a more careful way is crucial to our proof.

Corollary 1.3. *Let K be a finite dimensional compact metrizable space and G an infinite amenable group. The classical mean dimension of (K^G, σ_G) satisfies that*

$$\text{mdim}(K^G, \sigma_G) = \begin{cases} \text{dim}(K), & \text{if } \text{dim}(K \times K) = 2 \text{dim}(K) \\ \text{dim}(K) - 1, & \text{otherwise} \end{cases}.$$

Remark 1.4. It follows from Lemma 2.3 that if the acting group G is finite then the classical mean dimension of (K^G, σ_G) satisfies the following equality:

$$\text{mdim}(K^G, \sigma_G) = \begin{cases} \text{dim}(K), & \text{if } \text{dim}(K \times K) = 2 \text{dim}(K), \\ \text{dim}(K) - 1 + \frac{1}{|G|}, & \text{otherwise.} \end{cases}$$

In contrast to the above corollary which holds for actions of infinite amenable groups, Theorem 1.1 is more satisfactory as it applies to (and unifies the cases for) both finite and infinite sofic groups G .

1.4. Double finite actions. The simplest examples of group actions (probably) are *double finite actions*, namely, actions of *finite* groups on *finite dimensional* compact metrizable spaces.

²For all the actions of (infinite) amenable groups we would like to refer to the paper [JQ] by the authors, in which we established a satisfactory and unexpected formula for mean dimension of product actions. But we do not state this formula here as we shall not use it in the present paper.

For an action of an amenable group G on a compact metrizable space X we denote its *classical* mean dimension by $\text{mdim}(X, G)$. Li [Li13, Section 3] showed (for any sofic approximation sequence Σ for G) that

$$\text{mdim}_\Sigma(X, G) \leq \text{mdim}(X, G);$$

if in addition, G is infinite, then

$$\text{mdim}_\Sigma(X, G) = \text{mdim}(X, G).$$

However, for a finite group G (which is automatically amenable and sofic) this equality might be false. Indeed, Li [Li13, Section 3] provided a bunch of examples of double finite actions, for which, the above inequality is strict. So a natural and further question we may ask is to decide when (i.e. for which kind of sofic group actions) these two values will agree.

More precisely, we let a finite group G act continuously on a finite dimensional compact metrizable space X . Note that in this case, by the definition of classical mean dimension we have apparently that

$$\text{mdim}(X, G) = \frac{1}{|G|} \cdot \dim(X).$$

Nevertheless, the exact value of its sofic mean dimension (with respect to a sofic approximation sequence Σ for G) is not clear to us. In fact, Li [Li13, Lemma 3.9] gave an upper bound for $\text{mdim}_\Sigma(X, G)$, which is finer than $\text{mdim}(X, G)$, as follows:

$$\text{mdim}_\Sigma(X, G) \leq \frac{1}{|G|} \cdot \inf_{n \in \mathbb{N}} \frac{\dim(X^n)}{n}.$$

The second main result of this paper is to solve this problem completely, strengthening Li's estimate with an equality:

$$\text{mdim}_\Sigma(X, G) = \frac{1}{|G|} \cdot \inf_{n \in \mathbb{N}} \frac{\dim(X^n)}{n}.$$

Theorem 1.5 (Main theorem 2). *If a finite group G acts continuously on a finite dimensional compact metrizable space X and if Σ is a sofic approximation sequence for G , then*

$$\text{mdim}_\Sigma(X, G) = \begin{cases} \frac{1}{|G|} \cdot \dim(X), & \text{if } \dim(X \times X) = 2 \dim(X) \\ \frac{1}{|G|} \cdot (\dim(X) - 1), & \text{otherwise} \end{cases}.$$

Corollary 1.6 (Comparison theorem). *Suppose that a finite group G acts continuously on a finite dimensional compact metrizable space X and that Σ is a sofic approximation sequence for G . Then $\text{mdim}_\Sigma(X, G) = \text{mdim}(X, G)$ (resp. $\text{mdim}_\Sigma(X, G) < \text{mdim}(X, G)$) if and only if $\dim(X \times X) = 2 \dim(X)$ (resp. $\dim(X \times X) < 2 \dim(X)$).*

Remark 1.7. As we can see, Li’s theorem [Li13, Section 3], Theorem 1.1 and Theorem 1.5, in particular, present a typical class of sofic group actions, for which, sofic mean dimension does not depend on the sofic approximation sequences, while it has not yet been confirmed if this is still true for all sofic group actions.

1.5. Open questions. To end this section we pose the following open problems in this direction, which seem to be worth considering in order to have a complete understanding of sofic mean dimension of full shifts and amenable group actions.

- (i) Let K be an *infinite* dimensional compact metrizable space. Is the statement $\text{mdim}_\Sigma(K^G, \sigma_G) = +\infty$ true?
- (ii) Let a *finite* group G act continuously on an *infinite* dimensional compact metrizable space X . Is the statement $\text{mdim}_\Sigma(X, G) = +\infty$ true?
- (iii) Let an *infinite* sofic group G act continuously on a *finite* dimensional compact metrizable space X . Is the statement $\text{mdim}_\Sigma(X, G) \leq 0$ true?

We give some remarks here around this topic. The first problem is concerning sofic mean dimension of full shifts over infinite dimensional alphabets. For the case $G = \mathbb{Z}$ this problem was initially posed by Tsukamoto [Tsu19, Problem 1.3]. The second and third problems are closely related to sofic mean dimension of double finite actions, as they remove one of the finiteness restrictions, respectively. Moreover, regarding Question (iii), it is clear that $\text{mdim}_\Sigma(X, G) < +\infty$ (by definition); and if G is an infinite amenable group then the solution to Question (iii) is obviously affirmative (by the definition of classical mean dimension and by Li’s theorem [Li13, Section 3] stating that sofic mean dimension coincides with classical mean dimension provided the acting group is infinite and amenable). Finally, we conjecture that the answers to Questions (i) and (ii) are positive, while the answer to Question (iii) is negative. We hope to return to further research in relation to all the above questions in the future.

Acknowledgements. L. Jin was supported by Basal Funding AFB 170001 and Fondecyt Grant No. 3190127. Y. Qiao was supported by NNSF of China No. 11901206.

2. A BRIEF REVIEW OF SOFIC MEAN DIMENSION

2.1. Historical background. Both the notions of sofic groups and mean dimension originate with Misha Gromov [Gro99a, Gro99b] in 1999. Mean dimension is a fairly good numerical topological invariant of dynamical systems. It was investigated systematically within the framework of amenable group actions by Lindenstrauss and Weiss [LW00] (including wonderful applications to dynamical systems) around 2000. As an analogue

of topological dimension (i.e. the Lebesgue covering dimension), its advantage has now been shown excellently in the study of dynamical systems whose topological entropy is equal to infinity and whose phase space has infinite topological dimension. In 2013, Li [Li13] successfully extended the definition of mean dimension to the context of sofic group actions. Strictly speaking, Li [Li13] introduced a new concept with a fully novel view, called sofic mean dimension, which applies to all sofic group actions; and further, Li [Li13] established its connection with classical mean dimension as well, which is highly satisfactory and which we have mentioned in the previous section. It is worth pointing out that Li's generalization to the soficity setting is totally nontrivial. This section is dedicated to fundamental material on terminologies and notations. The writing of this section is mostly borrowed, for consistency, from the paper [JQ] by the authors, with necessary modifications adapting to the present paper.

2.2. Group actions. Throughout this subsection we let G be a topological group. The terminology “ G acts continuously on (or simply, acts on) a compact metrizable space X ” is planned to be understood a continuous mapping

$$\Phi : G \times X \rightarrow X, \quad (g, x) \mapsto gx$$

satisfying the following conditions:

$$\Phi(e, x) = x, \quad \Phi(gh, x) = \Phi(g, \Phi(h, x)), \quad \forall x \in X, \forall g, h \in G,$$

where e is the identity element of the group G . We generally omit the mapping Φ provided we have already gotten the action of G on X clear.

Full shifts are a typical class of group actions. This notion has appeared in Theorem 1.1. Let K be a compact metrizable space. The **shift action** of G on the product space K^G is defined as follows:

$$\sigma_G : G \times K^G \rightarrow K^G, \quad (g, (x_h)_{h \in G}) \mapsto (x_{hg})_{h \in G}.$$

Here K is usually called an *alphabet*.

Let G act continuously on compact metrizable spaces X_n , respectively, where n ranges over R which is some subset (finite or infinite) of \mathbb{N} . We define the **product action** of G on the product space $\prod_{n \in R} X_n$ as follows:

$$g(x_n)_{n \in R} = (gx_n)_{n \in R}, \quad \forall g \in G, \forall (x_n)_{n \in R} \in \prod_{n \in R} X_n.$$

Product actions will be considered in the sequel (e.g. the proofs of the main theorems).

2.3. Sofic groups. We denote by $|F|$ the cardinality of a set F . For every $d \in \mathbb{N}$ we write $[d]$ for the set $\{k \in \mathbb{N} : 1 \leq k \leq d\}$ and $\text{Sym}(d)$ for the group of permutations of $[d]$. A group G is **sofic** if there is a sequence

$$\Sigma = \{\sigma_i : G \rightarrow \text{Sym}(d_i)\}_{i \in \mathbb{N}}$$

together with a sequence $\{d_i\}_{i \in \mathbb{N}} \subset \mathbb{N}$ such that the following three conditions are satisfied:

- $\lim_{i \rightarrow \infty} \frac{1}{d_i} |\{k \in [d_i] : \sigma_i(st)(k) = \sigma_i(s)\sigma_i(t)(k)\}| = 1$ for all $s, t \in G$;
- $\lim_{i \rightarrow \infty} \frac{1}{d_i} |\{k \in [d_i] : \sigma_i(s)(k) \neq \sigma_i(t)(k)\}| = 1$ for all distinct $s, t \in G$;
- $\lim_{i \rightarrow \infty} d_i = +\infty$.

Such a sequence Σ is called a **sofic approximation sequence** for G .

Remark 2.1. Note that the third condition will be fulfilled automatically if we additionally assume the group G to be infinite.

Remark 2.2. The sofic groups form a fairly extensive class, which contain in particular all amenable groups and all residually finite groups. We notice that the existence of a non-sofic group is still unknown.

2.4. A dichotomy in dimension theory. We denote by $\dim(K)$ the topological dimension (i.e. the Lebesgue covering dimension) of a compact metrizable space K . If the space K is empty, then we set $\dim(K) = -\infty$. For a finite dimensional (nonempty) compact metrizable space K , since it was classically known [Dra01] that

$$2 \dim(K) - 1 \leq \dim(K \times K) \leq 2 \dim(K)$$

and since $\dim(K)$ must be a nonnegative integer, we have either

$$\dim(K \times K) = 2 \dim(K)$$

or

$$\dim(K \times K) = 2 \dim(K) - 1.$$

For a friendly treatment of the following result [Dra01] in dimension theory we refer to [Tsu19, Theorem 2.5].

Lemma 2.3. *Let K be a finite dimensional compact metrizable space. Then for every $n \in \mathbb{N}$*

$$\dim(K^n) = \begin{cases} n \dim(K), & \text{if } K \text{ satisfies } \dim(K \times K) = 2 \dim(K), \\ n \dim(K) - n + 1, & \text{otherwise.} \end{cases}$$

2.5. ϵ -embeddings, ρ_2 and ρ_∞ . Let X and P be two compact metrizable spaces, and let ρ be a compatible metric on X . For $\epsilon > 0$ a continuous mapping $f : X \rightarrow P$ is called an **ϵ -embedding** with respect to ρ if $f(x) = f(x')$ implies $\rho(x, x') < \epsilon$, for all $x, x' \in X$. Let $\text{Widim}_\epsilon(X, \rho)$ be the minimum (topological) dimension $\dim(P)$ of a compact metrizable space P which admits an ϵ -embedding $f : X \rightarrow P$ with respect to ρ .

Remark 2.4. We may verify that $\dim(X) = \lim_{\epsilon \rightarrow 0} \text{Widim}_\epsilon(X, \rho)$.

Let K be a compact metrizable space with a compatible metric ρ on it. For every $n \in \mathbb{N}$ we define on the product space K^n two compatible metrics ρ_2 and ρ_∞ as follows:

$$\rho_2 \left((x_i)_{i \in [n]}, (y_i)_{i \in [n]} \right) = \sqrt{\frac{1}{n} \sum_{i \in [n]} (\rho(x_i, y_i))^2},$$

$$\rho_\infty \left((x_i)_{i \in [n]}, (y_i)_{i \in [n]} \right) = \max_{i \in [n]} \rho(x_i, y_i).$$

We do not include $n \in \mathbb{N}$ in the notations ρ_2 and ρ_∞ because this does not cause any ambiguity.

The following result due to Tsukamoto [Tsu19, Lemma 3.1] is very practical. We shall employ it in our argument.

Theorem 2.5. *Let K be a finite dimensional compact metrizable space. Let ρ be a compatible metric on K . Then there is some $\delta > 0$ such that*

$$\text{Widim}_\epsilon(K^n, \rho_\infty) \geq n \cdot (\dim(K) - 1)$$

for all $n \in \mathbb{N}$ and all $0 < \epsilon < \delta$.

2.6. Amenable group actions and classical mean dimension. A group G is said to be **amenable** if there exists a sequence $\{F_n\}_{n \in \mathbb{N}}$ of nonempty finite subsets of G such that for any $g \in G$

$$\lim_{n \rightarrow \infty} \frac{|F_n \triangle gF_n|}{|F_n|} = 0.$$

Such a sequence $\{F_n\}_{n \in \mathbb{N}}$ is called a **Følner sequence** of the group G .

Let an amenable group G act continuously on a compact metrizable space X . Take a Følner sequence $\{F_n\}_{n \in \mathbb{N}}$ of G and a compatible metric ρ on X . For a nonempty finite subset F of G we set

$$\rho_F(x, x') = \rho_\infty \left((gx)_{g \in F}, (gx')_{g \in F} \right), \quad \forall x, x' \in X.$$

It is clear that ρ_F is also a compatible metric on X . The **mean dimension** of (X, G) is defined by

$$\text{mdim}(X, G) = \lim_{\epsilon \rightarrow 0} \lim_{n \rightarrow \infty} \frac{\text{Widim}_\epsilon(X, \rho_{F_n})}{|F_n|}.$$

It is well known that the limits in the above definition always exist. The value $\text{mdim}(X, G)$ is independent of the choices of a Følner sequence $\{F_n\}_{n \in \mathbb{N}}$ of G and a compatible metric ρ on X .

2.7. Sofic mean dimension and its relation to classical mean dimension. Suppose that a sofic group G acts continuously on a compact metrizable space X . Let

$$\Sigma = \{\sigma_i : G \rightarrow \text{Sym}(d_i)\}_{i \in \mathbb{N}}$$

be a sofic approximation sequence for G . We equip X with a compatible metric ρ . For a finite subset F of G , $\delta > 0$ and a map $\sigma : G \rightarrow \text{Sym}(d)$ (where $d \in \mathbb{N}$) we define

$$\text{Map}(\rho, F, \delta, \sigma) = \{\phi : [d] \rightarrow X : \rho_2(\phi \circ \sigma(s), s\phi) \leq \delta, \forall s \in F\}.$$

We consider the set $\text{Map}(\rho, F, \delta, \sigma)$ as a compact subspace of the product space X^d . In our context we usually write $\phi = (\phi_l)_{l \in [d]} \in X^d$ for $\phi : [d] \rightarrow X$ (i.e. for $l \in [d]$ we write ϕ_l for $\phi(l)$). The **sofic mean dimension** of (X, G) is defined by

$$\text{mdim}_\Sigma(X, G) = \sup_{\epsilon > 0} \inf_{F \subset G \text{ finite}, \delta > 0} \limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(\text{Map}(\rho, F, \delta, \sigma_i), \rho_\infty)}{d_i}.$$

Similar to the classical mean dimension, the definition of $\text{mdim}_\Sigma(X, G)$ does not depend on the compatible metrics ρ on X . Nevertheless, as we mentioned previously, it is not clear yet if there is an example of a sofic approximation sequence Σ' different from Σ , which leads to a different value $\text{mdim}_{\Sigma'}(X, G)$.

Li [Li13, Section 3] built a satisfying bridge between sofic mean dimension and classical mean dimension as follows.

Theorem 2.6. *If an infinite amenable group G acts continuously on a compact metrizable space X then $\text{mdim}_\Sigma(X, G) = \text{mdim}(X, G)$.*

For product actions of sofic groups Li [Li13, Section 2] showed a useful inequality for sofic mean dimension:

Lemma 2.7. *Let a sofic group G act continuously on compact metrizable spaces X_n , respectively, where n runs over some $R \subset \mathbb{N}$. Let Σ be a sofic approximation sequence for G . For the product action of G on the product space $\prod_{n \in R} X_n$ we have the following subadditivity $\text{mdim}_\Sigma(\prod_{n \in R} X_n, G) \leq \sum_{n \in R} \text{mdim}_\Sigma(X_n, G)$.*

3. PROOF OF THEOREM 1.1

Theorem 1.1 will follow directly from Lemma 2.3, Lemma 3.1 and Lemma 3.2. Let us start with settings. We let K be a finite dimensional compact metrizable space, G a sofic group and

$$\Sigma = \{\sigma_i : G \rightarrow \text{Sym}(d_i)\}_{i \in \mathbb{N}}$$

a sofic approximation sequence for G . What we are dealing with is the shift action σ_G of G on the product space K^G .

We denote by e the identity element of the group G . We fix a (countable) family $\{\alpha_g\}_{g \in G}$ of positive real numbers such that

$$\alpha_e = 1, \quad \sum_{g \in G} \alpha_g < 2.$$

We fix a compatible metric D on the alphabet K and take a metric ρ on K^G compatible with the product topology as follows:

$$\rho(x, y) = \sum_{g \in G} \alpha_g D(x_g, y_g), \quad (x = (x_g)_{g \in G}, y = (y_g)_{g \in G} \in K^G).$$

For simplicity we write $\text{diam}(K^G)$ for the diameter of the product space K^G with respect to the metric ρ . Without loss of generality we may assume $\text{diam}(K^G) = 1$.

Lemma 3.1 (Estimate from above).

$$\text{mdim}_\Sigma(K^G, \sigma_G) \leq \inf_{n \in \mathbb{N}} \frac{\dim(K^n)}{n}.$$

Proof. First of all, we note that

$$\inf_{n \in \mathbb{N}} \frac{\dim(K^n)}{n} = \lim_{i \rightarrow \infty} \frac{\dim(K^{d_i})}{d_i},$$

because $\dim(\cdot \times \cdot)$ (and in particular, $\dim(K^n)$ with respect to $n \in \mathbb{N}$) is subadditive and because $d_i \rightarrow +\infty$ as $i \rightarrow \infty$.

We take $\epsilon > 0$ and $\eta > 0$ arbitrarily and fix them temporarily. Since both of them are arbitrary, it suffices (by definition) to prove that there exist a finite (and nonempty) subset F_0 of G and some $\delta_0 > 0$ satisfying that

$$\limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(\text{Map}(\rho, F_0, \delta_0, \sigma_i), \rho_\infty)}{d_i} \leq \eta + \lim_{i \rightarrow \infty} \frac{\dim(K^{d_i})}{d_i}.$$

We choose a finite subset F_0 of G , containing the identity element e , such that if $\xi = (\xi_g)_{g \in G}$ and $\xi' = (\xi'_g)_{g \in G}$ coming from K^G satisfy $D(\xi_s, \xi'_s) \leq \epsilon/2$ for all $s \in F_0$, then they satisfy $\rho(\xi, \xi') < \epsilon$. We take $\delta_0 > 0$ sufficiently small such that

$$\delta_0 < \min \left\{ \frac{\eta}{(\dim(K) + 1) \cdot |F_0|^2}, \frac{\epsilon^2}{16} \right\}.$$

So now the main task is to estimate the term $\text{Widim}_\epsilon(\text{Map}(\rho, F_0, \delta_0, \sigma_i), \rho_\infty)$, for all sufficiently large $i \in \mathbb{N}$, from above.

We fix $i \in \mathbb{N}$ for the moment.

For every $j \in [d_i]$ we define a mapping

$$J_j : \text{Map}(\rho, F_0, \delta_0, \sigma_i) \rightarrow [0, 1]$$

by sending

$$\phi = (\phi_j)_{j \in [d_i]} \in \text{Map}(\rho, F_0, \delta_0, \sigma_i) \subset (K^G)^{d_i}$$

to

$$J_j(\phi) = \max \left\{ \max_{s \in F_0} \left(\rho(s\phi_j, \phi_{\sigma_i(s)(j)}) - \sqrt{\delta_0} \right), 0 \right\},$$

where $\phi_j = (\phi_j(g))_{g \in G} \in K^G$. Clearly, each mapping $J_j : \text{Map}(\rho, F_0, \delta_0, \sigma_i) \rightarrow [0, 1]$ is continuous.

Let $CK = ([0, 1] \times K) / \sim$ be the cone generated by K , where $(0, x) \sim (0, x')$ for all $x, x' \in K$. We denote by λx the equivalence class of $(\lambda, x) \in [0, 1] \times K$. We set $* = 0x$ and call the symbol $*$ the vertex of the cone. We notice that the following inequality is clear:

$$\dim(K) \leq \dim(CK) \leq \dim(K) + 1.$$

We construct a mapping H_i as follows:

$$H_i : \text{Map}(\rho, F_0, \delta_0, \sigma_i) \rightarrow K^{d_i} \times (CK)^{|F_0| \cdot d_i},$$

$$\phi = (\phi_j)_{j \in [d_i]} \mapsto \left((\phi_j(e))_{j \in [d_i]}, (J_j(\phi)\phi_j|_{F_0})_{j \in [d_i]} \right),$$

where we write $\phi_j|_{F_0}$ for $(\phi_j(s))_{s \in F_0} \in K^{F_0}$, which is the restriction of $\phi_j \in K^G$ to K^{F_0} , and where we write $J_j(\phi)\phi_j|_{F_0}$ for $(J_j(\phi)\phi_j(s))_{s \in F_0} \in (CK)^{|F_0|}$. Obviously, $H_i : \text{Map}(\rho, F_0, \delta_0, \sigma_i) \rightarrow K^{d_i} \times (CK)^{|F_0| \cdot d_i}$ is a continuous mapping (since every mapping $J_j : \text{Map}(\rho, F_0, \delta_0, \sigma_i) \rightarrow [0, 1]$ ($j \in [d_i]$) is continuous).

We now claim that $H_i : \text{Map}(\rho, F_0, \delta_0, \sigma_i) \rightarrow K^{d_i} \times (CK)^{|F_0| \cdot d_i}$ is an ϵ -embedding with respect to ρ_∞ .

To verify this claim, we take $\varphi = (\varphi_j)_{j \in [d_i]}$ and $\psi = (\psi_j)_{j \in [d_i]}$ in $\text{Map}(\rho, F_0, \delta_0, \sigma_i)$ and we assume that $H_i(\varphi) = H_i(\psi)$. We then need to show that $\rho_\infty(\varphi, \psi) < \epsilon$. In fact, it follows from $H_i(\varphi) = H_i(\psi)$ that for all $j \in [d_i]$

$$\varphi_j(e) = \psi_j(e), \quad J_j(\varphi)\varphi_j|_{F_0} = J_j(\psi)\psi_j|_{F_0}.$$

For each $j \in [d_i]$ we have two cases:

- either $J_j(\varphi)\varphi_j|_{F_0} = J_j(\psi)\psi_j|_{F_0} = *^{|F_0|}$, i.e. $J_j(\varphi) = J_j(\psi) = 0$;
- or $J_j(\varphi)\varphi_j|_{F_0} = J_j(\psi)\psi_j|_{F_0} \in (CK \setminus \{*\})^{|F_0|}$, i.e. $J_j(\varphi) = J_j(\psi) > 0$;

where we denote by the notation $*^{|F_0|}$ the unique element $(*, *, \dots, *)$ in the set $\{*\}^{|F_0|}$. If some $j \in [d_i]$ encounters the former case, then

$$\rho(s\varphi_j, \varphi_{\sigma_i(s)(j)}) \leq \sqrt{\delta_0}, \quad \rho(s\psi_j, \psi_{\sigma_i(s)(j)}) \leq \sqrt{\delta_0}, \quad \forall s \in F_0.$$

Since $s\varphi_j(e) = \varphi_j(s)$ and $s\psi_j(e) = \psi_j(s)$, this implies that

$$D(\varphi_j(s), \varphi_{\sigma_i(s)(j)}(e)) \leq \sqrt{\delta_0}, \quad D(\psi_j(s), \psi_{\sigma_i(s)(j)}(e)) \leq \sqrt{\delta_0}, \quad \forall s \in F_0.$$

We note that

$$\varphi_l(e) = \psi_l(e), \quad \forall l \in [d_i].$$

Hence we have

$$D(\varphi_j(s), \psi_j(s)) \leq 2\sqrt{\delta_0} < \epsilon/2, \quad \forall s \in F_0.$$

Thus, we deduce from the choice of F_0 that in this case

$$\rho(\varphi_j, \psi_j) < \epsilon.$$

If some $j \in [d_i]$ encounters the latter case, then it follows directly from the definition of CK that

$$\varphi_j|_{F_0} = \psi_j|_{F_0}.$$

By the choice of F_0 , this implies that in this case we also have

$$\rho(\varphi_j, \psi_j) < \epsilon.$$

Therefore we conclude that

$$\rho_\infty(\varphi, \psi) = \max_{j \in [d_i]} \rho(\varphi_j, \psi_j) < \epsilon.$$

This proves the claim.

By this claim we get an upper bound for $\text{Widim}_\epsilon(\text{Map}(\rho, F_0, \delta_0, \sigma_i), \rho_\infty)$:

$$\text{Widim}_\epsilon(\text{Map}(\rho, F_0, \delta_0, \sigma_i), \rho_\infty) \leq \dim(H_i(\text{Map}(\rho, F_0, \delta_0, \sigma_i))).$$

So it remains to deal with the term $\dim(H_i(\text{Map}(\rho, F_0, \delta_0, \sigma_i)))$.

Here let us put a slightly more detailed explanation in relation to the construction of H_i . We take an arbitrary $j \in [d_i]$. Intuitively speaking, for any $\varphi \in \text{Map}(\rho, F_0, \delta_0, \sigma_i)$ if $J_j(\varphi) = 0$ then we say that φ_j is *good*, because the family $\{\varphi_k(e) : k \in [d_i]\}$ is able to tell almost everything fairly well about $\varphi_j \in K^G$ (i.e. it covers information sufficiently close to $\{\varphi_j(s) : s \in F_0\}$), while if $J_j(\varphi) > 0$ then we say that φ_j is *bad*, because in this case the needed information indicated by the family $\{\varphi_k(e) : k \in [d_i]\}$ is far from necessary for admitting an ϵ -embedding unless we additionally record almost the whole φ_j (namely, $\{\varphi_j(s) : s \in F_0\}$ itself). But when we look at any fixed $j \in [d_i]$ it is always possible for φ_j to be good or bad as $\varphi \in \text{Map}(\rho, F_0, \delta_0, \sigma_i)$ changes. So seemingly we have to record a quite large rectangle $\{\varphi_k(s) : s \in F_0, k \in [d_i]\}$. However, if we simply record nearly everything then the dimension will be absolutely out of control. The point here is that for a given $j \in [d_i]$ we *cannot* expect φ_j to *move in a continuous way* between good and bad regions as φ moves within $\text{Map}(\rho, F_0, \delta_0, \sigma_i)$. This is why we need consider the cone CK when building H_i , as to recording every φ_j continuously in a more flexible and productive approach.

Next we estimate $\dim(H_i(\text{Map}(\rho, F_0, \delta_0, \sigma_i)))$ from above. For any $\phi = (\phi_j)_{j \in [d_i]}$ coming from $\text{Map}(\rho, F_0, \delta_0, \sigma_i)$ we set

$$\Omega_\phi(\rho, F_0, \delta_0, \sigma_i) = [d_i] \setminus \{j \in [d_i] : \rho(s\phi_j, \phi_{\sigma_i(s)(j)}) \leq \sqrt{\delta_0}, \forall s \in F_0\}.$$

This is a subset of $[d_i]$. On the one hand, since $\phi = (\phi_j)_{j \in [d_i]} \in \text{Map}(\rho, F_0, \delta_0, \sigma_i)$, we have

$$\sum_{j \in [d_i]} \rho(s\phi_j, \phi_{\sigma_i(s)(j)})^2 \leq \delta_0 \cdot d_i, \quad \forall s \in F_0$$

and therefore

$$\delta_0 \cdot |\Omega_\phi(\rho, F_0, \delta_0, \sigma_i)| \leq \sum_{j \in [d_i]} \sum_{s \in F_0} \rho(s\phi_j, \phi_{\sigma_i(s)(j)})^2 \leq |F_0| \cdot \delta_0^2 \cdot d_i.$$

It follows that

$$0 \leq |\Omega_\phi(\rho, F_0, \delta_0, \sigma_i)| \leq |F_0| \cdot \delta_0 \cdot d_i.$$

On the other hand, we notice that for all $j \in [d_i] \setminus \Omega_\phi(\rho, F_0, \delta_0, \sigma_i)$ we have $J_j(\phi) = 0$. This implies that for any given $\phi \in \text{Map}(\rho, F_0, \delta_0, \sigma_i)$ there are $(d_i - |\Omega_\phi(\rho, F_0, \delta_0, \sigma_i)|) \cdot |F_0|$ entries of $H_i(\phi)$, which will take the value $*$. More precisely, the compact metrizable space $H_i(\text{Map}(\rho, F_0, \delta_0, \sigma_i))$ is contained in the compact metrizable space

$$\bigcup_{l \in \mathbb{Z}, 0 \leq l \leq |F_0| \cdot \delta_0 \cdot d_i} \bigcup_{k_1, \dots, k_{d_i} \in \{0,1\}, k_1 + \dots + k_{d_i} = l} K^{d_i} \times \prod_{j=1}^{d_i} (\{*\}^{|F_0|})^{1-k_j} \times ((CK)^{|F_0|})^{k_j},$$

where $\{*\}^{|F_0|} \times ((CK)^{|F_0|})^0 = \{*\}^{|F_0|}$ and $(\{*\}^{|F_0|})^0 \times (CK)^{|F_0|} = (CK)^{|F_0|}$. Therefore we deduce that

$$\begin{aligned} \dim(H_i(\text{Map}(\rho, F_0, \delta_0, \sigma_i))) &\leq \max_{0 \leq l \leq |F_0| \cdot \delta_0 \cdot d_i} (\dim(K^{d_i}) + l \cdot |F_0| \cdot \dim(CK)) \\ &\leq \dim(K^{d_i}) + |F_0|^2 \cdot \delta_0 \cdot d_i \cdot (\dim(K) + 1). \end{aligned}$$

We note that here we have used the following classically known equality and inequality: For any compact metrizable spaces A and A'

$$\dim(A \cup A') = \max\{\dim(A), \dim(A')\}, \quad \dim(A \times A') \leq \dim(A) + \dim(A').$$

Thus, we obtain that

$$\text{Widim}_\epsilon(\text{Map}(\rho, F_0, \delta_0, \sigma_i), \rho_\infty) \leq \dim(K^{d_i}) + |F_0|^2 \cdot \delta_0 \cdot d_i \cdot (\dim(K) + 1).$$

Since $i \in \mathbb{N}$ is arbitrary, we conclude (from the choice of δ_0) that

$$\limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(\text{Map}(\rho, F_0, \delta_0, \sigma_i), \rho_\infty)}{d_i} \leq \eta + \lim_{i \rightarrow \infty} \frac{\dim(K^{d_i})}{d_i}.$$

This is as desired. □

Lemma 3.2 (Estimate from below).

$$\text{mdim}_\Sigma(K^G, \sigma_G) \geq \inf_{n \in \mathbb{N}} \frac{\dim(K^n)}{n}.$$

Proof. First of all, let us make a reduction. In order to show the statement it suffices to prove

$$\text{mdim}_\Sigma(K^G, \sigma_G) \geq \dim(K) - 1.$$

In fact, such an inequality will apply to all finite dimensional compact metrizable spaces K , in particular to K^n , for any $n \in \mathbb{N}$, which will indicate that

$$\text{mdim}_\Sigma((K^n)^G, \sigma_G) \geq \dim(K^n) - 1, \quad \forall n \in \mathbb{N}.$$

Since the action $((K^n)^G, \sigma_G)$ is (dynamically) isomorphic to the product action of σ_G on $(K^G)^n$, we will have, by Lemma 2.7, that

$$\text{mdim}_\Sigma(K^G, \sigma_G) \geq \frac{\dim(K^n) - 1}{n}, \quad \forall n \in \mathbb{N}$$

which will imply that

$$\text{mdim}_\Sigma(K^G, \sigma_G) \geq \lim_{n \rightarrow \infty} \frac{\dim(K^n)}{n} = \inf_{n \in \mathbb{N}} \frac{\dim(K^n)}{n}.$$

This is as required.

In the remaining part of the proof we prove

$$\text{mdim}_\Sigma(K^G, \sigma_G) \geq \dim(K) - 1.$$

By Theorem 2.5 we can find some $\epsilon > 0$ sufficiently small, which depends only on the finite dimensional compact metric space (K, D) , such that for all $n \in \mathbb{N}$

$$\text{Widim}_\epsilon(K^n, D_\infty) \geq n \cdot (\dim(K) - 1).$$

We fix it in the proof. We take $\delta > 0$ and a (nonempty) finite subset F of G arbitrarily, and fix them in the proof. By definition it suffices to show that

$$\limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(\text{Map}(\rho, F, \delta, \sigma_i), \rho_\infty)}{d_i} \geq \dim(K) - 1.$$

For every $i \in \mathbb{N}$ we define a mapping P_i as follows:

$$\begin{aligned} P_i : K^{d_i} &\rightarrow (K^G)^{d_i}, \quad p = (p_j)_{j \in [d_i]} \rightarrow \phi = (\phi_j)_{j \in [d_i]}, \\ \phi_j &= (\phi_j(g))_{g \in G} \in K^G, \quad \forall j \in [d_i], \\ \phi_j(g) &= \begin{cases} p_{\sigma_i(g)(j)}, & \text{if } g \in G \setminus \{e\} \\ p_j, & \text{if } g = e \end{cases}. \end{aligned}$$

Clearly, for any $i \in \mathbb{N}$ the mapping $P_i : K^{d_i} \rightarrow (K^G)^{d_i}$ is continuous and it is distance-increasing with respect to the metric D_∞ on K^{d_i} and the metric ρ_∞ on $(K^G)^{d_i}$, i.e.

$$D_\infty(p, p') \leq \rho_\infty(P_i(p), P_i(p')), \quad \forall p, p' \in K^{d_i}.$$

We observe that for all sufficiently large $i \in \mathbb{N}$ it will be true that $P_i(K^{d_i})$ is contained in $\text{Map}(\rho, F, \delta, \sigma_i)$. (We will verify this observation in a moment.) It follows that

$$\begin{aligned} \limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(\text{Map}(\rho, F, \delta, \sigma_i), \rho_\infty)}{d_i} &\geq \limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(P_i(K^{d_i}), \rho_\infty)}{d_i} \\ &\geq \limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(K^{d_i}, D_\infty)}{d_i} \\ &\geq \limsup_{i \rightarrow \infty} \frac{d_i \cdot (\dim(K) - 1)}{d_i} \\ &= \dim(K) - 1. \end{aligned}$$

This will end the proof.

Finally, it remains to verify that for all sufficiently large $i \in \mathbb{N}$ we have that $P_i(K^{d_i})$ is a subset of $\text{Map}(\rho, F, \delta, \sigma_i)$. We choose a (nonempty) finite subset E of G , containing the identity element $e \in G$, such that if $\xi = (\xi_g)_{g \in G}$ and $\xi' = (\xi'_g)_{g \in G}$ in K^G satisfy $\xi_g = \xi'_g$ for all $g \in E$, then they satisfy $\rho(\xi, \xi') < \delta/2$. For every $i \in \mathbb{N}$ we let

$$Q_i = \{j \in [d_i] : \sigma_i(s) \circ \sigma_i(t)(j) = \sigma_i(st)(j), \forall s \in E, \forall t \in F \cup \{e\}\}.$$

Since both E and F are finite, there is some $k \in \mathbb{N}$ sufficiently large such that for any integer $i > k$

$$\frac{|Q_i|}{d_i} > 1 - \frac{3\delta^2}{8}.$$

Now we show that

$$P_i(K^{d_i}) \subset \text{Map}(\rho, F, \delta, \sigma_i), \quad \forall i > k.$$

We fix an integer $i > k$ and take an arbitrary $p = (p_j)_{j \in [d_i]} \in K^{d_i}$. Let $\phi = P_i(p)$ and write $\phi = ((\phi_j(g))_{g \in G})_{j \in [d_i]} \in (K^G)^{d_i}$. We note that for any $j \in Q_i$ we have $\sigma_i(e)(j) = j$, and therefore it is clear that

$$\phi_j(g) = p_{\sigma_i(g)(j)}, \quad \forall g \in G, \forall j \in Q_i.$$

It follows that for any $j \in Q_i \cap (\sigma_i(t))^{-1}(Q_i)$ and any $t \in F \cup \{e\}$

$$t\phi_j(s) = \phi_j(st) = p_{\sigma_i(st)(j)} = p_{\sigma_i(s) \circ \sigma_i(t)(j)} = \phi_{\sigma_i(t)(j)}(s), \quad \forall s \in E.$$

By the choice of E we have

$$\rho(t\phi_j, \phi_{\sigma_i(t)(j)}) < \delta/2, \quad \forall t \in F, \forall j \in Q_i \cap (\sigma_i(t))^{-1}(Q_i).$$

Thus, we conclude that for all $t \in F$

$$\begin{aligned} \rho_2(\phi \circ \sigma_i(t), t\phi) &= \sqrt{\frac{1}{d_i} \cdot \sum_{j \in [d_i]} (\rho(\phi_{\sigma_i(t)(j)}, t\phi_j))^2} \\ &\leq \sqrt{\frac{\delta^2}{4} + \frac{|[d_i] \setminus (Q_i \cap (\sigma_i(t))^{-1}(Q_i))|}{d_i}} \\ &\leq \sqrt{\frac{\delta^2}{4} + 2 \cdot \left(1 - \frac{|Q_i|}{d_i}\right)} \\ &\leq \delta. \end{aligned}$$

This implies that $\phi \in \text{Map}(\rho, F, \delta, \sigma_i)$. The statement follows. \square

4. PROOF OF THEOREM 1.5

The proof of Theorem 1.5 will be fulfilled by Lemma 2.3, Lemma 4.1 and Lemma 4.2. To begin with, we put some settings. Let G be a finite group. Suppose that G acts continuously on a finite dimensional compact metrizable space X . We fix a compatible metric ρ on X and a sofic approximation sequence $\Sigma = \{\sigma_i : G \rightarrow \text{Sym}(d_i)\}_{i \in \mathbb{N}}$ for G . Without loss of generality, we assume $\text{diam}(X) = 1$.

Lemma 4.1 (Estimate from above).

$$\text{mdim}_\Sigma(X, G) \leq \frac{1}{|G|} \cdot \inf_{n \in \mathbb{N}} \frac{\dim(X^n)}{n}.$$

Proof. This was initially shown in [Li13]. We reproduce it here, with a slightly simplified proof (in terms of Widim), for completeness. The difference between Li's proof and our proof is that we shall consider ϵ -embeddings into a cone.

We fix $\epsilon > 0$ and $\theta > 0$ arbitrarily. It suffices to show that there is some $\delta_0 > 0$ such that

$$\limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(\text{Map}(\rho, G, \delta_0, \sigma_i), \rho_\infty)}{d_i} \leq \frac{1}{|G|} \cdot \inf_{n \in \mathbb{N}} \frac{\dim(X^n)}{n} + \theta.$$

For every $i \in \mathbb{N}$ we let $Q_i \subset [d_i]$ be the intersection of the following two sets:

$$\begin{aligned} & \{j \in [d_i] : \sigma_i(g)(j) \neq \sigma_i(h)(j), \forall g \neq h \in G\}, \\ & \{j \in [d_i] : \sigma_i(g)(\sigma_i(h)(j)) = \sigma_i(gh)(j), \forall g, h \in G\}. \end{aligned}$$

For any $j \in [d_i]$ and $A \subset [d_i]$ we set

$$\sigma_i(G)(j) = \{\sigma_i(g)(j) : g \in G\} \subset [d_i], \quad \sigma_i(G)(A) = \{\sigma_i(g)(a) : g \in G, a \in A\} \subset [d_i].$$

We note that by the definition of Q_i we have

$$|\sigma_i(G)(j)| = |G|, \quad \sigma_i(G)(\sigma_i(G)(j)) = \sigma_i(G)(j), \quad \forall j \in Q_i.$$

It follows that $\sigma_i(G)(j) = \sigma_i(G)(l)$, for all $l \in \sigma_i(G)(j)$. We put a $j \in Q_i$ into a set C_i and remove all the elements of $\sigma_i(G)(j)$ from Q_i . By dealing with the resulting sets finitely many times (with the same method) we can find³ some $C_i \subset Q_i$ of the maximum cardinality satisfying that the mapping

$$G \times C_i \rightarrow [d_i], \quad (g, j) \mapsto \sigma_i(g)(j)$$

is injective. We take $\tau > 0$ with $\tau \cdot \dim(X) < \theta/2$. Since G is finite, there exists some $i_0 \in \mathbb{N}$ such that

$$|\sigma_i(G)(C_i)| > (1 - \tau) \cdot d_i, \quad \forall i \geq i_0.$$

³We notice that the situation here is very simple because G is a finite group. In a much more complicated case, i.e. where the acting group is amenable but not necessarily finite, it is also possible to have such a one-to-one correspondence. But we do not need this in our proof. For the general statement we refer to [Li13, Section 3].

We take $\delta_0 > 0$ satisfying that $\delta_0 \cdot |G| \cdot (\dim(X) + 1) < \theta/2$ and that if two points $x, x' \in X$ satisfy $\rho(x, x') \leq 2\sqrt{\delta_0}$ then they satisfy $\max_{g \in G} \rho(gx, gx') < \epsilon$.

We fix $i \in \mathbb{N}$ for the moment. For each $j \in [d_i]$ we define a *continuous* mapping $J_j : \text{Map}(\rho, G, \delta_0, \sigma_i) \rightarrow [0, 1]$ by sending $\phi = (\phi_j)_{j \in [d_i]} \in \text{Map}(\rho, G, \delta_0, \sigma_i)$ to

$$J_j(\phi) = \max \left\{ \max_{g \in G} \left(\rho(g\phi_j, \phi_{\sigma_i(g)(j)}) - \sqrt{\delta_0} \right), 0 \right\}.$$

Now we define a *continuous* mapping

$$H_i : \text{Map}(\rho, G, \delta_0, \sigma_i) \rightarrow X^{C_i} \times (CX)^{d_i} \times X^{[d_i] \setminus \sigma_i(G)(C_i)},$$

$$\phi = (\phi_j)_{j \in [d_i]} \mapsto ((\phi_j)_{j \in C_i}, (J_j(\phi)\phi_j)_{j \in [d_i]}, (\phi_j)_{j \in [d_i] \setminus \sigma_i(G)(C_i)}).$$

We recall here that $CX = ([0, 1] \times X) / \sim$ denotes the cone generated by X , where $(0, x) \sim (0, x')$ for any $x, x' \in X$, and that λx is the equivalence class of $(\lambda, x) \in [0, 1] \times X$ (in particular, the symbol $* = 0x$ is to denote the vertex of CX).

We claim that H_i is an ϵ -embedding with respect to ρ_∞ . To verify this, we suppose that $H_i(\varphi) = H_i(\psi)$ for $\varphi, \psi \in \text{Map}(\rho, G, \delta_0, \sigma_i)$. We need to show $\rho(\varphi_j, \psi_j) < \epsilon$ for all $j \in [d_i]$. If $j \in [d_i]$ satisfies $j \in [d_i] \setminus \sigma_i(G)(C_i)$ or $J_j(\varphi) = J_j(\psi) > 0$, then it is clear that $\varphi_j = \psi_j$. So we assume $j \in \sigma_i(G)(C_i)$ and $J_j(\varphi) = J_j(\psi) = 0$. We write $j = \sigma_i(s)(c)$ for some $s \in G$ and $c \in C_i \subset Q_i$. Since $\rho(s^{-1}\varphi_j, \varphi_{\sigma_i(s^{-1})(j)}) \leq \sqrt{\delta_0}$, $\rho(s^{-1}\psi_j, \psi_{\sigma_i(s^{-1})(j)}) \leq \sqrt{\delta_0}$ and since $\varphi_{\sigma_i(s^{-1})(j)} = \varphi_c = \psi_c = \psi_{\sigma_i(s^{-1})(j)}$, we have $\rho(s^{-1}\varphi_j, s^{-1}\psi_j) \leq 2\sqrt{\delta_0}$. Thus, $\rho(\varphi_j, \psi_j) < \epsilon$. This proves the claim.

It follows from this claim that $\text{Widim}_\epsilon(\text{Map}(\rho, G, \delta_0, \sigma_i), \rho_\infty)$ is bounded from above by $\dim(H_i(\text{Map}(\rho, G, \delta_0, \sigma_i)))$. To estimate the latter term, we consider, for any $\phi \in \text{Map}(\rho, G, \delta_0, \sigma_i)$, the set $P_i(\phi) = \{j \in [d_i] : J_j(\phi) > 0\}$. Since $\phi \in \text{Map}(\rho, G, \delta_0, \sigma_i)$, we have $\delta_0 \cdot |P_i(\phi)| \leq \sum_{g \in G, j \in [d_i]} \rho(g\phi_j, \phi_{\sigma_i(g)(j)})^2 \leq |G| \cdot \delta_0^2 \cdot d_i$ and hence we deduce that $|P_i(\phi)| \leq |G| \cdot \delta_0 \cdot d_i$. This implies that any element in $H_i(\text{Map}(\rho, G, \delta_0, \sigma_i))$ has entries at most $(|C_i| + |G| \cdot \delta_0 \cdot d_i + |[d_i] \setminus \sigma_i(G)(C_i)|)$, which do not take the value $*$. Thus, $\dim(H_i(\text{Map}(\rho, G, \delta_0, \sigma_i)))$ is bounded from above by

$$\dim(X^{|C_i|}) + \dim(CX) \cdot |G| \cdot \delta_0 \cdot d_i + \dim(X) \cdot |[d_i] \setminus \sigma_i(G)(C_i)|$$

which, for all $i \geq i_0$, does not exceed

$$\dim(X^{|C_i|}) + (\dim(X) + 1) \cdot |G| \cdot \delta_0 \cdot d_i + \dim(X) \cdot \tau \cdot d_i \leq \dim(X^{|C_i|}) + \theta \cdot d_i.$$

Finally, we note that $|C_i| \rightarrow +\infty$ (because $d_i \rightarrow +\infty$) as $i \rightarrow \infty$ and that $|G| \cdot |C_i| \leq d_i$ (for all $i \in \mathbb{N}$). Thus, we conclude with

$$\limsup_{i \rightarrow \infty} \frac{\dim(X^{|C_i|})}{d_i} \leq \frac{1}{|G|} \cdot \lim_{i \rightarrow \infty} \frac{\dim(X^{|C_i|})}{|C_i|} = \frac{1}{|G|} \cdot \inf_{n \in \mathbb{N}} \frac{\dim(X^n)}{n}.$$

□

Lemma 4.2 (Estimate from below).

$$\text{mdim}_\Sigma(X, G) \geq \frac{1}{|G|} \cdot \inf_{n \in \mathbb{N}} \frac{\dim(X^n)}{n}.$$

Proof. First of all, we note that Lemma 2.7 allows us to reduce the statement to the following issue:

$$\text{mdim}_\Sigma(X^{|G|}, G) \geq \inf_{n \in \mathbb{N}} \frac{\dim(X^n)}{n}.$$

We fix a compatible metric $D = \rho_\infty$ on the product space $X^{|G|}$. We would like to remind the reader to keep in mind that the remaining part of this proof is concerned only with the product action $(X^{|G|}, G)$.

To show the reduced statement it suffices to prove

$$\text{mdim}_\Sigma(X^{|G|}, G) \geq \dim(X) - 1.$$

Indeed, because this inequality will apply to all those actions $(X^{|G|}, G)$, in particular, to $((X^n)^{|G|}, G)$ which is (dynamically) isomorphic to the product action $((X^{|G|})^n, G)$, for any $n \in \mathbb{N}$, we shall deduce from Lemma 2.7 that

$$\text{mdim}_\Sigma(X^{|G|}, G) \geq \frac{\text{mdim}_\Sigma((X^{|G|})^n, G)}{n} = \frac{\text{mdim}_\Sigma((X^n)^{|G|}, G)}{n} \geq \frac{\dim(X^n) - 1}{n}.$$

Since $n \in \mathbb{N}$ is arbitrary, this will imply that

$$\text{mdim}_\Sigma(X^{|G|}, G) \geq \lim_{n \rightarrow \infty} \frac{\dim(X^n) - 1}{n} = \inf_{n \in \mathbb{N}} \frac{\dim(X^n)}{n}.$$

This is as desired.

Next we prove

$$\text{mdim}_\Sigma(X^{|G|}, G) \geq \dim(X) - 1.$$

By Theorem 2.5 there exists some $\epsilon > 0$ (sufficiently small) which depends only on the finite dimensional compact metric space (X, ρ) , such that

$$\text{Widim}_\epsilon(X^n, \rho_\infty) \geq n \cdot (\dim(X) - 1), \quad \forall n \in \mathbb{N}.$$

We take $\delta > 0$ arbitrarily. We fix ϵ and δ in the proof. By definition we need to show that

$$\limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(\text{Map}(D, G, \delta, \sigma_i), D_\infty)}{d_i} \geq \dim(X) - 1.$$

For every $i \in \mathbb{N}$ we let $Q_i \subset [d_i]$ be the intersection of the following two sets:

$$\{j \in [d_i] : \sigma_i(g)(j) \neq \sigma_i(h)(j), \forall g \neq h \in G\},$$

$$\{j \in [d_i] : \sigma_i(g)(\sigma_i(h)(j)) = \sigma_i(gh)(j), \forall g, h \in G\}.$$

For any $j \in [d_i]$ and $A \subset [d_i]$ we set

$$\sigma_i(G)(j) = \{\sigma_i(g)(j) : g \in G\} \subset [d_i], \quad \sigma_i(G)(A) = \{\sigma_i(g)(a) : g \in G, a \in A\} \subset [d_i].$$

We note that by the definition of Q_i we have

$$|\sigma_i(G)(j)| = |G|, \quad \sigma_i(G)(\sigma_i(G)(j)) = \sigma_i(G)(j), \quad \forall j \in Q_i.$$

It follows that $\sigma_i(G)(j) = \sigma_i(G)(l)$, for all $l \in \sigma_i(G)(j)$. As what we have explained in the proof of Lemma 4.1, we can find some $C_i \subset Q_i$ satisfying that the mapping

$$G \times C_i \rightarrow [d_i], \quad (g, j) \mapsto \sigma_i(g)(j)$$

is injective. Since G is finite, there is some $i_0 \in \mathbb{N}$ (sufficiently large) such that

$$|\sigma_i(G)(C_i)| > (1 - \delta^2) \cdot d_i, \quad \forall i \geq i_0.$$

For every $i \in \mathbb{N}$ we define a mapping P_i as follows:

$$P_i : X^{d_i} \rightarrow (X^{|G|})^{d_i}, \quad p = (p_j)_{j \in [d_i]} \mapsto \phi = (\phi_j)_{j \in [d_i]} = ((\phi_j(g))_{g \in G})_{j \in [d_i]},$$

$$\phi_j(g) = \begin{cases} hg^{-1}p_{\sigma_i(g)(k)}, & \text{if } j = \sigma_i(h)(k), h \in G, k \in C_i \\ p_j, & \text{if } j \in [d_i] \setminus \sigma_i(G)(C_i) \end{cases}.$$

The mapping $P_i : X^{d_i} \rightarrow (X^{|G|})^{d_i}$ is well defined because the mapping

$$G \times C_i \rightarrow [d_i], \quad (g, j) \mapsto \sigma_i(g)(j)$$

is injective. Clearly, $P_i : X^{d_i} \rightarrow (X^{|G|})^{d_i}$ is continuous. Since for any $j \in [d_i]$ there is some $g \in G$ such that $\phi_j(g) = p_j$, it is distance-increasing with respect to the metric ρ_∞ on X^{d_i} and the metric D_∞ on $(X^{|G|})^{d_i}$, i.e.

$$\rho_\infty(p, p') \leq D_\infty(P_i(p), P_i(p')), \quad \forall p, p' \in X^{d_i}.$$

We claim that for all sufficiently large $i \in \mathbb{N}$ it will be true that $P_i(X^{d_i})$ is contained in $\text{Map}(D, G, \delta, \sigma_i)$. We will verify the claim in a moment. By this claim and by the choice of $\epsilon > 0$ we will obtain that

$$\begin{aligned} \limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(\text{Map}(D, G, \delta, \sigma_i), D_\infty)}{d_i} &\geq \limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(P_i(X^{d_i}), D_\infty)}{d_i} \\ &\geq \limsup_{i \rightarrow \infty} \frac{\text{Widim}_\epsilon(X^{d_i}, \rho_\infty)}{d_i} \\ &\geq \limsup_{i \rightarrow \infty} \frac{d_i \cdot (\dim(X) - 1)}{d_i} \\ &= \dim(X) - 1. \end{aligned}$$

This will end the proof.

Now we prove that $P_i(X^{d_i})$ is contained in $\text{Map}(D, G, \delta, \sigma_i)$ for all integers $i \geq i_0$. We fix an integer $i \geq i_0$. We take an arbitrary $p = (p_j)_{j \in [d_i]} \in X^{d_i}$. Let $\phi = P_i(p)$ and write $\phi = (\phi_j)_{j \in [d_i]} = ((\phi_j(g))_{g \in G})_{j \in [d_i]} \in (X^{|G|})^{d_i}$. We note that $C_i \subset Q_i$. By the construction

of the mapping $P_i : X^{d_i} \rightarrow (X^{|G|})^{d_i}$ we have obviously that if $j = \sigma_i(h)(c)$ for some $h \in G$ and $c \in C_i$ then for any $s \in G$ and any $g \in G$

$$s\phi_j(g) = shg^{-1}p_{\sigma_i(g)(c)} = \phi_{\sigma_i(sh)(c)}(g) = \phi_{\sigma_i(s)(\sigma_i(h)(c))}(g) = \phi_{\sigma_i(s)(j)}(g).$$

Thus, for any $j \in \sigma_i(G)(C_i)$

$$s\phi_j = \phi_{\sigma_i(s)(j)}, \quad \forall s \in G.$$

It follows that for all $s \in G$

$$\begin{aligned} D_2(s\phi, \phi \circ \sigma_i(s)) &= \sqrt{\frac{1}{d_i} \cdot \sum_{j \in [d_i]} (D(\phi_{\sigma_i(s)(j)}, s\phi_j))^2} \\ &= \sqrt{\frac{1}{d_i} \cdot \sum_{j \in [d_i] \setminus \sigma_i(G)(C_i)} (D(\phi_{\sigma_i(s)(j)}, s\phi_j))^2} \\ &\leq \sqrt{\frac{d_i - |\sigma_i(G)(C_i)|}{d_i}} \\ &\leq \delta. \end{aligned}$$

This implies that $\phi \in \text{Map}(D, G, \delta, \sigma_i)$. Thus, we conclude. \square

APPENDIX

The definition of *sofic mean dimension* was originally introduced by Li [Li13] (using open covers). However, we notice that the definition which we stated in the body of this paper (in terms of Widim) differs from Li's. The appendix is devoted to clarifying that these two definitions are equivalent for sofic mean dimension.

Let Y be a compact space and \mathcal{U} a finite open cover of Y . We denote

$$\text{ord}(\mathcal{U}) = \max_{y \in Y} \sum_{U \in \mathcal{U}} 1_U(y) - 1, \quad \mathcal{D}(\mathcal{U}) = \min_{\mathcal{V} \succ \mathcal{U}} \text{ord}(\mathcal{V}),$$

where $1_U(y)$ takes the value 1 (resp. 0) if $y \in U$ (resp. $y \notin U$) and where $\mathcal{V} \succ \mathcal{U}$ means that \mathcal{V} ranges over finite open covers of Y refining \mathcal{U} (i.e. every element of \mathcal{V} is contained in some element of \mathcal{U}). For $d \in \mathbb{N}$ we denote by \mathcal{U}^d the finite open cover of Y^d consisting of all those open sets in Y^d of the form $U_1 \times \cdots \times U_d$ (where $U_1, \dots, U_d \in \mathcal{U}$).

Let a sofic group G act continuously on a compact metrizable space X and let $\Sigma = \{\sigma_i : G \rightarrow \text{Sym}(d_i)\}_{i \in \mathbb{N}}$ be a sofic approximation sequence for G . We fix a compatible metric ρ on X . For a finite open cover \mathcal{U} of X , a nonempty finite subset F of G , $\delta > 0$, and $i \in \mathbb{N}$ we denote $\mathcal{D}(\mathcal{U}^{d_i} \cap \text{Map}(\rho, F, \delta, \sigma_i))$ by $\mathcal{D}(\mathcal{U}, \rho, F, \delta, \sigma_i)$. Recall that we have already given the definition of $\text{mdim}_\Sigma(X, G)$ in Section 2. Now what we want to see is

$$\text{mdim}_\Sigma(X, G) = \sup_{\mathcal{U}} \inf_{F \subset G \text{ finite}, \delta > 0} \limsup_{i \rightarrow \infty} \frac{\mathcal{D}(\mathcal{U}, \rho, F, \delta, \sigma_i)}{d_i},$$

where \mathcal{U} runs over finite open covers of X .

In fact, this can be shown with a simple and standard argument. The key point is to consider a Lebesgue number of a finite open cover \mathcal{U} with respect to ρ , and to use the following fact [LW00]: For a finite open cover \mathcal{U} of a compact metrizable space X , $\mathcal{D}(\mathcal{U}) \leq n$ (where n is a nonnegative integer) exactly when there is a compact metrizable space P of topological dimension n , which admits a continuous mapping $f : X \rightarrow P$ satisfying that there exists a finite open cover \mathcal{V} of P such that $f^{-1}(\mathcal{V})$ refines \mathcal{U} . Lastly, we note that although a compatible metric ρ on the space X is also involved in the right-hand side of the above equality, it is independent of the choice of a compatible metric ρ on X (for details see [Li13, Section 2]).

REFERENCES

- [Dra01] A. N. Dranishnikov. Cohomological dimension theory of compact metric spaces. *Topology Atlas Invited Contributions 6* (2001), 61 pp. <http://at.yorku.ca/t/a/i/c/43.htm>
- [Gro99a] Misha Gromov. Endomorphisms of symbolic algebraic varieties. *Journal of the European Mathematical Society* 1 (1999), 109–197.
- [Gro99b] Misha Gromov. Topological invariants of dynamical systems and spaces of holomorphic maps: I. *Math. Phys. Anal. Geom.* 2 (1999), 323–415.
- [JQ] Lei Jin, Yixiao Qiao. Mean dimension of product spaces: a fundamental formula. Submitted. Preprint on [arXiv:2102.10358](https://arxiv.org/abs/2102.10358).
- [Li13] Hanfeng Li. Sofic mean dimension. *Advances in Mathematics* 244 (2013), 570–604.
- [LW00] Elon Lindenstrauss, Benjamin Weiss. Mean topological dimension. *Israel Journal of Mathematics* 115 (2000), 1–24.
- [Tsu19] Masaki Tsukamoto. Mean dimension of full shifts. *Israel Journal of Mathematics* 230 (2019), 183–193.

LEI JIN: CENTER FOR MATHEMATICAL MODELING, UNIVERSITY OF CHILE AND UMI 2807 - CNRS
Email address: jinleim@impan.pl

YIXIAO QIAO (CORRESPONDING AUTHOR): SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA
 NORMAL UNIVERSITY, GUANGZHOU, GUANGDONG 510631, CHINA
Email address: yxqiao@mail.ustc.edu.cn